
LY6H [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000176956 | Nominal screening candidate



Conditional PTN–LY6H association: b = -0.121 per mediator SD, p = 0.000137, BH FDR = 0.00472 | Dx attenuation = 10.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Direct manuscript connection: neuronal SCZ expression and spatial eQTL/colocalization in Results 2.7 and Figure S42. This evidence concerns LY6H and does not genetically anchor PTN mediation.

TIMP2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000035862 | Nominal screening candidate



Conditional PTN–TIMP2 association: b = -0.070 per mediator SD, p = 0.00816, BH FDR = 0.0332 | Dx attenuation = 13.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Matrix-metalloproteinase regulation links this neuropil outcome to the manuscript ECM/PNN theme. This is an ECM hypothesis, not a PNN-target screen or a demonstrated PTN receptor interaction.

IGSF9B [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000080854 | Nominal screening candidate



Conditional PTN–IGSF9B association: b = -0.091 per mediator SD, p = 0.0283, BH FDR = 0.0851 | Dx attenuation = 12.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Synaptic adhesion and inhibitory-synapse biology fit the manuscript synaptic/GABAergic theme. Adult mouse localization is not restricted to inhibitory synapses.

GAP43 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000172020 | Nominal screening candidate



Conditional FGF1–GAP43 association: b = -0.084 per mediator SD, p = 0.00571, BH FDR = 0.0668 | Dx attenuation = 13.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Named in Results 2.4 for neuronal structure/outgrowth among PRS-associated genes. Its occurrence in both FGF neuropil-to-vascular screens merits comparison; a vascular SPG label does not establish endothelial origin.

GAP43 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000172020 | Nominal screening candidate



Conditional FGF2–GAP43 association: b = -0.099 per mediator SD, p = 4.3e-05, BH FDR = 0.000712 | Dx attenuation = 18.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Named in Results 2.4 for neuronal structure/outgrowth among PRS-associated genes. Its occurrence in both FGF neuropil-to-vascular screens merits comparison; a vascular SPG label does not establish endothelial origin.

APOLD1 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000178878 | Nominal screening candidate



Conditional FGF1–APOLD1 association: b = +0.231 per mediator SD, p = 0.0076, BH FDR = 0.143 | Dx attenuation = 10.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Endothelial junction and permeability biology provides a vascular-context example in the neuronal FGF1 screen. Functional literature does not establish an FGF1-to-APOLD1 mechanism here.

CNNM2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000148842 | Nominal screening candidate



Conditional PTN–CNNM2 association: b = -0.081 per mediator SD, p = 0.0176, BH FDR = 0.0598 | Dx attenuation = 12.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: External human and experimental evidence connects CNNM2 to schizophrenia-related cortical and synaptic phenotypes. This is a literature-based nomination, not a locus-specific finding from this mediation screen.

RGS4 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000117152 | Nominal screening candidate



Conditional FGF1–RGS4 association: b = -0.095 per mediator SD, p = 0.000136, BH FDR = 0.0257 | Dx attenuation = 14.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: A cortical G-protein signaling candidate with prior SCZ expression studies, including conflicting reports. The vascular-target result is a context comparison, not confirmation of a neuronal cellular origin.

ADCYAP1R1 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000078549 | Nominal screening candidate



Conditional FGF1–ADCYAP1R1 association: b = +0.127 per mediator SD, p = 0.0177, BH FDR = 0.209 | Dx attenuation = 8.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: PACAP receptor signaling offers a neurovascular signaling example. ADCYAP1R1 is not an FGF receptor; this is an outcome association with neuronal FGF1.

MT3 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000087250 | Nominal screening candidate



Conditional FGF1–MT3 association: b = +0.182 per mediator SD, p = 2.91e-06, BH FDR = 0.000283 | Dx attenuation = 19.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Representative of MT1G/MT1M/MT3-driven metal-response GO terms in the all-tested-background analysis, matching the manuscript vascular metal-homeostasis theme. Include both FGF1 and FGF2 pairs for comparison.

MT3 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000087250 | Nominal screening candidate



Conditional FGF2–MT3 association: b = +0.229 per mediator SD, p = 5.9e-14, BH FDR = 7.15e-12 | Dx attenuation = 28.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Representative of MT1G/MT1M/MT3-driven metal-response GO terms in the all-tested-background analysis, matching the manuscript vascular metal-homeostasis theme. Include both FGF1 and FGF2 pairs for comparison.

AGT [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000135744 | Nominal screening candidate



Conditional FGF1–AGT association: b = +0.430 per mediator SD, p = 2.31e-06, BH FDR = 0.000256 | Dx attenuation = 20.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Shared FGF1/FGF2 vascular outcome contributing to the regulation-of-growth GO term. Useful for comparing two ligands with the same target outcome; shared composition remains an alternative explanation.

AGT [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000135744 | Nominal screening candidate



Conditional FGF2–AGT association: b = +0.662 per mediator SD, p = 1.26e-21, BH FDR = 6.35e-19 | Dx attenuation = 37.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Shared FGF1/FGF2 vascular outcome contributing to the regulation-of-growth GO term. Useful for comparing two ligands with the same target outcome; shared composition remains an alternative explanation.

FAM107A [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF1–FAM107A association: b = +0.212 per mediator SD, p = 3.09e-06, BH FDR = 0.000283 | Dx attenuation = 23.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Shared across all three FGF-to-vascular screens, allowing comparison across both ligand and source microenvironment. Selection is based on overlap, not evidence of signaling specificity.

FAM107A [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF2–FAM107A association: b = +0.256 per mediator SD, p = 8.04e-13, BH FDR = 8.74e-11 | Dx attenuation = 32.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Shared across all three FGF-to-vascular screens, allowing comparison across both ligand and source microenvironment. Selection is based on overlap, not evidence of signaling specificity.

FAM107A [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF1–FAM107A association: b = +0.144 per mediator SD, p = 5.93e-05, BH FDR = 0.0182 | Dx attenuation = 13.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.

Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Selection: Shared across all three FGF-to-vascular screens, allowing comparison across both ligand and source microenvironment. Selection is based on overlap, not evidence of signaling specificity.