


lecture 11:
**stable mass transfer &
evolution of massive binaries**

response of mass-accreting star

- mass transfer stream from donor star:

- **direct impact** if $d_{\min} < R_a$
- **accretion disk** if $d_{\min} > R_a$

minimum distance of stream to
accretor's centre of mass ($d_{\min} \propto a$)





response of mass-accreting star

- effects of mass accretion on companion (MS) star:
 - kinetic energy of accretion flow \Rightarrow **accretion luminosity**, dissipated locally (hot spot or accretion disk)
 - transfer of angular momentum \Rightarrow **spin-up** of mass gainer (accretion of $\sim 0.1 M_a$ may already lead to break-up rotation) \Rightarrow this may lead to *non-conservative mass transfer*

response of mass-accreting star

- effects of mass accretion on companion (MS) star:
 - kinetic energy of accretion flow \Rightarrow **accretion luminosity**, dissipated locally (hot spot or accretion disk)
 - transfer of angular momentum \Rightarrow **spin-up** of mass gainer (accretion of $\sim 0.1 M_a$ may already lead to break-up rotation) \Rightarrow this may lead to *non-conservative mass transfer*
- further effects, to be illustrated next:
 - **thermal readjustment** of mass gainer \Rightarrow **over-luminosity** and **expansion** (when accretion occurs faster than gainer's thermal timescale) \Rightarrow *contact binary?*
 - if substantial accretion occurs: **rejuvenation** of mass gainer (due to larger convective core)
 - but also: higher mass causes increased rate of nuclear evolution \Rightarrow secondary can **overtake evolution** of primary \Rightarrow *contact binary*


case A: slow evolution into contact



example of *secondary overtaking primary*

- B – C: thermal-timescale mass transfer
- C – D: nuclear-timescale mass transfer \Rightarrow Algol-type binaries
- beyond D: both stars overfill RL \Rightarrow long-lived **contact binaries**

case A: rapid evolution into contact




example of *secondary accreting too fast to thermally readjust*

- B – C: thermal-timescale mass transfer, but $\tau_{\text{KH,acc}} \gg \tau_{\text{KH,don}} \sim \tau_{\text{mass-transfer}}$
- beyond C: both stars overflow RL during rapid mass transfer
 \Rightarrow outcome uncertain, likely **binary merger** \Rightarrow single star?



evolution into contact

- binary parameters for which contact occurs (example: $M_{1,i} = 12 M_{\text{sun}}$)





(*) if *spin-up of mass gainer* (see later) does not prevent accretion...

case A versus case B




- **case A:** rapid + slow phase of mass transfer during donor's MS phase
- mass transfer resumes after MS (“case AB”): remainder of donor's H-envelope transferred, remnant is an almost naked **helium star** (as in case B)

case A versus case B



- **case B:** mass transfer starts after MS phase *1
 - shorter thermal timescale \Rightarrow rapid mass transfer with higher dM/dt
 - rapid expansion during Hertzsprung gap \Rightarrow no long-lived, slow phase of mass transfer



case A versus case B



- **case B:** mass transfer starts after MS phase *1
 - shorter thermal timescale \Rightarrow rapid mass transfer with higher dM/dt
 - rapid expansion during Hertzsprung gap \Rightarrow no long-lived, slow phase of mass transfer


case A versus case B

case B evolution example



- **orbit expands** during (conservative) case B mass transfer (B–D)
 in this example (B–D): as $M_1 = 10 M_{\text{sun}} \rightarrow 2.5 M_{\text{sun}} \Rightarrow P = 12 \text{ d} \rightarrow 105 \text{ d}$

case B evolution example



- B – D: thermal timescale (rapid) mass transfer \Rightarrow few obs. counterparts
- D – E: detached, **He-burning** *1 (nuclear timescale)
 \Rightarrow expect many observed counterparts
- E: end He-burning *1, further RLOF phase possible


post-mass transfer binaries

- *1 stripped down to its (almost) bare He core:
 - a **He star** in long-lived He-burning state
 - if massive enough: He star appears as a **Wolf-Rayet star**
- *2 is still a **MS star** (O- or B-type)
- when $M_1 \ll M_2$ and large $P_{\text{orb}} \Rightarrow$ sign of **conservative** mass transfer (e.g. ϕ Per)
- however, many WR+O binaries with short P_{orb} and $M_1 \sim M_2 \Rightarrow$ must have lost substantial mass and AM

name	spectra	P (d)	M_1	M_2	R_1	R_2
CQ Cep	WN7 + O6	1.64	24	30	8.8	7.9
V398 Car	WN4 + O4-6	8.26	19	37		
GP Cep	WN6/WC + O3-6	6.69	15	27		
CV Ser	WC8 + O8-9	29.7	13	27		
V444 Cyg	WN5 + O6	4.21	9.3	28	2.9	8.5
→ ϕ Per	HeI em + B1IIIe	127	1.15	9.3	1.3:	5.5–8

post-mass transfer binaries



- in binaries like ϕ Per, the more massive **mass gainer outshines the He star**, which is hard to detect
 \Rightarrow mass gainers in massive binaries often **appear to be single stars**



Götberg+ 2022


post-mass transfer binaries

- in binaries like ϕ Per, the more massive **mass gainer outshines the He star**, which is hard to detect
 \Rightarrow mass gainers in massive binaries often **appear to be single stars**
- recently more such companions have been discovered using far-UV observations:



post-mass transfer binaries

- in binaries like ϕ Per, the more massive **mass gainer outshines the He star**, which is hard to detect
 \Rightarrow mass gainers in massive binaries often **appear to be single stars**




Götberg+ 2022

effects on the mass gainer:

- as a result of mass transfer, the mass gaining star is expected to:
 - **rotate** very rapidly
 - have modified **surface abundances**
- (qualitatively) similar effects expected for MS binary **mergers**

effects on the mass gainer

- evolution of **rotation velocities** of massive binaries:




case B

case A

de Mink+ 2013



effects on the mass gainer

- predicted distribution of rotation velocities:



effects on the mass gainer


- a (likely) manifestation of spin-up by accretion or merger: **Be stars**
 - ~20% of B-type stars ($3 - 20 M_{\text{sun}}$), rapidly rotating
 - equatorial mass loss into circumstellar disk (transient phenomenon)
- some Be stars have **stripped He-star companions** (e.g. ϕ Per) or **compact companions** (Be/X-ray binaries)
- **no** Be stars have normal B-star companions



credit: J. Bodensteiner

effects on the mass gainer

- predicted **nitrogen abundances** after mass accretion ($M_i = 16 + 12 M_{\text{sun}}$)



Wellstein+ 2001, Langer 2012

- mass transfer results in similar properties (N enrichment + rapid rotation) as expected from **rotational mixing in single stars**
- effects are very hard to disentangle...

effects of SN explosion

- next stage in the evolution of a massive binary: **core collapse** of the stripped **primary star** (if $M_{\text{He}} > \sim 2 M_{\text{sun}}$)
- resulting supernova explosion leads to **sudden mass loss**, and a possible “**kick**” imparted to the remnant (NS or BH)
⇒ both effects **change the binary orbit** (and possibly disrupt the binary)

effects of SN explosion

- next stage in the evolution of a massive binary: **core collapse** of the stripped **primary star** (if $M_{\text{He}} > \sim 2 M_{\text{sun}}$)
 - resulting supernova explosion leads to **sudden mass loss**, and a possible “**kick**” imparted to the remnant (NS or BH)
 - ⇒ both effects **change the binary orbit** (and possibly disrupt the binary)
 - resulting systems may be observed as
 - **X-ray binaries**: NS or BH accreting gas from a normal (MS, RG) companion star
 - **binary radio pulsars**: non-accreting NS with a stellar companion
 - normal stars with an unseen, very massive (non-accreting BH) companion
- ... and some of these eventually evolve into double NS, double BH or NS-BH binaries (GW sources)


X-ray binaries

- **high-mass X-ray binaries:**

OB-star + NS/BH

- often: **X-ray pulsations** \Rightarrow rotating, magnetized NS (allows measuring orbital RV variations of NS \Rightarrow mass function of OB star)
- OB-companion easily detectable
- eclipses of X-source + RV variations of OB star \Rightarrow measure mass of compact object

HMXB




- **low-mass X-ray binaries:**

G/K-star + NS/BH

- no X-ray pulsations, but often **X-ray bursts** (thermonuclear flashes of accreted gas on NS surface)
- companion often not observed (L is dominated by reprocessed X-rays in accretion disk), unless X-ray emission is **transient**

LMXB



X-ray binaries

two main sub-groups of HMXBs:

- **'standard' HMXBs:**



O-star + NS/BH, $P < \sim 10$ d, $e < \sim 0.1$

- O-star ($> \sim 20 M_{\text{sun}}$) often close to filling Roche lobe
- persistent X-ray emission
- accretion from stellar wind or beginning RLOF

- **Be/X-ray binaries:**

Be-star + NS, $P \sim 15\text{--}300$ d, $e < \sim 0.5$

- weak stellar winds, far from RLOF
- accretion from equatorial mass ejections of Be star ($\sim 8\text{--}20 M_{\text{sun}}$)
- transient X-ray emission (known population is incomplete)



mass loss in supernova

- special case: **symmetric explosion** (instantaneous, isotropic mass loss ΔM) in **circular** orbit \Rightarrow produces a wider, eccentric orbit with

$$e = \Delta M / M_{\text{tot},f}$$

$$a_f = a_i / (1 - e)$$

- binary is disrupted if $\Delta M > 0.5 M_{\text{tot},i}$

mass loss in supernova

mass loss in supernova

- special case: **symmetric explosion** (instantaneous, isotropic mass loss ΔM) in **circular** orbit \Rightarrow produces a wider, eccentric orbit with

$$e = \Delta M / M_{\text{tot},f}$$


$$a_f = a_i / (1 - e)$$

- binary is disrupted if $\Delta M > 0.5 M_{\text{tot},i}$
- general case:
 - initial orbit may be **eccentric** (if wide enough to avoid strong tides)
 - SN explosion likely to be slightly **asymmetric** \Rightarrow compact remnant receives a **kick** (recoil) velocity, up to few 100 km/s
- both introduce a **random effect**:
 - explosion may occur at any point on eccentric orbit
 - kick velocity can have an arbitrary direction
 - result is unpredictable, can only compute probabilities

mass loss in supernova

effect of supernova explosion

- probability that binary remains bound after SN explosion
- example: $M_1 = 10 M_{\text{sun}} \Rightarrow \text{SN} \Rightarrow M_{1f} = 1.35 M_{\text{sun}} \text{ (NS)}$
 - symmetric explosion in circular orbit: bound orbit if $M_2 > 7.3 M_{\text{sun}}$
 - **eccentric orbit** and/or **SN kick**: survival probability depends on M_2




finite probability of survival even for small M_2 , if explosion occurs near apastron

effect of supernova explosion

- probability that binary remains bound after SN explosion
- example: $M_1 = 10 M_{\text{sun}} \Rightarrow \text{SN} \Rightarrow M_{1f} = 1.35 M_{\text{sun}} \text{ (NS)}$
 - symmetric explosion in circular orbit: bound orbit if $M_2 > 7.3 M_{\text{sun}}$
 - **eccentric orbit** and/or **SN kick**: survival probability depends on M_2

SN kick in circular orbit:







- larger effect in wider orbits
- if $v_k \sim v_{\text{orb}}$, even a very small M_2 can remain bound if kick is opposite to orbital motion



formation scenario of HMXB

example:

conservative case B evolution

Age (Myr)	$M_1(M_\odot)$		$M_2(M_\odot)$	$P_{\text{orb}}(\text{d})$	
0.00	12.00		8.00	10.0	• MS + MS: detached binary
17.92	11.80		8.00	10.3	• start of RLOF
17.93	9.90		9.90	9.2	• minimum P_{orb}
17.96	2.70		17.10	71.0	• He-star + MS (Be star) • ϕ Per-like binary
		SN ↓			
21.01	1.35		17.00	83.6 $e=0.06$	• NS + MS (Be star) • Be/X-ray binary
				in case of symmetric SN	
26.21	1.35		16.60	87.0	• start of RLOF from *2 (unstable)

formation and evolution of a HMXB

another possible scenario, starting from a wider orbit:

