

Table 5. Triangulated evidence across the two top-25 candidate lists

Evidence source	GREmLN	GENIE3	Interpretation
CRISPRi perturbational evidence	3 candidates with a detected response among 20 represented	6 candidates with a detected response among 19 evaluable after QC	Broader perturbational evidence for GENIE3 in the related CD4 T cell screen
Audited literature evidence	7/25 strong or moderate	9/25 strong or moderate	Modest literature lean towards GENIE3; tiers remain provisional
TCR activation mRNA context	2/25	5/25	Activation-associated context was more frequent for GENIE3; no candidate showed a qualifying incremental BTLA-specific mRNA response
Independent BTLA molecular association	4/25	3/25	Slight numerical lean towards GREmLN, but protein and phosphosite coverage was sparse and unequal
Overall	Distinct candidates and molecular hypotheses	Broader perturbational and literature corroboration	No clear or consistent superiority across evidence sources

Descriptive comparison of evidence coverage across each model's top-25; no weighted score, Spearman correlation, bootstrap interval or superiority margin is implied.

CRISPRi: GREmLN denominator is candidates represented in the screen; GENIE3 denominator is candidates evaluable after on-target QC.

Literature tiers are audited provisional Paperclip v2 annotations pending documented human verification.

Candidate mRNA is contextual; independent molecular association is restricted to qualifying protein or phosphosite measurements. Assay coverage is incomplete and unequal across candidates.