

PHAS1202 – Astrophysics ICA : Solutions – Dec. 2016

1. (Students do *not* use the Stefan-Boltzmann constant)

$$\text{Luminosity, } L \propto R_*^2 T_{\text{eff}}^4 \quad [1]$$

$$\frac{L_{wd}}{L_{\odot}} = \frac{R_{wd}^2}{R_{\odot}^2} \times \frac{70000^4}{5800^4}$$

$$\frac{R_{wd}^2}{R_{\odot}^2} = 5 \times 4.713 \times 10^{-5} \quad [2]$$

$$R_{wd} = 0.015 R_{\odot} \quad [1]$$

$$\text{Wien's law (black-body approximation) gives: } \lambda_{\text{max}} = \frac{3 \times 10^{-3}}{T} \quad [1]$$

where λ_{max} is in m and T is in K.

$$\lambda_{\text{max}} = \frac{3 \times 10^{-3}}{70000} = 4.3 \times 10^{-8} \text{m} \quad [1]$$

$\sim 43\text{nm}$ corresponds to the ultraviolet waveband . [1]

[7 marks total]

2.

Sirius A luminosity is then

$$L = 23.5 \times 3.8 \times 10^{26} \text{ Watt} = 8.9 \times 10^{27} \text{ Watt}$$

Since fraction of mass liberated into energy by the fusion of H to He = 0.007 (= efficiency), $E_{\text{total}} = 0.007 mc^2$

The H mass burnt into He per sec will be

$$\frac{L}{\text{efficiency} \times c^2} = \frac{8.9 \times 10^{27}}{0.007 \times (3 \times 10^8)^2} = 1.4 \times 10^{13} \text{ kg s}^{-1} \quad [2]$$

If we assume that 10% of its H mass burns into He, then, since Sirius A mass is

$$M = 2.3 \text{ solar mass} = 2.3 \times 2 \times 10^{30} \text{ kg} = 4.6 \times 10^{30} \text{ kg}$$

Sirius A lifetime will be

$$\frac{0.1 \times M}{\text{Rate of burning}} = \frac{4.6 \times 10^{29}}{1.4 \times 10^{13}} \text{ s} = 3.3 \times 10^{16} \text{ s} = \frac{3 \times 10^{16}}{3 \times 10^7} \text{ years} = 1 \times 10^9 \text{ years}$$

[2]

while the Sun's lifetime is expected to be 10^{10} years. [1]

[7 marks total]

3. If the ionisation potential of the hydrogen atom is 13.6 eV, then $E(1) = -13.6 \text{ eV}$ and [1]

$$E(n) = -\frac{13.6}{n^2} \text{ eV}$$

Thus $E(2) - E(3) = -13.6 \left(\frac{1}{2^2} - \frac{1}{3^2} \right) \text{ eV} = -1.89 \text{ eV}$ and

[1]

$$\lambda = \frac{c}{\nu} = \frac{hc}{E} = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{1.89 \times 1.6 \times 10^{-19}} \text{ m} = 6.577 \times 10^{-7} \text{ m} = 657.7 \text{ nm}$$

[2]

Transitions whose lower level has $n = 2$ are called the *Balmer series*.

[1]

The transition between $n = 2$ and $n = 3$ is called $\text{H}\alpha$.

[1]

[6 marks total]

4.

Population I stars contain

- (i) a larger metal abundance, (ii) are distributed closer to the galactic disc, and
- (iii) are younger than Population II stars.

[3]

The Sun is a Population I star; Globular clusters are Population II

[1]

[4 marks total]

5

Velocity = distance/time = circumference/period:

$$\begin{aligned} T(\text{orbit}) &= \frac{2\pi 8000 \text{ pc} \times 3.1 \times 10^{13} \text{ km pc}^{-1}}{220 \text{ km s}^{-1} \times 3.16 \times 10^7 \text{ s yr}^{-1}} \\ &= 2.24 \times 10^8 \text{ yr} \end{aligned}$$

[2]

Approx. age of the solar system is $\sim 4.5 \times 10^9 \text{ yr}$
(Students expected to e.g. know the age of the Sun.)

[1]

Thus $4.5 \times 10^9 / 2.24 \times 10^8 \sim 20$ orbits completed.

[1]

The spiral arms cannot be fixed, permanent features, because they would quickly tighten ('wind up') after 20 revolutions. They instead represent a density wave phenomenon.

[2]

[6 marks total]

6. According to Hubble classification scheme:

SBa galaxy -- galaxy has a large nucleus, with a bar-like structure through it. The spiral arms emerge from the ends of the bar and are tightly wound. [2]

Sc galaxy -- galaxy has a relatively small nucleus, plus loosely wound spiral arms (no central bar). [2]

E6 galaxy -- elliptical galaxy, whose apparent shape is very elongated or flattened. [1]

[5 marks total]

7.

Distance to galaxy:
radial velocity

$$v = \left(\frac{(\lambda - \lambda_0)}{\lambda_0} \right) c$$

[1]

$$V = [(402.8 - 393.30)/393.30] \times 3 \times 10^5 = 7246 \text{ km s}^{-1} \quad [2]$$

$$\text{Hubble law, distance } d = v / H_0 = 7246 / 75 = 96.6 \text{ Mpc} \quad [2]$$

[5 marks total]