

Do you like Texas hold 'em

STAT230 Real World Assignment 2024W


Eason Li, Johnson Ji



Faculty of Math
University of Waterloo
Canada
April 8, 2024

In this report, we consider the case a player uses the best five-card poker hand out of seven cards.

For a 7-card hand to contain a Royal Flush, i.e.



it must contain the specific set of 5 cards (Ace, King, Queen, Jack, 10 of the same suit), with the other 2 cards being any of the remaining 47 cards in the deck. Therefore, the probability of getting a Royal Flush in a group of seven cards can be evaluated as

$$P(\text{Royal Flush in 7-card poker}) = \frac{4 \times C(47, 2)}{C(52, 7)}$$

With the similar idea, we can get discover the following probabilities:

Table 1: Probabilities of Being Dealt Specific Hands in Texas Hold'em Poker (7 cards)

Hand Rank	Absolute Frequency
Royal Flush	$\binom{4}{1} \binom{47}{2}$
Straight Flush (non-Royal)	$\binom{9}{1} \binom{4}{1} \binom{46}{2}$
Four of a Kind	$\binom{13}{1} \binom{48}{3}$
Full House	0.1441
Flush	0.197
Straight	0.392
Three of a Kind	2.112
Two Pair	4.753
One Pair	42.256
High Card	50.117