

Cover Page for Protocol

| | |
|--------------------------|--|
| Sponsor name: | Novo Nordisk A/S |
| NCT number | NCT03548935 |
| Sponsor trial ID: | NN9536-4373 |
| Official title of study: | Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity |
| Document date | 11 June 2020 |

16.1.1 Protocol and protocol amendments

List of contents

| | |
|----------------------------------|----------------------|
| Protocol | Link |
| Attachment I and II..... | Link |
| Amendment 2 - UK..... | Link |
| Amendment 3 - Global..... | Link |
| Amendment 4 - DE | Link |
| Amendment 5 - UK..... | Link |
| Amendment 6 - Global..... | Link |

*Redacted protocol
Includes redaction of personal identifiable information only.*

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

21 December 2017
2.0
Final
1 of 104

Novo Nordisk

Protocol

Protocol title: Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity

Substance name: semaglutide

Universal Trial Number: U1111-1200-8053

EudraCT Number: 2017-003436-36

Trial phase: 3a

In the following, Novo Nordisk A/S and its affiliates will be stated as “Novo Nordisk”.

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Table of Contents

| | Page |
|---|-----------|
| Table of Contents | 2 |
| 1 Synopsis | 5 |
| 2 Flowchart | 8 |
| 3 Introduction | 14 |
| 3.1 Trial rationale | 14 |
| 3.2 Background | 14 |
| 3.2.1 Semaglutide | 14 |
| 3.2.2 Trial population | 15 |
| 3.3 Benefit-risk assessment | 15 |
| 3.3.1 Benefits | 15 |
| 3.3.2 Risks and precautions | 16 |
| 3.3.3 Conclusion on benefit-risk profile | 17 |
| 4 Objectives and endpoints | 18 |
| 4.1 Primary, secondary and exploratory objective(s) | 18 |
| 4.2 Primary, secondary and exploratory endpoint(s) | 19 |
| 4.2.1 Primary endpoint | 19 |
| 4.2.2 Secondary endpoints | 19 |
| 4.2.2.1 Confirmatory secondary endpoints | 19 |
| 4.2.2.2 Supportive secondary endpoints | 19 |
| 4.2.3 Exploratory endpoints | 21 |
| 5 Trial design | 21 |
| 5.1 Overall design | 21 |
| 5.2 Subject and trial completion | 22 |
| 5.3 End of trial definition | 23 |
| 5.4 Scientific rationale for trial design | 23 |
| 5.5 Justification for dose | 23 |
| 6 Trial population | 23 |
| 6.1 Inclusion criteria | 23 |
| 6.2 Exclusion criteria | 24 |
| 6.3 Randomisation criteria | 25 |
| 6.4 Lifestyle restrictions | 26 |
| 6.4.1 Meals and dietary restrictions | 26 |
| 6.4.2 Caffeine and tobacco | 26 |
| 6.5 Screen failures | 27 |
| 7 Treatments | 27 |
| 7.1 Treatments administered | 27 |
| 7.1.1 Medical device | 29 |
| 7.1.2 Diet and physical activity counselling | 29 |
| 7.2 Dose modification | 30 |
| 7.3 Method of treatment assignment | 30 |
| 7.4 Blinding | 30 |
| 7.5 Preparation/Handling/Storage/Accountability | 31 |

| | | |
|-----------|---|-----------|
| 7.6 | Treatment compliance | 32 |
| 7.7 | Concomitant medication | 32 |
| 7.7.1 | Rescue medication | 32 |
| 7.8 | Treatment after the end of the trial | 32 |
| 8 | Discontinuation/Withdrawal criteria | 33 |
| 8.1 | Discontinuation of trial treatment | 33 |
| 8.1.1 | Temporary discontinuation of trial treatment | 34 |
| 8.2 | Withdrawal from the trial | 34 |
| 8.2.1 | Replacement of subjects | 35 |
| 8.3 | Lost to follow-up | 35 |
| 9 | Trial assessments and procedures | 35 |
| 9.1 | Efficacy assessments | 37 |
| 9.1.1 | Body measurements | 37 |
| 9.1.2 | Clinical outcome assessments | 37 |
| 9.1.3 | Clinical efficacy laboratory assessments | 38 |
| 9.2 | Adverse events | 38 |
| 9.2.1 | Time period and frequency for collecting AE and SAE information | 38 |
| 9.2.1.1 | Events for adjudication | 40 |
| 9.2.2 | Method of detecting AEs and SAEs | 41 |
| 9.2.3 | Follow-up on AEs and SAEs | 41 |
| 9.2.4 | Regulatory reporting requirements for SAEs | 41 |
| 9.2.5 | Hypoglycaemic, cardiovascular and death events | 41 |
| 9.2.6 | Disease-related events and/or disease-related outcomes not qualifying as an AE or SAE | 42 |
| 9.2.7 | Pregnancies and associated adverse events | 42 |
| 9.2.8 | Medical device incidents (including malfunctions) | 42 |
| 9.2.9 | Technical complaints | 42 |
| 9.3 | Treatment of overdose | 43 |
| 9.4 | Safety assessments | 43 |
| 9.4.1 | Mental health assessment instruments | 44 |
| 9.4.2 | Physical examinations | 45 |
| 9.4.3 | Vital signs | 45 |
| 9.4.4 | Electrocardiograms | 45 |
| 9.4.5 | Clinical safety laboratory assessments | 45 |
| 9.4.6 | Immunogenicity assessments | 46 |
| 9.5 | Pharmacokinetics | 46 |
| 9.6 | Pharmacodynamics | 46 |
| 9.7 | Genetics | 46 |
| 9.8 | Biomarkers | 46 |
| 9.9 | Severe hypersensitivity | 47 |
| 10 | Statistical considerations | 47 |
| 10.1 | Sample size determination | 48 |
| 10.2 | Definition of analysis sets | 50 |
| 10.3 | Statistical analyses | 51 |
| 10.3.1 | Primary endpoint | 51 |
| 10.3.2 | Secondary endpoints | 57 |
| 10.3.2.1 | Confirmatory secondary endpoints | 57 |

| | | |
|-------------------|---|-----------|
| 10.3.2.2 | Supportive secondary endpoints | 58 |
| 10.3.3 | Exploratory endpoints | 61 |
| 10.3.4 | Explorative statistical analysis for pharmacogenetics and biomarkers | 61 |
| 10.3.5 | Other analyses | 61 |
| 10.4 | Pharmacokinetic and/or pharmacodynamic modelling | 61 |
| 11 | Appendices | 62 |
| Appendix 1 | Abbreviations and Trademarks | 63 |
| Appendix 2 | Clinical laboratory tests | 66 |
| Appendix 3 | Trial governance considerations | 70 |
| Appendix 4 | Adverse events: definitions and procedures for recording, evaluation, follow-up, and reporting | 79 |
| Appendix 5 | Contraceptive guidance and collection of pregnancy information | 85 |
| Appendix 6 | Technical complaints: Definition and procedures for recording, evaluation, follow-up and reporting | 88 |
| Appendix 7 | Retention of human biosamples | 90 |
| Appendix 8 | Monitoring of calcitonin | 92 |
| Appendix 9 | Country-specific requirements | 95 |
| 12 | References | 98 |

Attachment I Global list of key staff and relevant departments and suppliers

Attachment II Country list of key staff and relevant departments

1 Synopsis

Rationale:

The prevalence of obesity has reached epidemic proportions in most countries around the world and the prevalence is still increasing at an alarming rate¹⁻⁷. The medical and societal impacts are considerable and obesity is one of the most significant public health challenges worldwide¹⁻⁷. Obesity is associated with increased risk of a variety of comorbidities, affects physical and mental health and reduces health related quality of life^{8,9}.

Pharmacotherapy may serve as a valuable adjunct to lifestyle intervention for individuals with obesity in order to achieve and sustain a clinically relevant weight loss, to improve comorbid conditions and to facilitate a healthier lifestyle. Few anti-obesity medications are currently available and there is a need for more effective and safe therapeutic options for treatment of obesity. The present trial is a 68-week trial designed to show reduction in body weight and compare the effect and safety of semaglutide subcutaneous (s.c.) 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity.

Objectives and endpoints:

Primary objective

To compare the effect of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity on body weight

Key secondary objectives

To compare the effect of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity on:

- Cardiovascular risk factors
- Clinical Outcome Assessments (COAs)

To compare the safety and tolerability of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity

Primary estimand

The estimand will quantify the average treatment effect of semaglutide relative to semaglutide placebo after 68 weeks, as an adjunct to a reduced-calorie diet and increased physical activity, in all randomised subjects regardless of adherence to treatment and regardless of starting rescue

interventions (weight management drugs or bariatric surgery) (“effectiveness”/“treatment policy” estimand). The estimand will cover all effect-related objectives.

Primary endpoint

The primary endpoints addressing the primary objective:

- Change from baseline at week 0 to week 68 in body weight (%)
- Subjects who after 68 weeks achieve (yes/no):
 - Body weight reduction $\geq 5\%$ from baseline at week 0

Confirmatory secondary endpoints

- Subjects who after 68 weeks achieve (yes/no):
 - Body weight reduction $\geq 10\%$ from baseline at week 0
 - Body weight reduction $\geq 15\%$ from baseline at week 0
- Change from baseline at week 0 to week 68 in:
 - Waist circumference (cm)
 - Systolic blood pressure (mmHg)
 - Physical functioning score (SF-36)
 - Physical function domain (5-items) score (IWQoL-Lite for CT)

Overall design:

This is a 68-week, randomised, double-blind, placebo-controlled, two-armed, parallel group, multi-centre, multinational clinical trial comparing semaglutide s.c. 2.4 mg once-weekly with semaglutide placebo once-weekly in subjects with overweight or obesity.

Key inclusion criteria

- Male or female, age ≥ 18 years at the time of signing informed consent
- Body mass index (BMI) $\geq 30 \text{ kg/m}^2$ or $\geq 27 \text{ kg/m}^2$ with the presence of at least one of the following weight-related comorbidities (treated or untreated): hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease
- History of at least one self-reported unsuccessful dietary effort to lose body weight

Key exclusion criteria

- HbA_{1c} $\geq 48 \text{ mmol/mol}$ (6.5%) as measured by the central laboratory at screening
- A self-reported change in body weight $> 5 \text{ kg}$ (11 lbs) within 90 days before screening irrespective of medical records

Number of subjects:

Approximately 2294 subjects will be screened to achieve 1950 subjects randomly assigned to trial product.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

21 December 2017
2.0
Final
7 of 104

Novo Nordisk

Treatment groups and duration:

Eligible subjects fulfilling all randomisation criteria at visit 2 will be randomised in a 2:1 manner to receive either semaglutide s.c 2.4 mg once-weekly or semaglutide placebo once-weekly as an adjunct to a reduced-calorie diet and increased physical activity.

The trial includes a screening visit to assess the subject's eligibility followed by visits/phone contacts every 2nd week for the first 20 weeks. For the remaining maintenance period until end of treatment (week 68), visits/phone contacts will take place every 4th week. A follow-up visit ('End of trial') for safety assessments is scheduled 7 weeks after end of treatment to account for the exposure and the long half-life of semaglutide.

The following trial products will be supplied by Novo Nordisk A/S:

- Semaglutide B 1.0 mg/mL PDS290 and semaglutide placebo, solution for injection, 3 mL PDS290 pre-filled pen-injector
- Semaglutide B 3.0 mg/mL PDS290 and semaglutide placebo, solution for injection, 3 mL PDS290 pre-filled pen-injector

2 Flowchart

[illegible]

[illegible]

[illegible]

| | Screening | Randomisation | Dose escalation period | | | | | | | | | | Maintenance period | | | | | | | | | | End of treatment | End of trial | |
|---|-----------|---------------|------------------------|----|----|----|----|----|----|-----|-----|-----|--------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------------------|--------------|---------|
| | V1 | V2 | V3 | V4 | V5 | V6 | V7 | V8 | V9 | V10 | P11 | V12 | P13 | V14 | P15 | V16 | P17 | V18 | P19 | V20 | P21 | V22 | P23 | V24 | V25 |
| Visit (V), Phone (P) | | | | | | | | | | | | | | | | | | | | | | | | V24 | V25 |
| Timing of Visit (Weeks) | -1 | 0 | 2 | 4 | 6 | 8 | 10 | 12 | 14 | 16 | 18 | 20 | 24 | 28 | 32 | 36 | 40 | 44 | 48 | 52 | 56 | 60 | 64 | 68 | 75 |
| Visit Window (Days) | -7 to 0 | 0 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | 0 to +5 |
| SAFETY | | | | | | | | | | | | | | | | | | | | | | | | | |
| Physical examination (9.4.2) | X | | | | | | | | | | | | | | | | | | | | | | | X | |
| Pregnancy test (9.4.5, Appendix 5) | X | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | X |
| ECG (9.4.4) | | X | | | | | | | | | | X | | | | | | | | | | | | X | |
| Adverse event (9.2, Appendix 4) | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X |
| Technical complaint (9.2.9, Appendix 4) | | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | |
| Breast neoplasms follow-up ^b (9.4) | | | | | | | | | | | | | | | | | | | | | | | | X | X |
| Colon neoplasms follow-up (9.4) | | | | | | | | | | | | | | | | | | | | | | | | X | X |
| Haematology (Appendix 2) | X | | | | | | | | | | | X | | | | | | | | X | | | | X | |
| Biochemistry (Appendix 2) | X | | | | | | | | | | | X | | | | | | | | X | | | | X | |
| Anti-Semaglutide Antibody (9.4.6) | | X | X | X | X | X | | X | | | | | | X | | | | | | X | | | | X | X |
| Vital Signs (9.4.3) | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pulse | X | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | | X | X |
| Clinical Outcome Assessments (9.4.1) | | | | | | | | | | | | | | | | | | | | | | | | | |
| Patient Health Questionnaire-9 (PHQ-9) (9.4.1) | X | X | | | | | | X | | | | X | | | | X | | | | X | | | | X | |
| Columbia-Suicide Severity Rating Scale (C-SSRS) (9.4.1) | X | X | | | | | | X | | | | X | | | | X | | | | X | | | | X | |

[illegible]

| | Screening | Randomisation | Dose escalation period | | | | | | | | Maintenance period | | | | | | | | | | | | End of treatment | End of trial | |
|---|-----------|---------------|------------------------|----|----|----|----|----|----|-----|--------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------------------|--------------|----------------|
| | V1 | V2 | V3 | V4 | P5 | V6 | P7 | V8 | P9 | V10 | P11 | V12 | P13 | V14 | P15 | V16 | P17 | V18 | P19 | V20 | P21 | V22 | P23 | V24 | V25 |
| Visit (V), Phone (P) | | | | | | | | | | | | | | | | | | | | | | | | | |
| Timing of Visit (Weeks) | -1 | 0 | 2 | 4 | 6 | 8 | 10 | 12 | 14 | 16 | 18 | 20 | 24 | 28 | 32 | 36 | 40 | 44 | 48 | 52 | 56 | 60 | 64 | 68 | 75 |
| Visit Window (Days) | -7 to 0 | ±0 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | ±3 | 0 to +5 |
| Hand out dose reminder card (7.1) | | X | X | X | X | X | X | X | | X | | | | | | | | | | | | | | | |
| Hand out and instruct in food diary (9) | X | | | | | | | | | | | | | | | | | | | | | | | | |
| Hand out and instruct in PK diary (9) | | X | X | X | X | X | | | | | | X | | | | | | X | | | | X | | | |
| Collect, review and transcribe diaries | | X | X | X | X | X | X | X | | X | | X | | X | | X | | X | | X | | X | | X | |
| Hand out ID card | X | | | | | | | | | | | | | | | | | | | | | | | | |
| Attend visit fasting (6.4.1) | | X | | | | | | | | | | X | | | | | | | | | | | | X | X ^g |

^a Demography consists of date of birth, sex, ethnicity, and race (according to local regulation).

^b For all female subjects.

^c Smoking is defined as smoking at least one cigarette or equivalent daily.

^d DEXA scan is performed in a sub-population.

^e DEXA scan should be performed in fasting state prior to randomisation (after informed consent is obtained).

^f Only for subjects where the separate informed consent for future research has been signed.

^g Fasting at V25 is defined as at least 2 hours without food and drink intake except water before attending the visit.

3 Introduction

3.1 Trial rationale

The prevalence of obesity has reached epidemic proportions in most countries around the world and the prevalence is still increasing at an alarming rate¹⁻⁷. The medical and societal impacts are considerable and obesity is one of the most significant public health challenges worldwide¹⁻⁷. Obesity is associated with increased risk of a variety of comorbidities including hyperglycaemia, type 2 diabetes (T2D), hypertension, dyslipidaemia, obstructive sleep apnoea, atherosclerosis, osteoarthritis, urinary incontinence, non-alcoholic steatohepatitis, cardiovascular diseases, certain types of cancer, and risk of early death¹⁰⁻²⁴. Moreover, obesity adversely affects physical and mental health and reduces health related quality of life^{8,9}. Obesity is also associated with decreased cardiorespiratory fitness, which also increases the risk of cardiovascular diseases and all-cause mortality²⁵.

The risk of obesity-related complications and comorbidities increases with increasing BMI, and a weight loss of 5–10 % has significant health benefits in terms of slowing progression to T2D²⁶⁻²⁹. Furthermore, a weight loss of 5–10% improves many other obesity related comorbidities as well as physical symptoms and quality of life³⁰⁻³⁷. Finally, studies suggest a beneficial impact of weight loss on cardiovascular risk and mortality in both people with diabetes and obesity³⁸⁻⁴⁰.

Lifestyle intervention in the form of diet and exercise is first line treatment for obesity, but most people with obesity struggle to achieve and maintain their weight loss⁴¹⁻⁵⁰. Surgical treatments offer an effective alternative for some people with severe obesity, but surgery carries a risk in connection with the procedure and is not without complications. Furthermore, surgery requires close follow-up of the individual which can be cumbersome and costly^{41-46, 51, 52}. Pharmacotherapy may therefore serve as a valuable adjunct to lifestyle intervention for individuals with obesity in order to achieve and sustain a clinically relevant weight loss, to improve comorbid conditions and to facilitate a healthier lifestyle. Few anti-obesity medications are currently available and there is a need for more safe and effective therapeutic options for treatment of obesity, especially treatments that also target weight maintenance, prevention and treatment of comorbidities^{41-45, 53, 54}.

3.2 Background

3.2.1 Semaglutide

Semaglutide is the next generation glucagon-like peptide -1 receptor agonist (GLP-1 RA) currently under development by Novo Nordisk for the treatment of weight management (NN9536). Semaglutide has been optimised resulting in a longer half-life of approximately 160 hours, making it suitable for once-weekly dosing⁵⁵. GLP-1 RA is a physiological regulator of appetite and GLP-1 receptors are present in several areas of the brain involved in appetite regulation⁵⁶.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 15 of 104

Novo Nordisk

A 52-week phase 2 dose-finding trial within weight management (NN9536-4153) has recently been completed. A total of 957 randomised subjects with obesity (without diabetes) were exposed to semaglutide (n=718), liraglutide 3.0 mg (n=103) or placebo (n=136). In this trial, an overall monotone dose-dependent weight loss was observed across the 5 semaglutide doses tested (0.05 to 0.4 mg once-daily). The estimated weight loss at week 52 was 13.8 % at the highest dose tested (0.4 mg once-daily) compared to the weight loss of 2.3% achieved by diet, exercise and placebo alone.⁵⁷

Clinical⁵⁸⁻⁶² and non-clinical data⁶³ indicate that the body weight-reducing effect of semaglutide is mainly mediated by a reduced energy intake. No unexpected safety findings were identified and the tolerability and safety profile was overall consistent with previous findings in the T2D development programme and the GLP-1 RA class in general.

A comprehensive review of results from the non-clinical and clinical studies of semaglutide can be found in the current edition of the investigator's brochure (IB)⁵⁷ and any updates hereof.

3.2.2 Trial population

The trial population will consist of subjects with obesity (BMI > 30 kg/m²) or with overweight (BMI > 27 kg/m²) and weight-related comorbidities. These subjects represent a clinically relevant population for pharmacotherapeutic weight management as they are at significant risk for weight-related morbidities and mortality, and are likely to benefit from weight reduction. Information about weight-related comorbidities, including hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease, will be collected systematically at screening by the investigators as part of the medical history.

Subjects with T2D are excluded from the trial. Subjects with overweight or obesity and T2D will be addressed in a dedicated trial.

First line treatment in weight management should always be lifestyle modification through a reduced-calorie diet and increased physical activity. Thus only subjects who have tried but failed a dietary weight loss intervention will be included in accordance with regulatory guidelines^{64, 65}.

3.3 Benefit-risk assessment

3.3.1 Benefits

Subjects will be treated with a regimen anticipated to be better than or equal to the weight management they receive at the time of entry into the trial. Results from the phase 2 trial (NN9536-4153) demonstrated that semaglutide s.c. once-daily as an adjunct to a reduced-calorie diet and increased physical activity was effective for weight loss in subjects with obesity, while displaying a satisfactory tolerability profile. Overall, a monotone dose-dependent weight loss was observed across all tested doses of semaglutide (0.05 to 0.4 mg once-daily). The weight loss was 11.55 percentage points larger for the 0.4 mg group compared with placebo. Weight loss was

accompanied by a consistent improvement in weight-related comorbidities, indicated by cardiovascular risk factors, lipid profile and glycaemic factors, as well as improvements in clinical outcome assessments.

In addition, it is expected that all subjects will benefit from participation through close contact with the trial site and counselling by a dietician or a similar qualified healthcare professional, all of which will most likely result in intensified weight management.

3.3.2 Risks and precautions

The sections below describe identified and potential risks associated with semaglutide treatment. The identified/potential risks are based on findings in non-clinical studies and clinical trials with semaglutide as well as other GLP-1 RAs. For each of these risks, mitigating actions have been implemented to minimise the risks for subjects enrolled in this trial.

Identified risks

- Gastrointestinal adverse events
 - Consistent with findings with other GLP-1 RAs, the most frequently reported adverse events (AE) in clinical trials with semaglutide were gastrointestinal AEs. A low starting dose and dose escalation steps will be implemented in the trial to mitigate the risk of gastrointestinal AEs.
- Cholelithiasis
 - Events of cholelithiasis were the most frequently reported gallbladder events in the phase 2 weight management trial (NN9536-4153) and were in a few instances co-reported with the event adjudication committee (EAC) confirmed acute pancreatitis. As a precaution, if cholelithiasis is suspected, appropriate clinical follow-up is to be initiated at the investigator's discretion.
- Acute pancreatitis
 - Acute pancreatitis has been observed with the use of GLP-1 RA drug class. As a precaution, subjects with a history of chronic pancreatitis or recent acute pancreatitis will not be enrolled in the trial. In addition, trial product should be discontinued in case of suspicion of acute pancreatitis in accordance to Section [8.1](#).

Potential risks

- Medullary thyroid cancer (MTC) (based on non-clinical data)
 - Expected proliferative thyroid C-cell changes were seen in the mouse and rat carcinogenicity studies after daily exposure to semaglutide for 2 years. No hyperplasia was observed in monkeys after 52 weeks exposure up to 13-fold above the clinical plasma exposure at 2.4 mg/week. In clinical trials with semaglutide, there have been no reports of MTC or clinically relevant changes in calcitonin levels. The C-cell changes in rodents are mediated by the GLP-1 receptor, which is not expressed in the normal

human thyroid. Accordingly, the risk of GLP-1 receptor-mediated C-cell changes in humans is considered to be low. However, as a precaution, exclusion and discontinuation criteria related to medical history of multiple endocrine neoplasia type 2 (MEN2) or MTC and elevated plasma levels of calcitonin (biomarker for MTC) have been implemented in the trial.

- Pancreatic cancer
 - There is currently no support from non-clinical studies, clinical trials or post-marketing data that GLP-1 RA based therapies increase the risk of pancreatic cancer, but pancreatic cancer has been classified as a potential class risk of GLP-1 RAs by European Medicines Agency. As a precaution, subjects with a history of malignant neoplasms within the past 5 years prior to screening will be excluded from the trial.
- Allergic reactions
 - As is the case with all protein-based pharmaceuticals, subjects treated with semaglutide are at risk of developing immunogenic and allergic reactions. As a precaution, subjects with known or suspected hypersensitivity to semaglutide or related products will not be enrolled in this trial.

Other risks

- Pregnancy and fertility (based on non-clinical data)
 - Studies in animals have shown reproductive toxicity. There are limited data from the use of semaglutide in pregnant women. Therefore, semaglutide should not be used during pregnancy. Exclusion and discontinuation criteria related to pregnancy have been implemented in the trial.

3.3.3 Conclusion on benefit-risk profile

Necessary precautions have been implemented in the design and planned conduct of the trial in order to minimise the risks and inconveniences of participation in the trial. The safety profile for semaglutide generated from the clinical and non-clinical development programme has not revealed any safety issues that would prohibit administration of semaglutide s.c. 2.4 mg once-weekly. The results of the phase 2 trial (NN9536-4153) indicate that semaglutide will provide a clinically meaningful weight loss.

In conclusion, the potential risk to the subjects in this trial is considered low and outweighed by the anticipated benefits that semaglutide would provide subjects included in this trial.

More detailed information about the known and expected benefits and risks and reasonably expected adverse events of semaglutide s.c. may be found in the IB⁵⁷ and any updates hereof.

4 Objectives and endpoints

4.1 Primary, secondary and exploratory objective(s)

Primary objective

To compare the effect of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity on body weight

Secondary objectives

To compare the effect of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity on:

- Cardiovascular risk factors
- Clinical Outcome Assessments (COAs)
- Glucose metabolism
- Other factors related to body weight

To compare the safety and tolerability of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity

Exploratory objectives

To compare the effect of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity in subjects with overweight or obesity on:

- Glycaemic status
- Use of medication for hypertension and dyslipidaemia
- Work productivity
- Treatment discontinuation
- Liver indices
- Urinary incontinence
- Diet and physical activity

Primary estimand

The estimand will quantify the average treatment effect of semaglutide relative to semaglutide placebo after 68 weeks, as an adjunct to a reduced-calorie diet and increased physical activity, in all randomised subjects regardless of adherence to treatment and regardless of starting rescue

interventions (weight management drugs or bariatric surgery) (“effectiveness”/“treatment policy” estimand). The estimand will cover all effect-related objectives.

Secondary estimand

The estimand will quantify the average treatment effect of semaglutide relative to semaglutide placebo after 68 weeks, as an adjunct to a reduced-calorie diet and increased physical activity, in all randomised subjects had they remained on their randomised treatment for the entire planned duration of the trial and not started any rescue intervention (weight management drugs or bariatric surgery) (“efficacy”/“hypothetical” estimand). The estimand will cover the primary objective.

4.2 Primary, secondary and exploratory endpoint(s)

4.2.1 Primary endpoint

The primary endpoints addressing the primary objective:

- Change from baseline at week 0 to week 68 in body weight (%)
- Subjects who after 68 weeks achieve (yes/no):
 - Body weight reduction $\geq 5\%$ from baseline at week 0

4.2.2 Secondary endpoints

The confirmatory and supportive secondary endpoints addressing the primary and secondary objectives are listed in Section [4.2.2.1](#) and [4.2.2.2](#).

4.2.2.1 Confirmatory secondary endpoints

- Subjects who after 68 weeks achieve (yes/no):
 - Body weight reduction $\geq 10\%$ from baseline at week 0
 - Body weight reduction $\geq 15\%$ from baseline at week 0
- Change from baseline at week 0 to week 68 in:
 - Waist circumference (cm)
 - Systolic blood pressure (mmHg)
 - Physical functioning score (SF-36)
 - Physical function domain (5-items) score (IWQoL-Lite for CT)

4.2.2.2 Supportive secondary endpoints

Effect endpoints:

- Change from baseline at week 0 to week 68 in:
 - Body weight (kg)
 - BMI (kg/m^2)
 - HbA_{1C} (% , mmol /mol)
 - FPG (mg/dL)

- Fasting serum insulin (mIU/L)
- Diastolic blood pressure (mmHg)
- Lipids (mg/dL)
 - Total cholesterol
 - High density lipoprotein (HDL) cholesterol
 - Low density lipoprotein (LDL) cholesterol
 - Very low density lipoprotein (VLDL) cholesterol
 - Free fatty acids
 - Triglycerides
- High sensitivity C-Reactive Protein (hsCRP) (mg/L)
- Plasminogen Activator Inhibitor-1 (PAI-1) Activity (AU/mL)
- Soluble Leptin Receptor (ng/mL)
- Leptin (ng/mL)
- Short Form-36 (SF-36)
 - role-physical score
 - bodily pain score
 - general health score
 - vitality score
 - social functioning score
 - role-emotional score
 - mental health score
 - physical component summary
 - mental component summary
- IWQoL-Lite for CT
 - pain/discomfort domain score
 - psychosocial domain score
 - total score
- Body composition (as assessed by Dual Energy X-ray Absorptometry (DEXA) in a subset of subjects)
 - Total fat mass (% , g)
 - Lean body mass (% , g)
 - Visceral fat mass (% , g)
- Subjects who after 68 weeks achieve (yes/no):
 - Responder definition value for SF-36 physical functioning score
 - Responder definition value for IWQoL-Lite for CT physical function domain (5-items) score

Safety endpoints:

- Number of treatment emergent adverse events (TEAEs) from baseline at week 0 to week 75

- Number of serious adverse events (SAEs) from baseline at week 0 to week 75
- Change from baseline at week 0 to week 68 in:
 - Pulse (bpm)
 - Amylase (U/L)
 - Lipase (U/L)
 - Calcitonin (ng/L)

4.2.3 Exploratory endpoints

The exploratory endpoints addressing the explorative objectives:

- Change from baseline at week 0 to week 68 in:
 - Glycaemic category (normo-glycaemia, pre-diabetes, T2D)
 - Antihypertensive medication (decrease, no change, increase)
 - Lipid-lowering medication (decrease, no change, increase)
 - The Stanford Presenteeism Scale (SPS-6), total score
 - Fatty liver Index (FLI) score category (<30 , ≥ 30 and <60 , ≥ 60)
 - International Consultation on Incontinence Questionnaire-Urinary Incontinence-Short Form (ICIQ-UI-SF), sum score (assessed in female subjects)
- Subjects who from randomisation at week 0 to week 68 have permanently discontinued randomised trial product (yes/no)
- Time to permanent discontinuation of randomised trial product (weeks)
- Number of days per week with at least one entry in the food diary from baseline at week 0 to week 68
- Number of minutes per week of physical activity from baseline at week 0 to week 68

5 Trial design

5.1 Overall design

This is a 68-week, randomised, double-blind, placebo-controlled, two-armed, parallel group, multi-centre, multinational clinical trial comparing semaglutide s.c. 2.4 mg once-weekly with semaglutide placebo once-weekly in subjects with overweight or obesity.

The trial includes a screening visit to assess the subject's eligibility followed by visits/phone contacts every 2nd week during dose escalation. From week 20, visits/phone contacts will take place every 4th week for the remaining maintenance period until end of treatment (week 68). A follow-up visit ('End of trial') for safety assessments is scheduled 7 weeks after end of treatment to account for the exposure and the long half-life of semaglutide.

The trial design is outlined in [Figure 5-1](#).




Figure 5-1 A schematic diagram of the trial design with the duration of the trial periods including follow-up period.

Eligible subjects fulfilling all randomisation criteria at visit 2 will be randomised in a 2:1 manner to receive either semaglutide s.c 2.4 mg once-weekly or semaglutide placebo once-weekly as an adjunct to a reduced-calorie diet and increased physical activity.

A sub-population of approximately 135 randomised subjects; corresponding to ~3% of the subjects in the phase 3a programme will have their body composition assessed by DEXA at screening and end-of-treatment to demonstrate that weight loss is primarily caused by reduction in fat mass, in accordance with the EMA and FDA guidelines^{66, 67}.

5.2 Subject and trial completion

Approximately 2294 subjects will be screened to achieve 1950 subjects randomly assigned to trial product.

Approximately 135 randomised subjects will be included in the DEXA sub-population.

Trial period completion for a subject:

Trial period completion is defined as when the randomised subject has completed the final scheduled visit ('end of trial' according to the flowchart).

'Date of trial completion' is the date the subject completed the final scheduled visit.

Treatment period completion for a subject:

Treatment period completion is defined as when the randomised subject has attended the 'end of treatment' visit according to the flowchart.

5.3 End of trial definition

The end of the trial is defined as the date of the last visit of the last subject in the trial.

5.4 Scientific rationale for trial design

The treatment duration of the trial is 68 weeks with an additional 7 weeks follow-up (without treatment). A 68-week treatment duration (including 52 weeks on target dose) is considered sufficient to assess weight loss, safety and tolerability in the phase 3 weight management development programme in accordance with regulatory guidelines^{64,65}. The 7 weeks follow-up period is included to account for the exposure and long half-life of semaglutide.

A randomised, double-blinded, placebo-controlled, multi-centre trial design is chosen to minimise bias in the assessment of the effect and safety of semaglutide s.c. 2.4 mg once-weekly versus semaglutide placebo, as an adjunct to a reduced calorie diet and increased physical activity.

5.5 Justification for dose

Results from the phase 2 dose-finding trial (NN9536-4153) showed that the semaglutide s.c. 0.4 mg once-daily dose was most effective in terms of weight loss while displaying an acceptable tolerability profile. Using population pharmacokinetic (PK) modelling, it was estimated that a once-weekly maintenance dose of 2.4 mg semaglutide s.c. will result in similar C_{max} at steady-state as that obtained by the once-daily 0.4 mg semaglutide s.c. dose in trial NN9536-4153.

A maintenance dose of 2.4 mg semaglutide s.c. once-weekly has been chosen for the phase 3 weight management development programme. The once-weekly dosing is anticipated to ease the burden of drug administration in clinical practice. Subjects will be initiated at a once-weekly dose of 0.25 mg and follow a fixed-dose escalation regimen, with dose increases every 4 weeks (to doses of 0.5, 1.0, 1.7 and 2.4 mg/week), until the target dose is reached after 16 weeks.

It is well known that to mitigate GI side effects with GLP-1 RA treatment, dose escalation to the target dose is required. Based on experience from the semaglutide T2D development programme, a similar fixed-dose escalation regimen was selected, with dose escalation every 4 weeks until the target dose of 2.4 mg is reached after 16 weeks.

Please refer to Section [7.1](#) for more details on treatment doses.

6 Trial population

Prospective approval of protocol deviations to recruitment and enrolment criteria, also known as protocol waivers or exemptions, is not permitted.

6.1 Inclusion criteria

Subjects are eligible to be included in the trial only if all of the following criteria apply:

1. Informed consent obtained before any trial-related activities. Trial-related activities are any procedures that are carried out as part of the trial, including activities to determine suitability for the trial
2. Male or female, age ≥ 18 years at the time of signing informed consent
3. Body mass index (BMI) ≥ 30 kg/m² or ≥ 27 kg/m² with the presence of at least one of the following weight-related comorbidities (treated or untreated): hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease
4. History of at least one self-reported unsuccessful dietary effort to lose body weight

The criteria will be assessed at the investigator's discretion unless otherwise stated.

For country specific requirements, see [Appendix 9](#).

6.2 Exclusion criteria

Subjects are excluded from the trial if any of the following criteria apply:

Glycaemia-related:

1. HbA1c ≥ 48 mmol/mol (6.5%) as measured by the central laboratory at screening
2. History of type 1 or type 2 diabetes mellitus
3. Treatment with glucose-lowering agent(s) within 90 days before screening
4. Treatment with a GLP-1 receptor agonist within 180 days before screening

Obesity-related:

5. A self-reported change in body weight > 5 kg (11 lbs) within 90 days before screening irrespective of medical records
6. Treatment with any medication for the indication of obesity within the past 90 days before screening
7. Previous or planned (during the trial period) obesity treatment with surgery or a weight loss device. However, the following are allowed: (1) liposuction and/or abdominoplasty, if performed > 1 year before screening, (2) lap banding, if the band has been removed > 1 year before screening, (3) intragastric balloon, if the balloon has been removed > 1 year before screening or (4) duodenal-jejunal bypass sleeve, if the sleeve has been removed > 1 year before screening
8. Uncontrolled thyroid disease, defined as thyroid stimulating hormone (TSH) > 6.0 mIU/L or < 0.4 mIU/L as measured by the central laboratory at screening

Mental health:

9. History of major depressive disorder within 2 years before screening
10. Diagnosis of other severe psychiatric disorder (e.g., schizophrenia, bipolar disorder)
11. A Patient Health Questionnaire-9 (PHQ-9) score of ≥ 15 at screening
12. A lifetime history of a suicidal attempt

13. Suicidal behaviour within 30 days before screening
14. Suicidal ideation corresponding to type 4 or 5 on the Columbia-Suicide Severity Rating Scale (C-SSRS) within the past 30 days before screening

General safety:

15. Use of non-herbal Chinese medicine or other non-herbal local medicine with unknown/unspecified content within 90 days before screening
16. Presence of acute pancreatitis within the past 180 days prior to the day of screening
17. History or presence of chronic pancreatitis
18. Calcitonin ≥ 100 ng/L as measured by the central laboratory at screening
19. Personal or first degree relative(s) history of multiple endocrine neoplasia type 2 or medullary thyroid carcinoma
20. Renal impairment measured as estimated Glomerular Filtration Rate (eGFR) value of eGFR < 15 ml/min/1.73 m² as defined by KDIGO 2012⁶⁸ by the central laboratory at screening
21. History of malignant neoplasms within the past 5 years prior to screening. Basal and squamous cell skin cancer and any carcinoma in-situ are allowed
22. Any of the following: myocardial infarction, stroke, hospitalisation for unstable angina or transient ischaemic attack within the past 60 days prior to screening
23. Subject presently classified as being in New York Heart Association (NYHA) Class IV
24. Surgery scheduled for the duration of the trial, except for minor surgical procedures, in the opinion of the investigator
25. Known or suspected abuse of alcohol or recreational drugs
26. Known or suspected hypersensitivity to trial product(s) or related products
27. Previous participation in this trial. Participation is defined as signed informed consent
28. Participation in another clinical trial within 90 days before screening
29. Other subject(s) from the same household participating in any semaglutide trial
30. Female who is pregnant, breast-feeding or intends to become pregnant or is of child-bearing potential and not using a highly effective contraceptive method
31. Any disorder, unwillingness or inability, not covered by any of the other exclusion criteria, which in the investigator's opinion, might jeopardise the subject's safety or compliance with the protocol

The criteria will be assessed at the investigator's discretion unless otherwise stated.

For country specific requirements, see [Appendix 9](#) and for contraceptive requirements, see [Appendix 5](#).

6.3 Randomisation criteria

Randomisation criteria applicable for all sites

Subjects are eligible to be randomised in the trial only if all of the following criteria apply:

1. Have kept a food diary with at least one entry per day between screening and randomisation. However, missed entries for a maximum of two days are allowed
2. A Patient health questionnaire-9 (PHQ-9) score of < 15 at randomisation
3. No suicidal behaviour in the period between screening and randomisation
4. No suicidal ideation corresponding to type 4 or 5 on the Columbia–Suicide Severity Rating Scale (C-SSRS) in the period between screening and randomisation

Subjects not fulfilling the randomisation criteria will be considered screen failure, see Section [6.5](#).

Randomisation criteria applicable only for DEXA sites

To be qualified for the DEXA sub-population, the following randomisation criteria must also be answered “yes”.

1. Evaluation of the quality of the baseline DEXA scan must be performed and found acceptable by the imaging laboratory before randomisation to the DEXA sub-study.
2. BMI ≤ 40 kg/m² at screening

Subjects who do not fulfil the above randomisation criteria can be randomised in the trial outside the DEXA sub-population.

6.4 Lifestyle restrictions

To ensure alignment in regards to performance of assessments across subjects and trial sites, the below restrictions apply.

6.4.1 Meals and dietary restrictions

- Subjects must attend the visits fasting according to the flowchart.
- Fasting is defined as at least 8 hours overnight before the visit, without food or liquids, except for water. Trial product and any medication which should be taken with or after a meal should be withheld on the day of the visit until blood samples have been obtained.
- At the ‘end of trial’ visit only 2 hours of fasting is required prior to the anti-semaglutide antibody sampling.
- If the subject is not fasting as required, the subject should be called in for a new visit within the visit window to have the fasting procedures done. Procedures requiring subject to fast include blood sampling of fasting plasma glucose (FPG), fasting serum insulin and free fatty acids.

6.4.2 Caffeine and tobacco

Subject should avoid caffeine and smoking at least 30 minutes prior to measuring the blood pressure.

6.5 Screen failures

Screen failures are defined as subjects who consent to participate in the clinical trial but are not eligible for participation according to in/exclusion criteria or randomisation criteria. A minimal set of screen failure information is required to ensure transparent reporting of screen failure subjects to meet requirements from regulatory authorities. Minimal information includes date of informed consent, date of visit, demography, screen failure details, eligibility criteria, and any serious adverse event (SAE). A screen failure session must be made in the interactive web response system (IWRS).

Individuals who do not meet the criteria for participation in this trial may not be rescreened. Re-sampling is not allowed if the subject has failed one of the inclusion criteria or fulfilled one of the exclusion criteria related to laboratory parameters. However, in case of technical issues (e.g. haemolysed or lost), re-sampling is allowed for the affected parameters.

7 Treatments

7.1 Treatments administered

All trial products listed in [Table 7-1](#) are considered investigational medicinal products (IMP). Trial product must only be used, if it appears clear and colourless.

Table 7-1 Trial products provided by Novo Nordisk A/S

| | | |
|---------------------------------|---|--|
| Trial product name: | Semaglutide B 1.0 mg/mL PDS290 or Semaglutide placebo* | Semaglutide B 3.0 mg/mL PDS290 or Semaglutide placebo |
| Dosage form: | Solution for injection | Solution for injection |
| Route of administration: | Subcutaneous | Subcutaneous |
| Dosing instructions: | Once-weekly | Once-weekly |
| Delivery device | 3 mL PDS290 pre-filled pen-injector | 3 mL PDS290 pre-filled pen-injector |

* Semaglutide B 1.0 mg/mL PDS290/semaglutide placebo will only be dispensed at the first dispensing visit.

- Dose escalation of semaglutide/semaglutide placebo should take place during the first 16 weeks after randomisation as described in [Table 7-2](#). All subjects should aim at reaching the recommended target dose of 2.4 mg semaglutide once-weekly or the corresponding volume of placebo.
- If a subject does not tolerate the recommended target dose of 2.4 mg once-weekly, the subject may stay at the lower dose level of 1.7 mg once-weekly. This should only be allowed if the subject would otherwise discontinue trial product completely and if considered safe to continue on trial product, as per the investigator's discretion. It is recommended that the subject makes at least one attempt to re-escalate to the recommended target dose of 2.4 mg once-weekly, as per the investigator's discretion.

- It is recommended that the investigator consults Novo Nordisk in case of persistent deviations from the planned escalation regimen.
- A dose reminder card will be handed out to the subjects at each site visit during the dose escalation period. This is to remind the subjects of the dose to be taken until next site visit and provide a conversion of the dose to value shown in the dose counter. Once the target dose has been reached, the dose reminder card is only handed out as needed.

Table 7-2 Dose escalation and maintenance of semaglutide/semaglutide placebo

| Trial product name | Dose | Volume | Value shown in dose counter* | Duration |
|---|---------|---------|------------------------------|----------|
| Dose escalation period | | | | |
| Semaglutide B 1.0 mg/mL PDS290 or semaglutide placebo | 0.25 mg | 0.25 ml | 25* | 4 weeks |
| Semaglutide B 1.0 mg/mL PDS290 or semaglutide placebo | 0.5 mg | 0.50 ml | 50* | 4 weeks |
| Semaglutide B 3.0 mg/mL PDS290 or semaglutide placebo | 1.0 mg | 0.34 ml | 34* | 4 weeks |
| Semaglutide B 3.0 mg/mL PDS290 or semaglutide placebo | 1.7 mg | 0.57 ml | 57* | 4 weeks |
| Maintenance period | | | | |
| Semaglutide B 3.0 mg/mL PDS290 or semaglutide placebo | 2.4 mg | 0.80 ml | 80* | 52 weeks |

* Conversion to dose is calculated based on 0.01 ml/value for both strengths.

- Subjects will be instructed to inject semaglutide/semaglutide placebo once-weekly at the same day of the week (to the extent possible) throughout the trial.
- Injections may be administered in the thigh, abdomen or upper arm, at any time of day irrespective of meals. Subjects should be encouraged to inject in the same area throughout the trial, but changing between left and right side is allowed.
- If a single dose of trial product is missed, it should be administered as soon as noticed, provided the time to the next scheduled dose is at least 2 days (48 hours). If a dose is missed and the next scheduled dose is less than 2 days (48 hours) away, the subject should not administer a dose until the next scheduled dose. A missed dose should not affect the scheduled dosing day of the week.
- If ≥ 2 consecutive doses of trial product are missed, the subject should be encouraged to re-commence the treatment if considered safe as per the investigator's discretion and if the subject does not meet any of the discontinuation criteria (Section [8.1](#)). The trial product should be continued as early as the situation allows. The missed doses should not affect the scheduled dosing day of the week. The start dose for re-initiation of trial product is at the investigator's discretion. In case of questions related to re-initiation of trial product, the investigator should consult Novo Nordisk global medical expert.

- Auxiliary supplies will be provided in accordance with the trial materials manual (TMM) please see [Table 7-3](#).

Table 7-3 Auxiliary supplies provided by Novo Nordisk A/S

| Auxiliary supply | Details |
|-------------------------|---|
| Needles | Needles for pre-filled pen system. Details provided in the TMM. Only needles provided and approved by Novo Nordisk must be used for administration of trial product. |
| Direction for use (DFU) | DFU for 3 ml PDS290 pre-filled pen-injector Not included in the dispensing unit and to be handed out separately |

7.1.1 Medical device

Information about the PDS290 pre-filled pen-injector may be found in the IB⁶⁹ and any updates hereof.

Information about the use of the PDS290 pre-filled pen-injector for semaglutide 1.0 mg/mL, semaglutide 3.0 mg/mL and semaglutide placebo can be found in the DFU.

Training in the PDS290 pre-filled pen-injector

The investigator must document that training in the DFU has been given to the subjects orally and in writing at the first dispensing visit. Training must be repeated, during the trial at regular intervals in order to ensure correct use of the medical device. Training is the responsibility of the investigator or a delegate.

7.1.2 Diet and physical activity counselling

All subjects in both treatment arms will receive counselling with regards to diet (500 kcal deficit per day relative to the estimated total energy expenditure (TEE) calculated once at randomisation) and physical activity (150 min of physical activity per week is encouraged, e.g. walking or use the stairs). Counselling should be done by a dietician or a similar qualified healthcare professional every 4th week via visits/phone contacts. Subjects will be instructed to record their food intake and physical activity daily (via paper diary, app or similar tool) in order to assist and evaluate their lifestyle intervention.

After randomisation the subjects can use a tool of their own choices (paper/app/other tool) for recording, ensuring it can be reviewed for content and number of entries during diet and physical activity counselling. Subjects must receive instructions in how to capture their physical activity and food intake. The subjects should keep diary of physical activity and food intake on a daily basis.

Calculation of estimated TEE

The TEE is calculated by multiplying the estimated Basal Metabolic Rate (BMR) ([Table 7-4](#)) with a Physical Activity Level value of 1.3.⁷⁰

$$TEE = BMR \times 1.3$$

Table 7-4 Equation for estimated BMR

| Sex | Age | BMR (kcal/day) |
|-------|-------------|--|
| Men | 18-30 years | $15.057 \times \text{weight at randomisation in kg} + 692.2$ |
| | 31-60 years | $11.472 \times \text{weight at randomisation in kg} + 873.1$ |
| | > 60 years | $11.711 \times \text{weight at randomisation in kg} + 587.7$ |
| Women | 18-30 years | $14.818 \times \text{weight at randomisation in kg} + 486.6$ |
| | 31-60 years | $8.126 \times \text{weight at randomisation in kg} + 845.6$ |
| | > 60 years | $9.082 \times \text{weight at randomisation in kg} + 658.5$ |

If a BMI ≤ 22.5 kg/m² is reached the recommended energy intake should be recalculated with no kcal deficit (maintenance diet) for the remainder of the trial. If deemed necessary the investigator could consult Novo Nordisk to discuss when maintenance diet can be initiated.

7.2 Dose modification

- Not applicable for this trial. Please refer to Section [7.1](#) for description of missed dose(s).

7.3 Method of treatment assignment

- All subjects will be centrally randomised using an IWRS and assigned to the next available treatment according to randomisation schedule. Trial product will be dispensed at the trial visits summarised in the flowchart.

7.4 Blinding

The active drug and placebo are visually identical for the following trial products:

- Semaglutide B 1.0 mg/mL PDS290/semaglutide placebo
- Semaglutide B 3.0 mg/mL PDS290/semaglutide placebo

The IWRS is used for blind-breaking instructions. The blind may be broken in a medical emergency if knowing the actual treatment would influence the treatment of the subject. Novo Nordisk will be notified immediately after breaking the blind. The date when and reason why the blind was broken must be recorded in the subject's medical record.

Whenever the blind is broken, the person breaking the blind must print the "code break confirmation" notification generated by the IWRS, and sign and date the document.

When the blind is broken, the treatment allocation will be accessible to the investigator and the Novo Nordisk Global Safety department. If IWRS is not accessible at the time of blind break, the IWRS helpdesk should be contacted. Contact details are listed in [Attachment I](#).

7.5 Preparation/Handling/Storage/Accountability

Only subjects enrolled in the trial may receive trial product and only authorised site staff may supply or administer trial product.

Table 7-5 Trial product storage conditions

| Trial product name | Storage conditions (not-in-use) | In-use conditions | In-use time ^a |
|-----------------------------------|--|--|--|
| Semaglutide B 1.0 mg/mL PDS290 | Store in refrigerator (2°C-8°C/36°F-46°F) | In-use conditions will be available on the trial product label | In-use time will be available on the trial product label |
| Semaglutide placebo | Do not freeze Protect from light | | |
| Semaglutide B 3.0 mg/mL PDS290 | Store in refrigerator (2°C-8°C/36°F-46°F) | In-use conditions will be available on the trial product label | In-use time will be available on the trial product label |
| Semaglutide placebo | Do not freeze Protect from light | | |

^aIn-use time starts when the product is taken out of the refrigerator in the subject's home.

- Each trial site will be supplied with sufficient trial products for the trial on an on-going basis controlled by the IWRS. Trial product will be distributed to the trial sites according to the number of subjects screened and randomised.
- The investigator must confirm that appropriate temperature conditions have been maintained during transit for all trial products received (see [Table 7-5](#)) and any discrepancies are reported and resolved before use of the trial products.
- All trial products must be stored in a secure, controlled, and monitored (manual or automated) area in accordance with the labelled storage conditions with access limited to the investigator and authorised site staff.
- The investigator must inform Novo Nordisk immediately if any trial product has been stored outside specified conditions. Additional details regarding handling of temperature deviations can be found in the TMM.
- Trial product that has been stored improperly must not be dispensed to any subject before it has been evaluated and approved for further use by Novo Nordisk.
- The investigator is responsible for drug accountability and record maintenance (i.e. receipt, accountability and final disposition records).
- Drug accountability should be performed on a pen level and must be documented in the IWRS.
- The subject must return all used, partly used and unused trial product including empty packaging materials during the trial as instructed by the investigator.
- Destruction of trial products can be performed on an ongoing basis and will be done according to local procedures after accountability is finalised by the site and reconciled by the monitor.
- Destruction of trial products must be documented in the IWRS.

- All returned, expired or damaged trial products (for technical complaint samples see [Appendix 6](#)) must be stored separately from non-allocated trial products. No temperature monitoring is required.
- Non-allocated trial products including expired or damaged products must be accounted as unused, at the latest at closure of the trial site.

7.6 Treatment compliance

Throughout the trial, the investigator will remind the subjects to follow the trial procedures and requirements to ensure subject compliance. If a subject is found to be non-compliant the investigator will remind the subject of the importance of following the instructions given including taking the trial products as prescribed.

Treatment compliance of trial product will be assessed by asking subject about missed doses and monitoring of PK dosing diaries. Information about compliance and missed doses should be described in the subject's medical record.

7.7 Concomitant medication

Any medication (including over-the-counter or prescription medicines) other than the trial product that the subject is receiving at the time of the first visit or receives during the trial must be recorded along with:

- Trade name or generic name
- Indication
- Dates of administration including start and stop dates
- Dose (only to be recorded for anti-hypertensive and lipid-lowering medication)

During the trial subjects should not initiate any anti-obesity treatment (e.g. medication) which is not part of the trial procedures. If such treatment is initiated, the subject should be instructed to stop the anti-obesity treatment.

Changes in concomitant medication must be recorded at each visit. If a change is due to an AE, then this must be reported according to Section [9.2](#).

7.7.1 Rescue medication

This section is not applicable for this trial.

7.8 Treatment after the end of the trial

After the end of the trial the subject should be treated at the discretion of the investigator.

8 Discontinuation/Withdrawal criteria

The subject may be discontinued at any time during the trial at the discretion of the investigator for safety, behavioural, compliance or administrative reasons.

Efforts must be made to have the subjects, who discontinue trial product, to continue in the trial. Subjects must be educated about the continued scientific importance of their data, even if they discontinue trial product. Only subjects who withdraw consent will be considered as withdrawn from the trial.

8.1 Discontinuation of trial treatment

- Discontinuation of treatment can be decided by either the investigator or the subject.
- Subjects who discontinue trial product should continue with the scheduled visits and assessments to ensure continued counselling and data collection.
 - If the subject does not wish to attend the scheduled clinic visits, efforts should be made to have the visits converted to phone contacts. However all efforts should be made to have the subject attend at least the 'end of treatment' clinic visit containing the final data collection of primary and confirmatory secondary efficacy endpoints, and the 'end of trial' visit.
- The 'end of trial' visit is scheduled approximately 7 weeks after the final data collection to ensure the safety of the subject.
 - If the subject refuses to attend the 'end of treatment' and/or 'end of trial' visit, information about the attempts to follow up with the subject must be documented in the subject's medical record.

The subject must be discontinued from trial product, if any of the following applies:

1. Included in the trial in violation of the inclusion and/or exclusion criteria and/or randomisation criteria
2. Safety concern as judged by the investigator
3. Calcitonin ≥ 100 ng/L (see [Appendix 8](#))
4. Suspicion of pancreatitis
5. Pregnancy
6. Intention of becoming pregnant
7. Simultaneous participation in another clinical trial of an approved or non-approved investigational medicinal product.

If acute pancreatitis is suspected, appropriate actions should be initiated, including local measurement of amylase and lipase (see [Appendix 4](#) for reporting).

Subjects meeting discontinuation of trial product criterion no. 4 are allowed to resume trial product if the Atlanta criteria⁷¹ are not fulfilled and thus the suspicion of acute pancreatitis is not confirmed.

Trial product may be resumed for subjects with a gallstone-induced pancreatitis in case of cholecystectomy.

Subjects meeting discontinuation of trial product criteria no. 2, 5 and 6 are allowed to resume trial product, if the criteria are no longer met (see Section [8.1.1](#)).

A subject who does not fulfil the eligibility criteria must not be randomised. If a subject is randomised in violation of inclusion, exclusion or randomisation criteria, this will be handled as an important protocol deviation, and the IEC/IRB and regulatory authorities must be notified according to local requirements.

The primary reason for discontinuation of trial product must be specified in the source data at the time of discontinuation, and subject should continue to follow the visit and assessment schedule. A change in 'treatment status' must be made in IWRS to discontinue trial product. If subject is not allowed to resume trial product, the reason for discontinuation will be recorded in the 'end of treatment' form in the CRF, and final drug accountability must be performed.

8.1.1 Temporary discontinuation of trial treatment

If a subject has discontinued trial product due to temporary safety concern not related to trial product and is allowed to resume, the subject should follow the guide for missed doses (Section [7.1](#)). Similarly, a subject who discontinues trial product on their own initiative should be encouraged to resume trial product (Section [7.1](#)).

Each missed dose should be recorded in the CRF, as per subject's recollection. If a 'treatment status' session previously has been made in IWRS to indicate discontinuation of trial product, a new "treatment status" session must be made to resume trial product.

8.2 Withdrawal from the trial

A subject may withdraw consent at any time at his/her own request.

If a subject withdraws consent, the investigator must ask the subject if he/she is willing, as soon as possible, to have assessments performed according to the 'end of treatment' visit. See the flowchart for data to be collected.

Final drug accountability must be performed even if the subject is not able to come to the trial site. The investigator must make a 'treatment status' session in IWRS to discontinue trial product.

If a subject withdraws from the trial, he/she may request destruction of any samples taken and not tested, and the investigator must document this in the medical record.

If the subject withdraws consent, Novo Nordisk may retain and continue to use any data collected before such a withdrawal of consent.

Although a subject is not obliged to give his/her reason(s) for withdrawing, the investigator must make a reasonable effort to ascertain the reason(s), while fully respecting the subject's rights. Where the reasons are obtained, the primary reason for withdrawal must be specified in the 'end of trial' form in the CRF.

8.2.1 Replacement of subjects

Subjects who discontinue trial product or withdraw from trial will not be replaced.

8.3 Lost to follow-up

A subject will be considered lost to follow-up if he or she repeatedly fails to return for scheduled visits and is unable to be contacted by the trial site.

The following actions must be taken if a subject fails to return to the trial site for a required visit:

- The site must attempt to contact the subject and reschedule the missed visit as soon as possible and counsel the subject on the importance of maintaining the assigned visit schedule and ascertain whether or not the subject wishes to and/or should continue in the trial.
- Before a subject is deemed lost to follow-up, the investigator must make every effort to regain contact with the subject (where possible, at least three telephone calls and, if necessary, a certified letter to the subject's last known mailing address or local equivalent methods). If attempts have failed, family members or other contacts consented by the subject can be contacted for alternative contact details. These contact attempts should be documented in the subject's source document.
- Should the subject continue to be unreachable at 'end of treatment' visit, he/she will be considered to have withdrawn from the trial with a primary reason of 'lost to follow-up'.

9 Trial assessments and procedures

- Trial procedures and their timing are summarised in the flowchart.
- Informed consent must be obtained before any trial related activity, see [Appendix 3](#).
- All screening evaluations must be completed and reviewed to confirm that potential subjects meet all eligibility criteria.
- The investigator will maintain a screening log to record details of all subjects screened and to confirm eligibility or record reason for screen failure, as applicable.
- At screening, subjects will be provided with a card stating that they are participating in a trial and giving contact details of relevant trial site staff.
- Adherence to the trial design requirements, including those specified in the flowchart, is essential and required for trial conduct.
- Assessments should be carried out according to the clinic's standard of practice unless otherwise specified in the current section. Efforts should be made to limit the bias between assessments. The suggested order of the assessments:

1. Electrocardiogram (ECG) and vital signs
 2. Blood samples
 3. Patient reported outcome questionnaires (see Section [9.1.2](#)) and mental health assessment instruments (see Section [9.4.1](#))
 4. Other assessments
- Source data of clinical assessments performed and recorded in the CRF must be available and will usually be the subject's medical records. Additional recording to be considered source data includes, but is not limited to laboratory reports, ECG, diary recordings, clinical outcome assessments. Ensure to transcribe into the CRF :
 - Diary:
 - Compliance with diet from screening to randomisation according to the randomisation criteria
 - Dosing information prior to PK sampling
 - Number of active minutes per week from physical activity diary
 - Number of days per week with at least one entry in the food diary
 - Evaluations of ECG
 - Subject must receive instructions in how to capture their daily food intake in the handed out food diary from screening to randomisation. Entries must be evaluated in accordance with the randomisation criteria.
 - Subjects must receive training in how to collect dosing information prior to PK sampling in a designated paper diary.
 - Only the subject can make entries and corrections in the diaries, unless the section is specified for site staff.
 - The barriers and motivation interview identifies barriers to and motivation for lifestyle change and compliance with the protocol. The interview will be conducted at screening to assist in identifying subjects who are unable or unwilling to comply with protocol procedures as per the exclusion criteria. In addition, the interview will ensure that any minor barriers are addressed during lifestyle counselling.
 - The results of the interview will not be entered into the CRF. It will be at the investigator's discretion to evaluate the motivation of the subject and related eligibility.
 - Subject's weight history must be recorded in the subject's medical record.
 - Review of diaries, mental health assessment instruments, ECG, and laboratory reports must be documented either on the documents or in the subject's medical record. If clarification of entries or discrepancies is needed, the subject must be questioned and a conclusion made in the subject's source documents. Care must be taken not to bias the subject.
 - Repeat laboratory samples may be taken for technical issues and unscheduled samples or assessments may be taken for safety reasons. Please refer to [Appendix 2](#) for further details on laboratory samples.
 - The fatty liver index is based on an algorithm including BMI, waist circumference, triglycerides and gamma-glutamyl-transferase (GGT).

- For subjects receiving antihypertensive or lipid-lowering treatment, the investigator should evaluate changes in the subject's treatment intensity within each therapeutic area. The evaluation should be based on whether an overall change from randomisation until the time of the evaluation has occurred (i.e., either increase, decrease or no change) after reviewing all available relevant information e.g., changes in drug dose, drug class, number of drugs or a combination of these.
- The investigator will evaluate the subject's glycaemic status periodically during the trial as detailed in the flowchart based on all available relevant information e.g. medical records, concomitant medication, blood glucose parameters (HbA1c, FPG) and AEs. The subject's glycaemic status will be categorised as normo-glycaemia, prediabetes or diagnosed with type 2 diabetes according to the American Diabetes Association's definitions⁷².
- For subjects in the DEXA sub-population, DEXA scan must be performed according to the manual from the supplier.

9.1 Efficacy assessments

Planned time points for all efficacy assessments are provided in the flowchart.

9.1.1 Body measurements

- Body weight should be measured at all site visits without shoes, on an empty bladder and only wearing light clothing. It should be measured on a digital scale and recorded in kilograms or pounds (one decimal) using the same scale throughout the trial.
- The scale must be calibrated yearly as a minimum.
- Height is measured without shoes in centimetres or inches (one decimal). BMI will be calculated by the CRF from screening data and must be in agreement with inclusion criterion no. 3.
- Waist circumference is defined as the abdominal circumference located midway between the lower rib margin and the iliac crest.
- Measures must be obtained in standing position with a non-stretchable measuring tape and to the nearest cm or inch.
- The tape should touch the skin but not compress soft tissue and twists in the tape should be avoided. The subject should be asked to breathe normally. The same measuring tape should be used throughout the trial. The measuring tape will be provided by Novo Nordisk to ensure standardisation.

9.1.2 Clinical outcome assessments

Subjects should be given the opportunity to complete the questionnaires by themselves without interruption. Each of the questionnaires takes approximately 10 minutes to complete.

The following patient reported outcome questionnaires will be used:

- Impact of Weight on Quality of Life-Lite for Clinical Trials version (IWQoL-Lite for CT), version 3.0
 - The IWQoL-Lite for CT is a 20-item modified version of a questionnaire tool designed to assess the weight-related quality of life.⁷³
- Patient Global Impression of Status (PGI-S) (IWQoL-Lite for CT) for physical function, version 1.0
- Patient Global Impression of Change (PGI-C) (IWQoL-Lite for CT) for physical function, version 1.0
- Short Form 36 v2.0 acute (SF-36)
 - SF-36 measures the subject's overall health related quality of life. It is a 36-item generic measure of health status that yields 2 summary scores for physical health and mental health, and 8 domain scores⁷⁴
- Stanford Presenteeism Scale (SPS-6), 2001 version
 - The SPS-6 evaluates workforce productivity. Absenteeism affects productivity; however, even when employees are physically present at their jobs, they may experience decreased productivity and below-normal work quality - a concept known as decreased presenteeism.⁷⁵
- International Consultation on Incontinence Questionnaire-Urinary Incontinence-Short Form (ICIQ-UI-SF)
 - The ICIQ-UI-SF assesses the symptoms and impact of urinary incontinence.⁷⁶

9.1.3 Clinical efficacy laboratory assessments

All protocol-required laboratory assessments, as defined in [Appendix 2](#), must be conducted in accordance with the flowchart and the laboratory manual.

9.2 Adverse events

The definitions of AEs and SAEs can be found in [Appendix 4](#).

The investigator is responsible for detecting, documenting, recording and following up on events that meet the definition of an AE or SAE.

9.2.1 Time period and frequency for collecting AE and SAE information

All AEs will be collected from the first trial-related activity after obtaining informed consent and until 'end of trial' visit, at the time points specified in the flowchart.


All SAEs will be recorded and reported to Novo Nordisk or designee within 24 hours, as indicated in [Appendix 4](#). The investigator must submit any updated SAE data to Novo Nordisk within 24 hours of it being available.

Investigators are not obligated to actively seek for AE or SAE in former trial subjects. However, if the investigator learns of any SAE, including a death, at any time after a subject has been discontinued from/completed the trial, and the investigator considers the event to be possibly/probably related to the investigational trial product or trial participation, the investigator must promptly notify Novo Nordisk.

The method of recording, evaluating and assessing causality of AE and SAE and the procedures for completing and transmitting SAE reports are provided in [Appendix 4](#).

Timelines for reporting of AEs including events for adjudication, Section [9.2.1.1](#), are listed in [Figure 9-1](#).

Some AEs require additional data collection via a specific event form. This includes medication errors observed during the trial. The relevant specific events are listed in [Table 9-1](#) and the reporting timelines in [Figure 9-1](#).



Timelines are from the awareness of an AE.
Queries to be resolved ≤ 14 calendar days.

AE: Adverse Events, SAE: Serious Adverse Events, SIF: Safety Information Form

Figure 9-1 Decision tree for determining the event type and the respective forms to complete with associated timelines

Table 9-1 AEs requiring additional data collection (via specific event form) and events for adjudication

| Event type | AE via specific event form | Event for adjudication (Section 9.2.1.1) |
|-----------------------------------|----------------------------|---|
| Medication error | X | |
| Death | | X |
| Cardiovascular events | | |
| Acute Coronary Syndrome | | X |
| Cerebrovascular event | | X |
| Heart failure | | X |
| Coronary artery revascularisation | | X |
| Acute pancreatitis | X | X |
| Acute gallbladder disease | X | |
| Malignant neoplasms | X | |
| Hepatic event | X | |

9.2.1.1 Events for adjudication

Event adjudication will be performed for adverse events in randomised subjects.

These events are reviewed by an independent external event adjudication committee in a blinded manner, refer to [Appendix 3](#) for further details.

There are 4 ways to identify events relevant for adjudication as described below:

- Investigator-reported events for adjudication: When reporting AEs, the investigator must select the appropriate AE category based on pre-defined criteria (see [Table 9-1](#) and [Appendix 4](#))
- Death
- AE Search (standardised screening): All AEs not directly reported by the investigator as requiring adjudication, will undergo screening to identify potential events for adjudication. The investigator can be queried to provide additional information related to the reported AE, e.g. alternative aetiology, underlying cause(s) and/or clinical details.
- EAC-identified events: When reviewing source documents provided for another event for adjudication, the EAC can identify additional events in scope for adjudication that were not initially reported by the investigator. In these instances, the investigator will be notified of the newly identified event and has the option to report the EAC-identified event. Regardless of whether the investigator decides to report the event, it will undergo adjudication. Occasionally, EAC-identified events may require the investigator to collect additional source documents, which should be provided by uploading to the event adjudication system (EAS).

With the exception of EAC-identified events, an event-specific adjudication form for the event in question should be completed in the CRF within 14 calendar days of the investigator's first knowledge of the event.

Copies of collected source documents should be labelled with trial ID, subject and AE number, redacted (anonymised of personal identifiers) and uploaded to the EAS within 4 weeks according to instructions outlined in the event adjudication site manual. If no, or insufficient source documents are provided to the adjudication supplier, the investigator can be asked to complete a clinical narrative to be uploaded to the EAS.

If new information becomes available for an event sent for adjudication, it is the responsibility of the investigator to ensure the new information is uploaded to the EAS.

9.2.2 Method of detecting AEs and SAEs

Care should be taken not to introduce bias when detecting AEs and/or SAEs. Open-ended and non-leading verbal questioning of the subject is the preferred method to inquire about events.

9.2.3 Follow-up on AEs and SAEs

After the initial AE/SAE report, the investigator is required to proactively follow each subject at subsequent visits/contacts. All SAEs will be followed until resolution, stabilization, or if the event is otherwise explained (e.g. chronic condition) or the subject is lost to follow-up (as defined in Section [8.3](#)). Further information on follow-up procedures is given in [Appendix 4](#).

9.2.4 Regulatory reporting requirements for SAEs

Prompt notification by the investigator to Novo Nordisk of a SAE is essential so that legal obligations and ethical responsibilities towards the safety of subjects and the safety of a trial product under clinical investigation are met.

Novo Nordisk has a legal responsibility to notify both the local regulatory authority and other regulatory agencies about the safety of a trial product under clinical investigation. Novo Nordisk will comply with country-specific regulatory requirements relating to safety reporting to the regulatory authority, IRB/IEC, and investigators.

Investigator safety reports must be prepared for SUSARs according to local regulatory requirements and Novo Nordisk policy and forwarded to investigators as necessary.

An investigator who receives an investigator safety report describing a SAE or other specific safety information (e.g. summary or listing of SAEs) from Novo Nordisk, will review and then file it along with the IB and will notify the IRB/IEC, if appropriate according to local requirements.

9.2.5 Hypoglycaemic, cardiovascular and death events

Hypoglycaemic, cardiovascular and death events will be handled and reported according to AE/SAEs description in Section [9.2.1](#).

9.2.6 Disease-related events and/or disease-related outcomes not qualifying as an AE or SAE

Not applicable for this trial.

9.2.7 Pregnancies and associated adverse events

Details of pregnancies in female subjects will be collected after the first-trial-related activity after obtaining informed consent and until the 'end of trial' visit.

If a pregnancy is reported in female subjects, the investigator should inform Novo Nordisk within 14 calendar days of learning of the pregnancy and should follow the procedures outlined in [Figure 9-2](#) and [Appendix 5](#).

Pregnancy outcome should be documented in the subject's medical record. Abnormal pregnancy outcome (e.g. spontaneous abortion, foetal death, stillbirth, congenital anomalies and ectopic pregnancy) is considered an SAE.




Figure 9-2 Decision tree for determining the forms to complete with associated timelines for pregnancy.

9.2.8 Medical device incidents (including malfunctions)


Section is not applicable for this trial. Refer to technical complaints in Section [9.2.9](#).

9.2.9 Technical complaints

The investigator must assess whether a technical complaint is related to an AE.

The definitions and reporting process for technical complaints can be found in [Appendix 6](#).

Timelines for reporting technical complaints are listed in [Figure 9-3](#).



AE: Adverse Event, SAE: Serious Adverse Event, SIF: Safety Information Form

Figure 9-3 Decision tree for determining the forms to complete with associated timelines for technical complaints.

9.3 Treatment of overdose

Overdoses of up to 4 mg in a single dose, and up to 4 mg in a week have been reported in clinical trials. The most commonly reported AE was nausea. All subjects recovered without complications.

There is no specific antidote for overdose with semaglutide. In the event of an overdose, appropriate supportive treatment should be initiated according to subject's clinical signs and symptoms.

The overdose must be reported as a medication error ([Appendix 4](#)) and for reporting timeline see Section [9.2.1](#).

In the event of an overdose, the investigator should closely monitor the subject for overdose-related AE/SAE and laboratory abnormalities. A prolonged period of observation and treatment may be necessary, taking into account the long half-life of semaglutide of approximately one week.

Decisions regarding dose interruptions or modifications will be made by the investigator based on the clinical evaluation of the subject.

For more information on overdose, also consult the current version of the IB^{S7} and any updates hereof.

9.4 Safety assessments

Planned time points for all safety assessments are provided in the flowchart.

A **concomitant illness** is any illness that is present at the start of the trial (i.e. at the first visit) or found as a result of a screening procedure or other trial procedures performed before exposure to trial product.

Medical history is a medical event that the subject has experienced in the past. Only relevant and significant medical history as judged by the investigator should be recorded. Findings of specific medical history should be described in designated forms.

As part of the medical history, information related to history of gallbladder disease, breast neoplasm, colon neoplasm, skin cancer, and psychiatric disorder will be recorded. Follow-up questions will be asked at the end of trial related to breast neoplasm and colon neoplasm

In case of an abnormal and clinically significant finding, the investigator must record the finding on the Medical History/Concomitant Illness form if it is present at screening. Any new finding fulfilling the AE definition (see [Appendix 4](#)) during the trial and any clinically significant worsening from baseline must be reported as an AE (see Section [9.2](#)).

9.4.1 Mental health assessment instruments

- PHQ-9⁷⁷ is a 9-item depression module of the patient health questionnaire, which is a self-administered diagnostic tool used for assessment of mental disorders. The questionnaire will be available in a linguistically validated translated version.
- C-SSRS⁷⁸ is a detailed questionnaire assessing both suicidal behaviour and suicidal ideation. The questionnaire will be administered as an interview by the investigator or a qualified delegate. The questionnaire (C-SSRS Baseline and C-SSRS Since Last Visit) will be available in a linguistically validated translated version.
 - Prior to administering the C-SSRS questionnaire, the investigator or qualified delegate must complete sufficient training.

If a subject has a PHQ-9 score of 10-14 (both inclusive) the subject should be referred to a mental health professional (MHP) if judged relevant by the investigator. If referral is not deemed relevant this along with the reason why must be documented in the subject's medical records.

A subject must be referred to a MHP if:

- the subject has a PHQ-9 score ≥ 15 or
- the subject has any suicidal behaviour or
- the subject has any suicidal ideation of type 4 or type 5 on any C-SSRS assessment or
- in the opinion of the investigator, it is necessary for the safety of the subject

If one or more of the referral criteria are met, the investigator should explain to the subject why the referral and psychiatric evaluation by a MHP is needed. If the subject refuses to be referred to a MHP, the subject's decision should be documented in subject's medical record and the investigator

must assess if it is safe for the subject to continue in the trial or if the subject should be discontinued from trial product.

If a subject's psychiatric disorder can be adequately treated with psychotherapy and/or pharmacotherapeutic treatment, then the subject, at the discretion of the investigator (and in agreement with the MHP), may continue in the trial. Otherwise, the subject must be discontinued from trial product due to safety concern as judged by the investigator.

9.4.2 Physical examinations

- A physical examination will include assessments of the general appearance, thyroid gland, breast (females), and abdomen, as well as the cardiovascular and respiratory systems.
- Investigators should pay special attention to clinical signs related to previous serious illnesses.

9.4.3 Vital signs

- The method for measuring systolic and diastolic blood pressure needs to follow the standard clinical practice at site.
- Blood pressure (diastolic and systolic) and pulse measurements should be preceded by at least 5 minutes of rest for the subject in a quiet setting without distractions (e.g. television, cell phones).
- Blood pressure and pulse measurements will be assessed in a sitting position with a completely automated device. Manual techniques will be used only if an automated device is not available.

9.4.4 Electrocardiograms

- 12-lead ECG will be obtained as outlined in the flowchart using an ECG machine that automatically calculates the heart rate and measures PR, QRS, QT, and QTc intervals.
- In a sub-population, ECGs will be collected for central storage at randomisation and 'end of treatment' visit for potential future assessment.

9.4.5 Clinical safety laboratory assessments

All protocol-required laboratory assessments, as defined in [Appendix 2](#), must be conducted in accordance with the flowchart and the laboratory manual.

- If the laboratory finding based on the results from central laboratory meets the criteria for laboratory outliers, a laboratory outlier form in the CRF should be completed. Please refer to [Appendix 2](#) for the criteria for the laboratory outliers.
- Urine pregnancy tests supplied by central laboratory must be performed for women of childbearing potential at screening and as specified in the flowchart. Urine pregnancy test must be repeated at any time during the trial if pregnancy is suspected. Further instructions can be found in the laboratory manual.

9.4.6 Immunogenicity assessments

Blood samples for determination of serum antibodies against semaglutide, including cross reactivity to endogenous GLP-1 will be taken during the trial at visits specified in the flowchart. Samples taken at the 'end of trial' visit which are positive for anti-semaglutide antibodies will be further characterised for in vitro neutralising effect towards semaglutide. In addition, if samples taken at the 'end of trial' visit are also positive for cross-reactivity against endogenous GLP-1, the samples will be analysed for in vitro neutralising effect towards endogenous GLP-1. The results of the analysis will only be disclosed after completion of the clinical trial report (CTR) if required by local regulation.

9.5 Pharmacokinetics

- Single blood samples for measuring plasma concentration of semaglutide will be drawn for both semaglutide and semaglutide placebo subjects on visits specified in the flowchart.
- Subject must be instructed to withhold their trial product dose in the morning of the clinic visit until blood sampling has been performed.
- The PK dosing information should be transcribed into the CRF for the last 2 doses of trial product, including time, date and injection site.
- The exact timing of obtaining the PK sample must be recorded on the laboratory form.
- The dual purpose of measuring plasma semaglutide levels is to perform population PK (Pop-PK) analyses and to assess the level of drug interference in the anti-semaglutide antibody analysis.

Samples will be used to evaluate the pharmacokinetics of semaglutide. Each plasma sample will be divided into 2 aliquots (e.g. one for pharmacokinetics and a backup,) and may also be used to evaluate safety or efficacy aspects that address concerns arising during or after the trial. Residual sample material may be used for exploratory investigation of metabolites and bioanalysis assay development and troubleshooting in relation to the pharmacokinetic assay.

9.6 Pharmacodynamics

Not applicable for this trial.

9.7 Genetics

Not applicable for this trial.

9.8 Biomarkers

Collection of samples for biomarker research is part of this trial to support the effect objectives. The following samples must be conducted in accordance with the laboratory manual and the flowchart:

Biomarkers linked to cardiovascular risk:

- PAI-1 Activity will be analysed by activity assay.

- hsCRP

Biomarkers linked to hunger and satiety:

- Leptin
- Soluble leptin receptor

9.9 Severe hypersensitivity

In the event of a severe immediate hypersensitivity reaction to trial product, blood sampling for assessment of anti-semaglutide IgE and binding antibodies should be conducted after 1–2 weeks and 7 weeks of trial product wash-out (i.e. after the subject had the last dose of the trial product).

In these cases, it is also recommended to test for tryptase (total and/or mature tryptase) within 3 hours of the hypersensitivity reaction. In case a tryptase sample was collected within 3 hours of the event of hypersensitivity reaction, a baseline tryptase sample should be taken at the same time as the IgE sample is obtained (after 1-2 weeks of trial product wash-out). Tryptase concentrations (if measured) as well as results of anti-semaglutide antibody and IgE isotype anti-semaglutide antibodies will be collected by Novo Nordisk and the results will be reported in the CTR.

10 Statistical considerations

Taxonomy of week 68 assessments

For each subject a given assessment at week 68 may be available or missing and [Table 10-1](#) describes the taxonomy for this. Note, this is done per assessment and per subject; subjects may be a different type for different assessments (a subject may have “available on randomised treatment (AT)” for body weight but “missing on randomised treatment (MT)” for waist circumference).

Table 10-1 Taxonomy for subjects based on week 68 assessments

| Assessment at week 68 | Subjects on randomised treatment at week 68 | Type description | Type Abbreviation |
|-----------------------|---|--|-------------------|
| Available | Yes | Available on randomised treatment: Subjects who complete the trial on randomised treatment with an assessment at week 68. Includes those that stop and restart trial product. | AT |
| | No | Available but discontinued Subjects who discontinued randomised treatment prematurely but returned to have an assessment at week 68. These are also called retrieved subjects | AD |
| Missing | Yes | Missing on randomised treatment: Subjects who complete the trial on randomised treatment without an assessment at week 68. Includes those that stop and restart trial product. | MT |
| | No | Missing and discontinued: Subjects who discontinued randomised treatment prematurely and did not return to have an assessment at week 68. These are also called non-retrieved subjects | MD |

10.1 Sample size determination

The sample size and thereby the power for this trial is primarily defined to support safety. However, no formal statistical inference is planned based on number of adverse events. Given the trial sample size, the power of statistical tests for effect endpoints is described below.

The tests of superiority of semaglutide 2.4 mg to semaglutide placebo for the primary and confirmatory secondary endpoints are performed using the fixed-sequence statistical strategy. This strategy tests the endpoints using a predefined hierarchical order, all at the significance level of 5%, moving to test the next endpoint only after a statistically significant superiority result (p -value < 5%) on the previous endpoint. The test hierarchy is given in [Table 10-2](#) with underlying assumptions, marginal power and effective power. The effective power is calculated under the assumption of independence of endpoints by multiplying the respective marginal powers successively. As the two primary endpoints are included in the statistical testing hierarchy, significant superiority of semaglutide 2.4 mg vs. semaglutide placebo must be demonstrated for each of the primary endpoints.

In the analysis approach addressing the primary estimand, week 68 assessments from retrieved subjects (AD) are used. These data are also used to impute missing measurements at week 68 for non-retrieved subjects (MD). The imputation is done separately within each treatment arm (see description below). However, for the power calculations missing values (MT and MD), regardless of treatment arm, are assumed to be similar to semaglutide placebo subjects. These assumptions are likely conservative with respect to the power, and correspond to the jump to reference sensitivity analysis planned below.

Assumptions

The common assumptions for the power calculations are:

- The significance level is 5%
- The randomisation ratio is 2:1
- For continuous endpoints the t-test on the mean difference assuming equal variances is used
- For binary endpoints the Pearson chi-square test for two independent proportions is used
- Based on data from NN9536-4153
 - 20% of subjects discontinue permanently and
 - 60% of these are retrieved (AD) at week 68
- All subjects in the semaglutide placebo arm are assumed to have same effect as subjects who complete the trial on semaglutide placebo (AT)
- Retrieved subjects (AD) in the semaglutide 2.4 mg arm are assumed to have an effect corresponding to half the treatment difference (compared to semaglutide placebo) of subjects who complete the trial on semaglutide 2.4 mg (AT)
- Non-retrieved subjects (MD) in the semaglutide 2.4 mg arm are assumed to have an effect corresponding to semaglutide placebo

Further assumptions made to calculate the power for each of the primary and confirmatory secondary endpoints are based on findings from other projects conducted by Novo Nordisk (NN8022 (SCALE), NN9535 (SUSTAIN), NN9924 (PIONEER)), and trial NN9536-4153 and are presented in [Table 10-2](#).

Given these assumptions, the sample size of 1950 subjects (1300 in the semaglutide 2.4 mg once-weekly and 650 in the semaglutide placebo arm), gives an effective power (marginal powers multiplied) of 99% for the first seven endpoints in the hierarchical testing procedure. As sample size is primarily driven by safety, additional scenarios for assumptions are not included due to the overall high power.

Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects

| Order | Endpoint | Assumed mean (\pm SD) or proportion for completers | | Expected mean (\pm SD) or proportion Semaglutide 2.4 mg | Expected difference or proportion ratio | Marginal power (%) | Effective power (%) |
|-------|-----------------------------|--|---------------------|--|---|--------------------|---------------------|
| | | Semaglutide 2.4 mg | Semaglutide placebo | | | | |
| 1 | % weight change # | 14 (\pm 10) | 3 (\pm 10) | 12.5 (\pm 11) | 9.5%-points | > 99 | > 99 |
| 2 | 5% responders | 82% | 42% | 76% | 1.8 | > 99 | > 99 |
| 3 | 10% responders | 66% | 24% | 60% | 2.5 | > 99 | > 99 |
| 4 | 15% responders | 46% | 12% | 41% | 3.4 | > 99 | > 99 |
| 5 | WC change (cm) # | 11 (\pm 10) | 4 (\pm 10) | 10 (\pm 10) | 6 cm | > 99 | > 99 |
| 6 | sBP change (mmHg) # | 6.2 (\pm 13) | 1.5 (\pm 13) | 5.5 (\pm 13) | 4 mmHg | > 99 | > 99 |
| 7 | SF-36 PF score change | 6 (\pm 10) | 2 (\pm 10) | 5.4 (\pm 11) | 3.4 score-points | > 99 | > 99 |
| 8 | IWQoL-Lite PFD score change | To be confirmed. Is currently being validated in trials N9536-4153 and NN9924-4233 | | | | | |

SD = standard deviation; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain; # shown as a positive number

10.2 Definition of analysis sets

Two analysis sets are defined:

- The *full analysis set* (FAS) includes all randomised subjects according to the intention-to-treat principle.
- The *safety analysis set* (SAS) includes all randomised subjects exposed to at least one dose of randomised treatment.

Any observation excluded from the analysis will be documented before database lock with the reason for exclusion provided.

Two observation periods are defined for each subject

- In-trial: The *in-trial period* is defined as the uninterrupted time interval from date of randomisation to date of last contact with trial site.
- On-treatment (with trial product): A time-point is considered as 'on-treatment' if any dose of trial product has been administered within the prior 2 weeks (14 days). The *on-treatment period* is defined as all times which are considered on-treatment.
 - In general, the on-treatment period will therefore be from the date of first trial product administration to date of last trial product administration excluding potential off-treatment time intervals triggered by at least two consecutive missed doses.

- For the evaluation of adverse events the lag time for each on-treatment time interval is 7 weeks (49 days).

The in-trial and on-treatment periods define the patient years of observation (PYO) and patient years of exposure (PYE), respectively, as the total time duration in the periods.

10.3 Statistical analyses

If necessary, a statistical analysis plan (SAP) may be written in addition to the protocol, including a more technical and detailed elaboration of the statistical analyses. The SAP will be finalised before database lock.

Effect endpoints will be analysed using the FAS; safety endpoints will be analysed using the SAS.

Results from statistical analyses will generally be accompanied by two-sided 95% confidence intervals and corresponding p-values. Superiority will be claimed if p-values are less than 5% and the estimated treatment contrasts favours semaglutide 2.4 mg.

Handling of missing baseline data

The last available and eligible observation at or before randomisation is used as the baseline value. If no assessments are available, the mean value at randomisation across all subjects is used as the baseline value.

10.3.1 Primary endpoint

Definition of primary endpoint: % weight change

Change from baseline (week 0) to week 68 in body weight (%) is defined as

$$\% \text{ weight change} = \frac{(\text{body weight at week 68} - \text{body weight at baseline})}{\text{body weight at baseline}} \times 100.$$

Definition of primary endpoint: 5% responders

A body weight reduction of at least 5% from baseline (week 0) to week 68 is defined as

$$5\% \text{ responder} = \begin{cases} 1 & \text{if } \% \text{ weight change} \leq -5\% \\ 0 & \text{if } \% \text{ weight change} > -5\% \end{cases}$$

Analyses addressing the primary estimand

The following statistical analyses and imputation methods are designed to address the primary estimand, i.e. to assess the effectiveness of semaglutide 2.4 mg.

The analysis model for % weight change is a linear regression (ANCOVA) of % weight change with randomised treatment as a factor and baseline body weight (kg) as covariate. The estimated treatment difference between semaglutide 2.4 mg and semaglutide placebo will be reported together with the associated two-sided 95% confidence interval (CI) and corresponding p-value.

The analysis model for the 5% responder endpoint is a logistic regression using randomised treatment as a factor and baseline body weight (kg) as covariate. The estimated odds ratio (OR) between semaglutide 2.4 mg and semaglutide placebo will be reported together with the associated two-sided 95% CI and corresponding p-value.

The superiority tests of semaglutide 2.4 mg vs. semaglutide placebo will be carried out as follows for the two analysis models.

Let $\mu_{\text{semaglutide}}$ and $\mu_{\text{semaglutide placebo}}$ denote the true mean of % weight change for semaglutide 2.4 mg and semaglutide placebo group, respectively. The null and alternative hypotheses tested are

$$\begin{aligned} H: \mu_{\text{semaglutide}} &\geq \mu_{\text{semaglutide placebo}} \text{ vs} \\ H_A: \mu_{\text{semaglutide}} &< \mu_{\text{semaglutide placebo}}. \end{aligned}$$

The hypothesis will be rejected and superiority claimed, if the upper limit of the estimated two-sided 95% CI is below 0.

Let $OR_{\text{semaglutide/semaglutide placebo}}$ denote the true odds ratio between semaglutide 2.4 mg and semaglutide placebo. The null and alternative hypotheses tested are

$$\begin{aligned} H: OR_{\text{semaglutide/semaglutide placebo}} &\leq 1 \text{ vs} \\ H_A: OR_{\text{semaglutide/semaglutide placebo}} &> 1. \end{aligned}$$

The hypothesis will be rejected and superiority claimed, if the lower limit of the estimated two-sided 95% CI is above 1.

Handling of missing week 68 values for the primary estimand

All available data at week 68 (AT and AD) are used and missing values (MT and MD) at week 68 will be imputed and the endpoints will be derived from the imputed values. Several approaches for imputation will be applied. First, a description of the primary imputation approach to address the primary estimand for the primary endpoints is given followed by a description of the sensitivity analyses used to assess the robustness of the primary analysis results. The sensitivity analyses investigate how assumptions on body weight development after discontinuation of randomised treatment impact the estimated treatment contrasts between semaglutide 2.4 mg and semaglutide placebo. An illustration of all imputation approaches for the effectiveness estimand is given in [Figure 10-1](#).




Figure 10-1 Illustration of imputation approaches for the effectiveness estimand

Primary imputation approach for the primary estimand

Multiple imputation approach using retrieved subjects (RD-MI): The primary imputation approach for the primary estimand is a multiple imputation similar to the one described by McEvoy⁷⁹. Missing body weight measurement at week 68 for non-retrieved subjects (MD) are imputed using assessments from retrieved subjects (AD) in each randomised treatment arm. This will be done according to the timing of last available observation (LAO) of body weight. Missing body weight measurements at week 68 for subjects on randomised treatment (MT) are imputed by sampling from available measurements at week 68 from subjects on randomised treatment (AT) in the relevant randomised treatment arm. The multiple imputation approach is done in three steps:

1. **Imputation:** Defines an imputation model using retrieved subjects (AD) from FAS and done within groups defined by randomised treatment and the timing of the LAO of body weight. The model will be a linear regression of body weight (kg) at week 68 with gender (male/female), baseline BMI (kg/m²) (in categories 27-<35, 35-<40, ≥40) as factors and baseline body weight (kg) and LAO of body weight (kg) as covariates. No interactions will be included. The grouping of timing will be done by quarters (intervals of 17 weeks). If timing by quarters is too restrictive, halves (intervals of 34 weeks) or excluding timing will be used. If any subjects are MT, an imputation model for missing body weight measurements at week 68 for MT subjects will also be defined using AT subjects in a similar way. The estimated posterior distribution for the parameters (regression coefficients and variances) in the imputation models are then used to impute missing week 68 body weight values for each randomised treatment arm. This will be done 1,000 times and results in 1,000 complete data sets.
2. **Analysis:** Analysis of each of the 1,000 complete data sets, using the analysis models (ANCOVA and logistic regression) results in 1,000 times 2 estimations.
3. **Pooling:** Integrates the 1,000 times 2 estimation results into two final results using Rubin's formula.

Based on NN9536-4153 phase 2 results 1,000 copies should be sufficient to establish stable results. If 1,000 copies are insufficient, 10,000 copies will be used. The multiple imputations will be generated using Novo Nordisk trial number 95364373 as seed number.

Sensitivity analyses

Jump to reference multiple imputation approach (J2R-MI): Missing values of body weight at week 68 (MT and MD) for both the semaglutide 2.4 mg and semaglutide placebo group are imputed by sampling among all available assessments at week 68 in the semaglutide placebo group (AT and AD). This approach makes the assumption that subjects instantly after discontinuation lose any effect of randomised treatment beyond what can be expected from semaglutide placebo treatment as adjunct to reduced-calorie diet and increased physical activity⁸⁰. The multiple imputation approach is done as above with the first step replaced by

1. **Imputation:** Defines an imputation model using semaglutide placebo subjects from FAS with a week 68 measurement (AT and AD). The model will be a linear regression of body weight (kg) at week 68 with gender (male/female), BMI (kg/m²) (in categories 27-<35, 35-<40, ≥40) as factors and baseline body weight (kg) as covariate. No interactions will be included. The estimated posterior distribution for the parameters (regression coefficients and variances) in the imputation models are then used to impute missing week 68 body weight values for each randomised treatment arm. This will be done 1,000 times and results in 1,000 complete data sets.

The jump to reference approach is the basis for the sample size calculations.

A single imputation approach as done by Sacks⁸¹ (S1-SI and S2-SI): Missing weight measurements at week 68 for non-retrieved subjects (MD) are imputed using a weight regain rate of 0.3 kg/month after LAO but truncated at no change from baseline whenever the extrapolation would lead to a positive weight gain relative to baseline. If a subject's weight at drug discontinuation represented a gain in weight relative to baseline, no additional gain will be imputed, and the unfavourable gain is carried forward to week 68. The weight regain imputation will be done for both randomised arms (S1-SI). Additionally, a version where only the semaglutide 2.4 mg arm uses the regain rate while the semaglutide placebo arm uses last available observation (corresponding to a weight regain rate of 0 kg/month) will be performed (S2-SI). For both versions, missing weight measurements at week 68 for subjects on randomised treatment (MT) are imputed by using LAO.

Tipping-point multiple imputation analysis (TP-MI): First, missing data are imputed according to the primary multiple imputation approach. Second, for the semaglutide 2.4 mg treatment arm a penalty will be added to the imputed values at week 68. The approach is to gradually increase this penalty until all confirmed conclusions from the primary analysis are reversed. For each hypothesis tested the specific value of the penalty that reverses the conclusion will be used to evaluate the robustness of the primary analysis results. This sensitivity analysis evaluates the robustness of the superiority conclusions.

Mixed model for repeated measurements (MMRM): This 'MMRM for effectiveness' will use all assessments regardless of adherence to randomised treatment, including assessments at week 68 for retrieved drop-outs (AD). The MMRM for effectiveness will be fitted using the same factor and covariate as for the primary analyses all nested within visit. An unstructured covariance matrix for measurements within the same subject will be employed, assuming that measurements for different subjects are independent.

Non-retrieved subjects as non-responders: For the 5% responder analysis an analysis using non-retrieved subjects as non-responders in the logistic regressions will be done.

Analysis addressing the secondary estimand

The secondary estimand for % weight change addresses the efficacy of semaglutide 2.4 mg and will be assessed using a 'MMRM for efficacy'. Week 68 assessments for retrieved drop-outs (AD) are not used in this analysis. The MMRM for efficacy will use assessments only from subjects who are taking the randomised treatment until end of treatment or until first discontinuing of randomised treatment. The derived date of the second consecutive missed dose will be used as the latest date for using assessments in this MMRM. The assessment closest in time and before the derived date of the second consecutive missed dose will be used as last assessment on randomised treatment. For subjects who initiate rescue interventions before completion or first discontinuing of randomised treatment, the date of starting weight management drugs or undergoing bariatric surgery will be used as latest date for using assessments in this MMRM. Similarly, the assessment closest in time and before the date of starting weight management drugs or undergoing bariatric surgery will be used as last assessment on randomised treatment. The MMRM for efficacy will be fitted using % weight change and the same factor and covariate as for the primary analyses all nested within visit. An unstructured covariance matrix for measurements within the same subject will be employed, assuming that measurements for different subjects are independent.

The secondary estimand for 5% responders will be assessed using the same MMRM for efficacy. From the MMRM individually predicted values for % weight change at week 68 will be used to classify each subject as 5% responder or not. This classification will then be analysed using a logistic regression model with treatment as the only factor.

An overview of all analysis and imputation methods to address the effectiveness and efficacy estimands for the primary endpoints is given in [Table 10-3](#).

Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy

| Objective | Endpoint | Test order | Endpoint type | Estimand | Analysis set | Statistical model | Imputation approach | Sensitivity analyses |
|----------------------------------|-----------------------------|------------|---------------|-----------|--------------|-------------------|---------------------|--|
| Primary endpoints | | | | | | | | |
| Primary | % weight change | 1 | Continuous | Primary | FAS | ANCOVA | RD-MI | J2R-MI S1-SI S2-SI TP-MI MMRM |
| | | | | Secondary | FAS | MMRM | - | - |
| Primary | 5% responders | 2 | Binary | Primary | FAS | LR | RD-MI | J2R-MI S1-SI S2-SI TP-MI MMRM Non-responder |
| | | | | Secondary | FAS | LR | MMRM | - |
| Confirmatory secondary endpoints | | | | | | | | |
| Primary | 10% responders | 3 | Binary | Primary | FAS | LR | RD-MI | Non-responders |
| | | | | Secondary | FAS | LR | MMRM | - |
| Primary | 15% responders | 4 | Binary | Primary | FAS | LR | RD-MI | Non-responders |
| | | | | Secondary | FAS | LR | MMRM | - |
| Primary | WC change (cm) | 5 | Continuous | Primary | FAS | ANCOVA | RD-MI | J2R-MI |
| | | | | Secondary | FAS | MMRM | - | - |
| Secondary | sBP change (mmHg) | 6 | Continuous | Primary | FAS | ANCOVA | RD-MI | J2R-MI |
| Secondary | SF-36 PF score change | 7 | Continuous | Primary | FAS | ANCOVA | RD-MI | J2R-MI |
| Secondary | IWQoL-Lite PFD score change | 8 | Continuous | Primary | FAS | ANCOVA | RD-MI | J2R-MI |

FAS = full analysis set; ANCOVA = analysis of covariance; RD-MI = multiple imputation using retrieved subjects; J2R-MI = jump to reference multiple imputation; S1-SI and S2-SI = single imputation as done by Sacks; TP-MI = tipping point multiple imputation; MMRM = mixed model for repeated measurements; LR = logistic regression; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain

10.3.2 Secondary endpoints

10.3.2.1 Confirmatory secondary endpoints

Confirmatory secondary endpoints are listed in Section 4 and are all included in the fixed-sequence statistical strategy, see above. All tests are tests of superiority of semaglutide 2.4 mg to semaglutide placebo.

Analyses addressing the primary estimand

All confirmatory secondary endpoints will be analysed using the same imputation approach as used for the primary endpoints and to address the primary estimand. The imputation model is the same as for the primary endpoints with body weight replaced by assessments of the endpoint to be analysed. The statistical model for continuous endpoints will be ANCOVA with factor and covariate as for the primary endpoint % weight change with baseline body weight replaced by the baseline

assessment of the endpoint to be analysed. The statistical model for body weight responder endpoints will be logistic regression with factor and covariate as for the primary endpoint 5% responders.

Analyses addressing the secondary estimand

The confirmatory secondary endpoints which relate to the primary objective will be analysed to address the secondary estimand using the same MMRM for efficacy described for the primary endpoints.

Sensitivity analyses for confirmatory secondary endpoints

For all continuous confirmatory secondary endpoints a sensitivity analysis using jump to reference as imputation approach will be carried out. For all binary confirmatory secondary endpoints a sensitivity analysis using non-retrieved subjects as non-responders will be carried out.

An overview of all analysis and imputation methods to address the effectiveness and efficacy estimands for confirmatory secondary endpoints is given in [Table 10-3](#).

10.3.2.2 Supportive secondary endpoints

Supportive secondary endpoints are listed in Section [4](#). All tests are tests of superiority of semaglutide 2.4 mg to semaglutide placebo.

Analyses addressing the primary estimand

The effect-related supportive secondary endpoints will be analysed using the same imputation approach as used for the primary endpoints and to address the primary estimand. The imputation model is the same as for the primary endpoints with body weight replaced by assessments of the endpoint to be analysed. The statistical model for continuous endpoints will be ANCOVA with factor and covariate as for the primary endpoint % weight change with baseline body weight replaced by the baseline assessment of the endpoint to be analysed.

The statistical model for responder endpoints relating to COAs will be logistic regression with randomised treatment as a factor and the baseline assessment of the endpoint to be analysed as covariate.

For body composition endpoints assessed by DEXA, missing data at week 68 will be imputed using jump to reference due to the low number of expected retrieved subjects in this subset. The analysis method will be ANCOVA with randomised treatment as a factor and the baseline assessment of the endpoint to be analysed as covariate. For all DEXA derived endpoints two-sided p-values for test of no difference between the two treatment groups will be presented.

For lipids and biomarkers a multiplicative model will be used, i.e. the ratio between post randomisation measurements and baseline will be calculated instead of differences, and both the dependent variable and covariate will be log-transformed.

Analyses addressing the secondary estimand

The supportive secondary endpoints which relate to the primary objective will be analysed to address the secondary estimand using the same MMRM for efficacy described for the primary endpoints.

Sensitivity analyses for supportive secondary endpoints

For supportive secondary endpoints no sensitivity analysis will be carried out.

Analysis of safety endpoints

The safety endpoint pulse will be analysed using an MMRM for efficacy as described in Section [10.3.1](#). For amylase, lipase and calcitonin descriptive statistics will be provided. The analysis of calcitonin will be stratified by gender.

Adverse events will be defined as “treatment-emergent” (TEAE), if the onset of the event occurs in the on-treatment period (see definition in Section [10.2](#)). TEAEs and SAEs will be summarised by descriptive statistics, such as frequencies and rates. No formal statistical inference will be carried out based on the number of TEAEs and SAEs.

An overview of all analysis and imputation methods to address the effectiveness and efficacy estimands for supportive secondary endpoints is given in [Table 10-4](#).

Protocol
Trial ID: NN9536-4373

CONFIDENTIAL

Date:
Version:
Status:
Page:

21 December 2017
2.0
Final
60 of 104

Novo Nordisk

Table 10-4 Analysis and imputation methods to address the effectiveness and efficacy estimands for supportive secondary endpoints

| Objective | Endpoint | Endpoint type | Estimand | Analysis set | Statistical model | Imputation approach | Sensitivity analyses |
|--|--|---------------|-----------|--------------|------------------------|---------------------|----------------------|
| Supportive secondary endpoints (effect related) | | | | | | | |
| Primary | Weight change (kg) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| | | | Secondary | FAS | MMRM | - | - |
| Primary | BMI change (kg/m ²) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| | | | Secondary | FAS | MMRM | - | - |
| Secondary | HbA _{1c} change (%; mmol/mol) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | FPG change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | Fasting insulin change (mIU/L) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | dBp change (mmHg) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | Total cholesterol change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | HDL change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | LDL change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | VLDL change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | FFA change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | Triglycerides change (mg/dL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | hsCRP change (mg/L) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | PAI-1 change (mg/L) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | sLR change (ng/mL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | Leptin change (ng/mL) | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 PF score responders # | Binary | Primary | FAS | LR | RD-MI | - |
| Secondary | SF-36 RP score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 BP score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 GH score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 VT score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 SF score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 RE score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 MH score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 PCS score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | SF-36 MCS score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | IWQoL-Lite PFD score responders## | Binary | Primary | FAS | LR | RD-MI | - |
| Secondary | IWQoL-Lite PDD score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | IWQoL-Lite PSD score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | IWQoL-Lite total score change | Continuous | Primary | FAS | ANCOVA | RD-MI | - |
| Secondary | Total fat mass change (%; g) | Continuous | Primary | FAS | ANCOVA | J2R-MI | - |
| Secondary | Lean body mass change (%; g) | Continuous | Primary | FAS | ANCOVA | J2R-MI | - |
| Secondary | Visceral fat mass change (%; g) | Continuous | Primary | FAS | ANCOVA | J2R-MI | - |
| Supportive secondary endpoints (safety related) | | | | | | | |
| Secondary | Number of TEAEs | Continuous | - | SAS | - | - | - |
| Secondary | Number of SAEs | Continuous | - | SAS | - | - | - |
| Secondary | Pulse change (bpm) | Continuous | - | SAS | MMRM | - | - |
| Secondary | Amylase change (U/L) | Continuous | - | SAS | Descriptive statistics | - | - |
| Secondary | Lipase change (U/L) | Continuous | - | SAS | Descriptive statistics | - | - |
| Secondary | Calcitonin change (ng/L) | Continuous | - | SAS | Descriptive statistics | - | - |

FAS = full analysis set; ANCOVA = analysis of covariance; RD-MI = multiple imputation using retrieved subjects; MMRM = mixed model for repeated measurements; BMI = body mass index; HbA1c = Hemoglobin A1c; FPG = fasting plasma glucose; dBP = diastolic blood pressure; HDL = high density lipoprotein; LDL = low density lipoprotein; VLDL = very low density lipoprotein; FFA = free fatty acids; hsCRP = high sensitivity C-Reactive Protein; PAI-1 = Plasminogen Activator Inhibitor-1; sLR = soluble Leptin Receptor; LR = logistic regression; SF-36 = Short Form 36 v2.0 acute; PF = Physical Functioning; RP = Role-Physical; BP = Bodily Pain; GH = General Health; VT = Vitality; SF = Social Functioning; RE = Role-Emotional; MH = Mental Health; PCS = Physical component summary; MCS = Mental component summary; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain; PDD = pain/discomfort domain; PSD = psychosocial domain; TEAEs = treatment emergent adverse events; SAEs = serious adverse events; # responder value = 4.3; ## responder value to be validated in trials NN9536-4153 and NN9924-4233

10.3.3 Exploratory endpoints

Exploratory endpoints are listed in Section 4. Observed data for exploratory endpoints will be summarised by descriptive statistics.

10.3.4 Explorative statistical analysis for pharmacogenetics and biomarkers

The statistical analysis of biomarker endpoints is described under Section [10.3.2.2](#).

10.3.5 Other analyses

All collected data that were not defined as endpoints will be summarised by descriptive statistics.

10.4 Pharmacokinetic and/or pharmacodynamic modelling

Population PK and exposure-response analyses will be used as supportive evidence for the evaluation of efficacy and safety and further to support the recommended dose of semaglutide in subjects with obesity. First, plasma semaglutide concentrations will be analysed using a population pharmacokinetic model, quantifying covariate (such as baseline body weight, age, gender, race, ethnicity, and injection site) effects on semaglutide exposure. Second, model based estimates of steady-state average concentrations will be derived for each subject, in order to facilitate subsequent exposure-response analyses. Relevant efficacy and safety endpoints will be related to steady-state average concentrations and subjected to model based analysis.

The analyses will be conducted separately for each trial and be combined into a meta-analysis, including the phase 2 trial and phase 3a trials with PK sampling. A modelling analysis plan will be prepared before first database lock in the semaglutide phase 3a programme for weight management, outlining details of the analyses. The modelling will be performed by Quantitative Clinical Pharmacology at Novo Nordisk and will be reported separately from the clinical trial reports.



Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

21 December 2017
2.0
Final
62 of 104

Novo Nordisk

11 Appendices

Appendix 1 Abbreviations and Trademarks

| | |
|-------------------|--|
| AD | available but discontinued |
| AE | adverse event |
| ALT | alanine aminotransferase |
| ANCOVA | analysis of covariance |
| AST | aspartate aminotransferase |
| AT | available on treatment |
| BMI | body mass index |
| BMR | basal metabolic rate |
| CI | confidence interval |
| CLAE | clinical laboratory adverse event |
| COA | Clinical Outcome Assessments |
| CRF | case report form |
| C-SSRS | Columbia-Suicide Severity Rating Scale |
| CTR | Clinical Trial Report |
| DEXA | Dual Energy X-ray Absorptometry |
| DFU | direction for use |
| DUN | dispensing unit number |
| EAC | event adjudication committee |
| EAS | event adjudication system |
| ECG | Electrocardiogram |
| eGFR | estimated glomerular filtration rate |
| FAS | full analysis set |
| FDA | U.S. Food and Drug Administration |
| FDAAA | FDA Amendments Act |
| FPG | fasting plasma glucose |
| FSH | follicle-stimulating hormone |
| GCP | Good Clinical Practice |
| GLP-1 RA | glucagon-like peptide-1 receptor agonist |
| HbA _{1c} | glycated haemoglobin |
| HDL | High density lipoprotein |

| | |
|-------------------|--|
| HRT | hormone replacement therapy |
| hsCRP | high sensitive C-reactive protein |
| IB | investigator's brochure |
| ICH | International Council for Harmonisation |
| ICIQ-UI-SF | International Consultation on Incontinence Questionnaire-Urinary Incontinence-Short Form |
| IEC | independent ethics committee |
| IMP | investigational medicinal product |
| IRB | institutional review board |
| IWQoL-Lite for CT | Impact of Weight on Quality of Life-Lite for Clinical Trials |
| IWRS | interactive web response system |
| KDIGO | Kidney Disease: Improving Global Outcomes |
| LAO | last available observation |
| LDL | low-density lipoprotein |
| MD | Missing and discontinued |
| MEN2 | multiple endocrine neoplasia type 2 |
| MHP | mental health professional |
| MMRM | mixed model for repeated measurements |
| MT | missing on treatment |
| MTC | medullary thyroid cancer |
| OR | odds ratio |
| PAI-1 | plasminogen activator inhibitor-1 |
| PCD | primary completion date |
| PGI-C | Patient Global Impression of Change |
| PGI-S | Patient Global Impression of Status |
| PHQ-9 | Patient Health Questionnaire-9 |
| PK | Pharmacokinetic |
| RA | receptor agonist |
| SAE | serious adverse event |
| SAP | statistical analysis plan |
| s.c. | Subcutaneous |

| | |
|-------|---|
| SD | standard deviation |
| SF-36 | Short Form-36 |
| SPS-6 | Stanford Presenteeism Scale-6 |
| SUSAR | suspected unexpected serious adverse reaction |
| T2D | type 2 diabetes |
| TEAE | treatment-emergent adverse event |
| TEE | total energy expenditure |
| TMM | trial materials manual |
| TSH | thyroid stimulating hormone |
| UNL | upper normal limit |
| VLDL | very low density lipoprotein |
| WOCBP | woman of child bearing potential |

Appendix 2 Clinical laboratory tests

- The tests detailed in [Table 11-1](#) and [Table 11-2](#) will be performed by the central laboratory.
- Laboratory samples specified in the protocol should be sent to the central laboratory for analysis.
- Additional tests may be performed at local laboratory at any time during the trial as determined necessary by the investigator or required by local regulations.
- The laboratory equipment may provide analyses not requested in the protocol but produced automatically in connection with the requested analyses according to specifications in the laboratory standard operating procedures. Such data will not be transferred to the trial database, but abnormal values will be reported to the investigator.
- The investigator must review all laboratory results for concomitant illnesses and AEs.
- Laboratory samples will be destroyed no later than at finalisation of the clinical trial report except, antibody samples which will be stored until marketing authorisation or destroyed at the latest 15 years from end of trial.
- For haematology samples (differential count) where the test result is not normal, then a part of the sample may be kept for up to 2 years or according to local regulations.

Table 11-1 Protocol-required efficacy laboratory assessments

| Laboratory assessments | Parameters |
|---|--|
| Glucose metabolism | <ul style="list-style-type: none"> • Fasting plasma glucose¹ • HbA1c • Fasting serum insulin |
| Lipids | <ul style="list-style-type: none"> • Cholesterol • High density lipoprotein (HDL) cholesterol • Low density lipoprotein (LDL) cholesterol • Triglycerides • Very-low-density lipoprotein (VLDL) cholesterol • Free fatty acids |
| Biomarkers | <ul style="list-style-type: none"> • PAI -1 Activity • hsCRP • leptin • Soluble leptin receptor |
| NOTES : ¹ A FPG result ≤ 3.9 mmol/L (70 mg/dL) in relation to planned fasting visits should not be reported as a hypoglycaemic episode but as a CLAE at the discretion of the investigator (Appendix 4). | |

Table 11-2 Protocol-required safety laboratory assessments

| Laboratory assessments | Parameters |
|---------------------------|---|
| Haematology | <ul style="list-style-type: none"> • Basophils • Eosinophils • Erythrocytes • Haematocrit • Haemoglobin • Leucocytes • Lymphocytes • Monocytes • Neutrophils • Thrombocytes |
| Biochemistry ¹ | <ul style="list-style-type: none"> • Alanine Aminotransferase (ALT)² • Albumin • Albumin corrected calcium • Alkaline phosphatase • Amylase³ • Aspartate Aminotransferase (AST)² • Calcitonin³ • Creatine kinase • Creatinine • Gamma Glutamyl Transferase • Lipase³ • Potassium • Sodium • Thyroid stimulating hormone (TSH)⁴ • Total bilirubin • Urea |
| Pregnancy Testing | <ul style="list-style-type: none"> • Urine human chorionic gonadotropin (hCG) pregnancy test (as needed for women of childbearing potential)⁵ |
| Other tests | <ul style="list-style-type: none"> • eGFR calculated by the central laboratory according to CKD-EPI creatinine equation as defined by KDIGO 2012⁶⁸ • Tryptase, in case of severe hypersensitivity (see Section 9.9) • Anti-semaglutide antibodies • Semaglutide plasma concentration |

Notes :

¹ Details of required actions and follow-up assessments for increased liver parameters including any discontinuation criteria are given in [Appendix 4](#) (Hy's Law) and Section [8.1](#).

² If ALT or AST > 3 upper normal limit (UNL), additional blood sample should be taken from the subject to analyse international normalised ratio (INR) by central laboratory (except at screening visit). Repeat testing of the abnormal laboratory assessments should be performed via central laboratory for the subject until abnormalities return to normal or baseline state.

³ Not collected at week 52.

⁴ If TSH level is out of normal range, additional testing will be performed by central lab: total and free T3 and T4 except at screening visit.

⁵ Local urine testing will be standard unless serum testing is required by local regulation or IRB/IEC.

- Anti-semaglutide antibodies, semaglutide plasma concentration, leptin and soluble leptin receptor analyses will be performed at a specialised laboratory.
- Laboratory/analyte results that could unblind the trial (e.g. antibodies) will only be disclosed after completion of the CTR if required by local regulation
- Laboratory outlier: If the following laboratory parameters are above/below the cut-off values in [Table 11-3](#), they are considered to be laboratory outliers and should be reported by completing a laboratory outlier form in the CRF:

Table 11-3 Criteria for laboratory outliers

| | Cut-off |
|---------------------------|------------------------------|
| Haematology | |
| Thrombocytes | < 25 x 10 ⁹ /L |
| Leucocytes | < 1 x 10 ⁹ /L |
| Lymphocytes | < 25 x 10 ⁹ /L |
| Biochemistry | |
| Albumin corrected calcium | < 1.5 mmol/L or > 3.4 mmol/L |
| Alkaline phosphatase | > 20 x UNL |
| Calcitonin | > 100 ng/L |
| Creatinine | > 6 x UNL |
| Creatine kinase | > 10 x UNL |
| Sodium | < 120 mmol/L or > 160 mmol/L |
| Potassium | < 2.5 mmol/L or > 7.0 mmol/L |

- Hepatic laboratory outlier: If the following hepatic laboratory parameters are above the cut-off values in [Table 11-4](#), it is considered to be hepatic laboratory outliers and should be reported by completing a hepatic event form in the CRF. It is at the investigator's discretion to determine whether it should also be reported as an adverse event (see [Appendix 4](#)).

Table 11-4 Criteria for hepatic laboratory outliers

| | Cut-off |
|----------------------|-----------|
| Alkaline phosphatase | >20 x UNL |
| ALT | >5 x UNL |
| AST | >5 x UNL |
| Total bilirubin | >10 x UNL |

Please note that in case of a hepatic event defined as ALT or AST >3 x UNL and total bilirubin > 2 x UNL, where no alternative aetiology exists (Hy's law), this must be reported as an SAE using the important medical event criterion if no other seriousness criteria are applicable.

Appendix 3 Trial governance considerations

1) Regulatory and ethical considerations

- This trial will be conducted in accordance with the protocol and with the following:
 - Consensus ethical principles derived from international guidelines including the Declaration of Helsinki⁸² and applicable ICH Good Clinical Practice (GCP) Guideline⁸³
 - Applicable laws and regulations
- The protocol, informed consent form, IB (as applicable) and other relevant documents (e.g. advertisements), must be submitted to an IRB/IEC and reviewed and approved by the IRB/IEC before the trial is initiated.
- Regulatory authorities will receive the clinical trial application, protocol amendments, reports on SAEs, and the clinical trial report according to national requirements.
- Any amendments to the protocol will require IRB/IEC approval before implementation of changes made to the trial design, except for changes necessary to eliminate an immediate safety hazard to trial subjects.
- Before a trial site is allowed to start screening subjects, written notification from Novo Nordisk must be received.
- The investigator will be responsible for:
 - providing written summaries of the status of the trial annually or more frequently in accordance with the requirements, policies, and procedures established by the IRB/IEC and/or regulatory authorities
 - notifying the IRB/IEC of SAEs or other significant safety findings as required by IRB/IEC procedures
 - providing oversight of the conduct of the trial at the site and adherence to requirements of ICH guidelines, the IRB/IEC, and all other applicable local regulations
 - ensuring submission of the clinical trial report synopsis to the IRB/IEC.

2) Financial disclosure

Investigators and subinvestigators will provide Novo Nordisk with sufficient, accurate financial information as requested to allow Novo Nordisk to submit complete and accurate financial certification or disclosure statements to the appropriate regulatory authorities. Investigators are responsible for providing information on financial interests during the course of the trial and one year after completion of the trial.

For US trial sites: verification under disclosures per Code of Federal Regulations (CFR) of Financial Conflict of Interest.

3) Informed consent process

- The investigator or his/her representative will explain the nature of the trial to the subject and answer all questions regarding the trial.

- The investigator must ensure the subject ample time to come to a decision whether or not to participate in the trial.
- Subjects must be informed that their participation is voluntary.
- Subjects will be required to sign and date a statement of informed consent that meets the requirements of local regulations, ICH guidelines⁸³, Declaration of Helsinki⁸² and the IRB/IEC or trial site.
- The medical record must include a statement that written informed consent was obtained before any trial related activity and the date when the written consent was obtained. The authorised person obtaining the informed consent must also sign and date the informed consent form before any trial related activity.
- The responsibility of seeking informed consent must remain with the investigator, but the investigator may delegate the task of informing to a medically qualified person, in accordance with local requirements.
- Subjects must be re-consented to the most current version of the informed consent form(s) during their participation in the trial
- A copy of the informed consent form(s) must be provided to the subject.

4) Information to subjects during trial

The site will be offered a communication package for the subject during the conduct of the trial. The package content is issued by Novo Nordisk. The communication package will contain written information intended for distribution to the subjects. The written information will be translated and adjusted to local requirements and distributed to the subject at the discretion of the investigator.

All written information to subjects must be sent to IRB/IEC for approval/favourable opinion and to regulatory authorities for approval or notification according to local regulations.

Different initiatives for subject retention will be implemented throughout this trial. Site retention activities may include cooking classes, group meetings, fitness memberships and others. Materials and items will be supplied if locally acceptable. The retention items will be relevant for the subjects' participation in the trial and/or their obesity and will not exceed local fair market value.

The initiatives for subject retention must be sent to IRB/IEC for approval/favourable opinion and to regulatory authorities for approval or notification according to local regulations.

5) Data protection

- Subjects will be assigned a 6-digit unique identifier, a subject number. Any subject records or datasets that are transferred to Novo Nordisk will contain the identifier only; subject names or any information which would make the subject identifiable will not be transferred.
- The subject and any biological material obtained from the subject will be identified by subject number, visit number and trial ID. Appropriate measures such as encryption or leaving out

certain identifiers will be enforced to protect the identity of subjects as required by local, regional and national requirements.

- The subject must be informed that his/her personal trial related data will be used by Novo Nordisk in accordance with local data protection law. The disclosure of the data must also be explained to the subject.
- The subject must be informed that his/her medical records may be examined by auditors or other authorised personnel appointed by Novo Nordisk, by appropriate IRB/IEC members, and by inspectors from regulatory authorities.

6) Committee structure

Novo Nordisk safety committee

Novo Nordisk will constitute an internal Semaglutide s.c. safety committee to perform ongoing safety surveillance. The Semaglutide s.c. safety committee may recommend unblinding of any data for further analysis, and in this case an independent ad hoc group will be established in order to maintain the blinding of the trial personnel.

Event adjudication committee

An independent external EAC is established to perform ongoing blinded adjudication of selected types of events and deaths (see [Table 9-1](#) and [Appendix 4](#)). The EAC will evaluate events sent for adjudication using pre-defined definitions and guidelines in accordance with the EAC Charter. The evaluation is based on review of pre-defined clinical data collected by the trial sites.

The EAC is composed of permanent members covering all required medical specialities. EAC members must disclose any potential conflicts of interest and must be independent of Novo Nordisk. The EAC will have no authorisations to impact on trial conduct, trial protocol or amendments.

The assessment made by both the EAC and the investigator will be presented in the clinical trial report.

In this trial, cardiovascular events will be adjudicated in order to adequately characterize the cardiovascular safety profile, since cardiovascular disease is an important and serious comorbidity of obesity⁸⁴. In addition, events of acute pancreatitis will be adjudicated because Novo Nordisk monitors these events closely, as treatment with GLP-1 RAs has been associated with acute pancreatitis.

7) Publication policy

The information obtained during the conduct of this trial is considered confidential, and may be used by or on behalf of Novo Nordisk for regulatory purposes as well as for the general development of the trial product. All information supplied by Novo Nordisk in connection with this trial shall remain the sole property of Novo Nordisk and is to be considered confidential information.

No confidential information shall be disclosed to others without prior written consent from Novo Nordisk. Such information shall not be used except in the performance of this trial.

The information obtained during this trial may be made available to other investigators who are conducting other clinical trials with the trial product, if deemed necessary by Novo Nordisk. Provided that certain conditions are fulfilled, Novo Nordisk may grant access to information obtained during this trial to researchers who require access for research projects studying the same disease and/or trial product studied in this trial.

Novo Nordisk may publish on its clinical trials website a redacted clinical trial report for this trial. One (or two) investigator(s) will be appointed by Novo Nordisk to review and sign the clinical trial report (signatory investigator) on behalf of all participating investigators. The signatory investigator(s) will be appointed based upon the criteria defined by the International Committee of Medical Journal Editors for research publications.⁸⁵

Communication of results

Novo Nordisk commits to communicate and disclose results of trials regardless of outcome. Disclosure includes publication of a manuscript in a peer-reviewed scientific journal, abstract submission with a poster or oral presentation at a scientific meeting or disclosure by other means.

The results of this trial will be subject to public disclosure on external web sites according to international and national regulations. Novo Nordisk reserves the right to defer the release of data until specified milestones are reached, for example when the clinical trial report is available. This includes the right not to release the results of interim analyses, because the release of such information may influence the results of the entire trial.

At the end of the trial, one or more scientific publications may be prepared collaboratively by the investigator(s) and Novo Nordisk. Novo Nordisk reserves the right to postpone publication and/or communication for up to 60 days to protect intellectual property.

In all cases the trial results will be reported in an objective, accurate, balanced and complete manner, with a discussion of the strengths and limitations. In the event of any disagreement on the content of any publication, both the investigators' and Novo Nordisk opinions will be fairly and sufficiently represented in the publication.

Authorship

Novo Nordisk will work with one or more investigator(s) and other experts who have contributed to the trial concept or design, acquisition, analysis or interpretation of data to report the results in one or more publications.

Authorship of publications should be in accordance with the Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals by the International Committee of Medical Journal Editors⁸⁵.

All authors will be provided with the relevant statistical tables, figures, and reports needed to evaluate the planned publication.

Where required by the journal, the investigator from each trial site will be named in an acknowledgement or in the supplementary material, as specified by the journal.

Site-specific publication(s) by investigator(s)

For a multicentre clinical trial, analyses based on single-site data usually have significant statistical limitations and frequently do not provide meaningful information for healthcare professionals or subjects, and therefore may not be supported by Novo Nordisk. Thus, Novo Nordisk may deny a request or ask for deferment of the publication of individual site results until the primary manuscript is accepted for publication. In line with Good Publication Practice, such individual reports should not precede the primary manuscript and should always reference the primary manuscript of the trial.

Investigator access to data and review of results

As owner of the trial database, Novo Nordisk has the discretion to determine who will have access to the database.

Individual investigators will have their own subjects' data, and will be provided with the randomisation code after results are available.

8) Dissemination of clinical trial data

Information of the trial will be disclosed at clinicaltrials.gov and novonordisk-trials.com. It will also be disclosed according to other applicable requirements such as those of the International Committee of Medical Journal Editors⁸⁶, the Food and Drug Administration Amendment Act (FDAAA)⁸⁷, European Commission Requirements^{88, 89} and other relevant recommendations or regulations. If a subject requests to be included in the trial via the Novo Nordisk e-mail contact at these web sites, Novo Nordisk may disclose the investigator's contact details to the subject. As a result of increasing requirements for transparency, some countries require public disclosure of investigator names and their affiliations.

The Primary Completion Date (PCD) is the last assessment of the primary endpoint, and is for this trial Last Subject First Treatment + 68 weeks corresponding to 'end of treatment' visit. If the last subject is withdrawn early, the PCD is considered the date when the last subject would have completed 'end of treatment' visit. The PCD determines the deadline for results disclosure at clinicaltrials.gov according to FDAAA.

9) Data quality assurance

Case Report Forms (CRFs)

- Novo Nordisk or designee is responsible for the data management of this trial including quality checking of the data.

- All subject data relating to the trial will be recorded on electronic CRFs unless transmitted electronically to Novo Nordisk or designee (e.g. laboratory data). The investigator is responsible for verifying that data entries are accurate and correct by physically or electronically signing the CRF.
- For some data both electronic and paper CRFs are used.
- The following will be provided as paper CRFs:
 - Pregnancy forms
- The following will be provided as paper CRFs to be used when access to the electronic CRF is revoked or temporarily unavailable:
 - AE forms
 - Safety information forms
 - Technical complaint forms (also to be used to report complaints that are not subject related, e.g. discovered at trial site before allocation)
- Corrections to the CRF data may be made by the investigator or the investigator's delegated staff. An audit trail will be maintained in the CRF application containing as a minimum: the old and the new data, identification of the person entering the data, date and time of the entry and reason for the correction. If corrections are made by the investigator's delegated staff after the date when the investigator signed the CRF, the CRF must be signed and dated again by the investigator.
- The investigator must ensure that data is recorded in the CRF as soon as possible, preferably within 5 working days after the visit. Once data has been entered, it will be available to Novo Nordisk for data verification and validation purposes.

Monitoring

- The investigator must permit trial-related monitoring, audits, IRB/IEC review, and regulatory agency inspections and provide direct access to source data documents (original documents, data and records). Direct access includes permission to examine, analyse, verify and reproduce any record(s) and report(s) that are important to the evaluation of the trial. If the electronic medical record does not have a visible audit trail, the investigator must provide the monitor with signed and dated printouts. In addition the relevant trial site staff should be available for discussions at monitoring visits and between monitoring visits (e.g. by telephone).
- Trial monitors will perform ongoing source data verification to confirm that data entered into the CRF by authorised site personnel are accurate, complete and verifiable from source documents; that the safety and rights of subjects are being protected, to monitor drug accountability and collect completed paper CRF pages, if applicable, and that the trial is being conducted in accordance with the currently approved protocol and any other trial agreements, ICH GCP, and all applicable regulatory requirements.

- Monitoring will be conducted using a risk based approach including risk assessment, monitoring plans, centralised monitoring (remote assessment of data by Novo Nordisk) and visits to trial sites.
- Monitors will review the subject's medical records and other source data e.g. the diaries and mental health assessment instruments, to ensure consistency and/or identify omissions compared to the CRF.

Protocol compliance

Deviations from the protocol should be avoided. If deviations do occur, the investigator must inform the monitor and the implications of the deviation must be reviewed and discussed.

Deviations must be documented and explained in a protocol deviation by stating the reason, date, and the action(s) taken. Some deviations, for which corrections are not possible, can be acknowledged and confirmed via edit checks in the CRF or via listings from the trial database.

10) Source documents

- All data entered in the CRF must be verifiable in source documentation other than the CRF.
- The original of the completed diaries must not be removed from the trial site, unless they form part of the CRF and a copy is kept at the site. For food- and physical activity diary, if the subject uses an app or a tool other than the paper diaries, the medical record or dietician's notes from the diet and physical activity counselling can be used as source document.
- Source documents provide evidence for the existence of the subject and substantiate the integrity of the data collected. Source documents are filed at the trial site.
- Data reported in the CRF that are transcribed from source documents must be consistent with the source documents or the discrepancies must be explained. The investigator may need to request previous medical records or transfer records. Also, current medical records must be available.
- It must be possible to verify subject's medical history in the source documents such as subject's medical record.
- Subjects completing electronic patient reported outcomes instruments are the data originators. Data will be transmitted to a technology service provider database, thus the service provider database is the source.
- The investigator must document any attempt to obtain external medical information by noting the date(s) when information was requested and who was contacted.
- Definition of what constitutes source data can be found in a source document agreement at each trial site. There will only be one source document defined at any time for any data element.

11) Retention of clinical trial documentation

- Records and documents, including signed informed consent forms, pertaining to the conduct of this trial must be retained by the investigator for 15 years after end of trial unless local regulations or institutional policies require a longer retention period. No records may be destroyed during the retention period without the written approval of Novo Nordisk. No records may be transferred to another location or party without written notification to Novo Nordisk.
- The investigator must be able to access his/her trial documents without involving Novo Nordisk in any way. If applicable, electronic CRF and other subject data will be provided in an electronic readable format to the investigator before access is revoked to the systems and/or electronic devices supplied by Novo Nordisk. Site-specific CRFs and other subject data (in an electronic readable format or as paper copies or prints) must be retained by the trial site. If the provided electronic data (e.g. the CD-ROM) is not readable during the entire storage period, the investigator can request a new copy. A copy of all data will be stored by Novo Nordisk.
- Subject's medical records must be kept for the maximum period permitted by the hospital, institution or private practice

12) Trial and site closure

Novo Nordisk reserves the right to close the trial site or terminate the trial at any time for any reason at the sole discretion of Novo Nordisk. If the trial is suspended or terminated, the investigator must inform the subjects promptly and ensure appropriate therapy and follow-up. The investigator and/or Novo Nordisk must also promptly inform the regulatory authorities and IRBs/IECs and provide a detailed written explanation.

Trial sites will be closed upon trial completion. A trial site is considered closed when all required documents and trial supplies have been collected and a trial site closure visit has been performed.

The investigator may initiate trial site closure at any time, provided there is reasonable cause and sufficient notice is given in advance of the intended termination.

Reasons for the early closure of a trial site by Novo Nordisk or investigator may include but are not limited to:

- failure of the investigator to comply with the protocol, the requirements of the IRB/IEC or local health authorities, Novo Nordisk procedures or GCP guidelines
- inadequate recruitment of subjects by the investigator
- discontinuation of further trial product development.

13) Responsibilities

The investigator is accountable for the conduct of the trial at his/her site and must ensure adequate supervision of the conduct of the trial at the trial site. If any tasks are delegated, the investigator must maintain a log of appropriately qualified persons to whom he/she has delegated specified trial-related duties. The investigator must ensure that there is adequate and documented training for all

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 78 of 104

Novo Nordisk

staff participating in the conduct of the trial. It is the investigator's responsibility to supervise the conduct of the trial and to protect the rights, safety, and well-being of the subjects.

A qualified physician, who is an investigator or a sub-investigator for the trial, must be responsible for all trial-related medical decisions.

The investigator is responsible for filing essential documents (i.e. those documents which individually and collectively permit evaluation of the conduct of a trial and the quality of the data produced) in the investigator trial master file. The documents, including the subject identification code list must be kept in a secure locked facility so that no unauthorized persons can get access to the data.

The investigator will take all necessary technical and organisational safety measures to prevent accidental or wrongful destruction, loss or deterioration of data. The investigator will prevent any unauthorised access to data or any other processing of data against applicable law. The investigator must be able to provide the necessary information or otherwise demonstrate to Novo Nordisk that such technical and organisational safety measures have been taken.

During any period of unavailability, the investigator must delegate responsibility for medical care of subjects to a specific qualified physician who will be readily available to subjects during that time.

If the investigator is no longer able to fulfil the role as investigator (e.g. if he/she moves or retires) a new investigator will be appointed in consultation with Novo Nordisk.

The investigator and other site personnel must have sufficient English skills according to their assigned task(s).

14) Indemnity statement

Novo Nordisk carries product liability for its products, and liability as assumed under the special laws, acts and/or guidelines for conducting clinical trials in any country, unless others have shown negligence.

Novo Nordisk assumes no liability in the event of negligence or any other liability of the sites or investigators conducting the trial or by persons for whom the said site or investigator are responsible.

Novo Nordisk may pay additional costs incurred in relation to assessments relevant for following the safety of the subject. Investigator must contact Novo Nordisk on a case by case basis for whether the costs will be covered.

Appendix 4 Adverse events: definitions and procedures for recording, evaluation, follow-up, and reporting

| AE definition |
|--|
| <ul style="list-style-type: none"> An AE is any untoward medical occurrence in a clinical trial subject that is temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. An AE can be any unfavourable and unintended sign, including an abnormal laboratory finding, symptom or disease (new or exacerbated) temporally associated with the use of a medicinal product. |

| Events <u>meeting</u> the AE definition |
|--|
| <ul style="list-style-type: none"> Any abnormal laboratory test results or safety assessments, including those that worsen from baseline, considered clinically significant in the medical and scientific judgment of the investigator. A CLAE: a clinical abnormal laboratory finding which is clinically significant, i.e. an abnormality that suggests a disease and/or organ toxicity and is of a severity that requires active management. Active management includes active treatment or further investigations, for example change of medicine dose or more frequent follow-up due to the abnormality. Exacerbation of a chronic or intermittent pre-existing condition including either an increase in frequency and/or intensity of the condition. Signs, symptoms or the clinical sequelae of a suspected drug-drug interaction. Signs, symptoms or the clinical sequelae of a suspected overdose of trial product regardless of intent. A "lack of efficacy" or "failure of expected pharmacological action" per se will not be reported as an AE or SAE. Such instances will be captured in the efficacy assessments. However, the signs, symptoms and/or clinical sequelae resulting from lack of efficacy will be reported as AE or SAE if they fulfil the definition. |
| Events <u>NOT</u> meeting the AE definition |
| <ul style="list-style-type: none"> Pre-existing conditions, anticipated day-to-day fluctuations of pre-existing conditions, including those identified during screening or other trial procedures performed before exposure to trial product. Note: pre-existing conditions should be recorded as medical history/concomitant illness. Pre-planned procedures, unless the condition for which the procedure was planned has worsened from the first trial related activity after the subject has signed the informed consent. |

| Definition of an SAE |
|---|
| An SAE is an AE that fulfils at least one of the following criteria: |
| <ul style="list-style-type: none"> Results in death Is life-threatening The term 'life-threatening' in the definition of 'serious' refers to an event in which the subject was at risk of death at the time of the event. It does not refer to an event which hypothetically might have caused death, if it were more severe. Requires inpatient hospitalisation or prolongation of existing hospitalisation Hospitalisation signifies that the subject has been detained at the hospital or emergency ward for observation and/or treatment that would not have been appropriate in the physician's office or outpatient setting. Complications that occur during hospitalisation are AEs. If a complication prolongs hospitalisation or fulfils any other serious criteria, the event is serious. When in doubt as to whether "hospitalisation" occurred or was necessary, the AE should be considered serious. Hospitalisation for elective treatment of a pre-existing condition that did not worsen from baseline is not considered an AE. Note: <ul style="list-style-type: none"> Hospitalisations for administrative, trial related and social purposes do not constitute AEs and should therefore not be reported as AEs or SAEs. |

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 80 of 104

Novo Nordisk

| |
|---|
| <ul style="list-style-type: none"> ▪ Hospital admissions for surgical procedures, planned before trial inclusion, are not considered AEs or SAEs. |
| <ul style="list-style-type: none"> • Results in persistent disability/incapacity • The term disability means a substantial disruption of a person's ability to conduct normal life functions. • This definition is not intended to include experience of relatively minor medical significance such as uncomplicated headache, nausea, vomiting, diarrhoea, influenza, and accidental trauma (e.g. sprained ankle), which may interfere with or prevent everyday life functions but do not constitute a substantial disruption. |
| <ul style="list-style-type: none"> • Is a congenital anomaly/birth defect |
| <ul style="list-style-type: none"> • Important medical event: • Medical or scientific judgment should be exercised in deciding whether SAE reporting is appropriate in other situations. This includes important medical events that may not be immediately life-threatening or result in death or hospitalisation, but may jeopardise the subject or may require medical or surgical intervention to prevent one of the other outcomes listed in the above definition. These events should usually be considered serious and reported as SAEs using the important medical event criterion. • The following AEs must always be reported as SAEs using the important medical event criterion, if no other seriousness criteria are applicable: <ul style="list-style-type: none"> ▪ suspicion of transmission of infectious agents via the trial product. ▪ risk of liver injury defined as ALT or AST >3 x UNL and total bilirubin >2 x UNL, where no alternative aetiology exists (Hy's law). |

Description of AEs requiring additional data collection (via specific event form) and events for adjudication.

AEs requiring additional data collection

AEs requiring additional data collection are AEs where the additional data will benefit the evaluation of the safety of the trial product (see [Table 9-1](#)). The selection of these events is based on the non-clinical and clinical data with semaglutide, knowledge from the GLP-1 RA drug class as well as regulatory requirements.

| Event type | Description |
|---------------------------|---|
| Acute gallbladder disease | Events of symptomatic acute gallbladder disease (including gallstones and cholecystitis) |
| Acute pancreatitis | The diagnosis of acute pancreatitis requires two of the following three features: <ol style="list-style-type: none"> 1. abdominal pain consistent with acute pancreatitis (acute onset of a persistent, severe, epigastric pain often radiating to the back) 2. serum lipase activity (and/or amylase activity) at least three times greater than the upper limit of normal 3. characteristic findings of acute pancreatitis on imaging. |
| Malignant neoplasm | Malignant neoplasm by histopathology or other substantial clinical evidence |

| | |
|-------------------|--|
| Hepatic event | <p>Hepatic event defined as:</p> <ul style="list-style-type: none"> Disorders of the liver including cholestatic conditions and liver related signs and symptoms ALT or AST > 3x UNL and total bilirubin > 2x UNL or INR>1.5* ALT or AST > 3x UNL with the appearance of fatigue, nausea, vomiting, right upper quadrant pain or tenderness, fever, rash, and/or eosinophilia (>5%) <p>*Please note that in case of a hepatic event defined as ALT or AST >3 x UNL and total bilirubin > 2 x UNL, where no alternative aetiology exists (Hy's law), this must be reported as an SAE using the important medical event criterion if no other seriousness criteria are applicable.</p> |
| Medication error: | <p>A medication error concerning trial products is defined as:</p> <ul style="list-style-type: none"> Administration of wrong drug. <ul style="list-style-type: none"> Note: Use of wrong dispensing unit number (DUN) is not considered a medication error unless it results in a confirmed administration of wrong drug. Wrong route of administration, such as intramuscular instead of subcutaneous. Accidental administration of more than 2.4 mg/week or a higher dose than intended during dose escalation, however, the administered dose must deviate from the intended dose to an extent where clinical consequences for the trial subject were likely to happen as judged by the investigator, although they did not necessarily occur. Administration of an overdose with the intention to cause harm, misuse or abuse of trial product. |

Events for adjudication

| Event type | Description | Adjudication outcome |
|-----------------------------------|---|--|
| Death | All-cause death | Cardiovascular death (including undetermined cause of death) Non-Cardiovascular death |
| Acute Coronary Syndrome | Acute Coronary Syndrome conditions include all types of acute myocardial infarction and hospitalisation for unstable angina pectoris | Acute myocardial infarction (including subgroup classifications) Hospitalisation for unstable angina pectoris |
| Cerebrovascular events | Episode of focal or global neurological dysfunction that could be caused by brain, spinal cord, or retinal vascular injury as a result of haemorrhage or infarction | Ischaemic stroke Haemorrhagic stroke Undetermined stroke Transient Ischaemic Attack |
| Coronary artery revascularisation | A Coronary revascularisation procedure is a catheter-based (PCI) or a surgical procedure (CABG) designed to improve myocardial blood flow | Coronary revascularisation procedure |
| Heart failure | Presentation of the patient for an urgent, unscheduled clinic/office/emergency department visit or hospital admission, with a | Heart failure hospitalisation Urgent heart failure visit |

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 82 of 104

Novo Nordisk

| | | | | |
|--|--------------------|---|--|--|
| | | primary diagnosis of heart failure (new episode or worsening of existing heart failure) | | |
| | Acute pancreatitis | <p>The diagnosis of acute pancreatitis requires two of the following three features:</p> <p>(1) abdominal pain consistent with acute pancreatitis (acute onset of a persistent, severe, epigastric pain often radiating to the back)</p> <p>(2) serum lipase activity (and/or amylase activity) at least three times greater than the upper limit of normal</p> <p>(3) characteristic findings of acute pancreatitis on imaging</p> | <p>Acute pancreatitis</p> <p>Mild</p> <p>Moderately severe</p> <p>Severe</p> | |

AE and SAE recording

- The investigator will record all relevant AE/SAE information in the CRF.
- The investigator will attempt to establish a diagnosis of the event based on signs, symptoms, and/or other clinical information. In such cases, the diagnosis (not the individual signs/symptoms) will be documented as the AE/SAE.
- When an AE/SAE occurs, it is the responsibility of the investigator to review all documentation (e.g. hospital progress notes, laboratory and diagnostics reports) related to the event.
- There may be instances when copies of source documents (e.g. medical records) for certain cases are requested by Novo Nordisk. In such cases, all subject identifiers, with the exception of the subject number, will be redacted on the copies of the source documents before submission to Novo Nordisk.
- For all non-serious AEs the applicable forms should be signed when the event is resolved or at the end of the trial at the latest. For sign-off of SAE related forms refer to “SAE reporting via paper CRF” later in this section.
- Novo Nordisk products used as concomitant medication: if an AE is considered to have a causal relationship with a Novo Nordisk marketed product used as concomitant medication in the trial, it is important that the suspected relationship is reported to Novo Nordisk, e.g. in the alternative aetiology section on the safety information form. Novo Nordisk may need to report this AE to relevant regulatory authorities.

Assessment of severity

The investigator will assess intensity for each event reported during the trial and assign it to one of the following categories:

- **Mild:** An event that is easily tolerated by the subject, causing minimal discomfort and not interfering with everyday activities.
- **Moderate:** An event that causes sufficient discomfort and interferes with normal everyday activities.
- **Severe:** An event that prevents normal everyday activities.

Note: Severe is a category used for rating the intensity of an event; and both an AE and SAE can be assessed as severe. An event is defined as ‘serious’ when it meets at least one of the outcomes described in the definition of an SAE and not when it is rated as severe.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 83 of 104

Novo Nordisk

Assessment of causality

The investigator is obligated to assess the relationship between trial product and the occurrence of each AE/SAE.

Relationship between an AE/SAE and the relevant trial product(s) should be assessed as:

- Probable - Good reason and sufficient documentation to assume a causal relationship.
- Possible - A causal relationship is conceivable and cannot be dismissed.
- Unlikely - The event is most likely related to aetiology other than the trial product.

Alternative aetiology, such as underlying disease(s), concomitant medication, and other risk factors, as well as the temporal relationship of the event to trial product administration will be considered and investigated.

The investigator should use the IB for the assessment. For each AE/SAE, the investigator must document in the medical records that he/she has reviewed the AE/SAE and has provided an assessment of causality.

There may be situations in which an SAE has occurred and the investigator has minimal information to include in the initial report. However, **it is important that the investigator always makes an assessment of causality for every event before the initial transmission of the SAE data.**

The investigator may change his/her opinion of causality in light of follow-up information and send a follow-up report with the updated causality assessment.

The causality assessment is one of the criteria used when determining regulatory reporting requirements.

Final outcome

The investigator will select the most appropriate outcome:

- **Recovered/resolved:** The subject has fully recovered, or by medical or surgical treatment the condition has returned to the level observed at the first trial-related activity after the subject signed the informed consent.
- **Recovering/resolving:** The condition is improving and the subject is expected to recover from the event. This term is only applicable if the subject has completed the trial or has died from another AE.
- **Recovered/resolved with sequelae:** The subject has recovered from the condition, but with lasting effect due to a disease, injury, treatment or procedure. If a sequelae meets an SAE criterion, the AE must be reported as an SAE.
- **Not recovered/not resolved:** The condition of the subject has not improved and the symptoms are unchanged or the outcome is not known.
- **Fatal:** This term is only applicable if the subject died from a condition related to the reported AE. Outcomes of other reported AEs in a subject before he/she died should be assessed as “recovered/resolved”, “recovering/resolving”, “recovered/resolved with sequelae” or “not recovered/not resolved”. An AE with a fatal outcome must be reported as an SAE.
- **Unknown:** This term is only applicable if the subject is lost to follow-up.

Follow-up of AE and SAE

The investigator is obligated to perform or arrange for the conduct of supplemental measurements and/or evaluations as medically indicated or as requested by Novo Nordisk to elucidate the nature and/or causality of the AE or SAE as fully as possible (e.g. severe hypersensitivity reactions). This may include additional laboratory tests (e.g. skin prick test) or investigations, histopathological examinations, or consultation with other health care professionals.

If a subject dies during participation in the trial or during a recognised follow-up period, the investigator should provide Novo Nordisk with a copy of autopsy report including histopathology.

New or updated information will be recorded in the CRF.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 84 of 104

Novo Nordisk

SAE reporting via electronic CRF

- Relevant forms (AE and safety information form) must be completed in the CRF.
- For reporting and sign-off timelines (see box below).
- If the CRF is unavailable for more than 24 hours, then the site will use the paper AE form and if the CRF is unavailable for more than 5 calendar days then the site will use the paper safety information form (see box below).
- The site will enter the SAE data into the CRF as soon as it becomes available, see Section [9.2.1](#).
- After the trial is completed at a given site, the CRF will be decommissioned to prevent the entry of new data or changes to existing data. If a site receives a report of a new SAE from a subject or receives updated data on a previously reported SAE after CRF decommission, then the site can report this information on a paper AE and safety information form (see box below) or to Novo Nordisk by telephone.

SAE reporting via paper CRF

- Relevant CRF forms (AE and safety information form) must be forwarded to Novo Nordisk either by fax, e-mail or courier.
- Initial notification via telephone is acceptable, although it does not replace the need for the investigator to complete the AE and safety information form within the designated reporting time frames (as illustrated in [Figure 9-1](#))
- AE form within 24 hours.
- Safety information form within 5 calendar days.
- Both forms must be signed within 7 calendar days.

Contact details for SAE reporting can be found in the investigator trial master file.

Appendix 5 Contraceptive guidance and collection of pregnancy information

It must be recorded in the CRF whether female subjects are of childbearing potential.

Definitions

Woman of Childbearing Potential (WOCBP)

A woman is considered fertile following menarche and until becoming postmenopausal unless permanently sterile.

Women in the following categories are not considered WOCBP

1. Premenarcheal
2. Premenopausal female with one of the following:

- Documented hysterectomy
- Documented bilateral salpingectomy
- Documented bilateral oophorectomy

Note: Documentation can come from the site personnel's review of subject's medical records, medical examination or medical history interview.

3. Postmenopausal female
 - . Otherwise, they must discontinue HRT to allow confirmation of postmenopausal A postmenopausal state is defined as no menses for 12 months without an alternative medical cause. A high Follicle Stimulating Hormone (FSH) level in the postmenopausal range may be used to confirm a postmenopausal state in women not using hormonal contraception or Hormonal Replacement Therapy (HRT). However, in the absence of 12 months of amenorrhea, a single FSH measurement is insufficient.
 - Females on HRT and whose menopausal status is in doubt will be required to use one of the non-hormonal highly effective contraception methods if they wish to continue their HRT during the trialstatus before trial enrolment.

Contraception guidance

Male subjects

No contraception measures are required for male subjects as the risk of teratogenicity/fetotoxicity caused by transfer of semaglutide in seminal fluid is unlikely.

Female subjects

Female subjects of childbearing potential are eligible to participate if they agree to use methods of contraception consistently and correctly as described in table(s) below:

Table 11-5 Highly effective contraceptive methods

| |
|--|
| Highly effective contraceptive methods that are user dependent^{a and b} Failure rate of <1% per year when used consistently and correctly. |
| Combined (oestrogen and progestogen containing) hormonal contraception associated with inhibition of ovulation <ul style="list-style-type: none"> oral intravaginal transdermal |
| Progestogen only hormonal contraception associated with inhibition of ovulation <ul style="list-style-type: none"> oral injectable |
| Highly effective methods that are user independent^b |
| Implantable progestogen only hormonal contraception associated with inhibition of ovulation <ul style="list-style-type: none"> Intrauterine Device Intrauterine hormone-releasing System Bilateral tubal occlusion |
| Vasectomised partner A vasectomised partner is a highly effective contraception method provided that the partner is the sole male sexual partner of the WOCBP and the absence of sperm has been confirmed. If not, an additional highly effective method of contraception should be used. |
| Sexual abstinence Sexual abstinence is considered a highly effective method only if defined as refraining from heterosexual intercourse during the entire period of risk associated with the trial product. The reliability of sexual abstinence needs to be evaluated in relation to the duration of the trial and the preferred and usual lifestyle of the subject. |
| Notes: ^a Typical use failure rates may differ from those when used consistently and correctly. Use should be consistent with local regulations regarding the use of contraceptive methods for subjects participating in clinical trials. ^b Contraception should be utilised during the treatment period and for at least 49 days after the last dose of trial product. |

In certain cases, it is accepted to use double barrier methods (a condom combined with an occlusive cap (e.g. diaphragm) with/without the use of spermicide). This should only be allowed in females with:

- 1) known intolerance to the highly effective methods mentioned in [Table 11-5](#) or where the use of any of the listed highly effective contraceptive measures are contraindicated in the individual subject, and/or
- 2) if the risk of initiating treatment with a specific highly effective method outweighs the benefit for the female.

Justification for accepting double barrier method should be at the discretion of the investigator taking into consideration his/hers knowledge about the female's obesity history, concomitant

illness, concomitant medication and observed AEs. The justification must be stated in the medical records.

Pregnancy testing

- WOCBP should only be included after a negative highly sensitive urine pregnancy test.
- Urine pregnancy testing should be performed at every site visit every 4 or 8 weeks during the treatment period, at end of treatment and after the 7 weeks off-drug follow-up period according to the flowchart.
- Additional urine pregnancy testing should be performed at monthly intervals during the treatment period, if required locally ([Appendix 9](#)).
- Pregnancy testing should be performed whenever a menstrual cycle is missed or when pregnancy is otherwise suspected.

Collection of pregnancy information

Female subjects who become pregnant

- Investigator will collect pregnancy information on any female subject, who becomes pregnant while participating in this trial.
- Information will be recorded on the appropriate form and submitted to Novo Nordisk within 14 calendar days of learning of a subject's pregnancy.
- Subject will be followed to determine the outcome of the pregnancy. The investigator will collect follow-up information on subject and neonate, which will be forwarded to Novo Nordisk. Generally, follow-up will not be required for longer than 1 month beyond the delivery date.
- Any termination of pregnancy will be reported, regardless of foetal status (presence or absence of anomalies) or indication for procedure.
- While pregnancy itself is not considered to be an AE or SAE, any pregnancy complication or elective termination of a pregnancy will be reported as an AE or SAE.
- A spontaneous abortion is always considered to be an SAE and will be reported as such.
- Any SAE occurring as a result of a post-trial pregnancy which is considered possibly/probably related to the trial product by the investigator will be reported to Novo Nordisk as described in [Appendix 4](#). While the investigator is not obligated to actively seek this information in former subjects, he or she may learn of an SAE through spontaneous reporting.

Any female subject who becomes pregnant while participating in the trial will discontinue trial product.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 88 of 104

Novo Nordisk

Appendix 6 Technical complaints: Definition and procedures for recording, evaluation, follow-up and reporting

Technical complaint definition

A technical complaint is any written, electronic or oral communication that alleges product (medicine or device) defects. The technical complaint may be associated with an AE, but does not concern the AE itself.

Examples of technical complaints:

- Problems with the physical or chemical appearance of trial products (e.g. discoloration, particles or contamination).
- Problems with packaging material including labelling.
- Problems related to medical devices (e.g. to the injection mechanism, dose setting mechanism, dose button or interface between the pen-injector and the needle).

Time period for detecting technical complaints

All technical complaints, which occur from the time of receipt of the product at trial site until the time of the last usage of the product, must be collected for products predefined on the technical complaint form.

Reporting of technical complaints to Novo Nordisk

Contact details (fax, e-mail and address) for Customer Complaint Center – refer to Attachment I

Technical complaints must be reported on a separate technical complaint form:

1. One technical complaint form must be completed for each affected DUN
2. If DUN is not available, a technical complaint form for each batch, code or lot number must be completed

Timelines for reporting of technical complaints to Novo Nordisk

The investigator must complete the technical complaint form in the CRF within the timelines specified in [Figure 9-3](#).

If the CRF is unavailable or when reporting a technical complaint that is not subject related, the information must be provided on a paper form by fax, e-mail or courier to Customer Complaint Center, Novo Nordisk, within the same timelines as stated above. When the CRF becomes available again, the investigator must enter the information on the technical complaint form in the CRF.

Follow-up of technical complaints

The investigator is responsible for ensuring that new or updated information will be recorded on the originally completed form.

Collection, storage and shipment of technical complaint samples

The investigator must collect the technical complaint sample and all associated parts that were packed in the same DUN and notify the monitor within 5 calendar days of obtaining the sample at trial site. The sample and all associated parts must be sent as soon as possible to Customer Complaint Center, Novo Nordisk, together with a copy of the completed technical complaint form. The technical complaint sample should contain the batch, code or lot number and, if available, the DUN. If the technical complaint sample is unobtainable, the reason must be stated on the technical complaint form. If several samples are shipped in one shipment, the sample and the corresponding technical complaint form should be kept together.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 89 of 104

Novo Nordisk

Storage of the technical complaint sample must be done in accordance with the conditions prescribed for the product.

Reporting of technical complaints for Novo Nordisk products not included in technical complaint form

Technical complaints on Novo Nordisk products not included in the technical complaint form should be reported to local Novo Nordisk affiliate with a reference to trial ID.

All technical complaints are handled by the Customer Complaint Center at Novo Nordisk. Only technical complaints related to adverse events will be included in the CTR.

Appendix 7 Retention of human biosamples

Antibody samples

- Antibody samples will be retained for potential later analysis for further characterisation of antibody responses towards drug, if required by health authorities or for safety reasons.
- Only Novo Nordisk staff and personnel from the specialised laboratory will have access to the stored specimens.
- The samples will be stored at the specialised laboratory or Novo Nordisk after end of trial and until marketing authorisation approval or until the research project terminates, but no longer than 15 years from end of trial after which they will be destroyed.
- Samples might be transferred to other countries, if not prohibited by local regulations.
- The subject's identity will remain confidential and the samples will be identified only by subject number, visit number and trial identification number. No direct identification of the subject will be stored together with the samples. The analyses will not have any medical consequences for the subjects or their relatives.
- Subjects can contact the investigator if they wish to be informed about results derived from stored biosamples obtained from their own body.

Biosamples for future analysis

- The trial will involve collection of human biosamples to be stored in a central archive.
- Subjects must sign and date a separate informed consent form before biosamples are collected to be stored for future analysis.
- The material to be collected is serum collected at baseline and end of treatment. A total of 5 aliquots will be stored for each subject at both time points.
- As new biomarkers related to the disease and/or safety, efficacy or mechanism of action of semaglutide may evolve during the conduct of the trial, the analyses of the stored biosamples may also include biomarkers that are unknown at present or have not been included in the scientific hypotheses at initiation of the trial.
- The biosamples will be stored at a central laboratory for 6 months and at a central storage facility contracted by Novo Nordisk up to 15 years after end of trial. Only Novo Nordisk and storage facility employees will be able to access the stored biosamples. The biosamples may be transferred to other countries for analysis and will be destroyed at the latest 15 years after end of trial.
- The subject may request the stored biosamples to be destroyed by withdrawing the designated informed consent.
- The results obtained from any already performed analyses of the samples will still be used.
- In the event that the collected biosamples (serum) will be used in the future, the investigator will become directly informed by Novo Nordisk about the results if the findings are deemed clinically relevant and analytically valid and quantifiable. In such case, a written summary of

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 21 December 2017
Version: 2.0
Status: Final
Page: 91 of 104

Novo Nordisk

the findings, including listings of subject specific values, will be provided once a firm conclusion from the results has been drawn by Novo Nordisk.

- Potentially, observations of neoplastic diseases, serious hereditary diseases, other un-treatable diseases, or any other abnormal findings could be part of the observations. Subjects can contact the investigator if they wish to be informed about results derived from stored biosamples obtained from their own body.

Appendix 8 Monitoring of calcitonin

Background

Treatment with GLP-1 RAs has been shown to be associated with thyroid C-cell changes in rodents but not in non-human primates. The human relevance of this finding is unknown. However, based on the findings in rodents, monitoring of serum calcitonin (a sensitive biomarker for C-cell activation) is currently being performed in clinical trials with semaglutide.

While there is general agreement on the clinical interpretation of substantially elevated calcitonin levels (> 100 ng/L) as likely indicative of C-cell neoplasia, the interpretation of values between upper normal range (5.0 and 8.4 ng/L for women and men, respectively) and 100 ng/L is less clear with regards to indication of disease.

There are several known confounding factors affecting calcitonin levels, e.g.:

- renal dysfunction
- tobacco use
- autoimmune thyroiditis
- several drug classes (e.g. proton pump inhibitors, beta-blockers, H₂-blockers and glucocorticoids)

Physiology of C-cell activation in various clinical conditions and in different patient populations (i.e. with various comorbidities) is poorly understood. There may be various clinical conditions not identified so far which mildly or moderately affect calcitonin secretion by C-cells.

Calcitonin monitoring

A blood sample will be drawn at pre-specified trial visits for measurement of calcitonin.

In case a subject has a calcitonin value ≥ 10 ng/L, the algorithm outlined in [Figure 11-1](#) and described below should be followed. The algorithm applies for all calcitonin values in the trial.




Figure 11-1 Flow of calcitonin monitoring

Calcitonin ≥ 100 ng/L

Action: The subject (even if a screen failure) must immediately be referred to a thyroid specialist for further evaluation and the trial product must be discontinued (see Section 8.1). The subject should remain in the trial; however, all medications suspected to relate to this condition must be discontinued until diagnosis has been established.

Background: These values were found in 9 (0.15%) of a population of 5817 patients with thyroid nodular disease⁹⁰. All of these patients were diagnosed with MTC, resulting in a positive predictive value of 100%.

Diagnostic evaluation should include:

- thyroid ultrasound examination
- fine needle aspiration of any nodules > 1 cm

- potentially, surgery with neck dissection

In case a subject is diagnosed with MTC, it is common clinical practice to explore the family history of MTC or MEN2 and perform a genetic test for RET proto-oncogene mutation.

Calcitonin ≥ 50 and < 100 ng/L

Action: The subject (even if a screen failure) should be referred to a thyroid specialist for further evaluation. The subject should remain in the trial and can continue on trial product.

Background: These values were found in 8 (0.14%) of the population of 5817 patients with thyroid nodular disease⁹⁰. Two of these subjects were diagnosed with MTC and two were diagnosed with C cell hyperplasia, resulting in a positive predictive value of a C-cell anomaly of 50%.

Diagnostic evaluation should include:

- thyroid ultrasound examination
- if available, and if there are no contraindications, a pentagastrin stimulation test should be done. For subjects with positive pentagastrin stimulation test, surgery should be considered.
- if pentagastrin stimulation test is not available, thyroid ultrasound and fine needle aspiration biopsy may add important clinical information about the need for surgery.

Calcitonin ≥ 10 and < 50 ng/L

Action: The subject can continue in the trial on trial product. Continue sampling of calcitonin according to the protocol.

If the subject is a screen failure, or if the value is from the last sample taken in the trial, the subject should be referred to a thyroid specialist for further evaluation.

Background: Calcitonin values from 20–50 ng/L were found in up to 1% of subjects of the population of 5817 patients with thyroid nodular disease⁹⁰. The predictive value of a C-cell anomaly for this calcitonin level was 8.3%. However, the likelihood of having a medullary carcinoma >1 cm with calcitonin in this range is extremely low.

For calcitonin values between 10-20 ng/L Costante et al.⁹⁰ identified 216 (3.7%) patients. One patient out of the 216 had a subsequent basal (unstimulated) calcitonin value of 33 ng/L, and had C-cell hyperplasia at surgery. Two other studies used a cut-off of calcitonin > 10 ng/L to screen for C-cell disease, but they do not provide sufficient information on patients with basal CT > 10 and < 20 ng/L to allow conclusions^{91,92}.

Appendix 9 Country-specific requirements

Section 5.2 Subject and trial completion

For Mexico: Approximately 60 subjects are planned to be randomised on trial product.

For Japan: Approximately 100 subjects are planned to be randomised on trial product.

Section 6.1 Inclusion criterion no. 2:

For Japan and Taiwan: age ≥ 20 years at the time of signing informed consent

Section 6.2 Exclusion criterion no. 29

Female who is pregnant, breast-feeding or intends to become pregnant or is of child-bearing potential and not using adequate contraceptive methods (highly effective contraceptive measures as required by local regulation or practice)

For Belgium: highly effective methods of birth control are defined as those, alone or in combination, that result in a low failure rate (i.e., less than 1% per year) when used consistently and correctly; such as implants, injectables, combined oral contraceptives, some IUDs, true sexual abstinence (i.e. refraining from heterosexual intercourse during the entire period of risk associated with the study treatments) or vasectomised partner.

For Denmark: Contraceptive measures considered adequate include intrauterine devices or hormonal contraception (oral contraceptive pills, implants, transdermal patches, vaginal rings or long-acting injections).

For Argentina: the following contraceptive measures are considered adequate: barrier methods (condoms or diaphragm) with spermicide, contraceptive pills or intrauterine device (IUD). The contraceptive methods will be reimbursed by the sponsor.

Section 7.5 Preparation/Handling/Storage/Accountability

For Japan: according to Japanese GCP, storage and drug accountability of the trial products at the study site is not in charge of Investigator, but in charge of the head of study site.

The head of study site should assign some or all of the responsibilities for accountability of the trial products at the sites to a trial product storage manager (a pharmacist in principle). The trial product storage manager should control and take accountability of the trial products in accordance with procedures specified by the sponsor. The head of study site or the trial product storage manager must ensure the availability of proper storage conditions, and record and evaluate the temperature.

For India: In-use conditions will be available on the Trial Material Manual.

Section 8.2 Withdrawal from the trial

For Mexico: should the subject his/her family members parents or legal representative decide to withdraw the consent for participation in the trial, the subject will be entitled to receive appropriate, free of charge medical care and/or trial drug during the follow up period of the protocol when it will be established with certainty that no untoward medical consequences of the subject's participation in the research occurred.

Appendix 3, 1) Regulatory and ethical considerations

For Bulgaria: According to Regulation 31 (for defining the rules of Good Clinical Practice, 12 Aug 2007), art. 4, par. (2) - The protocol is prepared according to the requirements of GCP and contains as a minimum the following: 1) assessment of the expected benefits and risks; 2) definition of inclusion and exclusion criteria; 3) rationale for trial population, especially when it is expected to include patients that cannot consent personally and other vulnerable groups; 4) description of the procedures for recruiting patients and getting informed consent when it is expected to include patients who are temporarily or constantly unable to consent personally and when it is expected to receive consent from an independent witness; 5) description of the plan and the procedures for assuring complementary medical cares for the participants after the trial is ended; 6) monitoring procedures; 7) publication policy.

For Mexico: In the case of Mexico, the above will include the following responsibilities for the head of the Institution/Health Care Establishment, Ethics, Research and, when applicable, Biosafety Committees and sponsor within their scope of responsibility:

- a) Investigation follow-up
- b) Damages to health arising from the investigation development; as well as those arising from interruption or advanced suspension of treatment due to non-attributable reasons to the Subject;
- c) Timely compliance of the terms in which the authorization of a research for health in human beings had been issued;
- d) To present in a timely manner the information required by the Health Authority.

For Russia: The trial should be conducted in compliance with the protocol and Ministry of Healthcare of Russian Federation' order #200H from April, 01, 2016"Approval of rules of good clinical practice".

For Japan: A seal is accepted as signature.

Appendix 3, 14) Indemnity statement

For Belgium: Law concerning experiments on the human person of 07 May 2004 - Article 29:

§1. Even if without fault, the sponsor is liable for the damage which the subject and/or his rightful claimants sustain and which shows either a direct or an indirect connection with the trial.

For France: The French Public Health Code article L 1121-10 (law n° 2004-806 of 9 August 2004 art. 88 I, IX, Journal Official of 11 August 2004. "The sponsor is responsible for identification of the harmful consequences of the biomedical the research for the person lending himself thereto and for indemnification of his beneficiaries, except in case of proof, incumbent on it, that the prejudice is not attributable to his fault of or the fault of any intervening party, without the sponsor's being entitled to call on acts by a third party or the voluntary withdrawal of the person who had initially consented to cooperating in the research".

For Mexico:

a) Novo Nordisk carries product liability for its products, and liability as assumed under the special laws, acts/and/or guidelines for conducting clinical trials in any country, including those applicable provisions on the Mexican United States. If the subject feels that something goes wrong during the course of this trial, the subject should contact the trial staff in the first instance.

b) If during their participation in the trial the subject experiences a disease or injury that, according to the trial doctor and the sponsor, is directly caused by the study medication and/or a study procedure that otherwise would not have been part of his/her regular medical care, the subject will receive from the institution or medical care establishment and free of charge, the appropriate medical treatment as required. In this case, the costs resulting from such treatment as well as the costs of any indemnification established by law will be covered by the trial sponsor in accordance with the terms provided by all applicable regulations; even if the subject discontinues his/her participation in the study by his own will or by a decision from the investigator.

c) By signing the informed consent, the subject will not renounce to any compensation or indemnification he/she may be entitled to by law, nor will he/she will incur any additional expense as a result of his/her participation in the trial; any additional expense resulting from the subject's participation in the trial will be covered by the trial sponsor.

For Poland:

Novo Nordisk carries liability for the Trial exclusively in the scope defined by the applicable laws and in particular by the Civil Code and the Pharmaceutical Law dated 6 September 2001 (uniform version Journal of Laws of 2008 No. 45 item 271 with amendments). In order to support potential claims for liability attributable to the Trial, Novo Nordisk and Investigator are covered by the Insurance Policy issued according to applicable Polish law.

Appendix 7 Retention of human biosamples

For Taiwan: No subjects from Taiwan will participate in the optional biobank part of the trial.

12 References

1. Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)*. 2008;32(9):1431-7.
2. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA*. 2014;311(8):806-14.
3. Stevens GA, Singh GM, Lu Y, Danaei G, Lin JK, Finucane MM, et al. National, regional, and global trends in adult overweight and obesity prevalences. *Popul Health Metr*. 2012;10(1):22.
4. NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19·2 million participants. *Lancet*. 2016;387(10026):1377-96.
5. Finkelstein EA, DiBonaventura M, Burgess SM, Hale BC. The costs of obesity in the workplace. *J Occup Environ Med*. 2010;52(10):971-6.
6. Van Nuys K, Globe D, Ng-Mak D, Cheung H, Sullivan J, Goldman D. The association between employee obesity and employer costs: evidence from a panel of U.S. employers. *Am J Health Promot*. 2014;28(5):277-85.
7. Wang YC, Pamplin J, Long MW, Ward ZJ, Gortmaker SL, Andreyeva T. Severe Obesity In Adults Cost State Medicaid Programs Nearly \$8 Billion In 2013. *Health Aff (Millwood)*. 2015;34(11):1923-31.
8. Garner RE, Feeny DH, Thompson A, Bernier J, McFarland BH, Huguet N, et al. Bodyweight, gender, and quality of life: a population-based longitudinal study. *Qual Life Res*. 2012;21(5):813-25.
9. Kearns B, Ara R, Young T, Relton C. Association between body mass index and health-related quality of life, and the impact of self-reported long-term conditions - cross-sectional study from the south Yorkshire cohort dataset. *BMC Public Health*. 2013;13:1009.
10. Cusi K. Role of obesity and lipotoxicity in the development of nonalcoholic steatohepatitis: pathophysiology and clinical implications. *Gastroenterology*. 2012;142(4):711-25.e6.
11. Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;9:88.
12. Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K, et al. General and abdominal adiposity and risk of death in Europe. *N Engl J Med*. 2008;359(20):2105-20.
13. Berghofer A, Pischon T, Reinhold T, Apovian CM, Sharma AM, Willich SN. Obesity prevalence from a European perspective: a systematic review. *BMC Public Health*. 2008;8:200.
14. Masters RK, Reither EN, Powers DA, Yang YC, Burger AE, Link BG. The impact of obesity on US mortality levels: the importance of age and cohort factors in population estimates. *Am J Public Health*. 2013;103(10):1895-901.
15. Arnold M, Pandeya N, Byrnes G, Renehan AG, Stevens GA, Ezzati M, et al. Global burden of cancer attributable to high body-mass index in 2012: a population-based study. *Lancet Oncol*. 2015;16(1):36-46.
16. World Health Organization. Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. World Health Organisation Geneva, Switzerland. 2000.

17. Thomsen M, Nordestgaard BG. Myocardial Infarction and Ischemic Heart Disease in Overweight and Obesity With and Without Metabolic Syndrome. *JAMA internal medicine*. 2013;174(1):15-22.
18. Eckel RH, Kahn SE, Ferrannini E, Goldfine AB, Nathan DM, Schwartz MW, et al. Obesity and type 2 diabetes: what can be unified and what needs to be individualized? *Journal of clinical endocrinology and metabolism*. 2011;96(6):1654-63.
19. Khaodhiar L, Cummings S, Apovian CM. Treating diabetes and prediabetes by focusing on obesity management. *Curr Diab Rep*. 2009;9(5):348-54.
20. Wheaton AG, Perry GS, Chapman DP, Croft JB. Sleep disordered breathing and depression among U.S. adults: National Health and Nutrition Examination Survey, 2005 -2008. *Sleep*. 2012;35(4):461-7.
21. Prospective Studies Collaboration, Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009;373(9669):1083-96.
22. Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA*. 1999;282(16):1523 -9.
23. Church TS, Kuk JL, Ross R, Priest EL, Biltoft E, Biltoff E, et al. Association of cardiorespiratory fitness, body mass index, and waist circumference to nonalcoholic fatty liver disease. *Gastroenterology*. 2006;130(7):2023-30.
24. Blagojevic M, Jinks C, Jeffery A, Jordan KP. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage*. 2010;18(1):24-33.
25. Wei M, Kampert JB, Barlow CE, Nichaman MZ, Gibbons LW, Paffenbarger RS, et al. Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men. *JAMA*. 1999;282(16):1547-53.
26. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, et al. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001;344(18):1343-50.
27. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346(6):393-403.
28. Li G, Zhang P, Wang J, Gregg EW, Yang W, Gong Q, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet*. 2008;371(9626):1783-9.
29. Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*. 2009;374(9702):1677-86.
30. Foster GD, Borradaile KE, Sanders MH, Millman R, Zammit G, Newman AB, et al. A Randomized Study on the Effect of Weight Loss on Obstructive Sleep Apnea Among Obese Patients With Type 2 Diabetes The Sleep AHEAD Study. *Archives of Internal Medicine*. 2009;169(17):1619-26.
31. Felson DT, Zhang Y, Anthony JM, Naimark A, Anderson JJ. Weight loss reduces the risk for symptomatic knee osteoarthritis in women. The Framingham Study. *Ann Intern Med*. 1992;116(7):535-9.

32. Dengo AL, Dennis EA, Orr JS, Marinik EL, Ehrlich E, Davy BM, et al. Arterial destiffening with weight loss in overweight and obese middle-aged and older adults. *Hypertension*. 2010;55(4):855-61.
33. Dattilo AM, Kris-Etherton PM. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr*. 1992;56(2):320-8.
34. Anderson JW, Konz EC. Obesity and disease management: effects of weight loss on comorbid conditions. *Obes Res*. 2001;9 Suppl 4:326S-34S.
35. Garber AJ, Abrahamson MJ, Barzilay JI, Blonde L, Bloomgarden ZT, Bush MA, et al. American Association of Clinical Endocrinologists' Comprehensive Diabetes Management Algorithm 2013 Consensus Statement - Executive Summary. *Endocrine Practice*. 2013;19(3):536-47.
36. Mertens IL, Van Gaal LF. Overweight, obesity, and blood pressure: the effects of modest weight reduction. *Obes Res*. 2000;8(3):270-8.
37. Deitel M, Stone E, Kassam HA, Wilk EJ, Sutherland DJ. Gynecologic-obstetric changes after loss of massive excess weight following bariatric surgery. *J Am Coll Nutr*. 1988;7(2):147-53.
38. Sjöström L. Review of the key results from the Swedish Obese Subjects (SOS) trial - a prospective controlled intervention study of bariatric surgery. *J Intern Med*. 2013;273(3):219-34.
39. Gregg EW, Jakicic JM, Blackburn G, Bloomquist P, Bray GA, Clark JM, et al. Association of the magnitude of weight loss and changes in physical fitness with long-term cardiovascular disease outcomes in overweight or obese people with type 2 diabetes: a post-hoc analysis of the Look AHEAD randomised clinical trial. *Lancet Diabetes Endocrinol*. 2016;4(11):913-21.
40. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, et al. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med*. 2017;377(1):13-27.
41. The American Society for Metabolic and Bariatric Surgery, The Obesity Society, The American Society of Bariatric Physicians and the American Association of Clinical Endocrinologists. Obesity is a Disease: Leading Obesity Groups Agree (Joint Press Release). 19 June 2013.
42. Mechanick JI, Garber AJ, Handelsman Y, Garvey WT. American Association of Clinical Endocrinologists' position statement on obesity and obesity medicine. *Endocr Pract*. 2012;18(5):642-8.
43. Toplak H, Woodward E, Yumuk V, Oppert JM, Halford JC, Frühbeck G. 2014 EASO Position Statement on the Use of Anti-Obesity Drugs. *Obes Facts*. 2015;8(3):166-74.
44. Frühbeck G, Toplak H, Woodward E, Yumuk V, Maislos M, Oppert JM, et al. Obesity: the gateway to ill health - an EASO position statement on a rising public health, clinical and scientific challenge in Europe. *Obes Facts*. 2013;6(2):117-20.
45. Apovian CM, Aronne LJ, Bessesen DH, McDonnell ME, Murad MH, Pagotto U, et al. Pharmacological management of obesity: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2015;100(2):342-62.
46. Ferguson C, David S, Divine L, Kahan S, Gallagher C, Gooding M, et al. Obesity Drug Outcome Measures. A Consensus Report of Considerations Regarding Pharmacologic Intervention. Available from: <https://publichealth.gwu.edu/pdf/obesitydrugmeasures.pdf>. 2012.

47. Sumithran P, Prendergast LA, Delbridge E, Purcell K, Shulkes A, Kriketos A, et al. Long-term persistence of hormonal adaptations to weight loss. *N Engl J Med*. 2011;365(17):1597-604.
48. Schwartz A, Doucet E. Relative changes in resting energy expenditure during weight loss: a systematic review. *Obes Rev*. 2010;11(7):531-47.
49. Pasman WJ, Saris WH, Westerterp-Plantenga MS. Predictors of weight maintenance. *Obes Res*. 1999;7(1):43-50.
50. Dombrowski SU, Knittle K, Avenell A, Araujo-Soares V, Sniehotta FF. Long term maintenance of weight loss with non-surgical interventions in obese adults: systematic review and meta-analyses of randomised controlled trials. *BMJ*. 2014;348:g2646.
51. Brethauer SA, Chand B, Schauer PR. Risks and benefits of bariatric surgery: current evidence. *Cleve Clin J Med*. 2006;73(11):993-1007.
52. Schauer PR, Ikramuddin S, Gourash W, Ramanathan R, Luketich J. Outcomes after laparoscopic Roux-en-Y gastric bypass for morbid obesity. *Ann Surg*. 2000;232(4):515-29.
53. Yumuk V, Tsigos C, Fried M, Schindler K, Busetto L, Micic D, et al. European Guidelines for Obesity Management in Adults. *Obes Facts*. 2015;8(6):402-24.
54. Clarke WL, Cox DJ, Gonder-Frederick LA, Julian D, Schlundt D, Polonsky W. Reduced awareness of hypoglycemia in adults with IDDM. A prospective study of hypoglycemic frequency and associated symptoms. *Diabetes Care*. 1995;18(4):517-22.
55. Lau J, Bloch P, Schäffer L, Pettersson I, Spetzler J, Kofoed J, et al. Discovery of the Once-Weekly Glucagon-Like Peptide-1 (GLP-1) Analogue Semaglutide. *J Med Chem*. 2015;58(18):7370-80.
56. Gutzwiller JP, Drewe J, Goke B, Schmidt H, Rohrer B, Lareida J, et al. Glucagon-like peptide-1 promotes satiety and reduces food intake in patients with diabetes mellitus type 2. *Am J Physiol*. 1999;276(5 Pt 2):R1541-4.
57. Novo Nordisk A/S. Investigator's Brochure, Semaglutide s.c. 2.4 mg, project NN9536, (edition 1, draft version 0.1). 2017.
58. Sorli C, Harashima SI, Tsoukas GM, Unger J, Karsbøl JD, Hansen T, et al. Efficacy and safety of once-weekly semaglutide monotherapy versus placebo in patients with type 2 diabetes (SUSTAIN 1): a double-blind, randomised, placebo-controlled, parallel-group, multinational, multicentre phase 3a trial. *Lancet Diabetes Endocrinol*. 2017;5(4):251-60.
59. Aroda VR, Bain SC, Cariou B, Piletič M, Rose L, Axelsen M, et al. Efficacy and safety of once-weekly semaglutide versus once-daily insulin glargine as add-on to metformin (with or without sulfonylureas) in insulin-naïve patients with type 2 diabetes (SUSTAIN 4): a randomised, open-label, parallel-group, multicentre, multinational, phase 3a trial. *Lancet Diabetes Endocrinol*. 2017;5(5):355-66.
60. Ahrén B, Masmiquel L, Kumar H, Sargin M, Karsbøl JD, Jacobsen SH, et al. Efficacy and safety of once-weekly semaglutide versus once-daily sitagliptin as an add-on to metformin, thiazolidinediones, or both, in patients with type 2 diabetes (SUSTAIN 2): a 56-week, double-blind, phase 3a, randomised trial. *Lancet Diabetes Endocrinol*. 2017;5(5):341-54.
61. Ahmann A, Capehorn M, Charpentier G, Dotta F, Henkel E, Lingvay I, et al. Efficacy and safety of once-weekly semaglutide vs exenatide ER after 56 Weeks in subjects with type 2 diabetes (SUSTAIN 3). European Association for the Study of Diabetes, 52nd meeting 2016, Oral Presentation #1472016.

62. Rodbard H, Lingvay I, Reed J, de la Rosa R, Rose L, Sugimoto D, et al. Efficacy and safety of semaglutide once-weekly vs placebo as add-on to basal insulin alone or in combination with metformin in subjects with type 2 diabetes (SUSTAIN 5) [abstract]. *Diabetologia*. 2016;59(Suppl 1):364-5.
63. Blundell J, Finlayson G, Axelsen M, Flint A, Gibbons C, Kvist T, et al. Effects of once-weekly semaglutide on appetite, energy intake, control of eating, food preference and body weight in subjects with obesity. *Diabetes Obes Metab*. 2017.
64. European Medicines Agency. EMA/CHMP/311805/2014; Guideline on clinical evaluation of medicinal products used in weight management. 01 Jan 2017.
65. Food and Drug Administration. FDA Guidance for Industry: Developing products for weight management. 2007.
66. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA*. 2013;309(1):71-82.
67. Prospective Studies C, Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009;373(9669):1083-96.
68. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int Suppl*. 2013;3(1):1-150.
69. Novo Nordisk A/S. Investigator's Brochure, Semaglutide s.c. 2.4 mg, project NN9536, (edition 1). 28 Nov 2017.
70. FAO/WHO/UNU. Human energy requirements. Report of a joint FAO/WHO/UNU expert consultation. FAO: food and nutrition technical report series 1. Rome: 2004 2004.
71. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, et al. Classification of acute pancreatitis-2012: revision of the Atlanta classification and definitions by international consensus. *Gut*. 2013;62(1):102-11.
72. American Diabetes Association. Standards of medical care in diabetes - 2017. *Diabetes Care*. 2017;40(Suppl 1):S1-135.
73. Kolotkin RL, Ervin CM, Meincke HH, Højbjerg L, Fehnel SE. Development of a clinical trials version of the Impact of Weight on Quality of Life-Lite questionnaire (IWQOL-Lite Clinical Trials Version): results from two qualitative studies. *Clin Obes*. 2017;7(5):290-9.
74. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*. 1992;30(6):473-83.
75. Koopman C, Pelletier KR, Murray JF, Sharda CE, Berger ML, Turpin RS, et al. Stanford presenteeism scale: health status and employee productivity. *J Occup Environ Med*. 2002;44(1):14-20.
76. Avery K, Donovan J, Peters TJ, Shaw C, Gotoh M, Abrams P. ICIQ: a brief and robust measure for evaluating the symptoms and impact of urinary incontinence. *Neurourol Urodyn*. 2004;23(4):322-30.
77. Kroenke K, Spitzer RL, Williams JB. The PHQ-9 - Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606-13.
78. Posner K, Oquendo MA, Gould M, Stanley B, Davies M. Columbia Classification Algorithm of Suicide Assessment (C-CASA): classification of suicidal events in the FDA's pediatric suicidal risk analysis of antidepressants. *Am J Psychiatry*. 2007;164(7):1035-43.

79. McEvoy BW. Missing data in clinical trials for weight management. *J Biopharm Stat.* 2016;26(1):30-6.
80. Carpenter JR, Roger JH, Kenward MG. Analysis of longitudinal trials with protocol deviation: a framework for relevant, accessible assumptions, and inference via multiple imputation. *J Biopharm Stat.* 2013;23(6):1352-71.
81. Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009;360(9):859-73.
82. World Medical Association. Declaration of Helsinki - Ethical Principles for Medical Research Involving Human Subjects. Last amended by the 64th WMA General Assembly, Fortaleza, Brazil. 1 Oct 2013.
83. ICH Harmonised Tripartite Guideline. Guideline for Good Clinical Practice E6(R2), Current Step 4 version. 09 Nov 2016.
84. Committee for Medicinal Products for Human Use (CHMP). Guideline on clinical evaluation of medicinal products used in weight management. EMA/CHMP/311805/2014. 2016.
85. International Committee of Medical Journal Editors. Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals; current version available at www.icmje.org.
86. De Angelis C, Drazen JM, Frizelle FA, Haug C, Hoey J, Horton R, et al. Clinical trial registration: a statement from the International Committee of Medical Journal Editors. *N Engl J Med.* 2004;351(12):1250-1.
87. U.S. Food and Drug Administration. Food and Drug Administration Amendments Act of 2007 as amended by the Final Rule "Clinical Trials Registration and Results Information Submission". 21 September 2016.
88. The European Parliament and the Council of the European Council. Directive 2001/20/EC of the European Parliament and of the Council of 4 April 2001 on the approximation of the laws, regulations and administrative provisions of the Member States relating to the implementation of good clinical practice in the conduct of clinical trials on medicinal products for human use, article 11. *Official Journal of the European Communities.* 01 May 2001.
89. The European Parliament and the Council of the European Council. Regulation (EC) No 726/2004 of the European Parliament and of the Council of 31 March 2004 laying down Community procedures for the authorisation and supervision of medicinal products for human and veterinary use and establishing a European Medicines Agency, article 57. 30 April 2004.
90. Costante G, Meringolo D, Durante C, Bianchi D, Nocera M, Tumino S, et al. Predictive value of serum calcitonin levels for preoperative diagnosis of medullary thyroid carcinoma in a cohort of 5817 consecutive patients with thyroid nodules. *J Clin Endocrinol Metab.* 2007;92(2):450-5.
91. Scheuba C, Kaserer K, Moritz A, Drosten R, Vierhapper H, Bieglmayer C, et al. Sporadic hypercalcitoninemia: clinical and therapeutic consequences. *Endocr Relat Cancer.* 2009;16(1):243-53.

Protocol
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

21 December 2017
2.0
Final
104 of 104

Novo Nordisk

92. Verga U, Ferrero S, Vicentini L, Brambilla T, Cirello V, Muzza M, et al. Histopathological and molecular studies in patients with goiter and hypercalcitoninemia: reactive or neoplastic C-cell hyperplasia? *Endocr Relat Cancer*. 2007;14(2):393-403.

Semaglutide s.c. 2.4 mg once weekly
Trial ID: NN9536-4373 (STEP 1)
Clinical Trial Report
Appendix 16.1.1

~~CONFIDENTIAL~~

Date:
Version:
Status:

11 Jun 2020
1.0
Final

Novo Nordisk

Global and country key Novo Nordisk staff

Attachments I and II (if applicable) to the protocol are located in the Trial Master File.

Content: Global key staff and Country key staff

Protocol Amendment no 2
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

15 March 2018
1.0
Final
1 of 3

Novo Nordisk

Protocol Amendment
no 2
to Protocol, version 2.0
dated 21 December 2017

Trial ID: NN9536-4373

**Title of protocol: Effect and safety of semaglutide 2.4 mg once-weekly in subjects with
overweight or obesity**

Trial phase: 3a
Applicable to: UK

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Protocol Amendment no 2
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

| | | |
|----------|---------------|---------------------|
| Date: | 15 March 2018 | Novo Nordisk |
| Version: | 1.0 | |
| Status: | Final | |
| Page: | 2 of 3 | |

Table of Contents

1 Introduction including rationale for the protocol amendment.....3

2 Changes3

1 Introduction including rationale for the protocol amendment

Section 9.4 has been updated to include location of flow chart

The use of double barrier methods mentioned in Appendix 5 is not applicable for UK. Therefore appendix 9 ‘country specific requirements’ is updated and must be followed in UK.

- Any new text is written *in italics*.
- Any text deleted from the protocol is written using strike through.

2 Changes

9.4 Safety assessments.

Planned time points for all safety assessments are provided in the flowchart. (*Section 2*)

Appendix 9 Country specific requirements will be updated with the following:

For UK: Contraceptive measures considered adequate include highly effective contraceptive methods as listed in Table 11-5 ‘Highly effective contraceptive methods’ in Appendix 5.

This means that the use of double barrier methods is not applicable for UK.

Protocol Amendment no 3
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

04 June 2018
1.0
Final
1 of 8

Novo Nordisk

Protocol Amendment
no 3
to Protocol, version 2.0
dated 21 Dec 2017

Trial ID:NN9536-4373

Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity

Trial phase: 3a
Applicable to all countries

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Table of Contents

| | | |
|----------|---|----------|
| 1 | Introduction including rationale for the protocol amendment..... | 3 |
| 2 | Changes | 3 |
| 2.1 | Section 2 Flowchart..... | 3 |
| 2.2 | Section 3.2 Background..... | 4 |
| 2.2.1 | Section 3.2.1 Semaglutide | 4 |
| 2.2.2 | 3.2.2 Trial population..... | 4 |
| 2.3 | 3.3 Benefit-risk assessment | 4 |
| 2.3.1 | 3.3.2 Risks and precautions | 4 |
| 2.4 | Section 7.1 Treatments administered | 5 |
| 2.5 | Section 8 Discontinuation/Withdrawal criteria..... | 5 |
| 2.5.1 | Section 8.1 Discontinuation of trial treatment | 5 |
| 2.6 | Section 9.7 Genetics..... | 6 |
| 2.7 | Section 9.8 Biomarkers <i>and biosamples for future analysis</i> | 6 |
| 2.8 | Section 10.1 Sample size determination..... | 7 |
| 2.8.1 | Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects | 7 |
| 2.9 | Section 10.3 Statistical analyses | 7 |
| 2.9.1 | Section 10.3.1 Primary endpoints | 7 |
| 2.9.1.1 | Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy | 7 |
| 2.10 | Appendix 2 Clinical laboratory tests | 7 |
| 2.10.1 | Table 11-3 Criteria for laboratory outliers | 7 |
| 2.11 | Appendix 3 Trial governance considerations | 7 |
| 2.11.1 | 9) Data quality assurance | 7 |
| 2.11.1.1 | Protocol compliance..... | 7 |
| 2.12 | Appendix 5 Contraceptive guidance and collection of pregnancy information..... | 8 |
| 2.12.1 | Table 11-5 Highly effective contraceptive methods | 8 |
| 2.13 | Appendix 7 Retention of human biosamples <i>and genetics</i> | 8 |
| 2.13.1 | Biosamples for future analysis | 8 |
| 2.14 | Section 12 References | 8 |

1 Introduction including rationale for the protocol amendment

This protocol is amended for the following reasons:

- To implement genetic biosamples for future analysis for those countries where it will be applicable.
The rationale for implementation is to ensure a better characterisation of the population in scope as new technologies and hypothesis develops. The collection of whole blood at baseline (visit 2) and end of treatment (visit 24) will allow for future genetic analysis thereby increasing the understanding of obesity, obesity treatment and related diseases in the future.
- As the scope of the biosampling for future analysis has broadened to cover obesity and - related diseases with this amendment, some paragraphs in appendix 7 have been revised. Furthermore, the informed consent has been updated to reflect the broader scope. A separate protocol will be developed before the biosamples from NN9536-4373 can be analysed in the future. This will include rationale for the analysis as well as reporting requirements.
- Removal of criteria for discontinuation of trial treatment for subjects included in the trial in violation of the inclusion and/or exclusion criteria and/or randomisation criteria. Subjects will not be discontinued from trial product if considered safe to continue. However, prospective approval of protocol deviations to recruitment and enrolment criteria, also known as protocol waivers or exemptions, is not permitted and deviations from the protocol should be avoided.
- Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment.
- Minor clarification of content and correction of minor errors and typos.

In this protocol amendment:

- Any new text is written *in italics*.
- Any text deleted from the protocol is written using ~~strike through~~.

2 Changes

2.1 Section 2 Flowchart

Tobacco Use^c (~~6.4.1~~)

Biosamples for future analysis^f (9.7, 9.8, Appendix 7)

2.2 Section 3.2 Background**2.2.1 Section 3.2.1 Semaglutide**

GLP-1 RA is a physiological regulator of appetite and GLP-1 receptors are present in several areas of the brain involved in appetite regulation⁵⁶.

Clinical⁵⁸⁻⁶²³ and non-clinical data⁶²⁶⁴ indicate that the body weight-reducing effect of semaglutide is mainly mediated by a reduced energy intake. No unexpected safety findings were identified and the tolerability and safety profile was overall consistent with previous findings in the T2D development programme and the GLP-1 RA class in general.

2.2.2 3.2.2 Trial population

The trial population will consist of subjects with obesity (BMI ≥ 30 kg/m²) or with overweight (BMI ≥ 27 kg/m²) and weight-related comorbidities.

2.3 3.3 Benefit-risk assessment**2.3.1 3.3.2 Risks and precautions**

The sections below describe identified and potential risks associated with semaglutide treatment. *For classification and further details of the risks, please refer to the current version of the IB or any updates hereof.* The identified/potential risks are based on findings in non-clinical studies and clinical trials with semaglutide as well as other GLP-1 RAs. For each of these risks, mitigating actions have been implemented to minimise the risks for subjects enrolled in this trial.

~~Identified risks~~**Gastrointestinal adverse events**

- Consistent with findings with other GLP-1 RAs, the most frequently reported adverse events (AE) in clinical trials with semaglutide were gastrointestinal AEs. A low starting dose and dose escalation steps will be implemented in the trial to mitigate the risk of gastrointestinal AEs.

~~Potential risks~~**Medullary thyroid cancer (MTC) (based on non-clinical data)**

- Expected proliferative thyroid C-cell changes were seen in the mouse and rat carcinogenicity studies after daily exposure to semaglutide for 2 years. No hyperplasia was observed in monkeys after 52 weeks exposure up to 13-fold above the clinical plasma exposure at 2.4 mg/week. In clinical trials with semaglutide, there have been no reports of MTC or clinically relevant changes in calcitonin levels. The C-cell changes in rodents are mediated by the GLP-1 receptor, which is not expressed in the normal human thyroid.

Accordingly, the risk of GLP-1 receptor-mediated C-cell changes in humans is considered to be low. However, as a precaution, exclusion and discontinuation criteria related to medical history of multiple endocrine neoplasia type 2 (MEN2) or MTC and elevated plasma levels of calcitonin (biomarker for MTC) have been implemented in the trial.

Other risks

Pregnancy and fertility (based on non-clinical data)

- Studies in animals have shown reproductive toxicity. There are limited data from the use of semaglutide in pregnant women. Therefore, semaglutide should not be used during pregnancy. Exclusion and discontinuation criteria related to pregnancy have been implemented in the trial.

2.4 Section 7.1 Treatments administered

- Injections may be administered in the thigh, abdomen or upper arm, at any time of day irrespective of meals. ~~Subjects should be encouraged to inject in the same area throughout the trial, but changing between left and right side is allowed.~~

2.5 Section 8 Discontinuation/Withdrawal criteria

2.5.1 Section 8.1 Discontinuation of trial treatment

The subject must be discontinued from trial product, if any of the following applies:

- ~~1. Included in the trial in violation of the inclusion and/or exclusion criteria and/or randomisation criteria~~
- ~~1.~~ Safety concern as judged by the investigator
- ~~2.~~ Calcitonin ≥ 100 ng/L (see Appendix 8)
- ~~3.~~ Suspicion of pancreatitis
- ~~4.~~ Pregnancy
- ~~5.~~ Intention of becoming pregnant
- ~~6.~~ Simultaneous participation in another clinical trial of an approved or non-approved Investigational medicinal product.

Subjects meeting discontinuation of trial product criterion no. ~~4~~ 3 are allowed to resume trial product if the Atlanta criteria⁷¹ are not fulfilled and thus the suspicion of acute pancreatitis is not confirmed.

Trial product may be resumed for subjects with a gallstone-induced pancreatitis in case of cholecystectomy.

Subjects meeting discontinuation of trial product criteria no. ~~2~~ 1, ~~5~~ 4 and ~~6~~ 5 are allowed to resume trial product, if the criteria are no longer met (see Section 8.1.1).

2.6 Section 9.7 Genetics

~~Not applicable for this trial.~~

A blood sample for genetic analysis will be collected from subjects who have consented to participate in the optional biobank component of the trial. Refer to section 9.8 and Appendix 7 for further details.

2.7 Section 9.8 Biomarkers and biosamples for future analysis

Collection of samples for biomarker research is part of this trial to support the efficacy objectives. The following samples must be collected in accordance with the laboratory manual and the flowchart:

Biomarkers linked to cardiovascular risk:

- *Plasminogen Activator Inhibitor-1 (PAI-1)* Activity will be analysed by activity assay.
- *High sensitive C-reactive protein (hsCRP)*

Collection of biosamples for future analysis (stored in a biobank) is a component of this trial. Participation in the biobank component is optional. Subjects who do not wish to participate in the biobank component may still participate in the trial. For the biobank, samples will be collected according to the flowchart and stored for future use.

The samples are collected for the purpose of allowing future analyses of biomarkers, both genetic and circulating, at a later point in time when new knowledge or improved measurement techniques may have become available. The analyses may include biomarkers currently known or discovered in the future.

Genetic analyses may include analysis of candidate genes or genetic markers throughout the genome with the purpose of understanding and predicting response to semaglutide as well as to understand obesity or other related diseases.

Analyses of circulating biomarkers will measure hormones, metabolites or other non-genetic serum entity with the purpose of understanding and predicting response to semaglutide as well as understanding obesity or other related diseases.

These samples need to be frozen and should be sent in batches to the central laboratory. The analyses are likely to be performed after the trial has come to an end, and results will therefore not be part of the clinical trial report. The biobank samples may be stored up to 15 years after end of trial at a central laboratory (see Appendix 7).

2.8 Section 10.1 Sample size determination**2.8.1 Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects**

Foot note: SD = standard deviation; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain; # shown as a positive number

All tests in the hierarchy are based on the primary estimand

2.9 Section 10.3 Statistical analyses**2.9.1 Section 10.3.1 Primary endpoints****2.9.1.1 Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy**

Foot note: FAS = full analysis set; ANCOVA = analysis of covariance; RD-MI = multiple imputation using retrieved subjects; J2R-MI = jump to reference multiple imputation; S1-SI and S2-SI = single imputation as done by Sacks; TP-MI = tipping point multiple imputation; MMRM = mixed model for repeated measurements; LR = logistic regression; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain

Test order refers to the order of the endpoint in the statistical test hierarchy outlined in Table 10-2

2.10 Appendix 2 Clinical laboratory tests**2.10.1 Table 11-3 Criteria for laboratory outliers**

Lymphocytes < ~~25~~ $0.2 \times 10^9/L$

2.11 Appendix 3 Trial governance considerations**2.11.1 9) Data quality assurance****2.11.1.1 Protocol compliance**

Deviations from the protocol should be avoided. If deviations do occur, the investigator must inform the monitor *without delay* and the implications of the deviation must be reviewed and discussed.

2.12 Appendix 5 Contraceptive guidance and collection of pregnancy information**2.12.1 Table 11-5 Highly effective contraceptive methods**Sexual abstinence^b**2.13 Appendix 7 Retention of human biosamples *and genetics*****2.13.1 Biosamples for future analysis**

In countries where applicable, the trial will involve collection of human biosamples to be stored in a central archive for future use.

~~The material to be collected is serum collected at baseline (visit 2) and end of treatment (visit 24) is: A total of 5 aliquots will be stored for each subject at both time points.~~

- *Whole blood (for genetic analysis)*
- *Serum (for analyses of circulating biomarkers)*

~~The results obtained from any already performed analyses of the samples will still be used.~~

~~In the event that the collected biosamples (serum) will be used in the future, the investigator will become directly informed by Novo Nordisk about the results if the findings are deemed clinically relevant and analytically valid and quantifiable. In such case, a written summary of the findings, including listings of subject specific values, will be provided once a firm conclusion from the results has been drawn by Novo Nordisk.~~

~~Potentially, observations of neoplastic diseases, serious hereditary diseases, other un-treatable diseases, or any other abnormal findings could be part of the observations. Subjects can contact the investigator if they wish to be informed about results derived from stored biosamples obtained from their own body.~~

In case the subject withdraws his/her informed consent for biosamples for future analysis and genetics, the monitor must contact the trial manager at Novo Nordisk as soon as possible in order to have the samples withdrawn from storage.

2.14 Section 12 References

64. Lu T, et al. Presented at the 77th scientific sessions of the American diabetes association, 9-13 June 2017, San Diego, CA, USA.

57. Novo Nordisk A/S. Investigator's Brochure, Semaglutide s.c. 2.4 mg, project NN9536, (edition 1, ~~draft version 0.1~~). 28 Nov 2017.

Protocol Amendment no 4-DE
Trial ID: NN9536-4373
EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

19 July 2018
1.0
Final
1 of 8

Novo Nordisk

Protocol Amendment

No 4

**to Protocol, version 2.0
dated 21 Dec 2017**

Trial ID: NN9536-4373

Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity

**Trial phase: 3a
Applicable to Germany**

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date:
 Version:
 Status:
 Page:

19 July 2018
 1.0
 Final
 2 of 8

Novo Nordisk

Table of Contents

| | | |
|----------|---|----------|
| 1 | Introduction including rationale for the protocol amendment..... | 3 |
| 2 | Changes | 3 |
| 2.1 | Section 2 Flowchart..... | 3 |
| 2.2 | Section 3.2 Background..... | 3 |
| 2.2.1 | Section 3.2.1 Semaglutide | 3 |
| 2.2.2 | 3.2.2 Trial population..... | 4 |
| 2.3 | 3.3 Benefit-risk assessment | 4 |
| 2.3.1 | 3.3.2 Risks and precautions | 4 |
| 2.4 | Section 7.1 Treatments administered | 5 |
| 2.5 | Section 9.7 Genetics | 5 |
| 2.6 | Section 9.8 Biomarkers and biosamples for future analysis | 5 |
| 2.7 | Section 10.1 Sample size determination..... | 6 |
| 2.7.1 | Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects | 6 |
| 2.8 | Section 10.3 Statistical analyses | 6 |
| 2.8.1 | Section 10.3.1 Primary endpoints | 6 |
| 2.8.1.1 | Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy | 6 |
| 2.9 | Appendix 2 Clinical laboratory tests | 6 |
| 2.9.1 | Table 11-3 Criteria for laboratory outliers | 6 |
| 2.10 | Appendix 3 Trial governance considerations | 7 |
| 2.10.1 | 9) Data quality assurance | 7 |
| 2.10.1.1 | Protocol compliance..... | 7 |
| 2.11 | Appendix 5 Contraceptive guidance and collection of pregnancy information..... | 7 |
| 2.11.1 | Table 11-5 Highly effective contraceptive methods | 7 |
| 2.12 | Appendix 7 Retention of human biosamples <i>and genetics</i> | 7 |
| 2.12.1 | Biosamples for future analysis | 7 |
| 2.13 | Section 12 References | 8 |

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date:
 Version:
 Status:
 Page:

19 July 2018
 1.0
 Final
 3 of 8

Novo Nordisk

1 Introduction including rationale for the protocol amendment

This protocol is amended for the following reasons:

- To implement genetic biosamples for future analysis for those countries where it will be applicable.

The rationale for implementation is to ensure a better characterisation of the population in scope as new technologies and hypothesis develops. The collection of whole blood at baseline (visit 2) and end of treatment (visit 24) will allow for future genetic analysis thereby increasing the understanding of obesity, obesity treatment and related diseases in the future.

- As the scope of the biosampling for future analysis has broadened to cover obesity and -related diseases with this amendment, some paragraphs in appendix 7 have been revised. Furthermore, the informed consent has been updated to reflect the broader scope. A separate protocol will be developed before the biosamples from NN9536-4373 can be analysed in the future. This will include rationale for the analysis as well as reporting requirements.
- Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment.
- Minor clarification of content and correction of minor errors and typos.

In this protocol amendment:

Any new text is written *in italics*.

Any text deleted from the protocol is written using ~~strike-through~~.

2 Changes

2.1 Section 2 Flowchart

Tobacco Use^c (~~6.4.1~~)

Biosamples for future analysis^f (9.7, 9.8, Appendix 7)

2.2 Section 3.2 Background

2.2.1 Section 3.2.1 Semaglutide

GLP-1 ~~RA~~ is a physiological regulator of appetite and GLP-1 receptors are present in several areas of the brain involved in appetite regulation⁵⁶.

Clinical⁵⁸⁻⁶²³ and non-clinical data⁶²⁶⁴ indicate that the body weight-reducing effect of semaglutide is mainly mediated by a reduced energy intake. No unexpected safety findings were identified and

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date: 19 July 2018
 Version: 1.0
 Status: Final
 Page: 4 of 8

Novo Nordisk

the tolerability and safety profile was overall consistent with previous findings in the T2D development programme and the GLP-1 RA class in general.

2.2.2 3.2.2 Trial population

The trial population will consist of subjects with obesity (BMI ≥ 30 kg/m²) or with overweight (BMI ≥ 27 kg/m²) and weight-related comorbidities.

2.3 3.3 Benefit-risk assessment

2.3.1 3.3.2 Risks and precautions

The sections below describe identified and potential risks associated with semaglutide treatment. *For classification and further details of the risks, please refer to the current version of the IB or any updates hereof.* The identified/potential risks are based on findings in non-clinical studies and clinical trials with semaglutide as well as other GLP-1 RAs. For each of these risks, mitigating actions have been implemented to minimise the risks for subjects enrolled in this trial.

~~Identified risks~~

Gastrointestinal adverse events

- Consistent with findings with other GLP-1 RAs, the most frequently reported adverse events (AE) in clinical trials with semaglutide were gastrointestinal AEs. A low starting dose and dose escalation steps will be implemented in the trial to mitigate the risk of gastrointestinal AEs.

~~Potential risks~~

Medullary thyroid cancer (MTC) (based on non-clinical data)

- Expected proliferative thyroid C-cell changes were seen in the mouse and rat carcinogenicity studies after daily exposure to semaglutide for 2 years. No hyperplasia was observed in monkeys after 52 weeks exposure up to 13-fold above the clinical plasma exposure at 2.4 mg/week. In clinical trials with semaglutide, there have been no ~~reports of MTC or~~ clinically relevant changes in calcitonin levels. The C-cell changes in rodents are mediated by the GLP-1 receptor, which is not expressed in the normal human thyroid. Accordingly, the risk of GLP-1 receptor-mediated C-cell changes in humans is considered to be low. However, as a precaution, exclusion and discontinuation criteria related to medical history of multiple endocrine neoplasia type 2 (MEN2) or MTC and elevated plasma levels of calcitonin (biomarker for MTC) have been implemented in the trial.

~~Other risks~~

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date: 19 July 2018
 Version: 1.0
 Status: Final
 Page: 5 of 8

Novo Nordisk

Pregnancy and fertility (based on non-clinical data)

- Studies in animals have shown reproductive toxicity. There are limited data from the use of semaglutide in pregnant women. Therefore, semaglutide should not be used during pregnancy. Exclusion and discontinuation criteria related to pregnancy have been implemented in the trial.

2.4 Section 7.1 Treatments administered

- Injections may be administered in the thigh, abdomen or upper arm, at any time of day irrespective of meals. ~~Subjects should be encouraged to inject in the same area throughout the trial, but changing between left and right side is allowed.~~

2.5 Section 9.7 Genetics

~~Not applicable for this trial.~~

A blood sample for genetic analysis will be collected from subjects who have consented to participate in the optional biobank component of the trial. Refer to section 9.8 and Appendix 7 for further details.

2.6 Section 9.8 Biomarkers and biosamples for future analysis

Collection of samples for biomarker research is part of this trial to support the efficacy objectives. The following samples must be collected in accordance with the laboratory manual and the flowchart:

Biomarkers linked to cardiovascular risk:

*Plasminogen Activator Inhibitor-1 (PAI-1) Activity will be analysed by activity assay.
 High sensitive C-reactive protein (hsCRP)*

Collection of biosamples for future analysis (stored in a biobank) is a component of this trial. Participation in the biobank component is optional. Subjects who do not wish to participate in the biobank component may still participate in the trial. For the biobank, samples will be collected according to the flowchart and stored for future use.

The samples are collected for the purpose of allowing future analyses of biomarkers, both genetic and circulating, at a later point in time when new knowledge or improved measurement techniques may have become available. The analyses may include biomarkers currently known or discovered in the future.

Genetic analyses may include analysis of candidate genes or genetic markers throughout the genome with the purpose of understanding and predicting response to semaglutide as well as to understand obesity or other related diseases.

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date: 19 July 2018
 Version: 1.0
 Status: Final
 Page: 6 of 8

Novo Nordisk

Analyses of circulating biomarkers will measure hormones, metabolites or other non-genetic serum entity with the purpose of understanding and predicting response to semaglutide as well as understanding obesity or other related diseases.

These samples need to be frozen and should be sent in batches to the central laboratory. The analyses are likely to be performed after the trial has come to an end, and results will therefore not be part of the clinical trial report. The biobank samples may be stored up to 15 years after end of trial at a central laboratory (see Appendix 7).

2.7 Section 10.1 Sample size determination

2.7.1 Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects

Foot note: SD = standard deviation; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain; # shown as a positive number

All tests in the hierarchy are based on the primary estimand

2.8 Section 10.3 Statistical analyses

2.8.1 Section 10.3.1 Primary endpoints

2.8.1.1 Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy

Foot note: FAS = full analysis set; ANCOVA = analysis of covariance; RD-MI = multiple imputation using retrieved subjects; J2R-MI = jump to reference multiple imputation; S1-SI and S2-SI = single imputation as done by Sacks; TP-MI = tipping point multiple imputation; MMRM = mixed model for repeated measurements; LR = logistic regression; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain

Test order refers to the order of the endpoint in the statistical test hierarchy outlined in Table 10-2

2.9 Appendix 2 Clinical laboratory tests

2.9.1 Table 11-3 Criteria for laboratory outliers

Lymphocytes < ~~25~~ $0.2 \times 10^9/L$

Protocol Amendment no 4-DE
 Trial ID: NN9536-4373
 EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

Date:
 Version:
 Status:
 Page:

19 July 2018
 1.0
 Final
 7 of 8

Novo Nordisk

2.10 Appendix 3 Trial governance considerations

2.10.1 9) Data quality assurance

2.10.1.1 Protocol compliance

Deviations from the protocol should be avoided. If deviations do occur, the investigator must inform the monitor *without delay* and the implications of the deviation must be reviewed and discussed.

2.11 Appendix 5 Contraceptive guidance and collection of pregnancy information

2.11.1 Table 11-5 Highly effective contraceptive methods

Sexual abstinence^b

2.12 Appendix 7 Retention of human biosamples *and genetics*

2.12.1 Biosamples for future analysis

In countries where applicable, the trial will involve collection of human biosamples to be stored in a central archive *for future use*.

The material to be collected is ~~serum collected~~ at baseline (*visit 2*) and end of treatment (*visit 24*) *is*:
~~A total of 5 aliquots will be stored for each subject at both time points.~~

Whole blood (for genetic analysis)

Serum (for analyses of circulating biomarkers)

~~The results obtained from any already performed analyses of the samples will still be used. In the event that the collected biosamples (serum) will be used in the future, the investigator will become directly informed by Novo Nordisk about the results if the findings are deemed clinically relevant and analytically valid and quantifiable. In such case, a written summary of the findings, including listings of subject specific values, will be provided once a firm conclusion from the results has been drawn by Novo Nordisk.~~

~~Potentially, observations of neoplastic diseases, serious hereditary diseases, other un-treatable diseases, or any other abnormal findings could be part of the observations. Subjects can contact the investigator if they wish to be informed about results derived from stored biosamples obtained from their own body.~~

In case the subject withdraws his/her informed consent for biosamples for future analysis and genetics, the monitor must contact the trial manager at Novo Nordisk as soon as possible in order to have the samples withdrawn from storage.

Protocol Amendment no 4-DE
Trial ID: NN9536-4373
EudraCT-No.: 2017-003436-36

~~CONFIDENTIAL~~

| | |
|----------|--------------|
| Date: | 19 July 2018 |
| Version: | 1.0 |
| Status: | Final |
| Page: | 8 of 8 |

Novo Nordisk

2.13 Section 12 References

64. *Lu T, et al. Presented at the 77th scientific sessions of the American diabetes association, 9-13 June 2017, San Diego, CA, USA.*

57. Novo Nordisk A/S. Investigator's Brochure, Semaglutide s.c. 2.4 mg, project NN9536, (edition 1, ~~draft version 0.1~~). 28 Nov 2017.

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
1 of 8

Novo Nordisk

Protocol Amendment
no 5
to Protocol, version 2.0
dated 21 Dec 2017

Trial ID:NN9536-4373

Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity

Trial phase: 3a
Applicable to the UK

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Table of Contents

| | | |
|----------|---|----------|
| 1 | Introduction including rationale for the protocol amendment..... | 3 |
| 2 | Changes | 3 |
| 2.1 | Section 2 Flowchart..... | 3 |
| 2.2 | Section 3.2 Background..... | 3 |
| 2.2.1 | Section 3.2.1 Semaglutide | 3 |
| 2.2.2 | 3.2.2 Trial population..... | 4 |
| 2.3 | 3.3 Benefit-risk assessment | 4 |
| 2.3.1 | 3.3.2 Risks and precautions | 4 |
| 2.4 | Section 7.1 Treatments administered | 5 |
| 2.5 | Section 9.7 Genetics | 5 |
| 2.6 | Section 9.8 Biomarkers <i>and biosamples for future analysis</i> | 5 |
| 2.7 | Section 10.1 Sample size determination..... | 6 |
| 2.7.1 | Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects | 6 |
| 2.8 | Section 10.3 Statistical analyses | 6 |
| 2.8.1 | Section 10.3.1 Primary endpoints | 6 |
| 2.8.1.1 | Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy | 6 |
| 2.9 | Appendix 2 Clinical laboratory tests | 7 |
| 2.9.1 | Table 11-3 Criteria for laboratory outliers | 7 |
| 2.10 | Appendix 3 Trial governance considerations | 7 |
| 2.10.1 | 9) Data quality assurance | 7 |
| 2.10.1.1 | Protocol compliance | 7 |
| 2.11 | Appendix 5 Contraceptive guidance and collection of pregnancy information..... | 7 |
| 2.11.1 | Table 11-5 Highly effective contraceptive methods | 7 |
| 2.12 | Appendix 7 Retention of human biosamples <i>and genetics</i> | 7 |
| 2.12.1 | Biosamples for future analysis | 7 |
| 2.13 | Section 12 References | 8 |

1 Introduction including rationale for the protocol amendment

This protocol is amended for the following reasons:

- To implement genetic biosamples for future analysis for those countries where it will be applicable.
The rationale for implementation is to ensure a better characterisation of the population in scope as new technologies and hypothesis develops. The collection of whole blood at baseline (visit 2) and end of treatment (visit 24) will allow for future genetic analysis thereby increasing the understanding of obesity, obesity treatment and related diseases in the future.
- As the scope of the biosampling for future analysis has broadened to cover obesity and - related diseases with this amendment, some paragraphs in appendix 7 have been revised. Furthermore, the informed consent has been updated to reflect the broader scope. A separate protocol will be developed before the biosamples from NN9536-4373 can be analysed in the future. This will include rationale for the analysis as well as reporting requirements.
- Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment.
- Minor clarification of content and correction of minor errors and typos.

In this protocol amendment:

- Any new text is written *in italics*.
- Any text deleted from the protocol is written using ~~strike through~~.

2 Changes

2.1 Section 2 Flowchart

Tobacco Use^c (~~6.4.1~~)

Biosamples for future analysis^f (9.7, 9.8, Appendix 7)

2.2 Section 3.2 Background

2.2.1 Section 3.2.1 Semaglutide

GLP-1 RA is a physiological regulator of appetite and GLP-1 receptors are present in several areas of the brain involved in appetite regulation⁵⁶.

Clinical⁵⁸⁻⁶²³ and non-clinical data⁶²⁶⁴ indicate that the body weight-reducing effect of semaglutide is mainly mediated by a reduced energy intake. No unexpected safety findings were identified and

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
4 of 8

Novo Nordisk

the tolerability and safety profile was overall consistent with previous findings in the T2D development programme and the GLP-1 RA class in general.

2.2.2 3.2.2 Trial population

The trial population will consist of subjects with obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$) or with overweight ($\text{BMI} \geq 27 \text{ kg/m}^2$) and weight-related comorbidities.

2.3 3.3 Benefit-risk assessment

2.3.1 3.3.2 Risks and precautions

The sections below describe identified and potential risks associated with semaglutide treatment. *For classification and further details of the risks, please refer to the current version of the IB or any updates hereof.* The identified/potential risks are based on findings in non-clinical studies and clinical trials with semaglutide as well as other GLP-1 RAs. For each of these risks, mitigating actions have been implemented to minimise the risks for subjects enrolled in this trial.

~~Identified risks~~

Gastrointestinal adverse events

- Consistent with findings with other GLP-1 RAs, the most frequently reported adverse events (AE) in clinical trials with semaglutide were gastrointestinal AEs. A low starting dose and dose escalation steps will be implemented in the trial to mitigate the risk of gastrointestinal AEs.

~~Potential risks~~

Medullary thyroid cancer (MTC) (based on non-clinical data)

- Expected proliferative thyroid C-cell changes were seen in the mouse and rat carcinogenicity studies after daily exposure to semaglutide for 2 years. No hyperplasia was observed in monkeys after 52 weeks exposure up to 13-fold above the clinical plasma exposure at 2.4 mg/week. In clinical trials with semaglutide, there have been no reports of MTC or clinically relevant changes in calcitonin levels. The C-cell changes in rodents are mediated by the GLP-1 receptor, which is not expressed in the normal human thyroid. Accordingly, the risk of GLP-1 receptor-mediated C-cell changes in humans is considered to be low. However, as a precaution, exclusion and discontinuation criteria related to medical history of multiple endocrine neoplasia type 2 (MEN2) or MTC and elevated plasma levels of calcitonin (biomarker for MTC) have been implemented in the trial.

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
5 of 8

Novo Nordisk

Other risks

Pregnancy and fertility (based on non-clinical data)

- Studies in animals have shown reproductive toxicity. There are limited data from the use of semaglutide in pregnant women. Therefore, semaglutide should not be used during pregnancy. Exclusion and discontinuation criteria related to pregnancy have been implemented in the trial.

2.4 Section 7.1 Treatments administered

- Injections may be administered in the thigh, abdomen or upper arm, at any time of day irrespective of meals. ~~Subjects should be encouraged to inject in the same area throughout the trial, but changing between left and right side is allowed.~~

2.5 Section 9.7 Genetics

~~Not applicable for this trial.~~

A blood sample for genetic analysis will be collected from subjects who have consented to participate in the optional biobank component of the trial. Refer to section 9.8 and Appendix 7 for further details.

2.6 Section 9.8 Biomarkers and biosamples for future analysis

Collection of samples for biomarker research is part of this trial to support the efficacy objectives. The following samples must be collected in accordance with the laboratory manual and the flowchart:

Biomarkers linked to cardiovascular risk:

- *Plasminogen Activator Inhibitor-1 (PAI-1)* Activity will be analysed by activity assay.
- *High sensitive C-reactive protein (hsCRP)*

Collection of biosamples for future analysis (stored in a biobank) is a component of this trial. Participation in the biobank component is optional. Subjects who do not wish to participate in the biobank component may still participate in the trial. For the biobank, samples will be collected according to the flowchart and stored for future use.

The samples are collected for the purpose of allowing future analyses of biomarkers, both genetic and circulating, at a later point in time when new knowledge or improved measurement techniques may have become available. The analyses may include biomarkers currently known or discovered in the future.

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
6 of 8

Novo Nordisk

Genetic analyses may include analysis of candidate genes or genetic markers throughout the genome with the purpose of understanding and predicting response to semaglutide as well as to understand obesity or other related diseases.

Analyses of circulating biomarkers will measure hormones, metabolites or other non-genetic serum entity with the purpose of understanding and predicting response to semaglutide as well as understanding obesity or other related diseases.

These samples need to be frozen and should be sent in batches to the central laboratory. The analyses are likely to be performed after the trial has come to an end, and results will therefore not be part of the clinical trial report. The biobank samples may be stored up to 15 years after end of trial at a central laboratory (see Appendix 7).

2.7 Section 10.1 Sample size determination

2.7.1 Table 10-2 Assumptions, marginal power and effective power for each endpoint in the hierarchical testing procedure given an anticipated number of 1950 randomised subjects

Foot note: SD = standard deviation; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain; # shown as a positive number

All tests in the hierarchy are based on the primary estimand

2.8 Section 10.3 Statistical analyses

2.8.1 Section 10.3.1 Primary endpoints

2.8.1.1 Table 10-3 Analysis and imputation methods to address the effectiveness and efficacy estimands for the primary and confirmatory secondary endpoints in the statistical testing hierarchy

Foot note: FAS = full analysis set; ANCOVA = analysis of covariance; RD-MI = multiple imputation using retrieved subjects; J2R-MI = jump to reference multiple imputation; S1-SI and S2-SI = single imputation as done by Sacks; TP-MI = tipping point multiple imputation; MMRM = mixed model for repeated measurements; LR = logistic regression; WC = waist circumference; sBP = systolic blood pressure; SF-36 = Short Form 36 v2.0 acute; PF = physical functioning; IWQoL-Lite = Impact of Weight on Quality of Life-Lite for Clinical Trials; PFD = physical function domain

Test order refers to the order of the endpoint in the statistical test hierarchy outlined in Table 10-2

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
7 of 8

Novo Nordisk

2.9 Appendix 2 Clinical laboratory tests

2.9.1 Table 11-3 Criteria for laboratory outliers

Lymphocytes < ~~25~~ $0.2 \times 10^9/L$

2.10 Appendix 3 Trial governance considerations

2.10.1 9) Data quality assurance

2.10.1.1 Protocol compliance

Deviations from the protocol should be avoided. If deviations do occur, the investigator must inform the monitor *without delay* and the implications of the deviation must be reviewed and discussed.

2.11 Appendix 5 Contraceptive guidance and collection of pregnancy information

2.11.1 Table 11-5 Highly effective contraceptive methods

Sexual abstinence^b

2.12 Appendix 7 Retention of human biosamples *and genetics*

2.12.1 Biosamples for future analysis

In countries where applicable, the trial will involve collection of human biosamples to be stored in a central archive *for future use*.

The material to be collected is ~~serum collected~~ at baseline (*visit 2*) and end of treatment (*visit 24*) is:
~~A total of 5 aliquots will be stored for each subject at both time points.~~

- *Whole blood (for genetic analysis)*
- *Serum (for analyses of circulating biomarkers)*

Protocol Amendment no 5
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

25 July 2018
1.0
Final
8 of 8

Novo Nordisk

~~The results obtained from any already performed analyses of the samples will still be used. In the event that the collected biosamples (serum) will be used in the future, the investigator will become directly informed by Novo Nordisk about the results if the findings are deemed clinically relevant and analytically valid and quantifiable. In such case, a written summary of the findings, including listings of subject specific values, will be provided once a firm conclusion from the results has been drawn by Novo Nordisk.~~

~~Potentially, observations of neoplastic diseases, serious hereditary diseases, other un-treatable diseases, or any other abnormal findings could be part of the observations. Subjects can contact the investigator if they wish to be informed about results derived from stored biosamples obtained from their own body.~~

In case the subject withdraws his/her informed consent for biosamples for future analysis and genetics, the monitor must contact the trial manager at Novo Nordisk as soon as possible in order to have the samples withdrawn from storage.

2.13 Section 12 References

64. Lu T, et al. Presented at the 77th scientific sessions of the American diabetes association, 9-13 June 2017, San Diego, CA, USA.

57. Novo Nordisk A/S. Investigator's Brochure, Semaglutide s.c. 2.4 mg, project NN9536, (edition 1, ~~draft version 0.1~~). 28 Nov 2017.

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

29 March 2019
1.0
Final
1 of 21

Novo Nordisk

Protocol Amendment
no 6
to Protocol, version 3.0
dated 04 Jun 2018

Trial ID: NN9536-4373

Effect and safety of semaglutide 2.4 mg once-weekly in subjects with overweight or obesity

Trial phase: 3a

Applicable to all countries

This ~~confidential~~ document is the property of Novo Nordisk. ~~No unpublished information contained herein may be disclosed without prior written approval from Novo Nordisk. Access to this document must be restricted to relevant parties.~~

Table of Contents

| | | |
|----------|---|-----------|
| 1 | Introduction including rationale for the protocol amendment..... | 4 |
| 2 | Changes for all countries | 5 |
| 2.1 | Section 2 Flowchart..... | 5 |
| 2.2 | Section 3 Introduction | 5 |
| 2.2.1 | Section 3.2 Background | 5 |
| 2.2.1.1 | Section 3.2.1 Semaglutide | 5 |
| 2.2.1.2 | Section 3.2.2 Trial population..... | 5 |
| 2.3 | Section 6 Trial population | 5 |
| 2.3.1 | Section 6.1 Inclusion criteria | 5 |
| 2.3.2 | Section 6.3 Randomisation criteria | 5 |
| 2.4 | Section 8 Discontinuation/Withdrawal criteria..... | 5 |
| 2.4.1 | Section 8.1 Discontinuation of trial treatment..... | 5 |
| 2.5 | Section 9 Trial assessments and procedures | 6 |
| 2.5.1 | Section 9.2 Adverse events | 6 |
| 2.5.1.1 | Section 9.2.1 Time period and frequency for collecting AE and SAE information | 6 |
| | Table 9-1 AEs requiring additional data collection (via specific event form) and events for adjudication | 6 |
| | (a) Section 9.2.1.1 Events for adjudication | 6 |
| 2.5.2 | Section 9.4 Safety assessments..... | 6 |
| 2.5.2.1 | Section 9.4.6 Immunogenicity assessments..... | 6 |
| 2.5.3 | Section 9.8 Biomarkers and biosamples for future analysis..... | 7 |
| 2.6 | Appendix 4 Adverse events: definitions and procedures for recording, evaluation, follow-up, and reporting | 8 |
| 2.6.1 | Description of AEs requiring additional data collection (via specific event form) and events for adjudication. | 8 |
| 2.7 | Appendix 5 Contraceptive guidance and collection of pregnancy information..... | 9 |
| 2.7.1 | Women in the following categories are not considered WOCBP..... | 9 |
| 2.7.1.1 | 3. Postmenopausal female | 9 |
| 2.7.2 | Table 11-5 Highly effective contraceptive methods | 9 |
| 2.7.2.1 | Highly effective methods that are user independent ^b | 9 |
| 2.7.3 | Collection of pregnancy information | 10 |
| 2.7.3.1 | Female subjects who become pregnant | 10 |
| 2.8 | Appendix 6 Technical complaints: Definition and procedures for recording, evaluation, follow-up and reporting..... | 10 |
| 2.9 | Appendix 9 Country-specific requirements | 10 |
| 2.10 | Appendix 10 Protocol amendment history..... | 11 |
| 3 | Changes for all countries participating in the extension phase | 14 |
| 3.1 | Section 1 Synopsis | 14 |
| 3.1.1 | Rationale | 14 |
| 3.1.2 | Overall design..... | 14 |
| 3.1.3 | Key inclusion criteria | 14 |
| 3.1.4 | Key exclusion criteria | 14 |
| 3.1.5 | Number of subjects: | 15 |

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

29 March 2019
1.0
Final
3 of 21

Novo Nordisk

| | | |
|---------|---|----|
| 3.1.6 | Treatment groups and duration..... | 15 |
| 3.2 | Section 2 Flowchart..... | 15 |
| 3.3 | Section 3 Introduction | 16 |
| 3.3.1 | Section 3.1 Trial rationale | 16 |
| 3.3.2 | Section 3.3 Benefit-risk assessment..... | 16 |
| 3.3.2.1 | Section 3.3.1 Benefits..... | 16 |
| 3.4 | Section 4 Objectives and endpoints | 16 |
| 3.4.1 | Section 4.1 Primary, secondary and exploratory objective(s) | 16 |
| 3.4.2 | Section 4.2 Primary, secondary and exploratory endpoint(s)..... | 16 |
| 3.4.2.1 | Section 4.2.3 Exploratory endpoints | 16 |
| 3.5 | Section 5 Trial design..... | 17 |
| 3.5.1 | Section 5.1 Overall design..... | 17 |
| 3.5.2 | Section 5.2 Subject and trial completion..... | 18 |
| 3.5.3 | Section 5.5 Justification for dose | 19 |
| 3.6 | Section 6 Trial population | 19 |
| 3.6.1 | Section 6.1 Inclusion criteria | 19 |
| 3.6.2 | Section 6.2 Exclusion criteria | 19 |
| 3.7 | Section 7 Treatments..... | 19 |
| 3.7.1 | Section 7.4 Blinding..... | 19 |
| 3.7.2 | Section 7.7 Concomitant medication | 19 |
| 3.7.3 | Section 8.3 Lost to follow-up | 20 |
| 3.8 | Section 9 Trial assessments and procedures | 20 |
| 3.8.1 | Section 9.2 Adverse events | 20 |
| 3.9 | Section 10 Statistical considerations | 20 |
| 3.9.1 | Section 10.2 Definition of analysis sets | 20 |
| 3.9.1.1 | Section 10.3.3 Exploratory endpoints | 21 |
| 3.10 | Appendix 1 Abbreviations..... | 21 |
| 3.11 | Appendix 2 Clinical laboratory tests..... | 21 |
| 3.11.1 | Table 11-1 Protocol-required efficacy laboratory assessments (<i>up to V25</i>)..... | 21 |
| 3.11.2 | Table 11-2 Protocol-required efficacy laboratory assessments (<i>V25_{ext}-V28_{ext}</i>) | 21 |

1 Introduction including rationale for the protocol amendment

This protocol is amended for the following reasons:

- To extend the trial for an additional 45-weeks off-treatment period after the last visit in the main phase (i.e. V25) without structured lifestyle intervention in the following countries only: Canada, Germany, the UK and selected sites in the US and Japan.
The rationale for extending the trial is to obtain a better understanding of how the weight, cardiovascular risk factors and glucose metabolism develops after completed pharmacotherapy intervention in people living with overweight or obesity. The extension of this trial mimics a real world setting and can thereby provide better guidance to health care professionals in the future management of people living with obesity as well as to these people.
Separate Clinical Trial Reports (CTR) will be prepared for the main and the extension phase of the trial.
- Addition of “Main phase” and “Extension phase” throughout the protocol to highlight which sections are related to the “main phase” and which are related to the “extension phase”. The protocol described the main phase in all sections unless it is explicitly described as referring to the extension phase.
- Minor clarification of content and correction of minor errors and typos.

In the following, the changes related to all countries are listed first in Section 2 followed by the changes related to the countries participating in the extension phase in Section 3.

| AMENDMENT HISTORY | | |
|--|-------------------------------|--|
| Document | Version | Applicable for |
| Updated protocol including amendment no. 4 (DE), 5(UK) and 6 (all countries) | Version 4.0, 29 March 2019 | All countries |
| Updated protocol including amendment no. 2 (UK) and 3 (all countries) | Version 3.0, 06 June 2018 | All countries, except Germany and the UK (local amendments 04 and 05, respectively, approved for protocol version 2.0) |
| Original protocol | Version 2.0, 21 December 2017 | All countries |

In this protocol amendment:

- Any new text is written *in italics*.
- Any text deleted from the protocol is written using ~~strike through~~.

2 Changes for all countries

2.1 Section 2 Flowchart

Main phase

Tobacco Use^c (6.4.24)

Concomitant medication X added at V25

b For all female subjects *of child bearing potential*.

2.2 Section 3 Introduction

2.2.1 Section 3.2 Background

2.2.1.1 Section 3.2.1 Semaglutide

Clinical⁵⁸⁹⁻⁶³⁴ and non-clinical data⁶⁴⁵ indicate that the body weight-reducing effect of semaglutide is mainly mediated by a reduced energy intake. No unexpected safety findings were identified and the tolerability and safety profile was overall consistent with previous findings in the T2D development programme and the GLP-1 RA class in general.

2.2.1.2 Section 3.2.2 Trial population

The trial population will consist of subjects with obesity (BMI ≥ 30.0 kg/m²) or with overweight (BMI ≥ 27.0 kg/m²) and weight-related comorbidities.

2.3 Section 6 Trial population

2.3.1 Section 6.1 Inclusion criteria

1. Body mass index (BMI) ≥ 30.0 kg/m² or ≥ 27.0 kg/m² with the presence of at least one of the following weight-related comorbidities (treated or untreated): hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease

2.3.2 Section 6.3 Randomisation criteria

Randomisation criteria applicable only for DEXA sites

2. BMI ≤ 40.0 kg/m² at screening

2.4 Section 8 Discontinuation/Withdrawal criteria

2.4.1 Section 8.1 Discontinuation of trial treatment

The subject must be discontinued from trial product, if any of the following applies:

1. Safety concern as judged by the investigator
2. Calcitonin ≥ 100 ng/L (see Appendix 8)
3. Suspicion of pancreatitis
4. Pregnancy
5. Intention of becoming pregnant
6. Simultaneous participation in another clinical trial of an approved or non-approved investigational medicinal product.

For country specific requirements for Germany and the UK refer to Appendix 9.

2.5 Section 9 Trial assessments and procedures

2.5.1 Section 9.2 Adverse events

2.5.1.1 Section 9.2.1 Time period and frequency for collecting AE and SAE information

Table 9-1 AEs requiring additional data collection (via specific event form) and events for adjudication

| Event type | AE via specific event form | Event for adjudication (Section 9.2.1.1) |
|-----------------------------------|----------------------------|--|
| Medication error | X | |
| Misuse/abuse | X | |
| Death | | X |
| Cardiovascular events | | |
| Acute Coronary Syndrome | | X |
| Cerebrovascular event | | X |
| Heart failure | | X |
| Coronary artery revascularisation | | X |
| Acute pancreatitis | X | X |
| Acute gallbladder disease | X | |
| Malignant neoplasms | X | |
| Hepatic event | X | |

(a) Section 9.2.1.1 Events for adjudication

Event adjudication will be performed for adverse events in randomised subjects *until Data Base Lock*.

2.5.2 Section 9.4 Safety assessments

2.5.2.1 Section 9.4.6 Immunogenicity assessments

Blood samples for determination of serum antibodies against semaglutide, including cross reactivity to endogenous GLP-1 will be taken during the trial at visits specified in the flowchart. Samples ~~taken at the 'end of trial' visit~~ which are positive for anti-semaglutide antibodies will be further

characterised for in vitro neutralising effect towards semaglutide. In addition, if samples ~~taken at the 'end of trial' visit~~ are also positive for cross-reactivity against endogenous GLP-1, the samples will be analysed for in vitro neutralising effect towards endogenous GLP-1. *Samples which are positive for anti-semaglutide antibodies will also be titrated.* The results of the analysis will only be disclosed after completion of the clinical trial report (CTR) if required by local regulation.

2.5.3 Section 9.8 Biomarkers and biosamples for future analysis

Collection of samples for biomarker research is part of this trial to support the effect objectives. The following samples must be conducted in accordance with the laboratory manual and the flowchart:

Biomarkers linked to cardiovascular risk:

- Plasminogen Activator Inhibitor (PAI-1) Activity will be analysed by activity assay.
- High sensitive C-reactive protein (hsCRP)

Biomarkers linked to hunger and satiety:

- *Leptin*
- *Soluble leptin receptor*

Collection of biosamples for future analysis (stored in a biobank) is a component of this trial. Participation in the biobank component is optional. Subjects who do not wish to participate in the biobank component may still participate in the trial. For the biobank, samples will be collected according to the flowchart and stored for future use.

The samples are collected for the purpose of allowing future analyses of biomarkers, both genetic and circulating, at a later point in time when new knowledge or improved measurement techniques may have become available. The analyses may include biomarkers currently known or discovered in the future.

Genetic analyses may include analysis of candidate genes or genetic markers throughout the genome with the purpose of understanding and predicting response to semaglutide as well as to understand obesity or other related diseases.

Analyses of circulating biomarkers will measure hormones, metabolites or other non-genetic serum entity with the purpose of understanding and predicting response to semaglutide as well as understanding obesity or other related diseases.

These samples need to be frozen and should be sent in batches to the central laboratory. The analyses are likely to be performed after the trial has come to an end, and results will therefore not be part of the clinical trial report. The biobank samples may be stored up to 15 years after end of trial at a central laboratory (see Appendix 7).

~~Biomarkers linked to hunger and satiety:~~

- ~~• Leptin~~
- ~~• Soluble leptin receptor~~ Collection of biosamples for future analysis (stored in a biobank) is a component of this trial. Participation in the biobank component is optional. Subjects who do not wish to participate in the biobank component may still participate in the trial. For the biobank, samples will be collected according to the flowchart and stored for future use.
- ~~• The samples are collected for the purpose of allowing future analyses of biomarkers, both genetic and circulating, at a later point in time when new knowledge or improved measurement techniques may have become available. The analyses may include biomarkers currently known or discovered in the future.~~

~~Genetic analyses may include analysis of candidate genes or genetic markers throughout the genome with the purpose of understanding and predicting response to semaglutide as well as to understand obesity or other related diseases.~~

~~Analyses of circulating biomarkers will measure hormones, metabolites or other non-genetic serum entity with the purpose of understanding and predicting response to semaglutide as well as understanding obesity or other related diseases.~~

~~These samples need to be frozen and should be sent in batches to the central laboratory. The analyses are likely to be performed after the trial has come to an end, and results will therefore not be part of the clinical trial report. The biobank samples may be stored up to 15 years after end of trial at a central laboratory (see Appendix 7).~~

2.6 Appendix 4 Adverse events: definitions and procedures for recording, evaluation, follow-up, and reporting

2.6.1 Description of AEs requiring additional data collection (via specific event form) and events for adjudication.

| | |
|-----------------------------------|---|
| Medication error: | <p>A medication error concerning trial products is defined as:</p> <ul style="list-style-type: none"> Administration of wrong drug. <ul style="list-style-type: none"> Note: Use of wrong dispensing unit number (DUN) is not considered a medication error unless it results in a confirmed administration of wrong drug. Wrong route of administration, such as intramuscular instead of subcutaneous. Accidental administration of more than 2.4 mg/week or a higher dose than intended during dose escalation, however, the administered dose must deviate from the intended dose to an extent where clinical consequences for the trial subject were likely to happen as judged by the investigator, although they did not necessarily occur. Administration of an overdose with the intention to cause harm, misuse or abuse of trial product. |
| Misuse or abuse of trial product* | <ul style="list-style-type: none"> Misuse is when the trial product is intentionally and inappropriately used. Abuse of trial product is persistent or sporadic, intentional excessive use, which is accompanied by harmful physical or psychological effects (e.g. overdose with the intention to cause harm). |

*Additional data for misuse or abuse of trial product is reported on the medication error event form.

2.7 Appendix 5 Contraceptive guidance and collection of pregnancy information

2.7.1 Women in the following categories are not considered WOCBP

2.7.1.1 3. Postmenopausal female

- ~~Otherwise, they must discontinue HRT to allow confirmation of postmenopausal~~ A postmenopausal state is defined as no menses for 12 months without an alternative medical cause. A high Follicle Stimulating Hormone (FSH) level in the postmenopausal range may be used to confirm a postmenopausal state in women not using hormonal contraception or Hormonal Replacement Therapy (HRT). However, in the absence of 12 months of amenorrhea, a single FSH measurement is insufficient.
- Females on HRT and whose menopausal status is in doubt will be required to use one of the non-hormonal highly effective contraception methods if they wish to continue their HRT during the trial ~~status before trial enrolment~~. *Otherwise, they must discontinue HRT to allow confirmation of postmenopausal status before trial enrolment.*

2.7.2 Table 11-5 Highly effective contraceptive methods

2.7.2.1 Highly effective methods that are user independent^b

Bullet added for “Implantable progestogen only hormonal contraception associated with inhibition of ovulation”.

2.7.3 Collection of pregnancy information

2.7.3.1 Female subjects who become pregnant

- Investigator will collect pregnancy information on any female subject, who becomes pregnant while participating in this trial.
- Information will be recorded on the appropriate form and submitted to Novo Nordisk within 14 calendar days of learning of a subject's pregnancy.
- Subject will be followed to determine the outcome of the pregnancy. The investigator will collect follow-up information on subject and neonate, which will be forwarded to Novo Nordisk. Generally, follow-up will not be required for longer than 1 month beyond the delivery date.
- Any termination of pregnancy will be reported, regardless of foetal status (presence or absence of anomalies) or indication for procedure.
- While pregnancy itself is not considered to be an AE or SAE, any pregnancy complication or elective termination of a pregnancy *for medical reasons* will be reported as an AE or SAE.
- A spontaneous abortion is always considered to be an SAE and will be reported as such.
- Any SAE occurring as a result of a post-trial pregnancy which is considered possibly/probably related to the trial product by the investigator will be reported to Novo Nordisk as described in Appendix 4. While the investigator is not obligated to actively seek this information in former subjects, he or she may learn of an SAE through spontaneous reporting

2.8 Appendix 6 Technical complaints: Definition and procedures for recording, evaluation, follow-up and reporting

~~All technical complaints are handled by the Customer Complaint Center at Novo Nordisk. Only technical complaints related to adverse events will be included in the CTR.~~

2.9 Appendix 9 Country-specific requirements

Section 5.2 Subject and trial completion

For Mexico: approximately ~~60~~70 subjects are planned to be randomised on trial product.

Section 6.2 Exclusion criterion no. ~~29~~30

Female who is pregnant, breast-feeding or intends to become pregnant or is of child-bearing potential and not using adequate contraceptive methods (highly effective contraceptive measures as required by local regulation or practice)

Section 8.1 Discontinuation of trial treatment

For Germany and the UK: According to local amendments for Germany and the UK (amendments 04 and 05 respectively), subjects who are included in the trial in violation of the inclusion and/or exclusion criteria and/or randomisation criteria will be discontinued from trial treatment.

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

29 March 2019
1.0
Final
11 of 21

Novo Nordisk

Appendix 7 Retention of human biosamples

For Taiwan: No subjects from Taiwan will participate in the optional biobank *incl. genetics* part of the trial.

2.10 Appendix 10 Protocol amendment history

The protocol amendment summary of changes table for the current updated protocol is located directly before the Table of Contents.

Protocol amendment 5: (UK, 25 July 2018 version 1.0)

This amendment is considered to be substantial based on the criteria set forth in Article 10(a) of Directive 2001/20/EC of the European Parliament and the Council of the European Union.

Overall rationale for the amendment

| Section # and name | Description of change | Rationale |
|--------------------------------------|--|--|
| Section 3.3: Benefit-risk assessment | Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment | To make the protocol more flexible and avoid repetitive information in the protocol and IB |
| N/A | Minor clarification of content and correction of minor errors and typos | To avoid misinterpretations |

Protocol amendment 4: (Germany, 19 July 2018 version 1.0)

This amendment is considered to be substantial based on the criteria set forth in Article 10(a) of Directive 2001/20/EC of the European Parliament and the Council of the European Union.

Overall rationale for the amendment

| Section # and name | Description of change | Rationale |
|--------------------------------------|--|--|
| Section 3.3: Benefit-risk assessment | Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment | To make the protocol more flexible and avoid repetitive information in the protocol and IB |
| N/A | Minor clarification of content and correction of minor errors and typos | To avoid misinterpretations |

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date: 29 March 2019
Version: 1.0
Status: Final
Page: 12 of 21

Novo Nordisk

Updated protocol including amendment 3: (Global - except Germany and the UK - 04 June 2018 version 1.0)

This amendment is considered to be substantial based on the criteria set forth in Article 10(a) of Directive 2001/20/EC of the European Parliament and the Council of the European Union.

Overall rationale for the amendment

| <i>Section # and name</i> | <i>Description of change</i> | <i>Rationale</i> |
|---|---|--|
| <i>Section 8.1: Discontinuation/Withdrawal criteria</i> | <i>Removal of criteria for discontinuation of trial treatment for subjects included in the trial in violation of the inclusion and/or exclusion criteria and/or randomisation criteria.</i> | <i>Subjects will not be discontinued from trial product if considered safe to continue. However, prospective approval of protocol deviations to recruitment and enrolment criteria, also known as protocol waivers or exemptions, is not permitted and deviations from the protocol should be avoided.</i> |
| <i>Section 3.3: Benefit-risk assessment</i> | <i>Classifications of risks have been removed from the protocol and instead a reference to the IB or any updates hereof has been added for further details of the risks associated with semaglutide treatment</i> | <i>To make the protocol more flexible and avoid repetitive information in the protocol and IB</i> |
| <i>N/A</i> | <i>Minor clarification of content and correction of minor errors and typos</i> | <i>To avoid misinterpretations</i> |

Protocol amendment 2: (UK, 15 March 2018 version 1.0)

This amendment is considered to be substantial based on the criteria set forth in Article 10(a) of Directive 2001/20/EC of the European Parliament and the Council of the European Union.

Overall rationale for the amendment

| <i>Section # and name</i> | <i>Description of change</i> | <i>Rationale</i> |
|--|---|-------------------------------------|
| <i>Appendix 9: Country specific requirements</i> | <i>The use of double barrier methods in Appendix 5 is not applicable for UK. Therefore appendix 9 'country specific requirements' is updated and must be followed in UK</i> | <i>Required by local regulation</i> |

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

29 March 2019
1.0
Final
13 of 21

Novo Nordisk

Protocol amendment 1: (Argentina, 22 January 2018 version 1.0)

This amendment is considered to be substantial based on the criteria set forth in Article 10(a) of Directive 2001/20/EC of the European Parliament and the Council of the European Union.

Overall rationale for the amendment

| <i>Section # and name</i> | <i>Description of change</i> | <i>Rationale</i> |
|--|---|-------------------------------------|
| <i>Appendix 5 & 9: Contraceptive methods and Country specific requirements</i> | <i>The local requirement on contraceptive methods in Appendix 9 in the protocol is less strict than the global requirements in Appendix 5, i.e. barrier methods are not allowed in this trial, therefore the local requirement is deleted and Appendix 5 must be followed in Argentina.</i> | <i>Required by local regulation</i> |

The amendment was superseded by commitment letter from the Sponsor.

3 Changes for all countries participating in the extension phase

3.1 Section 1 Synopsis

3.1.1 Rationale

Extension phase

The rationale for extending the trial is to obtain a better understanding of how the weight, cardiovascular risk factors and glucose metabolism develops after completed pharmacotherapy intervention in people living with overweight or obesity. The extension of this trial mimics a real world setting and can thereby provide better guidance to health care professionals in the future management of people living with obesity as well as to these people. Focus in the off-treatment extension phase is change in body weight, cardiovascular risk factors and glucose metabolism in a subset of subjects who have completed the main phase on the target dose of 2.4 mg semaglutide once-weekly or semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity.

The 2.4 mg dose has been selected as an inclusion criterion in the extension phase as this is the target dose in the main phase of NN9536-4373 as outlined in Section 5.5.

3.1.2 Overall design

Extension phase

The extension phase is a 45-week off-treatment period after the last visit in the main phase (i.e. V25) without structured lifestyle intervention. Subjects on the target dose of 2.4 mg semaglutide once-weekly or semaglutide placebo at the End of Treatment visit (V24) are considered eligible and will be offered continued follow-up at the investigational site for an additional 45 weeks. For subjects continuing in the extension phase, the total trial duration will be approximately 122 weeks (2 weeks screening, 68 weeks treatment and 52 weeks observation). The extension phase includes 5 visits to the clinic (week 68, 75, 80, 104 and week 120); the first two visits of the extension phase (V24_{ext} and V25_{ext}) coincide with the visits (V24 and V25) in the main phase of the trial.

3.1.3 Key inclusion criteria

Extension phase

- Informed consent for the extension phase obtained before any trial related activities for the extension phase*
- On randomised treatment on the target dose at week 68, i.e. treated with 2.4 mg semaglutide once-weekly or semaglutide placebo*

3.1.4 Key exclusion criteria

Extension phase

- Female who is pregnant or intends to become pregnant during the extension phase*

Protocol Amendment no. 6
Trial ID: NN9536-4373

~~CONFIDENTIAL~~

Date:
Version:
Status:
Page:

29 March 2019
1.0
Final
15 of 21

Novo Nordisk

- Any disorder, unwillingness or inability, not covered by any of the other exclusion criteria, which in the investigator's opinion, might jeopardise the subject's compliance with the extension of the trial

3.1.5 Number of subjects:

Extension phase: Approximately 300 subjects are planned to continue in the extension phase in the following countries only: Canada, Germany, the UK and selected sites in the US and Japan.

3.1.6 Treatment groups and duration

Extension phase:

The extension phase is a 45-week off-treatment period after the last visit in the main phase (i.e. V25) without structured lifestyle intervention

3.2 Section 2 Flowchart

| <i>Extension phase^a</i> | <i>Protocol Section</i> | <i>Extension</i> | | | | <i>End of trial_{ext}</i> |
|---|-----------------------------|--------------------------|--------------------------|--------------------------|--------------------------|-----------------------------------|
| <i>Visit (V)</i> | | <i>V24_{ext}</i> | <i>V25_{ext}</i> | <i>V26_{ext}</i> | <i>V27_{ext}</i> | <i>V28_{ext}</i> |
| <i>Timing of Visit from date of Randomisation (Weeks)</i> | | 68 | 75 | 80 | 104 | 120 |
| <i>Visit Window (Days)</i> | | ±3 | 0 to +5 | ±5 | ±5 | ±5 |
| <i>Informed Consent</i> | | X | | | | |
| <i>Inclusion and Exclusion Criteria</i> | 6.1 6.2 | X | X | | | |
| <i>Concomitant Medication</i> | 7.7 | | | X | X | X |
| <i>Body Measurements</i> | 9.1.1 | | | X | X | X |
| <i>Vital Signs</i> | 9.4.3 | | | X | X | X |
| <i>Laboratory Assessment</i> | Appendix 2, Table 11-2 | | X | X | X | X |

a The assessments in the extension phase flowchart are only to be completed by the subjects participating in the extension phase. These subjects must complete all visits and assessments in the main phase as well as all visits and assessments in the extension phase. A subject in the extension phase will complete the following visits in this order: V24, V24_{ext}, V25, V25_{ext}, V26_{ext}, V27_{ext} and finally V28_{ext}. The following visits coincide; V24 and V24_{ext} as well as V25 and V25_{ext}.

3.3 Section 3 Introduction**3.3.1 Section 3.1 Trial rationale*****Extension phase***

The rationale for extending the trial is to obtain a better understanding of how the weight, cardiovascular risk factors and glucose metabolism develops after completed pharmacotherapy intervention in people living with overweight or obesity. The extension of this trial mimics a real world setting and can thereby provide better guidance to health care professionals in the future management of people living with obesity as well as to these people. Focus in the off-treatment extension phase is change in body weight, cardiovascular risk factors and glucose metabolism in a subset of subjects who have completed the main phase on the target dose of 2.4 mg semaglutide once-weekly or semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity.

The 2.4 mg dose has been selected as an inclusion criterion in the extension phase as this is the target dose in the main phase of NN9536-4373 as outlined in Section 5.5.

3.3.2 Section 3.3 Benefit-risk assessment**3.3.2.1 Section 3.3.1 Benefits*****Extension phase***

Subjects who continue in the extension phase are expected to benefit from the continued contact with the trial site focusing on general health, including body weight, cardiovascular risk factors and glucose metabolism parameters. As the extension is an off-treatment period, there are no risks considered to the subjects' participation.

3.4 Section 4 Objectives and endpoints**3.4.1 Section 4.1 Primary, secondary and exploratory objective(s)****Exploratory objectives*****Extension phase***

To explore the change in body weight, cardiovascular risk factors and glucose metabolism in a subset of subjects after having completed treatment on the target dose of 2.4 mg semaglutide once-weekly or semaglutide placebo as an adjunct to a reduced-calorie diet and increased physical activity followed by an off-treatment period.

3.4.2 Section 4.2 Primary, secondary and exploratory endpoint(s)**3.4.2.1 Section 4.2.3 Exploratory endpoints**

The exploratory endpoints addressing the explorative objectives:

Main phase

~~The exploratory endpoints addressing the explorative objectives:~~

- Change from baseline at week 0 to week 68 in:
 - Glycaemic category (normo-glycaemia, pre-diabetes, T2D)
 - Antihypertensive medication (decrease, no change, increase)
 - Lipid-lowering medication (decrease, no change, increase)
 - The Stanford Presenteeism Scale (SPS-6), total score
 - Fatty liver Index (FLI) score category (<30, ≥30 and <60, ≥60)
 - International Consultation on Incontinence Questionnaire-Urinary Incontinence-Short Form (ICIQ-UI-SF), sum score (assessed in female subjects)
- Subjects who from randomisation at week 0 to week 68 have permanently discontinued randomised trial product (yes/no)
- Time to permanent discontinuation of randomised trial product (weeks)
- Number of days per week with at least one entry in the food diary from baseline at week 0 to week 68

Number of minutes per week of physical activity from baseline at week 0 to week 68

Extension phase

- *Change from week 68 to week 120 in:*
 - *Body weight (% , kg)*
 - *BMI (kg/m²)*
 - *HbA1c (% , mmol /mol)*
 - *Diastolic blood pressure (mmHg)*
 - *Systolic blood pressure (mmHg)*
 - *Lipids (mg/dL)*
 - *Total cholesterol*
 - *High density lipoprotein (HDL) cholesterol*
 - *Low density lipoprotein (LDL) cholesterol*
 - *Very low density lipoprotein (VLDL) cholesterol*
 - *Free fatty acids*
 - *Triglycerides*
- *High sensitivity C-Reactive Protein (hsCRP) (mg/L)*

3.5 Section 5 Trial design**3.5.1 Section 5.1 Overall design****Extension phase**

Eligible subjects participating in the extension phase will continue for an additional 45 weeks after the last visit in the main phase (i.e. V25). The week 68 visit will be the first visit in the extension

phase and the Subject Information and Informed Consent will be presented to the subject, followed by the week 75 visit where the first extension parameters will be collected (if subject has accepted participation). The first two visits in the extension phase ($V24_{ext}$ and $V25_{ext}$) coincide with the visits ($V24$ and $V25$) in the main phase of the trial. For subjects continuing in the extension phase, the total trial duration will be approximately 122 weeks (2 weeks screening, 68 weeks treatment, 52 weeks off-treatment period without structured lifestyle intervention).

Inserted new Figure 5-1:




Figure 5-1 A schematic diagram of the trial design with the duration of the trial periods including follow-up and extension period

3.5.2 Section 5.2 Subject and trial completion

Extension phase: Approximately 300 subjects are planned to continue in the extension phase in the following countries only: Canada, Germany, the UK and selected sites in the US and Japan.

The countries have been chosen as they together will represent the global trial population in the NN9636-4373 trial. The Japan and US sites will objectively be selected based on the highest recruitment numbers into the main phase. In case of equality in the recruitment numbers, the sites will be selected in a chronological order.

3.5.3 Section 5.5 Justification for dose***Extension phase***

The extension phase is a 45-week off-treatment period after the last visit in the main phase (i.e. V25) without structured lifestyle intervention.

3.6 Section 6 Trial population**3.6.1 Section 6.1 Inclusion criteria*****Extension phase***

- 1. Informed consent for the extension phase obtained before any trial related activities for the extension phase*
- 2. On randomised treatment on the target dose at week 68, i.e. 2.4 mg semaglutide once-weekly or semaglutide placebo*

3.6.2 Section 6.2 Exclusion criteria***Extension phase***

- 1. Female who is pregnant or intends to become pregnant during the extension phase*
- 2. Any disorder, unwillingness or inability, not covered by any of the other exclusion criteria, which in the investigator's opinion, might jeopardise the subject's compliance with the extension of the trial*

3.7 Section 7 Treatments**3.7.1 Section 7.4 Blinding*****Extension phase***

The blinding will be broken for the sponsor after completion of the main phase in preparation for the clinical trial report. Investigators and subjects participating in the extension phase should remain blinded in the extension phase until completion.

3.7.2 Section 7.7 Concomitant medication***Extension phase***

In the extension phase the following concomitant treatments and events that could have an impact on weight must be reported:

- Anti-obesity treatment, e.g. medications or surgery procedures*
- Anti-hypertensive medication*
- Lipid lowering medication*
- Anti-diabetes medication*
- Other, e.g. pregnancy*

3.7.3 Section 8.3 Lost to follow-up***Extension phase***

Efforts must be made to have the subjects, who agree to participate in the extension phase, also to continue in the extension phase. Subjects must be educated about the continued scientific importance of their data. Only subjects who withdraw consent will be considered as withdrawn from the extension phase of the trial.

A subject may withdraw consent at any time during the extension phase at his/her own request as described for the main phase in Section 8.2. If a subject is lost to follow-up all efforts should be made as described in Section 8.3 for the main phase.

3.8 Section 9 Trial assessments and procedures**3.8.1 Section 9.2 Adverse events*****Extension phase***

No systematic collection of AEs or SAEs will be performed, meaning investigators are not obliged to actively seek for AE or SAEs in trial subjects after the week 75 visit (V25); 7 weeks after last trial product administration.

If the investigator learns of any SAE, including death, at any time after week 75, and the investigator considers the event to be possibly/probably related to the investigational product or trial participation, the investigator must promptly notify Novo Nordisk by completing a paper AE- and Safety Information Form as described in Appendix 4; SAE reporting via paper CRF.

Adverse reactions associated with other products, including marketed Novo Nordisk products, should be reported via the national spontaneous reporting system to either the local competent authority or the marketing authorisation holder directly.

3.9 Section 10 Statistical considerations**3.9.1 Section 10.2 Definition of analysis sets**

ThreeTwo analysis sets are defined:

- The full analysis set (FAS) includes all randomised subjects according to the intention-to-treat principle.
- The safety analysis set (SAS) includes all randomised subjects exposed to at least one dose of randomised treatment.

- The extension analysis set (ExAS) includes all subjects eligible for the extension trial (as described in Section 6.1), who gave informed consent to participate and attended at least one of the following visits in the extension period: $V25_{ext}$, $V26_{ext}$, $V27_{ext}$ or $V28_{ext}$.

3.9.1.1 Section 10.3.3 Exploratory endpoints

For subjects in the ExAS, descriptive statistics will be presented for exploratory endpoints evaluating the change from week 68 to week 120. In addition, the change from baseline (week 0) to end of treatment (week 68) will be reported for the same set of parameters and subjects. Similarly, baseline data for the relevant parameters as well as demographics and baseline characteristics will be summarised for the ExAS.

3.10 Appendix 1 Abbreviations

The following items have been added to the list of abbreviations:

| | |
|-------------|------------------------|
| CTR_{ext} | CTR extension |
| ExAS | extension analysis set |
| EXT | extension |

3.11 Appendix 2 Clinical laboratory tests

- The tests detailed in Table 11-1, Table 11-2 and Table 11-23 will be performed by the central laboratory

3.11.1 Table 11-1 Protocol-required efficacy laboratory assessments (up to $V25$)

3.11.2 Table 11-2 Protocol-required efficacy laboratory assessments ($V25_{ext}$ - $V28_{ext}$)

| Laboratory assessments | Parameters |
|--|--|
| Glucose metabolism | <ul style="list-style-type: none"> • $HbA1c$ |
| Lipids | <ul style="list-style-type: none"> • Cholesterol • High density lipoprotein (HDL) cholesterol • Low density lipoprotein (LDL) cholesterol • Triglycerides • Very-low-density lipoprotein (VLDL) cholesterol • Free fatty acids |
| Biomarkers | <ul style="list-style-type: none"> • $hsCRP$ |
| Notes: All laboratory assessments in the extension phase are to be collected non-fasting. | |