

2023 February 1 Shift 2

1

AI24BTECH11004-Bheri Sai Likith Reddy

1 SECTION-A

1) The sum $\sum_{n=1}^{\infty} \frac{2n^2+3n+4}{(2n)!}$ is equal to :

- a) $\frac{11e}{2} + \frac{7}{2e}$
- b) $\frac{13e}{4} + \frac{5}{4e} - 4$
- c) $\frac{11e}{2} + \frac{7}{2e} - 4$
- d) $\frac{13e}{4} + \frac{5}{4e}$

2) Let $S = \left\{x \in R : 0 < x < 1 \text{ and } 2 \tan^{-1} \left(\frac{1-x}{1+x} \right) = \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)\right\}$. If $n(S)$ denotes the number of elements in S then :

- a) $n(S)=2$ and only one element in S is less than $\frac{1}{2}$
- b) $n(S)=2$ and only one element in S is less than $\frac{1}{2}$
- c) $n(S)=2$ and only one element in S is less than $\frac{1}{2}$
- d) $n(S)=0$

3) Let $\vec{a} = 2\hat{i} - 7\hat{j} + 5\hat{k}$, $\vec{b} = \hat{i} + \hat{k}$ and $\vec{c} = \hat{i} + 2\hat{j} - 3\hat{k}$ be three given vectors. If \vec{r} is a vector such that $\vec{r} \times \vec{a} = \vec{c} \times \vec{a}$ and $\vec{r} \cdot \vec{b} = 0$, then $|\vec{r}|$ is equal to :

- a) $\frac{11}{7} \sqrt{2}$
- b) $\frac{11}{7}$
- c) $\frac{11}{5} \sqrt{2}$
- d) $\frac{\sqrt{914}}{7}$

4) If $A = \frac{1}{2} \begin{pmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix}$, then :

- a) $A^{30} - A^{25} = 2I$
- b) $A^{30} + A^{25} + A = I$
- c) $A^{30} + A^{25} - A = I$
- d) $A^{30} = A^{25}$

5) Two dice are thrown independently. Let A be the event that the number appeared on the 1st die is less than the number appeared on the 2nd die, B be the event that the number appeared on the number appeared on the 1st die is even and that on the second die is odd, and C be the event that the number appeared on the 1st die is odd and that on the 2nd is even. Then

- a) the number of favourable cases of the event $(A \cup B) \cap C$ is 6
- b) A and B are mutually exclusive
- c) The number of favourable cases of the events A, B and C are 15, 6 and 6 respectively
- d) B and C are independent

6) Which of the following statements is a tautology ?

- a) $p \rightarrow (p \wedge (p \rightarrow q))$

- b) $(p \wedge q) \rightarrow (\neg(p) \rightarrow q)$
- c) $(p \wedge (p \rightarrow q)) \rightarrow \neg q$
- d) $p \vee (p \wedge q)$

7) The number of integral values of k , for which one root of the equation $2x^2 - 8x + k = 0$ lies in the interval $(2, 3)$, is:

- a) 2
- b) 0
- c) 1
- d) 3

8) Let $f : R - 0, 1 \rightarrow R$ be a function such that $f(x) + f\left(\frac{1}{1-x}\right) = 1 + x$. Then $f(2)$ is equal to :

- a) $\frac{9}{2}$
- b) $\frac{9}{4}$
- c) $\frac{7}{4}$
- d) $\frac{7}{3}$

9) Let the plane P pass through the intersection of the planes $2x + 3y - z = 2$ and $x + 2y + 3z = 6$, and be perpendicular to the plane $2x + y - z + 1 = 0$. If d is the distance of P from the point $(-7, 1, 1)$, then d_2 is equal to:

- a) $\frac{250}{83}$
- b) $\frac{15}{83}$
- c) $\frac{25}{83}$
- d) $\frac{250}{82}$

10) Let a, b be two real numbers such that $ab < 0$. If the complex number $\frac{1+ai}{b+i}$ is of unit modulus and $a+ib$ lies on the circle $|z - 1| = |2z|$, then a possible value of $\frac{1+|a|}{4b}$, where $[t]$ is greatest integer function, is :

- a) $-\frac{1}{2}$
- b) -1
- c) 1
- d) $\frac{1}{2}$

11) The sum of the absolute maximum and minimum values of the function $f(x) = |x^2 - 5x + 6| - 3x + 2$ in the interval $[-1, 3]$ is equal to:

- a) 10
- b) 12
- c) 13
- d) 24

12) Let $P(S)$ denote the power set of $S = \{1, 2, 3, \dots, 10\}$. Define the relations R_1 and R_2 on $P(S)$ as AR_1B if $(A \cap B^c) \cup (B \cap A^c) = \emptyset$ and AR_2B if $A \cup B^c = B \cup A^c$, $\forall A, B \in P(S)$. Then :

- a) both R_1 and R_2 are equivalence relations
- b) only R_1 is an equivalence relation
- c) only R_2 is an equivalence relation
- d) both R_1 and R_2 are not equivalence relations

13) The area of the region given by $\{(x, y) : xy \leq 8, 1 \leq y \leq x^2\}$ is :

- a) $8 \log_e^2 - \frac{13}{3}$
- b) $16 \log_e^2 - \frac{14}{3}$
- c) $8 \log_e^2 + \frac{7}{6}$
- d) $16 \log_e^2 + \frac{7}{3}$

14) Let $\alpha x = \exp(x^\beta y^\gamma)$ be the solution of the differential equation $2x^2 y dy - (1 - xy^2) dx =$

$0, x > 0, y(2) = \sqrt{\log_e^2}$. Then $\alpha + \beta - \gamma$ equals :

- a) 1
- b) -1
- c) 0
- d) 3

15) The value of the integral $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x + \frac{\pi}{4}}{2 - \cos 2x} dx$ is :

- a) $\frac{\pi^2}{6}$
- b) $\frac{\pi^2}{12\sqrt{3}}$
- c) $\frac{\pi^2}{3\sqrt{3}}$
- d) $\frac{\pi^2}{6\sqrt{3}}$