

BY ODHIAMBO DARREN GREGORY

+254704597085



KENYATTA UNIVERSITY

INSTITUTE OF OPEN LEARNING

SMA 210:
CALCULUS II

BY
D. SENGOTTAIYAN

DEPARTMENT OF MATHEMATICS

PREFACE

This **module** is designed primarily to provide the readers with the best preparation possible for the **Advanced** calculus examinations.

In its present form this **module** has developed from courses given by the author over the last thirty two years to the audience of mathematicians, physics and engineers in the University of Madras, Kenyatta University and the University of Nairobi.

BY KELVIN EKALIYO, THE NEXT.

It is hoped that it will be of great interest to students of pure and Applied Mathematics following **Advanced** calculus. Each lesson begins with a brief statement of definitions. Principles and important theorems followed by a set of solved and graded supplementary problems. Attention has been given to the lessons on applications of the theory.

The author is pleased to acknowledge Dr. L. O. Odongo who worked through the entire manuscript and checked all the problems in each and every lesson.

CONTENTS

CHAPTER 1: Integration of Functions Using Basic Formulae **1**

- 1.1 Introduction
- 1.2 Objectives of the Lesson
- 1.3 Meaning of Integration
- 1.4 Constant of Integration
- 1.5 The Notation
- 1.6 Basic Formula for Integration of Functions
- 1.7 Addition and Subtraction Rule in Integration
- 1.8 Integration of Constant Times a Function
- Worked examples
- Exercise
- Summary
- Further Reading

CHAPTER 2: Integration of Functions using the Extension of the Basic Formulae **8**

- 2.1 Introduction
- 2.2 Objective of the Lesson
- 2.3 Definition of Integrand
- 2.4 The Extension of the Basic Formulae
- 2.5 Proof of the Extension Formulae

BY KELVIN EKALIYO, THE NEXT.

Worked examples

Exercise

Summary

Further Reading

CHAPTER 3: Integration using Substitutions

16

3.1 Introduction

3.2 Objectives of the Lesson

3.3 Integration using Substitutions

3.4 Type I

3.5 Type II

3.6 Type III

3.7 Integrand containing u and u' in other forms, ($e^u u^1$, $\sin u u^1$) Worked examples

Exercise

Summary

Further Reading

CHAPTER 4: Integration by Parts

26

4.1 Introduction

4.2 Objectives of the Lesson

4.3 Formulae for Integration by Parts

4.4 Formula for Integration of $\ln x$

Worked examples

Exercise

Summary

Further Reading

CHAPTER 5: Integration using Trigonometric Identities

34

5.1 Introduction

5.2 Objectives of the Lesson

5.3 Review of important Trigonometric identities

5.4 Integration using the three fundamental identities

5.5 Integration using the formulae for $\cos 2x$

- 5.6 Integration using the product formulae
- 5.7 Integration using $\sin 3x$ and $\cos 3x$ formulae
- 5.8 Integrand containing quadratic expressions in the Denominator.

Worked examples

Exercise

Summary

CHAPTER 6: Integration using Partial Fractions

47

- 6.1 Introduction
- 6.2 Objectives of the Lesson
- 6.3 Integrand Containing only non-repeated linear Factors in the Denominator (Proper Fraction)
- 6.4 Integrand Containing linear Factors (non-repeated) in the Denominator (Proper Fraction)
- 6.5 Integrand Containing Quadratic Factors.
- 6.6 Integrand is an (Improper Fraction)
- Worked examples Exercise
- Summary
- Further Reading

CHAPTER 7: Definite Integrals and the Fundamental Theorem of the Integral Calculus.

53

- 7.1 Introduction
- 7.2 Objectives of the Lesson
- 7.3 Meaning of the Definite Integral
- 7.4 Important Properties of Definite Integral
- 7.5 Definition of Indefinite Integral
- 7.6 The Fundamental Theorem of the Integral Calculus
- 7.7 Evaluation of Definite Integrals
- Worked examples
- Exercise
- Summary
- Further Reading

CHAPTER 8: Area Under a Curve

61

- 8.1 Introduction
- 8.2 Objectives of the Lesson
- 8.3 Formula for the Area under a Curve
- 8.4 Area Below x – axis
- 8.5 Combined Area under a Curve
- 8.6 Area Between Two Intersecting Curves.
 - Worked examples
 - Exercise
 - Summary
 - Further Reading

CHAPTER 9: Improper Integrals

71

- 9.1 Introduction
- 9.2 Objectives of the Lesson
- 9.3 Definition of Improper Integrals
- 9.4 Convergence and Divergence of the Improper Integrals of the First Kind
- 9.5 Definition of Absolute and Conditional Convergence
- 9.6 Cauchy's Inequality for convergence of an Integral
- 9.7 Absolute Convergence of an Integral Implies Convergence
- 9.8 Convergence and Divergence of the Improper Integrals of the Second Kind.
- 9.9 The Cauchy's Principal value of Improper Integrals with Singularities. Worked examples
 - Exercise
 - Summary of the Chapter
 - Further Reading

CHAPTER 10: Length of arc of a Curve

83

- 10.1 Introduction
- 10.2 Objectives of the Lesson
- 10.3 Length of the Arc in Cartesian Form
- 10.4 Length of the Arc in Parametric Form
- 10.5 Length of the Arc in Polar Coordinates
 - Worked examples
 - Exercise
 - Summary
 - Further Reading

CHAPTER 11: Volume and Surface Area of Solids of Revolution

93

- 11.1 Introduction

- 11.2 Objectives of the Lesson
- 11.3 The Formula for the Volume of Solids of Revolution
- 11.4 The Formula for the Surface Area of Solids of Revolution
 - Worked examples
 - Exercise
 - Summary of the Chapter
 - Further Reading

CHAPTER 12: Numerical Integration (Trapezoidal and Simpsons Rules)

102

- 12.1 Introduction
- 12.2 Objectives of the Lesson
- 12.3 Area of a Trapezium
- 12.4 Trapezoidal Rule
- 12.5 Principle of Simpson's Rule
- 12.6 Simpsons Rule
- Worked example
 - Exercise
 - Summary of the chapter
 - Further Reading

**ANSWERS
INDEX**

115

CHAPTER 1

Integration of functions using Basic Formulae

1.1 Introduction

The process of integration is defined as the reverse or converse process of differentiation. In Differential Calculus, you have studied differentiation of functions and its applications. Now we shall learn the process of integration and its applications in the **Integral Calculus**. In this chapter we shall learn **TEN BASIC FORMULAE** for integration and integrate expressions using them.

1.2 Objectives of the chapter

By the end of this chapter you should be able to

- state the Ten Basic Formulae for integration.
- integrate simple functions using the Ten Basic Formulae only.

1.3 Meaning of Integration

Integration is defined as the reverse or the converse process of differentiation.

For example,

$\frac{d}{dx}$

We know that $\frac{d}{dx}(\sin x) = \cos x$

Now we say that the integration of $\cos x$ is $\sin x$.

We write this in symbols

$$\int \cos x dx = \sin x$$

(The symbol $\int$ is pronounced as “the integration of”. It was first used by the French Mathematician, Leibnitz. He used this symbol for the first letter ‘S’ of the word Summation)

1.4 Constant of Integration

$\frac{d}{dx}$

Now $\frac{d}{dx}(\sin x) = \cos x$ $\frac{d}{dx}(\sin x + c) = \cos x$. In fact any constant c , that is added dx

disappears in the process of differentiation. Hence we always put a constant c after the

process of integration. Hence we write $\int \cos x dx = \sin x + c$. Here c is called the **constant of integration**.

BY KELVIN EKALIYO, THE NEXT.

1.5 The Notation and the Integrand

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

If $f(x)$ is any function of x , $\int f(x)dx$ stands for integral of $f(x)$ with respect to x . the integral sign $\int$ cannot be divorced from dx if the integral is with respect to x .

The function $f(x)$ in $\int f(x)dx$ is called the integrand of the integral. For example in $\int x^2 dx$, x^2 is called the integrand of the integral.

1.6 (A) Basic Formulae for Integration of Algebraic functions.

$$\int x^{n-1} dx = \frac{x^n}{n} + c, \text{ if } n \neq 0$$

$$\int \frac{1}{x} dx = \int x^{-1} dx = \ln x + c$$

For example

$$\int x^{10} dx = \frac{x^{11}}{11} + c$$

$$\int x dx = \frac{x^2}{2} + c$$

1.7 (B) Basic Formulae for Integration of Exponential functions.

BY KELVIN EKALIYO, THE NEXT.

a_x

$$\int a^x dx = \frac{a^x}{\ln a} + c, a \text{ is any positive}$$

□

3. constant

 x
 x

$$4. \int e^x dx = e^x + c$$

For example

$$\int 2^x dx = \frac{2^x}{\ln 2} + c$$

 5_x

$$\int 5^x dx = \frac{5^x}{\ln 5} + c$$

1.8 (C) Basic Formulae for Integration of Trigonometric functions

$$5. \int \cos x dx = \sin x + c, \text{ since } \frac{d}{dx}(\sin x) = \cos x$$

MWALIMU GREG

$$7. \int \sec^2 x dx$$

$$8. \int \tan x dx = C, \text{ since } \frac{d}{dx} (\tan x) = \sec^2 x$$

$\int$

$$\int \sec x \tan x dx = \sec x + C, \text{ since } \frac{d}{dx} (\sec x) = \sec x \tan x$$

d

$$6. \int \sin x dx = -\cos x + C, \text{ since } \frac{d}{dx} (\cos x) = -\sin x$$

$$\int \cos x dx = \sin x + C, \text{ since } \frac{d}{dx} (\sin x) = \cos x$$

$$\int \sin x dx = -\cos x + C, \text{ since } \frac{d}{dx} (\cos x) = -\sin x$$

$$\int \cos x dx = \sin x + C, \text{ since } \frac{d}{dx} (\sin x) = \cos x$$

$$\int \sin x dx = -\cos x + C, \text{ since } \frac{d}{dx} (\cos x) = -\sin x$$

$$9. \int \sec x \tan x dx = \sec x + C, \text{ since } \frac{d}{dx} (\sec x) = \sec x \tan x$$

$$10. \int \cos x dx = \sin x + C, \text{ since } \frac{d}{dx} (\sin x) = \cos x$$

Note that we shall consider integration of $\tan x$, $\cot x$, $\sec x$ and $\csc x$ afterwards.

1.9 Addition and subtraction Rule in Integration

$$\text{Just like } \frac{d}{dx} (u \pm v \pm w) = \frac{d}{dx} u \pm \frac{d}{dx} v \pm \frac{d}{dx} w, \frac{d}{dx}$$

$$\text{we have } \int (u \pm v \pm w) dx = \int u dx \pm \int v dx \pm \int w dx,$$

where u , v , w are functions of x .

For example

$$\int (x^2 + \sin x + e^x) dx = \int x^2 dx + \int \sin x dx + \int e^x dx$$

$$= \frac{x^3}{3} - \cos x + e^x + C$$

BY KELVIN EKALIYO, THE NEXT.

1.10 Integration of constant times a function

$$\int kf(x)dx = k \int f(x)dx$$

For example $\int 5x dx = 5 \int x dx = \frac{5x^2}{2} + c$

$$\int 5x dx = 5 \int x dx = \frac{5x^2}{2} + c$$

$$\int 8 \cos x dx = 8 \int \cos x dx = 8 \sin x + c$$

Example 1

Find $\int (2x^5 + 3e^x + 4x + 5)dx$

Solution

$$\begin{aligned} \int (2x^5 + 3e^x + 4x + 5)dx &= 2 \int x^5 dx + 3 \int e^x dx + 4 \int x dx + 5 \int dx \\ &= \frac{2x^6}{6} + 3e^x + \frac{4x^2}{2} + 5x + c \\ &= \frac{x^6}{3} + 3e^x + 2x^2 + 5x + c \end{aligned}$$

We should put one constant c after the process of integration.

Note that

BY KELVIN EKALIYO, THE NEXT.

$$\int dx = x + c$$

$$\int 5dx = 5 \int dx = 5x + c$$

$$\int x dx = \frac{x^2}{2} + c$$

Example 2

Find $\int (2x + 3)(x + 2) dx$

Solution

First we find the product and then we integrate

$$(2x + 3)(x + 2)$$

$$= 2x^2 + 4x + 3x + 6$$

$$= 2x^2 + 7x + 6$$

$$\int (2x^2 + 7x + 6) dx = \frac{2x^3}{3} + \frac{7x^2}{2} + 6x + c$$

Example 3

$$\frac{2x^3 + 5x^2 + 8x + 9}{x}$$

Find $\int \frac{2x^3 + 5x^2 + 8x + 9}{x} dx$

Solution

First we divide each term by x and then we integrate

$$\frac{2x^3 + 5x^2 + 8x + 9}{x} = 2x^2 + 5x + 8 + \frac{9}{x}$$

$$\int (2x^2 + 5x + 8 + \frac{9}{x}) dx = \frac{2x^3}{3} + \frac{5x^2}{2} + 8x + 9 \ln|x| + c$$

BY KELVIN EKALIYO, THE NEXT.

$$\begin{aligned}
 &= \int (2x^3 - 5x - 8x + \frac{9}{x}) dx \\
 &= \frac{2x^4}{4} - \frac{5x^2}{2} - 8x + 9 \ln x + c \\
 &= \frac{1}{2} x^4 - \frac{5x^2}{2} - 8x + 9 \ln x + c
 \end{aligned}$$

Example 4

$$\int (x^2 - 5x + 6) dx$$

Find $\int (x^2 - 5x + 6) dx$

Solution

First we simplify the integrand $x^2 - 5x + 6$ and then we integrate. $x - 2$

$$\begin{aligned}
 \int (x^2 - 5x + 6) dx &= \int \frac{(x - 3)(x - 2)}{(x - 2)} dx \\
 &= \int (x - 3) dx \\
 &= \frac{x^2}{2} - 3x + c
 \end{aligned}$$

Example 5

$$\begin{aligned}
 &9x - \frac{1}{x^3} + x^9 - 2^x - 3e^x - x^{-1} \\
 &\text{Integrate}
 \end{aligned}$$

BY KELVIN EKALIYO, THE NEXT.

Solution $\int (9x^3 - 9 - 2x + 3e^x - x + 1) dx$

$$= \frac{9x^4}{4} - 9x - \frac{x^2}{2} + 3e^x - \frac{x^2}{2} + x + c$$

Example 6

Find $\int (2\cos x - 3\sin x + 4\sec^2 x + 5\cos e^x \cot x) dx$

Solution

$$\int (2\cos x - 3\sin x + 4\sec^2 x + 5\cos e^x \cot x)$$

$$dx = 2 \sin x - 3 \cos x + 4 \tan x + 5 \cos e^x$$

$+ c$ **Example 7**

Find $\int (\frac{2}{x} + 5\sec x \tan x + 6\operatorname{cosec}^2 x + 4^x + 8) dx$

Solution

$$\int (\frac{2}{x} + 5\sec x \tan x + 6\operatorname{cosec}^2 x + 4^x + 8) dx$$

$$= 2 \ln |x| + 5 \sec x + 6 \cot x + \frac{4^x}{\ln 4} + 8x + c$$

$$\operatorname{cosec}^2 x dx = -\cot x + c$$

$$= \frac{2}{x} + 5 \sec x + 6 \cot x + \frac{4^x}{\ln 4} + 8x + c$$

BY KELVIN EKALIYO, THE NEXT.

$$= 2 \ln x - 5 \sec x - 6 \cot x - \frac{x^4}{4} - 8x + c$$

Exercise 1

Using the ten basic formula for integration find

1. $\int (2x^5 - 3e^x + 4x - 1) dx$
2. $\int (3 \sin x - 4 \sec x \tan x + 5 \sec^2 x) dx$
3. $\int (3 \cos x - 4 \cos e^x \cot x + 5 \cos e^{2x}) dx$
4. $\int (x^{15} - x^{-15} + 8x^4 + 3e^x + 7x + 8) dx$
5. $\int \frac{(x^2 + 7x + 10)}{(x + 5)} dx$
6. $\int \frac{3x^2 + 4x^3 + 8x^5 + 9x + 2}{(x + 1)} dx$
7. $\int \frac{x^2}{(x^2 + 3x + 1)(x^3 + 7x^2 + 2)} dx$
8. $\int \frac{x^2}{x^2} dx$
9. If $\int (2x + 9) dx = 0$ find the values of x

BY KELVIN EKALIYO, THE NEXT.

10. If $\int (3x^2 + 2x + 12) dx = 0$ find the values of x
 $\int x^3 + 5x^2 + 6x + 12$

11. Find $\int (x^2 + 2x) e^{2x} dx$
 $\int e^{2x} + e^{5x} + e^x dx$

12. Find $\int e^x dx$

You have learnt the following from this chapter

- i). Integration is the reverse or converse process of differentiation.
- ii). The meaning of “integrand” and constant of integration. iii).

The Ten Basic Formula for Integration, namely

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \text{ if } n \neq -1$$

1. $\int x^n dx = \frac{x^{n+1}}{n+1} + C$

2. $\int \frac{1}{x} dx = \ln x + C = \ln Ax$

3. $\int a^x dx = \frac{a^x}{\ln a} + C$

4. $\int e^x dx = e^x + C$

5. $\int \cos x dx = \sin x + C$

6. $\int \sin x dx = -\cos x + C$

7. $\int \sec^2 x dx = \tan x + C$

BY KELVIN EKALIYO, THE NEXT.

$$8. \int \sec x \tan x dx = \sec x + c ,$$

$$9. \int \cos ecx \cot x dx = \cos ecx + c ,$$

$$10. \int \cos ec^2 x dx = \cot x + c ,$$

Further Reading

1. Pure mathematics (First Course)
By J. K Backhouse and others
Longman Group Ltd Harlow,
Essex, U.K

CHAPTER 2

Integration of Functions Containing First Degree Expressions – Extension of the Basic Formula.

2.1 Introduction

In chapter one you have learnt to integrate functions using the Ten Basic Formulae. Unlike differentiation of functions, integration is not so easy, however we have about six techniques of integration with which a given function can be integrated. One of the techniques is integration using the Extension formulae. Suppose that a first degree expression is present in the place of x in the Ten Basic formula, we can use conveniently the same Basic formulae and obtain the result. The new formulae are called “the extension formulae”. In this chapter we shall integrate expressions containing the first degree, using the Ten Extension Formulae.

2.2 Objectives of the chapter

By the end of this chapter you should be able to

- write the Ten Extension Formulae corresponding to the Ten Basic formulae.
- apply the Ten Extension Formulae to integrate expressions containing first degree in x .

2.3 Definition of the Integrand

The expression to be integrated is written between the integral symbol and the corresponding dx symbol. This expression or the function is called the **Integrand** of the integral.

BY KELVIN EKALIYO, THE NEXT.

For example in $\int x^2 e^x dx$, the expression $x^2 e^x$ is called the “integrand” of the integration.

Similarly in $\int \cos x dx$, $\cos x$ is called “the integrand” of the integral.

2.4 The extension of the Basic formulae

In each of the ten Basic formulae the integrand contains x . Suppose we replace x in both sides of the formula by a **first degree** expression in x of the form $(2x + 3)$, $(5x - 7)$, $(3 - 4x)$ or $(ax + b)$ in general. All the ten formulae hold good when we divide the final result by coefficient of x namely a . The Ten Extension Formulae are given in the following table:

Basic Formulae	Corresponding Extension formulae with an example – Applicable for first degree in x only
$\int x^n dx = \frac{x^{n+1}}{n+1} + c$ 1. $\int x^n dx$ $n \neq -1$	$\int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{a(n+1)} + c$ e.g. $\int (2x + 3)^{10} dx = \frac{(2x + 3)^{11}}{11 \cdot 2} + c$
$\int \frac{1}{x} dx = \ln x + c$ 2. $\int \frac{1}{x} dx$	$\int \frac{1}{ax + b} dx = \frac{1}{a} \ln ax + b + c$ e.g. $\int \frac{1}{2x + 3} dx = \frac{1}{2} \ln 2x + 3 + c$

BY KELVIN EKALIYO, THE NEXT.

p^x 3. $\int p^x dx = \frac{1}{\ln p} p^x + c$ (p is positive)	p^{ax+b} $\int p^{ax+b} dx = \frac{1}{a \ln p} p^{ax+b} + c$ $5x+1$ 3^{5x+1} e.g. $\int 3^{5x+1} dx = \frac{1}{5 \ln 3} 3^{5x+1} + c$
4. $\int e^x dx = e^x + c$	e^{ax+b} $\int e^{ax+b} dx = \frac{1}{a} e^{ax+b} + c$ $4x+7$ e^{4x+7} e.g. $\int e^{4x+7} dx = \frac{1}{4} e^{4x+7} + c$
5. $\int \cos x dx = \sin x + c$	$\int \cos(ax+b) dx = \frac{1}{a} \sin(ax+b) + c$ $\frac{\sin(ax+b)}{\sin 100x}$ e.g. $\int \cos 100x dx = \frac{1}{100} \sin 100x + c$
6. $\int \sin x dx = -\cos x + c$	$\int \sin(ax+b) dx = -\frac{1}{a} \cos(ax+b) + c$ e.g. $\int \sin(100x+5) dx = -\frac{1}{100} \cos(100x+5) + c$
7. $\int \sec^2 x dx = \tan x + c$	$\int \sec^2(ax+b) dx = \frac{1}{a} \tan(ax+b) + c$ a e.g. $\int \sec^2(3x+4) dx = \frac{1}{3} \tan(3x+4) + c$

BY KELVIN EKALIYO, THE NEXT.

8. $\int \sec^2 x dx$	$\int \sec^2(ax + b) dx = \frac{1}{a} \tan(ax + b) + c$ e.g. $\int \sec^2(20x + 1) dx = \frac{1}{20} \tan(20x + 1) + c$
9. $\int \sec x \tan x dx$	$\int \sec(ax + b) \tan(ax + b) dx = \frac{1}{a} \sec(ax + b) + c$ e.g. $\int \sec 5x \tan 5x dx = \frac{1}{5} \sec 5x + c$
10. $\int \sec x \cot x dx$	$\int \sec(ax + b) \cot(ax + b) dx = \frac{1}{a} \ln \sec(ax + b) + c$ e.g. $\int \sec x \cot x dx = \ln \sec x + c$

2.5 Proof of the Ten Extension Formulae

We shall prove some of the Ten Extension formulae using the substitution

$u = ax + b$ so that $dx = \frac{1}{a} du$

$$a \frac{1}{a} du = du$$

Example 1

$$\int (ax + b)^{n-1} dx$$

BY KELVIN EKALIYO, THE NEXT.

a). Prove that $\int (ax + b)^n dx = \frac{a(n+1)}{n+1} (ax + b)^{n+1} + c$

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

b). Hence find $\int (4x - 9)^{20} dx$

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

Solution

$$\frac{d}{dx} \int_a^b f(x) dx = f(b) - f(a)$$

Let $u = ax + b$ so that $du = a dx$

$$\int_a^b \frac{1}{u^n} du = \int_a^b u^{-n} du = \frac{u^{-n+1}}{-n+1} \Big|_a^b = \frac{u^{1-n}}{1-n} \Big|_a^b$$

a) Then $\int_a^b \frac{1}{(ax+b)^n} dx = \frac{1}{a} \int_a^b \frac{1}{u^n} du$

$\frac{1}{a} \int_a^b \frac{1}{u^n} du$ (we use the Basic formula to get) $= \frac{1}{a} \left[\frac{u^{1-n}}{1-n} \right]_a^b$

$$\begin{aligned} &= \frac{1}{a} \left[\frac{u^{1-n}}{1-n} \right]_a^b \\ &= \frac{1}{a} \left[\frac{(ax+b)^{1-n}}{1-n} \right]_a^b \\ &= \frac{1}{a(n-1)} \left[(ax+b)^{n-1} \right]_a^b \end{aligned}$$

b) Using $\int_a^b x^{20} dx = \frac{1}{21} x^{21} \Big|_a^b$,

$$\int_a^b (4x+9)^{20} dx = \frac{1}{21} (4x+9)^{21} \Big|_a^b$$

$$= \frac{(4x+9)^{21}}{84} \Big|_a^b$$

Example 2

$$\frac{d}{dx} \ln(ax+b)$$

a). Prove that $\frac{d}{dx} \ln(ax+b) = \frac{a}{ax+b}$

$$\frac{d}{dx} \ln(ax+b) = \frac{a}{ax+b}$$

b). Find _____

$$\frac{d}{dx} \ln(7 - 100x)$$

Solution

a). We shall use the Basic formula $\frac{d}{dx} \ln x = \frac{1}{x}$ to prove this result

$$\frac{d}{dx} \ln(7 - 100x)$$

Let $u = 7 - 100x$ so that $\frac{du}{dx} = -100$

$$\frac{d}{dx} \ln(7 - 100x) = \frac{d}{du} \ln u \cdot \frac{du}{dx}$$

$$= \frac{1}{u} \cdot (-100) = \frac{-100}{7 - 100x}$$

b). Using the extension formula

$$\frac{d}{dx} \ln(ax + b) = \frac{a}{ax + b}$$

$$\frac{d}{dx} \ln(7 - 100x) = \frac{-100}{7 - 100x}$$

unless you are asked to prove $\frac{d}{dx} \ln(7 - 100x) = \frac{-100}{7 - 100x}$ you can just write

$$\frac{d}{dx} \ln(7 - 100x) = \frac{-100}{7 - 100x}$$

down the result without proof.

Example 3

Prove that $\int e^{ax+b} dx = \frac{1}{a} e^{ax+b} + c$ where a and b are constants.

Proof

The integrand contains a linear function of x namely (a x + b)

$$\text{Let } u = a x + b \quad (1) \quad \text{so that } \frac{du}{dx} = a \quad (2)$$

$$\text{and } dx = \frac{du}{a} \quad (3)$$

Substitution of (1) and (3) in the given integral

$$\begin{aligned} \int e^{ax+b} dx &= \int e^u \frac{du}{a} \\ &= \frac{1}{a} \int e^u du \\ &= \frac{1}{a} e^u + c \\ &= \frac{e^{ax+b}}{a} + c \end{aligned}$$

In the same way by taking the linear function of x as u or u = a x + b we can establish all the ten extension formulae.

Example 4

Find $\int (2x+3)^5 dx$

Solution

We compare this integration with $\int x^5 dx$

$$\int x^5 dx = \frac{x^6}{6} + c$$

By the extension technique

$$\int (2x - 3)^6 dx = \frac{(2x - 3)^7}{7} + c$$

Example 5

Find $\int \frac{1}{(5x - 7)} dx$ (1)

Solution

We compare (1) with $\int \frac{1}{x} dx = \ln x + c$

By the extension technique to $\int \frac{1}{(ax + b)} dx = \frac{1}{a} \ln(ax + b) + c$

Example 6

Find $\int e^{3 - 4x} dx$ (1)

Solution

We compare (1) with the Basic formula

$$\int e^x dx = e^x + c$$

Then $\int e^{3 - 4x} dx = -\frac{1}{4} e^{3 - 4x} + c$

Example 7

Find $\int \sin(2 - 3x) dx$ (1)

Solution

Comparing (1) with the Basic formula

$$\int \sin x dx = -\cos x + c$$

$$\cos(2 - 3x)$$

$$\text{where } \int \sin(2 - 3x) dx = \frac{1}{3} \cos(2 - 3x) + c$$

Example 8

$$\text{Find } \int \cos(5 - 4x) \cot(5 - 4x) dx \quad (1)$$

Solution

Comparing (1) with the corresponding basic formula

BY KELVIN EKALIYO, THE NEXT.

$$\int \cos ecx \cot x dx = \int \cos ecx = c$$

$$\int \operatorname{cosec}(5-4x)$$

$$\text{We have } \int \cos ec(5-4x) \cot(5-4x) dx = \frac{1}{4} \int c$$

Note that the same $(5-4x)$ should be present in both the places in (1), or else we cannot use the formula.

Example 9

$$\text{Find } \int (3x-1)^{15} dx \quad (1)$$

Solution

$$\int (3x-1)^{15} dx = \frac{1}{16} \int (3x-1)^{16} dx \quad \text{we have}$$

$$\text{Using } \int (x)^n dx = \frac{x^{n+1}}{n+1}$$

$$\begin{aligned} \int (3x-1)^{15} dx &= \frac{1}{16} \int (3x-1)^{16} dx \\ &= \frac{1}{16} \cdot \frac{(3x-1)^{17}}{3} + c \\ &= \frac{(3x-1)^{17}}{48} + c \end{aligned}$$

Note:

You should note that the extension formula is applicable only when the integrand is of first degree in x , such as $2x+3$, $5-2x$ or $(ax+b)$ in general when a and b are constants.

Example 10

$$\text{Find } \int \frac{1}{3x-8} dx$$

$$\int \frac{1}{3x-8} dx$$

Solution

$$\text{Using the Basic formula } \int \frac{1}{x} dx = \ln x + c$$

BY KELVIN EKALIYO, THE NEXT.

$$\frac{1}{3} \frac{d}{dx} \ln(3x - 8) = \frac{1}{3} \cdot \frac{3}{3x - 8} = \frac{1}{3x - 8}$$

We have $\int \frac{1}{3x - 8} dx = \frac{1}{3} \ln(3x - 8) + c$

Example 11

Find $\int \sec 5x \tan 5x dx$

Solution

Using the Basic formula $\int \sec x \tan x dx = \sec x + c$

$$\text{We have } \int \sec 5x \tan 5x dx = \frac{\sec 5x}{5} + c$$

BY KELVIN EKALIYO, THE NEXT.

Example Find $\int (1 - 2x)^{10} e^{2-3x} (3 - \ln x) \sec 2x \, dx$

$$\int (1 - 2x)^{10} e^{2-3x} (3 - \ln x) \sec 2x \, dx$$

Solution

Using the corresponding Basic formula, we have

$$\int (1 - 2x)^{10} e^{2-3x} (3 - \ln x) \sec 2x \, dx = \frac{(1 - 2x)^{11} e^{2-3x} \ln(3 - x) \csc 7x}{11} + C$$

Exercise 2

1. $\cos(4x + 3)$

2. $\sin 3x$

3. $\sin \frac{1}{2}x$

4. e^{-2x}

5. e^{3x-2}

6. $\frac{1}{2x-5}$

7. $(4x - 3)^3$

8. $\cos(5x + 4)$

9. $\sin(3 - 4x)$

10. $\sec 3x \tan 3x$

11. $\operatorname{cosec} 2x \cot 2x$

Find the integrals of the following

12. $\int (\sin 13x \cos 4x + e^{2x}) \, dx$

13. $\int 2^{x-5} e^{x-5} \cos e^{25x} (2x - 3)^5 \, dx$

10. $\int \frac{1}{\cos(ax + b)} \, dx$

BY KELVIN EKALIYO, THE NEXT.

14. $\int \frac{ax + b}{(ax + b)^2} dx$

$\int$

$\int \frac{1}{x^2} dx = -\frac{1}{x} + C$

15. $\int \sec^2 x \tan^2 x \cos^2 x dx$

$\int \frac{1 - \sin^2(3x)}{(3 - 2x)^2} dx$

16.

$\int \frac{1}{(3 - 2x)(3 + 2x)} dx$

MWALIMU GREG

Summary

You have learnt the following from this chapter

- to integrate expressions using the Ten Extension Formulae

Further Reading

1. Advanced Calculus

By Watson Fulks

John Wiley & sons

New York Brisbane. Toronto

MWALIMU GREG

CHAPTER 3

Integration Using Substitutions.

3.1 Introduction

In the previous chapters you have learnt to integrate expressions using the Ten Basic formulae and the extension of the formulae. In this chapter you will learn one of the most important techniques of integration of expressions consisting of two functions one of which is the derivative of the other.

3.2 Objectives of the Chapter:

By the end of this chapter you should be able to

- recognize a function u and its derivative u' in the integrand & integrate the expression using the substitution u and u' .
- adjust the derivative so that the integrand consists of one function u and its derivative.
- derive the formulae for integration of $\tan x$ and $\cos x$.

3.3 Integration using substitution.

When the integrand consists of two functions one of which is the derivative of the other function we use the substitution technique of integration. We call the main du function as u and its derivative as u' or $\frac{du}{dx}$ so that $u' dx$ will become du . The dx substitution transforms the integral into one of the basic formula of integration in the new variable u only. Hence we integrate easily using the basic formula. The technique is illustrated in the following examples.

Type 1

$$\int \frac{u'}{u} dx = \ln u + c$$

Example 1

$$3x^2 + \cos x$$

Consider $\int (3x^2 + \cos x) dx$

$3x^2 + \cos x$ is the derivative of $x^3 + \sin x$.

Let us use the substitution

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

BY KELVIN EKALIYO, THE NEXT.

$$u = x^3 + \sin x \quad (1)$$

$$\text{Then } \frac{du}{dx} = 3x^2 + \cos x$$

$$\text{Then } du = (3x^2 + \cos x) dx. \quad (2)$$

Using (1) and (2) the integral is transformed into $\int \frac{du}{u}$

which comes under one of the basic formulae

$$\text{Hence } \int \frac{du}{u} = \ln |u| + c$$

$$= \ln u + c$$

$$= \ln (x^3 + \sin x) + c$$

Type 2

$$\int u^n du = \frac{u^{n+1}}{n+1} + c$$

$$\int u^n dx = \int \frac{u^n}{u^{n-1}} du = \int u^{n-(n-1)} du = \int u^1 du = \frac{u^2}{2} + c$$

Example 2

$$3x^2 + \cos x$$

$$\text{Consider } \int (3x^2 + \cos x) dx$$

Again there are two functions $x^3 + \sin x$ and its derivative $3x^2 + \cos x$ (leaving the index ten)

$$\text{Let } u = x^3 + \sin x \quad (1)$$

$$\text{Then } \frac{du}{dx} = 3x^2 + \cos x$$

$$\text{or } du = (3x^2 + \cos x) dx \quad (2)$$

Substitution of (1) and (2) in the given integral

$$\int (3x^2 + \cos x) dx$$

$$\int (x^3 + \sin x)^{10} dx = \int u^{10} du$$

$$= \frac{u^{11}}{11} + c \quad (\text{using the Basic formula we have})$$

$$= \frac{u^{11}}{11} + c$$

$$= \frac{1}{11} u^{11} + c$$

$$= \frac{1}{11} (x^3 + \sin x)^{11} + c$$

Type 3

$$\int e^{u(x)} dx = \int e^u du = e^u + C$$

Consider $\int e^{x^3 + \sin x} (3x^2 + \cos x) dx$

The integrand contains the function $u = x^3 + \sin x$ and its derivative $(3x^2 + \cos x)$

Let $u = x^3 + \sin x$ (1) $\frac{du}{dx} = 3x^2 + \cos x$

Then $du = (3x^2 + \cos x) dx$ (2)

Substitution of (1) and (2) in the given integral

$$\int e^{x^3 + \sin x} (3x^2 + \cos x) dx = \int e^u du$$

using the basic formula

BY KELVIN EKALIYO, THE NEXT.

$$\square e^u \square c$$

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

$$\int e^{x^3} \sin x \, dx = C$$

$$\int u^{n-1} \, du = \frac{u^n}{n} + C \quad n \neq 0$$

Type 4 $\int u^n \, dx = \int u \, du = \frac{u^{n+1}}{n+1} + C \quad n \neq -1$

Consider $\int (x^3 + 3x^2 + 6x + 2)^{15} (3x^2 + 6x + 6) \, dx$

The integrand contains the function $u = x^3 + 3x^2 + 6x + 2$
and its derivative $(3x^2 + 6x + 6)$

Let $u = x^3 + 3x^2 + 6x + 2$

$$\text{and } du = (3x^2 + 6x + 6) \, dx \quad (2)$$

Substitution of (1) and (2) in the given integral

$$\int (x^3 + 3x^2 + 6x + 2)^{15} (3x^2 + 6x + 6) \, dx = \int u^{15} \, du$$

$$\begin{aligned} &= \frac{u^{16}}{16} + C \\ &= \frac{(x^3 + 3x^2 + 6x + 2)^{16}}{16} + C \end{aligned}$$

Type 5

$$\int \sin u \, du = -\cos u + C$$

Consider $\int \sin (x^3 + x^2 + 5) (3x^2 + 2x) dx$.

The integrand contains a function $u = x^3 + x^2 + 5$

(within sine) and its derivative $u' = 3x^2 + 2x$

Let $u = x^3 + x^2 + 5$ (1) du

Then $\frac{du}{dx} = 3x^2 + 2x$

or $du = (3x^2 + 2x) dx$ (2)

Substitution of (1) and (2) in the given integral

BY KELVIN EKALIYO, THE NEXT.

$\int \sin (x^3 + x^2 + 5) (3x^2 + 2x) dx$.

$= \int \sin u du$ which is a standard formula

$= -\cos u + c = -\cos$

$(x^3 + x^2 + 5) + c$

Type 6

$\int \operatorname{cosec} u \cot u \cdot u' dx = \int \operatorname{cosec} u \cot u du = -\operatorname{cosec} u + c$

Consider $\int \operatorname{cosec} x^2 \cot x^2 (2x) dx$

Hence $u = x^2$ and $\frac{du}{dx} = 2x$ or $du = 2x dx$

Substitution of u in the given integral gives

$\int \operatorname{cosec} u \cot u du = -\operatorname{cosec} u + c$ using formula
 $= -\operatorname{cosec} x^2 + c$ **Example**

3

BY KELVIN EKALIYO, THE NEXT.

The integrand $\tan x \sec^2 x$ contains dx $u = \tan x$ and its derivative $\sec^2 x$

$$\text{Let } u = \tan x \quad (1) \quad du$$

$$\text{Then } \frac{du}{dx} = \sec^2 x$$

$$\text{or } du = \sec^2 x \, dx \quad (2)$$

Substitution of (1) and (2) in the given integral

$$\begin{aligned} \int \tan x \sec^2 x \, dx &= \int u \, du \\ &= \frac{u^2}{2} + c \\ &= \frac{\tan^2 x}{2} + c \end{aligned}$$

Example 4

Find

$$\int e^{\sin x} \cos x \, dx$$

BY KELVIN EKALIYO, THE NEXT.

Let $u = \sin x$, $du = \cos x dx$

$$\text{Hence } \int e^{\sin x} \cos x dx = \int e^u du \\ = e^u + c$$

$$= e^{\sin x} + c$$

Example 5

Find $\int (\tan x)^{10} \sec^2 x dx$

Let $u = \tan x$, then $du = \sec^2 x dx$

$$\text{Hence } \int (\tan x)^{10} \sec^2 x dx = \int u^{10} du \\ = \frac{u^{11}}{11} + c$$

$$= \frac{\tan^{11} x}{11} + c$$

4. Adjustment of the derivative

Sometimes the integrand may not have the exact derivative of the function u present. In such cases we shall adjust the derivative so that the function u and its exact derivative u^1 are present in the integrand. The following examples shall illustrate this principle.

Example 6

$$x^2 \int x + 1$$

$$\text{Find } \int \frac{x^2}{2x^3 + 3x^2 + 6x + 5} dx$$

Solution

The integrand has one function u
 $= 2x^3 + 3x^2 + 6x + 5$

Then $u^1 = 6x^2 + 6x + 6 = 6(x^2 + x + 1)$

The other function present in the integrand is $(x^2 + x + 1)$. We must multiply this second function by six and divide by six to get u^1 .

$$\text{Thus } \int \frac{x^2 + x + 1}{2x^3 + 3x^2 + 6x + 5} dx = \frac{1}{6} \int \frac{6x^2 + 6x + 6}{2x^3 + 3x^2 + 6x + 5} dx$$

Now the integrand has u and its derivative u^1 . Let
 $u = 2x^3 + 3x^2 + 6x + 5$

$$\frac{du}{dx} = 6x^2 + 6x + 6 = 6(x^2 + x + 1)$$

$$du = (6x^2 + 6x + 6) dx$$

Then the given integral is transformed into

$$\frac{1}{6} \int \frac{du}{u} = \frac{1}{6} \ln u + c$$

$$= \frac{1}{6} \ln(2x^3 + 3x^2 + 6x + 5) + c$$

Example 7

Find $\int x^3 \sqrt{x^4 - 1} dx$

Solution

$$\text{Let } u = x^4 - 1$$

$$\frac{du}{dx} = 4x^3$$

$$du = 4x^3 dx$$

$$\int x^3 \sqrt{x^4 - 1} dx = \frac{1}{4} \int \sqrt{u} du$$

$$\text{Then } \int x^3 \sqrt{x^4 - 1} dx = \frac{1}{4} \int \sqrt{u} du$$

$$= \frac{1}{4} \cdot \frac{2}{3} u^{3/2}$$

$$= \frac{1}{6} (x^4 - 1)^{3/2} + c$$

BY KELVIN EKALIYO, THE NEXT.

$$= \frac{u^{3/2}}{3/2} + C$$

$$= \frac{1}{6} (x^4 - 1)^{3/2} + C$$

Example 8

Find $\int \sec^5 x \tan x \, dx$

Solution

If $u = \sec x$, $\frac{du}{dx} = \sec x \tan x$

Now $\int \sec^5 x \tan x \, dx = \int \sec^4 x \sec x \tan x \, dx$.

Let $u = \sec x$ $du = \sec x \tan x \, dx$

Then $\int \sec^5 x \tan x \, dx = \int u^4 \, du$

$$= \frac{u^5}{5} + C$$

$$= \frac{\sec^5 x}{5} + C$$

Example 9

BY KELVIN EKALIYO, THE NEXT.

Find $\int \sin^3 8x \cos 8x dx$

Let $u = \sin 8x$, $\frac{du}{dx} = 8 \cos 8x$

$$du = 8 \cos 8x dx$$

Then $\int \sin^3 8x \cos 8x dx = \frac{1}{8} \int \sin^3 8x (8 \cos 8x) dx$

$$= \frac{1}{8} \int u^3 du$$

$$= \frac{1}{8} \cdot \frac{u^4}{4}$$

$$= \frac{1}{32} u^4 + C$$

$$= \frac{1}{32} \sin^4 8x + C$$

$$= \frac{\sin^4 8x}{32} + C$$

$$= \frac{\sin^4 8x}{32} + C$$

$$= \frac{\sin^4 8x}{32} + C$$

3.5 Integration of $\tan x$ and $\cot x$ (formulae)

a) $\int \tan x dx = \int \frac{\sin x}{\cos x} dx$

Let $u = \cos x$, $\frac{du}{dx} = -\sin x$

$$du = -\sin x dx$$

$$= \int \frac{-du}{u}$$

BY KELVIN EKALIYO, THE NEXT.

Then $\int \tan x dx = \int \frac{\sin x}{\cos x} dx$

$$= \int \frac{-du}{u}$$

$$= -\ln u + c$$

$$= \ln u^{-1} + c$$

$$= \ln \frac{1}{u} + c$$

$$= \ln \frac{1}{\cos x} + c$$

$$= \ln \frac{1}{\cos x} + c$$

$$= \ln (\sec x) + c$$

we write $\int \tan x dx = \ln |\sec x| + c$

showing that $\ln$ of negative values does not exist

b) Find $\int \cot x dx$

$$\int \cot x dx = \int \frac{\cos x}{\sin x} dx$$

Let $u = \sin x$ so that $du = \cos x dx$

$$\frac{du}{u}$$

Then $\int \cot x dx = \int \frac{du}{u} = \ln u + c$

$$\int \cot x dx = \ln (\sin x) + c$$

BY KELVIN EKALIYO, THE NEXT.

Exercise 3

Find the integrals of the following:

1. $\int (-x^3 + 2x + 1) dx$

$$\int \cos x + e^x$$

$$\int \frac{1}{x^2}$$

BY KELVIN EKALIYO, THE
NEXT.

2. $\int \overline{e^x - \sin x} \, dx$

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

3. $\int e^{x^2} x dx$

4. $\int x (x^2 - 3)^5 dx$

5. $\int \frac{21x}{x^2 - 5} dx$

$\int \frac{\sqrt{2x^3 - 5}}{x-1} dx$

6. $\int \frac{1}{(2x^2 - 1)^5} dx$

7. $\int 4x(1)^5 dx$

8. $\int \frac{e^{-1}}{(\ln x)^2} dx$

9. $\int \frac{1}{x} dx$

10. $\int \tan^5 x \sec^2 x dx$

11. $\int \sec^2 x \tan^2 x dx$

12. $\int \sec^4 x \tan x dx$

Summary

You have learnt the following from this chapter:

1. When the integrand contains a function and its derivative you can integrate using the substitution:

BY KELVIN EKALIYO, THE NEXT.

2. The following are the formulae learnt:

- $\frac{du}{dx}$
- a) $\int \frac{1}{u} dx = \ln u + c$
- $u^n \quad u^{-n+1}$
- b) $\int u^n dx = \frac{u^{n+1}}{n+1} + c$
- c) $\int u^{-n} dx = \frac{u^{-n+1}}{-n+1} + c$
- d) $\int e^u \frac{du}{dx} dx = e^u + c$
- e) $\int (\sin u) \frac{du}{dx} dx = -\cos u + c$
- and similar trigonometric results.
- f) $\int \tan x dx = \ln |\sec x| + c$
- g) $\int \cot x dx = \ln |\sin x| + c$

Further Reading

1. Advanced Calculus
By Watson Fulks
John Wiley & sons
New York Brisbane. Toronto

CHAPTER 4

Integration by parts

In chapter three you have learnt to integrate an expression containing two functions one of which is the derivative of the other. If the two functions in the integrand are not related we use an important technique called integration by parts. We shall establish a formula using the product rule of differentiation and apply the same to integrate product of two functions.

4.2 Objectives of the Chapter:

By the end of this chapter you should be able to

- derive the formula for integration by parts
- apply the formula to integrate product of two functions.

4.3 Integration by parts – Formula

We shall establish the formula

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx \quad (1)$$

Formula (1) is also written as $\int u dv = uv - \int v du$ (2)

In formula (1) the integrand contains the product of two functions namely

u and $\frac{dv}{dx}$

The result (1) will be helpful only when

$\int v \frac{du}{dx}$ is easier than $\int u \frac{dv}{dx}$

Formula for integration by parts

We know that $\frac{d(uv)}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$ (3)

Integrate both sides of (4) with respect to x

Then $u \frac{dv}{dx} = \frac{d(uv)}{dx} - v \frac{du}{dx}$ (4)

$$\int u \frac{dv}{dx} dx = \int \frac{d(uv)}{dx} dx - \int v \frac{du}{dx} dx$$

(5)

$$\text{or } \int u dv = uv - \int v du$$

Note: The functions u and $\frac{dv}{dx}$ can be Logarithmic, Inverse trigonometric, Algebraic, dx

Trigonometric or Exponential. We must arrange the two functions in proper order in the integrand before we proceed to apply the formula. You remember the order using the key word.: LIATE

where L, I, A, T and E represents the functions Logarithmic, Inverse trigonometric, Algebraic, Trigonometric or Exponential respectively.

The following examples illustrate the process of integration by parts.

Example 1

Find $\int e^x x dx$

Solution

The integrand consists of product of two functions e^x (Exponential) and x (Algebraic). Remember the order LIATE

Algebraic function should come first and then only Exponential function must be written

If we do not follow this principle

$$\int v \frac{du}{dx} \text{ will be more complicated than } \int u \frac{dv}{dx} \quad (\text{Remember LIATE !})$$

The required integration is

$$\int x e^x dx \text{ (in proper order) } =$$

$$u = x \quad \frac{du}{dx} = 1$$

$$v = e^x$$

$$\frac{d^2u}{dx^2} = 0$$

$$\frac{d^2u}{dx^2} = 0$$

$$\frac{d^2u}{dx^2} = 0$$

(constant we can put at the end)

Substitution of the results in the formula $\frac{dv}{dx} = \frac{du}{dx} \cdot u$

$$\frac{dv}{dx} = \frac{du}{dx} \cdot u$$

$$\frac{dv}{dx} = e^x \cdot x = x e^x$$

$$\begin{aligned} \text{We have } \int x e^x dx &= x e^x - \int e^x \cdot 1 dx \\ &= x e^x - e^x + c \end{aligned}$$

Example 2

Find $\int x \cos x dx$

Solution

(The order of LIATE is correct. Algebraic first and then comes Trigonometric; otherwise the formula will not be useful. Check yourself). dv

$$\text{Let } u = x \quad \frac{du}{dx} = 1, \quad dv = \cos x dx$$

$$\frac{du}{dx} = 1$$

$$\frac{du}{dx} = 1$$

$$\frac{du}{dx} = 1$$

$$v = \int \cos x dx$$

Using the formula

$$\frac{dv}{dx} = \frac{du}{dx} \cdot u$$

$$\frac{dv}{dx} = \frac{du}{dx} \cdot u$$

$$\int \frac{dv}{dx} \cdot u dx = \int u \cdot \frac{dv}{dx} dx - \int v \cdot \frac{du}{dx} dx$$

$$\begin{aligned} \text{We have } \int x \cos x dx &= x \sin x - \int \sin x \cdot 1 dx \\ &= x \sin x + \cos x + c \end{aligned}$$

BY KELVIN EKALIYO, THE NEXT.

Example 3

Find $\int x^3 \ln x \, dx$

Solution

$\int x^3 \ln x \, dx = \int \ln x \cdot x^3 \, dx$ writing in the order LIATE

Let $u = \ln x$

$$\underline{dv} = x^3$$

$$\underline{du} = \frac{1}{x}$$

$$v = \int x^3 \, dx = \frac{x^4}{4}$$

$$dx = x$$

Substituting in the formula

$$\underline{dv}$$

$$\underline{du}$$

$\int u \, dv = uv - \int v \, du$ we have

$$\int \ln x (x^3) \, dx = \ln x \cdot \frac{x^4}{4} - \int \frac{x^4}{4} \cdot \frac{1}{x} \, dx$$

$$x^4$$

$$x^3$$

$$\int x^3 \ln x \, dx = \frac{x^4}{4} \ln x - \int \frac{x^3}{4} \, dx$$

$$\frac{x^4}{4}$$

$$\frac{x^3}{16}$$

$$= \frac{x^4}{4} \ln x - \frac{x^4}{16} + C$$

Example 4

Find $\int x^2 e^{2x} dx$

Solution

The order of $x^2 e^{2x}$ satisfies "LIATE"

Let $u =$

$$x^2$$

$$du$$

$$=$$

$$2x$$

$$dx$$

Using the formula

$$\frac{dv}{dx} = e^{2x}$$

$$dv = e^{2x} dx$$

$$v = \int e^{2x} dx$$

$$v = \frac{1}{2} e^{2x}$$

$$e^{2x}$$

$$\frac{1}{2} e^{2x}$$

(constant can be put at the end)

$$\frac{dv}{dx} = uv = v \frac{du}{dx} = \frac{1}{2} e^{2x} x^2$$

$$dx = \frac{1}{2} e^{2x} x^2$$

we have $\int x^2 e^{2x} dx = \frac{1}{2} e^{2x} x^2 - \int \frac{1}{2} e^{2x} 2x dx$

$$\frac{1}{2}$$

$$= \frac{1}{2} x^2 e^{2x} - \int x e^{2x} dx$$

(1)

Now we must integrate $\int x e^{2x} dx$

Again let $u = x$ $\frac{dv}{dx} = 2x$ $=$

$$\frac{du}{dx} = 1 \quad v = e^{2x} \quad dx = \frac{1}{2} e^{-2x}$$

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

Using the formula

$\underline{dv}$

$\underline{du}$

$$\iint_U u \, dx \, dy = \iint_U uv \, dx \, dy + \iint_U v \, dx \, dy$$

$$\int x e^{2x} dx = \frac{1}{2} x e^{2x} - \int \frac{1}{2} e^{2x} dx$$

We have

2 2

$$\square \frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} \quad (2)$$

Using the result (2) in (1)

$$\int_{-\infty}^{\infty} \frac{1}{2} e^{-2x} \left(\frac{1}{x} e^{2x} - \frac{1}{e^{2x}} \right) x^2 e^{2x} dx = \frac{1}{24}$$

$$-\frac{1}{2}e^{2x} - \frac{1}{2}x e^{2x} + \frac{1}{4}e^{2x} + C$$

$$1 - 2x(2x^2 - 2x + 1) = e^{-2x}$$

BY KELVIN EKALIYO, THE NEXT.

Example 5Find $\int x^2 \cos x \, dx$ **Solution**

(In this problem Algebraic Function x^2 comes before the Trigonometric function $\cos x$. This satisfies the condition of LIATE).

$$\text{Let } u = x^2 \quad \frac{dv}{dx} = \cos x \quad \frac{du}{dx} = 2x \quad v = \sin x$$

$$\cos x \, dx = \sin x$$

Using the formula

$$\frac{dv}{dx} = uv - v \frac{du}{dx} \text{ we have}$$

$$\int \frac{dv}{dx} u \, dx = \int uv \, dx - \int v \frac{du}{dx} \, dx$$

$$\int \cos x \, dx = \int x^2 \sin x - \int \sin x \cdot 2x \, dx$$

$$= x^2 \sin x - 2 \int x \sin x \, dx \quad (1)$$

BY KELVIN EKALIYO, THE NEXT.Now we must integrate $\int x \sin x \, dx$

Using again integration by parts.

$$\text{Let } u = x \quad \frac{dv}{dx} = \sin x$$

$$\frac{du}{dx} = 1$$

$$v = -\cos x$$

Applying the formula

$$\frac{dv}{dx} = uv - v \frac{du}{dx} \text{ we have}$$

$$\int \int u \, dx \, dx = \int \int dx$$

$$\begin{aligned} \int x \sin x \, dx &= -x \cos x + \int \cos x \, 1 \, dx \\ &= -x \cos x + \sin x \end{aligned} \quad (2)$$

Using the result (2) in (1) we have

$$\begin{aligned} \int x^2 \cos x \, dx &= x^2 \sin x - 2(-x \cos x + \sin x) + c \\ &= x^2 \sin x + 2x \cos x - 2 \sin x + c \end{aligned}$$

Example 6

Find $\int x^3 \ln x \, dx$

Solution

To satisfy the condition for LIATE we write the integral as

$$\int \ln x (x^3) \, dx$$

Let $u = \ln x$

$$\frac{dv}{dx} = x^3$$

$$v = \frac{x^4}{4}$$

$$\frac{du}{dx} = \frac{1}{x}$$

$$\int u \, dv = uv - \int v \, du$$

$$\int \ln x \, x^3 \, dx = \frac{x^4}{4} \ln x - \int \frac{x^4}{4} \cdot \frac{1}{x} \, dx$$

Applying the formula

$$\int u \, dv = uv - \int v \, du \text{ we have}$$

$$\int \int u \, dx \, dx = \int \frac{1}{4} x^4 \, dx$$

$$\frac{1}{4} \int x^4 \, dx = \frac{1}{4} \cdot \frac{x^5}{5} = \frac{x^5}{20}$$

$$\int (\ln x) x^3 \, dx = \frac{x^4}{4} \ln x - \frac{x^4}{4}$$

BY KELVIN EKALIYO, THE NEXT.

1

d

x

x

MWALIMU GREG

BY KELVIN EKALIYO, THE NEXT.

$$\frac{1}{14} x^4 \ln x - \frac{1}{14} \int x^3 dx$$

$$= \frac{1}{14} x^4 \ln x - \frac{1}{14} \cdot \frac{x^4}{4} + c$$

$$= \frac{1}{14} x^4 \ln x - \frac{x^4}{56} + c$$

4.4 Formula for integration of $\ln x$, $x > 0$

$$\int \ln x \, dx = x \ln x - x + c$$

Proof

$$\text{Consider } \int \ln x \, dx = \int \ln x \cdot 1 \, dx$$

$$\text{Let } u = \ln x$$

$$\frac{du}{dx} = \frac{1}{x}$$

$$\text{Then } \frac{du}{dx} = \frac{1}{x}$$

$$\frac{dv}{dx} = 1$$

$$v = \int 1 \, dx = x$$

Using the formula

$$\int u \, dv = uv - \int v \, du$$

$$\int \ln x \, dx = x \ln x - \int \frac{1}{x} \cdot x \, dx$$

$$= x \ln x - \int 1 \, dx$$

$$= x \ln x - x + c$$

Example 7

$$\text{Find } \int (\ln x)^2 \, dx$$

$$\int (\ln x)^2 \, dx = \int (\ln x)^2 \cdot 1 \, dx$$

$$\text{Let } u = (\ln x)^2$$

$$\frac{du}{dx} = \frac{2 \ln x}{x}$$

Solution

BY KELVIN EKALIYO, THE NEXT.

$$\frac{dv}{dx} = \frac{2 \ln x}{x}$$

$$v = \int \frac{2 \ln x}{x} dx = x^2 \ln x - \frac{x^2}{2} + C$$

MWALIMU GREG

Substitution of the results in the formula $\frac{dv}{dx}$

$$\frac{d}{dx} uv = v \frac{du}{dx} + u \frac{dv}{dx}$$

$$\frac{d}{dx} u \frac{dv}{dx} = \frac{d}{dx} \left(u \frac{dv}{dx} \right)$$

$$= \frac{d}{dx} x(\ln x)^2 - \frac{d}{dx} 2x(\ln x) \cdot \frac{1}{x}$$

$$= (\ln x)$$

$$= x(\ln x)^2 - 2 \int (\ln x) dx$$

$$= x(\ln x)^2 - 2(x \ln x - x) + c$$

$$= x \ln x (\ln x - 2) + 2x + c$$

Exercise 4

Find

$$1 \int x e^{2x} dx$$

$$2 \int x \sin x dx$$

$$3 \int x \ln x dx$$

$$4 \int \ln(x-1) dx$$

$$5 \int x \cos 3x dx$$

$$6 \int e^x \sin x dx$$

$$7 \int x^2 \ln x dx$$

$$8 \int \frac{1}{\ln x x^2} dx$$

$$9 \int \ln 2x dx$$

$$10 \int \tan^{-1} x dx$$

$$11 \int x^2 \sin x dx$$

$$12 \int x^{-3} \ln x dx$$

Summary

You have learnt the following from this chapter:

1. Formula for integration by parts

$$\frac{dv}{dx}$$

$$\frac{du}{dx}$$

$$\frac{d}{dx} uv = u \frac{dv}{dx} + v \frac{du}{dx}$$

2. Apply the formula to integrate product of two functions.

BY KELVIN EKALIYO, THE NEXT.

3. The formula $\int \ln x \, dx = x \ln x - x + c$ where $x > 0$.

Further Reading

1. Advanced Calculus Watson
Fulks
John Wiley & Sons
New York. Brisbane. Toronto.

CHAPTER 5

Integration Using Trigonometric Identities.

5.1 Introduction

You have learnt some four important techniques of Integration so far. Now we shall learn in this chapter to integrate expressions using trigonometric identities and substitutions of trigonometric functions. Thus we transform the integrals to one of the basic formulae and then integrate easily.

5.2 Objectives of the chapter.

By the end of this chapter you should be able to: -

- use one of the important techniques of integration namely, integrating trigonometric expressions using trigonometric identities.

5.3 Review of some Important Trigonometric Identities Used in Integration.

- i). $\sin^2 x + \cos^2 x = 1$
- ii). $1 + \tan^2 x = \sec^2 x$
- iii). $1 + \cot^2 x = \operatorname{cosec}^2 x$
- iv). $\sin 2x = 2 \sin x \cos x$
- v). $\cos 2x = \cos^2 x - \sin^2 x = 2 \cos^2 x - 1 = 1 - 2 \sin^2 x$
- vi). $\sin (A + B) = \sin A \cos B + \cos A \sin B$ (1)
- vii). $\sin (A - B) = \sin A \cos B - \cos A \sin B$ Adding (1) and (2) we get the formulae:
 $2 \sin A \cos B = \sin (A + B) + \sin (A - B)$ (3)
Subtracting (2) from (1),
 $2 \cos A \sin B = \sin (A + B) - \sin (A - B)$ (4)
- viii). $\cos (A + B) = \cos A \cos B - \sin A \sin B$ (5)

BY KELVIN EKALIYO, THE NEXT.

$$\text{ix). } \cos(A - B) = \cos A \cos B + \sin A \sin B \text{ adding (5) and (6)} \quad (6)$$

$$2 \cos A \cos B = \cos(A + B) + \cos(A - B) \quad (7)$$

subtracting (6) from (5) we have

$$-2 \sin A \sin B = \cos(A + B) - \cos(A - B) \quad (8)$$

Formulae (3), (4), (7) and (8) are very useful to resolve the product of two trigonometric functions into sum of two functions and integrate the sum of the functions easily. (Anyone of the formulae (3) or (4) is enough).

$$\text{x). } \sin 3x = 3 \sin x - 4 \sin^3 x$$

$$\text{Hence } \sin^3 x = \frac{-\sin x}{4} \square \frac{-\sin 3x}{4}$$

$$\text{Similarly } \sin^3 30x = \frac{-\sin 10x}{4} \square \frac{\sin 30x}{4}$$

$$\text{xi). } \cos 3x = -3 \cos x + 4 \cos^3 x$$

$$\cos^3 x = \frac{-\cos x}{4} \square \frac{\cos 3x}{4}$$

5.4 Integration using the fundamental identities

$$\sin^2 x + \cos^2 x = 1 \quad (1)$$

$$1 + \tan^2 x = \sec^2 x \quad (2)$$

$$1 + \cot^2 x = \operatorname{cosec}^2 x \quad (3)$$

Example 1

$$\sin^2 x$$

Find $\int \sin^2 x \square 1 \square \cos x \, dx$

Solution

BY KELVIN EKALIYO, THE NEXT.

Using the identity $(1 - \sin^2 x) + \cos^2 x = 1$

$$\int (1 - \cos^2 x) dx$$

$$\int (1 - \cos^2 x) dx = \int (1 - \cos^2 x) dx$$

$$= \int (1 - \cos^2 x) dx = \int (1 - \cos^2 x) dx$$

$$= \int (1 - \cos^2 x) dx$$

$$= x - \sin x + c$$

Example 2

Find $\int \tan^2 x dx$

Solution

We use $1 + \tan^2 x = \sec^2 x$

$$\text{Then } \int \tan^2 x dx = \int (\sec^2 x - 1) dx$$

$$= \int \sec^2 x dx - \int 1 dx$$

$$= \tan x - x + c$$

Example 3

$$\cos^2 x$$

Find $\int (1 - \sin^2 x) dx$

Solution

We have the identity $\cos^2 x = 1 - \sin^2 x$

$$\int (1 - \sin^2 x) dx$$

BY KELVIN EKALIYO, THE NEXT.

Then $\int (1 - \sin x) dx = \int (1 - \sin x) dx$

$$= \int (1 - \sin x) dx = \int 1 dx - \int \sin x dx$$

$$= \int 1 dx - \int \sin x dx$$

$$= x - \cos x + c$$

$$= x - \cos x + c$$

Example 4

Find $\int \cot^2 x dx$

Solution

Using the identity $1 + \cot^2 x = \operatorname{cosec}^2 x$, we have $\int \cot^2 x dx = \int (\operatorname{cosec}^2 x - 1) dx$

$$= \int \operatorname{cosec}^2 x dx - \int 1 dx$$

$$= -\cot x - x + c$$

$$= -\cot x - x + c$$

5.5 Integration using the identity Cos

$$\begin{aligned} \cos^2 x &= \cos^2 x - \sin^2 x \\ &= 2 \cos^2 x - 1 \\ &= 1 - 2 \sin^2 x. \end{aligned}$$

Example 5

Find $\int \sin^2 x dx$

Solution

Using the identity $\cos 2x = 1 - 2 \sin^2 x$ or $\sin^2 x = \frac{1 - \cos 2x}{2}$ Using the identity $\cos 2x = 1 - 2 \sin^2 x$

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

BY KELVIN EKALIYO, THE NEXT.

$$\begin{aligned}
 \text{Hence } \int \sin^2 x dx &= \int \frac{1 - \cos 2x}{2} dx \\
 &= \frac{1}{2} \int (1 - \cos 2x) dx \\
 &= \frac{1}{2} \left(x - \frac{1}{2} \sin 2x \right) + c \\
 &= \frac{x}{2} - \frac{\sin 2x}{4} + c
 \end{aligned}$$

Example 6

Find $\int \cos^2 10x dx$

Solution

Using the identity $\cos 2x = 2 \cos^2 x - 1$

$$\cos 20x = 2 \cos^2 10x - 1, \text{ then } \cos^2 10x = \frac{1 + \cos 20x}{2}$$

$$\begin{aligned}
 \text{Then } \int \cos^2 10x dx &= \int \frac{1 + \cos 20x}{2} dx \\
 &= \frac{1}{2} \int (1 + \cos 20x) dx \\
 &= \frac{1}{2} \left(x + \frac{1}{20} \sin 20x \right) + c \\
 &= \frac{x}{2} + \frac{\sin 20x}{40} + c
 \end{aligned}$$

Example 7

$$\int \cos 6x \sin 3x dx$$

Find $\int (\cos 3x \sin 3x) dx$

Solution

BY KELVIN EKALIYO, THE NEXT.

Using the identity $\cos 2x = \cos^2 x - \sin^2 x$
 $\cos 6x = \cos^2 3x - \sin^2 3x$

$$\begin{aligned}
 \text{Hence } \int (\cos 3x \cos x - 6 \sin x \cos 3x) dx &= \int (\cos^2 3x - \sin^2 3x) dx \\
 &= \int (\cos 3x + \sin 3x)(\cos 3x - \sin 3x) dx \\
 &= \int (\cos^2 3x - \sin^2 3x) dx \\
 &= \int \frac{1 + \cos 6x}{2} - \frac{1 - \cos 6x}{2} dx \\
 &= \int \cos 6x dx \\
 &= \frac{1}{6} \sin 6x + c
 \end{aligned}$$

5.6 Integration using the identities,

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B)$$

$$2 \cos A \cos B = \cos (A + B) + \cos (A - B)$$

$$-2 \sin A \sin B = \cos (A+B) - \cos (A - B)$$

These identities convert the product into sums. The following examples illustrate the use of these identities.

Example 8

Find $\int \sin 10x \cos 8x dx$

Solution

Using the identity $2 \sin A \cos B = \sin (A + B) + \sin (A - B)$

1

BY KELVIN EKALIYO, THE NEXT.

We have $\int \sin 10x \cos 8x dx = \frac{1}{2} \int 2 \sin 10x \cos 8x dx$

$$= \frac{1}{2} \int (\sin 18x - \sin 2x) dx$$

$$= \frac{1}{2} \left[-\frac{1}{18} \cos 18x - \frac{1}{2} \cos 2x \right] + c$$

$$= -\frac{1}{36} \cos 18x - \frac{1}{4} \cos 2x + c$$

Example 9

Find $\int \cos 12x \sin 3x dx$

Solution

$$\int \cos 12x \sin 3x dx = \int \sin 3x \cos 12x dx$$

$$= \frac{1}{2} \int 2 \sin 3x \cos 12x dx$$

$$= \frac{1}{2} \int (\sin 15x - \sin(-9x)) dx$$

$$= \frac{1}{2} \int (\sin 15x + \sin 9x) dx$$

$$\frac{1}{2} \left[-\frac{1}{15} \cos 15x - \frac{1}{9} \cos 9x \right] + c$$

BY KELVIN EKALIYO, THE NEXT.

$$\begin{aligned}
 &= \frac{1}{2} (\cos 15x - \cos 9x) + c \\
 &= \frac{1}{30} \cos 15x - \frac{1}{18} \cos 9x + c
 \end{aligned}$$

Example 10

Find $\int \cos 5x \cos 7x dx$

Solution

$$\begin{aligned}
 \int \cos 5x \cos 7x dx &= 2 \int \frac{1}{2} \cos 5x \cos 7x dx \\
 &= 2 \int \frac{1}{2} [\cos(12x) + \cos(2x)] dx \\
 &= 2 \int \frac{1}{2} \cos 12x + \cos 2x dx \\
 &= \int \cos 12x + \cos 2x dx \\
 &= \frac{1}{12} \sin 12x + \frac{1}{2} \sin 2x + c \\
 &= \frac{1}{24} \sin 12x + \frac{1}{4} \sin 2x + c
 \end{aligned}$$

Example 11

Find $\int \sin 8x \sin 15x dx$

BY KELVIN EKALIYO, THE NEXT.

Solution

$$\int \sin 8x \sin 15x dx = \frac{1}{2} \int (\cos 7x - \cos 23x) dx$$

dx

$$= \frac{1}{2} \left(\frac{\sin 7x}{7} - \frac{\sin 23x}{23} \right) + C$$

$$= \frac{1}{2} \left(\frac{\sin 7x}{7} - \frac{\sin 23x}{23} \right) + C$$

2

BY KELVIN EKALIYO, THE NEXT.

$$\int (23x - \cos 7x) dx$$

MWALIMU GREG

$$\begin{aligned} &= \frac{1}{46} \sin 23x - \frac{1}{14} \sin 7x + c \\ &= \frac{1}{46} \sin 23x - \frac{1}{14} \sin 7x + c \end{aligned}$$

5.7 Integration using the identity

$$\begin{aligned} \sin 3x &= 3 \sin x - 4 \sin^3 x \\ \cos 3x &= -3 \cos x + 4 \cos^3 x \end{aligned}$$

Example 12

Find $\int \sin^3 6x \, dx$

Solution

Using the identity, $\sin 3x = 3 \sin x - 4 \sin^3 x$, $\sin 18x = 3 \sin 6x - 4 \sin^3 6x$

$$\begin{aligned} \sin^3 6x &= \frac{1}{4} \sin 6x - \frac{1}{4} \sin 18x \\ \int \sin^3 6x \, dx &= \frac{1}{4} \int \sin 6x \, dx - \frac{1}{4} \int \sin 18x \, dx \\ &= \frac{1}{4} \left(-\frac{1}{6} \cos 6x \right) - \frac{1}{4} \left(-\frac{1}{18} \cos 18x \right) + c \end{aligned}$$

Hence $\int \sin^3 6x \, dx = -\frac{1}{24} \cos 6x + \frac{1}{72} \cos 18x + c$

$$\begin{aligned} &= -\frac{1}{24} \cos 6x + \frac{1}{72} \cos 18x + c \\ &= -\frac{1}{24} \cos 6x + \frac{1}{72} \cos 18x + c \end{aligned}$$

Example 13

Find $\int \cos^3 9x \, dx$

Solution

Using the identity $\cos 3x = 4\cos^3 x - 3\cos x$
 $\cos 27x = 4\cos^3 9x - 3\cos 9x$

$$\text{Hence } \cos^3 9x = \frac{1}{4} \cos 27x + \frac{3}{4} \cos 9x$$

$$\int \cos^3 9x \, dx = \frac{1}{4} \int \cos 27x \, dx + \frac{3}{4} \int \cos 9x \, dx$$

$$\begin{aligned} &= \frac{1}{4} \left(\frac{1}{27} \sin 27x \right) + \frac{3}{4} \left(\frac{1}{9} \sin 9x \right) + c \\ &= \frac{1}{108} \sin 27x + \frac{1}{12} \sin 9x + c \end{aligned}$$

5.8 Formulae for integration when the integrand contains quadratic expressions in the denominator.

$$1. \int \frac{1}{x^2 + a^2} \, dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$$

$$2. \int \frac{1}{x^2 - a^2} \, dx = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c$$

$$3. \int \frac{1}{x^2 + a^2} \, dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$$

$$\begin{aligned}
 4. \quad & \frac{dx}{\sqrt{a^2 - x^2}} = \sinh^{-1} \frac{x}{a} \quad \text{if } x < a \\
 5. \quad & \frac{dx}{\sqrt{x^2 - a^2}} = \cosh^{-1} \frac{x}{a} \quad \text{if } x > a \\
 & = \cosh^{-1} \frac{x}{a} \quad \text{if } -x > a \\
 & \frac{dx}{\sqrt{x^2 - a^2}} = \sin^{-1} \frac{x}{a}
 \end{aligned}$$

Example 14 $\frac{dx}{\sqrt{x^2 - a^2}} = \frac{1}{a} \sinh^{-1} \frac{x}{a}$

Prove that $\frac{dx}{\sqrt{x^2 - a^2}} = \frac{1}{a} \sinh^{-1} \frac{x}{a}$

Proof Let $x = a \tan \theta$ so that $\theta =$

$\tan^{-1} \frac{x}{a}$

a

Then $x^2 + a^2 = a^2 \tan^2 \theta + a^2 = a^2 (1 + \tan^2 \theta) = a^2 \sec^2 \theta$
 $dx = d(a \tan \theta) = a \sec^2 \theta d\theta$

$$\begin{aligned}
 \frac{dx}{\sqrt{x^2 - a^2}} &= \frac{a \sec^2 \theta d\theta}{a \sec \theta} \\
 \text{Hence } \frac{dx}{\sqrt{x^2 - a^2}} &= \frac{1}{a} \sinh^{-1} \frac{x}{a}
 \end{aligned}$$

$$\begin{aligned}
 &= a \int \frac{1}{1-x^2} dx \\
 &= \frac{1}{2} a \int \frac{1}{1-x^2} dx \\
 &= \frac{1}{2} a \int \frac{1}{1-x^2} dx \\
 &= \frac{1}{2} a \int \frac{1}{1-x^2} dx
 \end{aligned}$$

Example 13

Prove that $\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} + C$

$$x = a \sin \theta \quad \Rightarrow \quad dx = a \cos \theta d\theta$$

Proof

Let $x = a \sin \theta$ so that $dx = a \cos \theta d\theta$

$$\begin{aligned}
 \text{and } \sqrt{a^2 - x^2} &= \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a \cos \theta \\
 &= a \cos \theta
 \end{aligned}$$

$$\begin{aligned}
 \text{Hence } \int \frac{1}{\sqrt{a^2 - x^2}} dx &= \int \frac{a \cos \theta d\theta}{a \cos \theta} \\
 &= \int d\theta \\
 &= \theta + C
 \end{aligned}$$

$$\begin{aligned}
 &= \theta + C \\
 &= \sin^{-1} \frac{x}{a} + C \\
 &\text{since } x = a \sin \theta = \sin^{-1} \frac{x}{a}
 \end{aligned}$$

Example 14 $\frac{dx}{x^2}$

Find $\int \frac{dx}{25x^2 - 4}$

Solution

$$\frac{dx}{25x^2 - 4}$$

$$\int \frac{dx}{25x^2 - 4} = \int \frac{dx}{25 \left(x^2 - \frac{4}{25} \right)}$$

$$= \frac{1}{25} \int \frac{dx}{x^2 - \frac{4}{25}}$$

$$= \frac{1}{25} \int \frac{dx}{x^2 - \left(\frac{2}{5} \right)^2}$$

$$= \frac{1}{25} \int \frac{dx}{x^2 - \left(\frac{2}{5} \right)^2}$$

$$= \frac{1}{25} \int \frac{dx}{x^2 - \left(\frac{2}{5} \right)^2}$$

$$= \frac{1}{25} \tan^{-1} \frac{x - \frac{2}{5}}{\frac{2}{5}} + c$$

Example 15

dx

Find $\int \frac{dx}{x^2 - 3x + 2}$

Solution

$$x^2 - 3x + 2 = x^2 - 3x + a^2 - a^2 - 2$$

$$= (x - a)^2 - a^2 + 2 = x^2$$

$$- 2ax + a^2 - a^2 + 2$$

$$\frac{3}{2}$$

Equating coefficient of x on both sides $2a = 3$ or $a =$

$$\text{Hence } x^2 - 3x + 2 = \frac{3}{2}x - \frac{1}{2}$$

$$= \frac{3}{2}x - \frac{1}{2}$$

dx

dx

$$\text{Then } \int (x^2 - 3x + 2) dx = \int \left(\frac{3}{2}x - \frac{1}{2} \right) dx$$

$$= \frac{3}{4}x^2 - \frac{1}{2}x + c$$

$$= \frac{3}{4}x^2 - \frac{1}{2}x + c$$

$$= \frac{3}{4}x^2 - \frac{1}{2}x + c$$

$$= \frac{3}{4}x^2 - \frac{1}{2}x + c$$

$\ln$

$$\text{using } \int \frac{1}{x} dx = \ln|x| + c$$

$$c$$

c

c

dx

$$\text{Hence } \int (x^2 - 3x + 2) dx = \frac{1}{3}x^3 - \frac{3}{2}x^2 + 2x + c$$

dx

BY KELVIN EKALIYO, THE NEXT.

Note: since $x^2 - 3x + 2 = (x - 2)(x - 1)$, we can resolve $\frac{2}{x^2 - 3x + 2}$

into partial fraction and then integrate to get the same answer. Try this method yourself.

Example 16

Find $\int \frac{2}{\sqrt{2x^2 + 5x + 1}} dx$

Solution

$$\begin{aligned} \text{Let } 2x^2 + 5x + 1 &= 2\left(x^2 + \frac{5}{2}x + \frac{1}{2}\right) \\ &= 2\left[\left(x + \frac{5}{4}\right)^2 - \left(\frac{5}{4}\right)^2 + \frac{1}{2}\right] = \\ &= 2\left[x^2 + 2ax + a^2 - a^2 + 1\right] \end{aligned}$$

Comparing the coefficient of x on both sides $4a = 5$ or $a = \frac{5}{4}$.

$$\text{Hence } 2x^2 + 5x + 1 = 2\left[\left(x + \frac{5}{4}\right)^2 - \left(\frac{5}{4}\right)^2 + \frac{1}{2}\right]$$

$$= 2\left[x^2 + 2\left(\frac{5}{4}\right)x + \left(\frac{5}{4}\right)^2 - \left(\frac{5}{4}\right)^2 + \frac{1}{2}\right]$$

$$= 2\left[x^2 + \frac{5}{2}x + \frac{25}{16} - \frac{25}{16} + \frac{8}{16}\right]$$

$$= 2\left[x^2 + \frac{5}{2}x + \frac{8}{16}\right]$$

$$= 2\left[x^2 + \frac{5}{2}x + \frac{1}{2}\right]$$

$$= 2 \int x \sqrt{4x^2 - 5} \, dx$$

$$= \frac{4x^2 - 5}{2} \sqrt{4x^2 - 5} - \int \frac{4x}{2} \sqrt{4x^2 - 5} \, dx$$

$$= \frac{4x^2 - 5}{2} \sqrt{4x^2 - 5} - \int 2x \sqrt{4x^2 - 5} \, dx$$

$$= \frac{4x^2 - 5}{2} \sqrt{4x^2 - 5} - \frac{2}{3} (4x^2 - 5)^{3/2} + C$$

$$= \frac{4x^2 - 5}{2} \sqrt{4x^2 - 5} - \frac{2}{3} (4x^2 - 5)^{3/2} + C$$

$$\frac{dx}{\sqrt{4x^2 - 5}} = \frac{1}{\sqrt{4x^2 - 5}} dx$$

$$\int \frac{dx}{\sqrt{4x^2 - 5}} = \int \frac{dx}{\sqrt{4x^2 - 5}} = \frac{1}{2} \int \frac{dx}{\sqrt{x^2 - \frac{5}{4}}}$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C \quad \text{if } x > \frac{\sqrt{5}}{2}$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C \quad \text{if } x > \frac{\sqrt{5}}{2}$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C$$

$$= \frac{1}{2} \ln \left| x + \sqrt{x^2 - \frac{5}{4}} \right| + C \quad \text{if } -x > \frac{\sqrt{5}}{2} \text{ or } x < -\frac{\sqrt{5}}{2}.$$

Exercise 5

Find the integrals of the following

1. $\int \sin^2 x \, dx$
2. $\int (1 - \cos x) \, dx$
3. $\int \cot^2 x \, dx$
4. $\int \cos^2 x \, dx$
5. $\int \cos^2 5x \, dx$
6. $\int \frac{\cos 8x}{\cos 4x - \sin 4x} \, dx$
7. $\int (\cos ax - \sin ax) \, dx$
8. $\int \cos 10x \sin 4x \, dx$
9. $\int \cos 4x \sin 9x \, dx$
10. $\int 5x^2 - 9x - \frac{8}{30} \, dx$
11. $\int \sin 5x \cos 7x \, dx$
12. $\int \cos 8x \cos 5x \, dx$
13. $\int \cos 3x \cos 11x \, dx$
14. $\int \sin 4x \sin 10x \, dx$
15. $\int \sin 12x \sin 5x \, dx$
16. $\int \sin^3 4x \sin dx$
17. $\int \sin^3 5x \, dx$
18. $\int \cos^3 5x \, dx$
19. $\int \cos^3 5x \, dx$
20. $\int \cos^3 5x \, dx$
21. $\int \cos^3 5x \, dx$
22. $\int \cos^3 5x \, dx$
23. $\int \cos^3 5x \, dx$
24. $\int \cos^3 5x \, dx$
25. $\int \cos^3 5x \, dx$
26. $\int \cos^3 5x \, dx$
27. $\int \cos^3 5x \, dx$
28. $\int \cos^3 5x \, dx$
29. $\int \cos^3 5x \, dx$

$$\int 8x^3 dx$$

$$\int x^2 - 16 dx$$

$$\int x^2 - 16 dx$$

$$\int x^2 - 25 dx$$

$$\int \sqrt{\quad} dx$$

$$\int \sqrt{x^2 - 25} dx$$

$$\int \frac{4 - x^2}{dx}$$

$$\int \sqrt{16 - x^2} dx$$

$$\int 3x^2 - 5x + 1 dx$$

Summary

You have learnt the following from this chapter:

- The technique of integration of trigonometric expressions using the following trigonometric identities.

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\int \frac{1}{4x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \frac{1}{x^2} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{9x^2 - 6x + 1} dx$$

$$\int \sqrt{25 - x^2} dx$$

$$2. \quad \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$3. \quad \tan^2 x = \sec^2 x - 1$$

$$4. \quad \cot^2 x = \operatorname{cosec}^2 x - 1$$

$$5. \quad 2 \sin A \cos B = \sin (A + B) + \sin (A - B)$$

$$6. \quad 2 \cos A \cos B = \cos (A + B) + \cos (A - B)$$

$$7. \quad -2 \sin A \sin B = \cos (A + B) - \cos (A - B)$$

$$8. \quad \sin 3x = 3 \sin x - 4 \sin^3 x$$

$$9. \quad \cos 3x = 4 \cos^3 x - 3 \cos x$$

ii). The technique of integration using the following formulae:

$$1. \quad \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$$

$$\int \frac{x}{x^2 + a^2} dx = \frac{1}{2} \ln |x^2 + a^2| + c$$

$$2. \quad \int \frac{1}{\sqrt{x^2 + a^2}} dx = \sinh^{-1} \frac{x}{a} + c$$

$$3. \quad \int \frac{1}{\sqrt{x^2 - a^2}} dx = \begin{cases} \cosh^{-1} \frac{x}{a} & \text{if } x > a \\ -\cosh^{-1} \frac{x}{a} & \text{if } x < -a \end{cases}$$

5. $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C$

Further Reading

Additional pure mathematics
 Harwood Clarke Heinemann
 Educational Books Ltd.
 London.

MWALIMU GREG

CHAPTER 6

Integration Using Partial Fractions

6.1 Introduction

Some algebraic expressions cannot be integrated directly. An important technique which we use for such expressions is to split it into partial fractions that can be integrated easily using the basic formulae.

6.2 Objectives of the chapter

By the end of this chapter you should be able to split an algebraic expression into partial fractions and integrate.

6.3 Expressions with only linear factors in the denominator

$$\frac{f(x)}{f(x)}$$

In this section we shall see $\int F(x) dx$ where $F(x)$ is a proper fraction (degree of x in f is less than that in F) and $F(x)$ has only non-repeated linear factors. The technique of integration of such integral is illustrated in the following examples:

Example 1

$$\text{Find } \int \frac{1}{(x-3)(x-4)} dx \quad (1)$$

Solution

$$\text{We write } \frac{1}{(x-3)(x-4)} = \frac{A}{x-3} + \frac{B}{x-4}$$

$$= \frac{A(x-4) + B(x-3)}{(x-3)(x-4)}$$

$$\text{Then } x-11 = A(x-4) + B(x+3) \quad (2)$$

We can put special values for x on both sides of (2)

Let $x=4$, then

$$4-11 = A(0) + B(7)$$

$$-7 = 7B, \quad \text{and } B=-1$$

Let $x = -3$, then (2) becomes

$$-3 - 11 = A(-3 - 4) + B(-3 + 3)$$

$$-14 = -7A \text{ and } A = 2$$

$$x + 11 = \frac{2}{x + 3} + \frac{1}{x - 4}$$

Hence
$$\frac{1}{(x + 3)(x - 4)} = \frac{2}{(x + 3)(x - 4)} - \frac{1}{(x + 3)(x - 4)}$$

Then
$$\int \frac{1}{(x + 3)(x - 4)} dx = \int \frac{2}{x + 3} dx - \int \frac{1}{x - 4} dx$$

We split the integrand into two partial fractions each can be integrated using the extension of the basic formulae for $(ax + b)$

$$\begin{aligned} \text{Then } \int \frac{1}{(x + 3)(x - 4)} dx &= 2 \ln(x + 3) - \ln(x - 4) + A \\ &= \ln(x + 3)^2 - \ln(x - 4) + A \\ &= \ln \frac{(x + 3)^2}{x - 4} + A \end{aligned}$$

6.4 Integration of expression with repeated linear factors in the denominator.

Consider
$$\int \frac{1}{(x + 2)(x + 1)(x - 1)} dx \quad (1)$$

The integrand is a proper fraction and with repeated linear factors It is written as

$$\begin{aligned} \frac{1}{(x + 2)(x + 1)(x - 1)} &= \frac{A}{x + 2} + \frac{B}{x + 1} + \frac{C}{(x - 1)^2} \\ &= \frac{A(x + 1)^2 + B(x + 2)(x + 1) + C(x + 2)}{(x + 2)(x + 1)(x - 1)} \end{aligned}$$

Then
$$1 = A(x - 1)^2 + B(x + 2)(x - 1) + C(x + 2) \quad (2)$$

Let us put $x = 1$ on both sides of (1) so that the terms containing A and B will become zero.

$$1$$

Then $1 = A(0) + B(0) + C(1+2)$, Hence $C =$

$\frac{1}{3}$

Similarly put $x = -2$ on both sides of (1) so that the terms containing B and C become zero.

Then $1 = A(-2 - 1)^2 + B(0) + C(0)$, Hence $A = \frac{1}{9}$.

To get the value of B either we can give one more value other than 1 and -2 for x or else we can compare the coefficients of x^2 on both sides of (2)

$$0 = A + B$$

$$1 = \frac{1}{9} + B$$

$$\frac{0}{9} = \frac{1}{9} + B \text{ then } B = -\frac{1}{9}$$

Hence the integral (1) becomes

$$\begin{aligned} \int \frac{1}{(x-2)(x-1)^2} dx &= \int \frac{1}{9(x-2)} dx - \int \frac{1}{9(x-1)} dx + \int \frac{1}{3(x-1)^2} dx \\ &= \frac{1}{9} \ln(x-2) - \frac{1}{9} \ln(x-1) - \frac{1}{3(x-1)} + C \\ &= \frac{1}{9} \ln \frac{x-2}{x-1} - \frac{1}{3(x-1)} + C \end{aligned}$$

6.5 Integration of expressions with quadratic factor in the denominator – proper fraction.

Example 2

Consider $\int \frac{3x+7}{(x-1)(x^2+1)} dx$ (1)

$$\int \frac{3x+7}{(x-1)(x^2+1)}$$

The integrand is a proper fraction and has a quadratic factor. It is written as

$$\frac{3x+7}{(x-1)(x^2+1)} = A + \frac{Bx+C}{x^2+1}$$

$$\frac{3x + 7}{(x - 1)(x^2 + 1)} = \frac{A(x^2 + 1) + (Bx + C)(x - 1)}{(x - 1)(x^2 + 1)}$$

Then $(3x + 7) = A(x^2 + 1) + (Bx + C)(x - 1)$ (2)

We can put $x = 1$ on both sides of (2) to get 3

$$(1) + 7 = A(1 + 1) + (B + C)(0)$$

$$10 = 2A \text{ or } A = 5$$

Putting $x = 0$, $7 = A - C$, $C = -2$

Equating coefficient of x^2 , $0 = A + B$ or $B = -A = -5$

Then $\frac{3x + 7}{(x - 1)(x^2 + 1)} = \frac{5}{x^2 + 1} - \frac{5x + 2}{x - 1}$

Hence $\int \frac{3x + 7}{(x - 1)(x^2 + 1)} dx = 5 \int \frac{dx}{x^2 + 1} - \int \frac{5x + 2}{x - 1} dx$

$$= 5 \tan^{-1} x + C - 5 \ln |x - 1| - \frac{5}{2} \ln |x^2 + 1|$$

6.6 Integrand is an improper fraction

If the degree of x in the Numerator is the same or greater than that of x in the denominator the Algebraic Fraction is called an improper fraction.

If the integrand $\frac{f(x)}{F(x)}$ is an improper fraction we shall divide $f(x)$ by $F(x)$

and the remainder is resolved into partial fractions. This principle is illustrated from the following example:

Example $\int \frac{3x^2 - 2x + 7}{(x-2)(x-1)} dx$

Find $\int \frac{3x^2 - 2x + 7}{(x-2)(x-1)} dx$

Solution

The integrand is an improper algebraic fraction. Hence using long division we find that the quotient is 3 and the remainder is $(x - 1)$

$$\begin{array}{r} 3x^2 - 2x + 7 \\ (x-2)(x-1) \overline{) 3x^2 - 2x + 7} \\ \underline{3x^2 - 5x + 2} \\ 3x - 5 \end{array}$$

Note: sometimes instead of long division it is easier to proceed as follows:

$$\frac{3x^2 - 2x + 7}{(x-2)(x-1)} = \frac{3(x^2 - x + 2) + x - 1}{(x-2)(x-1)}$$

$$\begin{array}{r} = 3 + \frac{x-1}{(x-2)(x-1)} \\ = 3 + \frac{1}{x-2} \end{array}$$

Now we can resolve $\frac{1}{(x-2)(x-1)}$ into partial fractions.

$$\frac{1}{(x-2)(x-1)} = \frac{A}{x-2} + \frac{B}{x-1}$$

$$\frac{A(x-1) + B(x-2)}{(x-2)(x-1)}$$

$$(x-1) = A(x+1) + B(x-2) \quad (2)$$

Putting $x = 2$ on both sides of (2) $A = \frac{1}{3}$.

Putting $x = -1$ on both sides of (2), we have $-2 = B(-3)$ or $\frac{2}{3} = B$

Then $\frac{1}{(x-2)(x+1)} = \frac{1}{3(x-2)} + \frac{2}{3(x+1)}$

Hence the integral (1), is written as

$$\int \frac{1}{(x-2)(x+1)} dx = \int \left(\frac{1}{3(x-2)} + \frac{2}{3(x+1)} \right) dx$$

□

□

$$= \frac{1}{3} \ln|x-2| + \frac{2}{3} \ln|x+1| + C$$

Exercise 6

Find

$$\int \frac{x-11}{(x-3)(x-4)} dx$$

1. $\int \frac{x}{(x-3)(x-4)} dx$

x

$$2. \int (25 - x^2) dx$$

$$\frac{3x^2 - 21x + 24}{x^2 - 1}$$

$$3. \int \frac{1}{(x-1)(x-2)(x-3)} dx$$

$$\frac{5x^2 - 10x + 11}{x^2 - 4}$$

$$4. \int \frac{1}{(x-3)(x^2-4)} dx$$

$$5. \int (xx^2 - 2) x^2 dx$$

$$6. \int \frac{5x + 4}{(x-1)(x-2)^2} dx$$

$$7. \int \frac{3x + 7}{x(x-2)(x-1)} dx$$

$$\frac{2x^3 - x + 1}{(x-3)(x^2-1)}$$

$$8. \int \frac{1}{(x-3)(x^2-1)} dx$$

$$\frac{x^3 - 2x^2 - 10x + 9}{(x^2 - 9)}$$

$$9. \int \frac{1}{(x^2 - 9)} dx$$

$$\frac{(x-2)}{(x^3-1)}$$

$$10. \int \frac{1}{(x^3-1)} dx$$

Hint: $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

$$\frac{(x^3 - 3)}{x^3 - 8} dx \quad a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

$$\frac{2x - 1}{x^3 - 1} dx$$

$$12. \int \frac{1}{x^3 - 1} dx$$

Summary

You have learnt the following from this chapter:

i). Splitting a fraction into partial fractions.

2x

$$\text{Type I: } \frac{1}{(x-a)(x-b)} = \frac{A}{(x-a)} + \frac{B}{(x-b)}$$

$$\text{Type II: } \frac{ax+b}{(x-a)(x-b)(x-c)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$$

$$\text{Type III: } \frac{ax+b}{(x-a)(px^2+qx+r)} = \frac{A}{x-a} + \frac{Bx+C}{px^2+qx+r}$$

Type IV: For improper algebraic fraction, we must divide and consider only the remainder for partial fraction.

ii). After partial fractions, integrand is transformed into the form of basic formula which can be integrated.

Further Reading

1. Additional pure mathematics
By L. Harwood Clarke.
Heinemann Educational Books Ltd London.

CHAPTER 7

Definite Integral and the Fundamental Theorem of Calculus

7.1 Introduction

You have learnt the integral of functions using seven techniques. In this chapter we shall learn the value of the integral between two limits for the variable. Such an integral is called a **Definite Integral**. The definite integral is evaluated using the

fundamental theorem of the integral calculus namely $\int_a^b f(x)dx = G(b) - G(a)$ where

$$\frac{d}{dx} \int_a^x f(x)dx = f(x)$$

7.2 Objectives of the Chapter

By the end of this chapter you should be able to

- define the definite integral
- state the fundamental Theorem of the Integral calculus & apply this theorem to evaluate the definite Integrals.

7.3. Definite Integral

Suppose that $f(x)$ is integrable in an interval $[a, b]$. Draw the curve $y = f(x)$




Figure 1

Divide the interval ab into n equal parts at $a(=a_0), a_1, a_2, \dots, a_n = b$

$$b - a$$

The width of each interval $h = \frac{b-a}{n}$ consider a point x_r in the interval

$$a_{r-1} \leq x \leq a_r$$

The integral $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n f(x_r) h$ is defined as the definite

integral of $f(x)$ in the interval $a \leq x \leq b$

□

Second definition of the definite Integral

If $f(x)$ is continuous in the interval $a \leq x \leq b$ we define the area enclosed by the curve $y = f(x)$, the ordinates $x = a, x = b$, and the x axis as the definite integral of b

$$\int_a^b f(x) dx$$

7.4 Some important properties of definite Integral:

If $f(x)$ is continuous in $a \leq x \leq b$

$$1. \int_a^a f(x) dx = 0$$

$$2. \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

3. If f is integrable in $[a, b]$ and c is a point between a and b

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

4. If $f(x)$ is integrable in $[a, b]$ and c is any constant then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

5. If $f(x)$ is integrable in $[a, b]$ and m is any constant then

$$\int_a^b mf(x) dx = m \int_a^b f(x) dx$$

$$\int_a^b f(x) dx = \int_a^b f(x) dx$$

$$\int_a^b f(x) dx = \int_a^b f(x) dx$$

$$\int_a^b f(x) dx = \int_a^b f(x) dx$$

$$\int_a^b f(x) dx = \lim_{m \rightarrow \infty} \sum_{i=1}^m f(x_i) \Delta x$$

7.5 Definition of an Indefinite Integral [or primitive of f(x)]

Suppose that $f(x)$ is integrable in the interval $[a, b]$ then

— $\int_a^x f(t) dt = F(x) - F(a)$ at each point x in the interval.

A function $F(x)$ with the property that $F'(x) = f(x)$ at every point x of the interval

$\int_a^x f(t) dt$ is called an indefinite integral or primitive of $f(x)$

Thus for a continuous function $f(x)$ we have $F(x) = \int_a^x f(t) dt$

Consider $\int_a^b f(x) dx$ if $\int_a^b f(x) dx = G(b) - G(a)$, we say in short, $G(x)$ is the

indefinite integral of $f(x)$

7.6 The fundamental Theorem of the integral calculus

If $f(x)$ is continuous in $[a, b]$ and G is a primitive (indefinite integral) of $f(x)$ then

$$\int_a^b f(x) dx = G(b) - G(a)$$

This is called the fundamental theorem of the integral calculus

Proof

Let F be given by

$$F(x) = \int_a^x f(t) dt$$

$$\text{Then } F(x) = G(x) + c$$

$$\text{Since } F(a) = \int_a^a f(t) dt = 0 \text{ we have}$$

$$F(a) = 0 = G(a) + c$$

$$c = -G(a)$$

$$\text{Hence } F(x) = G(x) - G(a)$$

$$\text{or } F(b) = \int_a^b f(x) dx = G(b) - G(a) \text{ where } G \text{ is the primitive of } f(x).$$

This is the Fundamental theorem of the Integral calculus. G

$(b) - G(a)$ is usually written as $\int_a^b f(x) dx$

7.7 Evaluation of definite integrals

Using the fundamental theorem of integral calculus we can evaluate definite integrals. The following examples will illustrate the evaluation of definite integrals.

Example 1

Evaluate $\int_2^3 (x^2 + 2x) dx$

Solution

$$\begin{aligned} \int_2^3 (x^2 + 2x) dx &= \left[\frac{x^3}{3} + x^2 \right]_2^3 \\ &= \left(\frac{3^3}{3} + 3^2 \right) - \left(\frac{2^3}{3} + 2^2 \right) \\ &= \left(\frac{27}{3} + 9 \right) - \left(\frac{8}{3} + 4 \right) \\ &= 9 + 9 - \frac{8}{3} - 4 \\ &= 18 - \frac{8}{3} \\ &= \frac{34}{3} \end{aligned}$$

Example 2

Evaluate $\int_0^{\frac{\pi}{2}} \cos^2 x dx$

Solution

$$\int_0^{\pi/2} \cos x \, dx = \left[\sin x \right]_0^{\pi/2} = \sin \frac{\pi}{2} - \sin 0 = 1 - 0 = 1$$

$$\int_0^{\pi/2} \cos^2 x \, dx = \int_0^{\pi/2} \frac{1 + \cos 2x}{2} \, dx$$

$$= \frac{1}{2} \int_0^{\pi/2} (1 + \cos 2x) \, dx$$

$$= \frac{1}{2} \left[x + \frac{\sin 2x}{2} \right]_0^{\pi/2}$$

$$= \frac{1}{2} \left[\frac{\pi}{2} + \frac{\sin \pi}{2} - \left(0 + \frac{\sin 0}{2} \right) \right] = \frac{1}{2} \left[\frac{\pi}{2} + 0 - 0 \right] = \frac{\pi}{4}$$

$$\int_0^{\pi/2} \cos^2 x \, dx = \frac{\pi}{4}$$

$$\int_0^{\pi/2} \sin^2 x \, dx = \frac{\pi}{4}$$

$$\int_0^{\pi/2} \sin^2 x \, dx = \frac{\pi}{4}$$

$$\int_0^{\pi/2} \sin^2 x \, dx = \frac{\pi}{4}$$

Example 3

$$\int_0^{\pi/2} \cos x \, dx$$

Evaluate $\int_0^{\pi/2} \sin x \, dx$

Solution

$\cos x$ is the derivative of $1 + \sin x$ $\frac{du}{dx} = \cos x$

Put $u = 1 + \sin x$, $\frac{du}{dx} = \cos x$

$$du = \cos x \, dx$$

$$\int_0^{\pi/2} \cos x \, dx = \int_0^{\pi/2} du$$

$\int_0^1 \ln(1 - \sin x) dx = \frac{1}{2} \ln 2$

Example 4
 Evaluate $\int_0^3 e^{2x} dx$

Solution
 $\int_0^3 e^{2x} dx = \left[\frac{1}{2} e^{2x} \right]_0^3 = \frac{1}{2} (e^6 - 1)$

□ □ □

10

$$\frac{1}{2} - \frac{1}{2}$$

$$1 - 6 + 1 = -4$$

$$\frac{1}{2} e$$

Example 5

$$\frac{1}{2}$$

Evaluate $\int_0^1 x \sin x dx$

Solution

Using integration by parts

$$\int_0^1 x \sin x dx = \left[-\frac{1}{2} x^2 \right]_0^1$$

$$\int_0^1 x \sin x dx = \left[-\frac{1}{2} x^2 \right]_0^1 + \int_0^1 \cos x dx$$

$$\begin{aligned} &= \left[-\frac{1}{2} x^2 + \sin x \right]_0^1 \\ &= (0 + 1) - (0 + 0) \\ &= 1 \end{aligned}$$

Example 6

$$\frac{1}{2}$$

Example 1
Evaluate $\int_0^1 \sin 8x \sin 3x dx$

Solution

$$\int_0^1 \sin 8x \sin 3x dx = \frac{1}{2} \int_0^1 (\cos 5x - \cos 11x) dx$$

$$\int_0^1 \sin 8x \sin 3x dx = \frac{1}{2} \left[\frac{\sin 5x}{5} - \frac{\sin 11x}{11} \right]_0^1$$

$$= \frac{1}{2} \left(\frac{\sin 5}{5} - \frac{\sin 11}{11} \right)$$

$$\int_0^{\pi/2} \cos 11x \cos 5x dx$$

$$= \frac{1}{2} \int_0^{\pi/2} (\cos 16x + \cos 6x) dx$$

$$= \frac{1}{2} \left[\frac{\sin 16x}{16} + \frac{\sin 6x}{6} \right]_0^{\pi/2} = 0$$

Exercise 7

Evaluate the following

$$\frac{x^2 - 3x + 8}{x^2 - 2x + 2}$$

1. $\int_0^1 x \, dx$

2. $\int_0^3 (x^2 - 3)(x^2 - 2) \, dx$

3. $\int_2^2 x^5 - 2x^2 \, dx$

4. $\int_0^{\pi} (2x - 3)^7 \, dx$

5. $\int_0^{\frac{\pi}{2}} \cos 7x \, dx$

6. $\int_0^{\frac{\pi}{2}} \tan(2x - 3) \, dx$

7. $\int_0^{\frac{\pi}{2}} \cos^3 x - 4x + 3x^2 \, dx$

8. $\int_0^1 x^2 - 2x + \sin x \, dx$

12. $\int_0^{\frac{\pi}{2}} \sin^2 5x \, dx$

13. $\int_0^{\frac{\pi}{2}} \sin^3 x \, dx$

14. $\int_0^{\frac{\pi}{2}} \cos^2 x \, dx$

15. $\int_1^2 x^2 e^x \, dx$

16. $\int_0^2 (x - 1)(dx - 2) \, dx$

17. $\int_0^1 (x - 1)(x^2 - dx) \, dx$

18. $\int_0^3 \frac{1}{(x - 1)^2(x - 2)} \, dx$

19. $\int_0^{\frac{\pi}{2}} (x^2 - 8x + 15) \, dx$

$$\int_0^1 \frac{x^2 \sin^2 x}{2} dx$$

$$8. \int_0^1 x \cos x dx$$

$$9. \int_0^2 e^{5x-3} dx$$

$$10. \int_1^2 \frac{7x-3}{x^2} dx$$

$$11. \int_0^1 \cos^2 3x dx$$

70

Summary

You have learnt the following from this chapter:

a). The fundamental Theorem of the Integral calculus If

$$\int_b^x f(x) dx = G(x) \text{ then}$$

$$\int_a^b f(x) dx = G(b) - G(a)$$

- b). $\int_a^b f(x) dx$ can be defined as the area enclosed by the four boundaries $y = f(x)$, the x -axis and the ordinates $x = a$ and $x = b$.

Further Reading

- i). Additional Pure mathematics
Harwood Clarke
Heinemann Educational Books Ltd
London

CHAPTER 8 Area Under a Curve

8.1 Introduction

In the previous chapter we have learnt the meaning of the definite integrals. In this chapter we shall learn one of the important applications of the definite integrals namely calculating the area under a curve, x – axis and the ordinates $x = a$ and $x = b$.

8.2 Objectives of the chapter

By the end of this chapter you should be able to: -

- establish the formula for finding the area under a curve the x – axis and the ordinates $x = a$ and $x = b$.
- apply the formula to calculate exact area under different types of curves and also the areas between two curves.

8.3 Formula for the area under a curve and enclosed by the lines the x – axis, and the ordinates $x = a$ and $x = b$.




Figure 1

Consider the area under a curve $y = f(x)$, and enclosed by the lines the x - axis and the ordinates $x = a$ and $x = b$.

Divide the area into n equal strips of width Δx and height y .

The area of the strip $PQRS = y_r \Delta x$. Consider the total area of all the strips.

Total area required = $\int_a^b y_r \Delta x$.

$x \rightarrow b$

b

If $\Delta x \rightarrow 0$ the area required is $A = \int_a^b y dx$ or $A = \int_a^b f(x) dx$

8.4 Area below x – axis

If the required area is below x – axis as in the figure 2




Figure 3




Figure 2

Then $A = \int_a^b y dx$ since the y coordinate is negative. Hence $\int_a^b f(x) dx$ will be negative. Since we are interested only in the magnitude of the area we can take only the magnitude if the area is negative.

8.5 Combined area under a curve

If we wish to find the area under a curve between $x = a$ and $x = c$ (figure 3). We must find the area above x-axis and the area below x – axis separately and add the magnitudes of the two areas.

For example, let the area under the curve between $x = a$ and $x = b$ be 15 units and in between $x = b$ to $x = c$ be –5 units. Then the total area under the curve between $x = a$ and $x = c$ is:

$$\begin{aligned} A &= |15| + |-5| \\ &= 15 + 5 \\ &= 20 \text{ sq units} \end{aligned}$$

8.6 Area between two intersecting curves




Figure 4

Let the equations of two curves be $y = f(x)$ and $y = g(x)$.

Suppose that they intersect at P and Q. Let the points of the intersection of the two curves be (a, y_1) and (b, y_2) .

The magnitude of the area between the two curves is:

$$A = \int_a^b |f(x) - g(x)| dx$$

Proof

The area bounded by the four boundaries $y = f(x)$, the x-axis, $x = a$ and $x = b$ is

$$\int_a^b f(x) dx$$

Similarly the area enclosed by the four boundaries $y = g(x)$, the x-axis, $x = a$ and $x = b$ is

$\int_a^b g(x) dx$. Hence the shaded area between the two curves is:

$$A = \int_a^b f(x) dx - \int_a^b g(x) dx$$

$$= \int_a^b (f(x) - g(x)) dx$$

If this is negative we can take only the magnitude or the modulus of A.

The following examples illustrate all the types of areas under the curves.

Example 1

Find the area of the rectangle bounded by the line $y = 2$, the x – axis, the ordinate $x = 1$ and $x = 5$ using the formula. $A = \int_a^b y dx$


Solution

Figure 5

The curve in this case is $y = 2$.

Using the formula $A = \int_a^b y dx$ we have

$$\begin{aligned} A &= \int_1^5 2 dx \\ &= 2x \Big|_1^5 \\ &= (10) - (2) \\ &= 8 \text{ sq units} \end{aligned}$$

The area in this case is very simple since the area of the rectangle of length 4 units and breadth 2 units is 8 sq units.

Example 2

Sketch the curve $y = 2x^2$ from $x = 0$ to $x = 5$.

Evaluate the area enclosed by the curve, the x – axis and the ordinates $x = 1$ to $x = 4$

Solution

Let us prepare a table showing some points on the curve.

X	0	1	3	5
---	---	---	---	---




Figure 6

The above figure shows the curve $y = 2x^2$.

By formula,

$$A = \int_a^b y \, dx$$

$$= \int_0^5 2x^2 \, dx$$

$$= 2x^3 \Big|_0^5$$

$$= 2(5^3 - 0^3)$$

$$= 2(125 - 0)$$

$$= 2(125)$$

$$= 250$$

Example 3

1. Sketch the curve $y = x^3 - 3x^2 + 2x$ from $x = 0$ to $x = 3$.
2. Find the area included between the curve $y = x^3 - 3x^2 + 2x$ and the axis of x from $x = 0$ to $x = 2$.

2

3. Evaluate $\int_0^3 (x^3 - 3x^2 + 2x) \, dx$. Interpret your result.

0

Solution

a) The following table shows some points on the curve.

x	0	1	2	0.5	1.5	2
y	0	0	0	0.375	-0.375	0

The graph is shown in the figure




Figure 7

b) Part of the area is positive and the other part is negative. Hence we should find each area separately.

Area from $x=0$ to $x=1$ is given by

$$A = \int_0^1 (x^3 - 3x^2 + 2x) dx$$

$$= \left[\frac{x^4}{4} - x^3 + x^2 \right]_0^1$$

$$= \left(\frac{1^4}{4} - 1^3 + 1^2 \right) - \left(\frac{0^4}{4} - 0^3 + 0^2 \right)$$

$$= \left(\frac{1}{4} - 1 + 1 \right) - 0$$

$$= \frac{1}{4}$$

= 1/4 sq units

Area from $x=1$ to $x=2$ is given by

$$A_2 = \int_1^2 (x^3 - 3x^2 + 2x) dx$$

$$= \int_0^1 (x^4 - x^3 - x^2) dx$$

$$= \left[\frac{x^5}{5} - \frac{x^4}{4} - \frac{x^3}{3} \right]_0^1$$

$$= \left(\frac{1}{5} - \frac{1}{4} - \frac{1}{3} \right) - \left(0 - 0 - 0 \right)$$

$$= -\frac{1}{4} \text{ sq units}$$

$$\text{Required area} = A_1 + A_2$$

$$= \frac{1}{4} + \frac{1}{4}$$

$$= \frac{1}{2} \text{ sq units}$$

$$\text{c) } A = \int_0^2 (x^3 - 3x^2 + 2x) dx$$

$$= \left[\frac{x^4}{4} - x^3 + x^2 \right]_0^2$$

$$= \left(\frac{2^4}{4} - 2^3 + 2^2 \right) - \left(0 - 0 + 0 \right)$$

$$= \left(\frac{16}{4} - 8 + 4 \right) - 0$$

$$= 0 \text{ sq unit}$$

If we evaluate the area directly from $x = 0$ to $x = 2$ we get the area = 0. This is because the area above x – axis is positive and that below the x - axis is negative. Hence we should not find the area directly from $x = 0$ to $x = 2$.

Example 4

Evaluate the area enclosed between the curve $y = x^2 - 5x + 4$ and the x - axis.

Solution

Here the ordinates are not given. Let us find the points of intersection of the curve with the x – axis.

Solving $y = x^2 - 5x + 4$ (the curve)


And $y = 0$ (the x - axis) we have

$$x^2 - 5x + 4 = 0$$

$$(x - 1)(x - 4) = 0$$

Hence $x = 1$ and $x = 4$.

If we sketch the curve $y = (x - 1)(x - 4)$ we get the following graph.



The entire area is below x - axis hence required area is:

$$A = \int_1^4 (x^2 - 5x + 4) dx$$

$$= \left[\frac{x^3}{3} - \frac{5x^2}{2} + 4x \right]_1^4$$

$$= \left(\frac{4^3}{3} - \frac{5 \cdot 4^2}{2} + 4 \cdot 4 \right) - \left(\frac{1^3}{3} - \frac{5 \cdot 1^2}{2} + 4 \cdot 1 \right)$$

$$= \left(\frac{64}{3} - 80 + 16 \right) - \left(\frac{1}{3} - \frac{5}{2} + 4 \right)$$

$$= \left(\frac{64}{3} - 64 \right) - \left(\frac{1}{3} - \frac{5}{2} + 4 \right)$$

$$= \frac{64}{3} - 64 - \frac{1}{3} + \frac{5}{2} - 4$$

$$= \frac{64}{3} - \frac{1}{3} - 64 + \frac{5}{2} - 4$$

$$= \frac{63}{3} - 64 + \frac{5}{2} - 4$$

$$= 21 - 64 + \frac{5}{2} - 4$$

$$= -47 + \frac{5}{2}$$

$$= -\frac{94}{2} + \frac{5}{2}$$

$$= -\frac{89}{2}$$

$$= -44.5$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

$$= 44.5 \text{ sq units}$$

Example 5

Find the area between the curve $x^2 = y$ and the straight line $y = 2$.

Solution

It is always better to sketch the curve to determine whether the required area is completely above the x - axis or below the x - axis or a portion of the area is above the x - axis and the other portion is below the x - axis.

The following figure shows the sketch of the curves:




Figure 9

Solving $y = x^2$ and $y = 2$ we get $x^2 = 2$ and $x = \sqrt{2}$ and $x = -\sqrt{2}$ and $y = 2$. Since the complete area required is above the x -axis we get the required area.

$$\begin{aligned}
 \text{Area} &= \int_{-\sqrt{2}}^{\sqrt{2}} y \, dx \\
 &= \int_{-\sqrt{2}}^{\sqrt{2}} x^2 \, dx \\
 &= \left[\frac{x^3}{3} \right]_{-\sqrt{2}}^{\sqrt{2}} \\
 &= \frac{(\sqrt{2})^3}{3} - \frac{(-\sqrt{2})^3}{3} \\
 &= \frac{2\sqrt{2}}{3} + \frac{2\sqrt{2}}{3} \\
 &= \frac{4\sqrt{2}}{3} \text{ sq units.}
 \end{aligned}$$

This is the area below the curve and the x -axis.

The shaded area = Area of the rectangle ABCD $-\frac{4\sqrt{2}}{3}$

$$= \frac{1}{2} \left(\frac{4\sqrt{2}}{3} \right)$$

$$= \frac{2\sqrt{2}}{3}$$

$$= \frac{2\sqrt{2}}{3} \text{ sq units}$$

Example 6

Find the area enclosed by the line $y = 4x$ and the curve $y = x^2$.

Solution

The graph is shown in the figure below:




Figure 10

The points of intersection of the curve $y = x^2$ and the line $y = 4x$ are given by solving the two equations.

Using $x^2 = 4x$ we have

$$x^2 - 4x = 0 \text{ or } x(x - 4) = 0.$$

Hence $x = 0$ or $x = 4$ correspondingly $y = 0$ or $y = 16$.

The required area (shaded) = Area under the line ($y = 4x$) – area under the curve ($y = x^2$).

$$= \int_0^4 4x \, dx - \int_0^4 x^2 \, dx$$

$$= \left[2x^2 - \frac{x^3}{3} \right]_0^4$$

$$\left[\frac{1}{2} \times \frac{1}{3} \times \frac{1}{3} \right]$$

$$\left[\frac{1}{2} \times \frac{1}{3} \times \frac{1}{3} \right]$$

$$= 32 - \frac{64}{3}$$

$$= \frac{32}{3} \text{ sq units}$$

Exercise 8

1. Find the area under the curve $y = x + 3x^2$ between $x = 1$ and $x = 2$.
2. Find the area enclosed by the curve $x^2y = 36$, the x - axis and the lines $x = 6$ and $x = 9$.
3. Find the area enclosed by the curve $\sqrt{x}$, the x - axis and the ordinates $x = 1$ and $x = 4$.
4. Find the area enclosed by the curve $y = x(x - 4)$ and the axis of x .
5. Find the area in the first quadrant enclosed by the line $y = x$ and the curve $y = x^3$.
6. Find the area between the curve $x^2 = y$ and the line $y = 2$.
7. Find the area in the positive quadrant enclosed by the curves $y = x^2$ and $y = x^3$.
8. Find the area common to the two curves $y^2 = 12x$ and $x^2 + y^2 = 24x$.
9. Find the area common to the circle $x^2 + y^2 = 25$ and the parabola $3x^2 = 16y$.
10. Find the area common to the circle $x^2 + y^2 = 16$ and the ellipse $x^2 + 3y^2 = 24$.

Summary

You have learnt the following from this chapter:

- i. The area enclosed by a curve $y = f(x)$, the x - axis, the ordinate $x = a$ and the ordinate $x = b$ is given by

$$\int_a^b f(x) dx$$

$$A = \int_a^b y dx$$

$$= \int_a^b f(x) dx$$

- ii. Area above the x – axis is positive and the area below the x – axis is negative.
- iii. If the required area consists of one portion above the x - axis and the other portion below the x – axis. We must find each area separately and add their magnitudes.
- iv. If $y = f(x)$ and $y = g(x)$ are two curves and if they intersect, the numerical value of the area between the two curves. $\int_a^b f(x) - g(x) dx$ provided the common points of intersection have their x coordinates a and b.

Further reading

Additional pure mathematics

By L. Harwood Clarke

Heinemann Educational Books Ltd

London

CHAPTER 9

Improper Integrals

9.1 Introduction

In the chapter seven, we have learnt to evaluate the definite integrals whose limits are finite ($a \leq x \leq b$). In this chapter we shall learn to evaluate some integrals whose limits are infinite or the integrands become infinite even though the interval is finite. Such integrals are called improper integrals. We shall define improper integrals in detail in the following sections.

9.2 Objectives of the chapter.

By the end of this chapter you should be able to

- define the improper integrals of two kinds
- test the convergence and divergence of improper integrals
- test the absolute convergence and conditional convergence of improper integrals
- evaluate the Cauchy's principal value of the improper integral of the second kind.

9.3 Definition of improper Integrals

There are two kinds of improper Integrals a).

Improper Integrals of the first kind

When one or both the limits of a definite integral is infinite the integral is called an improper integrals of the first kind.

b). Improper integrals of the second kind

When the integrand has a singular point (the integrand becomes infinite) in the interval of the limits, the definite integral is called an improper integral of the second kind.

$$x^3 \leq 2$$

Consider $\int_0^3 x^3 dx$

The interval of the limit is $0 \leq x \leq 3$

$$x^3 \leq 2$$

At the point $x = 1$, the integrand _____ becomes infinite and hence

$$x^3 \leq 2$$

$$x \leq 1$$

$\int_0^{\infty} \frac{1}{x^3 + 2} dx$ is an improper integral of the second kind

$\int_0^{\infty} \frac{1}{x} dx$ and $\int_0^{\infty} \frac{1}{(x-3)} dx$ are also improper integrals of the second kind having singularities at the lower limit $x = 0$ and at the upper limit $x = 3$ respectively.

9.4 Convergence or divergence of the improper integrals of the type

$\int_a^{\infty} f(x) dx$

$$\int_a^b f(x) dx$$

The improper integral $\int_a^{\infty} f(x) dx$ is said to be convergent if $\lim_{b \rightarrow \infty} \int_a^b f(x) dx$ exists (or converges) to a finite limit and the limit is called **the value of the improper integral**.

If the limit does not exist we say that the improper integral **does not exist (or diverges)**

Example 1

a). When will you say that the improper integral $\int_1^{\infty} \frac{1}{x} dx$ is convergent or divergent.

$$\int_1^{\infty} \frac{1}{x} dx$$

b). Show that $\int_1^{\infty} \frac{1}{x} dx$ is divergent.

Solution

$$\int_1^b \frac{1}{x} dx$$

$$\lim_{b \rightarrow \infty} \int_1^b \frac{1}{x} dx$$

a). If $\lim_{b \rightarrow \infty} \int_1^b \frac{1}{x} dx$ converges (or exists) we say that $\int_1^{\infty} \frac{1}{x} dx$ converges. Otherwise, we say that it diverges.

lim $\int_1^b \ln x dx = \lim_{b \rightarrow \infty} (b \ln b - b + 1)$

diverges. $\lim_{b \rightarrow \infty} (b \ln b - b + 1) = \infty$

$$= b \ln b - b + 1$$

$$b \ln b - b + 1$$

$$\begin{aligned} \lim_{b \rightarrow \infty} (b \ln b - b + 1) &= \lim_{b \rightarrow \infty} (b \ln b - b) + 1 \\ &= \lim_{b \rightarrow \infty} (b \ln b - b) \\ &= \lim_{b \rightarrow \infty} (b \ln b - b) \\ &= \lim_{b \rightarrow \infty} (b \ln b - b) \end{aligned}$$

$$\lim_{b \rightarrow \infty} (b \ln b - b) = \lim_{b \rightarrow \infty} (b \ln b - b)$$

$$\lim_{b \rightarrow \infty} (b \ln b - b) = \lim_{b \rightarrow \infty} (b \ln b - b)$$

Since $\lim_{b \rightarrow \infty} (b \ln b - b) = \infty$, the integral $\int_1^{\infty} \frac{1}{x} dx$ does not exist, it diverges.

Example 2

$$\int_1^{\infty} \frac{1}{x^2} dx$$

Show that the improper integral $\int_1^{\infty} \frac{1}{x^2} dx$ is convergent

Solution

$\int_1^b \frac{1}{x^2} dx$ converges or diverges if $\lim_{b \rightarrow \infty} \frac{1}{x^2}$ exists or does not.

$$\int_1^b \frac{1}{x^2} dx$$

$$\lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \left[-\frac{1}{x} \right]_1^b$$

$$\lim_{b \rightarrow \infty} \left[-\frac{1}{x} \right]_1^b = \lim_{b \rightarrow \infty} \left(-\frac{1}{b} + \frac{1}{1} \right)$$

$$= \lim_{b \rightarrow \infty} \left(-\frac{1}{b} + 1 \right)$$

$$= \lim_{b \rightarrow \infty} \left(1 - \frac{1}{b} \right)$$

$$= 1 - \lim_{b \rightarrow \infty} \frac{1}{b}$$

$$= 1 \quad (\text{Limit exists and finite})$$

Hence $\int_1^{\infty} \frac{1}{x^2} dx$ is convergent.

9.5 Definition of absolute convergence and conditional convergence of an improper integral

□□

□

i).The improper integral, $\int_a^\infty f(x) dx$ is said to be absolutely convergent if

□□

| |

$\int_a^\infty f(x) dx$ is convergent.

□□

ii).The improper integral $\int_a^\infty f(x) dx$ is said to be conditionally convergent if

□

$\int_a^\infty f(x) dx$ is convergent but $\int_a^\infty |f(x)| dx$ is not convergent.

□□

9.6

Cauchy's

inequality
for
definite

integrals $b b$

| |

Cauchy's inequality states that

□

$\int_a^\infty f(x) dx$ □

□□

$\int_a^\infty f(x) dx$

This inequality is very useful to determine the absolutely convergent and conditionally convergent improper integrals.

Example 3

$$\sum_{n=1}^{\infty} \sin nx$$

Show that $\sum_{n=1}^{\infty} \sin nx$ is absolutely convergent.

Solution

By Cauchy's inequality,

$$\left| \sum_{n=1}^N \sin nx \right| \leq \sqrt{\sum_{n=1}^N 1} \sqrt{\sum_{n=1}^N \sin^2 x} = \sqrt{N} \sqrt{\sum_{n=1}^N \sin^2 x}$$

$$\leq \sqrt{N} \sqrt{N} = N$$

$$\sum_{n=1}^{\infty} \sin nx \leq \sum_{n=1}^{\infty} \sin^2 nx$$

$$\sum_{n=1}^{\infty} \sin^2 nx \leq \sum_{n=1}^{\infty} \sin^2 nx$$

(1)

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N \sin^2 nx = \sum_{n=1}^{\infty} \sin^2 nx$$

$$\sum_{n=1}^{\infty} \sin^2 nx \leq \sum_{n=1}^{\infty} \sin^2 nx$$

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N \sin^2 nx = \sum_{n=1}^{\infty} \sin^2 nx$$

$$\lim_{b \rightarrow \infty} \int_b^{\infty} \frac{1}{e^x} dx$$

$$= \lim_{b \rightarrow \infty} \left[-e^{-x} \right]_b^{\infty}$$

$$= \lim_{b \rightarrow \infty} \left(0 - (-e^{-b}) \right)$$

□

$$= \lim_{b \rightarrow \infty} \left(\frac{1}{e^b} \right)$$

$$= \lim_{b \rightarrow \infty} \frac{1}{e^b} = 0$$

$$= 1 \text{ Since } \lim_{b \rightarrow \infty} \frac{1}{e^b} = 0$$

□

Hence $\int_0^{\infty} x e^{-x} dx$ is convergent.

b

9.7 Theorem: if $\int_a^b f(x) dx$ is an absolutely convergent improper integral it is also convergent, where b can be finite or infinite.

Proof

By Cauchy's inequality,

b

$$\left| \int_b^{\infty} f(x) dx \right| \leq \int_b^{\infty} |f(x)| dx$$

If $\int_a^{\infty} |f(x)| dx$ is convergent (or absolutely convergent) then it has an upper

bound M

$$\int_a^b |f(x)| dx \leq M(b-a)$$

Hence $\int_a^b f(x) dx \leq M(b-a)$

Then $\int_a^b f(x) dx \leq M(b-a)$ and hence it is convergent.

In other words $\int_a^b f(x) dx$ is convergent if $\int_a^b |f(x)| dx$ is convergent.

Example 5

Show that $\int_0^\infty e^{-x} \cos 2x \, dx$ is absolutely convergent.

Solution

We must show that $\int_0^\infty |e^{-x} \cos 2x| \, dx$ is convergent.

$$\int_0^b |e^{-x} \cos 2x| \, dx = \lim_{b \rightarrow \infty} \int_0^b |e^{-x} \cos 2x| \, dx$$

$$= \lim_{b \rightarrow \infty} \int_0^b |e^{-x}| \, dx$$

$$b > 1$$

$$\lim_{x \rightarrow \infty} e^{-x} = 0$$

$$b > 1$$

$$\lim_{x \rightarrow \infty} \frac{1}{e^b} = 0$$

$$\lim_{x \rightarrow \infty} \frac{1}{e^b} = 0$$

Since $\int_0^{\infty} e^{-x} \cos 2x \, dx$ is convergent since the limit exists $b > 1$

$$\int_0^{\infty} e^{-x} \cos 2x \, dx$$

$$\int_0^{\infty} e^{-x} \cos 2x \, dx$$

$$0$$

$$\int_0^{\infty}$$

Hence $\int_0^{\infty} e^{-x} \cos 2x \, dx$ is absolutely convergent.

9.8 Convergence and divergence of the improper integrals of the second kind.

$$b$$

suppose that $f(x)$ has a singularity at an

Consider the integral $\int_a^b f(x) dx$

interior point c of $[a, b]$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

(1)

The improper integral $\int_a^b f(x) dx$ is said to be convergent if both the improper integrals on the right side of (1) exist.

The sum of these is defined as **the value of the improper integral from a to b in (1)**

9.9 The Cauchy's principal value of $\int_a^b f(x) dx$

Let $f(x)$ be defined on $[a, b]$ except possibly at any interior point c , which is a singular point of $f(x)$.

If $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ does not exist but

$\lim_{\epsilon \rightarrow 0} \left(\int_a^{c-\epsilon} f(x) dx + \int_{c+\epsilon}^b f(x) dx \right)$ does exist. We call this limit

the Cauchy's principal value of $\int_a^b f(x) dx$

Thus the Cauchy's principal value of

$$\lim_{\epsilon \rightarrow 0} \left(\int_a^{c-\epsilon} f(x) dx + \int_{c+\epsilon}^b f(x) dx \right)$$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

□
□

Example 6

2 1

a) Show that $\int_1^{\infty} \frac{1}{x} dx$ is improper.

2 1

b) Find the Cauchy's principle value of $\int_1^{\infty} \frac{1}{x} dx$

Solution

1

a) The integrand $\frac{1}{x}$ has a singularity at $x = 0$ in the interval $[-1, \infty)$

2 □] x

2 1

Hence $\int_1^{\infty} \frac{1}{x} dx$ is an improper integral of the second kind.

1

0

b) $\int_1^{\infty} \frac{1}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx = \lim_{t \rightarrow \infty} [\ln x]_1^t = \lim_{t \rightarrow \infty} (\ln t - \ln 1) = \lim_{t \rightarrow \infty} \ln t = \infty$ Clearly $\int_0^1 \frac{1}{x} dx$ and $\int_{20}^{\infty} \frac{1}{x} dx$

both diverge;

1

$$\lim_{x \rightarrow 0^+} \ln x = -\infty$$

$\ln 0$ does not exist

$$\int_0^1 \frac{1}{x} dx = \lim_{\epsilon \rightarrow 0^+} \int_\epsilon^1 \frac{1}{x} dx$$

2

||

$$\int_0^1 \frac{1}{x} dx = \lim_{\epsilon \rightarrow 0^+} \left[\ln x \right]_\epsilon^1 = \lim_{\epsilon \rightarrow 0^+} (\ln 1 - \ln \epsilon) = -\lim_{\epsilon \rightarrow 0^+} \ln \epsilon = \infty$$

Hence $\int_0^1 \frac{1}{x} dx$ does not exist.

However, $\int_1^{\infty} \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \left[-\frac{1}{x} \right]_1^b = \lim_{b \rightarrow \infty} \left(-\frac{1}{b} + 1 \right) = 1$ exist when ϵ

$$\text{Since } \int_1^{\infty} \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \left[-\frac{1}{x} \right]_1^b = \lim_{b \rightarrow \infty} \left(-\frac{1}{b} + 1 \right) = 1$$

$$\text{and } \int_1^{\infty} \frac{1}{x} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x} dx = \lim_{b \rightarrow \infty} \left[\ln x \right]_1^b = \lim_{b \rightarrow \infty} (\ln b - \ln 1) = \infty$$

$$\text{Hence } \int_1^{\infty} \frac{1}{x} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x} dx = \lim_{b \rightarrow \infty} (\ln b - \ln 1) = \infty$$

Thus the Cauchy's principal value of $\int_{-1}^1 \frac{1}{x} dx = \ln 2$

Example 7

3

dx

$\int_{-2}^3$

Show that $\int_{-2}^3 \frac{1}{x^4} dx$

- i). does not converge
- ii). does not have a principal value

Solution

$\int_{-2}^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

The integral $\int_{-2}^3 \frac{1}{x^4} dx$ is improper since $x=0$ is a singularity on the interval $[-2 \leq x \leq 3]$

$\int_{-2}^0 \frac{1}{x^4} dx$

$\int_0^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

Now

$\int_{-2}^0 \frac{1}{x^4} dx$

$\int_0^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

$\int_{-2}^0 \frac{1}{x^4} dx$

$\int_0^3 \frac{1}{x^4} dx$

$\int_{-2}^3 \frac{1}{x^4} dx$

$\int_{-2}^0 \frac{1}{x^4} dx = \lim_{t \rightarrow 0^-} \int_{-2}^t \frac{1}{x^4} dx = \lim_{t \rightarrow 0^-} \left[-\frac{1}{3x^3} \right]_{-2}^t = \lim_{t \rightarrow 0^-} \left(-\frac{1}{3t^3} + \frac{1}{24} \right) = -\infty$

$\int_0^3 \frac{1}{x^4} dx = \lim_{t \rightarrow 0^+} \int_t^3 \frac{1}{x^4} dx = \lim_{t \rightarrow 0^+} \left[-\frac{1}{3x^3} \right]_t^3 = \lim_{t \rightarrow 0^+} \left(-\frac{1}{81} + \frac{1}{3t^3} \right) = \infty$

does not converge

The Cauchy's principal value of

$$\int_0^3 dx - \int_0^3 dx = \int_0^3 dx$$

$$\int_0^4 \frac{1}{x^2} dx - \int_0^4 \frac{1}{x^2} dx = \int_0^4 \frac{1}{x^2} dx \quad \text{where } x > 0$$

$$\int_0^3 \frac{1}{x^2} dx - \int_0^3 \frac{1}{x^2} dx = \int_0^3 \frac{1}{x^2} dx$$

$$\int_0^3 \frac{1}{x^2} dx = \int_0^3 \frac{1}{x^2} dx = \int_0^3 \frac{1}{x^2} dx$$

where

$$x > 0$$

$$\int_0^3 \frac{1}{x^2} dx = \int_0^3 \frac{1}{x^2} dx = \int_0^3 \frac{1}{x^2} dx$$

$$\int_0^3 \frac{1}{x^2} dx$$

$$= \int_0^3 \frac{1}{x^2} dx$$

Thus even the principal value does not exist.

Exercise 9

1. Classify the following into

Improper integrals of the first kind and Improper integrals of the second kind.

$$\int_0^1 \frac{1}{x^2} dx$$

$$\int_0^\infty e^{-x} \sin x dx$$

$$\text{a). } \int_0^1 \frac{1}{x^2} dx$$

$$\text{b). } \int_0^\infty \frac{1}{x^2} dx$$

$$\int_0^1 dx$$

$$\int_0^\infty \frac{1}{x^2} dx$$

3. When will you say that the improper integral $\int_0^\infty \frac{1}{x^2} dx$ converges or diverges.

$$\int_0^\infty \frac{1}{x^2} dx$$

$$\int_{-\infty}^{-3} x^{-3} dx$$

$$\int_2^{\infty} \frac{1}{x^4} dx$$

$$\int_5^{\infty} \frac{1}{x^3} dx$$

2. When will you say that the improper integral $\int_a^b f(x) dx$ converges or diverges.

4. Define the Cauchy's principal value of the improper integral $\int_a^b f(x) dx$ if $f(x)$ has a singularity at c where $a < c < b$.

5. Define absolute convergence and conditional convergence of an improper

$$\int_a^b f(x) dx$$

6. Show that $\int_0^{\infty} \frac{1}{x^5} dx$ is divergent.

7. Show that $\int_1^{\infty} \frac{1}{x^5} dx$ is divergent.

8. Show that $\int_0^{\infty} e^{-x} \cos 3x dx$ is absolutely convergent.

9. Test whether $\int_1^{\infty} \frac{\sin x}{x^2} dx$ is absolutely convergent or not.

10. Find Cauchy's principal value of $\int_{-1}^1 \frac{dx}{x^2}$.

11. Evaluate $\int_0^{\frac{\pi}{2}} \ln(\sin x) dx$

12. Prove that $\int_0^{\infty} \cos^2 x dx$ converges
(Hint: Make the substitution $u = x^2$)

13. Define $\int_a^b f(x) dx$ and $-\int_b^a f(x) dx$

14. Test the following integrals for convergence

a). $\int_0^{\frac{\pi}{2}} \sin x dx$

c). $\int_0^1 (1-x^2)^{-2} dx$

$\int_1^{\sqrt{x}} \frac{1}{x^3} dx$

b). $\int_1^{\infty} \ln x dx$

$\int_1^{\infty} \frac{1}{x^2} dx$

d). $\int_0^{\frac{\pi}{2}} \sin x dx$

15. Test whether the principal value for the integral $\int_0^5 \frac{dx}{x^4}$ exists.

16. Show that $\int_0^{\infty} \frac{\sin x \cos x}{x^4} dx$ converges.

Summary

You have learnt the following from the chapter:

- i. Definition of improper integrals of first and second kind.
- ii. Convergence and divergence of improper integrals.
- iii. Absolute convergence and conditional convergence of improper integrals.
- iv. Cauchy's inequality for absolute convergence.

v. The Cauchy's principal value of the improper integrals $\int_a^b f(x) dx$ when $f(x)$ has a singularity at c where $a \leq c \leq b$

Further reading

1. Advanced Calculus
By Watson Fulks
John Wiley and sons
New York. Brisbane Toronto.

CHAPTER 10

Length of an Arc of a Curve

10.1 Introduction

If you ask a secondary school student to measure the length of an arc of a curve, he will use a string or a flexible material like a string placed upon the arc of the curve and he measures the length of the string placing against a ruler. Also an approximation of length could be found by marking a number of points on the arc and finding the sum of the lengths of all the chords that are small. Any mathematician could do the same method unless he knows the equation of the curve either in Cartesian form or Parametric or any other form. In this chapter we shall learn the method of finding the length of an arc of a curve.

10.2 Objectives of the chapter

By the end of this chapter you should be able to.

- derive an expression or formula for S , the arc length of a curve if the Cartesian or parametric equation of the curve is known.
- calculate accurately the length of the curve using the formula.

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Proof

10.3 Length of an arc S of a curve in Cartesian equation is

$$S = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$




Figure 1

Suppose that an arc of a curve is divided into n parts by points $p_0, p_1, p_2, \dots, p_n$. We shall assume that the sum of the lengths of chords, $p_0 p_1 + p_1 p_2 + \dots + p_{n-1} p_n$ is equal to the length of the arc in the limit when the length of the chords $\rightarrow 0$.

If Δx and Δy are the increments in x and y from p_{r-1} to p_r .

$$\begin{aligned} (p_{r-1} p_r)^2 &= (\Delta x)^2 + (\Delta y)^2 \\ &= (\Delta x)^2 + (\Delta y)^2 \\ &= (\Delta x)^2 + (\Delta y)^2 \\ &= (\Delta x)^2 + (\Delta y)^2 \end{aligned}$$

□

$$\text{Then } p_{r-1} p_r = \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

Summing for all the chords and proceeding to the limit

$$\text{Length of arc } S = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

□

where a and b are the values of x corresponding to the end points of the arc.

10.4 The length of an arc S of a curve in the Parametric form is

$$S = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

□

□

where t_1 and t_2 are the values of t corresponding to the end points of the arc.

Proof

We have seen in the previous section

$$(p_r - p_r)^2 = (\Delta x)^2 + (\Delta y)^2$$

$$= \Delta x^2 + \Delta y^2 = (\Delta t)^2$$

$$\Delta x = \Delta t$$

□
□

$$\text{Then } p_r - p_r = \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta t)^2} = \Delta t$$

Summing for all the chords and proceeding to the limit.

$$\text{Length of the arc } S = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Example 1

Find the length of the arc of parabola $y^2 = 4ax$ cut off by its latus rectum.




Figure 2

Solution

The equation of a parabola in parametric equation is $x = at^2$, $y = 2at$.

Latus Rectum is the line perpendicular to the axis of the parabola (in the Figure 2) at a distance a from the vertex. If TS is the Latus Rectum, the coordinates of L is $(a, 0)$ and of T is $(a t^2, 2 a t)$. Hence $a t^2 = a$ and $t = 1$. At the vertex $0, a t^2 = 0$

Since arc SOT = Twice arc OT

$$2 \int_{t=0}^1 \sqrt{dx^2 + dy^2}$$

we have S

$$= \int_0^1 \sqrt{dx^2 + dy^2}$$

$$(SOT = 2OT) \int_0^1 \sqrt{dx^2 + dy^2}$$

$$\int_0^1$$

$$(\text{arc} = 2OT) = 2 \int_0^1 \sqrt{(2at)^2 + (2a)^2} dt$$

$$= 2 \int_0^1 \sqrt{4a^2 t^2 + 4a^2} dt$$

$$= 4a \int_0^1 \sqrt{1 + t^2} dt$$

$$= 4a \left[\frac{1}{2} \ln \left(\frac{1 + \sqrt{1 + t^2}}{t} \right) + \frac{1}{2} \sqrt{1 + t^2} \right]_0^1 \quad \text{using integration by parts.}$$

$$= 2a \left[2 \ln(2 + \sqrt{5}) - 2 \right]$$

Example 2

- a). Draw the curve $6y^2 = x(x-2)^2$
 b). Find the point of intersection of the curve with x-axis.

- c). Calculate the length of the loop of the curve $y^2 = \frac{1}{6}x(x-2)^2$




Figure 3

Solution

- b) Let $6y^2 = x(x-2)^2$
 Differentiating with respect to x dy
 $12y \frac{dy}{dx} = (x-2)^2 + 2x(x-2)$
 $= (x-2)(3x-2)$

$$\frac{dy}{dx} = \frac{(x-2)(3x-2)}{12y}$$

$$\int \frac{dy}{y} = \int \frac{(x-2)(3x-2)}{12x(x-2)^2} dx$$

$$= \int \frac{(3x-2)}{12x(x-2)} dx$$

$$= \frac{1}{12} \int \frac{(3x-2)}{x(x-2)} dx$$

$$= \frac{1}{12} \int \left(\frac{A}{x} + \frac{B}{x-2} \right) dx$$

$$= \frac{1}{12} \left(\int \frac{A}{x} dx + \int \frac{B}{x-2} dx \right)$$

$$= \frac{1}{12} \left(A \ln|x| + B \ln|x-2| \right) + C$$

$$= \frac{1}{12} (3x-2) \ln|x-2| - \frac{1}{12} (3x-2) \ln|x| + C$$

$$= \frac{1}{12} \int_0^2 \frac{(3x-2)}{x(x-2)} dx$$

$$= \frac{(3x - 2)^2}{24x}$$

$$\begin{aligned} \text{Length of the loop} &= 2 \int_0^a \sqrt{1 - \frac{(3x-2)^2}{24x}} dx \\ &= 2 \int_0^a \sqrt{\frac{24x - (3x-2)^2}{24x}} dx \end{aligned}$$

$$\begin{aligned} &= 2 \int_0^a \sqrt{\frac{24x - 9x^2 + 12x - 4}{24x}} dx \\ &= 2 \int_0^a \sqrt{\frac{-9x^2 + 36x - 4}{24x}} dx \\ &= \frac{2}{\sqrt{24}} \int_0^a \sqrt{-x^2 + 4x - \frac{1}{3}} dx \\ &= \frac{2}{\sqrt{24}} \left[\frac{1}{2} \left(-x^2 + 4x - \frac{1}{3} \right) \sqrt{-x^2 + 4x - \frac{1}{3}} + \frac{1}{2} \left(-x^2 + 4x - \frac{1}{3} \right) \right]_0^a \\ &= \frac{2}{\sqrt{24}} \left[\frac{1}{2} \left(-a^2 + 4a - \frac{1}{3} \right) \sqrt{-a^2 + 4a - \frac{1}{3}} + \frac{1}{2} \left(-a^2 + 4a - \frac{1}{3} \right) \right] \end{aligned}$$

□
□

Example 3 Show that the length of the circumference of a circle of radius a is $2\pi a$.

Solution




Figure 4

The equation of the circle of radius a is $x^2 + y^2 = a^2$. Differentiating with respect to x ,

$$\frac{dy}{dx} \cdot \frac{dy}{dx} + \frac{2x}{2y} = 0 \quad \text{we have } 2x + 2y \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$\text{length of arc APB} = S = \int_0^a \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^a \sqrt{1 + \frac{x^2}{y^2}} dx$$

$$= \int_0^a \sqrt{\frac{y^2 + x^2}{y^2}} dx = \int_0^a \frac{\sqrt{x^2 + y^2}}{y} dx = \int_0^a \frac{a}{y} dx = \frac{a}{y} \int_0^a \frac{1}{y} dx = \frac{a}{y} \left[\frac{x}{y} \right]_0^a = \frac{a}{y} \cdot \frac{a}{y} = \frac{a^2}{y^2}$$

$$\int_0^a y^2 dx$$

$$a^3$$

$$\int_0^a \sqrt{a^2 - x^2} dx$$

$$\int_0^a \sqrt{a^2 - x^2} dx$$

$$\int_0^a \sin x dx$$

$$S = \frac{\pi a^2}{2}$$

the length of the circumference of the circle
= 4 times S

$$= 4 \times \frac{\pi a^2}{2} = 2\pi a^2$$

Example 4

Find the length of the arc in the first quadrant of $y = 2x^{\frac{3}{2}}$ from $x = 0$ to $x = \frac{1}{3}$.

Solution

The equation of the curve is given in Cartesian form. The length of the arc of a curve in Cartesian form is given by

$$\int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$\int_0^2 \sqrt{1 + 9x} dx$$

$$\int_0^2 \sqrt{1 + 9x} dx$$

$$\int_0^2 \sqrt{1 + 9x} dx$$

$$\int_0^2 \sqrt{1 + 9x} dx$$

$$\int_0^2 \sqrt{1 + 9x} dx$$

using (ax + b) formula

$$\frac{2}{27} \int_0^2 \sqrt{1 + 9x} dx$$

$$\frac{2}{27} \left[\frac{2}{3} (1 + 9x)^{3/2} \right]_0^2$$

$$= 9 \text{ units}$$

Example 5

Find the length of the arc from $\theta = 0$ to $\theta = \frac{\pi}{4}$ of the curve given by $x = 3 \cos \theta$,

$$y = 3 \sin \theta.$$

Solution

The equation of the curve is given in parametric equation. In parametric equation, the length of arc is given by

$$S = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Here

$$\begin{aligned} S &= \int_0^{\frac{\pi}{4}} \sqrt{\left(\frac{d(3\cos\theta)}{d\theta}\right)^2 + \left(\frac{d(3\sin\theta)}{d\theta}\right)^2} d\theta \\ &= \int_0^{\frac{\pi}{4}} \sqrt{9\sin^2\theta + 9\cos^2\theta} d\theta \\ &= 3 \int_0^{\frac{\pi}{4}} \sqrt{\sin^2\theta + \cos^2\theta} d\theta \\ &= 3 \int_0^{\frac{\pi}{4}} 1 d\theta \\ &= 3 \left[\theta \right]_0^{\frac{\pi}{4}} \\ &= 3 \left(\frac{\pi}{4} - 0 \right) \\ &= \frac{3\pi}{4} \end{aligned}$$

$$= 3 \int_0^1 d\theta$$

θ

$$= 3(\theta)_0^1$$

$$= \frac{3\pi}{4} \text{ units}$$

Length of an arc S of a curve in polar coordinates (r, θ) is

$$S = \int_{\theta_1}^{\theta_2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

□
□
□
□

where values θ_1 and θ_2 are the values of θ corresponding to the end points of the curve.

Proof




Figure 5

Let $OP_{r-1} = r$ and $OP_r = r + \Delta r$

Angle $\angle P_{r-1}OP_r = \theta$ and angle $\angle P_{r-1}OP_r = \Delta \theta$

Let NP_{r-1} be perpendicular to OP_r .

Now $NP_{r-1} = r \Delta \theta$ and $NP_r = \Delta r$

$$\begin{aligned} \text{Hence } (P_{r-1}P_r)^2 &= (NP_r)^2 + (NP_{r-1})^2 \\ &= (\Delta r)^2 + (r \Delta \theta)^2 \end{aligned}$$

$$\begin{aligned} \lim_{\Delta \theta \rightarrow 0} \frac{r^2 \Delta \theta^2 + (\Delta r)^2}{(\Delta \theta)^2} &= \lim_{\Delta \theta \rightarrow 0} \left(r^2 + \frac{(\Delta r)^2}{(\Delta \theta)^2} \right) \\ &= r^2 \end{aligned}$$

□

$$S = \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$= \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$= \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$= \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$S = \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$= \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

□

where α and β are the values of θ corresponding to at the end points of the curve.

Example 6

Find the length of the equiangular spiral $r = a e^{k\theta}$ from $\theta = 0$ to $\theta = 2\pi$.

Solution

The equation of the curve is given in polar form. The length of arc of a curve in polar form is given by

$$S = \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} d\theta$$

$$S = \int_0^r r^2 \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$= \int_0^r \sqrt{1 + \left(\frac{dr}{dz}\right)^2} dz$$

$$k = 1 - k$$

□

Exercise 10

1. Find the length of the arc of $y = \ln \sec x$ from $x = -\frac{\pi}{6}$ to $x = \frac{\pi}{6}$ —
2. Find the length of the arc of the curve $y = \ln \sec x$ from $x = 0$ to $x = \frac{\pi}{3}$ □
3. Show that the length of the arc of the parabola $y^2 = 4ax$ cut off by the line $3y = 8x$ is $a(\ln 2 + \frac{15}{16})$. —
4. Sketch the astroid given by $x = a \cos^3 t$, $y = a \sin^3 t$ and find the length of its circumference.
5. Find the length of the arc of the curve $x = a(\cos \theta + \theta \sin \theta)$, $y = a(\sin \theta - \theta \cos \theta)$
6. Sketch the arc of the cycloid $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ from $\theta = 0$ to $\theta = 2\pi$. Find its length.
7. Find the length of the spiral of Archimedes $r = a\theta$ from $\theta = 0$ to $\theta = \pi$. 8. What is the length of the circumference of the cardioid $r = a(1 + \cos \theta)$.

Summary

You have learnt the following from this chapter. a).

The length of an arc S of a curve is

$$\int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

i). In Cartesian form $S = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$

$\int dx \int dy$ ii). In parametric form $S = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

$$dt \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

iii). In polar form $S = \int_{r_1}^{r_2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$

The limits being the end points of the curve

b). To apply the formula for various problems

Further Reading

Additional pure mathematics
Harwood Clarke
Heinemann Educational Books Ltd.
London.

CHAPTER 11

Volume and Surface Area of Solid of Revolution

11.1 Introduction

Consider an area under a curve between two ordinates. When this area is rotated about the axis of x the resulting solid is called a **Solid of Revolution**. A section of this solid by a plane perpendicular to the axis of x is a circle. When the area under a line of length l parallel to the axis of x is rotated you will obtain a right circular cylinder of height l . **The area under a line through the origin when rotated about the axis of x a right circular cone is obtained. We shall now find the volume and surface area of a solid of revolution, given the equation of the curve and the bounding ordinates.**

11.2 Objectives of the chapter

By the end of this chapter you shall be able to:

- derive the formulae for the volume and the surface area of the solid of revolution.
- apply these formulae for finding the volumes and surface areas of common solids.

11.3 The formula for the volume of a solid of revolution

The curve $y = f(x)$ bounded by the ordinate at $x = a$ and $x = b$ is rotated 360° or more about the x-axis a solid of revolution is generated. The volume of this solid is given by

$$V = \int_a^b \pi y^2 dx$$




Figure 1

Consider the curve $y = f(x)$ between the ordinates AN and BM. Let P be (x, y) and Q be $(x + \Delta x, y + \Delta y)$ on the curve. PS and QR are perpendicular to x-axis and the distance $RS = \Delta x$ where $\Delta x \rightarrow 0$.

When the area ANMBA is rotated about x axis, the elementary area PSRQ rotates about OX, forming a slice of radius y and thickness Δx thus an elementary cylinder of volume $\pi y^2 \Delta x$ is generated.

The volume generated by the area ANMB is

$V =$ Limits of the sum of the volume generated by such elementary areas as PSRQ. =

$$\lim_{\Delta x \rightarrow 0} \sum_{x=a}^b \pi y^2 \Delta x \text{ where } x \text{ varies from } x = a \text{ to } x = b.$$

$$\pi \int_a^b y^2 dx$$

$$V = \int_a^b \pi y^2 dx$$

b

$$\pi \int_a^b y^2 dx$$

$$V = \pi \int_a^b y^2 dx$$

Similarly if a curve is rotated about the y axis between the lines $y = c$ and $y = d$, the volume generated is given by

$$y \text{ from } c \text{ to } d$$

$$V = \pi \int_c^d x^2 dy$$

Example 1

Find the volume of a cone of height h and base radius r .

Solution




Figure 2

A cone is formed by the revolution of the area under the line OA about the x-axis.

The gradient of the line OA is $\frac{r}{h}$ and the equation of the line OA is $y = \frac{r}{h}x$

If V is the volume of the cone

$$x \text{ from } 0 \text{ to } h$$

$$V = \pi \int_0^h y^2 dx$$

$$\int_0^h \int_{-\sqrt{r^2-x^2}}^{\sqrt{r^2-x^2}} 2x \, dx \, h$$

$$\int_0^h \int_{-\sqrt{r^2-x^2}}^{\sqrt{r^2-x^2}} 2x \, dx \, h$$

$$\int_0^h \int_{-\sqrt{r^2-x^2}}^{\sqrt{r^2-x^2}} 2x \, dx \, h$$

$$\int_0^h \int_{-\sqrt{r^2-x^2}}^{\sqrt{r^2-x^2}} 2x \, dx \, h$$

$$= \frac{4}{3} \pi r^3$$

Example 2

Find the volume of a hemisphere of radius r units.

$$V = \frac{4}{3} \pi r^3$$

Hence show that the volume of a sphere of radius r is given by

Solution




Figure 3

A hemisphere is formed by the revolution of the area of a quadrant of a circle $x^2 + y^2 = r^2$ about the x-axis.

The volume of the hemisphere

$$= \int_0^r y^2 dx$$

$$= \int_0^r (r^2 - x^2) dx$$

$$= \left(r^2 x - \frac{x^3}{3} \right) \Big|_0^r$$

$$= \left(r^2 \cdot r - \frac{r^3}{3} \right) - \left(0 - \frac{0^3}{3} \right)$$

$$= \frac{2}{3} r^3$$

$$\frac{2}{3} r^3 \quad \frac{4}{3} r^3$$

Hence the volume of sphere = twice $\frac{2}{3} r^3 = \frac{4}{3} r^3$

Example 3

Find the volume obtained by rotating the area under the curve $y = 1 + x$ between $x = 1$ and $x = 2$ about the axis of x .

Solution

Volume

$$= \pi \int_1^2 y^2 dx$$

$$= \pi \int_1^2 (1+x)^2 dx$$

$$= \pi \int_1^2 (x^2 + 2x + 1) dx$$

$$= \pi \left[\frac{x^3}{3} + x^2 + x \right]_1^2$$

$$= \pi \left(\frac{8}{3} + 4 + 2 - \left(\frac{1}{3} + 1 + 1 \right) \right)$$

$$= \pi \left(\frac{8}{3} + 4 + 2 - \frac{1}{3} - 1 - 1 \right)$$

$$= \pi \left(\frac{8}{3} + 4 + 2 - \frac{1}{3} - 2 \right)$$

$$= \pi \left(\frac{8}{3} + 4 \right)$$

$$= \frac{19\pi}{3}$$

11.4 The Formula for the Surface Area of a Solid of Revolution

When an arc AB rotates about the axis of x a surface of revolution is generated. If arc AP = s and arc AQ = s + Δs, arc PQ = Δs.




Figure 4

When the elementary area PSRQ rotates about x axis the curved surface area of the cylinder of radius y and height Δs = 2πyΔs, where Δs → 0.

Hence the total surface area generated by the arc AB about x-axis.

= Limit $\sum 2\pi y \Delta s$ from x = a to x = b.

Limit $\Delta x \rightarrow 0$

$\Delta y \rightarrow 0$

2

$$\Delta s^2 = (\Delta x)^2 + (\Delta y)^2 \quad (\text{see 10.2})$$

$\Delta s = \sqrt{(\Delta x)^2 + (\Delta y)^2}$. From x = a to x = b since (Δs)

$x \rightarrow b$

$$S = \lim_{\Delta x \rightarrow 0} \sum_{x=a}^{x=b} 2\pi y \Delta s = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

This formula can be remembered as surface area is $S = 2\pi \int y ds$

In parametric form

$$\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Surface areas = $\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

Example 4

Find the total surface area of a sphere of radius r.

Solution

A sphere is obtained by the revolution of a semi circle about its diameter.

Total surface area of a hemisphere = $2\pi \int_0^r y ds$

The equation of the circle is $x^2 + y^2 = r^2$

Then $y^2 = r^2 - x^2$ differentiating with respect to x

$$2y \frac{dy}{dx} = -2x$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$dx = -\frac{y^2}{x} \frac{dy}{y} \text{ then } \int \frac{1}{x} dx = \int \frac{1}{y^2} dy$$

$$y^2 + y^2 = r^2, \text{ Hence } \int_0^r 1 dy = \int_0^r \frac{1}{y} dy$$

$$\int_0^r 1 dy = \int_0^r \frac{1}{y} dy$$

$$dy = \frac{r}{y}$$

But x

$$\int dx = \int y^2 \text{ Then}$$

surface area of the sphere

$$r^2 \int_0^{\pi} \sin^2 \theta d\theta$$

$$= 2 \int_0^a \int_{\sqrt{y}}^{\sqrt{1-y}} \sqrt{1-y} \, dx dy$$

$$= 2 \int_0^a \sqrt{1-y} \cdot y \, dy$$

$$= 2 \int_0^a \sqrt{1-y} \, dy$$

$$= 4 \int_0^a \sqrt{1-y} \, dy$$

$$= 4 \int_0^a \sqrt{1-y} \, dy$$

Example 5

Find the surface area generated by the loop of the curve $x=t^2$, $y=t^3$ about the y axis.

Solution

The curve cuts the x -axis at $y=0$ or $t=0$, Hence $t=0$ or $t=\sqrt[3]{3}$ or $x=0$ or $x=3$.

Eliminating t from $x=t^2$ and $y=t^3$ we have the Cartesian equation of the

curve, $9y^2 = x^3$

$x=3$ or $x=0$

$$ds^2 = dx^2 + dy^2$$

— since $(ds)^2 = (dx)^2 + (dy)^2$

$$dt = \frac{dx}{2t} = \frac{dy}{3t^2}$$

$$= 4t^2 \times 1 \times t^2 \times 2$$

$$= 8t^4$$

$$ds = 8t^4 dt$$

$$\frac{ds}{dt}$$

The required surface area = $2\pi \int_0^3 y ds$

$$= 2\pi \int_0^3 t^3 ds$$

$$= 2\pi \int_0^3 t^3 \times 8t^4 dt$$

$$= 16\pi \int_0^3 t^7 dt$$

$$= 16\pi \left[\frac{t^8}{8} \right]_0^3$$

$$= 2\pi \left[\frac{t^8}{4} \right]_0^3$$

$$= 2\pi \left[\frac{3^8}{4} - 0 \right]$$

$$= 2\pi \left[\frac{6561}{4} \right] = \frac{6561\pi}{2}$$

$$= \frac{6561\pi}{2} \text{ Sqr units}$$

$$= 3\pi \text{ Sqr units}$$

Example 6

Show that the curved surface area of a cone radius r , height h and slant height l is

$$\pi rl.$$

A




Figure 6

Solution

A cone is formed by the revolution of the area under the line OA. OB = height h, AB is the radius r and OA is the slant height l.

Gradient of OA is $\frac{r}{h}$ and hence the equation of OA is $y = \frac{r}{h}x$

$$2\pi \int_0^h y \sqrt{1 + \left(\frac{r}{h}\right)^2} dx$$

Surface area S =

$$= 2\pi \int_0^h \frac{r}{h}x \sqrt{1 + \frac{r^2}{h^2}} dx$$

$$= 2\pi \int_0^h \frac{r}{h}x \sqrt{h^2 + r^2} dx$$

$$= \frac{2\pi r}{h} \int_0^h x \sqrt{h^2 + r^2} dx$$

$$= \frac{2\pi r}{h} \left[\frac{1}{2} x^2 \sqrt{h^2 + r^2} \right]_0^h$$

$$= \frac{2\pi r}{h} \left[\frac{1}{2} h^2 \sqrt{h^2 + r^2} \right]$$

$$= \pi r l$$

Exercise 11

- Find the volume generated by revolving about the x – axis that part of the $y = x^2$ which lies between the origin and $x = 3$.

curve $y = x^3$

- The equation of the circle $x^2 + y^2 = 9$ is rotated about x- axis between the limits -3 and 3 . Find the volume of the sphere by integration.
- Sketch the curve $y = x^2 - x - 2$. The part of the curve lying between the points where it cuts the x- axis is rotated about that axis. Find the volume so generated.

- Find the volume of the spheroid generated by rotating the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$ about the minor axis.

- Find the volume when the loop of the curve $y^2 = x(2x - 1)^2$ revolves about the x – axis.

- Find the surface area of the solid generated by the revolution of the loop of the curve.

$x = t^2, y = \frac{t^3}{3}$ about the x – axis.

- Find the surface area of the solid generated by the revolution of the loop of the curve. $3ay^2 = x(x - a)^2$.

8. Prove that the area of the curved surface of a sphere of radius a intercepted between two parallel planes at distances b and c from the center is $2\pi a(b - c)$.
9. Find the surface area generated when the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ revolves about major axis.
10. The area between the line $y = x$ and the curve $y = x^2$ is rotated about the axis of x . Find the volume formed.

Summary

You have learnt the following from this chapter.

1. The formulae for the volume of a solid of revolution obtained by rotating a curve about x -axis between the ordinate at $x = a$ and $x = b$ is:

$$V = \pi \int_a^b y^2 dx$$

About y -axis between the lines $x = c$ and $x = d$ is:

$$V = \pi \int_c^d x^2 dy$$

2. The formula for the surface area of a solid of revolution about x -axis is:

$$S = 2\pi \int y ds \text{ if } s \text{ is known.}$$

$S = 2\pi \int_{t_1}^{t_2} y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$ if the equation of the curve is in parametric form.

3. Application of the formulae for any solid of revolution.

Further Reading

2. Additional pure mathematics By L. Harwood Clarke. Heinemann Educational Books Ltd London.
3. Advanced calculus By Watson Fulks.
John Wiley and sons New York. Brisbane. Toronto.

MWALIMU GREG

CHAPTER 12

Numerical integration:

Trapezoidal and Simpson Rule

12.1 Introduction

In chapter eight we have learnt how to find the area under a curve using definite integral. Suppose that the equation of the curve is not known or the known function of x cannot be integrated. In such cases an approximation for the area under a curve may be found by several numerical methods. In this chapter we shall consider two important methods namely Trapezoidal rule and Simpson rule to find approximate areas under a curve.


12.2 Objectives of the chapter:

By the end of this chapter you should be able to □
establish the Trapezoidal rule.

- derive the formula for the area under a parabola.
- establish Simpson's rule.
- apply the Trapezoidal rule and Simpson's rule.

12.3 : Area of a trapezium

A quadrilateral in which one pair of opposite sides are parallel is called a trapezium. The area of a trapezium is the mean of parallel sides multiplied by the perpendicular distance between them.




In the above figure A B C D is a trapezium in which A B is parallel to DC. If $AB=y_1$, $x = y_2$ and the distance $AD = h$ the area of the trapezium is

$$= \frac{h(y_1 + y_2)}{2} \text{ or } \frac{h}{2}(y_1 + y_2)$$

this

12.4: Trapezoidal rule



Consider a curve DC with end ordinates y_0 and y_n cutting x axis at A and B respectively. Divide the interval between A and B into n equal parts and erect ordinates at each point of division including the end points A and B. We wish to find

the area ABCD under the curve DC. Let $\frac{AB}{n} = h$.

The area under curve is found by joining the ends of consecutive ordinates and by treating each trapezium so formed as an approximation for the area under the corresponding portion of the graph.

The area of the first trapezium is $\frac{h}{2}(y_0 + y_1)$

of the second is $\frac{h}{2}(y_1 + y_2)$

of the third is $\frac{h}{2}(y_2 + y_3)$

.....

of the nth trapezium is $\frac{h}{2}(y_{n-1} + y_n)$

where h is the distance between consecutive ordinates.

Then an approximation for total area under the curve DC is

$$A = \frac{h}{2} (y_0 + y_1 + y_1 + y_2 + y_2 + y_3 + \dots + y_{n-1} + y_n)$$

$$A = \frac{h}{2} (y_0 + 2y_1 + 2y_2 + 2y_3 + \dots + 2y_{n-1} + y_n)$$

This is the trapezoidal rule

You can observe the following:


In finding the total sum,

- i) y_0 and y_n remain the same ii) $y_1, y_2, \dots, y_{n-1}$ are multiplied by two

$b - a$, to get the area under the curve. The total sum is multiplied by h , where h is the distance between consecutive ordinates.

curve

2 n



Given that OA is divided into 6 equal parts at and the corresponding ordinates are given below:

$x_0 = 0$	$y_0 = 8$
$x_1 = 1$	$y_1 = 12$
$x_2 = 2$	$y_2 = 14$
$x_3 = 3$	$y_3 = 11$
$x_4 = 4$	$y_4 = 9$
$x_5 = 5$	$y_5 = 3$
$x_6 = 6$	$y_6 = 1$

Solution

$$\frac{6-0}{6}$$

$x=0$ to 6 is divided into 6 equal parts; hence $h =$ h
 $= 1$

$$\text{Total area} = \frac{1}{2} y_0 + y_1 + y_2 + y_3 + y_4 + y_5 + \frac{1}{2} y_6$$

□
□

$$y_0 = 8 \quad y_1 = 12 \quad y_2 = 14 \quad y_3 = 11 \quad y_4 = 9 \quad y_5 = 6 \quad y_6 = 1$$

$$y_2 = 14 \quad y_3 = 11 \quad y_4 = 9 \quad y_5 = 6 \quad y_6 = 1$$

$$y_3 = 11 \quad y_4 = 9 \quad y_5 = 6 \quad y_6 = 1$$

$$y_4 = 9 \quad y_5 = 6 \quad y_6 = 1$$

$$y_5 = 6 \quad y_6 = 1$$

$$\text{Total Sum} = 107$$


□

□

$$\begin{aligned} \text{Total area} &= \frac{h}{2} (y_0 + y_1 + y_2 + y_3 + y_4 + y_5 + y_6) \\ &= \frac{1}{2} (107) \\ &= 53.5 \text{ sq units} \end{aligned}$$

Example 2

Consider the curve $y = \frac{1}{x}$ in the figure given below between $x = 1$ and $x = 2$. The x interval is divided into 5 equal parts. The ordinates at these points are $y_0, y_1, y_2, y_3, y_4, y_5$ which are given below:



y_0	y_1	y_2	y_3	y_4	y_5
-------	-------	-------	-------	-------	-------

1.0000	0.8333	0.7143	0.6250	0.5556	0.5000
--------	--------	--------	--------	--------	--------

- a) Calculate $\int_{1/2}^1 \frac{1}{x} dx$ using trapezoidal rule with 5 intervals (Assume $\int \frac{1}{x} dx$

is not known)

- b) Suppose that you know $\int \frac{1}{x} dx = \ln x + c$
Calculate the area under the curve.
- c) Calculate the percentage error.

Solution

$$y_0 = 1.0000$$

$$y_0 = 1.0000$$

$$y_1 = 0.8333$$

$$2y_1 = 1.6666$$

$$y_2 = 0.7143$$

$$2y_2 = 1.4286$$

$$y_3 = 0.6250$$

$$2y_3 = 1.2500$$

$$y_4 = 0.5556$$

$$2y_4 = 1.1112$$

$$y_5 = 0.5000$$

$$y_5 = 0.5000$$

Total Sum

$$= 6.9564$$

$$h = \frac{1 - 1/2}{5} = 0.2$$

$$5$$

$$2.1 \quad h = 0.2$$

Then $\int_1^{2.1} x dx = \frac{1}{2} y_0 + 2 y_1 + 2 y_2 + 2 y_3 + 2 y_4 + 2 y_5$

$$\begin{aligned} &= \frac{1}{2} \times 6.9564 \\ &= 0.69564 \\ &= 0.6956 \text{ to four decimals.} \end{aligned}$$

b) Now, $\int_1^2 x^{-1} dx = \ln x \Big|_1^2$

$$\begin{aligned} &= \ln 2 - \ln 1 \\ &= 0.6931 - 0 = 0.6931 \end{aligned}$$

c) To calculate the percentage error.

$$\begin{aligned} \text{The percentage error} &= \frac{0.6956 - 0.6931}{0.6931} \times 100 \\ &= 0.0025 \times 100 \\ &= 0.25\% \end{aligned}$$

$$\text{[Percentage error} = \frac{\text{True value} - \text{Approximat value}}{\text{True value}} \times 100 \%]$$


Example3

a) Find the distance traveled in a straight line by a particle in 6 seconds given that the velocity m per sec is connected with the time t sec as follows

t	0	1	2	3	4	5	6
v	10	12	15	16	11	5	3

b) Sketch the time velocity graph
(In velocity – time graph, area under graph represents the distance traveled)

Solution



Figure

We are required to find $\int_0^6 y dx$ or area under the curve between $t = 0$ and 6

$$y_0 = 10 \quad y_0 = 10$$

$$y_1 = 12 \quad 2y_1 = 24$$

$$y_2 = 15 \quad 2y_2 = 30$$

$$y_3 = 16 \quad 2y_3 = 32$$

$$y_4 = 11 \qquad 2y_4 = 22$$

$$y_5 = 5 \qquad 2y_5 = 10$$

$$y_6 = 3 \qquad y_6 = 3$$

$$\text{Total Sum} \qquad 131$$

$$h = \frac{6 - 0}{6} = 1 \text{ unit}$$


Total area under curve (distance traveled)

$$\begin{aligned} & \frac{h}{3} (y_0 + 2y_1 + 2y_2 + 2y_3 + 2y_4 + 2y_5 + y_6) \\ &= \frac{1}{3} (131) \\ &= 65.5 \text{ metre} \end{aligned}$$

12.5 Principle of Simpsons Rule

The Trapezoidal Rule we have learnt is obtained by treating each **part of the curve** as a straight line, $y = bx + c$. Suppose that we now find a better approximation for each part of the curve in the form $y = ax^2 + bx + c$ which is a parabola.

Formula for the area under the curve $y = ax^2 + bx + c$ between $x = -h$ and $x = h$



If the curve is $y = ax^2 + bx + c$ the **exact area** under the curve ABC is

$$\begin{aligned}
 A &= \int_{-h}^h (ax^2 + bx + c) dx \\
 &= \left[\frac{ax^3}{3} + \frac{bx^2}{2} + cx \right]_{-h}^h \\
 &= \left(\frac{ah^3}{3} + \frac{bh^2}{2} + ch \right) - \left(-\frac{ah^3}{3} + \frac{bh^2}{2} - ch \right) \\
 &= \frac{2ah^3}{3} + 2ch \quad (1)
 \end{aligned}$$

Now we shall find the area under the curve ABC in terms of the ordinates y_0, y_1, y_2

$$\begin{aligned}
 \text{The curve is} \quad & y = ax^2 + bx + c \\
 \text{When } x = -h \quad & y_0 = ah^2 - bh + c \\
 \text{When } x = 0 \quad & y_1 = c \\
 \text{When } x = h \quad & y_2 = ah^2 + bh + c \quad (2)
 \end{aligned}$$

$$\begin{aligned}
 \text{Now } y_0 + y_2 &= (ah^2 - bh + c) + (ah^2 + bh + c) \\
 &= 2ah^2 + 2c \\
 &= 2ah^2 + 2y_1 \text{ using (2)} \quad (3)
 \end{aligned}$$

$$\text{From (3) } 2ah^2 = y_0 + y_2 - 2y_1 \quad (4)$$

The actual area by integration, from (1) is

$$A = \frac{2ah^3}{3} + 2ch \quad (1)$$

$$= h \left(\frac{2ah^2}{3} + 2c \right) \quad (5)$$

$$= \frac{h}{3} (2ah^2 + 6c)$$

$$= \frac{h}{3} (y_0 - 2y_1 + y_2 + 6y_1) \text{ using (2) and (4) in (5)}$$


$$A = \frac{h}{3} (y_0 + 4y_1 + y_2)$$

3

Thus the area under the parabola ABC in terms of the ordinates is h

$$A = \frac{1}{3} (y_0 + 4y_1 + y_2)$$

12.6 Simpsons Rule for finding approximate area under a curve



As we have seen

The area between y_0 and y_2 $= \frac{1}{3} h(y_0 + 4y_1 + y_2)$

The area between y_2 and y_4 $= \frac{1}{3} h(y_2 + 4y_3 + y_4)$ The

area between y_4 and y_6 $= \frac{1}{3} h(y_4 + 4y_5 + y_6)$

.....
The area between y_{n-2} and y_n $= \frac{1}{3} h(y_{n-2} + 4y_{n-1} + y_n)$

The total area under the curve AB is therefore

$$A = \frac{1}{3} h(y_0 + 4y_1 + 2y_2 + 4y_3 + \dots + 4y_{n-1} + y_n)$$

This is Simpsons Rule

You must understand the following to apply Simpsons rule.

- i). Divide the interval $(a \leq x \leq b)$ into **even** number of equal intervals, $h = \frac{b - a}{n}$

n

- ii). In the addition, the first and the last ordinates y_0 and y_n remain the same
- iii). Add four times the ordinates $y_1, y_3, y_5, \dots$, and two times the ordinates y_2, y_4, y_6 , (alternatively four times odd and two times even ordinates respectively) are added.
- iv). The total sum is multiplied by one third of the width of equal interval (or multiply h the sum by $\frac{1}{3}$)

To apply Simsons Rule the number of divisions of the interval must be even In other words n should be an even positive integer to apply Simpsns Rule.

Example 4

6

Find $\int_0^6 y dx$, given the following values, using

- i) Simpsns Rule for the interval $(0 \leq x \leq 6)$
- ii) Trapezoidal Rule for the interval $(0 \leq x \leq 6)$

X	0	1	2	3	4	5	6
Y	8	12	14	11	9	3	1

Solution

- i) Area under curve using Simpsns Rule

$$Y_0 = 8 \quad Y_0 = 8$$

$$Y_1 = 12 \quad 4Y_1 = 48$$

$$Y_2 = 14 \quad 2Y_2 = 28$$

$$Y_3 = 11 \quad 4Y_3 = 44$$

$$Y_4 = 9 \quad 2Y_4 = 18$$

$$Y_5 = 3 \quad 4Y_5 = 12$$

$$Y_6 = 1 \quad Y_6 = 1$$

$$\text{Total Sum} \quad 159$$

$$h = \frac{6 - 0}{6} = 1$$

Approximate area is therefore.

$$\begin{aligned} A &= \frac{1}{3} h (y_0 + 4y_1 + 2y_2 + 4y_3 + 2y_4 + 4y_5 + y_6) \\ &= \frac{1}{3} (1)(159) \\ &= 53 \text{ sq units} \end{aligned}$$

ii) Area under curve using Trapezoidal Rule

(≤)

6)

$$Y_0 = 8$$

$$Y_0 = 8$$

$$Y_1 = 12$$

$$2Y_1 = 24$$

$$Y_2 = 14$$

$$2Y_2 = 28$$

$$Y_3 = 11$$

$$2Y_3 = 22$$

$$Y_4 = 9$$

$$2Y_4 = 18$$

$$Y_5 = 3$$

$$2Y_5 = 6$$

$$Y_6 = 1$$

$$Y_6 = 1$$


$$h = \frac{6 - 0}{6} = 1$$

Approximate area is therefore

$$\begin{aligned} A &= \frac{1}{2} h (y_0 + 2y_1 + 2y_2 + 2y_3 + 2y_4 + 4y_5 + y_6) \\ &= \frac{1}{2} (1) (107) \\ &= 53.5 \text{ sq units} \end{aligned}$$


Exercise 12

1. State the formula for
 - i). Trapezoidal Rule
 - ii). Simpsons Rule
2. What is the area of a trapezium whose ordinates are y_0 and y_1 and the distance between them is h .
3. Three consecutive ordinates are y_0 , y_1 , and y_2 , and each equal interval is h units. What is the area between the curve, x-axis and the ordinates y_0 and y_2



4. If $y_0 = 2$ units, $y_1 = 3$ units and $y_2 = 5$ units calculate the area included by the curve ABC, the ordinates y_0 , y_2 and x-axis.
 - i) If AB and BC are considered as straight lines ii) If the curve ABC is considered as a parabola $y = ax^2 + bx + c$.

5.



In the figure above, the equation of the curve is $y = \frac{1}{27x^2}$. Find the area under the curve between $x = 0$ and $x = 1$ using

under the curve between $x = 0$ and $x = 1$ using

- Trapezoidal Rule with ten equal parts
- Simpsons Rule with 10 equal parts

6. A car starts from rest and its velocity in meter per second over the first ten seconds is given by the following table:

t	0	1	2	3	4	5	6	7	8	9	10	v	2	2.4	2.8	3.2	4.0	4.2	4.5	6.0	6.2	6.3	6.4
---	---	---	---	---	---	---	---	---	---	---	----	---	---	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Find the distance traveled by the car at the end of 10 seconds.

3

7. Find an approximate value for $\int_0^1 x^2 dx$ using

0

- i) Simpsons Rule by taking 10 equal intervals
 ii) Trapezoidal Rule by taking 10 equal intervals
 iii) Integration

$$\int_1^2$$

$$\int_0^1 e^{-x} dx$$

8. Evaluate $\int_0^1 e^{-x} dx$ by Simpsons Rule taking ten intervals.
 9. The distance moved by a body starting from rest is given by the following table.

Time t sec	0	1	2	3	4	5	6
Distance s met	0	5	17	34	57	84	115

Plot a velocity-time graph and estimate the speed after 6 seconds.

128

Summary

You have learnt the following from this chapter: h

1. Area of trapezium = $\frac{h}{2} (y_0 + y_1)$ y_0 and y_1 are the ordinates and h

is the distance between y_1 and y_2 .

2. Area under a parabola $y = ax^2 + bx + c$ when there are two intervals bounded

$$\frac{h}{3} (y_0 + 4y_1 + y_2)$$

by y_0, y_1 and $y_2 = 3$

3. Area under a curve using Trapezoidal Rule is h

$$= \frac{h}{2} (y_0 + 2y_1 + 2y_2 + \dots + 2y_{n-1} + y_n)$$

4. Area under a curve using Simpson's Rule is h

$$A = \frac{h}{3} (y_0 + 4y_1 + 2y_2 + 4y_3 + 2y_4 + \dots + y_n) .$$

where n should be an even integer.

5. Applying the Trapezoidal Rule and Simpsons Rule for finding the approximate area under a curve between $x = a$ and $x = b$

Further Reading

1. Mathematics of Physics and Modern Engineering
By Sokolnikoff
Mc Graw – Hill Book company, INC New York Toronto London
2. Additional Pure Mathematics By L. Harwood Clarke
Heinmann Educational Books Ltd
London