

## Lesson 4: CLASSIFICATION

**Biological classification** is the scientific process of arranging organisms into hierarchical groups based on their shared biological characteristics, similarities, and evolutionary relationships. It is a branch of biology primarily managed through **Taxonomy** (the naming and defining of groups) and **Systematics** (the study of the diversification of living forms).

### Key Words in classification

- **Hierarchy:** Organisms are organized into a nested system of levels (taxa), moving from the most general to the most specific (Species).
- **Criteria:** Classification is based on various data points, including morphology (physical shape), phylogeny (evolutionary descent), and molecular data (DNA and proteins).
- **Nomenclature:** It provides a standardized system—**Binomial Nomenclature**—to ensure every species has a unique, universally recognized two-part name (e.g., *Panthera leo* for a lion).

### Taxonomy

Taxonomy is a sub-discipline of systematics. It focuses specifically on the practical aspects of identifying and categorizing organisms. The main aim of taxonomy is to provide a stable, universal system for naming and grouping organisms. It involves:

- **Identification:** Determining where an unknown specimen fits.
- **Description:** Listing the physical and genetic traits of a species.
- **Classification:** Placing the organism into the taxonomic hierarchy (Kingdom, Phylum, etc.).
- **Nomenclature:** Assigning a formal scientific name using the Binomial system.
- **Focus:** It is primarily concerned with the **rules and procedures** of classification.

### 2. Systematics


Systematics is a broader field that encompasses taxonomy but adds the critical layer of **evolutionary history**. Its main aim is to understand the diversification of life and the relationships between living things through time. It involves:

- **Phylogeny:** Constructing "evolutionary trees" to show how species diverged from common ancestors.
- **Comparative Biology:** Analyzing similarities and differences in DNA, anatomy, and behavior to find evolutionary links.
- **Biogeography:** Studying where organisms live and how they got there over time.
- **Focus:** It is concerned with the **evolutionary patterns** and the "why" behind the groupings.

## Phylogeny

Phylogeny is the actual historical path of evolution followed by a group of organisms. It represents the genealogical relationships between species, much like a massive, ancient family tree. It describes who is related to whom and how long ago they shared a common ancestor. A phylogeny is visually represented by a **Phylogenetic Tree**.

- **Root:** Represents the common ancestor of all organisms in the tree.
- **Nodes:** Points where a lineage splits (speciation), representing a common ancestor of the descending groups.
- **Branches:** Represent the evolution of lineages over time.
- **Tips (Taxa):** The groups or species currently being studied (usually at the right or top of the tree).



## Phylogenetics

Phylogenetics is the scientific discipline that uses data to reconstruct the phylogeny of a group. It is the analytical process of building the tree. Phylogeneticists use various "clues" to determine relationships: Physical traits like skeletal structure or leaf shape. **Molecular Data:** Comparing DNA, RNA, or protein sequences (the most common method today). **Fossil Records:** Using carbon dating and physical remains to place organisms in a timeline. **The main aim** is to create a **Cladogram** or tree that most accurately reflects the true evolutionary history of life.

## The Modern Taxonomic Rank

To define an organism's place in nature, scientists use the following ranks:

**Kingdom** (e.g., Animalia)

**Phylum** (e.g., Chordata)

**Class** (e.g., Mammalia)

**Order** (e.g., Primates)

**Family** (e.g., Hominidae)

**Genus** (e.g., *Homo*)

**Species** (e.g., *sapiens*)

Taxonomy is the part of science that focuses on naming and classifying or grouping of organisms based on their characteristics (similarities and dissimilarities). All modern classification systems have their roots in the Linnaean classification system. It was developed by Swedish botanist Carolus Linnaeus in the 1700s. He tried to classify all living things that were known at his time. He grouped together organisms that shared obvious physical traits, such as number of legs or shape of leaves. For his contribution, Linnaeus is known as the “father of taxonomy.”

Two important Linnaeus contributions to taxonomy are:

- Binomial nomenclature
- Linnaean Classification Hierarchy

## **BINOMIAL NOMENCLATURE**

Nomenclature is the naming of organisms and the rules governing the application of these names. In the nomenclature system, each organism is designated by two names, the first is the name of the genus; the second is the specific epithet (species). No two species can have the same name.

The names are always Latin (or Latinized) and the genus is capitalized while species name is not.

### **Rules in Zoological Nomenclature**

The rules of zoological nomenclature are governed by the **International Code of Zoological Nomenclature (ICZN)**. These rules ensure that every animal species has one unique, universally accepted scientific name to avoid confusion across different languages and regions.

#### **➤ Binomial Nomenclature:**

Every species name must consist of two parts, typically derived from Latin/Greek.

**Genus Name:** The first part, which is always capitalized (first letter).

**Specific Epithet:** The second part, which is always lowercase.

**Formatting:** Both names must be **italicized** in print (e.g., *Panthera leo*) or **underlined** when handwritten.

### ➤ The Principle of Priority

This is the "golden rule" of nomenclature. If a species is accidentally named twice by different scientists, the **earliest published name** (starting from Linnaeus's *Systema Naturae* in 1758) is the valid one. Any later names are considered "synonyms" and are discarded.

### ➤ Principle of Homonymy

The same name cannot be used for two different animals. A **Genus** name must be unique within the entire animal kingdom.

### ➤ Typification

Every named species must be linked to a physical specimen called a **Holotype**. This specimen is usually kept in a museum or herbarium. It serves as the permanent reference point to define exactly what that name represents if there is a future dispute.

### ➤ Standard Endings for Higher Taxa

In zoology, the names of families and subfamilies must have specific suffixes so they are easily recognizable:

**Family:** Ends in **-idae** (e.g., Felidae, the cat family).

**Subfamily:** Ends in **-inae** (e.g., Felinae).

### ➤ Publication Requirements

For a name to be valid it must be: Published in a work that is widely accessible to the scientific community. Accompanied by a description that points out the characters that differentiate the new taxon from others.

### ➤ Language and Grammar

Names must be treated as **Latin**, regardless of their origin. The specific epithet must agree grammatically with the genus name (for example, if the genus is feminine, the epithet should usually have a feminine ending).

The Latin word for white is *albus*. Watch how the ending changes:

| Genus (Gender)         | Scientific Name     | Common Name                |
|------------------------|---------------------|----------------------------|
| <b>Masculine</b> (-us) | <i>Lupus albus</i>  | Arctic Wolf                |
| <b>Feminine</b> (-a)   | <i>Egretta alba</i> | Great Egret                |
| <b>Neuter</b> (-um)    | <i>Chen album</i>   | White Goosefoot (a plant*) |

## Importance/ Significance of Classification

- **Standardization of Communication:** It provides a universal language (Binomial Nomenclature) that allows scientists across different countries and languages to identify the same organism without the confusion of local common names.
- **Systematic Organization:** It categorizes the millions of known animal species into a logical, hierarchical "filing system," making it easier to store, retrieve, and study biological data.
- **Mapping Evolutionary History:** Modern classification (systematics) reflects **phylogeny**, helping scientists understand how different species evolved from common ancestors.
- **Predictive Ability:** Because members of a group share similar traits, classification allows us to predict the anatomy, physiology, and behavior of a newly discovered species based on its relatives.
- **Identification of Biodiversity:** It serves as the baseline for identifying which species are unique, rare, or endangered, which is critical for environmental protection and conservation efforts.
- **Facilitating Research:** It helps in identifying "indicator species" for environmental health and assists in the search for medicinal compounds by targeting specific families or genera known for certain biochemical properties.
- **Understanding Ecological Roles:** By grouping animals with similar niche requirements, classification helps ecologists understand how energy and nutrients flow through an ecosystem.

## Construction of a Dichotomous Key

A dichotomous key is a scientific tool used to identify organisms or objects based on a series of paired, contrasting statements (couplets) that divide specimens into smaller groups. "Dichotomous" means "divided into two parts."

## **Taxonomic Hierarchy and Selection**

### **Types of Key Layouts**

There are two standard ways to format a zoological/dichotomous key:

#### **A. The Linked (Dichotomous) List**

The **Linked List** (also called the "Numbered" or "Sequential" key) is the industry standard. While branching diagrams are great for classrooms, the linked list is used in scientific journals and field guides because it can handle hundreds of species without becoming a cluttered mess of lines.

Let's use the **Vertebrate Key** to identify a **Frog**:

**Step 1:** Look at 1a (Feathers). The frog has none. Now look at **1b (No feathers)**. It matches! The instruction says **Go to 2**.


**Step 2:** Look at 2a (Hair/Fur). The frog has none. Now look at **2b (No hair/fur)**. It matches! The instruction says **Go to 3**.

**Step 3:** Look at 3a (Scales). The frog has none. Now look at **3b (Smooth, moist skin)**. This is the perfect description of a frog.

**Result:** The "Lead to" column for 3b gives you the final answer: **Class Amphibia**.

## B. The Branching (Flowchart) Diagram...mostly for beginners.

A **Branching Diagram** (or Flowchart) is often more intuitive than a list because it allows you to visualize the "taxonomic path" an animal takes. Instead of flipping back and forth between numbers, you simply follow the arrows based on the traits you observe.



When constructing a key for zoology, you must first decide the taxonomic scope. Are you identifying **Phyla** (broad), **Classes**, or **Species** (specific)? **Key focus areas include;**

➤ **Symmetry:** Radial (starfish) vs. Bilateral (insects/mammals).

**Radial Symmetry:** Body parts are arranged around a **central axis**; can be divided into two equal halves **along many planes passing through the central axis**. **Examples:** **Cnidaria** – jellyfish, sea anemones, corals.

### Characteristics:

- ✓ Typically seen in **sessile, drifting, or slow-moving animals**.
- ✓ Provides the ability to sense the environment equally in all directions.
- ✓ Usually cylindrical or umbrella-shaped bodies.
- **Bilateral Symmetry:** Body can be divided into two identical halves along only one plane (sagittal plane), giving right and left sides.

**Characteristics:** Found in actively moving organisms. Associated with **cephalization** (development of a head with sense organs and brain). Promotes streamlined movement and directional locomotion.

**Examples:** **Platyhelminthes** (flatworms). **Arthropoda** (insects, crustaceans, spiders). **Mollusca** (snails, octopus). **Echinoderm larvae**.

➤ **Body Cavity:** Acoelomate, Pseudocoelomate, or Coelomate.

**Coelom** = true body cavity with full mesoderm lining → more advanced animals.

**Pseudocoelom** = false body cavity partially lined by mesoderm → simpler animals.

### 1. What is a Coelom?

- ✓ Higher eukaryotic animals, generally have oogamous sexual reproduction/oogamy in which large eggs are fertilized by a tiny sperm.
- ✓ The zygote undergoes cleavage, and forms a solid ball of cells called morula.
- ✓ The morula undergoes blastulation to form a hollow ball of cells called a blastula.
- ✓ The blastula has a cavity called blastocoel/ blastocoelom and an opening called a blastopore (first opening in the embryo).
- ✓ The blastula undergoes gastrulation to form an embryo with germ/germinal layers called gastrula.
- ✓ The gastrula may have two layers (diploblastic condition) or three layers (triploblastic condition).


*Embryonic development with cell differentiation and specialization is the hallmark of higher eukaryotes (plants and animals).*

- Gastrulation converts the blastula into an embryo with two germ layers (diploblastic) or three germ layers (triploblastic).
- In the triploblasts, the inner layer, the endoderm, develops into lining that surrounds the digestive cavity.
- The outer layer, the ectoderm, will form external structures i.e.( skin and its associated structures such as mucus/slime glands, hair, fur, sense organs).
- The middle germ layer, the mesoderm, lies between these other two layers (endos=within, ektos=outside, mesos=middle, derma=skin) and is responsible for the formation of muscles and the rest of the organ systems e.g. respiratory, excretory, reproductive, nervous systems etc.

Animals with three body layers (tribloblastic) can be divided into three categories.

- In the primitive forms, there is no space/hollow/cavity between the layers, and they are acoelomate (a = without; koilos =hollow).
- The most advanced animals are coelomate. They contain a coelom, a fluid-filled space that develops in the mesoderm.
- Some middle level invertebrates are pseudocoelomates, with a fluid filled pseudocoel between the endoderm and mesoderm.

A pseudocoel forms from the remains of the blastocoel during gastrulation. whereas a coelom opens up in the mesoderm later in development.



➤ **Segmentation:** Present (earthworms) vs. Absent (roundworms).

Metamerism can be defined as repeated segmentation. Each segment (metamere) contains one pair of some or all of the organs of the body e.g. (nephridia, coelomoducts, gonads, ganglia etc).

Metamerism appears for instance in the annelids, arthropods, cestodes (flat worms-tapeworms) and chordates. Most cestodes e.g. (tapeworms) form many identical segments (proglottides) that are shed terminally when the reproductive capacity is fulfilled.

### The Comparison: Present vs. Absent

#### Segmentation Present (Phylum Annelida)

These animals have a body divided into a series of repeated segments (metameres). Both internally and externally, the body is "sectioned" like a row of train cars.

**The Example:** The Earthworm. If you look closely, you can see the distinct rings (annuli) along its body.

**In a Key:** This trait is often paired with the presence of **setae** (tiny bristles) or a **clitellum** (the "saddle" on an earthworm).

#### Segmentation Absent (Phylum Nematoda & Platyhelminthes)

The body is a continuous tube or a flat sheet with no internal partitions. It is smooth and uniform.

**The Example:** Roundworms (Nematodes) or Flatworms (Planaria). They may have a "head" and a "tail," but the space between is one solid unit.



**In a Key:** This leads the user toward traits like "Body shape: Cylindrical" (Roundworms) or "Body shape: Dorso-ventrally flattened" (Flatworms).

## 2. Incorporating Segmentation into a Lesson

If you were building a lesson for your students, you would use this trait to distinguish between the three main "Worm" phyla.

### Step-by-Step Selection Logic

1a. Body is segmented (looks like a series of rings) ..... **Phylum Annelida**

1b. Body is not segmented (smooth surface) ..... **Go to 2**

2a. Body is flattened (like a ribbon) ..... **Phylum Platyhelminthes**

2b. Body is cylindrical (tapered at both ends) ..... **Phylum Nematoda**

➤ **Skeletal Structure:** Exoskeleton (arthropods) vs. Endoskeleton (vertebrates).

When constructing a key based on skeletal structure, you are looking at where the "armor" or "support" is located in relation to the muscles and skin.

### Characteristic A: Exoskeleton (External)

The skeleton is a hard, outer shell made of **chitin**. It must be shed (molted) for the animal to grow.

**Key Indicator:** "Hard outer body covering with jointed appendages."

**Example:** Crabs, grasshoppers, spiders, and scorpions.

**Zoological Group:** Phylum Arthropoda.

### Characteristic B: Endoskeleton (Internal)

The skeleton is located inside the body, made of **bone or cartilage**. It grows along with the animal.

**Key Indicator:** "Internal framework of bone or cartilage; presence of a backbone."

**Example:** Fish, birds, humans, and crocodiles.

**Zoological Group:** Subphylum Vertebrata.

## 2. Incorporating into a Dichotomous Key

If you were teaching a lesson on broad animal classification, your key would look like this:

**The "Linked List" Format**

- 1a.** Skeleton is located outside the body (Exoskeleton)..... **Go to 2**
- 1b.** Skeleton is located inside the body (Endoskeleton)..... **Go to 3**
- 2a.** Has three pairs of legs (6 total)..... **Class Insecta**
- 2b.** Has four pairs of legs (8 total)..... **Class Arachnida**
- 3a.** Possesses scales and breathes with gills..... **Class Pisces**
- 3b.** Possesses hair or fur; produces milk..... **Class Mammalia**