

Piece-wise exponential models

Philipp Kopper

02 June 2020

Data preporcessing

As outlined in the previous section, the **PEM data structure** deals with the survival problem as a count data problem. The main feature of this representation is the partition of the follow-up time into a finite number of intervals. Within these intervals hazard rates are assumed to be piece-wise constant

Survival times & Events

Below, I show how to transform regular survival data into PEM format where the intervals are equidistant and of size 0.4. Still, it should be chosen with great care. The data set contains a case id, a variable indicating the observed event time (obs_times) and the observed event (status) As usual in survival analysis, the status 1 refers to the event while 0 to a censoring. A priori the selected interval cut points defining the baseline hazard in each interval are arbitrary.

```
library(pammttools)
library(tidyr)
library(dplyr)
data_set <- data.frame(id = 1:3, obs_times = c(1, 0.5, 2), status = c(0, 1, 1))
ped <- as_ped(data = data_set, Surv(obs_times, status) ~ ., id = "id",
  cut = seq(0, max(data_set$obs_times), 0.4))
data_set
```

```
##   id obs_times status
## 1  1      1.0      0
## 2  2      0.5      1
## 3  3      2.0      1
```

```
ped
```

```
##   id tstart tend interval      offset ped_status
## 1  1  0.0  0.4  (0,0.4] -0.9162907         0
## 2  1  0.4  0.8  (0.4,0.8] -0.9162907         0
## 3  1  0.8  1.2  (0.8,1.2] -1.6094379         0
## 4  2  0.0  0.4  (0,0.4] -0.9162907         0
## 5  2  0.4  0.8  (0.4,0.8] -2.3025851         1
## 6  3  0.0  0.4  (0,0.4] -0.9162907         0
## 7  3  0.4  0.8  (0.4,0.8] -0.9162907         0
## 8  3  0.8  1.2  (0.8,1.2] -0.9162907         0
## 9  3  1.2  1.6  (1.2,1.6] -0.9162907         0
## 10 3  1.6  2.0  (1.6,2]   -0.9162907         1
```

The offset reflects information on the observed survival time. Re-arranging the *Poisson likelihood* of the PEM one can derive the following definition:

$$\exp(o_{ij}) = t_{ij} \quad (1)$$

where o_{ij} reflects the offset of individual i at time j and t_{ij} to observed failure time. Effectively, this means that the offset o_{ij} is equal to the natural logarithm of interval length of the observation of interest i if there is an interval cut for each event time.

$$o_{ij} = \log(t_{ij} - t_{ij-1}) \quad (2)$$