

Logistic Regression Models with GEE for the Secondary Outcomes in the VQI FBVAR Dataset

Jennifer Ci, Thu Vu, Lily Hanyi Wang

Variables to adjust for

We used a correlation matrix to report the correlations between variables statistically. We also discussed with our collaborator on the choice of variables to adjust for. Here we did not report the correlation matrix.

- cluster on CENTERID
- adjust for AGECAT, GENDER, PREOP_SMOKING, PRIOR_AORSURG, PRIOR_CHF, PREOP_DIALYSIS
- adjust for PATHOLOGY, extent

In the model, we merge groups for the *extent*: merge “Juxtarenal AAA” with “Type 4 TAAA”; “Type 1 TAAA”, “Type 2 TAAA”, “Type 3 TAAA”, with “Type 5 TAAA”. Now *extent* is a binary variable, Juxtarenal or not.

- categorical secondary outcomes: POSTOP_CEREBROX, POSTOP_SPINAL_ISCHEMIA, POSTOP_DIALYSIS, POSTOP_LOS, POSTOP_COMPLICATIONS, POSTOP_LEGEMBO, POSTOP_RESPIRATORY, RETX_R_RTOR, BRANCH_POST, TOTAL_LOS, ICUSTAY, POSTOP_INTISCH

logistic regression with Generalized Estimating Equations for categorical secondary outcomes

- continuous secondary outcomes: POSTOP_PRBC

linear regression with Generalized Estimating Equations for continuous secondary outcomes

POSTOP_GFR

TOTAL_LOS: Length of stay in days between admission date and discharge date

TOTAL_LOS Length of Stay in days			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	5.14	4.11, 6.43	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

	TOTAL_LOS Length of Stay in days		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	3.79	2.99, 4.81	<0.001
AGECAT			
<50	—	—	
>79	1.91	0.88, 4.15	0.10
50-59	1.89	0.82, 4.36	0.13
60-69	1.32	0.63, 2.77	0.47
70-79	2.16	1.02, 4.55	0.043
GENDER			
female	—	—	
male	0.62	0.51, 0.75	<0.001
PREOP_SMOKING			
No	—	—	
Yes	0.99	0.76, 1.29	0.94
PRIOR_AORSURG			
Both	—	—	
Endo	1.59	0.93, 2.71	0.091
None	1.08	0.65, 1.80	0.76
Open	1.43	0.83, 2.47	0.20
PRIOR_CHF			
No	—	—	
Yes	1.51	1.21, 1.89	<0.001
PREOP_DIALYSIS			
No	—	—	
Yes	1.87	1.10, 3.16	0.020
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	1.83	1.16, 2.87	0.009
Dissection	1.77	1.02, 3.06	0.041
PAU/IMH extent	1.94	0.90, 4.20	0.093

¹OR = Odds Ratio, CI = Confidence Interval

TOTAL_LOS Length of Stay in days			
Characteristic	OR ¹	95% CI ¹	p-value
Juxtarenal	—	—	
No	2.10	1.72, 2.57	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

ICUSTAY: ICU Stay

ICU Stay			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	3.11	2.43, 3.97	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

ICU Stay			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.08	1.59, 2.71	<0.001
AGECAT			
<50	—	—	
>79	1.31	0.51, 3.37	0.58
50-59	1.30	0.47, 3.58	0.62
60-69	0.96	0.38, 2.42	0.93
70-79	1.28	0.50, 3.25	0.61
GENDER			
female	—	—	
male	0.60	0.49, 0.74	<0.001
PREOP_SMOKING			
No	—	—	
Yes	1.02	0.75, 1.38	0.92
PRIOR_AORSURG			
Both	—	—	
Endo	1.36	0.77, 2.41	0.29
None	0.75	0.43, 1.32	0.32
Open	1.31	0.72, 2.38	0.38
PRIOR_CHF			
No	—	—	
Yes	1.21	0.93, 1.57	0.17
PREOP_DIALYSIS			

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	ICU Stay		
	OR ¹	95% CI ¹	p-value
No	—	—	
Yes	1.84	1.07, 3.19	0.028
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	1.35	0.81, 2.25	0.25
Dissection	1.72	0.93, 3.19	0.084
PAU/IMH	1.25	0.50, 3.12	0.63
extent			
Juxtarenal	—	—	
No	2.23	1.78, 2.80	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_PRBC: Transfusion # Units PRBC

Transfusion # Units PRBC			
Characteristic	Beta	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	0.95	0.52, 1.4	<0.001

¹CI = Confidence Interval

POSTOP_INTISCH: Post-op Intestinal Ischemia

Intestinal Ischemia			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.13	0.56, 2.26	0.74

¹OR = Odds Ratio, CI = Confidence Interval

Intestinal Ischemia			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.07	0.50, 2.30	0.87
AGECAT			
<50	—	—	
>79	0.41	0.07, 2.37	0.32
50-59	0.49	0.06, 4.15	0.51
60-69	0.36	0.06, 2.15	0.26
70-79	0.45	0.08, 2.48	0.36
GENDER			
female	—	—	
male	0.62	0.37, 1.03	0.063
PREOP_SMOKING			
No	—	—	
Yes	0.86	0.43, 1.70	0.66
PRIOR_AORSURG			
Both	—	—	
Endo	1.94	0.23, 16.6	0.55
None	1.84	0.22, 15.8	0.58
Open	1.62	0.16, 15.9	0.68
PRIOR_CHF			
No	—	—	
Yes	1.05	0.53, 2.08	0.89
PREOP_DIALYSIS			

¹OR = Odds Ratio, CI = Confidence Interval

Intestinal Ischemia			
Characteristic	OR ¹	95% CI ¹	p-value
No	—	—	
Yes	0.69	0.09, 5.12	0.71
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.48	0.06, 3.66	0.48
Dissection	0.71	0.14, 3.71	0.69
PAU/IMH	1.28	0.16, 10.6	0.82
extent			
Juxtarenal	—	—	
No	0.94	0.53, 1.69	0.85

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_CEREBROSX: Post-op Cerebrovascular Stroke:

Also account for ARMNECK_ACCESS

Post-op Cerebrovascular Stroke			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	3.42	1.83, 6.38	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Post-op Cerebrovascular Stroke			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.81	0.91, 3.60	0.091
AGECAT			
<50	—	—	
>79	0.51	0.08, 3.29	0.48
50-59	0.79	0.13, 4.95	0.81
60-69	0.23	0.04, 1.45	0.12
70-79	0.62	0.11, 3.46	0.58
GENDER			
female	—	—	
male	0.49	0.26, 0.90	0.023
PREOP_SMOKING			
No	—	—	
Yes	0.73	0.34, 1.58	0.43
PRIOR_AORSURG			
Both	—	—	
Endo	1.20	0.24, 6.05	0.83
None	1.39	0.28, 6.92	0.69
Open	0.95	0.15, 5.88	0.95
PRIOR_CHF			
No	—	—	
Yes	2.14	1.07, 4.31	0.032

¹OR = Odds Ratio, CI = Confidence Interval

Post-op Cerebrovascular Stroke			
Characteristic	OR ¹	95% CI ¹	p-value
PREOP_DIALYSIS			
No	—	—	
Yes	1.97	0.46, 8.55	0.36
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	3.44	1.03, 11.6	0.045
Dissection	2.56	0.57, 11.5	0.22
PAU/IMH extent	1.46	0.21, 10.2	0.70
Juxtarenal	—	—	
No	1.44	0.68, 3.04	0.34
ARMNECK_ACCESS			
No	—	—	
Yes	2.31	1.26, 4.25	0.007

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_SPINAL_ISCHEMIA: Post-op Spinal Ischemia

Also account for POSTOP_SPINALDRAIN and OCCLUDED_CELIAC

Post-op Spinal Ischemia			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.56	1.70, 3.86	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Post-op Spinal Ischemia			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.61	1.00, 2.60	0.050
AGECAT			
<50	—	—	
>79	2.19	0.27, 17.7	0.46
50-59	1.99	0.23, 17.3	0.53
60-69	1.63	0.21, 13.0	0.64
70-79	2.13	0.27, 16.6	0.47
GENDER			
female	—	—	
male	0.88	0.58, 1.36	0.58
PREOP_SMOKING			
No	—	—	
Yes	1.22	0.64, 2.32	0.55
PRIOR_AORSURG			
Both	—	—	
Endo	1.23	0.52, 2.95	0.64
None	1.46	0.63, 3.40	0.37
Open	1.47	0.60, 3.63	0.40
PRIOR_CHF			
No	—	—	
Yes	1.06	0.61, 1.83	0.84

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Post-op Spinal Ischemia		
	OR ¹	95% CI ¹	p-value
PREOP_DIALYSIS			
No	—	—	
Yes	1.16	0.48, 2.83	0.75
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.98	0.43, 2.25	0.96
Dissection	1.08	0.36, 3.26	0.89
PAU/IMH extent	0.44	0.05, 3.63	0.45
Juxtarenal	—	—	
No	1.01	0.65, 1.57	0.97
POSTOP_SPINALDRAIN			
No	—	—	
Yes	19.0	11.1, 32.6	<0.001
OCCLUDED_CELIAC			
FALSE	—	—	
TRUE	1.57	0.83, 2.99	0.17

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_LOS: Length of stay in days between surgery date and discharge date

	Length of stay in days between surgery date and discharge date		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	3.39	2.69, 4.29	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

	Length of stay in days between surgery date and discharge date		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.46	1.91, 3.17	<0.001
AGECAT			
<50	—	—	
>79	2.28	0.88, 5.88	0.089
50-59	1.93	0.72, 5.21	0.19
60-69	1.55	0.60, 3.96	0.36
70-79	2.49	0.98, 6.35	0.055
GENDER			
female	—	—	
male	0.62	0.50, 0.75	<0.001
PREOP_SMOKING			
No	—	—	
Yes	0.97	0.74, 1.28	0.82
PRIOR_AORSURG			
Both	—	—	
Endo	1.26	0.73, 2.19	0.41
None	0.96	0.57, 1.62	0.88
Open	1.36	0.76, 2.41	0.30
PRIOR_CHF			
No	—	—	
Yes	1.41	1.11, 1.80	0.005

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Length of stay in days between surgery date and discharge date		
	OR ¹	95% CI ¹	p-value
PREOP_DIALYSIS			
No	—	—	
Yes	1.74	1.00, 3.00	0.048
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	1.59	0.96, 2.63	0.069
Dissection	1.96	1.11, 3.45	0.020
PAU/IMH extent	1.82	0.90, 3.67	0.094
Juxtarenal	—	—	
No	1.97	1.60, 2.44	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_COMPLICATIONS: Any Complications Post-op

	Any Complications Post-op		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.74	1.37, 2.21	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

	Any Complications Post-op		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.44	1.12, 1.86	0.004
AGECAT			
<50	—	—	
>79	1.17	0.46, 2.97	0.74
50-59	1.29	0.49, 3.42	0.61
60-69	0.97	0.39, 2.41	0.94
70-79	1.32	0.53, 3.31	0.55
GENDER			
female	—	—	
male	0.65	0.54, 0.78	<0.001
PREOP_SMOKING			
No	—	—	
Yes	1.01	0.79, 1.31	0.91
PRIOR_AORSURG			
Both	—	—	
Endo	0.95	0.56, 1.60	0.84
None	1.00	0.60, 1.65	0.99
Open	1.41	0.82, 2.42	0.21
PRIOR_CHF			
No	—	—	
Yes	1.28	1.04, 1.58	0.018
PREOP_DIALYSIS			

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Any Complications Post-op		
	OR ¹	95% CI ¹	p-value
No	—	—	
Yes	1.25	0.73, 2.15	0.41
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	1.32	0.82, 2.12	0.26
Dissection	0.94	0.51, 1.73	0.84
PAU/IMH	0.77	0.35, 1.70	0.52
extent			
Juxtarenal	—	—	
No	1.47	1.21, 1.78	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

RETX_R_RTOR: Re-intervention

Re-intervention			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.86	1.33, 2.62	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Re-intervention			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.48	1.04, 2.09	0.029
AGECAT			
<50	—	—	
>79	0.95	0.28, 3.27	0.93
50-59	1.41	0.38, 5.19	0.60
60-69	0.70	0.20, 2.44	0.58
70-79	0.94	0.28, 3.14	0.92
GENDER			
female	—	—	
male	0.59	0.45, 0.78	<0.001
PREOP_SMOKING			
No	—	—	
Yes	1.29	0.81, 2.05	0.28
PRIOR_AORSURG			
Both	—	—	
Endo	0.95	0.43, 2.09	0.90
None	1.01	0.47, 2.16	0.99
Open	0.67	0.27, 1.63	0.37
PRIOR_CHF			
No	—	—	
Yes	1.21	0.86, 1.72	0.27
PREOP_DIALYSIS			

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Re-intervention		
	OR ¹	95% CI ¹	p-value
No	—	—	
Yes	0.75	0.31, 1.81	0.52
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	1.68	0.88, 3.19	0.12
Dissection	0.64	0.23, 1.78	0.39
PAU/IMH	1.77	0.70, 4.47	0.23
extent			
Juxtarenal	—	—	
No	1.56	1.15, 2.12	0.004

¹OR = Odds Ratio, CI = Confidence Interval

BRANCH_POST: Post-treatment Status of All Branches

Post-treatment Status of All Branches			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.28	1.78, 2.91	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Post-treatment Status of All Branches			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.64	1.24, 2.16	<0.001
AGECAT			
<50	—	—	
>79	0.78	0.32, 1.90	0.58
50-59	1.00	0.39, 2.58	>0.99
60-69	0.95	0.40, 2.27	0.91
70-79	0.90	0.38, 2.14	0.81
GENDER			
female	—	—	
male	0.47	0.39, 0.58	<0.001
PREOP_SMOKING			
No	—	—	
Yes	1.08	0.79, 1.48	0.63
PRIOR_AORSURG			
Both	—	—	
Endo	0.79	0.46, 1.35	0.38
None	0.52	0.31, 0.86	0.011
Open	0.69	0.39, 1.23	0.21
PRIOR_CHF			
No	—	—	
Yes	1.43	1.12, 1.82	0.004

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Post-treatment Status of All Branches		
	OR ¹	95% CI ¹	p-value
PREOP_DIALYSIS			
No	—	—	
Yes	7.90	4.70, 13.3	<0.001
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.77	0.43, 1.37	0.37
Dissection	1.57	0.90, 2.74	0.12
PAU/IMH extent	0.45	0.16, 1.24	0.12
Juxtarenal	—	—	
No	1.56	1.25, 1.93	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

For the following variables, we need to merge age group “<50” and “50-59” so that there’s no 0 patients on one level.

POSTOP_DIALYSIS: Post-op Dialysis

Also account for OCCLUDED_RENAL

Post-op Dialysis			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.42	1.44, 4.07	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Post-op Dialysis			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.78	1.01, 3.13	0.047
AGECAT			
<59	—	—	
60-69	0.53	0.22, 1.27	0.16
70-79	0.40	0.17, 0.97	0.042
>79	0.39	0.15, 1.01	0.053
GENDER			
female	—	—	
male	0.62	0.38, 1.00	0.049
PREOP_SMOKING			
No	—	—	
Yes	1.31	0.60, 2.86	0.49
PRIOR_AORSURG			
Both	—	—	
Endo	1.04	0.26, 4.10	0.95
None	0.82	0.22, 3.04	0.76
Open	0.58	0.12, 2.80	0.50
PRIOR_CHF			
No	—	—	
Yes	0.74	0.37, 1.45	0.38
PREOP_DIALYSIS			

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Post-op Dialysis		
	OR ¹	95% CI ¹	p-value
No	—	—	
Yes	0.00	0.00, 0.00	<0.001
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.33	0.04, 2.96	0.32
Dissection	1.74	0.52, 5.82	0.37
PAU/IMH	2.33	0.51, 10.6	0.27
extent			
Juxtarenal	—	—	
No	0.99	0.58, 1.70	0.98
OCCLUDED_RENAL			
FALSE	—	—	
TRUE	4.82	2.33, 9.95	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_LEGEMBO: Leg Ischemia/Embol

	Leg Ischemia/Embol		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.97	1.13, 3.46	0.017

¹OR = Odds Ratio, CI = Confidence Interval

	Leg Ischemia/Embol		
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	1.74	0.94, 3.22	0.078
AGECAT			
<59	—	—	
60-69	0.80	0.28, 2.27	0.68
70-79	0.80	0.29, 2.23	0.67
>79	0.67	0.23, 2.00	0.48
GENDER			
female	—	—	
male	0.43	0.28, 0.67	<0.001
PREOP_SMOKING			
No	—	—	
Yes	2.61	0.98, 6.96	0.056
PRIOR_AORSURG			
Both	—	—	
Endo	139,944,960,652,840		
None	162,086,767,451,715		
Open	288,216,346,513,340		
PRIOR_CHF			
No	—	—	
Yes	0.85	0.44, 1.64	0.63
PREOP_DIALYSIS			
No	—	—	

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Leg Ischemia/Embol		
	OR ¹	95% CI ¹	p-value
Yes	0.58	0.08, 4.19	0.59
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.87	0.20, 3.86	0.86
Dissection	0.60	0.08, 4.59	0.62
PAU/IMH	2.28	0.49, 10.6	0.29
extent			
Juxtarenal	—	—	
No	1.01	0.61, 1.66	0.98

¹OR = Odds Ratio, CI = Confidence Interval

POSTOP_RESPIRATORY: Post-op Respiratory

Post-op Respiratory			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.65	1.87, 3.75	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Post-op Respiratory			
Characteristic	OR ¹	95% CI ¹	p-value
PRESENTATION			
Asymptomatic	—	—	
Symptomatic	2.17	1.47, 3.21	<0.001
AGECAT			
<59	—	—	
60-69	0.94	0.44, 1.98	0.86
70-79	1.06	0.50, 2.25	0.87
>79	0.94	0.42, 2.06	0.87
GENDER			
female	—	—	
male	0.66	0.47, 0.92	0.014
PREOP_SMOKING			
No	—	—	
Yes	1.08	0.68, 1.72	0.75
PRIOR_AORSURG			
Both	—	—	
Endo	0.59	0.27, 1.30	0.19
None	0.55	0.26, 1.16	0.11
Open	0.67	0.28, 1.60	0.36
PRIOR_CHF			
No	—	—	
Yes	1.23	0.82, 1.84	0.31
PREOP_DIALYSIS			
No	—	—	

¹OR = Odds Ratio, CI = Confidence Interval

Characteristic	Post-op Respiratory		
	OR ¹	95% CI ¹	p-value
Yes	0.78	0.27, 2.23	0.64
PATHOLOGY			
Aneurysm	—	—	
Aneurysm from dissection	0.60	0.23, 1.55	0.29
Dissection	0.83	0.29, 2.38	0.73
PAU/IMH	1.14	0.33, 4.02	0.83
extent			
Juxtarenal	—	—	
No	1.74	1.25, 2.41	<0.001

¹OR = Odds Ratio, CI = Confidence Interval

Code Appendix

```
knitr::opts_chunk$set(echo = FALSE,message = FALSE,warning = FALSE)
library(tidyverse)
library(table1)
library(dplyr)
library(geepack)
library(gtsummary)
library(broom.mixed)
## ----- working directories for Lily -----
wd_lily = '/Users/hanyiwang/Desktop/Comparative-analysis-of-treatments-of-CAA'
path_lily = c("../data/TEVAR_PROC.csv")

## ----- read data -----
setwd(wd_lily)
TEVAR_PROC = read.csv(path_lily)
# ### ----- Correlation matrix -----
# library(corrplot)
# library(tidyverse)
# library(caret)
# library(ggcorrplot)
#
# matrix <- PROC %>%
#   select_if(is.numeric) %>% subset(., select = -1)%>%
#   cor(.,use = "complete")
#
# corrplot(matrix, method="number")
#
# #select dataset that column are not numeric
# matrix <- PROC %>% select_if(negate(is.numeric))
#
# # find out the variables
# lapply(matrix[,], unique)
# sapply(lapply(matrix, unique), length)
#
# #select if more than one variable
# y<-matrix %>%select_if(function(col) length(unique(col))==2)
#
# # Convert all columns to factor
# data3 <- as.data.frame(unclass(y),
#                         stringsAsFactors = TRUE)
#
# model.matrix(~0 +., data=data3) %>%
#   cor(use="pairwise.complete.obs") %>%
#   ggcorrplot(show.diag = F, type="lower", lab=TRUE, lab_size=2)
#
# str(y)
## ----- modify dataset -----
TEVAR_PROC = TEVAR_PROC %>%
  mutate(extent = factor(extent,levels = c("Juxtarenal AAA","Type 1 TAAA","Type 2 TAAA","Type 3 TAAA","
  mutate(TOTAL_LOS = case_when(TOTAL_LOS>7 ~ '>7',
                                TOTAL_LOS<=7 ~ '<=7')) %>%
  mutate(ICUSTAY = case_when(ICUSTAY>4 ~ '>4',
```

```

        ICUSTAY<=4 ~ '<=4')) %>%
mutate(POSTOP_LOS = case_when(POSTOP_LOS>7 ~ '>7',
                             POSTOP_LOS<=7 ~ '<=7')) %>%

mutate(TOTAL_LOS = as.numeric(TOTAL_LOS != "<=7") ) %>%
mutate(ICUSTAY = as.numeric(ICUSTAY != "<=4") ) %>%
mutate(POSTOP_LOS = as.numeric(POSTOP_LOS != "<=7") ) %>%

mutate(POSTOP_INTISCH = as.numeric(POSTOP_INTISCH == "Yes")) %>%
mutate(POSTOP_CEREBROX = as.numeric(POSTOP_CEREBROX == "Yes")) %>%
mutate(POSTOP_SPINAL_ISCHEMIA = as.numeric(POSTOP_SPINAL_ISCHEMIA == "Yes")) %>%
mutate(POSTOP_DIALYSIS = as.numeric(POSTOP_DIALYSIS == "Yes")) %>%
mutate(POSTOP_COMPLICATIONS = as.numeric(POSTOP_COMPLICATIONS == "Yes")) %>%
mutate(POSTOP_LEGEMBO = as.numeric(POSTOP_LEGEMBO == "Yes")) %>%
mutate(POSTOP_RESPIRATORY = as.numeric(POSTOP_RESPIRATORY == "Yes")) %>%
mutate(RETX_R_RTOR = as.numeric(RETX_R_RTOR == "Yes")) %>%
mutate(BRANCH_POST = as.numeric(BRANCH_POST == "Yes"))

# remove missing values
TEVAR_PROC = subset(TEVAR_PROC, !is.na(extent))

## ----- change class of variables -----
names <- c('CENTERID', 'AGECAT', 'GENDER', 'PREOP_SMOKING', 'PRIOR_AORSURG', 'PRIOR_CHF', 'PREOP_DIALYSIS')
TEVAR_PROC[,names] <- lapply(TEVAR_PROC[,names] , factor)

## ----- Length of stay in days between admission date and discharge date -----

#table(PROC$extent)
#PROCNM[is.na(PROCNM) | PROCNM=="Inf"] = NA
# repeated patients, need id, geeglm needs complete data, extent is not complete

## ----- unadjusted -----
LOS1<-geeglm(TOTAL_LOS~ PRESENTATION, data=TEVAR_PROC, family=binomial(link="logit"), id = CENTERID, corstr = "independence")
t1<-LOS1 %>%
  tbl_regression(exponentiate=TRUE,
                tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**TOTAL_LOS Length of Stay in days**")%>%as_flex_table()

## ----- adjusted -----
LOS2<-geeglm(TOTAL_LOS ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS,
              data=TEVAR_PROC, family=binomial(link="logit"),id = CENTERID, corstr = "independence")
t2<-LOS2 %>%
  tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2),tab_spanner = "**TOTAL_LOS Length of Stay in days**")%>%as_flex_table()

```

```

#the difference in mean number of sessions attended comparing treatment to control

#adjust<-c('PRESENTATION','AGECAT', 'GENDER', 'PREOP_SMOKING','PRIOR_AORSURG', 'PRIOR_CHF','PATHOLOGY')
#PROC %>%
#  tbl_uvregression(
#    y = TOTAL_LOS,
#    x = PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+PATHOLOGY+NUM_TREATED_BRANCHES,
#    method = geepack::geeglm,
#    method.args = list(id = X+CENTERID, corstr = "independence"),
#    include = all_of(adjust)
#  ) %>%
#  as_kable()
## ----- ICU Stay -----

## ----- unadjusted -----
ICU1<-geeglm(ICUSTAY~ PRESENTATION, data=TEVAR_PROC, family=binomial(link="logit"), id = CENTERID, corstr="independence")

#summary(LOS2)
#summary(ICU1)

#ICU1<-geeglm(ICUSTAY ~ PRESENTATION, data=PROCNM,id = CENTERID, corstr = "independence")
t1<-ICU1 %>%tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                          pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**ICU Stay**")%>%as_flex_table()

## ----- adjusted -----
ICU2<-geeglm(ICUSTAY ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+
              , family=binomial(link="logit"), data=TEVAR_PROC, id = CENTERID, corstr = "independence")
t2<- ICU2 %>%tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                          pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2),tab_spanner = "**ICU Stay**")%>%as_flex_table()
## ----- Transfusion # Units PRBC -----
## ----- unadjusted -----
POSTOP_PRBC1<-geeglm(POSTOP_PRBC ~ PRESENTATION, data=TEVAR_PROC,id = CENTERID, corstr = "independence")
t1<-POSTOP_PRBC1 %>%tbl_regression(tidy_fun = broom.mixed::tidy,
                                  pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Transfusion # Units PRBC**")%>%as_flex_table()

## ----- adjusted -----
# POSTOP_PRBC2<-geeglm(POSTOP_PRBC ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+
# t2<- POSTOP_PRBC2 %>%tbl_regression(tidy_fun = broom.mixed::tidy,
#                                   pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
#   bold_p(t = 0.05)
# tbl_merge(tbls = list(t2),tab_spanner = "**Transfusion # Units PRBC**")%>%as_flex_table()

```

```

## ----- Post-op Intestinal Ischemia -----
## ----- unadjusted -----
POSTOP_INTISCH1<-geeglm(POSTOP_INTISCH ~ PRESENTATION, family=binomial(link="logit"), data=TEVAR_PROC,i

t1<-POSTOP_INTISCH1 %>%tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                                     pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Intestinal Ischemia**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_INTISCH2<-geeglm(POSTOP_INTISCH ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CH
, data=TEVAR_PROC,family=binomial(link="logit"), id = CENTERID, corstr = "independe
t2<- POSTOP_INTISCH2 %>%tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                                     pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner = "**Intestinal Ischemia**")%>%as_flex_table()

# model<-geeglm(as.numeric(POSTOP_DIALYSIS)~PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+PRIOR_AORSURG+PRIOR
# model<-geeglm(outcome~predictor+confounder, family=binomial(link = "logit"),
# data=na.omit(data), corstr='ar1', id=id, std.err="san.se")
# TOTAL_LOS
# ICUSTAY
# POSTOP_PRBC
# POSTOP_HIGHCREAT
# POSTOP_INTISCH
## ----- Post-op Cerebrovascular Stroke -----
## ----- unadjusted -----
POSTOP_CEREBROX1 = geeglm(POSTOP_CEREBROX ~ PRESENTATION,
                          data=TEVAR_PROC, family=binomial(link="logit"),
                          id = CENTERID, corstr = "independence")

t1 = POSTOP_CEREBROX1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Post-op Cerebrovascular Stroke**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_CEREBROX2 = geeglm(POSTOP_CEREBROX ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent+ARMNECK_ACCESS,
family=binomial(link="logit"), data=TEVAR_PROC,
id = CENTERID, corstr = "independence")

t2 = POSTOP_CEREBROX2 %>%
  tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

```

```

tbl_merge(tbls = list(t2),tab_spanner = "**Post-op Cerebrovascular Stroke**")%>%as_flex_table()
## ----- Post-op Spinal Ischemia -----
## ----- unadjusted -----
POSTOP_SPINAL_ISCHEMIA1 = geeglm(POSTOP_SPINAL_ISCHEMIA ~ PRESENTATION,
                                data=TEVAR_PROC,family=binomial(link="logit"),
                                id = CENTERID, corstr = "independence")

t1 = POSTOP_SPINAL_ISCHEMIA1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Post-op Spinal Ischemia**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_SPINAL_ISCHEMIA2 = geeglm(POSTOP_SPINAL_ISCHEMIA ~ PRESENTATION+AGECAT+GENDER+
                                PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+
                                PREOP_DIALYSIS+PATHOLOGY+extent+
                                POSTOP_SPINALDRAIN+OCCLUDED_CELIAC,
                                family=binomial(link="logit"), data=TEVAR_PROC,
                                id = CENTERID, corstr = "independence")

t2 = POSTOP_SPINAL_ISCHEMIA2 %>%
  tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2),tab_spanner = "**Post-op Spinal Ischemia**")%>%as_flex_table()

## ----- Length of stay in days between surgery date and discharge date -----
## ----- unadjusted -----
POSTOP_LOS1 = geeglm(POSTOP_LOS~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)

t1 = POSTOP_LOS1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Length of stay in days between surgery date and discharge date**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_LOS2 = geeglm(POSTOP_LOS ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                    PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                    family=binomial(link="logit"), data=TEVAR_PROC,
                    id = CENTERID, corstr = "independence")

t2 = POSTOP_LOS2 %>%
  tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,

```

```

        pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
bold_p(t = 0.05)

tbl_merge(tbls = list(t2),tab_spanner = "**Length of stay in days between surgery date and discharge date**")%>%
## ----- Any Complications Post-op -----
## ----- unadjusted -----
POSTOP_COMPLICATIONS1 = geeglm(POSTOP_COMPLICATIONS~ PRESENTATION,
                                data=TEVAR_PROC, family=binomial(link="logit"),
                                id = CENTERID, corstr = "independence")

t1 = POSTOP_COMPLICATIONS1 %>%
tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
               pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Any Complications Post-op**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_COMPLICATIONS2 = geeglm(POSTOP_COMPLICATIONS ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                                PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                                family=binomial(link="logit"), data=TEVAR_PROC,
                                id = CENTERID, corstr = "independence")

t2 = POSTOP_COMPLICATIONS2 %>%
tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
               pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
bold_p(t = 0.05)

tbl_merge(tbls = list(t2),tab_spanner = "**Any Complications Post-op**")%>%as_flex_table()
## ----- Re-intervention -----
## ----- unadjusted -----
RETX_R_RTOR1 = geeglm(RETX_R_RTOR~ PRESENTATION,
                      data=TEVAR_PROC, family=binomial(link="logit"),
                      id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)

t1 = RETX_R_RTOR1 %>%
tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
               pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
bold_p(t = 0.05)

tbl_merge(tbls = list(t1),tab_spanner = "**Re-intervention**")%>%as_flex_table()

## ----- adjusted -----
RETX_R_RTOR2 = geeglm(RETX_R_RTOR ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                      PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                      family=binomial(link="logit"), data=TEVAR_PROC,
                      id = CENTERID, corstr = "independence")

t2 = RETX_R_RTOR2 %>%
tbl_regression(exponentiate=TRUE,tidy_fun = broom.mixed::tidy,
               pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
bold_p(t = 0.05)

```



```

tbl_merge(tbls = list(t2), tab_spanner = "**Re-intervention**") %>% as_flex_table()
## ----- Post-treatment Status of All Branches -----
## ----- unadjusted -----
BRANCH_POST1 = geeglm(BRANCH_POST ~ PRESENTATION,
                      data = TEVAR_PROC, family = binomial(link = "logit"),
                      id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)

t1 = BRANCH_POST1 %>%
  tbl_regression(exponentiate = TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2)) %>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1), tab_spanner = "**Post-treatment Status of All Branches**") %>% as_flex_table()

## ----- adjusted -----
BRANCH_POST2 = geeglm(BRANCH_POST ~ PRESENTATION + AGE CAT + GENDER + PREOP_SMOKING +
                      PRIOR_AORSURG + PRIOR_CHF + PREOP_DIALYSIS + PATHOLOGY + extent,
                      family = binomial(link = "logit"), data = TEVAR_PROC,
                      id = CENTERID, corstr = "independence")

t2 = BRANCH_POST2 %>%
  tbl_regression(exponentiate = TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2)) %>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2), tab_spanner = "**Post-treatment Status of All Branches**") %>% as_flex_table()
## ----- modify dataset -----
TEVAR_PROC = TEVAR_PROC %>%
  mutate(AGE CAT = factor(AGE CAT, levels = c('<50', '50-59', '60-69', '70-79', '>79'),
                        labels = c('<59', '<59', '60-69', '70-79', '>79')))

## ----- Post-op Dialysis -----
# table1(~ AGE CAT | factor(POSTOP_DIALYSIS), data = subset(TEVAR_PROC, !is.na(POSTOP_DIALYSIS)))

## ----- unadjusted -----
POSTOP_DIALYSIS1 = geeglm(POSTOP_DIALYSIS ~ PRESENTATION,
                        data = TEVAR_PROC, family = binomial(link = "logit"),
                        id = CENTERID, corstr = "independence")

t1 = POSTOP_DIALYSIS1 %>%
  tbl_regression(exponentiate = TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2)) %>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1), tab_spanner = "**Post-op Dialysis**") %>% as_flex_table()

## ----- adjusted -----
POSTOP_DIALYSIS2 = geeglm(POSTOP_DIALYSIS ~ PRESENTATION + AGE CAT + GENDER +
                        PREOP_SMOKING + PRIOR_AORSURG + PRIOR_CHF +
                        PREOP_DIALYSIS + PATHOLOGY + extent +
                        OCCLUDED_RENAL,
                        family = binomial(link = "logit"), data = TEVAR_PROC,

```

```

      id = CENTERID, corstr = "independence")

t2 = POSTOP_DIALYSIS2 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                 pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2), tab_spanner = "**Post-op Dialysis**")%>%as_flex_table()

## ----- Leg Ischemia/Emboli -----
# table1(~ AGECAT | factor(POSTOP_LEGEMBO), data=subset(TEVAR_PROC, !is.na(POSTOP_LEGEMBO)))
## ----- unadjusted -----
POSTOP_LEGEMBO1 = geeglm(POSTOP_LEGEMBO~ PRESENTATION,
                        data=TEVAR_PROC, family=binomial(link="logit"),
                        id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)

t1 = POSTOP_LEGEMBO1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                 pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1), tab_spanner = "**Leg Ischemia/Embol**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_LEGEMBO2 = geeglm(POSTOP_LEGEMBO ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                        PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                        family=binomial(link="logit"), data=TEVAR_PROC,
                        id = CENTERID, corstr = "independence")

t2 = POSTOP_LEGEMBO2 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                 pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2), tab_spanner = "**Leg Ischemia/Embol**")%>%as_flex_table()
## ----- Post-op Respiratory -----
# table1(~ AGECAT | factor(POSTOP_RESPIRATORY), data=subset(TEVAR_PROC, !is.na(POSTOP_RESPIRATORY)))
## ----- unadjusted -----
POSTOP_RESPIRATORY1 = geeglm(POSTOP_RESPIRATORY~ PRESENTATION,
                            data=TEVAR_PROC, family=binomial(link="logit"),
                            id = CENTERID, corstr = "independence")

t1 = POSTOP_RESPIRATORY1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                 pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t1), tab_spanner = "**Post-op Respiratory**")%>%as_flex_table()

## ----- adjusted -----
POSTOP_RESPIRATORY2 = geeglm(POSTOP_RESPIRATORY ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+

```

```

        PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
        family=binomial(link="logit"), data=TEVAR_PROC,
        id = CENTERID, corstr = "independence")

t2 = POSTOP_RESPIRATORY2 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy,
                pvalue_fun = function(x) style_pvalue(x, digits = 2))%>%
  bold_p(t = 0.05)

tbl_merge(tbls = list(t2), tab_spanner = "**Post-op Respiratory**")%>%as_flex_table()

```