

preprocessing

Jennifer Ci, Thu Vu, Lily Hanyi Wang

2022-10-26

Dataset TEVAR_PROC_07 to start with.

population of interest: the asymptomatic and symptomatics groups.

PRESENTATION: 0 = Asymptomatic, 1 = Symptomatic, 2 = Rupture

Is this the correct variable?

What does rupture mean?

Should we just delete data missing symptom info?

	Overall
	(N=19564)
PRESENTATION	
Asymptomatic	9272 (47.4%)
Symptomatic	6624 (33.9%)
Rupture	1168 (6.0%)
Missing	2500 (12.8%)

potential outcome variables

Are these possible outcomes?

TOTAL_LOS: Length of Stay in days calculated by DISCHARGE_DT - ADMIT_DT \

POSTOP_LOS: Length of Stay in days calculated by DISCHARGE_DT - SURGERY_DT

	Asymptomatic	Symptomatic	Rupture	Overall
	(N=9272)	(N=6624)	(N=1168)	(N=19564)
factor(DEAD)				
0	8103 (87.4%)	5570 (84.1%)	741 (63.4%)	16254 (83.1%)
1	1168 (12.6%)	1053 (15.9%)	427 (36.6%)	3295 (16.8%)
Missing	1 (0.0%)	1 (0.0%)	0 (0%)	15 (0.1%)
PROC_SURVIVALDAYS				
Mean (SD)	719 (725)	657 (730)	522 (709)	798 (883)
Median [Min, Max]	456 [-355, 3360]	407 [0, 3200]	215 [0, 3410]	454 [-355, 3970]
Missing	1 (0.0%)	0 (0%)	0 (0%)	1 (0.0%)

patient condition variables:

AGECAT: 1 = <40, 2 = 40-49, 3 = 50-59, 4 = 60-69, 5 = 70-79, 6 = 80-89, 7 = >89

GENDER: 1 = Male, 2 = Female

PRIOR_CVD: 0 = None,1 = hx stroke, asymptomatic,2 = hx stroke, minor deficit,3 = hx stroke, major deficit

PRIOR_CAD: 0 = None,1 = hx MI but no sx,2 = Stable angina,3 = Unstable angina or MI < 6 mos (retired since 09/12/2012),4 = MI < 6 mos,5 = Unstable angina

PRIOR_CHF: 0 = None,1 = Asymp, hx CHF,2 = Mild,3 = Moderate,4 = Severe

COPD: 0 = No,1 = Not Treated,2 = On Meds,3 = On Home Oxygen

DIABETES: 0 = None,1 = Diet,2 = Non-insulin Meds,3 = Insulin

HTN: History of hypertension; 0 = No, 1 = Yes ($\geq 140/90$ or history) (retired since 11/15/2016), 2 = Yes, controlled [added on 04/13/2020] , 3 = Yes, uncontrolled [added on 04/13/2020]

PREOP_SMOKING: 0 = Never,1 = Prior,2 = Current

PRIOR_AORSURG: Any aortic procedures performed on a separate date prior to the index procedure; 0 = None,1 = Open,2 = Endo,3 = Both,4 = Other (retired since 09/30/2014)

PATHOLOGY: 1 = Aneurysm,2 = Dissection,3 = Aneurysm from dissection,4 = Trauma,5 = Penetrating Ulcer (PAU),6 = Intramural Hematoma (IMH),7 = PAU with IMH,8 = Aortic Thrombus,9 = Other (Retired) (retired since 09/30/2014),10 = Aorto-esophageal Fistula (Retired) (retired since 09/30/2014),11 = Aorto-bronchial Fistula (Retired) (retired since 09/30/2014).

Type of TAA?

PREOP_MAXAADI: “Maximum total aortic diameter within the last 6 months (including true and false lumen in dissection) within diseased segment being treated, measured at right angle to centerline, ideally with 3-D imaging software. Min/max range: 16 to 190 mm.”

surgery year?

	Asymptomatic	Symptomatic	Rupture	Overall
	(N=9272)	(N=6624)	(N=1168)	(N=19564)
GENDER				
male	6356 (68.6%)	3988 (60.2%)	736 (63.0%)	12667 (64.7%)
female	2916 (31.4%)	2636 (39.8%)	432 (37.0%)	6897 (35.3%)
AGE				
Mean (SD)	70.3 (11.5)	62.3 (16.1)	64.7 (19.5)	66.8 (14.6)
Median [Min, Max]	72.0 [0, 90.0]	65.0 [0, 90.0]	71.0 [0, 90.0]	70.0 [0, 90.0]
AGECAT				
<40	221 (2.4%)	662 (10.0%)	170 (14.6%)	1198 (6.1%)
40-49	252 (2.7%)	698 (10.5%)	71 (6.1%)	1185 (6.1%)
50-59	809 (8.7%)	1181 (17.8%)	123 (10.5%)	2411 (12.3%)
60-69	2423 (26.1%)	1633 (24.7%)	197 (16.9%)	4826 (24.7%)
70-79	3795 (40.9%)	1552 (23.4%)	323 (27.7%)	6543 (33.4%)
80-89	1663 (17.9%)	814 (12.3%)	241 (20.6%)	3138 (16.0%)
>89	109 (1.2%)	84 (1.3%)	43 (3.7%)	263 (1.3%)
factor(PREOP_SMOKING)				
0	1795 (19.4%)	2266 (34.2%)	458 (39.2%)	5132 (26.2%)
1	4818 (52.0%)	1975 (29.8%)	330 (28.3%)	8148 (41.6%)
2	2653 (28.6%)	2350 (35.5%)	340 (29.1%)	6145 (31.4%)
Missing	6 (0.1%)	33 (0.5%)	40 (3.4%)	139 (0.7%)
factor(PRIOR_CVD)				
0	8196 (88.4%)	5949 (89.8%)	1044 (89.4%)	15348 (78.5%)
1	661 (7.1%)	401 (6.1%)	65 (5.6%)	1135 (5.8%)
2	329 (3.5%)	197 (3.0%)	24 (2.1%)	551 (2.8%)
3	76 (0.8%)	55 (0.8%)	18 (1.5%)	150 (0.8%)
Missing	10 (0.1%)	22 (0.3%)	17 (1.5%)	2380 (12.2%)

	Asymptomatic	Symptomatic	Rupture	Overall
factor(PRIOR_CAD)				
0	7072 (76.3%)	5571 (84.1%)	991 (84.8%)	15560 (79.5%)
1	1627 (17.5%)	677 (10.2%)	111 (9.5%)	2777 (14.2%)
2	454 (4.9%)	213 (3.2%)	30 (2.6%)	798 (4.1%)
3	0 (0%)	0 (0%)	0 (0%)	7 (0.0%)
4	66 (0.7%)	68 (1.0%)	14 (1.2%)	168 (0.9%)
5	44 (0.5%)	73 (1.1%)	10 (0.9%)	153 (0.8%)
Missing	9 (0.1%)	22 (0.3%)	12 (1.0%)	101 (0.5%)
factor(PRIOR_CHF)				
0	7982 (86.1%)	5885 (88.8%)	1009 (86.4%)	17044 (87.1%)
1	786 (8.5%)	414 (6.3%)	78 (6.7%)	1432 (7.3%)
2	297 (3.2%)	166 (2.5%)	37 (3.2%)	574 (2.9%)
3	172 (1.9%)	103 (1.6%)	23 (2.0%)	329 (1.7%)
4	30 (0.3%)	36 (0.5%)	6 (0.5%)	90 (0.5%)
Missing	5 (0.1%)	20 (0.3%)	15 (1.3%)	95 (0.5%)
factor(COPD)				
0	6128 (66.1%)	5141 (77.6%)	898 (76.9%)	13976 (71.4%)
1	797 (8.6%)	432 (6.5%)	75 (6.4%)	1487 (7.6%)
2	1867 (20.1%)	837 (12.6%)	143 (12.2%)	3211 (16.4%)
3	474 (5.1%)	195 (2.9%)	41 (3.5%)	801 (4.1%)
Missing	6 (0.1%)	19 (0.3%)	11 (0.9%)	89 (0.5%)
factor(DIABETES)				
0	7698 (83.0%)	5614 (84.8%)	978 (83.7%)	16316 (83.4%)
1	371 (4.0%)	267 (4.0%)	53 (4.5%)	790 (4.0%)
2	921 (9.9%)	491 (7.4%)	75 (6.4%)	1698 (8.7%)
3	277 (3.0%)	233 (3.5%)	49 (4.2%)	671 (3.4%)
Missing	5 (0.1%)	19 (0.3%)	13 (1.1%)	89 (0.5%)
factor(HTN)				
0	1158 (12.5%)	1177 (17.8%)	323 (27.7%)	3040 (15.5%)
1	6319 (68.2%)	4107 (62.0%)	634 (54.3%)	13118 (67.1%)
2	1274 (13.7%)	706 (10.7%)	126 (10.8%)	2114 (10.8%)
3	483 (5.2%)	571 (8.6%)	63 (5.4%)	1117 (5.7%)
Missing	38 (0.4%)	63 (1.0%)	22 (1.9%)	175 (0.9%)
factor(PRIOR_AORSURG)				
0	6753 (72.8%)	5347 (80.7%)	954 (81.7%)	14839 (75.8%)
1	1377 (14.9%)	719 (10.9%)	108 (9.2%)	2549 (13.0%)
2	917 (9.9%)	479 (7.2%)	92 (7.9%)	1654 (8.5%)
3	216 (2.3%)	63 (1.0%)	10 (0.9%)	289 (1.5%)
4	0 (0%)	0 (0%)	1 (0.1%)	131 (0.7%)
Missing	9 (0.1%)	16 (0.2%)	3 (0.3%)	102 (0.5%)

Medical center info:

19 regions, 189 centers, 1094 physicians.

Most physicians only performed 1 or 2 procedures. Several performed over 100 procedures? Is that real?

There are regions and centers that performed many procedures. Would that affect the outcome?

Code Appendix

```
knitr::opts_chunk$set(echo = FALSE,message = FALSE,warning = FALSE)
library(tidyverse)
library(table1)

## ----- working directories for Lily -----
wd_lily = '/Users/hanyiwang/Desktop/Comparative-analysis-of-treatments-of-CAA'
path_lily = c(
  "../data/TEVAR_International_20210712/TEVAR_International_LTF_r12_2_14_20210701.csv",
  "../data/TEVAR_International_20210712/TEVAR_International_PROC_r12_2_14_20210701.csv",
  "../data/TEVAR_International_20210901/TEVAR_International_LTF_r12_2_14_20210901.csv",
  "../data/TEVAR_International_20210901/TEVAR_International_PROC_r12_2_14_20210901.csv")

## ----- read data -----
setwd(wd_lily)
TEVAR_LTF_07 = read.csv(path_lily[1])
TEVAR_PROC_07 = read.csv(path_lily[2])
#TEVAR_LTF_09 = read.csv(path_lily[3])
#TEVAR_PROC_09 = read.csv(path_lily[4])
## ----- data cleaning-----
TEVAR_PROC_07 = TEVAR_PROC_07 %>%
  mutate(DEAD=factor(DEAD)) %>%
  mutate(PRESENTATION = factor(PRESENTATION,levels = c(0,1,2),
                                labels = c('Asymptomatic','Symptomatic','Rupture')) %>%
  mutate(AGECAT = factor(AGECAT,levels = c(1,2,3,4,5,6,7),
                                labels = c('<40','40-49','50-59','60-69','70-79','80-89','>89')) %>%
  mutate(GENDER=factor(GENDER,levels=c(1,2),
                                labels=c('male','female'))))

## ----- population of interest -----
table1(~ PRESENTATION, data = TEVAR_PROC_07)
## ----- descriptive statistics table for outcomes-----
table1(~ factor(DEAD) + PROC_SURVIVALDAYS | PRESENTATION, data = TEVAR_PROC_07)
## ----- descriptive statistics table for patients conditions-----
table1(~ GENDER+AGE+AGECAT+factor(PREOP_SMOKING)+factor(PRIOR_CVD)+factor(PRIOR_CAD)+
  factor(PRIOR_CHF)+factor(COPD)+factor(DIABETES)+factor(HTN)+factor(PRIOR_AORSURG)
  | PRESENTATION, data = TEVAR_PROC_07)

## ----- descriptive statistics table for other variables of interest-----

#TEVAR_PROC_07 %>% select(REGIONID) %>% table()
#TEVAR_PROC_07 %>% select(CENTERID) %>% table()
#TEVAR_PROC_07 %>% select(PHYSICIANID) %>% table()
```