Logistic Regression Models with GEE for the VQI FBVAR Dataset

Jennifer Ci, Thu Vu, Lily Hanyi Wang

Variables to adjust for:

- cluster on CENTERID
- AGECAT, GENDER, PREOP_SMOKING, PRIOR_AORSURG, PRIOR_CHF, PREOP_DIALYSIS
- PATHOLOGY, extent

In the model, we merge groups for the extent: merge "Juxtarenal AAA" with "Type 4 TAAA"; "Type 1 TAAA", "Type 2 TAAA", "Type 3 TAAA", with "Type 5 TAAA". Now extent is a binary variable, Juxtarenal or not.

Outcome variables:

On this report GEE2:

• POSTOP_CEREBROSX, POSTOP_SPINAL_ISCHEMIA, POSTOP_DIALYSIS, POSTOP_LOS, POSTOP_COMPLICATIONS, POSTOP_LEGEMBO, POSTOP_RESPIRATORY, RETX_R_RTOR, BRANCH_POST

On the other report GEE:

• TOTAL_LOS,ICUSTAY, POSTOP_GFR, POSTOP_PRBC, POSTOP_INTISCH

POSTOP_CEREBROSX: Post-op Cerebrovascular Stroke:

Also account for ARMNECK_ACCESS

	Post-op Cerebrovascular Stroke			
Characteristic	OR^1	95% CI ¹	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	3.42	1.83, 6.38	< 0.001	
1				

¹OR = Odds Ratio, CI = Confidence Interval

	Post-op Cerebrovascular Str		
Characteristic	$ormalfont{OR}^1$	95% CI ¹	p-value
PRESENTATION			
Asymptomatic		_	
Symptomatic	1.81	0.91,3.60	0.091
AGECAT			

 $^{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

	Post-op Cerebrovascular Stroke			
Characteristic	$ormalsize{OR}^1$	$95\%~\mathrm{CI}^1$	p-value	
<50	_	_		
>79	0.51	0.08, 3.29	0.5	
50-59	0.79	0.13, 4.95	0.8	
60-69	0.23	0.04, 1.45	0.12	
70-79	0.62	0.11, 3.46	0.6	
GENDER				
female	_	_		
male	0.49	0.26, 0.90	0.023	
PREOP_SMOKING				
No				
Yes	0.73	0.34, 1.58	0.4	
PRIOR_AORSURG				
Both	_	_		
Endo	1.20	0.24, 6.05	0.8	
None	1.39	0.28, 6.92	0.7	
Open	0.95	0.15, 5.88	>0.9	
PRIOR_CHF				
No	_	_		
Yes	2.14	1.07, 4.31	0.032	
PREOP_DIALYSIS				
No		_		
Yes	1.97	0.46, 8.55	0.4	
PATHOLOGY				
Aneurysm	_	_		
Aneurysm from dissection	3.44	1.03, 11.6	0.045	
Dissection	2.56	0.57, 11.5	0.2	
PAU/IMH	1.46	$0.21,\ 10.2$	0.7	
extent				
Juxtarenal	_	_		
No	1.44	0.68, 3.04	0.3	
ARMNECK_ACCESS				
No				
Yes	2.31	1.26, 4.25	0.007	

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

POSTOP_SPINAL_ISCHEMIA: Post-op Spinal Ischemia

Also account for ${\tt POSTOP_SPINALDRAIN}$ and ${\tt OCCLUDED_CELIAC}$

	Post-op Spinal Ischemia			
Characteristic	OR^1	95% CI 1	p-value	
PRESENTATION			,	
Asymptomatic		_		
Symptomatic	2.56	1.70, 3.86	< 0.001	

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

	Post-op Spinal Ischemia				
Characteristic	\mathbf{OR}^1	$95\%~\mathrm{CI}^1$	p-value		
PRESENTATION					
Asymptomatic		_			
Symptomatic	1.61	1.00, 2.60	0.050		
AGECAT					
< 50		_			
>79	2.19	$0.27,\ 17.7$	0.5		
50-59	1.99	0.23, 17.3	0.5		
60-69	1.63	0.21,13.0	0.6		
70-79	2.13	0.27, 16.6	0.5		
GENDER					
female		_			
male	0.88	0.58, 1.36	0.6		
PREOP_SMOKING					
No		_			
Yes	1.22	0.64, 2.32	0.5		
PRIOR_AORSURG					
Both		_			
Endo	1.23	0.52, 2.95	0.6		
None	1.46	0.63, 3.40	0.4		
Open	1.47	0.60, 3.63	0.4		
PRIOR_CHF					
No		_			
Yes	1.06	0.61, 1.83	0.8		
$^{1}\mathrm{OR}=\mathrm{Odds}$ Ratio, $\mathrm{CI}=\mathrm{Confidence}$ Interval					

	Post-op Spinal Ischemia		
Characteristic	$ormalformath{\mathbf{R}}^1$	95% CI ¹	p-value
PREOP_DIALYSIS			
No	_	_	
Yes	1.16	0.48, 2.83	0.7
PATHOLOGY			
Aneurysm	_	_	
Aneurysm from dissection	0.98	0.43, 2.25	>0.9
Dissection	1.08	0.36,3.26	0.9
PAU/IMH	0.44	0.05, 3.63	0.4
extent			
Juxtarenal	_	_	
No	1.01	0.65, 1.57	>0.9
POSTOP_SPINALDRAIN			
No	_	_	
Yes	19.0	$11.1,\ 32.6$	< 0.001
OCCLUDED_CELIAC			
FALSE	_	_	
TRUE	1.57	0.83, 2.99	0.2

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

POSTOP_DIALYSIS: Post-op Dialysis

Also account for OCCLUDED_RENAL

	Post-op Dialysis			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	2.42	1.44,4.07	< 0.001	

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

	Post	t-op Dialysis	
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value
PRESENTATION			,
Asymptomatic			
10D Odda Datio CI Confidence	- T41		

 $^{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

	Post-o _l	Dialysis	
Characteristic	$ormalize{OR}^1$	95% CI ¹	p-value
Symptomatic	1.78	1.01, 3.15	0.047
AGECAT			
< 50	_	_	
>79	203,838,430,666,770		
50-59	607,435,459,760,613		
60-69	273,465,288,987,599		
70-79	210,692,025,743,388		
GENDER			
female	_	_	
male	0.62	0.39, 1.01	0.057
PREOP_SMOKING			
No	_	_	
Yes	1.29	0.60, 2.78	0.5
PRIOR_AORSURG			
Both	_	_	
Endo	1.05	0.26, 4.21	>0.9
None	0.85	$0.22,\ 3.24$	0.8
Open	0.58	0.12, 2.83	0.5
PRIOR_CHF			
No	_	_	
Yes	0.74	0.38, 1.47	0.4
PREOP_DIALYSIS			
No	_	_	
Yes	0.00	0.00, 0.00	< 0.001
PATHOLOGY			
Aneurysm	_	_	
Aneurysm from dissection	0.36	0.04,3.21	0.4
Dissection	2.06	0.63,6.68	0.2
PAU/IMH	2.41	0.53, 11.0	0.3
extent			
Juxtarenal	_	_	
No	1.03	0.61, 1.76	>0.9
OCCLUDED_RENAL			
$^{1}OR = Odds Ratio, CI = Conf$	idence Interval		

	op Dialysis		
Characteristic	OR^1	95% CI 1	p-value
FALSE	_	<u>—</u>	
TRUE	4.75	2.29, 9.83	< 0.001

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

POSTOP_LOS: Length of stay in days between surgery date and discharge date

	Length of stay in days between surgery date and discharge date				
Characteristic	OR^1	95% CI ¹	p-value		
PRESENTATION					
Asymptomatic		_			
Symptomatic	3.39	2.69, 4.29	< 0.001		

 $^{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

	Length of stay in days between surgery date and discharge date			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION	,			
Asymptomatic	_	_		
Symptomatic	2.46	1.91, 3.17	< 0.001	
AGECAT				
< 50	_	_		
>79	2.28	0.88, 5.88	0.089	
50-59	1.93	0.72, 5.21	0.2	
60-69	1.55	0.60, 3.96	0.4	
70-79	2.49	0.98,6.35	0.055	
GENDER				
female	_			
male	0.62	0.50, 0.75	< 0.001	
PREOP_SMOKING				
No	_			
Yes	0.97	0.74, 1.28	0.8	
PRIOR_AORSURG				
Both		_		
10P - Odda Patio CI - (Confidence Inter	,,,,,1		

¹OR = Odds Ratio, CI = Confidence Interval

		of stay in da date and dis	
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value
Endo	1.26	0.73, 2.19	0.4
None	0.96	0.57, 1.62	0.9
Open	1.36	0.76, 2.41	0.3
PRIOR_CHF			
No	_	_	
Yes	1.41	1.11, 1.80	0.005
PREOP_DIALYSIS			
No	_	_	
Yes	1.74	1.00, 3.00	0.048
PATHOLOGY			
Aneurysm	_		
Aneurysm from dissection	1.59	0.96, 2.63	0.069
Dissection	1.96	$1.11,\ 3.45$	0.020
PAU/IMH	1.82	0.90, 3.67	0.094
extent			
Juxtarenal			
No	1.97	1.60, 2.44	< 0.001

 $[\]frac{1.97}{1}$ OR = Odds Ratio, CI = Confidence Interval

${\bf POSTOP_COMPLICATIONS:\ Any\ Complications\ Post-op}$

	Any Complications Post-op			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	1.74	1.37, 2.21	< 0.001	

 $¹_{OR} = Odds Ratio, CI = Confidence Interval$

	Any (Complication	s Post-op
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value
PRESENTATION			
Asymptomatic	_	_	
$^{1}OR = Odds Ratio, CI = Co$	onfidence Inter	val	

	Any	Complication	s Post-op
Characteristic	$ormalfont{R}^1$	$95\%~\mathrm{CI}^1$	p-value
Symptomatic	1.44	1.12, 1.86	0.004
AGECAT			
< 50	_	_	
>79	1.17	0.46, 2.97	0.7
50-59	1.29	0.49, 3.42	0.6
60-69	0.97	0.39, 2.41	>0.9
70-79	1.32	0.53, 3.31	0.6
GENDER			
female		_	
male	0.65	0.54, 0.78	< 0.001
PREOP_SMOKING			
No	_	_	
Yes	1.01	0.79, 1.31	>0.9
PRIOR_AORSURG			
Both		_	
Endo	0.95	0.56, 1.60	0.8
None	1.00	0.60, 1.65	>0.9
Open	1.41	0.82, 2.42	0.2
PRIOR_CHF			
No	_	_	
Yes	1.28	1.04, 1.58	0.018
PREOP_DIALYSIS			
No	_	_	
Yes	1.25	0.73, 2.15	0.4
PATHOLOGY			
Aneurysm	_	_	
Aneurysm from dissection	1.32	0.82, 2.12	0.3
Dissection	0.94	0.51,1.73	0.8
PAU/IMH	0.77	0.35, 1.70	0.5
extent			
Juxtarenal		_	
No	1.47	1.21, 1.78	< 0.001

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

POSTOP_LEGEMBO: Leg Ischemia/Embol

	Leg Ischemia/Embol			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	1.97	1.13, 3.46	0.017	

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

	Leg Ischemia/Embol
Characteristic	OR^1
PRESENTATION	
Asymptomatic	_
Symptomatic	1.73
AGECAT	
< 50	_
>79	458, 932, 896, 998, 893, 117, 780, 708, 457, 649, 807, 168, 487, 458, 508, 727, 832, 855, 727, 852, 852, 852, 852, 852, 852, 852, 852
50-59	791, 185, 488, 625, 823, 058, 082, 792, 836, 217, 472, 083, 849, 201, 678, 512, 658, 724, 126, 126, 126, 126, 126, 126, 126, 126
60-69	544, 585, 817, 390, 864, 816, 453, 901, 583, 231, 375, 025, 203, 789, 035, 916, 660, 773, 496, 196, 196, 196, 196, 196, 196, 196, 1
70-79	545, 526, 688, 933, 917, 306, 277, 561, 448, 829, 344, 287, 639, 346, 295, 456, 815, 703, 326, 326, 326, 326, 326, 326, 326, 32
GENDER	
female	_
male	0.43
PREOP_SMOKING	
No	_
Yes	2.60
PRIOR_AORSURG	
Both	<u> </u>
Endo	11,407,653,425,563,561,885,696
None	$13,\!289,\!485,\!077,\!706,\!293,\!903,\!360$
Open	23,581,021,958,934,536,126,464
PRIOR_CHF	
No	_
Yes	0.86
PREOP_DIALYSIS	

 $^{1}\mathrm{OR}=\mathrm{Odds}$ Ratio, $\mathrm{CI}=\mathrm{Confidence}$ Interval

	${\bf Leg~Ischemia/Embol}$
Characteristic	OR^1
No	_
Yes	0.62
PATHOLOGY	
Aneurysm	_
Aneurysm from dissection	0.90
Dissection	0.65
PAU/IMH	2.36
extent	
Juxtarenal	_
No	1.02

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

${\bf POSTOP_RESPIRATORY:\ Post-op\ Respiratory}$

	Post-op Respiratory			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	2.65	1.87, 3.75	< 0.001	

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

		Post-op Respiratory	
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value
PRESENTATION			
Asymptomatic	_	_	
Symptomatic	2.16	$1.46,\ 3.20$	< 0.001
AGECAT			
< 50		_	
>79	1,443,137	$613,465,\ 3,394,888$	< 0.001
50-59	1,904,018	$772,645,\ 4,692,040$	< 0.001
60-69	1,430,807	$614,064,\ 3,333,866$	< 0.001
70-79	1,637,572	$687,735,\ 3,899,238$	< 0.001
GENDER			

 $^{^{1}\}mathrm{OR}=\mathrm{Odds}$ Ratio, $\mathrm{CI}=\mathrm{Confidence}$ Interval

	F	Post-op Respirator	y
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value
female		_	
male	0.66	0.47, 0.92	0.015
PREOP_SMOKING			
No	_	_	
Yes	1.07	0.67, 1.71	0.8
PRIOR_AORSURG			
Both	_	_	
Endo	0.59	0.26, 1.32	0.2
None	0.56	0.26, 1.18	0.13
Open	0.67	0.28, 1.60	0.4
PRIOR_CHF			
No	_	_	
Yes	1.24	0.83, 1.86	0.3
PREOP_DIALYSIS			
No	_	_	
Yes	0.78	0.27, 2.24	0.6
PATHOLOGY			
Aneurysm	_	_	
Aneurysm from dissection	0.62	0.24, 1.61	0.3
Dissection	0.94	0.33, 2.65	>0.9
PAU/IMH	1.19	0.34, 4.16	0.8
extent			
Juxtarenal	_	_	
No	1.77	1.27, 2.45	< 0.001

 $[\]frac{1.77}{1}$ OR = Odds Ratio, CI = Confidence Interval

 ${\tt RETX_R_RTOR} \hbox{: Re-intervention}$

	Re-intervention			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic		_		
Symptomatic	1.86	1.33, 2.62	< 0.001	

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

	Re-intervention			
Characteristic	OR^1	$95\%~\mathrm{CI}^1$	p-value	
PRESENTATION				
Asymptomatic	_	_		
Symptomatic	1.48	1.04, 2.09	0.029	
AGECAT				
< 50				
>79	0.95	$0.28,\ 3.27$	>0.9	
50-59	1.41	0.38, 5.19	0.6	
60-69	0.70	0.20, 2.44	0.6	
70-79	0.94	0.28, 3.14	>0.9	
GENDER				
female	_	_		
male	0.59	0.45, 0.78	< 0.001	
PREOP_SMOKING				
No	_	_		
Yes	1.29	0.81, 2.05	0.3	
PRIOR_AORSURG				
Both	_	_		
Endo	0.95	0.43, 2.09	0.9	
None	1.01	0.47, 2.16	>0.9	
Open	0.67	0.27, 1.63	0.4	
PRIOR_CHF				
No				
Yes	1.21	0.86, 1.72	0.3	
PREOP_DIALYSIS				
No	_	_		
Yes	0.75	0.31, 1.81	0.5	
PATHOLOGY				
Aneurysm	_	_		
Aneurysm from dissection	1.68	0.88, 3.19	0.12	
Dissection	0.64	0.23, 1.78	0.4	
PAU/IMH	1.77	0.70, 4.47	0.2	
extent				
Juxtarenal				
1 OR = Odds Ratio, CI = Con	fidence Inter	val		

	Re-intervention		
Characteristic	OR^1	95% CI ¹	p-value
No	1.56	1.15, 2.12	0.004

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

BRANCH_POST: Post-treatment Status of All Branches

	Post-treatment Status of All Branches		
Characteristic	OR^1	95% CI ¹	p-value
PRESENTATION			
Asymptomatic		_	
Symptomatic	2.28	1.78, 2.91	< 0.001

 $[\]overline{^{1}\text{OR} = \text{Odds Ratio, CI} = \text{Confidence Interval}}$

Characteristic	Post-treatment Status of All Branches		
	OR^1	$95\% \text{ CI}^1$	p-value
PRESENTATION			
Asymptomatic	_	_	
Symptomatic	1.64	1.24, 2.16	< 0.001
AGECAT			
< 50	_	_	
>79	0.78	0.32, 1.90	0.6
50-59	1.00	0.39, 2.58	>0.9
60-69	0.95	0.40, 2.27	>0.9
70-79	0.90	0.38, 2.14	0.8
GENDER			
female			
male	0.47	0.39, 0.58	< 0.001
PREOP_SMOKING			
No	_	_	
Yes	1.08	0.79, 1.48	0.6
PRIOR_AORSURG			
Both			
Endo	0.79	0.46, 1.35	0.4
10D Odda Datia CI C	anfidanca Inton	1	

¹OR = Odds Ratio, CI = Confidence Interval

	Post-treatment Status of All Branches		
Characteristic	OR^1	95% CI ¹	p-value
None	0.52	0.31, 0.86	0.011
Open	0.69	0.39, 1.23	0.2
PRIOR_CHF			
No			
Yes	1.43	1.12, 1.82	0.004
PREOP_DIALYSIS			
No	_	_	
Yes	7.90	4.70, 13.3	< 0.001
PATHOLOGY			
Aneurysm		_	
Aneurysm from dissection	0.77	0.43, 1.37	0.4
Dissection	1.57	0.90, 2.74	0.12
PAU/IMH	0.45	0.16, 1.24	0.12
extent			
Juxtarenal	_	_	
No	1.56	1.25, 1.93	< 0.001

 $[\]overline{^{1}}$ OR = Odds Ratio, CI = Confidence Interval

Code Appendix

```
knitr::opts_chunk$set(echo = FALSE,message = FALSE,warning = FALSE)
knitr::opts_chunk$set(fig.width=20, fig.height=20)
library(tidyverse)
library(table1)
library(dplyr)
library(geepack)
library(gtsummary)
library(broom.mixed)
## ----- working directories for Lily -----
wd_lily = '/Users/hanyiwang/Desktop/Comparative-analysis-of-treatments-of-CAA'
path_lily = c("../data/TEVAR_PROC.csv")
## ----- read data -----
setwd(wd_lily)
TEVAR_PROC = read.csv(path_lily)
## ----- modify some variables -----
TEVAR_PROC = TEVAR_PROC %>%
 mutate(extent = factor(extent,levels = c("Juxtarenal AAA", "Type 1 TAAA", "Type 2 TAAA", "Type 3 TAAA", "
 mutate(POSTOP_LOS = case_when(POSTOP_LOS>7 ~ '>7',
                               POSTOP LOS<=7 ~ '<=7')) %>%
 mutate(POSTOP_CEREBROSX = as.numeric(POSTOP_CEREBROSX == "Yes")) %>%
 mutate(POSTOP_SPINAL_ISCHEMIA = as.numeric(POSTOP_SPINAL_ISCHEMIA == "Yes")) %>%
 mutate(POSTOP_DIALYSIS = as.numeric(POSTOP_DIALYSIS == "Yes")) %>%
 mutate(POSTOP_LOS = as.numeric(POSTOP_LOS != "<=7") ) %>%
 mutate(POSTOP_COMPLICATIONS = as.numeric(POSTOP_COMPLICATIONS == "Yes")) %>%
 mutate(POSTOP_LEGEMBO = as.numeric(POSTOP_LEGEMBO == "Yes")) %>%
 mutate(POSTOP_RESPIRATORY = as.numeric(POSTOP_RESPIRATORY == "Yes")) %>%
 mutate(RETX_R_RTOR = as.numeric(RETX_R_RTOR == "Yes")) %>%
 mutate(BRANCH_POST = as.numeric(BRANCH_POST == "Yes"))
TEVAR_PROC = subset(TEVAR_PROC, !is.na(extent))
## ----- change class of variables -----
names <- c('CENTERID', 'AGECAT', 'GENDER', 'PREOP SMOKING', 'PRIOR AORSURG', 'PRIOR CHF', 'PREOP DIALYSI
TEVAR_PROC[,names] <- lapply(TEVAR_PROC[,names] , factor)</pre>
## ----- Post-op Cerebrovascular Stroke -----
## ----- unadjusted -----
POSTOP_CEREBROSX1 = geeglm(POSTOP_CEREBROSX ~ PRESENTATION,
                          data=TEVAR_PROC, family=binomial(link="logit"),
                          id = CENTERID, corstr = "independence")
t1 = POSTOP_CEREBROSX1 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
 bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Post-op Cerebrovascular Stroke**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_CEREBROSX2 = geeglm(POSTOP_CEREBROSX ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
```

```
PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent+ARMNECK_ACCESS,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP CEREBROSX2 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Post-op Cerebrovascular Stroke**")%>%as_flex_table()
## ----- Post-op Spinal Ischemia -----
## ----- unadjusted -----
POSTOP_SPINAL_ISCHEMIA1 = geeglm(POSTOP_SPINAL_ISCHEMIA ~ PRESENTATION,
                                data=TEVAR_PROC,family=binomial(link="logit"),
                                id = CENTERID, corstr = "independence")
t1 = POSTOP_SPINAL_ISCHEMIA1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Post-op Spinal Ischemia**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_SPINAL_ISCHEMIA2 = geeglm(POSTOP_SPINAL_ISCHEMIA ~ PRESENTATION+AGECAT+GENDER+
                                  PREOP_SMOKING+PRIOR_AORSURG+PRIOR_CHF+
                                  PREOP_DIALYSIS+PATHOLOGY+extent+
                                  POSTOP SPINALDRAIN+OCCLUDED CELIAC,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP_SPINAL_ISCHEMIA2 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Post-op Spinal Ischemia**")%>%as_flex_table()
## ----- Post-op Dialysis -----
## ----- unadjusted -----
POSTOP_DIALYSIS1 = geeglm(POSTOP_DIALYSIS ~ PRESENTATION,
                                data=TEVAR_PROC,family=binomial(link="logit"),
                                id = CENTERID, corstr = "independence")
t1 = POSTOP_DIALYSIS1 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Post-op Dialysis**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_DIALYSIS2 = geeglm(POSTOP_DIALYSIS ~ PRESENTATION+AGECAT+GENDER+
                                  PREOP SMOKING+PRIOR AORSURG+PRIOR CHF+
                                  PREOP_DIALYSIS+PATHOLOGY+extent+
```

```
OCCLUDED_RENAL,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP DIALYSIS2 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Post-op Dialysis**")%>%as_flex_table()
## ------ Length of stay in days between surgery date and discharge date -----
## ----- unadjusted -----
POSTOP_LOS1 = geeglm(POSTOP_LOS~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)
t1 = POSTOP LOS1 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Length of stay in days between surgery date and discharge dat
## ----- adjusted -----
POSTOP_LOS2 = geeglm(POSTOP_LOS ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP_LOS2 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Length of stay in days between surgery date and discharge dat
## ----- Any Complications Post-op -----
## ----- unadjusted -----
POSTOP_COMPLICATIONS1 = geeglm(POSTOP_COMPLICATIONS~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
t1 = POSTOP_COMPLICATIONS1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Any Complications Post-op**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_COMPLICATIONS2 = geeglm(POSTOP_COMPLICATIONS ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP_COMPLICATIONS2 %>%
```

```
tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Any Complications Post-op**")%>%as_flex_table()
## ----- Leg Ischemia/Emboli -----
## ----- unadjusted -----
POSTOP_LEGEMB01 = geeglm(POSTOP_LEGEMB0~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
# summary(POSTOP LOS1)
t1 = POSTOP LEGEMB01 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Leg Ischemia/Embol**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_LEGEMBO2 = geeglm(POSTOP_LEGEMBO ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP_LEGEMB02 %>%
  tbl regression(exponentiate=TRUE, tidy fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Leg Ischemia/Embol**")%>%as_flex_table()
## ----- Post-op Respiratory -----
## ----- unadjusted -----
POSTOP_RESPIRATORY1 = geeglm(POSTOP_RESPIRATORY~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
t1 = POSTOP_RESPIRATORY1 %>%
  tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Post-op Respiratory**")%>%as_flex_table()
## ----- adjusted -----
POSTOP_RESPIRATORY2 = geeglm(POSTOP_RESPIRATORY ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = POSTOP_RESPIRATORY2 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Post-op Respiratory**")%>%as_flex_table()
## ----- Re-intervention -----
## ----- unadjusted -----
```

```
RETX_R_RTOR1 = geeglm(RETX_R_RTOR~ PRESENTATION,
                    data=TEVAR_PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)
t1 = RETX R RTOR1 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Re-intervention**")%>%as_flex_table()
## ----- adjusted -----
RETX_R_RTOR2 = geeglm(RETX_R_RTOR ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = RETX_R_RTOR2 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Re-intervention**")%>%as_flex_table()
## ------ Post-treatment Status of All Branches ------
## ----- unadjusted -----
BRANCH_POST1 = geeglm(BRANCH_POST~ PRESENTATION,
                    data=TEVAR PROC, family=binomial(link="logit"),
                    id = CENTERID, corstr = "independence")
# summary(POSTOP_LOS1)
t1 = BRANCH_POST1 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold_p(t = 0.05)
tbl_merge(tbls = list(t1),tab_spanner ="**Post-treatment Status of All Branches**")%>%as_flex_table()
## ----- adjusted -----
BRANCH_POST2 = geeglm(BRANCH_POST ~ PRESENTATION+AGECAT+GENDER+PREOP_SMOKING+
                     PRIOR_AORSURG+PRIOR_CHF+PREOP_DIALYSIS+PATHOLOGY+extent,
                   family=binomial(link="logit"), data=TEVAR_PROC,
                   id = CENTERID, corstr = "independence")
t2 = BRANCH POST2 %>%
 tbl_regression(exponentiate=TRUE, tidy_fun = broom.mixed::tidy)%>%
  bold p(t = 0.05)
tbl_merge(tbls = list(t2),tab_spanner ="**Post-treatment Status of All Branches**")%>%as_flex_table()
```