

SIF3004 Final Year Project Proposal : RAdio Galaxy Environment Reference Survey (RAGERS) Project

Lim Ming Kang U2004991/1

November 2, 2022

1 Problem Statement

At the cosmological redshift of $2 < z < 3$ (observing the condition about 10.9 billion years ago), the star forming rates (SFRs) is the highest, known as "the cosmic noon". Those galaxies are usually "dusty", enriched with dust which serves as the materials to star forming.

A thorough understanding of those galaxies are important in understanding galaxies formation and evaluation of the early universe.

The growth and evolution of the aforementioned galaxy overdensities with redshift is to this day still not characterised. Nor is the effect the AGN activity has on the growth of the central overdensities.

Due to dust, those galaxies are not feasible to be observed in visible region, a telescope capable in observing in far infrared is needed to detect the galaxies.

Data directly from telescope is not suitable for analysis, as noise and false detections may be present, extensive data reduction and cleaning are essential to obtain a clean map for analysis.

The addition of far infrared data adds values to multiwavelength analysis, especially in obtaining photometric redshift.

2 Objectives

1. To reduce data from raw telescope data to obtain analysable data (eg. source count) for a single source field
2. To study statistically (eg. surface number density) of overdensities within a source field
3. To obtain photometric redshift for detected submm galaxies in a source field.

3 Background

Submillimetre Galaxies (SMGs) are rare galaxies with high star formation rates ($> 100 M_{\odot} yr^{-1}$) [?] populated in high redshift region. Suspected to be the progenitors for local giant galaxies, SMGs are good candidates to study the evolution of galaxies in high redshift. One challenge in observing SMGs is the amount of dust, that serves as the building blocks for star formation, which obscure visible light. However, far infrared/radio frequency are transparent to dust, thus the observation is allowed with submillimetre telescope.

Overdensities are regions in space where the density of matter is relatively higher than others. Events such as mergers of galaxies and active galactic nuclei (AGN) happen in overdensities region. At $z > 1$, High redshift radio galaxies (HzRG) are usually in the region of overdensities where SMGs are. The dynamic between SMGs and HzRGs can be studied so have a more robust understanding of galaxy evolution at high redshift.




Figure 1: General process of preparing useful data from raw telescope data

Telescope operating at submillimetre region can be used to observe SMGs. An example is Atacama Large Millimeter Array (ALMA) telescope located in northern Chile, it operates at around wavelength $0.4mm < \lambda < 2.73mm$. Other examples are Fred Young Submillimeter Telescope operating at $\lambda = 0.35mm$ and SCUBA-2 at James Clerk Maxwell Telescope (JCMT) operating at $\lambda = 0.35mm$ and $\lambda = 0.67mm$, whose data is used in the research of this proposal. Telescopes are usually associate with their propriety software for data handling. For instances, CASA for ALMA, Starlink for JCMT and AIPS for Very Large Baseline Array (VLBA)

Spectral energy distribution (SED): can derive photometric redshift, stellar mass and SFR

Spectral resolution (R) describes the ability of a sensor to define fine wavelength intervals

spectroscopy redshift is used to calibrate and optimize photometric redshift Spitzer/MIPS 24 micron

4 Research Methodology

The research will be carried out on raw data recorded on source 5C7269 detected by JCMT with SCUBA-2 under RAGERS Project, collaborate with RAGERS Malaysia Team (from where the raw data is acquired). Data Reduction is run on Starlink software. The process of data handling is shown in Figure 1

James Clerk Maxwell Telescope (JCMT) is a 15m telescope designed to run on submillimetre wavelength (far infrared region). It is positioned at Maunakea, Hawaii.

SCUBA-2 is a camera attached on JCMT to observe at $450\mu m$ and $850\mu m$ (666GHz and 353GHz) The data is formatted and stored in .sdf format, which is readable by starlink software.

The RAdio Galaxy Environment Reference Survey is a JCMT program to observe overdensities of dusty galaxies within the Mpc region of 33 radio galaxies at redshift range $1 < z < 3.5$ and mass range $M_* \geq 1010.8M_{\odot}$

5C7269 is a source field spanning across few Mpc centered around a HzRG, which is located at $1 < z < 3.5$

starlink is a software consisting of several packages used to reduce and analyse data recorded by SCUBA-2 telescope, including (but not limited to) SMURF for data reduction and GAIA for data visualisation.

5 Expected Results

From the final reduced data, surface number density as the function of radius from center HzRG can be calculated from the number counts. Spectral Energy Density (SED) of each detected submm galaxy

can be calculated by combining multiwavelength flux density to obtain photometric redshift.

6 Significance

Surface number density is essential in studying the effect of environment. Photometric redshift is necessary to study the evolution of galaxies with redshift.

References

- [1] E Da Cunha, JA Hodge, CM Casey, HSB Algera, M Kaasinen, I Smail, F Walter, WN Brandt, H Dannerbauer, and ROBERTO Decarli. Measurements of the dust properties in z 1–3 submillimeter galaxies with alma. *The Astrophysical Journal*, 919(1):30, 2021.