



TNAU AGRITECH PORTAL

AGRICULTURE



English | Tamil

[Home](#) | [Seasons & Varieties](#) | [Tillage](#) | [Nutrient Mgmnt](#) | [Irrigation Mgmnt](#) | [Weed Mgmnt](#) | [Crop Protection](#) | [Cost of Cultivation](#) | [Photobank](#)

Search

[Crop Production](#) :: [Cereals](#) :: [Rice](#) :: [Rainfed Rice](#)

After Cultivation

- Azospirillum inoculants 10 packets (2000g/ha) and Phosphobacteria 10 packets (2000g/ha) or 20 packets (4000g/ha) of Azophos mixed with 25 kg of FYM may be broadcasted uniformly over the field just after the receipt soaking rain / moisture.
- Thinning and gap filling should be done 14 - 21 days after sowing, taking advantage of the immediate rain
- Spray Cycocel 1000 ppm (1 ml of in one lit. of water) under water deficit situations to mitigate ill-effects.
- Foliar spray of KCl 1% to overcome moisture stress at different physiological stages of rice.

Updated on Jan 2023

[Home](#) | [Seasons & Varieties](#) | [Tillage](#) | [Nutrient Management](#) | [Irrigation Management](#) | [Weed Management](#) | [Crop Protection](#) | [Cost of Cultivation](#) | [Photobank](#) | [Disclaimer](#)

© All Rights Reserved. TNAU 2008 - 2023.