Home | Seasons & Varieties | Tillage | Nutrient Mgmnt | Irrigation Mgmnt | Weed Mgmnt | Crop Protection | Cost of Cultivation | Photobank

Search

Crop Production :: Cereals :: Rice :: Wet Seeded Rice

Weed Management

- Sowing with drum seeder and cono weeding at 10, 20 and 30 DAS.
- Hand weeding on 15 20 DAT and 45 DAT.
- Pre emergence (PE) application of Pretilachlor 0.75 kg/ha on 8 DAS followed by one hand weeding on 40 DAS.
- Pre emergence application of Pendimethalin 1.0 kg/ha at 8 DAS followed by one hand weeding on 45 DAT.
- Pre emergence application of Pyrazosulfuron ethyl at 20 g a.i /ha on 3 DAS followed by cono weeding on 25 DAS
- Pre-emergence application of Pendimethalin 1.0 kg/ha at 3 DAS followed by post emergence application of bispyribac sodium 25 g/ha at 25 DAS with one hand weeding at 45 DAS.







| Home | Seasons & Varieties | Tillage | Nutrient Management | Irrigation Management | Weed Management | Crop Protection | Cost of Cultivation | Photobank | Disclaimer

© All Rights Reserved. TNAU 2008-2024.