

# Metric Spaces

**Definition 1** A **metric space** is a pair  $(X, d)$ , where  $X$  is a set and  $d : X \times X \rightarrow [0, \infty)$  for all  $x, y, z \in X$  has the following properties:

- (Positivity)  $d(x, y) = 0 \iff x = y$ ,
- (Symmetry)  $d(x, y) = d(y, x)$ ,
- (Triangle inequality)  $d(x, y) \leq d(x, z) + d(z, y)$ .

A function  $d : X \times X \rightarrow [0, \infty)$  that satisfies these axioms is called a **distance function** on  $X$ . A subset  $U \subset X$  of a metric space  $(X, d)$  is called **open** if, for every  $x \in U$ , there exists a constant  $\epsilon > 0$  such that the open ball

$$B(\epsilon, x) := \{y \in X \mid d(x, y) < \epsilon\}$$

(centered at  $x$  with radius  $\epsilon$ ) is contained in  $U$ .

A subset  $F$  of a metric space  $(X, d)$  is **closed** if its complement  $F^c$  is open.

## Some basic facts about open sets

- Every ball  $B(x, r)$  is open, for if  $y \in B(x, r)$  and  $d(x, y) = s$  then  $B(y, r - s) \subset B(x, r)$ .
- $X$  and  $\emptyset$  are both open and closed.
- The union of any family of open sets is open, and hence the intersection of any family of closed sets is closed.
- The intersection (resp. union) of any finite family of open (resp. closed) sets is open (resp. closed). Indeed, if  $U_1, \dots, U_n$  are open and  $x \in \cap_{i=1}^n U_i$ , for each  $j$  there exists  $r_j > 0$  such that  $B(x, r_j) \subset U_j$ , and then  $B(x, r) \subset \cap_{i=1}^n U_i$  where  $r = \min(r_1, \dots, r_n)$ , so  $\cap_{i=1}^n U_i$  is open.

Let  $\mathcal{P}(X)$  be the collection of all subsets of  $X$ . Recall that a *topology* on  $X$  is a subfamily  $\tau \subset \mathcal{P}(X)$  satisfying

- $\emptyset \in \tau$  and  $X \in \tau$ ;
- if  $U_i \in \tau$  ( $i = 1, \dots, n$ ), then  $\cap_{i=1}^n U_i \in \tau$ ;
- if  $U_\alpha$  ( $\alpha \in \mathcal{I}$ ) is an arbitrary collection in  $\tau$ , then  $\cup_{\alpha \in \mathcal{I}} U_\alpha \in \tau$ .

The set of open subsets of  $(X, d)$  will be denoted by

$$U(X, d) := \{U \subset X \mid U \text{ is open}\}.$$

It follows from the definitions that the collection  $U(X, d)$  in a metric space  $(X, d)$  satisfies the axioms of a topology and so  $(X, d)$  is a topological space.

We will use  $\mathbb{F}$  to denote either  $\mathbb{C}$  or  $\mathbb{R}$ .

**Example 1.** The set  $\mathbb{R}$  of all real numbers endowed with the distance function  $d(x, y) = |x - y|$ , where  $|x|$  is the absolute value of  $x$ , is a metric space.

Similarly, the set of all complex numbers  $\mathbb{C}$  is a metric space with the distance function  $d(z, w) = |z - w|$ , where  $|z|$  is the modulus of  $z$  in  $\mathbb{C}$ .

**Example 2.** Let  $X$  be a nonempty set. The function

$$d(x, y) = \begin{cases} 0, & \text{if } x = y, \\ 1, & \text{if } x \neq y, \end{cases}$$

is a metric, called the discrete metric (also known as the trivial metric) on  $X$ . The space  $(X, d)$  is called the discrete metric space.

**Example 3.** Let  $C[a, b] = \{x(t) : x(t) \text{ is continuous on } [a, b]\}$  and define

$$d_1(x, y) := \max_{a \leq t \leq b} |x(t) - y(t)|, \quad d_2(x, y) := \int_a^b |x(t) - y(t)| dt.$$

Then  $d_1$  and  $d_2$  are metrics on  $C[a, b]$ .

**Example 4.** For any integer  $k \geq 1$ , the function  $d : \mathbb{F}^k \times \mathbb{F}^k \rightarrow [0, \infty)$  defined by

$$d(x, y) = \left( \sum_{j=1}^k |x_j - y_j|^2 \right)^{1/2},$$

is a metric on the set  $\mathbb{F}^k$ , called the standard metric on  $\mathbb{F}^k$ .

**Example 5.** More generally, take  $X = \mathbb{K}^n$  with any one of the metrics

$$d_{l^p}(x, y) = \begin{cases} \left( \sum_{j=1}^n |x_j - y_j|^p \right)^{1/p}, & 1 \leq p < \infty, \\ \max_{j=1, \dots, n} |x_j - y_j|, & p = \infty. \end{cases}$$

It is easy to see that  $d_{l^\infty}$  is a metric. For the case  $1 \leq p < \infty$ , we need only to use the

([Minkowski's inequality](#).) For arbitrary complex numbers  $x_1, \dots, x_n, y_1, \dots, y_n$  and a real number  $p \geq 1$ ,

$$\left( \sum_{j=1}^n |x_j + y_j|^p \right)^{1/p} \leq \left( \sum_{j=1}^n |x_j|^p \right)^{1/p} + \left( \sum_{j=1}^n |y_j|^p \right)^{1/p}.$$

**Proof.** We may assume that both real numbers

$$u = \left( \sum_{j=1}^n |x_j|^p \right)^{1/p} \quad \text{and} \quad v = \left( \sum_{j=1}^n |y_j|^p \right)^{1/p}$$

are positive. By the triangle inequality, we have

$$|x_k + y_k|^p \leq (|x_k| + |y_k|)^p = (u + v)^p \left( \frac{u}{u+v} \frac{|x_k|}{u} + \frac{v}{u+v} \frac{|y_k|}{v} \right)^p.$$

Since  $\frac{u}{u+v} + \frac{v}{u+v} = 1$  and  $x^p$  is convex for  $p \geq 1$ , we have

$$\left( \frac{u}{u+v} \frac{|x_k|}{u} + \frac{v}{u+v} \frac{|y_k|}{v} \right)^p \leq \frac{u}{u+v} \frac{|x_k|^p}{u^p} + \frac{v}{u+v} \frac{|y_k|^p}{v^p}.$$

Hence

$$|x_k + y_k|^p \leq (u + v)^p \left( \frac{u}{u+v} \frac{|x_k|^p}{u^p} + \frac{v}{u+v} \frac{|y_k|^p}{v^p} \right).$$

By summing both sides of the above inequality, we obtain

$$\sum_{j=1}^n |x_j + y_j|^p \leq (u + v)^p.$$

**Example 6.** If  $d$  is a metric on  $X$  and  $A \subset X$ , then  $d|_{(A \times A)}$  is a metric on  $A$ .

**Example 7.** If  $(X_1, d_1)$  and  $(X_2, d_2)$  are metric spaces, the product metric  $d$  on  $X_1 \times X_2$  is given by

$$d((x_1, x_2), (y_1, y_2)) = \max\{d(x_1, y_1), d(x_2, y_2)\}.$$

Other metrics are sometimes used on  $X_1 \times X_2$ , for instance,

$$d(x_1, y_1) + d(x_2, y_2) \text{ or } \sqrt{d(x_1, y_1)^2 + d(x_2, y_2)^2}.$$



**Definition 2.** A point  $x$  in a metric space  $X$  is said to be a **limit** of a sequence of points  $(x_n) \subset X$  if for every  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $d(x_n, x) < \epsilon$ , for all  $n \geq N$ .

If  $x$  is a limit of the sequence  $(x_n)$ , we say that  $(x_n)$  **converges** to  $x$  and write  $x_n \rightarrow x$ .

If a sequence has a limit, it is called **convergent**. Otherwise, it is called **divergent**. Observe that  $x_n \rightarrow x \Leftrightarrow d(x_n, x) \rightarrow 0$ .

A subset  $Y$  of a metric space  $(X, d)$  is **bounded** if there exists  $x \in X$  and  $r > 0$ , such that  $Y \subset B(r, x)$ . Otherwise,  $Y$  is **unbounded**.

**Any convergent sequence is bounded**, since if  $x_n \rightarrow x$ , then there exists  $N \in \mathbb{N}$  such that  $d(x_n, x) < 1$  for all  $n \geq N$  and so

$$d(x_n, x) \leq \max \left( 1, \max_{j=1, \dots, N-1} d(x_j, x) \right), \quad \forall n \in \mathbb{N}.$$

**Definition 3.** A point  $x \in E$  is said to be an **interior point** of  $E$  if

$$\exists r > 0, \text{ s.t. } B(x, r) \subset E.$$

The **interior** of  $E$  is the set of all its interior points and is denoted by  $E^\circ$ . A point  $x$  (not in  $E$ ) is an **exterior point** of  $E$  when

$$\exists r > 0, \text{ s.t. } B(x, r) \subset X \setminus E.$$

All other points are called **boundary points** of  $E$ .

The set of interior and boundary points of  $E$  is called the closure of  $E$  and denoted by  $\overline{E} = E^\circ \cup \partial E$ . Note that  $\overline{E}$  is also the intersection of all closed sets containing  $E$ .

The set  $X$  is partitioned into three parts: its interior  $E^\circ$ , its exterior  $(\overline{E})^c$ , and its boundary  $\partial E$ .

We call  $E$  **dense** in  $X$  if  $\overline{E} = X$ , and **nowhere dense** if  $\overline{E}$  has empty interior.

**Lemma 1.**  $x_n \rightarrow x \Leftrightarrow \forall$  open set  $U$  that contains  $x$  there exists an  $N$  such that  $x_n \in U$  for every  $n \geq N$ .

**Proof**  $\Rightarrow$ : Given any open set  $U$  that contains  $x$  there exists  $\epsilon > 0$  such that  $B(x, \epsilon) \subset U$ , and so  $\exists N$  such that  $x_n \in B(x, \epsilon) \subset U$  for all  $n \geq N$ .

$\Leftarrow$ : For any  $\epsilon > 0$ , since the set  $B(x, \epsilon)$  is open and contains  $x$ , there exists an  $N$  such that  $x_n \in B(x, \epsilon)$  for every  $n \geq N$ . Thus  $x_n \rightarrow x$ . □

**Lemma 2.** A subset  $A$  of  $(X, d)$  is closed  $\Leftrightarrow$  whenever  $(x_n) \subset A$  with  $x_n \rightarrow x$  it follows that  $x \in A$ .

**Proof**  $\Rightarrow$ : Let  $(x_n) \subset A$  with  $x_n \rightarrow x$ . If  $x \notin A$ , i.e.  $x \in A^c$ , there would exist  $B(x, r) \subset A^c$  since  $A^c$  is open. It follows that  $B(x, r) \cap A = \emptyset$ , which contradicts to  $x_n \rightarrow x$ .

$\Leftarrow$ : Take  $x \notin A$ . If for any  $\epsilon > 0$ ,  $B(x, r) \cap A \neq \emptyset$ , by taking  $\epsilon = \frac{1}{n}$ , we would get  $(x_n) \subset A$  such that  $d(x, x_n) < \frac{1}{n}$ , which implies that  $x_n \rightarrow x$  and so  $x \in A$ . This is a contradiction. Thus there is some  $r > 0$  such that  $B(x, r) \cap A = \emptyset$ , that is  $B(x, r) \subset A^c$ . Consequently,  $A^c$  is open. □

A set  $V \subset X$  is said to be a **neighborhood** of a point  $p \in X$  if there is an  $r > 0$  such that  $B(p, r) \subset V$ . In particular, **any open set  $D$  is a neighborhood of any  $p \in D$ .**

A point  $p$  is an **accumulation point** (or **limit point**) of a set  $A$  if every open ball around it contains other points of  $A$ ,

$$\forall \epsilon > 0, \exists q \neq p, q \in A \cap B(p, \epsilon).$$

Note that  $p$  is not necessarily an element of  $S$ . If  $q \in S$  and  $q$  is not an accumulation point of  $S$ , then  $q$  is an **isolated point** of  $S$ .

By Lemma 2,  **$S$  is closed  $\Leftrightarrow S$  contains all its accumulation points.**

The **boundary** of a set  $S \subset (X, d)$  is also the set  $\partial S = \overline{S} \cap \overline{S^c}$ . We note that  $p \in \partial S \Leftrightarrow B(p, \epsilon) \cap S \neq \emptyset$  and  $B(p, \epsilon) \cap S^c \neq \emptyset$  for any  $\epsilon > 0$ .

### Lemma 3

$$x \in \overline{A} \Leftrightarrow B(x, \epsilon) \cap A \neq \emptyset, \forall \epsilon > 0, \quad (0.1)$$

that is,  $x \notin \overline{A} \Leftrightarrow \exists \epsilon_0 > 0$  such that  $B(x, \epsilon_0) \cap A = \emptyset$ . It follows that  $x \in \overline{A} \Leftrightarrow \exists$  a sequence  $(x_n) \subset A$  such that  $x_n \rightarrow x$ .

**Proof** If  $x \notin \overline{A}$ , then there is some closed set  $K$  that contains  $A$  such that  $x \notin K$ . Since  $K$  is closed,  $X \setminus K$  is open, and so  $B(x, \epsilon_0) \cap K = \emptyset$  for some  $\epsilon_0 > 0$ , which shows that  $B(x, \epsilon_0) \cap A = \emptyset$  since  $K \supset A$ . On the other hand, if there exists  $\epsilon_0 > 0$  such that  $B(x, \epsilon_0) \cap A = \emptyset$ , then  $x$  is not contained in the closed set  $X \setminus B(x, \epsilon_0)$ , which contains  $A$ ; so  $x \notin \overline{A}$ .

Finally, if  $x \in \overline{A}$ , then (0.1) implies that for any  $n \in \mathbb{N}$  we have  $B(x, 1/n) \cap A \neq \emptyset$ , so  $\exists x_n \in A$  such that  $d(x_n, x) < 1/n$  and thus  $x_n \rightarrow x$ . Conversely, if  $(x_n) \in A$  with  $x_n \rightarrow x$ , then  $\forall \epsilon > 0, d(x_n, x) < \epsilon$  for  $n$  sufficiently large, and so  $B(x, \epsilon) \cap A \neq \emptyset$  which implies by (0.1) that  $x \in \overline{A}$ .  $\square$

**Lemma 4**  $(\overline{A^c})^c = A^o$  and so  $\overline{A^c} = X \Leftrightarrow A^o = \emptyset$ .

**Proof**  $x \in (\overline{A^c})^c \Leftrightarrow x \notin \overline{A^c} \Leftrightarrow B(x, \epsilon_0) \cap A^c = \emptyset$  for some  $\epsilon_0 > 0$   
 $\Leftrightarrow B(x, \epsilon_0) \subset A$  for some  $\epsilon_0 > 0 \Leftrightarrow x \in A^o$ .  $\square$

Recall that  $A$  is dense in  $X$  if and only if  $\overline{A} = X$ . It follows from Lemma 3 that  $A$  is dense in  $X \Leftrightarrow \forall x \in X, \epsilon > 0, B(x, \epsilon) \cap A \neq \emptyset$ , i.e.  $\exists p \in A$  such that  $d(p, x) < \epsilon$ .

A sequence  $(x_n)$  in a metric space  $(X, d)$  is called a **Cauchy sequence** if, for every  $\epsilon > 0$ , there exists an  $n_0 \in \mathbb{N}$  such that for any two integers  $n, m \geq n_0$ , we have  $d(x_n, x_m) < \epsilon$ . A metric space  $(X, d)$  is called **complete** if every Cauchy sequence in  $X$  converges.

For any  $n \in \mathbb{N}$ ,  $\mathbb{R}^n$ , equipped with the Euclidean metric, is complete, because a Cauchy sequence in  $\mathbb{R}^n$  is Cauchy in each coordinate.

**Theorem**  $\forall$  metric space  $X$ ,  $\exists$  a complete metric space  $Y$  and a map  $j : X \rightarrow Y$  s.t.

1)  $d_Y(j(x), j(w)) = d_X(x, w)$ ,  $\forall x, w \in X$ . 2)  $\overline{j(X)} = Y$ .

If  $Z$  is another complete metric space and  $k : X \rightarrow Z$  is a map satisfying the 1) and 2), then  $\exists$  a bijective map  $f : Y \rightarrow Z$  s.t.

$d_Z(f(y_1), f(y_2)) = d_Y(y_1, y_2) \forall y_1, y_2 \in Y$  and  
 $f(j(x)) = k(x) \forall x \in X$ .

The metric space  $Y$  is called the **completion** of  $X$ .

Let  $A$  be a non-empty set of the metric space  $(X, d)$ . The diameter  $A$  is defined as  $\text{diam}(A) = \sup_{x, y \in A} d(x, y)$ .

**Theorem 1** (Baire) *Let  $(X, d)$  be complete and let  $A_n \subset X, n \in \mathbb{N}$ , be closed satisfying  $A_n^o = \emptyset, \forall n \in \mathbb{N}$ . Then,*

$$(\cup_{n=1}^{\infty} A_n)^o = \emptyset. \quad (0.2)$$

**Proof**  $\overline{A_n^c} = X$ . Thus  $A_n^c$  is dense in  $X$  and is also open  $\forall n \in \mathbb{N}$ .  
 Let  $D_n = A_n^c$ ; by Lemma 4, we need to show that  $\overline{(\cup_{n=1}^{\infty} A_n)^c} = X$   
 or, that  $M := \cap_{n=1}^{\infty} D_n$  is dense in  $X$ , i.e., **for every open**  
 $U = B(x_0, r_0) \subset X, r_0 > 0$  **we have**  $U \cap M \neq \emptyset$ . Fix such an open  
 $U$ . Since  $D_1$  is open and dense in  $X$  there exist  $x_1 \in U \cap D_1$  and  
 $r_1 > 0$  such that

$$\overline{B(x_1, r_1)} \subset U \cap D_1, \quad 0 < r_1 < r_0/2.$$

By induction one can find sequences  $(x_n)$  and  $(r_n)$  such that


$$\overline{B(x_{n+1}, r_{n+1})} \subset B(x_n, r_n) \cap D_{n+1}, \quad 0 < r_{n+1} < r_n/2.$$

for  $n = 0, 1, 2, \dots$ . Since for  $l \geq k$ ,

$$d(x_k, x_l) \leq \sum_{i=k}^{l-1} d(x_i, x_{i+1}) \leq \frac{r_0}{2^{k-1}},$$

$(x_n)$  is Cauchy. Let  $y = \lim x_n$ . Since  $x_l \in B(x_k, r_k), \forall l \geq k$ ,  
 $\Rightarrow y \in \overline{B(x_k, r_k)} \subset D_k, \forall k \in \mathbb{N}$  and so  $y \in U \cap M$ .





**Theorem 2.** (Nested sets theorem). *Let  $A_1 \supset A_2 \supset \cdots$  be a decreasing chain of non-empty closed subsets of a complete metric space  $(X, d)$  and let  $\text{diam}(A_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then  $\bigcap_{n=1}^{\infty} A_n$  consists of exactly one point.*

**Proof.** Pick in each  $A_n$  a point  $a_n$ . If  $N \in \mathbb{N}$  and  $k, j > N$ , then since  $A_n \downarrow$ , the points  $a_k$  and  $a_j$  belong to  $A_N$ . Thus,  $d(a_j, a_k) \leq \text{diam}(A_N) \rightarrow 0$  as  $N \rightarrow \infty$ , i.e.,  $(a_n)$  is Cauchy. Let  $a = \lim a_n$ . For any  $N$  and any  $k > N$ ,  $a_k \in A_N$ . Hence,  $a = \lim a_k \in A_N$ , i.e.,  $a \in \bigcap_{N=1}^{\infty} A_N$ . Note that  $\bigcap_{n=1}^{\infty} A_n \subset A_N$  for all  $N$ , and so

$$\text{diam}(\bigcap_{n=1}^{\infty} A_n) \leq \text{diam}(A_N) \rightarrow 0, \quad N \rightarrow \infty.$$

But a set of diameter zero reduces to a single point. □

## Contracted mapping theorem

Suppose  $(X, \rho)$  is a complete MS,  $T : X \rightarrow X$  satisfies:  $\exists \theta \in [0, 1)$  s.t.

$$\forall x \in X, \forall y \in X, \quad \rho(Tx, Ty) \leq \theta \rho(x, y). \quad (\Rightarrow T \text{ is conti.})$$

Then  $\exists \bar{x} \in X$  s.t.  $T\bar{x} = \bar{x}$ .

*Proof.* Step 1. Let  $x_0 \in X$ ,  $x_1 := Tx_0, \dots, x_{n+1} := Tx_n, \dots$

$$\rho(x_n, x_{n+1}) \leq \theta \rho(x_n, x_{n-1}) \leq \theta^2 \rho(x_{n-1}, x_{n-2}) \leq \dots \leq \theta^n \rho(x_0, Tx_0)$$

$\Rightarrow$  for any  $n, p \in \mathbb{N}$ ,

$$\begin{aligned} \rho(x_n, x_{n+p}) &\leq \rho(x_n, x_{n+1}) + \rho(x_{n+1}, x_{n+2}) + \dots + \rho(x_{n+p-1}, x_{n+p}) \\ &\leq (\theta^n + \theta^{n+1} + \dots + \theta^{n+p-1}) \rho(x_0, Tx_0) \leq \frac{\theta^n}{1 - \theta} \rho(x_0, Tx_0) \end{aligned}$$

$\Rightarrow \{x_n\}_n$  is Cauchy.  $X$  is complete  $\Rightarrow \exists \bar{x} \in X$  s.t.  $\lim x_n = \bar{x}$ .  
 $\Rightarrow T\bar{x} = \bar{x}$ .

Step 2. Suppose  $\exists \bar{x}, \tilde{x} \in X$  s.t.  $\bar{x} = T\bar{x}$ ,  $\tilde{x} = T\tilde{x}$ . Then

$$\rho(T\bar{x}, T\tilde{x}) \leq \theta \rho(\bar{x}, \tilde{x}) = \theta \rho(T\bar{x}, T\tilde{x}) \Rightarrow \bar{x} = \tilde{x}$$

because  $\theta < 1$ .

**Def.** A metric space  $(X, d)$  is **separable** if it contains a countable dense subset, i.e., there exists a countable subset  $B \subset X$  such that  $\overline{B} = X$ .

Separability means that elements of  $X$  can be approximated arbitrarily closely by some countable collection  $\{x_1, x_2, \dots\}$ :

$$\forall x \in X \text{ and } \epsilon > 0, \exists j \in \mathbb{N} \text{ s.t. } d(x_j, x) < \epsilon.$$

**Examples.**  $\mathbb{R}$  and  $\mathbb{C}$  are separable, since  $\mathbb{Q}$  and  $\mathbb{Q} + i\mathbb{Q}$  are countable dense subsets of  $\mathbb{R}$  and  $\mathbb{C}$ , respectively. Since separability of  $(X, d_X)$  and  $(Y, d_Y)$  implies separability of  $X \times Y$  (with an appropriate metric), it follows that  $\mathbb{R}^n$  and  $\mathbb{C}^n$  are separable.

**Lemma 5** If  $(X, d)$  separable and  $Y \subset X$ , then  $(Y, d)$  is also separable.

**Proof.** Let's construct a countable dense subset  $A$  of  $Y$ . Suppose that  $\{x_1, x_2, \dots\}$  is dense in  $X$ . For each  $k, n \in \mathbb{N}$ , if  $B(x_n, \frac{1}{k}) \cap Y \neq \emptyset$ , then we choose one point from  $B(x_n, \frac{1}{k}) \cap Y$  and add it to  $A$ . Constructed in this way  $A$  is (at most) a countable set. Given  $y \in Y$  and  $\epsilon > 0$ , take  $k \in \mathbb{N}$  such that  $\frac{1}{k} < \frac{\epsilon}{2}$ . Let  $x_n$  be such that  $d(x_n, y) < \frac{1}{k}$ ; then  $B(x_n, \frac{1}{k}) \cap Y \neq \emptyset$ . It follows that there exists a  $z \in A \cap B(x_n, \frac{1}{k}) \cap Y$  and so

$$d(y, z) \leq d(y, x_n) + d(x_n, z) < \frac{2}{k} < \epsilon.$$

Hence  $\overline{A} = Y$ , i.e.,  $Y$  is separable.

## Continuous maps and compact sets

Def.  $(X, d), (Y, d_1)$  are metric spaces, the map

$$T : X \ni x \rightarrow y = Tx \in Y.$$

For  $x_0 \in X$ ,  $T$  is continuous at  $x_0$  iff  $\forall \epsilon > 0, \exists \delta > 0$ , s.t.  
 $d(x, x_0) < \delta \Rightarrow d_1(Tx, Tx_0) < \epsilon$ .

$T$  is continuous on  $D \subset X$  iff  $\forall x \in D$ , it is continuous at  $x$ .  $T$  is called uniformly continuous on  $D$  if  $\delta$  can be the same for all  $x_0 \in D$ , i.e.,  $\delta$  is independent of  $x_0 \in D$ .

$T$  is a homeomorphism iff it is bijective, and  $T$  and  $T^{-1}$  are continuous.

- $T$  is continuous at  $x_0 \Leftrightarrow$   
 $\forall \{x_n\} \subset X, x_n \rightarrow x_0 \Rightarrow Tx_n \rightarrow Tx_0$ .

Proof.  $\Rightarrow$  is obvious.

$\Leftarrow$ : if  $T$  is not continuous at  $x_0$ , then  $\exists \epsilon_0 > 0$ , s.t.  
 $\forall n \in \mathbb{N}, \exists x_n \in X, d(x_n, x_0) < \frac{1}{n}, d_1(Tx_n, Tx_0) \geq \epsilon_0$ .

Hence,  $x_n \rightarrow x_0, Tx_n \not\rightarrow Tx_0$ .

- $T$  is continuous

$\Leftrightarrow \forall \text{ open } G \subset Y, T^{-1}(G) \text{ is open in } X.$

Proof.  $\Rightarrow$ : Let  $G \subset Y$  be open,  $x_0 \in T^{-1}(G)$ , then  $T(x_0) \in G$ , so  $\exists B(Tx_0, \epsilon) \subset G$ . By continuity,  $\exists \delta > 0$  s.t.  $T(B(x_0, \delta)) \subset B(Tx_0, \epsilon)$ . Thus,  $B(x_0, \delta) \subset T^{-1}(B(Tx_0, \epsilon)) \subset T^{-1}(G)$ . Hence,  $T^{-1}(G)$  is open.

$\Leftarrow$ : Let  $\{x_n\} \subset X, x_n \rightarrow x_0$ . Fix  $\epsilon > 0$ . Since  $T^{-1}(B(Tx_0, \epsilon))$  is open,  $\exists \delta > 0$  s.t.  $B(x_0, \delta) \subset T^{-1}(B(Tx_0, \epsilon))$ . Thus,  $T(B(x_0, \delta)) \subset B(Tx_0, \epsilon)$ . From  $x_n \rightarrow x_0$ ,  $\exists N_0 \in \mathbb{N}$  s.t.  $d(x_n, x_0) < \delta, \forall n \geq N_0$ . Hence,  $d(Tx_n, Tx_0) < \epsilon, \forall n \geq N_0$ .

Def.  $T : (X, d) \rightarrow (Y, d_1)$  is an *isometry* if it preserves distances, i.e.,

$$d_1(Tx, Ty) = d(x, y) \quad \forall x, y \in X.$$

$X$  is said to be *isometric* to  $Y$  if there is a bijective isometry from  $X$  onto  $Y$ .

Def.  $(X, d)$  is *compact* if every sequence in  $X$  has a convergent subsequence. A subset  $Y$  of  $X$  is called *compact* if it is a compact subspace of  $X$ .

Hence a subset  $Y$  of a metric space  $X$  is compact  $\Leftrightarrow$  every sequence in  $Y$  has a subsequence that converges to a point in  $Y$ .

- A compact set in a metric space is closed and bounded.

Proof. Let  $E(\subset (X, d))$  be compact and  $x \in \bar{E}$ , then  $\exists (x_n) \subset E$  s.t.  $x_n \rightarrow x$ . Since  $E$  is compact,  $x \in E$ . Thus  $E$  is closed. If  $E$  is not bounded, then  $\forall n \in \mathbb{N}, \exists x_n \in E$  s.t.  $d(x_n, a) > n$ , where  $a \in X$  is a fixed point. Since every convergent sequence in  $X$  is bounded,  $(x_n)$  has no convergent subsequence. This contradicts the fact that  $E$  is compact.

Lemma. Let  $\{U_i\}_{i \in J}$  be an open covering of a compact space  $X$ . Then  $\exists r > 0$  s.t.  $\forall x \in X$ ,  $B(x, r)$  is contained in some  $U_i$  for some  $i \in J$ .



Proof. Assume  $\forall n \in \mathbb{N}, \exists x_n \in X$  s.t.

$B(x_n, \frac{1}{n}) \not\subseteq U_i, \forall i \in J$ . From the compactness of  $X$ ,  $\exists$

$(x_{n_k}) \subset (x_n)$  that converges to some  $x \in X$ . Let  $x \in U_{i_0}$ . Since  $U_{i_0}$  is open,  $\exists m$  s.t.  $B(x, 1/m) \subset U_{i_0}$ .

From  $x_{n_k} \rightarrow x$ ,  $\exists n_w \geq 2m$  s.t.  $x_{n_w} \in B(x, \frac{1}{2m})$ . Thus

$$B\left(x_{n_w}, \frac{1}{n_w}\right) \subset B\left(x_{n_w}, \frac{1}{2m}\right) \subset B\left(x, \frac{1}{m}\right) \subset U_{i_0},$$

which contradicts the assumption that

$B(x_n, 1/n) \not\subseteq U_i$  for all  $n \in \mathbb{N}$  and  $i \in J$ .

Theorem (Borel–Lebesgue)  $(X, d)$  is compact  $\Leftrightarrow$  every open covering  $\{U_i\}_{i \in J}$  of  $X$  contains a finite subcovering.

Proof.  $\Rightarrow$ : Let  $\{U_i\}_{i \in J}$  be an open covering of a compact  $X$ .  $\exists r > 0$  s.t. for every  $x \in X$ , we have  $B(x, r) \subset U_i$  for some  $i \in J$ . Let's prove that  $X$  can be covered by a finite number of  $B(x, r)$ .

If  $B(x_1, r) = X$  for some  $x_1 \in X$ , we are done.

Otherwise, choose  $x_2 \in X \setminus B(x_1, r)$ . If

$B(x_1, r) \cup B(x_2, r) = X$ , the proof is over. If by continuing this process we obtain  $X$  on some step, the proof is over. Otherwise,  $\exists (x_n) \subset X$  such that

$$x_{n+1} \notin B(x_1, r) \cup \dots \cup B(x_n, r)$$

for every  $n \in \mathbb{N}$ . Since  $d(x_n, x_m) \geq r, \forall m, n \in \mathbb{N}$ ,  $(x_n)$  has no convergent subsequence, a contradiction.

$\Leftarrow$ : Let  $(x_n) \subset X$  and assume that  $(x_n)$  has no convergent subsequence. Then  $\forall x \in X, \exists$  an open ball  $B(x, r_x)$  that contains no points of the sequence  $(x_n)$  except possibly  $x$  itself.  $\{B(x, r_x)\}_{x \in X}$  is an open covering of  $X$  and thus contains a finite subcovering. Hence,

$$X = B(x_1, r_1) \cup \dots \cup B(a_n, r_n),$$

for a finite set  $A = \{a_1, \dots, a_n\}$  in  $X$ . By the choice of  $B(x, r_x)$ , we have  $x_k \in A$  for all  $k \in \mathbb{N}$ , which contradicts the assumption that  $(x_n)$  has no convergent subsequence.

- A subset of  $\mathbb{K}^n$  (with the usual metric) is compact if and only if it is closed and bounded.

**Theorem** Suppose that  $K$  is a compact subset of  $(X, d_X)$  and that  $f : (X, d_X) \rightarrow (Y, d_Y)$  is continuous. Then  $f(K)$  is compact.

**Proof** Take  $\{f(x_n)\} \subset f(K)$ . There exists  $\{x_{n_k}\}$  such that  $x_{n_k} \rightarrow x_0 \in K$  since  $K$  is compact. Therefore,  $f(x_{n_k}) \rightarrow f(x_0) \in f(K)$  by the continuity of  $f$ . □

**Proposition** Let  $K$  be a compact subset of  $(X, d)$ . Then any continuous function  $f : K \rightarrow \mathbb{R}$  is bounded and attains its bounds.

**Proof**  $f(K) \subset \mathbb{R}$  is compact and so is bounded and closed. Let  $l = \sup\{f(x), x \in K\}$ ; then  $l < \infty$ . Take  $\{f(x_n)\} \subset f(K)$  so that  $f(x_n) \rightarrow l$ ; then  $l \in f(K)$  since  $f(K)$  is closed. Hence, there is a  $z \in K$  such that  $l = f(z)$ . Similarly, there is a  $y \in K$  such that  $f(y) = \inf\{f(x), x \in K\}$ . □

**Lemma** If  $f : (X, d_X) \rightarrow (Y, d_Y)$  is continuous and  $X$  is compact, then  $f$  is uniformly continuous on  $X$  :  $\forall \epsilon > 0 \exists \delta > 0$  such that

$$d_X(x, y) < \delta \Rightarrow d_Y(f(x), f(y)) < \epsilon, \quad x, y \in X. \quad (0.3)$$

**Proof** If  $f$  is not uniformly continuous then  $\exists \epsilon > 0$  s.t.  $\forall \delta > 0$ ,  $\exists x, y \in X$  with  $d_X(x, y) < \delta$  and  $d_Y(f(x), f(y)) \geq \epsilon$ . Taking  $\delta = 1/n$ , we can find  $x_n, y_n \in X$  such that

$$d_X(x_n, y_n) < 1/n \text{ and } d_Y(f(x_n), f(y_n)) \geq \epsilon. \quad (0.4)$$


Since  $X$  is compact,  $\exists \{x_{n_k}\} \subset \{x_n\}$  s.t.  $x_{n_k} \rightarrow x$ . It follows that  $y_{n_k} \rightarrow x$  also. Since  $f$  is continuous at  $x$ , we can find  $\delta > 0$  such that  $d_X(z, x) < \delta$  ensures that  $d_Y(f(z), f(x)) < \epsilon/2$ . Thus for  $j$  sufficiently large we have  $d_X(x_{n_j}, x) < \delta$ ,  $d_X(y_{n_k}, x) < \delta$ . Hence

$$d_Y(f(x_{n_j}), f(y_{n_j})) \leq d_Y(f(x_{n_j}), f(x)) + d_Y(f(y_{n_j}), f(x)) < \epsilon,$$

contradicting (0.4). □

A subset  $B \subseteq X$  is **totally bounded** when it can be covered by a finite number of  $\epsilon$ -balls, however small their radii  $\epsilon$ ,

$$\forall \epsilon > 0, \exists N \in \mathbb{N}, \exists a_1, \dots, a_N \in X, \quad B \subseteq \bigcup_{n=1}^N B_\epsilon(a_n).$$



► A totally bounded space  $X$  is separable.

*Proof* For each  $n = 1, 2, \dots$ , consider finite covers of  $X$  by balls  $B_{1/n}(a_{i,n})$  and let  $A_n := \{a_{i,n}\}$  be the finite set of the centers, so  $A := \bigcup_{n=1}^{\infty} A_n$  is countable. For any  $\epsilon > 0$  and any point  $x \in X$ , let  $n \geq 1/\epsilon$ , then  $x$  is covered by some ball  $B_{1/n}(a_{i,n})$ , i.e.,  $d(x, a_{i,n}) < \epsilon$ , thus  $\bar{A} = X$ .

**A uniformly continuous function maps totally bounded sets to totally bounded sets.**

*Proof* Let  $f: X \rightarrow Y$  be a uniformly continuous function,

$$\forall \epsilon > 0, \exists \delta > 0, \forall x \in X, \quad f B_\delta(x) \subseteq B_\epsilon(f(x)).$$

Let  $A$  be a totally bounded subset of  $X$ , covered by a finite number of balls  $A \subseteq \bigcup_{n=1}^N B_\delta(x_n)$ . Then

$$f A \subseteq \bigcup_{n=1}^N f B_\delta(x_n) \subseteq \bigcup_{n=1}^N B_\epsilon(f(x_n)).$$

A subset  $A$  of a topological space is said to be *relatively compact* or *precompact*, if its closure is compact.

*Every relatively compact subset of a metric space  $X$  is totally bounded; if the space  $X$  is complete then every totally bounded subset of  $X$  is relatively compact.*

*It is easy to show that a subset  $M$  of a metric space is relatively compact if and only if every sequence  $(x_n)$  has a convergent subsequence; in this case the limit of the subsequence need not be in  $M$ .*

**Proposition.** *A set  $A$  in a metric space  $X$  is totally bounded precisely when every infinite sequence of its elements contains a Cauchy subsequence.*

PROOF. Let  $A$  be totally bounded and let  $\{x_n\} \subset A$  be infinite. Let us cover  $A$  by finitely many balls of radius  $1$ . At least one ball  $U_1$  of this cover contains an infinite part of  $\{x_n\}$ . The set  $A \cap U_1$  can be covered by finitely many balls of radius  $1/2$ . We can find among them a ball  $U_2$  such that  $U_1 \cap U_2$  contains an infinite part of  $\{x_n\}$ . Continuing by induction, for every  $n$  we obtain a ball  $U_n$  of radius  $1/n$  with the property that  $V_n := U_1 \cap \cdots \cap U_n$  contains infinitely many points of the original sequence. Now we can find pairwise distinct elements  $x_{k_n} \in V_n$ . Clearly, we have obtained a Cauchy sequence.

Conversely, suppose that  $A$  possesses the indicated property. Suppose that for some  $\varepsilon > 0$  there is no finite  $\varepsilon$ -net in  $A$ . By induction we construct a sequence of points  $a_n \in A$  with mutual distances at least  $\varepsilon$ : for  $a_1$  we take an arbitrary element of  $A$ ; if points  $a_1, \dots, a_n$  are already constructed, there exists a point  $a_{n+1}$  with the distances at least  $\varepsilon$  to all of them, since otherwise the sets  $a_1, \dots, a_n$  would form an  $\varepsilon$ -net. Such a sequence does not contain a Cauchy subsequence.  $\square$