

# MA302 Homework 11

WANG Duolei, SID:12012727

wangdl2020@mail.sustech.edu.cn

## 13.5 Solution

1.  $T^*$  is invertible and  $(T^*)^{-1} = (T^{-1})^*$ .

Firstly,  $T$  is invertible, which means there exists  $S \in B(K, H)$  s.t.  $TS = id_K$ ,  $ST = id_H$ . And note the Hilbert Adjoint of  $S$  by  $S^*$ , we know that  $S^* \in B(H, K)$ . Claim that  $(T^*)^{-1} = S^*$ .

- (a)  $T^*S^* = id_H$ . Firstly,  $T^*S^* \in B(H)$  is clearly because of the combination of two bounded linear map is also bounded linear. And

$$(T^*S^*x, y)_H = (x, (T^*S^*)^*y) = (x, STy)_H = (x, y)_H, \forall x \in H, y \in K$$

thus,  $T^*S^* = id_H$ .

- (b)  $S^*T^* = id_K$ . One can prove this like (a).

Thus,  $(T^*)^{-1} = S^* = (T^{-1})^*$ .

2. Consider  $T \in B(H)$  is self-adjoint. According to 1's conclusion, we know that  $T^{-1} = (T^*)^{-1} = (T^{-1})^*$ . Thus  $T^{-1}$  is also a self-adjoint operator.

## 14. Solution

- (i) Considering that  $\{\alpha_i\}_{i=1}^\infty \subseteq \sigma_p(D_\alpha)$  is clearly. We consider the inverse. For every  $\alpha \in \sigma_p(D_\alpha)$ , we have some nonzero  $x_0$  such that

$$D_\alpha x_0 = (\alpha_1 x_1, \alpha_2 x_2, \dots) = \beta x_0$$

As  $x_0 \neq 0$ , there must exist some coordinates be nonzero, and consider the nonzero terms as  $x_{i_1}, x_{i_2}, \dots$ , we know that

$$\beta = \alpha_{i_1} = \alpha_{i_2} = \dots$$

thus,  $\beta \in \{\alpha_i\}_i^\infty$

1. Considering a  $\lambda \in \mathbb{C}$  and  $\lambda \notin \overline{\sigma_p(D_\alpha)}$ , then  $|\lambda - \alpha_i| \geq \delta$ . Thus consider every coordinates, we know it's  $(\alpha_i - \lambda)x_i$ , and the norm is also bigger than  $\delta$ . Thus, the  $D_\alpha - \lambda I$  is invertible, which has a bounded inverse. Thus  $D_\alpha - \lambda I$  is also invertible. And the spectrum is closed, the two set are same is clear.

## 14.4 Solution

Considering the Spectral Mapping Theorem, we know that

$$\lambda \in \sigma(T) \implies \lambda^n \in \sigma(T^n)$$

thus  $(r_\sigma(T))^n = r_\sigma(T^n)$ . And  $\sigma(T^n) \subset \{\lambda : \lambda \leq \|T^n\|\}$ , which means the upper bound of  $r_\sigma(T)$  is  $\|T^n\|^{1/n}$ . Thus

$$r_\sigma(T) \leq \liminf_{n \rightarrow \infty} \|T^n\|^{1/n}$$

## 14.5 Solution

Consider this operator is just a special case in 11.7, we know that the bound is  $\|T^n\|_{B(X)} \leq \frac{1}{n!}$ . And according to 14.4

$$r_\sigma(T) \leq \liminf_{n \rightarrow \infty} \|T^n\|^{1/n} = 0$$

Thus, all the eigenvalues must be zero. And we know that  $T$  is not invertible, as the 0 is not an eigenvalue, which is because

$$|(T - 0I)f| = 0 \implies Tf = 0 \implies f = 0$$

This also means  $T$  is also not surjective.