



## Antimicrobial activity of essential oils

**Bruna Fernanda Murbach Teles Andrade, Lidiane Nunes Barbosa, Isabella da Silva Probst & Ary Fernandes Júnior**

**To cite this article:** Bruna Fernanda Murbach Teles Andrade, Lidiane Nunes Barbosa, Isabella da Silva Probst & Ary Fernandes Júnior (2014) Antimicrobial activity of essential oils, Journal of Essential Oil Research, 26:1, 34-40, DOI: [10.1080/10412905.2013.860409](https://doi.org/10.1080/10412905.2013.860409)

**To link to this article:** <https://doi.org/10.1080/10412905.2013.860409>



Published online: 29 Nov 2013.



Submit your article to this journal [↗](#)



Article views: 4591



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 50 View citing articles [↗](#)

## Antimicrobial activity of essential oils

Bruna Fernanda Murbach Teles Andrade, Lidiane Nunes Barbosa, Isabella da Silva Probst and  
Ary Fernandes Júnior\*

Department of Microbiology and Immunology, Institute of Biosciences, Universidade Estadual Paulista Júlio  
de Mesquita Filho-UNESP, São Paulo, Brazil

(Received 11 July 2012; accepted 17 October 2013)

Natural products have been studied aiming to understand their biological properties. Thus, this study aimed to investigate the antimicrobial activity of twenty-seven essential oils (EOs) used in aromatherapy procedures, a natural therapy with great emphasis currently used against *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* strains. The agar dilution method was carried out and minimal inhibitory concentration against 50% and 90% of strains (MIC<sub>50%</sub> and MIC<sub>90%</sub> values) were reported. The *S. aureus* strains were highly susceptible with MIC<sub>90%</sub> from 0.21 mg/mL to black pepper (*Piper nigrum*) and tea tree (*Melaleuca alternifolia*) to 26.52 mg/mL with copaiba (*Copaifera officinalis*) EO. Cinnamon (*Cinnamomum cassia*) and clove (*Syzygium aromaticum*) EOs were effective against *E. coli* (2.0 mg/mL) while the *S. aromaticum* EO was against *P. aeruginosa* (8.29 mg/mL). Thus, the higher susceptibility of Gram-positive bacteria when compared with Gram-negative strains was found, and a large variability in the potential antibacterial has also been observed.

**Keywords:** antibacterial; aromatherapy; EO; minimal inhibitory concentration

### Introduction

Herbs, and their essential oils (EOs), have been used since the beginning of human history for flavored foods and beverages; they have been empirically used to disguise unpleasant odors, attract other individuals and control health problems, contributing to the welfare humans and animals, thus demonstrating the cultural and economic importance use of these products (1).

The EOs are typically liquid, clear and unusually colored, complex and the present compounds are volatile, characterized by a strong odor and synthesized by aromatic plants during secondary metabolites, which act to protect the plant against microorganisms and insects. They can be synthesized in several plant organs such as buds, flowers, leaves, stems, branches, seeds, berries, roots, wood or bark, being stored in secretory cells, cavities, channels, epidermal cells or trichomes (2). Temporal and spatial variations in the total content of secondary metabolites products from plants occur at different levels and, despite the existence of a genetic control, the expression may undergo changes resulting from biochemical, physiological, ecological and evolutionary interactions that represent an important interface between chemistry and the environment surrounding the plants (3).

As for industrial production, EOs are obtained by steam distillation, which is on the rise in food and

pharmaceutical applications, and pressurized supercritical fluid, especially carbon dioxide (4).

EOs have several biological properties, such as larvicidal action (5), antioxidant (6), analgesic and anti-inflammatory (7), fungicide (8) and antitumor activity (9).

The *in vitro* antimicrobial activity of EO has been researched extensively against a variety of microorganisms (10). Nevertheless, the emergence of multidrug-resistant bacteria poses a challenge to treating infections, so the need to find new substances with antimicrobial properties for use in the fight against these microorganisms is evident (11, 12). Historically, most antibiotics come from a small set of functional molecular structures whose lives were extended by generations of synthetic reorganizations and arrangements (13). Moreover, the food, pharmaceutical and cosmetic industries have shown great interest in the antimicrobial properties of EOs, as the use of natural additives has received importance as a trend in the replacement of synthetic preservatives (14).

The objective was to establish *in vitro* the antimicrobial activities of EOs that are normally used in natural therapies against *S. aureus*, *E. coli* and *P. aeruginosa* strains isolated from human clinical specimens and one standard ATCC (American Type Culture Collection) of each bacterial species; this was

---

\*Corresponding author. Email: [ary@ibb.unesp.br](mailto:ary@ibb.unesp.br)

Table 1. Density (mg/mL) and essential oils chemical compounds (%) obtained by chromatography–mass spectrometry (GC–MS) from the supplier of the essential oils (By Samia Aromatherapy/São Paulo/Brazil).

Essential oil	Density (mg/mL)	Compounds in essential oils (%)
Bergamot ( <i>Citrus aurantium bergamia</i> )	871	Limonene (35.24), linalina acetate (30.40), linalool (18.45), $\beta$ -pinene (5.42), $\gamma$ -terpinene (3.74), sabinene (0.92), $\alpha$ -pinene (0.89), myrcene (0.81), for-cymene (0.45)
Black pepper ( <i>Piper nigrum</i> )	846	Limonene (23.80), $\delta$ -3-carene (21.97), $\alpha$ -pinene (12.89), $\beta$ -caryophyllene (11.34), $\beta$ -pinene (3.91), sabinene (3.78), $\alpha$ -felandeno (3.76), myrcene (2.88), para-cymene (1.38), linalool (1.24), terpinolene (1.17), $\beta$ -selineno (1.11), 1.8 cineole (0.98), $\alpha$ -terpinene (0.97), $\alpha$ -humulene (0.77), $\alpha$ -copaene (0.71), eugenol (0.56), terpinen-4-ol (0.47), camphene (0.21), saffrole (0.17)
Brazil's spearmint ( <i>Mentha arvensis</i> )	849	Menthol (54.48), menthone (19.12), pulegone (5.57), isopulegol (2.02)
Cardamom ( <i>Elettaria cardamomum</i> )	869	n/d
Cedar ( <i>Cedrus atlantica</i> )	891	Widreno (27.75), $\alpha$ -cedrol (22.14), $\alpha$ -Cedrenus (19.84), $\alpha$ -muuroleno (4.55), widrol (3.79)
Cinnamon ( <i>Cinnamomum cassia</i> )	1008	Eugenol (72.13), eugenila acetate (3.87), $\beta$ -caryophyllene (3.48), benzyl benzoate (3.24), linalool (1.23), para-cymene (0.76), $\alpha$ -pinene (0.63), $\alpha$ -humulene (0.61), $\alpha$ -phellandrene (0.49), 1.8 cineole (0.27), limonene (0.22), camphene (0.21), $\beta$ -pinene (0.21)
Clary sage ( <i>Salvia sclarea</i> )	857	Linalina acetate (66.77), linalool (22.67), geranyl acetate (3.29), $\beta$ -caryophyllene (1.15), myrcene (0.18), limonene (0.15), 1.8 cineole (0.12)
Clove ( <i>Syzygium aromaticum</i> )	988	Eugenol (83.63), $\beta$ -caryophyllene (12.39), $\alpha$ -humulene (3.05), eugenol acetate (0.93)
Copaiba ( <i>Copaifera officinalis</i> )	884	$\beta$ -Caryophyllene (44.47), $\beta$ -bisabolene (8.0), germacrene B (8.0), $\alpha$ -copaene (7.98), germacrene d (5.95), $\alpha$ -humulene (5.40), $\delta$ -cadinene (4.57)
Cypress ( <i>Cupressus sempervirens</i> )	840	$\alpha$ -Pinene (52.26), $\delta$ -3-carene, $\alpha$ -terpinolene (2.65), $\alpha$ -terpinila acetate (2.63), limonene (2.60), myrcene (2.40), terpinen-4-ol (1.40), sabinene (1.24), $\beta$ -pinene (1.14), $\alpha$ -tujeno (0.96), $\alpha$ -fenchene (0.81), $\gamma$ -terpinene (0.79), p-cymene (0.70), geranyl acetate (0.41), $\alpha$ -terpinene (0.35), 1.8 cineole (0.34), camphene (0.28)
Eucalyptus ( <i>Eucalyptus globulus</i> )	883	1.8 Cineole (80.17), $\alpha$ -pinene (11.25), diacetone alcohol (4.32), p-cymene (2.28), $\alpha$ -terpineol (0.85), terpinen-4-ol (0.60), $\beta$ -pinene (0.53)
Fennel ( <i>Foeniculum vulgare</i> )	919	<i>trans</i> -Anethole (95.66), linalool (2.91), estragol (0.39), $\alpha$ -pinene (0.13)
Geranium ( <i>Pelargonium graveolens</i> )	848	Citronellol (31.58), geraniol (25.47), ferriato of citronelita (12.74), ferriato of geranyl (6.71), linalool (6.33); isomenthone (4.35), rose oxide (0.89), citronelita acetate (0.48)
Ginger ( <i>Zingiber officinalis</i> )	850	$\alpha$ -Zingiberene (22.85), curcumene (18.96), $\beta$ -sesquifilandro (13.12), $\beta$ -bisabolene (11.58), $\alpha$ -farnesene (4.28), camphene (1.77), $\beta$ -phellandrene (1.58), 1.8 cineole (1.35), $\alpha$ -pinene (0.43) <i>trans</i> - $\beta$ -farnesene (0.30), myrcene (0.20)
Lavender ( <i>Lavandula officinalis</i> )	853	1.8 Cineole (45.97), p-cymene (4.19), 1-terpinen-4-ol (2.30), $\alpha$ -pinene (1.48), limonene (1.46), gamma-terpinene (1.17), terpinolene (1.04)
Lemongrass ( <i>Cymbopogon schoenanthus</i> )	858	Geraniol (48.57), neral (32.86), geranyl acetate (3.98), $\beta$ -caryophyllene (1.59), linalool (1.23), camphene (1.19), caryophyllene oxide (0.67), eugenol (0.48), limonene (0.23), $\alpha$ -pinene (0.20), <i>trans</i> - $\beta$ -ocimene (0.12)
Marjoram ( <i>Origanum majorana</i> )	841	1.8 Cineole (48.05), linalool (22.69), limonene (8.10), $\alpha$ -pinene (4.42), $\beta$ -pinene (4.05), isobornyl acetate (2.82), para-cymene (2.21), estragol (1.02), $\gamma$ -terpinene (0.96), camphene (0.74), viridiflorol (0.73), myrcene (0.51), borneol (0.49) <i>trans</i> -linalool oxide (0.24), <i>cis</i> -linalool oxide (0.21)
Nutmeg ( <i>Myristica fragans</i> )	889	$\alpha$ -Pinene (18.35), myristicin (17.65), $\beta$ -pinene (12.29), sabinene (10.15), terpinen-4-ol (8.21), $\gamma$ -terpinene (4.18), limonene (3.63), para-cymene (3.15), $\alpha$ -terpinolene (2.91), saffrole (2.68), 1.8 cineole (2.16), terpinolene (1.84), methyl eugenol (1.59), $\alpha$ -terpineol (1.52), $\delta$ -3-carene (1.41), elemicin (0.74), eugenol (0.53)
Orange ( <i>Citrus aurantium dulcis</i> )	820	Limonene (96.25), myrcene (1.81), linalool (0.49), $\alpha$ -pinene (0.49), sabinene (0.32), $\beta$ -phellandrene (0.27)
Palmarosa ( <i>Cymbopogon martinii</i> )	874	Geraniol (57.49), geranyl acetate (13.56), linalool (1.71), $\beta$ -caryophyllene (1.07), ocimene (0.27)
Patchouli ( <i>Pogostemon patchouli</i> )	1009	Patchoulol (25.21), $\delta$ -guaiano (11.49); gurjunene- $\alpha$ (11.26); seicheleno (9.61), $\alpha$ -guaiano (9.56), benzyl alcohol (6.73), vidreno (3.12), aromadendrene (2.81), $\alpha$ -cedrol (2.63), $\beta$ -patchouleno (1.57)

(Continued)

Table 1. (Continued).

Essential oil	Density (mg/mL)	Compounds in essential oils (%)
Pine ( <i>Pinus sylvestris</i> )	874	Bornyl acetate (32.74), camphene (21.67), $\alpha$ -pinene (10.95), limonene (4.42), 1.8 cineole (3.15), borneol (3.11), $\beta$ -pinene (1.82), $\beta$ -caryophyllene (1.53), terpinolene (1.01), myrcene (0.54), geranyl acetate (12.34), camphor (0.22), para-cymene (0.14), $\gamma$ -terpinene (0.12)
Rosemary ( <i>Rosmarinus officinallis</i> )	885	1.8 Cineole (31.57), camphor (20.42), $\alpha$ -pinene (15.78), camphene (4.93), limonene (3.76), geraniol (2.43), myrcene (2.02), linalool (1.70), para-cymene (1.66), $\gamma$ -terpinene (1.14), $\alpha$ -terpinolene (0.99), bornyl acetate (0.41), borneol (0.15)
Tahiti lime ( <i>Citrus limonum</i> )	840	Limonene (62.34), $\gamma$ -terpinene (11.96), $\beta$ -pinene (10.23), $\beta$ -bisabolene (2.68), $\alpha$ -pinene (1.97), geraniol (1.84), myrcene (1.49), para-cymene (1.18), neral (1.04), <i>trans</i> - $\alpha$ -bergamotene (1.02), $\alpha$ -tujeno (0.50)
Tea tree ( <i>Melaleuca alternifolia</i> )	858	1-Terpinen-4-ol (53.40), p-cymene (8.9), gamma-terpinene (5.34), 1.8 cineole (3.18), $\alpha$ -pinene (1.40), terpinolene (1.05), limonene (0.70)
Vetiver ( <i>Vetiveria zizanioides</i> )	977	n/d
Ylang ylang ( <i>Cananga odorata</i> )	904	<i>trans</i> - $\beta$ -Caryophyllene (12.92), linalool (11.38), germacrene-d (11.21), benzyl acetate (10.34), geranyl acetate (9.87).

Note: n/d, information not obtained.

performed by the dilution of EOs onto Mueller Hinton agar (MHA) and the minimal inhibitory concentration (MIC) against each and the MIC<sub>50%</sub> and MIC<sub>90%</sub> values were recorded.

## Experimental

### Essential oils

Twenty-seven samples of EOs, from the supplier By Samia Aromatherapy (São Paulo-SP, Brazil) in amber glass vials with a capacity of 10 mL, were selected according to their frequent use in aromatherapy procedures. These EOs were: bergamot (*Citrus aurantium bergamia*), black pepper (*Piper nigrum*), Brazil's spearmint (*Mentha arvensis*), cardamom (*Elettaria cardamomum*), cedar (*Cedrus atlantica*), cinnamon (*Cinnamomum cassia*), clary sage (*Salvia sclarea*), clove (*Syzygium aromaticum*), copaiba (*Copaifera officinalis*), cypress (*Cupressus sempervirens*), eucalyptus (*Eucalyptus globulus*), fennel (*Foeniculum vulgare*), geranium (*Pelargonium graveolens*), ginger (*Zingiber officinalis*), lavender (*Lavandula officinalis*), lemongrass (*Cymbopogon schoenanthus*), marjoram (*Origanum majorana*), nutmeg (*Myristica fragans*), orange (*Citrus aurantium dulcis*), palmarosa (*Cymbopogon martinii*), patchouli (*Pogostemon patchouli*), pine (*Pinus sylvestris*), rosemary (*Rosmarinus officinallis*), Tahiti lime (*Citrus limonum*), tea tree (*Melaleuca alternifolia*), vetiver (*Vetiveria zizanioides*) and ylang ylang (*Cananga odorata*). The samples were kept at room temperature and the chemical characterization of the oils samples were provided by supplier, including the gas chromatography-mass spectrometry (GC-MS) analysis (Table 1). Density values from each studied oil were performed using methodology recommended by Fonseca and Librandi (15) in

Eppendorf tubes, which were weighed (P1) on an analytical balance and then weighed again (P2) after the addition of 1 mL (V) of oil. The density (D) was calculated using the formula below.

$$D = \frac{P_2 - P_1}{V} = \frac{mg}{mL}$$

### Bacterial strains

The bacterial strains from the American Type Culture Collection (ATCC) standard strains (*E. coli* ATCC 25922, *S. aureus* ATCC 25923 and *P. aeruginosa* ATCC 27853), as well as ten *S. aureus*, ten *E. coli* and nine *P. aeruginosa* isolated from human clinical specimens, were selected from strains stored at  $-80^{\circ}\text{C}$  in the Department of Microbiology and Immunology of Biosciences Institute of UNESP, Botucatu-SP. Prior to use, the strains were plated on blood agar medium to check viability and purity, and maintained on nutrient agar for use in bacterial susceptibility assays.

### Antimicrobial activity of EOs by the agar dilution method and MIC

Susceptibility tests for determining the MIC of EOs were carried out following the agar dilution method, adapted from CLSI (16) protocol, the plate discs were prepared and antimicrobial assays were performed during one day. Each EO was diluted alone in MHA plus 0.5% Tween 80 at  $45^{\circ}\text{C}$  in Petri dishes and equivalent concentrations of 0.025, 0.05, 0.1, 0.2, 0.5, 0.8, 1.0, 1.5, 2.0, 2.5 and 3.0% v/v were established. The strains were grown at  $35^{\circ}\text{C}/18\text{--}24$  hours in brain heart infusion (BHI) and standard suspensions were performed in sterile saline (0.85%) using scale 0.5 of MacFarland aiming a

bacterial concentration about  $1.5 \times 10^8$  colony forming units (CFU)/mL. The inoculation of thirty-two strains, from standardized suspensions, was made using a Sterr multi-inoculator using suspensions standardized at 0.5 MacFarland, with a second dilution performed onto BHI to obtain inoculum of an approximate concentration of  $10^5$ – $10^6$  CFU/mL. After Petri dishes were inoculated and incubated at 35°C/18–24 hours, bacterial growth was assessed and the MIC values were recorded for each strain. The conversion of values from % v/v to mg/mL, using the density values of each oil, and their calculations of the MIC<sub>50%</sub> and MIC<sub>90%</sub> for each tested bacterial strains were performed.

### Statistical analysis

The results obtained were used to compare three or more independent testing treatments via the Kruskal–Wallis test. For meaningful analysis ( $p \leq 0.001$ ), we apply the Student–Newman–Keuls test for multiple comparisons tests between treatments.

### Results and discussion

The research about antimicrobial activity, the action mechanism and potential use of volatile plant oils has received prominence in recent decades in parallel with advances in traditional approaches to protecting the health of humans, animals and food against the presence of pathogenic and spoilage microorganisms. Thus, investigations on the antimicrobial activity of plant extracts against different pathogens have been performed worldwide (17); our results have importance because they provide information about this subject.

The density (mg/mL), chemical compounds and their percentages in the total composition of the each EO (twenty-seven samples) were found; results are presented in Table 1. We emphasize that the data about chemical analysis of the oils studied were received from the company By Samia Aromatherapy who supplied the EO samples. All of the oils studied showed a density above 800 mg/mL, with orange oil having the lowest value (820 mg/mL) and patchouli (1009 mg/mL), cinnamon (1008 mg/mL) and clove (988mg/mL) presenting the highest density values. Although the chemical characterization of the oils plays a role in studies of this nature, according to some authors, it cannot be concluded that the major component is the biologically active compound of this study, so the effect can be attributed to a constituent or lesser extent a synergy between existing compounds in the oil (18–21). In general, the EO showed diversity in their chemical characterization, but these are in agreement with the literature in question.

A total of twenty-seven oils were assayed by the agar dilution method. The MIC<sub>90%</sub> values against bacterial strain tested (Table 2) show that *S. aureus* strains were susceptible to a high number of EOs, and eight of the twenty-seven oils tested showed inhibitory activity with MIC<sub>90%</sub> values below 0.30 mg/mL (e.g. eucalyptus, lemongrass, patchouli, black pepper, clary sage, tea tree, vetiver, ylang ylang).

With the tests using the agar dilution methodology at the concentrations tested, it was not possible to achieve MIC values for *P. aeruginosa* strains, except for cinnamon; these results corroborate those reported by Hammer et al. (21), who determined the MIC against *E. coli* and *S. aureus* but failed to find results for *P. aeruginosa* strains.

Thus, the *S. aureus* strains showed high susceptibility to natural products, which again confirms the results from the literature (22, 23), or in other words, Gram-positive species are more sensitive to natural products than Gram-negative bacteria.

These data are important for the treatment of infections caused by these bacteria; *S. aureus* is described as one of the main agents responsible for infection, as its virulence and ability to acquire antimicrobial resistance results in a serious problem throughout the world for hospitals and health professionals (24).

*Pseudomonas aeruginosa* is a Gram-negative bacterium that produces water-soluble pigments, which is widely distributed in soil and water, and is a hospital pathogen that grows in damp areas such as sinks, bathtubs and showers; it is also considered a resistant bacterium (25). Despite the fact that *E. coli* is a Gram-negative bacteria as well as *P. aeruginosa*, this bacterium showed sensitivity to fourteen oils at the highest concentrations tested. These results corroborate with those reported by Duarte et al. (26), who concluded that the *C. martini* (palmarosa) EO and its major component, geraniol, may be useful for the treatment of diarrhoea caused by *E. coli*.

EOs mainly include two biosynthetic groups, all characterized by low molecular weight, including aromatic and aliphatic constituents, and terpenes and terpenoids (27).

As a typical lipophilic compound, EOs cross the cell wall and cytoplasmic membrane and the cytotoxic activity appears to be linked to disruption of the structures of the different layers of polysaccharides, fatty acids and phospholipids, due to its mechanism of action that hits multiple targets at the same time (22). Permeability, composition and charge of the outer structures of the microorganisms mainly determined these differences; the lipophilic character of terpenes is associated with the antimicrobial mechanism (28). Numerous reports have been made about the mechanisms of antimicrobial action of the oils, and some cases have been partly elucidated,



Table 2. Minimal inhibitory concentration 50% (MIC<sub>50%</sub>) and 90% (MIC<sub>90%</sub>) (mg/mL) found on essential oils samples against ATCC standard and *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* strains isolated from clinical human specimens.

Essential oil	<i>S. aureus</i> (n=11)*, CIM <sub>50%</sub> –CIM <sub>90%</sub>	<i>E. coli</i> (n=11)*, CIM <sub>50%</sub> –CIM <sub>90%</sub>	<i>P. aeruginosa</i> (n=10)**, CIM <sub>50%</sub> –CIM <sub>90%</sub>
Bergamot ( <i>Citrus aurantium bergamia</i> )	10.50–19.81 <sup>w</sup>	>26.13–>26.13	>26.13–>26.13
Black pepper ( <i>Piper nigrum</i> )	0.21–0.21 <sup>a</sup>	>25.38–>25.38	>25.38–>25.38
Brazil's spearmint ( <i>Mentha arvensis</i> )	1.90–2.26 <sup>l</sup>	5.52–5.52	>25.47–>25.47
Cardamom ( <i>Elettaria cardamomum</i> )	7.58–7.58 <sup>s</sup>	>26.07–>26.07	>26.07–>26.07
Cedar ( <i>Cedrus atlantica</i> )	1.78–2.76 <sup>nl</sup>	22.27–26.73 <sup>k</sup>	>26.73–>26.73
Cinnamon ( <i>Cinnamomum cassia</i> )	1.00–1.14 <sup>ok</sup>	2.00–2.00 <sup>b</sup>	25.00–30.0 <sup>ba</sup>
Clary sage ( <i>Salvia sclarea</i> )	0.29–0.29 <sup>hf</sup>	>25.71–>25.71	>25.71–>25.71
Clove ( <i>Syzygium aromaticum</i> )	0.67–1.21 <sup>k</sup>	1.11–2.00 <sup>a</sup>	4.60–8.29 <sup>a</sup>
Copaiba ( <i>Copaifera officinalis</i> )	24.07–26.52 <sup>z</sup>	>26.52–>26.52	>26.52–>26.52
Cypress ( <i>Cupressus sempervirens</i> )	>25.2–>25.2	>25.20–>25.20	>25.20–>25.20
Eucalyptus ( <i>Eucalyptus globulus</i> )	0.22–0.22 <sup>c</sup>	11.00–14.35 <sup>h</sup>	>26.49–>26.49
Fennel ( <i>Foeniculum vulgare</i> )	7.81–7.81 <sup>us</sup>	13.08–20.22 <sup>l</sup>	>27.57–>27.57
Geranium ( <i>Pelargonium graveolens</i> )	0.20–0.31 <sup>ge</sup>	3.90–4.24 <sup>ec</sup>	>25.40–>25.40
Ginger ( <i>Zingiber officinalis</i> )	3.23–4.93 <sup>qp</sup>	>25.5–>25.5	>25.50–>25.50
Lavender ( <i>Lavandula officinalis officinalis</i> )	2.37–4.27 <sup>t</sup>	21.3–25.59 <sup>ml</sup>	>25.59–>25.59
Lemongrass ( <i>Cymbopogon schoenanthus</i> )	0.15–0.22 <sup>i</sup>	1.98–2.10 <sup>gc</sup>	>25.74–>25.74
Marjoram ( <i>Origanum majorana</i> )	4.21–4.21 <sup>p</sup>	4.21–4.21 <sup>dc</sup>	>25.23–>25.23
Nutmeg ( <i>Myristica fragrans</i> )	13.96–13.96 <sup>yx</sup>	18.52–18.52 <sup>j</sup>	>26.67–>26.67
Orange ( <i>Citrus aurantium dulcis</i> )	12.50–16.5 <sup>x</sup>	>24.63–>24.63	>24.63–>24.63
Palmarosa ( <i>Cymbopogon martinii</i> )	0.48–0.59 <sup>m</sup>	1.90–2.09 <sup>fc</sup>	>26.22–>26.22
Patchouli ( <i>Pogostemon patchouli</i> )	0.25–0.25 <sup>f</sup>	>30.27–>30.27	>30.27–>30.27
Pine ( <i>Pinus sylvestris</i> )	2.58–2.58 <sup>ji</sup>	>26.22–>26.22	>26.22–>26.22
Rosemary ( <i>Rosmarinus officinalis</i> )	6.40–7.26 <sup>t</sup>	17.70–22.12 <sup>i</sup>	>26.55–>26.55
Tahiti lime ( <i>Citrus limonum</i> )	10.0–14.91 <sup>v</sup>	>25.2–>25.2	>25.2–>25.2
Tea tree ( <i>Melaleuca alternifolia</i> )	0.21–0.21 <sup>b</sup>	4.29–4.29 <sup>c</sup>	>25.74–>25.74
Vetiver ( <i>Vetiveria zizanioides</i> )	0.24–0.24 <sup>c</sup>	>29.31–>29.31	>29.31–>29.31
Ylang ylang ( <i>Cananga odorata</i> )	0.23–0.23 <sup>d</sup>	>27.12–>27.12	>27.12–>27.12

Note: \*ATCC and plus ten clinical isolated; \*\*ATCC and plus nine clinical isolated. Values preceded by '>' were not considered in the statistical analysis because they did not show inhibitory capacity up to the maximum concentration tested in the trials. Different letters in columns represent statistical differences for antibacterial activities of essential oils (mg/mL) when  $p \leq 0.001$ .

e.g. the tea tree EO (*M. alternifolia*) and its major compound, terpinen-4-ol, which causes lysis and loss of membrane integrity due to output ions and cellular respiration inhibition (29, 30).

Cypress (*C. sempervirens*) was the only species that showed no antibacterial activity according to the susceptibility assay performed. According to Hammer et al. (21), this oil also showed no activity against *E. coli* and *P. aeruginosa*, although it has some effect on the growth of strains of *S. aureus* with an MIC value of 2% v/v. However, the authors report that a single standard NCTC strain (National Collection of Type Cultures) was used, which may explain the different results obtained, while this study tested strains isolated from human clinical cases, and therefore distinct phenotypes itself. However, *C. arizonica*, from the same family as *C. sempervirens*, has been attributed weak antimicrobial activity for this EO because of the high hydrocarbon content (31).

EO contain complex mixtures of components and thus have multiple antimicrobial properties; most of this action appears to derive from oxygenated terpenoids,

particularly phenolic terpenes, phenylpropanoids and alcohols; other constituents, e.g. hydrocarbons that typically showed low activities, can be used in combinations to increase their bioactivities (27).

In general, the oils of cinnamon and clove oils were those with the highest potential inhibitors against the three bacterial strains used. According to Prabuseenivasan et al. (32), these oils were able to inhibit the growth of both Gram-positive and Gram-negative species. Both oils showed eugenol, i.e. phenylpropanoid, to be the main compound.

The clove EO exhibited the best activity among the twenty-seven oils tested against both Gram-negative strains, but for the strains of *S. aureus*, this was black pepper oil with an MIC<sub>90%</sub> of 0.21 mg/mL.

Most of the EOs used in this study has terpinen-4-ol, linalool and eugenol as part of their compounds.

The antimicrobial mechanism involved with linalool is related to its high water solubility and to its ability to penetrate the bacteria cell wall (33). One hypothesis is that linalool has the potential to act as either a protein denaturing agent or as a solvent dehydrating agent,

which may also contribute to its antimicrobial activity (34).

The antimicrobial activity of some EOs could be explained by the significant amount of linalool, which is an oxygenated monoterpenoid (35). Knobloch et al. (36) related that linalool had a significantly increased antimicrobial activity when compared to eugenol. Overall, the antibacterial activity of the EOs can be related to the content of many of the compounds identified in the oils, including eugenol (17, 37).

Although linalool presents important antioxidant and antimicrobial effects (38), it must be noted that the antimicrobial effect of an EO depends on all of its chemical components (35).

Thus, we concluded that *P. aeruginosa* strains were highly resistant to the EOs, while the *S. aureus* strains were considerably sensitive, although the potential use of EOs can be applied to both Gram-positive and Gram-negative bacteria.

### Acknowledgements

The authors are grateful to the company By Samia Aromatherapy for providing the samples of essential oils as well as by chemical analysis of the samples. Also thank Prof. Dr. Luciano Barbosa of Department of Biostatistics/Biosciences Institute, UNESP, by statistics analysis of the results from the study.

### Conflict of interest

The authors have declared no conflict of interest

### References

- (1) C.M. Franz, *Essential oil research: Past, present and future*. Flavour Fragr. J., **25**, 112–113 (2010).
- (2) F. Bakkali, S. Averbeck, D. Averbeck and M. Idaomar, *Biological effects of essential oils – A review*. Food Chem. Toxicol., **46**, 446–475 (2008).
- (3) L. Gobbo-Neto and N.P. Lopes, *Plantas medicinais: Fatores de influência no conteúdo de metabólitos secundários*. Quim. Nova, **2**, 374–381 (2008).
- (4) M. Mazutti, B. Beledelli, A.J. Mossi, R.L. Cansian, C. Dariva, J.V. Oliveira and N. Paroul, *Caracterização química de extratos de Ocimum basilicum L. obtidos através de extração com CO<sub>2</sub> a altas pressões*. Quim. Nova, **6**, 1198–1202 (2006).
- (5) S. Rajkumar and A. Jebanesan, *Chemical composition and larvicidal activity of leaf essential oil from Clausena dentata (Willd) M. Roam. (Rutaceae) against the chikungunya vector, Aedes aegypti Linn. (Diptera: Culicidae)*. J. Asia Pac. Entomol., **13**, 107–109 (2010).
- (6) W.A. Wannes, B. Mhamdi, J. Sriti, M.B. Jemia, O. Ouchikh, G. Hamdaoui, M.E. Kchouk and B. Marzou, *Antioxidant activities of the essential oils and methanol extracts from myrtle (Myrtus communis var. italica L.) leaf, stem and flower*. Food Chem. Toxicol., **48**, 1362–1370 (2010).
- (7) S.S. Mendes, R.R. Bomfim, H.C.R. Jesus, P.B. Alves, A.F. Blank, C.S. Estevam, A.R. Antonioli and S.M. Thomazzi, *Evaluation of the analgesic and anti-inflammatory effects of the essential oil of Lippia gracilis leaves*. J. Ethnopharmacol., **129**, 391–397 (2010).
- (8) E.S. Carmo, E.O. Lima and E.L. Souza, *The potential of Origanum vulgare L. (Lamiaceae) essential oil in inhibiting the growth of some food-related Aspergillus species*. Braz. J. Microbiol., **39**, 362–367 (2008).
- (9) S.L. Silva, J.S. Chaar, P.M.S. Figueiredo and T. Yano, *Cytotoxic evaluation of essential oil from Casearia sylvestris Sw on human cancer cells and erythrocytes*. Acta Amazonica, **1**, 107–112 (2008).
- (10) P. López, C. Sánchez, R. Batlle and C. Nerín, *Solid- and vapor-phase antimicrobial activities of six essential oils: Susceptibility of selected foodborne bacterial and fungal strains*. J. Agric. Food Chem., **53**, 6939–6946 (2005).
- (11) R.S. Pereira, T.C. Sumita, M.R. Furlan, A.O.C. Jorge and M. Ueno, *Atividade antibacteriana de óleos essenciais em cepas isoladas de infecção urinária*. Rev. Saude Publica, **2**, 326–328 (2004).
- (12) S. Hemaiswarya, A.K. Kruthiventi and M. Doble, *Synergism between natural products and antibiotics against infectious diseases*. Phytomedicine, **15**, 639–652 (2008).
- (13) M.A. Fischbach and C.T. Walsh, *Antibiotics for emerging pathogens*. Science, **5944**, 1089–1093 (2009).
- (14) O.O. Okoh, A.P. Sadimenko and A.J. Afolayan, *Comparative evaluation of the antibacterial activities of the essential oils of Rosmarinus officinalis L. obtained by hydrodistillation and solvent free microwave extraction methods*. Food Chem., **120**, 308–312 (2010).
- (15) P. Fonseca and A.P.L. Librand, *Evaluation of physico-chemical and phytochemical characteristics of different tinctures of barbatimão (Stryphnodendron barbatiman)*. Braz. J. Pharm. Sci., **4**, 271–277 (2008).
- (16) Clinical and Laboratory Standards Institute/National Committee for clinical Laboratory Standards (CLSI/NCCLS). In: *Performance Standards for Antimicrobial Susceptibility Testing*; Fifteenth Information Supplement. CLSI/NCCLS document M 100–S15. Wayne, PA, (2005).
- (17) H.J.D. Dorman and S.G. Deans, *Antimicrobial agents from plants: Antibacterial activity of plant volatile oils*. J. Appl. Microbiol., **88**, 308–316 (2000).
- (18) M.M. Cowan, *Plant products as antimicrobial agents*. Clin. Microbiol. Rev., **12**, 564–582 (1999).
- (19) P.J. Houghton, M.J. Howes, C.C. Lee and G. Steventon, *Uses and abuses of in vitro tests in ethnopharmacology: Visualizing an elephant*. J. Ethnopharmacol., **110**, 391–400 (2007).
- (20) M.T. Pupo, M.B.C. Gallo and P.C. Vieira, *Biologia química: Uma estratégia moderna para a pesquisa em produtos naturais*. Quim. Nova, **6**, 1446–1455 (2007).
- (21) K.A. Hammer, C.F. Carson and T.V. Riley, *Antimicrobial activity of essential oils and other plant extracts*. J. Appl. Microbiol., **6**, 985–990 (1999).
- (22) J.E.C. Betoni, R.P. Mantovani, L.N. Barbosa, L.C. Di Stasi and A. Fernandes Júnior, *Synergism between plant extract and antimicrobial drugs used on Staphylococcus aureus diseases*. Mem. Inst. Oswaldo Cruz, **4**, 387–390 (2006).
- (23) N.C.C. Silva and A. Fernandes Junior, *Biological properties of medicinal plants: A review of their*

- antimicrobial activity*. J. Venom. Anim. Toxins Incl. Trop. Dis., **3**, 402–413 (2010).
- (24) M.J. Carvalho, F.C. Pimenta, M. Hayashida, E. Gir, A.M. Silva, C.P. Barbosa, S.R.M.S. Canini and S. Santiago, *Prevalence of methicillin-resistant and methicillin-susceptible S. aureus in the saliva of health professionals*. Clinics, **4**, 295–302 (2009).
- (25) E. Jawetz, J. Melnick and E. Aldelberg, *Microbiologia médica: Um livro médico Lange.*, McGraw-Hill Interamericana do Brasil, Rio de Janeiro (2005).
- (26) M.C.T. Duarte, E.E. Leme, C. Delarmelina, A.A. Soares, G.M. Figueira and A. Sartoratto, *Activity of essential oils from Brazilian medicinal plants on Escherichia coli*. J. Ethnopharmacol., **111**, 197–201 (2007).
- (27) I. Bassole and H. Juliani, *Essential oils in combination and their antimicrobial properties*. Molecules, **17**, 3989–4006 (2012).
- (28) P. Roman, K-S. Martyna, T. Mariusz and F. Jan, *Terpenes: Substances useful in human healthcare*. Arch. Immunol. Ther. Exp., **55**, 315–327 (2007).
- (29) S.D. Cox, J.L. Mann, H.C. Bell, J.E. Gustafson, J.R. Warmingtn and S.G. Wyllic, *The mode of antimicrobial action of the essential oils of Melaleuca alternifolia (tea tree oil)*. J. Appl. Microbiol., **88**, 170–175 (2000).
- (30) C.F. Carson, K.A. Hammer and T.V. Riley, *Melaleuca alternifolia (tea tree) oil: A review of antimicrobial and on the medicinal properties*. Clin. Microbiol. Rev., **1**, 50–62 (2006).
- (31) I. Cheraif, H. Ben Jannet, M. Hammami, M. Khouja and Z. Mighri, *Chemical composition and antimicrobial activity of essential oils of Cupressus arizonica Greene*. Biochem Systematics Ecol., **35**, 813–820 (2007).
- (32) S. Prabuseenivasan, M. Jayakumar and S. Ignacimuthu, *In vitro antibacterial activity of some plant essential oils*. BMC Complement. Altern. Med., **39**, 1–8 (2006).
- (33) P. Suppakul, J. Miltz, K. Sonneveld and S.W. Bigger, *Antimicrobial properties of basil and its possible application in food packaging*. J. Agric. Food. Chem., **51**, 3197–3207 (2003).
- (34) M.J. Pelczar, E.C.S. Chan and N.R. Krieg, *Control of Microorganism: Chemical Agents in Microbiology: Concepts and Applications*. McGraw-Hill, New York (1993).
- (35) M. Stefan, M. Zamfirache, C. Padurariu, E. Truta and I. Gostin, *The composition and antibacterial activity of essential oils in three Ocimum species growing in Romania*. Cent. Eur. J. Biol., **8**, 600–608 (2013).
- (36) K. Knobloch, A. Pauli, B. Iberl, H. Weigand and N. Weis, *Antibacterial and antifungal properties of essential oil components*. J. Essent. Oil Res., 1989 (1), 118–119 (1989).
- (37) B. Ouattara, R.E. Simard, R.A. Holley, G.J.P. Piette and A. Bégin, *Antibacterial activity of selected fatty acids and essential oils against six meat spoilage organisms*. Int. J. Food Microbiol., **37**, 155–162 (1997).
- (38) C.L. Queiroga, M.C. Teixeira Duarte, B. Baesa Ribeiro and P.M. de Magalhães, *Linalool production from the leaves of Bursera aloexylon and its antimicrobial activity*. Fitoterapia, **78**, 327–328 (2007).