




Table 1

URL	Title	Table Link	Table Pic (top)		Table Header	Table footer																																																																																																																																			
https://europepmc.org/article/MED/35807419	Effects of Frankincense Compounds on Infection, Inflammation, and Oral Health.	https://europepmc.org/articles/PMC9268443/table/molecules-27-04174-t003/	<p>Table 3 Boswellic acid extracts or chemical constituents with antimicrobial activity against oral pathogens in vitro.</p> <table border="1"> <thead> <tr> <th>Microbe</th> <th>Extract or Chemical Constituent</th> <th>MIC (μg/mL)</th> <th>References</th> </tr> </thead> <tbody> <tr> <td>Aggregatibacter actinomycetemcomitans ATCC 33384 (Gram-negative)</td> <td>Boswellia serrata</td> <td>512</td> <td>[113]</td> </tr> <tr> <td>Aggregatibacter actinomycetemcomitans JP2 NOV99 (Gram-negative)</td> <td>HAE of Boswellia serrata</td> <td>78</td> <td>[112]</td> </tr> <tr> <td>Streptococcus mutans PTCC 25175 (Gram-positive)</td> <td>KBA</td> <td>16</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>2</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>32</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Streptococcus mutans PTCC 1688 (Gram-positive)</td> <td>HAE of Boswellia serrata</td> <td>50,000</td> <td>[111]</td> </tr> <tr> <td>Enterococcus faecalis ATCC 29212 (Gram-positive)</td> <td>KBA</td> <td>16</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>4</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>8</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Enterococcus faecium ATCC 8042 (Gram-positive)</td> <td>KBA</td> <td>16</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>4</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>8</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Actinomyces viscosus ATCC 15987 (Gram-positive)</td> <td>KBA</td> <td>8</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>2</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>64</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Streptococcus sanguinis ATCC 10556 (Gram-positive)</td> <td>KBA</td> <td>8</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>2</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>128</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Fusobacterium nucleatum ATCC 10953 (Gram-negative)</td> <td>KBA</td> <td>>128</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>>128</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>>128</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Prevotella intermedia ATCC 25611 (Gram-negative)</td> <td>KBA</td> <td>16</td> <td>[109]</td> </tr> <tr> <td></td> <td>AKBA</td> <td>4</td> <td></td> </tr> <tr> <td></td> <td>BA</td> <td>32</td> <td></td> </tr> <tr> <td></td> <td>ABA</td> <td>>128</td> <td></td> </tr> <tr> <td>Bacillus cereus ATCC 33777</td> <td>KBA</td> <td>0</td> <td>[109]</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>MIC: minimum inhibitory concentration; KBA: 11-keto-β-boswellic acid; AKBA: acetyl-11-keto-β-boswellic acid; BA: β-boswellic acid; ABA: acetyl-β-boswellic acid; HAE: hydro-alcoholic extract.</p>	Microbe	Extract or Chemical Constituent	MIC (μ g/mL)	References	Aggregatibacter actinomycetemcomitans ATCC 33384 (Gram-negative)	Boswellia serrata	512	[113]	Aggregatibacter actinomycetemcomitans JP2 NOV99 (Gram-negative)	HAE of Boswellia serrata	78	[112]	Streptococcus mutans PTCC 25175 (Gram-positive)	KBA	16	[109]		AKBA	2			BA	32			ABA	>128		Streptococcus mutans PTCC 1688 (Gram-positive)	HAE of Boswellia serrata	50,000	[111]	Enterococcus faecalis ATCC 29212 (Gram-positive)	KBA	16	[109]		AKBA	4			BA	8			ABA	>128		Enterococcus faecium ATCC 8042 (Gram-positive)	KBA	16	[109]		AKBA	4			BA	8			ABA	>128		Actinomyces viscosus ATCC 15987 (Gram-positive)	KBA	8	[109]		AKBA	2			BA	64			ABA	>128		Streptococcus sanguinis ATCC 10556 (Gram-positive)	KBA	8	[109]		AKBA	2			BA	128			ABA	>128		Fusobacterium nucleatum ATCC 10953 (Gram-negative)	KBA	>128	[109]		AKBA	>128			BA	>128			ABA	>128		Prevotella intermedia ATCC 25611 (Gram-negative)	KBA	16	[109]		AKBA	4			BA	32			ABA	>128		Bacillus cereus ATCC 33777	KBA	0	[109]	Table 3 Boswellic acid extracts or chemical constituents with antimicrobial activity against oral pathogens in vitro.	MIC: minimum inhibitory concentration; KBA: 11-keto- β -boswellic acid; AKBA: acetyl-11-keto- β -boswellic acid; BA: β -boswellic acid; ABA: acetyl- β -boswellic acid; HAE: hydro-alcoholic extract.
Microbe	Extract or Chemical Constituent	MIC (μ g/mL)	References																																																																																																																																						
Aggregatibacter actinomycetemcomitans ATCC 33384 (Gram-negative)	Boswellia serrata	512	[113]																																																																																																																																						
Aggregatibacter actinomycetemcomitans JP2 NOV99 (Gram-negative)	HAE of Boswellia serrata	78	[112]																																																																																																																																						
Streptococcus mutans PTCC 25175 (Gram-positive)	KBA	16	[109]																																																																																																																																						
	AKBA	2																																																																																																																																							
	BA	32																																																																																																																																							
	ABA	>128																																																																																																																																							
Streptococcus mutans PTCC 1688 (Gram-positive)	HAE of Boswellia serrata	50,000	[111]																																																																																																																																						
Enterococcus faecalis ATCC 29212 (Gram-positive)	KBA	16	[109]																																																																																																																																						
	AKBA	4																																																																																																																																							
	BA	8																																																																																																																																							
	ABA	>128																																																																																																																																							
Enterococcus faecium ATCC 8042 (Gram-positive)	KBA	16	[109]																																																																																																																																						
	AKBA	4																																																																																																																																							
	BA	8																																																																																																																																							
	ABA	>128																																																																																																																																							
Actinomyces viscosus ATCC 15987 (Gram-positive)	KBA	8	[109]																																																																																																																																						
	AKBA	2																																																																																																																																							
	BA	64																																																																																																																																							
	ABA	>128																																																																																																																																							
Streptococcus sanguinis ATCC 10556 (Gram-positive)	KBA	8	[109]																																																																																																																																						
	AKBA	2																																																																																																																																							
	BA	128																																																																																																																																							
	ABA	>128																																																																																																																																							
Fusobacterium nucleatum ATCC 10953 (Gram-negative)	KBA	>128	[109]																																																																																																																																						
	AKBA	>128																																																																																																																																							
	BA	>128																																																																																																																																							
	ABA	>128																																																																																																																																							
Prevotella intermedia ATCC 25611 (Gram-negative)	KBA	16	[109]																																																																																																																																						
	AKBA	4																																																																																																																																							
	BA	32																																																																																																																																							
	ABA	>128																																																																																																																																							
Bacillus cereus ATCC 33777	KBA	0	[109]																																																																																																																																						
https://europepmc.org/article/MED/35011355	Phytochemical Profile, Antimicrobial, Cytotoxic, and Antioxidant Activities of Fresh and Air-Dried Satureja nabateorum Essential Oils.	https://europepmc.org/articles/PMC8746579/table/molecules-27-00125-t002/	<p>Table 2 Antibacterial and antifungal properties of the essential oils of <i>S. nabateorum</i> (MIC values in μg/mL).</p> <table border="1"> <thead> <tr> <th></th> <th><i>S. aureus</i></th> <th><i>E. faecium</i></th> <th><i>E. coli</i></th> <th><i>P. aeruginosa</i></th> <th><i>K. pneumoniae</i></th> <th><i>P. vulgaris</i></th> <th>MRSA</th> <th><i>C. albicans</i></th> </tr> <tr> <th>Source</th> <th>ATCC 25923</th> <th>ATCC 700221</th> <th>ATCC 25922</th> <th>ATCC 27853</th> <th>ATCC 13883</th> <th>ATCC 700221</th> <th>Clinically diagnosed</th> <th>ATCC 90028</th> </tr> </thead> <tbody> <tr> <td>Fluconazole</td> <td>ND</td> <td>ND</td> <td>ND</td> <td>ND</td> <td>ND</td> <td>ND</td> <td>ND</td> <td>1.56</td> </tr> <tr> <td>Ampicillin</td> <td>3.12 ± 0.06</td> <td>1.56 ± 0.01</td> <td>3.12 ± 0.01</td> <td>ND</td> <td>1 ± 0.01</td> <td>18 ± 0.91</td> <td>ND</td> <td>ND</td> </tr> <tr> <td>Ciprofloxacin</td> <td>0.78 ± 0.01</td> <td>0.78 ± 0.01</td> <td>1.56 ± 0.08</td> <td>3.12 ± 0.1</td> <td>0.125 ± 0.02</td> <td>15 ± 0.68</td> <td>12.50 ± 0.58</td> <td>ND</td> </tr> <tr> <td>Fresh <i>S. nabateorum</i> EO</td> <td>0.80 ± 0.01</td> <td>1.25 ± 0.03</td> <td>2.25 ± 0.01</td> <td>6.25 ± 0.61</td> <td>0.135 ± 0.04</td> <td>22 ± 1.02</td> <td>12.50 ± 0.61</td> <td>0.75 ± 0.05</td> </tr> <tr> <td>Air-dried <i>S. nabateorum</i> EO</td> <td>0.85 ± 0.01</td> <td>1.25 ± 0.05</td> <td>2.25 ± 0.02</td> <td>12.50 ± 1.31</td> <td>0.135 ± 0.01</td> <td>22 ± 1.13</td> <td>6.25 ± 0.97</td> <td>0.75 ± 0.05</td> </tr> </tbody> </table> <p>MIC: minimum inhibitory concentration (μg/mL); ND: Not detected.</p>		<i>S. aureus</i>	<i>E. faecium</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>K. pneumoniae</i>	<i>P. vulgaris</i>	MRSA	<i>C. albicans</i>	Source	ATCC 25923	ATCC 700221	ATCC 25922	ATCC 27853	ATCC 13883	ATCC 700221	Clinically diagnosed	ATCC 90028	Fluconazole	ND	ND	ND	ND	ND	ND	ND	1.56	Ampicillin	3.12 ± 0.06	1.56 ± 0.01	3.12 ± 0.01	ND	1 ± 0.01	18 ± 0.91	ND	ND	Ciprofloxacin	0.78 ± 0.01	0.78 ± 0.01	1.56 ± 0.08	3.12 ± 0.1	0.125 ± 0.02	15 ± 0.68	12.50 ± 0.58	ND	Fresh <i>S. nabateorum</i> EO	0.80 ± 0.01	1.25 ± 0.03	2.25 ± 0.01	6.25 ± 0.61	0.135 ± 0.04	22 ± 1.02	12.50 ± 0.61	0.75 ± 0.05	Air-dried <i>S. nabateorum</i> EO	0.85 ± 0.01	1.25 ± 0.05	2.25 ± 0.02	12.50 ± 1.31	0.135 ± 0.01	22 ± 1.13	6.25 ± 0.97	0.75 ± 0.05	Table 2 Antibacterial and antifungal properties of the essential oils of <i>S. nabateorum</i> (MIC values in μ g/mL).	MIC: minimum inhibitory concentration (μ g/mL); ND: Not detected.																																																																					
	<i>S. aureus</i>	<i>E. faecium</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>K. pneumoniae</i>	<i>P. vulgaris</i>	MRSA	<i>C. albicans</i>																																																																																																																																	
Source	ATCC 25923	ATCC 700221	ATCC 25922	ATCC 27853	ATCC 13883	ATCC 700221	Clinically diagnosed	ATCC 90028																																																																																																																																	
Fluconazole	ND	ND	ND	ND	ND	ND	ND	1.56																																																																																																																																	
Ampicillin	3.12 ± 0.06	1.56 ± 0.01	3.12 ± 0.01	ND	1 ± 0.01	18 ± 0.91	ND	ND																																																																																																																																	
Ciprofloxacin	0.78 ± 0.01	0.78 ± 0.01	1.56 ± 0.08	3.12 ± 0.1	0.125 ± 0.02	15 ± 0.68	12.50 ± 0.58	ND																																																																																																																																	
Fresh <i>S. nabateorum</i> EO	0.80 ± 0.01	1.25 ± 0.03	2.25 ± 0.01	6.25 ± 0.61	0.135 ± 0.04	22 ± 1.02	12.50 ± 0.61	0.75 ± 0.05																																																																																																																																	
Air-dried <i>S. nabateorum</i> EO	0.85 ± 0.01	1.25 ± 0.05	2.25 ± 0.02	12.50 ± 1.31	0.135 ± 0.01	22 ± 1.13	6.25 ± 0.97	0.75 ± 0.05																																																																																																																																	
https://europepmc.org/article/MED/35011355	Phytochemical Profile, Antimicrobial, Cytotoxic, and Antioxidant Activities of Fresh and Air-Dried Satureja nabateorum Essential Oils.	https://europepmc.org/articles/PMC8746579/table/molecules-27-00125-t003/	<p>Table 3 Cytotoxic activity IC₅₀ values of the EOs obtained from fresh and air-dried <i>S. nabateorum</i> compared with Doxorubicin.</p> <table border="1"> <thead> <tr> <th colspan="5">IC₅₀ (μg/mL) Values against Different Cancer Cell Lines</th> </tr> <tr> <th>Cells</th> <th>MCF-7</th> <th>COLO-205</th> <th>HeLa</th> <th>HepG2</th> </tr> </thead> <tbody> <tr> <td>Air-dried <i>S. nabateorum</i></td> <td>1090 ± 3.25</td> <td>134.3 ± 1.2</td> <td>89 ± 0.74</td> <td>ND</td> </tr> <tr> <td>Fresh <i>S. nabateorum</i></td> <td>930 ± 1.25</td> <td>256 ± 1.95</td> <td>82 ± 0.98</td> <td>90 ± 0.77</td> </tr> <tr> <td>DOX</td> <td>3 ± 0.89</td> <td>5 ± 0.78</td> <td>9.5 ± 1.0</td> <td>25.43 ± 1.73</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p>	IC ₅₀ (μ g/mL) Values against Different Cancer Cell Lines					Cells	MCF-7	COLO-205	HeLa	HepG2	Air-dried <i>S. nabateorum</i>	1090 ± 3.25	134.3 ± 1.2	89 ± 0.74	ND	Fresh <i>S. nabateorum</i>	930 ± 1.25	256 ± 1.95	82 ± 0.98	90 ± 0.77	DOX	3 ± 0.89	5 ± 0.78	9.5 ± 1.0	25.43 ± 1.73	Table 3 Cytotoxic activity IC ₅₀ values of the EOs obtained from fresh and air-dried <i>S. nabateorum</i> compared with Doxorubicin.																																																																																																												
IC ₅₀ (μ g/mL) Values against Different Cancer Cell Lines																																																																																																																																									
Cells	MCF-7	COLO-205	HeLa	HepG2																																																																																																																																					
Air-dried <i>S. nabateorum</i>	1090 ± 3.25	134.3 ± 1.2	89 ± 0.74	ND																																																																																																																																					
Fresh <i>S. nabateorum</i>	930 ± 1.25	256 ± 1.95	82 ± 0.98	90 ± 0.77																																																																																																																																					
DOX	3 ± 0.89	5 ± 0.78	9.5 ± 1.0	25.43 ± 1.73																																																																																																																																					

URL	Title	Table Link	Table Pic (top)		Table Header	Table footer																																																																																																																							
https://europepmc.org/article/MED/35604913#free-full-text	Using nanoemulsions of the essential oils of a selection of medicinal plants from Jazan, Saudi Arabia, as a green larvicultural against <i>Culex pipiens</i> .	https://europepmc.org/articles/PMC9126372/table/pone.0267150.t002/	<p>Table 2</p> <p>LC50 and LC95 of the used essential oils after 24 hours of application.</p> <table border="1"> <thead> <tr> <th>Treatments</th><th>LC50 (95 CI)</th><th>LC95 (95 CI)</th><th>Slope±SE</th></tr> </thead> <tbody> <tr> <td>Basil ordinary form</td><td>81.07 (64.2–102.6)</td><td>227.9 (163.3–430.03)</td><td>3.6±0.6</td></tr> <tr> <td>Basil nanoemulsion</td><td>65.19 (50.5–82.7)</td><td>193.3(137.2–378.7)</td><td>3.48±0.6</td></tr> <tr> <td>Cuminum ordinary form</td><td>96.29 (77.7–120.7)</td><td>242.36(177.8–434.8)</td><td>4.1±0.7</td></tr> <tr> <td>Cuminum nanoemulsion</td><td>64.5(48.3–83.4)</td><td>222.9(151.5–489.09)</td><td>3.05±0.6</td></tr> <tr> <td>Clove ordinary form</td><td>394(111–2096)</td><td>1658(651–4882348)</td><td>2.6±0.3</td></tr> <tr> <td>Clove nanoemulsion</td><td>379(198–810)</td><td>1815(838–19457)</td><td>2.4±0.3</td></tr> <tr> <td>Henna ordinary form</td><td>306(168–718)</td><td>1703(724–31076)</td><td>2.2±0.3</td></tr> <tr> <td>Henna nanoemulsion</td><td>288(153–712)</td><td>1869(742–47639)</td><td>2.02±0.3</td></tr> <tr> <td>Ginger ordinary form</td><td>494(388–636)</td><td>1682(1171–3117)</td><td>3.9±0.4</td></tr> <tr> <td>Ginger nanoemulsion</td><td>479(376–615)</td><td>1623(1134–2935)</td><td>3.1±0.4</td></tr> </tbody> </table> <p>Open in a separate window</p>	Treatments	LC50 (95 CI)	LC95 (95 CI)	Slope±SE	Basil ordinary form	81.07 (64.2–102.6)	227.9 (163.3–430.03)	3.6±0.6	Basil nanoemulsion	65.19 (50.5–82.7)	193.3(137.2–378.7)	3.48±0.6	Cuminum ordinary form	96.29 (77.7–120.7)	242.36(177.8–434.8)	4.1±0.7	Cuminum nanoemulsion	64.5(48.3–83.4)	222.9(151.5–489.09)	3.05±0.6	Clove ordinary form	394(111–2096)	1658(651–4882348)	2.6±0.3	Clove nanoemulsion	379(198–810)	1815(838–19457)	2.4±0.3	Henna ordinary form	306(168–718)	1703(724–31076)	2.2±0.3	Henna nanoemulsion	288(153–712)	1869(742–47639)	2.02±0.3	Ginger ordinary form	494(388–636)	1682(1171–3117)	3.9±0.4	Ginger nanoemulsion	479(376–615)	1623(1134–2935)	3.1±0.4	Table 2	LC50 and LC95 of the used essential oils after 24 hours of application.																																																																												
Treatments	LC50 (95 CI)	LC95 (95 CI)	Slope±SE																																																																																																																										
Basil ordinary form	81.07 (64.2–102.6)	227.9 (163.3–430.03)	3.6±0.6																																																																																																																										
Basil nanoemulsion	65.19 (50.5–82.7)	193.3(137.2–378.7)	3.48±0.6																																																																																																																										
Cuminum ordinary form	96.29 (77.7–120.7)	242.36(177.8–434.8)	4.1±0.7																																																																																																																										
Cuminum nanoemulsion	64.5(48.3–83.4)	222.9(151.5–489.09)	3.05±0.6																																																																																																																										
Clove ordinary form	394(111–2096)	1658(651–4882348)	2.6±0.3																																																																																																																										
Clove nanoemulsion	379(198–810)	1815(838–19457)	2.4±0.3																																																																																																																										
Henna ordinary form	306(168–718)	1703(724–31076)	2.2±0.3																																																																																																																										
Henna nanoemulsion	288(153–712)	1869(742–47639)	2.02±0.3																																																																																																																										
Ginger ordinary form	494(388–636)	1682(1171–3117)	3.9±0.4																																																																																																																										
Ginger nanoemulsion	479(376–615)	1623(1134–2935)	3.1±0.4																																																																																																																										
https://europepmc.org/article/MED/35755940	Antioxidant, Antidiabetic, and Antibacterial Potentials and Chemical Composition of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> Grown Wild in Morocco.	https://europepmc.org/articles/PMC9217590/table/tab4/	<p>Table 4</p> <p>Minimum inhibitory concentration (MIC), minimum bactericidal concentration (MBC), and MBC/MIC values of the essential oils (EOs) of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> against the ten tested bacterial strains.</p> <table border="1"> <thead> <tr> <th rowspan="2">Bacterial strain</th><th colspan="3"><i>S. officinalis</i></th><th colspan="3"><i>M. suaveolens</i></th><th colspan="3">Chloramphenicol</th></tr> <tr> <th>MIC</th><th>MBC</th><th>MBC/MIC</th><th>MIC</th><th>MBC</th><th>MBC/MIC</th><th>MIC</th><th>MBC</th><th>MBC/MIC</th></tr> </thead> <tbody> <tr> <td><i>Escherichia coli</i></td><td>3.12^{a,b}</td><td>6.25^{a,b}</td><td>2</td><td>1.56^b</td><td>1.56^b</td><td>1</td><td>4.0</td><td>4.0</td><td>1</td></tr> <tr> <td><i>Salmonella typhimurium</i></td><td>6.25^{a,b}</td><td>12.5^{a,b}</td><td>2</td><td>3.12^b</td><td>6.25^b</td><td>2</td><td>4.0</td><td>8.0</td><td>2</td></tr> <tr> <td><i>Klebsiella pneumoniae</i></td><td>3.12^{a,b}</td><td>6.25^{a,b}</td><td>2</td><td>1.56^b</td><td>1.56^b</td><td>1</td><td>4.0</td><td>4.0</td><td>1</td></tr> <tr> <td><i>Proteus mirabilis</i></td><td>3.12^{a,b}</td><td>6.25^{a,b}</td><td>2</td><td>1.56^b</td><td>3.12^b</td><td>2</td><td>4.0</td><td>8.0</td><td>2</td></tr> <tr> <td><i>Pseudomonas aeruginosa</i></td><td>25.0^b</td><td>50.0^{a,b}</td><td>2</td><td>25.0^b</td><td>25.0^b</td><td>1</td><td>64</td><td>64</td><td>1</td></tr> <tr> <td><i>Staphylococcus aureus</i></td><td>0.78^{a,b}</td><td>1.56^{a,b}</td><td>2</td><td>0.39^b</td><td>0.39^b</td><td>1</td><td>2.0</td><td>4.0</td><td>2</td></tr> <tr> <td><i>Staphylococcus epidermidis</i></td><td>1.56^{a,b}</td><td>1.56^{a,b}</td><td>1</td><td>0.78^b</td><td>1.56^b</td><td>2</td><td>4.0</td><td>4.0</td><td>1</td></tr> <tr> <td><i>Enterococcus faecalis</i></td><td>3.12^{a,b}</td><td>3.12^{a,b}</td><td>1</td><td>1.56^b</td><td>1.56^b</td><td>1</td><td>2.0</td><td>4.0</td><td>2</td></tr> <tr> <td><i>Listeria monocytogenes</i></td><td>0.78^{a,b}</td><td>1.56^{a,b}</td><td>2</td><td>0.78^b</td><td>0.78^b</td><td>1</td><td>2.0</td><td>2.0</td><td>1</td></tr> <tr> <td><i>Bacillus cereus</i></td><td>1.56^{a,b}</td><td>1.56^{a,b}</td><td>1</td><td>0.78^b</td><td>1.56^b</td><td>2</td><td>4.0</td><td>4.0</td><td>1</td></tr> </tbody> </table> <p>Values of MIC and MBC for the tested EO are in mg/mL, while for chloramphenicol are in µg/mL. ^{a,b}P < 0.05 vs. <i>M. suaveolens</i> EO and chloramphenicol, respectively.</p>	Bacterial strain	<i>S. officinalis</i>			<i>M. suaveolens</i>			Chloramphenicol			MIC	MBC	MBC/MIC	MIC	MBC	MBC/MIC	MIC	MBC	MBC/MIC	<i>Escherichia coli</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	1.56 ^b	1	4.0	4.0	1	<i>Salmonella typhimurium</i>	6.25 ^{a,b}	12.5 ^{a,b}	2	3.12 ^b	6.25 ^b	2	4.0	8.0	2	<i>Klebsiella pneumoniae</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	1.56 ^b	1	4.0	4.0	1	<i>Proteus mirabilis</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	3.12 ^b	2	4.0	8.0	2	<i>Pseudomonas aeruginosa</i>	25.0 ^b	50.0 ^{a,b}	2	25.0 ^b	25.0 ^b	1	64	64	1	<i>Staphylococcus aureus</i>	0.78 ^{a,b}	1.56 ^{a,b}	2	0.39 ^b	0.39 ^b	1	2.0	4.0	2	<i>Staphylococcus epidermidis</i>	1.56 ^{a,b}	1.56 ^{a,b}	1	0.78 ^b	1.56 ^b	2	4.0	4.0	1	<i>Enterococcus faecalis</i>	3.12 ^{a,b}	3.12 ^{a,b}	1	1.56 ^b	1.56 ^b	1	2.0	4.0	2	<i>Listeria monocytogenes</i>	0.78 ^{a,b}	1.56 ^{a,b}	2	0.78 ^b	0.78 ^b	1	2.0	2.0	1	<i>Bacillus cereus</i>	1.56 ^{a,b}	1.56 ^{a,b}	1	0.78 ^b	1.56 ^b	2	4.0	4.0	1	Table 4	Values of MIC and MBC for the tested EO are in mg/mL, while for chloramphenicol are in µg/mL. ^{a,b} P < 0.05 vs. <i>M. suaveolens</i> EO and chloramphenicol, respectively.	Minimum inhibitory concentration (MIC), minimum bactericidal concentration (MBC), and MBC/MIC values of the essential oils (EOs) of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> against the ten tested bacterial strains.
Bacterial strain	<i>S. officinalis</i>				<i>M. suaveolens</i>			Chloramphenicol																																																																																																																					
	MIC	MBC	MBC/MIC	MIC	MBC	MBC/MIC	MIC	MBC	MBC/MIC																																																																																																																				
<i>Escherichia coli</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	1.56 ^b	1	4.0	4.0	1																																																																																																																				
<i>Salmonella typhimurium</i>	6.25 ^{a,b}	12.5 ^{a,b}	2	3.12 ^b	6.25 ^b	2	4.0	8.0	2																																																																																																																				
<i>Klebsiella pneumoniae</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	1.56 ^b	1	4.0	4.0	1																																																																																																																				
<i>Proteus mirabilis</i>	3.12 ^{a,b}	6.25 ^{a,b}	2	1.56 ^b	3.12 ^b	2	4.0	8.0	2																																																																																																																				
<i>Pseudomonas aeruginosa</i>	25.0 ^b	50.0 ^{a,b}	2	25.0 ^b	25.0 ^b	1	64	64	1																																																																																																																				
<i>Staphylococcus aureus</i>	0.78 ^{a,b}	1.56 ^{a,b}	2	0.39 ^b	0.39 ^b	1	2.0	4.0	2																																																																																																																				
<i>Staphylococcus epidermidis</i>	1.56 ^{a,b}	1.56 ^{a,b}	1	0.78 ^b	1.56 ^b	2	4.0	4.0	1																																																																																																																				
<i>Enterococcus faecalis</i>	3.12 ^{a,b}	3.12 ^{a,b}	1	1.56 ^b	1.56 ^b	1	2.0	4.0	2																																																																																																																				
<i>Listeria monocytogenes</i>	0.78 ^{a,b}	1.56 ^{a,b}	2	0.78 ^b	0.78 ^b	1	2.0	2.0	1																																																																																																																				
<i>Bacillus cereus</i>	1.56 ^{a,b}	1.56 ^{a,b}	1	0.78 ^b	1.56 ^b	2	4.0	4.0	1																																																																																																																				

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																																					
https://europepmc.org/article/MED/35755940	Antioxidant, Antidiabetic, and Antibacterial Potentials and Chemical Composition of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> Grown Wild in Morocco.	https://europepmc.org/articles/PMC9217590/table/tab3/	<p>Table 3</p> <p>In vitro antibacterial activities of the essential oils (EOs) of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> via the disc-diffusion method. Chloramphenicol and DMSO were used as a positive and negative control, respectively.</p> <table border="1"> <thead> <tr> <th></th> <th><i>S. officinalis</i> EO (10 µl/disc)</th> <th><i>M. suaveolens</i> EO (10 µl/disc)</th> <th>Chloramphenicol (30 µg/disc)</th> <th>10% DMSO (10 µl/disc)</th> </tr> </thead> <tbody> <tr> <td>Gram-positive bacteria</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>Staphylococcus aureus</i></td> <td>28.0 ± 0.6</td> <td>31.6 ± 0.6^a</td> <td>30.8 ± 0.6</td> <td>—</td> </tr> <tr> <td><i>Staphylococcus epidermidis</i></td> <td>25.1 ± 0.5</td> <td>27.5 ± 0.3^a</td> <td>28.7 ± 0.3</td> <td>—</td> </tr> <tr> <td><i>Enterococcus faecalis</i></td> <td>20.2 ± 0.4</td> <td>22.7 ± 0.4^a</td> <td>23.4 ± 0.3</td> <td>—</td> </tr> <tr> <td><i>Listeria monocytogenes</i></td> <td>25.0 ± 0.3</td> <td>29.6 ± 0.4^a</td> <td>27.5 ± 0.2</td> <td>—</td> </tr> <tr> <td><i>Bacillus cereus</i></td> <td>26.8 ± 0.4</td> <td>30.4 ± 0.3^a</td> <td>29.0 ± 0.2</td> <td>—</td> </tr> <tr> <td>Gram-negative bacteria</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>Escherichia coli</i></td> <td>18.6 ± 0.3</td> <td>21.7 ± 0.2^a</td> <td>28.1 ± 0.3</td> <td>—</td> </tr> <tr> <td><i>Salmonella typhimurium</i></td> <td>16.3 ± 0.2</td> <td>18.7 ± 0.1^{a,b}</td> <td>21.0 ± 0.3</td> <td>—</td> </tr> <tr> <td><i>Klebsiella pneumoniae</i></td> <td>17.6 ± 0.2</td> <td>19.4 ± 0.2^{a,b}</td> <td>23.1 ± 0.2</td> <td>—</td> </tr> <tr> <td><i>Proteus mirabilis</i></td> <td>17.1 ± 0.3</td> <td>21.3 ± 0.1^{a,b}</td> <td>25.3 ± 0.5</td> <td>—</td> </tr> <tr> <td><i>Pseudomonas aeruginosa</i></td> <td>8.1 ± 0.5^b</td> <td>9.2 ± 0.1^b</td> <td>—</td> <td>—</td> </tr> </tbody> </table> <p>Values are the mean zone of inhibition (mm) ± SD. (—), undetected activity. ^{a,b}<i>P</i> < 0.05 vs. <i>S. officinalis</i> EO and chloramphenicol, respectively.</p>		<i>S. officinalis</i> EO (10 µl/disc)	<i>M. suaveolens</i> EO (10 µl/disc)	Chloramphenicol (30 µg/disc)	10% DMSO (10 µl/disc)	Gram-positive bacteria					<i>Staphylococcus aureus</i>	28.0 ± 0.6	31.6 ± 0.6 ^a	30.8 ± 0.6	—	<i>Staphylococcus epidermidis</i>	25.1 ± 0.5	27.5 ± 0.3 ^a	28.7 ± 0.3	—	<i>Enterococcus faecalis</i>	20.2 ± 0.4	22.7 ± 0.4 ^a	23.4 ± 0.3	—	<i>Listeria monocytogenes</i>	25.0 ± 0.3	29.6 ± 0.4 ^a	27.5 ± 0.2	—	<i>Bacillus cereus</i>	26.8 ± 0.4	30.4 ± 0.3 ^a	29.0 ± 0.2	—	Gram-negative bacteria					<i>Escherichia coli</i>	18.6 ± 0.3	21.7 ± 0.2 ^a	28.1 ± 0.3	—	<i>Salmonella typhimurium</i>	16.3 ± 0.2	18.7 ± 0.1 ^{a,b}	21.0 ± 0.3	—	<i>Klebsiella pneumoniae</i>	17.6 ± 0.2	19.4 ± 0.2 ^{a,b}	23.1 ± 0.2	—	<i>Proteus mirabilis</i>	17.1 ± 0.3	21.3 ± 0.1 ^{a,b}	25.3 ± 0.5	—	<i>Pseudomonas aeruginosa</i>	8.1 ± 0.5 ^b	9.2 ± 0.1 ^b	—	—	Table 3	Values are the mean zone of inhibition (mm) ± SD. (—), undetected activity. ^{a,b} <i>P</i> < 0.05 vs. <i>S. officinalis</i> EO and chloramphenicol, respectively.																				
	<i>S. officinalis</i> EO (10 µl/disc)	<i>M. suaveolens</i> EO (10 µl/disc)	Chloramphenicol (30 µg/disc)	10% DMSO (10 µl/disc)																																																																																						
Gram-positive bacteria																																																																																										
<i>Staphylococcus aureus</i>	28.0 ± 0.6	31.6 ± 0.6 ^a	30.8 ± 0.6	—																																																																																						
<i>Staphylococcus epidermidis</i>	25.1 ± 0.5	27.5 ± 0.3 ^a	28.7 ± 0.3	—																																																																																						
<i>Enterococcus faecalis</i>	20.2 ± 0.4	22.7 ± 0.4 ^a	23.4 ± 0.3	—																																																																																						
<i>Listeria monocytogenes</i>	25.0 ± 0.3	29.6 ± 0.4 ^a	27.5 ± 0.2	—																																																																																						
<i>Bacillus cereus</i>	26.8 ± 0.4	30.4 ± 0.3 ^a	29.0 ± 0.2	—																																																																																						
Gram-negative bacteria																																																																																										
<i>Escherichia coli</i>	18.6 ± 0.3	21.7 ± 0.2 ^a	28.1 ± 0.3	—																																																																																						
<i>Salmonella typhimurium</i>	16.3 ± 0.2	18.7 ± 0.1 ^{a,b}	21.0 ± 0.3	—																																																																																						
<i>Klebsiella pneumoniae</i>	17.6 ± 0.2	19.4 ± 0.2 ^{a,b}	23.1 ± 0.2	—																																																																																						
<i>Proteus mirabilis</i>	17.1 ± 0.3	21.3 ± 0.1 ^{a,b}	25.3 ± 0.5	—																																																																																						
<i>Pseudomonas aeruginosa</i>	8.1 ± 0.5 ^b	9.2 ± 0.1 ^b	—	—																																																																																						
https://europepmc.org/article/MED/35755940	Antioxidant, Antidiabetic, and Antibacterial Potentials and Chemical Composition of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> Grown Wild in Morocco.	https://europepmc.org/articles/PMC9217590/table/tab2/	<p>Table 2</p> <p>In vitro inhibitory effects of <i>Salvia officinalis</i> and <i>Mentha suaveolens</i> essential oils (EOs) on α-amylase and α-glucosidase enzymatic activities.</p> <table border="1"> <thead> <tr> <th>Assay</th> <th>Acarbose</th> <th><i>S. officinalis</i> EO</th> <th><i>M. suaveolens</i> EO</th> </tr> </thead> <tbody> <tr> <td>α-Amylase (IC_{50} in µg/mL)</td> <td>396.42 ± 5.16</td> <td>81.91 ± 0.03^a</td> <td>94.30 ± 0.06^a</td> </tr> <tr> <td>α-Glucosidase (IC_{50} in µg/mL)</td> <td>199.53 ± 1.12</td> <td>113.17 ± 0.02^a</td> <td>141.16 ± 0.2^a</td> </tr> </tbody> </table> <p>Open in a separate window</p> <p>^a <i>P</i> < 0.05 vs. the positive control (acarbose).</p>	Assay	Acarbose	<i>S. officinalis</i> EO	<i>M. suaveolens</i> EO	α -Amylase (IC_{50} in µg/mL)	396.42 ± 5.16	81.91 ± 0.03 ^a	94.30 ± 0.06 ^a	α -Glucosidase (IC_{50} in µg/mL)	199.53 ± 1.12	113.17 ± 0.02 ^a	141.16 ± 0.2 ^a	Table 2	^a <i>P</i> < 0.05 vs. the positive control (acarbose).																																																																									
Assay	Acarbose	<i>S. officinalis</i> EO	<i>M. suaveolens</i> EO																																																																																							
α -Amylase (IC_{50} in µg/mL)	396.42 ± 5.16	81.91 ± 0.03 ^a	94.30 ± 0.06 ^a																																																																																							
α -Glucosidase (IC_{50} in µg/mL)	199.53 ± 1.12	113.17 ± 0.02 ^a	141.16 ± 0.2 ^a																																																																																							
https://europepmc.org/article/MED/35889245	Essential Oil of the Plants Growing in the Brazilian Amazon: Chemical Composition, Antioxidants, and Biological Applications.	https://europepmc.org/articles/PMC9318482/table/molecules-27-0437-3-t002/	<p>Table 2</p> <p>Essential oils of the Amazon and their antioxidant activities.</p> <table border="1"> <thead> <tr> <th>Species (Plants Part)</th> <th>Family</th> <th>Method</th> <th>Results</th> <th>References</th> </tr> </thead> <tbody> <tr> <td><i>Aniba parviflora</i> (Leaves)</td> <td>Lauraceae</td> <td>DPPH</td> <td>TEAC = 90.1–287.9 mg TE/mL</td> <td>[18]</td> </tr> <tr> <td><i>A. parviflora</i> (Branches)</td> <td>Lauraceae</td> <td>DPPH</td> <td>TEAC = 94.1–358.4 mg TE/mL</td> <td>[18]</td> </tr> <tr> <td><i>A. roseodora</i> (Aerial parts)</td> <td>Lauraceae</td> <td>ABTS</td> <td>EC₅₀ = 15.46 µg/mL</td> <td>[19]</td> </tr> <tr> <td><i>Endlicheria arenosa</i> (Leaves)</td> <td>Lauraceae</td> <td>DPPH</td> <td>TEAC = 334.1 ± 41.6 mg TE/mL</td> <td>[28]</td> </tr> <tr> <td><i>E. arenosa</i> (Twigs)</td> <td>Lauraceae</td> <td>DPPH</td> <td>TEAC = 252.6 ± 24.4 mg TE/mL</td> <td>[28]</td> </tr> <tr> <td><i>Eugenia eugenia</i> (Aerial parts)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>TEAC = 216.5 ± 11.6 mg TE/mL</td> <td>[30]</td> </tr> <tr> <td><i>E. flavescens</i> (Aerial parts)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>TEAC = 122.6 ± 6.8 mg TE/mL</td> <td>[30]</td> </tr> <tr> <td><i>E. patrisii</i> (Aerial parts)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>TEAC = 111.2 ± 12.4 mg TE/mL</td> <td>[30]</td> </tr> <tr> <td><i>E. patrisii</i> (Leaves)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>Inhibition = 28.9 ± 4.8%</td> <td>[32]</td> </tr> <tr> <td><i>E. patrisii</i> (Dry leaves)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>Inhibition = 99.0 ± 0.099% (Specimen A) Inhibition = 204.0 ± 0.877% (Specimen B)</td> <td>[31]</td> </tr> <tr> <td></td> <td></td> <td>ABTS</td> <td>Inhibition = 31.4 ± 0.1% (Specimen A) Inhibition = 17.9 ± 0.069% (Specimen B)</td> <td></td> </tr> <tr> <td><i>E. puniceifolia</i> (Dry leaves)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>Inhibition = 408.0 ± 0.10% (Specimen A) Inhibition = 285.0 ± 0.028% (Specimen B)</td> <td>[31]</td> </tr> <tr> <td></td> <td></td> <td>ABTS</td> <td>Inhibition = 9.5 ± 0.034% (Specimen A) Inhibition = 37.7 ± 0.035% (Specimen B)</td> <td></td> </tr> <tr> <td><i>E. uniflora</i> (Leaves)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>Inhibition = 42.6 ± 0.3 to 64.2 ± 0.3%</td> <td>[34]</td> </tr> <tr> <td><i>E. uniflora</i> (Dry leaves)</td> <td>Myrtaceae</td> <td>DPPH</td> <td>Inhibition = 30.3 ± 3.3 to 40.6 ± 1.9%</td> <td>[48]</td> </tr> <tr> <td></td> <td></td> <td>β-Carotene</td> <td>Inhibition = 153.5 ± 16.5 to 228.3 ± 19.2%</td> <td></td> </tr> </tbody> </table> <p>DPPH, 2,2-Diphenyl-1-picrylhydrazyl; ABTS, 2,2-azinobis-(3-ethylbenzothiazoline-6-sulfonate); EC₅₀ (concentration required to obtain 50% antioxidant effect).</p> <p>Open in a separate window</p>	Species (Plants Part)	Family	Method	Results	References	<i>Aniba parviflora</i> (Leaves)	Lauraceae	DPPH	TEAC = 90.1–287.9 mg TE/mL	[18]	<i>A. parviflora</i> (Branches)	Lauraceae	DPPH	TEAC = 94.1–358.4 mg TE/mL	[18]	<i>A. roseodora</i> (Aerial parts)	Lauraceae	ABTS	EC ₅₀ = 15.46 µg/mL	[19]	<i>Endlicheria arenosa</i> (Leaves)	Lauraceae	DPPH	TEAC = 334.1 ± 41.6 mg TE/mL	[28]	<i>E. arenosa</i> (Twigs)	Lauraceae	DPPH	TEAC = 252.6 ± 24.4 mg TE/mL	[28]	<i>Eugenia eugenia</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 216.5 ± 11.6 mg TE/mL	[30]	<i>E. flavescens</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 122.6 ± 6.8 mg TE/mL	[30]	<i>E. patrisii</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 111.2 ± 12.4 mg TE/mL	[30]	<i>E. patrisii</i> (Leaves)	Myrtaceae	DPPH	Inhibition = 28.9 ± 4.8%	[32]	<i>E. patrisii</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 99.0 ± 0.099% (Specimen A) Inhibition = 204.0 ± 0.877% (Specimen B)	[31]			ABTS	Inhibition = 31.4 ± 0.1% (Specimen A) Inhibition = 17.9 ± 0.069% (Specimen B)		<i>E. puniceifolia</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 408.0 ± 0.10% (Specimen A) Inhibition = 285.0 ± 0.028% (Specimen B)	[31]			ABTS	Inhibition = 9.5 ± 0.034% (Specimen A) Inhibition = 37.7 ± 0.035% (Specimen B)		<i>E. uniflora</i> (Leaves)	Myrtaceae	DPPH	Inhibition = 42.6 ± 0.3 to 64.2 ± 0.3%	[34]	<i>E. uniflora</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 30.3 ± 3.3 to 40.6 ± 1.9%	[48]			β-Carotene	Inhibition = 153.5 ± 16.5 to 228.3 ± 19.2%		Table 2	Essential oils of the Amazon and their antioxidant activities.
Species (Plants Part)	Family	Method	Results	References																																																																																						
<i>Aniba parviflora</i> (Leaves)	Lauraceae	DPPH	TEAC = 90.1–287.9 mg TE/mL	[18]																																																																																						
<i>A. parviflora</i> (Branches)	Lauraceae	DPPH	TEAC = 94.1–358.4 mg TE/mL	[18]																																																																																						
<i>A. roseodora</i> (Aerial parts)	Lauraceae	ABTS	EC ₅₀ = 15.46 µg/mL	[19]																																																																																						
<i>Endlicheria arenosa</i> (Leaves)	Lauraceae	DPPH	TEAC = 334.1 ± 41.6 mg TE/mL	[28]																																																																																						
<i>E. arenosa</i> (Twigs)	Lauraceae	DPPH	TEAC = 252.6 ± 24.4 mg TE/mL	[28]																																																																																						
<i>Eugenia eugenia</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 216.5 ± 11.6 mg TE/mL	[30]																																																																																						
<i>E. flavescens</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 122.6 ± 6.8 mg TE/mL	[30]																																																																																						
<i>E. patrisii</i> (Aerial parts)	Myrtaceae	DPPH	TEAC = 111.2 ± 12.4 mg TE/mL	[30]																																																																																						
<i>E. patrisii</i> (Leaves)	Myrtaceae	DPPH	Inhibition = 28.9 ± 4.8%	[32]																																																																																						
<i>E. patrisii</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 99.0 ± 0.099% (Specimen A) Inhibition = 204.0 ± 0.877% (Specimen B)	[31]																																																																																						
		ABTS	Inhibition = 31.4 ± 0.1% (Specimen A) Inhibition = 17.9 ± 0.069% (Specimen B)																																																																																							
<i>E. puniceifolia</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 408.0 ± 0.10% (Specimen A) Inhibition = 285.0 ± 0.028% (Specimen B)	[31]																																																																																						
		ABTS	Inhibition = 9.5 ± 0.034% (Specimen A) Inhibition = 37.7 ± 0.035% (Specimen B)																																																																																							
<i>E. uniflora</i> (Leaves)	Myrtaceae	DPPH	Inhibition = 42.6 ± 0.3 to 64.2 ± 0.3%	[34]																																																																																						
<i>E. uniflora</i> (Dry leaves)	Myrtaceae	DPPH	Inhibition = 30.3 ± 3.3 to 40.6 ± 1.9%	[48]																																																																																						
		β-Carotene	Inhibition = 153.5 ± 16.5 to 228.3 ± 19.2%																																																																																							

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																						
https://europepmc.org/article/MED/35889245	Essential Oil of the Plants Growing in the Brazilian Amazon: Chemical Composition, Antioxidants, and Biological Applications.	https://europepmc.org/articles/PMC9318482/table/molecules-27-0437-3-t003/	<p>Table 3 Antibacterial activity of essential oils from species found in the Amazon.</p> <table border="1"> <thead> <tr> <th>Species</th> <th>Family</th> <th>Methods</th> <th>Microrganisms (Results)</th> <th>References</th> </tr> </thead> <tbody> <tr> <td><i>Anaxagorea brevipes</i> (Leaves)</td> <td>Annonaceae</td> <td>Microbroth dilution</td> <td><i>Kocuria rhizophila</i> (MIC = 50.00 µg/mL) <i>Staphylococcus aureus</i> (MIC = 250.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-negative (8-) (MIC = 25.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-positive (7+) (MIC = 250.00 µg/mL) <i>Enterococcus faecalis</i> (MIC = 250.00 µg/mL)</td> <td>[15]</td> </tr> <tr> <td><i>Aniba parviflora</i> (Aerial parts)</td> <td>Lauraceae</td> <td>Agar disk diffusion/The plate microdilution</td> <td><i>Klebsiella pneumoniae</i> (DDM = 9.20 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 15.44 mm/MIC = >10 µL/mL) <i>Enterococcus faecalis</i> (DDM = 11.2 mm/MIC = >10 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 13.3 mm/MIC = >10 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 1.3 mm/MIC = 1.3 µL/mL) <i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL) <i>Staphylococcus epidermidis</i> (MIC = 250.00 µg/mL)</td> <td>[17]</td> </tr> <tr> <td><i>A. roseodora</i> (Aerial parts)</td> <td>Lauraceae</td> <td>Agar disk diffusion</td> <td><i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL)</td> <td>[17]</td> </tr> <tr> <td><i>Bacogeopsis pleiosperma</i> (Barks)</td> <td>Annonaceae</td> <td>Microbroth dilution</td> <td><i>Staphylococcus aureus</i> (MIC = 0.19 mg/mL)</td> <td>[22]</td> </tr> <tr> <td><i>B. multiflora</i> (Leaves)</td> <td>Annonaceae</td> <td>Microdilution</td> <td><i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.09 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 0.19 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL) <i>Salmonella enterica</i> (MIC = 1.5 mg/mL) <i>Escherichia coli</i> (MIC = 4.68 µg/mL)</td> <td>[23]</td> </tr> <tr> <td><i>B. multiflora</i> (Aerial parts)</td> <td>Annonaceae</td> <td>Microdilution</td> <td><i>Pseudomonas aeruginosa</i> (MIC = 4.68 µg/mL) <i>Enterococcus faecalis</i> (MIC = 4.68 µg/mL) <i>Streptococcus mutan</i> (MIC = 4.68 µg/mL) <i>streptococcus pyogenes</i> (MIC = 4.68 µg/mL) <i>MRSA</i> (MIC = 4.68 µg/mL)</td> <td>[24]</td> </tr> <tr> <td><i>Duguetia quinarensis</i> (Aerial parts)</td> <td>Annonaceae</td> <td>Microdilution</td> <td><i>Streptococcus mutan</i> (MIC = 37.5 µg/mL) <i>Streptococcus pyogenes</i> (MIC = 37.5 µg/mL)</td> <td>[24]</td> </tr> <tr> <td><i>Ephedranthus amazonicus</i> (Leaves)</td> <td>Annonaceae</td> <td>Microdilution</td> <td><i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.19 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 2.50 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL)</td> <td>[25]</td> </tr> <tr> <td><i>Endlicheria arenosa</i> (Leaves)</td> <td>Lauraceae</td> <td>Microbroth dilution</td> <td><i>Pseudomonas aeruginosa</i> (MIC = 125.0 µg/mL) <i>Escherichia coli</i> (MIC = 19.5 µg/mL)</td> <td>[26]</td> </tr> <tr> <td><i>E. arenosa</i> (Twigs)</td> <td>Lauraceae</td> <td>Microbroth dilution</td> <td><i>Staphylococcus aureus</i> (MIC = 625.0 µg/mL)</td> <td>[26]</td> </tr> <tr> <td><i>Fuscoa longifolia</i> (Aerial parts)</td> <td>Annonaceae</td> <td>Microdilution</td> <td><i>Pseudomonas aeruginosa</i> (MIC = 37.5 µg/mL) <i>Streptococcus mutan</i> (MIC = 37.5 µg/mL)</td> <td>[24]</td> </tr> </tbody> </table> <p>MIC, minimum inhibitory concentration; DDM, disk diffusion method.</p>	Species	Family	Methods	Microrganisms (Results)	References	<i>Anaxagorea brevipes</i> (Leaves)	Annonaceae	Microbroth dilution	<i>Kocuria rhizophila</i> (MIC = 50.00 µg/mL) <i>Staphylococcus aureus</i> (MIC = 250.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-negative (8-) (MIC = 25.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-positive (7+) (MIC = 250.00 µg/mL) <i>Enterococcus faecalis</i> (MIC = 250.00 µg/mL)	[15]	<i>Aniba parviflora</i> (Aerial parts)	Lauraceae	Agar disk diffusion/The plate microdilution	<i>Klebsiella pneumoniae</i> (DDM = 9.20 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 15.44 mm/MIC = >10 µL/mL) <i>Enterococcus faecalis</i> (DDM = 11.2 mm/MIC = >10 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 13.3 mm/MIC = >10 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 1.3 mm/MIC = 1.3 µL/mL) <i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL) <i>Staphylococcus epidermidis</i> (MIC = 250.00 µg/mL)	[17]	<i>A. roseodora</i> (Aerial parts)	Lauraceae	Agar disk diffusion	<i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL)	[17]	<i>Bacogeopsis pleiosperma</i> (Barks)	Annonaceae	Microbroth dilution	<i>Staphylococcus aureus</i> (MIC = 0.19 mg/mL)	[22]	<i>B. multiflora</i> (Leaves)	Annonaceae	Microdilution	<i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.09 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 0.19 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL) <i>Salmonella enterica</i> (MIC = 1.5 mg/mL) <i>Escherichia coli</i> (MIC = 4.68 µg/mL)	[23]	<i>B. multiflora</i> (Aerial parts)	Annonaceae	Microdilution	<i>Pseudomonas aeruginosa</i> (MIC = 4.68 µg/mL) <i>Enterococcus faecalis</i> (MIC = 4.68 µg/mL) <i>Streptococcus mutan</i> (MIC = 4.68 µg/mL) <i>streptococcus pyogenes</i> (MIC = 4.68 µg/mL) <i>MRSA</i> (MIC = 4.68 µg/mL)	[24]	<i>Duguetia quinarensis</i> (Aerial parts)	Annonaceae	Microdilution	<i>Streptococcus mutan</i> (MIC = 37.5 µg/mL) <i>Streptococcus pyogenes</i> (MIC = 37.5 µg/mL)	[24]	<i>Ephedranthus amazonicus</i> (Leaves)	Annonaceae	Microdilution	<i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.19 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 2.50 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL)	[25]	<i>Endlicheria arenosa</i> (Leaves)	Lauraceae	Microbroth dilution	<i>Pseudomonas aeruginosa</i> (MIC = 125.0 µg/mL) <i>Escherichia coli</i> (MIC = 19.5 µg/mL)	[26]	<i>E. arenosa</i> (Twigs)	Lauraceae	Microbroth dilution	<i>Staphylococcus aureus</i> (MIC = 625.0 µg/mL)	[26]	<i>Fuscoa longifolia</i> (Aerial parts)	Annonaceae	Microdilution	<i>Pseudomonas aeruginosa</i> (MIC = 37.5 µg/mL) <i>Streptococcus mutan</i> (MIC = 37.5 µg/mL)	[24]	Table 3 Antibacterial activity of essential oils from species found in the Amazon.	MIC, minimum inhibitory concentration; DDM, disk diffusion method.										
Species	Family	Methods	Microrganisms (Results)	References																																																																							
<i>Anaxagorea brevipes</i> (Leaves)	Annonaceae	Microbroth dilution	<i>Kocuria rhizophila</i> (MIC = 50.00 µg/mL) <i>Staphylococcus aureus</i> (MIC = 250.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-negative (8-) (MIC = 25.00 µg/mL) <i>Staphylococcus aureus</i> penicillinase-positive (7+) (MIC = 250.00 µg/mL) <i>Enterococcus faecalis</i> (MIC = 250.00 µg/mL)	[15]																																																																							
<i>Aniba parviflora</i> (Aerial parts)	Lauraceae	Agar disk diffusion/The plate microdilution	<i>Klebsiella pneumoniae</i> (DDM = 9.20 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 15.44 mm/MIC = >10 µL/mL) <i>Enterococcus faecalis</i> (DDM = 11.2 mm/MIC = >10 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 13.3 mm/MIC = >10 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 1.3 mm/MIC = 1.3 µL/mL) <i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL) <i>Staphylococcus epidermidis</i> (MIC = 250.00 µg/mL)	[17]																																																																							
<i>A. roseodora</i> (Aerial parts)	Lauraceae	Agar disk diffusion	<i>Escherichia coli</i> (DDM = 13.2 mm/MIC = >10 µL/mL) <i>Klebsiella pneumoniae</i> (DDM = 11.6 mm/MIC = >10 µL/mL) <i>Staphylococcus aureus</i> (DDM = 26.7 mm/MIC = 1.3 µL/mL) <i>Enterococcus faecalis</i> (DDM = 8.80 mm/MIC = 5 µL/mL) <i>Staphylococcus epidermidis</i> (DDM = 38.4 mm/MIC = 5 µL/mL) <i>Streptococcus pyogenes</i> (DDM = 40.00/MIC = 1.3 µL/mL)	[17]																																																																							
<i>Bacogeopsis pleiosperma</i> (Barks)	Annonaceae	Microbroth dilution	<i>Staphylococcus aureus</i> (MIC = 0.19 mg/mL)	[22]																																																																							
<i>B. multiflora</i> (Leaves)	Annonaceae	Microdilution	<i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.09 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 0.19 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL) <i>Salmonella enterica</i> (MIC = 1.5 mg/mL) <i>Escherichia coli</i> (MIC = 4.68 µg/mL)	[23]																																																																							
<i>B. multiflora</i> (Aerial parts)	Annonaceae	Microdilution	<i>Pseudomonas aeruginosa</i> (MIC = 4.68 µg/mL) <i>Enterococcus faecalis</i> (MIC = 4.68 µg/mL) <i>Streptococcus mutan</i> (MIC = 4.68 µg/mL) <i>streptococcus pyogenes</i> (MIC = 4.68 µg/mL) <i>MRSA</i> (MIC = 4.68 µg/mL)	[24]																																																																							
<i>Duguetia quinarensis</i> (Aerial parts)	Annonaceae	Microdilution	<i>Streptococcus mutan</i> (MIC = 37.5 µg/mL) <i>Streptococcus pyogenes</i> (MIC = 37.5 µg/mL)	[24]																																																																							
<i>Ephedranthus amazonicus</i> (Leaves)	Annonaceae	Microdilution	<i>Staphylococcus aureus</i> (MIC = 0.09 mg/mL) <i>Enterococcus faecalis</i> (MIC = 0.19 mg/mL) <i>Streptococcus sanguinis</i> (MIC = 2.50 mg/mL) <i>Pseudomonas aeruginosa</i> (MIC = 3.0 mg/mL) <i>Escherichia coli</i> (MIC = 1.5 mg/mL)	[25]																																																																							
<i>Endlicheria arenosa</i> (Leaves)	Lauraceae	Microbroth dilution	<i>Pseudomonas aeruginosa</i> (MIC = 125.0 µg/mL) <i>Escherichia coli</i> (MIC = 19.5 µg/mL)	[26]																																																																							
<i>E. arenosa</i> (Twigs)	Lauraceae	Microbroth dilution	<i>Staphylococcus aureus</i> (MIC = 625.0 µg/mL)	[26]																																																																							
<i>Fuscoa longifolia</i> (Aerial parts)	Annonaceae	Microdilution	<i>Pseudomonas aeruginosa</i> (MIC = 37.5 µg/mL) <i>Streptococcus mutan</i> (MIC = 37.5 µg/mL)	[24]																																																																							
https://europepmc.org/article/MED/35889245	Essential Oil of the Plants Growing in the Brazilian Amazon: Chemical Composition, Antioxidants, and Biological Applications.	https://europepmc.org/articles/PMC9318482/table/molecules-27-0437-3-t004/	<p>Table 4 Antifungal activity of essential oils from the Amazon.</p> <table border="1"> <thead> <tr> <th>Species</th> <th>Family</th> <th>Methods</th> <th>Microrganisms (Results)</th> <th>References</th> </tr> </thead> <tbody> <tr> <td><i>Copaifera multijuga</i> (resin)</td> <td>Fabaceae</td> <td>ASD</td> <td><i>Aspergillus flavus</i> (MIC = 0.08 mg/mL—19.5 ± 2.1) <i>Aspergillus niger</i> (MIC = 0.1 mg/mL—9.5 ± 0.7) <i>Aspergillus tamarii</i> (MIC = 0.5 mg/mL—9.0 ± 0.0) <i>Aspergillus tamarii</i> (MIC = 0.3 mg/mL—12.5 ± 3.5) <i>Aspergillus terreus</i> (MIC = 0.3 mg/mL—11.5 ± 2.1) <i>Candida guilliermondii</i> (MIC = 0.1 mg/mL—9.5 ± 1.1) <i>Candida tropicalis</i> (MIC = 0.5 mg/mL—10.0 ± 0.0) <i>Candida parapsilosis</i> (MIC = 0.1 mg/mL—16.0 ± 1.4)</td> <td>[26]</td> </tr> <tr> <td><i>Ocimum compechianum</i> (leaves/stems)</td> <td>Lamiaceae</td> <td>PDA</td> <td>Growth (%) <i>Fusarium oxysporum</i> (IC₅₀ 0.25 µL/mL—23.9 ± 3.8) (IC₅₀ 0.50 µL/mL—47.1 ± 6.2) (IC₅₀ 0.75 µL/mL—59.4 ± 1.2) (IC₅₀ 1.00 µL/mL—60.8 ± 3.7) (IC₅₀ 2.50 µL/mL—70.3 ± 8.7)</td> <td>[48]</td> </tr> <tr> <td><i>O. compechianum</i> (leaves/stems)</td> <td>Lamiaceae</td> <td>PDA</td> <td>Germination (%) <i>Fusarium oxysporum</i> (IC₅₀ 0.50 µL/mL—22.6 ± 1.6) (IC₅₀ 0.75 µL/mL—38.1 ± 11.6) (IC₅₀ 1.00 µL/mL—33.0 ± 1.7) (IC₅₀ 2.50 µL/mL—58.7 ± 0.0)</td> <td>[48]</td> </tr> <tr> <td><i>O. compechianum</i> (leaves/stems)</td> <td>Lamiaceae</td> <td>PDA</td> <td>Growth (%) <i>Colletotrichum gossypii</i> (IC₅₀ 0.25 µL/mL—0.0 ± 0.0) (IC₅₀ 0.50 µL/mL—31.5 ± 1.5) (IC₅₀ 0.75 µL/mL—50.7 ± 8.7) (IC₅₀ 1.00 µL/mL—55.0 ± 3.3) (IC₅₀ 2.50 µL/mL—100.0 ± 0.0)</td> <td>[48]</td> </tr> <tr> <td><i>Ocotea longifolia</i> (leaves)</td> <td>Lauraceae</td> <td>PDA</td> <td><i>Fusarium oxysporum</i> f. sp. <i>dianthi</i>—Inhibition: 31.2 ± 0.45%</td> <td>[83]</td> </tr> <tr> <td><i>O. macrophylla</i> (leaves)</td> <td>Lauraceae</td> <td>PDA</td> <td><i>Botryosphaeriaceae</i>—Inhibition: 32.8 ± 0.21% <i>Fusarium oxysporum</i> f. sp. <i>dianthi</i>—Inhibition: 13.2 ± 0.32%</td> <td>[83]</td> </tr> <tr> <td><i>Piper aduncum</i> (aerial parts)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>Cladosporium cladosporioides</i> (DL = 100 µg) <i>Cladosporium sphaerospermum</i> (DL = 100 µg)</td> <td>[82]</td> </tr> <tr> <td><i>P. alegrenum</i> (aerial parts)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>Cladosporium cladosporioides</i> (DL = <0.1) <i>Cladosporium sphaerospermum</i> (DL = <0.1)</td> <td>[12]</td> </tr> <tr> <td><i>P. divaricatum</i> (aerial parts)</td> <td>Piperaceae</td> <td>MIC</td> <td><i>F. solani</i> f. sp. <i>piperis</i> (MIC = 0.75 mg/mL = 63.36 ± 0.00) (MIC = <1.00 mg/mL = 77.10 ± 10.49) (MIC = <2.50 mg/mL = 92.37 ± 3.50)</td> <td>[81]</td> </tr> <tr> <td><i>P. hispidum</i> (aerial parts)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>C. cladosporioides</i> (MIC = 0.5 µg) <i>C. sphaerospermum</i> (MIC = 5.0 µg)</td> <td>[82]</td> </tr> <tr> <td><i>P. krueppelii</i> (twigs)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>Cladosporium cladosporioides</i> (DL = 0.1) <i>C. cladosporioides</i> (MIC = 0.1 µg/mL)</td> <td>[84]</td> </tr> <tr> <td><i>P. krueppelii</i> (leaves)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>C. cladosporioides</i> (MIC = 0.5 µg/mL) <i>C. sphaerospermum</i> (MIC = 0.5 µg/mL)</td> <td>[84]</td> </tr> <tr> <td><i>P. marginatum</i> (aerial parts)</td> <td>Piperaceae</td> <td>TLC plates</td> <td><i>C. cladosporioides</i> (DL = 10 µg/mL) <i>C. sphaerospermum</i> (DL = 25 µg/mL)</td> <td>[85]</td> </tr> </tbody> </table> <p>MIC, minimum inhibitory concentration; DDM, disk diffusion method.</p>	Species	Family	Methods	Microrganisms (Results)	References	<i>Copaifera multijuga</i> (resin)	Fabaceae	ASD	<i>Aspergillus flavus</i> (MIC = 0.08 mg/mL—19.5 ± 2.1) <i>Aspergillus niger</i> (MIC = 0.1 mg/mL—9.5 ± 0.7) <i>Aspergillus tamarii</i> (MIC = 0.5 mg/mL—9.0 ± 0.0) <i>Aspergillus tamarii</i> (MIC = 0.3 mg/mL—12.5 ± 3.5) <i>Aspergillus terreus</i> (MIC = 0.3 mg/mL—11.5 ± 2.1) <i>Candida guilliermondii</i> (MIC = 0.1 mg/mL—9.5 ± 1.1) <i>Candida tropicalis</i> (MIC = 0.5 mg/mL—10.0 ± 0.0) <i>Candida parapsilosis</i> (MIC = 0.1 mg/mL—16.0 ± 1.4)	[26]	<i>Ocimum compechianum</i> (leaves/stems)	Lamiaceae	PDA	Growth (%) <i>Fusarium oxysporum</i> (IC ₅₀ 0.25 µL/mL—23.9 ± 3.8) (IC ₅₀ 0.50 µL/mL—47.1 ± 6.2) (IC ₅₀ 0.75 µL/mL—59.4 ± 1.2) (IC ₅₀ 1.00 µL/mL—60.8 ± 3.7) (IC ₅₀ 2.50 µL/mL—70.3 ± 8.7)	[48]	<i>O. compechianum</i> (leaves/stems)	Lamiaceae	PDA	Germination (%) <i>Fusarium oxysporum</i> (IC ₅₀ 0.50 µL/mL—22.6 ± 1.6) (IC ₅₀ 0.75 µL/mL—38.1 ± 11.6) (IC ₅₀ 1.00 µL/mL—33.0 ± 1.7) (IC ₅₀ 2.50 µL/mL—58.7 ± 0.0)	[48]	<i>O. compechianum</i> (leaves/stems)	Lamiaceae	PDA	Growth (%) <i>Colletotrichum gossypii</i> (IC ₅₀ 0.25 µL/mL—0.0 ± 0.0) (IC ₅₀ 0.50 µL/mL—31.5 ± 1.5) (IC ₅₀ 0.75 µL/mL—50.7 ± 8.7) (IC ₅₀ 1.00 µL/mL—55.0 ± 3.3) (IC ₅₀ 2.50 µL/mL—100.0 ± 0.0)	[48]	<i>Ocotea longifolia</i> (leaves)	Lauraceae	PDA	<i>Fusarium oxysporum</i> f. sp. <i>dianthi</i> —Inhibition: 31.2 ± 0.45%	[83]	<i>O. macrophylla</i> (leaves)	Lauraceae	PDA	<i>Botryosphaeriaceae</i> —Inhibition: 32.8 ± 0.21% <i>Fusarium oxysporum</i> f. sp. <i>dianthi</i> —Inhibition: 13.2 ± 0.32%	[83]	<i>Piper aduncum</i> (aerial parts)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = 100 µg) <i>Cladosporium sphaerospermum</i> (DL = 100 µg)	[82]	<i>P. alegrenum</i> (aerial parts)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = <0.1) <i>Cladosporium sphaerospermum</i> (DL = <0.1)	[12]	<i>P. divaricatum</i> (aerial parts)	Piperaceae	MIC	<i>F. solani</i> f. sp. <i>piperis</i> (MIC = 0.75 mg/mL = 63.36 ± 0.00) (MIC = <1.00 mg/mL = 77.10 ± 10.49) (MIC = <2.50 mg/mL = 92.37 ± 3.50)	[81]	<i>P. hispidum</i> (aerial parts)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (MIC = 0.5 µg) <i>C. sphaerospermum</i> (MIC = 5.0 µg)	[82]	<i>P. krueppelii</i> (twigs)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = 0.1) <i>C. cladosporioides</i> (MIC = 0.1 µg/mL)	[84]	<i>P. krueppelii</i> (leaves)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (MIC = 0.5 µg/mL) <i>C. sphaerospermum</i> (MIC = 0.5 µg/mL)	[84]	<i>P. marginatum</i> (aerial parts)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (DL = 10 µg/mL) <i>C. sphaerospermum</i> (DL = 25 µg/mL)	[85]	Table 4 Antifungal activity of essential oils from the Amazon.	MIC, minimum inhibitory concentration; DDM, disk diffusion method.
Species	Family	Methods	Microrganisms (Results)	References																																																																							
<i>Copaifera multijuga</i> (resin)	Fabaceae	ASD	<i>Aspergillus flavus</i> (MIC = 0.08 mg/mL—19.5 ± 2.1) <i>Aspergillus niger</i> (MIC = 0.1 mg/mL—9.5 ± 0.7) <i>Aspergillus tamarii</i> (MIC = 0.5 mg/mL—9.0 ± 0.0) <i>Aspergillus tamarii</i> (MIC = 0.3 mg/mL—12.5 ± 3.5) <i>Aspergillus terreus</i> (MIC = 0.3 mg/mL—11.5 ± 2.1) <i>Candida guilliermondii</i> (MIC = 0.1 mg/mL—9.5 ± 1.1) <i>Candida tropicalis</i> (MIC = 0.5 mg/mL—10.0 ± 0.0) <i>Candida parapsilosis</i> (MIC = 0.1 mg/mL—16.0 ± 1.4)	[26]																																																																							
<i>Ocimum compechianum</i> (leaves/stems)	Lamiaceae	PDA	Growth (%) <i>Fusarium oxysporum</i> (IC ₅₀ 0.25 µL/mL—23.9 ± 3.8) (IC ₅₀ 0.50 µL/mL—47.1 ± 6.2) (IC ₅₀ 0.75 µL/mL—59.4 ± 1.2) (IC ₅₀ 1.00 µL/mL—60.8 ± 3.7) (IC ₅₀ 2.50 µL/mL—70.3 ± 8.7)	[48]																																																																							
<i>O. compechianum</i> (leaves/stems)	Lamiaceae	PDA	Germination (%) <i>Fusarium oxysporum</i> (IC ₅₀ 0.50 µL/mL—22.6 ± 1.6) (IC ₅₀ 0.75 µL/mL—38.1 ± 11.6) (IC ₅₀ 1.00 µL/mL—33.0 ± 1.7) (IC ₅₀ 2.50 µL/mL—58.7 ± 0.0)	[48]																																																																							
<i>O. compechianum</i> (leaves/stems)	Lamiaceae	PDA	Growth (%) <i>Colletotrichum gossypii</i> (IC ₅₀ 0.25 µL/mL—0.0 ± 0.0) (IC ₅₀ 0.50 µL/mL—31.5 ± 1.5) (IC ₅₀ 0.75 µL/mL—50.7 ± 8.7) (IC ₅₀ 1.00 µL/mL—55.0 ± 3.3) (IC ₅₀ 2.50 µL/mL—100.0 ± 0.0)	[48]																																																																							
<i>Ocotea longifolia</i> (leaves)	Lauraceae	PDA	<i>Fusarium oxysporum</i> f. sp. <i>dianthi</i> —Inhibition: 31.2 ± 0.45%	[83]																																																																							
<i>O. macrophylla</i> (leaves)	Lauraceae	PDA	<i>Botryosphaeriaceae</i> —Inhibition: 32.8 ± 0.21% <i>Fusarium oxysporum</i> f. sp. <i>dianthi</i> —Inhibition: 13.2 ± 0.32%	[83]																																																																							
<i>Piper aduncum</i> (aerial parts)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = 100 µg) <i>Cladosporium sphaerospermum</i> (DL = 100 µg)	[82]																																																																							
<i>P. alegrenum</i> (aerial parts)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = <0.1) <i>Cladosporium sphaerospermum</i> (DL = <0.1)	[12]																																																																							
<i>P. divaricatum</i> (aerial parts)	Piperaceae	MIC	<i>F. solani</i> f. sp. <i>piperis</i> (MIC = 0.75 mg/mL = 63.36 ± 0.00) (MIC = <1.00 mg/mL = 77.10 ± 10.49) (MIC = <2.50 mg/mL = 92.37 ± 3.50)	[81]																																																																							
<i>P. hispidum</i> (aerial parts)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (MIC = 0.5 µg) <i>C. sphaerospermum</i> (MIC = 5.0 µg)	[82]																																																																							
<i>P. krueppelii</i> (twigs)	Piperaceae	TLC plates	<i>Cladosporium cladosporioides</i> (DL = 0.1) <i>C. cladosporioides</i> (MIC = 0.1 µg/mL)	[84]																																																																							
<i>P. krueppelii</i> (leaves)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (MIC = 0.5 µg/mL) <i>C. sphaerospermum</i> (MIC = 0.5 µg/mL)	[84]																																																																							
<i>P. marginatum</i> (aerial parts)	Piperaceae	TLC plates	<i>C. cladosporioides</i> (DL = 10 µg/mL) <i>C. sphaerospermum</i> (DL = 25 µg/mL)	[85]																																																																							
https://europepmc.org/article/MED/35889245	Essential Oil of the Plants Growing in the Brazilian Amazon: Chemical Composition, Antioxidants, and Biological Applications.	https://europepmc.org/articles/PMC9318482/table/molecules-27-0437-3-t005/	<p>Table 5 Cytotoxic activity of essential oils from species found in the Amazon.</p> <table border="1"> <thead> <tr> <th>Species</th> <th>Botanic Family</th> <th>Methods</th> <th>Results</th> <th>References</th> </tr> </thead> <tbody> <tr> <td><i>Anaxagorea brevipes</i></td> <td>Annonaceae</td> <td>SRB assay</td> <td>MCF-7 = TG1 12.8 µg/mL NCH460 = (TG1 13.0 µg/mL)</td> <td>[15]</td> </tr> <tr> <td><i>Eugenia cuspidiifolia</i></td> <td>Myrtaceae</td> <td>Alamar blue assay</td> <td>PC3 = 9.6 µg/mL (HCT116) = IC₅₀ 18.1 µg mL⁻¹ (SK-MEL-19) = IC₅₀ 26.17 µg mL⁻¹ (ACP02) = IC₅₀ > 50 µg mL⁻¹ (MRC-5) = IC₅₀ 25.51 µg mL⁻¹</td> <td>[29]</td> </tr> <tr> <td><i>E. egensis</i></td> <td>Myrtaceae</td> <td>MTT colorimetric assay</td> <td>HCT-116 = IC₅₀ 25 µg/mL SKMEL19 = IC₅₀ 25 µg/mL AGP-01 = IC₅₀ > 25 µg/mL</td> <td>[30]</td> </tr> <tr> <td><i>E. florescens</i></td> <td>Myrtaceae</td> <td></td> <td>HCT-116 = IC₅₀ 13.9 µg/mL SKMEL19 = ****</td> <td>[30]</td> </tr> <tr> <td><i>E. patrisii</i></td> <td>Myrtaceae</td> <td></td> <td>AGP-01 = ****</td> <td>[32]</td> </tr> <tr> <td><i>E. polystachya</i></td> <td>Myrtaceae</td> <td></td> <td>HCT-116 = IC₅₀ 10.3 µg/mL SKMEL19 = ****</td> <td>[30]</td> </tr> <tr> <td><i>E. stipitata</i></td> <td>Myrtaceae</td> <td></td> <td>MCF7 = IC₅₀ 19.10 µg/mL SKMEL19 = IC₅₀ 17.20 µg/mL AGP-01 = **** HCT116 = ****</td> <td>[32]</td> </tr> <tr> <td><i>E. tapacumensis</i></td> <td>Myrtaceae</td> <td>Alamar blue assay</td> <td>MCF7 = IC₅₀ 24.35 µg mL⁻¹ (HCT116) = IC₅₀ 12.37 µg mL⁻¹ (SK-MEL-19) = IC₅₀ > 50 µg mL⁻¹ (ACP02) = IC₅₀ > 50 µg mL⁻¹ (MRC-5) = IC₅₀ 36.12 µg mL⁻¹</td> <td>[29]</td> </tr> <tr> <td><i>E. uniflora</i></td> <td>Myrtaceae</td> <td>MTT colorimetric assay</td> <td>HCT-116 (IC₅₀ E2: 16.26 µg/mL; IC₅₀ E4: 9.28 µg/mL) AGP-01, (IC₅₀ E2: 12.60 µg/mL; IC₅₀ E4: 8.73 µg/mL) SKMEL-19 (IC₅₀ E2: 12.20 µg/mL; IC₅₀ E4: 15.42 µg/mL) MRC-5 (IC₅₀ E2: 10.27 µg/mL; IC₅₀ E4: 14.95 µg/mL)</td> <td>[90]</td> </tr> <tr> <td><i>Iryanthera polynera</i></td> <td>Myristicaceae</td> <td>SRB assay</td> <td>PC-3 = IC₅₀ 14.69 ± 4.33 µg/mL MCF-7 = IC₅₀ 13.63 ± 3.23 µg/mL</td> <td>[39]</td> </tr> <tr> <td><i>Myrcia splendens</i></td> <td>Myrtaceae</td> <td>MTT colorimetric assay</td> <td>SKMEL-19 = IC₅₀ 8.50 µg/mL AGP01 = IC₅₀ 4.70 µg/mL HCT116 = IC₅₀ 8.80 µg/mL MRC5 = IC₅₀ 6.11 µg/mL</td> <td>[32]</td> </tr> </tbody> </table> <p>MTT(3-(4,5-dimethylthiazolol-2-yl)-2,5-diphenyltetrazolium bromide). *** = statistically similar at 95% confidence level by Tukey's test.</p>	Species	Botanic Family	Methods	Results	References	<i>Anaxagorea brevipes</i>	Annonaceae	SRB assay	MCF-7 = TG1 12.8 µg/mL NCH460 = (TG1 13.0 µg/mL)	[15]	<i>Eugenia cuspidiifolia</i>	Myrtaceae	Alamar blue assay	PC3 = 9.6 µg/mL (HCT116) = IC ₅₀ 18.1 µg mL ⁻¹ (SK-MEL-19) = IC ₅₀ 26.17 µg mL ⁻¹ (ACP02) = IC ₅₀ > 50 µg mL ⁻¹ (MRC-5) = IC ₅₀ 25.51 µg mL ⁻¹	[29]	<i>E. egensis</i>	Myrtaceae	MTT colorimetric assay	HCT-116 = IC ₅₀ 25 µg/mL SKMEL19 = IC ₅₀ 25 µg/mL AGP-01 = IC ₅₀ > 25 µg/mL	[30]	<i>E. florescens</i>	Myrtaceae		HCT-116 = IC ₅₀ 13.9 µg/mL SKMEL19 = ****	[30]	<i>E. patrisii</i>	Myrtaceae		AGP-01 = ****	[32]	<i>E. polystachya</i>	Myrtaceae		HCT-116 = IC ₅₀ 10.3 µg/mL SKMEL19 = ****	[30]	<i>E. stipitata</i>	Myrtaceae		MCF7 = IC ₅₀ 19.10 µg/mL SKMEL19 = IC ₅₀ 17.20 µg/mL AGP-01 = **** HCT116 = ****	[32]	<i>E. tapacumensis</i>	Myrtaceae	Alamar blue assay	MCF7 = IC ₅₀ 24.35 µg mL ⁻¹ (HCT116) = IC ₅₀ 12.37 µg mL ⁻¹ (SK-MEL-19) = IC ₅₀ > 50 µg mL ⁻¹ (ACP02) = IC ₅₀ > 50 µg mL ⁻¹ (MRC-5) = IC ₅₀ 36.12 µg mL ⁻¹	[29]	<i>E. uniflora</i>	Myrtaceae	MTT colorimetric assay	HCT-116 (IC ₅₀ E2: 16.26 µg/mL; IC ₅₀ E4: 9.28 µg/mL) AGP-01, (IC ₅₀ E2: 12.60 µg/mL; IC ₅₀ E4: 8.73 µg/mL) SKMEL-19 (IC ₅₀ E2: 12.20 µg/mL; IC ₅₀ E4: 15.42 µg/mL) MRC-5 (IC ₅₀ E2: 10.27 µg/mL; IC ₅₀ E4: 14.95 µg/mL)	[90]	<i>Iryanthera polynera</i>	Myristicaceae	SRB assay	PC-3 = IC ₅₀ 14.69 ± 4.33 µg/mL MCF-7 = IC ₅₀ 13.63 ± 3.23 µg/mL	[39]	<i>Myrcia splendens</i>	Myrtaceae	MTT colorimetric assay	SKMEL-19 = IC ₅₀ 8.50 µg/mL AGP01 = IC ₅₀ 4.70 µg/mL HCT116 = IC ₅₀ 8.80 µg/mL MRC5 = IC ₅₀ 6.11 µg/mL	[32]	Table 5 Cytotoxic activity of essential oils from species found in the Amazon.	MTT(3-(4,5-dimethylthiazolol-2-yl)-2,5-diphenyltetrazolium bromide). *** = statistically similar at 95% confidence level by Tukey's test.										
Species	Botanic Family	Methods	Results	References																																																																							
<i>Anaxagorea brevipes</i>	Annonaceae	SRB assay	MCF-7 = TG1 12.8 µg/mL NCH460 = (TG1 13.0 µg/mL)	[15]																																																																							
<i>Eugenia cuspidiifolia</i>	Myrtaceae	Alamar blue assay	PC3 = 9.6 µg/mL (HCT116) = IC ₅₀ 18.1 µg mL ⁻¹ (SK-MEL-19) = IC ₅₀ 26.17 µg mL ⁻¹ (ACP02) = IC ₅₀ > 50 µg mL ⁻¹ (MRC-5) = IC ₅₀ 25.51 µg mL ⁻¹	[29]																																																																							
<i>E. egensis</i>	Myrtaceae	MTT colorimetric assay	HCT-116 = IC ₅₀ 25 µg/mL SKMEL19 = IC ₅₀ 25 µg/mL AGP-01 = IC ₅₀ > 25 µg/mL	[30]																																																																							
<i>E. florescens</i>	Myrtaceae		HCT-116 = IC ₅₀ 13.9 µg/mL SKMEL19 = ****	[30]																																																																							
<i>E. patrisii</i>	Myrtaceae		AGP-01 = ****	[32]																																																																							
<i>E. polystachya</i>	Myrtaceae		HCT-116 = IC ₅₀ 10.3 µg/mL SKMEL19 = ****	[30]																																																																							
<i>E. stipitata</i>	Myrtaceae		MCF7 = IC ₅₀ 19.10 µg/mL SKMEL19 = IC ₅₀ 17.20 µg/mL AGP-01 = **** HCT116 = ****	[32]																																																																							
<i>E. tapacumensis</i>	Myrtaceae	Alamar blue assay	MCF7 = IC ₅₀ 24.35 µg mL ⁻¹ (HCT116) = IC ₅₀ 12.37 µg mL ⁻¹ (SK-MEL-19) = IC ₅₀ > 50 µg mL ⁻¹ (ACP02) = IC ₅₀ > 50 µg mL ⁻¹ (MRC-5) = IC ₅₀ 36.12 µg mL ⁻¹	[29]																																																																							
<i>E. uniflora</i>	Myrtaceae	MTT colorimetric assay	HCT-116 (IC ₅₀ E2: 16.26 µg/mL; IC ₅₀ E4: 9.28 µg/mL) AGP-01, (IC ₅₀ E2: 12.60 µg/mL; IC ₅₀ E4: 8.73 µg/mL) SKMEL-19 (IC ₅₀ E2: 12.20 µg/mL; IC ₅₀ E4: 15.42 µg/mL) MRC-5 (IC ₅₀ E2: 10.27 µg/mL; IC ₅₀ E4: 14.95 µg/mL)	[90]																																																																							
<i>Iryanthera polynera</i>	Myristicaceae	SRB assay	PC-3 = IC ₅₀ 14.69 ± 4.33 µg/mL MCF-7 = IC ₅₀ 13.63 ± 3.23 µg/mL	[39]																																																																							
<i>Myrcia splendens</i>	Myrtaceae	MTT colorimetric assay	SKMEL-19 = IC ₅₀ 8.50 µg/mL AGP01 = IC ₅₀ 4.70 µg/mL HCT116 = IC ₅₀ 8.80 µg/mL MRC5 = IC ₅₀ 6.11 µg/mL	[32]																																																																							

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer
https://europepmc.org/article/MED/35747375	Determination of Volatile Compounds of <i>Mentha piperita</i> and <i>Lavandula multifida</i> and Investigation of Their Antibacterial, Antioxidant, and Antidiabetic Properties.	https://europepmc.org/articles/PMC9213120/table/tab2/	Table 2 MIC and MBC values of MPEO and LMEO. 	Table 2 MIC and MBC values of MPEO and LMEO.	
https://europepmc.org/article/MED/35747375	Determination of Volatile Compounds of <i>Mentha piperita</i> and <i>Lavandula multifida</i> and Investigation of Their Antibacterial, Antioxidant, and Antidiabetic Properties.	https://europepmc.org/articles/PMC9213120/table/tab3/	Table 3 Antioxidant activity using DPPH, H ₂ O ₂ , and xanthine oxidase (XO) methods of MPEO and LMEO.  * Different superscript letters in the same column indicate significant difference ($p < 0.05$).	Table 3 Antioxidant activity using DPPH, H ₂ O ₂ , and xanthine oxidase (XO) methods of MPEO and LMEO.	Different superscript letters in the same column indicate significant difference ($p < 0.05$).
https://europepmc.org/article/MED/35747375	Determination of Volatile Compounds of <i>Mentha piperita</i> and <i>Lavandula multifida</i> and Investigation of Their Antibacterial, Antioxidant, and Antidiabetic Properties.	https://europepmc.org/articles/PMC9213120/table/tab4/	Table 4 The IC ₅₀ values of MPEO and LMEO on α -amylase, α -glucosidase, and lipase inhibition.  * Different superscript letters in the same column indicate significant difference ($p < 0.05$).	Table 4 The IC ₅₀ values of MPEO and LMEO on α -amylase, α -glucosidase, and lipase inhibition.	Different superscript letters in the same column indicate significant difference ($p < 0.05$).

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																																																																																			
https://europepmc.org/article/MED/35630653	Investigation and Biological Assessment of <i>Rumex vesicarius</i> L. Extract: Characterization of the Chemical Components and Antioxidant, Antimicrobial, Cytotoxic, and Anti-Dengue Vector Activity	https://europepmc.org/articles/PMC9147116/table/molecules-27-0317-7-t002/	<p>Table 2</p> <p>Radical scavenging activity (%) and IC₅₀ values (mg/L) at various concentrations of the methanol extract of <i>R. vesicarius</i> and the standard ascorbic acid according to DPPH assay.</p> <table border="1"> <thead> <tr> <th>Treatment</th><th>Conc. (mg/L)</th><th>Radical Scavenging Activity (%)</th><th>IC₅₀ (mg/L)</th></tr> </thead> <tbody> <tr> <td rowspan="6"><i>Rumex vesicarius</i> L</td><td>5</td><td>10.64 ± 0.51 F</td><td>28.89</td></tr> <tr><td>10</td><td>33.05 ± 1.41 E</td><td></td></tr> <tr><td>20</td><td>46.04 ± 2.26 D</td><td></td></tr> <tr><td>30</td><td>53.80 ± 2.60 C</td><td></td></tr> <tr><td>40</td><td>59.83 ± 3.01 B</td><td></td></tr> <tr><td>50</td><td>74.28 ± 3.51 A</td><td></td></tr> <tr> <td colspan="2">LSD_{0.05}</td><td>1.81 ***</td><td></td></tr> <tr> <td rowspan="6">Ascorbic acid</td><td>1</td><td>2.52 ± 0.01 F</td><td>12.48</td></tr> <tr><td>2.5</td><td>10.52 ± 0.02 E</td><td></td></tr> <tr><td>5</td><td>36.77 ± 0.17 D</td><td></td></tr> <tr><td>10</td><td>49.62 ± 0.31 C</td><td></td></tr> <tr><td>15</td><td>59.33 ± 1.12 B</td><td></td></tr> <tr><td>20</td><td>69.11 ± 1.43 A</td><td></td></tr> <tr> <td colspan="2">LSD_{0.05}</td><td>1.61 ***</td><td></td></tr> </tbody> </table> <p>Values are the mean ($n = 3$) ± standard deviation. LSD_{0.05} is the least significant difference between two means, as each test was run in duplicate (calculated by factorial ANOVA). Different superscript letters within each treatment (column) express significant variation at a probability level of 0.05 (Duncan's test). ***: significant at $p \leq 0.001$.</p>	Treatment	Conc. (mg/L)	Radical Scavenging Activity (%)	IC ₅₀ (mg/L)	<i>Rumex vesicarius</i> L	5	10.64 ± 0.51 F	28.89	10	33.05 ± 1.41 E		20	46.04 ± 2.26 D		30	53.80 ± 2.60 C		40	59.83 ± 3.01 B		50	74.28 ± 3.51 A		LSD _{0.05}		1.81 ***		Ascorbic acid	1	2.52 ± 0.01 F	12.48	2.5	10.52 ± 0.02 E		5	36.77 ± 0.17 D		10	49.62 ± 0.31 C		15	59.33 ± 1.12 B		20	69.11 ± 1.43 A		LSD _{0.05}		1.61 ***		Table 2	Values are the mean ($n = 3$) ± standard deviation. LSD _{0.05} is the least significant difference between two means, as each test was run in duplicate (calculated by factorial ANOVA). Different superscript letters within each treatment (column) express significant variation at a probability level of 0.05 (Duncan's test). ***: significant at $p \leq 0.001$.																																																																																	
Treatment	Conc. (mg/L)	Radical Scavenging Activity (%)	IC ₅₀ (mg/L)																																																																																																																																					
<i>Rumex vesicarius</i> L	5	10.64 ± 0.51 F	28.89																																																																																																																																					
	10	33.05 ± 1.41 E																																																																																																																																						
	20	46.04 ± 2.26 D																																																																																																																																						
	30	53.80 ± 2.60 C																																																																																																																																						
	40	59.83 ± 3.01 B																																																																																																																																						
	50	74.28 ± 3.51 A																																																																																																																																						
LSD _{0.05}		1.81 ***																																																																																																																																						
Ascorbic acid	1	2.52 ± 0.01 F	12.48																																																																																																																																					
	2.5	10.52 ± 0.02 E																																																																																																																																						
	5	36.77 ± 0.17 D																																																																																																																																						
	10	49.62 ± 0.31 C																																																																																																																																						
	15	59.33 ± 1.12 B																																																																																																																																						
	20	69.11 ± 1.43 A																																																																																																																																						
LSD _{0.05}		1.61 ***																																																																																																																																						
https://europepmc.org/article/MED/35745024	Phytochemical Composition of <i>Commiphora</i> Oleogum Resins and Their Cytotoxicity against Skin Cancer Cells.	https://europepmc.org/articles/PMC9229828/table/molecules-27-0390-3-t005/	<p>Table 5</p> <p>Cytotoxic efficacies of <i>Commiphora</i> extracts and essential oils against the epidermoid carcinoma cell line A431 and the malignant melanoma cells lines RPMI-7951 and SK-MEL-28. Half maximal inhibitory concentrations (IC₅₀) are given in µg/mL, and for pure compounds, additionally in brackets as µM. XTT assay, 72 h. All data were mean ± SEM of three biological samples, each analyzed in triplicates.</p> <table border="1"> <thead> <tr> <th rowspan="2">Samples</th><th colspan="3">IC₅₀, µg/mL (µM)</th></tr> <tr> <th>A431</th><th>RPMI-7951</th><th>SK-MEL-28</th></tr> </thead> <tbody> <tr> <td>Extracts</td><td></td><td></td><td></td></tr> <tr> <td><i>C. myrrha</i></td><td>11.7 ± 0.9</td><td>6.1 ± 0.5</td><td>11.9 ± 0.8</td></tr> <tr> <td><i>C. erythraea</i></td><td>17.3 ± 1.7</td><td>11.5 ± 0.7</td><td>20.4 ± 1.1</td></tr> <tr> <td><i>C. mukul</i></td><td>7.6 ± 1.0</td><td>2.9 ± 0.7</td><td>10.9 ± 1.5</td></tr> <tr> <td><i>C. kataf</i></td><td>8.4 ± 0.7</td><td>3.7 ± 1.5</td><td>11.4 ± 2.7</td></tr> <tr> <td><i>C. holtziana</i></td><td>21.4 ± 1.6</td><td>7.5 ± 1.6</td><td>15.4 ± 3.2</td></tr> <tr> <td><i>C. confusa</i></td><td>11.1 ± 1.2</td><td>5.6 ± 0.3</td><td>13.2 ± 0.6</td></tr> <tr> <td><i>C. kua</i></td><td>14.6 ± 2.4</td><td>10.6 ± 1.3</td><td>14.3 ± 2.1</td></tr> <tr> <td><i>C. from Tarraxo</i></td><td>18.6 ± 1.9</td><td>4.3 ± 0.7</td><td>23.4 ± 1.6</td></tr> <tr> <td><i>C. from Ogaden</i></td><td>13.1 ± 1.5</td><td>6.6 ± 0.8</td><td>14.4 ± 2.1</td></tr> <tr> <td>Essential oils</td><td></td><td></td><td></td></tr> <tr> <td><i>C. myrrha</i></td><td>6.4 ± 0.2</td><td>6.3 ± 0.8</td><td>9.5 ± 1.7</td></tr> <tr> <td><i>C. erythraea</i></td><td>9.7 ± 0.5</td><td>8.7 ± 1.0</td><td>12.5 ± 2.4</td></tr> <tr> <td><i>C. mukul</i></td><td>20.2 ± 0.8</td><td>20.2 ± 4.2</td><td>37.2 ± 4.5</td></tr> <tr> <td><i>C. kataf</i></td><td>10.5 ± 0.5</td><td>9.9 ± 2.0</td><td>19.2 ± 1.0</td></tr> <tr> <td><i>C. holtziana</i></td><td>10.1 ± 0.9</td><td>9.0 ± 0.8</td><td>14.9 ± 1.7</td></tr> <tr> <td><i>C. confusa</i></td><td>15.9 ± 1.5</td><td>17.0 ± 2.3</td><td>22.7 ± 1.9</td></tr> <tr> <td><i>C. kua</i></td><td>14.2 ± 3.1</td><td>9.3 ± 0.7</td><td>22.3 ± 0.6</td></tr> <tr> <td><i>C. from Ogaden</i></td><td>8.7 ± 1.1</td><td>5.2 ± 0.2</td><td>20.3 ± 2.5</td></tr> <tr> <td>Lead compounds</td><td></td><td></td><td></td></tr> <tr> <td>(E)-Guggulsterone</td><td>12.6 ± 1.6 (40.3 ± 5.3)</td><td>3.7 ± 0.6 (11.7 ± 1.9)</td><td>11.1 ± 1.1 (35.5 ± 3.4)</td></tr> <tr> <td>(Z)-Guggulsterone</td><td>18.9 ± 1.2 (60.4 ± 3.7)</td><td>6.4 ± 0.6 (20.5 ± 2.0)</td><td>10.7 ± 0.8 (34.2 ± 2.6)</td></tr> <tr> <td>Curzerenone</td><td>23.9 ± 5.1 (103.7 ± 22.0)</td><td>17.5 ± 5.2 (76.1 ± 22.6)</td><td>23.1 ± 3.0 (100.5 ± 13.0)</td></tr> <tr> <td>Furanoeudesma-1,3-diene</td><td>9.9 ± 0.1 (46.2 ± 0.2)</td><td>7.1 ± 0.8 (33.3 ± 3.6)</td><td>12.0 ± 0.3 (55.9 ± 1.2)</td></tr> <tr> <td>Myrrhone</td><td>35.4 ± 4.3 (155.0 ± 19.0)</td><td>6.8 ± 2.5 (29.9 ± 11.1)</td><td>14.8 ± 11.8 (65.0 ± 51.8)</td></tr> <tr> <td>Botanical drugs</td><td></td><td></td><td></td></tr> <tr> <td>Myrrhinil-Intest®</td><td>317.9 ± 22.1</td><td>204.2 ± 5.8</td><td>300.8 ± 6.2</td></tr> <tr> <td>Guglipid®</td><td>41.3 ± 0.3</td><td>27.3 ± 3.3</td><td>35.7 ± 0.9</td></tr> <tr> <td>Positive controls</td><td></td><td></td><td></td></tr> <tr> <td>Cisplatin</td><td>10.7 ± 7.0 (35.8 ± 23.3)</td><td>10.7 ± 3.8 (35.8 ± 12.7)</td><td>49.6 ± 6.2 (165.5 ± 28.3)</td></tr> <tr> <td>5-Fluorouracil</td><td>1.0 ± 0.3 (7.4 ± 2.1)</td><td>12.9 ± 4.1 (99.1 ± 31.7)</td><td>161.9 ± 81.3 (1245.0 ± 624.9)</td></tr> </tbody> </table> <p>Open in a separate window</p>	Samples	IC ₅₀ , µg/mL (µM)			A431	RPMI-7951	SK-MEL-28	Extracts				<i>C. myrrha</i>	11.7 ± 0.9	6.1 ± 0.5	11.9 ± 0.8	<i>C. erythraea</i>	17.3 ± 1.7	11.5 ± 0.7	20.4 ± 1.1	<i>C. mukul</i>	7.6 ± 1.0	2.9 ± 0.7	10.9 ± 1.5	<i>C. kataf</i>	8.4 ± 0.7	3.7 ± 1.5	11.4 ± 2.7	<i>C. holtziana</i>	21.4 ± 1.6	7.5 ± 1.6	15.4 ± 3.2	<i>C. confusa</i>	11.1 ± 1.2	5.6 ± 0.3	13.2 ± 0.6	<i>C. kua</i>	14.6 ± 2.4	10.6 ± 1.3	14.3 ± 2.1	<i>C. from Tarraxo</i>	18.6 ± 1.9	4.3 ± 0.7	23.4 ± 1.6	<i>C. from Ogaden</i>	13.1 ± 1.5	6.6 ± 0.8	14.4 ± 2.1	Essential oils				<i>C. myrrha</i>	6.4 ± 0.2	6.3 ± 0.8	9.5 ± 1.7	<i>C. erythraea</i>	9.7 ± 0.5	8.7 ± 1.0	12.5 ± 2.4	<i>C. mukul</i>	20.2 ± 0.8	20.2 ± 4.2	37.2 ± 4.5	<i>C. kataf</i>	10.5 ± 0.5	9.9 ± 2.0	19.2 ± 1.0	<i>C. holtziana</i>	10.1 ± 0.9	9.0 ± 0.8	14.9 ± 1.7	<i>C. confusa</i>	15.9 ± 1.5	17.0 ± 2.3	22.7 ± 1.9	<i>C. kua</i>	14.2 ± 3.1	9.3 ± 0.7	22.3 ± 0.6	<i>C. from Ogaden</i>	8.7 ± 1.1	5.2 ± 0.2	20.3 ± 2.5	Lead compounds				(E)-Guggulsterone	12.6 ± 1.6 (40.3 ± 5.3)	3.7 ± 0.6 (11.7 ± 1.9)	11.1 ± 1.1 (35.5 ± 3.4)	(Z)-Guggulsterone	18.9 ± 1.2 (60.4 ± 3.7)	6.4 ± 0.6 (20.5 ± 2.0)	10.7 ± 0.8 (34.2 ± 2.6)	Curzerenone	23.9 ± 5.1 (103.7 ± 22.0)	17.5 ± 5.2 (76.1 ± 22.6)	23.1 ± 3.0 (100.5 ± 13.0)	Furanoeudesma-1,3-diene	9.9 ± 0.1 (46.2 ± 0.2)	7.1 ± 0.8 (33.3 ± 3.6)	12.0 ± 0.3 (55.9 ± 1.2)	Myrrhone	35.4 ± 4.3 (155.0 ± 19.0)	6.8 ± 2.5 (29.9 ± 11.1)	14.8 ± 11.8 (65.0 ± 51.8)	Botanical drugs				Myrrhinil-Intest®	317.9 ± 22.1	204.2 ± 5.8	300.8 ± 6.2	Guglipid®	41.3 ± 0.3	27.3 ± 3.3	35.7 ± 0.9	Positive controls				Cisplatin	10.7 ± 7.0 (35.8 ± 23.3)	10.7 ± 3.8 (35.8 ± 12.7)	49.6 ± 6.2 (165.5 ± 28.3)	5-Fluorouracil	1.0 ± 0.3 (7.4 ± 2.1)	12.9 ± 4.1 (99.1 ± 31.7)	161.9 ± 81.3 (1245.0 ± 624.9)	Table 5	Cytotoxic efficacies of <i>Commiphora</i> extracts and essential oils against the epidermoid carcinoma cell line A431 and the malignant melanoma cells lines RPMI-7951 and SK-MEL-28. Half maximal inhibitory concentrations (IC ₅₀) are given in µg/mL, and for pure compounds, additionally in brackets as µM. XTT assay, 72 h. All data were mean ± SEM of three biological samples, each analyzed in triplicates.
Samples	IC ₅₀ , µg/mL (µM)																																																																																																																																							
	A431	RPMI-7951	SK-MEL-28																																																																																																																																					
Extracts																																																																																																																																								
<i>C. myrrha</i>	11.7 ± 0.9	6.1 ± 0.5	11.9 ± 0.8																																																																																																																																					
<i>C. erythraea</i>	17.3 ± 1.7	11.5 ± 0.7	20.4 ± 1.1																																																																																																																																					
<i>C. mukul</i>	7.6 ± 1.0	2.9 ± 0.7	10.9 ± 1.5																																																																																																																																					
<i>C. kataf</i>	8.4 ± 0.7	3.7 ± 1.5	11.4 ± 2.7																																																																																																																																					
<i>C. holtziana</i>	21.4 ± 1.6	7.5 ± 1.6	15.4 ± 3.2																																																																																																																																					
<i>C. confusa</i>	11.1 ± 1.2	5.6 ± 0.3	13.2 ± 0.6																																																																																																																																					
<i>C. kua</i>	14.6 ± 2.4	10.6 ± 1.3	14.3 ± 2.1																																																																																																																																					
<i>C. from Tarraxo</i>	18.6 ± 1.9	4.3 ± 0.7	23.4 ± 1.6																																																																																																																																					
<i>C. from Ogaden</i>	13.1 ± 1.5	6.6 ± 0.8	14.4 ± 2.1																																																																																																																																					
Essential oils																																																																																																																																								
<i>C. myrrha</i>	6.4 ± 0.2	6.3 ± 0.8	9.5 ± 1.7																																																																																																																																					
<i>C. erythraea</i>	9.7 ± 0.5	8.7 ± 1.0	12.5 ± 2.4																																																																																																																																					
<i>C. mukul</i>	20.2 ± 0.8	20.2 ± 4.2	37.2 ± 4.5																																																																																																																																					
<i>C. kataf</i>	10.5 ± 0.5	9.9 ± 2.0	19.2 ± 1.0																																																																																																																																					
<i>C. holtziana</i>	10.1 ± 0.9	9.0 ± 0.8	14.9 ± 1.7																																																																																																																																					
<i>C. confusa</i>	15.9 ± 1.5	17.0 ± 2.3	22.7 ± 1.9																																																																																																																																					
<i>C. kua</i>	14.2 ± 3.1	9.3 ± 0.7	22.3 ± 0.6																																																																																																																																					
<i>C. from Ogaden</i>	8.7 ± 1.1	5.2 ± 0.2	20.3 ± 2.5																																																																																																																																					
Lead compounds																																																																																																																																								
(E)-Guggulsterone	12.6 ± 1.6 (40.3 ± 5.3)	3.7 ± 0.6 (11.7 ± 1.9)	11.1 ± 1.1 (35.5 ± 3.4)																																																																																																																																					
(Z)-Guggulsterone	18.9 ± 1.2 (60.4 ± 3.7)	6.4 ± 0.6 (20.5 ± 2.0)	10.7 ± 0.8 (34.2 ± 2.6)																																																																																																																																					
Curzerenone	23.9 ± 5.1 (103.7 ± 22.0)	17.5 ± 5.2 (76.1 ± 22.6)	23.1 ± 3.0 (100.5 ± 13.0)																																																																																																																																					
Furanoeudesma-1,3-diene	9.9 ± 0.1 (46.2 ± 0.2)	7.1 ± 0.8 (33.3 ± 3.6)	12.0 ± 0.3 (55.9 ± 1.2)																																																																																																																																					
Myrrhone	35.4 ± 4.3 (155.0 ± 19.0)	6.8 ± 2.5 (29.9 ± 11.1)	14.8 ± 11.8 (65.0 ± 51.8)																																																																																																																																					
Botanical drugs																																																																																																																																								
Myrrhinil-Intest®	317.9 ± 22.1	204.2 ± 5.8	300.8 ± 6.2																																																																																																																																					
Guglipid®	41.3 ± 0.3	27.3 ± 3.3	35.7 ± 0.9																																																																																																																																					
Positive controls																																																																																																																																								
Cisplatin	10.7 ± 7.0 (35.8 ± 23.3)	10.7 ± 3.8 (35.8 ± 12.7)	49.6 ± 6.2 (165.5 ± 28.3)																																																																																																																																					
5-Fluorouracil	1.0 ± 0.3 (7.4 ± 2.1)	12.9 ± 4.1 (99.1 ± 31.7)	161.9 ± 81.3 (1245.0 ± 624.9)																																																																																																																																					
https://europepmc.org/article/MED/35351963	Repellency and insecticidal activity of seven Mugwort (<i>Artemisia argyi</i>) essential oils against the malaria vector <i>Anopheles sinensis</i> .	https://europepmc.org/articles/PMC8964668/table/Tab2/	<p>Table 2</p> <p>The percentages of repellencies of seven <i>Ar. argyi</i> essential oils and DEET against the four-day old <i>An. sinensis</i> adults during different exposed times.</p> <table border="1"> <thead> <tr> <th rowspan="2">Oil treatments (1.5 µL/cm²)^a</th><th colspan="5">Minutes post application*</th></tr> <tr> <th>5 min</th><th>20 min</th><th>35 min</th><th>50 min</th><th>65 min</th></tr> </thead> <tbody> <tr> <td>CQ</td><td>88.32 ± 3.19^b</td><td>79.18 ± 1.63^b</td><td>66.82 ± 3.20^b</td><td>57.49 ± 5.85^b</td><td>34.55 ± 5.02^b</td></tr> <tr> <td>SC</td><td>72.53 ± 3.95^e</td><td>61.41 ± 1.33^c</td><td>42.81 ± 2.37^d</td><td>27.44 ± 1.38^d</td><td>14.78 ± 1.05^c</td></tr> <tr> <td>YN</td><td>68.62 ± 2.93^e</td><td>58.91 ± 1.15^c</td><td>37.56 ± 0.67^d</td><td>22.30 ± 1.58^d</td><td>9.02 ± 2.33^c</td></tr> <tr> <td>HB</td><td>80.09 ± 3.03^c</td><td>74.98 ± 1.74^b</td><td>57.97 ± 5.65^c</td><td>40.04 ± 1.73^c</td><td>28.94 ± 2.68^b</td></tr> <tr> <td>HN</td><td>90.21 ± 3.62^b</td><td>80.51 ± 2.02^b</td><td>69.30 ± 4.40^b</td><td>59.20 ± 2.51^b</td><td>37.29 ± 2.34^b</td></tr> <tr> <td>SD</td><td>91.90 ± 2.42^b</td><td>81.48 ± 2.47^b</td><td>71.67 ± 5.95^b</td><td>61.31 ± 1.49^b</td><td>39.43 ± 2.94^b</td></tr> <tr> <td>GS</td><td>100 ± 0.00^a</td><td>99.57 ± 0.97^a</td><td>94.28 ± 3.21^a</td><td>76.01 ± 1.06^a</td><td>65.68 ± 3.26^a</td></tr> <tr> <td>DEET (10%)</td><td>100 ± 0.00^a</td><td>99.59 ± 0.91^a</td><td>95.04 ± 3.21^a</td><td>83.27 ± 3.69^a</td><td>79.81 ± 4.28^a</td></tr> </tbody> </table> <p>*The repellent rates with different superscript letters in the same column are significantly different at $p < 0.05$. The rates were determined with three replications. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.</p>	Oil treatments (1.5 µL/cm ²) ^a	Minutes post application*					5 min	20 min	35 min	50 min	65 min	CQ	88.32 ± 3.19 ^b	79.18 ± 1.63 ^b	66.82 ± 3.20 ^b	57.49 ± 5.85 ^b	34.55 ± 5.02 ^b	SC	72.53 ± 3.95 ^e	61.41 ± 1.33 ^c	42.81 ± 2.37 ^d	27.44 ± 1.38 ^d	14.78 ± 1.05 ^c	YN	68.62 ± 2.93 ^e	58.91 ± 1.15 ^c	37.56 ± 0.67 ^d	22.30 ± 1.58 ^d	9.02 ± 2.33 ^c	HB	80.09 ± 3.03 ^c	74.98 ± 1.74 ^b	57.97 ± 5.65 ^c	40.04 ± 1.73 ^c	28.94 ± 2.68 ^b	HN	90.21 ± 3.62 ^b	80.51 ± 2.02 ^b	69.30 ± 4.40 ^b	59.20 ± 2.51 ^b	37.29 ± 2.34 ^b	SD	91.90 ± 2.42 ^b	81.48 ± 2.47 ^b	71.67 ± 5.95 ^b	61.31 ± 1.49 ^b	39.43 ± 2.94 ^b	GS	100 ± 0.00 ^a	99.57 ± 0.97 ^a	94.28 ± 3.21 ^a	76.01 ± 1.06 ^a	65.68 ± 3.26 ^a	DEET (10%)	100 ± 0.00 ^a	99.59 ± 0.91 ^a	95.04 ± 3.21 ^a	83.27 ± 3.69 ^a	79.81 ± 4.28 ^a	Table 2	*The repellent rates with different superscript letters in the same column are significantly different at $p < 0.05$. The rates were determined with three replications. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.																																																																								
Oil treatments (1.5 µL/cm ²) ^a	Minutes post application*																																																																																																																																							
	5 min	20 min	35 min	50 min	65 min																																																																																																																																			
CQ	88.32 ± 3.19 ^b	79.18 ± 1.63 ^b	66.82 ± 3.20 ^b	57.49 ± 5.85 ^b	34.55 ± 5.02 ^b																																																																																																																																			
SC	72.53 ± 3.95 ^e	61.41 ± 1.33 ^c	42.81 ± 2.37 ^d	27.44 ± 1.38 ^d	14.78 ± 1.05 ^c																																																																																																																																			
YN	68.62 ± 2.93 ^e	58.91 ± 1.15 ^c	37.56 ± 0.67 ^d	22.30 ± 1.58 ^d	9.02 ± 2.33 ^c																																																																																																																																			
HB	80.09 ± 3.03 ^c	74.98 ± 1.74 ^b	57.97 ± 5.65 ^c	40.04 ± 1.73 ^c	28.94 ± 2.68 ^b																																																																																																																																			
HN	90.21 ± 3.62 ^b	80.51 ± 2.02 ^b	69.30 ± 4.40 ^b	59.20 ± 2.51 ^b	37.29 ± 2.34 ^b																																																																																																																																			
SD	91.90 ± 2.42 ^b	81.48 ± 2.47 ^b	71.67 ± 5.95 ^b	61.31 ± 1.49 ^b	39.43 ± 2.94 ^b																																																																																																																																			
GS	100 ± 0.00 ^a	99.57 ± 0.97 ^a	94.28 ± 3.21 ^a	76.01 ± 1.06 ^a	65.68 ± 3.26 ^a																																																																																																																																			
DEET (10%)	100 ± 0.00 ^a	99.59 ± 0.91 ^a	95.04 ± 3.21 ^a	83.27 ± 3.69 ^a	79.81 ± 4.28 ^a																																																																																																																																			

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																																				
https://europepmc.org/article/MED/35351963	Repellency and insecticidal activity of seven Mugwort (<i>Artemisia argyi</i>) essential oils against the malaria vector <i>Anopheles sinensis</i> .	https://europepmc.org/articles/PMC8964668/table/Tab3/	<p>Table 3</p> <p>Median lethal knockdown activity (mins) of the essential oils extracted from seven geographical regions of <i>Ar. argyi</i> against female adults of <i>An. sinensis</i>.</p> <table border="1"> <thead> <tr> <th>Oil</th> <th>KT₅₀ (min)^a</th> <th>95% Confidence limits</th> <th>Toxicity regression equation (y = a + bx)</th> <th>χ²</th> </tr> </thead> <tbody> <tr> <td>CQ</td> <td>13.85 ± 3.5</td> <td>13.08–14.62</td> <td>y = -4.33 + 0.31x</td> <td>0.29*</td> </tr> <tr> <td>SC</td> <td>10.18 ± 3.4</td> <td>9.43–10.91</td> <td>y = -3.71 + 0.36x</td> <td>0.83*</td> </tr> <tr> <td>YN</td> <td>50.62 ± 8.6</td> <td>48.19–53.52</td> <td>y = -2.69 + 0.05x</td> <td>12.95*</td> </tr> <tr> <td>HB</td> <td>16.77 ± 4.7</td> <td>15.39–18.08</td> <td>y = -1.87 + 0.11x</td> <td>5.55*</td> </tr> <tr> <td>HN</td> <td>10.02 ± 2.8</td> <td>9.02–10.96</td> <td>y = -2.21 + 0.22x</td> <td>3.02*</td> </tr> <tr> <td>SD</td> <td>9.32 ± 2.2</td> <td>8.61–10.05</td> <td>y = -3.36 + 0.36x</td> <td>2.84*</td> </tr> <tr> <td>GS</td> <td>4.71 ± 1.2</td> <td>3.50–5.32</td> <td>y = -2.45 + 0.51x</td> <td>0.33*</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>^aKT₅₀ represents the knockdown time of half individuals, which was determined by log-probit analysis. Three experimental repeats with the essential oil dosage of 2 μL. *Significant at p < 0.05. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.</p>	Oil	KT ₅₀ (min) ^a	95% Confidence limits	Toxicity regression equation (y = a + bx)	χ ²	CQ	13.85 ± 3.5	13.08–14.62	y = -4.33 + 0.31x	0.29*	SC	10.18 ± 3.4	9.43–10.91	y = -3.71 + 0.36x	0.83*	YN	50.62 ± 8.6	48.19–53.52	y = -2.69 + 0.05x	12.95*	HB	16.77 ± 4.7	15.39–18.08	y = -1.87 + 0.11x	5.55*	HN	10.02 ± 2.8	9.02–10.96	y = -2.21 + 0.22x	3.02*	SD	9.32 ± 2.2	8.61–10.05	y = -3.36 + 0.36x	2.84*	GS	4.71 ± 1.2	3.50–5.32	y = -2.45 + 0.51x	0.33*	Table 3	<p>Median lethal knockdown activity (mins) of the essential oils extracted from seven geographical regions of <i>Ar. argyi</i> against female adults of <i>An. sinensis</i>.</p> <p>^aKT₅₀ represents the knockdown time of half individuals, which was determined by log-probit analysis. Three experimental repeats with the essential oil dosage of 2 μL. *Significant at p < 0.05. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.</p>																																												
Oil	KT ₅₀ (min) ^a	95% Confidence limits	Toxicity regression equation (y = a + bx)	χ ²																																																																																					
CQ	13.85 ± 3.5	13.08–14.62	y = -4.33 + 0.31x	0.29*																																																																																					
SC	10.18 ± 3.4	9.43–10.91	y = -3.71 + 0.36x	0.83*																																																																																					
YN	50.62 ± 8.6	48.19–53.52	y = -2.69 + 0.05x	12.95*																																																																																					
HB	16.77 ± 4.7	15.39–18.08	y = -1.87 + 0.11x	5.55*																																																																																					
HN	10.02 ± 2.8	9.02–10.96	y = -2.21 + 0.22x	3.02*																																																																																					
SD	9.32 ± 2.2	8.61–10.05	y = -3.36 + 0.36x	2.84*																																																																																					
GS	4.71 ± 1.2	3.50–5.32	y = -2.45 + 0.51x	0.33*																																																																																					
https://europepmc.org/article/MED/35351963	Repellency and insecticidal activity of seven Mugwort (<i>Artemisia argyi</i>) essential oils against the malaria vector <i>Anopheles sinensis</i> .	https://europepmc.org/articles/PMC8964668/table/Tab4/	<p>Table 4</p> <p>Fumigation toxicity of the essential oils extracted from seven geographical regions of <i>Ar. argyi</i> in China against female adults of <i>An. sinensis</i>.</p> <table border="1"> <thead> <tr> <th>Oil</th> <th>LD₅₀ (μL/L)^a</th> <th>95% confidence limit</th> <th>Toxicity regression equation (y = a + bx)</th> <th>χ²</th> </tr> </thead> <tbody> <tr> <td>CQ</td> <td>14.95 ± 7.8</td> <td>13.32–16.43</td> <td>y = -1.65 + 0.11x</td> <td>2.15*</td> </tr> <tr> <td>SC</td> <td>22.90 ± 5.5</td> <td>20.82–25.08</td> <td>y = -1.56 + 0.06x</td> <td>2.91*</td> </tr> <tr> <td>YN</td> <td>51.26 ± 6.1</td> <td>42.87–71.69</td> <td>y = -2.36 + 0.04x</td> <td>2.77*</td> </tr> <tr> <td>HB</td> <td>48.74 ± 2.1</td> <td>39.09–76.29</td> <td>y = -1.39 + 0.02x</td> <td>4.15*</td> </tr> <tr> <td>HN</td> <td>26.49 ± 4.5</td> <td>24.25–29.13</td> <td>y = -1.70 + 0.06x</td> <td>3.00*</td> </tr> <tr> <td>SD</td> <td>18.66 ± 1.2</td> <td>16.75–20.44</td> <td>y = -1.54 + 0.08x</td> <td>3.55*</td> </tr> <tr> <td>GS</td> <td>9.40 ± 3.6</td> <td>5.15–12.08</td> <td>y = -1.51 + 0.16x</td> <td>8.62*</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>^aDetermined with at least 6 concentrations at μL/L. *Significant at p < 0.05. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.</p>	Oil	LD ₅₀ (μL/L) ^a	95% confidence limit	Toxicity regression equation (y = a + bx)	χ ²	CQ	14.95 ± 7.8	13.32–16.43	y = -1.65 + 0.11x	2.15*	SC	22.90 ± 5.5	20.82–25.08	y = -1.56 + 0.06x	2.91*	YN	51.26 ± 6.1	42.87–71.69	y = -2.36 + 0.04x	2.77*	HB	48.74 ± 2.1	39.09–76.29	y = -1.39 + 0.02x	4.15*	HN	26.49 ± 4.5	24.25–29.13	y = -1.70 + 0.06x	3.00*	SD	18.66 ± 1.2	16.75–20.44	y = -1.54 + 0.08x	3.55*	GS	9.40 ± 3.6	5.15–12.08	y = -1.51 + 0.16x	8.62*	Table 4	<p>Fumigation toxicity of the essential oils extracted from seven geographical regions of <i>Ar. argyi</i> in China against female adults of <i>An. sinensis</i>.</p> <p>^aDetermined with at least 6 concentrations at μL/L. *Significant at p < 0.05. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively.</p>																																												
Oil	LD ₅₀ (μL/L) ^a	95% confidence limit	Toxicity regression equation (y = a + bx)	χ ²																																																																																					
CQ	14.95 ± 7.8	13.32–16.43	y = -1.65 + 0.11x	2.15*																																																																																					
SC	22.90 ± 5.5	20.82–25.08	y = -1.56 + 0.06x	2.91*																																																																																					
YN	51.26 ± 6.1	42.87–71.69	y = -2.36 + 0.04x	2.77*																																																																																					
HB	48.74 ± 2.1	39.09–76.29	y = -1.39 + 0.02x	4.15*																																																																																					
HN	26.49 ± 4.5	24.25–29.13	y = -1.70 + 0.06x	3.00*																																																																																					
SD	18.66 ± 1.2	16.75–20.44	y = -1.54 + 0.08x	3.55*																																																																																					
GS	9.40 ± 3.6	5.15–12.08	y = -1.51 + 0.16x	8.62*																																																																																					
https://europepmc.org/article/MED/35448753#free-full-text	Essential Oils and Extracts of <i>Juniperus macrocarpa</i> Sm. and <i>Juniperus oxycedrus</i> L.: Comparative Phytochemical Composition and Anti-Proliferative and Antioxidant Activities	https://europepmc.org/articles/PMC9031627/table/plants-11-01025-t003/	<p>Table 3</p> <p>Antioxidant activity of essential oils and extracts of <i>J. macrocarpa</i> and <i>J. oxycedrus</i>.</p> <table border="1"> <thead> <tr> <th>Sample</th> <th>ABTS IC₅₀ (μg/mL)</th> <th>DPPH IC₅₀ (μg/mL)</th> <th>FRAP Test μM Fe(II)/g^c</th> <th>β-Carotene Bleaching IC₅₀ (μg/mL)</th> <th></th> </tr> <tr> <th></th> <th></th> <th></th> <th></th> <th></th> <th>30 min 60 min</th> </tr> </thead> <tbody> <tr> <td colspan="6" style="text-align: center;"><i>J. macrocarpa</i></td> </tr> <tr> <td>Essential oil</td> <td>20.4% ^a</td> <td>34.1% ^b</td> <td>2.4 ± 0.2</td> <td>54.8 ± 3.4</td> <td>49.4 ± 2.8</td> </tr> <tr> <td>Ethyl acetate extract</td> <td>147.6 ± 4.8</td> <td>40.9 ± 2.4</td> <td>26.4 ± 1.8</td> <td>84.9 ± 3.5</td> <td>95.7 ± 3.9</td> </tr> <tr> <td>Methanol extract</td> <td>39.1 ± 1.7</td> <td>29.3 ± 1.5</td> <td>23.6 ± 1.5</td> <td>65.1 ± 2.2</td> <td>62.5 ± 2.8</td> </tr> <tr> <td colspan="6" style="text-align: center;"><i>J. oxycedrus</i></td> </tr> <tr> <td>Essential oil</td> <td>5.2% ^a</td> <td>31.6% ^b</td> <td>3.8 ± 0.3</td> <td>47.5 ± 2.6</td> <td>5.9 ± 3.4</td> </tr> <tr> <td>Ethyl acetate extract</td> <td>9.3 ± 1.3</td> <td>20.6 ± 2.3</td> <td>99.5 ± 3.7</td> <td>15.1 ± 1.1</td> <td>13.2 ± 0.8</td> </tr> <tr> <td>Methanol extract</td> <td>6.2 ± 1.1</td> <td>19.7 ± 2.5</td> <td>101.9 ± 3.9</td> <td>23.1 ± 1.2</td> <td>17.1 ± 0.9</td> </tr> <tr> <td colspan="6" style="text-align: center;">Positive control</td> </tr> <tr> <td>Ascorbic acid</td> <td>1.7 ± 0.4</td> <td>5.1 ± 0.8</td> <td></td> <td></td> <td></td> </tr> <tr> <td>BHT</td> <td></td> <td></td> <td>63.2 ± 4.4</td> <td></td> <td></td> </tr> <tr> <td>Propyl gallate</td> <td></td> <td></td> <td></td> <td>1.1 ± 0.05</td> <td>1.0 ± 0.06</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>Data are expressed as mean ± SD (n = 3). ^a at the maximum concentration tested (500 μg/mL). ^b at the maximum concentration tested (1000 μg/mL). ^c at 2.5 mg/mL.</p>	Sample	ABTS IC ₅₀ (μg/mL)	DPPH IC ₅₀ (μg/mL)	FRAP Test μM Fe(II)/g ^c	β-Carotene Bleaching IC ₅₀ (μg/mL)							30 min 60 min	<i>J. macrocarpa</i>						Essential oil	20.4% ^a	34.1% ^b	2.4 ± 0.2	54.8 ± 3.4	49.4 ± 2.8	Ethyl acetate extract	147.6 ± 4.8	40.9 ± 2.4	26.4 ± 1.8	84.9 ± 3.5	95.7 ± 3.9	Methanol extract	39.1 ± 1.7	29.3 ± 1.5	23.6 ± 1.5	65.1 ± 2.2	62.5 ± 2.8	<i>J. oxycedrus</i>						Essential oil	5.2% ^a	31.6% ^b	3.8 ± 0.3	47.5 ± 2.6	5.9 ± 3.4	Ethyl acetate extract	9.3 ± 1.3	20.6 ± 2.3	99.5 ± 3.7	15.1 ± 1.1	13.2 ± 0.8	Methanol extract	6.2 ± 1.1	19.7 ± 2.5	101.9 ± 3.9	23.1 ± 1.2	17.1 ± 0.9	Positive control						Ascorbic acid	1.7 ± 0.4	5.1 ± 0.8				BHT			63.2 ± 4.4			Propyl gallate				1.1 ± 0.05	1.0 ± 0.06	Table 3	<p>Antioxidant activity of essential oils and extracts of <i>J. macrocarpa</i> and <i>J. oxycedrus</i>.</p> <p>Data are expressed as mean ± SD (n = 3). ^a at the maximum concentration tested (500 μg/mL). ^b at the maximum concentration tested (1000 μg/mL). ^c at 2.5 mg/mL.</p>
Sample	ABTS IC ₅₀ (μg/mL)	DPPH IC ₅₀ (μg/mL)	FRAP Test μM Fe(II)/g ^c	β-Carotene Bleaching IC ₅₀ (μg/mL)																																																																																					
					30 min 60 min																																																																																				
<i>J. macrocarpa</i>																																																																																									
Essential oil	20.4% ^a	34.1% ^b	2.4 ± 0.2	54.8 ± 3.4	49.4 ± 2.8																																																																																				
Ethyl acetate extract	147.6 ± 4.8	40.9 ± 2.4	26.4 ± 1.8	84.9 ± 3.5	95.7 ± 3.9																																																																																				
Methanol extract	39.1 ± 1.7	29.3 ± 1.5	23.6 ± 1.5	65.1 ± 2.2	62.5 ± 2.8																																																																																				
<i>J. oxycedrus</i>																																																																																									
Essential oil	5.2% ^a	31.6% ^b	3.8 ± 0.3	47.5 ± 2.6	5.9 ± 3.4																																																																																				
Ethyl acetate extract	9.3 ± 1.3	20.6 ± 2.3	99.5 ± 3.7	15.1 ± 1.1	13.2 ± 0.8																																																																																				
Methanol extract	6.2 ± 1.1	19.7 ± 2.5	101.9 ± 3.9	23.1 ± 1.2	17.1 ± 0.9																																																																																				
Positive control																																																																																									
Ascorbic acid	1.7 ± 0.4	5.1 ± 0.8																																																																																							
BHT			63.2 ± 4.4																																																																																						
Propyl gallate				1.1 ± 0.05	1.0 ± 0.06																																																																																				

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																									
https://europepmc.org/article/MED/35448753#free-full-text	Essential Oils and Extracts of Juniperus macrocarpa Sm. and Juniperus oxycedrus L.: Comparative Phytochemical Composition and Anti-Proliferative and Antioxidant Activities	https://europepmc.org/articles/PMC9031627/table/plants-11-01025-t004/	<p>Table 4</p> <p>Anti-proliferative activity (IC_{50} µg/mL) of essential oils and extracts of <i>J. macrocarpa</i> and <i>J. oxycedrus</i> against four cancer cell lines (MCF-7, MDA-MB-231, A549, and COR-L23).</p> <table border="1"> <thead> <tr> <th>Sample</th><th>MCF-7</th><th>MDA-MB-231</th><th>A549</th><th>COR-L23</th></tr> </thead> <tbody> <tr> <td align="center" colspan="5"><i>J. macrocarpa</i></td></tr> <tr> <td>Essential oil</td><td>85.4 ± 3.2 **</td><td>96.4 ± 3.8 **</td><td>>200</td><td>101.0 ± 3.9 **</td></tr> <tr> <td>Ethyl acetate extract</td><td>163.4 ± 4.9 **</td><td>186.2 ± 5.1 **</td><td>>200</td><td>>200</td></tr> <tr> <td>Methanol extract</td><td>>200</td><td>>200</td><td>>200</td><td>>200</td></tr> <tr> <td align="center" colspan="5"><i>J. oxycedrus</i></td></tr> <tr> <td>Essential oil</td><td>>200</td><td>>200</td><td>>200</td><td>>200</td></tr> <tr> <td>Ethyl acetate extract</td><td>147.9 ± 4.6 **</td><td>158.1 ± 5.1 **</td><td>>200</td><td>39.1 ± 1.4 **</td></tr> <tr> <td>Methanol extract</td><td>>200</td><td>>200</td><td>87.9 ± 4.7 **</td><td>26.0 ± 1.3 **</td></tr> <tr> <td align="center" colspan="5"><i>Positive control</i></td></tr> <tr> <td>Taxol</td><td>0.08 ± 0.004</td><td>1.6 ± 0.03</td><td></td><td></td></tr> <tr> <td>Vinblastine sulfate</td><td></td><td></td><td>67.3 ± 2.0</td><td>45.5 ± 0.7</td></tr> </tbody> </table> <p>Open in a separate window</p> <p>Data are expressed as median ± S.D. (n = 3). A549: human lung adenocarcinoma cell line; MCF-7: human breast cancer ER+ cell line; MDA-MB-231: triple negative breast adenocarcinoma cell line; COR-L23: human lung large cell carcinoma cell line. ** p < 0.01 vs. positive control.</p>	Sample	MCF-7	MDA-MB-231	A549	COR-L23	<i>J. macrocarpa</i>					Essential oil	85.4 ± 3.2 **	96.4 ± 3.8 **	>200	101.0 ± 3.9 **	Ethyl acetate extract	163.4 ± 4.9 **	186.2 ± 5.1 **	>200	>200	Methanol extract	>200	>200	>200	>200	<i>J. oxycedrus</i>					Essential oil	>200	>200	>200	>200	Ethyl acetate extract	147.9 ± 4.6 **	158.1 ± 5.1 **	>200	39.1 ± 1.4 **	Methanol extract	>200	>200	87.9 ± 4.7 **	26.0 ± 1.3 **	<i>Positive control</i>					Taxol	0.08 ± 0.004	1.6 ± 0.03			Vinblastine sulfate			67.3 ± 2.0	45.5 ± 0.7	Table 4	Data are expressed as median ± S.D. (n = 3). A549: human lung adenocarcinoma cell line; MCF-7: human breast cancer ER+ cell line; MDA-MB-231: triple negative breast adenocarcinoma cell line; COR-L23: human lung large cell carcinoma cell line. ** p < 0.01 vs. positive control.													
Sample	MCF-7	MDA-MB-231	A549	COR-L23																																																																										
<i>J. macrocarpa</i>																																																																														
Essential oil	85.4 ± 3.2 **	96.4 ± 3.8 **	>200	101.0 ± 3.9 **																																																																										
Ethyl acetate extract	163.4 ± 4.9 **	186.2 ± 5.1 **	>200	>200																																																																										
Methanol extract	>200	>200	>200	>200																																																																										
<i>J. oxycedrus</i>																																																																														
Essential oil	>200	>200	>200	>200																																																																										
Ethyl acetate extract	147.9 ± 4.6 **	158.1 ± 5.1 **	>200	39.1 ± 1.4 **																																																																										
Methanol extract	>200	>200	87.9 ± 4.7 **	26.0 ± 1.3 **																																																																										
<i>Positive control</i>																																																																														
Taxol	0.08 ± 0.004	1.6 ± 0.03																																																																												
Vinblastine sulfate			67.3 ± 2.0	45.5 ± 0.7																																																																										
https://europepmc.org/article/MED/35721323#free-full-text	Chemical Composition, Antioxidant, and Mosquito Larvicidal Activity of Essential Oils from <i>Hyptis capitata</i> Jacq.	https://europepmc.org/articles/PMC9205432/table/t002/	<p>Table 2</p> <p>Larvicidal Activity of the Essential Oils from Leaf and the Inflorescence of <i>H. capitata</i> Against <i>A. stephensi</i></p> <table border="1"> <thead> <tr> <th>Source of Essential Oil</th><th>Concentration (µg/mL)</th><th>% of Mortality at 24 h±SD</th><th>LC₅₀ µg/mL (LCL-UCL)</th><th>LC₉₀ µg/mL (LCL-UCL)</th><th>X²</th></tr> </thead> <tbody> <tr> <td rowspan="6">Inflorescence</td><td>Control</td><td>0±0</td><td>33.19 (28.74–37.00)</td><td>81.69 (76.70–87.89)</td><td>49.66</td></tr> <tr><td>20</td><td>42±2.73</td><td></td><td></td><td></td></tr> <tr><td>40</td><td>54±4.18</td><td></td><td></td><td></td></tr> <tr><td>60</td><td>71±4.18</td><td></td><td></td><td></td></tr> <tr><td>80</td><td>87±2.73</td><td></td><td></td><td></td></tr> <tr><td>100</td><td>100±0</td><td></td><td></td><td></td></tr> <tr> <td rowspan="10">Leaves</td><td>Control</td><td>0±0</td><td>39.08 (35.26–42.51)</td><td>87.06 (82.56–92.41)</td><td>59.85</td></tr> <tr><td>20</td><td>37±2.73</td><td></td><td></td><td></td></tr> <tr><td>40</td><td>47±2.73</td><td></td><td></td><td></td></tr> <tr><td>60</td><td>67±4.47</td><td></td><td></td><td></td></tr> <tr><td>80</td><td>83±2.73</td><td></td><td></td><td></td></tr> <tr><td>100</td><td>97±4.47</td><td></td><td></td><td></td></tr> <tr><td>120</td><td>100±0</td><td></td><td></td><td></td></tr> </tbody> </table> <p>Open in a separate window</p> <p>Abbreviations: SD, standard deviations; LCL & UCL, lower confidence limits and upper confidence limits; X², chi-square test.</p>	Source of Essential Oil	Concentration (µg/mL)	% of Mortality at 24 h±SD	LC ₅₀ µg/mL (LCL-UCL)	LC ₉₀ µg/mL (LCL-UCL)	X ²	Inflorescence	Control	0±0	33.19 (28.74–37.00)	81.69 (76.70–87.89)	49.66	20	42±2.73				40	54±4.18				60	71±4.18				80	87±2.73				100	100±0				Leaves	Control	0±0	39.08 (35.26–42.51)	87.06 (82.56–92.41)	59.85	20	37±2.73				40	47±2.73				60	67±4.47				80	83±2.73				100	97±4.47				120	100±0				Table 2	Abbreviations: SD, standard deviations; LCL & UCL, lower confidence limits and upper confidence limits; X ² , chi-square test.
Source of Essential Oil	Concentration (µg/mL)	% of Mortality at 24 h±SD	LC ₅₀ µg/mL (LCL-UCL)	LC ₉₀ µg/mL (LCL-UCL)	X ²																																																																									
Inflorescence	Control	0±0	33.19 (28.74–37.00)	81.69 (76.70–87.89)	49.66																																																																									
	20	42±2.73																																																																												
	40	54±4.18																																																																												
	60	71±4.18																																																																												
	80	87±2.73																																																																												
	100	100±0																																																																												
Leaves	Control	0±0	39.08 (35.26–42.51)	87.06 (82.56–92.41)	59.85																																																																									
	20	37±2.73																																																																												
	40	47±2.73																																																																												
	60	67±4.47																																																																												
	80	83±2.73																																																																												
	100	97±4.47																																																																												
	120	100±0																																																																												
	https://europepmc.org/article/MED/35807578#free-full-text	Chemical Diversity and Anti-Insect Activity Evaluation of Essential Oils Extracted from Five <i>Artemisia</i> Species.	https://europepmc.org/articles/PMC9269011/table/plants-11-01627-t003/	<p>Table 3</p> <p>Contact toxicity of the essential oils from five <i>Artemisia</i> species and major compounds against <i>L. bostrychophila</i> adults.</p> <table border="1"> <thead> <tr> <th>Samples</th><th>LD₅₀ (µg/cm²)</th><th>FL (µg/cm²)</th><th>Slope ± SE</th><th>p-Value</th><th>X²</th></tr> </thead> <tbody> <tr> <td><i>A. dalai-lamae</i></td><td>115.94</td><td>104.58–129.58</td><td>6.42 ± 0.95</td><td>0.193</td><td>13.58</td></tr> <tr> <td><i>A. tangutica</i></td><td>70.48</td><td>68.20–73.89</td><td>12.46 ± 1.45</td><td>0.374</td><td>19.30</td></tr> <tr> <td><i>A. sieversiana</i></td><td>195.51</td><td>177.33–215.97</td><td>6.87 ± 0.94</td><td>0.851</td><td>7.89</td></tr> <tr> <td><i>A. tanacetifolia</i></td><td colspan="5">less than 50% mortality at concentration of 50%</td></tr> <tr> <td><i>A. ordosica</i></td><td>52.11</td><td>51.55–53.87</td><td>4.88 ± 0.50</td><td>0.664</td><td>19.62</td></tr> <tr> <td>1,8-Cineole ¹</td><td>1048.74</td><td>1021.95–1096.85</td><td>9.50 ± 0.91</td><td>-</td><td>11.76</td></tr> <tr> <td>Camphor ¹</td><td>207.26</td><td>199.78–214.99</td><td>13.81 ± 1.47</td><td>-</td><td>15.87</td></tr> <tr> <td>3-Carene ²</td><td>223.62</td><td>205.65–243.00</td><td>5.92 ± 0.62</td><td>-</td><td>8.52</td></tr> <tr> <td>Caryophyllene ²</td><td>52.52</td><td>43.52–60.83</td><td>2.77 ± 0.39</td><td>-</td><td>9.62</td></tr> <tr> <td>Pyrethrins ³</td><td>18.72</td><td>17.60–19.92</td><td>2.98 ± 0.40</td><td>0.99</td><td>10.56</td></tr> </tbody> </table> <p>Open in a separate window</p> <p>¹ Data from Liu et al. [21]; ² Data from Cao et al. [22]; ³ Data from Liu et al. [23].</p>	Samples	LD ₅₀ (µg/cm ²)	FL (µg/cm ²)	Slope ± SE	p-Value	X ²	<i>A. dalai-lamae</i>	115.94	104.58–129.58	6.42 ± 0.95	0.193	13.58	<i>A. tangutica</i>	70.48	68.20–73.89	12.46 ± 1.45	0.374	19.30	<i>A. sieversiana</i>	195.51	177.33–215.97	6.87 ± 0.94	0.851	7.89	<i>A. tanacetifolia</i>	less than 50% mortality at concentration of 50%					<i>A. ordosica</i>	52.11	51.55–53.87	4.88 ± 0.50	0.664	19.62	1,8-Cineole ¹	1048.74	1021.95–1096.85	9.50 ± 0.91	-	11.76	Camphor ¹	207.26	199.78–214.99	13.81 ± 1.47	-	15.87	3-Carene ²	223.62	205.65–243.00	5.92 ± 0.62	-	8.52	Caryophyllene ²	52.52	43.52–60.83	2.77 ± 0.39	-	9.62	Pyrethrins ³	18.72	17.60–19.92	2.98 ± 0.40	0.99	10.56	Table 3	1 Data from Liu et al. [21]; 2 Data from Cao et al. [22]; 3 Data from Liu et al. [23].						
	Samples	LD ₅₀ (µg/cm ²)	FL (µg/cm ²)	Slope ± SE	p-Value	X ²																																																																								
	<i>A. dalai-lamae</i>	115.94	104.58–129.58	6.42 ± 0.95	0.193	13.58																																																																								
<i>A. tangutica</i>	70.48	68.20–73.89	12.46 ± 1.45	0.374	19.30																																																																									
<i>A. sieversiana</i>	195.51	177.33–215.97	6.87 ± 0.94	0.851	7.89																																																																									
<i>A. tanacetifolia</i>	less than 50% mortality at concentration of 50%																																																																													
<i>A. ordosica</i>	52.11	51.55–53.87	4.88 ± 0.50	0.664	19.62																																																																									
1,8-Cineole ¹	1048.74	1021.95–1096.85	9.50 ± 0.91	-	11.76																																																																									
Camphor ¹	207.26	199.78–214.99	13.81 ± 1.47	-	15.87																																																																									
3-Carene ²	223.62	205.65–243.00	5.92 ± 0.62	-	8.52																																																																									
Caryophyllene ²	52.52	43.52–60.83	2.77 ± 0.39	-	9.62																																																																									
Pyrethrins ³	18.72	17.60–19.92	2.98 ± 0.40	0.99	10.56																																																																									

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																					
https://europepmc.org/article/MED/35807578#free-full-text	Chemical Diversity and Anti-Insect Activity Evaluation of Essential Oils Extracted from Five <i>Artemisia</i> Species.	https://europepmc.org/articles/PMC9269011/table/plants-11-01627-t004/	<p>Table 4</p> <p>Contact toxicities of the essential oils from five <i>Artemisia</i> species and major compounds against <i>T. castaneum</i> adults.</p> <table border="1"> <thead> <tr> <th>Samples</th> <th>LD₅₀ (µg/adult)</th> <th>FL (µg/cm²)</th> <th>Slope ± SE</th> <th>p-Value</th> <th>χ²</th> </tr> </thead> <tbody> <tr> <td><i>A. dalai-lamae</i></td> <td>25.70</td> <td>23.09–28.49</td> <td>4.19 ± 0.45</td> <td>0.681</td> <td>19.34</td> </tr> <tr> <td><i>A. tangutica</i></td> <td>17.42</td> <td>15.17–19.98</td> <td>3.38 ± 0.45</td> <td>0.553</td> <td>16.57</td> </tr> <tr> <td><i>A. sieversiana</i></td> <td></td> <td>less than 50% mortality at concentration of 50%</td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>A. tanacetifolia</i></td> <td>41.90</td> <td>37.08–47.57</td> <td>3.88 ± 0.45</td> <td>0.995</td> <td>6.17</td> </tr> <tr> <td><i>A. ordosica</i></td> <td>21.68</td> <td>19.86–23.59</td> <td>5.41 ± 5.06</td> <td>0.944</td> <td>13.34</td> </tr> <tr> <td>1,8-Cineole ¹</td> <td>18.83</td> <td>17.13–20.69</td> <td>4.86 ± 0.50</td> <td>-</td> <td>16.56</td> </tr> <tr> <td>Camphor ²</td> <td></td> <td>less than 50% mortality at concentration of 50%</td> <td></td> <td></td> <td></td> </tr> <tr> <td>3-Carene ³</td> <td>63.43</td> <td>57.16–70.75</td> <td>4.11 ± 0.45</td> <td>-</td> <td>11.67</td> </tr> <tr> <td>Caryophyllene ³</td> <td>25.86</td> <td>22.61–30.24</td> <td>2.97 ± 0.39</td> <td>-</td> <td>13.13</td> </tr> <tr> <td>Pyrethrins ⁴</td> <td>0.26</td> <td>0.22–0.30</td> <td>3.34 ± 0.32</td> <td>0.95</td> <td>13.11</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>¹ Data from Wang et al. [24]; ² Data from Guo et al. [25]; ³ Data from Cao et al. [22]; ⁴ Data from Guo et al. [26].</p>	Samples	LD ₅₀ (µg/adult)	FL (µg/cm ²)	Slope ± SE	p-Value	χ ²	<i>A. dalai-lamae</i>	25.70	23.09–28.49	4.19 ± 0.45	0.681	19.34	<i>A. tangutica</i>	17.42	15.17–19.98	3.38 ± 0.45	0.553	16.57	<i>A. sieversiana</i>		less than 50% mortality at concentration of 50%				<i>A. tanacetifolia</i>	41.90	37.08–47.57	3.88 ± 0.45	0.995	6.17	<i>A. ordosica</i>	21.68	19.86–23.59	5.41 ± 5.06	0.944	13.34	1,8-Cineole ¹	18.83	17.13–20.69	4.86 ± 0.50	-	16.56	Camphor ²		less than 50% mortality at concentration of 50%				3-Carene ³	63.43	57.16–70.75	4.11 ± 0.45	-	11.67	Caryophyllene ³	25.86	22.61–30.24	2.97 ± 0.39	-	13.13	Pyrethrins ⁴	0.26	0.22–0.30	3.34 ± 0.32	0.95	13.11	Table 4	Contact toxicities of the essential oils from five <i>Artemisia</i> species and major compounds against <i>T. castaneum</i> adults.			1 Data from Wang et al. [24]; 2 Data from Guo et al. [25]; 3 Data from Cao et al. [22]; 4 Data from Guo et al. [26].
Samples	LD ₅₀ (µg/adult)	FL (µg/cm ²)	Slope ± SE	p-Value	χ ²																																																																					
<i>A. dalai-lamae</i>	25.70	23.09–28.49	4.19 ± 0.45	0.681	19.34																																																																					
<i>A. tangutica</i>	17.42	15.17–19.98	3.38 ± 0.45	0.553	16.57																																																																					
<i>A. sieversiana</i>		less than 50% mortality at concentration of 50%																																																																								
<i>A. tanacetifolia</i>	41.90	37.08–47.57	3.88 ± 0.45	0.995	6.17																																																																					
<i>A. ordosica</i>	21.68	19.86–23.59	5.41 ± 5.06	0.944	13.34																																																																					
1,8-Cineole ¹	18.83	17.13–20.69	4.86 ± 0.50	-	16.56																																																																					
Camphor ²		less than 50% mortality at concentration of 50%																																																																								
3-Carene ³	63.43	57.16–70.75	4.11 ± 0.45	-	11.67																																																																					
Caryophyllene ³	25.86	22.61–30.24	2.97 ± 0.39	-	13.13																																																																					
Pyrethrins ⁴	0.26	0.22–0.30	3.34 ± 0.32	0.95	13.11																																																																					
https://europepmc.org/article/MED/35684405#free-full-text	Chemical Constituents, Antioxidant, Anti-Tyrosinase, Cytotoxicity, and Anti-Melanogenesis Activities of <i>Etlingera elatior</i> (Jack) Leaf Essential Oils.	https://europepmc.org/articles/PMC9182332/table/molecules-27-03469-t002/	<p>Table 2</p> <p>Antioxidant activities of <i>E. elatior</i> leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD).</p> <table border="1"> <thead> <tr> <th></th> <th>HD</th> <th>MAHD</th> </tr> </thead> <tbody> <tr> <td>DPPH (IC₅₀, mg/mL)</td> <td>16.25 ± 0.09 ^a</td> <td>39.29 ± 0.27 ^b</td> </tr> <tr> <td>FRAP (mg TEAC/g)</td> <td>0.91 ± 0.01 ^a</td> <td>0.43 ± 0.00 ^b</td> </tr> <tr> <td>Tyrosinase inhibition (IC₅₀, mg/mL)</td> <td>2.34 ± 0.04 ^a</td> <td>2.97 ± 0.03 ^b</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>The superscribed small letter displayed the significant differences (<i>p</i> < 0.05) between the extraction methods.</p>		HD	MAHD	DPPH (IC ₅₀ , mg/mL)	16.25 ± 0.09 ^a	39.29 ± 0.27 ^b	FRAP (mg TEAC/g)	0.91 ± 0.01 ^a	0.43 ± 0.00 ^b	Tyrosinase inhibition (IC ₅₀ , mg/mL)	2.34 ± 0.04 ^a	2.97 ± 0.03 ^b	Table 2	Antioxidant activities of <i>E. elatior</i> leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD).	The superscribed small letter displayed the significant differences (<i>p</i> < 0.05) between the extraction methods.																																																								
	HD	MAHD																																																																								
DPPH (IC ₅₀ , mg/mL)	16.25 ± 0.09 ^a	39.29 ± 0.27 ^b																																																																								
FRAP (mg TEAC/g)	0.91 ± 0.01 ^a	0.43 ± 0.00 ^b																																																																								
Tyrosinase inhibition (IC ₅₀ , mg/mL)	2.34 ± 0.04 ^a	2.97 ± 0.03 ^b																																																																								
https://europepmc.org/article/MED/35889501	Preliminary Study on Phytochemical Constituents and Biological Activities of Essential Oil from <i>Myriactis nepalensis</i> Less.	https://europepmc.org/articles/PMC9324352/table/molecules-27-04631-t002/	<p>Table 2</p> <p>Diameter of the inhibition zones (DIZ), minimum inhibitory concentrations (MIC) and minimum bactericidal concentration (MBC) of the essential oil of <i>M. nepalensis</i> (MNEO).</p> <table border="1"> <thead> <tr> <th rowspan="2">Strain</th> <th colspan="2">DIZ (mm) ± SD</th> <th colspan="2">MIC (mg/mL)</th> <th colspan="2">MBC (mg/mL)</th> </tr> <tr> <th>MNEO</th> <th>Chl</th> <th>MNEO</th> <th>Chl</th> <th>MNEO</th> <th>Chl</th> </tr> </thead> <tbody> <tr> <td>Gram positive</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>B. subtilis</i> ATCC 6633</td> <td>14.0 ± 1.2</td> <td>25.1 ± 1.0</td> <td>0.039</td> <td>0.002</td> <td>0.156</td> <td>0.004</td> </tr> <tr> <td><i>S. aureus</i> ATCC 6538</td> <td>16.7 ± 1.7</td> <td>23.6 ± 1.3</td> <td>0.078</td> <td>0.002</td> <td>0.078</td> <td>0.016</td> </tr> <tr> <td><i>P. larvae</i> ATCC 9545</td> <td>13.4 ± 1.1</td> <td>28.3 ± 1.7</td> <td>0.039</td> <td>0.001</td> <td>0.039</td> <td>0.002</td> </tr> <tr> <td>Gram negative</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>E. coli</i> ATCC 25922</td> <td>8.3 ± 0.9</td> <td>25.9 ± 0.8</td> <td>1.250</td> <td>0.002</td> <td>>2.500</td> <td>0.004</td> </tr> <tr> <td><i>P. aeruginosa</i> ATCC 27853</td> <td>10.5 ± 1.3</td> <td>16.7 ± 0.4</td> <td>0.625</td> <td>0.031</td> <td>2.500</td> <td>0.250</td> </tr> </tbody> </table> <p>DIZ, diameter of the inhibition zones (mm) is given as the mean ± SD of triplicate experiments; positive control: Chl, chloramphenicol.</p>	Strain	DIZ (mm) ± SD		MIC (mg/mL)		MBC (mg/mL)		MNEO	Chl	MNEO	Chl	MNEO	Chl	Gram positive							<i>B. subtilis</i> ATCC 6633	14.0 ± 1.2	25.1 ± 1.0	0.039	0.002	0.156	0.004	<i>S. aureus</i> ATCC 6538	16.7 ± 1.7	23.6 ± 1.3	0.078	0.002	0.078	0.016	<i>P. larvae</i> ATCC 9545	13.4 ± 1.1	28.3 ± 1.7	0.039	0.001	0.039	0.002	Gram negative							<i>E. coli</i> ATCC 25922	8.3 ± 0.9	25.9 ± 0.8	1.250	0.002	>2.500	0.004	<i>P. aeruginosa</i> ATCC 27853	10.5 ± 1.3	16.7 ± 0.4	0.625	0.031	2.500	0.250	Table 2	Diameter of the inhibition zones (DIZ), minimum inhibitory concentrations (MIC) and minimum bactericidal concentration (MBC) of the essential oil of <i>M. nepalensis</i> (MNEO).	DIZ, diameter of the inhibition zones (mm) is given as the mean ± SD of triplicate experiments; positive control: Chl, chloramphenicol.						
Strain	DIZ (mm) ± SD		MIC (mg/mL)		MBC (mg/mL)																																																																					
	MNEO	Chl	MNEO	Chl	MNEO	Chl																																																																				
Gram positive																																																																										
<i>B. subtilis</i> ATCC 6633	14.0 ± 1.2	25.1 ± 1.0	0.039	0.002	0.156	0.004																																																																				
<i>S. aureus</i> ATCC 6538	16.7 ± 1.7	23.6 ± 1.3	0.078	0.002	0.078	0.016																																																																				
<i>P. larvae</i> ATCC 9545	13.4 ± 1.1	28.3 ± 1.7	0.039	0.001	0.039	0.002																																																																				
Gram negative																																																																										
<i>E. coli</i> ATCC 25922	8.3 ± 0.9	25.9 ± 0.8	1.250	0.002	>2.500	0.004																																																																				
<i>P. aeruginosa</i> ATCC 27853	10.5 ± 1.3	16.7 ± 0.4	0.625	0.031	2.500	0.250																																																																				

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																									
https://europepmc.org/article/MED/35889501	Preliminary Study on Phytochemical Constituents and Biological Activities of Essential Oil from <i>Myriactis nepalensis</i> Less.	https://europepmc.org/articles/PMC9324352/table/molecules-27-0463-1-t003/	<p>Table 3</p> <p>Fractional inhibitory concentration index (FICI) values of the essential oil of <i>M. nepalensis</i> (MNEO) and chloramphenicol combinations.</p> <table border="1"> <thead> <tr> <th>Microorganism</th> <th></th> <th>MICa ($\mu\text{g/mL}$)</th> <th>MICc ($\mu\text{g/mL}$)</th> <th>FICI</th> </tr> </thead> <tbody> <tr> <td rowspan="2"><i>Bacillus subtilis</i></td> <td>MNEO</td> <td>39.10</td> <td>9.76</td> <td>0.37 (S)</td> </tr> <tr> <td>Chl</td> <td>3.90</td> <td>0.48</td> <td></td> </tr> <tr> <td rowspan="2"><i>Staphylococcus aureus</i></td> <td>MNEO</td> <td>78.13</td> <td>4.88</td> <td>0.56 (A)</td> </tr> <tr> <td>Chl</td> <td>3.90</td> <td>1.95</td> <td></td> </tr> <tr> <td rowspan="2"><i>Escherichia coli</i></td> <td>MNEO</td> <td>1250.00</td> <td>312.50</td> <td>0.31 (S)</td> </tr> <tr> <td>Chl</td> <td>3.90</td> <td>0.24</td> <td></td> </tr> <tr> <td rowspan="2"><i>Pseudomonas aeruginosa</i></td> <td>MNEO</td> <td>625.00</td> <td>39.10</td> <td>0.13 (S)</td> </tr> <tr> <td>Chl</td> <td>15.60</td> <td>0.98</td> <td></td> </tr> </tbody> </table> <p>MICa: MIC alone; MICc: MIC combined; Chl: chloramphenicol. S, synergy; A, additivity.</p>	Microorganism		MICa ($\mu\text{g/mL}$)	MICc ($\mu\text{g/mL}$)	FICI	<i>Bacillus subtilis</i>	MNEO	39.10	9.76	0.37 (S)	Chl	3.90	0.48		<i>Staphylococcus aureus</i>	MNEO	78.13	4.88	0.56 (A)	Chl	3.90	1.95		<i>Escherichia coli</i>	MNEO	1250.00	312.50	0.31 (S)	Chl	3.90	0.24		<i>Pseudomonas aeruginosa</i>	MNEO	625.00	39.10	0.13 (S)	Chl	15.60	0.98		Table 3	Fractional inhibitory concentration index (FICI) values of the essential oil of <i>M. nepalensis</i> (MNEO) and chloramphenicol combinations.
Microorganism		MICa ($\mu\text{g/mL}$)	MICc ($\mu\text{g/mL}$)	FICI																																										
<i>Bacillus subtilis</i>	MNEO	39.10	9.76	0.37 (S)																																										
	Chl	3.90	0.48																																											
<i>Staphylococcus aureus</i>	MNEO	78.13	4.88	0.56 (A)																																										
	Chl	3.90	1.95																																											
<i>Escherichia coli</i>	MNEO	1250.00	312.50	0.31 (S)																																										
	Chl	3.90	0.24																																											
<i>Pseudomonas aeruginosa</i>	MNEO	625.00	39.10	0.13 (S)																																										
	Chl	15.60	0.98																																											
https://europepmc.org/article/MED/35889501	Preliminary Study on Phytochemical Constituents and Biological Activities of Essential Oil from <i>Myriactis nepalensis</i> Less.	https://europepmc.org/articles/PMC9324352/table/molecules-27-0463-1-t004/	<p>Table 4</p> <p>Fractional inhibitory concentration index (FICI) values of the essential oil of <i>M. nepalensis</i> (MNEO) and streptomycin combinations.</p> <table border="1"> <thead> <tr> <th>Microorganism</th> <th></th> <th>MICa ($\mu\text{g/mL}$)</th> <th>MICc ($\mu\text{g/mL}$)</th> <th>FICI</th> </tr> </thead> <tbody> <tr> <td rowspan="2"><i>Bacillus subtilis</i></td> <td>MNEO</td> <td>39.00</td> <td>4.88</td> <td>0.37 (S)</td> </tr> <tr> <td>SM</td> <td>0.49</td> <td>0.12</td> <td></td> </tr> <tr> <td rowspan="2"><i>Staphylococcus aureus</i></td> <td>MNEO</td> <td>78.13</td> <td>4.88</td> <td>0.12 (S)</td> </tr> <tr> <td>SM</td> <td>1.95</td> <td>0.12</td> <td></td> </tr> <tr> <td rowspan="2"><i>Escherichia coli</i></td> <td>MNEO</td> <td>1250.00</td> <td>312.50</td> <td>0.50 (S)</td> </tr> <tr> <td>SM</td> <td>3.90</td> <td>0.98</td> <td></td> </tr> <tr> <td rowspan="2"><i>Pseudomonas aeruginosa</i></td> <td>MNEO</td> <td>625.00</td> <td>39.00</td> <td>0.12 (S)</td> </tr> <tr> <td>SM</td> <td>1.95</td> <td>0.12</td> <td></td> </tr> </tbody> </table> <p>MICa: MIC alone; MICc: MIC combined; SM: streptomycin. S, synergy.</p>	Microorganism		MICa ($\mu\text{g/mL}$)	MICc ($\mu\text{g/mL}$)	FICI	<i>Bacillus subtilis</i>	MNEO	39.00	4.88	0.37 (S)	SM	0.49	0.12		<i>Staphylococcus aureus</i>	MNEO	78.13	4.88	0.12 (S)	SM	1.95	0.12		<i>Escherichia coli</i>	MNEO	1250.00	312.50	0.50 (S)	SM	3.90	0.98		<i>Pseudomonas aeruginosa</i>	MNEO	625.00	39.00	0.12 (S)	SM	1.95	0.12		Table 4	MICa: MIC alone; MICc: MIC combined; SM: streptomycin. S, synergy.
Microorganism		MICa ($\mu\text{g/mL}$)	MICc ($\mu\text{g/mL}$)	FICI																																										
<i>Bacillus subtilis</i>	MNEO	39.00	4.88	0.37 (S)																																										
	SM	0.49	0.12																																											
<i>Staphylococcus aureus</i>	MNEO	78.13	4.88	0.12 (S)																																										
	SM	1.95	0.12																																											
<i>Escherichia coli</i>	MNEO	1250.00	312.50	0.50 (S)																																										
	SM	3.90	0.98																																											
<i>Pseudomonas aeruginosa</i>	MNEO	625.00	39.00	0.12 (S)																																										
	SM	1.95	0.12																																											
https://europepmc.org/article/MED/35889501	Preliminary Study on Phytochemical Constituents and Biological Activities of Essential Oil from <i>Myriactis nepalensis</i> Less.	https://europepmc.org/articles/PMC9324352/table/molecules-27-0463-1-t005/	<p>Table 5</p> <p>IC₅₀ ($\mu\text{g/mL}$) \pm SD values to the essential oil of <i>M. nepalensis</i> and positive control doxorubicin against cell lines.</p> <table border="1"> <thead> <tr> <th>Cell Line</th> <th>Essential Oil</th> <th>Doxorubicin</th> </tr> </thead> <tbody> <tr> <td>HepG2</td> <td>19.53 \pm 2.84</td> <td>0.46 \pm 0.02</td> </tr> <tr> <td>MCF-7</td> <td>13.13 \pm 1.90</td> <td>0.70 \pm 0.05</td> </tr> <tr> <td>HL-7702</td> <td>19.14 \pm 0.63</td> <td>0.60 \pm 0.13</td> </tr> <tr> <td>A-549</td> <td>19.19 \pm 3.08</td> <td>0.48 \pm 0.01</td> </tr> <tr> <td>HCT-116</td> <td>35.22 \pm 8.36</td> <td>0.57 \pm 0.03</td> </tr> </tbody> </table> <p>Open in a separate window</p>	Cell Line	Essential Oil	Doxorubicin	HepG2	19.53 \pm 2.84	0.46 \pm 0.02	MCF-7	13.13 \pm 1.90	0.70 \pm 0.05	HL-7702	19.14 \pm 0.63	0.60 \pm 0.13	A-549	19.19 \pm 3.08	0.48 \pm 0.01	HCT-116	35.22 \pm 8.36	0.57 \pm 0.03	Table 5	IC ₅₀ ($\mu\text{g/mL}$) \pm SD values to the essential oil of <i>M. nepalensis</i> and positive control doxorubicin against cell lines.																							
Cell Line	Essential Oil	Doxorubicin																																												
HepG2	19.53 \pm 2.84	0.46 \pm 0.02																																												
MCF-7	13.13 \pm 1.90	0.70 \pm 0.05																																												
HL-7702	19.14 \pm 0.63	0.60 \pm 0.13																																												
A-549	19.19 \pm 3.08	0.48 \pm 0.01																																												
HCT-116	35.22 \pm 8.36	0.57 \pm 0.03																																												

URL	Title	Table Link	Table Pic (top)		Table Header	Table footer																																																																																																																																																																																																										
https://europepmc.org/article/MED/35406974#free-full-text	The Phytochemical Profile and Anticancer Activity of Anthemis tinctoria and Angelica sylvestris Used in Estonian Ethnomedicine.	https://europepmc.org/articles/PMC9003001/table/plants-11-00994-t007/	Table 7 Anticancer activity of essential oils and methanolic extracts of <i>A. tinctoria</i> and <i>A. sylvestris</i> . <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">% Inhibition</th> </tr> <tr> <th colspan="5">Essential oil of <i>A. tinctoria</i> (Aerial Parts)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>91.80</td> <td>97.07</td> <td>81.11</td> <td>93.85</td> <td>96.32</td> </tr> <tr> <td>20</td> <td>25.33</td> <td>28.47</td> <td>23.85</td> <td>38.51</td> <td>40.18</td> </tr> <tr> <td>4</td> <td>12.88</td> <td>19.99</td> <td>17.81</td> <td>28.31</td> <td>25.99</td> </tr> <tr> <td>0.8</td> <td>0.13</td> <td>8.81</td> <td>8.59</td> <td>12.23</td> <td>10.03</td> </tr> <tr> <td>IC₅₀</td> <td>44.98 ± 2.96</td> <td>43.04 ± 4.50</td> <td>55.45 ± 5.70</td> <td>29.96 ± 2.25</td> <td>27.75 ± 1.86</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">Essential oil of <i>A. sylvestris</i> (aerial parts)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>87.08</td> <td>88.79</td> <td>103.03</td> <td>90.21</td> <td>96.76</td> </tr> <tr> <td>20</td> <td>33.14</td> <td>32.31</td> <td>38.47</td> <td>42.23</td> <td>52.30</td> </tr> <tr> <td>4</td> <td>11.61</td> <td>15.34</td> <td>13.35</td> <td>20.35</td> <td>18.36</td> </tr> <tr> <td>0.8</td> <td>-2.02</td> <td>1.93</td> <td>4.45</td> <td>7.21</td> <td>4.92</td> </tr> <tr> <td>IC₅₀</td> <td>37.46 ± 2.33</td> <td>38.06 ± 2.09</td> <td>30.72 ± 1.81</td> <td>27.78 ± 1.28</td> <td>19.84 ± 2.35</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">Essential oil of <i>A. sylvestris</i> (roots)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>97.82</td> <td>99.34</td> <td>75.35</td> <td>95.08</td> <td>97.22</td> </tr> <tr> <td>20</td> <td>44.11</td> <td>36.14</td> <td>40.70</td> <td>39.34</td> <td>49.19</td> </tr> <tr> <td>4</td> <td>20.33</td> <td>12.44</td> <td>27.66</td> <td>21.86</td> <td>28.60</td> </tr> <tr> <td>0.8</td> <td>5.14</td> <td>6.00</td> <td>16.85</td> <td>10.53</td> <td>14.28</td> </tr> <tr> <td>IC₅₀</td> <td>24.69 ± 1.96</td> <td>34.09 ± 2.08</td> <td>33.36 ± 2.25</td> <td>30.37 ± 2.35</td> <td>19.73 ± 2.18</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">Methanolic extract of <i>A. tinctoria</i> (aerial parts)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>21.29</td> <td>30.50</td> <td>29.89</td> <td>29.83</td> <td>25.36</td> </tr> <tr> <td>20</td> <td>14.28</td> <td>8.76</td> <td>5.09</td> <td>8.83</td> <td>10.46</td> </tr> <tr> <td>IC₅₀</td> <td>>100</td> <td>>100</td> <td>>100</td> <td>>100</td> <td>>100</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">Methanolic extract of <i>A. sylvestris</i> (aerial parts)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>14.50</td> <td>29.69</td> <td>20.61</td> <td>36.32</td> <td>19.62</td> </tr> <tr> <td>20</td> <td>2.91</td> <td>9.65</td> <td>7.00</td> <td>7.85</td> <td>8.12</td> </tr> <tr> <td>IC₅₀</td> <td>>100</td> <td>>100</td> <td>>100</td> <td>>100</td> <td>>100</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th rowspan="2">Concentration (µg/mL)</th> <th colspan="5">Methanolic extract of <i>A. sylvestris</i> (roots)</th> </tr> <tr> <th>HepG2</th> <th>MKN7</th> <th>SW480</th> <th>LNCaP</th> <th>KB</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>67.48</td> <td>68.94</td> <td>63.43</td> <td>76.48</td> <td>73.07</td> </tr> </tbody> </table> <p>HepG2, human hepatocellular carcinoma; MKN7, human gastric carcinoma; SW480, human colon carcinoma; LNCaP, human prostate carcinoma; KB, human carcinoma in the mouth. Ellipticine: the positive control sample; * the positive control sample.</p>	Concentration (µg/mL)	% Inhibition					Essential oil of <i>A. tinctoria</i> (Aerial Parts)					HepG2	MKN7	SW480	LNCaP	KB	100	91.80	97.07	81.11	93.85	96.32	20	25.33	28.47	23.85	38.51	40.18	4	12.88	19.99	17.81	28.31	25.99	0.8	0.13	8.81	8.59	12.23	10.03	IC ₅₀	44.98 ± 2.96	43.04 ± 4.50	55.45 ± 5.70	29.96 ± 2.25	27.75 ± 1.86	Concentration (µg/mL)	Essential oil of <i>A. sylvestris</i> (aerial parts)					HepG2	MKN7	SW480	LNCaP	KB	100	87.08	88.79	103.03	90.21	96.76	20	33.14	32.31	38.47	42.23	52.30	4	11.61	15.34	13.35	20.35	18.36	0.8	-2.02	1.93	4.45	7.21	4.92	IC ₅₀	37.46 ± 2.33	38.06 ± 2.09	30.72 ± 1.81	27.78 ± 1.28	19.84 ± 2.35	Concentration (µg/mL)	Essential oil of <i>A. sylvestris</i> (roots)					HepG2	MKN7	SW480	LNCaP	KB	100	97.82	99.34	75.35	95.08	97.22	20	44.11	36.14	40.70	39.34	49.19	4	20.33	12.44	27.66	21.86	28.60	0.8	5.14	6.00	16.85	10.53	14.28	IC ₅₀	24.69 ± 1.96	34.09 ± 2.08	33.36 ± 2.25	30.37 ± 2.35	19.73 ± 2.18	Concentration (µg/mL)	Methanolic extract of <i>A. tinctoria</i> (aerial parts)					HepG2	MKN7	SW480	LNCaP	KB	100	21.29	30.50	29.89	29.83	25.36	20	14.28	8.76	5.09	8.83	10.46	IC ₅₀	>100	>100	>100	>100	>100	Concentration (µg/mL)	Methanolic extract of <i>A. sylvestris</i> (aerial parts)					HepG2	MKN7	SW480	LNCaP	KB	100	14.50	29.69	20.61	36.32	19.62	20	2.91	9.65	7.00	7.85	8.12	IC ₅₀	>100	>100	>100	>100	>100	Concentration (µg/mL)	Methanolic extract of <i>A. sylvestris</i> (roots)					HepG2	MKN7	SW480	LNCaP	KB	100	67.48	68.94	63.43	76.48	73.07	Table 7 Anticancer activity of essential oils and methanolic extracts of <i>A. tinctoria</i> and <i>A. sylvestris</i> .	HepG2, human hepatocellular carcinoma; MKN7, human gastric carcinoma; SW480, human colon carcinoma; LNCaP, human prostate carcinoma; KB, human carcinoma in the mouth. Ellipticine: the positive control sample; * the positive control sample.
Concentration (µg/mL)	% Inhibition																																																																																																																																																																																																															
	Essential oil of <i>A. tinctoria</i> (Aerial Parts)																																																																																																																																																																																																															
HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																												
100	91.80	97.07	81.11	93.85	96.32																																																																																																																																																																																																											
20	25.33	28.47	23.85	38.51	40.18																																																																																																																																																																																																											
4	12.88	19.99	17.81	28.31	25.99																																																																																																																																																																																																											
0.8	0.13	8.81	8.59	12.23	10.03																																																																																																																																																																																																											
IC ₅₀	44.98 ± 2.96	43.04 ± 4.50	55.45 ± 5.70	29.96 ± 2.25	27.75 ± 1.86																																																																																																																																																																																																											
Concentration (µg/mL)	Essential oil of <i>A. sylvestris</i> (aerial parts)																																																																																																																																																																																																															
	HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																											
100	87.08	88.79	103.03	90.21	96.76																																																																																																																																																																																																											
20	33.14	32.31	38.47	42.23	52.30																																																																																																																																																																																																											
4	11.61	15.34	13.35	20.35	18.36																																																																																																																																																																																																											
0.8	-2.02	1.93	4.45	7.21	4.92																																																																																																																																																																																																											
IC ₅₀	37.46 ± 2.33	38.06 ± 2.09	30.72 ± 1.81	27.78 ± 1.28	19.84 ± 2.35																																																																																																																																																																																																											
Concentration (µg/mL)	Essential oil of <i>A. sylvestris</i> (roots)																																																																																																																																																																																																															
	HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																											
100	97.82	99.34	75.35	95.08	97.22																																																																																																																																																																																																											
20	44.11	36.14	40.70	39.34	49.19																																																																																																																																																																																																											
4	20.33	12.44	27.66	21.86	28.60																																																																																																																																																																																																											
0.8	5.14	6.00	16.85	10.53	14.28																																																																																																																																																																																																											
IC ₅₀	24.69 ± 1.96	34.09 ± 2.08	33.36 ± 2.25	30.37 ± 2.35	19.73 ± 2.18																																																																																																																																																																																																											
Concentration (µg/mL)	Methanolic extract of <i>A. tinctoria</i> (aerial parts)																																																																																																																																																																																																															
	HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																											
100	21.29	30.50	29.89	29.83	25.36																																																																																																																																																																																																											
20	14.28	8.76	5.09	8.83	10.46																																																																																																																																																																																																											
IC ₅₀	>100	>100	>100	>100	>100																																																																																																																																																																																																											
Concentration (µg/mL)	Methanolic extract of <i>A. sylvestris</i> (aerial parts)																																																																																																																																																																																																															
	HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																											
100	14.50	29.69	20.61	36.32	19.62																																																																																																																																																																																																											
20	2.91	9.65	7.00	7.85	8.12																																																																																																																																																																																																											
IC ₅₀	>100	>100	>100	>100	>100																																																																																																																																																																																																											
Concentration (µg/mL)	Methanolic extract of <i>A. sylvestris</i> (roots)																																																																																																																																																																																																															
	HepG2	MKN7	SW480	LNCaP	KB																																																																																																																																																																																																											
100	67.48	68.94	63.43	76.48	73.07																																																																																																																																																																																																											
https://europepmc.org/article/MED/35707466#free-full-text	Comparative Analysis of Chemical Composition and Antibacterial and Anti-Inflammatory Activities of the Essential Oils from Chrysanthemum morifolium of Different Flowering Stages and Different Parts.	https://europepmc.org/articles/PMC9192287/table/tab2/	Table 2 Antibacterial activities (MIC mg/mL) of six essential oils of <i>C. morifolium</i> . <table border="1"> <thead> <tr> <th rowspan="2">Number</th> <th rowspan="2">Samples</th> <th><i>Propionibacterium acacia</i> (ATCC6919)</th> <th><i>Staphylococcus aureus</i> (ATCC25923)</th> </tr> <tr> <th>MIC^a (mg/mL)</th> <th>MIC (mg/mL)</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>JM</td> <td>25</td> <td>10</td> </tr> <tr> <td>2</td> <td>TJ</td> <td>25</td> <td>10</td> </tr> <tr> <td>3</td> <td>JH</td> <td>25</td> <td>10</td> </tr> <tr> <td>4</td> <td>SLC</td> <td>25</td> <td>20</td> </tr> <tr> <td>5</td> <td>RC</td> <td>50</td> <td>50</td> </tr> <tr> <td>6</td> <td>Solvent control</td> <td>+^b</td> <td>+</td> </tr> <tr> <td>7</td> <td>Negative control</td> <td>-^c</td> <td>-</td> </tr> <tr> <td>8</td> <td>Penicillin</td> <td>/^d</td> <td>0.0035</td> </tr> <tr> <td>9</td> <td>Erythromycin lactobionate</td> <td>0.061</td> <td>—</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>^aMIC: minimum inhibitory concentration. ^b+: no bacteria grew, indicating that there were no other miscellaneous bacteria in the blank solvent and medium. ^c-: there is bacterial growth, showing that the solvent has no inhibitory effect on bacteria. ^d/: the minimum inhibitory concentration (MIC) of this bacterium was not tested.</p>	Number	Samples	<i>Propionibacterium acacia</i> (ATCC6919)	<i>Staphylococcus aureus</i> (ATCC25923)	MIC ^a (mg/mL)	MIC (mg/mL)	1	JM	25	10	2	TJ	25	10	3	JH	25	10	4	SLC	25	20	5	RC	50	50	6	Solvent control	+ ^b	+	7	Negative control	- ^c	-	8	Penicillin	/ ^d	0.0035	9	Erythromycin lactobionate	0.061	—	Table 2 Antibacterial activities (MIC mg/mL) of six essential oils of <i>C. morifolium</i> .	^a MIC: minimum inhibitory concentration. ^b +: no bacteria grew, indicating that there were no other miscellaneous bacteria in the blank solvent and medium. ^c -: there is bacterial growth, showing that the solvent has no inhibitory effect on bacteria. ^d /: the minimum inhibitory concentration (MIC) of this bacterium was not tested.																																																																																																																																																																	
Number	Samples	<i>Propionibacterium acacia</i> (ATCC6919)	<i>Staphylococcus aureus</i> (ATCC25923)																																																																																																																																																																																																													
		MIC ^a (mg/mL)	MIC (mg/mL)																																																																																																																																																																																																													
1	JM	25	10																																																																																																																																																																																																													
2	TJ	25	10																																																																																																																																																																																																													
3	JH	25	10																																																																																																																																																																																																													
4	SLC	25	20																																																																																																																																																																																																													
5	RC	50	50																																																																																																																																																																																																													
6	Solvent control	+ ^b	+																																																																																																																																																																																																													
7	Negative control	- ^c	-																																																																																																																																																																																																													
8	Penicillin	/ ^d	0.0035																																																																																																																																																																																																													
9	Erythromycin lactobionate	0.061	—																																																																																																																																																																																																													
https://europepmc.org/article/MED/35883828	Valeriana pilosa Roots Essential Oil: Chemical Composition, Antioxidant Activities, and Molecular Docking Studies on Enzymes Involved in Redox Biological Processes.	https://europepmc.org/articles/PMC9311991/table/antioxidants-11-01337-t002/	Table 2 Antioxidant activities of essential oil of <i>Valeriana pilosa</i> . <table border="1"> <thead> <tr> <th rowspan="2">Samples</th> <th>FRAP (mM TEAC)</th> <th>ABTS⁺ IC₅₀</th> <th>DPPH IC₅₀</th> </tr> <tr> <th></th> <th></th> <th></th> </tr> </thead> <tbody> <tr> <td>VPEO</td> <td>0.0421 ± 0.02</td> <td>0.30 ± 0.05</td> <td>0.38 ± 0.07</td> </tr> <tr> <td>Quercetin</td> <td>143.00 ± 0.04</td> <td>0.07 ± 0.03</td> <td>0.06 ± 0.02</td> </tr> <tr> <td>Trolox®</td> <td>-</td> <td>0.012 ± 0.07</td> <td>0.011 ± 0.04</td> </tr> </tbody> </table> <p>FRAP = ferric-reducing antioxidant power; ABTS⁺ = 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid); DPPH = 2,2-diphenyl-1-picrylhydrazyl radical; GAE = gallic acid equivalent; TEAC = Trolox® equivalent antioxidant capacity. Results are expressed as mean values ± SEM (n = 3).</p>	Samples	FRAP (mM TEAC)	ABTS ⁺ IC ₅₀	DPPH IC ₅₀				VPEO	0.0421 ± 0.02	0.30 ± 0.05	0.38 ± 0.07	Quercetin	143.00 ± 0.04	0.07 ± 0.03	0.06 ± 0.02	Trolox®	-	0.012 ± 0.07	0.011 ± 0.04	Table 2 Antioxidant activities of essential oil of <i>Valeriana pilosa</i> .	FRAP = ferric-reducing antioxidant power; ABTS ⁺ = 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid); DPPH = 2,2-diphenyl-1-picrylhydrazyl radical; GAE = gallic acid equivalent; TEAC = Trolox® equivalent antioxidant capacity. Results are expressed as mean values ± SEM (n = 3).																																																																																																																																																																																								
Samples	FRAP (mM TEAC)	ABTS ⁺ IC ₅₀	DPPH IC ₅₀																																																																																																																																																																																																													
VPEO	0.0421 ± 0.02	0.30 ± 0.05	0.38 ± 0.07																																																																																																																																																																																																													
Quercetin	143.00 ± 0.04	0.07 ± 0.03	0.06 ± 0.02																																																																																																																																																																																																													
Trolox®	-	0.012 ± 0.07	0.011 ± 0.04																																																																																																																																																																																																													

URL	Title	Table Link	Table Pic (top)	Table Header	Table footer																																																																							
https://europepmc.org/article/MED/35630757#free-full-text	Phytochemical Analysis, Antibacterial Activity and Modulating Effect of Essential Oil from <i>Syzygium cumini</i> (L.) Skeels.	https://europepmc.org/articles/PMC9145283/table/molecules-27-0328-1-t002/	<p>Table 2</p> <p>Minimum inhibitory concentration ($\mu\text{g}/\text{mL}$) of the <i>S. cumini</i> EO against the standard multidrug-resistant strains.</p> <table border="1"> <thead> <tr> <th>Strains</th> <th><i>E. coli</i></th> <th><i>P. aeruginosa</i></th> <th><i>S. aureus</i></th> </tr> </thead> <tbody> <tr> <td>Standard strains</td> <td>ATCC 25922</td> <td>ATCC 25853</td> <td>ATCC 25923</td> </tr> <tr> <td></td> <td>512</td> <td>≥ 1024</td> <td>≥ 1024</td> </tr> <tr> <td>Multidrug-resistant strains</td> <td>EC06</td> <td>PA24</td> <td>SA10</td> </tr> <tr> <td>EOSC</td> <td>≥ 1024</td> <td>≥ 1024</td> <td>≥ 1024</td> </tr> <tr> <td>Gentamicin</td> <td>18</td> <td>4</td> <td>35</td> </tr> <tr> <td>Erythromycin</td> <td>25</td> <td>32</td> <td>23</td> </tr> <tr> <td>Norfloxacin</td> <td>290</td> <td>56</td> <td>300</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p>	Strains	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>	Standard strains	ATCC 25922	ATCC 25853	ATCC 25923		512	≥ 1024	≥ 1024	Multidrug-resistant strains	EC06	PA24	SA10	EOSC	≥ 1024	≥ 1024	≥ 1024	Gentamicin	18	4	35	Erythromycin	25	32	23	Norfloxacin	290	56	300	Table 2	Minimum inhibitory concentration ($\mu\text{g}/\text{mL}$) of the <i>S. cumini</i> EO against the standard multidrug-resistant strains.																																							
Strains	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>																																																																									
Standard strains	ATCC 25922	ATCC 25853	ATCC 25923																																																																									
	512	≥ 1024	≥ 1024																																																																									
Multidrug-resistant strains	EC06	PA24	SA10																																																																									
EOSC	≥ 1024	≥ 1024	≥ 1024																																																																									
Gentamicin	18	4	35																																																																									
Erythromycin	25	32	23																																																																									
Norfloxacin	290	56	300																																																																									
https://europepmc.org/article/MED/35140337#free-full-text	Chemical composition, antioxidant, and antimicrobial activity of <i>Elsholtzia beddomei</i> C. B. Clarke ex Hook. f. essential oil.	https://europepmc.org/articles/PMC8828703/table/Tab2/	<p>Table 2</p> <p>Antioxidant activities of essential oil and standards.</p> <table border="1"> <thead> <tr> <th rowspan="2">Sample</th> <th>Yield</th> <th colspan="2">IC_{50} ($\mu\text{g}/\text{mL}$)</th> </tr> <tr> <th>(%w/w)</th> <th>DPPH</th> <th>ABTS</th> </tr> </thead> <tbody> <tr> <td><i>E. beddomei</i> essential oil</td> <td>1.38 \pm 0.21</td> <td>148.31 \pm 0.23^b</td> <td>172.22 \pm 0.32^b</td> </tr> <tr> <td>trolox</td> <td>-</td> <td>7.34 \pm 0.12^a</td> <td>11.31 \pm 0.15^a</td> </tr> <tr> <td>linalool</td> <td></td> <td>201.23 \pm 0.26^d</td> <td>226.28 \pm 0.18^d</td> </tr> <tr> <td>perillaldehyde</td> <td></td> <td>186.43 \pm 0.27^c</td> <td>197.61 \pm 0.63^c</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>The data are mean \pm standard deviation. Different letters indicate significant differences ($p < 0.05$).</p>	Sample	Yield	IC_{50} ($\mu\text{g}/\text{mL}$)		(%w/w)	DPPH	ABTS	<i>E. beddomei</i> essential oil	1.38 \pm 0.21	148.31 \pm 0.23 ^b	172.22 \pm 0.32 ^b	trolox	-	7.34 \pm 0.12 ^a	11.31 \pm 0.15 ^a	linalool		201.23 \pm 0.26 ^d	226.28 \pm 0.18 ^d	perillaldehyde		186.43 \pm 0.27 ^c	197.61 \pm 0.63 ^c	Table 2	The data are mean \pm standard deviation. Different letters indicate significant differences ($p < 0.05$).																																																
Sample	Yield	IC_{50} ($\mu\text{g}/\text{mL}$)																																																																										
	(%w/w)	DPPH	ABTS																																																																									
<i>E. beddomei</i> essential oil	1.38 \pm 0.21	148.31 \pm 0.23 ^b	172.22 \pm 0.32 ^b																																																																									
trolox	-	7.34 \pm 0.12 ^a	11.31 \pm 0.15 ^a																																																																									
linalool		201.23 \pm 0.26 ^d	226.28 \pm 0.18 ^d																																																																									
perillaldehyde		186.43 \pm 0.27 ^c	197.61 \pm 0.63 ^c																																																																									
https://europepmc.org/article/MED/35140337#free-full-text	Chemical composition, antioxidant, and antimicrobial activity of <i>Elsholtzia beddomei</i> C. B. Clarke ex Hook. f. essential oil.	https://europepmc.org/articles/PMC8828703/table/Tab3/	<p>Table 3</p> <p>Antimicrobial activity of <i>E. beddomei</i> essential oil and chloramphenicol.</p> <table border="1"> <thead> <tr> <th rowspan="2">Microorganism</th> <th colspan="2">Chloramphenicol</th> <th colspan="3"><i>E. beddomei</i> essent</th> </tr> <tr> <th>MIC ($\mu\text{g}/\text{mL}$)</th> <th>MMC ($\mu\text{g}/\text{mL}$)</th> <th>Zone of inhibition diameter (mm)</th> <th>MIC ($\mu\text{g}/\text{mL}$)</th> <th>MMC ($\mu\text{g}/\text{mL}$)</th> </tr> </thead> <tbody> <tr> <td>Gram-positive bacteria</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>S. aureus</i> ATCC 25923</td> <td>3.91^a</td> <td>15.62^a</td> <td>35.13 \pm 1.44^a</td> <td>3.91^a</td> <td>31.25^b</td> </tr> <tr> <td><i>S. epidermidis</i> ATCC 12228</td> <td>3.91^a</td> <td>15.62^a</td> <td>34.94 \pm 0.81^a</td> <td>3.91^a</td> <td>15.62^a</td> </tr> <tr> <td><i>B. subtilis</i> ATCC 6051</td> <td>3.91^a</td> <td>15.62^a</td> <td>34.88 \pm 1.21^a</td> <td>3.91^a</td> <td>15.62^a</td> </tr> <tr> <td>Gram-negative bacteria</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>E. coli</i> ATCC 25922</td> <td>7.81^b</td> <td>62.50^c</td> <td>28.91 \pm 0.52^c</td> <td>7.81^b</td> <td>62.50^c</td> </tr> <tr> <td><i>P. aeruginosa</i> ATCC 27853</td> <td>3.91^a</td> <td>31.25^b</td> <td>32.24 \pm 0.52^b</td> <td>7.81^b</td> <td>62.50^c</td> </tr> <tr> <td><i>E. aerogenes</i> ATCC 13048</td> <td>7.81^b</td> <td>62.50^c</td> <td>28.81 \pm 0.48^c</td> <td>7.81^b</td> <td>62.50^c</td> </tr> <tr> <td>Fungus</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>C. albicans</i> ATCC 10231</td> <td>7.81^b</td> <td>62.50^c</td> <td>28.01 \pm 0.21^c</td> <td>15.62^c</td> <td>125.00^d</td> </tr> </tbody> </table> <p style="text-align: right;">Open in a separate window</p> <p>The data are mean \pm standard deviation. Different letters indicate significant differences ($p < 0.05$).</p>	Microorganism	Chloramphenicol		<i>E. beddomei</i> essent			MIC ($\mu\text{g}/\text{mL}$)	MMC ($\mu\text{g}/\text{mL}$)	Zone of inhibition diameter (mm)	MIC ($\mu\text{g}/\text{mL}$)	MMC ($\mu\text{g}/\text{mL}$)	Gram-positive bacteria						<i>S. aureus</i> ATCC 25923	3.91 ^a	15.62 ^a	35.13 \pm 1.44 ^a	3.91 ^a	31.25 ^b	<i>S. epidermidis</i> ATCC 12228	3.91 ^a	15.62 ^a	34.94 \pm 0.81 ^a	3.91 ^a	15.62 ^a	<i>B. subtilis</i> ATCC 6051	3.91 ^a	15.62 ^a	34.88 \pm 1.21 ^a	3.91 ^a	15.62 ^a	Gram-negative bacteria						<i>E. coli</i> ATCC 25922	7.81 ^b	62.50 ^c	28.91 \pm 0.52 ^c	7.81 ^b	62.50 ^c	<i>P. aeruginosa</i> ATCC 27853	3.91 ^a	31.25 ^b	32.24 \pm 0.52 ^b	7.81 ^b	62.50 ^c	<i>E. aerogenes</i> ATCC 13048	7.81 ^b	62.50 ^c	28.81 \pm 0.48 ^c	7.81 ^b	62.50 ^c	Fungus						<i>C. albicans</i> ATCC 10231	7.81 ^b	62.50 ^c	28.01 \pm 0.21 ^c	15.62 ^c	125.00 ^d	Table 3	The data are mean \pm standard deviation. Different letters indicate significant differences ($p < 0.05$).
Microorganism	Chloramphenicol		<i>E. beddomei</i> essent																																																																									
	MIC ($\mu\text{g}/\text{mL}$)	MMC ($\mu\text{g}/\text{mL}$)	Zone of inhibition diameter (mm)	MIC ($\mu\text{g}/\text{mL}$)	MMC ($\mu\text{g}/\text{mL}$)																																																																							
Gram-positive bacteria																																																																												
<i>S. aureus</i> ATCC 25923	3.91 ^a	15.62 ^a	35.13 \pm 1.44 ^a	3.91 ^a	31.25 ^b																																																																							
<i>S. epidermidis</i> ATCC 12228	3.91 ^a	15.62 ^a	34.94 \pm 0.81 ^a	3.91 ^a	15.62 ^a																																																																							
<i>B. subtilis</i> ATCC 6051	3.91 ^a	15.62 ^a	34.88 \pm 1.21 ^a	3.91 ^a	15.62 ^a																																																																							
Gram-negative bacteria																																																																												
<i>E. coli</i> ATCC 25922	7.81 ^b	62.50 ^c	28.91 \pm 0.52 ^c	7.81 ^b	62.50 ^c																																																																							
<i>P. aeruginosa</i> ATCC 27853	3.91 ^a	31.25 ^b	32.24 \pm 0.52 ^b	7.81 ^b	62.50 ^c																																																																							
<i>E. aerogenes</i> ATCC 13048	7.81 ^b	62.50 ^c	28.81 \pm 0.48 ^c	7.81 ^b	62.50 ^c																																																																							
Fungus																																																																												
<i>C. albicans</i> ATCC 10231	7.81 ^b	62.50 ^c	28.01 \pm 0.21 ^c	15.62 ^c	125.00 ^d																																																																							

URL	Title	Table Link	Table Pic (top)		Table Header	Table footer																																																																																																																																																																															
https://europepmc.org/article/MED/34950215#free-full-text	Lamiaceae Essential Oils, Phytochemical Profile, Antioxidant, and Biological Activities.	https://europepmc.org/articles/PMC8692021/table/tabc2/	<p>Table 2 Antioxidant activity of essential oils from <i>Lamiaceae</i> species.</p> <table border="1"> <thead> <tr> <th>Species and plant part</th><th>Technique</th><th>Results</th><th>Reference</th></tr> </thead> <tbody> <tr> <td><i>Cedronella canariensis</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 615.5 ± 76.5 µg/mL</td><td>[114]</td></tr> <tr> <td></td><td>ABTS</td><td>IC₅₀ = 10.5 ± 0.6 µg/mL</td><td></td></tr> <tr> <td></td><td>FRAP</td><td>IC₅₀ = 3.8 ± 1.46 µmol TE/g</td><td></td></tr> <tr> <td><i>Mentha piperita</i> (leaves)</td><td>DPPH</td><td>Radical scavenging = 92.6 ± 6.86%</td><td>[115]</td></tr> <tr> <td></td><td>Reducing power</td><td>Reducing power = 0.9 ± 0.3</td><td></td></tr> <tr> <td><i>M. pulegium</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 321.41 ± 2.53 µg/mL</td><td>[116]</td></tr> <tr> <td></td><td>FRAP</td><td>IC₅₀ = 58.27 ± 2.72 µg/mL</td><td></td></tr> <tr> <td><i>M. rotundifolia</i> (leaves)</td><td>DPPH</td><td>IC₅₀ = 2222.2 ± 25.2 µg/mL</td><td>[117]</td></tr> <tr> <td></td><td>ABTS</td><td>IC₅₀ = 133.8 ± 4.8 µg/mL</td><td></td></tr> <tr> <td></td><td>Reducing power</td><td>IC₅₀ = 166.6 ± 1.9 µg/mL</td><td></td></tr> <tr> <td></td><td>Phosphomolybdate</td><td>IC₅₀ = 45.2 ± 1.2 µg/mL</td><td></td></tr> <tr> <td><i>M. spicata</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 3450 ± 172.5 µg/mL</td><td>[118]</td></tr> <tr> <td></td><td>ABTS</td><td>IC₅₀ = 40.2 ± 0.2 µg/mL</td><td></td></tr> <tr> <td></td><td>FRAP</td><td>IC₅₀ = 215 ± 4.50 µg/mL</td><td></td></tr> <tr> <td><i>M. spicata</i> (leaves)</td><td>DPPH</td><td>IC₅₀ = 41.23 µg/mL</td><td>[119]</td></tr> <tr> <td><i>O. basilicum</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 4.04 ± 0.09–0.21 ± 0.02 mg/mL</td><td>[120]</td></tr> <tr> <td></td><td>β-carotene</td><td>Bleaching = 23.8 ± 0.6–85.3 ± 1.0%</td><td></td></tr> <tr> <td><i>Origanum dictamnus</i> (flowers)</td><td>DPPH</td><td>IC₅₀ = 0.0459 ± 0.0042% (v/v)</td><td>[121]</td></tr> <tr> <td><i>O. floribundum</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 369.9 ± 3.1–1091.7 ± 4.5 µg/mL</td><td>[122]</td></tr> <tr> <td></td><td>Reducing power</td><td>IC₅₀ = 230 ± 5.2–315 ± 3.9 µg/mL</td><td></td></tr> <tr> <td></td><td>ABTS</td><td>IC₅₀ = 33.6 ± 0.3–95.5 ± 2.2 µg/mL</td><td></td></tr> <tr> <td><i>O. vulgare</i> (aerial parts)</td><td>ABTS</td><td>IC₅₀ = 14,00257 mg/mL</td><td>[123]</td></tr> <tr> <td><i>O. vulgare</i> (flowers)</td><td>DPPH</td><td>EC₅₀ = 0.68 mL/mL</td><td>[124]</td></tr> <tr> <td><i>O. vulgare</i> (stems)</td><td>DPPH</td><td>EC₅₀ = 1.82 mL/mL</td><td>[124]</td></tr> <tr> <td><i>R. officinalis</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 523.41 ± 8.25 µg/mL</td><td>[116]</td></tr> <tr> <td></td><td>FRAP</td><td>IC₅₀ = 85.74 ± 7.57 µg/mL</td><td></td></tr> <tr> <td><i>R. officinalis</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 10.08 ± 0.15 µg/mL</td><td>[125]</td></tr> <tr> <td></td><td>TBARS</td><td>IC₅₀ = 1.76 ± 0.02 µg/mL</td><td></td></tr> <tr> <td><i>R. officinalis</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 13.00 ± 0.51 µg/mL</td><td>[126]</td></tr> <tr> <td><i>Satureja hortensis</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 13.45 ± 0.35 µg/mL</td><td>[127]</td></tr> <tr> <td><i>Scutellaria immaculata</i> (aerial parts)</td><td>DPPH</td><td>IC₅₀ = 82.8 ± 3.1 µg/mL</td><td>[128]</td></tr> </tbody> </table>	Species and plant part	Technique	Results	Reference	<i>Cedronella canariensis</i> (aerial parts)	DPPH	IC ₅₀ = 615.5 ± 76.5 µg/mL	[114]		ABTS	IC ₅₀ = 10.5 ± 0.6 µg/mL			FRAP	IC ₅₀ = 3.8 ± 1.46 µmol TE/g		<i>Mentha piperita</i> (leaves)	DPPH	Radical scavenging = 92.6 ± 6.86%	[115]		Reducing power	Reducing power = 0.9 ± 0.3		<i>M. pulegium</i> (aerial parts)	DPPH	IC ₅₀ = 321.41 ± 2.53 µg/mL	[116]		FRAP	IC ₅₀ = 58.27 ± 2.72 µg/mL		<i>M. rotundifolia</i> (leaves)	DPPH	IC ₅₀ = 2222.2 ± 25.2 µg/mL	[117]		ABTS	IC ₅₀ = 133.8 ± 4.8 µg/mL			Reducing power	IC ₅₀ = 166.6 ± 1.9 µg/mL			Phosphomolybdate	IC ₅₀ = 45.2 ± 1.2 µg/mL		<i>M. spicata</i> (aerial parts)	DPPH	IC ₅₀ = 3450 ± 172.5 µg/mL	[118]		ABTS	IC ₅₀ = 40.2 ± 0.2 µg/mL			FRAP	IC ₅₀ = 215 ± 4.50 µg/mL		<i>M. spicata</i> (leaves)	DPPH	IC ₅₀ = 41.23 µg/mL	[119]	<i>O. basilicum</i> (aerial parts)	DPPH	IC ₅₀ = 4.04 ± 0.09–0.21 ± 0.02 mg/mL	[120]		β-carotene	Bleaching = 23.8 ± 0.6–85.3 ± 1.0%		<i>Origanum dictamnus</i> (flowers)	DPPH	IC ₅₀ = 0.0459 ± 0.0042% (v/v)	[121]	<i>O. floribundum</i> (aerial parts)	DPPH	IC ₅₀ = 369.9 ± 3.1–1091.7 ± 4.5 µg/mL	[122]		Reducing power	IC ₅₀ = 230 ± 5.2–315 ± 3.9 µg/mL			ABTS	IC ₅₀ = 33.6 ± 0.3–95.5 ± 2.2 µg/mL		<i>O. vulgare</i> (aerial parts)	ABTS	IC ₅₀ = 14,00257 mg/mL	[123]	<i>O. vulgare</i> (flowers)	DPPH	EC ₅₀ = 0.68 mL/mL	[124]	<i>O. vulgare</i> (stems)	DPPH	EC ₅₀ = 1.82 mL/mL	[124]	<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 523.41 ± 8.25 µg/mL	[116]		FRAP	IC ₅₀ = 85.74 ± 7.57 µg/mL		<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 10.08 ± 0.15 µg/mL	[125]		TBARS	IC ₅₀ = 1.76 ± 0.02 µg/mL		<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 13.00 ± 0.51 µg/mL	[126]	<i>Satureja hortensis</i> (aerial parts)	DPPH	IC ₅₀ = 13.45 ± 0.35 µg/mL	[127]	<i>Scutellaria immaculata</i> (aerial parts)	DPPH	IC ₅₀ = 82.8 ± 3.1 µg/mL	[128]	<p>Table 2 Antioxidant activity of essential oils from <i>Lamiaceae</i> species.</p>	<p>Table 2 Antioxidant activity of essential oils from <i>Lamiaceae</i> species.</p>																																																
Species and plant part	Technique	Results	Reference																																																																																																																																																																																		
<i>Cedronella canariensis</i> (aerial parts)	DPPH	IC ₅₀ = 615.5 ± 76.5 µg/mL	[114]																																																																																																																																																																																		
	ABTS	IC ₅₀ = 10.5 ± 0.6 µg/mL																																																																																																																																																																																			
	FRAP	IC ₅₀ = 3.8 ± 1.46 µmol TE/g																																																																																																																																																																																			
<i>Mentha piperita</i> (leaves)	DPPH	Radical scavenging = 92.6 ± 6.86%	[115]																																																																																																																																																																																		
	Reducing power	Reducing power = 0.9 ± 0.3																																																																																																																																																																																			
<i>M. pulegium</i> (aerial parts)	DPPH	IC ₅₀ = 321.41 ± 2.53 µg/mL	[116]																																																																																																																																																																																		
	FRAP	IC ₅₀ = 58.27 ± 2.72 µg/mL																																																																																																																																																																																			
<i>M. rotundifolia</i> (leaves)	DPPH	IC ₅₀ = 2222.2 ± 25.2 µg/mL	[117]																																																																																																																																																																																		
	ABTS	IC ₅₀ = 133.8 ± 4.8 µg/mL																																																																																																																																																																																			
	Reducing power	IC ₅₀ = 166.6 ± 1.9 µg/mL																																																																																																																																																																																			
	Phosphomolybdate	IC ₅₀ = 45.2 ± 1.2 µg/mL																																																																																																																																																																																			
<i>M. spicata</i> (aerial parts)	DPPH	IC ₅₀ = 3450 ± 172.5 µg/mL	[118]																																																																																																																																																																																		
	ABTS	IC ₅₀ = 40.2 ± 0.2 µg/mL																																																																																																																																																																																			
	FRAP	IC ₅₀ = 215 ± 4.50 µg/mL																																																																																																																																																																																			
<i>M. spicata</i> (leaves)	DPPH	IC ₅₀ = 41.23 µg/mL	[119]																																																																																																																																																																																		
<i>O. basilicum</i> (aerial parts)	DPPH	IC ₅₀ = 4.04 ± 0.09–0.21 ± 0.02 mg/mL	[120]																																																																																																																																																																																		
	β-carotene	Bleaching = 23.8 ± 0.6–85.3 ± 1.0%																																																																																																																																																																																			
<i>Origanum dictamnus</i> (flowers)	DPPH	IC ₅₀ = 0.0459 ± 0.0042% (v/v)	[121]																																																																																																																																																																																		
<i>O. floribundum</i> (aerial parts)	DPPH	IC ₅₀ = 369.9 ± 3.1–1091.7 ± 4.5 µg/mL	[122]																																																																																																																																																																																		
	Reducing power	IC ₅₀ = 230 ± 5.2–315 ± 3.9 µg/mL																																																																																																																																																																																			
	ABTS	IC ₅₀ = 33.6 ± 0.3–95.5 ± 2.2 µg/mL																																																																																																																																																																																			
<i>O. vulgare</i> (aerial parts)	ABTS	IC ₅₀ = 14,00257 mg/mL	[123]																																																																																																																																																																																		
<i>O. vulgare</i> (flowers)	DPPH	EC ₅₀ = 0.68 mL/mL	[124]																																																																																																																																																																																		
<i>O. vulgare</i> (stems)	DPPH	EC ₅₀ = 1.82 mL/mL	[124]																																																																																																																																																																																		
<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 523.41 ± 8.25 µg/mL	[116]																																																																																																																																																																																		
	FRAP	IC ₅₀ = 85.74 ± 7.57 µg/mL																																																																																																																																																																																			
<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 10.08 ± 0.15 µg/mL	[125]																																																																																																																																																																																		
	TBARS	IC ₅₀ = 1.76 ± 0.02 µg/mL																																																																																																																																																																																			
<i>R. officinalis</i> (aerial parts)	DPPH	IC ₅₀ = 13.00 ± 0.51 µg/mL	[126]																																																																																																																																																																																		
<i>Satureja hortensis</i> (aerial parts)	DPPH	IC ₅₀ = 13.45 ± 0.35 µg/mL	[127]																																																																																																																																																																																		
<i>Scutellaria immaculata</i> (aerial parts)	DPPH	IC ₅₀ = 82.8 ± 3.1 µg/mL	[128]																																																																																																																																																																																		
https://europepmc.org/article/MED/34950215#free-full-text	Lamiaceae Essential Oils, Phytochemical Profile, Antioxidant, and Biological Activities.	https://europepmc.org/articles/PMC8692021/table/tabc3/	<p>Table 3 Antibacterial activity of <i>Lamiaceae</i> essential oils.</p> <table border="1"> <thead> <tr> <th>Species</th><th>Bacteria</th><th>Method applied</th><th>Results</th><th>Reference</th></tr> </thead> <tbody> <tr> <td><i>Mentha spicata</i></td><td><i>E. coli</i></td><td>Disc-diffusion</td><td>11.8–21 mm</td><td>[119]</td></tr> <tr> <td></td><td><i>S. enterica</i></td><td></td><td>8–18 mm</td><td></td></tr> <tr> <td></td><td><i>P. aeruginosa</i></td><td></td><td>10–16 mm</td><td></td></tr> <tr> <td></td><td><i>S. aureus</i></td><td></td><td>8–13 mm</td><td></td></tr> <tr> <td></td><td><i>S. epidermidis</i></td><td></td><td>10.1–11.2 mm</td><td></td></tr> <tr> <td></td><td><i>B. subtilis</i></td><td></td><td>9–11.5 mm</td><td></td></tr> <tr> <td><i>Melissa officinalis</i></td><td><i>P. aeruginosa</i></td><td>Agar-disc-diffusion</td><td>16.0 ± 1.2 mm</td><td>[141]</td></tr> <tr> <td></td><td><i>K. pneumonia</i></td><td></td><td>3.0 ± 0.6 mm</td><td></td></tr> <tr> <td></td><td><i>S. aureus</i></td><td></td><td>20.0 ± 1.6 mm</td><td></td></tr> <tr> <td></td><td><i>C. koseri</i></td><td></td><td>14.0 ± 1.0 mm</td><td></td></tr> <tr> <td><i>Origanum compactum</i></td><td><i>E. coli</i> K12</td><td>Microdilution</td><td>29.00 ± 0.35 mm</td><td>[140]</td></tr> <tr> <td></td><td><i>L. innocua</i> 4030</td><td></td><td>49.00 ± 1.00 mm</td><td></td></tr> <tr> <td></td><td><i>S. aureus</i> 25.923</td><td></td><td>43.00 ± 0.35 mm</td><td></td></tr> <tr> <td><i>O. vulgare</i></td><td><i>M. luteus</i></td><td>Microdilution</td><td>270 mg/mL</td><td>[142]</td></tr> <tr> <td></td><td><i>S. aureus</i></td><td></td><td>263 mg/mL</td><td></td></tr> <tr> <td></td><td><i>E. coli</i></td><td></td><td>214 mg/mL</td><td></td></tr> <tr> <td></td><td><i>P. aeruginosa</i></td><td></td><td>383 mg/mL</td><td></td></tr> <tr> <td><i>Salvia ringens</i></td><td><i>E. coli</i></td><td>Microdilution</td><td>14.25</td><td>[144]</td></tr> <tr> <td></td><td><i>S. typhimurium</i></td><td></td><td>14.25</td><td></td></tr> <tr> <td></td><td><i>S. enteritidis</i></td><td></td><td>11.40</td><td></td></tr> <tr> <td></td><td><i>P. tolaasii</i></td><td></td><td>14.25</td><td></td></tr> <tr> <td></td><td><i>P. aeruginosa</i></td><td></td><td>17.10</td><td></td></tr> <tr> <td></td><td><i>P. mirabilis</i></td><td></td><td>17.10</td><td></td></tr> <tr> <td></td><td><i>S. aureus</i></td><td></td><td>9.50</td><td></td></tr> <tr> <td></td><td><i>B. cereus</i></td><td></td><td>9.50</td><td></td></tr> <tr> <td></td><td><i>M. flavus</i></td><td></td><td>9.50</td><td></td></tr> <tr> <td></td><td><i>S. lutea</i></td><td></td><td>11.40</td><td></td></tr> <tr> <td></td><td><i>L. monocytogenes</i></td><td></td><td>9.50</td><td></td></tr> <tr> <td><i>Teucrium africanum</i></td><td><i>S. pyogenes</i> (ATCC)</td><td>Microdilution</td><td>0.16 mg/mL</td><td>[139]</td></tr> <tr> <td><i>T. trifidum</i></td><td><i>S. aureus</i></td><td>Microdilution</td><td>2 mg/mL</td><td></td></tr> <tr> <td><i>Thymus pulegioides</i></td><td>.</td><td>Turbidity measurements</td><td>0.5 mg/mL</td><td>[143]</td></tr> <tr> <td><i>T. serpyllum</i></td><td><i>S. mutans</i></td><td>Turbidity measurements</td><td>0.9 mg/mL</td><td></td></tr> <tr> <td></td><td></td><td>CFU</td><td>1.750.000 bacterial/mL</td><td></td></tr> <tr> <td><i>T. vulgaris</i></td><td><i>S. mutans</i></td><td>Turbidity measurements</td><td>0.75 mg/mL</td><td></td></tr> </tbody> </table>	Species	Bacteria	Method applied	Results	Reference	<i>Mentha spicata</i>	<i>E. coli</i>	Disc-diffusion	11.8–21 mm	[119]		<i>S. enterica</i>		8–18 mm			<i>P. aeruginosa</i>		10–16 mm			<i>S. aureus</i>		8–13 mm			<i>S. epidermidis</i>		10.1–11.2 mm			<i>B. subtilis</i>		9–11.5 mm		<i>Melissa officinalis</i>	<i>P. aeruginosa</i>	Agar-disc-diffusion	16.0 ± 1.2 mm	[141]		<i>K. pneumonia</i>		3.0 ± 0.6 mm			<i>S. aureus</i>		20.0 ± 1.6 mm			<i>C. koseri</i>		14.0 ± 1.0 mm		<i>Origanum compactum</i>	<i>E. coli</i> K12	Microdilution	29.00 ± 0.35 mm	[140]		<i>L. innocua</i> 4030		49.00 ± 1.00 mm			<i>S. aureus</i> 25.923		43.00 ± 0.35 mm		<i>O. vulgare</i>	<i>M. luteus</i>	Microdilution	270 mg/mL	[142]		<i>S. aureus</i>		263 mg/mL			<i>E. coli</i>		214 mg/mL			<i>P. aeruginosa</i>		383 mg/mL		<i>Salvia ringens</i>	<i>E. coli</i>	Microdilution	14.25	[144]		<i>S. typhimurium</i>		14.25			<i>S. enteritidis</i>		11.40			<i>P. tolaasii</i>		14.25			<i>P. aeruginosa</i>		17.10			<i>P. mirabilis</i>		17.10			<i>S. aureus</i>		9.50			<i>B. cereus</i>		9.50			<i>M. flavus</i>		9.50			<i>S. lutea</i>		11.40			<i>L. monocytogenes</i>		9.50		<i>Teucrium africanum</i>	<i>S. pyogenes</i> (ATCC)	Microdilution	0.16 mg/mL	[139]	<i>T. trifidum</i>	<i>S. aureus</i>	Microdilution	2 mg/mL		<i>Thymus pulegioides</i>	.	Turbidity measurements	0.5 mg/mL	[143]	<i>T. serpyllum</i>	<i>S. mutans</i>	Turbidity measurements	0.9 mg/mL				CFU	1.750.000 bacterial/mL		<i>T. vulgaris</i>	<i>S. mutans</i>	Turbidity measurements	0.75 mg/mL		<p>Table 3 Antibacterial activity of <i>Lamiaceae</i> essential oils.</p>	<p>Table 3 Antibacterial activity of <i>Lamiaceae</i> essential oils.</p>	
Species	Bacteria	Method applied	Results	Reference																																																																																																																																																																																	
<i>Mentha spicata</i>	<i>E. coli</i>	Disc-diffusion	11.8–21 mm	[119]																																																																																																																																																																																	
	<i>S. enterica</i>		8–18 mm																																																																																																																																																																																		
	<i>P. aeruginosa</i>		10–16 mm																																																																																																																																																																																		
	<i>S. aureus</i>		8–13 mm																																																																																																																																																																																		
	<i>S. epidermidis</i>		10.1–11.2 mm																																																																																																																																																																																		
	<i>B. subtilis</i>		9–11.5 mm																																																																																																																																																																																		
<i>Melissa officinalis</i>	<i>P. aeruginosa</i>	Agar-disc-diffusion	16.0 ± 1.2 mm	[141]																																																																																																																																																																																	
	<i>K. pneumonia</i>		3.0 ± 0.6 mm																																																																																																																																																																																		
	<i>S. aureus</i>		20.0 ± 1.6 mm																																																																																																																																																																																		
	<i>C. koseri</i>		14.0 ± 1.0 mm																																																																																																																																																																																		
<i>Origanum compactum</i>	<i>E. coli</i> K12	Microdilution	29.00 ± 0.35 mm	[140]																																																																																																																																																																																	
	<i>L. innocua</i> 4030		49.00 ± 1.00 mm																																																																																																																																																																																		
	<i>S. aureus</i> 25.923		43.00 ± 0.35 mm																																																																																																																																																																																		
<i>O. vulgare</i>	<i>M. luteus</i>	Microdilution	270 mg/mL	[142]																																																																																																																																																																																	
	<i>S. aureus</i>		263 mg/mL																																																																																																																																																																																		
	<i>E. coli</i>		214 mg/mL																																																																																																																																																																																		
	<i>P. aeruginosa</i>		383 mg/mL																																																																																																																																																																																		
<i>Salvia ringens</i>	<i>E. coli</i>	Microdilution	14.25	[144]																																																																																																																																																																																	
	<i>S. typhimurium</i>		14.25																																																																																																																																																																																		
	<i>S. enteritidis</i>		11.40																																																																																																																																																																																		
	<i>P. tolaasii</i>		14.25																																																																																																																																																																																		
	<i>P. aeruginosa</i>		17.10																																																																																																																																																																																		
	<i>P. mirabilis</i>		17.10																																																																																																																																																																																		
	<i>S. aureus</i>		9.50																																																																																																																																																																																		
	<i>B. cereus</i>		9.50																																																																																																																																																																																		
	<i>M. flavus</i>		9.50																																																																																																																																																																																		
	<i>S. lutea</i>		11.40																																																																																																																																																																																		
	<i>L. monocytogenes</i>		9.50																																																																																																																																																																																		
<i>Teucrium africanum</i>	<i>S. pyogenes</i> (ATCC)	Microdilution	0.16 mg/mL	[139]																																																																																																																																																																																	
<i>T. trifidum</i>	<i>S. aureus</i>	Microdilution	2 mg/mL																																																																																																																																																																																		
<i>Thymus pulegioides</i>	.	Turbidity measurements	0.5 mg/mL	[143]																																																																																																																																																																																	
<i>T. serpyllum</i>	<i>S. mutans</i>	Turbidity measurements	0.9 mg/mL																																																																																																																																																																																		
		CFU	1.750.000 bacterial/mL																																																																																																																																																																																		
<i>T. vulgaris</i>	<i>S. mutans</i>	Turbidity measurements	0.75 mg/mL																																																																																																																																																																																		
https://europepmc.org/article/MED/34950215#free-full-text	Lamiaceae Essential Oils, Phytochemical Profile, Antioxidant, and Biological Activities.	https://europepmc.org/articles/PMC8692021/table/tabc4/	<p>Table 4 Antifungal activity of <i>Lamiaceae</i> essential oils.</p> <table border="1"> <thead> <tr> <th>Species</th><th>Fungi</th><th>Method applied</th><th>Results</th><th>Reference</th></tr> </thead> <tbody> <tr> <td><i>Lepechinia mutica</i></td><td><i>C. albicans</i></td><td>Broth microdilution</td><td>MIC > 9 mg/mL</td><td>[148]</td></tr> <tr> <td></td><td><i>M. canis</i></td><td></td><td>2.2 < MIC ≤ 4.5 mg/mL</td><td></td></tr> <tr> <td></td><td><i>T. rubrum</i></td><td></td><td>2.2 < MIC ≤ 4.5 mg/mL</td><td></td></tr> <tr> <td></td><td><i>F. graminearum</i></td><td></td><td>MIC > 9 mg/mL</td><td></td></tr> <tr> <td></td><td><i>P. oryzae</i></td><td></td><td>MIC > 9 mg/mL</td><td></td></tr> <tr> <td><i>O. basilicum</i></td><td><i>A. flavus</i></td><td>Potato dextrose agar (PDA)</td><td>500 ppm: 30% 750 pp : 50% 1000 ppm: 70%</td><td>[150]</td></tr> <tr> <td></td><td></td><td></td><td></td><td></td></tr> <tr> <td><i>O. basilicum</i></td><td><i>C. albicans</i></td><td>Sabouraud dextrose agar (SDA)</td><td>MIC: 1.25 µL/mL MLC: 2.5 µL/mL</td><td>[151]</td></tr> <tr> <td></td><td><i>C. tropicalis</i></td><td></td><td>MIC: 2.5–1.25 µL/mL MLC: 2.5 µL/mL</td><td></td></tr> <tr> <td></td><td><i>C. krusei</i></td><td></td><td>MIC: 1.25 µL/mL MLC: 2.5 µL/mL</td><td></td></tr> <tr> <td></td><td><i>C. guilliermondii</i></td><td></td><td>MIC: 1.25 µL/mL MLC: 1.25 µL/mL</td><td></td></tr> <tr> <td></td><td><i>C. parapsilosis</i></td><td></td><td>MIC: 1.25 µL/mL MLC: 2.5 µL/mL</td><td></td></tr> <tr> <td></td><td><i>C. neoformans</i></td><td></td><td>MIC: 0.16–0.32 µL/mL MLC: 0.64–0.32 µL/mL</td><td></td></tr> <tr> <td></td><td><i>T. mentagrophytes</i></td><td></td><td>MIC: 0.64 µL/mL MLC: 1.25</td><td></td></tr> </tbody> </table>	Species	Fungi	Method applied	Results	Reference	<i>Lepechinia mutica</i>	<i>C. albicans</i>	Broth microdilution	MIC > 9 mg/mL	[148]		<i>M. canis</i>		2.2 < MIC ≤ 4.5 mg/mL			<i>T. rubrum</i>		2.2 < MIC ≤ 4.5 mg/mL			<i>F. graminearum</i>		MIC > 9 mg/mL			<i>P. oryzae</i>		MIC > 9 mg/mL		<i>O. basilicum</i>	<i>A. flavus</i>	Potato dextrose agar (PDA)	500 ppm: 30% 750 pp : 50% 1000 ppm: 70%	[150]						<i>O. basilicum</i>	<i>C. albicans</i>	Sabouraud dextrose agar (SDA)	MIC: 1.25 µL/mL MLC: 2.5 µL/mL	[151]		<i>C. tropicalis</i>		MIC: 2.5–1.25 µL/mL MLC: 2.5 µL/mL			<i>C. krusei</i>		MIC: 1.25 µL/mL MLC: 2.5 µL/mL			<i>C. guilliermondii</i>		MIC: 1.25 µL/mL MLC: 1.25 µL/mL			<i>C. parapsilosis</i>		MIC: 1.25 µL/mL MLC: 2.5 µL/mL			<i>C. neoformans</i>		MIC: 0.16–0.32 µL/mL MLC: 0.64–0.32 µL/mL			<i>T. mentagrophytes</i>		MIC: 0.64 µL/mL MLC: 1.25		<p>Table 4 Antifungal activity of <i>Lamiaceae</i> essential oils.</p>	<p>Table 4 Antifungal activity of <i>Lamiaceae</i> essential oils.</p>																																																																																																					
Species	Fungi	Method applied	Results	Reference																																																																																																																																																																																	
<i>Lepechinia mutica</i>	<i>C. albicans</i>	Broth microdilution	MIC > 9 mg/mL	[148]																																																																																																																																																																																	
	<i>M. canis</i>		2.2 < MIC ≤ 4.5 mg/mL																																																																																																																																																																																		
	<i>T. rubrum</i>		2.2 < MIC ≤ 4.5 mg/mL																																																																																																																																																																																		
	<i>F. graminearum</i>		MIC > 9 mg/mL																																																																																																																																																																																		
	<i>P. oryzae</i>		MIC > 9 mg/mL																																																																																																																																																																																		
<i>O. basilicum</i>	<i>A. flavus</i>	Potato dextrose agar (PDA)	500 ppm: 30% 750 pp : 50% 1000 ppm: 70%	[150]																																																																																																																																																																																	
<i>O. basilicum</i>	<i>C. albicans</i>	Sabouraud dextrose agar (SDA)	MIC: 1.25 µL/mL MLC: 2.5 µL/mL	[151]																																																																																																																																																																																	
	<i>C. tropicalis</i>		MIC: 2.5–1.25 µL/mL MLC: 2.5 µL/mL																																																																																																																																																																																		
	<i>C. krusei</i>		MIC: 1.25 µL/mL MLC: 2.5 µL/mL																																																																																																																																																																																		
	<i>C. guilliermondii</i>		MIC: 1.25 µL/mL MLC: 1.25 µL/mL																																																																																																																																																																																		
	<i>C. parapsilosis</i>		MIC: 1.25 µL/mL MLC: 2.5 µL/mL																																																																																																																																																																																		
	<i>C. neoformans</i>		MIC: 0.16–0.32 µL/mL MLC: 0.64–0.32 µL/mL																																																																																																																																																																																		
	<i>T. mentagrophytes</i>		MIC: 0.64 µL/mL MLC: 1.25																																																																																																																																																																																		