

Methods in Psycholinguistics

— Mixed effects linear regression —

05/04/2017

Judith Degen
Stanford University

Generalized Linear Mixed Models


Experiments typically have more than one **participant** and more than one **item**

—> violation of the assumption of independence!


We would like to take these idiosyncrasies into account, and maybe even investigate them.

Solution: Generalized Linear Mixed or Multilevel Models (a.k.a. hierarchical, mixed-effects)


Recall: Generalized Linear Model



Generalized Linear Mixed Model




Generalized Linear Mixed Model



Cluster-specific parameters
("random effects")

Shared parameters
("fixed effects")

Generalized Linear Mixed Model



Mixed linear model

...back to our lexical decision experiment.

A number of predictors seem to affect RTs:

- frequency
- family size
- native language
- interactions

Additionally, different participants may have

- different overall decision speeds
- differing sensitivity to predictors

We want to draw inferences about all of this at the same time!

Mixed linear model

Let's allow each participant i to have idiosyncratic differences in response times:

$$RT_{ij} = \alpha + \beta x_{ij} + \overset{\sim N(0, \sigma_b)}{\underbrace{b_i}} + \epsilon_{ij}$$

Mixed linear model

Let's allow each participant i to have idiosyncratic differences in response times:

$$RT_{ij} = \alpha + \beta x_{ij} + \overbrace{b_i}^{\sim N(0, \sigma_b)} + \epsilon_{ij}$$

- Idea: model distribution of participant differences as deviation from grand mean
- Mixed models approximate deviation by fitting a normal distribution
 - with mean 0
 - only additional parameter to fit is variance

Mixed linear model

Let's allow each participant i to have idiosyncratic differences in response times:

$$RT_{ij} = \alpha + \beta x_{ij} + \overset{\sim N(0, \sigma_b)}{\underbrace{b_i}} + \epsilon_{ij}$$

- Idea: model distribution of participant differences as deviation from grand mean
- Mixed models approximate deviation by fitting a normal distribution
 - with mean 0
 - only additional parameter to fit is variance

Let's translate this into R!

Interpretation of the output

Random
effects
estimates



Random effects:

Groups	Name	Variance	Std.Dev.
Subject	(Intercept)	0.02373	0.1541
Residual		0.03274	0.1809

Number of obs: 1659, groups: Subject, 21

Fixed
effects
estimates
(same)



Fixed effects:

	Estimate	Std. Error	t value
(Intercept)	6.588778	0.037737	174.6
Frequency	-0.042872	0.003485	-12.3

2 notable differences:

1. random effects estimates
2. no more p-values!

Mixed models with one intercept

Let's look at the by-subject adjustments to the intercept:
Best Unbiased Linear Predictors (BLUPs)

BLUPs are *not* fitted parameters. Only one degree of freedom added to the model. BLUPs are estimated a posteriori based on the fitted model.


$$P(b_i | \hat{\alpha}, \hat{\beta}, \hat{\sigma}_b, \hat{\sigma}_\epsilon, X, Y)$$

The BLUPs are chosen to maximize this probability.


The effect

Predicted by-subject means

Linear model




Mixed linear model




The effect

Predicted RTs


Linear model




Mixed linear model



A second random intercept



A second random intercept



Mixed models with random slopes

Not just the intercept, but **any of the slopes** (of the predictors) may differ between individuals.

For example, subjects may show different sensitivity to frequency:

```
> head(ranef(m)$Subject)
      (Intercept)  Frequency
A1 -0.113081677  0.002001991
A2 -0.237501826  0.015898141
A3 -0.005238541  0.003482887
C  -0.132056269  0.014382981
D   0.001134412  0.003810049
I  -0.141643431  0.002989325
```


The BLUPs of the random slope reflect the by-subject adjustments to the overall frequency effect.

Random intercept, slope, and covariance

Random effects (e.g., intercepts and slopes) may be correlated.

By default, R fits these covariances, introducing additional degrees of freedom (parameters).

What do correlations
between random
effects mean?



Where did our p-values go?

- t-value anti-conservative, so no p-values!
- there are many different options for computing p-values if you really need them. You can check these in R directly by saying `?pvalues`
- some reasonably easy to use ones for **fixed effects**:
 - likelihood ratio tests via `anova`
 - `lmerTest` package (uses Satterthwaite or Kenward-Roger approximations)