# Open Science and Preregistration

Sebastian Schuster

LINGUIST 245 May 3, 2018

- 1. Number of subjects per condition
  - a. Run 10 subjects per condition
  - b. Perform a t-test
  - c. If p < .05: Publish paper!
    - Otherwise: Go to step a.

- 2. Have multiple dependent variables
  - a. Run tests to predict each of the variables
  - b. Pick the dependent variable that gives you a significant p-value

- 3. Run models with many different independent variables
  - a. Have a set of many independent variables
  - b. Run models with various combinations and interactions until your manipulation is significant

- 4. Have conditions that you don't report on
  - a. Run n > 2 conditions
  - b. Pick 2 conditions which differ significantly and don't tell anybody about the other conditions

#### **DON'T DO ANY OF THESE THINGS!!!**


Researcher degrees of freedom	Significance level		
	p < .1	p < .05	p < .01
Situation A: two dependent variables $(r = .50)$	17.8%	9.5%	2.2%

	Significance level		
Researcher degrees of freedom	p < .1	p < .05	p < .01
Situation A: two dependent variables $(r = .50)$	17.8%	9.5%	2.2%
Situation B: addition of 10 more observations per cell	14.5%	7.7%	1.6%

Researcher degrees of freedom	Significance level		
	p < .1	p < .05	p < .01
Situation A: two dependent variables (r = .50)	17.8%	9.5%	2.2%
Situation B: addition of 10 more observations per cell	14.5%	7.7%	1.6%
Situation C: controlling for gender or interaction of gender with treatment	21.6%	11.7%	2.7%

Researcher degrees of freedom	Significance level		
	p < .1	p < .05	p < .01
Situation A: two dependent variables $(r = .50)$	17.8%	9.5%	2.2%
Situation B: addition of 10 more observations per cell	14.5%	7.7%	1.6%
Situation C: controlling for gender or interaction of gender with treatment	21.6%	11.7%	2.7%
Situation D: dropping (or not dropping) one of three conditions	23.2%	12.6%	2.8%

Researcher degrees of freedom	Significance level		
	p < .1	p < .05	10. > q
Situation A: two dependent variables $(r = .50)$	17.8%	9.5%	2.2%
Situation B: addition of 10 more observations per cell	14.5%	7.7%	1.6%
Situation C: controlling for gender or interaction of gender with treatment	21.6%	11.7%	2.7%
Situation D: dropping (or not dropping) one of three conditions	23.2%	12.6%	2.8%
Combine Situations A and B	26.0%	14.4%	3.3%
Combine Situations A, B, and C	50. <del>9</del> %	30.9%	8.4%
Combine Situations A, B, C, and D	81.5%	60.7%	21.5%




#### Pre-registration

- To keep p-value meaningful, fix the following things before collecting data
  - 1. number of subjects you'll run
  - exclusion criteria:
    Which data points are you going to exclude from your analysis
  - 3. dependent variable
  - 4. independent variables
  - 5. experimental conditions

#### Preregistration

- Preregistering provides you (and reviewers and readers of your paper) with proof that you actually fixed all these things
- Just requires filling out a short questionnaire which is permanently stored on a pre-registration platform



### My current workflow

- 1. Come up with and implement experiment
- 2. Run pilot study with 2-4 subjects
- 3. Write analysis scripts and test them with pilot data
- 4. Preregister study and upload analysis scripts to OSF
- 5. Run actual study
- 6. Analyze data with pre-registered analysis script
- 7. (optional) Do **exploratory** post-hoc analyses

# What you'll do today

- 1. Come up with and implement experiment
- 2. Run pilot study with 2-4 subjects
- 3. Write analysis scripts and test them with pilot data
- 4. Preregister study and upload analysis scripts to OSF
- 5. Run actual study
- 6. Analyze data with pre-registered analysis script
- 7. (optional) Do **exploratory** post-hoc analyses

### **Tutorial**

# www.osf.io