

Counting

Chapter 6

Edited by: Dr. Meshal Alfarhood

Permutations and Combinations

Section 6.3

Section Summary

Permutations (without Repetition)

Combinations (without Repetition)

Permutations

Definition: A *permutation* of a set of distinct objects is an ordered arrangement of these objects.

- An ordered arrangement of r elements of a set is called an r -permutation.
- $P(n, r) = n(n - 1)(n - 2) \cdots (n - r + 1)$
- $P(n, r) = nPr$

$$P(n, r) = \frac{n!}{(n - r)!}$$

Example: How many ways can we select three students from a group of five students to stand in line for a picture?

- #ways = $5 \times 4 \times 3 = 60$.
- Using the formula: $P(5, 3) = \frac{5!}{(5-3)!} = \frac{5!}{2!} = \frac{5 \times 4 \times 3 \times 2 \times 1}{2 \times 1} = 5 \times 4 \times 3 = 60$.

Counting Permutations

Examples:

1. How many ways are there to select a first-prize winner, a second prize winner, and a third-prize winner from 100 different people who have entered a contest?
 - $P(100,3) = 100 \cdot 99 \cdot 98 = 970,200$
2. Suppose that a saleswoman has to visit eight different cities. She must begin her trip in a specified city, but she can visit the other seven cities in any order she wishes. How many possible orders can the saleswoman use when visiting these cities?
 - The first city is chosen. Thus we have $P(7,7) = 7! = 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5040$ paths.
3. How many permutations of the letters **ABCDEFGH** contain the string **ABC** ?
 - We solve this problem by counting the permutations of six objects, ABC, D, E, F, G, and H. Thus, we have $P(6,6) = 6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 720$

Combinations

Definition: An ***r-combination*** of elements of a set is an unordered selection of r elements from the set.

$$C(n, r) = nCr$$

$$C(n, r) = \frac{n!}{(n-r)!r!}.$$

$C(n, r)$ is also denoted by $\binom{n}{r}$

Example: Let S be the set $\{a, b, c, d\}$.

- The 3-combination of S is all 3-element subsets of S . It is the same as $C(4, 3) = 4 \rightarrow \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}$.
- The 2-combination of S is all 2-element subsets of S . It is the same as $C(4, 2) = 6 \rightarrow \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\}$.

Counting Combinations

Examples:

1. How many ways are there to select five players from a 10-member tennis team to make a trip to a match at another school.

$$C(10,5) = \frac{10!}{5!5!} = 252.$$

2. How many different committees of three students can be formed from a group of four students?

$$C(4,3) = \frac{4!}{1!3!} = 4.$$

3. Suppose that there are 9 faculty members in the mathematics department and 11 in the computer science department. How many ways are there to select a committee to develop a discrete mathematics course at a school if the committee is to consist of three faculty members from the mathematics department and four from the computer science department?

$$C(9,3) \cdot C(11,4) = \frac{9!}{3!6!} \cdot \frac{11!}{4!7!} = 84 \cdot 330 = 27,720.$$

Exercise

- Find the value of the following:

1. $P(n, 0) = 1$
2. $P(n, 1) = n$
3. $P(n, n) = n!$
4. $C(n, 0) = 1$
5. $C(n, 1) = n$
6. $C(n, n) = 1$

- True or False:

1. $P(n, r) = C(n, r) \cdot r!$ True
2. $C(n, r) = C(n, n-r)$ True
3. $P(n, n) = 2 \cdot P(n, n-2)$ True

More Examples

Examples:

1. How many bit strings of length 10 that has exactly 3 zeros?

$$C(10,3) = \binom{10}{3}$$

2. How many bit strings of length 10 that has at least 3 zeros?

$$\binom{10}{3} + \binom{10}{4} + \cdots + \binom{10}{10} = \sum_{k=3}^{10} \binom{10}{k}$$

3. Suppose we have 4 books in Arabic, 3 books in Math, and 3 books in CS. How many ways to arrange them?

$$10!$$

4. Suppose we have 4 books in Arabic, 3 books in Math, and 3 books in CS. How many ways to arrange if we want each subject books together?

$$3! (4! \cdot 3! \cdot 3!)$$

Generalized Permutations and Combinations

Section 6.5

Section Summary

Permutations (with Repetition)

Combinations (with Repetition)

Permutations with Indistinguishable Objects

Permutations with Repetition

Definition: The number of r -permutations of a set of n objects with repetition allowed is n^r .

Examples:

1. How many strings of length five can be formed from the uppercase letters of the English alphabet?
 - **Solution:** The number of such strings is 26^5 .
2. From the set of first 10 natural numbers (0-9), you are asked to make a four-digit number (PIN). How many different permutations are possible?
 - **Solution:** The total number of such PINs is 10^4 .

Combinations with Repetition

Definition: The number of r -combinations from a set with n elements when repetition is allowed is:

$$C(n + r - 1, r) = C(n + r - 1, n - 1).$$

Examples:

1. How many ways are there to select five bills from a box containing at least five of each of the following: \$1, \$2, \$5, \$10, \$20, \$50, \$100?

- **Solution:** $[n=7, r=5]$

$$C(11, 5) = \frac{11!}{5!6!} = 462$$

2. Suppose that a cookie shop has four different kinds of cookies. How many different ways can six cookies be chosen?

- **Solution:** The number of ways to choose six cookies is the number of 6-combinations of a set with four elements. $[n=4, r=6]$

$$C(9, 6) = C(9, 3) = \frac{9 \cdot 8 \cdot 7}{1 \cdot 2 \cdot 3} = 84$$

Combinations with Repetition₂

Examples:

1. How many solutions does the equation have, where x_1 , x_2 and x_3 are nonnegative integers? $x_1 + x_2 + x_3 = 11$

- **Solution:** Each solution corresponds to a way to select 11 items from a set with three elements; x_1 items of type one, x_2 items of type two, and x_3 items of type three. [$n=3$, $r=11$]

$$C(3+11-1, 11) = C(13, 11) = C(13, 2) = \frac{13 \cdot 12}{1 \cdot 2} = 78$$

2. How many solutions for the same equation in question 1, where $x_1 \geq 1$, $x_2 \geq 2$ and $x_3 \geq 3$?

- **Solution:** $1+2+3=6 \rightarrow 11-6=5$. Therefore, each solution corresponds to a way to select 5 items from a set with three elements. [$n=3$, $r=5$]

$$C(3+5-1, 5) = C(7, 5) = C(7, 2) = \frac{7 \cdot 6}{1 \cdot 2} = 21$$

Summarizing the Formulas for Counting Permutations and Combinations with and without Repetition

TABLE 1 Combinations and Permutations With and Without Repetition.

<i>Type</i>	<i>Repetition Allowed?</i>	<i>Formula</i>
<i>r</i> -permutations	No	$\frac{n!}{(n-r)!}$
<i>r</i> -combinations	No	$\frac{n!}{r!(n-r)!}$
<i>r</i> -permutations	Yes	n^r
<i>r</i> -combinations	Yes	$\frac{(n+r-1)!}{r!(n-1)!}$

Permutations with Indistinguishable Objects

Theorem: The number of different permutations of n objects, where there are n_1 indistinguishable objects of type 1, n_2 indistinguishable objects of type 2, ..., and n_k indistinguishable objects of type k , is:

$$\frac{n!}{n_1! n_2! \cdots n_k!}.$$

Example:

- How many different strings can be made by reordering the letters of the word **SUCCESS**.
 - $n = 7$ (#letters)
 - # strings = $\frac{7!}{3! 2!} = 420$
- Same as (1), but must start with letter "S".
 - $n = 6$ (#letters)
 - # strings = $\frac{6!}{2! 2!} = 180$