

A Read fate per locus, grouped by indel rate
BICF2S23758182 81% 6%
BICF2S2305942 80% 8%
BICF2P1262794 78% 4%
BICF2P57528 78% 12%
BICF2P363248 75% 15%
BICF2P400484 74% 12%
BICF2P1165732 73% 13%
BICF2G630126293 71% 14%
BICF2S22912279 71% 14%
BICF2P223300 71% 11%
BICF2P239653 68% 20%
BICF2P878735 72% 18%
BICF2P153324 71% 12%
BICF2S23219794 70% 11%
BICF2G630165989 70% 21%
BICF2P866415 68% 13%
BICF2S23626182 67% 19%
BICF2G630320565 ◆ 67% 18%
BICF2P1060759 66% 17%
BICF2S23033793 65% 15%
BICF2P285168 63% 18%
BICF2G630340139 48% 34%
BICF2G630703836 67% 18%
BICF2G630128194 ◆ 67% 19%
BICF2S23225775 ◆ 67% 23%
BICF2S23526960 67% 22%
TIGRP2P247920 66% 22%
BICF2G630409723 64% 25%
BICF2P1138616 64% 22%
BICF2G630580879 62% 25%
BICF2G630276750 62% 26%
BICF2P1448058 55% 31%
TIGRP2P116705 ◆ 43% 21%
0% 25% 50% 75% 100%
Share of reads
0.1 0.2 0.3
Indels per read
B Indels vs reads used
Spearman ρ = -0.80, Holm p < 0.001, n = 33
Reads used
80% 70% 60% 50%
0.10 0.15 0.20 0.25 0.30
Indels per read
C Primer length vs indels
Spearman ρ = 0.76, Holm p < 0.001, n = 33
Indels per read
0.30 0.25 0.20 0.15 0.10
10 15 20 25
Scored primer length (nt)
D Homopolymer at SNP vs ties
Spearman ρ = 0.49, Holm p = 0.029, n = 33
TIGRP2P116705
No allele signal
20% 10% 0%
1 2 3 4 5 6
Homopolymer at SNP (nt)
Used for genotyping
Low-quality SNP base
Flagging filters
No allele signal
Poor alignment
Homopolymer ≥5 nt in primer