



南方科技大学  
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

## Tutorial Questions (Week 6)

Shihang Lu (卢仕航), TA

Southern University of Science and Technology

Email: [lush2021@mail.sustech.edu.cn](mailto:lush2021@mail.sustech.edu.cn)


- Review
- Quiz Problem
- Basic Problems with Answers 3.8
- Basic Problems 3.34
- Advanced Problems 3.40
- Q&A

- Basic knowledge on signal computation
- Exponential signals: Euler's relation, periodic, integral
- CT/DT unit impulse/step function
- System Properties
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability
  5. Time-invariance
  6. Linearity

- CT/DT LTI systems
- Convolution operation procedure
  1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
  2. Some known or **typical convolution results**
  3. **Properties** of convolution
- Unit impulse response and properties of LTI systems
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability

- CT/DT LTI systems
- Convolution operation procedure
  1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
  2. Some known or **typical convolution results**
  3. **Properties** of convolution
- Unit impulse response and properties of LTI systems
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability

# Example 1



$$\triangleright x[n] = a^n u[n]$$

$$\triangleright h[n] = u[n]$$

$$\triangleright y[n] = x[n] * h[n] ?$$


$$y[n] = \begin{cases} \frac{1-a^{n+1}}{1-a}, & n \geq 0 \\ 0, & n < 0 \end{cases}$$

$$h[n-k]$$

$$x[k]h[n-k] = \begin{cases} a^k, & 0 \leq k \leq n \\ 0, & k < 0, k > n \end{cases}$$

$$y[n] = \sum_{k=-\infty}^{\infty} x[k]h[n-k] = \sum_{k=0}^n a^k = \frac{1-a^{n+1}}{1-a}$$

## Example 2



$$x(t) = e^{-at}u(t)$$

$$h(t) = u(t)$$

$$y(t) = x(t) * h(t)$$

$$y(t) = \begin{cases} \frac{1 - e^{-at}}{a}, & t \geq 0 \\ 0, & t < 0 \end{cases} \quad \Rightarrow \quad \frac{1 - e^{-at}}{a} u(t)$$

- CT/DT LTI systems
- Convolution operation procedure
  1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
  2. Some known or **typical convolution results**
  3. **Properties** of convolution
- Unit impulse response and properties of LTI systems
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability



表 3.4 基本信号的卷积表

连续时间卷积积分			离散时间卷积和		
$x(t)$	$h(t)$	$x(t) * h(t)$	$x[n]$	$h[n]$	$x[n] * h[n]$
$x(t)$	$\delta(t)$	$x(t)$	$x[n]$	$\delta[n]$	$x[n]$
$x(t)$	$u(t)$	$\int_{-\infty}^t x(\tau) d\tau$	$x[n]$	$u[n]$	$\sum_{k=-\infty}^n x[k]$
$x(t)$	$\delta'(t)$	$x'(t)$	$x[n]$	$\Delta\delta[n]$	$x[n] - x[n-1]$
$u(t)$	$u(t)$	$tu(t)$	$u[n]$	$u[n]$	$(n+1)u[n]$
$e^{-at}u(t)$	$u(t)$	$\frac{1-e^{-at}}{a}u(t)$	$a^n u[n]$	$u[n]$	$\frac{1-a^{n+1}}{1-a}u[n]$
$\sin(\omega t)u(t)$	$u(t)$	$\frac{1-\cos(\omega t)}{\omega}u(t)$	$\sin(\Omega n)u[n]$	$u[n]$	
$\cos(\omega t)u(t)$	$u(t)$	$\frac{\sin(\omega t)}{\omega}u(t)$	$\cos(\Omega n)u[n]$	$u[n]$	
$e^{-at}u(t)$	$e^{-at}u(t)$	$te^{-at}u(t)$	$a^n u[n]$	$a^n u[n]$	$(n+1)a^n u[n]$
$e^{-at}u(t)$	$e^{-bt}u(t)$	$\frac{e^{-at}-e^{-bt}}{b-a}u(t)$	$a^n u[n]$	$b^n u[n]$	$\frac{b^{n+1}-a^{n+1}}{b-a}u[n]$

说明：表 3.4 中空着的卷积和运算结果，感兴趣的读者可自行补上。

- CT/DT LTI systems
- Convolution operation procedure
  1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
  2. Some known or **typical convolution results**
  3. **Properties** of convolution
- Unit impulse response and properties of LTI systems
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability

## □ Commutative property

$$x(t) * h(t) = h(t) * x(t)$$

$$\int_{-\infty}^{\infty} x(\tau)h(t-\tau)d\tau = \int_{-\infty}^{\infty} h(\tau)x(t-\tau)d\tau$$


$$x[n] * h[n] = h[n] * x[n]$$

$$\sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

## □ Distributive property

$$x(t) * [h_1(t) + h_2(t)] = x(t) * h_1(t) + x(t) * h_2(t)$$

$$x[n] * \{h_1[n] + h_2[n]\} = x[n] * h_1[n] + x[n] * h_2[n]$$



## □ Associative property

$$[x(t) * h_1(t)] * h_2(t) = x(t) * [h_1(t) * h_2(t)]$$

$$\{x[n] * h_1[n]\} * h_2[n] = x[n] * \{h_1[n] * h_2[n]\}$$

## □ Time-invariant property (Collect the time shift)

$$y(t) = x(t) * h(t)$$

$$x(t) * h(t - t_0) = y(t - t_0)$$

$$x(t - t_1) * h(t - t_2) = y(t - t_1 - t_2)$$

$$x[n] * h[n] = y[n]$$

$$x[n] * h[n - m] = y[n - m]$$

$$x[n - m_1] * h[n - m_2] = y[n - m_1 - m_2]$$

## □ Difference property

$$\frac{d}{dt}[x(t) * h(t)] = x(t) * \frac{dh(t)}{dt} = \frac{dx(t)}{dt} * h(t) = \frac{dy(t)}{dt}$$

$$\nabla \{x[n] * h[n]\} = \nabla x[n] * h[n] = x[n] * \nabla h[n] = \nabla y[n]$$

## □ Integral property

$$\int_{-\infty}^t [x(\tau) * h(\tau)] d\tau = x(t) * \int_{-\infty}^t h(\tau) d\tau = \int_{-\infty}^t x(\tau) d\tau * h(t) = \int_{-\infty}^t y(\tau) d\tau$$

$$\sum_{k=-\infty}^n \{x[k] * h[k]\} = x[n] * \left\{ \sum_{k=-\infty}^n h[k] \right\} = \left\{ \sum_{k=-\infty}^n x[k] \right\} * h[n] = \sum_{k=-\infty}^n y[k]$$

□ For unit impulse/step signal

□ More unit impulse/step signals, more simple

$$x(t) * \delta(t) = x(t)$$

$$x(t) * \delta(t - t_0) = x(t - t_0)$$


$$x(t) * u(t) = \int_{-\infty}^t x(\tau) d\tau$$

$$u(t) = \int_{-\infty}^t \delta(\tau) d\tau$$

$$x[n] * \delta[n] = x[n]$$

$$x[n] * \delta[n - m] = x[n - m]$$

$$x[n] * u[n] = \sum_{m=-\infty}^n x[m]$$




$$y(t) = r_T(t) * r_T(t) = \frac{d}{dt} r_T(t) * \int_{-\infty}^t r_T(\tau) d\tau$$

$$h_{-1}(t) = \int_{-\infty}^t r_T(\tau) d\tau$$

$$\begin{aligned} y(t) &= [\delta(t + T/2) - \delta(t - T/2)] * h_{-1}(t) \\ &= h_{-1}(t + T/2) - h_{-1}(t - T/2) \end{aligned}$$

1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
2. Some known or **typical convolution results**
3. **Properties** of convolution





- CT/DT LTI systems
- Convolution operation procedure
  1. **Figure computation** based on “Flip-slide-multiply-sum/integral”
  2. Some known or **typical convolution results**
  3. **Properties** of convolution
- Unit impulse response and properties of LTI systems
  1. Memoryless or with memory
  2. Causality
  3. Invertibility
  4. Stability

System properties:

- With memory or memoryless

$$y(n) = f(x(n))$$

- Invertible

For a system  $x \rightarrow y$ , if  $x_1 \neq x_2$ , then  $y_1 \neq y_2$

- Causal

... up to that time  $n$  ...

- Stable (BIBO)

either prove the system is stable, or find a specific counterexample

## LTI System properties:

- With memory or memoryless
  - A linear, time-invariant, causal system is memoryless only

$$\text{if } h[n] = K\delta[n] \quad h(t) = K\delta(t)$$

$$y[n] = Kx[n] \quad y(t) = Kx(t)$$


if  $K=1$  further, they are identity systems

$$y[n] = x[n] = \sum_{k=-\infty}^{\infty} x[k]\delta[n-k] = x[n] * \delta[n]$$


$$y(t) = x(t) = \int_{-\infty}^{\infty} x(\tau)\delta(t-\tau)d\tau = x(t) * \delta(t)$$

## LTI System properties:

- Invertible



(a)



(b)

## LTI System properties:

- Causal

Causality: CT LTI system is causal  $\Leftrightarrow h(t) = 0$ , at  $t < 0$

- This is because that the input unit impulse function  $\delta(t)=0$  at  $t < 0$

As a result:

$$y(t) = \int_{-\infty}^{+\infty} x(\tau)h(t-\tau)d\tau = \int_{-\infty}^t x(\tau)h(t-\tau)d\tau$$

$t - \tau \geq 0$ , or  $\tau \leq t$

$y(t)$  only depends on  $x(\tau < t)$ .

## LTI System properties:

- Stable (BIBO)

**BIBO** Stability: CT LTI system is stable  $\Leftrightarrow \int_{-\infty}^{+\infty} |h(\tau)| d\tau < \infty$

→ Sufficient condition:

For  $|x(t)| \leq x_{\max} < \infty$ ,

Cauchy-Schwarz Inequation

$$|y(t)| = \left| \int_{-\infty}^{+\infty} x(\tau) h(t - \tau) d\tau \right| \leq x_{\max} \left| \int_{-\infty}^{+\infty} h(t - \tau) d\tau \right| < \infty.$$

→ Necessary condition:


Suppose  $\int_{-\infty}^{+\infty} |h(\tau)| d\tau = \infty$

Contradiction Case

Let  $x(t) = h^*(-t)/|h^*(-t)|$ , then  $|x(t)| \equiv 1$  bounded


$$\text{But } y(0) = \int_{-\infty}^{+\infty} x(\tau) h(-\tau) d\tau = \int_{-\infty}^{+\infty} \frac{h^*(-\tau) h(-\tau)}{|h(-\tau)|} d\tau = \int_{-\infty}^{+\infty} |h(-\tau)| d\tau = \infty$$

$$f_1(t) * f_2(t)$$



**Figure computation** based on “Flip-slide-multiply-sum/integral”


Step 1°. Flip:  $f_1(t) * f_2(t) = \int_{-\infty}^{+\infty} f_1(\tau) \cdot f_2(t-\tau) d\tau$



Please note that:


$t \geq 0$ : right shift

$t < 0$ : left shift  $\Rightarrow$  result zero



Step 2°. Slide

Step 3°. Multiply




①  $3 \leq t < 6$   $f_1(t) * f_2(t) = \int_1^{-2+t} \frac{1}{2} \cdot 2 d\tau = t-3$

②  $6 \leq t < 9$   $f_1(t) * f_2(t) = \int_{-5+t}^{-2+t} \frac{1}{2} \cdot 2 d\tau = 3$

③  $9 \leq t < 12$   $f_1(t) * f_2(t) = \int_{-5+t}^7 \frac{1}{2} \cdot 2 d\tau = 12-t$

$t > 12 \Rightarrow 0$



Step 4°. Integral.

$$\textcircled{1} \quad 3 \leq t < 6 \quad f_1(t) * f_2(t) = \int_1^{-2+t} \frac{1}{2} \cdot 2 d\tau = t-3$$

$$\textcircled{2} \quad 6 \leq t < 9 \quad f_1(t) * f_2(t) = \int_{-5+t}^{-2+t} \frac{1}{2} \cdot 2 d\tau = 3$$

$$\textcircled{3} \quad 9 \leq t < 12 \quad f_1(t) * f_2(t) = \int_{-5+t}^7 \frac{1}{2} \cdot 2 d\tau = 12-t$$


$$\therefore f_1(t) * f_2(t) = \begin{cases} 0, & t < 3 \\ t-3, & 3 \leq t < 6 \\ 3, & 6 \leq t < 9 \\ 12-t, & 9 \leq t \leq 12 \\ 0, & t > 12 \end{cases}$$




Some known or **typical convolution results**


$$\begin{aligned}
 & f_1(t) * f_2(t) \\
 &= 2 \cdot [u(t-1) - u(t-7)] * \frac{1}{2} [(t-2) - u(t-5)] \\
 &= 2 \cdot \frac{1}{2} \cdot u(t) * [\delta(t-1) - \delta(t-7)] * u(t) * [\delta(t-2) - \delta(t-5)] \\
 &= u(t) * u(t) * [\delta(t-1) - \delta(t-7)] * [\delta(t-2) - \delta(t-5)] \\
 &= tu * [\delta(t-3) - \delta(t-6) - \delta(t-9) + \delta(t-12)] \\
 &= (t-3)u(t-3) - (t-6)u(t-6) - (t-9)u(t-9) + (t-12)u(t-12) \\
 &(1) t < 3, f_1(t) * f_2(t) = 0 \\
 &(2) 3 \leq t < 6, f_1(t) * f_2(t) = t - 3 \\
 &(3) 6 \leq t < 9, f_1(t) * f_2(t) = (t-3) - (t-6) = 3 \\
 &(4) 9 \leq t < 12, f_1(t) * f_2(t) = (t-3) - (t-6) - (t-9) = 12 - t \\
 &(5) 12 \leq t, f_1(t) * f_2(t) = (t-3) - (t-6) - (t-9) + (t-12) = 0
 \end{aligned}$$

**Properties** of convolution






$$f_1(t) = g_1(t) * \delta(t - 4)$$




$$f_2(t) = g_2(t) * \delta(t - \frac{7}{2})$$



$$\therefore f_1(t) * f_2(t) = g_1(t) * g_2(t) * \delta(t - \frac{15}{2})$$

$$= G(t) * \delta(t - \frac{15}{2})$$

$$= G(t - \frac{15}{2})$$



- Fourier Analysis
  1. Response for Complex Exponentials
  2. Fourier Series: Synthesis Equation & Analysis Equation
  3. Properties
  4. Computation on  $a_k$ 
    - ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)
    - ② By the signal Fourier transform in  $[-T/2, T/2]$
    - ③ Formula

- Fourier Analysis

1. Response for Complex Exponentials

2. Fourier Series: Synthesis Equation & Analysis Equation

3. Properties

4. Computation on  $a_k$

- ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)

- ② By the signal Fourier transform in  $[-T/2, T/2]$

- ③ Formula

## • CT LTI systems

$$\begin{aligned}
 x(t) = e^{st} &\longrightarrow \boxed{h(t)} \longrightarrow y(t) = \int_{-\infty}^{+\infty} h(\tau) e^{s(t-\tau)} d\tau \\
 &\quad \boxed{y(t) = x(t) * h(t) = h(t) * x(t) = \int_{-\infty}^{\infty} h(\tau) x(t-\tau) d\tau} \\
 &\quad = \left[ \int_{-\infty}^{+\infty} h(\tau) e^{-s\tau} d\tau \right] e^{st} \\
 &\quad = \underbrace{H(s)}_{\text{eigenvalue}} \underbrace{e^{st}}_{\text{eigenfunction}}
 \end{aligned}$$

$$e^{st} \rightarrow H(s) \cdot e^{st}$$

$$A(\text{常数}) \rightarrow H(0) \cdot A$$

## • DT LTI systems

$$\begin{aligned}
 x[n] = z^n &\longrightarrow \boxed{h[n]} \longrightarrow y[n] = \sum_{m=-\infty}^{\infty} h[m] z^{n-m} \\
 &\quad = \left[ \sum_{m=-\infty}^{\infty} h[m] z^{-m} \right] z^n \\
 &\quad = \underbrace{H(z)}_{\text{eigenvalue}} \underbrace{z^n}_{\text{eigenfunction}}
 \end{aligned}$$

- Fourier Analysis
  1. Response for Complex Exponentials
  2. Fourier Series: Synthesis Equation & Analysis Equation
  3. Properties
  4. Computation on  $a_k$ 
    - ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)
    - ② By the signal Fourier transform in  $[-T/2, T/2]$
    - ③ Formula

## CT Fourier Series Pair

$$(\omega_o = 2\pi/T)$$

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_o t} \quad (\text{Synthesis equation})$$

$$a_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_o t} dt \quad (\text{Analysis equation})$$

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_o t} \xrightarrow{\times e^{-jn\omega_o t}} x(t) e^{-jn\omega_o t} = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_o t} e^{-jn\omega_o t} \xrightarrow{\text{Integral}} \int_0^T x(t) e^{-jn\omega_o t} dt = \int_0^T \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_o t} e^{-jn\omega_o t} dt$$

$$= \sum_{k=-\infty}^{+\infty} a_k \int_0^T e^{jk\omega_o t} e^{-jn\omega_o t} dt$$

Since

$$\int_0^T e^{jk\omega_o t} e^{-jn\omega_o t} dt = \begin{cases} T, & k = n \\ 0, & k \neq n \end{cases}$$

$$\longrightarrow a_n = \frac{1}{T} \int_0^T x(t) e^{-jn\omega_o t} dt \quad (k = n)$$

Please note that (0,T) can be replaced  
by any interval of length T

- Fourier Analysis
  1. Response for Complex Exponentials
  2. Fourier Series: Synthesis Equation & Analysis Equation
  - 3. Properties**
  4. Computation on  $a_k$ 
    - ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)
    - ② By the signal Fourier transform in  $[-T/2, T/2]$
    - ③ Formula



- Linearity  $x(t) \longleftrightarrow a_k, y(t) \longleftrightarrow b_k \Rightarrow \alpha x(t) + \beta y(t) \longleftrightarrow \alpha a_k + \beta b_k$

- Conjugate Symmetry

Based on the same  
period among x and y!

$$x(t) \Rightarrow a_{-k} = a_k^*$$

Proof:

$$a_{-k} = \frac{1}{T} \int_T x(t) e^{jk\omega_o t} dt = \left[ \frac{1}{T} \int_T x^*(t) e^{-jk\omega_o t} dt \right]^* = a_k^*$$

$$\Downarrow a_k = \text{Re}\{a_k\} + j\text{Im}\{a_k\}$$

$$= |a_k| e^{j\angle a_k}$$

$\text{Re}\{a_k\}$  is even,  $\text{Im}\{a_k\}$  is odd

or

$|a_k|$  is even,  $\angle a_k$  is odd

- Time shift

$$x(t) \longleftrightarrow a_k$$

$$x(t - t_o) \longleftrightarrow a_k e^{-jk\omega_o t_o} = a_k e^{-jk2\pi t_o / T}$$

Introduce a linear phase shift  $\propto t_o$

- Time Reversal

$$x(-t) \xleftrightarrow{FS} a_{-k}$$

the effect of sign change for  $x(t)$  and  $a_k$  are **identical**

*Example:*  $x(t)$ : ...  $a_{-2}$   $a_{-1}$   $a_0$   $a_1$   $a_2$  ...

$x(-t)$ : ...  $a_2$   $a_1$   $a_0$   $a_{-1}$   $a_{-2}$  ...

- Time Scaling

$\alpha$  : positive real number

$x(\alpha t)$ : periodic with period  $T/\alpha$  and fundamental frequency  $\alpha\omega_0$

$$x(\alpha t) = \sum_{k=-\infty}^{\infty} a_k e^{jk(\alpha\omega_0)t}$$

$a_k$  **unchanged**, but  $x(\alpha t)$  and each harmonic component are different

## • Multiplication Property

$x(t) \longleftrightarrow a_k$ ,  $y(t) \longleftrightarrow b_k$  (Both  $x(t)$  and  $y(t)$  are  
 $\Downarrow$  periodic with the same period  $T$ )

$$x(t) \cdot y(t) \longleftrightarrow c_k = \sum_{l=-\infty}^{+\infty} a_l b_{k-l} = a_k * b_k$$

**Proof**

$$\begin{aligned}
 x(t)y(t) &= \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t} \cdot \sum_{l=-\infty}^{+\infty} b_l e^{jl\omega_0 t} = \sum_{k=-\infty}^{+\infty} \sum_{l=-\infty}^{+\infty} a_k b_l e^{j(l+k)\omega_0 t} \\
 &= \sum_{k=-\infty}^{+\infty} \sum_{p=-\infty}^{+\infty} a_k b_{p-k} e^{jp\omega_0 t} \quad (l+k=p, l=p-k) \\
 &= \sum_{p=-\infty}^{+\infty} \left( \sum_{k=-\infty}^{+\infty} a_k b_{p-k} \right) e^{jp\omega_0 t} \\
 &= \sum_{k=-\infty}^{+\infty} \left( \sum_{l=-\infty}^{+\infty} a_l b_{k-l} \right) e^{jk\omega_0 t} \quad (p=k, k=l)
 \end{aligned}$$

- Parseval Relation

$$\frac{1}{T} \int_T |x(t)|^2 dt = \sum_{k=-\infty}^{+\infty} |a_k|^2$$

**Observation:** power is the same whether measured in the time-domain or the frequency-domain

**Proof**

$$|x(t)|^2 = x(t)x^*(t)$$

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t}, x^*(t) = \sum_{l=-\infty}^{+\infty} a_{-l}^* e^{jk\omega_0 t} = \sum_{l=-\infty}^{+\infty} a_l^* e^{-jk\omega_0 t}$$

$$\frac{1}{T} \int_T |x(t)|^2 dt = \frac{1}{T} \int_T \sum_{k=-\infty}^{+\infty} \sum_{l=-\infty}^{+\infty} a_k a_l^* e^{j(k-l)\omega_0 t} dt = \frac{1}{T} \sum_{k=-\infty}^{+\infty} \sum_{l=-\infty}^{+\infty} a_k a_l^* \int_T e^{j(k-l)\omega_0 t} dt$$

$$\int_T e^{jk\omega_0 t} e^{-jl\omega_0 t} dt = \begin{cases} T, k = l \\ 0, k \neq l \end{cases}$$

$$\frac{1}{T} \int_T |x(t)|^2 dt = \sum_{k=-\infty}^{+\infty} |a_k|^2$$

- Fourier Analysis
  1. Response for Complex Exponentials
  2. Fourier Series: Synthesis Equation & Analysis Equation
  3. Properties
  4. Computation on  $a_k$ 
    - ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)
    - ② By the signal Fourier transform in  $[-T/2, T/2]$
    - ③ Formula

$$x(t) = \sin(\omega_0 t) = \frac{1}{2j} \left( e^{j\omega_0 t} - e^{-j\omega_0 t} \right)$$

$$a_1 = \frac{1}{2j}, a_{-1} = -\frac{1}{2j}$$

**Conjugate Symmetry**

$$x(t) = 1 + \sin(\omega_0 t) + 2 \cos(\omega_0 t) + \cos\left(2\omega_0 t + \frac{\pi}{4}\right)$$

$$x(t) = 1 + \frac{1}{2j} \left( e^{j\omega_0 t} - e^{-j\omega_0 t} \right) + \frac{2}{2} \left( e^{j\omega_0 t} + e^{-j\omega_0 t} \right) + \frac{1}{2} \left( e^{j\left(2\omega_0 t + \frac{\pi}{4}\right)} + e^{-j\left(2\omega_0 t + \frac{\pi}{4}\right)} \right)$$

$$= 1 + \left( 1 + \frac{1}{2j} \right) e^{j\omega_0 t} + \left( 1 - \frac{1}{2j} \right) e^{-j\omega_0 t} + \frac{1}{2} e^{j\frac{\pi}{4}} e^{j2\omega_0 t} + \frac{1}{2} e^{-j\frac{\pi}{4}} e^{-j2\omega_0 t}$$

$$a_0 = 1, a_1 = \left( 1 + \frac{1}{2j} \right), a_2 = \frac{\sqrt{2}}{4} (1 + j), a_{-1} = \left( 1 - \frac{1}{2j} \right), a_{-2} = \frac{\sqrt{2}}{4} (1 - j)$$

- **Conjugate Symmetry**

- **Please think about**  $x(t) = 1 + \sin(\omega_0 t) + 2 \cos(\omega_0 t) + \cos\left(\frac{\omega_0}{2} t + \frac{\pi}{4}\right)$

- **Hint: You should find the period and the fundamental frequency!**

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\frac{\omega_0}{2}t}$$

$$a_0 = 1, a_2 = \left( 1 + \frac{1}{2j} \right), a_1 = \frac{\sqrt{2}}{4} (1 + j), a_{-2} = \left( 1 - \frac{1}{2j} \right), a_{-1} = \frac{\sqrt{2}}{4} (1 - j)$$

- Fourier Analysis
  1. Response for Complex Exponentials
  2. Fourier Series: Synthesis Equation & Analysis Equation
  3. Properties
  4. Computation on  $a_k$ 
    - ① When  $x(t)$  are sin, cos or complex exponential signals (Euler Equation)
    - ② By the signal Fourier transform in  $[-T/2, T/2]$
    - ③ Formula





$$a_k = \frac{1}{T} X_0(j\omega) \Big|_{\omega=k\omega_0}$$

$$x_0(t) \xleftrightarrow{\mathcal{F}} X_0(j\omega) \quad x_0(t) = x(t), -\frac{T}{2} \leq t \leq \frac{T}{2}$$

$\mathcal{F} : \text{Fourier Transform}$

Please refer to (4.10) on page 289 for more details



## Example 4.4


Consider the rectangular pulse signal

$$x(t) = \begin{cases} 1, & |t| < T_1 \\ 0, & |t| > T_1 \end{cases}, \quad (4.16)$$


as shown in Figure 4.8(a). Applying eq. (4.9), we find that the Fourier transform of this signal is

$$X(j\omega) = \int_{-T_1}^{T_1} e^{-j\omega t} dt = 2 \frac{\sin \omega T_1}{\omega}, \quad (4.17)$$

as sketched in Figure 4.8(b).




(a)



(b)

## Recall Example 3.5 on Textbook



$$\begin{aligned}
 x_0(t) &\xleftrightarrow{\mathcal{F}} X_0(j\omega) = \frac{2 \sin(\omega T_1)}{\omega} \\
 a_k &= \frac{1}{T} X_0(j\omega) \Big|_{\omega=k\omega_0} = \frac{1}{T} \frac{2 \sin(\omega T_1)}{\omega} \Big|_{\omega=k\omega_0} \\
 &= \frac{1}{T} \frac{2 \sin(k\omega_0 T_1)}{k\omega_0} \\
 &= \frac{1}{T} \frac{2 \sin(k\omega_0 T_1)}{k \frac{2\pi}{T}} \\
 &= \frac{\sin(k\omega_0 T_1)}{k\pi}
 \end{aligned}$$

By the signal Fourier transform in  $[-T/2, T/2]$

Rewrite

$$x(t) = \sum_{k=-\infty}^{+\infty} x_0(t - kT), \quad x_0(t) = \begin{cases} 1, & |t| < T_1 \\ 0, & |t| > T_1 \end{cases}$$

$$\begin{aligned}
 x(t) &= \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t}, \quad \omega_0 = \frac{2\pi}{T} \\
 a_k &= \frac{1}{T} \int_{-T_1}^{T_1} e^{-jk\omega_0 t} dt = \frac{\sin(k\omega_0 T_1)}{\pi k}, \quad k \neq 0
 \end{aligned}$$

Based on analysis equation

**3.8.** Suppose we are given the following information about a signal  $x(t)$ :

1.  $x(t)$  is real and odd.
2.  $x(t)$  is periodic with period  $T = 2$  and has Fourier coefficients  $a_k$ .
3.  $a_k = 0$  for  $|k| > 1$ .
4.  $\frac{1}{2} \int_0^2 |x(t)|^2 dt = 1$ .

Specify two different signals that satisfy these conditions.

- 3.8. Since  $x(t)$  is real and odd (clue 1), its Fourier series coefficients  $a_k$  are purely imaginary and odd (See Table 3.1). Therefore,  $a_k = -a_{-k}$  and  $a_0 = 0$ . Also, since it is given that  $a_k = 0$  for  $|k| > 1$ , the only unknown Fourier series coefficients are  $a_1$  and  $a_{-1}$ . Using Parseval's relation,

$$\frac{1}{T} \int_{\langle T \rangle} |x(t)|^2 dt = \sum_{k=-\infty}^{\infty} |a_k|^2,$$

for the given signal we have

$$\frac{1}{2} \int_0^2 |x(t)|^2 dt = \sum_{k=-1}^1 |a_k|^2.$$

Using the information given in clue (4) along with the above equation,

$$|a_1|^2 + |a_{-1}|^2 = 1 \quad \Rightarrow \quad 2|a_1|^2 = 1$$

Therefore,

$$a_1 = -a_{-1} = \frac{1}{\sqrt{2}j} \quad \text{or} \quad a_1 = -a_{-1} = -\frac{1}{\sqrt{2}j}$$

The two possible signals which satisfy the given information are

$$x_1(t) = \frac{1}{\sqrt{2}j} e^{j(2\pi/2)t} - \frac{1}{\sqrt{2}j} e^{-j(2\pi/2)t} = -\sqrt{2} \sin(\pi t)$$

and

$$x_2(t) = -\frac{1}{\sqrt{2}j} e^{j(2\pi/2)t} + \frac{1}{\sqrt{2}j} e^{-j(2\pi/2)t} = \sqrt{2} \sin(\pi t)$$

**3.34.** Consider a continuous-time LTI system with impulse response


$$h(t) = e^{-4|t|}.$$

Find the Fourier series representation of the output  $y(t)$  for each of the following inputs:

(a)  $x(t) = \sum_{n=-\infty}^{+\infty} \delta(t - n)$

(b)  $x(t) = \sum_{n=-\infty}^{+\infty} (-1)^n \delta(t - n)$

(c)  $x(t)$  is the periodic wave depicted in Figure P3.34.



**Figure P3.34**

3.34. The frequency response of the system is given by

$$H(j\omega) = \int_{-\infty}^{\infty} e^{-4|t|} e^{-j\omega t} dt = \frac{1}{4 + j\omega} + \frac{1}{4 - j\omega}.$$

(a) Here,  $T = 1$  and  $\omega_0 = 2\pi$  and  $a_k = 1$  for all  $k$ . The FS coefficients of the output are

$$b_k = a_k H(jk\omega_0) = \frac{1}{4 + j2\pi k} + \frac{1}{4 - j2\pi k}.$$

(b) Here,  $T = 2$  and  $\omega_0 = \pi$  and

$$a_k = \begin{cases} 0, & k \text{ even} \\ 1, & k \text{ odd} \end{cases}.$$

Therefore, the FS coefficients of the output are

$$b_k = a_k H(jk\omega_0) = \begin{cases} 0, & k \text{ even} \\ \frac{1}{4 + j\pi k} + \frac{1}{4 - j\pi k}, & k \text{ odd} \end{cases}.$$

(c) Here,  $T = 1$ ,  $\omega_0 = 2\pi$  and

$$a_k = \begin{cases} 1/2, & k = 0 \\ 0, & k \text{ even}, k \neq 0 \\ \frac{\sin(\pi k/2)}{\pi k}, & k \text{ odd} \end{cases}.$$

Therefore, the FS coefficients of the output are

$$b_k = a_k H(jk\omega_0) = \begin{cases} 1/4, & k = 0 \\ 0, & k \text{ even}, k \neq 0 \\ \frac{\sin(\pi k/2)}{\pi k} \left[ \frac{1}{4+j2\pi k} + \frac{1}{4-j2\pi k} \right], & k \text{ odd} \end{cases}.$$



**3.40.** Let  $x(t)$  be a periodic signal with fundamental period  $T$  and Fourier series coefficients  $a_k$ . Derive the Fourier series coefficients of each of the following signals in terms of  $a_k$ :

(a)  $x(t - t_0) + x(t + t_0)$

(b)  $\mathcal{E}\{x(t)\}$

(c)  $\mathcal{R}\{x(t)\}$

(d)  $\frac{d^2 x(t)}{dt^2}$

(e)  $x(3t - 1)$  [for this part, first determine the period of  $x(3t - 1)$ ]

$$\begin{aligned} \text{Even } \{x(t)\} &= \frac{x(t) + x(-t)}{2} \\ \text{Odd } \{x(t)\} &= \frac{x(t) - x(-t)}{2} \end{aligned}$$

(a)  $x(t - t_0)$  is also periodic with period  $T$ . The Fourier series coefficients  $b_k$  of  $x(t - t_0)$  are

$$\begin{aligned} b_k &= \frac{1}{T} \int_T x(t - t_0) e^{-jk(2\pi/T)t} dt \\ &= \frac{e^{-jk(2\pi/T)t_0}}{T} \int_T x(\tau) e^{-jk(2\pi/T)\tau} d\tau \\ &= e^{-jk(2\pi/T)t_0} a_k \end{aligned}$$

Similarly, the Fourier series coefficients of  $x(t + t_0)$  are

$$c_k = e^{jk(2\pi/T)t_0} a_k.$$

Finally, the Fourier series coefficients of  $x(t - t_0) + x(t + t_0)$  are

$$d_k = b_k + c_k = e^{-jk(2\pi/T)t_0} a_k + e^{jk(2\pi/T)t_0} a_k = 2 \cos(k2\pi t_0/T) a_k.$$

(b) Note that  $\mathcal{E}v\{x(t)\} = [x(t) + x(-t)]/2$ . The FS coefficients of  $x(-t)$  are

$$\begin{aligned} b_k &= \frac{1}{T} \int_T x(-t) e^{-jk(2\pi/T)t} dt \\ &= \frac{1}{T} \int_T x(\tau) e^{jk(2\pi/T)\tau} d\tau \\ &= a_{-k} \end{aligned}$$

Therefore, the FS coefficients of  $\mathcal{E}v\{x(t)\}$  are

$$c_k = \frac{a_k + b_k}{2} = \frac{a_k + a_{-k}}{2}.$$

(c) Note that  $\mathcal{Re}\{x(t)\} = [x(t) + x^*(t)]/2$ . The FS coefficients of  $x^*(t)$  are

$$b_k = \frac{1}{T} \int_T x^*(t) e^{-jk(2\pi/T)t} dt.$$

Conjugating both sides, we get

$$b_k^* = \frac{1}{T} \int_T x(t) e^{jk(2\pi/T)t} dt = a_{-k}.$$

Therefore, the FS coefficients of  $\mathcal{Re}\{x(t)\}$  are

$$c_k = \frac{a_k + b_k}{2} = \frac{a_k + a_{-k}^*}{2}.$$

(d) The Fourier series synthesis equation gives

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{j(2\pi/T)kt}.$$

Differentiating both sides wrt  $t$  twice, we get

$$\frac{d^2 x(t)}{dt^2} = \sum_{k=-\infty}^{\infty} -k^2 \frac{4\pi^2}{T^2} a_k e^{j(2\pi/T)kt}.$$

By inspection, we know that the Fourier series coefficients of  $d^2 x(t)/dt^2$  are  $-k \frac{4\pi^2}{T^2} a_k$ .

(e) The period of  $x(3t)$  is a third of the period of  $x(t)$ . Therefore, the signal  $x(3t - 1)$  is periodic with period  $T/3$ . The Fourier series coefficients of  $x(3t)$  are still  $a_k$ . Using the analysis of part (a), we know that the Fourier series coefficients of  $x(3t - 1)$  is  $e^{-jk(6\pi/T)} a_k$ .



南方科技大学  
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Thanks for Your Attendance**

**Q&A**

Shihang Lu (卢仕航), TA

Southern University of Science and Technology

Email: [lush2021@mail.sustech.edu.cn](mailto:lush2021@mail.sustech.edu.cn)

工学院南楼220 (周五16:30-17:30)