

## CSE 331 Final Exam Preparation

This is in no way a substitute for exam preparation, merely a compilation of all the key talking points.

Chandra Neppalli

May 3, 2021

# Counter Example

Best proof to use to *disprove* universally true propositions.

# Counter Example

Best proof to use to *disprove* universally true propositions.

Ex: Every day is a Wednesday, where a counter example would be Monday is not Wednesday.

# Contradiction

Best proof to use if you want to assert something is true.

# Contradiction

Best proof to use if you want to assert something is true.

Assume what you want to prove is false, then show this leads to a contradiction.

# Contradiction

Best proof to use if you want to assert something is true.

Assume what you want to prove is false, then show this leads to a contradiction.

Therefore, the original assumption has to be true.

# Contraposition

Best proof for proving causality. Define two propositions E and F.

# Contraposition

Best proof for proving causality. Define two propositions  $E$  and  $F$ .

If you want to prove that  $E \rightarrow F$ , it might be more doable to prove  $\neg F \rightarrow \neg E$ , as they are both logically equivalent.



# Contraposition

Best proof for proving causality. Define two propositions  $E$  and  $F$ .

If you want to prove that  $E \rightarrow F$ , it might be more doable to prove  $\neg F \rightarrow \neg E$ , as they are both logically equivalent.

This is especially useful if the **scope** of  $F$  is smaller than the scope of  $E$ .

# Direct Proof

If the proof is *simple*, consider directly proving it.

# Direct Proof

If the proof is *simple*, consider directly proving it.

Remember though, that you must maintain *W.L.O.G*, that your proof can never be too specific and must be arbitrary.

# Proof by Induction

Proof by Induction is a really nice proof technique when you reduce your proof to a known correct base case.

# Proof by Induction

Proof by Induction is a really nice proof technique when you reduce your proof to a known correct base case.

If proof needs to be correct for all numbers  $\in \mathbb{N}$ , and each step is dependant on the previous step, then *every* step can be reduced to a definitive base case that is easy to directly prove.

## Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

- $P(0) = I$



# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

- $P(0) = I$
- $P(i)$  is an accumulator. This means that  $P(i + 1) > P(i)$

# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

- $P(0) = l$
- $P(i)$  is an accumulator. This means that  $P(i+1) > P(i)$
- $\forall i, P(i) \leq k$

# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

- $P(0) = l$
- $P(i)$  is an accumulator. This means that  $P(i+1) > P(i)$
- $\forall i, P(i) \leq k$

From these 3 properties, the number of iterations is bounded by  $k - l + 1$

# Extra: Progress Measure

This is useful for proving an algorithm with a loop terminates.

Let  $P(i)$  denote an integer such that:

- $P(0) = l$
- $P(i)$  is an accumulator. This means that  $P(i+1) > P(i)$
- $\forall i, P(i) \leq k$

From these 3 properties, the number of iterations is bounded by  $k - l + 1$

Note: This isn't a runtime analysis, rather a proof that the algorithm terminates.

# Greedy Stays Ahead

This technique is used to prove that a greedy algorithm returns an optimal solution.

# Greedy Stays Ahead

This technique is used to prove that a greedy algorithm returns an optimal solution.

At every step of a greedy algorithm, it will stay *at least* as far as the optimal solution at that step.

# Greedy Stays Ahead

This technique is used to prove that a greedy algorithm returns an optimal solution.

At every step of a greedy algorithm, it will stay *at least* as far as the optimal solution at that step.

HW4 “Attack on Alarms” and Interval Scheduling are examples of problems with greedy solutions.

# Introduction

Let's say there are two groups: Group A and Group B.



# Introduction

Let's say there are two groups: Group A and Group B.

How do we generate a **stable** matching between each member of the two groups efficiently?

# Introduction

Let's say there are two groups: Group A and Group B.

How do we generate a **stable** matching between each member of the two groups efficiently?

Moreover, what is a **stable** matching?

# Perfect Matchings

A **perfect matching** is a bijective matching between  $A$  and  $B$ .

# Perfect Matchings

A **perfect matching** is a bijective matching between A and B.

**Every member** in group A is matched with exactly one member in group B.

# Perfect Matchings

A **perfect matching** is a bijective matching between A and B.

**Every member** in group A is matched with exactly one member in group B.

Conversely, every member in group B is matched with **exactly** one member in group A.

# Perfect Matchings

A **perfect matching** is a bijective matching between A and B.

**Every member** in group A is matched with exactly one member in group B.

Conversely, every member in group B is matched with **exactly** one member in group A.

With  $n$  members in each group, there are  $n!$  perfect matchings.

# Instability

For a particular matching, define a member  $m$  from group A and  $n$  from group B such that  $(m, n)$  is not in the matching.

# Instability

For a particular matching, define a member  $m$  from group A and  $n$  from group B such that  $(m, n)$  is not in the matching.

If  $m$  prefers  $n$  over their current matching **and**  $n$  prefers  $m$  over their current matching, the entire match is an instability.



# Stable Matching

A stable matching **is** a perfect matching with no instability.

# Stable Matching

A stable matching **is** a perfect matching with no instability.

It therefore follows that the number of stable matchings is *at least* the number of perfect matchings, or  $n!$

# Stable Matching

A stable matching **is** a perfect matching with no instability.

It therefore follows that the number of stable matchings is *at least* the number of perfect matchings, or  $n!$

The Gale Shapely Algorithm is an  $O(n^3)$  time algorithm that can output a stable matching.

# Stable Matching

A stable matching **is** a perfect matching with no instability.

It therefore follows that the number of stable matchings is *at least* the number of perfect matchings, or  $n!$

The Gale Shapely Algorithm is an  $O(n^3)$  time algorithm that can output a stable matching.

With the right data structures, the runtime can be reduced to  $O(n^2)$ .

# Stable Matching

A stable matching **is** a perfect matching with no instability.

It therefore follows that the number of stable matchings is *at least* the number of perfect matchings, or  $n!$

The Gale Shapely Algorithm is an  $O(n^3)$  time algorithm that can output a stable matching.

With the right data structures, the runtime can be reduced to  $O(n^2)$ .

Even though the runtime isn't linear, because the input size is  $2n^2 \rightarrow \Theta(n^2)$ <sup>1</sup>, the runtime **with respect** to the input size is  $O(N)$ , or linear time.

---

<sup>1</sup>This comes from  $n$  Group A members and  $n$  Group B members with their  $2n$  preference lists

# Stable Matching

Code:

---

```
Initially all  $m \in M$  and  $w \in W$  are free
While there is a man  $m$  who is free and hasn't proposed to every
woman
    Choose such a man  $m$ 
    Let  $w$  be the highest-ranked woman in  $m$ 's
        preference list to whom  $m$  has not yet proposed
    If  $w$  is free then
         $(m, w)$  become engaged
    Else  $w$  is currently engaged to  $m'$ 
        If  $w$  prefers  $m'$  to  $m$  then
             $m$  remains free
        If  $w$  prefers  $m$  to  $m'$ 
             $(m, w)$  become engaged
             $m'$  becomes free
        Endif
    Endif
EndWhile
Return the set  $S$  of engaged pairs.
```