

Stat 346 Homework 2

Nathan Jarus

Feb. 13, 2014

1 Problem 1, KNN #2.10

2 Problem 2

2.1 Part a

The confidence interval for β_1 is defined as follows:

$$\beta_1 \pm t_{1-\alpha/2, n-1}(s(b_1)) \quad (1)$$

Substituting the provided values, we get:

$$3 \pm 2.09302 \quad (2)$$

2.2 Part b

$$H_0 : \beta_1 = 4 \quad (3)$$

$$H_a : \beta_1 \neq 4 \quad (4)$$

Based on this, we can set up a test statistic:

$$t^* = \frac{b_1 - 4}{s(b_1)} = -1 \quad (5)$$

How does one calculate the p value?

Ultimately, I think we fail to reject H_0 .


3 Problem 3

3.1 Part a

The scatterplot is shown in Figure 3.1 With sufficient smoothing, the relationship is vaguely linear.

16:58 Wednesday, February 12, 2014 1

GPA and ACT
Scatter plot of ACT Score vs. GPA with Smoothing Line N=80



3.2 Part b

GPA and ACT**Scatter plot of ACT Score vs. GPA with Smoothing Line N=80****The REG Procedure****Model: MODEL1****Dependent Variable: gpa**

| | |
|------------------------------------|-----|
| Number of Observations Read | 120 |
| Number of Observations Used | 120 |

| Analysis of Variance | | | | | |
|-----------------------------|-----------|-----------------------|--------------------|----------------|------------------|
| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
| Model | 1 | 3.58785 | 3.58785 | 9.24 | 0.0029 |
| Error | 118 | 45.81761 | 0.38828 | | |
| Corrected Total | 119 | 49.40545 | | | |

| | | | |
|-----------------------|----------|-----------------|--------|
| Root MSE | 0.62313 | R-Square | 0.0726 |
| Dependent Mean | 3.07405 | Adj R-Sq | 0.0648 |
| Coeff Var | 20.27049 | | |

| Parameter Estimates | | | | | | | |
|----------------------------|-----------|---------------------------|-----------------------|----------------|--------------------|------------------------------|---------|
| Variable | DF | Parameter Estimate | Standard Error | t Value | Pr > t | 95% Confidence Limits | |
| Intercept | 1 | 2.11405 | 0.32089 | 6.59 | <.0001 | 1.47859 | 2.74951 |
| act | 1 | 0.03883 | 0.01277 | 3.04 | 0.0029 | 0.01353 | 0.06412 |

The estimated regression equation is

$$\hat{Y} = 2.11405 + 0.03883 * \hat{X} \quad (6)$$

For the slope, the 95% confidence limits are 0.01353 and 0.06412. Since they are both positive, we can conclude with confidence > 95% that there is a positive linear relationship between GPA and ACT scores.

3.3 Part c

A significance test for the slope, b_1 uses the hypothesis

$$H_0 : b_1 = 0 \quad (7)$$

$$H_a : b_1 \neq 0 \quad (8)$$

Our test statistic is 3.04 with 118 DOF and the resulting p-value is 0.0029. Since $p < 0.05$, we reject H_0 , meaning that there is a significant linear relationship between GPA and ACT scores.

3.4 Part d

The lowest X value in the data set is 14, so an ACT score of 0 (the intercept) would not be within the scope of the model.

3.5 Part e

Our mean response for an ACT score of 21 is a GPA of 2.92948. The 95% confidence interval is (2.7826, 3.0763).

3.6 Part f

The prediction is the same as for Part E, 2.92948. However, the confidence interval is (1.6868, 4.1721).

3.7 Part g

Percent of variance accounted for by ACT score:

$$\frac{3.58785}{49.40545} * 100 = 7.26205\% \quad (9)$$

Correlation: 0.26948

3.8 Part h


I do not think that ACT scores make a good predictor for GPA. The confidence intervals on predictions are very wide, and ACT score accounts for only 7% of the variance in GPA.

4 Problem 4

4.1 Part a

As shown in Figure 4.1, the data definitely follows a linear pattern, but seems to be sorted into buckets. This may demonstrate a nonlinear relationship between hardness and time, or it may be a limitation of the measurement method for hardness.

Plastic Hardness
Scatter plot of hardness vs time with regression line



4.2 Part b

Plastic Hardness

Scatter plot of hardness vs time with regression line

The REG Procedure

Model: MODEL1

Dependent Variable: hardness

| | |
|------------------------------------|----|
| Number of Observations Read | 16 |
| Number of Observations Used | 16 |

| Analysis of Variance | | | | | |
|------------------------|----|----------------|-------------|---------|--------|
| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
| Model | 1 | 1245.57198 | 1245.57198 | 506.51 | <.0001 |
| Error | 14 | 34.42802 | 2.45914 | | |
| Corrected Total | 15 | 1280.00000 | | | |

| | | | |
|-----------------------|----------|-----------------|--------|
| Root MSE | 1.56817 | R-Square | 0.9731 |
| Dependent Mean | 28.00000 | Adj R-Sq | 0.9712 |
| Coeff Var | 5.60059 | | |

| Parameter Estimates | | | | | | | |
|---------------------|----|--------------------|----------------|---------|---------|-----------------------|-----------|
| Variable | DF | Parameter Estimate | Standard Error | t Value | Pr > t | 95% Confidence Limits | |
| Intercept | 1 | -79.89337 | 4.81005 | -16.61 | <.0001 | -90.20990 | -69.57683 |
| hours | 1 | 0.47833 | 0.02125 | 22.51 | <.0001 | 0.43275 | 0.52392 |

Estimated regression equation:

$$\hat{Y} = -79.89337 + 0.47833 * X \quad (10)$$


The slope 95% confidence interval is (0.43275, 0.52392).

4.3 Part c

As shown in Figure 4.3, the prediction band is wider to account for the extra variation due to the fact that new observations will probably not fall on the regression line.

12:44 Thursday, February 13, 2014 1

Plastic Hardness
Confidence and Prediction Bands



| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|--------|----|----------------|-------------|---------|---------|
| Model | 1 | 45.20 | 45.20 | 43.88 | <0.0001 |
| Error | 70 | 72.1 | 1.03 | | |
| Total | 71 | 117.3 | | | |

Tab. 9: Problem 5 ANOVA Table

4.4 Part d

97.31031% of the variance in hardness is due to time. The correlation coefficient between hardness and time is 0.98646.

4.5 Part e

Time seems to be a very good predictor for plastic hardness.

5 Problem 5

$$H_0 : b_1 = 0 \quad (11)$$

$$H_a : b_1 \neq 0 \quad (12)$$

From the values in table 9, we can reject the null hypothesis and conclude that we have a reasonably good fit.