A Diagrammatic Proof of the Haldane Gap via Tensor Networks

Sirui Lu
Max-Planck-Institut für Quantenoptik, Garching, Germany and
Munich Center for Quantum Science and Technology (MCQST), Schellingstrasse 4, D-80799 München, Germany
(Dated: November 7, 2026)

We give a diagrammatic exposition of OpenAI's computer-assisted proof of a uniform gap in the periodic spin-1 Heisenberg chain. By reading the partition-function tensor network along space and imaginary time, we show how bounds for finite rings imply a gap for all sufficiently long even rings. We improve the spin-1 gap bound and extend the argument to certify uniform gaps for seven additional reflection-positive spin chains and ladders. We formalize the general argument in a reusable Lean library and give a complete Lean proof for a mixed-field Ising example.

Haldane predicted a gap for integer-spin antiferromagnetic Heisenberg chains, in contrast to their half-integer counterparts [1, 2]. For spin one, numerical work places the bulk gap near 0.4105 [3]. Unlike the Affleck–Kennedy–Lieb–Tasaki model [4], the Heisenberg chain is not frustration free: neighboring bonds cannot simultaneously attain their singlet ground states. Finite-size criteria based on frustration freeness [5, 6] therefore do not apply directly.

OpenAI proved a unique ground state and a uniform gap $\Delta_L > (\log 20)/784$ for every even periodic length $L \geq 2304$ using certified finite partition-function bounds [7]. Here we explain the proof using tensor-network diagrams. We extend it to other spin models under a condition that we identify as reflection positivity, and give Lean proofs. We also improve the spin-1 gap bound using OpenAI's finite trace calculations and trial states; see details [8].

Thermal purity and the gap.— Consider


$$H_L = \sum_{j=1}^{L} \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}, \qquad Z_L(\beta) = \operatorname{Tr} e^{-\beta H_L}, \tag{1}$$

with periodic indices, spin one, even $L \geq 4$, and exchange coupling one. Order energies with multiplicity and put $\Delta_L = E_1(L) - E_0(L)$, illustrated by  . For $\beta > 0$, the Gibbs state $\rho_L = e^{-\beta H_L}/Z_L(\beta)$ has ordered weights $w_0 \geq w_1 \geq \cdots$. Its thermal purity $P_\tau(L,\beta) = \operatorname{Tr} \rho_L^2 \leq w_0$ gives

$$e^{-\beta\Delta_L} = \frac{w_1}{w_0} \leq \frac{1-w_0}{w_0} \leq \frac{1-P_\tau(L,\beta)}{P_\tau(L,\beta)}. \tag{2}$$

Purity close to one forces a single dominant Gibbs weight,  . Thus $P_\tau > 1/2$ forces a unique ground state, and $1 - P_\tau \leq q < 1/2$, with $q > 0$, gives $\Delta_L \geq \beta^{-1} \log[(1-q)/q]$.

Proof strategy.— To keep the gap bound positive as we lower the temperature, $\log(1/q)$ must grow proportionally to β. Squaring and normalizing the Gibbs state doubles β, but does not extend the chain. The key idea from the OpenAI proof is to construct a spatial state whose square doubles the length, so that we can iterate both operations:



If each step squares the purity errors, their negative logarithms grow as 2^j, matching $\beta_j = 2^j\beta_0$. Taking the logarithm and dividing by β_j therefore leaves a positive constant in the gap bound. We must establish this estimate and cover the lengths between steps. First we construct the spatial state.


Spatial transfer matrix.— Set $\tau = \beta/M$ and let $S_L(\tau)$ be the product of $I - \tau h_{j,j+1}$ over even bonds followed by odd bonds, where $h = \boldsymbol{S} \cdot \boldsymbol{S}$. Contracting the resulting finite network in time or space gives [9]

$$\underbrace{\operatorname{Tr} S_L(\beta/M)^M}_{\text{time}} = Z_L^{(M)}(\beta) = \underbrace{\operatorname{Tr} T_{\beta,M}^{L/2}}_{\text{space}}. \tag{3}$$



In the Cartesian basis $|x\rangle, |y\rangle, |z\rangle$, the generators $G_\alpha = iS^\alpha$ are real antisymmetric matrices. Hence

$$-h = \sum_{\alpha=x,y,z} G_\alpha \otimes G_\alpha, \qquad I - \tau h = \sum_{\gamma=0,x,y,z} A_\gamma \otimes A_\gamma, \tag{4}$$

where $A_0 = I$ and $A_\alpha = \sqrt{\tau}\, G_\alpha$. Diagrammatically,  . Matching real bond factors express reflection positivity [10–12].

For bond-label sequences $a, b \in \{0, x, y, z\}^M$, define the real matrix $\mathcal{F}_{ba} = \operatorname{Tr}[(A_{b_1} A_{a_1}) \cdots (A_{b_M} A_{a_M})]$. The two sites sharing an even bond have identical factor orderings. Contracting their shared sequence gives

$$T_{\beta,M} = \mathcal{F}^{\mathsf{T}}\mathcal{F} \geq 0, \qquad \text{}. \tag{5}$$


This transfer matrix advances by two sites and is independent of L.


For even $n \geq 4$, normalize $T_{\beta,M}^{n/2}$ by its trace to obtain the density operator $\sigma_n^{(M)}$. The trace is nonzero because

$\mathcal{F}_{00} = 3$ for the all-zero sequences. Nonnegative transfer eigenvalues give spatial purity, squaring, and power-sum monotonicity inequalities. At each fixed L, β, the matrix product limit $S_L(\beta/M)^M \to e^{-\beta H_L}$ passes these closed scalar inequalities to exact partition functions. We take this limit before the doubling argument, requiring neither a limiting transfer matrix nor a Trotter error uniform in L. The finite-Trotter section of the Supplemental Material [8] proves these statements. Thermal identities below concern the exact Gibbs operator, not finite Trotter products.


Coupling and squaring.— The two purities are

$$P_{\mathrm{s}}(n,\beta) = \lim_{M\to\infty} \mathrm{Tr}[(\sigma_n^{(M)})^2] = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2},$$
$$P_\tau(n,\beta) = \mathrm{Tr}\,\rho_n(\beta)^2 = \frac{Z_n(2\beta)}{Z_n(\beta)^2}. \tag{6}$$

Both lie in $(0,1]$. Their contractions, 

and 

, join normalized copies in space and time. The rectangle  denotes $Z_n(\beta)$, with width n and height β. Joining four copies in either order gives



The two paths therefore give the exact identity


$$P_{\mathrm{s}}(n,2\beta)P_\tau(n,\beta)^2 = P_\tau(2n,\beta)P_{\mathrm{s}}(n,\beta)^2. \tag{7}$$

It holds without reflection positivity; positivity is needed to use its factors as purities.

For probabilities u_i, write $P = \sum_i u_i^2$. Their normalized squares have purity P', with

$$1 - P' = \frac{2\sum_{i<j}(u_i u_j)^2}{P^2} \le \frac{(1-P)^2}{2P^2} = f(1-P), \tag{8}$$

where $f(z) = z^2/[2(1-z)^2]$. Thus squaring concentrates weights,  . The same operation acts in the two directions:




(9)

Suppose $1 - P_{\mathrm{s}}(n,\beta) \le p$ and $1 - P_\tau(2n,\beta) \le q$. Dropping the favorable denominator $P_\tau(n,\beta)^2 \le 1$ in Eq. (7), then squaring in space, gives p'. Repeating the identity at length $2n$, then squaring in time, gives q':

$$p' = f(1-(1-p)^2(1-q)),$$
$$q' = f(1-(1-q)^2(1-p')). \tag{10}$$

For $0 \le p, q, p', q' < 1$, these bound $1 - P_{\mathrm{s}}(2n,2\beta)$ and $1 - P_\tau(4n,2\beta)$. The thermal update uses the new spatial bound:



The finite seed and uniform gap.— The existing computation supplies an independent input.

Lemma 1 (Finite seed [7])**.** *At $\beta_0 = 49/4$, the periodic spin-1 chain satisfies $P_{\mathrm{s}}(36,\beta_0) \ge 1 - 47916/10^6$ and $P_\tau(72,\beta_0) \ge 1 - 181521/10^6$.*

Accepting this lemma suffices for the following argument. Six rationally verified, upward-rounded updates give $p_6 = 0.0084709$ and $q_6 = 0.0055149$ at $(n_6,\beta_6) = (2304, 784)$, as shown in Fig. 1. The doubling-argument section of the Supplemental Material [8] records the exact orbit and rounding rule.

For $p, q \le \varepsilon \le 0.01$, both updates are at most $5\varepsilon^2$; the Supplemental Material [8] proves this scalar estimate. Restarting at $(2304, 784)$ gives bounds $\varepsilon_j = 20^{-2^j}/5$ at $n_j = 2304\,2^j$, $\beta_j = 784\,2^j$. To cover every even $L \ge 2304$, choose j with $n_j \le L \le 2n_j$. Spatial power-sum monotonicity at β_j and $2\beta_j$ yields

$$P_\tau(L,\beta_j) \ge [P_\tau(2n_j,\beta_j)P_{\mathrm{s}}(n_j,\beta_j)^2]^{L/(2n_j)}$$
$$\ge (1-\varepsilon_j)^3 \ge 1 - 3\varepsilon_j. \tag{11}$$

Consequently the ground state is unique. Moreover, $3\varepsilon_j/(1-3\varepsilon_j) \le (60/97)20^{-2^j} < 20^{-2^j}$. Equation (2) therefore proves

$$\Delta_L > \frac{\log 20}{784} \simeq 0.003821 \qquad \text{for every even } L \ge 2304. \tag{12}$$

Finite seed bounds.—
The starting lemma requires two ratios at $\beta_0 = 49/4$:

$$P_{\mathrm{s}}(36,\beta_0) = \frac{Z_{72}(\beta_0)}{Z_{36}(\beta_0)^2},$$
$$P_\tau(72,\beta_0) = \frac{Z_{72}(2\beta_0)}{Z_{72}(\beta_0)^2}. \tag{13}$$




FIG. 1. The asymmetric recurrence and its spin-1 seed. The shaded boundary is numerical, not a rigorous region certificate. The plotted orbit is rationally verified; labels give selected lengths $n_k = 36\,2^k$, with $\beta_k = (49/4)2^k$. After six steps it enters the gray square $[0, 0.01]^2$. No certified inference uses the shaded region.

Each numerator needs a lower bound; each denominator needs an upper bound. Exact trace calculations on 4–12 sites and a 72-site trial state supply them [7]:

	Quantity	Bound	Method
	$Z_{72}(\beta_0)$	lower	MPS + small-ring traces
	$Z_{36}(\beta_0)$	upper	MPS + small-ring traces
	$Z_{72}(2\beta_0)$	lower	MPS
	$Z_{72}(\beta_0)$	upper	small-ring traces

Readers may skip this explanation and use Lemma 1 directly. Defining decimals below are exact rationals; approximate values are only explanatory.

Step 1: Spatial trace moments. A continuous-time, one-site spatial transfer operator gives

$$Z_n(\beta) = \operatorname{Tr} K_\beta^n = \quad , \qquad n \geq 4. \tag{14}$$

It is self-adjoint but need not be positive, unlike the positive two-site transfer matrix used in the doubling argument. Anticipating the trial energy below, set $\kappa = 700741/500000$ and

$$\widehat{Z}_n(\beta) = e^{-n\beta\kappa} Z_n(\beta), \qquad \widehat{K} = e^{-\beta_0\kappa} K_{\beta_0}, \qquad \widehat{Z}_n(\beta_0) = \sum_i \lambda_i^n. \tag{15}$$

The λ_i are real eigenvalues of $\widehat{K}$, possibly negative. The shift cancels from both purities. Changing n changes only the power: short-chain traces constrain the same spectrum needed at 36 and 72 sites.

Step 2: Separate the moments by symmetry. Spin rotations preserve the Heisenberg Hamiltonian and induce transfer symmetries. The commuting π rotations split four sign sectors: an invariant sector J and three sectors with identical spectra, denoted N, exchanged by axis permutations. Thus

$$\widehat{Z}_n = J_n + 3N_n, \qquad \widehat{Z}_n(P) = J_n - N_n. \tag{16}$$

Here $J_n = \sum_{i \in J} \lambda_i^n$, and N_n sums one N list; all traces are at β_0. The twist P rotates one end of the physical closing bond by π, rather than inserting a global rotation into the untwisted physical trace.

Integer trace recursions with explicit truncation and rounding bounds enclose both traces for $4 \leq n \leq 12$. Inverting Eq. (16) gives sector moments, including

$$J_{12} \leq m_J = 1.0657205, \qquad N_{12} \leq m_N = 0.2783155. \tag{17}$$

Step 3: A trial state gives a lower bound. Integer tensors of bond dimension 26 define a periodic 72-site MPS ψ. Its norm and one-bond expectation are exact contractions:


$$\mathcal{N} = \langle \psi, \psi \rangle = \quad , \qquad \mathcal{E} = \langle \psi, h_{1,2}\psi \rangle = \quad . \tag{18}$$

Each diagram contains 72 tensors per layer. Translation invariance gives $\langle \psi, H_{72}\psi \rangle/(72\mathcal{N}) = \mathcal{E}/\mathcal{N} < -\kappa$. The variational principle gives $E_0(72) < -72\kappa$. The shift adds κ on each of the 72 sites, so the shifted ground energy is negative and its Boltzmann weight exceeds one:

$$\widehat{Z}_{72}(\beta) \geq e^{-\beta(E_0(72)+72\kappa)} > 1 \qquad (\beta > 0). \tag{19}$$

This calculation supplies both $\sum_i \lambda_i^{72} > 1$ at β_0 and the thermal numerator $\widehat{Z}_{72}(2\beta_0) > 1$.

Step 4: A lower bound on the leading spatial eigenvalue. Suppose every $|\lambda_i| \leq \ell = 0.999$. Write $u_i = \lambda_i^{12} \geq 0$, so $\lambda_i^{72} = u_i^6$. In J, the total mass is at most m_J and each entry is at most ℓ^{12}. Convexity increases the sixth-power sum when mass passes from a smaller entry to a larger one, up to the cap. Since $\ell^{12} < m_J < 2\ell^{12}$, the extremum has one entry ℓ^{12} and remainder $m_J - \ell^{12}$:



Taking sixth powers suppresses the remainder. Each N list contributes at most m_N^6, giving

$$\widehat{Z}_{72}(\beta_0) \leq \ell^{72} + (m_J - \ell^{12})^6 + 3m_N^6 < 0.932 < 1. \tag{20}$$

This contradicts Eq. (19). Hence some $|\lambda_0| > \ell$; it belongs to J because $m_N < \ell^{12}$.

Step 5: Polynomial bounds on the remaining spectrum. A nonnegative polynomial $F(x) = \sum_{j=4}^{12} f_j x^j$ converts known sector moments into individual eigenvalue bounds:

$$F(\lambda_i) \le \sum_{a\in\mathcal{S}} F(\lambda_a) = \sum_{j=4}^{12} f_j \mathcal{S}_j \le B, \qquad i \in \mathcal{S}. \tag{21}$$

Here $\mathcal{S}$ is J or N. Where $F(x) > B$, no eigenvalue can occur. Two filters give $|\lambda| < U = 1.00139$ in J and $|\lambda| < V = 0.872$ in N. The Supplemental Material [8] gives their coefficients and checks both exterior rays.

Removing λ_0 leaves J mass at most $m_J - \ell^{12}$. Since $V^{12} < m_N < 2V^{12}$, convexity bounds each N list by one entry at V^{12} and the remainder. For $k = 36, 72$, the residual moment $\sum_{i\neq 0} \lambda_i^k$ is therefore at most

$$R_k = (m_J - \ell^{12})^{k/12} + 3\left[V^k + (m_N - V^{12})^{k/12}\right]. \tag{22}$$

Retain λ_0^{72} in the spatial numerator and bound its denominator by $(\lambda_0^{36} + R_{36})^2$. For thermal purity, use the trial numerator and $\widehat{Z}_{72}(\beta_0) \le U^{72} + R_{72}$. Thus

$$\begin{aligned} P_{\mathrm{s}}(36, \beta_0) &\ge [1 + R_{36}/\ell^{36}]^{-2} = 1 - p, \\ P_{\tau}(72, \beta_0) &\ge [U^{72} + R_{72}]^{-2} = 1 - q. \end{aligned} \tag{23}$$

Exact rational comparisons give the starting bounds

$$p \le \frac{47916}{10^6}, \qquad q \le \frac{181521}{10^6}. \tag{24}$$

For orientation, $p \simeq 0.0479157710062$ and $q \simeq 0.181520168033$; these approximations are not recurrence inputs.

The same trace moments, the existing 120-site trial state, and temperature lifting give $\Delta_L > 0.07290$ for every even $L \ge 36$. The lifted bound covers $L \ge 60$; an unlifted estimate bridges $36 \le L \le 72$. The Supplemental Material [8] gives the proof, and the repository provides the certificate.

Reflection-positive models.— The argument requires locality and reflection positivity [11, 12], not spin-rotation symmetry. Consider a translation-invariant ring of L cells with Hamiltonian

$$H_L = \sum_{j=1}^{L} h_{j,j+1} + \sum_{j=1}^{L} o_j, \qquad j + L \equiv j. \tag{25}$$

Here h couples neighboring cells and o acts within a cell. Let the real matrix $F = F^{\mathsf{T}}$, $F^2 = I$, represent reflection within a cell; for a dimer, F exchanges the two spins. We take o real symmetric and require $FoF = o$, so the cell Hamiltonian respects this reflection. Across a bond, we require

$$-h = \sum_{\alpha} C_\alpha \otimes F C_\alpha F, \tag{26}$$

where each C_α is real and either symmetric or antisymmetric. This pairs each bond factor with its reflected copy. Contracting two neighboring cells therefore gives a positive Gram transfer matrix, as in Eq. (5); see details [8]. We can then use the same spatial purity inequalities and doubling argument as for the spin-1 chain.

An admissible seed at even n_0 then gives a unique ground state and a uniform gap for every even $L \ge n_0$, using finite-prefix interpolation and an analytic tail bound ✓L. Table I includes staggered-field chains [13], bond-alternating chains [14, 15], chains with single-ion anisotropy [16], and Heisenberg ladders [17]. Each model needs its own finite seed; the analytical argument is unchanged.

Formal verification.— The marker ✓L denotes a formalized implication with its stated assumptions. The analytic criterion and recurrence are formalized in Lean. The path-space kernel construction and trace identities needed for the original proof are not yet available in mathlib. We therefore use finite matrices. A positive transfer matrix at finite M gives closed scalar inequalities; a controlled product limit at fixed L, β passes them to exact quantities before iteration ✓L.

We prove the gap theorem in Lean for general reflection-positive Hamiltonians and apply the same theorem to the models in Table I. We provide the proofs as a library so that others can reproduce the results and add models.

To obtain a complete Lean proof, we must also verify the finite calculations in Lean. We have done so for the mixed-field Ising model at $(g, h) = (1/4, 1/20)$. For spin one, we check the MPS contractions in Lean and supply the remaining externally certified finite bounds as hypotheses.

Discussion.— Using diagrams and visual summaries [18] to expose the structure of the proof let us certify gaps for seven other spin-chain and ladder models (Table I), and improve the spin-1 bound using the existing finite inputs. We also explain the boundary-field proof for open chains using diagrams [8, 19].

A next target is the spin-2 Heisenberg chain, whose smaller gap and longer correlation length make the finite inputs harder to certify [20]. In two spatial dimensions, an analogous argument would track three purities and require additional spatial inequalities as well as certified finite inputs. Establishing these inputs for the honeycomb AKLT model [21, 22] remains open for this method.

The OpenAI proof combines simple ideas with a substantial computation. This work also explores numerical certification in Lean and calls for improvements in the numerical efficiency of proof assistants, making them comparable with symbolic computation systems such as Mathematica and numerical linear algebra engines such as those in MATLAB and Julia. The fully checked Ising seed shows that the finite calculation can be included in the formal proof; the larger spin-1 calculations provide

TABLE I. Uniform gap bounds for reflection-positive models. Every even length at or above the stated threshold is covered. A cell or rung contains two spins; L counts the units shown. Spin operators have $\boldsymbol{S}^2 = S(S+1)I$; σ^α are Pauli matrices. Numerical values are finite-ring estimates on 12–20 spins, not certified uniform bounds, except for the spin-1 Heisenberg bulk estimate [3]. The mixed-field Ising row gives the kernel-checked bound.

Model	Hamiltonian	Even lengths	Gap $\geq$	Numerical
Staggered field, $S = \frac{1}{2}$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1} - \sum_j (-1)^j S^z_j$	$L \geq 8$ sites	1.1489	1.874
Dimerized, $S = \frac{1}{2}$	$\sum_j [1 + (-1)^j/2]\, \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 4$ cells	0.5347	1.189
Dimerized, $S = 1$	$\sum_j [1 + 9(-1)^j/10]\, \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 2$ cells	0.5288	1.763
Single-ion anisotropy, $S = 1$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1} + 4\sum_j (S^z_j)^2$	$L \geq 12$ sites	0.8994	2.269
Heisenberg ladder, $S = \frac{1}{2}$	$\sum_{j,a=1,2} \boldsymbol{S}_{j,a} \cdot \boldsymbol{S}_{j+1,a} + 4\sum_j \boldsymbol{S}_{j,1} \cdot \boldsymbol{S}_{j,2}$	$L \geq 4$ rungs	1.1024	3.138
Ising ladder	$-\sum_j (\sigma^z_{j,1}\sigma^z_{j+1,1} + \sigma^z_{j,2}\sigma^z_{j+1,2} + \sigma^z_{j,1}\sigma^z_{j,2}) - 3\sum_{j,a=1,2} \sigma^x_{j,a}$	$L \geq 4$ rungs	0.6368	2.662
Mixed-field Ising	$-\sum_j \sigma^z_j \sigma^z_{j+1} - \frac{1}{4}\sum_j \sigma^x_j - \frac{1}{20}\sum_j \sigma^z_j$	$L \geq 6$ sites	0.3922	1.190
Heisenberg, $S = 1$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 36$ sites	0.0729	0.410

a further test of how much computation can be checked directly in the kernel.

We thank Xiao-Liang Qi and Ignacio Cirac for helpful discussions. Part of this work was carried out at the Kavli Institute for Theoretical Physics and during a visit to Xiao-Liang Qi at Stanford University; we thank both institutions for their hospitality. An initial version of this work [23] was circulated and discussed among colleagues at Stanford on October 7, 2026, and we thank Shengqi Sang, Yimu Bao, and Junkai Dong for helpful discussions. A presentation of this work was given at KITP [24]. The code, gap certificates, Lean 4 formalization and blueprint, interactive explainer, and open challenges are available at https://github.com/LionSR/CertifiedGaps (see also https://sirui-lu.com/CertifiedGaps/).

AI-use declaration.—S. L. prompted GPT-6-Astra (OpenAI) to recast the proofs of Refs. [7, 19] in tensor-network diagrams, and used TeXRA [25] and OpenAI Codex with GPT-6-Astra and GPT-6-Sol for subsequent calculations, formalization, and manuscript preparation. All mathematical arguments, numerical certificates, and formal proofs were verified by the author.

[1] F. D. M. Haldane, Phys. Lett. A **93**, 464 (1983).
[2] F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).
[3] S. R. White and D. A. Huse, Phys. Rev. B **48**, 3844 (1993).
[4] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).
[5] S. Knabe, J. Stat. Phys. **52**, 627 (1988).
[6] D. Gosset and E. Mozgunov, J. Math. Phys. **57**, 091901 (2016), arXiv:1512.00088 [quant-ph].
[7] OpenAI, The periodic spin-one Haldane gap, OpenAI Math Release, result 268, preprint (2026).
[8] See Supplemental Material for the open-chain proof, the finite-dimensional transfer construction and periodic bootstrap, and the finite seed bounds and temperature-lifting estimates.
[9] M. Suzuki, Phys. Rev. B **31**, 2957 (1985).
[10] F. J. Dyson, E. H. Lieb, and B. Simon, J. Stat. Phys. **18**, 335 (1978).
[11] J. Fröhlich, R. Israel, E. H. Lieb, and B. Simon, Commun. Math. Phys. **62**, 1 (1978).
[12] A. Jaffe and B. Janssens, Commun. Math. Phys. **346**, 1021 (2016), arXiv:1506.04197 [math-ph].
[13] M. Oshikawa and I. Affleck, Phys. Rev. Lett. **79**, 2883 (1997).
[14] T. Barnes, J. Riera, and D. A. Tennant, Phys. Rev. B **59**, 11384 (1999).
[15] S. Yamamoto, Phys. Rev. B **51**, 16128 (1995).
[16] W. Chen, K. Hida, and B. C. Sanctuary, Phys. Rev. B **67**, 104401 (2003).
[17] E. Dagotto and T. M. Rice, Science **271**, 618 (1996).
[18] E. R. Tufte, *Envisioning Information* (Graphics Press, Cheshire, Connecticut, 1990).
[19] OpenAI, A boundary-field gap for the spin-one Heisenberg chain, OpenAI Math Release, result 268, preprint (2026).
[20] S. Todo and K. Kato, Phys. Rev. Lett. **87**, 047203 (2001).
[21] M. Lemm, A. W. Sandvik, and L. Wang, Phys. Rev. Lett. **124**, 177204 (2020), arXiv:1910.11810 [quant-ph].
[22] N. Pomata and T.-C. Wei, Phys. Rev. Lett. **124**, 177203 (2020), arXiv:1911.01410 [cond-mat.stat-mech].
[23] S. Lu and GPT-6-Astra, A short diagrammatic explanation of the spin-1 Haldane gap proof (2026), October 7, 2026.
[24] S. Lu, Bridging gaps with tensor networks: The Haldane gap proof and beyond, Talk at the KITP program *Intermediate-Scale AI for Physics* (2026), October 20, 2026 (video recording).
[25] TeXRA, TeXRA: A multi-agent research assistant for theorists (2025).
[26] T. Kennedy, J. Phys.: Condens. Matter **2**, 5737 (1990).
[27] W. Tang, H.-H. Tu, and L. Wang, Phys. Rev. Lett. **125**,

170604 (2020), arXiv:2004.12928 [cond-mat.str-el].

Supplemental Material

CONTENTS

SI. Open chains with boundary fields 7
SI.A. The boundary-field chain and the additional estimate 7
SI.B. Boundary vectors and boundary conditions 9
SI.C. Finite inputs and ring estimates 10
SI.D. Three steps for the open chain 11
a. Step 1: control the new boundary remainder 12
b. Step 2: double the length 13
c. Step 3: square in imaginary time 13
SI.E. Initialization and the small-defect induction 13
SI.F. Equal fields, odd lengths, and the gap 14

SII. Finite Trotter networks and the periodic doubling argument 15
SII.A. Finite Trotter slices and a two-site transfer matrix 15
SII.B. Finite-transfer moment inequalities 17
SII.C. Controlled product limit 18
SII.D. Torus identities and asymmetric doubling 18
SII.E. Six exact rational steps 19
SII.F. Scalar tail, interpolation, and the strict gap 20

SIII. Finite seed certificates and improved bounds 21
SIII.A. The base seed at length 36 21
a. Hamiltonians, twists, and the spatial transfer operator 21
b. Exact finite trace and trial inputs 22
c. Sector moments and polynomial filters 24
d. Capped masses and the two base defects 25
SIII.B. Temperature lifting and the improved uniform bound 25
a. Lifted moments and a dominant spatial weight 25
b. The physical thermal residual 26
c. Finite-prefix coverage and the doubling tail 27
SIII.C. Reproducibility and arithmetic assumptions 28

Appendix SI: Open chains with boundary fields

SI.A. The boundary-field chain and the additional estimate

We now extend the diagrammatic doubling argument from periodic rings to open chains with endpoint magnetic fields. Consider the spin-1 Heisenberg chain with equal endpoint fields,

$$H_n^{\mathrm{s}} = \sum_{i=1}^{n-1} \boldsymbol{S}_i \cdot \boldsymbol{S}_{i+1} - h(S_1^z + S_n^z), \qquad h = \frac{3}{5}, \tag{S1}$$

where $\sum_\alpha (S^\alpha)^2 = 2I$ and the exchange coupling is one. We give a diagrammatic proof of the following theorem from Ref. [19], using its certified finite inputs.

Theorem S1 (Boundary-field gap)**.** *For every odd $n \geq 1921$, H_n^{s} has a unique ground state and*

$$\Delta_n > \frac{\log 10}{392} \simeq 0.005874.$$

Without the endpoint fields, the effective spin-1/2 modes at the two ends produce four low-lying states whose splitting is exponentially small in the chain length [26]. Applying equal fields on an odd chain lifts this boundary degeneracy [19].

Our strategy is to bound the thermal purity of the open chain at a scale β proportional to the length, just as in the periodic argument. List the energies of H_n^{s} with multiplicity and let $w_0 \geq w_1 \geq \cdots$ be their Gibbs weights. With $P_\tau^{\mathrm{s}}(n,\beta) = \sum_m w_m^2$ and $\Delta_n = E_1(n) - E_0(n)$, we have


$$w_0 \geq P_\tau^{\mathrm{s}}(n,\beta), \qquad e^{-\beta\Delta_n} \leq \frac{1 - P_\tau^{\mathrm{s}}(n,\beta)}{P_\tau^{\mathrm{s}}(n,\beta)}. \tag{S2}$$

A thermal purity above 1/2 forces a unique ground state and bounds the gap from below. What changes in the open-chain setting is how we propagate the thermal purity across doubled lengths and inverse temperatures.

To see where the new difficulty enters, consider the spatial transfer operator at a fixed temperature. Suppose that X_β has a simple positive leading eigenvalue λ and write the normalized transfer operator as

$$T := X_\beta/\lambda = \Pi + R, \qquad \Pi = |\psi\rangle\langle\psi|, \qquad \Pi R = R\Pi = 0, \quad \|\psi\| = 1. \tag{S3}$$

In a periodic ring, the trace weights the leading projector Π by one. In an open chain, the chain ends are contracted with boundary vectors, so the leading contribution is weighted instead by a boundary overlap:



(S4)

Because c is not fixed by normalization and could a priori vanish, squaring in space no longer closes on its own. For an even length m, the normalized boundary matrix element splits into the leading overlap and a nonnegative remainder,

$$\begin{aligned} B_m &= \bar{v} \text{—} T^m \text{—} v \\ &= \bar{v} \text{—} \Pi \text{—} v + \bar{v} \text{—} R^m \text{—} v = c + D_m, \\ D_m &= \|R^{m/2} v\|^2 \geq 0. \end{aligned} \tag{S5}$$

To show that the leading eigenvector dominates B_m, we must bound the relative boundary remainder

$$e_m := \frac{D_m}{B_m} = 1 - \frac{c}{B_m}, \tag{S6}$$

rather than the absolute remainder D_m alone. Increasing the chain length suppresses D_m by powers of $\|R\|$ while leaving c unchanged.

When we double the inverse temperature, however, both the boundary vector v and the leading eigenvector ψ change. The overlap c at the old temperature does not by itself bound the new overlap c'. The key idea is to compare two even lengths at the new temperature. Writing primes for quantities at 2β, the difference

$$B'_k - B'_m = D'_k - D'_m$$

cancels the unknown overlap c' because c' is independent of length. An exact thermal-purity bridge then relates the ratio B'_k/B'_m back to known quantities at the old temperature.

The proof therefore proceeds in four stages:

1. In Sec. SI SI.B, we express the open-chain partition functions as boundary contractions of the continuous-time transfer operator. Putting the same boundary vector at both ends produces an auxiliary opposite-field chain with nonnegative remainders at even lengths, while a π spin rotation at one end restores equal fields.

2. In Sec. SI SI.C, we use the periodic doubling recurrence to control the bulk transfer spectrum and spectral ratio across all scales, and we state the four finite seed bounds needed to start the open-chain recurrence.

3. In Secs. SI SI.D and SI SI.E, we combine the two-length boundary comparison, length doubling, and thermal squaring into a coupled recurrence for e_m and the even-length opposite-field thermal purity, and we run three explicit initialization steps before entering the small-defect induction.

4. In Sec. SI SI.F, we pass from even opposite-field chains to odd equal-field chains by bounding the signed remainder in absolute value, completing the proof of Theorem S1.

SI.B. Boundary vectors and boundary conditions

We first represent the open-chain partition functions as transfer-operator matrix elements. We use the continuous-time bond expansion of Ref. [19]: we sum over the times of bond insertions instead of placing them on a finite Trotter grid. In the Cartesian basis, $G_\alpha = iS^\alpha$ is real antisymmetric and

$$-\boldsymbol{S}_i \cdot \boldsymbol{S}_{i+1} = \sum_{\alpha=x,y,z} G_{\alpha,i} G_{\alpha,i+1}.$$

A bond history a is an ordered list of events (τ, α) with $0 < \tau < \beta$. The measure μ_β sums the labels and integrates the ordered times; its total mass is $e^{3\beta}$, including unit mass on the empty history. The local tensor is

$$W_\beta^{ss'}(a,b) = e^{-\kappa\beta} \langle s| \mathcal{T}_\tau \prod_{(\tau,\alpha)\in a\cup b} G_\alpha |s'\rangle,$$
$$\kappa = \frac{700741}{500000}. \tag{S7}$$

Later events are placed to the left, so $W_\beta(a,b)$ depends on the relative event times in the two histories, not only on their event counts or on the measure. Tracing the physical indices gives the real symmetric transfer kernel $X_\beta(a,b)$ on $L^2(\mu_\beta)$.

The energy shift κ is the one used in Ref. [19]. Throughout this section, Z_n and all open-chain partition functions include the shift $\kappa n I$; it places the leading transfer eigenvalue near one at the seed scales and cancels from every purity and spectral gap.

To represent a boundary field along z, we cap the virtual bond history with the boundary vector

$$v_\beta(a) = \begin{cases} (ih)^{|a|}, & \text{all event labels are } z, \\ 0, & \text{otherwise,} \end{cases} \qquad \|v_\beta\|^2 = e^{\beta h^2}. \tag{S8}$$

Its value on the empty history is one, and its norm follows by summing $(\beta h^2)^k/k!$ over the number k of z events.

The cap inserts boundary events into the same time-ordered product as the bond events. Summing their counts and integrating their times adds the corresponding field term to the Hamiltonian. We now check the signs of the two endpoint fields produced by v_β. The conjugated left cap supplies $(-ih)G_z = +hS^z$ in the exponential, which corresponds to the Hamiltonian term $-hS_1^z$. The unconjugated right cap supplies $(ih)G_z = -hS^z$, corresponding to $+hS_n^z$. Contracting with the same boundary vector v_β at both ends therefore gives the auxiliary opposite-field chain

$$H_n^{\mathrm{o}} = H_n^{\mathrm{s}} + 2hS_n^z. \tag{S9}$$

To obtain equal endpoint fields, let $P = \mathrm{diag}(1,-1,-1)$ be the spin rotation by π about the x axis. Its virtual action U_P negates each z event, which reverses the right endpoint field and produces H_n^{s}. In diagram form,

$$e^{-\beta(H_n^{\mathrm{o}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } W_\beta \text{ — } \cdots \text{ — } W_\beta \text{ — } v_\beta,$$
$$Y_n(\beta) := \mathrm{Tr}\, e^{-\beta(H_n^{\mathrm{o}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } X_\beta^n \text{ — } v_\beta,$$
$$Y_n^{\mathrm{s}}(\beta) := \mathrm{Tr}\, e^{-\beta(H_n^{\mathrm{s}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } X_\beta^n \text{ — } U_P v_\beta. \tag{S10}$$

Tracing the physical legs in the first line gives the opposite-field partition function $Y_n(\beta)$ in the second; inserting U_P at the right cap gives the equal-field partition function $Y_n^{\mathrm{s}}(\beta)$ in the third.

Spin rotations act on bond histories by real orthogonal operators that commute with X_β, so $[U_P, X_\beta] = 0$. Regrouping the expansion into these site-by-site contractions is exact: before the energy shift, the absolute traced series is bounded by $3^n e^{\beta[3(n-1)+2h]} < \infty$, so Fubini's theorem justifies contracting the time-ordered expansion site by site.

Having expressed both open chains through X_β and v_β, we work first with the opposite-field partition function $Y_m(\beta)$ at even lengths m, where Eq. (S5) decomposes the boundary contraction into a sum of nonnegative terms. We will return to the odd equal-field chain $Y_n^{\mathrm{s}}(\beta)$ in Sec. SI SI.F, using an absolute remainder bound that does not require odd powers of the transfer operator to be positive.

SI.C. Finite inputs and ring estimates

We now identify the finite inputs required to drive the open-chain recurrence across the dyadic scales

$$\beta_j = 49\,2^{j-2}, \qquad n_j = 120\,2^{j-2}, \qquad M_j = 2n_j, \qquad j \geq 2, \tag{S11}$$

where n_j is a ring length and M_j is an even open-chain length. At each scale j, the two-length boundary comparison and the length-doubling step require three bulk properties of the transfer operator X_{β_j}: a simple positive leading eigenvalue λ_j invariant under U_P, a bound on the subleading norm $\|R_j\|^{n_j}$ to suppress the boundary remainder across the length difference $M_j - k_j$, and a lower bound on the bulk normalization ratio $r_j = \lambda_{j+1}/\lambda_j^2$ relating consecutive temperatures. All three bulk properties follow at every $j \geq 2$ from the periodic spatial and thermal purities $P_{\mathrm{s}}(n_2, \beta_2)$ and $P_\tau(M_2, \beta_2)$ at the initial scale $j = 2$ ($\beta_2 = 49$, $n_2 = 120$, $M_2 = 240$). To start the open-chain recurrence itself, we also need an initial bound on the boundary defect $e_{k_2}(2)$ at a shorter even length $k_2 = 142 < M_2$, together with a lower bound on the opposite-field thermal purity $P_\tau^{\mathrm{o}}(M_2, \beta_2) = Y_{M_2}(2\beta_2)/Y_{M_2}(\beta_2)^2$.

Ref. [19] certifies these four finite seed bounds at $j = 2$:

$$\begin{aligned} P_{\mathrm{s}}(120, 49) \geq .948, \qquad & P_\tau(240, 49) \geq .948, \\ e_{142}(2) \leq .14, \qquad & P_\tau^{\mathrm{o}}(240, 49) \geq .79. \end{aligned} \tag{S12}$$

Here $P_{\mathrm{s}}(n, \beta) = Z_{2n}(\beta)/Z_n(\beta)^2$ and $P_\tau(n, \beta) = Z_n(2\beta)/Z_n(\beta)^2$ concern the periodic ring, while $e_m(j)$, defined by Eq. (S6) at β_j, and $P_\tau^{\mathrm{o}}(m, \beta) = Y_m(2\beta)/Y_m(\beta)^2$ concern the opposite-field open chain.

We first use the two ring inputs in Eq. (S12) to control the bulk transfer operator at every scale $j \geq 2$. By the periodic doubling argument, the ring purities satisfy

$$\begin{aligned} P_{\mathrm{s}}(n_j, \beta_j), \ P_\tau(M_j, \beta_j) \geq 1 - \varepsilon_j, \\ \varepsilon_2 = .052, \qquad \varepsilon_{j+1} = 6\varepsilon_j^2. \end{aligned} \tag{S13}$$

To verify the factor of 6 in this scalar recurrence, recall the squaring inequality: if a positive trace-one operator of purity P is squared and renormalized, its new purity P' satisfies

$$1 - P' \leq f(1 - P), \qquad f(z) = \frac{z^2}{2(1-z)^2}. \tag{S14}$$

The periodic coupling identity first gives $P_{\mathrm{s}}(n, 2\beta) \geq (1-\varepsilon)^3$. For $0 \leq \varepsilon \leq .052$, we have

$$\begin{aligned} f(1 - (1-\varepsilon)^3) &= \frac{\varepsilon^2}{2}\left[\sum_{s=1}^{3}(1-\varepsilon)^{-s}\right]^2 \\ &\leq 6\varepsilon^2 \leq \varepsilon. \end{aligned} \tag{S15}$$

After spatial squaring bounds the new spatial defect by $6\varepsilon^2$, the second periodic coupling identity gives $P_\tau(4n, \beta) \geq (1-\varepsilon)^2(1-6\varepsilon^2) \geq (1-\varepsilon)^3$. Thermal squaring then bounds the new thermal defect by the same $6\varepsilon^2$, proving Eq. (S13) by induction on $j \geq 2$ without waiting for the smaller open-chain threshold established below.

From the ring purities in Eq. (S13), we now extract the bulk spectral properties of X_{β_j} needed for the open chain. Write λ_j for the leading eigenvalue of X_{β_j}, and decompose $T_j = X_{\beta_j}/\lambda_j = \Pi_j + R_j$ as in Eq. (S3). We claim that for every $j \geq 2$,

$$\begin{aligned} \lambda_j > 0 \text{ is simple}, \qquad & \|R_j\|^{n_j} \leq d(\varepsilon_j), \\ & d(\varepsilon) = \frac{\varepsilon}{2 - 3\varepsilon}, \qquad \|R_j\| < 1, \\ & r_j := \frac{\lambda_{j+1}}{\lambda_j^2}, \qquad 0 < r_j \leq 1, \\ & r_j^{M_j} \geq (1-\varepsilon_j)(1-\varepsilon_{j+1}), \qquad U_P \Pi_j = \Pi_j. \end{aligned} \tag{S16}$$

We verify each assertion in Eq. (S16) in turn. For any probability distribution of purity at least $1 - \varepsilon > 1/2$, let p_0 be its largest weight and put $z = 1 - p_0$. Since p_0 bounds the purity from above, $z < 1/2$, and $1 - \varepsilon \leq p_0^2 + (1-p_0)^2 = 1 - 2z(1-z)$ gives $2z(1-z) \leq \varepsilon$. Hence

$$z \leq \varepsilon, \qquad z \leq \frac{\varepsilon}{2(1-\varepsilon)}, \qquad \frac{z}{1-z} \leq d(\varepsilon).$$

Applying this estimate to the normalized weights $p_i = |\lambda_i(\beta_j)|^{n_j}/Z_{n_j}(\beta_j)$ yields the subleading tail bound

$$\sum_{i\geq 1}\left|\frac{\lambda_i(\beta_j)}{\lambda_j}\right|^{n_j} \leq d(\varepsilon_j). \tag{S17}$$

Because $d(\varepsilon_j) < 1$, the largest-modulus eigenvalue λ_j is simple and $\|R_j\|^{n_j} \leq d(\varepsilon_j) < 1$. Moreover, λ_j is positive: because X_{β_j} is Hilbert–Schmidt and the remaining spectrum has strictly smaller spectral radius, a negative isolated leading eigenvalue would make $\mathrm{Tr_v}\, X_{\beta_j}^N$ negative for all sufficiently large odd N, contradicting $Z_N(\beta_j) > 0$.

Next, to see that $U_P\Pi_j = \Pi_j$, note that the three π rotations about the coordinate axes and the cyclic permutation of those axes commute with the real transfer kernel X_{β_j}. Each acts by a sign on the real one-dimensional leading eigenspace. The order-three cyclic permutation must act by $+1$, and it conjugates the three π rotations into one another. Their signs are therefore equal; since the product of two π rotations is the third, the common sign is $+1$, which gives $U_P\Pi_j = \Pi_j$.

Finally, to bound the bulk normalization ratio $r_j = \lambda_{j+1}/\lambda_j^2$, observe that $P_\tau(N, \beta_j) \leq 1$ and taking even Nth roots as $N \to \infty$ give $\lambda_{j+1} \leq \lambda_j^2$, so $r_j \leq 1$. At length $M_j = n_{j+1}$, combining the leading spatial weight at β_{j+1} with the ring thermal purity at β_j gives the lower bound

$$\begin{aligned}\lambda_{j+1}^{M_j} &\geq (1-\varepsilon_{j+1})Z_{M_j}(\beta_{j+1}) \\ &\geq (1-\varepsilon_{j+1})(1-\varepsilon_j)Z_{M_j}(\beta_j)^2 \\ &\geq (1-\varepsilon_{j+1})(1-\varepsilon_j)\lambda_j^{2M_j}.\end{aligned} \tag{S18}$$

Dividing by $\lambda_j^{2M_j}$ proves $r_j^{M_j} \geq (1-\varepsilon_j)(1-\varepsilon_{j+1})$. With the bulk transfer estimates in Eq. (S16) established, we turn to the open-chain recurrence.

SI.D. Three steps for the open chain

We now set up the open-chain quantities at scale j and show how to advance them to scale $j+1$. Set $v_j = v_{\beta_j}$ and, for even $m \geq 2$, define the normalized opposite-field partition function, its remainder, and the boundary defect by

$$\begin{aligned} c_j &= |\langle \psi_j, v_j\rangle|^2, \\ B_m(j) &= \frac{Y_m(\beta_j)}{\lambda_j^m} = c_j + D_m(j) > 0, \\ D_m(j) &= \|R_j^{m/2} v_j\|^2, \qquad e_m(j) = \frac{D_m(j)}{B_m(j)}. \end{aligned} \tag{S19}$$

For even lengths $m \geq k$, absorbing $R_j^{k/2}$ into each boundary cap gives the spatial decay of the remainder,


$$\begin{aligned} D_m(j) &= \bar{y} \text{ — } \boxed{R_j^{m-k}} \text{ — } y, \qquad y = R_j^{k/2} v_j, \\ 0 \leq D_m(j) &\leq \|R_j\|^{m-k} D_k(j). \end{aligned} \tag{S20}$$

In particular, $D_m(j)$ is nonincreasing in even m, so $c_j \leq B_m(j) \leq B_k(j)$.

To connect quantities at β_j and $\beta_{j+1} = 2\beta_j$, we use the opposite-field thermal purity. By the definition of $B_m(j)$ and $r_j = \lambda_{j+1}/\lambda_j^2$, we have the exact thermal-purity bridge

$$\begin{aligned} P_\tau^{\mathrm{o}}(m, \beta_j) &= \frac{Y_m(\beta_{j+1})}{Y_m(\beta_j)^2} \in (0, 1], \\ B_m(j+1) &= P_\tau^{\mathrm{o}}(m, \beta_j) r_j^{-m} B_m(j)^2. \end{aligned} \tag{S21}$$

This identity comes from multiplying the full open-chain Gibbs operators in imaginary time, not from a sitewise fusion of local MPO tensors. Writing $\rho_m^{\mathrm{o}}(\beta) = e^{-\beta(H_m^{\mathrm{o}} + \kappa m I)}/Y_m(\beta)$, the imaginary-time contraction is




(S22)

Assuming bounds on a boundary defect $e_k(j)$ and on the thermal purity $P^{\circ}_{\tau}(m,\beta_j)$ at scale j, we advance to scale $j+1$ in three steps: first we bound the boundary defect $e_m(j+1)$ at the new temperature; second we use that new boundary defect to double the chain length from m to $2m$ in the thermal purity at β_j; third we square in imaginary time to obtain $P^{\circ}_{\tau}(2m,\beta_{j+1})$.

a. Step 1: control the new boundary remainder

Our first task is to bound the boundary defect $e_m(j+1)$ at the new inverse temperature β_{j+1} without knowing the new leading overlap c_{j+1} in advance. Fix even lengths $m > k \geq 2$, assume $e_k(j) < 1$, and write primes for quantities at scale $j+1$. Choose a suppression factor $q < 1$ satisfying $\|R_{j+1}\|^{m-k} \leq q$, so that Eq. (S20) at scale $j+1$ gives $D'_m \leq qD'_k$. Subtracting the normalized boundary contractions at lengths k and m at the same temperature β_{j+1} cancels their common leading term c':



(S23)

That is, $B'_k - B'_m = D'_k - D'_m$. Combining $D'_k = D'_m + B'_k - B'_m$ with $D'_m \leq qD'_k$ gives

$$\begin{aligned}(1-q)D'_m &\leq q(B'_k - B'_m),\\ e'_m &\leq \frac{q}{1-q}\left(\frac{B'_k}{B'_m} - 1\right).\end{aligned} \tag{S24}$$

The unknown overlap c' has dropped out of the difference $B'_k - B'_m$.

We can now bound the new-temperature ratio B'_k/B'_m in terms of scale-j quantities by applying the bridge identity (S21) at both lengths k and m:

$$\begin{aligned}\frac{B'_k}{B'_m} &= \frac{P^{\circ}_{\tau}(k,\beta_j)}{P^{\circ}_{\tau}(m,\beta_j)} r_j^{m-k}\left[\frac{B_k(j)}{B_m(j)}\right]^2\\ &\leq \frac{1}{P^{\circ}_{\tau}(m,\beta_j)[1-e_k(j)]^2}.\end{aligned} \tag{S25}$$

Here we used $P^{\circ}_{\tau}(k,\beta_j) \leq 1$, $r_j \leq 1$, and the lower bound

$$B_m(j) \geq c_j = [1-e_k(j)]B_k(j) > 0$$

at the old temperature. Equation (S25) makes no assumption on the new overlap c'; once Eqs. (S24) and (S25) yield $e'_m < 1$, it follows automatically that $c' = (1-e'_m)B'_m > 0$.

Notice why Eq. (S24) produces a quadratic improvement in the small-defect regime. It is a product of two small factors: the spatial decay factor $q/(1-q)$ over the length difference $m-k$, and the excess of the ratio B'_k/B'_m above one. Specifically, when $m = M_j$, $k = M_j/2$, and $\varepsilon_j \leq x \leq .01$, the ring bound (S16) at scale $j+1$ gives

$$\|R_{j+1}\|^{M_j} \leq d(\varepsilon_{j+1}) \leq d(6x^2).$$

Across the half-chain $m-k = M_j/2$, we may therefore take

$$q = \sqrt{d(\varepsilon_{j+1})} \leq \sqrt{\frac{6x^2}{2-18x^2}} \leq 2x. \tag{S26}$$

Separately, when $e_k(j) \leq x$ and $P^{\circ}_{\tau}(m,\beta_j) \geq 1-x$, Eq. (S25) bounds the excess above one by

$$\frac{B'_k}{B'_m} - 1 \leq (1-x)^{-3} - 1 = O(x). \tag{S27}$$

Multiplying Eqs. (S26) and (S27) bounds e'_m by $O(x^2)$. The subtraction of one in Eq. (S24), coming from the cancellation of c' in Eq. (S23), supplies this second factor of x.

b. *Step 2: double the length*

Having bounded the boundary defect $e_M(j+1)$ at the new temperature, we next double the open-chain length from M to $2M$ in the thermal purity at the old temperature β_j. For any even M, dividing the bridge identity (S21) at length $2M$ by that at length M gives

$$\begin{aligned}\frac{P_\tau^o(2M,\beta_j)}{P_\tau^o(M,\beta_j)} &= r_j^M \frac{B_{2M}(j+1)}{B_M(j+1)}\left[\frac{B_M(j)}{B_{2M}(j)}\right]^2 \\ &\geq r_j^M[1-e_M(j+1)]. \end{aligned} \tag{S28}$$

In the last factor, monotonicity at the old temperature gives $B_M(j) \geq B_{2M}(j)$, so $[B_M(j)/B_{2M}(j)]^2 \geq 1$. In the middle factor, however, $B_{2M}(j+1) \leq B_M(j+1)$, so we must use the new-temperature boundary defect bound from Step 1:

$$B_{2M}(j+1) \geq c_{j+1} = [1-e_M(j+1)]B_M(j+1).$$

Equation (S28) therefore retains both the bulk spectral-ratio factor r_j^M, controlled by Eq. (S16), and the new-temperature boundary factor $1-e_M(j+1)$.

c. *Step 3: square in imaginary time*

Once Step 2 bounds $P_\tau^o(2M,\beta_j)$ at the doubled length $2M$, we double the inverse temperature from β_j to β_{j+1} by squaring the open-chain Gibbs state in imaginary time. Just as in the periodic argument, Eqs. (S22) and (S14) give

$$1-P_\tau^o(2M,\beta_{j+1}) \leq f(1-P_\tau^o(2M,\beta_j)). \tag{S29}$$

Notice why the three steps must be performed in this order: because $P_\tau^o(M,\beta_j)$ already connects β_j and β_{j+1}, doubling the length in Step 2 requires the new-temperature boundary defect $e_M(j+1)$ from Step 1 before we can square in imaginary time in Step 3.

SI.E. Initialization and the small-defect induction

We now assemble the three steps of Sec. SI SI.D into an explicit recurrence across the scales $j \geq 2$. We track the boundary defect at the even lengths

$$k_2 = 142, \qquad k_{j+1} = M_j. \tag{S30}$$

Since $M_{j+1} = 2M_j$, this choice gives $k_j = M_j/2$ for all $j \geq 3$. Let E_j and η_j denote upper bounds on $e_{k_j}(j)$ and $1-P_\tau^o(M_j,\beta_j)$, respectively. By Eqs. (S16), (S24), (S25), (S28), and (S29), the updates from scale j to scale $j+1$ take the form

$$\begin{aligned} F(q,e,t) &= \frac{q}{1-q}\left[\frac{1}{(1-e)^2 t}-1\right], \\ q_j &= d(\varepsilon_{j+1})^{(M_j-k_j)/M_j}, \\ E_{j+1} &\geq F(q_j,E_j,1-\eta_j), \\ A_j &= (1-\eta_j)(1-\varepsilon_j)(1-\varepsilon_{j+1})(1-E_{j+1}), \\ \eta_{j+1} &\geq f(1-A_j), \end{aligned} \tag{S31}$$

valid whenever $E_j, \eta_j, E_{j+1}, \eta_{j+1} \in [0,1)$. The product A_j bounds $P_\tau^o(2M_j,\beta_j)$ from below by Eq. (S28); thermal squaring then gives the update for η_{j+1}. Here the exponent $(M_j-k_j)/M_j$ in q_j comes from $n_{j+1} = M_j$ and the new-temperature ring remainder bound $\|R_{j+1}\|^{M_j} \leq d(\varepsilon_{j+1})$. Because $F(q,e,t)$ increases with q and e and decreases with t, any upper bound on q_j may be substituted into the update for E_{j+1}.

At the initial scale $j=2$, the seed defects $E_2 = .14$ and $\eta_2 = .21$ from Eq. (S12) are too large to enter the analytical small-defect induction ($x \leq .01$) directly. We therefore carry out three explicit rational steps from $j=2$ to $j=5$. Starting with $\varepsilon_2 = .052$, $E_2 = .14$, and $\eta_2 = .21$, Eq. (S31) produces the outward-rounded rational bounds in the following table:

j	E_j	η_j	upper bound for q_j
2	.14	.21	.142
3	.12	.155	.0282
4	.016	.026	.0028
5	.001	.001	—

All decimals in this table denote exact rationals. At $j = 2$, the first suppression bound uses $k_2 = 142$ and $M_2 = 240$, so that

$$\frac{M_2 - k_2}{M_2} = \frac{49}{120}, \qquad d(\varepsilon_3)^{49} < .142^{120}.$$

At $j = 3$ and $j = 4$, we have $(M_j - k_j)/M_j = 1/2$, so we verify the upper bounds on q_j by squaring .0282 and .0028 and comparing with $d(\varepsilon_4)$ and $d(\varepsilon_5)$. Substituting these values into Eq. (S31) verifies each row of the table by exact rational arithmetic. Together with $\varepsilon_5 < .000015$, the $j = 5$ row brings all three defects below .01, establishing the base case of the small-defect induction:

$$\varepsilon_j,\ e_{k_j}(j),\ 1 - P^{\mathrm{o}}_\tau(M_j, \beta_j) \le x_j, \qquad j \ge 5,$$
$$x_5 = .01, \qquad x_{j+1} = 10x_j^2, \qquad x_j = 10^{-1-2^{j-5}}. \tag{S32}$$

To prove the induction step in Eq. (S32), assume that the three bounds hold at scale $j \ge 5$ and write $x = x_j \le .01$. For Step 1, Eqs. (S26) and (S27) give $q_j \le 2x$ and


$$F(2x, x, 1-x) \le \frac{6x^2}{(1-2x)(1-x)^3} < 7x^2. \tag{S33}$$

Thus $E_{j+1} = 10x^2$ is a valid upper bound on $e_{M_j}(j+1)$. For Step 2, substituting $\eta_j \le x$, $\varepsilon_j \le x$, $\varepsilon_{j+1} \le 6x^2$, and $E_{j+1} \le 10x^2$ into A_j gives $A_j \ge 1 - 2x - 16x^2$. For Step 3, since $0 < x \le .01$, thermal squaring yields

$$f(2x + 16x^2) < 3x^2. \tag{S34}$$

Hence $\eta_{j+1} = 10x^2$ is a valid upper bound on $1 - P^{\mathrm{o}}_\tau(2M_j, \beta_{j+1})$, and $\varepsilon_{j+1} \le 6x^2 \le 10x^2$ closes the induction at scale $j+1$.

The following diagram summarizes one induction step from scale j to scale $j+1 = j'$ ($M_{j+1} = 2M_j$):




SI.F. Equal fields, odd lengths, and the gap

We have established the small-defect bounds (S32) for even-length opposite-field chains at every scale $j \ge 5$. It remains to transfer these bounds to the target equal-field Hamiltonian H^{s}_n at arbitrary odd lengths $n \ge M_5 + 1 = 1921$ and extract the spectral gap.

Define the normalized equal-field partition function by


$$Q_j(n) = \frac{Y^{\mathrm{s}}_n(\beta_j)}{\lambda_j^n}.$$

Because U_P commutes with X_{β_j} and satisfies $U_P\Pi_j = \Pi_j$ by Eq. (S16), inserting U_P at the right cap preserves the leading contribution c_j:



(S35)

For any even $M \le n$, we split off $R_j^{M/2}$ on each side of the remainder contraction to obtain



$$\begin{aligned} Q_j(n) - c_j &= \\ &= \langle y, R_j^{n-M} U_P y \rangle, \qquad y = R_j^{M/2} v_j. \end{aligned} \tag{S36}$$

Because $\|R_j\| \le 1$ and U_P is orthogonal, we have $\|R_j^{n-M}U_P\| \le 1$, and Cauchy–Schwarz bounds the signed remainder in absolute value by the even-length opposite-field remainder $D_M(j)$:

$$|Q_j(n) - c_j| \le \|y\|^2 = D_M(j). \tag{S37}$$

This inequality controls $Q_j(n)$ at odd n without assuming that odd powers of R_j are positive.

Now fix a scale $j \ge 5$ and an odd length n in the interval $M_j \le n \le 2M_j$, and set $M = M_j = k_{j+1}$. Applying Eq. (S37) at β_j and β_{j+1} bounds $Q_j(n)$ from above and $Q_{j+1}(n)$ from below in terms of B_M:

$$\begin{aligned} Q_j(n) &\le c_j + D_M(j) = B_M(j), \\ Q_{j+1}(n) &\ge c_{j+1} - D_M(j+1) \\ &= [1 - 2e_M(j+1)] B_M(j+1) \\ &\ge (1 - 2x_{j+1}) B_M(j+1) > 0. \end{aligned} \tag{S38}$$

Combining Eq. (S38) with the bridge identity (S21) at length M, and using $r_j \le 1$ and $n - M \le M$ together with the bulk ratio bound $r_j^M \ge (1-\varepsilon_j)(1-\varepsilon_{j+1})$ from Eq. (S16), we obtain the equal-field thermal purity bound

$$\begin{aligned} P_\tau^{\mathrm{s}}(n, \beta_j) &= r_j^n \frac{Q_{j+1}(n)}{Q_j(n)^2} \\ &\ge (1 - 2x_{j+1}) P_\tau^{\mathrm{o}}(M, \beta_j) r_j^{n-M} \\ &\ge (1 - 2x_{j+1})(1 - x_j)(1 - \varepsilon_j)(1 - \varepsilon_{j+1}) \\ &\ge 1 - 2x_j - 26x_j^2 \\ &\ge 1 - 4x_j, \end{aligned} \tag{S39}$$

where the coefficient $26 = 20 + 6$ collects $2x_{j+1} = 20x_j^2$ and $\varepsilon_{j+1} \le 6x_j^2$.

Proof of Theorem S1. The intervals $[M_j, 2M_j]$ for $j \ge 5$ cover every integer $n \ge M_5 = 1920$. Given any odd $n \ge 1921$, choose $j \ge 5$ with $M_j \le n \le 2M_j$. Since $x_j \le x_5 = .01$, Eq. (S39) gives $w_0 \ge P_\tau^{\mathrm{s}}(n, \beta_j) \ge 1 - 4x_j > 1/2$, so H_n^{s} has a unique ground state. At this length n and finite inverse temperature β_j, Eq. (S2) gives

$$\begin{aligned} e^{-\beta_j \Delta_n} &\le \frac{4x_j}{1 - 4x_j} \\ &\le \frac{5}{12} 10^{-2^{j-5}} < 10^{-2^{j-5}}, \end{aligned} \tag{S40}$$

where we used $x_j = 10^{-1}10^{-2^{j-5}}$ and $1 - 4x_j \ge .96$, which yields the prefactor $.4/.96 = 5/12 < 1$. Taking logarithms and dividing by $\beta_j = 392\, 2^{j-5}$ proves $\Delta_n > (\log 10)/392$. □

Appendix SII: Finite Trotter networks and the periodic doubling argument

SII.A. Finite Trotter slices and a two-site transfer matrix

OpenAI's proof uses a continuous-time spatial integral kernel [7]. For our Lean formalization we instead prove the spatial inequalities for finite matrices and pass them to exact partition functions by a controlled product limit.

Consider the periodic spin-1 Heisenberg Hamiltonian

$$H_L = \sum_{j=0}^{L-1} h_{j,j+1}, \qquad h = \sum_{\alpha=x,y,z} S^\alpha \otimes S^\alpha, \qquad Z_L(\beta) = \operatorname{Tr} e^{-\beta H_L}, \tag{S1}$$

with indices taken modulo L, exchange coupling equal to one, and $\beta > 0$. In the Cartesian basis $s, s' \in \{x, y, z\}$, set

$$(S^\alpha)_{ss'} = -i\epsilon_{\alpha ss'}, \qquad G_\alpha = iS^\alpha, \qquad -h = \sum_\alpha G_\alpha \otimes G_\alpha. \tag{S2}$$

Here $\epsilon_{\alpha ss'}$ is the Levi-Civita symbol, Tr denotes the matrix trace, and $\mathbb{I}$ denotes the identity on the relevant space. Each G_α is real antisymmetric and has operator norm one. Equation (S2) is the real bond-reflection-positive factorization used below [10, 11].

To replace the continuous-time path integral by a finite-dimensional trace, we discretize imaginary time into M brickwork slices. For $L = 2N$, $N \geq 1$, and a positive integer M, let $\tau = \beta/M$. Define the linearized brickwork slice and its trace by

$$\begin{aligned} S_{2N}(\tau) &= \prod_{k=0}^{N-1} (\mathbb{I} - \tau h_{2k,2k+1}) \prod_{k=0}^{N-1} (\mathbb{I} - \tau h_{2k+1,2k+2}), \\ Z_{2N}^{(M)}(\beta) &= \operatorname{Tr} S_{2N}(\beta/M)^M. \end{aligned} \tag{S3}$$

Products are ordered from left to right in increasing index. The trace is real because all factors have real entries.

Let $\Gamma = \{\varnothing, x, y, z\}$, with $C_\varnothing = \mathbb{I}$ and $C_\alpha = \sqrt{\tau}\, G_\alpha$. The bond expansion is the exact finite identity

$$\mathbb{I} - \tau h = \sum_{\gamma \in \Gamma} C_\gamma \otimes C_\gamma. \tag{S4}$$

An imaginary-time bond sequence is an element $p = (p_0, \ldots, p_{M-1})$ of $\mathcal{D}_M = \Gamma^M$. Define the single-site trace matrix by

$$\mathcal{F}_{pq} = \operatorname{Tr}\left[\prod_{t=0}^{M-1} (C_{p_t} C_{q_t}) \right], \qquad p, q \in \mathcal{D}_M. \tag{S5}$$

The product over t follows the multiplication order of the M slices. The matrix $\mathcal{F}$ depends on M and τ, but not on N.

To contract the network in space, assign sequences b_k and c_k to the even and odd bonds starting at $2k$ and $2k+1$, respectively. In slice t, site $2k$ carries $C_{b_{k,t}} C_{c_{k-1,t}}$, whereas site $2k+1$ carries $C_{b_{k,t}} C_{c_{k,t}}$. Expanding all bond sums, multiplying sitewise tensor products, and taking the physical trace gives

$$\begin{aligned} Z_{2N}^{(M)}(\beta) &= \sum_{\substack{b_0,\ldots,b_{N-1} \in \mathcal{D}_M \\ c_0,\ldots,c_{N-1} \in \mathcal{D}_M}} \prod_{k=0}^{N-1} \mathcal{F}_{b_k c_{k-1}} \mathcal{F}_{b_k c_k} \\ &= \sum_{c_0,\ldots,c_{N-1} \in \mathcal{D}_M} \prod_{k=0}^{N-1} (\mathcal{F}^{\mathsf{T}} \mathcal{F})_{c_{k-1} c_k} = \operatorname{Tr}(\mathcal{F}^{\mathsf{T}} \mathcal{F})^N, \end{aligned} \tag{S6}$$

where $c_{-1} = c_{N-1}$. The middle equality sums each even-bond sequence b_k independently; the last equality is the closed-path expansion of a matrix trace.

Thus the spatial transfer matrix

$$T_{M,\tau} = \mathcal{F}^{\mathsf{T}} \mathcal{F} = \mathcal{F}^\dagger \mathcal{F} \succeq 0 \tag{S7}$$

advances *two sites*, not one. Because $\mathcal{F}$ is real, its transpose equals its adjoint, and $v^\dagger T_{M,\tau} v = \|\mathcal{F} v\|^2 \geq 0$. Neither symmetry nor positivity of $\mathcal{F}$ itself is required. This finite two-site Gram matrix is distinct from the continuous-time one-site self-adjoint kernel used in a thermal-MPO description [27], and we do not construct a limiting kernel.

We extend the same contraction to any real reflection-positive cell model. Its assumptions are a finite-dimensional local cell space, a finite family of real matrices C_α, each symmetric or antisymmetric, a real symmetric on-site matrix o, and a real symmetric involution F satisfying

$$F^2 = \mathbb{I}, \qquad FoF = o, \qquad h = -\sum_\alpha C_\alpha \otimes FC_\alpha F. \tag{S8}$$

The Hamiltonian is $H_L = \sum_j h_{j,j+1} + \sum_j o_j$, and lengths count cells. Insert the on-site layer $\prod_i(\mathbb{I} - \tau o_i)$ between the even and odd bond layers. Writing $A_\varnothing = \mathbb{I}$, $A_\alpha = \sqrt{\tau}\, C_\alpha$, and $O = \mathbb{I} - \tau o$, the bond expansion becomes $\mathbb{I} - \tau h = \sum_\gamma A_\gamma \otimes FA_\gamma F$. Replace Eq. (S5) by

$$\mathcal{F}_{pq} = \mathrm{Tr} \prod_{t=0}^{M-1} (A_{p_t} O F A_{q_t} F), \tag{S9}$$

where the sequences now range over the enlarged bond-label alphabet. An odd cell carries $FA_{p_t}FOA_{q_t}$. Since $FO = OF$, this factor is the conjugate by F of the even-cell factor. Conjugating the entire product leaves its trace unchanged. The two sublattices therefore contribute the same $\mathcal{F}_{pq}$, and Eqs. (S6) and (S7) follow without change.

SII.B. Finite-transfer moment inequalities

Positivity of $T_{M,\tau}$ expresses $Z_L^{(M)}(\beta)$ as an even moment sum of nonnegative weights, from which we derive the spatial purity, squaring, and interpolation inequalities at finite M. At fixed M, β, let $\lambda_i \geq 0$ be the eigenvalues of $T_{M,\beta/M}$, and put $\mu_i = \sqrt{\lambda_i}$. For every even $L \geq 2$, Eq. (S6) gives

$$Z_L^{(M)}(\beta) = \sum_i \mu_i^L. \tag{S10}$$

For any finite nonnegative family a_i, let $S_k = \sum_i a_i^k$. Then

$$S_2 \leq S_1^2, \qquad 2(S_2^2 - S_4) \leq (S_1^2 - S_2)^2. \tag{S11}$$

Indeed, $S_1^2 - S_2 = 2\sum_{i<j} a_i a_j$, whereas $S_2^2 - S_4 = 2\sum_{i<j} a_i^2 a_j^2$. Squaring the first sum and retaining only its diagonal terms proves the second inequality.

If $\sum_i a_i = 1$, the purity is $P = \sum_i a_i^2 > 0$. Squaring and normalizing replaces the distribution by a_i^2/P, of purity $P' = S_4/P^2$. Thus

$$1 - P' \leq f(1 - P), \qquad f(z) = \frac{z^2}{2(1-z)^2}. \tag{S12}$$

The map f is nondecreasing on $0 \leq z < 1$, since $z/(1-z)$ is nonnegative and increasing there.

For $0 < u \leq v$, nonnegative weights also satisfy

$$\sum_i \mu_i^v \leq \left(\sum_i \mu_i^u \right)^{v/u}. \tag{S13}$$

If the sum on the right is zero, the claim is immediate. Otherwise normalize μ_i^u to probabilities a_i; then $\sum_i a_i^{v/u} \leq \sum_i a_i = 1$.

Applying Eqs. (S11) and (S13) to Eq. (S10) yields, at fixed β, the closed scalar inequalities

$$\begin{aligned} Z_{2n}^{(M)} &\leq (Z_n^{(M)})^2, \\ 2((Z_{2n}^{(M)})^2 - Z_{4n}^{(M)}) &\leq ((Z_n^{(M)})^2 - Z_{2n}^{(M)})^2, \\ Z_v^{(M)} &\leq (Z_u^{(M)})^{v/u}, \end{aligned} \tag{S14}$$

for even $n \geq 2$ and even $v \geq u \geq 2$. We suppress the temperature argument only in this display.

SII.C. Controlled product limit

To pass the finite-M inequalities in Eq. (S14) to the exact partition functions $Z_L(\beta)$, we prove the matrix product limit $S_L(\beta/M)^M \to e^{-\beta H_L}$ at fixed L and β. Use the operator norm, which is submultiplicative and satisfies $\|\mathbb{I}\| = 1$. For $\|X\|, \|Y\| \le K$, $K \ge 1$, telescoping powers gives

$$\left\|X^k - Y^k\right\| \le kK^k \|X - Y\|. \tag{S15}$$

For $k \ge 1$, expand the difference as $\sum_{r=0}^{k-1} X^{k-1-r}(X-Y)Y^r$; each term is bounded by $K^{k-1}\|X-Y\| \le K^k \|X-Y\|$. The case $k = 0$ is immediate.

Suppose $M(X_M - \mathbb{I}) \to A$, and set $Y_M = e^{A/M}$. Then $M(Y_M - \mathbb{I}) \to A$ and $Y_M^M = e^A$. With $C = \|A\| + 1$, convergence implies, for sufficiently large M,

$$\|X_M\|, \|Y_M\| \le 1 + C/M.$$

Equation (S15) now gives the quantitative comparison

$$\begin{aligned} \left\|X_M^M - e^A\right\| &\le (1 + C/M)^M \|M(X_M - Y_M)\| \\ &\le e^C \|M(X_M - Y_M)\| \longrightarrow 0. \end{aligned} \tag{S16}$$

In particular, if $g(0) = \mathbb{I}$ and $g'(0) = A$, then $g(1/M)^M \to e^A$.

For fixed L, β, list the negated bond operators in the order of Eq. (S3), including the negated on-site operators when present, as $B_1, \ldots, B_k$. Their sum is $-H_L$. The polynomial

$$g(t) = \prod_{i=1}^{k} (\mathbb{I} + \beta t B_i)$$

has derivative $g'(0) = -\beta H_L$. Hence

$$S_L(\beta/M)^M \longrightarrow e^{-\beta H_L}, \qquad Z_L^{(M)}(\beta) \longrightarrow Z_L(\beta). \tag{S17}$$

More explicitly, Taylor expansion gives $g(1/M) - e^{-\beta H_L/M} = O(M^{-2})$. Equation (S16) therefore bounds the operator error by $e^C K_{L,\beta}/M$ for some finite $K_{L,\beta}$ and sufficiently large M, with $C = \|\beta H_L\| + 1$. Trace convergence follows from $|\mathrm{Tr}\, A| \le 3^L \|A\|$ in the spin-1 model.

We take this limit separately at each length occurring in Eq. (S14). Continuity preserves all three inequalities, so they hold for $Z_L(\beta)$. We take this limit *before* iterating the doubling argument, which requires neither an error bound uniform in L, nor eigenvector convergence, nor a single limiting transfer operator. The finite slices need not form an exact Gibbs semigroup; thermal squaring below uses only the exact exponential after the limit.

SII.D. Torus identities and asymmetric doubling

With the exact partition-function inequalities established, we now set up the coupled doubling recurrence for the spatial and thermal purities:

$$P_{\mathrm{s}}(n, \beta) = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2}, \qquad P_\tau(L, \beta) = \frac{Z_L(2\beta)}{Z_L(\beta)^2}. \tag{S18}$$

All partition functions are strictly positive. The thermal purity is the purity of the Gibbs state $\rho_L(\beta) = e^{-\beta H_L}/Z_L(\beta)$. The limiting spatial inequalities and the Boltzmann power sums give $0 < P_{\mathrm{s}}, P_\tau \le 1$ at even lengths and

$$\begin{aligned} 1 - P_{\mathrm{s}}(2n, \beta) &\le f(1 - P_{\mathrm{s}}(n, \beta)), \\ 1 - P_\tau(L, 2\beta) &\le f(1 - P_\tau(L, \beta)). \end{aligned} \tag{S19}$$

For the spatial bound, divide the second limiting inequality in Eq. (S14) by $2Z_{2n}(\beta)^2$. For the thermal bound, apply Eq. (S12) to the normalized Boltzmann weights.

A torus of width n and height β represents $Z_n(\beta)$. Gluing four copies in either order gives the exact cancellation

$$P_{\mathrm{s}}(n,2\beta)P_\tau(n,\beta)^2 = \frac{Z_{2n}(2\beta)}{Z_n(\beta)^4} = P_{\mathrm{s}}(n,\beta)^2 P_\tau(2n,\beta). \tag{S20}$$

Solving for one factor and repeating at length $2n$ yields

$$\begin{aligned} P_{\mathrm{s}}(n,2\beta) &= \frac{P_{\mathrm{s}}(n,\beta)^2 P_\tau(2n,\beta)}{P_\tau(n,\beta)^2}, \\ P_\tau(4n,\beta) &= \frac{P_\tau(2n,\beta)^2 P_{\mathrm{s}}(2n,2\beta)}{P_{\mathrm{s}}(2n,\beta)^2}. \end{aligned} \tag{S21}$$

Both identities follow directly by substituting Eq. (S18); they concern exact partition functions.

Let $n \geq 2$ be even, and let $0 \leq p, q < 1$ bound the seed defects:

$$P_{\mathrm{s}}(n,\beta) \geq 1-p, \qquad P_\tau(2n,\beta) \geq 1-q. \tag{S22}$$

To double both n and β, we alternate the coupling identity with squaring in space and imaginary time, feeding the updated spatial defect into the thermal step. Write $z(a,b) = 1-(1-a)^2(1-b)$. The asymmetric update is

$$p' = f(z(p,q)), \qquad q' = f(z(q,p')). \tag{S23}$$

Assuming $p' < 1$, its four steps are:

1. The first identity in Eq. (S21), with $P_\tau(n,\beta) \leq 1$, gives $P_{\mathrm{s}}(n,2\beta) \geq (1-p)^2(1-q) = 1-z(p,q)$.
2. Spatial squaring and monotonicity of f give $P_{\mathrm{s}}(2n,2\beta) \geq 1-p'$.
3. The second identity in Eq. (S21), with $P_{\mathrm{s}}(2n,\beta) \leq 1$, gives $P_\tau(4n,\beta) \geq (1-q)^2(1-p') = 1-z(q,p')$.
4. Thermal squaring gives $P_\tau(4n,2\beta) \geq 1-q'$.

Thus (p',q') bounds the defects at $(2n,2\beta)$. Any upper bound below one substituted for p' in the third step remains valid; we use this rounding rule next.

SII.E. Six exact rational steps

The analytic tail estimate in Sec. SII SII.F requires both defects to be at most $1/100$. Because the initial spin-1 seed defects at $(n_0,\beta_0) = (36, 49/4)$ are larger, we first iterate the asymmetric update for six steps in exact rational arithmetic. The finite-size input is

$$P_{\mathrm{s}}(36,49/4) \geq 1 - \frac{47916}{10^6}, \qquad P_\tau(72,49/4) \geq 1 - \frac{181521}{10^6}. \tag{S24}$$

Ref. [7] provides these certified seed bounds for the periodic spin-1 chain; their finite-size spectral computation is not repeated here. The following calculation verifies their propagation under the recurrence.

All terminating decimals defining bounds denote exact rationals. Use the ceiling grid $D^{-1} = 10^{-7}$, $D = 10^7$, and update sequentially:

$$\begin{aligned} p_{j+1} &= \frac{\lceil D f(z(p_j,q_j)) \rceil}{D}, \\ q_{j+1} &= \frac{\lceil D f(z(q_j,p_{j+1})) \rceil}{D}. \end{aligned} \tag{S25}$$

We round the spatial bound upward before the thermal update. Table S1 lists the resulting orbit.

To check the table in integer arithmetic, write $a_j = Dp_j$, $b_j = Dq_j$, and set

$$V_j = (D-a_j)^2(D-b_j), \qquad W_j = (D-b_j)^2(D-a_{j+1}). \tag{S26}$$

TABLE S1. Upward-rounded defect bounds at $n_j = 36\,2^j$, $\beta_j = (49/4)2^j$.

j	n_j	β_j	p_j	q_j
0	36	49/4	0.0479160	0.1815210
1	72	49/2	0.0605003	0.1733838
2	144	49	0.0686630	0.1632461
3	288	98	0.0713693	0.1447318
4	576	196	0.0633146	0.1055663
5	1152	392	0.0376144	0.0446518
6	2304	784	0.0084709	0.0055149

For each $j = 0, \ldots, 5$, the entries in Table S1 satisfy

$$\begin{aligned} 2(a_{j+1}-1)V_j^2 &< D(D^3 - V_j)^2 \le 2a_{j+1}V_j^2, \\ 2(b_{j+1}-1)W_j^2 &< D(D^3 - W_j)^2 \le 2b_{j+1}W_j^2. \end{aligned} \tag{S27}$$

Multiplying the ceiling comparisons in Eq. (S25) by their positive denominators gives Eq. (S27), since $f(z(p_j, q_j)) = (D^3 - V_j)^2/(2V_j^2)$. The right-hand inequalities alone suffice to verify the upper bounds. Because all entries lie below one, the four-step update applies at every row and yields

$$\begin{aligned} P_{\mathrm{s}}(2304, 784) &\ge 1 - 0.0084709 > 0.99, \\ P_\tau(4608, 784) &\ge 1 - 0.0055149 > 0.99. \end{aligned} \tag{S28}$$

Both defects at $(2304, 784)$ now lie below the $1/100$ threshold.

SII.F. Scalar tail, interpolation, and the strict gap

Once both defects are at most $1/100$, a single scalar recurrence bounds all subsequent doubling steps. For $0 \le \varepsilon \le 1/100$,

$$f(1 - (1-\varepsilon)^3) \le \frac{9\varepsilon^2}{2(1-3\varepsilon)^2} \le 5\varepsilon^2 \le \varepsilon. \tag{S29}$$

The first inequality uses $1 - (1-\varepsilon)^3 \le 3\varepsilon$ and monotonicity of f; the second uses $9/[2(97/100)^2] < 5$. If both input defects are at most ε, the spatial update is therefore at most $5\varepsilon^2 \le \varepsilon$. Consequently, the thermal coupling input is also at most $1 - (1-\varepsilon)^3$, and its squared defect is at most $5\varepsilon^2$.

Starting from Eq. (S28), reset the scale index and put

$$n_j = 2304\,2^j, \qquad \beta_j = 784\,2^j, \qquad \varepsilon_j = \frac{20^{-2^j}}{5}, \qquad j \ge 0. \tag{S30}$$

Here $\varepsilon_0 = 1/100$ and $\varepsilon_{j+1} = 5\varepsilon_j^2$. Induction using Eq. (S29) proves

$$P_{\mathrm{s}}(n_j, \beta_j) \ge 1 - \varepsilon_j, \qquad P_\tau(2n_j, \beta_j) \ge 1 - \varepsilon_j. \tag{S31}$$

To cover every even length $L \ge 2304$ between the dyadic scales n_j, choose j with $n_j \le L \le 2n_j$. The limiting power-sum inequality in Eq. (S14), applied at β_j and $2\beta_j$, gives

$$\begin{aligned} Z_L(\beta_j)^2 &\le Z_{n_j}(\beta_j)^{2L/n_j}, \\ Z_L(2\beta_j) &\ge Z_{2n_j}(2\beta_j)^{L/(2n_j)}. \end{aligned} \tag{S32}$$

Set $r = L/(2n_j) \in [1/2, 1]$ and

$$x = \frac{Z_{2n_j}(2\beta_j)}{Z_{n_j}(\beta_j)^4} = P_\tau(2n_j, \beta_j) P_{\mathrm{s}}(n_j, \beta_j)^2.$$

Since $(1-\varepsilon_j)^3 \le x \le 1$, dividing the second inequality in Eq. (S32) by the first yields

$$P_\tau(L,\beta_j) \ge x^r \ge x \ge (1-\varepsilon_j)^3 \ge 1-3\varepsilon_j > \frac{1}{2}. \tag{S33}$$

Finally, we convert this thermal purity bound at β_j into a lower bound on the spectral gap. Let $E_0(L) \le E_1(L) \le \cdots$ include multiplicities, and let $w_m = e^{-\beta_j E_m(L)}/Z_L(\beta_j)$. The largest Gibbs weight satisfies $P_\tau(L,\beta_j) = \sum_m w_m^2 \le w_0$. Equation (S33) implies $w_0 > 1/2$, which proves that the ground state is unique. Moreover, $w_1 \le 1 - w_0$. With $\Delta_L = E_1(L) - E_0(L)$, we obtain

$$\begin{aligned} e^{-\beta_j \Delta_L} = \frac{w_1}{w_0} &\le \frac{3\varepsilon_j}{1-3\varepsilon_j} \\ &\le \frac{60}{97}\, 20^{-2^j} < 20^{-2^j}, \end{aligned} \tag{S34}$$

where we used $1 - 3\varepsilon_j \ge 97/100$. Taking logarithms and dividing by $\beta_j = 784\, 2^j$ at this finite, length-dependent temperature gives

$$\boxed{\Delta_L > \frac{\log 20}{784} \quad \text{for every even } L \ge 2304}. \tag{S35}$$

The strict inequality comes from the finite prefactor $60/97 < 1$, without taking a fixed-length limit $\beta \to \infty$. Given the two seed bounds in Eq. (S24), the finite-matrix transfer inequalities, rational orbit, scalar tail, and dyadic interpolation establish Eq. (S35).

Appendix SIII: Finite seed certificates and improved bounds

In this section we explain the finite inputs and scalar estimates from Ref. [7] that certify the starting pair

$$\begin{aligned} 1 - P_{\mathrm{s}}(36, 49/4) &\le \frac{47916}{10^6}, \\ 1 - P_\tau(72, 49/4) &\le \frac{181521}{10^6}. \end{aligned} \tag{S1}$$

These two bounds form Lemma 1 in the main text. We then present the temperature-lifting calculation, which combines the same short-ring trace bounds with the 120-site trial state of Ref. [7] to prove a unique ground state and

$$\Delta_L > 0.072901771 \quad \text{for every even } L \ge 36. \tag{S2}$$

We have reproduced the finite trace and trial contractions of Ref. [7] in exact arithmetic; the stronger gap bound uses only additional scalar inequalities on those inputs. Every terminating decimal that defines a bound below denotes an exact rational number, and we mark numerical approximations explicitly.

SIII.A. The base seed at length 36

To prove Eq. (S1) at $b_0 = 49/4$, we need lower bounds on the numerators $Z_{72}(b_0)$ and $Z_{72}(2b_0)$ and upper bounds on the denominators $Z_{36}(b_0)^2$ and $Z_{72}(b_0)^2$. A 72-site trial state bounds $Z_{72}(b)$ from below for every $b > 0$. Traces on $4 \le n \le 12$ sites constrain the eigenvalues of the one-site spatial transfer operator at b_0, which lets us bound the subleading contributions to $Z_{36}(b_0)$ and $Z_{72}(b_0)$.

a. Hamiltonians, twists, and the spatial transfer operator

We consider the periodic spin-1 Heisenberg Hamiltonian

$$H_n = \sum_{j=1}^{n} \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}, \qquad \boldsymbol{S}_j^2 = 2I, \qquad \boldsymbol{S}_{n+1} = \boldsymbol{S}_1, \tag{S3}$$

with unit exchange coupling. Lengths count sites, and $\Delta_L = E_1(L) - E_0(L)$, with eigenvalues ordered with multiplicity. Anticipating the trial energy per site, we shift the Hamiltonian by κ per site:

$$\kappa = \frac{700741}{500000}, \qquad \widehat{H}_n = H_n + \kappa n I, \qquad Z_n(\beta) = \operatorname{Tr} e^{-\beta \widehat{H}_n}. \tag{S4}$$

This extensive shift cancels in both purities:

$$P_{\mathrm{s}}(n,\beta) = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2}, \qquad P_\tau(n,\beta) = \frac{Z_n(2\beta)}{Z_n(\beta)^2}. \tag{S5}$$

To separate the spatial transfer spectrum by spin-rotation symmetry, we also use rings with a twisted closing bond. For an on-site spin rotation u_g representing g, we define

$$\begin{aligned} H_n(g) &= \sum_{j=1}^{n-1} h_{j,j+1} + h_{n,1}(g), \\ h_{n,1}(g) &= \sum_{\alpha=x,y,z} S_n^\alpha (u_g S^\alpha u_g^\dagger)_1, \\ Z_n(\beta, g) &= \operatorname{Tr} e^{-\beta(H_n(g) + \kappa n I)}, \end{aligned} \tag{S6}$$


where $h_{j,j+1} = \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$. The twist modifies the spatial seam rather than inserting $u_g^{\otimes n}$ into the untwisted Gibbs trace.

The continuous-time construction of Ref. [7] provides a self-adjoint one-site spatial transfer operator K_β for the shifted Hamiltonian and a virtual symmetry operator U_g such that

$$Z_n(\beta, g) = \operatorname{Tr}(K_\beta^n U_g), \qquad n \geq 4. \tag{S7}$$

At $g = 1$, $Z_n(\beta) = \sum_i \lambda_i(\beta)^n$ is a power sum of the real eigenvalues $\lambda_i(\beta)$ of K_β. Because the same eigenvalues govern every length $n \geq 4$, trace bounds on $4 \leq n \leq 12$ constrain the power sums at $n = 36$ and $n = 72$. Unlike the positive two-site Trotter matrix of Section SII SII.A, the one-site operator K_β can have negative eigenvalues, so we must bound $|\lambda_i(\beta)|$ on both sides of zero.

In our diagrams, horizontal wires carry spatial virtual indices, vertical double wires carry physical indices, and closed loops denote traces. Commutation of K_β with U_g gives



(S8)

where the ring contains n transfer vertices and U_g implements the twisted closing bond of Eq. (S6).

b. *Exact finite trace and trial inputs*

We now state the two numerical inputs from Ref. [7] used at $b_0 = 49/4$. The first input is the table of shifted traces $Z_n(b_0)$ and $Z_n(b_0, P)$ for $4 \leq n \leq 12$, where P is a π rotation about a Cartesian axis. Table S2 lists the rational centers, each certified within the absolute error

$$\epsilon_Z = 3 \times 10^{-5}. \tag{S9}$$

These enclosures come from integer Horner recursions for the half-Gibbs operator rather than dense diagonalization. For an unshifted Hamiltonian $H = H_n(g)$, we set

$$R = I - \frac{(3n/2)I + H}{16}, \tag{S10}$$

which satisfies $\|R\| \leq 1$ on each ring in Table S2. At $b_0 = 49/4$, the shifted half-Gibbs operator is

$$e^{-b_0(H + \kappa n I)/2} = e^{-b_0(\kappa - 3/2)n/2} e^{98(R-I)}. \tag{S11}$$

TABLE S2. Exact rational centers of the shifted traces at $b_0 = 49/4$ from Ref. [7]. Each entry has certified absolute error at most 3×10^{-5}.

n	$Z_n(b_0)$ center	$Z_n(b_0, P)$ center
4	124.885379	0.065757
5	0.002925	2.960742
6	12.870614	0.286599
7	0.054069	1.920414
8	4.639283	0.510140
9	0.195455	1.514846
10	2.654823	0.675419
11	0.377629	1.314544
12	1.900547	0.787405

We approximate $e^{98(R-I)}$ by the normalized Poisson polynomial

$$\sum_{j=0}^{J} p_j R^j, \qquad p_j = \frac{98^j/j!}{\sum_{\ell=0}^{J} 98^\ell/\ell!}, \qquad J = 280, \tag{S12}$$

whose truncation and normalization error in operator norm is at most

$$2e^{-98} \sum_{j>280} \frac{98^j}{j!} < 2^{-84}. \tag{S13}$$

Setting $Q = 2^{55}$, we replace Qp_j by integers c_j with $|c_j - Qp_j| \le 1$. Writing $R = D/s$ with an integer matrix D and integer divisor $s > 0$, we evaluate each basis column e_r by the downward recursion

$$X_{J+1} = 0, \qquad X_j = \left\lfloor \frac{DX_{j+1}}{s} \right\rfloor + c_j e_r, \tag{S14}$$

for $j = J, \dots, 0$, where the floor acts entrywise, including on negative entries. In dimension $d = 3^n$, each step introduces a Euclidean rounding error of at most $\sqrt{d} + 1$. Because $\|R\| \le 1$, these errors do not amplify, so the total Euclidean error in X_0 is at most $(J+1)(\sqrt{d}+1)$ before adding the Poisson tail (S13). The trace $Z_n(b_0, g)$ equals the squared Frobenius norm of the half-Gibbs operator, which we obtain by summing the squared column norms with their exact symmetry multiplicities and multiplying by a rational enclosure of the scalar prefactor in Eq. (S11).

The second input is a periodic matrix-product trial state defined by real integer matrices A_s of bond dimension 26, used at $N = 72$ for the base seed and at $N = 120$ for temperature lifting. Setting

$$\begin{aligned} \psi_N(s_1, \dots, s_N) &= \operatorname{Tr}(A_{s_1} \cdots A_{s_N}), \\ T &= \sum_s A_s \otimes A_s, \\ O &= \sum_{s,t,s',t'} h_{st,s't'} A_s A_t \otimes A_{s'} A_{t'}, \end{aligned} \tag{S15}$$

where T is the $26^2 \times 26^2$ trial transfer matrix, we express the norm and energy expectation as exact finite-matrix traces:

$$\|\psi_N\|^2 = \quad = \operatorname{Tr} T^N,$$

$$\langle \psi_N, H_N \psi_N \rangle = N \quad = N \operatorname{Tr}(T^{N-2} O). \tag{S16}$$

Each layer has N tensors, and the two-site operator h acts on one bond. Exact integer arithmetic verifies that for $N \in \{72, 120\}$,

$$\begin{aligned} &\operatorname{Tr} T^N > 0, \\ &500000 \operatorname{Tr}(T^{N-2}O) + 700741 \operatorname{Tr} T^N < 0. \end{aligned} \tag{S17}$$

By the variational principle, the ground energy satisfies $E_0(N) < -\kappa N$, so the shifted ground energy $E_0(N) + \kappa N$ is negative. Retaining only the ground-state Boltzmann weight gives the lower bounds

$$Z_{72}(\beta) > 1, \qquad Z_{120}(\beta) > 1 \quad (\beta > 0). \tag{S18}$$

In the base seed, $Z_{72}(2b_0) > 1$ bounds the thermal numerator directly, whereas $Z_{72}(b_0) > 1$ forces a large leading spatial eigenvalue.

c. *Sector moments and polynomial filters*

We next split the spectrum of K_{b_0} into spin-rotation sectors and bound its individual eigenvalues. The four virtual π rotations $D_2 = \{1, P_x, P_y, P_z\}$ commute with K_{b_0} and define the character projectors

$$\Pi_\chi = \frac{1}{4} \sum_{g \in D_2} \chi(g) U_g, \qquad \operatorname{Tr}(K_{b_0}^n \Pi_\chi) =$$ 

. (S19)

Cyclic permutation of the coordinate axes conjugates the three nontrivial characters into one another, so their three sectors have identical eigenvalue lists, which we denote by N; we denote the D_2-invariant eigenvalue list by J. Writing $J_n = \sum_{\lambda \in J} \lambda^n$ and $N_n = \sum_{\lambda \in N} \lambda^n$ at b_0, we have

$$\begin{aligned} Z_n &= J_n + 3N_n, & Z_n(P) &= J_n - N_n, \\ J_n &= \frac{Z_n + 3Z_n(P)}{4}, & N_n &= \frac{Z_n - Z_n(P)}{4}. \end{aligned} \tag{S20}$$

From Eq. (S9), the sector moments J_n and N_n for $4 \le n \le 12$ have absolute errors at most ϵ_Z and $\epsilon_Z/2$ around their rational centers. In particular, at $n = 12$ these enclosures yield the twelfth-moment upper bounds

$$J_{12} \le m_J = \frac{2131441}{2000000}, \qquad N_{12} \le m_N = \frac{556631}{2000000}. \tag{S21}$$

To bound individual eigenvalues from these finite power sums, we evaluate the nonnegative polynomials of Ref. [7],

$$\begin{aligned} F_J(x) &= x^4(7 + 17x - 280x^2 - 185x^3 + 1000x^4)^2, \\ F_N(x) &= x^4(12 - 394x^2 + 1000x^4)^2. \end{aligned} \tag{S22}$$

Pairing each coefficient of F_J and F_N with the upper or lower endpoint of the corresponding moment interval for $4 \le n \le 12$ gives

$$\begin{aligned} \sum_{\lambda \in J} F_J(\lambda) &\le B_J = 318604.54848175, \\ \sum_{\lambda \in N} F_N(\lambda) &\le B_N = 48169.880699. \end{aligned} \tag{S23}$$

Because F_J and F_N are nonnegative on $\mathbb{R}$, each single eigenvalue satisfies $F_J(\lambda) \le B_J$ for $\lambda \in J$ and $F_N(\lambda) \le B_N$ for $\lambda \in N$.

For $U = 1.00139$ and $V = 0.872$, we expand $F_J(\pm(U + y))$ and $F_N(\pm(V + y))$ in powers of $y \ge 0$. Exact rational arithmetic shows that their constant terms exceed B_J and B_N and all higher coefficients are nonnegative, which rules out eigenvalues on the exterior rays $|x| \ge U$ and $|x| \ge V$:

$$|\lambda| < U \quad (\lambda \in J), \qquad |\lambda| < V \quad (\lambda \in N). \tag{S24}$$

Checking both signs $\pm(U + y)$ and $\pm(V + y)$ is required because K_{b_0} is self-adjoint rather than positive.

d. Capped masses and the two base defects

To pass from the twelfth moments (S21) and the eigenvalue caps (S24) to $n = 36$ and $n = 72$, we bound higher power sums under a mass constraint and an entry cap. For $m \geq 0$, $a > 0$, and $r \geq 1$, set

$$\mathcal{C}(m, a, r) = qa^r + (m - qa)^r, \qquad q = \lfloor m/a \rfloor. \tag{S25}$$

Whenever $0 \leq t_i \leq a$ and $\sum_i t_i \leq m$, convexity of t^r implies $\sum_i t_i^r \leq \mathcal{C}(m, a, r)$: shifting mass from a smaller unsaturated entry to a larger one increases $\sum_i t_i^r$, so the maximum is achieved by placing q entries at the cap a and the remaining mass $m - qa$ in a single entry. Applied to $t_i = \lambda_i^{12} \geq 0$, this bounds higher moments of the 12th-power weights.

We first show that J contains a dominant eigenvalue with $|\lambda_0| > \ell = 0.999$. If every $\lambda \in J$ satisfied $|\lambda| \leq \ell$, then applying Eq. (S25) with $r = 6$ to J (with cap ℓ^{12}) and to each of the three N sectors (with cap V^{12}) would give

$$Z_{72}(b_0) \leq \mathcal{C}(m_J, \ell^{12}, 6) + 3\mathcal{C}(m_N, V^{12}, 6) < 0.931 < 1, \tag{S26}$$

contradicting the 72-site trial bound $Z_{72}(b_0) > 1$ in Eq. (S18). Because $m_N < \ell^{12}$, no N eigenvalue can exceed ℓ in modulus, so J has at least one eigenvalue λ_0 with $|\lambda_0| > \ell$.

Removing $\lambda_0^{12} > \ell^{12}$ from J_{12} leaves residual twelfth moment at most $m_J - \ell^{12} < U^{12}$ in J. For even $k \geq 12$, the subleading moment $\sum_{i \neq 0} \lambda_i^k$ is therefore bounded by

$$R_k = (m_J - \ell^{12})^{k/12} + 3\mathcal{C}(m_N, V^{12}, k/12). \tag{S27}$$

At $k = 36$ and $k = 72$, we obtain the two base purity bounds:

$$\begin{aligned} P_{\mathrm{s}}(36, b_0) &\geq \left(1 + \frac{R_{36}}{\ell^{36}}\right)^{-2} = 1 - p, \\ P_{\tau}(72, b_0) &\geq \left(U^{72} + R_{72}\right)^{-2} = 1 - q. \end{aligned} \tag{S28}$$

For $P_{\mathrm{s}}(36, b_0)$, we bound the numerator from below by $Z_{72}(b_0) \geq \lambda_0^{72}$, bound $Z_{36}(b_0) \leq |\lambda_0|^{36} + R_{36}$, and use $|\lambda_0| > \ell$. For $P_{\tau}(72, b_0)$, we bound the numerator from below by $Z_{72}(2b_0) > 1$ from Eq. (S18) and bound the denominator by $Z_{72}(b_0) = \lambda_0^{72} + \sum_{i \neq 0} \lambda_i^{72} \leq U^{72} + R_{72}$.

Because $k/12 \in \{3, 6\}$ is an integer, Eqs. (S26)–(S28) involve only rational arithmetic. Exact evaluation verifies

$$p \leq \frac{47916}{10^6}, \qquad q \leq \frac{181521}{10^6}, \tag{S29}$$

which completes the proof of Eq. (S1). Numerically, $p \simeq 0.0479157710062$ and $q \simeq 0.181520168033$.

SIII.B. Temperature lifting and the improved uniform bound

Because the gap bound from the doubling argument scales inversely with the seed inverse temperature, certifying smaller defects at a moderate inverse temperature and stopping after a finite prefix of doubling steps gives a much larger gap constant. We now explain the temperature-lifting method, which uses the same trace table at $b_0 = 49/4$ and the 120-site trial bound $Z_{120}(\beta) > 1$ to certify a seed at $(n_0, b) = (60, rb_0)$ with $r \geq 1$. Combining that lifted seed with a tighter unlifted bound at $(36, b_0)$ covers all even lengths $L \geq 36$.

a. Lifted moments and a dominant spatial weight

We keep m_J, m_N from Eq. (S21) and sharpen the filter caps (S24) to the exact rationals

$$U = 1.001350644012, \qquad V = 0.871255704653, \tag{S30}$$

verified by the same exterior-ray expansions of F_J and F_N. Since $U^{12} < m_J < 2U^{12}$ and $V^{12} < m_N < 2V^{12}$, Eq. (S25) bounds the full sector moments at any even length $k \geq 12$ by

$$\begin{aligned} \mathcal{J}_k &= U^k + (m_J - U^{12})^{k/12}, \\ \mathcal{N}_k &= V^k + (m_N - V^{12})^{k/12}, \\ M_k &= \mathcal{J}_k + 3\mathcal{N}_k. \end{aligned} \tag{S31}$$

Because $N_k \geq 0$ at even k, we have $Z_k(b_0) = J_k + 3N_k \leq M_k$ and $Z_k(b_0, P) = J_k - N_k \leq \mathcal{J}_k$.

Now fix a rational factor $r \geq 1$, an even length $12 \leq k \leq 60$, and the lifted inverse temperature $b = rb_0$. Both H_k and $H_k(P)$ are self-adjoint, so their physical Boltzmann weights $z_i > 0$ at b_0 satisfy $\sum_i z_i^r \leq (\sum_i z_i)^r$. This transfers the upper bounds at b_0 to b:

$$Z_k(b) \leq D := M_k^r, \qquad Z_k(b, P) \leq E := \mathcal{J}_k^r. \tag{S32}$$

In rational arithmetic we replace D and E by outward rational upper bounds satisfying $E \leq D$.

At the lifted inverse temperature b, let K_b be the one-site spatial transfer operator and define the kth-power weights

$$w_i = |\lambda_i(b)|^k, \qquad h = \frac{60}{k}, \qquad \tau = \frac{D + 3E}{4}. \tag{S33}$$

Since k is even, $w_i = \lambda_i(b)^k \geq 0$. Thus $\sum_i w_i = Z_k(b) \leq D$, the J-sector sum satisfies $\sum_{\lambda_i \in J} w_i = [Z_k(b) + 3Z_k(b, P)]/4 \leq \tau$, and the 120-site trial bound (S18) gives

$$\sum_i w_i^{2h} = Z_{120}(b) > 1. \tag{S34}$$

If a rational threshold $t > 0$ satisfies

$$Dt^{2h-1} < 1, \qquad 3t > D, \tag{S35}$$

then at least one weight must satisfy $w_0 > t$; otherwise $\sum_i w_i^{2h} \leq t^{2h-1} \sum_i w_i \leq Dt^{2h-1} < 1$, contradicting Eq. (S34). Moreover, $3t > D$ forces w_0 to lie in J, because any weight in N is triply degenerate and would contribute more than $3t > D$ to $\sum_i w_i$.

Removing w_0 leaves residual J mass $x \in [0, \tau - t]$, while each of the three identical N sectors has mass at most $(D - t - x)/3$. For any $s \geq 1$, the residual moment $\sum_{i \neq 0} w_i^s$ is at most $x^s + 3[(D - t - x)/3]^s$. By convexity on $x \in [0, \tau - t]$, this sum attains its maximum at one of the two endpoints:

$$\mathcal{R}_s(t) = \max\left\{3\left(\frac{D - t}{3}\right)^s, (\tau - t)^s + 3\left(\frac{D - \tau}{3}\right)^s\right\}. \tag{S36}$$

All choices below satisfy $0 < t \leq \tau \leq D$.

Once $w_0 > t$ is established, any larger rational t' with

$$(t')^{2h} + \mathcal{R}_{2h}(t) < 1 \tag{S37}$$

implies $w_0 > t'$, since $w_0 \leq t'$ would give $Z_{120}(b) = w_0^{2h} + \sum_{i \neq 0} w_i^{2h} \leq (t')^{2h} + \mathcal{R}_{2h}(t) < 1$. Iterating Eq. (S37) tightens the lower bound t on w_0. Because $w_0^h = |\lambda_0(b)|^{60}$ and $\sum_{i \neq 0} |\lambda_i(b)|^{60} \leq \mathcal{R}_h(t)$, we obtain the lifted spatial purity bound at $n = 60$:

$$P_{\mathrm{s}}(60, b) \geq \left(1 + \frac{\mathcal{R}_h(t)}{t^h}\right)^{-2}. \tag{S38}$$

b. *The physical thermal residual*

To bound $P_\tau(120, b)$ at $b = rb_0$, we work directly with the physical Boltzmann weights $z_i = e^{-b_0 \widehat{E}_i(120)}$ of $\widehat{H}_{120}$ rather than the spatial eigenvalues of K_b. Ordering $z_0 \geq z_1 \geq \cdots$, we have $z_0 > 1$ from the 120-site trial state and $\sum_i z_i = Z_{120}(b_0) \leq M_{120}$ from Eq. (S31). Hence

$$\sum_{i>0} \frac{z_i}{z_0} \leq M_{120} - 1, \qquad \sum_{i>0} \left(\frac{z_i}{z_0}\right)^r \leq (M_{120} - 1)^r. \tag{S39}$$

At $b = rb_0$, the physical partition functions are $Z_{120}(b) = \sum_i z_i^r$ and $Z_{120}(2b) = \sum_i z_i^{2r} \geq z_0^{2r}$. Dividing numerator and denominator by z_0^{2r} gives the lifted thermal purity bound:

$$P_\tau(120, rb_0) \geq [1 + (M_{120} - 1)^r]^{-2}. \tag{S40}$$

The transfer identity $\mathrm{Tr}\, K_\beta^L = Z_L(\beta)$ equates the full traces at each β; it does not identify individual spatial eigenvalues with physical Boltzmann weights.

TABLE S3. Temperature-lifting certificates at $(60, rb_0)$ with $b_0 = 49/4$. Defect bounds at $j = 0$ are rounded upward; gap bounds hold for every even $L \geq 60$, and for every even $L \geq 36$ once combined with the unlifted bridge on $36 \leq L \leq 72$.

k	r	Spatial defect	Thermal defect	$\Delta_L >$
30	2	0.0307608679	0.0590740709	0.070450002
28	17/8	0.0305302052	0.0479556776	0.072901771

c. *Finite-prefix coverage and the doubling tail*

We now convert the seed bounds at (n_0, b_0) into a uniform gap bound by combining a finite prefix of rational doubling steps with an analytic tail estimate.

Proposition S1 (Finite-prefix coverage)**.** *Let n_0 be even and $b_0 > 0$. Suppose $0 \leq p_j, q_j < 1$ bound respectively $1 - P_{\mathrm{s}}(n_j, b_j)$ and $1 - P_\tau(2n_j, b_j)$ at $(n_j, b_j) = (2^j n_0, 2^j b_0)$ for $0 \leq j \leq K$. Put $A_j = (1-p_j)^2(1-q_j)$ and assume $A_j > 1/2$ for $j < K$. Let $u = \max(p_K, q_K)$ and choose C such that*

$$0 < u < 1/6, \qquad Cu < 1, \qquad C(1-3u) > 3,$$
$$G(u) := \frac{1}{2}\left((1-u)^{-1} + (1-u)^{-2} + (1-u)^{-3}\right)^2 \leq C. \tag{S41}$$

If g is strictly smaller than every member of

$$\left\{ \frac{\log(A_j/(1-A_j))}{b_j} : 0 \leq j < K \right\} \cup \left\{ \frac{\log(1/(Cu))}{b_K} \right\}, \tag{S42}$$

then every even $L \geq n_0$ has a unique ground state and $\Delta_L > g$. An entry with $A_j = 1$ is interpreted as $+\infty$.

Proof. For even $n_j \leq L \leq 2n_j$ with $0 \leq j < K$, spatial power-sum interpolation (S33) gives $P_\tau(L, b_j) \geq A_j^{L/(2n_j)} \geq A_j > 1/2$. The largest normalized Boltzmann weight a_0 satisfies $a_0 \geq P_\tau(L, b_j) > 1/2$, so the ground state is unique, and the next weight $a_1 \leq 1 - a_0$ obeys

$$e^{-b_j \Delta_L} = \frac{a_1}{a_0} \leq \frac{1-a_0}{a_0} \leq \frac{1-A_j}{A_j}, \tag{S43}$$

which yields $\Delta_L \geq \log(A_j/(1-A_j))/b_j > g$. For $j \geq K$, since $f(1-(1-x)^3) = x^2 G(x)$ and G is increasing on $[0, u]$, Eq. (S41) implies that $u_m = C^{-1}(Cu)^{2^{m-K}}$ bounds both defects at (n_m, b_m) by induction. Interpolating on $n_m \leq L \leq 2n_m$ gives $P_\tau(L, b_m) \geq 1 - 3u_m > 1/2$ and

$$e^{-b_m \Delta_L} \leq \frac{3u_m}{1-3u_m} \leq \frac{3}{C(1-3u)}(Cu)^{2^{m-K}} < (Cu)^{2^{m-K}},$$

so $\Delta_L > \log(1/(Cu))/b_K > g$. Together, the intervals $[n_j, 2n_j]$ for $0 \leq j < K$ and the tail $L \geq n_K$ cover every even $L \geq n_0$. □

We generate the prefix (p_j, q_j) from the asymmetric recurrence with $f(x) = x^2/[2(1-x)^2]$:

$$p_{j+1} = \left\lceil f(1-(1-p_j)^2(1-q_j)) \right\rceil_\delta,$$
$$q_{j+1} = \left\lceil f(1-(1-q_j)^2(1-p_{j+1})) \right\rceil_\delta, \tag{S44}$$

where $\lceil x \rceil_\delta = \delta \lceil x/\delta \rceil$. For the lifted seeds at $(60, rb_0)$, we take $K = 10$ steps with $\delta = 10^{-2048}$ and apply Proposition S1.

Table S3 lists two parameter choices. The integer choice $(k, r) = (30, 2)$ has $h = 2$, so its spatial residual (S36) uses only second and fourth powers and already gives $\Delta_L > 0.07045$ for even $L \geq 60$. The fractional choice $(k, r) = (28, 17/8)$ gives $\Delta_L > 0.072901771$ for even $L \geq 60$.

To cover the remaining even lengths $36 \leq L < 60$, we evaluate the unlifted base-seed formulas (S27)–(S28) at $(36, b_0)$ using the sharper caps (S30) and $\ell = 0.999997937629$, validated by the capped-mass inequality $\mathcal{C}(m_J, \ell^{12}, 6) + 3\mathcal{C}(m_N, V^{12}, 6) < 1$. At $j = 0$, the resulting factor $A_0 = (1-p_0)^2(1-q_0)$ satisfies $\log(A_0/(1-A_0))/b_0 > 0.09$, which by Eq. (S43) bounds $\Delta_L > 0.09$ on every even $36 \leq L \leq 72$. Combining this bridge on $[36, 72]$ with the lifted bound on $[60, \infty)$ proves Eq. (S2) for every even $L \geq 36$.

SIII.C. Reproducibility and arithmetic assumptions

The repository provides the stored certificates and standalone verifiers for the finite inputs and scalar inequalities above.

To keep every scalar check in exact rational arithmetic, we enclose rational powers and logarithms by rational endpoints. For $x \geq 0$ and a rational exponent $a/b \geq 0$ with $b > 0$, nonnegative rational bounds l, u satisfy $l \leq x^{a/b} \leq u$ if and only if the integer powers satisfy $l^b \leq x^a \leq u^b$. To enclose $\log x$ for $x > 0$, we write $x = 2^k y$ with $k \in \mathbb{Z}$ and $1 \leq y < 2$, and set $z = (y-1)/(y+1) \in [0, 1/3)$. The series

$$\log y = 2 \sum_{j=0}^{M-1} \frac{z^{2j+1}}{2j+1} + E_M,$$
$$0 \leq E_M \leq \frac{2z^{2M+1}}{(2M+1)(1-z^2)} \tag{S45}$$

at $M = 40$ encloses $\log y$ and $\log 2$ in rational intervals, which we round outward. Floating-point calculations are used only to select parameters, never in the certificate checks.

Our trusted base consists of exact integer and rational arithmetic, the explicit Poisson and Horner error bounds for the short-ring traces, and the analytical transfer, interpolation, and doubling inequalities proved in Sections SII SII.A–SII SII.F and above. Together with the finite trace and trial inputs of Ref. [7], these checks establish Eq. (S2) for the periodic spin-1 chain (S3).