
How the Initial Estimate Is Certified

(Dated: October 15, 2026)

The finite-scale input $P_{\mathrm{s}}(2304,784), P_{\tau}(4608,784) \geq 0.99$ of Ref. [1] is established without direct tensor contraction at that scale. Ref. [3] certifies a pair of bounds at length 36 and $\beta = 49/4$, which are then propagated to $(2304,784)$ through six scale doublings with decoupled defects. The underlying verification reduces to three closed tensor networks: twisted rings of at most twelve sites, symmetry-projected rings, and an integer matrix-product trial state of bond dimension 26 on 72 sites.

Appendix I: Architecture of the Certificate

Throughout, X_b is the spatial transfer operator of Ref. [1] at inverse temperature b, with the energy shift $\kappa = 700741/500000$ absorbed into H. Its eigenvalues $\lambda_i(b)$ are real, and $Z_n(b) = \sum_i \lambda_i^n$. The certificate proceeds in four steps, from twisted partition functions to the initial estimate:




(I1)

- *Twisted partition functions* (Sec. J): certified values of $Z_n(b,g)$ for $4 \leq n \leq 12$, $b = 49/4$, and two twists g.
- *Sectors and filters* (Sec. K): bounds on every eigenvalue of X_b, sector by sector. A trial vector then forces one eigenvalue above $\ell = 0.999$.
- *Initial defects* (Sec. K): $1 - P_{\mathrm{s}}(36,b) \leq p_0$ and $1 - P_{\tau}(72,b) \leq q_0$.
- *Six doublings* (Sec. L): the coupled update reaches $(2304,784)$.

Every number is an exact rational or an integer; floating-point arithmetic is used nowhere in the proof.

Appendix J: Twisted boundary conditions

Twists and symmetries. A proper signed permutation g of the Cartesian axes acts on histories by U_g, which relabels events and multiplies by their signs. Rotating every matrix inside a site trace shows that U_g slides through a transfer vertex,



(J1)

so a single U_g may be placed on any virtual bond of the ring. On the physical side it conjugates one end of a single bond, the seam:


$$Z_n(b,g) = \operatorname{Tr} X_b^n U_g = \quad = \operatorname{Tr} e^{-bH_n(g)}. \tag{J2}$$



The certificate needs $g = 1$ and the π rotation $g = P$.

Purification. The physical side is computed as a squared Frobenius norm, which splits the imaginary-time circle at half its length,

$$Z_n(b,g) = \quad = \|F\|_F^2, \qquad F = e^{-bH/2}, \tag{J3}$$



so that only columns Fe_r are needed, one basis state e_r at a time.

Uniformization. Shift and scale H so that $R = 1 - (\frac{3}{2}n + H)/16$ is a contraction. Uniformization (Jensen's method) then writes F, up to the scalar $e^{-b(\kappa-3/2)n/2}$, as a Poisson mixture of powers of R:

$$F \propto e^{98(R-1)} = \sum_{j\geq 0} e^{-98} \frac{98^j}{j!} \quad , \tag{J4}$$



with j factors of R in the jth term. This is the discrete counterpart of the event expansion behind the thermal MPO: the events of W_β arrive as a Poisson process, and here each factor of R is one event.

The sum is cut at $j = 280$. The truncation error is below 2^{-84} in operator norm, because $\|R\| \leq 1$ [4]. The truncated sum is evaluated by Horner's rule on integer vectors scaled by $Q = 2^{55}$, taking the floor after every multiplication by R. Each multiplication adds at most $\sqrt{\dim} + 1$ to the Euclidean error, and the contraction carries earlier errors forward unamplified. Each column is therefore known to within $281(\sqrt{3^n} + 2)$ in units of $1/Q$.

Orbits. The trace in Eq. (J3) needs one column per basis word. Cyclic shifts, reversal and global spin flip permute the columns without changing their norms. For the twisted ring, a shift moves the seam, and a diagonal sign gauge moves it back. Only one representative per orbit is computed, weighted by the orbit size. The results are eighteen integers $W_{n,g}$ of about 30 digits. Turning them into rational enclosures gives the table of $Z_n(49/4, g)$, each value to within $\pm 3 \times 10^{-5}$. As an independent sanity check, not part of the proof, floating-point exact diagonalization reproduces all ten centers for $4 \leq n \leq 8$ to within 10^{-6}.

Appendix K: From partition functions to the transfer spectrum

Sector projectors. The four twists $g \in D_2 = \{1, P_x, P_y, P_z\}$ commute with X_b, so their character sums are projectors that can be inserted into the ring:

$$\Pi_\chi = \tfrac{1}{4}\textstyle\sum_{g\in D_2} \chi(g) U_g, \qquad \operatorname{Tr} X_b^n \Pi_\chi = \tag{K1}$$




A cyclic rotation of the axes permutes the three nontrivial characters, so those three sectors carry identical eigenvalue lists, called N. Write J for the list of the invariant sector. Inverting the characters gives both lists from the two twists alone:

$$J_n = \tfrac{1}{4}\big[Z_n + 3Z_n(P)\big], \qquad N_n = \tfrac{1}{4}\big[Z_n - Z_n(P)\big], \tag{K2}$$

with $Z_n = J_n + 3N_n$. The coefficients sum to at most one in absolute value, so each sector moment inherits the 3×10^{-5} error. This holds for odd n too; odd moments carry sign information.

Polynomial filters (the term of Ref. [3]). For any polynomial $F \geq 0$ on $\mathbb{R}$, the closed network $\operatorname{Tr} F(X_b)$ is a finite combination of ring partition functions:

$$\sum_i F(\lambda_i) = \sum_j f_j \quad \leq \quad B. \tag{K3}$$



Every term on the left is nonnegative, so no single eigenvalue can have $F(\lambda) > B$. Two degree-12 filters of the form $x^4 Q(x)^2$ give

$$|\lambda| < v_J = 1.00139 \text{ in } J, \qquad |\lambda| < v_N = 0.872 \text{ in } N. \tag{K4}$$


The filter needs $F \geq 0$ on the whole real line, not only on $[0, \infty)$, because X_b is self-adjoint but not positive. Reflection positivity is what makes the eigenvalues real [2].

Capped masses (also Ref. [3]'s term; a convexity bound). The twelfth moments are bounded by $m_J = 1.0657205$ and $m_N = 0.2783155$. Weights $t_i \leq h$ with total mass at most m have their rth powers maximized by filling bars to the cap,  :


$$\sum_i t_i^r \leq \mathcal{C}(m, h, r) = qh^r + (m - qh)^r, \qquad q = \lfloor m/h \rfloor. \tag{K5}$$

The trial vector. One matrix-product vector on 72 sites, with real integer tensors A_s of bond dimension 26, has energy per site below $-\kappa$. Its norm and energy are two double-layer rings,

$$\langle\psi|\psi\rangle = \quad = \operatorname{Tr} T^{72}, \tag{K6}$$



$$\langle\psi|H|\psi\rangle = 72$$



The virtual space carries four spin-$\frac{1}{2}$, three spin-$\frac{3}{2}$ and one spin-$\frac{5}{2}$ multiplets, $4{\cdot}2{+}3{\cdot}4{+}6 = 26$. The contraction splits into blocks of dimension at most 162, evaluated in exact integers. The outcome, $E_0(72) < -72\kappa$, means that one Boltzmann factor of $Z_{72}(b)$ exceeds one, so

$$Z_{72}(b) > 1 \quad \text{for every } b > 0. \tag{K7}$$

A dominant eigenvalue. Suppose every eigenvalue in J had $|\lambda| \leq \ell = 0.999$. The capped-mass bound applied to J and to the three copies of N would give $Z_{72}(49/4) \leq 0.931 < 1$, contradicting Eq. (K7). Hence some $\lambda_0 \in J$ has $|\lambda_0| > \ell$: the spectrum looks like  , with one tall bar.

Removing λ_0 leaves the following bound on the kth moment of the rest:

$$R_A(k) = 3\,\mathcal{C}\big(m_N, v_N^{12}, \tfrac{k}{12}\big) + \big(m_J - \ell^{12}\big)^{k/12}. \tag{K8}$$

Initial defects. The two purities at the base scale now follow, each keeping only the dominant term:

$$P_{\mathrm{s}}(36, b) = \quad \geq \Big(1 + \frac{R_A(36)}{\ell^{36}}\Big)^{-2} = 1 - p_0,$$



$$P_\tau(72, b) = \frac{Z_{72}(2b)}{Z_{72}(b)^2} \geq \big(v_J^{72} + R_A(72)\big)^{-2} = 1 - q_0. \tag{K9}$$


The first line uses $Z_{72} \geq \lambda_0^{72}$. The second uses $Z_{72}(2b) > 1$ from the trial vector. Numerically $p_0 = 0.0479$ and $q_0 = 0.1815$.

Appendix L: Six doublings

The thermal defect q_0 exceeds the diagonal threshold $\varepsilon_* \simeq 0.127$ of Ref. [2]. A single common ε would therefore not start the iteration. The coupled update keeps the two defects apart instead:

$$p_{j+1} = \big\lceil f\big(1 - (1 - p_j)^2(1 - q_j)\big)\big\rceil_7, \qquad q_{j+1} = \big\lceil f\big(1 - (1 - q_j)^2(1 - p_{j+1})\big)\big\rceil_7, \tag{L1}$$

These are moves 1–4 of the doubling diagram of Ref. [1], with each output rounded up to a multiple of 10^{-7}.



The figure plots the defects on a logarithmic scale against the length $n_j = 36 \cdot 2^j$; the inverse temperature doubles alongside.

- The thermal defect falls from the first step.
- The spatial defect first *rises*, to 0.071 at $j = 3$: at small n the coupling identity costs more than squaring gains.

- Both then collapse together and cross 10^{-2} at the last step,

$$1 - P_{\mathrm{s}}(2304, 784) \le 0.0084513,$$
$$1 - P_{\tau}(4608, 784) \le 0.0054986. \tag{L2}$$

This recovers the initial estimate of Ref. [1] (Proposition 5.5 of Ref. [3]); re-evaluating the recurrence from the rational definitions of p_0 and q_0 confirms agreement digit for digit.

The weaker initial estimate and tighter seed chains. Ref. [3] also certifies $P_{\mathrm{s}}(60, 105/4)$, $P_{\tau}(120, 105/4) > 7/8$. That estimate uses a table at $b = 21/2$ for $n \le 10$, three twists, the trial vector at $N = 120$, and a temperature lift $\sum z_i^{5/2} \le (\sum z_i)^{5/2}$. It gives the weaker bound $\Delta_L > \frac{4}{105} \log \frac{80}{79}$ for every even $L \ge 60$. Evaluating the asymmetric rational recurrence directly from $(36, 49/4)$ with temperature lifting sharpens the uniform bound to $\Delta_L > 0.0729$ for every even $L \ge 36$ (`models/spin1`), while restricting to even-length traces $n \le 12$ with real SU(2) twists yields $\Delta_L > 0.039$ for every even $L \ge 60$ (`models/lean_seeds`).

Appendix M: Model-Dependent vs Universal Components

Each ingredient is a closed network, and only a few of them depend on the model.

- *Model independent:* the uniformization of Eq. (J4), the orbit reduction, the filters of Eq. (K3), the capped masses, and the doubling of Eq. (L1).
- *Model specific:*
  - the symmetry group that splits the spectrum into sectors, which for spin 1 makes N appear three times;
  - one good trial vector;
  - the sizes at which the filters isolate one eigenvalue.

Multi-level spectral cutout and trace reduction. In Eq. (K9), bounding $Z_{72}(2b) > 1$ from a single trial vector retains only the ground bar and discards the thermally populated single-magnon band, leaving $q_0 = 0.1815 > \varepsilon_*$ and requiring six pre-doublings from $n_0 = 36$ to 2304 (and twisted traces through $n = 12$, with 1.31×10^8 sparse nonzeros). At the seed scale ($n_0 = 60$, $2n_0 = 120$, $b = 49/4$), retaining the lowest 7 eigenvalues (27 states with multiplicity) in $Z_{120}(2b)$ raises the base-scale thermal purity from 0.761 to $0.9613 > 1 - \varepsilon_*$, eliminating all six pre-doublings and allowing the Poisson trace table to stop at $n \le 10$ (8.92×10^6 nonzeros, a 14.7× reduction; `models/lean_seeds`). Certifying those 7 levels at $N = 120$ requires a subspace deflation inequality $H_{120} \succeq E_{\mathrm{cut}}(I - QQ^\dagger) + Q\Lambda Q^\dagger$, because periodic-cluster Temple–Kato bounds have a negative denominator at $N = 120$ and ungrouped Krylov residual disks overlap across the 2.6×10^{-3} multiplet splitting.

Fast exact traces and the spin-2 starting-point barrier. For spin 1, rings of twelve sites have $3^{12} \simeq 5.3 \times 10^5$ states; fused three-step integer Horner kernels and packed 64-bit orbit lookup tables evaluate all eighteen twisted traces through $n = 12$ in under one minute on ten CPU threads (`models/spin1`). For spin 2, twelve-site rings have $5^{12} \simeq 2.4 \times 10^8$ states and the correlation length is roughly eight times longer ($\xi \approx 49$). Beyond that dimension growth, certified thermal MPO squaring (`CertifiedGaps.MPO`) at the estimated spin-2 seed ($n_0 = 256, \beta_0 = 64$) ($\beta_0 \|H\| \approx 9.8 \times 10^4$) faces a scalar starting-point barrier: initializing from a global quadratic polynomial $I - (\beta_0/2m)H + c(\beta_0/2m)^2 H^2$ requires $m \sim 10^6$–10^7 (≥ 20 squarings) before bond truncation enters, whereas local-gate Trotterization reduces the initial slice count by two orders of magnitude alongside a certified rational $D = 4$ matrix-product upper bound $E_0(256)/256 \le -4.725068$ (`models/spin2`).

[1] Companion note, *A diagrammatic proof of the Haldane gap via spacetime tensor networks*, `notes/thermal-mpo-gap-note.tex` (2026).

[2] Companion note, *Scope of the spacetime bootstrap*, `notes/thermal-mpo-gap-scope.tex` (2026).

[3] OpenAI, *The periodic spin-one Haldane gap*, manuscript dated September 24, 2026, Secs. 3–5 and Appendices A–C.

[4] B. L. Fox and P. W. Glynn, Computing Poisson probabilities, Commun. ACM **31**, 440 (1988).