A Diagrammatic Proof of the Haldane Gap via Tensor Networks

Sirui Lu
Max-Planck-Institut für Quantenoptik, Garching, Germany and
Munich Center for Quantum Science and Technology (MCQST), Schellingstrasse 4, D-80799 München, Germany
(Dated: November 7, 2026)

We give a diagrammatic exposition of OpenAI's computer-assisted proof of a uniform gap in the periodic spin-1 Heisenberg chain. Reading the partition-function tensor network along space and imaginary time shows how finite-ring bounds propagate across doubled lengths and inverse temperatures to all longer even rings. From the same finite inputs, we sharpen the spin-1 bound to $\Delta_L > 0.07290$ for all even $L \geq 36$, and extend the method to seven additional reflection-positive spin chains and ladders. We formalize the general theorem in Lean and verify a mixed-field Ising seed in the kernel.

Haldane predicted a gap for integer-spin antiferromagnetic Heisenberg chains, in contrast to their half-integer counterparts [1, 2]. For spin one, numerical work places the bulk gap near 0.4105 [3]. Unlike the Affleck–Kennedy–Lieb–Tasaki model [4], the Heisenberg chain is not frustration free: neighboring bonds cannot simultaneously attain their singlet ground states. Finite-size criteria based on frustration freeness [5, 6] therefore do not apply directly.

OpenAI recently proved a unique ground state and a uniform gap $\Delta_L > (\log 20)/784 \simeq 0.003821$ for every even periodic length $L \geq 2304$ from certified finite partition-function bounds [7]. Here we recast the proof in tensor-network diagrams so that each step—positivity of the spatial transfer matrix, torus gluing, and purity squaring—is read directly from a contraction identity. This formulation isolates reflection positivity as the condition behind the spatial inequalities, allowing us to certify uniform gaps for seven additional spin chains and ladders and to formalize the general theorem in Lean. Reanalyzing OpenAI's finite trace and trial-state inputs at a higher inverse temperature also sharpens the periodic spin-1 bound to $\Delta_L > 0.07290$ for every even $L \geq 36$ [8].

Thermal purity and the gap.— Consider


$$H_L = \sum_{j=1}^{L} \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}, \qquad Z_L(\beta) = \operatorname{Tr} e^{-\beta H_L}, \tag{1}$$

with periodic indices, spin one, even $L \geq 4$, and exchange coupling one. Ordering the eigenvalues of H_L with multiplicity defines the gap $\Delta_L = E_1(L) - E_0(L)$ (). For $\beta > 0$, the Gibbs state $\rho_L = e^{-\beta H_L}/Z_L(\beta)$ has ordered weights $w_0 \geq w_1 \geq \cdots$. Its thermal purity $P_\tau(L,\beta) = \operatorname{Tr} \rho_L^2 \leq w_0$ gives

$$e^{-\beta\Delta_L} = \frac{w_1}{w_0} \leq \frac{1-w_0}{w_0} \leq \frac{1-P_\tau(L,\beta)}{P_\tau(L,\beta)}. \tag{2}$$



When $P_\tau(L,\beta)$ is close to one, a single weight dominates the Gibbs spectrum,  : $P_\tau > 1/2$ forces $w_0 > 1/2$ and hence a unique ground state, while $0 < 1 - P_\tau \leq q < 1/2$ gives $\Delta_L \geq \beta^{-1} \log[(1-q)/q]$.

Proof strategy.— To keep this gap bound positive as β grows with L, $\log(1/q)$ must scale proportionally to β. Squaring and normalizing the Gibbs state doubles β at fixed L; OpenAI's key idea is to pair ρ_L with a positive spatial state σ_n whose square doubles the ring length n. Starting from an even seed ring (n_0, β_0), we alternate the two doublings along the staircase:




Because each step squares the purity defects while doubling $\beta_j = 2^j \beta_0$, their negative logarithms grow as 2^j and dividing by β_j leaves a uniform positive lower bound on the gap. Carrying out this staircase requires constructing σ_n, coupling space and imaginary time, and interpolating across the even lengths between n_j and $2n_j$.

Spatial transfer matrix.— Set $\tau = \beta/M$ and let $S_L(\tau)$ be the product of $I - \tau h_{j,j+1}$ over even bonds followed by odd bonds, where $h = \boldsymbol{S} \cdot \boldsymbol{S}$. Contracting the resulting brickwork network in imaginary time or in space gives [9]

$$\underbrace{\operatorname{Tr} S_L(\beta/M)^M}_{\text{  }} = Z_L^{(M)}(\beta) = \underbrace{\operatorname{Tr} T_{\beta,M}^{L/2}}_{\text{  }} . \tag{3}$$

In the Cartesian basis $|x\rangle, |y\rangle, |z\rangle$, the generators $G_\alpha = iS^\alpha$ are real antisymmetric matrices. Hence

$$-h = \sum_{\alpha=x,y,z} G_\alpha \otimes G_\alpha, \qquad I - \tau h = \sum_{\gamma=0,x,y,z} A_\gamma \otimes A_\gamma, \tag{4}$$

where $A_0 = I$ and $A_\alpha = \sqrt{\tau}\, G_\alpha$; diagrammatically,  $= \sum_\gamma$  . These matching real bond factors express reflection positivity [10–12].

For bond-label sequences $a, b \in \{0, x, y, z\}^M$, define the real trace matrix $\mathcal{F}_{ba} = \operatorname{Tr}[(A_{b_1} A_{a_1}) \cdots (A_{b_M} A_{a_M})]$.

Because the two sites sharing an even bond carry identical factor orderings, summing their shared bond sequence forms a positive two-site Gram transfer matrix independent of L:


$$T_{\beta,M} = \mathcal{F}^{\mathsf{T}}\mathcal{F} \geq 0, \qquad \mathcal{F}^{\mathsf{T}}\,\mathcal{F} = T_{\beta,M}. \tag{5}$$

For even $n \geq 4$, normalizing $T_{\beta,M}^{n/2}$ by its positive trace ($\mathcal{F}_{00} = 3$ on the all-zero sequence) defines a density matrix $\sigma_n^{(M)}$. Its nonnegative eigenvalues yield closed inequalities for spatial purity, purity squaring, and power-sum interpolation at finite M. Taking $M \to \infty$ at each fixed (L, β) sends $S_L(\beta/M)^M \to e^{-\beta H_L}$ and passes those scalar inequalities to the exact partition functions $Z_L(\beta)$ before the doubling recurrence begins (Sec. SII.A of Ref. [8]).

Coupling and squaring.— Representing $Z_n(\beta)$ by the $n \times \beta$ torus, the spatial and thermal purities join two normalized copies side by side ($\sigma_n^{(M)}\,\sigma_n^{(M)}$ at finite M) or stack them in imaginary time (ρ_n / ρ_n):

$$P_{\mathrm{s}}(n,\beta) = \lim_{M\to\infty} \mathrm{Tr}[(\sigma_n^{(M)})^2] = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2},$$
$$P_\tau(n,\beta) = \mathrm{Tr}\,\rho_n(\beta)^2 = \frac{Z_n(2\beta)}{Z_n(\beta)^2}. \tag{6}$$

Both lie in $(0, 1]$. Assembling the $2n \times 2\beta$ torus from four copies of $n \times \beta$ in either order,



gives the exact coupling identity

$$P_{\mathrm{s}}(n,2\beta)P_\tau(n,\beta)^2 = P_\tau(2n,\beta)P_{\mathrm{s}}(n,\beta)^2. \tag{7}$$


For probabilities u_i with purity $P = \sum_i u_i^2$, squaring and renormalizing concentrates the weights, $\mapsto$, giving a new purity P' that obeys

$$1 - P' = \frac{2\sum_{i<j}(u_i u_j)^2}{P^2} \leq \frac{(1-P)^2}{2P^2} = f(1-P), \tag{8}$$

where $f(z) = z^2/[2(1-z)^2]$. This squaring map acts identically along space and imaginary time:

$$\sigma_n^{(M)}\,\sigma_n^{(M)} = \mathrm{Tr}[(\sigma_n^{(M)})^2]\,\sigma_{2n}^{(M)},$$
$$\begin{matrix}\rho_n(\beta)\\ \rho_n(\beta)\end{matrix} = P_\tau(n,\beta)\,\rho_n(2\beta). \tag{9}$$

To advance defect bounds $1 - P_{\mathrm{s}}(n,\beta) \leq p$ and $1 - P_\tau(2n,\beta) \leq q$ from (n,β) to $(2n,2\beta)$, we alternate torus gluing with squaring along the two directions:



Dropping $P_\tau(n,\beta)^2 \leq 1$ in Eq. (7) gives $P_{\mathrm{s}}(n,2\beta) \geq (1-p)^2(1-q)$, and spatial squaring yields p'. Feeding p' into Eq. (7) at length $2n$ with $P_{\mathrm{s}}(2n,\beta)^2 \leq 1$ gives $P_\tau(4n,\beta) \geq (1-q)^2(1-p')$, and thermal squaring yields q':

$$p' = f(1-(1-p)^2(1-q)),$$
$$q' = f(1-(1-q)^2(1-p')). \tag{10}$$

Whenever $0 \leq p, q, p', q' < 1$, the pair (p', q') bounds $1 - P_{\mathrm{s}}(2n,2\beta)$ and $1 - P_\tau(4n,2\beta)$.

The finite seed and uniform gap.— Once both defects are at most 0.01, the scalar inequality $f(1-(1-\varepsilon)^3) \leq 5\varepsilon^2$ drives all further doubling steps analytically (Sec. SII.F of Ref. [8]). To reach that regime, we start from OpenAI's certified finite seed:

Lemma 1 (Finite seed [7]). *At $\beta_0 = 49/4$, the periodic spin-1 chain satisfies $P_{\mathrm{s}}(36,\beta_0) \geq 1 - 47916/10^6$ and $P_\tau(72,\beta_0) \geq 1 - 181521/10^6$.*

Although $q_0 = 0.181521$ starts above the symmetric fixed point $\varepsilon_* \simeq 0.1273$, the smaller spatial defect $p_0 = 0.047916$ pulls q' down in Eq. (10). Six rationally verified, upward-rounded updates carry the seed into $[0, 0.01]^2$ with $p_6 = 0.0084709$ and $q_6 = 0.0055149$ at $(n_6, \beta_6) = (2304, 784)$ (Fig. 1; exact orbit in Sec. SII.E of Ref. [8]).

Restarting at $(2304, 784)$ with $\varepsilon_0 = 0.01$ and $\varepsilon_{j+1} = 5\varepsilon_j^2$ gives $\varepsilon_j = 20^{-2^j}/5$ at $n_j = 2304\,2^j$ and $\beta_j = 784\,2^j$. To cover every even $L \geq 2304$, choose j with $n_j \leq L \leq 2n_j$. Power-sum monotonicity of the nonnegative spatial weights gives $Z_v(\beta) \leq Z_u(\beta)^{v/u}$ for even $v \geq u$; applying this bound at $2\beta_j$ in the numerator and at β_j in the denominator of $P_\tau(L,\beta_j)$ yields

$$P_\tau(L,\beta_j) \geq [P_\tau(2n_j,\beta_j)P_{\mathrm{s}}(n_j,\beta_j)^2]^{L/(2n_j)}$$
$$\geq (1-\varepsilon_j)^3 \geq 1 - 3\varepsilon_j. \tag{11}$$

Thus the ground state is unique, and $\varepsilon_j \leq 1/100$ gives $3\varepsilon_j/(1-3\varepsilon_j) \leq (60/97)20^{-2^j} < 20^{-2^j}$. Equation (2)




FIG. 1. The asymmetric recurrence (10) and its spin-1 seed orbit at $n_k = 36\,2^k$ and $\beta_k = (49/4)2^k$. Six rationally verified, upward-rounded steps carry the seed at $n_0 = 36$ into the analytical target square $[0, 0.01]^2$ (gray) at $n_6 = 2304$; the teal region shows the numerical basin of attraction.

therefore proves

$$\Delta_L > \frac{\log 20}{784} \simeq 0.003821 \qquad \text{for every even } L \geq 2304. \tag{12}$$

Finite seed bounds.— Proving Lemma 1 at $\beta_0 = 49/4$ reduces to bounding the numerators of

$$P_s(36, \beta_0) = \frac{Z_{72}(\beta_0)}{Z_{36}(\beta_0)^2}, \qquad P_\tau(72, \beta_0) = \frac{Z_{72}(2\beta_0)}{Z_{72}(\beta_0)^2} \tag{13}$$

from below and their denominators from above. Ref. [7] supplies these four bounds from short-ring traces on $4 \leq n \leq 12$ sites and a 72-site matrix-product trial state:

	Quantity	Bound	Method
	$Z_{72}(\beta_0)$	lower	MPS + small-ring traces
	$Z_{36}(\beta_0)$	upper	MPS + small-ring traces
	$Z_{72}(2\beta_0)$	lower	MPS
	$Z_{72}(\beta_0)$	upper	small-ring traces

We summarize the five steps that turn these finite inputs into Lemma 1 (terminating decimals defining bounds denote exact rationals; full details in Sec. SIII of Ref. [8]).

Step 1: Spatial trace moments. To use both odd and even short-ring traces ($4 \leq n \leq 12$) to constrain the partition functions at $n = 36$ and $n = 72$, we use the continuous-time self-adjoint one-site transfer operator K_β of Ref. [7]:

$$Z_n(\beta) = \operatorname{Tr} K_\beta^n = \quad , \qquad n \geq 4. \tag{14}$$

Shifting the energy per site by $\kappa = 700741/500000$ (chosen so that the 72-site trial energy per site in Step 3 is below $-\kappa$) defines

$$\widehat{Z}_n(\beta) = e^{-n\beta\kappa} Z_n(\beta), \qquad \widehat{K} = e^{-\beta_0\kappa} K_{\beta_0}, \qquad \widehat{Z}_n(\beta_0) = \sum_i \lambda_i^n, \tag{15}$$

with real eigenvalues λ_i of either sign. The shift cancels from both purities, and every ring length $n \geq 4$ samples the same eigenvalue list $\{\lambda_i\}$.

Step 2: Separate the moments by symmetry. The commuting π spin rotations split the spectrum of $\widehat{K}$ into four sign sectors: a rotation-invariant sector J and three sectors with identical spectra N, permuted by cyclic axis rotations. Twisting the closing bond of the ring by a π rotation P negates two of the three N sectors, so the untwisted and twisted traces decompose as

$$\widehat{Z}_n = J_n + 3N_n, \qquad \widehat{Z}_n(P) = J_n - N_n, \tag{16}$$

where $J_n = \sum_{i \in J} \lambda_i^n$ and $N_n = \sum_{i \in N} \lambda_i^n$ at β_0. Integer Poisson–Horner trace recursions enclose both traces for $4 \leq n \leq 12$; inverting Eq. (16) gives the twelfth-moment upper bounds

$$J_{12} \leq m_J = 1.0657205, \qquad N_{12} \leq m_N = 0.2783155. \tag{17}$$

Step 3: A trial state gives a lower bound. Integer tensors of bond dimension 26 define a periodic 72-site MPS ψ whose norm and one-bond energy expectation contract exactly as 72-step matrix traces:


$$\mathcal{N} = \langle \psi, \psi \rangle = \quad , \qquad \mathcal{E} = \langle \psi, h_{1,2}\psi \rangle = \quad . \tag{18}$$

Translation invariance gives $\langle \psi, H_{72}\psi \rangle/(72\mathcal{N}) = \mathcal{E}/\mathcal{N} < -\kappa$, so $E_0(72) < -72\kappa$ by the variational principle. Retaining the ground-state Boltzmann weight gives

$$\widehat{Z}_{72}(\beta) \geq e^{-\beta(E_0(72)+72\kappa)} > 1 \qquad (\beta > 0), \tag{19}$$

which supplies both $\sum_i \lambda_i^{72} > 1$ at β_0 and the thermal numerator bound $\widehat{Z}_{72}(2\beta_0) > 1$.

Step 4: A lower bound on the leading spatial eigenvalue. To bound the spatial numerator $\widehat{Z}_{72}(\beta_0) \geq \lambda_0^{72}$ from below, we show that J contains a leading eigenvalue with $|\lambda_0| > \ell = 0.999$. Write $u_i = \lambda_i^{12} \geq 0$, so $\lambda_i^{72} = u_i^6$. If every $|\lambda_i| \leq \ell$, then each $u_i \leq \ell^{12}$; because $u \mapsto u^6$ is convex and $\ell^{12} < m_J < 2\ell^{12}$, the sum $\sum_{i \in J} u_i^6$ at total mass at most m_J is maximized by placing one entry at the cap ℓ^{12} and the remainder at $m_J - \ell^{12}$:



Raising to the sixth power suppresses the remainder and bounds each N sector by m_N^6, yielding

$$\begin{aligned}\widehat{Z}_{72}(\beta_0) &\le \ell^{72} + (m_J - \ell^{12})^6 + 3m_N^6 \\ &< 0.932 < 1,\end{aligned} \tag{20}$$

which contradicts Eq. (19). Because $m_N < \ell^{12}$, the eigenvalue with $|\lambda_0| > \ell$ must lie in J.

Step 5: Polynomial bounds on the remaining spectrum. To bound the denominators $\widehat{Z}_{36}(\beta_0)$ and $\widehat{Z}_{72}(\beta_0)$ from above, we cap the remaining eigenvalues in each sector $\mathcal{S} \in \{J, N\}$ using a nonnegative polynomial $F(x) = \sum_{j=4}^{12} f_j x^j$:

$$F(\lambda_i) \le \sum_{a\in\mathcal{S}} F(\lambda_a) = \sum_{j=4}^{12} f_j \mathcal{S}_j \le B, \qquad i \in \mathcal{S}. \tag{21}$$

No eigenvalue in $\mathcal{S}$ can lie where $F(x) > B$; two explicit filters rule out $|\lambda| \ge U = 1.00139$ in J and $|\lambda| \ge V = 0.872$ in N (Sec. SIII.A.c of Ref. [8]).

Removing $\lambda_0^{12} > \ell^{12}$ leaves residual twelfth moment at most $m_J - \ell^{12}$ in J, while convexity with cap V^{12} ($V^{12} < m_N < 2V^{12}$) bounds each N sector by one entry at V^{12} and the remainder $m_N - V^{12}$. For $k = 36, 72$, the subleading moment $\sum_{i\neq 0} \lambda_i^k$ is therefore at most

$$\begin{aligned}R_k = &(m_J - \ell^{12})^{k/12} \\ &+ 3\left[V^k + (m_N - V^{12})^{k/12}\right].\end{aligned} \tag{22}$$

Using $\widehat{Z}_{72}(\beta_0) \ge \lambda_0^{72}$ and $\widehat{Z}_{36}(\beta_0) \le |\lambda_0|^{36} + R_{36}$ with $|\lambda_0| > \ell$ for $P_{\rm s}(36, \beta_0)$, and using $\widehat{Z}_{72}(2\beta_0) > 1$ and $\widehat{Z}_{72}(\beta_0) \le U^{72} + R_{72}$ for $P_\tau(72, \beta_0)$, gives

$$\begin{aligned}P_{\rm s}(36, \beta_0) &\ge [1 + R_{36}/\ell^{36}]^{-2} = 1 - p, \\ P_\tau(72, \beta_0) &\ge [U^{72} + R_{72}]^{-2} = 1 - q,\end{aligned} \tag{23}$$

and exact rational evaluation verifies Lemma 1:

$$p \le \frac{47916}{10^6}, \qquad q \le \frac{181521}{10^6}. \tag{24}$$

For physical Boltzmann weights z_i at β_0, the inequality $\sum_i z_i^r \le (\sum_i z_i)^r$ transfers the short-ring trace bounds to $r\beta_0$, with $r = 17/8$. Together with the 120-site trial state of Ref. [7], these give smaller initial defects at $n_0 = 60$. A finite prefix of asymmetric updates then certifies $\Delta_L > 0.07290$ for all even $L \ge 60$; a sharpened unlifted bound at $(36, \beta_0)$ bridges $36 \le L \le 72$, covering every even $L \ge 36$ (Sec. SIII.B of Ref. [8]; certificates in the repository).

Reflection-positive models.— The spatial Gram factorization (5) relies on locality and bond reflection positivity [11, 12], not on SU(2) symmetry. Consider a periodic ring of $L \ge 2$ unit cells with Hamiltonian

$$H_L = \sum_{j=1}^{L} h_{j,j+1} + \sum_{j=1}^{L} o_j, \qquad j + L \equiv j, \tag{25}$$

where h couples neighboring cells and o acts within a cell (for $L = 2$, the sum retains both periodic bonds $h_{1,2} + h_{2,1}$). Suppose that in some local spin basis there is a real symmetric involution F ($F^2 = I$, such as spin exchange within a dimer) for which o is real symmetric with $FoF = o$ and

$$-h = \sum_\alpha C_\alpha \otimes FC_\alpha F, \tag{26}$$

with each C_α real symmetric or antisymmetric. Conjugating odd cells by F matches their single-cell trace matrix $\mathcal{F}$ to that of even cells, so the two-cell transfer matrix again takes the positive Gram form $T_{\beta,M} = \mathcal{F}^{\mathsf{T}}\mathcal{F} \ge 0$ of Eq. (5) (Sec. SII.A of Ref. [8]), and the same spatial purity inequalities and doubling recurrence apply.

An admissible seed at even n_0 then certifies a unique ground state and a uniform gap for every even $L \ge n_0$ via finite-prefix interpolation and an analytic tail bound ✓L. Table I applies this theorem to staggered-field chains [13], bond-alternating chains [14, 15], chains with single-ion anisotropy [16], Heisenberg ladders [17], and Ising models, with finite seeds certified from verified short-ring spectra or exact matrix arithmetic.

Formal verification.— The marker ✓L links to a Lean theorem with its stated hypotheses. Formalizing the continuous-time path-space kernel of Ref. [7] in Lean would require measure-theoretic trace identities not yet in `mathlib`. Our finite-Trotter proof replaces that kernel by the finite Gram matrix $T_{\beta,M} = \mathcal{F}^{\mathsf{T}}\mathcal{F} \ge 0$ and passes the resulting moment inequalities to $Z_L(\beta)$ through the fixed-(L, β) matrix product limit before iterating the recurrence ✓L.

We formalize the general reflection-positive gap theorem in a Lean library and instantiate it for every model in Table I. For the mixed-field Ising chain at $(g, h) = (1/4, 1/20)$, we also verify the finite seed arithmetic directly in the Lean kernel to obtain an unconditional theorem; for the spin-1 chain, we check the MPS trial contractions in Lean and take the remaining certified short-ring trace bounds as hypotheses.

Discussion.— Recasting the proof in tensor-network diagrams and visual summaries [18] separates its universal contraction and squaring steps from the model-specific seed bounds. That separation let us certify uniform gaps for seven additional reflection-positive chains and ladders (Table I), sharpen the periodic spin-1 bound from the same finite inputs, and give a diagrammatic

TABLE I. Uniform gap bounds for reflection-positive models. Every even length at or above the stated threshold is covered. A cell or rung contains two spins; L counts the units shown. Spin operators have $\boldsymbol{S}^2 = S(S+1)I$; σ^α are Pauli matrices. The last column lists uncertified finite-ring gap estimates on 12–20 spins (bulk DMRG estimate [3] for spin-1 Heisenberg); the mixed-field Ising bound is verified end to end in the Lean kernel.

Model	Hamiltonian	Even lengths	Gap $\geq$	Numerical
Staggered field, $S = \frac{1}{2}$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1} - \sum_j (-1)^j S^z_j$	$L \geq 8$ sites	1.1489	1.874
Dimerized, $S = \frac{1}{2}$	$\sum_j [1 + (-1)^j/2]\, \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 4$ cells	0.5347	1.189
Dimerized, $S = 1$	$\sum_j [1 + 9(-1)^j/10]\, \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 2$ cells	0.5288	1.763
Single-ion anisotropy, $S = 1$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1} + 4\sum_j (S^z_j)^2$	$L \geq 12$ sites	0.8994	2.269
Heisenberg ladder, $S = \frac{1}{2}$	$\sum_{j,a=1,2} \boldsymbol{S}_{j,a} \cdot \boldsymbol{S}_{j+1,a} + 4\sum_j \boldsymbol{S}_{j,1} \cdot \boldsymbol{S}_{j,2}$	$L \geq 4$ rungs	1.1024	3.138
Ising ladder	$-\sum_j (\sigma^z_{j,1}\sigma^z_{j+1,1} + \sigma^z_{j,2}\sigma^z_{j+1,2} + \sigma^z_{j,1}\sigma^z_{j,2}) - 3\sum_{j,a=1,2} \sigma^x_{j,a}$	$L \geq 4$ rungs	0.6368	2.662
Mixed-field Ising	$-\sum_j \sigma^z_j \sigma^z_{j+1} - \frac{1}{4}\sum_j \sigma^x_j - \frac{1}{20}\sum_j \sigma^z_j$	$L \geq 6$ sites	0.3922	1.190
Heisenberg, $S = 1$	$\sum_j \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$	$L \geq 36$ sites	0.0729	0.410

proof of the open-chain boundary-field theorem by tracking the overlap of the boundary vectors with the leading spatial eigenvector [8, 19].

A natural next target in one dimension is the spin-2 Heisenberg chain, whose smaller gap and longer correlation length push the seed scale to larger rings and lower temperatures [20]. In two spatial dimensions, torus gluing links one thermal and two spatial purities, requiring spatial moment inequalities on a torus and certified finite-torus inputs; certifying such a seed for the honeycomb AKLT model [21, 22] remains open.

Finally, our Lean formalization highlights the gap between proof assistants and scientific computing environments: while our mixed-field Ising proof verifies its finite seed inside the Lean kernel, scaling in-kernel verification to the spin-1 trace recursions calls for tighter integration between formal kernels, symbolic algebra systems, and verified numerical linear algebra.

We thank Xiao-Liang Qi and Ignacio Cirac for helpful discussions. Part of this work was carried out at the Kavli Institute for Theoretical Physics and during a visit to Xiao-Liang Qi at Stanford University; we thank both institutions for their hospitality. An initial version of this work [23] was circulated and discussed among colleagues at Stanford on October 7, 2026, and we thank Shengqi Sang, Yimu Bao, and Junkai Dong for helpful discussions. A presentation of this work was given at KITP [24]. The code, gap certificates, Lean 4 formalization and blueprint, interactive explainer, and open challenges are available at https://github.com/LionSR/CertifiedGaps (see also https://sirui-lu.com/CertifiedGaps/).

AI-use declaration.—S. L. prompted GPT-6-Astra (OpenAI) to recast the proofs of Refs. [7, 19] in tensor-network diagrams, and used TeXRA [25] and OpenAI Codex with GPT-6-Astra and GPT-6-Sol for subsequent calculations, formalization, and manuscript preparation. All mathematical arguments, numerical certificates, and formal proofs were verified by the author.

[1] F. D. M. Haldane, Phys. Lett. A **93**, 464 (1983).
[2] F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).
[3] S. R. White and D. A. Huse, Phys. Rev. B **48**, 3844 (1993).
[4] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).
[5] S. Knabe, J. Stat. Phys. **52**, 627 (1988).
[6] D. Gosset and E. Mozgunov, J. Math. Phys. **57**, 091901 (2016), arXiv:1512.00088 [quant-ph].
[7] OpenAI, The periodic spin-one Haldane gap, OpenAI Math Release, result 268, preprint (2026).
[8] See Supplemental Material for the open-chain proof, the finite-dimensional transfer construction and periodic bootstrap, and the finite seed bounds and temperature-lifting estimates.
[9] M. Suzuki, Phys. Rev. B **31**, 2957 (1985).
[10] F. J. Dyson, E. H. Lieb, and B. Simon, J. Stat. Phys. **18**, 335 (1978).
[11] J. Fröhlich, R. Israel, E. H. Lieb, and B. Simon, Commun. Math. Phys. **62**, 1 (1978).
[12] A. Jaffe and B. Janssens, Commun. Math. Phys. **346**, 1021 (2016), arXiv:1506.04197 [math-ph].
[13] M. Oshikawa and I. Affleck, Phys. Rev. Lett. **79**, 2883 (1997).
[14] T. Barnes, J. Riera, and D. A. Tennant, Phys. Rev. B **59**, 11384 (1999).
[15] S. Yamamoto, Phys. Rev. B **51**, 16128 (1995).
[16] W. Chen, K. Hida, and B. C. Sanctuary, Phys. Rev. B **67**, 104401 (2003).
[17] E. Dagotto and T. M. Rice, Science **271**, 618 (1996).
[18] E. R. Tufte, *Envisioning Information* (Graphics Press, Cheshire, Connecticut, 1990).
[19] OpenAI, A boundary-field gap for the spin-one Heisen-

berg chain, OpenAI Math Release, result 268, preprint (2026).
[20] S. Todo and K. Kato, Phys. Rev. Lett. **87**, 047203 (2001).
[21] M. Lemm, A. W. Sandvik, and L. Wang, Phys. Rev. Lett. **124**, 177204 (2020), arXiv:1910.11810 [quant-ph].
[22] N. Pomata and T.-C. Wei, Phys. Rev. Lett. **124**, 177203 (2020), arXiv:1911.01410 [cond-mat.stat-mech].
[23] S. Lu and GPT-6-Astra, A short diagrammatic explanation of the spin-1 Haldane gap proof (2026), October 7, 2026.
[24] S. Lu, Bridging gaps with tensor networks: The Haldane gap proof and beyond, Talk at the KITP program *Intermediate-Scale AI for Physics* (2026), October 20, 2026 (video recording).
[25] TeXRA, TeXRA: A multi-agent research assistant for theorists (2025).
[26] T. Kennedy, J. Phys.: Condens. Matter **2**, 5737 (1990).
[27] W. Tang, H.-H. Tu, and L. Wang, Phys. Rev. Lett. **125**, 170604 (2020), arXiv:2004.12928 [cond-mat.str-el].

Supplemental Material

CONTENTS

SI. Open chains with boundary fields 7
SI.A. The boundary-field chain and the additional estimate 7
SI.B. Boundary vectors and boundary conditions 9
SI.C. Finite inputs and ring estimates 10
SI.D. Three steps for the open chain 11
a. Step 1: control the new boundary remainder 12
b. Step 2: double the length 13
c. Step 3: square in imaginary time 13
SI.E. Initialization and the small-defect induction 13
SI.F. Equal fields, odd lengths, and the gap 14

SII. Finite Trotter networks and the periodic doubling argument 15
SII.A. Finite Trotter slices and a two-site transfer matrix 15
SII.B. Finite-transfer moment inequalities 17
SII.C. Controlled product limit 18
SII.D. Torus identities and asymmetric doubling 18
SII.E. Six exact rational steps 19
SII.F. Scalar tail, interpolation, and the strict gap 20

SIII. Finite seed certificates and improved bounds 22
SIII.A. The base seed at length 36 22
a. Hamiltonians, twists, and the spatial transfer operator 22
b. Exact finite trace and trial inputs 23
c. Sector moments and polynomial filters 24
d. Capped masses and the two base defects 25
SIII.B. Temperature lifting and the improved uniform bound 26
a. Lifted moments and a dominant spatial weight 26
b. The physical thermal residual 27
c. Finite-prefix coverage and the doubling tail 27
SIII.C. Reproducibility and arithmetic assumptions 28

Appendix SI: Open chains with boundary fields

SI.A. The boundary-field chain and the additional estimate

We now extend the diagrammatic doubling argument from periodic rings to open chains with endpoint magnetic fields. Consider the spin-1 Heisenberg chain with equal endpoint fields, ,

$$H_n^{\mathrm{s}} = \sum_{i=1}^{n-1} \boldsymbol{S}_i \cdot \boldsymbol{S}_{i+1} - h(S_1^z + S_n^z), \qquad h = \frac{3}{5}, \tag{S1}$$

where $\sum_\alpha (S^\alpha)^2 = 2I$ and the exchange coupling is one. We give a diagrammatic proof of the following theorem from Ref. [19], using its certified finite inputs.

Theorem S1 (Boundary-field gap)**.** *For every odd $n \geq 1921$, H_n^{s} has a unique ground state and*

$$\Delta_n > \frac{\log 10}{392} \simeq 0.005874.$$

Without the endpoint fields, the effective spin-1/2 modes at the two ends produce four low-lying states whose splitting is exponentially small in the chain length [26]. Applying equal fields on an odd chain lifts this boundary degeneracy [19].

Our strategy is to bound the thermal purity of the open chain at a scale β proportional to the length, just as in the periodic argument. List the energies of H_n^{s} with multiplicity and let $w_0 \geq w_1 \geq \cdots$ be their Gibbs weights. With $P_\tau^{\mathrm{s}}(n,\beta) = \sum_m w_m^2$ and $\Delta_n = E_1(n) - E_0(n)$, we have


$$w_0 \geq P_\tau^{\mathrm{s}}(n,\beta), \qquad e^{-\beta\Delta_n} \leq \frac{1 - P_\tau^{\mathrm{s}}(n,\beta)}{P_\tau^{\mathrm{s}}(n,\beta)}. \tag{S2}$$

A thermal purity above 1/2 forces a unique ground state and bounds the gap from below. What changes in the open-chain setting is how we propagate the thermal purity across doubled lengths and inverse temperatures.

To see where the new difficulty enters, consider the spatial transfer operator at a fixed temperature. Suppose that X_β has a simple positive leading eigenvalue λ,  , and write the normalized transfer operator as

$$T := X_\beta/\lambda = \Pi + R, \qquad \Pi = |\psi\rangle\langle\psi|, \qquad \Pi R = R\Pi = 0, \quad \|\psi\| = 1. \tag{S3}$$

In a periodic ring, the trace weights the leading projector Π by one. In an open chain, the chain ends are contracted with boundary vectors, so the leading contribution is weighted instead by how much of v lies along ψ,  :



(S4)

Because c is not fixed by normalization and could a priori vanish, squaring in space no longer closes on its own. For an even length m, the normalized boundary matrix element splits into the leading overlap and a nonnegative remainder,

$$\begin{aligned} B_m &= \bar{v} \text{—} T^m \text{—} v \\ &= \bar{v} \text{—} \Pi \text{—} v + \bar{v} \text{—} R^m \text{—} v = c + D_m, \\ D_m &= \|R^{m/2} v\|^2 \geq 0. \end{aligned} \tag{S5}$$

To show that the leading eigenvector dominates B_m, we must bound the remainder's share of B_m,

$$e_m := \frac{D_m}{B_m} = 1 - \frac{c}{B_m}, \tag{S6}$$

the grey fraction of at the seed value $e_{142} \leq .14$ of Eq. (S12), rather than the absolute remainder D_m alone. Increasing the chain length suppresses D_m by powers of $\|R\|$ while leaving c unchanged.

When we double the inverse temperature, however, both the boundary vector v and the leading eigenvector ψ change. The overlap c at the old temperature does not by itself bound the new overlap c'. The key idea is to compare two even lengths at the new temperature. Writing primes for quantities at 2β, the difference

$$B'_k - B'_m = D'_k - D'_m$$

cancels the unknown overlap c' because c' is independent of length. An exact thermal-purity bridge then relates the ratio B'_k/B'_m back to known quantities at the old temperature.

The proof therefore proceeds in four stages:

1. In Sec. SI.B, we express the open-chain partition functions as boundary contractions of the continuous-time transfer operator. Putting the same boundary vector at both ends produces an auxiliary opposite-field chain with nonnegative remainders at even lengths, while a π spin rotation at one end restores equal fields.

2. In Sec. SI.C, we use the periodic doubling recurrence to control the bulk transfer spectrum and spectral ratio across all scales, and we state the four finite seed bounds needed to start the open-chain recurrence.

3. In Secs. SI.D and SI.E, we combine the two-length boundary comparison, length doubling, and thermal squaring into a coupled recurrence for e_m and the even-length opposite-field thermal purity, and we run three explicit initialization steps before entering the small-defect induction.

4. In Sec. SI.F, we pass from even opposite-field chains to odd equal-field chains by bounding the signed remainder in absolute value, completing the proof of Theorem S1.

SI.B. Boundary vectors and boundary conditions

We first represent the open-chain partition functions as transfer-operator matrix elements. We use the continuous-time bond expansion of Ref. [19]: we sum over the times of bond insertions instead of placing them on a finite Trotter grid. In the Cartesian basis, $G_\alpha = iS^\alpha$ is real antisymmetric and

$$-\boldsymbol{S}_i \cdot \boldsymbol{S}_{i+1} = \sum_{\alpha=x,y,z} G_{\alpha,i} G_{\alpha,i+1}.$$

A bond history a is an ordered list of events (τ, α) with $0 < \tau < \beta$. The measure μ_β sums the labels and integrates the ordered times; its total mass is $e^{3\beta}$, including unit mass on the empty history. The local tensor is

$$W_\beta^{ss'}(a,b) = e^{-\kappa\beta} \langle s | \mathcal{T}_\tau \prod_{(\tau,\alpha)\in a\cup b} G_\alpha | s' \rangle,$$
$$\kappa = \frac{700741}{500000}. \tag{S7}$$

Time ordering $\mathcal{T}_\tau$ places later events to the left according to the interleaved times in $a \cup b$, and tracing the physical indices defines the real symmetric transfer kernel $X_\beta(a,b)$ on $L^2(\mu_\beta)$.

The energy shift κ is the one used in Ref. [19]. Throughout this section, Z_n and all open-chain partition functions include the shift $\kappa n I$; it places the leading transfer eigenvalue near one at the seed scales and cancels from every purity and spectral gap.

To represent a boundary field along z, we cap the virtual bond history with the boundary vector

$$v_\beta(a) = \begin{cases} (ih)^{|a|}, & \text{all event labels are } z, \\ 0, & \text{otherwise,} \end{cases} \qquad \|v_\beta\|^2 = e^{\beta h^2}. \tag{S8}$$

Its value on the empty history is one, and its norm follows by summing $(\beta h^2)^k/k!$ over the number k of z events.

The cap inserts boundary events into the same time-ordered product as the bond events. Summing their counts and integrating their times adds the corresponding field term to the Hamiltonian. We now check the signs of the two endpoint fields produced by v_β. The conjugated left cap supplies $(-ih)G_z = +hS^z$ in the exponential, which corresponds to the Hamiltonian term $-hS_1^z$. The unconjugated right cap supplies $(ih)G_z = -hS^z$, corresponding to $+hS_n^z$. Contracting with the same boundary vector v_β at both ends therefore gives the auxiliary opposite-field chain $\uparrow\!\cdots\!\downarrow$,

$$H_n^{\mathrm{o}} = H_n^{\mathrm{s}} + 2hS_n^z. \tag{S9}$$

To obtain equal endpoint fields, let $P = \mathrm{diag}(1,-1,-1)$ be the spin rotation by π about the x axis. Its virtual action U_P negates each z event, which reverses the right endpoint field and produces H_n^{s}, $\uparrow\!\cdots\!\uparrow$. In diagram form,

$$e^{-\beta(H_n^{\mathrm{o}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } W_\beta \text{ — } \cdots \text{ — } W_\beta \text{ — } v_\beta,$$
$$Y_n(\beta) := \mathrm{Tr}\, e^{-\beta(H_n^{\mathrm{o}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } X_\beta^n \text{ — } v_\beta,$$
$$Y_n^{\mathrm{s}}(\beta) := \mathrm{Tr}\, e^{-\beta(H_n^{\mathrm{s}}+\kappa n I)} = \quad \bar{v}_\beta \text{ — } X_\beta^n \text{ — } U_P v_\beta. \tag{S10}$$

For an n-site chain, site i carries $W_\beta(a_{i-1}, a_i)$, with internal bond histories $a_1, \ldots, a_{n-1}$ and boundary histories a_0, a_n contracted with $\bar{v}_\beta, v_\beta$. Taking the physical trace at each of the n sites produces X_β^n and gives $Y_n(\beta)$ in the second line; inserting U_P at the right cap gives the equal-field partition function $Y_n^{\mathrm{s}}(\beta)$ in the third.

Spin rotations act on bond histories by real orthogonal operators that commute with X_β, so $[U_P, X_\beta] = 0$. Regrouping the expansion into these site-by-site contractions is exact: before the energy shift, the absolute traced series is bounded by $3^n e^{\beta[3(n-1)+2h]} < \infty$, so Fubini's theorem justifies contracting the time-ordered expansion site by site.

Having expressed both open chains through X_β and v_β, we work first with the opposite-field partition function $Y_m(\beta)$ at even lengths m, where Eq. (S5) decomposes the boundary contraction into a sum of nonnegative terms. We will return to the odd equal-field chain $Y_n^{\mathrm{s}}(\beta)$ in Sec. SI.F, using an absolute remainder bound that does not require odd powers of the transfer operator to be positive.

SI.C. Finite inputs and ring estimates

We now identify the finite inputs required to drive the open-chain recurrence across the dyadic scales

$$\beta_j = 49\,2^{j-2}, \qquad n_j = 120\,2^{j-2}, \qquad M_j = 2n_j, \qquad j \geq 2, \tag{S11}$$

where n_j is a ring length and M_j is an even open-chain length. We retain the index $j = 2$ for the certified starting point of Ref. [19], at $\beta_2 = 49$. At each scale j, the two-length boundary comparison and the length-doubling step require three bulk properties of the transfer operator X_{β_j}: a simple positive leading eigenvalue λ_j invariant under U_P, a bound on the subleading norm $\|R_j\|^{n_j}$ to suppress the boundary remainder across the length difference $M_j - k_j$, and a lower bound on the bulk normalization ratio $r_j = \lambda_{j+1}/\lambda_j^2$ relating consecutive temperatures. All three bulk properties follow at every $j \geq 2$ from the periodic spatial and thermal purities $P_{\mathrm{s}}(n_2, \beta_2)$ and $P_\tau(M_2, \beta_2)$ at the initial scale $j = 2$ ($\beta_2 = 49$, $n_2 = 120$, $M_2 = 240$). To start the open-chain recurrence itself, we also need an initial bound on the boundary defect $e_{k_2}(2)$ at a shorter even length $k_2 = 142 < M_2$, together with a lower bound on the opposite-field thermal purity $P_\tau^{\mathrm{o}}(M_2, \beta_2) = Y_{M_2}(2\beta_2)/Y_{M_2}(\beta_2)^2$.

Ref. [19] certifies these four finite seed bounds at $j = 2$:

$$\begin{aligned} P_{\mathrm{s}}(120, 49) \geq .948, \qquad & P_\tau(240, 49) \geq .948, \\ e_{142}(2) \leq .14, \qquad & P_\tau^{\mathrm{o}}(240, 49) \geq .79. \end{aligned} \tag{S12}$$

Here $P_{\mathrm{s}}(n, \beta) = Z_{2n}(\beta)/Z_n(\beta)^2$ and $P_\tau(n, \beta) = Z_n(2\beta)/Z_n(\beta)^2$ concern the periodic ring, while $e_m(j)$, defined by Eq. (S6) at β_j, and $P_\tau^{\mathrm{o}}(m, \beta) = Y_m(2\beta)/Y_m(\beta)^2$ concern the opposite-field open chain.

We first use the two ring inputs in Eq. (S12) to control the bulk transfer operator at every scale $j \geq 2$. By the periodic doubling argument, the ring purities satisfy

$$\begin{aligned} P_{\mathrm{s}}(n_j, \beta_j), \ P_\tau(M_j, \beta_j) \geq 1 - \varepsilon_j, \\ \varepsilon_2 = .052, \qquad \varepsilon_{j+1} = 6\varepsilon_j^2. \end{aligned} \tag{S13}$$

To verify the factor of 6 in this scalar recurrence, recall the squaring inequality: if a positive trace-one operator of purity P is squared and renormalized, which concentrates its weights,  its new purity P' satisfies

$$1 - P' \leq f(1 - P), \qquad f(z) = \frac{z^2}{2(1-z)^2}. \tag{S14}$$

The periodic coupling identity first gives $P_{\mathrm{s}}(n, 2\beta) \geq (1-\varepsilon)^3$. For $0 \leq \varepsilon \leq .052$, we have

$$\begin{aligned} f(1 - (1-\varepsilon)^3) &= \frac{\varepsilon^2}{2}\left[\sum_{s=1}^{3}(1-\varepsilon)^{-s}\right]^2 \\ &\leq 6\varepsilon^2 \leq \varepsilon. \end{aligned} \tag{S15}$$

After spatial squaring bounds the new spatial defect by $6\varepsilon^2$, the second periodic coupling identity gives $P_\tau(4n, \beta) \geq (1-\varepsilon)^2(1-6\varepsilon^2) \geq (1-\varepsilon)^3$. Thermal squaring then bounds the new thermal defect by the same $6\varepsilon^2$, proving Eq. (S13) by induction on $j \geq 2$ without waiting for the smaller open-chain threshold established below.

From the ring purities in Eq. (S13), we now extract the bulk spectral properties of X_{β_j} needed for the open chain. Write λ_j for the leading eigenvalue of X_{β_j}, and decompose $T_j = X_{\beta_j}/\lambda_j = \Pi_j + R_j$ as in Eq. (S3). We claim that for every $j \geq 2$,

$$\begin{aligned} \lambda_j > 0 \text{ is simple}, \qquad & \|R_j\|^{n_j} \leq d(\varepsilon_j), \\ & d(\varepsilon) = \frac{\varepsilon}{2 - 3\varepsilon}, \qquad \|R_j\| < 1, \\ & r_j := \frac{\lambda_{j+1}}{\lambda_j^2}, \qquad 0 < r_j \leq 1, \\ & r_j^{M_j} \geq (1-\varepsilon_j)(1-\varepsilon_{j+1}), \qquad U_P \Pi_j = \Pi_j. \end{aligned} \tag{S16}$$

We verify each assertion in Eq. (S16) in turn. For any probability distribution of purity at least $1 - \varepsilon > 1/2$, let p_0 be its largest weight and put $z = 1 - p_0$. Since p_0 bounds the purity from above, $z < 1/2$, and $1 - \varepsilon \leq p_0^2 + (1-p_0)^2 = 1 - 2z(1-z)$ gives $2z(1-z) \leq \varepsilon$. Hence

$$z \leq \varepsilon, \qquad z \leq \frac{\varepsilon}{2(1-\varepsilon)}, \qquad \frac{z}{1-z} \leq d(\varepsilon).$$

Applying this estimate to the normalized weights $p_i = |\lambda_i(\beta_j)|^{n_j}/Z_{n_j}(\beta_j)$ yields the subleading tail bound

$$\sum_{i\geq 1}\left|\frac{\lambda_i(\beta_j)}{\lambda_j}\right|^{n_j} \leq d(\varepsilon_j). \tag{S17}$$

Because $d(\varepsilon_j) < 1$, the largest-modulus eigenvalue λ_j is simple and $\|R_j\|^{n_j} \leq d(\varepsilon_j) < 1$. Moreover, λ_j is positive: because X_{β_j} is Hilbert–Schmidt and the remaining spectrum has strictly smaller spectral radius, a negative isolated leading eigenvalue would make $\mathrm{Tr_v}\, X_{\beta_j}^N$ negative for all sufficiently large odd N, contradicting $Z_N(\beta_j) > 0$.

Next, to see that $U_P\Pi_j = \Pi_j$, note that the three π rotations about the coordinate axes and the cyclic permutation of those axes commute with the real transfer kernel X_{β_j}. Each acts by a sign on the real one-dimensional leading eigenspace. The order-three cyclic permutation must act by $+1$, and it conjugates the three π rotations into one another. Their signs are therefore equal; since the product of two π rotations is the third, the common sign is $+1$, which gives $U_P\Pi_j = \Pi_j$.

Finally, to bound the bulk normalization ratio $r_j = \lambda_{j+1}/\lambda_j^2$, observe that $P_\tau(N, \beta_j) \leq 1$ and taking even Nth roots as $N \to \infty$ give $\lambda_{j+1} \leq \lambda_j^2$, so $r_j \leq 1$. At length $M_j = n_{j+1}$, combining the leading spatial weight at β_{j+1} with the ring thermal purity at β_j gives the lower bound

$$\begin{aligned}\lambda_{j+1}^{M_j} &\geq (1-\varepsilon_{j+1})Z_{M_j}(\beta_{j+1}) \\ &\geq (1-\varepsilon_{j+1})(1-\varepsilon_j)Z_{M_j}(\beta_j)^2 \\ &\geq (1-\varepsilon_{j+1})(1-\varepsilon_j)\lambda_j^{2M_j}.\end{aligned} \tag{S18}$$

Dividing by $\lambda_j^{2M_j}$ proves $r_j^{M_j} \geq (1-\varepsilon_j)(1-\varepsilon_{j+1})$. With the bulk transfer estimates in Eq. (S16) established, we turn to the open-chain recurrence.

SI.D. Three steps for the open chain

We now set up the open-chain quantities at scale j and show how to advance them to scale $j+1$. Set $v_j = v_{\beta_j}$ and, for even $m \geq 2$, define the normalized opposite-field partition function, its remainder, and the boundary defect by

$$\begin{aligned} c_j &= |\langle \psi_j, v_j\rangle|^2, \\ B_m(j) &= \frac{Y_m(\beta_j)}{\lambda_j^m} = c_j + D_m(j) > 0, \\ D_m(j) &= \|R_j^{m/2} v_j\|^2, \qquad e_m(j) = \frac{D_m(j)}{B_m(j)}. \end{aligned} \tag{S19}$$

For even lengths $m \geq k$, absorbing $R_j^{k/2}$ into each boundary cap gives the spatial decay of the remainder,

$$\begin{aligned} D_m(j) &= \quad , \qquad y = R_j^{k/2} v_j, \\ 0 \leq D_m(j) &\leq \|R_j\|^{m-k} D_k(j). \end{aligned} \tag{S20}$$

The remainder therefore decays geometrically with the extra length $m-k$, , and in particular $D_m(j)$ is nonincreasing in even m, so $c_j \leq B_m(j) \leq B_k(j)$.

To connect quantities at β_j and $\beta_{j+1} = 2\beta_j$, we use the opposite-field thermal purity. By the definition of $B_m(j)$ and $r_j = \lambda_{j+1}/\lambda_j^2$, we have the exact thermal-purity bridge

$$\begin{aligned} P_\tau^{\mathrm{o}}(m, \beta_j) &= \frac{Y_m(\beta_{j+1})}{Y_m(\beta_j)^2} \in (0, 1], \\ B_m(j+1) &= P_\tau^{\mathrm{o}}(m, \beta_j) r_j^{-m} B_m(j)^2. \end{aligned} \tag{S21}$$


Multiplying the normalized open-chain Gibbs operators $\rho_m^{\mathrm{o}}(\beta) = e^{-\beta(H_m^{\mathrm{o}} + \kappa m I)}/Y_m(\beta)$ in imaginary time gives

$$= P_\tau^{\mathrm{o}}(m, \beta_j) \tag{S22}$$

Assuming bounds on a boundary defect $e_k(j)$ and on the thermal purity $P^{\circ}_{\tau}(m,\beta_j)$ at scale j, we advance to scale $j+1$ in three steps: first we bound the boundary defect $e_m(j+1)$ at the new temperature; second we use that new boundary defect to double the chain length from m to $2m$ in the thermal purity at β_j; third we square in imaginary time to obtain $P^{\circ}_{\tau}(2m,\beta_{j+1})$.

a. Step 1: control the new boundary remainder

Our first task is to bound the boundary defect $e_m(j+1)$ at the new inverse temperature β_{j+1} without knowing the new leading overlap c_{j+1} in advance. Fix even lengths $m > k \geq 2$, assume $e_k(j) < 1$, and write primes for quantities at scale $j+1$. Choose a suppression factor $q < 1$ satisfying $\|R_{j+1}\|^{m-k} \leq q$, so that Eq. (S20) at scale $j+1$ gives $D'_m \leq qD'_k$. Subtracting the normalized boundary contractions at lengths k and m at the same temperature β_{j+1} cancels their common leading term c':



(S23)

That is, $B'_k - B'_m = D'_k - D'_m$. Combining $D'_k = D'_m + B'_k - B'_m$ with $D'_m \leq qD'_k$ gives

$$\begin{aligned}(1-q)D'_m &\leq q(B'_k - B'_m),\\ e'_m &\leq \frac{q}{1-q}\left(\frac{B'_k}{B'_m} - 1\right).\end{aligned} \tag{S24}$$

The unknown overlap c' has dropped out of the difference $B'_k - B'_m$.

We can now bound the new-temperature ratio B'_k/B'_m in terms of scale-j quantities by applying the bridge identity (S21) at both lengths k and m:

$$\begin{aligned}\frac{B'_k}{B'_m} &= \frac{P^{\circ}_{\tau}(k,\beta_j)}{P^{\circ}_{\tau}(m,\beta_j)} r_j^{m-k}\left[\frac{B_k(j)}{B_m(j)}\right]^2\\ &\leq \frac{1}{P^{\circ}_{\tau}(m,\beta_j)[1-e_k(j)]^2}.\end{aligned} \tag{S25}$$

Here we used $P^{\circ}_{\tau}(k,\beta_j) \leq 1$, $r_j \leq 1$, and the lower bound

$$B_m(j) \geq c_j = [1-e_k(j)]B_k(j) > 0$$

at the old temperature. Equation (S25) makes no assumption on the new overlap c'; once Eqs. (S24) and (S25) yield $e'_m < 1$, it follows automatically that $c' = (1-e'_m)B'_m > 0$.

Notice why Eq. (S24) produces a quadratic improvement in the small-defect regime. It is a product of two small factors: the spatial decay factor $q/(1-q)$ over the length difference $m-k$, and the excess of the ratio B'_k/B'_m above one. Specifically, when $m = M_j$, $k = M_j/2$, and $\varepsilon_j \leq x \leq .01$, the ring bound (S16) at scale $j+1$ gives

$$\|R_{j+1}\|^{M_j} \leq d(\varepsilon_{j+1}) \leq d(6x^2).$$

Across the half-chain $m-k = M_j/2$, we may therefore take

$$q = \sqrt{d(\varepsilon_{j+1})} \leq \sqrt{\frac{6x^2}{2-18x^2}} \leq 2x. \tag{S26}$$

Separately, when $e_k(j) \leq x$ and $P^{\circ}_{\tau}(m,\beta_j) \geq 1-x$, Eq. (S25) bounds the excess above one by

$$\frac{B'_k}{B'_m} - 1 \leq (1-x)^{-3} - 1 = O(x). \tag{S27}$$

Multiplying Eqs. (S26) and (S27) bounds e'_m by $O(x^2)$. The subtraction of one in Eq. (S24), coming from the cancellation of c' in Eq. (S23), supplies this second factor of x.

b. *Step 2: double the length*

Having bounded the boundary defect $e_M(j+1)$ at the new temperature, we next double the open-chain length from M to $2M$ in the thermal purity at the old temperature β_j. For any even M, dividing the bridge identity (S21) at length $2M$ by that at length M gives

$$\begin{aligned}\frac{P_\tau^{\mathrm{o}}(2M,\beta_j)}{P_\tau^{\mathrm{o}}(M,\beta_j)} &= r_j^M \frac{B_{2M}(j+1)}{B_M(j+1)}\left[\frac{B_M(j)}{B_{2M}(j)}\right]^2 \\ &\geq r_j^M[1-e_M(j+1)]. \end{aligned} \tag{S28}$$

In the last factor, monotonicity at the old temperature gives $B_M(j) \geq B_{2M}(j)$, so $[B_M(j)/B_{2M}(j)]^2 \geq 1$. In the middle factor, however, $B_{2M}(j+1) \leq B_M(j+1)$, so we must use the new-temperature boundary defect bound from Step 1:

$$B_{2M}(j+1) \geq c_{j+1} = [1-e_M(j+1)]B_M(j+1).$$

Equation (S28) therefore retains both the bulk spectral-ratio factor r_j^M, controlled by Eq. (S16), and the new-temperature boundary factor $1-e_M(j+1)$.

c. *Step 3: square in imaginary time*

Once Step 2 bounds $P_\tau^{\mathrm{o}}(2M,\beta_j)$ at the doubled length $2M$, we double the inverse temperature from β_j to β_{j+1} by squaring the open-chain Gibbs state in imaginary time. Just as in the periodic argument, Eqs. (S22) and (S14) give

$$1-P_\tau^{\mathrm{o}}(2M,\beta_{j+1}) \leq f(1-P_\tau^{\mathrm{o}}(2M,\beta_j)). \tag{S29}$$

Notice why the three steps must be performed in this order: because $P_\tau^{\mathrm{o}}(M,\beta_j)$ already connects β_j and β_{j+1}, doubling the length in Step 2 requires the new-temperature boundary defect $e_M(j+1)$ from Step 1 before we can square in imaginary time in Step 3.

SI.E. Initialization and the small-defect induction

We now assemble the three steps of Sec. SI.D into a recurrence across $j \geq 2$. The initial boundary estimate in Eq. (S12) is certified at 142 sites. We use that length for the first comparison and the preceding target length thereafter:

$$k_2 = 142, \qquad k_{j+1} = M_j. \tag{S30}$$

Since $M_{j+1} = 2M_j$, this choice gives $k_j = M_j/2$ for all $j \geq 3$. Let E_j and η_j denote upper bounds on $e_{k_j}(j)$ and $1-P_\tau^{\mathrm{o}}(M_j,\beta_j)$, respectively. By Eqs. (S16), (S24), (S25), (S28), and (S29), the updates from scale j to scale $j+1$ take the form

$$\begin{aligned} F(q,e,t) &= \frac{q}{1-q}\left[\frac{1}{(1-e)^2 t}-1\right], \\ q_j &= d(\varepsilon_{j+1})^{(M_j-k_j)/M_j}, \\ E_{j+1} &\geq F(q_j, E_j, 1-\eta_j), \\ A_j &= (1-\eta_j)(1-\varepsilon_j)(1-\varepsilon_{j+1})(1-E_{j+1}), \\ \eta_{j+1} &\geq f(1-A_j), \end{aligned} \tag{S31}$$

valid whenever $E_j, \eta_j, E_{j+1}, \eta_{j+1} \in [0,1)$. The product A_j bounds $P_\tau^{\mathrm{o}}(2M_j,\beta_j)$ from below by Eq. (S28); thermal squaring then gives the update for η_{j+1}. Here the exponent $(M_j-k_j)/M_j$ in q_j comes from $n_{j+1} = M_j$ and the new-temperature ring remainder bound $\|R_{j+1}\|^{M_j} \leq d(\varepsilon_{j+1})$. Because $F(q,e,t)$ increases with q and e and decreases with t, any upper bound on q_j may be substituted into the update for E_{j+1}.

At the initial scale $j=2$, the seed defects $E_2 = .14$ and $\eta_2 = .21$ from Eq. (S12) are too large to enter the analytical small-defect induction ($x \leq .01$) directly. We therefore carry out three explicit rational steps from $j=2$ to $j=5$. Starting with $\varepsilon_2 = .052$, $E_2 = .14$, and $\eta_2 = .21$, Eq. (S31) produces the outward-rounded rational bounds in the following table:

j	E_j	η_j	upper bound for q_j
2	.14	.21	.142
3	.12	.155	.0282
4	.016	.026	.0028
5	.001	.001	—


All decimals in this table denote exact rationals. Both defects collapse along the three steps, E_j as  and η_j as  ($j = 2, \dots, 5$, each relative to its value at $j = 2$): the first step gains little because $q_2 \le .142$ is still large, whereas once q_j is small the defects square. At $j = 2$, the first suppression bound uses $k_2 = 142$ and $M_2 = 240$, so that

$$\frac{M_2 - k_2}{M_2} = \frac{49}{120}, \qquad d(\varepsilon_3)^{49} < .142^{120}.$$

At $j = 3$ and $j = 4$, we have $(M_j - k_j)/M_j = 1/2$, so we verify the upper bounds on q_j by squaring .0282 and .0028 and comparing with $d(\varepsilon_4)$ and $d(\varepsilon_5)$. Substituting these values into Eq. (S31) verifies each row of the table by exact rational arithmetic. Together with $\varepsilon_5 < .000015$, the $j = 5$ row brings all three defects below .01, establishing the base case of the small-defect induction:

$$\begin{aligned} &\varepsilon_j,\ e_{k_j}(j),\ 1 - P^{\mathrm{o}}_\tau(M_j, \beta_j) \le x_j, \qquad j \ge 5, \\ &\qquad x_5 = .01, \qquad x_{j+1} = 10x_j^2, \qquad x_j = 10^{-1-2^{j-5}}. \end{aligned} \tag{S32}$$

To prove the induction step in Eq. (S32), assume that the three bounds hold at scale $j \ge 5$ and write $x = x_j \le .01$. One step from scale j to scale $j + 1 = j'$ ($M_{j+1} = 2M_j$) proceeds along two coupled lanes:



For Step 1, Eqs. (S26) and (S27) give $q_j \le 2x$ and

$$F(2x, x, 1 - x) \le \frac{6x^2}{(1 - 2x)(1 - x)^3} < 7x^2, \tag{S33}$$

so $E_{j+1} = 10x^2$ bounds $e_{M_j}(j + 1)$. For Step 2, substituting $\eta_j \le x$, $\varepsilon_j \le x$, $\varepsilon_{j+1} \le 6x^2$, and $E_{j+1} \le 10x^2$ into A_j gives $A_j \ge 1 - 2x - 16x^2$. For Step 3, since $0 < x \le .01$, thermal squaring yields

$$f(2x + 16x^2) < 3x^2, \tag{S34}$$

so $\eta_{j+1} = 10x^2$ bounds $1 - P^{\mathrm{o}}_\tau(2M_j, \beta_{j+1})$, and $\varepsilon_{j+1} \le 6x^2 \le 10x^2$ closes the induction at scale $j + 1$.


SI.F. Equal fields, odd lengths, and the gap

We have established the small-defect bounds (S32) for even-length opposite-field chains at every scale $j \ge 5$. It remains to transfer these bounds to the target equal-field Hamiltonian H^{s}_n at arbitrary odd lengths $n \ge M_5 + 1 = 1921$ and extract the spectral gap.

Define the normalized equal-field partition function by


$$Q_j(n) = \frac{Y^{\mathrm{s}}_n(\beta_j)}{\lambda_j^n}.$$

Because U_P commutes with X_{β_j} and satisfies $U_P\Pi_j = \Pi_j$ by Eq. (S16), inserting U_P at the right cap preserves the leading contribution c_j:



(S35)

For any even $M \le n$, we split off $R_j^{M/2}$ on each side of the remainder contraction to obtain



$$\begin{aligned} Q_j(n) - c_j &= \\ &= \langle y, R_j^{n-M} U_P y \rangle, \qquad y = R_j^{M/2} v_j. \end{aligned} \tag{S36}$$

Because $\|R_j\| \le 1$ and U_P is orthogonal, we have $\|R_j^{n-M}U_P\| \le 1$, and Cauchy–Schwarz bounds the signed remainder in absolute value by the even-length opposite-field remainder $D_M(j)$:

$$|Q_j(n) - c_j| \le \|y\|^2 = D_M(j). \tag{S37}$$

This inequality controls $Q_j(n)$ at odd n without assuming that odd powers of R_j are positive.

Now fix a scale $j \ge 5$ and an odd length n in the interval $M_j \le n \le 2M_j$, and set $M = M_j = k_{j+1}$. Applying Eq. (S37) at β_j and β_{j+1} bounds $Q_j(n)$ from above and $Q_{j+1}(n)$ from below in terms of B_M:

$$\begin{aligned} Q_j(n) &\le c_j + D_M(j) = B_M(j), \\ Q_{j+1}(n) &\ge c_{j+1} - D_M(j+1) \\ &= [1 - 2e_M(j+1)] B_M(j+1) \\ &\ge (1 - 2x_{j+1}) B_M(j+1) > 0. \end{aligned} \tag{S38}$$

Combining Eq. (S38) with the bridge identity (S21) at length M, and using $r_j \le 1$ and $n - M \le M$ together with the bulk ratio bound $r_j^M \ge (1-\varepsilon_j)(1-\varepsilon_{j+1})$ from Eq. (S16), we obtain the equal-field thermal purity bound

$$\begin{aligned} P_\tau^{\mathrm{s}}(n, \beta_j) &= r_j^n \frac{Q_{j+1}(n)}{Q_j(n)^2} \\ &\ge (1 - 2x_{j+1}) P_\tau^{\mathrm{o}}(M, \beta_j) r_j^{n-M} \\ &\ge (1 - 2x_{j+1})(1 - x_j)(1 - \varepsilon_j)(1 - \varepsilon_{j+1}) \\ &\ge 1 - 2x_j - 26x_j^2 \\ &\ge 1 - 4x_j, \end{aligned} \tag{S39}$$

where the coefficient $26 = 20 + 6$ collects $2x_{j+1} = 20x_j^2$ and $\varepsilon_{j+1} \le 6x_j^2$.

Proof of Theorem S1. The intervals $[M_j, 2M_j]$ for $j \ge 5$ cover every integer $n \ge M_5 = 1920$. Given any odd $n \ge 1921$, choose $j \ge 5$ with $M_j \le n \le 2M_j$. Since $x_j \le x_5 = .01$, Eq. (S39) gives $w_0 \ge P_\tau^{\mathrm{s}}(n, \beta_j) \ge 1 - 4x_j > 1/2$, so H_n^{s} has a unique ground state. At this length n and finite inverse temperature β_j, Eq. (S2) gives

$$\begin{aligned} e^{-\beta_j \Delta_n} &\le \frac{4x_j}{1 - 4x_j} \\ &\le \frac{5}{12} 10^{-2^{j-5}} < 10^{-2^{j-5}}, \end{aligned} \tag{S40}$$

where we used $x_j = 10^{-1}10^{-2^{j-5}}$ and $1 - 4x_j \ge .96$, which yields the prefactor $.4/.96 = 5/12 < 1$. Taking logarithms and dividing by $\beta_j = 392\, 2^{j-5}$ proves $\Delta_n > (\log 10)/392$. □

Appendix SII: Finite Trotter networks and the periodic doubling argument

SII.A. Finite Trotter slices and a two-site transfer matrix

OpenAI's proof uses a continuous-time spatial integral kernel [7]. For our Lean formalization we instead prove the spatial inequalities for finite matrices and pass them to exact partition functions by a controlled product limit.

Consider the periodic spin-1 Heisenberg Hamiltonian

$$H_L = \sum_{j=0}^{L-1} h_{j,j+1}, \qquad h = \sum_{\alpha=x,y,z} S^\alpha \otimes S^\alpha, \qquad Z_L(\beta) = \operatorname{Tr} e^{-\beta H_L}, \tag{S1}$$

with indices taken modulo L, exchange coupling equal to one, and $\beta > 0$. In the Cartesian basis $s, s' \in \{x, y, z\}$, set


$$(S^\alpha)_{ss'} = -i\epsilon_{\alpha ss'}, \qquad G_\alpha = iS^\alpha, \qquad -h = \sum_\alpha G_\alpha \otimes G_\alpha. \tag{S2}$$

Here $\epsilon_{\alpha ss'}$ is the Levi-Civita symbol, Tr denotes the matrix trace, and $\mathbb{I}$ denotes the identity on the relevant space. Each G_α is real antisymmetric and has operator norm one. Equation (S2) is the real bond-reflection-positive factorization used below [10, 11].

To replace the continuous-time path integral by a finite-dimensional trace, we discretize imaginary time into M brickwork slices. For $L = 2N$, $N \geq 1$, and a positive integer M, let $\tau = \beta/M$. Define the linearized brickwork slice and its trace by

$$\begin{aligned} S_{2N}(\tau) &= \prod_{k=0}^{N-1} (\mathbb{I} - \tau h_{2k,2k+1}) \prod_{k=0}^{N-1} (\mathbb{I} - \tau h_{2k+1,2k+2}), \\ Z_{2N}^{(M)}(\beta) &= \operatorname{Tr} S_{2N}(\beta/M)^M. \end{aligned} \tag{S3}$$

Products are ordered from left to right in increasing index. The trace is real because all factors have real entries. Equation (S3) multiplies M slices. The same network read along space multiplies two-site columns,



(S4)


and we now construct the column $T_{M,\tau}$ by expanding each gate into bond factors.

Let $\Gamma = \{\varnothing, x, y, z\}$, with $C_\varnothing = \mathbb{I}$ and $C_\alpha = \sqrt{\tau}\, G_\alpha$. The bond expansion is the exact finite identity

$$\mathbb{I} - \tau h = \sum_{\gamma \in \Gamma} C_\gamma \otimes C_\gamma. \tag{S5}$$

An imaginary-time bond sequence is an element $p = (p_0, \ldots, p_{M-1})$ of $\mathcal{D}_M = \Gamma^M$. Define the single-site trace matrix by

$$\mathcal{F}_{pq} = \operatorname{Tr}\left[\prod_{t=0}^{M-1} (C_{p_t} C_{q_t})\right] = \qquad p, q \in \mathcal{D}_M. \tag{S6}$$



The diagram shows one site through $M = 3$ slices, with the odd-bond factors C_{q_t} on the left, the even-bond factors C_{p_t} on the right, and the physical trace closing the worldline. The product over t follows the multiplication order of the M slices. The matrix $\mathcal{F}$ depends on M and τ, but not on N.

To contract the network in space, assign sequences b_k and c_k to the even and odd bonds starting at $2k$ and $2k+1$, respectively. In slice t, site $2k$ carries $C_{b_{k,t}} C_{c_{k-1,t}}$, whereas site $2k+1$ carries $C_{b_{k,t}} C_{c_{k,t}}$. Expanding all bond sums, multiplying sitewise tensor products, and taking the physical trace gives

$$\begin{aligned} Z_{2N}^{(M)}(\beta) &= \sum_{\substack{b_0,\ldots,b_{N-1} \in \mathcal{D}_M \\ c_0,\ldots,c_{N-1} \in \mathcal{D}_M}} \prod_{k=0}^{N-1} \mathcal{F}_{b_k c_{k-1}} \mathcal{F}_{b_k c_k} \\ &= \sum_{c_0,\ldots,c_{N-1} \in \mathcal{D}_M} \prod_{k=0}^{N-1} (\mathcal{F}^{\mathsf{T}} \mathcal{F})_{c_{k-1} c_k} = \operatorname{Tr}(\mathcal{F}^{\mathsf{T}} \mathcal{F})^N, \end{aligned} \tag{S7}$$

where $c_{-1} = c_{N-1}$. The middle equality sums each even-bond sequence b_k independently; the last equality is the closed-path expansion of a matrix trace.

Thus the spatial transfer matrix

$$T_{M,\tau} = \mathcal{F}^{\mathsf{T}}\mathcal{F} = \mathcal{F}^{\dagger}\mathcal{F} \succeq 0 \tag{S8}$$

advances *two sites*: because $\mathcal{F}$ is real, its transpose equals its adjoint and $v^{\dagger}T_{M,\tau}v = \|\mathcal{F}v\|^2 \geq 0$ without requiring $\mathcal{F}$ itself to be symmetric or positive. Unlike a continuous-time one-site thermal-MPO kernel [27], $T_{M,\tau}$ is a finite Gram matrix at each M.

We extend the same contraction to any real reflection-positive cell model. Its assumptions are a finite-dimensional local cell space, a finite family of real matrices C_α, each symmetric or antisymmetric, a real symmetric on-site matrix o, and a real symmetric involution F satisfying

$$F^2 = \mathbb{I}, \qquad FoF = o, \qquad h = -\sum_{\alpha} C_\alpha \otimes FC_\alpha F. \tag{S9}$$

The transpose signs cancel between the two factors of each bond term, so $h^{\mathsf{T}} = h$. Together with $o^{\mathsf{T}} = o$, this makes $H_L = \sum_j h_{j,j+1} + \sum_j o_j$ self-adjoint. Lengths count cells. Insert the on-site layer $\prod_i(\mathbb{I} - \tau o_i)$ between the even and odd bond layers. Writing $A_\varnothing = \mathbb{I}$, $A_\alpha = \sqrt{\tau}\, C_\alpha$, and $O = \mathbb{I} - \tau o$, the bond expansion becomes $\mathbb{I} - \tau h = \sum_\gamma A_\gamma \otimes FA_\gamma F$. Replace Eq. (S6) by

$$\mathcal{F}_{pq} = \mathrm{Tr} \prod_{t=0}^{M-1} (A_{p_t} O F A_{q_t} F), \tag{S10}$$

where the sequences now range over the enlarged bond-label alphabet. An odd cell carries $FA_{p_t}FOA_{q_t}$. Since $FO = OF$, this factor is the conjugate by F of the even-cell factor. Conjugating the entire product leaves its trace unchanged. The two sublattices therefore contribute the same $\mathcal{F}_{pq}$, and Eqs. (S7) and (S8) follow without change.

SII.B. Finite-transfer moment inequalities

Positivity of $T_{M,\tau}$ expresses $Z_L^{(M)}(\beta)$ as an even moment sum of nonnegative weights, from which we derive the spatial purity, squaring, and interpolation inequalities at finite M. At fixed M, β, let $\lambda_i \geq 0$ be the eigenvalues of $T_{M,\beta/M}$, and put $\mu_i = \sqrt{\lambda_i}$. For every even $L \geq 2$, Eq. (S7) gives

$$Z_L^{(M)}(\beta) = \sum_i \mu_i^L. \tag{S11}$$

For any finite nonnegative family a_i, let $S_k = \sum_i a_i^k$. Then

$$S_2 \leq S_1^2, \qquad 2(S_2^2 - S_4) \leq (S_1^2 - S_2)^2. \tag{S12}$$

Indeed, $S_1^2 - S_2 = 2\sum_{i<j} a_i a_j$, whereas $S_2^2 - S_4 = 2\sum_{i<j} a_i^2 a_j^2$. Squaring the first sum and retaining only its diagonal terms proves the second inequality.

If $\sum_i a_i = 1$, the purity is $P = \sum_i a_i^2 > 0$. Squaring and normalizing replaces the distribution by a_i^2/P, of purity $P' = S_4/P^2$. Thus

$$1 - P' \leq f(1 - P), \qquad f(z) = \frac{z^2}{2(1-z)^2}. \tag{S13}$$

The map f is nondecreasing on $0 \leq z < 1$, since $z/(1-z)$ is nonnegative and increasing there. For $0 < z < 1/2$, $f(z) < z$,  , and squaring makes a small defect quadratically smaller.

For $0 < u \leq v$, nonnegative weights also satisfy

$$\sum_i \mu_i^v \leq \left(\sum_i \mu_i^u\right)^{v/u}. \tag{S14}$$

If the sum on the right is zero, the claim is immediate. Otherwise normalize μ_i^u to probabilities a_i; then $\sum_i a_i^{v/u} \leq \sum_i a_i = 1$.

Applying Eqs. (S12) and (S14) to Eq. (S11) yields, at fixed β, the closed scalar inequalities

$$\begin{aligned} Z_{2n}^{(M)} &\leq (Z_n^{(M)})^2, \\ 2((Z_{2n}^{(M)})^2 - Z_{4n}^{(M)}) &\leq ((Z_n^{(M)})^2 - Z_{2n}^{(M)})^2, \\ Z_v^{(M)} &\leq (Z_u^{(M)})^{v/u}, \end{aligned} \tag{S15}$$

for even $n \geq 2$ and even $v \geq u \geq 2$. We suppress the temperature argument only in this display.

SII.C. Controlled product limit

To pass the finite-M inequalities in Eq. (S15) to the exact partition functions $Z_L(\beta)$, we prove the matrix product limit $S_L(\beta/M)^M \to e^{-\beta H_L}$ at fixed L and β. Use the operator norm, which is submultiplicative and satisfies $\|\mathbb{I}\| = 1$. For $\|X\|, \|Y\| \leq K$, $K \geq 1$, telescoping powers gives

$$\left\|X^k - Y^k\right\| \leq kK^k \|X - Y\|. \tag{S16}$$

For $k \geq 1$, expand the difference as $\sum_{r=0}^{k-1} X^{k-1-r}(X-Y)Y^r$; each term is bounded by $K^{k-1}\|X-Y\| \leq K^k\|X-Y\|$. The case $k = 0$ is immediate.

Suppose $M(X_M - \mathbb{I}) \to A$, and set $Y_M = e^{A/M}$. Then $M(Y_M - \mathbb{I}) \to A$ and $Y_M^M = e^A$. With $C = \|A\| + 1$, convergence implies, for sufficiently large M,

$$\|X_M\|, \|Y_M\| \leq 1 + C/M.$$

Equation (S16) now gives the quantitative comparison

$$\begin{aligned} \left\|X_M^M - e^A\right\| &\leq (1 + C/M)^M \|M(X_M - Y_M)\| \\ &\leq e^C \|M(X_M - Y_M)\| \longrightarrow 0. \end{aligned} \tag{S17}$$

In particular, if $g(0) = \mathbb{I}$ and $g'(0) = A$, then $g(1/M)^M \to e^A$.

For fixed L, β, list the negated bond operators in the order of Eq. (S3), including the negated on-site operators when present, as $B_1, \ldots, B_k$. Their sum is $-H_L$. The polynomial

$$g(t) = \prod_{i=1}^{k} (\mathbb{I} + \beta t B_i)$$

has derivative $g'(0) = -\beta H_L$. Hence

$$S_L(\beta/M)^M \longrightarrow e^{-\beta H_L}, \qquad Z_L^{(M)}(\beta) \longrightarrow Z_L(\beta). \tag{S18}$$

More explicitly, Taylor expansion gives $g(1/M) - e^{-\beta H_L/M} = O(M^{-2})$. Equation (S17) therefore bounds the operator error by $e^C K_{L,\beta}/M$ for some finite $K_{L,\beta}$ and sufficiently large M, with $C = \|\beta H_L\| + 1$. Trace convergence follows from $|\mathrm{Tr}\, A| \leq 3^L \|A\|$ in the spin-1 model.

Taking $M \to \infty$ at each fixed (L, β) occurring in Eq. (S15) passes all three inequalities to the exact partition functions $Z_L(\beta)$. Because this limit precedes the doubling recurrence, the proof needs only convergence at individual pairs (L, β), and all subsequent thermal identities act on the exact Gibbs operators $e^{-\beta H_L}$.

SII.D. Torus identities and asymmetric doubling

With the exact partition-function inequalities established, we now set up the coupled doubling recurrence for the spatial and thermal purities. A torus of width n and height β,  , represents $Z_n(\beta)$; the spatial purity joins two copies side by side and the thermal purity stacks them in imaginary time:

$$P_{\mathrm{s}}(n,\beta) = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2} = \frac{\square}{\square^2}, \qquad P_\tau(L,\beta) = \frac{Z_L(2\beta)}{Z_L(\beta)^2} = \frac{\square}{\square^2}. \tag{S19}$$

All partition functions are strictly positive. The thermal purity is the purity of the Gibbs state $\rho_L(\beta) = e^{-\beta H_L}/Z_L(\beta)$. The limiting spatial inequalities and the Boltzmann power sums give $0 < P_{\mathrm{s}}, P_\tau \le 1$ at even lengths and

$$\begin{aligned} 1 - P_{\mathrm{s}}(2n,\beta) &\le f(1 - P_{\mathrm{s}}(n,\beta)), \\ 1 - P_\tau(L,2\beta) &\le f(1 - P_\tau(L,\beta)). \end{aligned} \tag{S20}$$

For the spatial bound, divide the second limiting inequality in Eq. (S15) by $2Z_{2n}(\beta)^2$. For the thermal bound, apply Eq. (S13) to the normalized Boltzmann weights.

Gluing four copies into the doubled torus, first in time and then in space or the other way round, gives the exact cancellation

$$P_{\mathrm{s}}(n,2\beta)P_\tau(n,\beta)^2 = \frac{Z_{2n}(2\beta)}{Z_n(\beta)^4} = \frac{\boxplus}{\square^4} = P_{\mathrm{s}}(n,\beta)^2 P_\tau(2n,\beta). \tag{S21}$$


Solving for one factor and repeating at length $2n$ yields

$$\begin{aligned} P_{\mathrm{s}}(n,2\beta) &= \frac{P_{\mathrm{s}}(n,\beta)^2 P_\tau(2n,\beta)}{P_\tau(n,\beta)^2}, \\ P_\tau(4n,\beta) &= \frac{P_\tau(2n,\beta)^2 P_{\mathrm{s}}(2n,2\beta)}{P_{\mathrm{s}}(2n,\beta)^2}. \end{aligned} \tag{S22}$$

Both identities follow directly by substituting Eq. (S19).

Let $n \ge 2$ be even, and let $0 \le p, q < 1$ bound the seed defects:

$$P_{\mathrm{s}}(n,\beta) \ge 1 - p, \qquad P_\tau(2n,\beta) \ge 1 - q. \tag{S23}$$

To double both n and β,  , we alternate the coupling identity with squaring in space and imaginary time, feeding the updated spatial defect into the thermal step. Write $z(a,b) = 1 - (1-a)^2(1-b)$. The asymmetric update is

$$p' = f(z(p,q)), \qquad q' = f(z(q,p')). \tag{S24}$$

Assuming $p' < 1$, its four steps are:

1. The first identity in Eq. (S22), with $P_\tau(n,\beta) \le 1$, gives $P_{\mathrm{s}}(n,2\beta) \ge (1-p)^2(1-q) = 1 - z(p,q)$.
2. Spatial squaring and monotonicity of f give $P_{\mathrm{s}}(2n,2\beta) \ge 1 - p'$.
3. The second identity in Eq. (S22), with $P_{\mathrm{s}}(2n,\beta) \le 1$, gives $P_\tau(4n,\beta) \ge (1-q)^2(1-p') = 1 - z(q,p')$.
4. Thermal squaring gives $P_\tau(4n,2\beta) \ge 1 - q'$.

Thus (p', q') bounds the defects at $(2n, 2\beta)$. Any upper bound below one substituted for p' in the third step remains valid; we use this rounding rule next.

SII.E. Six exact rational steps


The analytic tail estimate in Sec. SII.F requires both defects to be at most $1/100$. Because the initial spin-1 seed defects certified in Sec. SIII at $(n_0, \beta_0) = (36, 49/4)$,

$$P_{\mathrm{s}}(36,49/4) \ge 1 - \frac{47916}{10^6}, \qquad P_\tau(72,49/4) \ge 1 - \frac{181521}{10^6}, \tag{S25}$$

are larger, we first iterate the asymmetric update for six steps in exact rational arithmetic. Using the ceiling grid $D^{-1} = 10^{-7}$ ($D = 10^7$), we round the spatial bound upward before substituting it into the thermal update:

$$\begin{aligned} p_{j+1} &= \frac{\lceil D f(z(p_j,q_j)) \rceil}{D}, \\ q_{j+1} &= \frac{\lceil D f(z(q_j,p_{j+1})) \rceil}{D}. \end{aligned} \tag{S26}$$

Table S1 lists the resulting exact rational orbit, drawn here on a logarithmic scale:



While the thermal defect falls from the first step, the spatial defect initially rises to 0.0714 at $j = 3$ because the coupling step costs more than spatial squaring gains until q_j becomes small; both then collapse together and cross 10^{-2} at $j = 6$. The dotted line marks $\varepsilon_* \simeq 0.1273$, the fixed point of the symmetric one-step map $\varepsilon \mapsto f(1-(1-\varepsilon)^3)$. Because $q_0 = 0.1815$ starts above ε_*, a symmetric bound could not start the iteration; feeding the smaller updated spatial bound p_{j+1} into q_{j+1} pulls the orbit below 10^{-2}.

TABLE S1. Upward-rounded defect bounds at $n_j = 36\,2^j$, $\beta_j = (49/4)2^j$.

j	n_j	β_j	p_j	q_j
0	36	49/4	0.0479160	0.1815210
1	72	49/2	0.0605003	0.1733838
2	144	49	0.0686630	0.1632461
3	288	98	0.0713693	0.1447318
4	576	196	0.0633146	0.1055663
5	1152	392	0.0376144	0.0446518
6	2304	784	0.0084709	0.0055149

To check the table in integer arithmetic, write $a_j = Dp_j$, $b_j = Dq_j$, and set

$$V_j = (D-a_j)^2(D-b_j), \qquad W_j = (D-b_j)^2(D-a_{j+1}). \tag{S27}$$

For each $j = 0, \dots, 5$, the entries in Table S1 satisfy

$$\begin{aligned} 2(a_{j+1}-1)V_j^2 &< D(D^3-V_j)^2 \le 2a_{j+1}V_j^2, \\ 2(b_{j+1}-1)W_j^2 &< D(D^3-W_j)^2 \le 2b_{j+1}W_j^2. \end{aligned} \tag{S28}$$

Multiplying the ceiling comparisons in Eq. (S26) by their positive denominators gives Eq. (S28), since $f(z(p_j,q_j)) = (D^3-V_j)^2/(2V_j^2)$. The right-hand inequalities alone suffice to verify the upper bounds. Because all entries lie below one, the four-step update applies at every row and yields

$$\begin{aligned} P_{\mathrm{s}}(2304, 784) &\ge 1 - 0.0084709 > 0.99, \\ P_{\tau}(4608, 784) &\ge 1 - 0.0055149 > 0.99. \end{aligned} \tag{S29}$$

Both defects at $(2304, 784)$ now lie below the $1/100$ threshold.

SII.F. Scalar tail, interpolation, and the strict gap

Once both defects are at most $1/100$, a single scalar recurrence bounds all subsequent doubling steps. For $0 \le \varepsilon \le 1/100$,

$$f(1-(1-\varepsilon)^3) \le \frac{9\varepsilon^2}{2(1-3\varepsilon)^2} \le 5\varepsilon^2 \le \varepsilon. \tag{S30}$$

The first inequality uses $1-(1-\varepsilon)^3 \le 3\varepsilon$ and monotonicity of f; the second uses $9/[2(97/100)^2] < 5$. If both input defects are at most ε, the spatial update is therefore at most $5\varepsilon^2 \le \varepsilon$. Consequently, the thermal coupling input is also at most $1-(1-\varepsilon)^3$, and its squared defect is at most $5\varepsilon^2$.


Starting from Eq. (S29), reset the scale index and put

$$n_j = 2304\,2^j, \qquad \beta_j = 784\,2^j, \qquad \varepsilon_j = \frac{20^{-2^j}}{5}, \qquad j \geq 0. \tag{S31}$$

Here $\varepsilon_0 = 1/100$ and $\varepsilon_{j+1} = 5\varepsilon_j^2$, so $\log_{10}\varepsilon_j$ falls doubly exponentially, , from -2 at $j = 0$ to -21.5 at $j = 4$. Induction using Eq. (S30) proves

$$P_{\mathrm{s}}(n_j, \beta_j) \geq 1 - \varepsilon_j, \qquad P_\tau(2n_j, \beta_j) \geq 1 - \varepsilon_j. \tag{S32}$$

To cover every even length $L \geq 2304$ between the dyadic scales n_j, choose j with $n_j \leq L \leq 2n_j$. The known tori sit at the two ends of the doubling step; the one on the left bounds Z_L from above at β_j, and the one on the right bounds it from below at $2\beta_j$:



The limiting power-sum inequality in Eq. (S15), applied at β_j and $2\beta_j$, gives

$$\begin{aligned} Z_L(\beta_j)^2 &\leq Z_{n_j}(\beta_j)^{2L/n_j}, \\ Z_L(2\beta_j) &\geq Z_{2n_j}(2\beta_j)^{L/(2n_j)}. \end{aligned} \tag{S33}$$

Set $r = L/(2n_j) \in [1/2, 1]$ and

$$x = \frac{Z_{2n_j}(2\beta_j)}{Z_{n_j}(\beta_j)^4} = P_\tau(2n_j, \beta_j) P_{\mathrm{s}}(n_j, \beta_j)^2.$$

Since $(1 - \varepsilon_j)^3 \leq x \leq 1$, dividing the second inequality in Eq. (S33) by the first yields

$$P_\tau(L, \beta_j) \geq x^r \geq x \geq (1 - \varepsilon_j)^3 \geq 1 - 3\varepsilon_j > \frac{1}{2}. \tag{S34}$$

Finally, we convert this thermal purity bound at β_j into a lower bound on the spectral gap. Let $E_0(L) \leq E_1(L) \leq \cdots$ include multiplicities, and let $w_m = e^{-\beta_j E_m(L)}/Z_L(\beta_j)$. The largest Gibbs weight satisfies $P_\tau(L, \beta_j) = \sum_m w_m^2 \leq w_0$. Equation (S34) implies $w_0 > 1/2$, which proves that the ground state is unique. Moreover, $w_1 \leq 1 - w_0$. With $\Delta_L = E_1(L) - E_0(L)$, we obtain

$$\begin{aligned} e^{-\beta_j \Delta_L} = \frac{w_1}{w_0} &\leq \frac{3\varepsilon_j}{1 - 3\varepsilon_j} \\ &\leq \frac{60}{97}\, 20^{-2^j} < 20^{-2^j}, \end{aligned} \tag{S35}$$

where we used $1 - 3\varepsilon_j \geq 97/100$. Taking logarithms and dividing by $\beta_j = 784\,2^j$ at this finite, length-dependent temperature gives

$$\boxed{\Delta_L > \frac{\log 20}{784} \quad \text{for every even } L \geq 2304}. \tag{S36}$$

The strict inequality comes from the finite prefactor $60/97 < 1$, without taking a fixed-length limit $\beta \to \infty$. On the jth dyadic interval, Eq. (S35) gives the larger bound $\log[(1 - 3\varepsilon_j)/(3\varepsilon_j)]/\beta_j$, which decreases toward $(\log 20)/784$, (from 4.43×10^{-3} at $j = 0$ to 3.86×10^{-3} at $j = 4$); the uniform bound is their infimum. Given the two seed bounds in Eq. (S25), the finite-matrix transfer inequalities, rational orbit, scalar tail, and dyadic interpolation establish Eq. (S36).

Appendix SIII: Finite seed certificates and improved bounds

In this section we explain the finite inputs and scalar estimates from Ref. [7] that certify the starting pair

$$1 - P_{\mathrm{s}}(36, 49/4) \le \frac{47916}{10^6},$$
$$1 - P_{\tau}(72, 49/4) \le \frac{181521}{10^6}. \tag{S1}$$

These two bounds form Lemma 1 in the main text. We then present the temperature-lifting calculation, which combines the same short-ring trace bounds with the 120-site trial state of Ref. [7] to prove a unique ground state and

$$\Delta_L > 0.072901771 \quad \text{for every even } L \ge 36. \tag{S2}$$

We have reproduced the finite trace and trial contractions of Ref. [7] in exact arithmetic; the stronger gap bound uses only additional scalar inequalities on those inputs. Every terminating decimal that defines a bound below denotes an exact rational number, and we mark numerical approximations explicitly.

SIII.A. The base seed at length 36

To prove Eq. (S1) at $b_0 = 49/4$, we need lower bounds on the numerators $Z_{72}(b_0)$ and $Z_{72}(2b_0)$ and upper bounds on the denominators $Z_{36}(b_0)^2$ and $Z_{72}(b_0)^2$. A 72-site trial state bounds $Z_{72}(b)$ from below for every $b > 0$. Traces on $4 \le n \le 12$ sites constrain the eigenvalues of the one-site spatial transfer operator at b_0, which lets us bound the subleading contributions to $Z_{36}(b_0)$ and $Z_{72}(b_0)$.

a. Hamiltonians, twists, and the spatial transfer operator

We consider the periodic spin-1 Heisenberg Hamiltonian

$$H_n = \sum_{j=1}^{n} \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}, \qquad \boldsymbol{S}_j^2 = 2I, \qquad \boldsymbol{S}_{n+1} = \boldsymbol{S}_1, \tag{S3}$$

with unit exchange coupling. Lengths count sites, and $\Delta_L = E_1(L) - E_0(L)$, with eigenvalues ordered with multiplicity. Anticipating the trial energy per site, we shift the Hamiltonian by κ per site:

$$\kappa = \frac{700741}{500000}, \qquad \widehat{H}_n = H_n + \kappa n I, \qquad Z_n(\beta) = \operatorname{Tr} e^{-\beta \widehat{H}_n}. \tag{S4}$$

We omit hats on shifted quantities in this section: $Z_n(\beta)$ denotes $\widehat{Z}_n(\beta)$ in Eq. (15), and K_{b_0} below denotes the shifted operator $\widehat{K}$. This extensive shift cancels in both purities:

$$P_{\mathrm{s}}(n, \beta) = \frac{Z_{2n}(\beta)}{Z_n(\beta)^2}, \qquad P_{\tau}(n, \beta) = \frac{Z_n(2\beta)}{Z_n(\beta)^2}. \tag{S5}$$

To separate the spatial transfer spectrum by spin-rotation symmetry, we also use rings whose closing bond is twisted by a spin rotation, 

, alongside the plain ring  . For an on-site spin rotation u_g representing g, we define

$$H_n(g) = \sum_{j=1}^{n-1} h_{j,j+1} + h_{n,1}(g),$$
$$h_{n,1}(g) = \sum_{\alpha=x,y,z} S_n^{\alpha} (u_g S^{\alpha} u_g^{\dagger})_1,$$
$$Z_n(\beta, g) = \operatorname{Tr} e^{-\beta(H_n(g) + \kappa n I)}, \tag{S6}$$

where $h_{j,j+1} = \boldsymbol{S}_j \cdot \boldsymbol{S}_{j+1}$ and u_g acts on one side of the closing bond $(n, 1)$.

TABLE S2. Exact rational centers of the shifted traces at $b_0 = 49/4$ from Ref. [7]. Each entry has certified absolute error at most 3×10^{-5}.


n	$Z_n(b_0)$ center	$Z_n(b_0, P)$ center
4	124.885379	0.065757
5	0.002925	2.960742
6	12.870614	0.286599
7	0.054069	1.920414
8	4.639283	0.510140
9	0.195455	1.514846
10	2.654823	0.675419
11	0.377629	1.314544
12	1.900547	0.787405

The continuous-time construction of Ref. [7] provides a self-adjoint one-site spatial transfer operator K_β for the shifted Hamiltonian and a virtual symmetry operator U_g such that

$$Z_n(\beta, g) = \mathrm{Tr}(K_\beta^n U_g), \qquad n \geq 4. \tag{S7}$$

At $g = 1$, $Z_n(\beta) = \sum_i \lambda_i(\beta)^n$ is a power sum of the real eigenvalues $\lambda_i(\beta)$ of K_β (which can have either sign). Because the same eigenvalues govern every length $n \geq 4$, trace bounds on $4 \leq n \leq 12$ constrain the power sums at $n = 36$ and $n = 72$.

In our diagrams, horizontal wires carry spatial virtual indices, vertical double wires carry physical indices, and closed loops denote traces. Commutation of K_β with U_g gives



(S8)

where the ring contains n transfer vertices and U_g implements the twisted closing bond of Eq. (S6).

b. *Exact finite trace and trial inputs*

We now state the two numerical inputs from Ref. [7] used at $b_0 = 49/4$. The first input is the table of shifted traces $Z_n(b_0)$ and $Z_n(b_0, P)$ for $4 \leq n \leq 12$, where P is a π rotation about a Cartesian axis. Table S2 lists the rational centers, each certified within the absolute error

$$\epsilon_Z = 3 \times 10^{-5}. \tag{S9}$$

These enclosures come from integer Horner recursions for the half-Gibbs operator rather than dense diagonalization. For an unshifted Hamiltonian $H = H_n(g)$, we set

$$R = I - \frac{(3n/2)I + H}{16}, \tag{S10}$$

which satisfies $\|R\| \leq 1$ on each ring in Table S2. To check this, remove the local twist on each three-site path by a unitary rotation. Coupling the outer spins first shows that the two-bond cluster has spectrum in $[-3, 2]$. Averaging over the cyclic clusters gives $-(3n/2)I \leq H \leq nI$. For $4 \leq n \leq 12$, we therefore have $0 \leq (3n/2)I + H \leq (5n/2)I \leq 32I$, placing the spectrum of R in $[-1, 1]$. At $b_0 = 49/4$, the shifted half-Gibbs operator is

$$e^{-b_0(H+\kappa n I)/2} = e^{-b_0(\kappa - 3/2)n/2} e^{98(R-I)}. \tag{S11}$$

We approximate $e^{98(R-I)}$ by the normalized Poisson polynomial

$$\sum_{j=0}^{J} p_j R^j, \qquad p_j = \frac{98^j/j!}{\sum_{\ell=0}^{J} 98^\ell/\ell!}, \qquad J = 280, \tag{S12}$$

whose truncation and normalization error in operator norm is at most

$$2e^{-98}\sum_{j>280}\frac{98^j}{j!}<2^{-84}. \tag{S13}$$

Setting $Q=2^{55}$, we replace Qp_j by integers c_j with $|c_j-Qp_j|\le 1$. Writing $R=D/s$ with an integer matrix D and integer divisor $s>0$, we evaluate each basis column e_r by the downward recursion



$$X_{J+1}=0,\qquad X_j=\left\lfloor\frac{DX_{j+1}}{s}\right\rfloor+c_je_r, \tag{S14}$$

for $j=J,\dots,0$, where the floor acts entrywise, including on negative entries. In dimension $d=3^n$, each step introduces a Euclidean rounding error of at most $\sqrt{d}+1$. Because $\|R\|\le 1$, these errors do not amplify, so the total Euclidean error in X_0 is at most $(J+1)(\sqrt{d}+1)$ before adding the Poisson tail (S13). The trace $Z_n(b_0,g)$ equals the squared Frobenius norm of the half-Gibbs operator, which we obtain by summing the squared column norms with their exact symmetry multiplicities and multiplying by a rational enclosure of the scalar prefactor in Eq. (S11).

The second input is a periodic matrix-product trial state defined by real integer matrices A_s of bond dimension 26, used at $N=72$ for the base seed and at $N=120$ for temperature lifting. Setting

$$\begin{aligned}\psi_N(s_1,\dots,s_N)&=\operatorname{Tr}(A_{s_1}\cdots A_{s_N}),\\ T&=\sum_s A_s\otimes A_s,\\ O&=\sum_{s,t,s',t'}h_{st,s't'}A_sA_t\otimes A_{s'}A_{t'},\end{aligned} \tag{S15}$$

where T is the $26^2\times 26^2$ trial transfer matrix, we express the norm and energy expectation as exact finite-matrix traces:

$$\begin{aligned}\|\psi_N\|^2&=\quad\text{  }\quad=\operatorname{Tr}T^N,\\ \langle\psi_N,H_N\psi_N\rangle&=N\quad\text{  }\quad=N\operatorname{Tr}(T^{N-2}O).\end{aligned} \tag{S16}$$

Each layer has N tensors, and the two-site operator h acts on one bond. Exact integer arithmetic verifies that for $N\in\{72,120\}$,

$$\begin{aligned}&\operatorname{Tr}T^N>0,\\ &500000\operatorname{Tr}(T^{N-2}O)+700741\operatorname{Tr}T^N<0.\end{aligned} \tag{S17}$$


By the variational principle, the ground energy satisfies $E_0(N)<-\kappa N$, so the shifted ground energy $E_0(N)+\kappa N$ is negative. Retaining only the ground-state Boltzmann weight gives the lower bounds

$$Z_{72}(\beta)>1,\qquad Z_{120}(\beta)>1\quad(\beta>0). \tag{S18}$$

In the base seed, $Z_{72}(2b_0)>1$ bounds the thermal numerator directly, whereas $Z_{72}(b_0)>1$ forces a large leading spatial eigenvalue.

c. *Sector moments and polynomial filters*

We next split the spectrum of K_{b_0} into spin-rotation sectors and bound its individual eigenvalues. The four virtual π rotations $D_2=\{1,P_x,P_y,P_z\}$ commute with K_{b_0} and define the character projectors

$$\Pi_\chi=\frac{1}{4}\sum_{g\in D_2}\chi(g)U_g,\qquad \operatorname{Tr}(K_{b_0}^n\Pi_\chi)=\quad\text{  }. \tag{S19}$$

Cyclic permutation of the coordinate axes conjugates the three nontrivial characters into one another, so their three sectors have identical eigenvalue lists, which we denote by N; we denote the D_2-invariant eigenvalue list by J. Writing $J_n = \sum_{\lambda\in J}\lambda^n$ and $N_n = \sum_{\lambda\in N}\lambda^n$ at b_0, we have

$$\begin{aligned} Z_n &= J_n + 3N_n, & Z_n(P) &= J_n - N_n, \\ J_n &= \frac{Z_n + 3Z_n(P)}{4}, & N_n &= \frac{Z_n - Z_n(P)}{4}. \end{aligned} \tag{S20}$$

From Eq. (S9), the sector moments J_n and N_n for $4 \le n \le 12$ have absolute errors at most ϵ_Z and $\epsilon_Z/2$ around their rational centers. At $n = 12$, we add the common tolerance ϵ_Z to both centers to obtain the twelfth-moment upper bounds

$$J_{12} \le m_J = \frac{2131441}{2000000}, \qquad N_{12} \le m_N = \frac{556631}{2000000}. \tag{S21}$$

To bound individual eigenvalues from these finite power sums, we evaluate the nonnegative polynomials of Ref. [7],

$$\begin{aligned} F_J(x) &= x^4(7 + 17x - 280x^2 - 185x^3 + 1000x^4)^2, \\ F_N(x) &= x^4(12 - 394x^2 + 1000x^4)^2. \end{aligned} \tag{S22}$$

Pairing each coefficient of F_J and F_N with the upper or lower endpoint of the corresponding moment interval for $4 \le n \le 12$ gives

$$\begin{aligned} \sum_{\lambda\in J} F_J(\lambda) &\le B_J = 318604.54848175, \\ \sum_{\lambda\in N} F_N(\lambda) &\le B_N = 48169.880699. \end{aligned} \tag{S23}$$

Because F_J and F_N are nonnegative on $\mathbb{R}$, each single eigenvalue satisfies $F_J(\lambda) \le B_J$ for $\lambda \in J$ and $F_N(\lambda) \le B_N$ for $\lambda \in N$. Since F_N is small on most of $[-1, 1]$ and exceeds B_N for $|x| \ge 0.872$, no N eigenvalue can lie on the shaded exterior rays,  .

For $U = 1.00139$ and $V = 0.872$, we expand $F_J(\pm(U + y))$ and $F_N(\pm(V + y))$ in powers of $y \ge 0$. Exact rational arithmetic shows that their constant terms exceed B_J and B_N and all higher coefficients are nonnegative, ruling out eigenvalues on both exterior rays $|x| \ge U$ and $|x| \ge V$:

$$|\lambda| < U \quad (\lambda \in J), \qquad |\lambda| < V \quad (\lambda \in N). \tag{S24}$$

d. *Capped masses and the two base defects*

To pass from the twelfth moments (S21) and the eigenvalue caps (S24) to $n = 36$ and $n = 72$, we bound higher power sums under a mass constraint and an entry cap. For $m \ge 0$, $a > 0$, and $r \ge 1$, set

$$\mathcal{C}(m, a, r) = qa^r + (m - qa)^r, \qquad q = \lfloor m/a \rfloor. \tag{S25}$$

Whenever $0 \le t_i \le a$ and $\sum_i t_i \le m$, convexity of t^r implies $\sum_i t_i^r \le \mathcal{C}(m, a, r)$: shifting mass from a smaller unsaturated entry to a larger one increases $\sum_i t_i^r$, so the maximum is achieved by filling q entries to the cap a and putting the remaining mass $m - qa$ in a single entry,  . Applied to $t_i = \lambda_i^{12} \ge 0$, this bounds higher moments of the 12th-power weights.

We first show that J contains a dominant eigenvalue with $|\lambda_0| > \ell = 0.999$. If every $\lambda \in J$ satisfied $|\lambda| \le \ell$, then applying Eq. (S25) with $r = 6$ to J, whose extremal configuration  has one entry at the cap $\ell^{12} \simeq 0.988$ and 0.078 left over, and to each of the three N sectors, with cap $V^{12} \simeq 0.193$, would give

$$Z_{72}(b_0) \le \mathcal{C}(m_J, \ell^{12}, 6) + 3\mathcal{C}(m_N, V^{12}, 6) < 0.931 < 1, \tag{S26}$$

contradicting the 72-site trial bound $Z_{72}(b_0) > 1$ in Eq. (S18). Because $m_N < \ell^{12}$, no N eigenvalue can exceed ℓ in modulus, so J has at least one eigenvalue λ_0 with $|\lambda_0| > \ell$: the twelfth-power weights look like  , one tall bar in J next to a small J remainder and three identical N lists.

Removing $\lambda_0^{12} > \ell^{12}$ from J_{12} leaves residual twelfth moment at most $m_J - \ell^{12} < U^{12}$ in J. For even $k \geq 12$, the subleading moment $\sum_{i\neq 0} \lambda_i^k$ is therefore bounded by

$$R_k = (m_J - \ell^{12})^{k/12} + 3\mathcal{C}(m_N, V^{12}, k/12). \tag{S27}$$

At $k = 36$ and $k = 72$, we obtain the two base purity bounds:

$$\begin{aligned} P_{\mathrm{s}}(36, b_0) &\geq \left(1 + \frac{R_{36}}{\ell^{36}}\right)^{-2} = 1 - p, \\ P_\tau(72, b_0) &\geq \left(U^{72} + R_{72}\right)^{-2} = 1 - q. \end{aligned} \tag{S28}$$

For $P_{\mathrm{s}}(36, b_0)$, we bound the numerator from below by $Z_{72}(b_0) \geq \lambda_0^{72}$, bound $Z_{36}(b_0) \leq |\lambda_0|^{36} + R_{36}$, and use $|\lambda_0| > \ell$. For $P_\tau(72, b_0)$, we bound the numerator from below by $Z_{72}(2b_0) > 1$ from Eq. (S18) and bound the denominator by $Z_{72}(b_0) = \lambda_0^{72} + \sum_{i\neq 0} \lambda_i^{72} \leq U^{72} + R_{72}$.

Because $k/12 \in \{3, 6\}$ is an integer, Eqs. (S26)–(S28) involve only rational arithmetic. Exact evaluation verifies

$$p \leq \frac{47916}{10^6}, \qquad q \leq \frac{181521}{10^6}, \tag{S29}$$

which completes the proof of Eq. (S1). Numerically, $p \simeq 0.0479157710062$ and $q \simeq 0.181520168033$.

SIII.B. Temperature lifting and the improved uniform bound

Because the gap bound from the doubling argument scales inversely with the seed inverse temperature, certifying smaller defects at a moderate inverse temperature and stopping after a finite prefix of doubling steps gives a much larger gap constant. We now explain the temperature-lifting method, which uses the same trace table at $b_0 = 49/4$ and the 120-site trial bound $Z_{120}(\beta) > 1$ to certify a seed at $(n_0, b) = (60, rb_0)$ with $r \geq 1$. Combining that lifted seed with a tighter unlifted bound at $(36, b_0)$ covers all even lengths $L \geq 36$.

a. Lifted moments and a dominant spatial weight

We keep m_J, m_N from Eq. (S21) and sharpen the filter caps (S24) to the exact rationals

$$U = 1.001350644012, \qquad V = 0.871255704653, \tag{S30}$$

verified by the same exterior-ray expansions of F_J and F_N. Since $U^{12} < m_J < 2U^{12}$ and $V^{12} < m_N < 2V^{12}$, Eq. (S25) bounds the full sector moments at any even length $k \geq 12$ by

$$\begin{aligned} \mathcal{J}_k &= U^k + (m_J - U^{12})^{k/12}, \\ \mathcal{N}_k &= V^k + (m_N - V^{12})^{k/12}, \\ M_k &= \mathcal{J}_k + 3\mathcal{N}_k. \end{aligned} \tag{S31}$$

Because $N_k \geq 0$ at even k, we have $Z_k(b_0) = J_k + 3N_k \leq M_k$ and $Z_k(b_0, P) = J_k - N_k \leq \mathcal{J}_k$.

Now fix a rational factor $r \geq 1$, an even length $12 \leq k \leq 60$, and the lifted inverse temperature $b = rb_0$. Both H_k and $H_k(P)$ are self-adjoint, so their physical Boltzmann weights $z_i > 0$ at b_0 satisfy $\sum_i z_i^r \leq (\sum_i z_i)^r$. This transfers the upper bounds at b_0 to b:

$$Z_k(b) \leq D := M_k^r, \qquad Z_k(b, P) \leq E := \mathcal{J}_k^r. \tag{S32}$$

In rational arithmetic we replace D and E by outward rational upper bounds satisfying $E \leq D$.

At the lifted inverse temperature b, let K_b be the one-site spatial transfer operator and define the kth-power weights

$$w_i = |\lambda_i(b)|^k, \qquad h = \frac{60}{k}, \qquad \tau = \frac{D + 3E}{4}. \tag{S33}$$

Since k is even, $w_i = \lambda_i(b)^k \geq 0$. Thus $\sum_i w_i = Z_k(b) \leq D$, the J-sector sum satisfies $\sum_{\lambda_i \in J} w_i = [Z_k(b) + 3Z_k(b,P)]/4 \leq \tau$, and the 120-site trial bound (S18) gives

$$\sum_i w_i^{2h} = Z_{120}(b) > 1. \tag{S34}$$

If a rational threshold $t > 0$ satisfies

$$Dt^{2h-1} < 1, \qquad 3t > D, \tag{S35}$$

then at least one weight must satisfy $w_0 > t$; otherwise $\sum_i w_i^{2h} \leq t^{2h-1} \sum_i w_i \leq Dt^{2h-1} < 1$, contradicting Eq. (S34). Moreover, $3t > D$ forces w_0 to lie in J, because any weight in N is triply degenerate and would contribute more than $3t > D$ to $\sum_i w_i$.

Removing w_0 leaves residual J mass $x \in [0, \tau - t]$, while each of the three identical N sectors has mass at most $(D - t - x)/3$. For any $s \geq 1$, the residual moment $\sum_{i \neq 0} w_i^s$ is at most $x^s + 3[(D - t - x)/3]^s$. By convexity on $x \in [0, \tau - t]$, this sum attains its maximum at one of the two endpoints:

$$\mathcal{R}_s(t) = \max\left\{3\left(\frac{D-t}{3}\right)^s, (\tau - t)^s + 3\left(\frac{D-\tau}{3}\right)^s\right\}. \tag{S36}$$

All choices below satisfy $0 < t \leq \tau \leq D$.

Once $w_0 > t$ is established, any larger rational t' with

$$(t')^{2h} + \mathcal{R}_{2h}(t) < 1 \tag{S37}$$

implies $w_0 > t'$, since $w_0 \leq t'$ would give $Z_{120}(b) = w_0^{2h} + \sum_{i \neq 0} w_i^{2h} \leq (t')^{2h} + \mathcal{R}_{2h}(t) < 1$. Iterating Eq. (S37) tightens the lower bound t on w_0. Because $w_0^h = |\lambda_0(b)|^{60}$ and $\sum_{i \neq 0} |\lambda_i(b)|^{60} \leq \mathcal{R}_h(t)$, we obtain the lifted spatial purity bound at $n = 60$:

$$P_{\mathrm{s}}(60, b) \geq \left(1 + \frac{\mathcal{R}_h(t)}{t^h}\right)^{-2}. \tag{S38}$$

b. *The physical thermal residual*

Scaling the inverse temperature from b_0 to $b = rb_0$ changes the spatial transfer operator K_b, whereas it simply raises each physical Boltzmann weight $z_i = e^{-b_0 \widehat{E}_i(120)}$ of $\widehat{H}_{120}$ to the rth power. To bound $P_\tau(120, b)$, we therefore work directly with the ordered Boltzmann weights $z_0 \geq z_1 \geq \cdots$ at b_0. We have $z_0 > 1$ from the 120-site trial state and $\sum_i z_i = Z_{120}(b_0) \leq M_{120}$ from Eq. (S31). Hence

$$\sum_{i>0} \frac{z_i}{z_0} \leq M_{120} - 1, \qquad \sum_{i>0} \left(\frac{z_i}{z_0}\right)^r \leq (M_{120} - 1)^r. \tag{S39}$$

At $b = rb_0$, the partition functions are $Z_{120}(b) = \sum_i z_i^r$ and $Z_{120}(2b) = \sum_i z_i^{2r} \geq z_0^{2r}$: raising the weights to the power $r > 1$ concentrates them on the ground state,  at $r = 17/8$. Dividing numerator and denominator by z_0^{2r} gives the lifted thermal purity bound:

$$P_\tau(120, rb_0) \geq [1 + (M_{120} - 1)^r]^{-2}. \tag{S40}$$

c. *Finite-prefix coverage and the doubling tail*

We now convert the seed bounds at (n_0, b_0) into a uniform gap bound by combining a finite prefix of rational doubling steps with an analytic tail estimate.

Proposition S1 (Finite-prefix coverage)**.** *Let n_0 be even and $b_0 > 0$. Suppose $0 \leq p_j, q_j < 1$ bound respectively $1 - P_{\mathrm{s}}(n_j, b_j)$ and $1 - P_\tau(2n_j, b_j)$ at $(n_j, b_j) = (2^j n_0, 2^j b_0)$ for $0 \leq j \leq K$. Put $A_j = (1 - p_j)^2(1 - q_j)$ and assume $A_j > 1/2$ for $j < K$. Let $u = \max(p_K, q_K)$ and choose C such that*

$$0 < u < 1/6, \qquad Cu < 1, \qquad C(1 - 3u) > 3,$$
$$G(u) := \frac{1}{2}\left((1-u)^{-1} + (1-u)^{-2} + (1-u)^{-3}\right)^2 \leq C. \tag{S41}$$

TABLE S3. Temperature-lifting certificates at $(60, rb_0)$ with $b_0 = 49/4$. Defect bounds at $j = 0$ are rounded upward; gap bounds hold for every even $L \geq 60$, and for every even $L \geq 36$ once combined with the unlifted bridge on $36 \leq L \leq 72$.

k	r	Spatial defect	Thermal defect	$\Delta_L >$
30	2	0.0307608679	0.0590740709	0.070450002
28	17/8	0.0305302052	0.0479556776	0.072901771

If g is strictly smaller than every member of

$$\left\{ \frac{\log(A_j/(1-A_j))}{b_j} : 0 \leq j < K \right\} \cup \left\{ \frac{\log(1/(Cu))}{b_K} \right\}, \tag{S42}$$

then every even $L \geq n_0$ has a unique ground state and $\Delta_L > g$. An entry with $A_j = 1$ is interpreted as $+\infty$.

Proof. For even $n_j \leq L \leq 2n_j$ with $0 \leq j < K$, spatial power-sum interpolation (S34) gives $P_\tau(L, b_j) \geq A_j^{L/(2n_j)} \geq A_j > 1/2$. The largest normalized Boltzmann weight a_0 satisfies $a_0 \geq P_\tau(L, b_j) > 1/2$, so the ground state is unique, and the next weight $a_1 \leq 1 - a_0$ obeys

$$e^{-b_j \Delta_L} = \frac{a_1}{a_0} \leq \frac{1-a_0}{a_0} \leq \frac{1-A_j}{A_j}, \tag{S43}$$

which yields $\Delta_L \geq \log(A_j/(1-A_j))/b_j > g$. For $j \geq K$, since $f(1-(1-x)^3) = x^2 G(x)$ and G is increasing on $[0, u]$, Eq. (S41) implies that $u_m = C^{-1}(Cu)^{2^{m-K}}$ bounds both defects at (n_m, b_m) by induction. Interpolating on $n_m \leq L \leq 2n_m$ gives $P_\tau(L, b_m) \geq 1 - 3u_m > 1/2$ and

$$e^{-b_m \Delta_L} \leq \frac{3u_m}{1-3u_m} \leq \frac{3}{C(1-3u)} (Cu)^{2^{m-K}} < (Cu)^{2^{m-K}},$$

so $\Delta_L > \log(1/(Cu))/b_K > g$. Together, the intervals $[n_j, 2n_j]$ for $0 \leq j < K$ and the tail $L \geq n_K$ cover every even $L \geq n_0$. □

We generate the prefix (p_j, q_j) from the asymmetric recurrence with $f(x) = x^2/[2(1-x)^2]$:

$$\begin{aligned} p_{j+1} &= \lceil f(1-(1-p_j)^2(1-q_j)) \rceil_\delta, \\ q_{j+1} &= \lceil f(1-(1-q_j)^2(1-p_{j+1})) \rceil_\delta, \end{aligned} \tag{S44}$$

where $\lceil x \rceil_\delta = \delta \lceil x/\delta \rceil$. For the lifted seeds at $(60, rb_0)$, we take $K = 10$ steps with $\delta = 10^{-2048}$ and apply Proposition S1.

Table S3 lists two parameter choices. The integer choice $(k, r) = (30, 2)$ has $h = 2$, so its spatial residual (S36) uses only second and fourth powers and already gives $\Delta_L > 0.07045$ for even $L \geq 60$. The fractional choice $(k, r) = (28, 17/8)$ gives $\Delta_L > 0.072901771$ for even $L \geq 60$.

To cover the remaining even lengths $36 \leq L < 60$, we evaluate the unlifted base-seed formulas (S27)–(S28) at $(36, b_0)$ using the sharper caps (S30) and $\ell = 0.999997937629$, validated by the capped-mass inequality $\mathcal{C}(m_J, \ell^{12}, 6) + 3\mathcal{C}(m_N, V^{12}, 6) < 1$. At $j = 0$, the resulting factor $A_0 = (1-p_0)^2(1-q_0)$ satisfies $\log(A_0/(1-A_0))/b_0 > 0.09$, which by Eq. (S43) bounds $\Delta_L > 0.09$ on every even $36 \leq L \leq 72$. Combining this bridge on $[36, 72]$ with the lifted bound on $[60, \infty)$ proves Eq. (S2) for every even $L \geq 36$.

SIII.C. Reproducibility and arithmetic assumptions

The repository provides the stored certificates and standalone verifiers for the finite inputs and scalar inequalities above.

To keep every scalar check in exact rational arithmetic, we enclose rational powers and logarithms by rational endpoints. For $x \geq 0$ and a rational exponent $a/b \geq 0$ with $b > 0$, nonnegative rational bounds l, u satisfy $l \leq x^{a/b} \leq u$ if and only if the integer powers satisfy $l^b \leq x^a \leq u^b$. To enclose $\log x$ for $x > 0$, we write $x = 2^k y$ with $k \in \mathbb{Z}$ and

$1 \le y < 2$, and set $z = (y-1)/(y+1) \in [0, 1/3)$. The series

$$\log y = 2 \sum_{j=0}^{M-1} \frac{z^{2j+1}}{2j+1} + E_M,$$
$$0 \le E_M \le \frac{2z^{2M+1}}{(2M+1)(1-z^2)} \tag{S45}$$

at $M = 40$ encloses $\log y$ and $\log 2$ in rational intervals, which we round outward. Floating-point calculations are used only to select parameters, never in the certificate checks.

Our trusted base consists of exact integer and rational arithmetic, the explicit Poisson and Horner error bounds for the short-ring traces, and the analytical transfer, interpolation, and doubling inequalities proved in Sections SII.A–SII.F and above. Together with the finite trace and trial inputs of Ref. [7], these checks establish Eq. (S2) for the periodic spin-1 chain (S3).