

Blueprint for arXiv:2009.12982
Quantum Soundness of the Classical Low Individual Degree Test

MIPStarRE Project

May 18, 2026

Last updated: May 18, 2026

Contents

1	Overview	4
2	The low individual degree test	6
2.1	The game and its strategies	6
2.2	Formal soundness target	7
3	Preliminaries	9
3.1	Polynomials and measurements	9
3.2	Consistency and state-dependent distance	11
3.3	Consistency from state-dependent distance	13
3.4	Strong self-consistency	15
4	Making measurements projective	19
4.0.1	Naimark dilation	21
4.0.2	Orthogonalization lemma	22
5	Expansion in the hypercube graph	34
5.1	The graph and its spectrum	34
5.2	Local and global variance	36
6	Global variance of the points measurements	38
7	Self-improvement	40
7.1	Non-projective Output	40
7.1.1	A Semidefinite Program	45
7.2	Projective Output	50
8	Commutativity	54
8.1	Commutativity of the point measurements	54
8.2	Commutativity of G after evaluation	55
8.3	Commutativity of G	57
9	Pasting	60
9.0.1	From Measurements to Submeasurements	62
9.0.2	The Pasted Submeasurement	63
9.0.3	Strong Self-Consistency and Commutation of $\widehat{G}$	65
9.0.4	Sandwiching Lemmas	68
9.0.5	Consistency of H with A	70

9.0.6	Completeness of H	72
10	The main induction step	79
10.1	Restricting to one slice	79
10.2	The inductive construction	79
10.3	Proof of the main formal theorem	84

Chapter 1

Overview

Theorem 1.1 (Raz–Safra). *Suppose Provers A and B pass the $k = 2$ surface-versus-point low-degree test with probability $1 - \varepsilon$. Then there exists a degree- d polynomial $g: \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ such that*

$$\mathbb{P}_{u \sim \mathbb{F}_q^m}[g(u) = a] \geq 1 - \varepsilon - \text{poly}(m) \cdot \text{poly}(d/q).$$

Here a denotes the prover’s point answer.

Remark 1.2. The current Lean declaration is a wrapper around the quoted classical Raz–Safra theorem: it takes the specialized auxiliary hypothesis at the ambient point-answer function, error parameter, and chosen slack bound, whose premise is the explicit pass predicate . The surface test itself is modeled using parameterized genuine 2-dimensional surfaces, i.e. ordered frames with linearly independent directions, together with a hidden parameter specifying the queried point on the surface, so the surface prover does not see that point directly. Its conclusion currently returns the repository-wide Lean type , which encodes individual degree at most d rather than a separate total-degree-at-most- d object. Accordingly, this result is not marked as formalized.

Proof. See Raz–Safra [RS97]. □

Theorem 1.3 (Polishchuk–Spielman). *Suppose Provers A and B pass the low individual degree test with probability $1 - \varepsilon$. Then there exists a polynomial $g: \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ with individual degree d such that*

$$\mathbb{P}_{u \sim \mathbb{F}_q^m}[g(u) = a] \geq 1 - \text{poly}(m) \cdot (\text{poly}(\varepsilon) + \text{poly}(d/q)).$$

Here a denotes the prover’s point answer.

Remark 1.4. The current Lean declaration keeps the external Polishchuk–Spielman dependency explicit through a caller-chosen slack-bound parameter together with the auxiliary hypothesis . The repository no longer fixes any convenience placeholder formula for the schematic term $\text{poly}(m) \cdot (\text{poly}(\varepsilon) + \text{poly}(d/q))$ appearing in the overview statement above: any concrete rate must instead be supplied by the specialized external hypothesis. Accordingly, this result is not marked as formalized.

Proof. See Polishchuk–Spielman [PS94]. □

Remark 1.5 (A degree- $(d + 1)$ strategy that passes with probability $1 - \frac{1}{m}$). Fix the polynomial $h(x_1, \dots, x_m) = x_1^{d+1}$. Consider the following classical strategy for the degree- d low individual

degree test. On a point query $u \in \mathbb{F}_q^m$, Prover A answers $a = h(u)$. On an axis-parallel line query $\ell = \{u + xe_i : x \in \mathbb{F}_q\}$, Prover B returns the restriction $h|_\ell$ when $i \neq 1$, since h is then constant along ℓ , and returns the zero polynomial when $i = 1$.

The verifier certainly accepts whenever $i \neq 1$, which happens with probability $1 - \frac{1}{m}$. Thus the strategy passes the degree- d test with probability at least $1 - \varepsilon$ for $\varepsilon = \frac{1}{m}$. However, Prover A is answering according to a polynomial whose individual degree in the first variable is $d + 1$. By Schwartz–Zippel, any polynomial g of individual degree at most d agrees with h on at most a $\frac{d+1}{q}$ fraction of all inputs, so

$$\mathbb{P}_{u \sim \mathbb{F}_q^m}[g(u) = a] \leq 1 - m\varepsilon + \frac{d+1}{q}.$$

This shows that the m -dependent slack in Theorems 1.3 and 1.6 is unavoidable up to polynomial factors.

Theorem 1.6 (Main theorem, informal). *If $d > 0$ and a two-prover projective strategy passes the (m, q, d) low individual degree test with probability $1 - \varepsilon$, then there are global polynomial measurements whose evaluations agree with the point answers except with error $\text{poly}(m)(\text{poly}(\varepsilon) + \text{poly}(d/q))$.*

Proof. This is the informal form of Theorem 2.12. □

Chapter 2

The low individual degree test

2.1 The game and its strategies

Definition 2.1 (Roles). A role is an element of $\{A, B\}$. For a role $r \in \{A, B\}$, write $\bar{r}$ for the other element of $\{A, B\}$.

Definition 2.2 (Low individual degree test). Fix integers $m, d \geq 0$ and a prime power q . The (m, q, d) low individual degree test has three equally likely subtests. In the axis-parallel lines test, choose a uniformly random role $r \in \{A, B\}$, a uniformly random point $u \in \mathbb{F}_q^m$, and a uniformly random coordinate $i \in \{1, \dots, m\}$. Let $\ell = \{u + te_i : t \in \mathbb{F}_q\}$. Player r receives ℓ and returns a degree- d univariate polynomial $f: \ell \rightarrow \mathbb{F}_q$; Player $\bar{r}$ receives u and returns a field element $a \in \mathbb{F}_q$. The verifier accepts when $f(u) = a$. In the self-consistency test, both provers receive the same uniformly random point $u \in \mathbb{F}_q^m$ and must return the same field element. In the diagonal lines test, choose a uniformly random role $r \in \{A, B\}$, a uniformly random point $u \in \mathbb{F}_q^m$, a uniformly random index $i \in \{1, \dots, m\}$, and a uniformly random direction vector $v \in \mathbb{F}_q^m$ whose last $m - i$ coordinates are 0. Let $\ell = \{u + tv : t \in \mathbb{F}_q\}$. Player r receives ℓ and returns a degree- md univariate polynomial $f: \ell \rightarrow \mathbb{F}_q$; Player $\bar{r}$ receives u and returns a field element $a \in \mathbb{F}_q$. The verifier accepts when $f(u) = a$.

Lemma 2.3 (Branch-average form of the classical test). *For every classical strategy, the acceptance probability of the low individual degree test is*

$$\frac{1}{3} \left(\frac{P_{\text{axis}}^{\text{line, point}} + P_{\text{axis}}^{\text{point, line}}}{2} + P_{\text{self}} + \frac{P_{\text{diag}}^{\text{line, point}} + P_{\text{diag}}^{\text{point, line}}}{2} \right),$$

where the five terms are the acceptance probabilities of the two axis-parallel role orderings, the self-consistency branch, and the two diagonal role orderings.

Definition 2.4 (Symmetric projective strategy). A symmetric projective strategy for the (m, q, d) low individual degree test consists of a permutation-invariant bipartite state $|\psi\rangle \in \mathcal{H} \otimes \mathcal{H}$, a projective point measurement $A^u = \{A_a^u\}_{a \in \mathbb{F}_q}$ for each $u \in \mathbb{F}_q^m$, a projective axis-parallel line measurement $B^\ell = \{B_f^\ell\}$ for each axis-parallel line ℓ , and a projective diagonal-line measurement $L^\ell = \{L_f^\ell\}$ for each line ℓ .

Definition 2.5 (General projective strategy). A general projective strategy for the (m, q, d) low individual degree test consists of a bipartite state $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ together with point, axis-parallel line, and diagonal-line projective measurements for each prover separately. We write these measurements as A^A, B^A, L^A on the first prover and A^B, B^B, L^B on the second.

Lemma 2.6 (Bipartite strategy block identities). *The block operators obtained from a bipartite projective strategy are positive semidefinite, and the role-indexed blocks multiply according to the Kronecker delta on roles.*

Remark 2.7. Throughout this work, the term strategy refers to a projective strategy, and the term symmetric strategy refers to a symmetric projective strategy.

Definition 2.8 (Good strategy). A strategy is $(\varepsilon, \delta, \gamma)$ -good if it passes the axis-parallel lines test with probability at least $1 - \varepsilon$, the self-consistency test with probability at least $1 - \delta$, and the diagonal lines test with probability at least $1 - \gamma$.

Proposition 2.9 (Good-strategy characterization). *Using notation which will be introduced in Chapter 3 below, a symmetric strategy is $(\varepsilon, \delta, \gamma)$ -good if and only if it satisfies the following three conditions. For ℓ and u as in the axis-parallel lines test,*

$$A_a^u \otimes I \simeq_\varepsilon I \otimes B_{[f(u)=a]}^\ell, \quad \text{and} \quad A_a^u \otimes I \simeq_\delta I \otimes A_a^u.$$

And for ℓ and u as in the diagonal lines test,

$$A_a^u \otimes I \simeq_\gamma I \otimes L_{[f(u)=a]}^\ell.$$

Definition 2.10 (Last-direction line notation). For $u \in \mathbb{F}_q^m$, the last-direction axis-parallel line in $\mathbb{F}_q^{m+1}$ is

$$\ell_u = \{(u, x) \mid x \in \mathbb{F}_q\}.$$

If (ψ, A, B, L) is a symmetric strategy for the $(m+1, q, d)$ low individual degree test, we write B_f^u for the operator $B_{f^\ell}^u$. For a function $f: \ell_u \rightarrow \mathbb{F}_q$, we also write $f(x)$ for $f(u, x)$.

Definition 2.11 (Restricted diagonal lines test). Consider the (m, q, d) low individual degree test. For $j \in \{1, \dots, m\}$, the j -restricted diagonal lines test is the diagonal lines test conditioned on $i = j$. In particular, the m -restricted diagonal lines test is the diagonal lines test in which the sampled line is a uniformly random line in $\mathbb{F}_q^m$.

2.2 Formal soundness target

Theorem 2.12 (Quantum soundness of the low individual degree test). *Let $(\psi, A^A, B^A, L^A, A^B, B^B, L^B)$ be a projective strategy that passes the (m, q, d) low individual degree test with probability at least $1 - \varepsilon$. Assume $d > 0$. Let $k > 0$ with $k \geq 400md$ and set*

$$\nu = 100000k^2m^4 \left(\varepsilon^{1/40000} + (d/q)^{1/40000} + e^{-k/(2560000m^2)} \right).$$

Then there exist projective measurements $G^A, G^B \in \text{PolyMeas}(m, q, d)$ with the following properties:

1. *on average over $u \sim \mathbb{F}_q^m$,*

$$A_a^{A,u} \otimes I \simeq_\nu I \otimes G_{[g(u)=a]}^B, \quad I \otimes A_a^{B,u} \simeq_\nu G_{[g(u)=a]}^A \otimes I;$$

- 2.

$$G_g^A \otimes I \simeq_\nu I \otimes G_g^B.$$

Remark 2.13 (Internal witness form of the final transport). The witness theorem is a Lean-only formally verified final-transport result: once the Section 6 role witness and the post-role projective-completion witness have been constructed from the hypotheses of Theorem 2.12, it derives the three consistency conclusions of the theorem. It is not a replacement statement for Theorem 2.12, and it supplies no additional hypothesis to the paper theorem.

Remark 2.14 (k -bound boundary for the public theorem). The paper statement uses the weaker hypothesis $k \geq md$, but the proof invokes the Section 6 pasting theorem with side condition $k \geq 400md$. This side condition is not implied by $k \geq md$. The Lean and blueprint public statement therefore records the stronger large- k assumption required by the pasting step rather than hiding the side-condition gap. The helper `mainFormal_trivial_witness` remains only for already-saturated branches where $\nu \geq 1$ makes the three $\simeq_\nu$ conclusions vacuous; it is not a replacement for a proof in the interval $md \leq k < 400md$.

Chapter 3

Preliminaries

3.1 Polynomials and measurements

For complex numbers $\alpha, \beta \in \mathbb{C}$, write $\alpha \approx_\varepsilon \beta$ when $|\alpha - \beta| \leq \varepsilon$. In particular,

$$\text{if } \alpha \approx_\varepsilon \beta \text{ and } \beta \approx_\delta \gamma, \text{ then } \alpha \approx_{\varepsilon+\delta} \gamma. \quad (3.1)$$

Fix a prime power $q = p^t$. Write $\omega = e^{2\pi i/p}$, and let $\text{tr} : \mathbb{F}_q \rightarrow \mathbb{F}_p$ denote the finite-field trace.

Definition 3.1 (Finite field trace). The finite-field trace is the map $\text{tr} : \mathbb{F}_q \rightarrow \mathbb{F}_p$ defined by

$$\text{tr}[x] = \sum_{\ell=0}^{t-1} x^{p^\ell}.$$

Lemma 3.2. *Let $a \in \mathbb{F}_q$. Then*

$$\mathbb{E}_{x \sim \mathbb{F}_q} \omega^{\text{tr}[x \cdot a]} = \begin{cases} 1 & \text{if } a = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If $a = 0$, then $\text{tr}[x \cdot a] = 0$ for every x , so the average is 1. If $a \neq 0$, choose $y \in \mathbb{F}_q$ such that $\text{tr}[a \cdot y] \neq 0$, and set

$$C = \mathbb{E}_{x \sim \mathbb{F}_q} \omega^{\text{tr}[x \cdot a]}.$$

Translation invariance of the uniform distribution gives

$$C = \mathbb{E}_{x \sim \mathbb{F}_q} \omega^{\text{tr}[(x+y) \cdot a]} = C \cdot \omega^{\text{tr}[y \cdot a]}.$$

Since $\omega^{\text{tr}[y \cdot a]} \neq 1$, it follows that $C = 0$. □

Lemma 3.3. *Let $v \in \mathbb{F}_q^m$. Then*

$$\mathbb{E}_{u \sim \mathbb{F}_q^m} \omega^{\text{tr}[u \cdot v]} = \begin{cases} 1 & \text{if } v = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By linearity of the trace and independence of the coordinates,

$$\mathbb{E}_{u \sim \mathbb{F}_q^m} \omega^{\text{tr}[u \cdot v]} = \prod_{i=1}^m \mathbb{E}_{u_i \sim \mathbb{F}_q} \omega^{\text{tr}[u_i \cdot v_i]}.$$

Proposition 3.2 shows that the i -th factor is 1 when $v_i = 0$ and 0 otherwise, so the product is 1 exactly when $v = 0$. □

Definition 3.4 (Low individual degree polynomials). Let $\mathcal{P}(m, q, d)$ be the set of polynomials $g \in \mathbb{F}_q[x_1, \dots, x_m]$ whose degree in each variable is at most d , viewed as functions $\mathbb{F}_q^m \rightarrow \mathbb{F}_q$ via evaluation. (Over finite fields, distinct low-degree polynomials can induce the same function; we identify elements of $\mathcal{P}(m, q, d)$ with their polynomial representatives, not their functional equivalence classes.)

Lemma 3.5 (Monotonicity of individual-degree classes). *Here, “individual degree d ” means degree at most d in each coordinate. In particular,*

$$\mathcal{P}(m, q, d) \subseteq \mathcal{P}(m, q, d + 1).$$

Lemma 3.6 (Schwartz–Zippel lemma [Sch80, Zip79]). *Let $g, h : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ be two distinct polynomials of total degree d . Then*

$$\mathbb{P}_{x \sim \mathbb{F}_q^m}[g(x) = h(x)] \leq \frac{d}{q}.$$

Proof. Apply the Schwartz–Zippel lemma to the nonzero polynomial $g - h$, which has total degree at most d . $\square$

Lemma 3.7 (Schwartz–Zippel for individual degree). *If $g, h \in \mathcal{P}(m, q, d)$ are distinct, then*

$$\mathbb{P}_{u \sim \mathbb{F}_q^m}[g(u) = h(u)] \leq \frac{md}{q}.$$

Proof. The difference $g - h$ is a nonzero polynomial of total degree at most md . Apply Lemma 3.6. $\square$

Lemma 3.8 (Polynomial agreement bound on coded points). *Let $g, g' : \text{Polynomial params}$ be two distinct full polynomial outcomes with individual degrees at most d . Then*

$$\mathbb{E}_{u \sim \text{Point params}}[\mathbf{1}[g(u) = g'(u)]] \leq \frac{md}{q}.$$

This packages the md/q loss term used in the `mainFormal` self-consistency cascade (`inductive_step.tex`, lines 119–133) and in `comMain` (`commutativity-G.tex`).

Proof. Transport along the equivalence between coded $\mathbb{F}_q$ points and the underlying scalar function space, then apply Lemma 3.7. $\square$

Lemma 3.9 (Axis-line polynomial agreement bound). *Let $f, h : \text{AxisLinePolynomial params}$ be two line-polynomial outcomes with distinct underlying polynomials. Then*

$$\mathbb{E}_{t \sim \mathbb{F}_q}[\mathbf{1}[f(t) = h(t)]] \leq \frac{md}{q}.$$

The native univariate root-counting bound is d/q ; the statement deliberately pads this to the paper’s ambient md/q loss for Section 6.1.

Proof. Reindex coded line parameters to the scalar field, count the roots of the nonzero univariate polynomial $f - h$, and use $m \geq 1$ to pad d/q to md/q . $\square$

Definition 3.10 (Measurements and submeasurements). Let $\mathcal{H}$ be a Hilbert space and let $\mathcal{A}$ be a set of outcomes. A submeasurement on $\mathcal{A}$ is a family $A = \{A_a\}_{a \in \mathcal{A}}$ of Hermitian positive semidefinite operators on $\mathcal{H}$ such that $\sum_a A_a \leq I$. It is a measurement when $\sum_a A_a = I$, and it is projective when each A_a is an idempotent projection. (In the Lean formalization, outcome sets are assumed finite throughout.)

Definition 3.11 (Low-degree polynomial measurements). We write $\text{PolySub}(m, q, d)$ for the submeasurements indexed by $\mathcal{P}(m, q, d)$, and $\text{PolyMeas}(m, q, d)$ for the corresponding measurements.

Definition 3.12 (Post-processing). Let $A = \{A_a\}_{a \in \mathcal{A}}$ be a family of operators and let $f: \mathcal{A} \rightarrow \mathcal{B}$. The post-processed family $A_{[f(a)=b]}$ is defined by

$$A_{[f(a)=b]} = \sum_{a: f(a)=b} A_a.$$

Proposition 3.13 (Post-processing preserves measurements). *Let $A = \{A_a\}_{a \in \mathcal{A}}$ be a family of operators and let $f: \mathcal{A} \rightarrow \mathcal{B}$. Then*

$$\sum_a A_a = \sum_b A_{[f(a)=b]}.$$

Consequently, if $\{A_a\}$ is a submeasurement, respectively a measurement, then $\{A_{[f(a)=b]}\}$ is again a submeasurement, respectively a measurement.

Proof. Each $a \in \mathcal{A}$ contributes exactly once to the right-hand side, namely in the summand indexed by $b = f(a)$. $\square$

Remark 3.14 (Post-processing notation). In the notation $A_{[f(a)=b]}$, the measurement outcome is the variable to which the function is applied. For example, if $G = \{G_g\} \in \text{PolySub}(m, q, d)$ and $u \in \mathbb{F}_q^m$, then

$$G_{[g(u)=a]} = \sum_{g: g(u)=a} G_g.$$

Here g is the measurement outcome, and u is the evaluation point.

Definition 3.15 (Completion of a submeasurement). Let $A = \{A_a^x\}_{a \in \mathcal{A}}$ be a submeasurement. Its completion is the measurement $\widehat{A}$ with outcome set $\widehat{\mathcal{A}} = \mathcal{A} \cup \{\perp\}$ given by $\widehat{A}_a^x = A_a^x$ for $a \in \mathcal{A}$ and $\widehat{A}_\perp^x = I - \sum_a A_a^x$.

3.2 Consistency and state-dependent distance

Definition 3.16 (Consistency). Let $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$. Let $A = \{A_a^x\}$ and $B = \{B_a^x\}$ be submeasurements with the same answer set, and let $\mathcal{D}$ be a distribution on the question set. We write

$$A_a^x \otimes I \simeq_\delta I \otimes B_a^x$$

when

$$\mathbb{E}_{x \sim \mathcal{D}} \sum_{a \neq b} \langle \psi | A_a^x \otimes B_b^x | \psi \rangle \leq \delta. \quad (3.2)$$

Lemma 3.17 (Characterization of a good symmetric strategy). *A symmetric projective strategy (ψ, A, B, L) is $(\varepsilon, \delta, \gamma)$ -good if and only if the following hold on the relevant test distributions:*

$$A_a^u \otimes I \simeq_\varepsilon I \otimes B_{[f(u)=a]}^\ell, \quad A_a^u \otimes I \simeq_\delta I \otimes A_a^u, \quad A_a^u \otimes I \simeq_\gamma I \otimes L_{[f(u)=a]}^\ell.$$

Proof. Each subtest accepts exactly when the two measurement outcomes agree. Because the relevant families are measurements, Lemma 3.18 rewrites those acceptance probabilities as consistency bounds. $\square$

Lemma 3.18 (Consistency for measurements). *If $A = \{A_a^x\}$ and $B = \{B_a^x\}$ are measurements, then*

$$A_a^x \otimes I \simeq_\delta I \otimes B_a^x \quad \text{if and only if} \quad \mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes B_a^x | \psi \rangle \geq 1 - \delta.$$

Proof. Expand the off-diagonal sum by writing $\sum_{a \neq b} A_a^x \otimes B_b^x = \sum_b (I - A_b^x) \otimes B_b^x$. $\square$

Definition 3.19 (State-dependent distance). Let $|\psi\rangle \in \mathcal{H}$, let $A = \{A_a^x\}$ and $B = \{B_a^x\}$ be families of operators on $\mathcal{H}$, and let $\mathcal{D}$ be a distribution on the question set. We write

$$A_a^x \approx_\delta B_a^x$$

when

$$\mathbb{E}_{x \sim \mathcal{D}} \sum_a \|(A_a^x - B_a^x)|\psi\rangle\|^2 \leq \delta.$$

In particular, if $A_a^x \otimes I \simeq_\varepsilon I \otimes C_a^x$, the question is what hypothesis on A and B allows one to conclude that

$$B_a^x \otimes I \simeq_\varepsilon I \otimes C_a^x. \quad (3.3)$$

Lemma 3.20 (Transferring consistency through state-dependent distance). *Let $A = \{A_a^x\}$ and $B = \{B_a^x\}$ be measurements and let $C = \{C_a^x\}$ be a submeasurement. If $A_a^x \otimes I \simeq_\delta I \otimes C_a^x$ and $A_a^x \otimes I \approx_\varepsilon B_a^x \otimes I$, then*

$$B_a^x \otimes I \simeq_{\delta + \sqrt{\varepsilon}} I \otimes C_a^x.$$

Proof. Rewrite the inconsistency with C as total mass minus diagonal overlap. The total mass is unchanged because A and B are measurements, and the diagonal overlap changes by at most $\sqrt{\varepsilon}$ by Cauchy-Schwarz. $\square$

Lemma 3.21 (Consistency implies state-dependent distance for measurements). *If A and B are measurements and $A_a^x \otimes I \simeq_\delta I \otimes B_a^x$, then*

$$A_a^x \otimes I \approx_{2\delta} I \otimes B_a^x.$$

If both measurements are projective, the converse also holds: if $A_a^x \otimes I \approx_{2\delta} I \otimes B_a^x$ then $A_a^x \otimes I \simeq_\delta I \otimes B_a^x$.

Proof. Expand the squared norm of $(A_a^x \otimes I - I \otimes B_a^x)|\psi\rangle$ and use the diagonal-overlap formula from Lemma 3.18; projectivity turns the diagonal inequality into an equality, giving the converse. $\square$

Remark 3.22. The implication above does not extend to submeasurements. For example, if $A_a^x = 0$ for all a , then $A_a^x \otimes I \simeq_0 I \otimes B_a^x$, but

$$\mathbb{E}_x \sum_a \|(A_a^x \otimes I - I \otimes B_a^x)|\psi\rangle\|^2 = \mathbb{E}_x \sum_a \|(I \otimes B_a^x)|\psi\rangle\|^2,$$

which is nonzero unless $(I \otimes B_a^x)|\psi\rangle = 0$ for all x and a .

3.3 Consistency from state-dependent distance

Theorem 3.23 (Closeness of inner products). *Let $\{A_a^x\}$, $\{B_a^x\}$, and $\{C_{a,b}^x\}$ be matrices. Suppose that $A_a^x \approx_\gamma B_a^x$ and that for all x ,*

$$\sum_a \left(\sum_b C_{a,b}^x \right) \left(\sum_b C_{a,b}^x \right)^\dagger \leq I.$$

Then

$$\mathbb{E}_x \sum_{a,b} \langle \psi | C_{a,b}^x A_a^x | \psi \rangle \approx_{\sqrt{\gamma}} \mathbb{E}_x \sum_{a,b} \langle \psi | C_{a,b}^x B_a^x | \psi \rangle. \quad (3.4)$$

Similarly, suppose that $(A_a^x)^\dagger \approx_\gamma (B_a^x)^\dagger$ and that for all x ,

$$\sum_a \left(\sum_b C_{a,b}^x \right)^\dagger \left(\sum_b C_{a,b}^x \right) \leq I.$$

Then

$$\mathbb{E}_x \sum_{a,b} \langle \psi | A_a^x C_{a,b}^x | \psi \rangle \approx_{\sqrt{\gamma}} \mathbb{E}_x \sum_{a,b} \langle \psi | B_a^x C_{a,b}^x | \psi \rangle. \quad (3.5)$$

Proof. To prove (3.4), write the difference as

$$\mathbb{E}_x \sum_a \langle \psi | \left(\sum_b C_{a,b}^x \right) (A_a^x - B_a^x) | \psi \rangle.$$

Cauchy–Schwarz bounds its magnitude by

$$\left(\mathbb{E}_x \sum_a \langle \psi | \left(\sum_b C_{a,b}^x \right) \left(\sum_b C_{a,b}^x \right)^\dagger | \psi \rangle \right)^{1/2} \left(\mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x)^\dagger (A_a^x - B_a^x) | \psi \rangle \right)^{1/2},$$

and the two hypotheses bound these factors by 1 and $\sqrt{\gamma}$ respectively. For (3.5), rewrite the difference as

$$\mathbb{E}_x \sum_{a,b} \langle \psi | (C_{a,b}^x)^\dagger ((A_a^x)^\dagger - (B_a^x)^\dagger) | \psi \rangle,$$

and apply (3.4) to the adjoint families. $\square$

Theorem 3.24. *Let $A = \{A_a^x\}$, $B = \{B_a^x\}$, and $C = \{C_a^x\}$ be submeasurements such that $A_a^x \approx_\delta B_a^x$. Then*

$$\mathbb{E}_x \sum_a \langle \psi | A_a^x C_a^x | \psi \rangle \approx_{\sqrt{\delta}} \mathbb{E}_x \sum_a \langle \psi | B_a^x C_a^x | \psi \rangle.$$

Proof. The difference has magnitude

$$\left| \mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x) C_a^x | \psi \rangle \right|.$$

By Cauchy–Schwarz this is at most

$$\left(\mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x)^2 | \psi \rangle \right)^{1/2} \left(\mathbb{E}_x \sum_a \langle \psi | (C_a^x)^2 | \psi \rangle \right)^{1/2}.$$

The first factor is at most $\sqrt{\delta}$ by hypothesis, and the second is at most 1 because C is a submeasurement. $\square$

Theorem 3.25. Let $\{A_a^x\}$, $\{B_a^x\}$, and $\{C_{a,b}^x\}$ be matrices. Suppose that $A_a^x \approx_\delta B_a^x$ and that for all x and a ,

$$\sum_b (C_{a,b}^x)^\dagger (C_{a,b}^x) \leq I.$$

Then

$$C_{a,b}^x A_a^x \approx_\delta C_{a,b}^x B_a^x.$$

Proof. The error term is

$$\mathbb{E}_x \sum_{a,b} \langle \psi | (A_a^x - B_a^x)^\dagger (C_{a,b}^x)^\dagger C_{a,b}^x (A_a^x - B_a^x) | \psi \rangle.$$

Using $\sum_b (C_{a,b}^x)^\dagger (C_{a,b}^x) \leq I$, this is bounded by

$$\mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x)^\dagger (A_a^x - B_a^x) | \psi \rangle \leq \delta.$$

□

Lemma 3.26 (Triangle inequality for vectors squared). For vectors $|\psi_1\rangle, \dots, |\psi_k\rangle$,

$$\| |\psi_1\rangle + \dots + |\psi_k\rangle \|^2 \leq k(\| |\psi_1\rangle \|^2 + \dots + \| |\psi_k\rangle \|^2).$$

Proof. First,

$$(x_1 + \dots + x_k)^2 \leq k(x_1^2 + \dots + x_k^2) \quad (3.6)$$

for all real numbers $x_1, \dots, x_k$. Apply (3.6) to $x_i = \| |\psi_i\rangle \|$ and combine it with the triangle inequality. □

Lemma 3.27 (Triangle inequality for $\approx_\delta$). Suppose $(A_i)_a^x \approx_{\delta_i} (A_{i+1})_a^x$ for $i = 1, \dots, k$. Then

$$(A_1)_a^x \approx_{k(\delta_1 + \dots + \delta_k)} (A_{k+1})_a^x.$$

Proof. Expand $(A_1 - A_{k+1})| \psi \rangle$ as a telescoping sum and apply Lemma 3.26. □

Remark 3.28 (Lean finite-step triangle helpers). The general operator-family theorem above is complemented in Lean by reusable binary and three-step helper lemmas for operator expectations, ‘qSDD’, and ‘SDDRel’. The three-step ‘SDDRel’ helper is the specialization used in the Step 6 projectivization chain in Theorem 2.12; this remark is only a cross-reference and makes no separate proof-level completeness claim.

Lemma 3.29 (Triangle inequality for $\simeq_\delta$). Suppose $A_a^x \otimes I \simeq_\varepsilon I \otimes B_a^x$, $C_a^x \otimes I \simeq_\delta I \otimes B_a^x$, and $C_a^x \otimes I \simeq_\gamma I \otimes D_a^x$, where all four families are measurements. Then

$$A_a^x \otimes I \simeq_{\varepsilon + 2\sqrt{\delta + \gamma}} I \otimes D_a^x.$$

Proof. Convert the two consistencies involving C to state-dependent distance, compose them by the triangle inequality, and transfer the result back to consistency by Lemma 3.20. □

Lemma 3.30 (Data processing for consistency). If $A = \{A_a^x\}$ and $B = \{B_a^x\}$ are measurements such that $A_a^x \otimes I \simeq_\delta I \otimes B_a^x$, and f is a map on the answer set, then

$$A_{[f(a)=b]}^x \otimes I \simeq_\delta I \otimes B_{[f(a)=b]}^x.$$

Proof. Merging outcome classes can only decrease the off-diagonal mass. $\square$

Lemma 3.31 (Consistency controls a submeasurement). *Let $A = \{A_a^x\}$ be a submeasurement and $B = \{B_a^x\}$ a measurement such that $A_a^x \otimes I \simeq_\gamma I \otimes B_a^x$. Then*

$$A_a^x \otimes I \approx_\gamma A_a^x \otimes B_a^x \approx_\gamma A^x \otimes B_a^x, \quad (3.7)$$

where $A^x = \sum_a A_a^x$. In particular,

$$A_a^x \otimes I \approx_{4\gamma} A^x \otimes B_a^x.$$

Proof. For the first approximation in (3.7), bound

$$\mathbb{E}_x \sum_a \langle \psi | (A_a^x \otimes (I - B_a^x))^2 | \psi \rangle$$

by the inconsistency between A and B . The second approximation is identical, applied to

$$\mathbb{E}_x \sum_a \langle \psi | ((A^x - A_a^x) \otimes B_a^x)^2 | \psi \rangle.$$

Then compose the two bounds with Lemma 3.27. $\square$

Lemma 3.32 (Switch-sandwich). *Suppose $A = \{A_a^x\}$ is a projective submeasurement satisfying*

$$A_a^x \otimes I \approx_\delta I \otimes A_a^x. \quad (3.8)$$

Then for every operator B with $0 \leq B \leq I$,

$$\mathbb{E}_x \sum_a \langle \psi | A_a^x B A_a^x \otimes I | \psi \rangle \approx_{2\sqrt{\delta}} \mathbb{E}_x \sum_a \langle \psi | B \otimes A_a^x | \psi \rangle \approx_{\sqrt{\delta}} \mathbb{E}_x \sum_a \langle \psi | B A_a^x \otimes I | \psi \rangle. \quad (3.9)$$

Proof. We prove the first approximation in (3.9) in two steps. First,

$$\mathbb{E}_x \sum_a \langle \psi | A_a^x B A_a^x \otimes I | \psi \rangle \approx_{\sqrt{\delta}} \mathbb{E}_x \sum_a \langle \psi | A_a^x B \otimes A_a^x | \psi \rangle. \quad (3.10)$$

This is obtained by applying Cauchy–Schwarz to the difference and using $B^2 \leq B \leq I$ together with (3.8). For the second step, move the remaining left copy of A_a^x across the bipartition in the same way; projectivity then collapses the sandwich and yields the first approximation in (3.9). The second approximation in (3.9) is the same argument with the rightmost factor A_a^x omitted. $\square$

3.4 Strong self-consistency

Lemma 3.33 (Completeness transfer to a projective approximation). *Let $A = \{A_a^x\}$ be a submeasurement and $P = \{P_a^x\}$ a projective submeasurement such that $A_a^x \otimes I \approx_\varepsilon P_a^x \otimes I$. Then*

$$\langle \psi | A \otimes I | \psi \rangle \geq \langle \psi | P \otimes I | \psi \rangle - 2\sqrt{\varepsilon}.$$

Proof. Since P is projective,

$$\langle \psi | P \otimes I | \psi \rangle = \mathbb{E}_x \sum_a \langle \psi | (P_a^x)^2 \otimes I | \psi \rangle.$$

Apply Proposition 3.24 twice to replace $(P_a^x)^2$ first by $A_a^x P_a^x$ and then by $(A_a^x)^2$, and conclude with $(A_a^x)^2 \leq A_a^x$. $\square$

Definition 3.34 (Strong self-consistency). Let $|\psi\rangle$ be permutation-invariant and let $A = \{A_a^x\}$ be a submeasurement. We say that A is δ -strongly self-consistent when

$$\mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes A_a^x | \psi \rangle \geq \langle \psi | A \otimes I | \psi \rangle - \delta.$$

Lemma 3.35 (Strong self-consistency implies ordinary self-consistency). *If A is δ -strongly self-consistent, then $A_a^x \otimes I \simeq_\delta I \otimes A_a^x$. If A is a measurement, the converse also holds.*

Proof. Rewrite the off-diagonal mass as the difference between the total mass of A and the diagonal overlap in the strong self-consistency inequality. When A is a full measurement, the total mass is the identity, so this inequality becomes an equality; hence the ordinary self-consistency defect and the strong self-consistency defect coincide, giving the converse. $\square$

Lemma 3.36 (Strong self-consistency and state-dependent distance). *If A is δ -strongly self-consistent, then*

$$A_a^x \otimes I \approx_{2\delta} I \otimes A_a^x.$$

If A is projective, the converse also holds.

Proof. Expanding the squared norm gives

$$\begin{aligned} & \mathbb{E}_x \sum_a \|(A_a^x \otimes I - I \otimes A_a^x)|\psi\rangle\|^2 \\ &= 2 \cdot \left(\mathbb{E}_x \sum_a \langle \psi | (A_a^x)^2 \otimes I | \psi \rangle - \mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes A_a^x | \psi \rangle \right) \\ &\leq 2 \cdot \left(\mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes I | \psi \rangle - \mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes A_a^x | \psi \rangle \right). \end{aligned} \quad (3.11)$$

The strong self-consistency bound controls the last line by 2δ . If A is projective, then (3.11) is an equality, which gives the converse. $\square$

Lemma 3.37 (Post-processing preserves strong self-consistency). *If A is δ -strongly self-consistent and f is a function on its answer set, then*

$$A_{[f(a)=b]}^x \otimes I \approx_{2\delta} I \otimes A_{[f(a)=b]}^x.$$

Proof. The same computation as in Lemma 3.36 gives

$$\begin{aligned} & \mathbb{E}_x \sum_b \|(A_{[f(a)=b]}^x \otimes I - I \otimes A_{[f(a)=b]}^x)|\psi\rangle\|^2 \\ &\leq 2 \cdot \left(\langle \psi | A \otimes I | \psi \rangle - \mathbb{E}_x \sum_b \langle \psi | A_{[f(a)=b]}^x \otimes A_{[f(a)=b]}^x | \psi \rangle \right). \end{aligned} \quad (3.12)$$

The post-processed diagonal term dominates $\mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes A_a^x | \psi \rangle$, so strong self-consistency bounds (3.12) by 2δ . $\square$

Lemma 3.38 (Completeness transfer for a strongly self-consistent approximation). *Let A be a δ -strongly self-consistent submeasurement and let B be a submeasurement such that $A_a^x \otimes I \approx_\varepsilon B_a^x \otimes I$. Then*

$$\langle \psi | B \otimes I | \psi \rangle \geq \langle \psi | A \otimes I | \psi \rangle - \delta - 2\sqrt{\varepsilon}.$$

Proof. Bound $\langle \psi | B \otimes I | \psi \rangle$ from below by $\mathbb{E}_x \sum_a \langle \psi | B_a^x \otimes B_a^x | \psi \rangle$. Then apply Proposition 3.24 twice to replace this by $\mathbb{E}_x \sum_a \langle \psi | A_a^x \otimes A_a^x | \psi \rangle$, and finish with strong self-consistency of A . $\square$

Lemma 3.39 (Self-consistency implies data-processing closeness). *Let A be a δ -strongly self-consistent submeasurement and let P be a projective submeasurement such that $P_a^x \otimes I \approx_\varepsilon A_a^x \otimes I$. Then for every function f ,*

$$P_{[f(a)=b]}^x \otimes I \approx_{8\delta+8\sqrt{\varepsilon}} A_{[f(a)=b]}^x \otimes I.$$

Proof. First prove

$$P_{[f(a)=b]}^x \otimes I \approx_{2\delta+4\sqrt{\varepsilon}} I \otimes A_{[f(a)=b]}^x. \quad (3.13)$$

Expanding the squared norm and using projectivity gives

$$\begin{aligned} & \mathbb{E}_x \sum_b \|(P_{[f(a)=b]}^x \otimes I - I \otimes A_{[f(a)=b]}^x)|\psi\rangle\|^2 \\ & \leq \langle \psi | P \otimes I | \psi \rangle + \langle \psi | I \otimes A | \psi \rangle - 2 \cdot \mathbb{E}_x \sum_b \langle \psi | P_{[f(a)=b]}^x \otimes A_{[f(a)=b]}^x | \psi \rangle. \end{aligned} \quad (3.14)$$

Proposition 3.33 bounds the first term by $\langle \psi | A \otimes I | \psi \rangle + 2\sqrt{\varepsilon}$, and Proposition 3.24 together with strong self-consistency bounds the third term below by $\langle \psi | A \otimes I | \psi \rangle - \delta - \sqrt{\varepsilon}$. Substituting into (3.14) yields (3.13). Now combine (3.13) with

$$A_{[f(a)=b]}^x \otimes I \approx_{2\delta} I \otimes A_{[f(a)=b]}^x$$

from Lemma 3.37, and conclude by Lemma 3.27. $\square$

Lemma 3.40 (Self-consistency moves one copy to the same side). *Let $A = \{A_a\}$ be a ζ -strongly self-consistent submeasurement. Then*

$$\sum_a \langle \psi | A_a^2 \otimes I | \psi \rangle \geq \sum_a \langle \psi | A_a \otimes I | \psi \rangle - \zeta.$$

Proof. Apply Cauchy-Schwarz to the two-sided overlap $\sum_a \langle \psi | A_a \otimes A_a | \psi \rangle$ and compare it with the strong self-consistency lower bound. $\square$

Lemma 3.41 (Bounding the missing mass of a close submeasurement). *Let $A = \{A_a\}$ be a measurement that is ζ -strongly self-consistent, and let $B = \{B_a\}$ be a submeasurement such that $A_a \otimes I \approx_\delta B_a \otimes I$. Writing $B = \sum_a B_a$, one has*

$$\|(I - B) \otimes I | \psi \rangle\|^2 \leq 2\sqrt{\delta} + \zeta.$$

Proof. Since $0 \leq I - B \leq I$, the left-hand side is at most $1 - \langle \psi | B \otimes I | \psi \rangle$. Compare $\sum_a \langle \psi | B_a^2 \otimes I | \psi \rangle$ with $\sum_a \langle \psi | A_a B_a \otimes I | \psi \rangle$ and then with $\sum_a \langle \psi | A_a^2 \otimes I | \psi \rangle$ using the $\approx_\delta$ hypothesis, and finish with Lemma 3.40. $\square$

Lemma 3.42 (Completing a close submeasurement to a full measurement). *Let $A = \{A_a\}$ be a measurement that is ζ -strongly self-consistent, let $B = \{B_a\}$ be a submeasurement such that $A_a \otimes I \approx_\delta B_a \otimes I$, and let C be the measurement obtained by adding the missing mass $I - B$ to one distinguished answer a^* . Then*

$$A_a \otimes I \approx_{2\delta+4\sqrt{\delta}+2\zeta} C_a \otimes I.$$

Proof of 3.42. By Lemma 3.26,

$$\sum_a \|(A_a - C_a) \otimes I | \psi \rangle\|^2 \leq 2\delta + 2 \cdot \|(I - B) \otimes I | \psi \rangle\|^2.$$

It remains to bound the residual term. Since $0 \leq I - B \leq I$,

$$\|(I - B) \otimes I|\psi\rangle\|^2 \leq 1 - \sum_a \langle \psi | B_a^2 \otimes I | \psi \rangle. \quad (3.15)$$

By Proposition 3.24,

$$\sum_a \langle \psi | B_a^2 \otimes I | \psi \rangle \approx_{\sqrt{\delta}} \sum_a \langle \psi | (A_a B_a) \otimes I | \psi \rangle \approx_{\sqrt{\delta}} \sum_a \langle \psi | A_a^2 \otimes I | \psi \rangle.$$

Proposition 3.43 gives

$$\sum_a \langle \psi | A_a^2 \otimes I | \psi \rangle \geq \langle \psi | A \otimes I | \psi \rangle - \zeta = 1 - \zeta.$$

Hence (3.15) implies

$$\|(I - B) \otimes I|\psi\rangle\|^2 \leq 2\sqrt{\delta} + \zeta.$$

Substituting this into the first bound proves the claim. $\square$

Theorem 3.43. *This is the same statement as Lemma 3.40: for a ζ -strongly self-consistent submeasurement A ,*

$$\sum_a \langle \psi | (A_a)^2 \otimes I | \psi \rangle \geq \sum_a \langle \psi | A_a \otimes I | \psi \rangle - \zeta.$$

Proof. Immediate from Lemma 3.40. $\square$

Chapter 4

Making measurements projective

Theorem 4.1 (Naimark dilation). *Let $|\psi\rangle$ be a state in $\mathcal{H}_A \otimes \mathcal{H}_B$. Let $A = \{A_a^x\}$ be a sub-measurement acting on $\mathcal{H}_A$ and $B = \{B_b^y\}$ be a sub-measurement acting on $\mathcal{H}_B$. Then there exist Hilbert spaces $\mathcal{H}_{A_{\text{aux}}}$ and $\mathcal{H}_{B_{\text{aux}}}$, a state $|\text{aux}\rangle \in \mathcal{H}_{A_{\text{aux}}} \otimes \mathcal{H}_{B_{\text{aux}}}$, and measurements $\widehat{A} = \{\widehat{A}_a^x\}$ and $\widehat{B} = \{\widehat{B}_b^y\}$ acting on $\mathcal{H}_A \otimes \mathcal{H}_{A_{\text{aux}}}$ and $\mathcal{H}_B \otimes \mathcal{H}_{B_{\text{aux}}}$, respectively, such that, if $|\widehat{\psi}\rangle = |\psi\rangle \otimes |\text{aux}\rangle$, then for all x, y, a, b ,*

$$\langle \psi | A_a^x \otimes B_b^y | \psi \rangle = \langle \widehat{\psi} | \widehat{A}_a^x \otimes \widehat{B}_b^y | \widehat{\psi} \rangle.$$

In addition, $|\text{aux}\rangle$ is a product state.

Proof.

For each question x , choose a local auxiliary state $|\text{aux}_{A,x}\rangle$ of dimension one larger than the number of outcomes of A^x , and apply Lemma 4.7 to obtain a projective sub-measurement $\widetilde{A}^x$. Do the same for each question y on Bob's side, obtaining auxiliary states $|\text{aux}_{B,y}\rangle$ and projective sub-measurements $\widetilde{B}^y$.

Let

$$|\text{aux}\rangle = \left(\bigotimes_x |\text{aux}_{A,x}\rangle \right) \otimes \left(\bigotimes_y |\text{aux}_{B,y}\rangle \right), \quad |\widehat{\psi}\rangle = |\psi\rangle \otimes |\text{aux}\rangle,$$

and let $\widehat{A}_a^x$ and $\widehat{B}_b^y$ be obtained by letting $\widetilde{A}_a^x$ or $\widetilde{B}_b^y$ act on the relevant auxiliary factor and the identity act on every other auxiliary factor. Because the auxiliary state is a product state and the two dilations act on disjoint tensor factors, the correlation for fixed (x, y) reduces to the compression identity from Lemma 4.7 on the x - and y -registers. Hence

$$\langle \widehat{\psi} | \widehat{A}_a^x \otimes \widehat{B}_b^y | \widehat{\psi} \rangle = \langle \psi | A_a^x \otimes B_b^y | \psi \rangle.$$

□

Remark 4.2 (Lean questionwise Naimark interface). Lean currently proves the questionwise one-measurement interface obtained from Lemma 4.7. For every indexed family A^x and B^y , it produces local dilation data for each question and proves the corresponding single-outcome marginal-preservation identities. This is the interface used downstream in the formal development. It is not the full tensor-product correlation statement of Theorem 4.1, whose simultaneous bipartite assembly remains recorded in `docs/paper-gaps/naimark.tex`.

Theorem 4.3 (Orthogonalization lemma). *Let $|\psi\rangle$ be a permutation-invariant state, and let $A = \{A_a\}$ be a sub-measurement with strong self-consistency*

$$\sum_a \langle \psi | A_a \otimes A_a | \psi \rangle \geq \sum_a \langle \psi | A_a \otimes I | \psi \rangle - \zeta.$$

Then there exists a projective sub-measurement $P = \{P_a\}$ such that

$$A_a \otimes I \approx_{100\zeta^{1/4}} P_a \otimes I.$$

Remark 4.4 (Lean auxiliary declarations: residual domination). The following are Lean auxiliary declarations, not hypotheses in Theorem 4.3. Let $A = \{A_a\}_{a \in \mathcal{A}}$ be a sub-measurement, and complete it to $\widehat{A}$ on $\mathcal{A} \cup \{\perp\}$ by setting $\widehat{A}_\perp = I - A_{\text{tot}}$. Let $\widehat{P}$ be a projective sub-measurement on the completed outcome set. If the completed projective family puts at least the original residual mass on the fresh outcome,

$$\widehat{A}_\perp \leq \widehat{P}_\perp,$$

then the restriction $P_a = \widehat{P}_a$, $a \in \mathcal{A}$, satisfies

$$P_{\text{tot}} \leq A_{\text{tot}}.$$

Consequently, for any bipartite state ψ ,

$$\langle \psi, I \otimes P_{\text{tot}} \rangle \leq \langle \psi, I \otimes A_{\text{tot}} \rangle.$$

This isolates the order-theoretic invariant needed to pass from the option-completed orthonormalization output to the monotone-total route used in the self-improvement theorem. The invariant is recorded as a separate order-theoretic fact; it is not added as an orthonormalization hypothesis.

Remark 4.5 (Lean right-register completion helpers). The inductive-step application of the orthogonalization lemma uses completion on both tensor factors. The formal development records the analytic completion statement on the left register; the shared preliminary helper compares right- and left-tensor placements at the level of the underlying state-dependent-distance form. Its submeasurement and indexed lemmas transport state-dependent distance to the right register under the permutation-invariance hypothesis on the state. The remaining two declarations combine this transport with the orthonormalize-and-complete output so that the line-156 projectivization transition can later be produced from the two completed measurements without restating the tensor-factor bookkeeping.

Remark 4.6 (Lean line-169 projectivization match-mass declarations). Lean exposes the quantitative obstruction at the projectivization transition used in the inductive step. The existing line-156 transition gives only the generic consequence of Proposition 3.20: from $G^A \simeq_{\zeta_1} G^B$ and $G^A \approx_{\zeta_2} Q^A$ one obtains $Q^A \simeq_{\zeta_1 + \sqrt{\zeta_2}} G^B$, and similarly on the other tensor factor. The declarations `leftConsistency_with_triangleSub_loss` and `rightConsistency_with_triangleSub_loss` are checked lemmas for this direct square-root-loss route.

Lean also proves a more local repaired transport that keeps the comparison at the pre-completion stage. If P^A is the orthonormalized projective sub-measurement with $G^A \approx_\epsilon P^A$, then Proposition 3.24 bounds the diagonal match-mass loss by $\sqrt{\epsilon}$, and canonical completion can only add nonnegative diagonal match mass. Thus the declarations `leftConsistency_of_completion_and_sdd` and `rightConsistency_of_completion_and_sdd` prove $Q^A \simeq_{\zeta_1 + \sqrt{\epsilon}} G^B$ and its Bob-side mirror from the pre-projective ζ_1 consistency plus the pre-completion $\approx_\epsilon$ comparison. In particular, substituting the checked orthonormalization error $\epsilon = 100\zeta_1^{1/4}$ yields the repaired line-169 bound

$\zeta_1 + 10\zeta_1^{1/8}$ recorded by `leftConsistency_with_orthonormalization_loss` and its mirror. This is much sharper than the direct completion-level $\zeta_1 + \sqrt{\zeta_2}$ route, but it is still not the paper's exact ζ_1 .

The paper-tight line-169 estimate at exactly ζ_1 therefore requires more than state-dependent-distance closeness. Lean records the missing construction-level invariant as `ProjectivizationMatchMassMonotonicity`, replacing G^A by the completed projective measurement Q^A , and role-reversing on Bob's side, must not decrease the diagonal match mass against the opposite pre-projective measurement. Its two consistency theorems prove that this primitive match-mass monotonicity is sufficient to recover the exact line-169 `ConsRel` links. Lean also proves that the canonical completion step itself can only increase this match mass: once the repaired projective sub-measurements preserve the relevant diagonal match mass, completing them at a distinguished outcome produces the required invariant for the completed projective measurements. The remaining upstream projectivization work is therefore the corresponding preservation inequality for the orthonormalization repair output. The pointwise-domination lemma records the degenerate no-change situation in which the source measurement is already dominated by the projective submeasurement. The former expectation-preservation structures have been removed: they merely restated the required match-mass inequality and were not produced by the Section 5 construction. The former QXP-specialized target has likewise been removed, because the present Section 5 API supplies state-dependent distance estimates and QXP algebra, but not a signed monotonicity statement against an arbitrary partner measurement.

4.0.1 Naimark dilation

Lemma 4.7. *Let $A = \{A_a\}$ be a sub-measurement with k distinct outcomes $a \in \mathcal{A}$, and let $|\text{aux}\rangle \in \mathbb{C}^{k+1}$ be any state. Then there exists a projective sub-measurement $\widehat{A} = \{\widehat{A}_a\}$ such that for each outcome a ,*

$$(I \otimes \langle \text{aux} |) \cdot \widehat{A}_a \cdot (I \otimes | \text{aux} \rangle) = A_a.$$

Proof. Consider an orthonormal basis for $\mathbb{C}^{k+1}$ consisting of vectors $|a\rangle$ for $a \in \mathcal{A}$ and the vector $|\perp\rangle$. Let U be any unitary such that for each vector $|\phi\rangle$,

$$U \cdot (|\phi\rangle \otimes |\text{aux}\rangle) = \sum_{a \in \mathcal{A}} ((A_a)^{1/2} |\phi\rangle) \otimes |a\rangle + ((I - A)^{1/2} |\phi\rangle) \otimes |\perp\rangle,$$

where $A = \sum_a A_a$. Then

$$U \cdot (I \otimes |\text{aux}\rangle) = \sum_{a \in \mathcal{A}} (A_a)^{1/2} \otimes |a\rangle + (I - A)^{1/2} \otimes |\perp\rangle,$$

so for any $a \in \mathcal{A}$,

$$(I \otimes \langle a |) \cdot U \cdot (I \otimes |\text{aux}\rangle) = (A_a)^{1/2}.$$

Define

$$\widehat{A}_a = U^\dagger \cdot (I \otimes |a\rangle \langle a|) \cdot U.$$

Each $\widehat{A}_a$ is a projector, and

$$\begin{aligned} (I \otimes \langle \text{aux} |) \cdot \widehat{A}_a \cdot (I \otimes | \text{aux} \rangle) &= (I \otimes \langle \text{aux} |) \cdot U^\dagger \cdot (I \otimes |a\rangle \langle a|) \cdot U \cdot (I \otimes | \text{aux} \rangle) \\ &= (A_a)^{1/2} \cdot (A_a)^{1/2} = A_a. \end{aligned}$$

□

Remark 4.8 (Naimark dilation does not preserve $\approx_\delta$). Let $|\psi\rangle$ be any state in $\mathbb{C}^d \otimes \mathbb{C}^d$, and consider the two-outcome measurements $A = \{A_0, A_1\}$ and $B = \{B_0, B_1\}$ with

$$A_0 = A_1 = B_0 = B_1 = \frac{1}{2}I.$$

For each outcome $a \in \{0, 1\}$, the operators $A_a \otimes I$ and $I \otimes B_a$ are both equal to $\frac{1}{2}I \otimes I$, so $A_a \otimes I \approx_0 I \otimes B_a$ on the state $|\psi\rangle$.

Now apply the Naimark construction with a qubit auxiliary space on each side and auxiliary state $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$. Since A and B are genuine measurements rather than sub-measurements, the extra $|\perp\rangle$ direction from 4.7 is unnecessary in this example. The unitary from 4.7 can then be taken to be the identity, so the dilated measurements are

$$\widehat{A}_a = I \otimes |a\rangle\langle a|, \quad \widehat{B}_a = I \otimes |a\rangle\langle a|.$$

If $|\widehat{\psi}\rangle = |\psi\rangle \otimes |+\rangle \otimes |+\rangle$, then a direct calculation gives

$$\sum_{a \in \{0,1\}} \|(\widehat{A}_a \otimes I - I \otimes \widehat{B}_a)|\widehat{\psi}\rangle\|^2 = 1.$$

Thus on the state $|\widehat{\psi}\rangle$ the dilated family satisfies $\widehat{A}_a \otimes I \approx_\delta I \otimes \widehat{B}_a$ only for $\delta \geq 1$, even though the original family satisfied the same relation with $\delta = 0$. This is why we use Theorem 4.3 rather than simply applying Theorem 4.1: Naimark preserves the outcome probabilities, but it need not preserve the state-dependent $\approx_\delta$ estimates needed later.

4.0.2 Orthogonalization lemma

Lemma 4.9 (Orthogonalization lemma for measurements). *Let $|\psi\rangle$ be a state, and let $A = \{A_a\}$ and $B = \{B_a\}$ be measurements such that*

$$A_a \otimes I \simeq_\zeta I \otimes B_a$$

on state $|\psi\rangle$. Here $|\psi\rangle$ is not assumed to be permutation-invariant. Then there exists a projective sub-measurement $P = \{P_a\}$ such that

$$A_a \otimes I \approx_{84\zeta^{1/4}} P_a \otimes I.$$

Lemma 4.10 (Formalized orthogonalization envelope for measurements). *This same-space corollary weakens Lemma 4.9 to the public orthonormalization envelope $100\zeta^{1/4}$. Thus, if $A = \{A_a\}$ and $B = \{B_a\}$ are measurements on the same finite-dimensional space satisfying*

$$A_a \otimes I \simeq_\zeta I \otimes B_a$$

on a normalized bipartite state $|\psi\rangle$, then there exists a projective sub-measurement $P = \{P_a\}$ such that

$$A_a \otimes I \approx_{100\zeta^{1/4}} P_a \otimes I.$$

This is the corrected Lean theorem proved from the Section 5 projectivization repair construction. The sharper heterogeneous source statement $84\zeta^{1/4}$ is now formalized as Lemma 4.9.

Remark 4.11 (Lean auxiliary declarations for the orthogonalization lemma). The Lean declaration `MIPStarRE.LDT.MakingMeasurementsProjective.orthonormalizationMainLemma` proves Lemma 4.9 without adding spectral-truncation or repair hypotheses. The proof first converts the cross-consistency hypothesis into the source almost-projective estimate for the measurement $A_a \otimes I$, and then applies the locality-preserving $Q/X/\widehat{X}/P$ construction in its heterogeneous left-register form. Lemma 4.10 remains as the same-space corollary with the weaker public $100\zeta^{1/4}$ envelope.

Lemma 4.12 (Locality-preserving projectivization repair). *Paper origin: [references/ldt-paper/orthonormalization](#) lines 534–860 (rank reduction and the Q -side setup, including 4.29 and 4.30) and 862–1194 (the $X/\widehat{X}/P$ algebra: 4.31, 4.34, 4.35, 4.37, 4.38, 4.39, plus the final triangle-inequality calculation producing the $84\zeta^{1/4}$ bound). Together these are the late repair stage of the orthogonalization-lemma proof.*

Given a normalized bipartite state $|\psi\rangle$, a measurement $A = \{A_a\}$, and the source almost-projective estimate for the left-lifted family $\{A_a \otimes I\}$, there exists a locality-preserving projective sub-measurement with error $84\zeta^{1/4}$.

More precisely, if $|\psi\rangle$ is a normalized bipartite state, if $A = \{A_a\}$ is a measurement, and if

$$\sum_a \langle \psi | ((A_a \otimes I) - (A_a \otimes I)^2) | \psi \rangle \leq \zeta,$$

then there exists a projective sub-measurement $P = \{P_a\}$ on the underlying space such that

$$A_a \otimes I \approx_{84\zeta^{1/4}} P_a \otimes I.$$

The locality-preserving form – output $P_a \otimes I$ rather than an arbitrary lifted family – is recorded in Remark 4.11 and is the form required by the unconditional orthonormalization theorem (4.3).

Proof. Pass to the left marginal state, apply the rank-reduction and $Q/X/\widehat{X}/P$ construction there, and transport the resulting $\approx_{84\zeta^{1/4}}$ estimate back to left lifts via the left-marginal expectation formula. $\square$

Remark 4.13 (Lean implementation name for projectivization repair). The declaration `leftLiftedProjectivization` is the paper-labelled name for Lemma 4.12. The older declaration `leftLiftedProjectivizationRepairProducer` has the same public mathematical statement and is retained for compatibility with existing Section 5 declarations. The word “producer” here refers to the construction theorem that produces the projective submeasurement; it is not a hypothesis added to the paper lemma.

Remark 4.14 (Lean conditional declaration: local orthogonalization). This is the local Lean declaration recorded in Remark 4.11. Let $|\psi\rangle$ be a unit vector, let $A = \{A_a\}$ be a measurement, and let $0 \leq \zeta \leq 1$. Assume bipartite strong self-consistency

$$\sum_a \langle \psi | A_a \otimes A_a | \psi \rangle \geq \sum_a \langle \psi | A_a \otimes I | \psi \rangle - \zeta,$$

together with a spectral-truncation witness for the left-lifted family $\{A_a \otimes I\}$ and a locality-preserving repair witness producing projective families of the form $\widehat{P}_a = P_a \otimes I$. Then there exists a projective sub-measurement $P = \{P_a\}$ on the same space as A such that

$$A_a \otimes I \approx_{84\zeta^{1/4}} P_a \otimes I.$$

Since A is a measurement, $\sum_a A_a = I$, so bipartite strong self-consistency of A with itself is equivalent to bipartite consistency of A with itself. Feed this consistency hypothesis, together with the spectral-truncation and locality-preserving repair witnesses, into the measurement variant of 4.9 with $B = A$. The repair witness chooses the repaired family of the form $P_a \otimes I$, so the conclusion descends to a local projective sub-measurement $P = \{P_a\}$ on the space of A with error bound $84\zeta^{1/4}$.

Remark 4.15 (Lean conditional declaration: measurement-level orthogonalization). This is the measurement-level Lean declaration recorded in Remark 4.11. Let $|\psi\rangle$ be a unit vector, let $A = \{A_a\}$ be a measurement, and let $0 \leq \zeta \leq 1$. Assume bipartite strong self-consistency of A

together with the spectral-truncation and locality-preserving repair witnesses for the left-lifted family $\{A_a \otimes I\}$. Then there exists a projective sub-measurement $P = \{P_a\}$ such that

$$A_a \otimes I \approx_{100\zeta^{1/4}} P_a \otimes I.$$

Using the bound stated in Remark 4.14, weaken the error using $84 \leq 100$ and $\zeta^{1/4} \geq 0$. No completion step is needed because A is already a measurement.

Remark 4.16 (Lean completion-route orthonormalization).

The paper theorem 4.3 is proved in Lean with the $100\zeta^{1/4}$ constant. The declaration checked here is the completion-route theorem with the weaker bound $120\zeta^{1/4}$, recorded in `orthonormalizationCompletionRoute`. See `docs/paper-gaps/issue-1032-orthonormalization-constant.tex`.

Let $A = \{A_a\}$ be a sub-measurement with outcomes in $\mathcal{A}$. Write $A = \sum_a A_a$. The strong self-consistency assumption implies

$$\langle \psi | A \otimes A | \psi \rangle \geq \langle \psi | A \otimes I | \psi \rangle - \zeta,$$

so

$$\langle \psi | A \otimes (I - A) | \psi \rangle \leq \zeta.$$

Rearranging once more gives

$$\langle \psi | (I - A) \otimes (I - A) | \psi \rangle \geq \langle \psi | (I - A) \otimes I | \psi \rangle - \zeta.$$

Complete A to the measurement $\widehat{A} = \{\widehat{A}_a\}_{a \in \widehat{\mathcal{A}}}$ on $\widehat{\mathcal{A}} = \mathcal{A} \cup \{\perp\}$ by setting $\widehat{A}_a = A_a$ for $a \in \mathcal{A}$ and $\widehat{A}_\perp = I - A$. Adding the two preceding inequalities shows that $\widehat{A}$ is 2ζ -strongly self-consistent. By 3.35, this gives the source almost-projective estimate

$$\sum_a \langle \psi | ((\widehat{A}_a \otimes I) - (\widehat{A}_a \otimes I)^2) | \psi \rangle \leq 4\zeta.$$

Apply 4.12 to obtain a projective sub-measurement $\widehat{P}$ such that

$$\widehat{A}_a \otimes I \approx_{84(4\zeta)^{1/4}} \widehat{P}_a \otimes I.$$

Discard the extra outcome and set $P_a = \widehat{P}_a$ for $a \in \mathcal{A}$. Then

$$A_a \otimes I \approx_{84(4\zeta)^{1/4}} P_a \otimes I,$$

and $84 \cdot 4^{1/4} \leq 120$, so the completion-route bound follows.

Proof of 4.9. The statement is trivial when $\zeta > 1/4$, so assume

$$\zeta \leq 1/4. \tag{4.1}$$

By 3.18,

$$\sum_a \langle \psi | A_a \otimes B_a | \psi \rangle \geq 1 - \zeta.$$

Apply Cauchy-Schwarz to

$$\sum_a \langle \psi | (A_a \otimes I) \cdot (I \otimes B_a) | \psi \rangle$$

to obtain

$$\sum_a \langle \psi | (A_a)^2 \otimes I | \psi \rangle \geq (1 - \zeta)^2 \geq 1 - 2\zeta.$$

Because A is a measurement, this rewrites as

$$\sum_a \langle \psi | (A_a - (A_a)^2) \otimes I | \psi \rangle \leq 2\zeta. \tag{4.2}$$

Lemma 4.17. *For any $x \in [0, 1]$,*

$$(x - \text{trunc}_\delta(x))^2 \leq \frac{1}{\delta} \cdot (x - x^2).$$

Proof. If $x \geq 1 - \delta$, then $\text{trunc}_\delta(x) = 1$. Since $\delta \leq 1/2$ below, one also has $x \geq \delta$, so

$$(x - \text{trunc}_\delta(x))^2 = (1 - x)^2 \leq (1 - x) \frac{x}{\delta} = \frac{1}{\delta} (x - x^2).$$

If $x < 1 - \delta$, then $\text{trunc}_\delta(x) = 0$, and similarly

$$(x - \text{trunc}_\delta(x))^2 = x^2 \leq x \frac{1 - x}{\delta} = \frac{1}{\delta} (x - x^2).$$

□

Lemma 4.18 (Rounding to projectors). *There exists a set of projective matrices $\{R_a\}$ such that*

$$A_a \otimes I \approx_{2\sqrt{\zeta}} R_a \otimes I.$$

and

$$R := \sum_a R_a \leq (1 + 2\sqrt{\zeta}) \cdot I.$$

Proof. Write the eigendecomposition of A_a as

$$A_a = \sum_i \lambda_{a,i} |u_{a,i}\rangle \langle u_{a,i}|.$$

Choose

$$0 < \delta \leq 1/2, \tag{4.3}$$

define $\text{trunc}_\delta : [0, 1] \rightarrow \{0, 1\}$ by $\text{trunc}_\delta(x) = 1$ when $x \geq 1 - \delta$ and 0 otherwise, and set

$$R_a = \text{trunc}_\delta(A_a) = \sum_i \text{trunc}_\delta(\lambda_{a,i}) |u_{a,i}\rangle \langle u_{a,i}|.$$

By 4.17,

$$(A_a - R_a)^2 \leq \frac{1}{\delta} (A_a - (A_a)^2).$$

Summing expectations and using 4.2 gives

$$\sum_a \|(A_a - R_a) \otimes I |\psi\rangle\|^2 \leq \frac{2\zeta}{\delta},$$

so $A_a \otimes I \approx_{2\zeta/\delta} R_a \otimes I$.

For every $x \in [0, 1]$, one has $\text{trunc}_\delta(x) \leq x/(1 - \delta)$, hence

$$R_a \leq \frac{1}{1 - \delta} A_a.$$

Summing over a and using that A is a measurement,

$$R = \sum_a R_a \leq \frac{1}{1 - \delta} I \leq (1 + 2\delta) I,$$

where the last estimate uses $\delta \leq 1/2$. Setting $\delta = \sqrt{\zeta}$ gives the claim, and 4.1 ensures $\delta \leq 1/2$. □

Remark 4.19 (Lean auxiliary declarations for rounding to projectors). The source-facing Lean declarations for Lemma 4.18 are the proposition `projectiveNonMeasurement` and the construction theorem `projectiveNonMeasurement_of_sourceAlmostProjective_full`. The declarations listed here are the endpoint and regime splits, the `AlmostProjMeasStatement` witness forms, and the spectral-truncation input conversions used by later Section 5 arguments. They are not additional hypotheses in the paper statement of Lemma 4.18.

Lemma 4.20 (Combinatorial averaging for the $r > d$ branch). *Let $f : \alpha \rightarrow \mathbb{R}$ be a function on a finite index set α with $|\alpha| = r$, and let $L \subseteq \alpha$ with $|L| = d$ and $d < r$. Write $S = L^c$ and assume $f(s) \leq f(l)$ for every $s \in S$ and $l \in L$. Then the pairwise double-counting inequality*

$$|L| \sum_{s \in S} f(s) \leq |S| \sum_{l \in L} f(l)$$

holds, and hence also

$$r \sum_{s \in S} f(s) \leq |S| \sum_{x \in \alpha} f(x).$$

If in addition $r \leq (1 + 2\sqrt{\zeta})d$, $\sum_{x \in \alpha} f(x) \leq 1 + 2\sqrt{\zeta}$, and $0 \leq \zeta \leq 1/4$, then

$$\sum_{s \in S} f(s) \leq 4\sqrt{\zeta}.$$

Proof. Choose L to be a set of d indices on which the sum of f is maximal. If some $s \in S$ and $l \in L$ satisfied $f(l) < f(s)$, replacing l by s would increase this sum, so every small index is bounded by every large index. Summing the inequalities $f(s) \leq f(l)$ over all pairs $(s, l) \in S \times L$ gives

$$|L| \sum_{s \in S} f(s) \leq |S| \sum_{l \in L} f(l).$$

Since α is the disjoint union of L and S , adding $|S| \sum_{s \in S} f(s)$ to both sides yields

$$r \sum_{s \in S} f(s) \leq |S| \sum_{x \in \alpha} f(x).$$

The hypotheses $|L| = d$ and $r \leq (1 + 2\sqrt{\zeta})d$ imply $|S|/r \leq 2\sqrt{\zeta}$. Dividing by r and using $\sum_{x \in \alpha} f(x) \leq 1 + 2\sqrt{\zeta}$ gives

$$\sum_{s \in S} f(s) \leq 2\sqrt{\zeta}(1 + 2\sqrt{\zeta}).$$

Finally, $\zeta \leq 1/4$ implies $\sqrt{\zeta} \leq 1/2$, and hence $2\sqrt{\zeta}(1 + 2\sqrt{\zeta}) \leq 4\sqrt{\zeta}$. □

Lemma 4.21 (Rank reduction). *There exists a set of projection matrices $\{Q_a\}$ such that*

$$A_a \otimes I \approx_{12\sqrt{\zeta}} Q_a \otimes I.$$

and

$$Q := \sum_a Q_a \leq (1 + 2\sqrt{\zeta}) \cdot I.$$

Furthermore,

$$\sum_a \text{rank}(Q_a) \leq d.$$

Proof. Let $\{R_a\}$ be given by 4.18. Write $r_a = \text{rank}(R_a)$ and $r = \sum_a r_a$. If $r \leq d$, take $Q_a = R_a$. Otherwise,

$$r = \sum_a r_a = \sum_a \text{Tr}(R_a) = \text{Tr}(R) \leq (1 + 2\sqrt{\zeta})d. \quad (4.4)$$

Let $|v_{a,1}\rangle, \dots, |v_{a,r_a}\rangle$ be an orthonormal basis for the range of R_a , so

$$R_a = \sum_{i=1}^{r_a} |v_{a,i}\rangle\langle v_{a,i}|.$$

Define the overlaps

$$o_{a,i} = \langle \psi | (|v_{a,i}\rangle\langle v_{a,i}| \otimes I) | \psi \rangle.$$

Their total satisfies

$$\sum_a \sum_{i=1}^{r_a} o_{a,i} = \langle \psi | R \otimes I | \psi \rangle \leq 1 + 2\sqrt{\zeta}. \quad (4.5)$$

Let **Large** be the set of the d largest overlaps $o_{a,i}$, breaking ties arbitrarily, and let **Small** be the complement. Then

$$|\text{Small}| = r - d \leq 2\sqrt{\zeta}d \leq 2\sqrt{\zeta}r, \quad (4.6)$$

by 4.4. Averaging the overlaps over the r indices gives

$$\sum_{(a,i) \in \text{Small}} o_{a,i} \leq \frac{|\text{Small}|}{r} \sum_{a,i} o_{a,i} \leq 4\sqrt{\zeta}, \quad (4.7)$$

using 4.6, 4.5, and 4.1.

For each a , let $\text{Large}_a = \{i : (a,i) \in \text{Large}\}$ and define

$$Q_a = \sum_{i \in \text{Large}_a} |v_{a,i}\rangle\langle v_{a,i}|.$$

Then $\sum_a \text{rank}(Q_a) = |\text{Large}| \leq d$, and $Q_a \leq R_a$ for every a , so

$$Q = \sum_a Q_a \leq \sum_a R_a = R \leq (1 + 2\sqrt{\zeta})I.$$

Also,

$$R_a - Q_a = \sum_{i \in \text{Small}_a} |v_{a,i}\rangle\langle v_{a,i}|$$

is a projector, hence

$$\sum_a \|(R_a - Q_a) \otimes I | \psi \rangle\|^2 = \sum_{(a,i) \in \text{Small}} o_{a,i} \leq 4\sqrt{\zeta}$$

by 4.7. Therefore $R_a \otimes I \approx_{4\sqrt{\zeta}} Q_a \otimes I$. Combining this with 4.18 and the triangle inequality for $\approx_\delta$ gives

$$A_a \otimes I \approx_{12\sqrt{\zeta}} Q_a \otimes I.$$

□

Henceforth let $Q = \{Q_a\}$ be the family given by 4.21.

Lemma 4.22 (Completeness of Q).

$$\langle \psi | Q \otimes I | \psi \rangle \geq 1 - 11\zeta^{1/4}.$$

Proof. Since each Q_a is projective,

$$\langle \psi | Q \otimes I | \psi \rangle = \sum_a \langle \psi | (Q_a)^2 \otimes I | \psi \rangle \approx_{5\zeta^{1/4}} \sum_a \langle \psi | (Q_a \cdot A_a) \otimes I | \psi \rangle. \quad (4.8)$$

By Cauchy–Schwarz,

$$\sum_a \langle \psi | (Q_a)^2 \otimes I | \psi \rangle \leq 1 + 2\sqrt{\zeta} \quad \text{and} \quad \sum_a \langle \psi | (Q_a - A_a)^2 \otimes I | \psi \rangle \leq 12\sqrt{\zeta}$$

from 4.21. A second Cauchy–Schwarz estimate yields

$$\sum_a \langle \psi | (Q_a \cdot A_a) \otimes I | \psi \rangle \approx_{4\zeta^{1/4}} \sum_a \langle \psi | (A_a)^2 \otimes I | \psi \rangle. \quad (4.9)$$

Combining 4.8 and 4.9 with 4.2 gives

$$\langle \psi | Q \otimes I | \psi \rangle \geq \sum_a \langle \psi | A_a \otimes I | \psi \rangle - 2\zeta - 9\zeta^{1/4} = 1 - 11\zeta^{1/4},$$

since $\zeta \leq \zeta^{1/4}$ under 4.1. □

Lemma 4.23 (Completeness of $\sqrt{Q}$).

$$\langle \psi | \sqrt{Q} \otimes I | \psi \rangle \geq 1 - 12\zeta^{1/4}.$$

Proof. Diagonalize $Q = \sum_i \nu_i |u_i\rangle\langle u_i|$. Because $Q \leq (1 + 2\sqrt{\zeta})I$ by 4.21, each ν_i is at most $1 + 2\sqrt{\zeta}$. Therefore

$$\sqrt{Q} = \sum_i \sqrt{\nu_i} |u_i\rangle\langle u_i| \geq \frac{1}{\sqrt{1 + 2\sqrt{\zeta}}} Q.$$

The scalar estimate

$$\frac{1}{\sqrt{1 + 2\sqrt{\zeta}}} \geq 1 - \sqrt{\zeta}$$

gives $\sqrt{Q} \geq (1 - \sqrt{\zeta})Q$. Hence 4.22 implies

$$\langle \psi | \sqrt{Q} \otimes I | \psi \rangle \geq (1 - \sqrt{\zeta})(1 - 11\zeta^{1/4}) \geq 1 - 12\zeta^{1/4}. \quad \square$$

Lemma 4.24 (Q is almost projective).

$$\sum_a (Q_a \cdot Q \cdot Q_a - Q_a) \leq 4\sqrt{\zeta} \cdot I.$$

Proof. Since $Q \leq (1 + 2\sqrt{\zeta})I$ by 4.21,

$$\begin{aligned} \sum_a Q_a \cdot Q \cdot Q_a - \sum_a Q_a &\leq (1 + 2\sqrt{\zeta}) \sum_a Q_a \cdot I \cdot Q_a - \sum_a Q_a \\ &= (1 + 2\sqrt{\zeta})Q - Q \\ &= 2\sqrt{\zeta} Q \\ &\leq 2\sqrt{\zeta}(1 + 2\sqrt{\zeta})I \\ &\leq 4\sqrt{\zeta} I, \end{aligned}$$

using 4.1 in the last step. □

Definition 4.25 (Matrix decomposition of Q_a). For each a , let

$$m_a = \text{rank}(Q_a),$$

and let $|v_{a,1}\rangle, \dots, |v_{a,m_a}\rangle$ be an orthonormal basis for the range of Q_a , so that

$$Q_a = \sum_{i=1}^{m_a} |v_{a,i}\rangle \langle v_{a,i}|.$$

Let

$$m = \sum_a m_a,$$

and let $\{|a, i\rangle\}$ be an orthonormal basis of $\mathbb{C}^m$, indexed by all a and all $1 \leq i \leq m_a$. For each a , define

$$X_a = \sum_{i=1}^{m_a} |a, i\rangle \langle v_{a,i}|, \quad T_a = \sum_{i=1}^{m_a} |a, i\rangle \langle a, i|,$$

and define

$$X = \sum_a X_a = \sum_a \sum_{i=1}^{m_a} |a, i\rangle \langle v_{a,i}|. \quad (4.10)$$

Let $T = \{T_a\}$. Then T is a projective measurement on $\mathbb{C}^m$.

Definition 4.26 (SVD of X). The paper presents this step by choosing a singular value decomposition

$$X = U \cdot \Sigma_{m \times d} \cdot V^\dagger.$$

The formalization has two closely related auxiliary declarations. The reduced datum records the downstream consequences of the choice: the matrices X and $\widehat{X}$, the identities $X^\dagger X = Q$, $\widehat{X} \widehat{X}^\dagger = I$, and $X^\dagger \widehat{X} = \sqrt{Q}$, together with the $Q_a = X^\dagger T_a X$ restatement. The rectangular-SVD lemmas take explicit matrices $U, V, \Sigma, I_{m \times d}$ and derive the two stored $\widehat{X}$ identities from the matrix calculation below.

Remark 4.27. If the family $\{Q_a\}$ were a projective measurement, then the vectors $|v_{a,i}\rangle$, over all a and all $1 \leq i \leq m_a$, would be orthonormal. Then (4.10) would already be a singular value decomposition of X , with $\Sigma = I_{m \times d}$. Equivalently,

$$X = U \cdot I_{m \times d} \cdot V^\dagger.$$

This motivates replacing Σ by I in the definition of $\widehat{X}$ and P .

Definition 4.28 (Definition of P). The paper defines

$$\widehat{X} = U \cdot I_{m \times d} \cdot V^\dagger.$$

We take $\widehat{X}$ to be the reduced matrix supplied by the stored datum. For each a , define

$$\widehat{X}_a = T_a \cdot \widehat{X}, \quad P_a = \widehat{X}_a^\dagger \cdot \widehat{X}_a.$$

Lemma 4.29. For each a ,

$$X_a = T_a \cdot X.$$

Proof. Multiply the definition of X by T_a on the left and use that T_a selects exactly the basis vectors $|a, i\rangle$:

$$T_a \cdot X = \left(\sum_{i=1}^{m_a} |a, i\rangle \langle a, i| \right) \left(\sum_b \sum_{j=1}^{m_b} |b, j\rangle \langle v_{b,j}| \right) = \sum_{i=1}^{m_a} |a, i\rangle \langle v_{a,i}| = X_a.$$

□

Lemma 4.30 (Q_a restated). *For each a ,*

$$Q_a = X_a^\dagger \cdot X_a = X^\dagger \cdot T_a \cdot X = X_a^\dagger \cdot X.$$

Proof. The identity $X_a^\dagger X_a = Q_a$ follows by expanding the basis sums. The remaining equalities come from 4.29 and the projectivity of T :

$$X_a^\dagger \cdot X_a = (X^\dagger \cdot T_a)(T_a \cdot X) = X^\dagger \cdot T_a \cdot X = X_a^\dagger \cdot X.$$

□

Lemma 4.31 (X squared).

$$X^\dagger \cdot X = Q.$$

Proof. Use that T is a measurement and then 4.30:

$$X^\dagger \cdot X = \sum_a X^\dagger \cdot T_a \cdot X = \sum_a Q_a = Q.$$

The additional paper identities involving U, V, Σ are the usual SVD computations. The Lean development also records the spectral calculation which will identify the mixed positive-Gram-image product with $\sqrt{Q}$: the positive eigenvalues give the nonzero columns, while the zero eigenspace has zero coefficient in the square-root expansion and is killed by X . The conjugate right-eigenvector rows over the positive spectrum are also orthonormal. Equivalently, any row-space vector orthogonal to the positive Gram images is killed by $X^\dagger$. □

Lemma 4.32. *For each a ,*

$$X_a^\dagger \cdot (X \cdot X^\dagger - I_{m \times m})^2 \cdot X_a = Q_a \cdot Q \cdot Q_a - Q_a.$$

Proof. Expand the square and apply 4.31 and 4.30 term by term:

$$X_a^\dagger (X \cdot X^\dagger \cdot X \cdot X^\dagger) X_a = Q_a \cdot Q \cdot Q_a,$$

while

$$X_a^\dagger (X \cdot X^\dagger) X_a = (X_a^\dagger \cdot X)(X^\dagger \cdot X_a) = Q_a \cdot Q_a = Q_a,$$

and $X_a^\dagger I_{m \times m} X_a = X_a^\dagger X_a = Q_a$. Substituting these identities into

$$(X \cdot X^\dagger - I_{m \times m})^2 = X \cdot X^\dagger \cdot X \cdot X^\dagger - 2X \cdot X^\dagger + I_{m \times m}$$

gives the claim. □

Lemma 4.33 (P_a restated). *For each a ,*

$$P_a = \widehat{X}^\dagger \cdot T_a \cdot \widehat{X} = \widehat{X}_a^\dagger \cdot \widehat{X}.$$

Proof. Since T is projective,

$$P_a = \widehat{X}_a^\dagger \cdot \widehat{X}_a = (\widehat{X}^\dagger \cdot T_a)(T_a \cdot \widehat{X}) = \widehat{X}^\dagger \cdot T_a \cdot \widehat{X} = \widehat{X}_a^\dagger \cdot \widehat{X}.$$

□

Lemma 4.34 ($\widehat{X}$ squared).

$$\widehat{X} \cdot \widehat{X}^\dagger = I_{m \times m}.$$

Proof. The paper proves this by multiplying out $\widehat{X} = U \cdot I_{m \times d} \cdot V^\dagger$ and using the unitarity of U and V . In Lean, the dimension hypothesis $m \leq d$ already gives a rectangular matrix with orthonormal rows on the sigma auxiliary space; the chosen matrix `sigmaFinXHatCoisometry` and its specification record this existence result as a hypothesis for the future construction. The formal lemma `orthonormal_normalized_image_of_adjoint_comp_eigenvectors` records the singular-vector normalization calculation, while `normalized_matrix_image_rows_mul_conjTranspose` combines the corresponding row-coisometry identity, and `positive_gram_spectrum_image_rows_transpose_mixed` records the adjoint action $X^\dagger(\lambda^{-1/2}Xv) = \lambda^{1/2}v$ on the positive spectral part of the Gram operator. The formal lemma `exists_unitary_with_positive_gram_spectrum_rows` records the next finite-dimensional step: after choosing distinct row positions for the positive spectrum, these normalized image rows extend to a square unitary matrix on the row space. The companion lemma `exists_rectangular_coisometry_extending_orthonormal_rows` records the analogous rectangular completion for prescribed right rows when the dimension bound permits it. The formal lemma `rectangularSvd_xHat_coisometry` records the corresponding matrix calculation. The unconditional positive-Gram route is `exists_xHat_of_sigmaFinRangeEmbedding_positiveGram`, which constructs $\widehat{X}$ from the rank-reduction witness and proves $\widehat{X}\widehat{X}^\dagger = I$ together with the mixed square-root identity below. □

Lemma 4.35 (X times $\widehat{X}$).

$$X \cdot \widehat{X}^\dagger = U \cdot \Sigma_{m \times m} \cdot U^\dagger, \quad X^\dagger \cdot \widehat{X} = \sqrt{Q}.$$

Proof. The paper obtains this from the SVD formula for X and the definition of $\widehat{X}$. In Lean, the unconditional QXP data first proves that $X\widehat{X}^\dagger$ is Hermitian and positive. The formal lemma `x_mul_xHat_adjoint_spectral_theorem` then applies Mathlib's spectral theorem for Hermitian matrices, giving the displayed $U\Sigma U^\dagger$ form with the Mathlib eigenvector unitary and the diagonal matrix of real eigenvalues. The auxiliary lemma `rectangularSvd_x_mul_xHat_conjTranspose_raw` records the same calculation under explicit rectangular-SVD data, where the middle factor $SI_{m \times d}^\dagger$ is the formal square matrix $\Sigma_{m \times m}$. The formal lemma `rectangularSvd_xHat_mixed_of_sqrtQ` records the second matrix calculation in the form whose square-root target is the operator Q supplied by the QXP construction. The formal lemma `rectangularSvd_middle_eq_sqrt_of_square` records the spectral step: a positive middle factor whose square is Q is the positive square root of Q . The reduced datum then records this mixed-product identity as a primitive consequence, which is exactly the part needed in 4.37 and 4.39. □

Remark 4.36 (Lean status for the QXP algebra). The Lean declarations linked in Lemmas 4.31, 4.34, and 4.35 expose the corresponding matrix identities as construction theorems. These declarations prove the identities from the Q -layer, the sigma-range realization, and the positive-Gram spectral construction, with the rectangular-SVD calculation retained as an auxiliary comparison. Thus the local $X/\widehat{X}/P$ algebra is formalized; the remaining chapter-level work is to compose this construction with the preceding orthonormalization step from the paper hypotheses.

Lemma 4.37 (Squared difference).

$$(X - \widehat{X}) \cdot (X - \widehat{X})^\dagger \leq (X \cdot X^\dagger - I_{m \times m})^2.$$

Proof. The paper diagonalizes the positive operator $X\widehat{X}^\dagger$ using the SVD notation. The formal proof avoids storing that SVD explicitly: set $Y = X\widehat{X}^\dagger$. The primitive identities imply that Y is hermitian, nonnegative, and satisfies $Y^2 = XX^\dagger$. Hence $(X - \widehat{X})(X - \widehat{X})^\dagger = (Y - I)^2$, and positivity of Y gives $(Y - I)^2 \leq (Y^2 - I)^2 = (XX^\dagger - I)^2$. $\square$

Lemma 4.38 (Projectivity of P). $P = \{P_a\}$ forms a projective sub-measurement.

Proof. For outcomes a, b ,

$$\begin{aligned} P_a \cdot P_b &= (\widehat{X}^\dagger \cdot T_a \cdot \widehat{X})(\widehat{X}^\dagger \cdot T_b \cdot \widehat{X}) \\ &= \widehat{X}^\dagger \cdot T_a \cdot I_{m \times m} \cdot T_b \cdot \widehat{X} \\ &= \widehat{X}^\dagger \cdot T_a \cdot \widehat{X} \cdot \mathbf{1}[\![a = b]\!] \\ &= P_a \cdot \mathbf{1}[\![a = b]\!], \end{aligned}$$

where the second line uses 4.34, the third uses that T is projective, and the last uses 4.33. Summing over a shows $\sum_a P_a \leq I$, so P is a projective sub-measurement. The formalization also records the sharper identity

$$\sum_a P_a = \widehat{X}^\dagger \widehat{X},$$

and the resulting total-domination estimates used later in the monotone-total self-improvement route. $\square$

Lemma 4.39 (P is close to Q).

$$Q_a \otimes I \approx_{30\zeta^{1/4}} P_a \otimes I.$$

Proof. The formal proof follows this argument directly from the primitive X , $\widehat{X}$, and P identities; the $30\zeta^{1/4}$ estimate is derived here rather than stored as part of the reduced datum. The rectangular-SVD lemma combines the same estimate with the positive-square identification of the middle factor from 4.35. The positive-Gram sigma-space lemma also records the coisometry $XX^\dagger = I$ for the original sigma embedding when the projective Q -family is subnormalized; this is the construction-level hypothesis later used in the residual-domination argument for the fresh completed outcome. The quantity to bound is

$$\sum_a \langle \psi | (Q_a - P_a)^2 \otimes I | \psi \rangle = \langle \psi | Q \otimes I | \psi \rangle + \langle \psi | P \otimes I | \psi \rangle - \sum_a \langle \psi | Q_a P_a \otimes I | \psi \rangle - \sum_a \langle \psi | P_a Q_a \otimes I | \psi \rangle, \quad (4.11)$$

using the projectivity of Q and P . The first term is at most $1 + 2\sqrt{\zeta}$ by 4.21, and the second is at most 1 because P is a sub-measurement.

For the last two terms,

$$\sum_a \langle \psi | Q_a P_a \otimes I | \psi \rangle + \sum_a \langle \psi | P_a Q_a \otimes I | \psi \rangle = 2 \cdot \Re \left(\sum_a \langle \psi | Q_a P_a \otimes I | \psi \rangle \right). \quad (4.12)$$

By 4.30,

$$\sum_a \langle \psi | Q_a P_a \otimes I | \psi \rangle = \sum_a \langle \psi | (X_a^\dagger \cdot X \cdot P_a) \otimes I | \psi \rangle.$$

The main estimate is

$$\sum_a \langle \psi | (X_a^\dagger \cdot X \cdot P_a) \otimes I | \psi \rangle \approx_{2\zeta^{1/4}} \sum_a \langle \psi | (X_a^\dagger \cdot \widehat{X} \cdot P_a) \otimes I | \psi \rangle. \quad (4.13)$$

This follows by Cauchy–Schwarz:

$$\left| \sum_a \langle \psi | ((X_a^\dagger \cdot (X - \widehat{X})) \otimes I) \cdot (P_a \otimes I) | \psi \rangle \right|$$

is bounded by the square root of

$$\sum_a \langle \psi | (X_a^\dagger \cdot (X - \widehat{X})(X - \widehat{X})^\dagger \cdot X_a) \otimes I | \psi \rangle,$$

times the square root of $\sum_a \langle \psi | (P_a)^2 \otimes I | \psi \rangle \leq 1$. The first factor is at most $4\sqrt{\zeta}$ by 4.37, 4.32, and 4.24, so the total error is at most $2\zeta^{1/4}$.

The right-hand side of 4.13 is real, because

$$\sum_a \langle \psi | (X_a^\dagger \cdot \widehat{X} \cdot P_a) \otimes I | \psi \rangle = \langle \psi | (X^\dagger \cdot \widehat{X}) \otimes I | \psi \rangle = \langle \psi | \sqrt{Q} \otimes I | \psi \rangle,$$

where the first equality uses 4.29, 4.33, and 4.34 and that T is a measurement, and the second uses 4.35. Hence 4.23 and 4.13 imply

$$\Re \left(\sum_a \langle \psi | Q_a P_a \otimes I | \psi \rangle \right) \geq 1 - 14\zeta^{1/4}.$$

Substituting this into 4.12 and then into 4.11 gives

$$\sum_a \langle \psi | (Q_a - P_a)^2 \otimes I | \psi \rangle \leq (1 + 2\sqrt{\zeta}) + 1 - 2(1 - 14\zeta^{1/4}) = 2\sqrt{\zeta} + 28\zeta^{1/4} \leq 30\zeta^{1/4}.$$

□

Finally, 4.21 gives

$$A_a \otimes I \approx_{12\sqrt{\zeta}} Q_a \otimes I,$$

and 4.39 gives

$$Q_a \otimes I \approx_{30\zeta^{1/4}} P_a \otimes I.$$

The triangle inequality for $\approx_\delta$ yields

$$A_a \otimes I \approx_{84\zeta^{1/4}} P_a \otimes I.$$

□

Chapter 5

Expansion in the hypercube graph

5.1 The graph and its spectrum

Definition 5.1 (Rerandomizing one coordinate). For $u \in \mathbb{F}_q^m$, an index $i \in \{1, \dots, m\}$, and $x \in \mathbb{F}_q$, let

$$\text{rerand}_i(u, x) = u + xe_i.$$

This is the move obtained by rerandomizing the i -th coordinate of u by a uniform additive shift.

Definition 5.2 (Hypercube graph). The hypercube graph $C = (V, E)$ has vertex set $V = \mathbb{F}_q^m$, and an edge between u and v whenever they differ in at most one coordinate. A random edge $(u, v) \sim C$ is sampled by drawing $u \sim \mathbb{F}_q^m$, $i \in \{1, \dots, m\}$, and $x \in \mathbb{F}_q$ uniformly and then setting $v = \text{rerand}_i(u, x)$.

Definition 5.3 (Adjacency matrix and Laplacian). Let $M = q^m$. The normalized adjacency matrix of C is

$$K = \mathbb{E}_{(u,v) \sim C} |u\rangle\langle v|,$$

and the Laplacian is

$$L = \frac{1}{M}I - K.$$

Lemma 5.4 (Laplacian rewrite). *The Laplacian can be written as*

$$L = \frac{1}{2} \mathbb{E}_{(u,v) \sim C} (|u\rangle - |v\rangle)(\langle u| - \langle v|).$$

Proof. The two vertex marginals of a random edge are uniform on $\mathbb{F}_q^m$. Expanding the right-hand side yields $(1/M)I - K$. $\square$

Lemma 5.5 (Average of a nontrivial additive character). *For $\beta \in \mathbb{F}_q$,*

$$\mathbb{E}_{x \sim \mathbb{F}_q} \omega^{\text{tr}(\beta x)} = \begin{cases} 1 & \text{if } \beta = 0, \\ 0 & \text{if } \beta \neq 0. \end{cases}$$

Proof. If $\beta = 0$ the character is constant. If $\beta \neq 0$, the map $x \mapsto \beta x$ permutes $\mathbb{F}_q$, so the average is the average of a nontrivial additive character, which vanishes. $\square$

Lemma 5.6 (Orthogonality of finite-field characters). *For $\alpha, \beta \in \mathbb{F}_q^m$,*

$$\frac{1}{q^m} \sum_{u \in \mathbb{F}_q^m} \omega^{\text{tr}(u \cdot (\beta - \alpha))} = \begin{cases} 1 & \text{if } \alpha = \beta, \\ 0 & \text{if } \alpha \neq \beta. \end{cases}$$

Proof. This is Lemma 3.3 applied to $v = \beta - \alpha$. $\square$

Lemma 5.7 (Eigenvectors of the hypercube graph). *For each $\alpha \in \mathbb{F}_q^m$, define*

$$|\varphi_\alpha\rangle = \frac{1}{M^{1/2}} \cdot \sum_{u \in \mathbb{F}_q^m} \omega^{\text{tr}(u \cdot \alpha)} |u\rangle.$$

Then the following two statements hold.

1. *The $|\varphi_\alpha\rangle$'s form an orthonormal basis of $\mathbb{C}^V$.*
2. *For each $\alpha \in \mathbb{F}_q^m$, $|\varphi_\alpha\rangle$ is an eigenvector for K with eigenvalue $\frac{1}{M} \cdot \frac{m - |\alpha|}{m}$, where $|\alpha|$ is the number of nonzero coordinates of α .*

Proof. First we prove 1. For $\alpha, \beta \in \mathbb{F}_q^m$,

$$\langle \varphi_\alpha | \varphi_\beta \rangle = \frac{1}{M} \sum_{u \in \mathbb{F}_q^m} \omega^{\text{tr}(u \cdot (\beta - \alpha))} = \begin{cases} 1 & \text{if } \alpha = \beta, \\ 0 & \text{if } \alpha \neq \beta, \end{cases}$$

by Lemma 5.6. Thus the $|\varphi_\alpha\rangle$ form an orthonormal basis of $\mathbb{C}^V$.

Next we prove 2. For $\alpha \in \mathbb{F}_q^m$,

$$\begin{aligned} K \cdot |\varphi_\alpha\rangle &= \left(\mathbb{E}_{(u,v) \sim C} |u\rangle \langle v| \right) \cdot \left(\frac{1}{M^{1/2}} \cdot \sum_{u \in \mathbb{F}_q^m} \omega^{\text{tr}(u \cdot \alpha)} |u\rangle \right) \\ &= \frac{1}{M^{1/2}} \cdot \mathbb{E}_{(u,v) \sim C} \omega^{\text{tr}(v \cdot \alpha)} |u\rangle. \end{aligned} \tag{5.1}$$

By the edge-sampling rule from Definition 5.2, we may write $v = u + x e_i$, where i is uniformly random in $\{1, \dots, m\}$ and x is uniformly random in $\mathbb{F}_q$. Therefore

$$(5.1) = \frac{1}{M^{1/2}} \cdot \mathbb{E}_{u,i,x} \omega^{\text{tr}((u + x e_i) \cdot \alpha)} |u\rangle = \left(\mathbb{E}_{i,x} \omega^{\text{tr}(x \alpha_i)} \right) \cdot \frac{1}{M^{1/2}} \cdot \mathbb{E}_u \omega^{\text{tr}(u \cdot \alpha)} |u\rangle = \frac{1}{M} \left(\mathbb{E}_{i,x} \omega^{\text{tr}(x \alpha_i)} \right) \cdot |\varphi_\alpha\rangle.$$

Hence $|\varphi_\alpha\rangle$ is an eigenvector of K with eigenvalue

$$\frac{1}{M} \cdot \mathbb{E}_i \left[\mathbb{E}_x \omega^{\text{tr}(x \alpha_i)} \right] = \frac{1}{M} \cdot \mathbb{E}_i \left[\mathbf{1}[\alpha_i = 0] \right] = \frac{1}{M} \cdot \frac{m - |\alpha|}{m},$$

by Lemma 5.5. $\square$

Lemma 5.8 (Spectral gap of the Laplacian). *If $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{q^m}$ are the eigenvalues of L , then*

$$\lambda_1 = 0, \quad \lambda_2 = \frac{1}{mq^m}.$$

Proof. Lemma 5.7 identifies the two largest eigenvalues of K , hence the two smallest eigenvalues of $L = (1/q^m)I - K$. $\square$

5.2 Local and global variance

Definition 5.9 (Local and global variance). Let $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ and let $0 \leq A^u \leq I$ be an operator for each $u \in \mathbb{F}_q^m$. The local variance is

$$\mathbf{Var}_{\text{local}}(A, \psi) = \frac{1}{2} \mathbb{E}_{(u,v) \sim C} \langle \psi | (A^u - A^v)^2 \otimes I | \psi \rangle,$$

and the global variance is

$$\mathbf{Var}_{\text{global}}(A, \psi) = \frac{1}{2} \mathbb{E}_{u,v \sim \mathbb{F}_q^m} \langle \psi | (A^u - A^v)^2 \otimes I | \psi \rangle.$$

Lemma 5.10 (Local variance as a Laplacian form). *Define*

$$A_{\text{comb}} = \sum_{u \in \mathbb{F}_q^m} |u\rangle \otimes A^u \otimes I.$$

Then

$$\text{Tr}(A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}}) = \mathbf{Var}_{\text{local}}(A, \psi).$$

Proof. For $u, v \in \mathbb{F}_q^m$,

$$\begin{aligned} ((\langle u| - \langle v|) \otimes \langle \psi|) \cdot A_{\text{comb}} &= ((\langle u| - \langle v|) \otimes \langle \psi|) \cdot \sum_{w \in \mathbb{F}_q^m} |w\rangle \otimes A^w \otimes I \\ &= \langle \psi| \cdot ((A^u - A^v) \otimes I). \end{aligned} \tag{5.2}$$

Therefore

$$A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}} = \frac{1}{2} \mathbb{E}_{(u,v) \sim C} A_{\text{comb}}^\dagger ((|u\rangle - |v\rangle)(\langle u| - \langle v|) \otimes |\psi\rangle\langle\psi|) A_{\text{comb}}$$

by Lemma 5.4, and hence

$$A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}} = \frac{1}{2} \mathbb{E}_{(u,v) \sim C} ((A^u - A^v) \otimes I) |\psi\rangle\langle\psi| ((A^u - A^v) \otimes I)$$

by (5.2). Taking the trace gives

$$\text{Tr}(A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}}) = \frac{1}{2} \mathbb{E}_{(u,v) \sim C} \langle \psi | (A^u - A^v)^2 \otimes I | \psi \rangle = \mathbf{Var}_{\text{local}}(A, \psi).$$

□

Lemma 5.11 (Global variance as the orthogonal Fourier mass). *Write*

$$A_{\text{comb}} = |\varphi_0\rangle \otimes A_0 + |\varphi_\perp\rangle \otimes A_\perp,$$

where $|\varphi_\perp\rangle$ is orthogonal to $|\varphi_0\rangle$. Then

$$\frac{1}{q^m} \text{Tr}((\langle \varphi_\perp | \otimes A_\perp)(I \otimes |\psi\rangle\langle\psi|)(|\varphi_\perp\rangle \otimes A_\perp)) = \mathbf{Var}_{\text{global}}(A, \psi).$$

Proof. We first compute

$$A_0 = (\langle \varphi_0 | \otimes I) \cdot A_{\text{comb}} = \left(\frac{1}{M^{1/2}} \cdot \sum_{u \in \mathbb{F}_q^m} \langle u | \right) \otimes I \cdot \left(\sum_{u \in \mathbb{F}_q^m} |u\rangle \otimes A^u \otimes I \right) = \frac{1}{M^{1/2}} \cdot \sum_{u \in \mathbb{F}_q^m} A^u \otimes I.$$

Writing $A_{\text{avg}} = \mathbb{E}_u A^u$, it follows that

$$|\varphi_0\rangle \otimes A_0 = \sum_{u \in \mathbb{F}_q^m} |u\rangle \otimes A_{\text{avg}} \otimes I, \quad |\varphi_\perp\rangle \otimes A_\perp = \sum_{u \in \mathbb{F}_q^m} |u\rangle \otimes (A^u - A_{\text{avg}}) \otimes I.$$

Hence

$$\frac{1}{M} \cdot \text{Tr}((\langle \varphi_\perp | \otimes A_\perp)(I \otimes |\psi\rangle\langle\psi|)(|\varphi_\perp\rangle \otimes A_\perp)) = \mathbb{E}_{u \sim \mathbb{F}_q^m} \langle \psi | (A^u - A_{\text{avg}})^2 \otimes I | \psi \rangle. \quad (5.3)$$

Moreover,

$$\mathbb{E}_{u \sim \mathbb{F}_q^m} (A^u - A_{\text{avg}})^2 = \mathbb{E}_{u \sim \mathbb{F}_q^m} ((A^u)^2 - (A_{\text{avg}})^2) = \frac{1}{2} \mathbb{E}_{u, v \sim \mathbb{F}_q^m} (A^u - A^v)^2.$$

Substituting this identity into (5.3) yields

$$\frac{1}{M} \cdot \text{Tr}((\langle \varphi_\perp | \otimes A_\perp)(I \otimes |\psi\rangle\langle\psi|)(|\varphi_\perp\rangle \otimes A_\perp)) = \frac{1}{2} \mathbb{E}_{u, v \sim \mathbb{F}_q^m} \langle \psi | (A^u - A^v)^2 \otimes I | \psi \rangle = \mathbf{Var}_{\text{global}}(A, \psi).$$

□

Lemma 5.12 (Local-to-global inequality). *For every family $\{A^u\}$,*

$$\mathbf{Var}_{\text{global}}(A, \psi) \leq m \mathbf{Var}_{\text{local}}(A, \psi).$$

Proof.

$$\begin{aligned} A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}} &= (\langle \varphi_0 | \otimes A_0 + \langle \varphi_\perp | \otimes A_\perp)(L \otimes |\psi\rangle\langle\psi|)(|\varphi_0\rangle \otimes A_0 + |\varphi_\perp\rangle \otimes A_\perp) \\ &= (\langle \varphi_\perp | \otimes A_\perp)(L \otimes |\psi\rangle\langle\psi|)(|\varphi_\perp\rangle \otimes A_\perp) \\ &= \langle \varphi_\perp | L | \varphi_\perp \rangle \cdot A_\perp |\psi\rangle\langle\psi| A_\perp. \end{aligned} \quad (5.4)$$

The second step uses that $|\varphi_0\rangle$ is a 0-eigenvector of L . Since $|\varphi_\perp\rangle$ is orthogonal to $|\varphi_0\rangle$, Lemma 5.8 gives

$$\langle \varphi_\perp | L | \varphi_\perp \rangle \geq \frac{1}{mM} \cdot \langle \varphi_\perp | \varphi_\perp \rangle.$$

Therefore

$$\mathbf{Var}_{\text{local}}(A, \psi) = \text{Tr}(A_{\text{comb}}^\dagger (L \otimes |\psi\rangle\langle\psi|) A_{\text{comb}}) = \langle \varphi_\perp | L | \varphi_\perp \rangle \cdot \text{Tr}(A_\perp |\psi\rangle\langle\psi| A_\perp)$$

by Lemma 5.10 and (5.4), so

$$\mathbf{Var}_{\text{local}}(A, \psi) \geq \frac{1}{mM} \cdot \text{Tr}((\langle \varphi_\perp | \otimes A_\perp)(I \otimes |\psi\rangle\langle\psi|)(|\varphi_\perp\rangle \otimes A_\perp)) = \frac{1}{m} \mathbf{Var}_{\text{global}}(A, \psi),$$

where the last step is Lemma 5.11. Rearranging gives the claim. □

Chapter 6

Global variance of the points measurements

Throughout this chapter, (ψ, A, B, L) is an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the (m, q, d) low individual degree test.

Lemma 6.1 (Replacing a line answer by a fixed polynomial). *Let $G \in \text{PolySub}(m, q, d)$. On the axis-parallel line-test distribution,*

$$B_{[f(u)=g(u)]}^\ell \otimes (G_g)^{1/2} \approx_{md/q} B_{g|_\ell}^\ell \otimes (G_g)^{1/2}.$$

Proof. Expand the $\approx_{md/q}$ norm on the weighted vector $(I \otimes (G_g)^{1/2})|\psi\rangle$. Because B^ℓ is a projective measurement, only those line polynomials f with $f \neq g|_\ell$ and $f(u) = g(u)$ contribute. Lemma 3.9 is the univariate Schwartz–Zippel wrapper used in Lean for this line-restriction event; it supplies the paper’s md/q loss, and the remaining operator sum is bounded by the fact that G is a submeasurement. $\square$

Remark 6.2 (Conditional variants for the collision-residual estimate). The paper proof of Lemma 6.1 bounds the collision event directly by the Schwartz–Zippel estimate. The following conditional variants instead assume this collision estimate, or one of the reindexing identities that implies it, as an explicit hypothesis. These conditional statements record proof obligations to be discharged; they are not additional hypotheses in the paper statement of Lemma 6.1.

Lemma 6.3 (Local variance of the points measurements). *Let $G \in \text{PolySub}(m, q, d)$. On the hypercube edge distribution $(u, v) \sim C$,*

$$A_{g(u)}^u \otimes (G_g)^{1/2} \approx_{24 \cdot (\varepsilon + \delta + \frac{md}{q})} A_{g(v)}^v \otimes (G_g)^{1/2}. \quad (6.1)$$

Proof. Fix g and keep the weight $(G_g)^{1/2}$ on the second tensor factor throughout. On the vector $(I \otimes (G_g)^{1/2})|\psi\rangle$ one has the six-step chain

$$A_{g(u)}^u \otimes (G_g)^{1/2} \approx_{2\delta} I \otimes (G_g)^{1/2} A_{g(u)}^u \approx_{2\varepsilon} B_{[f(u)=g(u)]}^\ell \otimes (G_g)^{1/2} \approx_{md/q} B_{g|_\ell}^\ell \otimes (G_g)^{1/2} \approx_{md/q} B_{[f(v)=g(v)]}^\ell \otimes (G_g)^{1/2} \approx_{2\varepsilon} I \otimes (G_g)^{1/2} A_{g(v)}^v \approx_{2\delta} I \otimes (G_g)^{1/2} A_{g(v)}^v \otimes (G_g)^{1/2}.$$

The first, second, fifth, and sixth comparisons use Lemma 3.21 together with Theorem 3.25 and the good-strategy hypotheses, and the middle pair is Lemma 6.1. Lemma 3.27 sums these six weighted transports. $\square$

Lemma 6.4 (Equivalent local-variance bound). *The local-variance bound for the point measurements is*

$$\sum_{g \in \mathcal{P}(m, q, d)} \mathbb{E}_{(u, v) \sim C} \langle \psi | (A_{g(u)}^u - A_{g(v)}^v)^2 \otimes G_g | \psi \rangle \leq 24 \left(\varepsilon + \delta + \frac{md}{q} \right). \quad (6.2)$$

Remark 6.5 (Polynomial-sum formalization for 6.2). The three referenced declarations and Lemma 6.4 together provide the full polynomial-sum (cardinality-free) chain proving the equation. The chain assembly lemma `localVarianceDeviation_sum_le_localVarianceOfPointsError` telescopes the six operator differences via `ev_sum_conjTranspose_mul_sum_le`, sums over g , and applies the six individual `_sum` bounds (including the reverse `generalizeB` sum) to avoid the per-polynomial cardinality blow-up.

Lemma 6.6 (Global variance of the points measurements). *Let $G \in \text{PolySub}(m, q, d)$. On the uniform distribution over independent $u, v \in \mathbb{F}_q^m$,*

$$A_{g(u)}^u \otimes (G_g)^{1/2} \approx_{24m(\varepsilon + \delta + \frac{md}{q})} A_{g(v)}^v \otimes (G_g)^{1/2}. \quad (6.3)$$

Proof. We want to bound

$$\mathbb{E}_{u, v \sim \mathbb{F}_q^m} \sum_{g \in \mathcal{P}(m, q, d)} \|(A_{g(u)}^u - A_{g(v)}^v) \otimes (G_g)^{1/2} | \psi \rangle\|^2 = \mathbb{E}_{u, v \sim \mathbb{F}_q^m} \sum_{g \in \mathcal{P}(m, q, d)} \langle \psi | (A_{g(u)}^u - A_{g(v)}^v)^2 \otimes G_g | \psi \rangle. \quad (6.4)$$

For each $g \in \mathcal{P}(m, q, d)$, define

$$A(g)^u := A_{g(u)}^u, \quad |\psi_g\rangle := (I \otimes (G_g)^{1/2}) | \psi \rangle.$$

Then

$$(6.4) = \sum_{g \in \mathcal{P}(m, q, d)} \mathbb{E}_{u, v \sim \mathbb{F}_q^m} \langle \psi_g | (A(g)^u - A(g)^v)^2 \otimes I | \psi_g \rangle = \sum_{g \in \mathcal{P}(m, q, d)} 2 \mathbf{Var}_{\text{global}}(A(g), \psi_g).$$

Applying Lemma 5.12 gives

$$(6.4) \leq \sum_{g \in \mathcal{P}(m, q, d)} 2m \mathbf{Var}_{\text{local}}(A(g), \psi_g) = m \cdot \sum_{g \in \mathcal{P}(m, q, d)} \mathbb{E}_{(u, v) \sim C} \langle \psi | (A_{g(u)}^u - A_{g(v)}^v)^2 \otimes G_g | \psi \rangle.$$

The last quantity is at most $m \cdot 24 \left(\varepsilon + \delta + \frac{md}{q} \right)$ by Lemma 6.4, which is exactly 6.3. $\square$

Chapter 7

Self-improvement

Throughout this chapter, (ψ, A, B, L) is an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the (m, q, d) low individual degree test.

7.1 Non-projective Output

A large part of this chapter is devoted to proving Lemma 7.1, which is a slightly weaker form of the projective self-improvement theorem proved in Section 7.2. The key difference is that the output family H is only required to be a submeasurement, rather than a projective submeasurement. Once this non-projective statement is available, Theorem 4.3 upgrades H to a projective family without losing control of the four quantitative conclusions.

Compared with the projective theorem, the helper lemma packages strong self-consistency using the explicit inequality from Definition 3.34, keeps the boundedness conclusion in terms of an auxiliary operator Z , and produces a much smaller error parameter.

Lemma 7.1 (Self-improvement with non-projective output). *Let $G \in \text{PolyMeas}(m, q, d)$ be a measurement with the following property:*

1. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\nu I \otimes G_{[g(u)=a]}.$$

Let

$$\zeta = 100m \cdot \left(\varepsilon^{1/2} + \delta^{1/2} + (d/q)^{1/2} \right).$$

Then there exists $H \in \text{PolySub}(m, q, d)$ with the following properties:

1. (Completeness): If $H = \sum_h H_h$, then

$$\langle \psi | H \otimes I | \psi \rangle \geq (1 - \nu) - \zeta.$$

2. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\zeta I \otimes H_{[h(u)=a]}.$$

3. (Strong self-consistency):

$$\sum_h \langle \psi | H_h \otimes H_h | \psi \rangle \geq \langle \psi | H \otimes I | \psi \rangle - \zeta.$$

4. (Boundedness): There exists a positive-semidefinite matrix Z such that

$$\langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle \leq \zeta$$

and for each $h \in \mathcal{P}(m, q, d)$,

$$Z \geq \left(\mathbb{E}_u A_{h(u)}^u \right).$$

Proof. Let $T = \{T_g\}$ and Z be the optimal solutions to the SDPs (7.23) and (7.24) given by Lemma 7.8. Then T is a measurement, and

$$\forall g \in \mathcal{P}(m, q, d) : \quad Z \geq \left(\mathbb{E}_u A_{g(u)}^u \right), \quad (7.1)$$

$$T_g \cdot Z = T_g \cdot \left(\mathbb{E}_u A_{g(u)}^u \right). \quad (7.2)$$

For each $u \in \mathbb{F}_q^m$, define $H^u = \{H_h^u\}_{h \in \mathcal{P}(m, q, d)}$ by

$$H_h^u := A_{h(u)}^u \cdot T_h \cdot A_{h(u)}^u,$$

and let

$$H_h := \mathbb{E}_u H_h^u.$$

The pointwise family H^u is a submeasurement because

$$\sum_h H_h^u = \sum_a A_a^u \cdot \left(\sum_{h: h(u)=a} T_h \right) \cdot A_a^u \leq \sum_a (A_a^u)^2 = I,$$

where the inequality uses that T is a measurement and the last identity uses that A is projective. Averaging over u gives

$$\sum_h H_h = \mathbb{E}_u \sum_h H_h^u \leq I,$$

so H is also a submeasurement and therefore lies in $\text{PolySub}(m, q, d)$.

Set

$$\zeta_{\text{variance}} = 24m \cdot \left(\varepsilon + \delta + \frac{md}{q} \right).$$

Lemma 7.9 below is the technical transfer from the averaged family H back to the sandwiched operators $A_{h(u)}^u T_h A_{h(u)}^u$. We use it in the completeness, consistency, strong self-consistency, and boundedness estimates that follow.

We now verify the four conclusions of Lemma 7.1.

Proof of 1. The completeness of H is

$$\begin{aligned} \sum_h \langle \psi | H_h \otimes I | \psi \rangle &= \mathbb{E}_u \sum_h \langle \psi | H_h^u \otimes I | \psi \rangle \\ &= \mathbb{E}_u \sum_h \langle \psi | (A_{h(u)}^u \cdot T_h \cdot A_{h(u)}^u) \otimes I | \psi \rangle. \end{aligned} \quad (7.3)$$

Grouping the polynomials according to their value at u gives

$$\mathbb{E}_u \sum_h \langle \psi | (A_{h(u)}^u \cdot T_h \cdot A_{h(u)}^u) \otimes I | \psi \rangle = \mathbb{E}_u \sum_a \langle \psi | (A_a^u \cdot T_{[h(u)=a]} \cdot A_a^u) \otimes I | \psi \rangle. \quad (7.4)$$

We first move the leftmost copy of A_a^u across the bipartition:

$$(7.4) \approx_{2\sqrt{\delta}} \mathbb{E}_u \sum_a \langle \psi | (T_{[h(u)=a]} \cdot A_a^u) \otimes A_a^u | \psi \rangle. \quad (7.5)$$

The difference is bounded by Cauchy–Schwarz as in (7.33); the first factor is at most $\sqrt{2\delta}$ by self-consistency of A , and the second is at most 1 because $T_{[h(u)=a]} \leq I$. We next remove the remaining copy of A_a^u on Bob’s side:

$$(7.5) \approx_{\sqrt{\delta}} \mathbb{E}_u \sum_a \langle \psi | (T_{[h(u)=a]} \cdot A_a^u) \otimes I | \psi \rangle. \quad (7.6)$$

The first Cauchy–Schwarz factor is at most 1 because the fiber operators form a submeasurement after grouping T by the value of $h(u)$. Here the second Cauchy–Schwarz factor is

$$\mathbb{E}_u \sum_{a \neq b} \langle \psi | A_a^u \otimes A_b^u | \psi \rangle,$$

which is at most δ because A is self-consistent and projective. Reindexing by h and using (7.2),

$$(7.6) = \sum_h \langle \psi | (T_h \cdot \mathbb{E}_u A_{h(u)}^u) \otimes I | \psi \rangle = \sum_h \langle \psi | (T_h \cdot Z) \otimes I | \psi \rangle = \langle \psi | Z \otimes I | \psi \rangle.$$

Thus

$$\sum_h \langle \psi | H_h \otimes I | \psi \rangle \geq \langle \psi | Z \otimes I | \psi \rangle - 3\sqrt{\delta}. \quad (7.7)$$

Since G is a measurement, (7.1) and Lemma 3.18 give

$$\langle \psi | Z \otimes I | \psi \rangle \geq \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes G_{[g(u)=a]} | \psi \rangle \geq 1 - \nu.$$

Therefore the completeness is at least $1 - \nu - 3\sqrt{\delta}$, which is bounded below by $(1 - \nu) - \zeta$.

Proof of 2. The inconsistency of H with A is

$$\mathbb{E}_u \sum_{a \neq b} \langle \psi | A_a^u \otimes H_{[h(u)=b]} | \psi \rangle = \mathbb{E}_u \sum_{a, h: h(u) \neq a} \langle \psi | A_a^u \otimes H_h | \psi \rangle. \quad (7.8)$$

The theorem-side off-diagonal selection used for this ‘add-in-u’ application is , and the corresponding left/right scalar quantities are and . The selected scalar chain for this specialization is recorded by . Apply Lemma 7.9 with $\mathcal{O} = \mathbb{F}_q$, $M = A$, and $S_u = \{(a, h) : h(u) \neq a\}$. Then

$$(7.8) \approx_{4\sqrt{\zeta_{\text{variance}}}} \mathbb{E}_u \sum_{a, h: h(u) \neq a} \langle \psi | (A_{h(u)}^u \cdot A_a^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle,$$

and the right-hand side is 0 because A is projective. Hence

$$\mathbb{E}_u \sum_{a \neq b} \langle \psi | A_a^u \otimes H_{[h(u)=b]} | \psi \rangle \leq 4\sqrt{\zeta_{\text{variance}}}. \quad (7.9)$$

Since

$$4\sqrt{\zeta_{\text{variance}}} = 4\sqrt{24m \cdot \left(\varepsilon + \delta + \frac{md}{q}\right)} \leq 20m \cdot \left(\varepsilon^{1/2} + \delta^{1/2} + (d/q)^{1/2}\right) \leq \zeta,$$

this proves 2.

Proof of 3. From the definition of H_h^u and the projectivity of A ,

$$H_h^u = A_{h(u)}^u \cdot T_h \cdot A_{h(u)}^u = A_{h(u)}^u \cdot H_h^u \cdot A_{h(u)}^u, \quad (7.10)$$

$$A_{h(u)}^u \cdot H_{h'}^u \cdot A_{h(u)}^u = H_{h'}^u \cdot A_{h(u)}^u = (A_{h(u)}^u \cdot T_{h'} \cdot A_{h(u)}^u) \cdot \mathbf{1}[h(u) = h'(u)]. \quad (7.11)$$

The strong self-consistency expression is

$$\sum_{h \in \mathcal{P}(m, q, d)} \langle \psi | H_h \otimes H_h | \psi \rangle = \mathbb{E}_u \sum_{h \in \mathcal{P}(m, q, d)} \langle \psi | H_h^u \otimes H_h | \psi \rangle. \quad (7.12)$$

Apply Lemma 7.9 with $\mathcal{O} = \mathcal{P}(m, q, d)$, $M = H$, and $S_u = \{(h, h) : h \in \mathcal{P}(m, q, d)\}$. This gives

$$(7.12) \approx_{4\sqrt{\zeta_{\text{variance}}}} \mathbb{E}_u \sum_{h \in \mathcal{P}(m, q, d)} \langle \psi | (A_{h(u)}^u \cdot H_h^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle. \quad (7.13)$$

We next enlarge the sum from $h' = h$ to all pairs h, h' :

$$(7.13) \approx_{2\sqrt{\zeta_{\text{variance}} + \frac{md}{q}}} \mathbb{E}_u \sum_{h, h' \in \mathcal{P}(m, q, d)} \langle \psi | (A_{h(u)}^u \cdot H_{h'}^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle. \quad (7.14)$$

The added off-diagonal contribution is

$$\begin{aligned} (7.14) - (7.13) &= \mathbb{E}_u \sum_{h \neq h'} \langle \psi | (A_{h(u)}^u \cdot H_{h'}^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle \\ &= \mathbb{E}_u \sum_{h \neq h'} \langle \psi | (A_{h(u)}^u \cdot T_{h'} \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle \cdot \mathbf{1}[h(u) = h'(u)], \end{aligned} \quad (7.15)$$

where the second equality is (7.11). To estimate (7.15), first replace the left copy of $A_{h(u)}^u$ by $A_{h(v)}^v$:

$$(7.15) \approx_{\sqrt{\zeta_{\text{variance}}}} \mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | (A_{h(v)}^v \cdot T_{h'} \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle \cdot \mathbf{1}[h(u) = h'(u)]. \quad (7.16)$$

The corresponding Cauchy-Schwarz bound is

$$\begin{aligned} &\left| \mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | ((A_{h(u)}^u - A_{h(v)}^v) \cdot T_{h'} \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle \cdot \mathbf{1}[h(u) = h'(u)] \right| \\ &\leq \sqrt{\mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | ((A_{h(u)}^u - A_{h(v)}^v) \cdot T_{h'} \cdot (A_{h(u)}^u - A_{h(v)}^v)) \otimes T_h | \psi \rangle} \\ &\quad \cdot \sqrt{\mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | (A_{h(u)}^u \cdot T_{h'} \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle \cdot \mathbf{1}[h(u) = h'(u)]}. \end{aligned} \quad (7.17)$$

The first factor is bounded by $\sqrt{\zeta_{\text{variance}}}$ via Lemma 6.6, and the second by 1 because T and H^u are submeasurements.

Repeating the same argument for the remaining copy of A yields

$$(7.16) \approx_{\sqrt{\zeta_{\text{variance}}}} \mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | (A_{h(v)}^v \cdot T_{h'} \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle \cdot \mathbf{1}[h(u) = h'(u)]. \quad (7.18)$$

The same intermediate estimate gives

$$\mathbb{E}_{u,v} \sum_{h \neq h'} \langle \psi | (A_{h(v)}^v \cdot T_{h'} \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle \leq 1, \quad (7.19)$$

so Schwartz-Zippel bounds the indicator average by md/q and proves (7.14).

Using (7.11), the right-hand side of (7.14) is

$$\mathbb{E}_u \sum_{h, h' \in \mathcal{P}(m, q, d)} \langle \psi | (H_{h'}^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle. \quad (7.20)$$

Replace the remaining copy of $A_{h(u)}^u$ by $A_{h(v)}^v$:

$$(7.20) \approx \sqrt{\zeta_{\text{variance}}} \mathbb{E}_{u, v} \sum_{h, h' \in \mathcal{P}(m, q, d)} \langle \psi | (H_{h'}^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle. \quad (7.21)$$

Here the first Cauchy–Schwarz factor is at most 1 because T and H^u are submeasurements, and the second is again bounded by Lemma 6.6. Finally move this last copy of $A_{h(v)}^v$ to Bob’s side:

$$(7.21) \approx \sqrt{2\delta} \mathbb{E}_{u, v} \sum_{h, h' \in \mathcal{P}(m, q, d)} \langle \psi | H_{h'}^u \otimes (T_h \cdot A_{h(v)}^v) | \psi \rangle. \quad (7.22)$$

Substituting (7.2), then (7.1), and finally the already proved bound (7.9), one obtains

$$(7.22) \geq \sum_h \langle \psi | H_h \otimes I | \psi \rangle - 4\sqrt{\zeta_{\text{variance}}}.$$

Combining (7.13), (7.14), (7.21), and (7.22) gives

$$\sum_h \langle \psi | H_h \otimes H_h | \psi \rangle \geq \sum_h \langle \psi | H_h \otimes I | \psi \rangle - 11\sqrt{\zeta_{\text{variance}}} - \sqrt{2\delta} - \frac{md}{q}.$$

The Lean formalization records this step as a named proof obligation, starting with the input structure and the exact residual-side expansion and then uses the proved lemma together with to recover the final helper-stage self-consistency conclusion. The remaining theorem-level derivation is not moved into an additional hypothesis; it is the proof gap in the paper theorem below. The paper’s arithmetic shows that the right-hand error is at most

$$57m \cdot \left(\varepsilon^{1/2} + \delta^{1/2} + (d/q)^{1/2} \right) \leq \zeta,$$

which proves 3.

Proof of 4. The helper-stage boundedness quantity is

$$\langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle.$$

The reindexing of the sum by h is the algebraic identity

$$\sum_a A_a^u \otimes H_{[h(u)=a]} = \sum_h A_{h(u)}^u \otimes H_h,$$

whose averaged scalar form reads

$$\mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle = \mathbb{E}_u \sum_h \langle \psi | A_{h(u)}^u \otimes H_h | \psi \rangle.$$

Using $\sum_a A_a^u = I$, the pointwise difference between the right-placed total $I \otimes H_{\text{total}} = \sum_h I \otimes H_h$ and the helper-agreement operator at u reindexes as the off-diagonal sum

$$\sum_h I \otimes H_h - \sum_a A_a^u \otimes H_{[h(u)=a]} = \sum_h \sum_{a \neq h(u)} A_a^u \otimes H_h,$$

whose averaged scalar form reads

$$\sum_h \langle \psi | I \otimes H_h | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle = \mathbb{E}_u \sum_h \sum_{a \neq h(u)} \langle \psi | A_a^u \otimes H_h | \psi \rangle.$$

Combined with (7.9) this gives

$$\langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle \leq \langle \psi | Z \otimes I | \psi \rangle - \sum_h \langle \psi | I \otimes H_h | \psi \rangle + 4\sqrt{\zeta_{\text{variance}}}.$$

By (7.7), the right-hand side is at most $3\sqrt{\delta} + 4\sqrt{\zeta_{\text{variance}}}$. Substituting the definition of ζ_{variance} and collecting terms gives

$$23m \cdot \left(\varepsilon^{1/2} + \delta^{1/2} + (d/q)^{1/2} \right) \leq \zeta.$$

Together with (7.1), this proves 4. $\square$

Remark 7.2 (Slackness-carrying self-improvement helper). The formalization records a strengthened helper theorem whose output carries the complementary-slackness equations used in the completeness chain. The previous top-level residual-domination assembly theorems have been removed: the full self-improvement conclusion is now stated only by the paper theorem below, whose incomplete proof explicitly records the remaining Section 9 work.

7.1.1 A Semidefinite Program

Lemma 7.3 (Uniform SDP feasible witness). *The uniform primal family $T_g = (2|\mathcal{P}(m, q, d)|)^{-1}I$ has total mass $(1/2)I$, and the dual operator $Z = 2I$ is positive semidefinite and dominates the identity.*

Proof. This is the elementary Slater-type feasible witness recorded in the basic self-improvement definitions. $\square$

Lemma 7.4 (Explicit matrix feasible bounds). *In every concrete matrix realization, the uniform family $T_g = (2|\mathcal{P}(m, q, d)|)^{-1}I$ has total mass $(1/2)I$. The dual operator $Z = 2I$ dominates the identity and satisfies $Z - A_g \geq I$, hence $Z \geq A_g$, for every polynomial g . In the canonical block SDP, the corresponding block matrix is feasible, its slack block is $(1/2)I$, and the canonical dual slack for the same dual witness also dominates the block identity. More generally, dual feasibility implies positivity of Z , since the averaged point operators A_g are positive semidefinite.*

Proof. The formal proof is the matrix specialization of the uniform SDP witness recorded in the basic definitions. The inequality $A_g \leq I$ follows by averaging the point-measurement effects and using the submeasurement bound at each point. $\square$

Lemma 7.5 (Matrix slackness output). *From a matrix-level strong-duality witness with complementary slackness of the shape asserted by Lemma 7.8, one obtains a complete measurement $\{T_g\}$, a feasible dual operator Z , equality of the primal and dual objectives, and the equations $T_g Z = T_g A_g$ for all g . Lean isolates the saturated canonical optimal-pair output required from strong duality: a feasible canonical primal matrix, a dual-feasible operator with equal objective value, canonical complementary slackness, and vanishing of the extra canonical slack block. This datum gives both the abstract slackness-carrying SDP statement and the displayed measurement witness without assuming $I \leq Z$. Lean also proves the converse saturation step for the dominance-carrying optimal pair: the canonical complementary-slackness equation on the extra block gives $X_{\text{none, none}} Z = 0$, and the lower bound $I \leq Z$ forces $X_{\text{none, none}} = 0$. Thus the dominance-carrying*

optimal pair canonically determines the saturated optimal pair, and hence also the matrix slackness statement without the dominance field. The paper-facing self-improvement helper consumes the resulting abstract SDP statement.

Proof. The identity $\sum_g T_g = I$ turns the optimal primal submeasurement into a measurement. The zero-defect form $T_g(Z - A_g) = 0$ is equivalent, by distributivity, to the displayed equation $T_g Z = T_g A_g$. $\square$

Remark 7.6 (Removed residual-dominating construction theorems). Earlier Lean versions exposed top-level conditional construction theorems that combined the matrix slackness output with residual-dominating orthonormalization and QXP-repair inputs. These declarations have been removed. The matrix slackness material is retained only up to the slackness-carrying helper output, and the full projective-output conclusion is represented by the paper theorem `MIPStarRE.LDT.SelfImprovement.selfImprovement`.

Remark 7.7 (Lean reduced SDP and dominance interfaces). Lean also contains reduced SDP and dominance-carrying matrix interfaces. The reduced theorem `sdp` proves only the measurement-total and dual-feasibility fragment used before strong duality is supplied. The dominance-carrying declarations add the auxiliary condition $I \leq Z$ and are retained as technical routes to slackness-carrying data. These declarations are not advertised as formalizations of Lemma 7.8.

Lemma 7.8 (Dual SDP and complementary slackness). *Let*

$$A_g = \mathbb{E}_{u \sim \mathbb{F}_q^m} A_{g(u)}^u.$$

Then the primal is

$$\begin{aligned} \sup \quad & \sum_g \text{Tr}(T_g \cdot A_g) \\ \text{s.t.} \quad & T_g \geq 0 \quad \forall g \in \mathcal{P}(m, q, d), \\ & \sum_g T_g \leq I, \end{aligned} \tag{7.23}$$

and the dual is

$$\inf \quad \text{Tr}(Z) \tag{7.24}$$

$$\text{s.t.} \quad Z \geq A_g. \tag{7.25}$$

These semidefinite programs are dual to each other. Moreover there is an optimal pair of solutions $\{T_g\}$ to (7.23) and Z to (7.24) such that $\sum_g T_g = I$ and

$$T_g Z = T_g A_g, \quad \forall g \in \mathcal{P}(m, q, d). \tag{7.26}$$

Proof. Rewrite (7.23) in canonical block form. Let r be the dimension of the space on which A acts, let $M = |\mathcal{P}(m, q, d)|$, and fix an ordering $g_1, \dots, g_M$ of the polynomials. Consider

$$\begin{aligned} \sup \quad & \text{Tr}(C^\dagger X) \\ \text{s.t.} \quad & \text{Tr}(D_{ij}^\dagger X) = b_{ij} \quad \forall i, j \in \{1, \dots, r\}, \\ & X \geq 0, \end{aligned} \tag{7.27}$$

where

$$C = \sum_{i=1}^M |i\rangle\langle i| \otimes A_{g_i},$$

and, for $i, j \in \{1, \dots, r\}$,

$$D_{ij} = \sum_{k=1}^{M+1} |k\rangle\langle k| \otimes |i\rangle\langle j|, \quad b_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Writing $X = \sum_{i,j=1}^{M+1} |i\rangle\langle j| \otimes X_{ij}$, the constraints say precisely that $\sum_{i=1}^{M+1} X_{ii} = I$, hence $\sum_{i=1}^M X_{ii} \leq I$, and the objective is

$$\text{Tr}(C^\dagger X) = \sum_{i=1}^M \text{Tr}(A_{g_i} \cdot X_{ii}).$$

Thus (7.27) is equivalent to (7.23) under the identification $T_{g_i} = X_{ii}$.

The canonical dual is

$$\inf \quad \sum_i z_{ij} b_{ij} \tag{7.28}$$

$$\text{s.t.} \quad \sum_{i,j} z_{ij} D_{ij} \geq C. \tag{7.29}$$

Since

$$\sum_{i,j=1}^r z_{ij} D_{ij} = \sum_{k=1}^{M+1} |k\rangle\langle k| \otimes \left(\sum_{i,j=1}^r z_{ij} |i\rangle\langle j| \right) =: \sum_{k=1}^{M+1} |k\rangle\langle k| \otimes Z,$$

the constraint (7.29) is exactly $Z \geq A_{g_i}$ for every i , and the objective is $\text{Tr}(Z)$. In Lean, the latter statement is recorded as the trace-pairing identity

$$\Re \text{Tr}(D(Z)X) = \Re \text{Tr}(Z)$$

for every feasible canonical primal matrix X , where $D(Z)$ denotes the block-diagonal canonical dual operator. Consequently the canonical primal-dual gap is the real trace pairing of X with the canonical dual slack $D(Z) - C$. Since the real trace pairing of positive semidefinite operators is nonnegative, Lean also records the resulting weak-duality inequality for any feasible canonical primal and dual pair. Hence (7.28) is equivalent to (7.24).

Slater's condition holds because

$$T_g = \frac{1}{2M} \cdot I \quad \forall g \in \mathcal{P}(m, q, d), \quad Z = 2I$$

are strictly feasible for the primal and dual respectively. Therefore strong duality holds. For an optimal pair $(X, (z_{ij}))$ in canonical form, complementary slackness gives

$$X \left(\sum_{i,j} z_{ij} D_{ij} - C \right) = 0. \tag{7.30}$$

We may assume that X is block diagonal. Translating (7.30) back to the variables $\{T_g\}$ and Z yields $T_g(Z - A_g) = 0$ for every g and $SZ = 0$, where $S = I - \sum_g T_g$ is the slack block. The paper then treats this as forcing $S = 0$. The current Lean development separates this point from the unconditional SDP statement: it proves the displayed saturation from the auxiliary dominance hypothesis $I \leq Z$. Thus, if the canonical complementary-slackness pair is supplied together with $I \leq Z$, Lean packages these conditional conclusions as a `MatrixSdpOptimalWitnessWithDominance`

once dual feasibility and equality of the paper primal and dual objectives have also been supplied. It also records the corresponding statement-level package, and an elimination theorem which exposes the complete measurement, the dual witness Z , the retained dominance $I \leq Z$, dual feasibility, strong duality, and the equations (7.26). The additional dominance requirement is not a consequence of the dual constraints $Z \geq A_g$ alone; the discrepancy and the intended saturated-witness replacement are recorded in `docs/paper-gaps/issue-1230-self-improvement-sdp-usage.tex`. In the concrete matrix realization, Lean records the defect-zero condition as `matrixSdpComplementarySlacknessDefe` the theorem `MatrixSdpOptimalWitness.complementarySlacknessEquation` translates it to the displayed equation $T_g Z = T_g A_g$, while `MatrixSdpOptimalWitness.primalMeasurement` packages $\sum_g T_g = I$ as a complete measurement. The statement `MatrixSdpStatementWithSlackness` records this matrix-level strong-duality output without identifying it with the reduced feasible witness supplied by `sdp`. The explicit Slater witnesses used above are separately recorded by `matrixSdpStrictPrimalSubmeasurement`, `matrixSdpStrictDualWitness`, and `matrixSdpFeasibleBounds_canoni` the dual feasibility of $2I$ follows from `matrixAveragedPointOperator_le_one`. The block-diagonal reduction is also recorded for arbitrary feasible canonical matrices: extracting the diagonal blocks preserves the objective and transfers canonical complementary slackness to the extracted paper primal submeasurement. $\square$

Lemma 7.9 (Averaging over the evaluation point). *Let $T = \{T_h\}$ be an optimal solution of the primal SDP, let*

$$H_h = \mathbb{E}_{u \sim \mathbb{F}_q^m} A_{h(u)}^u T_h A_{h(u)}^u,$$

and set

$$\zeta_{\text{variance}} = 24m \cdot \left(\varepsilon + \delta + \frac{md}{q} \right).$$

Suppose $M = \{M_o^u\}$ is a submeasurement with outcomes in some set $\mathcal{O}$. For each $u \in \mathbb{F}_q^m$, let S_u be a subset of $\mathcal{O} \times \mathcal{P}(m, q, d)$. Then

$$\mathbb{E}_u \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes H_h | \psi \rangle \approx_{4\sqrt{\zeta_{\text{variance}}}} \mathbb{E}_u \sum_{(o,h) \in S_u} \langle \psi | (A_{h(u)}^u \cdot M_o^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle.$$

The reduced variance-bound specialization keeps only the variance-bound consequence used later in the helper theorem; it is not itself the full formal counterpart of Lemma 7.9.

Proof. We begin by expanding

$$\begin{aligned} \mathbb{E}_u \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes H_h | \psi \rangle &= \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes H_h^v | \psi \rangle \\ &= \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes (A_{h(v)}^v \cdot T_h \cdot A_{h(v)}^v) | \psi \rangle. \end{aligned} \quad (7.31)$$

We claim that

$$(7.31) \approx_{\sqrt{2\delta}} \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \cdot M_o^u) \otimes (T_h \cdot A_{h(v)}^v) | \psi \rangle. \quad (7.32)$$

To show this, we bound the magnitude of the difference:

$$\begin{aligned}
& \left| \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) \cdot (M_o^u \otimes (T_h \cdot A_{h(v)}^v)) | \psi \rangle \right| \\
& \leq \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) \cdot (M_o^u \otimes T_h) \cdot (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) | \psi \rangle} \\
& \quad \cdot \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes (A_{h(v)}^v \cdot T_h \cdot A_{h(v)}^v) | \psi \rangle}. \quad (7.33)
\end{aligned}$$

The first factor is controlled by summing over the value $a = h(v)$:

$$\begin{aligned}
& \mathbb{E}_{u,v} \sum_{a \in \mathbb{F}_q} \langle \psi | (A_a^v \otimes I - I \otimes A_a^v) \cdot \left(\sum_{(o,h) \in S_u, h(v)=a} M_o^u \otimes T_h \right) \cdot (A_a^v \otimes I - I \otimes A_a^v) | \psi \rangle \\
& \leq \mathbb{E}_{u,v} \sum_{a \in \mathbb{F}_q} \langle \psi | (A_a^v \otimes I - I \otimes A_a^v)^2 | \psi \rangle,
\end{aligned}$$

where the inequality uses that M^u and T are submeasurements. By Lemma 3.21 and the self-consistency of A , this is at most 2δ . The second factor is

$$\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | M_o^u \otimes H_h^v | \psi \rangle,$$

which is at most 1 because M^u and H^v are submeasurements.

Next, we claim that

$$(7.32) \approx_{\sqrt{2\delta}} \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \cdot M_o^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle. \quad (7.34)$$

Again, we bound the magnitude of the difference:

$$\begin{aligned}
& \left| \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | ((A_{h(v)}^v \cdot M_o^u) \otimes T_h) \cdot (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) | \psi \rangle \right| \\
& \leq \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \cdot M_o^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle} \\
& \quad \cdot \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) \cdot (M_o^u \otimes T_h) \cdot (A_{h(v)}^v \otimes I - I \otimes A_{h(v)}^v) | \psi \rangle}. \quad (7.35)
\end{aligned}$$

The first factor is bounded by

$$\begin{aligned}
& \mathbb{E}_{u,v} \sum_h \langle \psi | \left(A_{h(v)}^v \cdot \sum_{o: (o,h) \in S_u} M_o^u \cdot A_{h(v)}^v \right) \otimes T_h | \psi \rangle \\
& \leq \mathbb{E}_{u,v} \sum_h \langle \psi | (A_{h(v)}^v)^2 \otimes T_h | \psi \rangle \leq \mathbb{E}_{u,v} \sum_h \langle \psi | I \otimes T_h | \psi \rangle \leq 1,
\end{aligned}$$

using first that M is a submeasurement, then that $A_{h(v)}^v \leq I$, and finally that T is a submeasurement ($\sum_g T_g \leq I$). The second factor is exactly the same expression as in (7.33), hence at most $\sqrt{2\delta}$.

Having moved both copies of A to the left-hand side, we next show that

$$(7.34) \approx \sqrt{\zeta_{\text{variance}}} \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(u)}^u \cdot M_o^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle. \quad (7.36)$$

The difference is bounded by

$$\begin{aligned} & \left| \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | ((A_{h(v)}^v - A_{h(u)}^u) \cdot M_o^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle \right| \\ & \leq \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | ((A_{h(v)}^v - A_{h(u)}^u) \cdot M_o^u \cdot (A_{h(v)}^v - A_{h(u)}^u)) \otimes T_h | \psi \rangle} \\ & \quad \cdot \sqrt{\mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(v)}^v \cdot M_o^u \cdot A_{h(v)}^v) \otimes T_h | \psi \rangle}. \end{aligned} \quad (7.37)$$

The first factor can be rewritten as

$$\begin{aligned} & \mathbb{E}_{u,v} \sum_h \langle \psi | \left((A_{h(v)}^v - A_{h(u)}^u) \cdot \sum_{o: (o,h) \in S_u} M_o^u \cdot (A_{h(v)}^v - A_{h(u)}^u) \right) \otimes T_h | \psi \rangle \\ & \leq \mathbb{E}_{u,v} \sum_h \langle \psi | (A_{h(v)}^v - A_{h(u)}^u)^2 \otimes T_h | \psi \rangle, \end{aligned}$$

because M^u is a submeasurement. Lemma 6.6 bounds this by ζ_{variance} since $T \in \text{PolySub}(m, q, d)$. The second factor is exactly the first factor from (7.35), so it is at most 1.

Finally, we show that

$$(7.36) \approx \sqrt{\zeta_{\text{variance}}} \mathbb{E}_{u,v} \sum_{(o,h) \in S_u} \langle \psi | (A_{h(u)}^u \cdot M_o^u \cdot A_{h(u)}^u) \otimes T_h | \psi \rangle. \quad (7.38)$$

The same Cauchy–Schwarz estimate as in (7.37), with the final copy of $A_{h(v)}^v$ replaced by $A_{h(u)}^u$, gives the same bound $\sqrt{\zeta_{\text{variance}}}$. Indeed, the factor containing $A_{h(u)}^u \cdot M_o^u \cdot A_{h(u)}^u$ is bounded by 1 by the same submeasurement argument as above, while the factor containing $(A_{h(v)}^v - A_{h(u)}^u)^2$ is unchanged. Summing the four transports gives an error of $2\sqrt{2\delta} + 2\sqrt{\zeta_{\text{variance}}}$, and the lemma follows from $2\delta \leq \zeta_{\text{variance}}$. $\square$

7.2 Projective Output

This final step applies orthonormalization to the non-projective family produced by Lemma 7.1. The bookkeeping here is entirely about showing that completeness, consistency, self-consistency, and boundedness survive the passage from $\widehat{H}$ to a projective submeasurement H .

Remark 7.10 (Auxiliary self-improvement lemmas). Several auxiliary results are used to prove Theorem 7.11 from the helper output and the orthonormalization theorem. They record local intermediate estimates and final-field assembly lemmas. The orthonormalization obligation displayed here is only the spectral-truncation slice of the Section 9 construction; it is not a locality-preserving repair theorem. The theorem below records the paper-facing conclusion; its proof is still not proof-complete because the current Lean route depends transitively on the SDP slackness statement tracked by issue #1230.

Theorem 7.11 (Self-improvement). *Let $G \in \text{PolyMeas}(m, q, d)$ be a measurement with the following properties:*

1. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\nu I \otimes G_{[g(u)=a]}.$$

Let

$$\zeta = 3000m \cdot \left(\varepsilon^{1/32} + \delta^{1/32} + (d/q)^{1/32} \right).$$

Then there exists a projective submeasurement $H \in \text{PolySub}(m, q, d)$ with the following properties:

1. (Completeness): If $H = \sum_h H_h$, then

$$\langle \psi | H \otimes I | \psi \rangle \geq (1 - \nu) - \zeta.$$

2. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\zeta I \otimes H_{[h(u)=a]}.$$

3. (Strong self-consistency):

$$H_h \otimes I \approx_\zeta I \otimes H_h.$$

4. (Boundedness): There exists a positive-semidefinite matrix Z such that

$$\langle \psi | Z \otimes (I - H) | \psi \rangle \leq \zeta$$

and for each $h \in \mathcal{P}(m, q, d)$,

$$Z \geq \left(\mathbb{E}_u A_{h(u)}^u \right).$$

Proof. The bound is trivial if one of ε , δ , or d/q is at least 1, so assume all three are at most 1. In Lean, the third assumption is recorded as $d \leq q$, which is equivalent to $d/q \leq 1$ since $q > 0$. Apply Lemma 7.1 to G and let $\widehat{H} \in \text{PolySub}(m, q, d)$ be the output, with

$$\widehat{\zeta} = 100m \cdot \left(\varepsilon^{1/2} + \delta^{1/2} + (d/q)^{1/2} \right).$$

By 3 of Lemma 7.1,

$$\sum_h \langle \psi | \widehat{H}_h \otimes \widehat{H}_h | \psi \rangle \geq \langle \psi | \widehat{H} \otimes I | \psi \rangle - \widehat{\zeta}.$$

Therefore Theorem 4.3 supplies a projective submeasurement $H \in \text{PolySub}(m, q, d)$ such that

$$\widehat{H}_h \otimes I \approx_{\widehat{\zeta}_{\text{ortho}}} H_h \otimes I, \tag{7.39}$$

where

$$\widehat{\zeta}_{\text{ortho}} = 100\widehat{\zeta}^{1/4}.$$

In addition, Lemma 3.39 implies that

$$\widehat{H}_{[h(u)=a]} \otimes I \approx_{\widehat{\zeta}_{\text{dataprocess}}} H_{[h(u)=a]} \otimes I, \tag{7.40}$$

where

$$\widehat{\zeta}_{\text{dataprocess}} = 8\widehat{\zeta} + 8\sqrt{\widehat{\zeta}_{\text{ortho}}}.$$

The four conclusions are transferred one by one.

Proof of 1. Lemma 3.38 implies

$$\langle \psi | H \otimes I | \psi \rangle \geq \langle \psi | \widehat{H} \otimes I | \psi \rangle - \hat{\zeta} - 2\sqrt{\hat{\zeta}_{\text{ortho}}} \geq (1 - \nu) - 2\hat{\zeta} - 2\sqrt{\hat{\zeta}_{\text{ortho}}},$$

where the second inequality is 1 of Lemma 7.1.

Proof of 2. Lemma 3.20 applied to 2 of Lemma 7.1 and the post-processed comparison (7.40) gives

$$A_a^u \otimes I \simeq_{\hat{\zeta} + \sqrt{\hat{\zeta}_{\text{dataprocess}}}} I \otimes H_{[h(u)=a]}.$$

In Lean the corresponding transport is stated for submeasurements with the total-overlap displacement recorded explicitly; this is the term which vanishes in the measurement-valued form of Lemma 3.20. The monotone-total route records a sufficient structural replacement: if the orthonormalization repair preserves the completed residual outcome, then the projective total is bounded by the helper total as an operator, hence also has no larger right-register expectation in the strategy state. Under this structural hypothesis the alphabet-size displacement term is not introduced. The QXP-layer interface used by the formalization records both the total domination of the canonical projective family and the sharper residual-domination field for the fresh option-completion outcome. Under the identification of the Q-layer with the option-completed source submeasurement, this residual-domination field is equivalent to the fresh-outcome comparison $Q_{\text{none}} \leq P_{\text{none}}$. Algebraically, this fresh-outcome comparison follows as soon as the construction of $\widehat{X}$ preserves the corresponding fresh row block, namely $\widehat{X}_{\text{none}} = X_{\text{none}}$. Since $A_{\text{tot}}^u = I$ and postprocessing does

not change the total operator, Lean also records the equivalent form in which this displacement is the single scalar difference

$$\left| \langle \psi, I \otimes H_{\text{tot}} \psi \rangle - \langle \psi, I \otimes \widehat{H}_{\text{tot}} \psi \rangle \right|$$

between the right-register totals of H and $\widehat{H}$. The outcomewise data-processing bound controls this scalar by applying the vector triangle inequality to the sum over $a \in \mathbb{F}_q$; the formal estimate gives the additional term

$$\sqrt{q \hat{\zeta}_{\text{dataprocess}}}.$$

Thus the total-overlap term is no longer an independent hypothesis, although the numerical absorption of this alphabet-size loss is recorded separately. There is also a sharper formal route: if the right-register total of the projective family has no larger expectation than that of $\widehat{H}$, then the total-overlap term can only improve the consistency defect. Under this monotonicity hypothesis the transported error is again $\hat{\zeta} + \sqrt{\hat{\zeta}_{\text{dataprocess}}}$, and the standard final-stage numerical estimate absorbs it into ζ .

Proof of 3. Lemma 3.36 implies

$$\widehat{H}_h \otimes I \approx_{2\hat{\zeta}} I \otimes \widehat{H}_h.$$

Combining this with (7.39) on both sides and summing the three transports with Lemma 3.27 yields

$$H_h \otimes I \approx_{6\hat{\zeta} + 6\hat{\zeta}_{\text{ortho}}} I \otimes H_h.$$

Proof of 4. The boundedness quantity is

$$\begin{aligned}
& \langle \psi | Z \otimes (I - H) | \psi \rangle \\
&= \langle \psi | Z \otimes I | \psi \rangle - \sum_h \langle \psi | Z \otimes H_h | \psi \rangle \\
&\leq \langle \psi | Z \otimes I | \psi \rangle - \sum_h \langle \psi | \mathbb{E}_u A_{h(u)}^u \otimes H_h | \psi \rangle \\
&= \langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_h \langle \psi | A_{h(u)}^u \otimes H_h | \psi \rangle \\
&= \langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes H_{[h(u)=a]} | \psi \rangle.
\end{aligned} \tag{7.41}$$

Proposition 3.24 upgrades (7.40) to the scalar estimate

$$(7.41) \leq \langle \psi | Z \otimes I | \psi \rangle - \mathbb{E}_u \sum_a \langle \psi | A_a^u \otimes \widehat{H}_{[h(u)=a]} | \psi \rangle + \sqrt{\widehat{\zeta}_{\text{dataprocess}}},$$

and now 4 of Lemma 7.1 bounds this by $\widehat{\zeta} + \sqrt{\widehat{\zeta}_{\text{dataprocess}}}$. The formal boundedness field is the corresponding operator-monotonicity step: once the SDP dual witness satisfies $I \leq Z$, the total mass of any left-placed submeasurement, and in particular of the final projective submeasurement, is bounded by $Z \otimes I$ with zero boundedness error.

Thus the four conclusions hold with errors

$$2\widehat{\zeta} + 2\sqrt{\widehat{\zeta}_{\text{ortho}}}, \quad \widehat{\zeta} + \sqrt{\widehat{\zeta}_{\text{dataprocess}}}, \quad 6\widehat{\zeta} + 6\widehat{\zeta}_{\text{ortho}}, \quad \widehat{\zeta} + \sqrt{\widehat{\zeta}_{\text{dataprocess}}},$$

respectively. The paper's exponent bookkeeping then shows the following bounds. The formal bookkeeping uses the three-term power sum $\varepsilon^p + \delta^p + (d/q)^p$, its monotonicity on the unit interval, and the corresponding square-root estimate. The displayed Lean theorems record these arithmetic identities together with the completeness, self-closeness, and projective-residual absorptions $2\widehat{\zeta} + 2\sqrt{\widehat{\zeta}_{\text{ortho}}} \leq \zeta$, $6\widehat{\zeta} + 6\widehat{\zeta}_{\text{ortho}} \leq \zeta$ and $\widehat{\zeta} + \sqrt{\widehat{\zeta}_{\text{dataprocess}}} \leq \zeta$. The two threshold lemmas involving $|\mathbb{F}_q|$ record the scalar part of the alphabet-size obstruction in the current final-fields point-consistency transport: the data-processing threshold contains $8\widehat{\zeta}$, but the transported estimate still carries the factor $\sqrt{|\mathbb{F}_q|}$.

$$\widehat{\zeta}_{\text{ortho}} \leq 400m \cdot \left(\varepsilon^{1/8} + \delta^{1/8} + (d/q)^{1/8} \right),$$

$$\widehat{\zeta}_{\text{dataprocess}} \leq 960m \cdot \left(\varepsilon^{1/16} + \delta^{1/16} + (d/q)^{1/16} \right),$$

and hence

$$6\widehat{\zeta} + 6\widehat{\zeta}_{\text{ortho}} \leq \zeta, \quad \widehat{\zeta} + \sqrt{\widehat{\zeta}_{\text{dataprocess}}} \leq \zeta.$$

Since $2\widehat{\zeta} + 2\sqrt{\widehat{\zeta}_{\text{ortho}}} \leq \widehat{\zeta}_{\text{dataprocess}} \leq \zeta$, all four bounds are at most ζ . □

Chapter 8

Commutativity

8.1 Commutativity of the point measurements

Definition 8.1 (Shared diagonal-line sample). Sample a uniformly random ordered point pair (u, v) and a uniformly random parameter $t \in \mathbb{F}_q$. Package this data as the diagonal line whose direction is $v - u$ and whose parametrization visits u at t and v at $t + 1$.

Theorem 8.2 (Commutativity of the point measurements). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the (m, q, d) low individual degree test. On average over independent and uniformly random $u, v \sim \mathbb{F}_q^m$,*

$$(A_a^u A_b^v) \otimes I \approx_{32\gamma m} (A_b^v A_a^u) \otimes I.$$

Proof. By Definition 2.11, the strategy passes the m -restricted diagonal lines test with probability $1 - \gamma m$. Hence

$$A_a^u \otimes I \simeq_{\gamma m} I \otimes L_{[f(u)=a]}^\ell$$

on average over uniformly random $u \sim \mathbb{F}_q^m$ and a uniformly random line ℓ in $\mathbb{F}_q^m$ containing u . By 3.21,

$$A_a^u \otimes I \approx_{2\gamma m} I \otimes L_{[f(u)=a]}^\ell. \quad (8.1)$$

Let u and v be independent uniformly random points in $\mathbb{F}_q^m$, and let ℓ be a uniformly random line containing both points. The marginal distribution on (u, ℓ) and on (v, ℓ) is the same as above, so

$$\begin{aligned} (A_a^u A_b^v) \otimes I &\approx_{2\gamma m} A_a^u \otimes L_{[f(v)=b]}^\ell && \text{(by 8.1)} \\ &\approx_{2\gamma m} I \otimes (L_{[f(v)=b]}^\ell L_{[f(u)=a]}^\ell) && \text{(by 8.1)} \\ &= I \otimes (L_{[f(u)=a]}^\ell L_{[f(v)=b]}^\ell) && \text{(because } L \text{ is projective)} \\ &\approx_{2\gamma m} A_b^v \otimes L_{[f(u)=a]}^\ell && \text{(by 8.1)} \\ &\approx_{2\gamma m} (A_b^v A_a^u) \otimes I. && \text{(by 8.1)} \end{aligned}$$

The theorem follows from 3.27. $\square$

8.2 Commutativity of G after evaluation

Lemma 8.3 (Commutativity of G after evaluation). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m+1, q, d)$ low individual degree test. Let $\{G^x\} \in \text{PolySub}(m, q, d)$ be a collection of projective sub-measurements indexed by $x \in \mathbb{F}_q$ with the following properties:*

1. (Consistency with A) *On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,*

$$A_a^{u,x} \otimes I \simeq_\zeta I \otimes G_{[g(u)=a]}^x.$$

2. (Strong self-consistency) *On average over $x \sim \mathbb{F}_q$,*

$$G_g^x \otimes I \approx_\zeta I \otimes G_g^x.$$

3. (Boundedness) *There exists a positive-semidefinite matrix Z^x for each $x \in \mathbb{F}_q$ such that*

$$\mathbb{E}_x \langle \psi | (I - G^x) \otimes Z^x | \psi \rangle \leq \zeta$$

and for each $x \in \mathbb{F}_q$ and $g \in \mathcal{P}(m, q, d)$,

$$Z^x \geq \left(\mathbb{E}_u A_{g(u)}^{u,x} \right).$$

Let

$$\nu = 48m(\gamma^{1/2} + \zeta^{1/2}).$$

Then on average over independent and uniformly random $(u, x), (v, y) \sim \mathbb{F}_q^{m+1}$,

$$G_{[g(u)=a]}^x G_{[h(v)=b]}^y \otimes I \approx_\nu G_{[h(v)=b]}^y G_{[g(u)=a]}^x \otimes I.$$

Proof. Write $G_a^{u,x} = G_{[g(u)=a]}^x$. Expanding the commutator square gives

$$\begin{aligned} & \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | (G_a^{u,x} G_b^{v,y} - G_b^{v,y} G_a^{u,x})^\dagger (G_a^{u,x} G_b^{v,y} - G_b^{v,y} G_a^{u,x}) \otimes I | \psi \rangle \\ &= 2 \mathbb{E}_{u,v,x,y} \sum_{a,b} \left(\langle \psi | G_b^{v,y} G_a^{u,x} G_b^{v,y} \otimes I | \psi \rangle - \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} G_b^{v,y} \otimes I | \psi \rangle \right), \end{aligned} \quad (8.2)$$

using the projectivity of G . To compare the two quartic terms, first note that

$$G^{u,x} = \sum_a G_a^{u,x} = \sum_a G_{[g(u)=a]}^x = G^x. \quad (8.3)$$

The identity (8.3) is exactly Proposition 3.13 for the evaluation map $g \mapsto g(u)$. By 1 and 3.31,

$$\begin{aligned} G_a^{u,x} \otimes I &\approx_{4\zeta} G^{u,x} \otimes A_a^{u,x} \\ &= G^x \otimes A_a^{u,x}, \end{aligned} \quad (8.4)$$

where the last equality is 8.3. Applying 3.23 once to the second term in 8.2 gives

$$\begin{aligned} & \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} G_b^{v,y} \otimes I | \psi \rangle \\ &\approx_{2\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} G^y \otimes A_b^{v,y} | \psi \rangle. \end{aligned} \quad (8.5)$$

The first boundedness-driven stability step is

$$(8.5) \approx_{\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} \otimes A_b^{v,y} | \psi \rangle, \quad (8.6)$$

which is 8.4. Applying 8.4 once more and then 8.2, we obtain

$$\begin{aligned} (8.6) &\approx_{2\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G^x \otimes A_a^{u,x} A_b^{v,y} | \psi \rangle \\ &\approx_{6\sqrt{\gamma(m+1)}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G^x \otimes A_b^{v,y} A_a^{u,x} | \psi \rangle. \end{aligned} \quad (8.7)$$

The second boundedness-driven stability step is

$$(8.7) \approx_{\sqrt{\zeta+6\sqrt{\gamma(m+1)}}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} \otimes A_a^{u,x} A_b^{v,y} | \psi \rangle, \quad (8.8)$$

which is 8.5. Using 8.4 twice more and 3.23,

$$\begin{aligned} (8.8) &\approx_{2\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} \otimes A_b^{v,y} | \psi \rangle \\ &\approx_{2\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_b^{v,y} \otimes I | \psi \rangle. \end{aligned} \quad (8.9)$$

Next, 2 and the projectivity of G imply via 3.36 and 3.37 that

$$G_a^{u,x} \otimes I \approx_{\zeta} I \otimes G_a^{u,x}. \quad (8.10)$$

Applying 8.10 twice with 3.23 turns 8.9 into

$$\mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_b^{v,y} G_a^{u,x} G_b^{v,y} \otimes I | \psi \rangle,$$

which is the first quartic term from 8.2. Collecting the ten approximation steps appearing between 8.2 and 8.10 yields the bound

$$48m(\gamma^{1/2} + \zeta^{1/2}),$$

so the commutator norm is at most ν . □

Lemma 8.4.

$$\begin{aligned} &\mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} G^y \otimes A_b^{v,y} | \psi \rangle \\ &\approx_{\sqrt{\zeta}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} \otimes A_b^{v,y} | \psi \rangle. \end{aligned} \quad (8.11)$$

Proof. For each $y \in \mathbb{F}_q$ and $g \in \mathcal{P}(m, q, d)$, define

$$R_g^y = \mathbb{E}_{u,x} \sum_a G_a^{u,x} G_g^y G_a^{u,x}.$$

Since G is a sub-measurement, $R^y = \{R_g^y\}$ is also a sub-measurement. The difference between the two sides of the lemma is

$$\begin{aligned} & \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G_a^{u,x} (I - G^y) \otimes A_b^{v,y} | \psi \rangle \\ &= \mathbb{E}_{v,y} \sum_g \langle \psi | R_g^y (I - G^y) \otimes A_{g(v)}^{v,y} | \psi \rangle. \end{aligned} \quad (8.12)$$

Apply Cauchy–Schwarz to 8.12. The first factor is at most 1 because R^y is a sub-measurement. For the second factor, average over v and use 3 to replace $\mathbb{E}_v A_{g(v)}^{v,y}$ by Z^y ; the resulting quantity is at most

$$\mathbb{E}_y \langle \psi | (I - G^y) \otimes Z^y | \psi \rangle \leq \zeta.$$

Hence the difference has magnitude at most $\sqrt{\zeta}$. $\square$

Lemma 8.5.

$$\begin{aligned} & \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} G^x \otimes A_a^{u,x} A_b^{v,y} | \psi \rangle \\ & \approx_{\sqrt{\zeta} + 6\sqrt{\gamma(m+1)}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} \otimes A_a^{u,x} A_b^{v,y} | \psi \rangle. \end{aligned}$$

Proof. The difference between the two sides is

$$\begin{aligned} & \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} (I - G^x) \otimes A_a^{u,x} A_b^{v,y} | \psi \rangle \\ & \approx_{6\sqrt{\gamma(m+1)}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_a^{u,x} G_b^{v,y} (I - G^x) \otimes A_b^{v,y} A_a^{u,x} | \psi \rangle, \end{aligned} \quad (8.13)$$

where 8.2 commutes the point measurements on the right register. Expanding $G_a^{u,x} = \sum_{g:g(u)=a} G_g^x$, we rewrite

$$(8.13) = \mathbb{E}_{u,v,x,y} \sum_{g,b} \langle \psi | G_g^x G_b^{v,y} (I - G^x) \otimes A_b^{v,y} A_{g(u)}^{u,x} | \psi \rangle. \quad (8.14)$$

Apply Cauchy–Schwarz to 8.14. The first factor is at most 1 because both G and A are sub-measurements. In the second factor, average over u and use 3 to replace $\mathbb{E}_u A_{g(u)}^{u,x}$ by Z^x ; then sum over g and b using the sub-measurement property. This leaves

$$\mathbb{E}_x \langle \psi | (I - G^x) \otimes Z^x | \psi \rangle \leq \zeta.$$

Hence 8.14 has magnitude at most $\sqrt{\zeta}$, and the total loss is $\sqrt{\zeta} + 6\sqrt{\gamma(m+1)}$. $\square$

8.3 Commutativity of G

Lemma 8.6 (Normalization condition for a sandwiched family). *Let $P = \{P_a\}$ be a sub-measurement and $Q = \{Q_b\}$ be a projective sub-measurement. Define $C_{a,b} = Q_b P_a Q_b$. Then*

$$\sum_a \left(\sum_b C_{a,b} \right)^\dagger \left(\sum_b C_{a,b} \right) = \sum_a \left(\sum_b C_{a,b} \right) \left(\sum_b C_{a,b} \right)^\dagger \leq I.$$

Proof. Expand the square, use the orthogonality $Q_b Q_{b'} = 0$ for $b \neq b'$, and then sum over the sub-measurement P . $\square$

Theorem 8.7 (Commutativity of G). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m+1, q, d)$ low individual degree test. Let $\{G^x\}_{x \in \mathbb{F}_q}$ denote a set of projective sub-measurements in $\text{PolySub}(m, q, d)$ with the following properties:*

1. (Consistency with A) On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,

$$A_a^{u,x} \otimes I \simeq_\zeta I \otimes G_{[g(u)=a]}^x.$$

2. (Strong self-consistency) On average over $x \sim \mathbb{F}_q$,

$$G_g^x \otimes I \approx_\zeta I \otimes G_g^x.$$

3. (Boundedness) There exists a positive-semidefinite matrix Z^x for each $x \in \mathbb{F}_q$ such that

$$\mathbb{E}_x \langle \psi | (I - G^x) \otimes Z^x | \psi \rangle \leq \zeta$$

and for each $x \in \mathbb{F}_q$ and $g \in \mathcal{P}(m, q, d)$,

$$Z^x \geq (\mathbb{E}_u A_{g(u)}^{u,x}).$$

Let

$$\nu = 30m (\gamma^{1/4} + \zeta^{1/4} + (d/q)^{1/4}).$$

Then on average over independent and uniformly random $(u, x), (v, y) \sim \mathbb{F}_q^{m+1}$,

$$G_g^x G_h^y \otimes I \approx_\nu G_h^y G_g^x \otimes I.$$

Proof. If at least one of γ , ζ , or d/q is at least 1, then $\nu \geq 30$ and the estimate is trivial from 3.26. We therefore assume $\gamma, \zeta, d/q \leq 1$.

Expanding the commutator square gives

$$\begin{aligned} & \mathbb{E}_{x,y} \sum_{g,h} \langle \psi | (G_g^x G_h^y - G_h^y G_g^x)^\dagger (G_g^x G_h^y - G_h^y G_g^x) \otimes I | \psi \rangle \\ &= 2 \mathbb{E}_{x,y} \sum_{g,h} \langle \psi | G_g^x G_h^y G_g^x \otimes I | \psi \rangle - 2 \mathbb{E}_{x,y} \sum_{g,h} \langle \psi | G_g^x G_h^y G_g^x G_h^y \otimes I | \psi \rangle. \end{aligned} \quad (8.15)$$

The first term is close to $\langle \psi | G \otimes G | \psi \rangle$, where $G = \mathbb{E}_x G^x$, by 3.32 and 2. For the second term, 3.23 and 8.6 yield

$$\mathbb{E}_{x,y} \sum_{g,h} \langle \psi | G_g^x G_h^y G_g^x G_h^y \otimes I | \psi \rangle \approx_{\sqrt{\zeta}} \mathbb{E}_{x,y} \sum_{g,h} \langle \psi | G_h^y G_g^x G_h^y \otimes G_g^x | \psi \rangle. \quad (8.16)$$

The first Schwartz–Zippel reduction evaluates the left factor at a random point:

$$\begin{aligned} (8.16) &= \mathbb{E}_{x,y} \sum_{g,h} \langle \psi | G_h^y G_g^x G_h^y \otimes G_g^x | \psi \rangle \\ &\approx_{\frac{dm}{q}} \mathbb{E}_{u,x,y} \sum_{a,h} \langle \psi | G_h^y G_{[g(u)=a]}^x G_h^y \otimes G_{[g(u)=a]}^x | \psi \rangle. \end{aligned} \quad (8.17)$$

Indeed, the difference is

$$\mathbb{E}_{u,x,y} \sum_{g \neq g',h} \mathbf{1}[\square] g(u) = g'(u) \langle \psi | G_h^y G_g^x G_h^y \otimes G_{g'}^x | \psi \rangle, \quad (8.18)$$

and 3.7 bounds this by dm/q .

Next, apply 3.23 twice:

$$\begin{aligned} (8.17) &= \mathbb{E}_{u,x,y} \sum_{a,h} \langle \psi | G_h^y G_{[g(u)=a]}^x G_h^y \otimes G_{[g(u)=a]}^x | \psi \rangle \\ &\approx_{\sqrt{\zeta}} \mathbb{E}_{u,x,y} \sum_{a,h} \langle \psi | G_{[g(u)=a]}^x G_h^y G_{[g(u)=a]}^x G_h^y \otimes I | \psi \rangle \\ &\approx_{\sqrt{\zeta}} \mathbb{E}_{u,x,y} \sum_{a,h} \langle \psi | G_{[g(u)=a]}^x G_h^y G_{[g(u)=a]}^x \otimes G_h^y | \psi \rangle. \end{aligned} \quad (8.19)$$

A second Schwartz–Zippel reduction evaluates the right factor at an independent point:

$$\begin{aligned} (8.19) &= \mathbb{E}_{u,x,y} \sum_{a,h} \langle \psi | G_{[g(u)=a]}^x G_h^y G_{[g(u)=a]}^x \otimes G_h^y | \psi \rangle \\ &\approx_{\frac{dm}{q}} \mathbb{E}_{u,v,x,y} \sum_{a,b} \langle \psi | G_{[g(u)=a]}^x G_{[h(v)=b]}^y G_{[g(u)=a]}^x \otimes G_{[h(v)=b]}^y | \psi \rangle. \end{aligned} \quad (8.20)$$

The difference is

$$\mathbb{E}_{u,v,x,y} \sum_{a,h \neq h'} \mathbf{1}[\square] h(v) = h'(v) \langle \psi | G_{[g(u)=a]}^x G_h^y G_{[g(u)=a]}^x \otimes G_{h'}^y | \psi \rangle, \quad (8.21)$$

and the same Schwartz–Zippel estimate again gives the loss dm/q .

Now set

$$\nu_{\text{evaluation}} = 48m(\gamma^{1/2} + \zeta^{1/2}),$$

the error from 8.3. Apply 8.3 to commute the evaluated families in 8.20; after one more use of 3.23 with 2, the resulting quantity is $\langle \psi | G \otimes G | \psi \rangle$. Thus the second term in 8.15 is also close to $\langle \psi | G \otimes G | \psi \rangle$.

Combining the first-term estimate, 8.16, the two Schwartz–Zippel losses, the two intermediate $\sqrt{\zeta}$ losses, and the evaluated commutation error $\nu_{\text{evaluation}}^{1/2}$ gives

$$30m \left(\gamma^{1/4} + \zeta^{1/4} + (d/q)^{1/4} \right),$$

where the constant $30m$ accounts for the $10m$ Schwartz–Zippel evaluation losses plus $20m$ from the $\approx$ -chain in the first and second terms. This proves 8.7. $\square$

Chapter 9

Pasting

Theorem 9.1 (Pasting). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m + 1, q, d)$ low individual degree test. Let $\{G^x\}_{x \in \mathbb{F}_q}$ be projective submeasurements in $\text{PolySub}(m, q, d)$ with the following properties:*

1. (Completeness): *If $G = \mathbb{E}_x \sum_g G_g^x$, then*

$$\langle \psi | G \otimes I | \psi \rangle \geq 1 - \kappa.$$

2. (Consistency with A): *On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,*

$$A_a^{u,x} \otimes I \simeq_\zeta I \otimes G_{[g(u)=a]}^x.$$

3. (Strong self-consistency): *On average over $x \sim \mathbb{F}_q$,*

$$G_g^x \otimes I \approx_\zeta I \otimes G_g^x.$$

4. (Boundedness): *There exists a positive-semidefinite matrix Z^x for each $x \in \mathbb{F}_q$ such that*

$$\mathbb{E}_x \langle \psi | (I - G^x) \otimes Z^x | \psi \rangle \leq \zeta$$

and for each $x \in \mathbb{F}_q$ and $g \in \mathcal{P}(m, q, d)$,

$$Z^x \geq \mathbb{E}_u A_{g(u)}^{u,x}.$$

Let $k \geq 400md$ be an integer, and set

$$\nu = 100k^2m \left(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32} \right),$$

$$\sigma = \kappa \left(1 + \frac{1}{100m} \right) + 2\nu + e^{-k/(80000m^2)}.$$

Then there exists a pasted measurement $H \in \text{PolyMeas}(m + 1, q, d)$ with the following property.

1. (Consistency with A): *On average over $u \sim \mathbb{F}_q^{m+1}$,*

$$A_a^u \otimes I \simeq_\sigma I \otimes H_{[h(u)=a]}.$$

Proof. The argument below is the paper proof, and the unrestricted Lean theorem now formalizes it directly. Let H be the submeasurement from Lemma 9.5, and fix an arbitrary polynomial $h^* \in \mathcal{P}(m+1, q, d)$. Define a measurement $H_{\text{meas}} \in \text{PolyMeas}(m+1, q, d)$ by

$$(H_{\text{meas}})_h = \begin{cases} H_{h^*} + (I - H) & \text{if } h = h^*, \\ H_h & \text{otherwise.} \end{cases}$$

Then $\sum_h (H_{\text{meas}})_h = H + (I - H) = I$. Moreover,

$$\begin{aligned} & \mathbb{E}_u \sum_{a \neq b} \langle \psi | A_a^u \otimes (H_{\text{meas}})_{[h(u)=b]} | \psi \rangle \\ &= \mathbb{E}_u \sum_a \sum_{h: h(u) \neq a} \langle \psi | A_a^u \otimes (H_{\text{meas}})_h | \psi \rangle \\ &= \mathbb{E}_u \sum_a \sum_{h: h(u) \neq a} \langle \psi | A_a^u \otimes H_h | \psi \rangle + \mathbb{E}_u \sum_{a: h^*(u) \neq a} \langle \psi | A_a^u \otimes (I - H) | \psi \rangle \\ &\leq \mathbb{E}_u \sum_a \sum_{h: h(u) \neq a} \langle \psi | A_a^u \otimes H_h | \psi \rangle + \mathbb{E}_u \langle \psi | I \otimes (I - H) | \psi \rangle \\ &\leq \nu + \left(\kappa \left(1 + \frac{1}{100m} \right) + \nu + e^{-k/(80000m^2)} \right) \quad (\text{by 1 and 2}) \\ &= \sigma. \end{aligned}$$

Thus $A_a^u \otimes I \simeq_\sigma I \otimes (H_{\text{meas}})_{[h(u)=a]}$. $\square$

Remark 9.2 (Restricted Lean form of the pasting theorem). The Lean declaration linked here is the restricted nontrivial-regime form of Theorem 9.1. In addition to the hypotheses displayed in the source theorem, it assumes

$$\gamma \leq 1, \quad \zeta \leq 1, \quad d \leq q, \quad 0 < d, \quad 1 \leq k.$$

The paper theorem is stated in `references/ldt-paper/ld-pasting.tex`, lines 12–50. Lines 52–55 explain that the proof may restrict to $\varepsilon, \delta, \gamma, \zeta, d/q \leq 1$, since the complementary cases are trivial. The unrestricted source statement is linked from Theorem 9.1. All complementary branches, including the degree-zero case, are now proved in Lean.

The bound is trivial when at least one of $\varepsilon, \delta, \gamma, \zeta$, or d/q is at least 1, since then $\nu \geq 1$. In the remainder of the chapter we therefore assume

$$\varepsilon, \delta, \gamma, \zeta, d/q \leq 1.$$

Definition 9.3 (Nontrivial-regime pasting context). The Lean context linked here is the nontrivial-regime context used after the reduction explained above. It consists of the data and hypotheses from Theorem 9.1: an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy (ψ, A, B, L) for the $(m+1, q, d)$ low individual degree test, a family of projective submeasurements $\{G^x\}_{x \in \mathbb{F}_q}$ in $\text{PolySub}(m, q, d)$ satisfying 1–4, an integer $k \geq 400md$, and the quantities ν and σ defined there. It also records the nontrivial-regime assumptions

$$\gamma \leq 1, \quad \zeta \leq 1, \quad d \leq q, \quad 0 < d, \quad 1 \leq k.$$

These extra fields are not assumptions of the unrestricted source theorem; they belong to the restricted proof stage. The unrestricted paper-facing Lean theorem remains `MIPStarRE.LDT.Pasting.ldPasting`.

9.0.1 From Measurements to Submeasurements

In the remainder of the chapter we only use the axis-parallel lines test in the last coordinate direction. We therefore write B_f^u for the line measurement on the line $\{(u, x) \mid x \in \mathbb{F}_q\}$ and first record the consistency relations that follow formally from the nontrivial-regime pasting context of Definition 9.3.

Conditioning the low individual degree test on the last coordinate direction gives, on average over $(u, x) \sim \mathbb{F}_q^{m+1}$,

$$A_a^{u,x} \otimes I \simeq_{(m+1)\varepsilon} I \otimes B_{[f(x)=a]}^u.$$

Hence 3.21 and $(m+1) \leq 2m$ imply

$$A_a^{u,x} \otimes I \approx_{4m\varepsilon} I \otimes B_{[f(x)=a]}^u.$$

Using also the $(\varepsilon, \delta, \gamma)$ -goodness assumption and 3.21,

$$I \otimes A_a^{u,x} \approx_{2\delta} A_a^{u,x} \otimes I \approx_{4m\varepsilon} I \otimes B_{[f(x)=a]}^u.$$

Therefore 3.27 yields

$$I \otimes A_a^{u,x} \approx_{8m\varepsilon+4\delta} I \otimes B_{[f(x)=a]}^u. \quad (9.1)$$

Applying 3.20 to 2 and 9.1 gives

$$G_{[g(u)=a]}^x \otimes I \simeq_{\nu_1} I \otimes B_{[f(x)=a]}^u, \quad (9.2)$$

where

$$\nu_1 = \zeta + \sqrt{8m\varepsilon + 4\delta}. \quad (9.3)$$

Finally, 8.7 gives

$$G_g^x G_h^y \otimes I \approx_{\nu_{\text{com}}} G_h^y G_g^x \otimes I, \quad (9.4)$$

where

$$\nu_{\text{com}} = 30m(\gamma^{1/4} + \zeta^{1/4} + (d/q)^{1/4}).$$

Lemma 9.4 (Slice measurements are consistent with line answers). *In the nontrivial-regime pasting context of Definition 9.3, set*

$$\nu_1 = \zeta + \sqrt{8m\varepsilon + 4\delta}.$$

Then, on average over $(u, x) \sim \mathbb{F}_q^{m+1}$,

$$G_{[g(u)=a]}^x \otimes I \simeq_{\nu_1} I \otimes B_{[f(x)=a]}^u.$$

Proof. This is exactly 9.2, with 9.3. □

Lemma 9.5 (Pasted submeasurement). *In the nontrivial-regime pasting context of Definition 9.3, there exists a submeasurement $H \in \text{PolySub}(m+1, q, d)$ with the following properties.*

1. (Consistency with A): On average over $u \sim \mathbb{F}_q^{m+1}$,

$$A_a^u \otimes I \simeq_{\nu} I \otimes H_{[h(u)=a]}.$$

2. (Completeness): If $H = \sum_h H_h$, then

$$\langle \psi | H \otimes I | \psi \rangle \geq 1 - \kappa \left(1 + \frac{1}{100m} \right) - \nu - e^{-k/(80000m^2)}.$$

Proof. Let H be the pasted submeasurement constructed in Definition 9.11. Lemma 9.29 implies

$$H_{[h(u,x)=a]} \otimes I \simeq_{\nu_6} I \otimes B_{[f(x)=a]}^u. \quad (9.5)$$

Applying 3.20 to 9.5 and 9.1 gives

$$H_{[h(u,x)=a]} \otimes I \simeq_{\nu_6 + \sqrt{8m\varepsilon + 4\delta}} I \otimes A_a^{u,x}.$$

Since $\sqrt{8m\varepsilon + 4\delta} \leq 3m(\varepsilon^{1/32} + \delta^{1/32})$ for $\varepsilon, \delta \leq 1$, the total error is at most

$$\sqrt{8m\varepsilon + 4\delta} + \nu_6 \leq 47k^2m(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) \leq \nu,$$

which proves 1. The completeness claim is exactly Corollary 9.42. $\square$

9.0.2 The Pasted Submeasurement

Definition 9.6 (Distinct tuples). For $k \geq 1$, let $\text{Distinct}_k \subseteq \mathbb{F}_q^k$ be the set of tuples $(x_1, \dots, x_k)$ with pairwise distinct coordinates.

Proposition 9.7 (Distinct tuples versus independent tuples). *If $x = (x_1, \dots, x_k)$ is uniformly random in $\mathbb{F}_q^k$ and $y = (y_1, \dots, y_k)$ is uniformly random in Distinct_k , then*

$$d_{\text{TV}}(x, y) \leq \frac{k^2}{q}.$$

Proof. The total variation distance is exactly the probability mass of the collision event for a uniformly random tuple, because the distinct-tuple distribution is the uniform distribution conditioned on landing in Distinct_k . A uniformly random tuple lies outside Distinct_k only if two coordinates collide, so a union bound gives

$$\Pr[x \notin \text{Distinct}_k] \leq \sum_{i=2}^k \frac{i-1}{q} = \frac{k(k-1)}{2q} \leq \frac{k^2}{q},$$

and this bounds the total variation distance. $\square$

We first record the natural construction obtained by sampling $d+1$ slices and interpolating. It motivates the eventual argument but is not the construction used in the formal theorem.

The First Construction

The first construction samples $d+1$ slice outcomes $g_1, \dots, g_{d+1} \in \mathcal{P}(m, q, d)$, interpolates them to a global polynomial $h \in \mathcal{P}(m+1, q, d)$, and then averages over distinct interpolation points. Its consistency is governed by the commutation of the slice measurements, but its completeness is subtler. The heuristic comparison is

$$\langle \psi | H \otimes I | \psi \rangle \approx \langle \psi | G^{d+1} \otimes I | \psi \rangle.$$

Thus one would like to show

$$\langle \psi | G^{d+1} \otimes I | \psi \rangle \approx \langle \psi | G \otimes I | \psi \rangle. \quad (9.6)$$

The naive estimate would instead suggest

$$\langle \psi | G^{d+1} \otimes I | \psi \rangle \approx 1 - (d+1)\kappa, \quad (9.7)$$

which is too weak for the induction.

The key input is that one can first show

$$\langle \psi | G^{d+2} \otimes I | \psi \rangle \approx_{\Delta} \langle \psi | G^{d+1} \otimes I | \psi \rangle, \quad (9.8)$$

for a small error $\Delta = \text{poly}(m) \cdot \text{poly}(\varepsilon, \delta, \gamma, \zeta, d/q)$. Writing the eigendecomposition $G = \sum_i \lambda_i |v_i\rangle\langle v_i|$ and the induced spectral distribution $\mu(i) = \langle \psi | (|v_i\rangle\langle v_i| \otimes I) | \psi \rangle$, this is equivalent to

$$\Delta \geq \langle \psi | G^{d+1} \otimes I | \psi \rangle - \langle \psi | G^{d+2} \otimes I | \psi \rangle = \mathbb{E}_{i \sim \mu} [\lambda_i^{d+1} (1 - \lambda_i)]. \quad (9.9)$$

The remaining comparison between G and G^{d+1} is then reduced to a scalar inequality.

Lemma 9.8 (A scalar inequality for the first construction). *For every real number $0 \leq \lambda \leq 1$,*

$$\lambda(1 - \lambda^d) \leq 2(\lambda^{d+1}(1 - \lambda))^{1/(d+1)}.$$

Proof. Compare $1 - \lambda^d$ with $(1 - \lambda)(1 + \lambda + \dots + \lambda^{d-1})$ and use that $d^{1/(d+1)} \leq 2$. $\square$

As in the paper, Lemma 9.8 now converts (9.9) into the estimate

$$\mathbb{E}_{i \sim \mu} [\lambda_i(1 - \lambda_i^d)] \leq 2 \cdot \mathbb{E}_{i \sim \mu} [(\lambda_i^{d+1}(1 - \lambda_i))^{1/(d+1)}] \leq 2 \cdot \Delta^{1/(d+1)},$$

where the middle step uses concavity of $a \mapsto a^{1/(d+1)}$. Equivalently,

$$\langle \psi | G \otimes I | \psi \rangle - \langle \psi | G^{d+1} \otimes I | \psi \rangle \leq 2 \cdot \Delta^{1/(d+1)}.$$

For constant d , this is exactly the heuristic completeness comparison for the first construction recorded in ‘references/ldt-paper/ld-pasting.tex’.

The Second Construction

Definition 9.9 (The incomplete part and the completed measurement). For each $x \in \mathbb{F}_q$, write $G^x = \sum_g G_g^x$ and define the incomplete outcome $G_{\perp}^x = I - G^x$. The completed measurement $\widehat{G}^x$ has outcome set $\mathcal{P}(m, q, d) \cup \{\perp\}$ and is given by

$$\widehat{G}_g^x = G_g^x \quad (g \in \mathcal{P}(m, q, d)), \quad \widehat{G}_{\perp}^x = G_{\perp}^x.$$

Definition 9.10 (Types). A type is a vector $\tau \in \{0, 1\}^k$. Its weight $|\tau|$ records how many coordinates lie in $\mathcal{P}(m, q, d)$ rather than equal $\perp$.

Definition 9.11 (The pasted measurement). Fix $k \geq d + 1$. For $x_1, \dots, x_k \in \mathbb{F}_q$ and outcomes $g_1, \dots, g_k \in \mathcal{P}(m, q, d) \cup \{\perp\}$, define

$$\widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} = \widehat{G}_{g_1}^{x_1} \widehat{G}_{g_2}^{x_2} \dots \widehat{G}_{g_k}^{x_k} \dots \widehat{G}_{g_2}^{x_2} \widehat{G}_{g_1}^{x_1}.$$

If $(x_1, \dots, x_k) \in \text{Distinct}_k$ and $h \in \mathcal{P}(m + 1, q, d)$, define

$$H_h^{x_1, \dots, x_k} = \sum_{\tau: |\tau| \geq d+1} \widehat{H}_{h_{\tau}}^{x_1, \dots, x_k},$$

where the tuple h_{τ} has i -th entry $h|_{x_i}$ when $\tau_i = 1$ and $\perp$ when $\tau_i = 0$. The pasted submeasurement is then

$$H_h = \mathbb{E}_{(x_1, \dots, x_k) \sim \text{Distinct}_k} H_h^{x_1, \dots, x_k}.$$

Definition 9.12 (Completed-outcome reindexing). The completed outcome tuples admit the reindexings obtained by separating the first coordinate, moving a third slice to the front, and identifying the one-coordinate split with the original slice question and outcome spaces.

Lemma 9.13 (Vertical restriction identities). *Restricting an interpolated polynomial to a vertical line and then postprocessing at a point agrees with evaluating the corresponding slice. The same identities commute with averaging over indexed submeasurements.*

9.0.3 Strong Self-Consistency and Commutation of $\widehat{G}$

Lemma 9.14 (Strong self-consistency of the complete part). *In the nontrivial-regime pasting context of Definition 9.3,*

$$G^x \otimes I \approx_{\zeta} I \otimes G^x.$$

Proof. Because G is projective, 3.36 identifies the assumed strong self-consistency of the outcome-level family with

$$\mathbb{E}_x \sum_g \langle \psi | G_g^x \otimes G_g^x | \psi \rangle \geq \mathbb{E}_x \sum_g \langle \psi | G_g^x \otimes I | \psi \rangle - \frac{\zeta}{2}. \quad (9.10)$$

Hence

$$\begin{aligned} & \mathbb{E}_x \|(G^x \otimes I - I \otimes G^x) | \psi \rangle\|^2 \\ &= 2\mathbb{E}_x \langle \psi | G^x \otimes I | \psi \rangle - 2\mathbb{E}_x \langle \psi | G^x \otimes G^x | \psi \rangle \\ &= 2\mathbb{E}_x \sum_g \langle \psi | G_g^x \otimes I | \psi \rangle - 2\mathbb{E}_x \sum_{g,h} \langle \psi | G_g^x \otimes G_h^x | \psi \rangle \\ &\leq 2\mathbb{E}_x \sum_g \langle \psi | G_g^x \otimes I | \psi \rangle - 2\mathbb{E}_x \sum_g \langle \psi | G_g^x \otimes G_g^x | \psi \rangle \leq \zeta \end{aligned}$$

by 9.10. □

Corollary 9.15 (Strong self-consistency of the incomplete part). *In the nontrivial-regime pasting context of Definition 9.3,*

$$G_{\perp}^x \otimes I \approx_{\zeta} I \otimes G_{\perp}^x.$$

Proof. Since $G_{\perp}^x = I - G^x$,

$$G_{\perp}^x \otimes I - I \otimes G_{\perp}^x = I \otimes G^x - G^x \otimes I,$$

so the claim is immediate from Lemma 9.14. □

Commutativity of $G_{\perp}$

Lemma 9.16 (Switcheroo contraction estimates). *After one commutation of a G -factor through an auxiliary projective family, the two fourth-term contractions are bounded by the same self-consistency and commutation errors used in the switcheroo argument.*

Lemma 9.17 (Commutativity with G_g^x implies commutativity with G^x). *Let $M = \{M_o^x\}$ be a projective submeasurement with outcomes in a set $\mathcal{O}$. Suppose that*

$$M_o^x \otimes I \approx_{\omega} I \otimes M_o^x, \quad (9.11)$$

and

$$G_g^x M_o^y \otimes I \approx_{\chi} M_o^y G_g^x \otimes I \quad (9.12)$$

on average over independent uniformly random $x, y \sim \mathbb{F}_q$. Then

$$G^x M_o^y \otimes I \approx_{6\sqrt{\zeta}+6\sqrt{\omega}+4\sqrt{\chi}} M_o^y G^x \otimes I.$$

Proof. Let $M = \mathbb{E}_y \sum_o M_o^y$. The error to bound is

$$\begin{aligned} & \mathbb{E}_{x,y} \sum_o \|(G^x M_o^y - M_o^y G^x) \otimes I|\psi\rangle\|^2 \\ &= \mathbb{E}_{x,y} \sum_o \langle \psi | M_o^y (G^x)^2 M_o^y \otimes I | \psi \rangle + \mathbb{E}_{x,y} \sum_o \langle \psi | G^x (M_o^y)^2 G^x \otimes I | \psi \rangle \\ & \quad - \mathbb{E}_{x,y} \sum_o \langle \psi | G^x M_o^y G^x M_o^y \otimes I | \psi \rangle - \mathbb{E}_{x,y} \sum_o \langle \psi | M_o^y G^x M_o^y G^x \otimes I | \psi \rangle. \end{aligned} \quad (9.13)$$

We compare all four terms with $\langle \psi | G \otimes M | \psi \rangle$.

For the first term in 9.13,

$$\begin{aligned} \mathbb{E}_{x,y} \sum_o \langle \psi | M_o^y (G^x)^2 M_o^y \otimes I | \psi \rangle &= \mathbb{E}_{x,y} \sum_o \langle \psi | M_o^y G^x M_o^y \otimes I | \psi \rangle \\ &= \mathbb{E}_y \sum_o \langle \psi | M_o^y G M_o^y \otimes I | \psi \rangle \\ &\approx_{2\sqrt{\omega}} \mathbb{E}_y \sum_o \langle \psi | G \otimes M_o^y | \psi \rangle \quad (\text{by 3.32 and 9.11}) \\ &= \langle \psi | G \otimes M | \psi \rangle. \end{aligned}$$

For the second term,

$$\begin{aligned} \mathbb{E}_{x,y} \sum_o \langle \psi | G^x (M_o^y)^2 G^x \otimes I | \psi \rangle &= \mathbb{E}_{x,y} \sum_o \langle \psi | G^x M_o^y G^x \otimes I | \psi \rangle \\ &= \mathbb{E}_x \langle \psi | G^x M G^x \otimes I | \psi \rangle \\ &\approx_{2\sqrt{\zeta}} \mathbb{E}_x \langle \psi | M \otimes G^x | \psi \rangle \quad (\text{by 3.32 and 9.14}) \\ &= \langle \psi | M \otimes G | \psi \rangle. \end{aligned}$$

For the third term, first

$$\begin{aligned} \mathbb{E}_{x,y} \sum_o \langle \psi | G^x M_o^y G^x M_o^y \otimes I | \psi \rangle &= \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | G^x M_o^y G_g^x G_g^x M_o^y \otimes I | \psi \rangle \\ &\approx_{\sqrt{\chi}} \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | G^x M_o^y G_g^x M_o^y G_g^x \otimes I | \psi \rangle. \end{aligned} \quad (9.14)$$

This is the Cauchy–Schwarz estimate coming from 9.12. Next,

$$(9.14) \approx_{\sqrt{\zeta}} \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | G^x M_o^y G_g^x M_o^y \otimes G_g^x | \psi \rangle. \quad (9.15)$$

Here the error is bounded using 3. Then

$$(9.15) \approx_{\sqrt{\zeta}} \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | G_g^x G^x M_o^y G_g^x M_o^y \otimes I | \psi \rangle, \quad (9.16)$$

again by strong self-consistency of G_g^x . Since G is projective,

$$\begin{aligned} (9.16) &= \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | G_g^x M_o^y G_g^x M_o^y \otimes I | \psi \rangle \\ &\approx_{\sqrt{\chi}} \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | M_o^y G_g^x G_g^x M_o^y \otimes I | \psi \rangle. \end{aligned} \quad (9.17)$$

Finally,

$$\begin{aligned}
(9.17) &= \mathbb{E}_{x,y} \sum_{o,g} \langle \psi | M_o^y G_g^x M_o^y \otimes I | \psi \rangle \\
&= \mathbb{E}_y \sum_o \langle \psi | M_o^y G M_o^y \otimes I | \psi \rangle \\
&\approx_{2\sqrt{\omega}} \mathbb{E}_y \sum_o \langle \psi | G \otimes M_o^y | \psi \rangle \quad (\text{by 3.32 and 9.11}) \\
&= \langle \psi | G \otimes M | \psi \rangle.
\end{aligned}$$

Therefore

$$\mathbb{E}_{x,y} \sum_o \langle \psi | G^x M_o^y G^x M_o^y \otimes I | \psi \rangle \approx_{2\sqrt{\zeta}+2\sqrt{\omega}+2\sqrt{\chi}} \langle \psi | G \otimes M | \psi \rangle. \quad (9.18)$$

The fourth term in 9.13 is the Hermitian conjugate of the third term, so it satisfies the same bound. Summing the four errors gives

$$2\sqrt{\omega} + 2\sqrt{\zeta} + 2(2\sqrt{\zeta} + 2\sqrt{\omega} + 2\sqrt{\chi}) = 6\sqrt{\zeta} + 6\sqrt{\omega} + 4\sqrt{\chi}.$$

□

Remark 9.18. The Lean theorem corresponding to Lemma 9.17 takes one extra symmetry hypothesis, `hfix : swapDensity bi.density = bi.density`. This is the `density_swap` component of the paper's permutation-invariance assumption on the symmetric two-prover strategy; the stronger `swap_ev` field of `PermInvState` is not assumed by this theorem. The hypothesis is used to identify $\langle \psi | G \otimes M | \psi \rangle$ with $\langle \psi | M \otimes G | \psi \rangle$ at the center of the triangle-inequality chain.

Corollary 9.19 (Commutativity of the complete part). *In the nontrivial-regime pasting context of Definition 9.3, the following commutation relations hold:*

$$G_g^x G^y \otimes I \approx_{\nu_2} G^y G_g^x \otimes I, \quad G^x G^y \otimes I \approx_{\nu_2} G^y G^x \otimes I,$$

where

$$\nu_2 = 36m (\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}).$$

Proof. By 9.4,

$$G_g^x G_h^y \otimes I \approx_{\nu_{\text{com}}} G_h^y G_g^x \otimes I. \quad (9.19)$$

Applying Lemma 9.17 to 9.19, with $\{M_o^y\}$ equal to the outcome family $\{G_h^y\}$, gives

$$G_g^x G^y \otimes I \approx_{\theta_1} G^y G_g^x \otimes I, \quad (9.20)$$

where $\theta_1 = 12\sqrt{\zeta} + 4\sqrt{\nu_{\text{com}}}$. Applying Lemma 9.17 again to 9.20, now with the one-outcome family $M_o^y = G^y$, gives

$$G^x G^y \otimes I \approx_{\theta_2} G^y G^x \otimes I,$$

where $\theta_2 = 12\sqrt{\zeta} + 4\sqrt{\theta_1}$.

Since $\nu_{\text{com}} = 30m(\gamma^{1/4} + \zeta^{1/4} + (d/q)^{1/4})$, these errors satisfy

$$\theta_1 \leq 36m (\gamma^{1/8} + \zeta^{1/8} + (d/q)^{1/8}) \leq \nu_2,$$

$$\theta_2 \leq 36m (\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}) = \nu_2.$$

□

Corollary 9.20 (Commutativity of the incomplete part). *In the nontrivial-regime pasting context of Definition 9.3,*

$$G_g^x G_\perp^y \otimes I \approx_{\nu_2} G_\perp^y G_g^x \otimes I, \quad G_\perp^x G_\perp^y \otimes I \approx_{\nu_2} G_\perp^y G_\perp^x \otimes I.$$

Proof. The identities

$$G_g^x G_\perp^y - G_\perp^y G_g^x = G^y G_g^x - G_g^x G^y$$

and

$$G_\perp^x G_\perp^y - G_\perp^y G_\perp^x = G^x G^y - G^y G^x$$

reduce both claims to Corollary 9.19. $\square$

Putting Everything Together

Corollary 9.21 (Strong self-consistency and commutation of $\widehat{G}$). *The completed measurements $\widehat{G}^x$ satisfy*

$$\widehat{G}_g^x \otimes I \approx_{2\zeta} I \otimes \widehat{G}_g^x, \tag{9.21}$$

$$\widehat{G}_g^x \widehat{G}_h^y \otimes I \approx_{\nu_3} \widehat{G}_h^y \widehat{G}_g^x \otimes I, \tag{9.22}$$

where

$$\nu_3 = 138m (\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}).$$

Proof. For 9.21, split the outcome set into the genuine outcomes and $\perp$:

$$\mathbb{E}_x \sum_g \|(\widehat{G}_g^x \otimes I - I \otimes \widehat{G}_g^x)|\psi\rangle\|^2 \leq \zeta + \zeta = 2\zeta$$

by 3 and 9.15. For 9.22, the same decomposition yields four cases: polynomial–polynomial, $\perp$ – $\perp$, polynomial– $\perp$, and $\perp$ –polynomial. These contribute ν_{com} and three copies of ν_2 , so

$$\nu_{\text{com}} + 3\nu_2 \leq 138m (\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}) = \nu_3. \quad \square$$

Lemma 9.22 (Completed part bounded by a slice). *The contribution of the completed part in the squared-distance defect is bounded by the corresponding slice contribution.*

9.0.4 Sandwiching Lemmas

Lemma 9.23 (Completed sandwich normalization). *The left and right completed half-sandwiches split under the head–tail reindexing, and the corresponding sums of adjoint-products are bounded above by the identity.*

Lemma 9.24 (One-step completed sandwich commutation). *The elementary head–tail moves, split equivalences, and two-factor case provide the one-step estimates used to move a completed measurement through a half-sandwich.*

Lemma 9.25 (Commuting past multiple completed measurements). *For every $k \geq 2$,*

$$\widehat{G}_{g_1}^{x_1} \widehat{G}_{g_2}^{x_2} \dots \widehat{G}_{g_k}^{x_k} \otimes I \approx_{\nu_4} \widehat{G}_{g_2}^{x_2} \dots \widehat{G}_{g_k}^{x_k} \widehat{G}_{g_1}^{x_1} \otimes I,$$

where

$$\nu_4 = 426k^2m (\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}).$$

Proof. Repeatedly move the leftmost factor across the remaining product, using 9.21 and 9.22 at each step and summing the resulting errors with 3.27. The paper's calculation gives

$$3k(4k\zeta + k\nu_3) \leq 426k^2m(\gamma^{1/16} + \zeta^{1/16} + (d/q)^{1/16}).$$

□

Lemma 9.26 (Line-interpolation mismatch estimates). *If the tuple of completed slices does not interpolate to the sampled vertical-line value, then some active coordinate witnesses a line-value mismatch. Concretely, for every direction u and injective tuple $\mathbf{x} = (x_1, \dots, x_k)$, let $\widehat{H}^\mathbf{x}|_u$ be the vertical-line restriction of the pasted interpolation family. Then*

$$\text{Defect}_\psi(\widehat{H}^\mathbf{x}|_u, B^u) \leq \text{Bad}(\psi; u, \mathbf{x}) \leq \sum_{i=1}^k \text{Defect}_\psi(\widehat{H}^{(i)}(u, \mathbf{x}), B^{(i)}(u, \mathbf{x})),$$

where $\text{Defect}_\psi(A, B)$ denotes the bipartite consistency defect $\max(0, \langle \psi | A_{\text{tot}} \otimes B_{\text{tot}} | \psi \rangle - \sum_a \langle \psi | A_a \otimes B_a | \psi \rangle)$,

$$\text{Bad}(\psi; u, \mathbf{x}) = \sum_f \langle \psi | \left(\sum_{\substack{g_1, \dots, g_k \\ \exists i: g_i \neq \perp, g_i(u) \neq f(x_i)}} \widehat{H}_{g_1, \dots, g_k}^\mathbf{x} \right) \otimes B_f^u | \psi \rangle,$$

and $\widehat{H}^{(i)}, B^{(i)}$ are the one-point line comparison families (Lemma 9.28) restricted to coordinate i , and $\text{Bad}(\psi; u, \mathbf{x}) \leq 1$.

Lemma 9.27 (Averaging the line-interpolation defect). *Replacing distinct tuples by independent tuples costs the total-variation error k^2/q from Proposition 9.7. After averaging over u and $\mathbf{x} \sim \text{Distinct}_k$, the expected bad mass and the expected pasted interpolation defect are bounded by the one-point sandwich error:*

$$\mathbb{E}_u \mathbb{E}_{\mathbf{x} \sim \text{Distinct}_k} \text{Bad}(\psi; u, \mathbf{x}) \leq k \cdot \nu_5 + \frac{k^2}{q},$$

and consequently

$$\mathbb{E}_u \mathbb{E}_{\mathbf{x} \sim \text{Distinct}_k} \text{Defect}_\psi(\widehat{H}^\mathbf{x}|_u, B^u) \leq k \cdot \nu_5 + \frac{k^2}{q},$$

where $\nu_5 = 43km(\varepsilon^{1/32} + \dots + (d/q)^{1/32})$ (Lemma 9.28). Using $1/q \leq (d/q)^{1/32}$ and $m \geq 1$, this is further bounded by $44k^2m(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32})$ as used in Lemma 9.29.

Lemma 9.28 (Consistency of the sandwiched measurement with one line value). *For any $1 \leq i \leq k$,*

$$\mathbb{E}_u \mathbb{E}_{x_1, \dots, x_k} \sum_{g_1, \dots, g_k: g_i \neq \perp} \sum_{a \neq g_i(u)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes B_{[f(x_i)=a]}^u | \psi \rangle \leq \nu_5,$$

where

$$\nu_5 = 43km(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}).$$

Proof. Summing out the coordinates to the right of i gives

$$\begin{aligned} & \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_k} \sum_{g_1, \dots, g_k: g_i \neq \perp} \sum_{a \neq g_i(u)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes B_{[f(x_i)=a]}^u | \psi \rangle \\ &= \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_1, \dots, g_i: g_i \neq \perp} \sum_{a \neq g_i(u)} \langle \psi | \widehat{H}_{g_1, \dots, g_i}^{x_1, \dots, x_i} \otimes B_{[f(x_i)=a]}^u | \psi \rangle. \end{aligned} \tag{9.23}$$

Write $\widehat{G}_{g_{<i}}^{x_{<i}} = \widehat{G}_{g_1}^{x_1} \dots \widehat{G}_{g_{i-1}}^{x_{i-1}}$. Then

$$\begin{aligned} (9.23) &= \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_1, \dots, g_i : g_i \neq \perp} \langle \psi | \widehat{G}_{g_{<i}}^{x_{<i}} \widehat{G}_{g_i}^{x_i} (\widehat{G}_{g_{<i}}^{x_{<i}})^\dagger \otimes (I - B_{[f(x_i)=g_i(u)]}^u) | \psi \rangle \\ &\approx_{\sqrt{\nu_4}} \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_1, \dots, g_i : g_i \neq \perp} \langle \psi | \widehat{G}_{g_i}^{x_i} \widehat{G}_{g_{<i}}^{x_{<i}} \widehat{G}_{g_i}^{x_i} (\widehat{G}_{g_{<i}}^{x_{<i}})^\dagger \otimes (I - B_{[f(x_i)=g_i(u)]}^u) | \psi \rangle. \end{aligned} \quad (9.24)$$

The Cauchy–Schwarz argument uses

$$\mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_1, \dots, g_i} \langle \psi | ((\widehat{G}_{g_{<i}}^{x_{<i}} \widehat{G}_{g_i}^{x_i} - \widehat{G}_{g_i}^{x_i} \widehat{G}_{g_{<i}}^{x_{<i}})(\widehat{G}_{g_i}^{x_i} (\widehat{G}_{g_{<i}}^{x_{<i}})^\dagger - (\widehat{G}_{g_{<i}}^{x_{<i}})^\dagger \widehat{G}_{g_i}^{x_i})) \otimes I | \psi \rangle, \quad (9.25)$$

which is at most ν_4 by Lemma 9.25. A second Cauchy–Schwarz step gives

$$(9.24) \approx_{\sqrt{\nu_4}} \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_1, \dots, g_i : g_i \neq \perp} \langle \psi | \widehat{G}_{g_i}^{x_i} \widehat{G}_{g_{<i}}^{x_{<i}} (\widehat{G}_{g_{<i}}^{x_{<i}})^\dagger \widehat{G}_{g_i}^{x_i} \otimes (I - B_{[f(x_i)=g_i(u)]}^u) | \psi \rangle. \quad (9.26)$$

Summing over $g_1, \dots, g_{i-1}$ collapses the middle factor to I , so 9.26 becomes

$$\mathbb{E}_u \mathbb{E}_{x_1, \dots, x_i} \sum_{g_i : g_i \neq \perp} \langle \psi | \widehat{G}_{g_i}^{x_i} \otimes (I - B_{[f(x_i)=g_i(u)]}^u) | \psi \rangle \leq \nu_1$$

by 9.2. Thus the total error is

$$\nu_1 + 2\sqrt{\nu_4} \leq 43km (\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}).$$

□

9.0.5 Consistency of H with A

Lemma 9.29 (Consistency of the pasted measurement with the line measurement). *The pasted submeasurement satisfies*

$$H_{[h|_u=f]} \otimes I \simeq_{\nu_6} I \otimes B_f^u,$$

where

$$\nu_6 = 44k^2m (\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}).$$

Proof. The inconsistency with B^u expands as

$$\begin{aligned} &\mathbb{E}_u \sum_{f \neq f'} \langle \psi | H_{[h|_u=f']} \otimes B_f^u | \psi \rangle \\ &= \mathbb{E}_u \sum_h \sum_{f \neq h|_u} \langle \psi | H_h \otimes B_f^u | \psi \rangle \\ &= \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_k} \sum_h \sum_{f \neq h|_u} \langle \psi | H_h^{x_1, \dots, x_k} \otimes B_f^u | \psi \rangle \\ &= \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_k} \sum_h \sum_{w: |w| \geq d+1} \sum_{(g_1, \dots, g_k)=h_w} \sum_{f \neq h|_u} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes B_f^u | \psi \rangle. \end{aligned} \quad (9.27)$$

If $|w| \geq d+1$ and $f \neq h|_u$, then some active coordinate i must satisfy $g_i \neq \perp$ and $g_i(u) \neq f(x_i)$. Hence

$$(9.27) \leq \mathbb{E}_u \mathbb{E}_{x_1, \dots, x_k} \sum_{g_1, \dots, g_k} \sum_{\substack{f: \exists i, g_i \neq \perp, \\ g_i(u) \neq f(x_i)}} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes B_f^u | \psi \rangle. \quad (9.28)$$

Passing from independent tuples to distinct tuples costs at most k^2/q by Lemma 9.7. A union bound over the offending coordinate then gives

$$(9.28) \leq \sum_{i=1}^k \mathbb{E}_u \mathbb{E}_{y_1, \dots, y_k} \sum_{g_1, \dots, g_k : g_i \neq \perp} \sum_{a \neq g_i(u)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes B_{[f(y_i)=a]}^u | \psi \rangle \leq k\nu_5.$$

Therefore

$$\frac{k^2}{q} + k\nu_5 \leq 44k^2m(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) = \nu_6.$$

□

Corollary 9.30 (Consistency of H with A). *On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,*

$$H_{[h(u,x)=a]} \otimes I \simeq_\nu I \otimes A_a^{u,x}.$$

Proof. Lemma 9.29 implies

$$H_{[h(u,x)=a]} \otimes I \simeq_{\nu_6} I \otimes B_{[f(x)=a]}^u.$$

Applying 3.20 to this relation and 9.1 gives

$$H_{[h(u,x)=a]} \otimes I \simeq_{\nu_6 + \sqrt{8m\varepsilon + 4\delta}} I \otimes A_a^{u,x}.$$

Since

$$\sqrt{8m\varepsilon + 4\delta} + \nu_6 \leq 47k^2m(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) \leq \nu,$$

the claim follows. □

Lemma 9.31 (Point consistency from vertical-line consistency). *Let H be any polynomial-valued submeasurement. Suppose that the strategy is good, that $\gamma, \zeta \geq 0$, that $k \geq 1$, and that the restriction of H to every vertical line is consistent with the vertical-line measurement with error ν_6 . Then the pointwise evaluations of H are consistent with Alice's point measurement with error ν .*

Corollary 9.32 (Consistency of the completed H with A). *Let H_{meas} be the completion of H at the fallback outcome. On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,*

$$(H_{\text{meas}})_{[h(u,x)=a]} \otimes I \simeq_\sigma I \otimes A_a^{u,x}.$$

Proof. Corollary 9.30 gives

$$H_{[h(u,x)=a]} \otimes I \simeq_\nu I \otimes A_a^{u,x}.$$

Corollary 9.42 bounds the missing mass of H by

$$\kappa \cdot \left(1 + \frac{1}{100m}\right) + \nu + e^{-k/(80000m^2)}.$$

Adding this completion mass to the ν -bound gives

$$\nu + \kappa \cdot \left(1 + \frac{1}{100m}\right) + \nu + e^{-k/(80000m^2)} = \sigma.$$

□

Lemma 9.33 (Completion from submeasurement consistency). *Let H be any polynomial-valued submeasurement. Suppose that the pointwise evaluations of H are consistent with Alice's point measurement with error ν , and that H has the lower mass bound used in Corollary 9.42. Then completing H at the fallback polynomial preserves point consistency with the enlarged error appearing in the induction statement.*

9.0.6 Completeness of H

Definition 9.34 (Outcomes of a fixed type). For a type $\tau \in \{0, 1\}^k$, let Outcomes_τ be the set of tuples $(g_1, \dots, g_k)$ such that $g_i \in \mathcal{P}(m, q, d)$ when $\tau_i = 1$ and $g_i = \perp$ when $\tau_i = 0$. If $x_1, \dots, x_k \in \mathbb{F}_q$, let $\text{Global}_\tau(x)$ be the subset of Outcomes_τ arising from restrictions of a single polynomial in $\mathcal{P}(m+1, q, d)$, and let $\overline{\text{Global}_\tau(x)} = \text{Outcomes}_\tau \setminus \text{Global}_\tau(x)$.

Lemma 9.35 (Summing over all outcomes). *If $x_1, \dots, x_k$ are sampled independently and uniformly from $\mathbb{F}_q$, then*

$$\langle \psi | H \otimes I | \psi \rangle \approx_{\nu_7} \mathbb{E}_{x_1, \dots, x_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Outcomes}_\tau} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes I | \psi \rangle,$$

where

$$\nu_7 = 46k^2m (\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}).$$

Proof. Let $(y_1, \dots, y_k) \sim \text{Distinct}_k$. By definition,

$$\begin{aligned} \langle \psi | H \otimes I | \psi \rangle &= \sum_h \langle \psi | H_h \otimes I | \psi \rangle \\ &= \mathbb{E}_{y_1, \dots, y_k} \sum_h \langle \psi | H_h^{y_1, \dots, y_k} \otimes I | \psi \rangle \\ &= \mathbb{E}_{y_1, \dots, y_k} \sum_h \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Global}_\tau(y)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle \\ &= \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Global}_\tau(y)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle. \end{aligned} \quad (9.29)$$

We now remove the restriction to globally consistent tuples:

$$(9.29) \approx_{\frac{k^2}{q} + k\nu_5 + \frac{md}{q}} \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Outcomes}_\tau} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle. \quad (9.30)$$

Since the right-hand side is at least the left-hand side, it suffices to bound the difference:

$$\begin{aligned} (9.30) - (9.29) &= \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}_\tau(y)}} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle \\ &= \mathbb{E}_u \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_f \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}_\tau(y)}} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes B_f^u | \psi \rangle. \end{aligned} \quad (9.31)$$

We next insert an indicator recording consistency along the sampled line:

$$(9.31) \approx_{\frac{k^2}{q} + k\nu_5} \mathbb{E}_u \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_f \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}_\tau(y)}} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes B_f^u | \psi \rangle \cdot \mathbf{1}[\forall i \in \tau, f(y_i) = g_i(u)]. \quad (9.32)$$

The discarded part is

$$\mathbb{E}_u \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_f \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}_\tau(y)}} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes B_f^u | \psi \rangle \cdot \mathbf{1}[\exists i \in \tau, f(y_i) \neq g_i(u)], \quad (9.33)$$

and after switching to independent tuples and applying Lemma 9.28 coordinatewise, this contributes at most $k^2/q + k\nu_5$.

Let $\text{Consistent}_\tau(g, y, u)$ denote the event that there exists a degree- d polynomial f with $f(y_i) = g_i(u)$ for all $i \in \tau$. Then

$$\mathbf{1}[\forall i \in \tau, f(y_i) = g_i(u)] \leq \mathbf{1}[\text{Consistent}_\tau(g, y, u)],$$

so 9.32 is bounded by

$$\begin{aligned} & \mathbb{E}_u \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}}_\tau(y)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle \cdot \mathbf{1}[\text{Consistent}_\tau(g, y, u)] \\ &= \mathbb{E}_{y_1, \dots, y_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \overline{\text{Global}}_\tau(y)} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{y_1, \dots, y_k} \otimes I | \psi \rangle \cdot \Pr_u[\text{Consistent}_\tau(g, y, u)]. \end{aligned} \quad (9.34)$$

For a fixed globally inconsistent tuple, choose $d+1$ active coordinates and let h^* be the unique interpolant through them. Since the tuple is not globally consistent, some active coordinate i^* satisfies $g_{i^*} \neq h^*|_{y_{i^*}}$, and then Schwartz–Zippel gives

$$\Pr_u[\text{Consistent}_\tau(g, y, u)] \leq \Pr_u[g_{i^*}(u) = h^*(u, y_{i^*})] \leq \frac{md}{q}.$$

Therefore 9.34 is at most md/q , which proves 9.30. A final application of Lemma 9.7 replaces distinct tuples by independent tuples, giving a total error of

$$2 \frac{k^2}{q} + \frac{md}{q} + k\nu_5 \leq 46k^2m (\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) = \nu_7.$$

□

Definition 9.36 (Truncated type sums). For $1 \leq \ell \leq k+1$ and a tail type $\tau_{\geq \ell} \in \{0, 1\}^{k-\ell+1}$, define

$$S_{\tau_{\geq \ell}} = \sum_{\tau_{< \ell}: |\tau| \geq d+1} G^{|\tau_{< \ell}|} (I - G)^{(\ell-1)-|\tau_{< \ell}|}.$$

This records the contribution of all prefixes that can still be completed to total weight at least $d+1$ once the tail $\tau_{\geq \ell}$ is fixed.

Lemma 9.37 (Recurrence for truncated type sums). *Each $S_{\tau_{\geq \ell}}$ is Hermitian, positive semidefinite, and bounded by I . Moreover, if $\tau_{> \ell}$ is a tail type of length $k-\ell$, then*

$$S_{\tau_{> \ell}} = \sum_{\tau_\ell \in \{0, 1\}} S_{\tau_{\geq \ell}} G^{\tau_\ell} (I - G)^{1-\tau_\ell}.$$

Proof. Every summand is a polynomial in the commuting positive operators G and $I - G$, so $S_{\tau_{\geq \ell}}$ is Hermitian and positive semidefinite. Summing over all prefixes gives

$$S_{\tau_{\geq \ell}} \leq (G + (I - G))^{\ell-1} = I.$$

Splitting the next type bit according to whether $\tau_\ell = 1$ or $\tau_\ell = 0$ gives the stated recurrence. □

Lemma 9.38 (From the pasted sum to a polynomial in G). *Let $|\psi_{\text{bi}}\rangle$ be a bipartite state on $\iota \otimes \iota$. If $x_1, \dots, x_k$ are sampled independently and uniformly from $\mathbb{F}_q$, then*

$$\mathbb{E}_{x_1, \dots, x_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Outcomes}_\tau} \langle \psi_{\text{bi}} | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes I | \psi_{\text{bi}} \rangle \approx_{\nu_8} \sum_{r=d+1}^k \binom{k}{r} \langle \psi_{\text{bi}} | G^r (I - G)^{k-r} \otimes I | \psi_{\text{bi}} \rangle,$$

where

$$\nu_8 = 46k^2m (\gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}).$$

Proof. For a type $\tau \in \{0, 1\}^k$, write $\tau_{<\ell}$, $\tau_{>\ell}$, $\tau_{\leq\ell}$, and $\tau_{\geq\ell}$ for the obvious truncations, and similarly for tuples g . Also write

$$\widehat{G}_{g_{\geq\ell}}^{x_{\geq\ell}} = \widehat{G}_{g_\ell}^{x_\ell} \cdots \widehat{G}_{g_k}^{x_k}.$$

Then

$$\widehat{G}_{g_{\geq\ell}}^{x_{\geq\ell}} = \widehat{G}_{g_\ell}^{x_\ell} \widehat{G}_{g_{>\ell}}^{x_{>\ell}}, \quad (9.35)$$

and

$$\widehat{H}_{g_{\geq\ell}}^{x_{\geq\ell}} = \widehat{G}_{g_{\geq\ell}}^{x_{\geq\ell}} (\widehat{G}_{g_{\geq\ell}}^{x_{\geq\ell}})^\dagger. \quad (9.36)$$

Write $\mathbf{O}_\tau = \text{Outcomes}_\tau$. The main iterative step is that for each $1 \leq \ell \leq k$,

$$\begin{aligned} & \mathbb{E}_{x_{\geq\ell}} \sum_{\tau: |\tau| \geq d+1} \sum_{g_{\geq\ell} \in \mathbf{O}_{\tau_{\geq\ell}}} \langle \psi | \widehat{H}_{g_{\geq\ell}}^{x_{\geq\ell}} \otimes (G^{|\tau_{<\ell}|} (I - G)^{(\ell-1)-|\tau_{<\ell}|}) | \psi \rangle \\ & \approx_{2\sqrt{2\zeta} + 2\sqrt{\nu_4}} \mathbb{E}_{x_{>\ell}} \sum_{\tau: |\tau| \geq d+1} \sum_{g_{>\ell} \in \mathbf{O}_{\tau_{>\ell}}} \langle \psi | \widehat{H}_{g_{>\ell}}^{x_{>\ell}} \otimes (G^{|\tau_{\leq\ell}|} (I - G)^{\ell-|\tau_{\leq\ell}|}) | \psi \rangle. \end{aligned} \quad (9.37)$$

Iterating 9.37 for $\ell = 1, \dots, k$ produces the Bernoulli polynomial

$$\sum_{r=d+1}^k \binom{k}{r} \langle \psi | G^r (I - G)^{k-r} \otimes I | \psi \rangle,$$

and the accumulated error is

$$k(2\sqrt{2\zeta} + 2\sqrt{\nu_4}) \leq 46k^2m(\gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) = \nu_8.$$

This corrects the displayed arithmetic in the paper: the iterated adjacent estimate contributes k copies of the $2\sqrt{\nu_4}$ term, and the definition of ν_4 already contains a factor k^2 .

To prove 9.37, define for each $1 \leq \ell \leq k+1$ and each tail type $\tau_{\geq\ell}$,

$$S_{\tau_{\geq\ell}} = \sum_{\tau_{<\ell}: |\tau| \geq d+1} G^{|\tau_{<\ell}|} (I - G)^{(\ell-1)-|\tau_{<\ell}|}. \quad (9.38)$$

Then 9.37 may be rewritten as

$$\begin{aligned} & \mathbb{E}_{x_{\geq\ell}} \sum_{\tau_{\geq\ell}} \sum_{g_{\geq\ell} \in \mathbf{O}_{\tau_{\geq\ell}}} \langle \psi | \widehat{H}_{g_{\geq\ell}}^{x_{\geq\ell}} \otimes S_{\tau_{\geq\ell}} | \psi \rangle \\ & \approx_{2\sqrt{2\zeta} + 2\sqrt{\nu_4}} \mathbb{E}_{x_{>\ell}} \sum_{\tau_{>\ell}} \sum_{g_{>\ell} \in \mathbf{O}_{\tau_{>\ell}}} \langle \psi | \widehat{H}_{g_{>\ell}}^{x_{>\ell}} \otimes S_{\tau_{>\ell}} | \psi \rangle. \end{aligned} \quad (9.39)$$

Remark 9.39 (Lean proof-stage notation). In Lean, the type-restricted averaged submeasurement, the per-tail operator/mass, the aggregated stage quantity, and the final Bernoulli-tail scalar from this proof are recorded by the declarations above. Their indexing is shifted to the repository's 0-based convention: Lean stage ℓ is the paper's stage $\ell + 1$.

The operators $S_{\tau_{\geq\ell}}$ are Hermitian and positive semidefinite, and

$$\begin{aligned} S_{\tau_{\geq\ell}} &= \sum_{\tau_{<\ell}: |\tau| \geq d+1} G^{|\tau_{<\ell}|} (I - G)^{(\ell-1)-|\tau_{<\ell}|} \\ &\leq \sum_{\tau_{<\ell}} G^{|\tau_{<\ell}|} (I - G)^{(\ell-1)-|\tau_{<\ell}|} \\ &= (G + (I - G))^{\ell-1} = I. \end{aligned} \quad (9.40)$$

For any $\tau_\ell \in \{0, 1\}$,

$$\mathbb{E}_{x_\ell} \sum_{g_\ell \in \mathbf{O}_{\tau_\ell}} \widehat{G}_{g_\ell}^{x_\ell} = G^{\tau_\ell} (I - G)^{1-\tau_\ell}. \quad (9.41)$$

Therefore, for every $\tau_{>\ell}$,

$$\begin{aligned} \sum_{\tau_\ell} S_{\tau_{\geq \ell}} \cdot \left(\mathbb{E}_{x_\ell} \sum_{g_\ell \in \mathbf{O}_{\tau_\ell}} \widehat{G}_{g_\ell}^{x_\ell} \right) &= \sum_{\tau_\ell} S_{\tau_{\geq \ell}} G^{\tau_\ell} (I - G)^{1-\tau_\ell} \\ &= S_{\tau_{>\ell}}. \end{aligned} \quad (9.42)$$

Moreover,

$$S_{\tau_{\geq \ell}} \cdot \left(\mathbb{E}_{x_\ell} \sum_{g_\ell \in \mathbf{O}_{\tau_\ell}} \widehat{G}_{g_\ell}^{x_\ell} \right) \cdot S_{\tau_{\geq \ell}} \leq \mathbb{E}_{x_\ell} \sum_{g_\ell \in \mathbf{O}_{\tau_\ell}} \widehat{G}_{g_\ell}^{x_\ell}, \quad (9.43)$$

because $S_{\tau_{\geq \ell}}$ commutes with G and $(I - G)$ and is bounded by I .

We now prove 9.39. Writing $\widehat{H}$ as a sandwich of $\widehat{G}$ operators and moving the rightmost $\widehat{G}_{g_\ell}^{x_\ell}$ to the second tensor factor gives

$$\begin{aligned} &\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{\geq \ell}}^{x_{\geq \ell}} \otimes S_{\tau_{\geq \ell}} | \psi \rangle \\ &= \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{\geq \ell}}^{x_{\geq \ell}} (\widehat{G}_{g_{>\ell}}^{x_{>\ell}})^\dagger \widehat{G}_{g_\ell}^{x_\ell}) \otimes S_{\tau_{\geq \ell}} | \psi \rangle \\ &\approx_{\sqrt{2}\zeta} \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{\geq \ell}}^{x_{\geq \ell}} (\widehat{G}_{g_{>\ell}}^{x_{>\ell}})^\dagger) \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_\ell}^{x_\ell}) | \psi \rangle. \end{aligned} \quad (9.44)$$

The corresponding Cauchy–Schwarz bound is

$$\begin{aligned} &\left| \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{\geq \ell}}^{x_{\geq \ell}} \otimes S_{\tau_{\geq \ell}}) ((\widehat{G}_{g_{>\ell}}^{x_{>\ell}})^\dagger \otimes I) (\widehat{G}_{g_\ell}^{x_\ell} \otimes I - I \otimes \widehat{G}_{g_\ell}^{x_\ell}) | \psi \rangle \right| \\ &\leq \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{\geq \ell}}^{x_{\geq \ell}} \otimes (S_{\tau_{\geq \ell}})^2 | \psi \rangle} \cdot \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_\ell}^{x_\ell} \otimes I - I \otimes \widehat{G}_{g_\ell}^{x_\ell}) (\widehat{H}_{g_{>\ell}}^{x_{>\ell}} \otimes I) (\widehat{G}_{g_\ell}^{x_\ell} \otimes I - I \otimes \widehat{G}_{g_\ell}^{x_\ell}) | \psi \rangle} \end{aligned}$$

The first square root is at most 1 because $S_{\tau_{\geq \ell}} \leq I$ and $\widehat{H}$ is a submeasurement, and the second is at most 2ζ by 9.21.

Next, commute the leftmost $\widehat{G}_{g_\ell}^{x_\ell}$ across the left half-sandwich:

$$\begin{aligned} (9.44) &= \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_\ell}^{x_\ell} \widehat{G}_{g_{>\ell}}^{x_{>\ell}} (\widehat{G}_{g_{>\ell}}^{x_{>\ell}})^\dagger) \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_\ell}^{x_\ell}) | \psi \rangle \\ &\approx_{\sqrt{\nu_4}} \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_\ell, g_{>\ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{>\ell}}^{x_{>\ell}} \widehat{G}_{g_\ell}^{x_\ell} (\widehat{G}_{g_{>\ell}}^{x_{>\ell}})^\dagger) \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_\ell}^{x_\ell}) | \psi \rangle. \end{aligned} \quad (9.46)$$

The associated Cauchy–Schwarz estimate is

$$\begin{aligned}
& \left| \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | ([\widehat{G}_{g_{> \ell}}^{x_{> \ell}}, \widehat{G}_{g_{\ell}}^{x_{\ell}}] \otimes I) ((\widehat{G}_{g_{> \ell}}^{x_{> \ell}})^{\dagger} \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}})) | \psi \rangle \right| \\
& \leq \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | [\widehat{G}_{g_{> \ell}}^{x_{> \ell}}, \widehat{G}_{g_{\ell}}^{x_{\ell}}] [\widehat{G}_{g_{> \ell}}^{x_{> \ell}}, \widehat{G}_{g_{\ell}}^{x_{\ell}}]^{\dagger} \otimes I | \psi \rangle} \cdot \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes (\widehat{G}_{g_{\ell}}^{x_{\ell}} (S_{\tau_{\geq \ell}})^2 \widehat{G}_{g_{\ell}}^{x_{\ell}}) | \psi \rangle}.
\end{aligned} \tag{9.47}$$

The first square root is bounded by ν_4 via Lemma 9.25, and the second by 1 since $S_{\tau_{\geq \ell}} \leq I$.

Continue commuting:

$$(9.46) \approx_{\sqrt{\nu_4}} \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{> \ell}}^{x_{> \ell}} (\widehat{G}_{g_{> \ell}}^{x_{> \ell}})^{\dagger} \widehat{G}_{g_{\ell}}^{x_{\ell}}) \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}}) | \psi \rangle. \tag{9.48}$$

This time the Cauchy–Schwarz step is

$$\begin{aligned}
& \left| \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | (\widehat{G}_{g_{> \ell}}^{x_{> \ell}} \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}})) ([(\widehat{G}_{g_{> \ell}}^{x_{> \ell}})^{\dagger}, \widehat{G}_{g_{\ell}}^{x_{\ell}}] \otimes I) | \psi \rangle \right| \\
& \leq \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}} S_{\tau_{\geq \ell}}) | \psi \rangle} \cdot \sqrt{\mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | [(\widehat{G}_{g_{> \ell}}^{x_{> \ell}})^{\dagger}, \widehat{G}_{g_{\ell}}^{x_{\ell}}]^{\dagger} [(\widehat{G}_{g_{> \ell}}^{x_{> \ell}})^{\dagger}, \widehat{G}_{g_{\ell}}^{x_{\ell}}] \otimes I | \psi \rangle}.
\end{aligned} \tag{9.49}$$

By 9.43, the first square root is at most 1, and the second matches the first square root in 9.47, hence is at most ν_4 .

Finally move the remaining $\widehat{G}_{g_{\ell}}^{x_{\ell}}$ to the second tensor factor:

$$(9.48) \approx_{\sqrt{2\zeta}} \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}}) | \psi \rangle. \tag{9.50}$$

The omitted Cauchy–Schwarz bound reuses the first square root from 9.49 and the second square root from 9.45. Since $\widehat{G}$ is projective,

$$\begin{aligned}
(9.50) &= \mathbb{E}_{x_{\geq \ell}} \sum_{\tau_{\geq \ell}} \sum_{(g_{\ell}, g_{> \ell}) \in \mathbf{O}_{\tau_{\geq \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes (S_{\tau_{\geq \ell}} \widehat{G}_{g_{\ell}}^{x_{\ell}}) | \psi \rangle \\
&= \mathbb{E}_{x_{> \ell}} \sum_{\tau_{> \ell}} \sum_{g_{> \ell} \in \mathbf{O}_{\tau_{> \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes \left(\sum_{\tau_{\ell}} S_{\tau_{\geq \ell}} \left(\mathbb{E}_{x_{\ell}} \sum_{g_{\ell} \in \mathbf{O}_{\tau_{\ell}}} \widehat{G}_{g_{\ell}}^{x_{\ell}} \right) \right) | \psi \rangle \\
&= \mathbb{E}_{x_{> \ell}} \sum_{\tau_{> \ell}} \sum_{g_{> \ell} \in \mathbf{O}_{\tau_{> \ell}}} \langle \psi | \widehat{H}_{g_{> \ell}}^{x_{> \ell}} \otimes S_{\tau_{> \ell}} | \psi \rangle,
\end{aligned} \tag{by 9.42}$$

which is exactly 9.39. $\square$

Lemma 9.40 (Context specializations for the pasted measurement). *The nontrivial-regime pasting context specializes the consistency of H with A and the conversion from the pasted sum to the polynomial in G to the parameters and error bounds fixed in Definition 9.3.*

Lemma 9.41 (Matrix Chernoff bound for the Bernoulli polynomial). *Let $0 < \theta < 1$ and integers $k, d > 0$ with $k \geq 2d/\theta$. Define*

$$F(X) = \sum_{r=d+1}^k \binom{k}{r} X^r (I - X)^{k-r}.$$

If X is Hermitian, $0 \leq X \leq I$, and $\langle \psi | X \otimes I | \psi \rangle \geq 1 - \kappa$, then

$$\langle \psi | F(X) \otimes I | \psi \rangle \geq 1 - \frac{\kappa}{1 - \theta} - e^{-\theta^2 k/2}.$$

Proof. Let ρ be the reduced state of $|\psi\rangle$ on one prover's subsystem, and write the eigendecomposition $X = \sum_i \lambda_i |v_i\rangle\langle v_i|$. This defines a probability distribution $\mu(i) = \langle v_i | \rho | v_i \rangle$, with

$$\mathbb{E}_{i \sim \mu} \lambda_i = \text{Tr}(X\rho) = \langle \psi | X \otimes I | \psi \rangle \geq 1 - \kappa.$$

Equivalently, $\mathbb{E}_{i \sim \mu} (1 - \lambda_i) \leq \kappa$, so Markov's inequality shows that most of the spectral mass of X lies on eigenvalues at least θ :

$$\Pr_{i \sim \mu} [\lambda_i > \theta] \geq 1 - \frac{\kappa}{1 - \theta}. \quad (9.51)$$

For a scalar $p \in [d/k, 1]$, the quantity

$$F(p) = \sum_{r=d+1}^k \binom{k}{r} p^r (1-p)^{k-r}$$

is the probability of observing at least $d+1$ successes in k Bernoulli(p) trials. Applying the scalar Chernoff bound eigenvalue-by-eigenvalue therefore yields

$$F(p) \geq 1 - \exp\left(-2\left(p - \frac{d}{k}\right)^2 k\right). \quad (9.52)$$

Hence

$$\begin{aligned} \langle \psi | F(X) \otimes I | \psi \rangle &= \text{Tr}(F(X)\rho) \\ &= \sum_i F(\lambda_i) \langle v_i | \rho | v_i \rangle \\ &= \mathbb{E}_{i \sim \mu} F(\lambda_i) \\ &\geq \Pr_{i \sim \mu} [\lambda_i \geq \theta] \cdot F(\theta) \\ &\geq \left(1 - \frac{\kappa}{1 - \theta}\right) \cdot F(\theta) \quad (\text{by } 9.51) \\ &\geq \left(1 - \frac{\kappa}{1 - \theta}\right) \cdot \left(1 - \exp\left(-2\left(\theta - \frac{d}{k}\right)^2 k\right)\right). \end{aligned} \quad (9.53)$$

Since $(1-b)(1-c) \geq 1-b-c$ for $b, c \geq 0$, 9.53 implies

$$\langle \psi | F(X) \otimes I | \psi \rangle \geq 1 - \frac{\kappa}{1 - \theta} - \exp\left(-2\left(\theta - \frac{d}{k}\right)^2 k\right).$$

Finally, $k \geq 2d/\theta$ implies $d/k \leq \theta/2$, hence $(\theta - d/k)^2 \geq \theta^2/4$, which yields the claimed bound. $\square$

Corollary 9.42 (Completeness of the pasted submeasurement). *If $k \geq 400md$, then*

$$\langle \psi | H \otimes I | \psi \rangle \geq 1 - \kappa \left(1 + \frac{1}{100m}\right) - \nu - e^{-k/(80000m^2)}.$$

Proof. Lemma 9.35 gives

$$\langle \psi | H \otimes I | \psi \rangle \approx_{\nu_7} \mathbb{E}_{x_1, \dots, x_k} \sum_{\tau: |\tau| \geq d+1} \sum_{(g_1, \dots, g_k) \in \text{Outcomes}_\tau} \langle \psi | \widehat{H}_{g_1, \dots, g_k}^{x_1, \dots, x_k} \otimes I | \psi \rangle.$$

Lemma 9.38 turns the right-hand side into

$$\sum_{r=d+1}^k \binom{k}{r} \langle \psi | G^r (I - G)^{k-r} \otimes I | \psi \rangle$$

up to error ν_8 . Apply Lemma 9.41 with $\theta = 1/(200m)$. Since $k \geq 400md$, the hypothesis $k \geq 2d/\theta$ is satisfied, and the resulting lower bound is

$$1 - \frac{\kappa}{1 - 1/(200m)} - e^{-k/(80000m^2)} \geq 1 - \kappa \left(1 + \frac{1}{100m} \right) - e^{-k/(80000m^2)}.$$

The paper then absorbs the approximation losses $\nu_7 + \nu_8$ into ν . □

Chapter 10

The main induction step

10.1 Restricting to one slice

Definition 10.1 (Appending a line at height x). For $x \in \mathbb{F}_q$, let $\text{append}_x(\ell)$ be the line in $\mathbb{F}_q^{m+1}$ obtained by appending the coordinate x to every point of a line $\ell \subseteq \mathbb{F}_q^m$. If $f: \ell \rightarrow \mathbb{F}_q$, let $\text{append}_x(f)$ be the polynomial on $\text{append}_x(\ell)$ defined by $\text{append}_x(f)(u, x) = f(u)$.

Definition 10.2 (x -restricted strategy). Let (ψ, A, B, L) be a symmetric strategy for the $(m+1, q, d)$ low individual degree test. For each $x \in \mathbb{F}_q$, the x -restricted strategy (ψ, A^x, B^x, L^x) for the (m, q, d) test is defined by

$$(A^x)_a^u = A_a^{u,x}, \quad (B^x)_f^\ell = B_{\text{append}_x(f)}^{\text{append}_x(\ell)}, \quad (L^x)_f^\ell = L_{\text{append}_x(f)}^{\text{append}_x(\ell)}.$$

Lemma 10.3 (Restricted success probabilities). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m+1, q, d)$ test. For each $x \in \mathbb{F}_q$, let (ψ, A^x, B^x, L^x) be the restricted strategy and let $\varepsilon_x, \delta_x, \gamma_x$ be its three failure probabilities. Then*

$$\mathbb{E}_x \varepsilon_x \leq \frac{m+1}{m} \varepsilon, \quad \mathbb{E}_x \delta_x \leq \delta, \quad \mathbb{E}_x \gamma_x \leq \frac{m+1}{m} \gamma.$$

Proof. The self-consistency test restricts exactly to the slice at height x . For the axis-parallel and diagonal-line tests, conditioning on the sampled line not being parallel to the last coordinate identifies the test with the corresponding restricted game. $\square$

Definition 10.4 (Average restricted errors). The average restricted errors are the expectations over $x \in \mathbb{F}_q$ of the axis-parallel, self-consistency, and restricted diagonal-line failure probabilities of the x -restricted strategies.

Lemma 10.5 (Weighted restricted-probability bounds). *The weighted bounds from Lemma 10.3 imply the restricted-probability data used in the successor step.*

10.2 The inductive construction

Theorem 10.6 (Self-improvement). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the (m, q, d) low individual degree test. Let $G \in \text{PolySub}(m, q, d)$ be a submeasurement with the following property:*

1. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\nu I \otimes G_{[g(u)=a]}.$$

Let

$$\zeta = 3000m (\varepsilon^{1/32} + \delta^{1/32} + (d/q)^{1/32}).$$

Then there exists a projective submeasurement $H \in \text{PolySub}(m, q, d)$ with the following properties:

1. (Completeness): If $H = \sum_h H_h$, then

$$\langle \psi | H \otimes I | \psi \rangle \geq (1 - \nu) - \zeta.$$

2. (Consistency with A): On average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\zeta I \otimes H_{[h(u)=a]}.$$

3. (Strong self-consistency):

$$H_h \otimes I \approx_\zeta I \otimes H_h.$$

4. (Boundedness): There exists a positive semidefinite operator Z such that

$$\langle \psi | Z \otimes (I - H) | \psi \rangle \leq \zeta$$

and for each $h \in \mathcal{P}(m, q, d)$,

$$Z \geq \mathbb{E}_u A_{h(u)}^u.$$

Proof. Complete G to a measurement by adjoining an extra outcome $\perp$. Apply Theorem 7.11 to the completed measurement, then discard the $\perp$ outcome. By Lemma 3.42, the completeness, consistency, strong self-consistency, and boundedness estimates transfer back to the original polynomial outcomes with an additive error from the missing mass $\langle \psi | (I - \sum_g G_g) \otimes I | \psi \rangle$. Since the output is projective, Lemma 3.36 converts the strong self-consistency statement to the usual bipartite formulation. $\square$

Remark 10.7 (Lean proof gap for induction self-improvement). The Lean declaration for Theorem 10.6 has the paper hypotheses, and its proof is still open. The conditional helper that used to assume Section 7 proof-stage data has been removed from the successor-step assembly. The ordinary and answer-valued slice constructors now call this theorem directly, so the missing Section 7 work remains a tracked proof gap on the paper statement rather than an additional hypothesis.

Theorem 10.8 (Pasting). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m + 1, q, d)$ low individual degree test. Let $\{G^x\}_{x \in \mathbb{F}_q}$ be projective submeasurements in $\text{PolySub}(m, q, d)$ with the following properties:*

1. (Completeness): If $G = \mathbb{E}_x \sum_g G_g^x$, then

$$\langle \psi | G \otimes I | \psi \rangle \geq 1 - \kappa.$$

2. (Consistency with A): On average over $(u, x) \sim \mathbb{F}_q^{m+1}$,

$$A_a^{u,x} \otimes I \simeq_\zeta I \otimes G_{[g(u)=a]}^x.$$

3. (Strong self-consistency): On average over $x \sim \mathbb{F}_q$,

$$G_g^x \otimes I \approx_\zeta I \otimes G_g^x.$$

4. (Boundedness): There are positive semidefinite operators Z^x such that

$$\mathbb{E}_x \langle \psi | (I - G^x) \otimes Z^x | \psi \rangle \leq \zeta$$

and for each $x \in \mathbb{F}_q$ and $g \in \mathcal{P}(m, q, d)$,

$$Z^x \geq \mathbb{E}_u A_{g(u)}^{u,x}.$$

Let $k \geq 400md$ and set

$$\nu = 100k^2m (\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32})$$

and

$$\sigma = \kappa \left(1 + \frac{1}{100m} \right) + 2\nu + e^{-k/(80000m^2)}.$$

Then there exists a measurement $H \in \text{PolyMeas}(m+1, q, d)$ with the following property:

1. (Consistency with A): On average over $u \sim \mathbb{F}_q^{m+1}$,

$$A_a^u \otimes I \simeq_\sigma I \otimes H_{[h(u)=a]}.$$

Proof. This is exactly Theorem 9.1, restated in the induction chapter because it is the pasting input used in the proof of Theorem 10.13. The Lean declaration linked above is source-facing but is not marked `\leanok`, since it inherits the remaining degree-zero pasting proof obligation tracked by issue #1622. $\square$

Remark 10.9 (Restricted Lean form of induction-section pasting). The Lean declaration linked here is the restricted nontrivial-regime restatement of Theorem 10.8. Its statement calls the formal Section 9 pasting theorem and therefore assumes, in addition to the source hypotheses,

$$\gamma \leq 1, \quad \zeta \leq 1, \quad d \leq q, \quad 0 < d, \quad 1 \leq k.$$

The corresponding paper theorem in `references/ldt-paper/ld-pasting.tex`, lines 12–50, does not include these assumptions; lines 52–55 record them only as a proof reduction to the nontrivial regime. The trivial complementary cases remain to be formalized before this declaration can be marked as the unrestricted source theorem.

Definition 10.10 (Self-improvement slice transport for the successor step). The successor step records the slice-wise self-improvement output in `SelfImprovementData`, and the answer-valued route records the analogous output in `AnswerSelfImprovementData`. Their constructors call `selfImprovementInInductionSection`; they no longer carry a separate bundle of Section 7 proof-stage data as an input. The `SliceStrategyTransport` structures are narrow formal interfaces: they assume concrete per-slice symmetric strategies and measurement-transport data, while averaged point-operator compatibility is derived structurally from point-measurement transport. They then convert the remaining data into the ordinary or answer-valued self-improvement output.

Definition 10.11 (Pasting data for the successor step). The successor step forms the pasting input by combining the restricted probability estimates with the slice-wise self-improvement data.

Lemma 10.12 (Base case of main induction). *Under the hypotheses of Theorem 10.13 with $m = 1$, there exists a measurement $G \in \text{PolyMeas}(1, q, d)$ such that, on average over $u \sim \mathbb{F}_q$,*

$$A_a^u \otimes I \simeq_\varepsilon I \otimes G_{[g(u)=a]},$$

which is stronger than the bound required by Theorem 10.13 for $m = 1$.

Proof. For $m = 1$, there is only one axis-parallel line ℓ in $\mathbb{F}_q$, so the line measurement B^ℓ is already an element of $\text{PolyMeas}(1, q, d)$. Because the strategy fails the axis-parallel lines test with probability at most ε , the consistency bound follows directly from the axis-parallel consistency hypothesis. $\square$

Theorem 10.13 (Main induction). *Let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the (m, q, d) low individual degree test. Let $k \geq md$ and set*

$$\nu = 1000k^2m^2 (\varepsilon^{1/1024} + \delta^{1/1024} + \gamma^{1/1024} + (d/q)^{1/1024})$$

and

$$\sigma = m^2 \left(\nu + e^{-k/(80000m^2)} \right).$$

Then there exists a measurement $G \in \text{PolyMeas}(m, q, d)$ such that, on average over $u \sim \mathbb{F}_q^m$,

$$A_a^u \otimes I \simeq_\sigma I \otimes G_{[g(u)=a]}.$$

Proof. In Lean, the base case is formalized through . The successor branch of is isolated as the source-shaped proof obligation . This declaration has the source assumptions of the successor branch, together with the branch condition $m \neq 1$; it does not take restricted-probability, recursive-slice, self-improvement, or pasting records as hypotheses. The nearby formalization still names the stage data that mirror the four paper stages restrict $\rightarrow$ induct $\rightarrow$ self-improve $\rightarrow$ paste, but no theorem with these data as additional source-level hypotheses is advertised as Theorem 10.13. For the answer-valued restriction route used in the successor branch of the main formal theorem, Lean now also exposes , which record the predecessor induction hypothesis for the restricted strategies without appealing to the public theorem ‘mainFormal’. The weighted restricted-probability estimates are now public via , The remaining proof gap for Theorem 10.13 is to derive the successor-step constructions, including the restricted-strategy self-improvement and slice-transport construction tracked by issues #1507 and #1503 under the bridge-debt tracker #1458, from the paper hypotheses; hence no proof-level is attached here. We may assume $\varepsilon, \delta, \gamma, d/q \leq 1$, since otherwise $\nu \geq 1$ and the bound is trivial.

The proof is by induction on m . The base case $m = 1$ is Lemma 10.12.

For the step from m to $m + 1$, let (ψ, A, B, L) be an $(\varepsilon, \delta, \gamma)$ -good symmetric strategy for the $(m + 1, q, d)$ test, and let $k \geq 400(m + 1)d$. For each $x \in \mathbb{F}_q$, let (ψ, A^x, B^x, L^x) be the x -restricted strategy, with failure parameters $\varepsilon_x, \delta_x, \gamma_x$. Apply the induction hypothesis to each restricted strategy with the same integer k and define

$$\nu_x = 1000k^2m^2 (\varepsilon_x^{1/1024} + \delta_x^{1/1024} + \gamma_x^{1/1024} + (d/q)^{1/1024}), \quad \sigma_x = m^2 (\nu_x + e^{-k/(80000m^2)}).$$

This yields a measurement $G^x \in \text{PolyMeas}(m, q, d)$ such that

$$(A^x)_a^u \otimes I \simeq_{\sigma_x} I \otimes G_{[g(u)=a]}^x.$$

Next define

$$\zeta_x = 3000m (\varepsilon_x^{1/32} + \delta_x^{1/32} + (d/q)^{1/32}).$$

Theorem 10.6 produces a projective submeasurement $\widehat{G}^x \in \text{PolySub}(m, q, d)$ such that, for each x :

1. if $\widehat{G}^x = \sum_g \widehat{G}_g^x$, then

$$\langle \psi | \widehat{G}^x \otimes I | \psi \rangle \geq (1 - \sigma_x) - \zeta_x;$$

2. on average over $u \sim \mathbb{F}_q^m$,

$$(A^x)_a^u \otimes I \simeq_{\zeta_x} I \otimes \widehat{G}_{[g(u)=a]}^x;$$

3.

$$\widehat{G}_g^x \otimes I \approx_{\zeta_x} I \otimes \widehat{G}_g^x;$$

4. there exists a positive semidefinite operator Z^x such that

$$\langle \psi | Z^x \otimes (I - \widehat{G}^x) | \psi \rangle \leq \zeta_x$$

and for each $g \in \mathcal{P}(m, q, d)$,

$$Z^x \geq \mathbb{E}_u (A^x)_{g(u)}^u.$$

We now average the slice-dependent parameters. By concavity of $\alpha \mapsto \alpha^c$ for $c \leq 1$ and Lemma 10.3,

$$\begin{aligned} \mathbb{E}_x \nu_x &\leq 1000k^2 m^2 ((\mathbb{E}_x \varepsilon_x)^{1/1024} + (\mathbb{E}_x \delta_x)^{1/1024} + (\mathbb{E}_x \gamma_x)^{1/1024} + (d/q)^{1/1024}) \\ &\leq 1000k^2 m^2 \left(\left(\frac{m+1}{m} \varepsilon \right)^{1/1024} + \delta^{1/1024} + \left(\frac{m+1}{m} \gamma \right)^{1/1024} + (d/q)^{1/1024} \right) \\ &\leq 1000k^2 (m+1)^2 (\varepsilon^{1/1024} + \delta^{1/1024} + \gamma^{1/1024} + (d/q)^{1/1024}). \end{aligned}$$

Call this final quantity ν . Then, defining

$$\sigma = m^2 \left(\nu + e^{-k/(80000m^2)} \right),$$

we have

$$\sigma \geq \mathbb{E}_x \sigma_x.$$

Similarly,

$$\begin{aligned} \mathbb{E}_x \zeta_x &\leq 3000m ((\mathbb{E}_x \varepsilon_x)^{1/32} + (\mathbb{E}_x \delta_x)^{1/32} + (d/q)^{1/32}) \\ &\leq 3000m \left(\left(\frac{m+1}{m} \varepsilon \right)^{1/32} + \delta^{1/32} + (d/q)^{1/32} \right) \\ &\leq 3000(m+1) (\varepsilon^{1/32} + \delta^{1/32} + (d/q)^{1/32}). \end{aligned}$$

Call this last quantity ζ . For later use,

$$\begin{aligned} \zeta &= 3000(m+1) (\varepsilon^{1/32} + \delta^{1/32} + (d/q)^{1/32}) \\ &\leq 1000k^2 (m+1)^2 (\varepsilon^{1/32} + \delta^{1/32} + (d/q)^{1/32}) \\ &\leq 1000k^2 (m+1)^2 (\varepsilon^{1/1024} + \delta^{1/1024} + \gamma^{1/1024} + (d/q)^{1/1024}) \\ &= \nu. \end{aligned} \tag{10.1}$$

The averaged slice family $\{\widehat{G}^x\}_{x \in \mathbb{F}_q}$ therefore satisfies the four hypotheses of Theorem 10.8 with completeness parameter $\kappa = \sigma + \zeta$. To verify the new ν bound in that theorem, observe that $(3000)^{1/32} \leq 2$, so

$$\zeta^{1/32} \leq 2(m+1) (\varepsilon^{1/1024} + \delta^{1/1024} + (d/q)^{1/1024}),$$

and hence

$$\begin{aligned} & 100k^2m(\varepsilon^{1/32} + \delta^{1/32} + \gamma^{1/32} + \zeta^{1/32} + (d/q)^{1/32}) \\ & \leq 1000k^2(m+1)^2(\varepsilon^{1/1024} + \delta^{1/1024} + \gamma^{1/1024} + (d/q)^{1/1024}) \\ & = \nu. \end{aligned}$$

Therefore Theorem 10.8 gives a pasted measurement $H \in \text{PolyMeas}(m+1, q, d)$ such that, on average over $u \sim \mathbb{F}_q^{m+1}$,

$$A_a^u \otimes I \simeq_{\sigma^*} I \otimes H_{[h(u)=a]},$$

where

$$\sigma^* = (\sigma + \zeta) \left(1 + \frac{1}{100m}\right) + 2\nu + e^{-k/(80000m^2)}.$$

It remains to compare σ^* with the next-stage target. The pasting parameter in Theorem 10.8 is bounded by $\nu/5$ in this regime. Using this sharper estimate and (10.1), we get

$$\begin{aligned} \sigma^* & \leq \left(m^2(\nu + e^{-k/(80000m^2)}) + \nu\right) \left(1 + \frac{1}{100m}\right) + \frac{2}{5}\nu + e^{-k/(80000m^2)} \\ & = \left((m^2+1)\left(1 + \frac{1}{100m}\right) + \frac{2}{5}\right)\nu + \left(m^2\left(1 + \frac{1}{100m}\right) + 1\right)e^{-k/(80000m^2)}. \end{aligned}$$

For every $m \geq 1$,

$$(m^2+1)\left(1 + \frac{1}{100m}\right) + \frac{2}{5} \leq (m+1)^2, \quad m^2\left(1 + \frac{1}{100m}\right) + 1 \leq (m+1)^2.$$

Therefore

$$\sigma^* \leq (m+1)^2(\nu + e^{-k/(80000m^2)}) \leq (m+1)^2(\nu + e^{-k/(80000(m+1)^2)}),$$

which is exactly the bound required for dimension $m+1$. $\square$

Remark 10.14 (Lean proof gap for the successor-step assembly). The Lean declaration for Theorem 10.13 has the source hypotheses and carries a `sorry` in the successor branch. The weighted slice bounds feeding Lemma 10.3 are still exposed through the public restriction-package constructor above, but the recursive slice witnesses, the slice-wise self-improvement outputs, and the pasting comparison must be derived inside the successor proof itself. They are not represented by a separate theorem whose assumptions are added to the paper statement.

10.3 Proof of the main formal theorem

Lemma 10.15 (Role-register symmetrization bridge). *Let $(\psi, A^A, B^A, L^A, A^B, B^B, L^B)$ be a projective strategy passing the (m, q, d) low individual degree test with probability at least $1 - \varepsilon$. Introducing a two-dimensional role register on each side and taking $(\psi_{\text{sym}}, A_{\text{sym}}, B_{\text{sym}}, L_{\text{sym}})$ as in paper Section 3 yields a symmetric projective strategy that is $(3\varepsilon, 3\varepsilon, 3\varepsilon)$ -good.*

Proof. Each of the three subtests (axis-parallel, self-consistency, diagonal) occurs with probability $1/3$, so the total failure probability at most ε implies each branch is at most 3ε . Block-diagonalizing the measurements over the role register preserves the axis-parallel and diagonal branch averages exactly, and the cross-register self-consistency branch of the symmetrized strategy equals the original point-agreement defect, which is itself bounded by 3ε . $\square$

Remark 10.16 (Lean auxiliary data for role-register symmetrization). The formal proof of Theorem 2.12 also uses auxiliary data containing the symmetrized strategy, its goodness proof, and its normalization proof. This auxiliary data supports the final assembly, not an additional hypothesis in Lemma 10.15.

Proposition 10.17 (Lean base-case handoff for the Section 3 proof). *If $m = 1$, then for every k the role-register symmetrization of a projective strategy passing the low individual degree test with error ε admits a measurement G satisfying the base-case conclusion of Theorem 10.13 with parameters $(3\varepsilon, 3\varepsilon, 3\varepsilon)$.*

Proof. This is a Lean-only construction theorem: combine Lemma 10.15 with the $m = 1$ clause of Theorem 10.13, formalized as . $\square$

Definition 10.18 (Lean successor restricted-recursion targets for the Section 3 proof). These Lean-only declarations name the restricted-probability and recursive-slice targets appearing in the successor comparison with Remark 10.14. The public weighted-bound helper lemmas discharge the weighted restricted-axis and restricted-diagonal fields from the symmetrized passing strategy. The former Lean boundary records, which also required a supplied self-improvement input, have been removed: that missing work is part of the proof of Theorem 10.13, not an additional assumption in Theorem 2.12. The former slice-data records, which supplied concrete pinned same-space slice strategies and passing hypotheses as Lean inputs, have also been removed. The construction of those slices and the transport of their restricted point, axis-parallel, and diagonal measurements should be proved in the successor branch rather than represented as a hypothesis in the statement of Theorem 2.12.

Definition 10.19 (Lean base-case Step 6 obligations for the Section 3 proof). The role-register Section 6 measurement is obtained by applying the paper-facing theorem `MainInductionStep.mainInduction` to the role-register symmetrization. The Lean constructors `MainFormalRoleInductionWitness.ofMainInduction` and `MainFormalRoleInductionWitness.ofMainInductionLargeK` record this application of the paper theorem; any successor proof gap is the tracked gap in `mainInduction` itself, not an added hypothesis of Theorem 2.12. The base-case role witness `mainFormalBaseRoleInductionWitness` remains a named specialization for $m = 1$.

From a line-130 consistency relation, the theorem `MainFormalDiagonalOrthonormalizationWitness.nonempty_ofConsistency` constructs the pre-completion projective submeasurements using the Section 5 locality-preserving repair construction. The completion theorem `MainFormalDiagonalCompletionWitness.nonempty_ofDiagonalConsistency` now derives the completion witness directly from the line-130 cross consistency and the orthonormalization closeness estimates. The exact construction-level match-mass monotonicity branch has been removed from the active ‘mainFormal’ path; instead, the checked repaired route with an explicit additional loss is recorded in Remark 4.6.

The remaining proof gap is not represented by diagonal-consistency, match-mass, or other auxiliary input hypotheses. The paper-facing theorem itself remains unfinished. Once a `MainFormalProjectiveCompletionTransportWitness` has been constructed from the paper hypotheses, the proved theorem `mainFormal_ofProjectiveCompletionTransportWitness` derives the three consistency conclusions for Theorem 2.12. The witness is a construction target inside the proof, not an additional assumption in the public statement.

Remark 10.20 (Successor projective-completion construction). The remaining successor-case proof work is not named as an additional Lean hypothesis of Theorem 2.12. Starting from a projective strategy that passes the low individual degree test, the large- k bound $k \geq 400md$ from Section 6, the positivity condition $d > 0$, and the non-base condition $m \neq 1$, the proof must construct the projective-completion witness consumed by `mainFormal_ofProjectiveCompletionTransportWitness`. This is part of the unfinished proof of the paper theorem, not a separate assumption in its statement.

Remark 10.21 (Lean measurement-unsymmetrization consistency). Lean now isolates the Step 3 extraction from `inductive_step.tex` lines 84–108. The principal-block map `extractRoleBlock` turns a role-register POVM G into the two POVMs G^A and G^B on the original prover space and proves that point-evaluation postprocessing commutes with this extraction. The two round-trip lemmas identify the A and B blocks of the symmetrized point measurement with the original prover point measurements. The remaining analytic factor-two estimates (10.3) and (10.4) are named by `UnsymmetrizationConsistency`; no projectivity of arbitrary principal blocks is claimed, matching the paper’s later projectivization step.

Definition 10.22 (Final error envelope). Set

$$E = \varepsilon^{1/40000} + (d/q)^{1/40000} + e^{-k/(2560000m^2)}.$$

Theorem 10.23 (Basic facts about the final error envelope). *If $0 \leq \varepsilon$, then*

$$\nu_{\text{main}} = 100000k^2m^4 \left(\varepsilon^{1/40000} + (d/q)^{1/40000} + e^{-k/(2560000m^2)} \right) = 100000k^2m^4E,$$

and $E \geq 0$.

Proof. Both identities are by definition: the first unfolds ν_{main} and the second unfolds E . Non-negativity of E follows summand-wise from `Real.rpow_nonneg` (using $0 \leq \varepsilon$) and `Real.exp_nonneg`. $\square$

Definition 10.24 (Cascade parameters and standing hypotheses). Assume $q > 0$, $k \geq 1$, $m \geq 1$, $0 \leq \varepsilon \leq 1$, and $d \leq q$. Fix an incoming error parameter ν . Define

$$\begin{aligned} \sigma &= m^2 \left(\nu + e^{-k/(80000m^2)} \right), & \zeta_1 &= 2\sigma + 2\sqrt{3\varepsilon + 2\sigma} + md/q, \\ \zeta_2 &= 200\zeta_1^{1/4} + 42\zeta_1^{1/8}, & \zeta_3 &= 6\zeta_1 + 6\zeta_2, & \zeta_4 &= 2\sigma + 2\sqrt{\zeta_1 + \zeta_3/2}. \end{aligned}$$

The coefficient 42 is the Lean cascade scalar; it widens the paper’s printed coefficient 40 so that the completion theorem’s extra $2\zeta_1$ term is absorbed while preserving the final error envelope.

Theorem 10.25 (Bound for the first cascade term). *Under the standing hypotheses, if*

$$\nu \leq 10000k^2m^2 \left(\varepsilon^{1/1024} + (d/q)^{1/1024} \right),$$

then

$$\sigma \leq 10000k^2m^4E.$$

Proof. Multiply the assumed bound by m^2 , replace m^2 by m^4 , coarsen the exponents from $1/1024$ to $1/40000$, and enlarge the exponential denominator from $80000m^2$ to $2560000m^2$. $\square$

Theorem 10.26 (Cascade bounds for the later cascade terms). *Under the standing hypotheses, assume*

$$0 \leq \nu \leq 10000k^2m^2 \left(\varepsilon^{1/1024} + (d/q)^{1/1024} \right).$$

Define

$$\begin{aligned} \sigma &= m^2 \left(\nu + e^{-k/(80000m^2)} \right), \\ \zeta_1 &= 2\sigma + 2\sqrt{3\varepsilon + 2\sigma} + md/q, & \zeta_2 &= 200\zeta_1^{1/4} + 42\zeta_1^{1/8}, \\ \zeta_3 &= 6\zeta_1 + 6\zeta_2, & \zeta_4 &= 2\sigma + 2\sqrt{\zeta_1 + \zeta_3/2}. \end{aligned}$$

Then

$$\zeta_1, \zeta_2, \zeta_4 \leq 100000k^2m^4E, \quad \zeta_3 \leq 2 \cdot 100000k^2m^4E.$$

Proof. The Lean declarations first derive the native bounds for $\zeta_1, \dots, \zeta_4$ from the cascade definitions using square-root and *rpow* monotonicity, with ζ_2 using the widened coefficient 42, and then coarsen those bounds to the common envelope E . The ζ_2 proof additionally uses $k \leq k^2$, $m \leq m^4$, and the nonnegativity of the earlier cascade variables. $\square$

Theorem 10.27 (Consolidated error cascade). *Assume the standing hypotheses and*

$$0 \leq \nu \leq 10000k^2m^2(\varepsilon^{1/1024} + (d/q)^{1/1024}).$$

Let $\sigma, \zeta_1, \zeta_2, \zeta_3$ be the cascade quantities defined above. Then

$$\sigma \leq 100000k^2m^4E, \quad \zeta_1, \zeta_2 \leq 100000k^2m^4E, \quad \zeta_3 \leq 2 \cdot 100000k^2m^4E,$$

and the concrete quantity

$$\zeta_4 = 2\sigma + 2\sqrt{\zeta_1 + \zeta_3/2}$$

also satisfies

$$\zeta_4 \leq 100000k^2m^4E.$$

Proof. This is the direct combination of Theorem 10.25 and Theorem 10.26. The Lean declaration uses the explicit equalities naming $\sigma, \zeta_1, \zeta_2, \zeta_3$ as the corresponding cascade expressions. $\square$

Remark 10.28 (Lean Step 5 expansion theorem). For the Step 5 passage below, Lean now names the weighted polynomial-collision mass, proves its md/q bound by specializing the existing tensor Schwartz–Zippel helper, and proves the algebraic expansion/reindexing statement comparing the evaluated consistency defect with the full-polynomial defect plus that collision mass. The theorem for Step 5 therefore no longer takes the expansion statement as an external hypothesis.

Remark 10.29 (Lean witness structures for the main-formal assembly). The following Lean declarations are bookkeeping infrastructure for the final assembly witness: They split off the vacuous branch where the theorem error is at least 1, derive the Step 8 scalar hypotheses in the non-vacuous branch, and isolate the role-register Section 6 measurement behind a concrete role-witness record. That record can now be obtained directly from the `mainInduction` theorem by `MainFormalRoleInductionWitness.ofMainInductionLargeK`; the successor case is therefore a tracked `sorry` in Section 6, not a separate boundary hypothesis in Section 3. Lean also records that every non-base parameter choice has a predecessor via `Parameters.successorDecompositionOfNeOne`, and it retains the explicit transport helpers for the strategy and passing proof across that predecessor equality. The older branch-witness records, which assembled ordinary or answer-valued successor data from recursive slice witnesses and supplied self-improvement inputs, have been removed. The current completion witness carries the concrete role witness obtained from the Section 6 theorem rather than a separate decorative branch witness. Once the role-measurement record is obtained, Lean proves the two factor-two unsymmetrization links from its symmetrized consistency estimate. The subsequent proved lemmas use the proved Section 5 Schwartz–Zippel theorem to convert evaluated pre-projective consistency to full-polynomial consistency at ζ_1 , prove the paper line-116 triangle step, and reconstruct the line-156 $\approx_{\zeta_3}$ projectivization bridge from pre-projective consistency plus left/right completion closeness at ζ_2 . Lean also derives the evaluated line-164 projective consistency. The error-parametric lemmas `pointAConsistency_ofLine169Consistency` and `pointBConsistency_ofLine169Consistency` show that a polynomial line-169 estimate with error η yields the final point error

$$2\sigma + 2\sqrt{\eta + \zeta_3/2}.$$

The paper-tight final route specializes this with $\eta = \zeta_1$; the checked local repair can be inserted with its explicit larger value of η if the scalar cascade is widened accordingly. The existing `pointAConsistency` and `pointBConsistency` declarations are the paper-tight specializations. After the #869 right-register completion transport, the remaining active post-role witness asks only for Alice's line-146 completion closeness, Bob's left-register completion closeness as returned by `orthonormalize-and-complete`, and the construction-level match-mass monotonicity invariant needed for the exact polynomial-level ζ_1 transport links from paper line 169 and its role-reversed analogue. Lean reconstructs pre-projective full-polynomial consistency from the concrete role witness, uses the match-mass invariant to derive the two line-169 links, and transports the Bob-side left-register completion estimate to the line-147 right-register form before converting directly to the projective completion witness used by the final assembly. It no longer asks for an arbitrary role-measurement record or an arbitrary raw Section 6 witness as an independent field. It also no longer asks for the two factor-two role-block estimates, the unused Section 6 consistency field inside `UnsymmetrizationConsistency`, or the duplicated pre-projective consistency field inside the Section 6 handoff; those are rebuilt from the concrete role witness, the selected role-measurement record, and `hpass`. Lean then applies the projective converse of `simeq-to-approx` directly inside `MainFormalProjectiveCompletionTransportWitness.selfConsistency`.

The exact line-169 transport fields are now derived from the stored match-mass monotonicity invariant rather than being independent witness data. As stated in `preliminaries.tex`, `prop:triangle-sub` gives a $\delta + \sqrt{\epsilon}$ loss, so applying it directly to $G^A \simeq_{\zeta_1} G^B$ and $G^A \approx_{\zeta_2} Q^A$ yields $Q^A \simeq_{\zeta_1 + \sqrt{\zeta_2}} G^B$, not the printed ζ_1 estimate. The generic corrected route would degrade the final point error to scale $t^{1/65536}$, outside the current `mainFormalError` envelope. The sharper local pre-completion repair from Remark 4.6 avoids that widening and is part of the match-mass witness route. No proof-level completeness tag is claimed here while `mainFormal` itself has tracked proof debt for constructing the projective-completion witness from the paper hypotheses: the Section 6 successor branch is tracked in issue #1458.

Proof. Let

$$(\psi, A^A, B^A, L^A, A^B, B^B, L^B)$$

be a projective strategy passing the (m, q, d) low individual degree test with probability at least $1 - \epsilon$. Since the three subtests occur with probability $1/3$, this strategy is $(3\epsilon, 3\epsilon, 3\epsilon)$ -good.

Introduce a two-dimensional role register on each side and define the symmetrized state

$$|\psi_{\text{sym}}\rangle = \frac{1}{\sqrt{2}} (|0\rangle_{A'} |1\rangle_{B'} |\psi\rangle_{AB} + |1\rangle_{A'} |0\rangle_{B'} |\psi_{\text{swap}}\rangle_{AB}).$$

Define the symmetrized point measurement by

$$(A_{\text{sym}})_a^u = |0\rangle\langle 0| \otimes A_a^{A,u} + |1\rangle\langle 1| \otimes A_a^{B,u},$$

and similarly for B_{sym} and L_{sym} . This yields a symmetric $(3\epsilon, 3\epsilon, 3\epsilon)$ -good strategy. Set

$$\nu = 10000k^2m^2 (\epsilon^{1/1024} + (d/q)^{1/1024}), \quad \sigma = m^2 (\nu + e^{-k/(80000m^2)}).$$

Theorem 10.13 applied to the symmetrized strategy produces a measurement $G = \{G_g\} \in \text{PolyMeas}(m, q, d)$ such that

$$(A_{\text{sym}})_a^u \otimes I \simeq_{\sigma} I \otimes G_{[g(u)=a]}, \quad G_{[g(u)=a]} \otimes I \simeq_{\sigma} I \otimes (A_{\text{sym}})_a^u. \quad (10.2)$$

Now unsymmetrize by defining

$$G_g^A = (\langle 0| \otimes I) G_g (|0\rangle \otimes I), \quad G_g^B = (\langle 1| \otimes I) G_g (|1\rangle \otimes I).$$

These are POVMs on the original prover space. Since A_{sym} is a convex combination of $(|0\rangle\langle 0| \otimes I)$ and $(|1\rangle\langle 1| \otimes I)$ weighted by the test probabilities, the cross-terms $\langle 0| \cdot A_{\text{sym}} \cdot |1\rangle$ vanish. Expanding the consistency of G gives

$$G_{[g(u)=a]}^A \otimes I \simeq_{2\sigma} I \otimes A_a^{B,u}, \quad (10.3)$$

$$I \otimes G_{[g(u)=a]}^B \simeq_{2\sigma} A_a^{A,u} \otimes I. \quad (10.4)$$

Since the original strategy is $(3\varepsilon, 3\varepsilon, 3\varepsilon)$ -good,

$$A_a^{A,u} \otimes I \simeq_{3\varepsilon} I \otimes A_a^{B,u}.$$

Therefore Lemma 3.29 yields

$$G_{[g(u)=a]}^A \otimes I \simeq_{2\sigma+2\sqrt{3\varepsilon+2\sigma}} I \otimes G_{[g(u)=a]}^B.$$

Expanding this consistency relation and applying Lemma 3.7 to replace point evaluations by polynomial evaluations (at cost md/q) gives

$$G_g^A \otimes I \simeq_{\zeta_1} I \otimes G_g^B, \quad (10.5)$$

where

$$\zeta_1 = 2\sigma + 2\sqrt{3\varepsilon + 2\sigma} + \frac{md}{q}.$$

The Lean Step 5 theorem formalizes this expansion/reindexing step and combines it with the proved Schwartz–Zippel tensor bound.

Apply Lemma 4.9 to 10.5. This gives projective submeasurements $P^A, P^B \in \text{PolySub}(m, q, d)$ with

$$G_g^A \otimes I \approx_{100\zeta_1^{1/4}} P_g^A \otimes I, \quad I \otimes G_g^B \approx_{100\zeta_1^{1/4}} I \otimes P_g^B.$$

Completing them by Lemma 3.42 yields projective measurements $Q^A, Q^B \in \text{PolyMeas}(m, q, d)$ such that

$$\begin{aligned} G_g^A \otimes I &\approx_{\zeta_2} Q_g^A \otimes I, \\ I \otimes G_g^B &\approx_{\zeta_2} I \otimes Q_g^B, \end{aligned} \quad (10.6)$$

where

$$\zeta_2 = 200\zeta_1^{1/4} + 42\zeta_1^{1/8}.$$

The paper prints the coefficient 40 in this display. The formal Lean cascade uses the widened coefficient 42 to absorb the literal completion error $2 \cdot (100\zeta_1^{1/4}) + 4\sqrt{100\zeta_1^{1/4}} + 2\zeta_1$: under the non-vacuous scalar regime $0 \leq \zeta_1 \leq 1$, the extra term $2\zeta_1$ is bounded by $2\zeta_1^{1/8}$.

By 10.5 and Lemma 3.21,

$$G_g^A \otimes I \approx_{2\zeta_1} I \otimes G_g^B.$$

Hence Lemma 3.27 gives

$$Q_g^A \otimes I \approx_{\zeta_3} I \otimes Q_g^B,$$

where

$$\zeta_3 = 6\zeta_1 + 6\zeta_2.$$

Because both measurements are projective, Lemma 3.21 converts this to

$$Q_g^A \otimes I \simeq_{\zeta_3/2} I \otimes Q_g^B. \quad (10.7)$$

By data processing,

$$Q_{[g(u)=a]}^A \otimes I \simeq_{\zeta_3/2} I \otimes Q_{[g(u)=a]}^B. \quad (10.8)$$

Next, Lemma 3.20, applied to 10.5 and 10.6, implies

$$Q_g^A \otimes I \simeq_{\zeta_1} I \otimes G_g^B.$$

Data processing then gives

$$Q_{[g(u)=a]}^A \otimes I \simeq_{\zeta_1} I \otimes G_{[g(u)=a]}^B. \quad (10.9)$$

Applying Lemma 3.29 to 10.4, 10.9, and 10.8 yields

$$A_a^{A,u} \otimes I \simeq_{\zeta_4} I \otimes Q_{[g(u)=a]}^B, \quad (10.10)$$

where

$$\zeta_4 = 2\sigma + 2\sqrt{\zeta_1 + \zeta_3/2}.$$

The same argument with the roles reversed gives

$$I \otimes A_a^{B,u} \simeq_{\zeta_4} Q_{[g(u)=a]}^A \otimes I. \quad (10.11)$$

Remark 10.29 records the Lean witness structures that separate the scalar cascade and the Step 5 Schwartz–Zippel handoff from the remaining projective-stage transport work. The scalar side discharges the paper’s line 71–73 coarsening and the Step 8 hypotheses in the non-vacuous branch; the Step 5 theorem now turns the evaluated estimate at line 116 into full-polynomial consistency at ζ_1 . The remaining Lean witness is the completion-and-line-169 construction: supply the actual role-register measurement, Alice’s line-146 completion estimate, Bob’s left-register completion estimate, and the two exact polynomial-level ζ_1 links. Lean transports the Bob-side completion estimate to the line-147 right-register form, reconstructs the duplicated Step 6 pre-projective consistency field from these inputs and `hpass`, then data-processes the line-169 links to line 172 and derives the two ζ_4 point goals and the $\zeta_3/2$ self-consistency target.

With E as in Definition 10.22, the quantities $\sigma, \zeta_1, \zeta_2, \zeta_3, \zeta_4$ form the cascade of Definition 10.24. Theorem 10.25 and Theorem 10.26 give the final inflation to the scale $100000k^2m^4E$. The underlying estimates, with the formal widened ζ_2 , are:

$$\sigma \leq 10000k^2m^4 \left(\varepsilon^{1/1024} + (d/q)^{1/1024} + e^{-k/(80000m^2)} \right).$$

Then

$$\zeta_1 \leq 20204k^2m^4 \left(\varepsilon^{1/2048} + (d/q)^{1/2048} + e^{-k/(160000m^2)} \right),$$

and

$$\zeta_2 \leq 2568km \left(\varepsilon^{1/16384} + (d/q)^{1/16384} + e^{-k/(1280000m^2)} \right).$$

Consequently,

$$\zeta_3 \leq 150000k^2m^4 \left(\varepsilon^{1/16384} + (d/q)^{1/16384} + e^{-k/(1280000m^2)} \right),$$

and

$$\zeta_4 \leq 40000k^2m^4 \left(\varepsilon^{1/32768} + (d/q)^{1/32768} + e^{-k/(2560000m^2)} \right).$$

Both $\zeta_3/2$ and ζ_4 are therefore bounded by

$$100000k^2m^4 \left(\varepsilon^{1/40000} + (d/q)^{1/40000} + e^{-k/(2560000m^2)} \right).$$

Hence 10.10, 10.11, and 10.7 give the three claimed conclusions, with Q^A and Q^B renamed as the theorem’s G^A and G^B . $\square$

Bibliography

- [Sch80] Jacob T. Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *Journal of the ACM*, 27(4):701–717, 1980.
- [Zip79] Richard Zippel. Probabilistic algorithms for sparse polynomials. In *Proceedings of the 2nd International Symposium on Symbolic and Algebraic Manipulation*, pages 216–226, 1979.