

Fundamental Theorem of Matrix Product States

A Formalization Blueprint

Authors

March 16, 2026

Contents

1	Introduction	4
2	Matrix Product Vectors	5
2.1	Basic definitions	5
2.2	Gauge equivalence and same MPV	6
2.3	Injectivity and normality	7
2.4	Canonical form	7
2.5	Blocking	8
2.6	MPV overlap	9
3	The Single-Block Fundamental Theorem	10
3.1	Trace pairing	10
3.2	Linear extension	11
3.3	Simplicity and Skolem–Noether	11
3.4	Assembly	12
4	Quantum Channels and Positive Maps	13
4.1	Positive maps and channels	13
4.2	Irreducibility	14
4.3	Cesàro fixed-point theorem	14
4.4	Fixed-point projection	15
4.5	Peripheral spectrum and primitivity	15
4.5.1	Periodicity removal by powering	16
4.5.2	Peripheral closure via adjoint fixed point	17
4.6	Representations: infrastructure	18
4.7	Choi–Jamiołkowski isomorphism (Wolf Prop. 2.1)	19
4.8	Kraus representation theorem (Wolf Thm. 2.1)	19
4.9	Stinespring dilation (Wolf Thm. 2.2)	20
4.10	Quantum dynamical semigroups	20
5	Schwarz Inequalities and Multiplicative Domains	21
5.1	Kadison–Schwarz inequality	21
5.2	Multiplicative domain	22
5.2.1	Full multiplicative-domain structure	23
5.2.2	Schwarz inequality for normal operators	24
5.2.3	Positive maps preserve order and spectral intervals	24
5.2.4	Wolf Example 5.3	25

6	Perron–Frobenius Theory for Channels and Transfer Maps	26
6.1	Positive definiteness	26
6.2	Uniqueness	27
6.3	Existence and assembly	28
6.4	Right- and left-canonical gauges	28
6.5	Similarity preserves irreducibility	28
6.6	Perron–Frobenius eigenvector existence	29
6.7	Exponential positivity for irreducible CP maps	30
6.8	Ergodicity of irreducible channels	30
6.9	Spectral radius at the Perron eigenvalue	31
6.10	Spectral characterization of irreducibility	31
7	Spectral Gap and Block Separation	33
7.1	Mixed transfer operator	33
7.2	MPV overlap as transfer trace	34
7.3	Eigenvalue bound and spectral gap	35
7.4	Spectral gap — rectangular case	35
7.5	MPV overlap decay	36
7.6	Block separation	36
7.7	Primitive overlap convergence (spectral-gap form)	36
8	Wielandt Bound	38
8.1	Cumulative span	38
8.2	Fitting decomposition	39
8.3	Nonzero trace product	39
8.4	Eigenvector extraction	40
8.5	Eigenvector spreading	40
8.6	Assembly	41
8.7	Lemma 2(b): fixed-length matrix spanning	42
8.8	Blocked tensor and rectangular span	43
8.9	Primitive MPS tensors	44
8.10	Primitivity and normality	45
	8.10.1 Primitivity-to-normality bridge	45
	8.10.2 Primitive implies irreducible	46
	8.10.3 Primitivity bridge to strong irreducibility and normality	46
8.11	Cumulative span to word span	47
9	Canonical Form Reduction	49
9.1	Block-diagonal assembly	49
9.2	Transfer map normalization	50
9.3	Invariant subspace decomposition	50
9.4	Iterated reduction	51
	9.4.1 Burnside bridge: irreducibility and spanning	52
9.5	Proportional single-block theorem	53
9.6	Canonical form from peripheral primitivity	53
9.7	Normal canonical form	54
9.8	Steps toward canonical form existence	55
9.9	Zero-block separation and honest block decomposition	56
9.10	Arbitrary-input TP-gauge pipeline	57

9.11	Blocking infrastructure	57
9.12	Cyclic sector decomposition	58
9.13	Assembly: arbitrary tensor to primitive block decomposition	58
9.14	Open directions	59
10	Block Permutation and Separation	60
10.1	Block permutation algebra	60
10.2	Product algebra linear extension	61
10.3	Block separation	61
11	Bases of Normal Tensors	64
11.1	Bases of normal tensors	64
11.2	Permutation rigidity for bases of normal tensors	65
11.3	Coefficient convergence	66
11.4	Newton–Girard identities	67
12	Full Assembly	68
12.1	Weak Fundamental Theorem: pipeline end-to-end	68
12.2	Proportional Fundamental Theorem	69
12.3	Equal-MPV Fundamental Theorem	70
12.4	Periodic Fundamental Theorem (de las Cuevas–Cirac–Schuch–Pérez-García)	71
12.4.1	Irreducible form	71
12.4.2	The Z -gauge degree of freedom	72
12.4.3	Statement of the periodic Fundamental Theorem	72
13	Quantum Dynamical Semigroups	74
13.1	Dynamical semigroups and the exponential form	74
13.2	Perturbation theory	75
13.3	GKSL/Lindblad generators	75
13.3.1	Generator decomposition and conditional complete positivity	76
13.3.2	The Lindblad form	76
13.3.3	Prop. 7.2: Characterization of conditional complete positivity	77
13.3.4	Prop. 7.3: CP semigroups and CCP generators	77
13.3.5	Prop. 7.4: Freedom in generator representation	77
13.3.6	Thm. 7.1: The GKSL/Lindblad theorem	77
13.3.7	Kossakowski matrix form	78

Chapter 1

Introduction

This blueprint describes the fundamental theorem for periodic, translation-invariant matrix product vectors. The basic object is a family of matrices $\{A^i\}$, viewed as a tensor that generates vectors $|V^{(N)}(A)\rangle$ on chains of length N .

Chapters 2 and 3 describe the algebraic core. Chapter 2 introduces word evaluation, MPV coefficients, gauge equivalence, injectivity, blocking, overlap, and the block-injective canonical-form data. Chapter 3 proves the single-block fundamental theorem by a purely algebraic route through trace pairings, linear extension, simplicity of matrix algebras, and Skolem–Noether.

The later chapters develop the spectral and canonical-form machinery needed for the full multi-block theorem. Chapter 4 studies positive maps and quantum channels. Chapters 6 and 7 supply Perron–Frobenius theory and overlap decay. Chapters 8 and 9 connect blocking, primitivity, irreducibility, and the normal/canonical-form reductions. Chapters 10 and 11 carry out block permutation and BNT theory; Chapter 12 states the final assembly.

Chapter 2

Matrix Product Vectors

Fix a physical dimension d and a bond dimension D . This chapter studies the tensors that generate matrix product vector (MPV) families. All tensors in this blueprint are translation-invariant with periodic boundary conditions (PBC).

2.1 Basic definitions

Definition 2.1 (MPS tensor). A (translation-invariant, PBC) *MPS tensor* with physical dimension d and bond dimension D is a collection of matrices

$$\{A^i\}_{i=0}^{d-1}, \quad A^i \in M_D(\mathbb{C}),$$

indexed by a physical index $i \in \{0, \dots, d-1\}$. Such a tensor defines an MPV family.

Definition 2.2 (Word evaluation). Given a word $w = (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$, the *word evaluation* is the matrix product

$$A^w := A^{i_1} A^{i_2} \dots A^{i_L} \in M_D(\mathbb{C}),$$

with the convention that the empty word evaluates to the identity: $A^\emptyset = \mathbb{1}_D$.

Lemma 2.3 (Word evaluation respects concatenation). *For any words w_1, w_2 , $A^{w_1 w_2} = A^{w_1} A^{w_2}$.*

Proof. Immediate from the definition of A^w . $\square$

Definition 2.4 (Matrix product vector). The *matrix product vector* (MPV) of a tensor A at system size N is the vector

$$|V^{(N)}(A)\rangle = \sum_{i_1, \dots, i_N=0}^{d-1} \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}) |i_1, \dots, i_N\rangle \in (\mathbb{C}^d)^{\otimes N}.$$

Equivalently, for a configuration $\sigma = (i_1, \dots, i_N) \in \{0, \dots, d-1\}^N$, we write

$$V^{(N)}(A)_\sigma := \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}).$$

The coefficient function $\sigma \mapsto V^{(N)}(A)_\sigma$ gives the components; the displayed ket is the corresponding vector in $(\mathbb{C}^d)^{\otimes N}$. For a general word w , we also write

$$c_w(A) := \text{tr}(A^w),$$

The *MPV family* generated by A is the collection $\mathcal{V}(A) = \{|V^{(N)}(A)\rangle\}_{N \geq 1}$.

Definition 2.5 (Transfer map). The *transfer map* associated to a tensor A is the linear map $\mathcal{E}_A : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$\mathcal{E}_A(X) = \sum_{i=0}^{d-1} A^i X (A^i)^\dagger.$$

Remark 2.6. The transfer map is automatically positive: if $X \geq 0$ then $\mathcal{E}_A(X) \geq 0$, because each summand $A^i X (A^i)^\dagger$ is positive semidefinite. The abstract positive-map framework appears in Chapter 4.

2.2 Gauge equivalence and same MPV

Definition 2.7 (Gauge equivalence). Two tensors A and B of the same bond dimension D are *gauge equivalent* if there exists an invertible matrix $X \in \text{GL}_D(\mathbb{C})$ such that

$$B^i = X A^i X^{-1} \quad \text{for all } i \in \{0, \dots, d-1\}.$$

Definition 2.8 (Same MPV). Two tensors A and B (of the same bond dimension) *generate the same MPV family* if $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every system size $N \geq 1$.

Definition 2.9 (Same MPV — different bond dimensions). For tensors A and B of possibly different bond dimensions D_1 and D_2 , we write $\mathcal{V}(A) = \mathcal{V}(B)$ if $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every $N \geq 1$.

Remark 2.10. Definition 2.8 is the same-bond-dimension specialization of Definition 2.9. We keep both because later results use both the same-dimension and different-dimension forms.

Definition 2.11 (Proportional MPV). Two tensors A and B (of possibly different bond dimensions) *generate proportional MPV families* if for every N there exists $c_N \in \mathbb{C}$ such that $|V^{(N)}(A)\rangle = c_N |V^{(N)}(B)\rangle$.

Lemma 2.12 (Word evaluation under scaling). *Let ζA denote the tensor with matrices $\{\zeta A^i\}_{i=0}^{d-1}$. Then for any word w ,*

$$(\zeta A)^w = \zeta^{|w|} A^w,$$

where $|w|$ is the length of w .

Proof. Induction on w : the base case gives $\zeta^0 = 1$, and each step picks up one factor of ζ . $\square$

Definition 2.13 (Gauge-phase equivalence). Two tensors A and B are *gauge-phase equivalent* if there exist $X \in \text{GL}_D(\mathbb{C})$ and $\zeta \in \mathbb{C} \setminus \{0\}$ such that

$$B^i = \zeta X A^i X^{-1} \quad \text{for all } i.$$

Remark 2.14. After spectral-radius normalization one may restrict ζ to a unit-modulus phase $e^{i\phi}$. We allow an arbitrary nonzero scalar because later canonical-form statements keep the block-scaling factors explicit.

Lemma 2.15 (Word evaluation under conjugation). *If $B^i = X A^i X^{-1}$ for all i , then for every word w ,*

$$B^w = X A^w X^{-1}.$$

Proof. Induction on the word length, using $XX^{-1} = \mathbb{1}$. $\square$

Theorem 2.16 (Gauge invariance of MPVs). *If A and B are gauge equivalent, then they generate the same MPV family.*

Proof. Let $B^i = X A^i X^{-1}$. By Lemma 2.15, $B^w = X A^w X^{-1}$ for every word w . Then $\text{tr}(B^w) = \text{tr}(X A^w X^{-1}) = \text{tr}(A^w)$ by cyclicity of the trace. $\square$

2.3 Injectivity and normality

Definition 2.17 (Injectivity). A tensor A is *injective* if its matrices span the full matrix algebra: $\text{span}_{\mathbb{C}}\{A^i : i = 0, \dots, d-1\} = M_D(\mathbb{C})$.

Definition 2.18 (L -block injectivity). A tensor A is *L -block injective* if $\text{span}_{\mathbb{C}}\{A^{i_1} \dots A^{i_L} : (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L\} = M_D(\mathbb{C})$. An injective tensor is 1-block injective.

Definition 2.19 (Normal tensor). A tensor A is *normal* [CPGSV21] if there exists a finite L_0 such that A is L_0 -block injective.

Remark 2.20. The algebraic characterization of normality in Definition 2.19 — eventual full Kraus rank, equivalently eventual block injectivity — is equivalent to the spectral characterization used in [CPGSV21, Definition IV.1]: after TP normalization, the transfer map $\mathcal{E}_A$ is a primitive channel, equivalently irreducible with trivial peripheral spectrum. This equivalence is proved in [SPGWC10, Proposition 3]. Within this blueprint, the directions of the equivalence are supplied by Theorem 8.49, Theorem 8.59, and Theorem 9.27. In particular, every subsequent use of “normal” in this blueprint refers to this single notion; Definition 9.29 employs the spectral formulation, but it describes the same class of tensors.

2.4 Canonical form

Definition 2.21 (Block-injective canonical form). In this blueprint, a *block-injective canonical form* for a tensor consists of:

1. a number of blocks $r \geq 1$,
2. bond dimensions $D_1, \dots, D_r$,
3. injective tensors $\{A_k^i\}_{i=0}^{d-1}$ with $A_k^i \in M_{D_k}(\mathbb{C})$ for each block $k \in \{1, \dots, r\}$,
4. scaling factors $\mu_1, \dots, \mu_r \in \mathbb{C}$.

The associated tensor is the block-diagonal tensor

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

This is the block-injective version of the canonical-form data; see [PGVWC07] and [CPGSV21].

Remark 2.22. This blueprint uses the block-injective (biCF-style) canonical form: each block is injective in the sense of Definition 2.17. This is stronger than the NT-block canonical form of [CPGSV21], where the blocks are only normal in the paper’s spectral sense. The paper-faithful NT/BNT/CFII route is developed later in Chapter 9; the present section records the stronger block-injective package used in the later block-injective assembly.

Definition 2.23 (Block-diagonal tensor from blocks). Given r blocks with bond dimensions $(D_k)_{k=1}^r$, per-block tensors $\{A_k^i\}$, and scaling factors $(\mu_k)_{k=1}^r$, the associated *block-diagonal tensor* is the tensor of total bond dimension $D = \sum_{k=1}^r D_k$ defined by

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

Theorem 2.24 (MPV decomposition). *The MPV of a block-diagonal tensor decomposes as*

$$|V^{(N)}(A)\rangle = \sum_{k=1}^r \mu_k^N |V^{(N)}(A_k)\rangle.$$

Equivalently, for each configuration σ one has

$$V^{(N)}(A)_\sigma = \sum_{k=1}^r \mu_k^N V^{(N)}(A_k)_\sigma.$$

Proof. For each physical index i , the block-diagonal tensor $A^i = \bigoplus_k \mu_k A_k^i$ is a block-diagonal matrix. By Lemma 2.12, the word evaluation of each block picks up a factor $\mu_k^{|w|}$, and the full word evaluation is block-diagonal with blocks $\mu_k^N A_k^w$. Taking the trace of a block-diagonal matrix gives the sum of the block traces: $\text{tr}(A^w) = \sum_k \mu_k^N \text{tr}(A_k^w)$. $\square$

2.5 Blocking

Definition 2.25 (Blocked tensor). Given a tensor A and a blocking length L , the *L -blocked tensor* is the tensor with physical index set $\{0, \dots, d-1\}^L$ (hence physical dimension d^L) and the same bond dimension D , whose matrices are indexed by words $(i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$:

$$(A^{[L]})^{(i_1, \dots, i_L)} = A^{i_1} A^{i_2} \dots A^{i_L}.$$

Lemma 2.26 (Blocked word evaluation). *If $w = (J_1, \dots, J_m)$ is a word in the blocked alphabet, where each J_k is an L -tuple in $\{0, \dots, d-1\}^L$, then concatenating the block words gives a word $\tilde{w}$ of length mL in the original alphabet and one has*

$$(A^{[L]})^w = A^{\tilde{w}}.$$

Proof. Induct on the blocked word w and split off the first block word. Lemma 2.3 turns concatenation of blocks into multiplication of the corresponding evaluations. $\square$

Lemma 2.27 (Blocking preserves MPV coefficients). *For each basis state $\sigma = (\sigma_1, \dots, \sigma_N)$ of the blocked system, where each σ_k is an L -tuple in $\{0, \dots, d-1\}^L$, there exists a basis state $\tilde{\sigma} = (\tilde{\sigma}_1, \dots, \tilde{\sigma}_{NL})$ of the original system such that*

$$V^{(N)}(A^{[L]})_\sigma = V^{(NL)}(A)_{\tilde{\sigma}}.$$

Proof. Concatenate the block words and apply the blocked word evaluation lemma (Lemma 2.26). $\square$

Theorem 2.28 (Blocking preserves same MPV). *If A and B generate the same MPV family, then for any blocking length L , the blocked tensors $A^{[L]}$ and $B^{[L]}$ also generate the same MPV family.*

Proof. By Lemma 2.27, each MPV coefficient of the blocked tensor $A^{[L]}$ equals a coefficient of A at the corresponding concatenated configuration. The same holds for $B^{[L]}$ (with the same concatenated configuration). Since A and B agree on all configurations, so do $A^{[L]}$ and $B^{[L]}$. $\square$

2.6 MPV overlap

Definition 2.29 (MPV as Hilbert-space vector). We regard $|V^{(N)}(A)\rangle$ as an element of $\mathbb{C}^{d^N}$ indexed by configurations $\sigma \in \{0, \dots, d-1\}^N$, with σ -component $V^{(N)}(A)_\sigma$ as in Definition 2.4. This allows us to use the standard Euclidean inner product on $\mathbb{C}^{d^N}$.

Definition 2.30 (MPV inner product). The *MPV inner product* between two tensors A and B at system size N is

$$\langle V^{(N)}(A) | V^{(N)}(B) \rangle := \sum_{\sigma \in \{0, \dots, d-1\}^N} \overline{V^{(N)}(A)_\sigma} V^{(N)}(B)_\sigma,$$

i.e. the Euclidean inner product on $\mathbb{C}^{d^N}$ (conjugate-linear in the first argument).

Definition 2.31 (MPV overlap). The *MPV overlap* between two tensors A (of bond dimension D_1) and B (of bond dimension D_2) at system size N is

$$O_{AB}(N) := \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma}.$$

This is the bilinear physics-oriented overlap used later. By Lemma 2.32, it is the complex conjugate of the Hilbert-space inner product from Definition 2.30.

Lemma 2.32 (Overlap equals conjugate inner product). $O_{AB}(N) = \overline{\langle V^{(N)}(A) | V^{(N)}(B) \rangle}$.

Proof. The overlap sums $V_\sigma \overline{W_\sigma}$ while the inner product sums $\overline{V_\sigma} W_\sigma$. Complex conjugation swaps these. $\square$

Chapter 3

The Single-Block Fundamental Theorem

This chapter proves that if an injective MPS tensor A generates the same MPV family as another tensor B of the same bond dimension, then A and B are gauge equivalent. The proof is purely algebraic and does not use the later overlap or normal-form reductions. The trace-pairing maps introduced below are auxiliary devices attached to the chosen generating families $\{A^i\}$ and $\{B^i\}$; they serve only to construct the linear map that is later shown to be a gauge transform.

3.1 Trace pairing

Theorem 3.1 (Nondegeneracy of the trace pairing). *For $M \in M_D(\mathbb{C})$, $\text{tr}(MN) = 0$ for all $N \in M_D(\mathbb{C})$ if and only if $M = 0$.*

Proof. Testing against the matrix units E_{ji} (the matrix with 1 in position (j, i) and 0 elsewhere) gives $\text{tr}(E_{ji}M) = M_{ij}$. Hence if $\text{tr}(MN) = 0$ for all N , then every entry of M vanishes. $\square$

Lemma 3.2 (Same MPV implies trace agreement). *If A and B generate the same MPV family, then $\text{tr}(A^w) = \text{tr}(B^w)$ for every word w .*

Proof. Viewing the word w as a configuration of length $|w|$, Definition 2.4 gives the corresponding MPV coefficient as the trace coefficient $c_w(A) = \text{tr}(A^w)$. The same-MPV hypothesis gives equality at every system size. $\square$

Definition 3.3 (Trace pairing map). For a tensor A , attach to the chosen family $\{A^i\}$ the auxiliary linear map $\Phi_A : M_D(\mathbb{C}) \rightarrow \mathbb{C}^d$ defined by

$$\Phi_A(M)_i = \text{tr}(M \cdot A^i).$$

Theorem 3.4 (Injectivity of the trace pairing map). *If A is injective, then $\ker \Phi_A = \{0\}$.*

Proof. If $\Phi_A(M) = 0$, then $\text{tr}(M \cdot A^i) = 0$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, this gives $\text{tr}(MN) = 0$ for all N , so $M = 0$ by Theorem 3.1. $\square$

Theorem 3.5 (Nonvanishing trace on injective tensors). *If $D \geq 1$, A is injective, A and B generate the same MPV family, and $B^i = 0$ for all i , then we have a contradiction.*

Proof. If $B^i = 0$ for all i , then Lemma 3.2 gives $\text{tr}(A^w) = 0$ for every word w , in particular for all length-1 words. Since the $\{A^i\}$ span $M_D(\mathbb{C})$, the trace functional vanishes identically, contradicting $\text{tr}(\mathbb{1}_D) = D \neq 0$. $\square$

3.2 Linear extension

Theorem 3.6 (Existence and uniqueness of the linear extension). *If A is injective and A, B generate the same MPV family, then there exists a unique linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ satisfying $T(A^i) = B^i$ for all i .*

Proof. Write $\ell_A : \mathbb{C}^d \rightarrow M_D(\mathbb{C})$ for the map that sends a coefficient vector c to $\sum_i c_i A^i$ (linear combination with the A^i). Applying Lemma 3.2 to all words of length 2 shows $\Phi_A \circ \ell_A = \Phi_B \circ \ell_B$. Since A is injective, ℓ_A is surjective, giving $\text{range}(\Phi_A) \subseteq \text{range}(\Phi_B)$. By Theorem 3.4, $\ker \Phi_A = \{0\}$, so rank-nullity gives $\dim(\text{range}(\Phi_A)) = \dim(M_D(\mathbb{C}))$. Hence $\dim(\text{range}(\Phi_B)) \geq \dim(M_D(\mathbb{C}))$, and rank-nullity again gives $\ker \Phi_B = \{0\}$. A left inverse g of Φ_B gives $T := g \circ \Phi_A$, and uniqueness follows because the $\{A^i\}$ span $M_D(\mathbb{C})$. $\square$

Theorem 3.7 (Multiplicativity of the linear extension). *If A is injective and A, B generate the same MPV family, and $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is the unique linear map satisfying $T(A^i) = B^i$ for all i , then $T(MN) = T(M)T(N)$ for all $M, N \in M_D(\mathbb{C})$.*

Proof. **Step 1** ($\Phi_B \circ T = \Phi_A$). Applying Lemma 3.2 to words of length 2 gives $\Phi_B(T(A^i)) = \Phi_B(B^i) = \Phi_A(A^i)$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, $\Phi_B \circ T = \Phi_A$.

Step 2 ($T(A^i A^j) = B^i B^j$). Lemma 3.2 applied to the word (i, j, k) gives $\text{tr}(A^i A^j A^k) = \text{tr}(B^i B^j B^k)$ for all i, j, k . So $\Phi_B(T(A^i A^j))_k = \Phi_A(A^i A^j)_k = \text{tr}(A^i A^j A^k) = \text{tr}(B^i B^j B^k) = \Phi_B(B^i B^j)_k$. The same rank-nullity argument used in the proof of Theorem 3.6 gives $\ker \Phi_B = \{0\}$. Hence $T(A^i A^j) = B^i B^j$.

Step 3 (bilinear extension). For fixed j , the maps $M \mapsto T(M \cdot A^j)$ and $M \mapsto T(M) \cdot B^j$ agree on the spanning set $\{A^i\}$, hence on all of $M_D(\mathbb{C})$. Applying this once more with j replaced by a general matrix gives full multiplicativity. $\square$

3.3 Simplicity and Skolem–Noether

Theorem 3.8 (Simplicity of $M_D(\mathbb{C})$). *A nonzero multiplicative $\mathbb{C}$ -linear endomorphism of $M_D(\mathbb{C})$ is bijective.*

Proof. The kernel of a multiplicative linear map is a two-sided ideal. Since $M_D(\mathbb{C})$ is a simple ring (it has no nontrivial two-sided ideals), the kernel is either $\{0\}$ or $M_D(\mathbb{C})$. The latter would force the map to be zero, contradicting the hypothesis. Hence the map is injective, and surjectivity follows from finite-dimensionality. $\square$

Theorem 3.9 (Skolem–Noether). *Every $\mathbb{C}$ -algebra automorphism of $M_D(\mathbb{C})$ is inner: for any $f \in \text{Aut}_{\mathbb{C}\text{-alg}}(M_D(\mathbb{C}))$, there exists $X \in \text{GL}_D(\mathbb{C})$ such that $f(M) = XMX^{-1}$ for all M .*

Proof. Via the canonical identification $M_D(\mathbb{C}) \cong \text{End}_{\mathbb{C}}(\mathbb{C}^D)$, the automorphism f induces an automorphism of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$. Every algebra automorphism of $\text{End}(V)$ is conjugation by an invertible linear map (this is a standard result in the theory of central simple algebras). Translating back to matrices gives the invertible X . $\square$

Lemma 3.10 (Promotion to algebra homomorphism). *If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a multiplicative surjective $\mathbb{C}$ -linear map, then $T(\mathbb{1}) = \mathbb{1}$. Equivalently, the same underlying function defines a $\mathbb{C}$ -algebra homomorphism.*

Proof. By surjectivity, there exists X with $T(X) = \mathbb{1}$. Then $T(\mathbb{1}) = T(X) \cdot T(\mathbb{1}) = T(X \cdot \mathbb{1}) = T(X) = \mathbb{1}$, using multiplicativity and $T(X) = \mathbb{1}$. With $T(\mathbb{1}) = \mathbb{1}$ established, T preserves the unit and hence is a $\mathbb{C}$ -algebra homomorphism. $\square$

3.4 Assembly

Theorem 3.11 (Single-block Fundamental Theorem). *Let A be an injective MPV tensor and let B be an MPV tensor of the same bond dimension. If A and B generate the same MPV family, then they are gauge equivalent: there exists $X \in \text{GL}_D(\mathbb{C})$ such that $B^i = XA^iX^{-1}$ for all i .*

Proof. By Theorem 3.6, there is a unique linear T with $T(A^i) = B^i$. By Theorem 3.7, T is multiplicative. T is nonzero: if $T = 0$ then $B^i = 0$ for all i , contradicting Theorem 3.5. By Theorem 3.8, T is bijective. By Lemma 3.10, T is a $\mathbb{C}$ -algebra automorphism. Theorem 3.9 gives an invertible X with $T(M) = XMX^{-1}$, and evaluating at $M = A^i$ yields $B^i = XA^iX^{-1}$. $\square$

Chapter 4

Quantum Channels and Positive Maps

This chapter develops the theory of positive maps and quantum channels on matrix algebras, following [Wol12].

4.1 Positive maps and channels

Definition 4.1 (Positive map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *positive* if $E(X) \geq 0$ whenever $X \geq 0$, where $X \geq 0$ means that X is positive semidefinite.

Remark 4.2. We use the Loewner order on matrices: $X \leq Y$ means $Y - X \geq 0$.

Definition 4.3 (Trace-preserving map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *trace-preserving* if $\text{tr}(E(X)) = \text{tr}(X)$ for all X .

Definition 4.4 (Completely positive map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *completely positive* (CP) if it admits a *Kraus representation*: there exist operators $\{K_i\}_{i=0}^{r-1}$ with $K_i \in M_D(\mathbb{C})$ such that

$$E(X) = \sum_{i=0}^{r-1} K_i X K_i^\dagger \quad \text{for all } X \in M_D(\mathbb{C}).$$

We use the Kraus characterisation because it matches the later Kadison–Schwarz arguments. By Choi’s theorem, this is equivalent to $E \otimes \text{id}_n$ being positive for all n .

Theorem 4.5 (CP maps are positive). *Every completely positive map is positive.*

Proof. If $X \geq 0$, then each summand $K_i X K_i^\dagger \geq 0$ (conjugation preserves positive semidefiniteness), so the sum is PSD. $\square$

Definition 4.6 (Quantum channel). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a *quantum channel* if it is both completely positive and trace-preserving (CPTP), following [Wol12].

Definition 4.7 (Density matrices). The set of *density matrices* in $M_D(\mathbb{C})$ is

$$\mathcal{D}_D = \{\rho \in M_D(\mathbb{C}) : \rho \geq 0, \text{tr}(\rho) = 1\}.$$

Theorem 4.8 (Density matrices are compact). $\mathcal{D}_D$ is compact (in the entry-wise topology on $M_D(\mathbb{C})$).

Proof. The PSD cone is closed (preimage of the closed nonnegative cone under continuous quadratic forms), and if $\rho \geq 0$ with $\text{tr}(\rho) = 1$ then each entry satisfies $|\rho_{ij}|^2 \leq \rho_{ii}\rho_{jj} \leq 1$: the diagonal entries are nonnegative and sum to 1, and the off-diagonal bound is the PSD Cauchy–Schwarz inequality. Hence the matrix entries are uniformly bounded, and Heine–Borel gives compactness. $\square$

Theorem 4.9 (Density matrices are convex). $\mathcal{D}_D$ is convex.

Proof. Convex combination of PSD matrices is PSD, and trace is linear. $\square$

Theorem 4.10 (Density matrices are nonempty). For $D \geq 1$, $\mathcal{D}_D \neq \emptyset$.

Proof. $\frac{1}{D}\mathbb{1}_D$ is a density matrix. $\square$

Theorem 4.11 (Channels preserve density matrices). If E is a quantum channel, then $E(\mathcal{D}_D) \subseteq \mathcal{D}_D$.

Proof. Complete positivity implies positivity (Theorem 4.5), so $E(\rho) \geq 0$; trace preservation gives $\text{tr}(E(\rho)) = \text{tr}(\rho) = 1$. $\square$

4.2 Irreducibility

Definition 4.12 (Irreducible map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *irreducible* if whenever P is an orthogonal projection satisfying $E(PM_D(\mathbb{C})P) \subseteq PM_D(\mathbb{C})P$, then $P = 0$ or $P = \mathbb{1}$. This is [Wol12, Thm. 6.2, condition (1)]. The definition applies to any linear map; complete positivity is not required.

Theorem 4.13 (Injectivity implies irreducibility). If A is an injective MPS tensor, then its transfer map $\mathcal{E}_A$ is irreducible.

Proof. If P is an invariant projection for $\mathcal{E}_A$, then $(\mathbb{1} - P)A^i P = 0$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, $(\mathbb{1} - P)MP = 0$ for all M , so $P = 0$ or $P = \mathbb{1}$. $\square$

Remark 4.14 (Transfer map is CP). The transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is completely positive: it is already written in Kraus form, with Kraus operators $\{A^i\}$.

Theorem 4.15 (Transfer map is a channel). If $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, then the transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is a quantum channel (CPTP).

Proof. Complete positivity is the observation of Remark 4.14. Trace preservation: $\text{tr}(\mathcal{E}_A(X)) = \sum_i \text{tr}(A^i X (A^i)^\dagger) = \text{tr}((\sum_i (A^i)^\dagger A^i)X) = \text{tr}(X)$ by cyclicity and the normalization hypothesis. $\square$

4.3 Cesàro fixed-point theorem

Definition 4.16 (Cesàro mean). The *Cesàro mean* of a linear map E at order N , applied to a starting matrix ρ_0 , is

$$\sigma_N = \frac{1}{N} \sum_{n=0}^{N-1} E^n(\rho_0).$$

Theorem 4.17 (Cesàro telescope). $E(\sigma_N) - \sigma_N = \frac{1}{N}(E^N(\rho_0) - \rho_0)$.

Proof. Telescoping: $E(\sigma_N) = \frac{1}{N} \sum_{n=1}^N E^n(\rho_0)$ and subtracting σ_N cancels all but the first and last terms. $\square$

Theorem 4.18 (Hermitian fixed-point decomposition). *Let E be a quantum channel and $X = E(X)$ a Hermitian fixed point with decomposition $X = P_+ - P_-$ into orthogonal positive and negative parts ($P_{\pm} \geq 0$, $\text{tr}(P_+ P_-) = 0$). Then $E(P_+) = P_+$ and $E(P_-) = P_-$. This is [Wol12, Prop. 6.8].*

Proof. Let Q be the support projection of P_+ . Trace preservation and positivity give $\text{tr}(QE(P_-)) = 0$ and $E(P_-) \geq 0$, so $QE(P_-) = 0$. The fixed-point equation $E(P_+) - E(P_-) = P_+ - P_-$ then identifies $E(P_{\pm}) = P_{\pm}$. $\square$

Theorem 4.19 (Cesàro fixed-point theorem). *Every quantum channel $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ (with $D \geq 1$) has a nonzero positive semidefinite fixed point: there exists $\rho \geq 0$, $\rho \neq 0$, with $E(\rho) = \rho$.*

Proof. Start with any $\rho_0 \in \mathcal{D}_D$ (Theorem 4.10). Each Cesàro mean σ_N lies in $\mathcal{D}_D$ (by convexity and channel preservation). By compactness (Theorem 4.8), extract a convergent subsequence $\sigma_{N_k} \rightarrow \rho$. The telescope identity (Theorem 4.17) gives $\|E(\sigma_N) - \sigma_N\| = \frac{1}{N}\|E^N(\rho_0) - \rho_0\| \rightarrow 0$, so $E(\rho) = \rho$ by continuity. In [Wol12, Thm. 6.11], existence is proved via Brouwer's fixed-point theorem; we use the Cesàro argument instead. $\square$

4.4 Fixed-point projection

Definition 4.20 (Fixed-point projection). Given a linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ and a fixed point ρ with $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$, the *fixed-point projection* is the rank-one map

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho.$$

This projects onto the span of ρ along the kernel of the trace functional. When the fixed point is unique up to scaling, this span is the full fixed-point space.

Theorem 4.21 (Power decomposition). *If E is trace-preserving and $E(\rho) = \rho$, then for $n \geq 1$,*

$$E^n = P + (E - P)^n,$$

where P is the fixed-point projection.

Proof. P is idempotent ($P^2 = P$), the fixed-point relation $E(\rho) = \rho$ gives $E \circ P = P$, and trace preservation gives $P \circ E = P$. These imply $(E - P) \circ P = 0$ and $P \circ (E - P) = 0$, so in the binomial expansion of $(P + (E - P))^n$ all cross terms vanish, leaving $E^n = P^n + (E - P)^n = P + (E - P)^n$. $\square$

4.5 Peripheral spectrum and primitivity

Definition 4.22 (Peripheral spectrum). The *peripheral spectrum* of a bounded linear operator T is the set of spectral values μ with $|\mu|$ equal to the spectral radius of T .

Definition 4.23 (Peripheral eigenvalues). The *peripheral eigenvalues* of a linear map E on a finite-dimensional space are the eigenvalues λ on the unit circle: $|\lambda| = 1$. For a channel, the spectral radius is 1 [Wol12, Prop. 6.1], so these coincide with the eigenvalues of maximal modulus.

Note: The condition $|\lambda| = 1$ is appropriate for channels, since the spectral radius of a channel is 1. For a general linear map with spectral radius $r \neq 1$ the condition would generalise to $|\lambda| = r$.

Theorem 4.24 (1 is a peripheral eigenvalue). *If E has a nonzero fixed point $\rho \neq 0$ with $E(\rho) = \rho$, then 1 is a peripheral eigenvalue of E .*

Proof. ρ is an eigenvector with eigenvalue 1, and $|1| = 1$. □

Remark 4.25 (Peripheral eigenvalues lie on the unit circle). Every peripheral eigenvalue λ satisfies $|\lambda| = 1$. This is simply the defining condition in Definition 4.23.

Theorem 4.26 (Finiteness of peripheral eigenvalues). *On a finite-dimensional space, the set of peripheral eigenvalues is finite.*

Proof. Peripheral eigenvalues are a subset of all eigenvalues, which is a finite set in finite dimensions. □

Theorem 4.27 (Power-stable peripheral eigenvalues are roots of unity). *Let E be a linear endomorphism on a finite-dimensional space. If μ is a peripheral eigenvalue of E and μ^n is an eigenvalue of E for every $n \geq 1$, then μ is a root of unity.*

Proof. Since μ^n is an eigenvalue for every n , the pigeonhole principle on the finite eigenvalue set gives $\mu^a = \mu^b$ for some $a \neq b$, hence $\mu^{|a-b|} = 1$. □

Theorem 4.28 (Peripheral powers imply root of unity). *If the set of peripheral eigenvalues of a linear endomorphism E on a finite-dimensional space is closed under powers ($\mu \in \text{peripheral}(E)$ and $n \geq 1$ imply $\mu^n \in \text{peripheral}(E)$), then every peripheral eigenvalue is a root of unity.*

Proof. The closure hypothesis provides $\mu^n \in \text{peripheral}(E)$ for all n , which in particular means μ^n is an eigenvalue. Theorem 4.27 then gives $\mu^p = 1$ for some $p > 0$. □

4.5.1 Periodicity removal by powering

Remark 4.29. The results in this subsection are purely spectral: they use only finiteness of a set of complex numbers and the spectral mapping theorem for powers. In particular, they do not rely on complete positivity. For transfer maps, replacing a tensor by its p -blocked tensor replaces $\mathcal{E}_A$ by $\mathcal{E}_A^p$, so this powering step is the operator-theoretic form of blocking.

Lemma 4.30 (Common exponent for a finite set of roots of unity). *Let $s \subseteq \mathbb{C}$ be a finite set. Assume that for every $\mu \in s$ there exists an exponent $p_\mu > 0$ with $\mu^{p_\mu} = 1$. Then there exists $p > 0$ such that $\mu^p = 1$ for all $\mu \in s$.*

Proof. Take p to be the least common multiple of the exponents p_μ . □

Lemma 4.31 (Powering collapses peripheral eigenvalues). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a linear map, and assume that E has a nonzero fixed point $\rho \neq 0$. Let $p > 0$. If every peripheral eigenvalue μ of E satisfies $\mu^p = 1$, then $\text{peripheral}(E^p) = \{1\}$.*

Proof. The fixed point ρ gives $1 \in \text{peripheral}(E^p)$. For the reverse inclusion, let $\nu \in \text{peripheral}(E^p)$. Spectral mapping gives $\nu = \mu^p$ for some $\mu \in \sigma(E)$. From $|\nu| = 1$ and $p > 0$ we obtain $|\mu| = 1$, hence $\mu \in \text{peripheral}(E)$. The hypothesis $\mu^p = 1$ then forces $\nu = 1$. □

For a normalized MPS tensor, the transfer map is naturally trace-preserving but need not be unital. In general one should not expect a single gauge choice to make it both unital and trace-preserving.

Remark 4.32 (Bi-canonical special case). When a Kraus map happens to be both unital and trace-preserving (the “bi-canonical” case), the Kadison–Schwarz equality at peripheral eigenvectors (Theorem 5.10) shows directly that the peripheral eigenvalues are closed under powers, and hence are roots of unity by Theorem 4.28. This is the simplest route, but it requires both normalizations simultaneously. The more general adjoint-fixed-point formulation below covers the case where only unitality and a positive definite adjoint fixed point are available.

4.5.2 Peripheral closure via adjoint fixed point

The following results replace the trace-preservation hypothesis by the existence of a positive definite fixed point for the *adjoint* Kraus map, matching the weighted-trace argument in [CPGSV17, Appendix A].

Lemma 4.33 (Peripheral closure via adjoint fixed point). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital ($\sum_i K_i K_i^\dagger = \mathbb{1}$), and assume:*

1. *the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point $\rho > 0$, and*
2. *E is irreducible.*

Then $\text{peripheral}(E)$ is closed under powers: if $\mu \in \text{peripheral}(E)$ then $\mu^n \in \text{peripheral}(E)$ for every $n \in \mathbb{N}$.

Proof. Let $E(X) = \mu X$ with $|\mu| = 1$, and set $G := E(X^\dagger X) - E(X)^\dagger E(X)$. By Kadison–Schwarz, $G \geq 0$. Since $E^*(\rho) = \rho$,

$$\text{tr}(\rho G) = \text{tr}(\rho E(X^\dagger X)) - \text{tr}(\rho E(X)^\dagger E(X)) = \text{tr}(E^*(\rho) X^\dagger X) - \text{tr}(\rho E(X)^\dagger E(X)) = \text{tr}(\rho X^\dagger X) - |\mu|^2 \text{tr}(\rho X^\dagger X) = 0.$$

Thus the trace of the Kadison–Schwarz gap against the adjoint fixed point vanishes. Because $\rho > 0$ and $G \geq 0$, this forces $G = 0$, so $E(X^\dagger X) = E(X)^\dagger E(X)$. Hence $X^\dagger X$ is a PSD fixed point, which is positive definite by irreducibility (Theorem 6.3). So X is invertible, and multiplicative-domain powers give $E(X^n) = \mu^n X^n$, so μ^n is peripheral. $\square$

Theorem 4.34 (Roots of unity via adjoint fixed point). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital, and assume that the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point and that E is irreducible. Then every peripheral eigenvalue of E is a root of unity.*

Proof. Combine the closure-under-powers result (Lemma 4.33) with Theorem 4.28. $\square$

Remark 4.35. Lemma 4.33 and Theorem 4.34 cover the bi-canonical (unital + TP) case as well: trace preservation gives $E^*(\mathbb{1}) = \mathbb{1}$, which is a positive definite fixed point of the adjoint. The adjoint-fixed-point formulation matches the argument in [CPGSV17, Appendix A], which uses a faithful invariant state rather than bi-canonicity.

Definition 4.36 (Channel period). The *period* of a channel E is the number of peripheral eigenvalues (counted without multiplicity): $p(E) := |\text{peripheral}(E)|$.

Definition 4.37 (Primitive map). A linear map is *primitive* if its only peripheral eigenvalue is $\lambda = 1$, i.e. $\text{peripheral}(E) = \{1\}$. For channels this corresponds to [Wol12, Thm. 6.7]. This definition applies to any linear endomorphism, not only to channels.

Theorem 4.38 (Primitivity equals period one). *A linear map with a nonzero fixed point is primitive if and only if its period is 1.*

Proof. If $\text{peripheral}(E) = \{1\}$, then $|\text{peripheral}(E)| = 1$. Conversely, if $|\text{peripheral}(E)| = 1$, then since $1 \in \text{peripheral}(E)$ (Theorem 4.24), the only element is 1. $\square$

Theorem 4.39 (Primitive maps have a spectral gap on the complement). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be trace-preserving, and let $\rho \neq 0$ satisfy $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$. Let*

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$$

be the fixed-point projection associated to ρ . Assume that:

1. *E is primitive;*
2. *every eigenvalue μ of E satisfies $|\mu| \leq 1$;*
3. *whenever $E(X) = X$ and $\text{tr}(X) = 0$, one has $X = 0$.*

Then every eigenvalue ν of $E - P$ satisfies $|\nu| < 1$.

Proof. Let $(E - P)(X) = \nu X$ with $X \neq 0$. Then

$$E(X) = \nu X + P(X).$$

If $\text{tr}(X) = 0$, then $P(X) = 0$, so $E(X) = \nu X$ and hence ν is an eigenvalue of E with $|\nu| \leq 1$. If $|\nu| = 1$, primitivity forces $\nu = 1$, so X is a trace-zero fixed point of E ; by assumption this implies $X = 0$, a contradiction. Thus $|\nu| < 1$.

If instead $\text{tr}(X) \neq 0$, trace preservation and $\text{tr}(P(X)) = \text{tr}(X)$ give

$$\text{tr}(X) = \text{tr}(E(X)) = \nu \text{tr}(X) + \text{tr}(P(X)) = \nu \text{tr}(X) + \text{tr}(X),$$

so $\nu \text{tr}(X) = 0$ and therefore $\nu = 0$. In either case, $|\nu| < 1$. $\square$

4.6 Representations: infrastructure

The following infrastructure supports the Choi–Jamiołkowski isomorphism (Theorem 4.44) and the representation theorems of [Wol12, Ch. 2].

Definition 4.40 (Partial traces). Let $X \in M_{d,d'}(\mathbb{C})$ be a bipartite matrix indexed by $(\text{Fin } d \times \text{Fin } d')^2$. The *left partial trace* $\text{tr}_A(X)$ and *right partial trace* $\text{tr}_B(X)$ are the $d' \times d'$ and $d \times d$ matrices defined by

$$(\text{tr}_A(X))_{ij} = \sum_k X_{(k,i),(k,j)}, \quad (\text{tr}_B(X))_{ij} = \sum_k X_{(i,k),(j,k)}.$$

Definition 4.41 (Maximally entangled state). The *maximally entangled state* on $\mathbb{C}^d \otimes \mathbb{C}^d$ is the rank-one projector

$$|\Omega\rangle\langle\Omega|, \quad |\Omega\rangle = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |j, j\rangle.$$

Concretely, $(|\Omega\rangle\langle\Omega|)_{(i_1,i_2),(j_1,j_2)} = \frac{1}{d} \delta_{i_1 j_1} \delta_{i_2 j_2}$ and $\text{tr}(|\Omega\rangle\langle\Omega|) = 1$.

Definition 4.42 (Tensor product of a map with identity). Given a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the *tensor extension* $T \otimes \text{id}$ acts on bipartite matrices $X \in M_{D \times D}(\mathbb{C})$ by applying T to each “slice”:

$$(T \otimes \text{id})(X)_{(i_1, i_2), (j_1, j_2)} = (T(X^{(i_2, j_2)}))_{i_1 j_1},$$

where $X_{ab}^{(i_2, j_2)} = X_{(a, i_2), (b, j_2)}$ is the bipartite slice.

4.7 Choi–Jamiołkowski isomorphism (Wolf Prop. 2.1)

Definition 4.43 (Choi matrix). The *Choi matrix* of a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is

$$\tau = (T \otimes \text{id})(|\Omega\rangle\langle\Omega|) \in M_{D \times D}(\mathbb{C}).$$

Concretely, $\tau_{(i_1, i_2), (j_1, j_2)} = \frac{1}{D} (T(E_{i_2 j_2}))_{i_1 j_1}$ where $E_{i_2 j_2}$ is the matrix unit. This is the Choi–Jamiołkowski convention of [Wol12, Prop. 2.1].

Theorem 4.44 (Prop. 2.1: CP iff Choi matrix is PSD). *A linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is completely positive (in the Kraus sense) if and only if its Choi matrix $\tau \geq 0$.*

Proof. ($\Rightarrow$) If $T(X) = \sum_i K_i X K_i^\dagger$, then $\tau = \sum_i |v_i\rangle\langle v_i|$ where $(v_i)_{(a, b)} = c K_i(a, b)$ and $c = 1/\sqrt{D}$. This is a sum of rank-one PSD matrices.

($\Leftarrow$) If $\tau \geq 0$, write $\tau = \sum_m |v_m\rangle\langle v_m|$ by spectral decomposition, define $K_m(a, b) = v_m(a, b)/c$, and recover $T(X) = \sum_m K_m X K_m^\dagger$ by linearity on matrix units. $\square$

Theorem 4.45 (Prop. 2.1: TP implies partial trace condition). *If T is trace-preserving, then $\text{tr}_A(\tau) = \frac{1}{D} \mathbb{1}_D$.*

Proof. $(\text{tr}_A(\tau))_{ij} = \text{tr}(T(\Omega^{(ij)})) = \text{tr}(\Omega^{(ij)})$ by trace preservation, where $\Omega^{(ij)} = \frac{1}{D} E_{ij}$ is the (i, j) -slice of $|\Omega\rangle\langle\Omega|$. Taking the trace gives $\frac{1}{D} \delta_{ij}$. $\square$

Theorem 4.46 (Prop. 2.1: Hermiticity correspondence). *The Choi matrix τ is Hermitian if and only if $T(B^\dagger) = T(B)^\dagger$ for all $B \in M_D(\mathbb{C})$.*

Proof. The Choi matrix entries satisfy $\tau_{(a, i), (b, j)} = \frac{1}{D} (T(E_{ij}))_{ab}$. Hermiticity of τ says $\overline{\tau_{(b, j), (a, i)}} = \tau_{(a, i), (b, j)}$, which unravels to $T(E_{ji})_{ba} = \overline{T(E_{ij})_{ab}}$, i.e., $T(E_{ij}^\dagger) = T(E_{ij})^\dagger$. By linearity this extends to all B . $\square$

Theorem 4.47 (Prop. 2.1: Trace of Choi matrix for TP map). *If T is trace-preserving, then $\text{tr}(\tau) = 1$.*

Proof. $\text{tr}(\tau) = \text{tr}(\text{tr}_A(\tau)) = \text{tr}(\frac{1}{D} \mathbb{1}_D) = 1$. $\square$

4.8 Kraus representation theorem (Wolf Thm. 2.1)

Theorem 4.48 (Thm. 2.1: TP implies Kraus normalization). *If $T(X) = \sum_i K_i X K_i^\dagger$ is trace-preserving, then $\sum_i K_i^\dagger K_i = \mathbb{1}$.*

Proof. For any N , $\text{tr}((\sum_i K_i^\dagger K_i) N) = \sum_i \text{tr}(K_i^\dagger K_i N) = \sum_i \text{tr}(K_i N K_i^\dagger) = \text{tr}(T(N)) = \text{tr}(N)$. Non-degeneracy of the trace pairing forces $\sum_i K_i^\dagger K_i = \mathbb{1}$. $\square$

Theorem 4.49 (Thm. 2.1: Kraus normalization implies TP). *If $\sum_i K_i^\dagger K_i = \mathbb{1}$, then $T(X) = \sum_i K_i X K_i^\dagger$ is trace-preserving.*

Proof. $\text{tr}(T(X)) = \sum_i \text{tr}(K_i X K_i^\dagger) = \sum_i \text{tr}(K_i^\dagger K_i X) = \text{tr}((\sum_i K_i^\dagger K_i) X) = \text{tr}(X)$. $\square$

Theorem 4.50 (Thm. 2.1: Unitary freedom in Kraus operators). *Let U be a unitary $r \times r$ matrix ($U^\dagger U = \mathbb{1}$) and suppose $K_j = \sum_\ell U_{j\ell} \tilde{K}_\ell$. Then $\{K_j\}$ and $\{\tilde{K}_\ell\}$ define the same Kraus map:*

$$\sum_j K_j X K_j^\dagger = \sum_\ell \tilde{K}_\ell X \tilde{K}_\ell^\dagger \quad \forall X \in M_D(\mathbb{C}).$$

Proof. Substitute the combination formula, expand the triple sum over j, ℓ, ℓ' , swap summation order, and use the entry-wise form of $U^\dagger U = \mathbb{1}$ to collapse to $\ell = \ell'$ diagonal terms. $\square$

4.9 Stinespring dilation (Wolf Thm. 2.2)

Definition 4.51 (Stinespring isometry). Given Kraus operators $\{K_j\}_{j=0}^{r-1}$ with $K_j \in M_D(\mathbb{C})$, the *Stinespring isometry* $V : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^r$ is the $(D \cdot r) \times D$ matrix

$$V_{(i,j),k} = (K_j)_{ik}.$$

This implements $V = \sum_j K_j \otimes |j\rangle$.

Theorem 4.52 (Thm. 2.2: Stinespring dual representation). *For Kraus operators $\{K_j\}$ and $A \in M_D(\mathbb{C})$,*

$$V^\dagger (A \otimes \mathbb{1}_r) V = \sum_j K_j^\dagger A K_j = T^*(A).$$

This is the Stinespring representation of the dual (Heisenberg picture) map T^ .*

Proof. Compute $(V^\dagger (A \otimes \mathbb{1}_r) V)_{ab}$ by summing over the product index (i, j) ; the δ_{jl} from $\mathbb{1}_r$ collapses the sum over l , giving $\sum_j K_j^\dagger A K_j$. $\square$

Theorem 4.53 (Thm. 2.2: Isometry condition iff TP). *$V^\dagger V = \mathbb{1}_D$ if and only if $\sum_j K_j^\dagger K_j = \mathbb{1}_D$. In particular, V is an isometry precisely when the Kraus map is trace-preserving.*

Proof. $(V^\dagger V)_{ab} = \sum_{(i,j)} \overline{V_{(i,j),a}} V_{(i,j),b} = \sum_j \sum_i \overline{(K_j)_{ia}} (K_j)_{ib} = (\sum_j K_j^\dagger K_j)_{ab}$. $\square$

Theorem 4.54 (Thm. 2.2: Stinespring Schrödinger representation). *The Kraus map $T(\rho) = \sum_j K_j \rho K_j^\dagger$ equals the partial trace over the dilation space:*

$$T(\rho)_{ij} = \sum_{k=0}^{r-1} (V \rho V^\dagger)_{(i,k),(j,k)} = \text{tr}_r(V \rho V^\dagger).$$

This is the Schrödinger-picture Stinespring representation.

Proof. Expand $(V \rho V^\dagger)_{(i,k),(j,k)}$ and sum over k to recover $\sum_k K_k \rho K_k^\dagger$. $\square$

4.10 Quantum dynamical semigroups

The one-parameter semigroup theory of Wolf Chapter 7 is treated in Chapter 13. That later chapter records the exponential representation of norm-continuous semigroups, perturbation bounds, and the GKSL/Lindblad description of generators. We do not duplicate that material here.

Chapter 5

Schwarz Inequalities and Multiplicative Domains

This chapter collects the Schwarz-inequality material from Wolf's Chapter 5 [Wol12]. We begin with the Kadison–Schwarz inequality for Kraus maps, then describe the multiplicative domain and its algebraic structure. We also record the extension to normal operators for positive subunital maps, the resulting order and spectral contractivity statements, and Wolf's standard example of a Schwarz map that is not completely positive.

5.1 Kadison–Schwarz inequality

A Kraus map is the concrete realisation of a completely positive map (Definition 4.4). The Kadison–Schwarz inequality and the multiplicative-domain characterisation are proved first for Kraus maps. The transfer map is a Kraus map by construction (Remark 4.14), so these results apply directly to transfer maps.

Definition 5.1 (Kraus map). Given operators $\{K_i\}_{i=0}^{d-1}$ with $K_i \in M_D(\mathbb{C})$, the *Kraus map* is $E(X) = \sum_{i=0}^{d-1} K_i X K_i^\dagger$. A linear map is completely positive (Definition 4.4) if and only if it can be written in this form.

Definition 5.2 (Adjoint Kraus map). The *adjoint Kraus map* is $E^*(X) = \sum_{i=0}^{d-1} K_i^\dagger X K_i$. When the $\{K_i\}$ are the matrices of an MPS tensor A , this is the transfer map of the conjugate-transposed family $i \mapsto (A^i)^\dagger$.

Definition 5.3 (Unital Kraus map). A Kraus map is *unital* if $\sum_{i=0}^{d-1} K_i K_i^\dagger = \mathbb{1}$. Equivalently, $E(\mathbb{1}) = \mathbb{1}$. In the later gauge language this is the right-canonical condition.

Definition 5.4 (Trace-preserving Kraus map). A Kraus map is *trace-preserving* if $\sum_{i=0}^{d-1} K_i^\dagger K_i = \mathbb{1}$. Equivalently, the adjoint Kraus map is unital. This is the standard MPS normalization condition. In the later gauge language it is the left-canonical condition, so Kadison–Schwarz arguments are often applied to the adjoint map.

Theorem 5.5 (Kadison–Schwarz inequality). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a unital Kraus map (i.e. $\sum_i K_i K_i^\dagger = \mathbb{1}$). Then for all $X \in M_D(\mathbb{C})$,*

$$E(X^\dagger X) \geq E(X)^\dagger E(X).$$

This is [Wol12, Eq. (5.2)].

Proof. The 2×2 block matrix $\begin{bmatrix} X^\dagger X & X^\dagger \\ X & \mathbb{1} \end{bmatrix}$ is PSD ($= vv^\dagger$ with $v = [X^\dagger, \mathbb{1}]^T$). Applying the unital CP map blockwise preserves positive semidefiniteness (each block $K_i(\cdot)K_i^\dagger$ preserves PSD, and unitality ensures the $(2, 2)$ -block maps to $\mathbb{1}$). The Schur complement of the $(2, 2)$ -block $\mathbb{1}$ gives the result. $\square$

Theorem 5.6 (Kadison–Schwarz for the adjoint channel). *If $\sum_i K_i^\dagger K_i = \mathbb{1}$ (trace-preserving), then the adjoint Kraus map satisfies $E^*(X^\dagger X) \geq E^*(X)^\dagger E^*(X)$.*

Proof. The adjoint operators $\{K_i^\dagger\}$ satisfy $\sum_i K_i^\dagger (K_i^\dagger)^\dagger = \sum_i K_i^\dagger K_i = \mathbb{1}$, so they form a unital family. Apply Theorem 5.5 to $\{K_i^\dagger\}$. $\square$

Theorem 5.7 (Hilbert–Schmidt contraction). *If a Kraus map is both unital and trace-preserving, then $\text{tr}(E(X)^\dagger E(X)) \leq \text{tr}(X^\dagger X)$ for all $X \in M_D(\mathbb{C})$.*

Proof. By Theorem 5.5, $E(X^\dagger X) - E(X)^\dagger E(X) \geq 0$, so $\text{tr}(E(X^\dagger X)) \geq \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. $\square$

5.2 Multiplicative domain

Theorem 5.8 (KS gap decomposition). *For a unital Kraus map E , the Kadison–Schwarz gap decomposes as*

$$E(X^\dagger X) - E(X)^\dagger E(X) = \sum_{i=0}^{d-1} R_i^\dagger R_i,$$

where $R_i := XK_i^\dagger - K_i^\dagger E(X)$.

Proof. Expand the right-hand side $\sum_i (XK_i^\dagger - K_i^\dagger E(X))^\dagger (XK_i^\dagger - K_i^\dagger E(X))$, distribute, and use unitality $\sum_i K_i K_i^\dagger = \mathbb{1}$ to identify the result with the KS gap. $\square$

Theorem 5.9 (KS equality implies Kraus commutation). *Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$ for some X , then $XK_i^\dagger = K_i^\dagger E(X)$ for all i .*

Proof. By Theorem 5.8, the KS gap is $\sum_i R_i^\dagger R_i$ with $R_i = XK_i^\dagger - K_i^\dagger E(X)$. If the gap vanishes, $\sum_i R_i^\dagger R_i = 0$. Each summand $R_i^\dagger R_i$ is PSD, so each must be zero, giving $R_i = 0$. $\square$

Theorem 5.10 (KS equality for peripheral eigenvectors). *Let E be a Kraus map that is both unital and trace-preserving. If $E(X) = \mu X$ with $|\mu| = 1$, then the KS gap vanishes: $E(X^\dagger X) = E(X)^\dagger E(X)$.*

Proof. The gap $G := E(X^\dagger X) - E(X)^\dagger E(X)$ is PSD by Theorem 5.5. Its trace satisfies $\text{tr}(G) = \text{tr}(E(X^\dagger X)) - \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. Using $E(X) = \mu X$ and $|\mu| = 1$, $\text{tr}(E(X)^\dagger E(X)) = |\mu|^2 \text{tr}(X^\dagger X) = \text{tr}(X^\dagger X)$. So $\text{tr}(G) = 0$, and a PSD matrix with zero trace must be zero. $\square$

Theorem 5.11 (Left multiplicative identity from KS equality). *Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then $E(X^\dagger Y) = E(X)^\dagger E(Y)$ for all $Y \in M_D(\mathbb{C})$.*

Proof. From Theorem 5.9, $XK_i^\dagger = K_i^\dagger E(X)$ for all i . Taking conjugate transposes: $K_i X^\dagger = E(X)^\dagger K_i$. Then

$$\begin{aligned} E(X^\dagger Y) &= \sum_i K_i X^\dagger Y K_i^\dagger \\ &= \sum_i E(X)^\dagger K_i Y K_i^\dagger \\ &= E(X)^\dagger E(Y). \end{aligned}$$

□

5.2.1 Full multiplicative-domain structure

Definition 5.12 (Right and left multiplicative domains). For a Kraus map E , define

$$\mathcal{A}_R(E) := \{X \in M_D(\mathbb{C}) : E(XY) = E(X)E(Y) \text{ for all } Y \in M_D(\mathbb{C})\}$$

and

$$\mathcal{A}_L(E) := \{X \in M_D(\mathbb{C}) : E(YX) = E(Y)E(X) \text{ for all } Y \in M_D(\mathbb{C})\}.$$

We follow the convention that $\mathcal{A}_R(E)$ controls right multiplication and $\mathcal{A}_L(E)$ controls left multiplication. These are the two one-sided multiplicative domains from [Wol12, Eqs. (5.11)–(5.12)].

Definition 5.13 (Full multiplicative domain). The *multiplicative domain* of E is

$$\mathcal{A}(E) := \mathcal{A}_R(E) \cap \mathcal{A}_L(E).$$

Theorem 5.14 (Multiplicative-domain characterization). *Let E be a unital Kraus map. Then*

$$X \in \mathcal{A}_R(E) \iff E(XX^\dagger) = E(X)E(X)^\dagger$$

and

$$X \in \mathcal{A}_L(E) \iff E(X^\dagger X) = E(X)^\dagger E(X).$$

Together these are the two one-sided characterizations from [Wol12, Thm. 5.7].

Proof. If X lies in a one-sided multiplicative domain, evaluate the defining identity at $Y = X^\dagger$. Conversely, the left-domain implication is exactly Theorem 5.11, and the right-domain implication follows by applying Theorem 5.11 to $X^\dagger$. □

Definition 5.15 (One-sided multiplicative domains are subalgebras). For a unital Kraus map E , both $\mathcal{A}_R(E)$ and $\mathcal{A}_L(E)$ are unital subalgebras of $M_D(\mathbb{C})$.

Definition 5.16 (Multiplicative domain as a $*$ -subalgebra). For a unital Kraus map E , the full multiplicative domain $\mathcal{A}(E)$ is a $*$ -subalgebra of $M_D(\mathbb{C})$. This is the algebraic content of [Wol12, Thm. 5.7].

Definition 5.17 (Restriction to the multiplicative domain). If E is unital, then the restricted map $E|_{\mathcal{A}(E)} : \mathcal{A}(E) \rightarrow M_D(\mathbb{C})$ is a $*$ -algebra homomorphism. This is the homomorphism conclusion of [Wol12, Thm. 5.7].

5.2.2 Schwarz inequality for normal operators

Theorem 5.18 (CP Schwarz inequality for normal operators). *Let $E^*(X) = \sum_i K_i^\dagger X K_i$ be the adjoint Kraus map of a trace-preserving Kraus family, and let $A \in M_D(\mathbb{C})$ be normal. Then the Schwarz gap*

$$E^*(A^\dagger A) - E^*(A^\dagger)E^*(A)$$

is positive semidefinite. Equivalently,

$$E^*(A^\dagger)E^*(A) \leq E^*(A^\dagger A).$$

This is the completely positive case of [Wol12, Prop. 5.1].

Proof. The positivity statement is exactly Theorem 5.6. The second formulation is just the corresponding Loewner-order reformulation. $\square$

Theorem 5.19 (Prop. 5.1: Schwarz inequality for normal operators). *Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $T(\mathbb{1}) \leq \mathbb{1}$. If $A \in M_D(\mathbb{C})$ is normal, then*

$$T(A^\dagger A) - T(A^\dagger)T(A) \geq 0.$$

This is [Wol12, Prop. 5.1].

Proof. Restrict T to the commutative $*$ -subalgebra generated by $\mathbb{1}$, A , $A^\dagger$, and $A^\dagger A$. On this domain the map is completely positive, so the Kraus-form Kadison–Schwarz inequality from Theorem 5.5 applies and yields the claim. $\square$

5.2.3 Positive maps preserve order and spectral intervals

Theorem 5.20 (Positive maps are monotone). *If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive and $A \leq B$, then $T(A) \leq T(B)$.*

Proof. Since $B - A \geq 0$, positivity gives $T(B - A) \geq 0$. By linearity, this is $T(B) - T(A) \geq 0$. $\square$

Theorem 5.21 (Positive maps preserve adjoints). *If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive, then $T(A^\dagger) = T(A)^\dagger$ for all $A \in M_D(\mathbb{C})$.*

Proof. Write $A = B + iC$ with B and C Hermitian. Positive maps send Hermitian matrices to Hermitian matrices, so they preserve this decomposition and commute with taking adjoints. $\square$

Theorem 5.22 (Order intervals are preserved). *Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be positive with $T(\mathbb{1}) \leq \mathbb{1}$. If $a \leq 0 \leq b$ and $a\mathbb{1} \leq A \leq b\mathbb{1}$, then*

$$a\mathbb{1} \leq T(A) \leq b\mathbb{1}.$$

This is the matrix form of [Wol12, Eq. (5.21)].

Proof. Apply monotonicity to the inequalities $a\mathbb{1} \leq A \leq b\mathbb{1}$. The scalar bounds are preserved because $a \leq 0 \leq b$ and $T(\mathbb{1}) \leq \mathbb{1}$. $\square$

Theorem 5.23 (Spectral interval contractivity). *Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be positive with $T(\mathbb{1}) \leq \mathbb{1}$, and let $A \in M_D(\mathbb{C})$ be Hermitian. If $\text{spec}(A) \subseteq [a, b]$ with $a \leq 0 \leq b$, then*

$$\text{spec}(T(A)) \subseteq [a, b].$$

This is [Wol12, Eq. (5.21)].

Proof. For Hermitian matrices, the inclusion $\text{spec}(A) \subseteq [a, b]$ is equivalent to the order bounds $a\mathbb{1} \leq A \leq b\mathbb{1}$. Apply Theorem 5.22 and translate the resulting inequalities back into spectral bounds. $\square$

5.2.4 Wolf Example 5.3

Definition 5.24 (Wolf Example 5.3 map). Consider the linear map $T : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$ defined by

$$T(A) = \frac{1}{2}A^T + \frac{1}{4}\text{tr}(A)\mathbb{1}.$$

This is the map from [Wol12, Ex. 5.3].

Theorem 5.25 (Wolf Example 5.3 is positive). *The map from Definition 5.24 is positive.*

Proof. It is a convex combination of the transpose map and the map $A \mapsto \text{tr}(A)\mathbb{1}$, and both of these maps send positive semidefinite matrices to positive semidefinite matrices. $\square$

Theorem 5.26 (Wolf Example 5.3 satisfies the Schwarz inequality). *For every $A \in M_2(\mathbb{C})$,*

$$T(A^\dagger A) - T(A^\dagger)T(A) \geq 0,$$

where T is the map from Definition 5.24.

Proof. Write the Schwarz gap as $F(A)$. One checks that $F(A + \lambda\mathbb{1}) = F(A)$ for every scalar λ , so it is enough to treat the case $\text{tr}(A) = 0$. In that case a direct 2×2 computation shows that $F(A)$ dominates $\frac{1}{2}(A^\dagger A)^T$, hence is positive semidefinite. $\square$

Theorem 5.27 (Wolf Example 5.3 is not completely positive). *The map from Definition 5.24 is not completely positive.*

Proof. Compute the Choi matrix of T and test it on the antisymmetric vector $|01\rangle - |10\rangle$. The resulting expectation value is negative, so the Choi matrix is not positive semidefinite. $\square$

Chapter 6

Perron–Frobenius Theory for Channels and Transfer Maps

This chapter establishes the Perron–Frobenius theory used throughout the blueprint: existence, positive definiteness, and uniqueness of fixed points for injective and irreducible transfer maps, canonical gauge constructions, ergodicity of irreducible channels, the spectral-radius identification of Perron eigenvalues, and Wolf’s spectral characterization of irreducibility. The conceptual source is Wolf’s treatment of irreducible positive maps [Wol12, Ch. 6, especially Thms. 6.2–6.5 and Cor. 6.3] and [EHK78].

6.1 Positive definiteness

Remark 6.1. Wolf’s general theory treats irreducible positive maps first ([Wol12, Thm. 6.2]), then derives non-degeneracy and positive definiteness ([Wol12, Thm. 6.3]). The injective-tensor version (Theorem 6.2) uses a shorter direct kernel argument; the irreducible CP-map version (Theorem 6.3) follows via the support-projection route of [Wol12, Thm. 6.2].

Theorem 6.2 (PSD fixed point is positive definite). *Let A be an injective MPS tensor and let $\rho \geq 0$, $\rho \neq 0$, be a fixed point of the transfer map $\mathcal{E}_A(\rho) = \rho$. Then ρ is positive definite.*

Proof. Suppose $v^\dagger \rho v = 0$ for some $v \neq 0$. Since $\rho \geq 0$, this implies $\rho v = 0$. The fixed-point equation gives $v^\dagger \rho v = \sum_i ((A^i)^\dagger v)^\dagger \rho ((A^i)^\dagger v)$. Each term is nonneg; if the sum is zero, then $\rho (A^i)^\dagger v = 0$ for all i : the kernel of ρ is invariant under all $(A^i)^\dagger$. Since the $\{A^i\}$ span $M_D(\mathbb{C})$, we have $\rho M v = 0$ for every matrix M . Given any vector w , choose M with $M v = w$. Then $\rho w = 0$ for every w , so $\rho = 0$, contradicting $\rho \neq 0$. In [Wol12, Thm. 6.3, item 2], the same conclusion is reached via the growth condition $(\mathbb{1} + T)^{d-1}(X) > 0$ of [Wol12, Thm. 6.2, condition (2)]; here the direct kernel argument is shorter under injectivity. $\square$

Theorem 6.3 (PSD fixed point is PD — irreducible version). *If $\mathcal{E}_A$ is irreducible and $\rho \geq 0$, $\rho \neq 0$ is a fixed point, then ρ is positive definite.*

Proof. Suppose ρ is not positive definite. Then ρ has a nontrivial kernel; let Q be the orthogonal projection onto the support of ρ (the complement of the kernel). A direct computation using the fixed-point equation $\mathcal{E}_A(\rho) = \rho$ shows that Q is an invariant projection for $\mathcal{E}_A$: it satisfies $Q \mathcal{E}_A(Q X Q) Q = \mathcal{E}_A(Q X Q)$ for all X . By irreducibility of $\mathcal{E}_A$ (Definition 4.12), every invariant projection is 0 or $\mathbb{1}$. If $Q = 0$ then $\rho = 0$, contradicting $\rho \neq 0$. If $Q = \mathbb{1}$ then ρ is positive definite,

contradicting our assumption. Hence ρ must be positive definite. This is the support-projection direction of [Wol12, Thm. 6.2, condition (1)]. $\square$

Theorem 6.4 (Orthogonal trace condition). *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, and let $A, B \geq 0$ be nonzero positive semidefinite matrices with $\text{tr}(BA) = 0$. Then there exists t with $1 \leq t \leq D - 1$ such that*

$$\text{tr}(B \cdot E^t(A)) > 0.$$

Proof. The growth theorem for irreducible CP maps ([Wol12, Thm. 6.2, item 2]) gives $(\mathbb{1} + E)^{D-1}(A) > 0$ (positive definite). Since $B \geq 0$ is nonzero, $\text{tr}(B \cdot (\mathbb{1} + E)^{D-1}(A)) > 0$. Expanding by the binomial theorem:

$$\text{tr}\left(B \cdot \sum_{k=0}^{D-1} \binom{D-1}{k} E^k(A)\right) > 0.$$

Each term $\text{tr}(B \cdot E^k(A)) \geq 0$ since both factors are PSD. The $k = 0$ term vanishes by hypothesis ($\text{tr}(BA) = 0$). Hence at least one $k \in \{1, \dots, D - 1\}$ contributes a strictly positive term. This is [Wol12, Thm. 6.2, item 4]: item (2) (growth condition) implies item (4) by a binomial expansion and PSD positivity argument. $\square$

6.2 Uniqueness

Theorem 6.5 (Uniqueness of PSD fixed point). *Let A be an injective MPS tensor. If $\rho, \sigma \geq 0$ are both nonzero fixed points of $\mathcal{E}_A$, then $\sigma = c\rho$ for some $c \in \mathbb{C}$.*

Proof. Both ρ and σ are positive definite by Theorem 6.2. Write $\rho = SS^\dagger$ via a spectral square root and set $H = (S^\dagger)^{-1}\sigma S^{-1}$, which is Hermitian and positive definite. Let c_0 be the minimal eigenvalue of H . Then $\tau := \sigma - c_0\rho$ is a PSD fixed point which is *not* positive definite (it has a zero eigenvalue by construction of c_0). By Theorem 6.2, τ must be zero, giving $\sigma = c_0\rho$.

Relation to Wolf: In [Wol12, Thm. 6.3, item 2], uniqueness is proved via the growth condition of [Wol12, Thm. 6.2]. The same minimal-eigenvalue argument applies here, using the injective-tensor positive-definiteness (Theorem 6.2) in place of the growth condition. $\square$

Remark 6.6. The proof produces a positive real scalar $c_0 > 0$: it is the minimal eigenvalue of the positive definite Hermitian matrix $H = (S^\dagger)^{-1}\sigma S^{-1}$.

Theorem 6.7 (Uniqueness of positive eigenvalue for irreducible CP maps). *Let $D \geq 1$ and let E be an irreducible completely positive map on $M_D(\mathbb{C})$. Suppose $\rho, \sigma \geq 0$ are nonzero positive semidefinite matrices satisfying $E(\rho) = r_1\rho$ and $E(\sigma) = r_2\sigma$ for real scalars $r_1, r_2 > 0$. Then $r_1 = r_2$.*

Proof. Extract Kraus operators K for E . Irreducibility of E passes to the adjoint transfer map $E^\dagger(\cdot) = \sum_i K_i^\dagger(\cdot) K_i$, which is a positive CP map. The Perron–Frobenius existence theorem (Theorem 6.15) together with the irreducible PSD-to-PosDef upgrade (Theorem 6.3) gives a positive definite eigenvector $\tau > 0$ with eigenvalue $t > 0$: $E^\dagger(\tau) = t\tau$. For any nonzero PSD X with $E(X) = sX$, the trace pairing identity yields

$$s \cdot \text{tr}(\tau X) = \text{tr}(\tau \cdot E(X)) = \text{tr}(E^\dagger(\tau) \cdot X) = t \cdot \text{tr}(\tau X).$$

Since $\tau > 0$ and $X \geq 0$, $X \neq 0$, the scalar $\text{tr}(\tau X) > 0$, so $s = t$. Applying this to both (ρ, r_1) and (σ, r_2) gives $r_1 = t = r_2$. **Relation to Wolf:** This is [Wol12, Thm. 6.3, item 3], establishing non-degeneracy of the spectral-radius eigenvalue for irreducible CP maps. The proof follows Wolf’s dual-map trace argument. $\square$

6.3 Existence and assembly

Theorem 6.8 (Existence of PSD fixed point). *Assume $D \geq 1$. Let A be an MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ (so that $\mathcal{E}_A$ is trace-preserving). Then there exists $\rho \geq 0$, $\rho \neq 0$, with $\mathcal{E}_A(\rho) = \rho$.*

Proof. Under the normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, the transfer map is a quantum channel (Theorem 4.15). The Cesàro fixed-point theorem (Theorem 4.19) then provides the fixed point; injectivity is not needed for this step. $\square$

Theorem 6.9 (Quantum Perron–Frobenius). *Assume $D \geq 1$. Let A be an injective MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then the transfer map $\mathcal{E}_A$ has a unique PSD fixed point ρ (up to scaling), and ρ is positive definite.*

Proof. Combine Theorems 6.8, 6.2, and 6.5. $\square$

6.4 Right- and left-canonical gauges

Theorem 6.10 (Right-canonical (unital) gauge). *Let $\mathcal{E}_A(\rho) = \rho$ and let S be an invertible matrix with $SS^\dagger = \rho$. Define the gauged operators $A'^i = S^{-1}A^iS$. Then $\sum_i A'^i(A'^i)^\dagger = \mathbb{1}$, i.e. the gauged Kraus map is unital. This is the right-canonical normalization used later in the blueprint.*

Proof. Expand each summand: $(S^{-1}A^iS)(S^{-1}A^iS)^\dagger = S^{-1}A^i \underbrace{SS^\dagger}_\rho (A^i)^\dagger (S^\dagger)^{-1}$. Summing over i and using $\sum_i A^i(A^i)^\dagger = \rho$ gives $S^{-1}\rho(S^\dagger)^{-1} = S^{-1}SS^\dagger(S^\dagger)^{-1} = \mathbb{1}$. $\square$

Theorem 6.11 (Left-canonical (trace-preserving) gauge). *Let σ be a fixed point of the adjoint transfer map $\sum_i (A^i)^\dagger \sigma A^i = \sigma$, and let S be invertible with $S^\dagger S = \sigma$. Define $A'^i = SA^iS^{-1}$. Then $\sum_i (A'^i)^\dagger A'^i = \mathbb{1}$, i.e. the gauged Kraus map is trace-preserving. This is the left-canonical normalization used later in the blueprint.*

Proof. Expand each summand: $(SA^iS^{-1})^\dagger (SA^iS^{-1}) = (S^\dagger)^{-1}(A^i)^\dagger \underbrace{S^\dagger S}_\sigma A^i S^{-1}$. Summing over i and using $\sum_i (A^i)^\dagger \sigma A^i = \sigma$ gives $(S^\dagger)^{-1}\sigma S^{-1} = (S^\dagger)^{-1}S^\dagger S S^{-1} = \mathbb{1}$. $\square$

Remark 6.12. The right-canonical gauge of Theorem 6.10 and the left-canonical gauge of Theorem 6.11 are generally different similarity transforms: one is built from a fixed point of $\mathcal{E}_A$, the other from a fixed point of the adjoint transfer map. This section does not claim that a single gauge makes the transfer map simultaneously unital and trace-preserving.

6.5 Similarity preserves irreducibility

Lemma 6.13 (Similarity transform preserves irreducibility). *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $C \in M_D(\mathbb{C})$ be invertible ($\det C \neq 0$), and let $c > 0$. Define the similarity-transformed map*

$$E'(X) = c \cdot C^{-1} E(C X C^\dagger) (C^\dagger)^{-1}.$$

Then E' is also irreducible.

Proof. Suppose Q is an invariant projection for E' : $Q \neq 0$, $Q \neq \mathbb{1}$, and $(\mathbb{1} - Q)E'(QXQ)Q = 0$ for all X . Setting $R = CQC^\dagger$, the support projection P of R is an invariant projection for E . Irreducibility of E forces $P = 0$ or $P = \mathbb{1}$. The invertibility of C then forces $Q = 0$ or $Q = \mathbb{1}$, a contradiction. This is the full similarity version of [Wol12, Prop. 6.6]; the scalar case ($C = \mathbb{1}$) gives the scaling result stated next. $\square$

Theorem 6.14 (Scaling preserves irreducibility). *If E is an irreducible map and $c \neq 0$, then cE is also irreducible. This is the scalar special case ($C = \mathbb{1}$) of [Wol12, Prop. 6.6].*

Proof. An invariant projection for cE is an invariant projection for E (scale out the nonzero constant from the commutation relation). $\square$

6.6 Perron–Frobenius eigenvector existence

This section establishes the existence of a positive definite eigenvector for the adjoint transfer map of an irreducible MPS tensor, and uses it to construct a TP-gauge normalization. The PSD-eigenvector step corresponds to [Wol12, Thm. 6.5] (general positive maps) and [EHK78]; the upgrade to positive definiteness uses the irreducible-map theory of [Wol12, Thm. 6.3]. The TP-gauge application follows [CPGSV17, Appendix A].

Theorem 6.15 (PSD eigenvector for positive maps). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $D > 0$, and assume that $E(\sigma) \neq 0$ for every nonzero PSD matrix σ . Then there exist a nonzero PSD matrix ρ and a positive real $r > 0$ such that $E(\rho) = r\rho$.*

This is the finite-dimensional Perron–Frobenius theorem for positive maps on matrix algebras ([Wol12, Thm. 6.5]; see also [EHK78]). The proof applies Brouwer’s fixed-point theorem to the normalization map $\rho \mapsto E(\rho)/\text{tr}(E(\rho))$ on the compact convex set of density matrices.

Theorem 6.16 (Adjoint transfer map is nonzero). *If some Kraus operator $A^i \neq 0$, then the adjoint transfer map $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is nonzero.*

Proof. If $\mathcal{E}_A^\dagger = 0$ then $\mathcal{E}_A^\dagger(\mathbb{1}) = 0$, so $\sum_i (A^i)^\dagger A^i = 0$. Since each summand $(A^i)^\dagger A^i$ is PSD, each $(A^i)^\dagger = 0$ and hence each $A^i = 0$. $\square$

Theorem 6.17 (PosDef adjoint eigenvector for irreducible tensors). *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive definite matrix σ and a positive real $r > 0$ such that $\mathcal{E}_A^\dagger(\sigma) = r\sigma$.*

This combines the general PF existence ([Wol12, Thm. 6.5]) with the irreducible upgrade to positive definiteness ([Wol12, Thm. 6.3, item 2]); the application to the adjoint transfer map follows [CPGSV17, Appendix A].

Proof. By Theorem 6.16, $\mathcal{E}_A^\dagger \neq 0$. Since $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is a Kraus map, it is positive. Irreducibility of A transfers to the adjoint transfer map, and together with $\mathcal{E}_A^\dagger \neq 0$ this implies $\mathcal{E}_A^\dagger(\tau) \neq 0$ for every nonzero PSD matrix τ . Hence Theorem 6.15 applies to $\mathcal{E}_A^\dagger$, giving $\sigma_0 \geq 0$ (PSD, nonzero) and $r > 0$ with $\mathcal{E}_A^\dagger(\sigma_0) = r\sigma_0$. Rescale: set $T^i = r^{-1/2}(A^i)^\dagger$. Then $\mathcal{E}_T(\sigma_0) = \sigma_0$, and $\mathcal{E}_T$ is irreducible (by Theorem 6.14, since irreducibility of $\mathcal{E}_A$ transfers to the adjoint and is preserved under scaling). Since $\mathcal{E}_T$ is irreducible with a PSD fixed point, the PSD-to-PosDef upgrade (Theorem 6.3) gives σ_0 positive definite. $\square$

Theorem 6.18 (TP-gauge existence for irreducible tensors). *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive real r , a positive definite matrix σ , and a gauge-equivalent tensor B such that*

$$B^i = \sigma^{1/2} \cdot (r^{-1/2} A^i) \cdot \sigma^{-1/2}$$

is trace-preserving: $\sum_i (B^i)^\dagger B^i = \mathbb{1}$.

This is the “spectral rescaling + TP gauge” step of [CPGSV17, Appendix A]. The similarity is generally non-unitary.

Proof. By Theorem 6.17, obtain σ positive definite and $r > 0$ with $\mathcal{E}_A^\dagger(\sigma) = r\sigma$. Set $c = r^{-1/2}$ and $A'^i = cA^i$. Then $\mathcal{E}_{A'}^\dagger(\sigma) = \sigma$. By the TP gauge construction (Theorem 6.11), $B^i = \sigma^{1/2} A'^i \sigma^{-1/2}$ satisfies $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. $\square$

6.7 Exponential positivity for irreducible CP maps

Theorem 6.19 (Exponential truncation is positive definite). *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $A \geq 0$ be nonzero, and let $t > 0$. Then the finite exponential truncation*

$$\sum_{k=0}^{D-1} \frac{t^k}{k!} E^k(A)$$

is positive definite. This is the finite-sum core of [Wol12, Thm. 6.2, item (3)].

Proof. If a nonzero vector v annihilated this quadratic form, then every term $E^k(A)$ with $0 \leq k \leq D-1$ would annihilate v . The growth theorem for irreducible CP maps then forces $(\mathbb{1} + E)^{D-1}(A)$ to annihilate v , contradicting its positive definiteness. $\square$

Theorem 6.20 (Exponential positivity for irreducible CP maps). *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $A \geq 0$ be nonzero, and let $t > 0$. Then*

$$\exp(tE)(A) = \sum_{k=0}^{\infty} \frac{t^k}{k!} E^k(A)$$

is positive definite. This is [Wol12, Thm. 6.2, item (3)].

Proof. By Theorem 6.19, the first D terms already form a positive definite matrix. Every remaining term is positive semidefinite, so the full exponential series is the sum of a positive definite matrix and a positive semidefinite tail. $\square$

6.8 Ergodicity of irreducible channels

Theorem 6.21 (Unique density-matrix fixed point for an irreducible channel). *Let E be an irreducible quantum channel on $M_D(\mathbb{C})$ with $D > 0$. Then there exists a unique density matrix σ such that $E(\sigma) = \sigma$. Moreover σ is positive definite. This is the qualitative form of [Wol12, Cor. 6.3].*

Proof. Every Cesàro mean of a density matrix stays inside the compact convex set of density matrices, so a subsequence converges. Its limit is a fixed point by the telescoping argument, positive definiteness follows from irreducibility, and uniqueness comes from the uniqueness of positive semidefinite Perron eigenvectors at eigenvalue 1. $\square$

Theorem 6.22 (Cesàro convergence for irreducible channels). *Let E be an irreducible quantum channel on $M_D(\mathbb{C})$ with $D > 0$, and let ρ be any density matrix. Then the Cesàro means*

$$\frac{1}{N} \sum_{n=0}^{N-1} E^n(\rho)$$

converge to the unique positive definite density-matrix fixed point of E . This is [Wol12, Cor. 6.3].

Proof. Every subsequential limit of the Cesàro means is again a density-matrix fixed point. By Theorem 6.21, there is only one such fixed point, so every convergent subsequence has the same limit. Compactness then forces the whole sequence to converge to that limit. $\square$

6.9 Spectral radius at the Perron eigenvalue

Theorem 6.23 (Spectral radius is the Perron eigenvalue). *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$. Suppose $\rho > 0$ and $E(\rho) = r\rho$ with $r > 0$. Then the spectral radius of E is r . This is [Wol12, Thm. 6.3, item (4)].*

Proof. Choose a positive definite eigenvector for the adjoint map with the same eigenvalue, gauge by its square root, and rescale by r^{-1} . The resulting map is trace-preserving and has a positive definite fixed point, so its spectral radius is 1. Undoing the similarity and scalar rescaling recovers the spectral radius of E . $\square$

Theorem 6.24 (Real spectral-radius identity). *Under the hypotheses of Theorem 6.23, the real-valued spectral radius of E is also equal to r .*

Proof. This is the real-valued reformulation of Theorem 6.23. $\square$

6.10 Spectral characterization of irreducibility

Definition 6.25 (Wolf spectral properties). A completely positive map E on $M_D(\mathbb{C})$ has the *spectral properties* of [Wol12, Thm. 6.4] if there exist a positive real number r , a positive definite right eigenvector ρ , and a positive definite left eigenvector σ for the adjoint map such that $E(\rho) = r\rho$, $E^\dagger(\sigma) = r\sigma$, every positive semidefinite right eigenvector for eigenvalue r is a scalar multiple of ρ , and the spectral radius of E is equal to r .

Theorem 6.26 (Irreducibility gives the spectral properties). *Every nonzero irreducible completely positive map on $M_D(\mathbb{C})$ has the spectral properties of Definition 6.25.*

Proof. Combine Perron–Frobenius existence for the map and its adjoint with the uniqueness theorem for positive semidefinite Perron eigenvectors and the spectral-radius identity of Theorem 6.23. $\square$

Theorem 6.27 (Spectral properties imply irreducibility). *If a completely positive map on $M_D(\mathbb{C})$ has the spectral properties of Definition 6.25, then it is irreducible.*

Proof. Gauge by the positive definite left eigenvector and rescale by the Perron eigenvalue to obtain a trace-preserving map with a positive definite fixed point. A nontrivial invariant projection would then produce, by Cesàro averaging in its corner algebra, a second positive semidefinite fixed point not proportional to the Perron vector, contradicting the uniqueness clause in Definition 6.25. $\square$

Theorem 6.28 (Irreducibility iff Wolf's spectral properties). *Let E be a nonzero completely positive map on $M_D(\mathbb{C})$. Then E is irreducible if and only if it has the spectral properties of Definition 6.25. This is [Wol12, Thm. 6.4].*

Proof. Combine Theorems 6.26 and 6.27. □

Chapter 7

Spectral Gap and Block Separation

This chapter proves that the MPV overlaps between distinct (non-gauge-phase-equivalent) MPS blocks decay to zero as system size grows. It also proves the corresponding self-overlap convergence to 1 under the complementary spectral-gap hypothesis that encodes primitivity in the cases of interest. The argument is based on a spectral analysis of the *mixed transfer operator*, following [PGVWC07].

The algebraic mixed-transfer identities in the first two sections do not use normalization. Starting with the spectral-radius estimates, we assume the trace-preserving normalization

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}.$$

7.1 Mixed transfer operator

Definition 7.1 (Mixed transfer operator). The *mixed transfer operator* (or *cross transfer operator*) for two MPS tensors A and B of the same bond dimension D is the linear map $F_{AB} : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$F_{AB}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Definition 7.2 (Rectangular mixed transfer operator). For MPS tensors A of bond dimension D_1 and B of bond dimension D_2 (possibly $D_1 \neq D_2$), the *rectangular mixed transfer operator* is $F_{AB}^{\text{rect}} : M_{D_1 \times D_2}(\mathbb{C}) \rightarrow M_{D_1 \times D_2}(\mathbb{C})$ defined by

$$F_{AB}^{\text{rect}}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Remark 7.3 (Self-transfer). Setting $B = A$ recovers the ordinary transfer map: $F_{AA} = \mathcal{E}_A$.

Theorem 7.4 (Iterated mixed transfer). For all $X \in M_D(\mathbb{C})$ and $N \geq 0$,

$$F_{AB}^N(X) = \sum_{\sigma \in \{0, \dots, d-1\}^N} A^\sigma X (B^\sigma)^\dagger,$$

where $A^\sigma = A^{\sigma_1} \dots A^{\sigma_N}$ denotes the word evaluation (Definition 2.2).

Proof. Induction on N : the base case is $F^0(X) = X$; the inductive step applies F_{AB} to each summand and re-indexes via $\{0, \dots, d-1\}^{N+1} \cong \{0, \dots, d-1\} \times \{0, \dots, d-1\}^N$. $\square$

Theorem 7.5 (Matrix trace of iterated mixed transfer at the identity). *For $N \geq 0$,*

$$\text{tr}(F_{AB}^N(\mathbb{1})) = \sum_{\sigma} \text{tr}(A^{\sigma}(B^{\sigma})^{\dagger}).$$

This is the matrix trace of F_{AB}^N evaluated at the identity matrix. It should not be confused with the operator trace $\text{Tr}(F_{AB}^N)$, which equals the MPV overlap (Theorem 7.7).

Proof. Apply the matrix trace to each summand in Theorem 7.4 with $X = \mathbb{1}$: $\text{tr}(F_{AB}^N(\mathbb{1})) = \sum_{\sigma} \text{tr}(A^{\sigma} \cdot \mathbb{1} \cdot (B^{\sigma})^{\dagger}) = \sum_{\sigma} \text{tr}(A^{\sigma}(B^{\sigma})^{\dagger})$. $\square$

7.2 MPV overlap as transfer trace

Recall from Definition 2.31 that

$$O_{AB}(N) = \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_{\sigma} \overline{V^{(N)}(B)_{\sigma}}.$$

This section rewrites that scalar as the operator trace of the mixed transfer power.

Lemma 7.6 (Trace expansion over matrix units). *For any linear endomorphism T on $M_D(\mathbb{C})$, the operator trace can be expanded as*

$$\text{Tr}(T) = \sum_{p,q=0}^{D-1} (T(E_{pq}))_{pq},$$

where E_{pq} is the matrix with 1 in position (p, q) and 0 elsewhere.

Proof. The operator trace $\text{Tr}(T)$ equals the sum of $\text{tr}(T(E_{pq}) \cdot E_{qp})$ over all (p, q) , and $\text{tr}(M \cdot E_{qp}) = M_{pq}$. $\square$

Theorem 7.7 (Overlap equals transfer trace). *For tensors A, B of bond dimension $D \geq 1$,*

$$O_{AB}(N) = \text{Tr}(F_{AB}^N).$$

Proof. Expand the operator trace using Lemma 7.6 as a sum over matrix units E_{pq} . Apply Theorem 7.4 to evaluate $F_{AB}^N(E_{pq})$, extract the (p, q) -entry, and reassemble. The resulting double sum over (p, q) and σ equals $\sum_{\sigma} V^{(N)}(A)_{\sigma} \overline{V^{(N)}(B)_{\sigma}} = O_{AB}(N)$. $\square$

Theorem 7.8 (Rectangular overlap as transfer trace). *For tensors A of bond dimension $D_1 \geq 1$ and B of bond dimension $D_2 \geq 1$,*

$$O_{AB}(N) = \text{Tr}((F_{AB}^{\text{rect}})^N).$$

Proof. Expand the operator trace over the rectangular matrix-unit basis. Then apply the rectangular iterated-transfer formula and sum over the matrix-unit indices exactly as in Theorem 7.7. $\square$

Lemma 7.9 (Word trace as entrywise inner product). *For a word $\sigma \in \{0, \dots, d-1\}^N$,*

$$\text{tr}(A^{\sigma}(B^{\sigma})^{\dagger}) = \sum_{j,k=0}^{D-1} (A^{\sigma})_{jk} \overline{(B^{\sigma})_{jk}}.$$

Proof. Expand the matrix trace and the (j, j) -entry of $A^{\sigma}(B^{\sigma})^{\dagger}$. $\square$

7.3 Eigenvalue bound and spectral gap

The eigenvalue bound comes from a uniform Frobenius-norm estimate obtained directly from the trace-preserving normalization. For the rigidity theorem, one gauges A and B separately using positive fixed points of their transfer maps to obtain unital Kraus families, and then applies the Kadison–Schwarz equality argument to a block embedding of a mixed-transfer eigenvector.

Theorem 7.10 (Eigenvalue bound). *Let A, B be normalized MPS tensors. Every eigenvalue μ of F_{AB} satisfies $|\mu| \leq 1$.*

Proof. Expand $F_{AB}^n(X)$ as a sum over words and use Cauchy–Schwarz, together with $\sum_i (A^i)^\dagger A^i = \sum_i (B^i)^\dagger B^i = \mathbb{1}$, to obtain a uniform Frobenius-norm bound $\|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for all n , where C_D depends only on D . If $F_{AB}(X) = \mu X$ and $X \neq 0$, then $|\mu|^n \|X\|_{\text{HS}} = \|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for every n . Hence $|\mu| \leq 1$.

The argument uses only the trace-preserving normalization and does not appeal to Perron–Frobenius theory. $\square$

Theorem 7.11 (Spectral radius bound). *$\rho(F_{AB}) \leq 1$ for normalized tensors.*

Proof. The spectral radius is the supremum of $|\mu|$ over all eigenvalues, and each satisfies $|\mu| \leq 1$ by Theorem 7.10. $\square$

Theorem 7.12 (Spectral radius ≥ 1 implies gauge-phase equivalence). *Let A, B be injective normalized MPS tensors. If $\rho(F_{AB}) \geq 1$, then A and B are gauge-phase equivalent.*

Proof. Since $\rho(F_{AB}) \leq 1$ (Theorem 7.11) and $\rho(F_{AB}) \geq 1$ by hypothesis, there exists an eigenvalue μ with $|\mu| = 1$ and eigenvector $X \neq 0$: $F_{AB}(X) = \mu X$.

Choose positive fixed points of the transfer maps of A and B (Theorem 6.9) and gauge the two tensors separately to unital Kraus families (Theorem 6.10). Gauge X accordingly and embed it as an off-diagonal block matrix for the block Kraus family with diagonal blocks A^i and B^i . The KS equality for peripheral eigenvectors (Theorem 5.10) and the resulting Kraus commutation relation (Theorem 5.9) yield intertwining identities between A^i , B^i , and the gauged eigenvector. Injectivity and the multiplicative-domain argument force this intertwiner to be invertible; Skolem–Noether (Theorem 3.9) then yields $B^i = \bar{\mu} X A^i X^{-1}$ for all i , establishing gauge-phase equivalence (Definition 2.13).

The crucial point is that one gauges the individual transfer maps to unital form; one does not require the mixed transfer map itself to be simultaneously unital and trace-preserving. $\square$

Theorem 7.13 (Strict spectral gap). *Let A, B be injective normalized MPS tensors that are not gauge-phase equivalent. Then $\rho(F_{AB}) < 1$.*

Proof. $\rho(F_{AB}) \leq 1$ by Theorem 7.11. If $\rho(F_{AB}) = 1$, Theorem 7.12 gives gauge-phase equivalence, contradicting the hypothesis. $\square$

7.4 Spectral gap — rectangular case

Theorem 7.14 (Rectangular spectral gap). *Let A be an injective normalized tensor of bond dimension D_1 and B an injective normalized tensor of bond dimension D_2 , with $D_1 \neq D_2$. Then $\rho(F_{AB}^{\text{rect}}) < 1$.*

Proof. The same word-expansion argument as in Theorem 7.10 gives $|\mu| \leq 1$ for every eigenvalue of F_{AB}^{rect} . If $\rho(F_{AB}^{\text{rect}}) = 1$, choose a modulus-one eigenvector X .

Gauge A and B separately to unital Kraus families and apply the same block-embedding Kadison–Schwarz argument as in the square case. It shows that $\ker(X)$ is invariant under all matrices on $\mathbb{C}^{D_2}$, so X is injective and therefore $D_2 \leq D_1$. Applying the same argument to $X^\dagger$ gives $D_1 \leq D_2$. Hence $D_1 = D_2$, contradicting the hypothesis. $\square$

Theorem 7.15 (Rectangular overlap decay). *If $D_1 \neq D_2$ and both tensors are injective and normalized, then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. The spectral gap $\rho(F_{AB}^{\text{rect}}) < 1$ (Theorem 7.14) implies $(F_{AB}^{\text{rect}})^N \rightarrow 0$, so the operator trace converges to 0. By Theorem 7.8, $O_{AB}(N) = \text{Tr}((F_{AB}^{\text{rect}})^N) \rightarrow 0$. $\square$

7.5 MPV overlap decay

Theorem 7.16 (Transfer powers converge to zero). *Let A, B be injective normalized tensors that are not gauge-phase equivalent. Then $F_{AB}^n(X) \rightarrow 0$ for every $X \in M_D(\mathbb{C})$.*

Proof. The spectral gap $\rho(F_{AB}) < 1$ (Theorem 7.13) implies $\|F_{AB}^n\|_{\text{op}} \rightarrow 0$ by the Gelfand formula, so $F_{AB}^n(X) \rightarrow 0$ for every X . $\square$

Theorem 7.17 (Overlap decay). *Let A, B be injective normalized tensors of the same bond dimension that are not gauge-phase equivalent. Then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. By Theorem 7.7, $O_{AB}(N) = \text{Tr}(F_{AB}^N)$. Expand the operator trace over matrix units (Lemma 7.6): $\text{Tr}(F_{AB}^N) = \sum_{p,q} (F_{AB}^N(E_{pq}))_{pq}$. By Theorem 7.16, each $F_{AB}^N(E_{pq}) \rightarrow 0$ entry-wise. Since the sum is finite, $O_{AB}(N) \rightarrow 0$. $\square$

7.6 Block separation

These asymptotic facts are the ingredients used later in the canonical-form separation arguments.

Remark 7.18 (Cross-correlation decay). If A and B are injective normalized tensors that are not gauge-phase equivalent, then for every $X \in M_D(\mathbb{C})$ one has $\text{tr}(F_{AB}^N(X)) \rightarrow 0$ as $N \rightarrow \infty$. This is the scalar trace consequence of the operator convergence $F_{AB}^N(X) \rightarrow 0$ from Theorem 7.16.

Remark 7.19 (Self-correlation persists). If ρ is a fixed point of $\mathcal{E}_A$, then $\text{tr}(\mathcal{E}_A^N(\rho)) = \text{tr}(\rho)$ for all $N \geq 0$. In particular this applies to the distinguished QPF fixed point from Theorem 6.9.

7.7 Primitive overlap convergence (spectral-gap form)

The theorems in this section are stated directly in terms of the complementary map $E - P$. This is the analytic form needed for the later overlap argument. For primitive normalized transfer maps, Chapter 4 supplies this spectral-gap input under explicit hypotheses.

Theorem 7.20 (Complementary spectral gap implies trace convergence to 1). *Let E be trace-preserving with fixed point ρ ($\text{tr}(\rho) \neq 0$) and let P be the fixed-point projection (Definition 4.20). If $\rho_{\text{spec}}(E - P) < 1$, then $\text{Tr}(E^n) \rightarrow 1$ as $n \rightarrow \infty$.*

Proof. By the power decomposition (Theorem 4.21), $E^n = P + (E - P)^n$ for $n \geq 1$. Since $\rho_{\text{spec}}(E - P) < 1$, the Gelfand formula gives $(E - P)^n \rightarrow 0$, so $\text{Tr}(E^n) \rightarrow \text{Tr}(P)$. The operator trace of P equals 1: since $P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$, we have $\text{Tr}(P) = \frac{\text{tr}(\rho)}{\text{tr}(\rho)} = 1$. $\square$

Theorem 7.21 (Self-overlap convergence from a complementary spectral gap). *Let A be a normalized MPS tensor with a nonzero PSD fixed point ρ of the transfer map. If $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$ (where P is the fixed-point projection), then $O_{AA}(N) \rightarrow 1$ as $N \rightarrow \infty$.*

Proof. By Theorem 7.7 (with $A = B$), $O_{AA}(N) = \text{Tr}(\mathcal{E}_A^N)$. By Theorem 7.20, $\text{Tr}(\mathcal{E}_A^N) \rightarrow 1$. $\square$

Chapter 8

Wielandt Bound

This chapter proves a quantum analog of Wielandt's inequality: for a normal MPS tensor with bond dimension D , the cumulative span of matrix products stabilises after at most D^2 steps. The results follow [SPGWC10].

8.1 Cumulative span

Definition 8.1 (Word span). The *word span* at length n is the subspace

$$S_n(A) = \text{span}_{\mathbb{C}}\{A^w : |w| = n\} \subseteq M_D(\mathbb{C}).$$

This is [SPGWC10, §II].

Definition 8.2 (Cumulative span). The *cumulative span* at level n is

$$T_n(A) = S_0(A) + S_1(A) + \cdots + S_n(A).$$

Lemma 8.3 (Monotonicity of cumulative span). $T_n(A) \leq T_{n+1}(A)$ for all n .

Proof. T_{n+1} includes all words of length $\leq n+1$, which contains all words of length $\leq n$. $\square$

Lemma 8.4 (Dimension bound). $\dim T_n(A) \leq D^2$ for all n .

Proof. $T_n(A) \subseteq M_D(\mathbb{C})$ and $\dim M_D(\mathbb{C}) = D^2$. $\square$

Theorem 8.5 (Cumulative span stabilisation). If $T_n(A) = T_{n+1}(A)$, then $T_m(A) = T_n(A)$ for all $m \geq n$.

Proof. If $T_n = T_{n+1}$, then $S_{n+1} \subseteq T_n$. Left-multiplying any element of S_{n+1} by A^i gives an element of S_{n+2} , so $S_{n+2} \subseteq T_{n+1} = T_n$. By induction, $S_m \subseteq T_n$ for all $m > n$, hence $T_m = T_n$. $\square$

Theorem 8.6 (Dimension growth). If $T_n(A) \neq T_{n+1}(A)$, then $\dim T_{n+1}(A) > \dim T_n(A)$.

Proof. $T_n \leq T_{n+1}$ (Lemma 8.3) and $T_n \neq T_{n+1}$ imply strict submodule inclusion, hence strict rank increase. $\square$

Lemma 8.7 (Word span characterises block injectivity). $S_L(A) = M_D(\mathbb{C})$ if and only if A is L -block injective.

Proof. Both conditions assert that the span of all products of length L equals $M_D(\mathbb{C})$. $\square$

8.2 Fitting decomposition

Definition 8.8 (Fitting decomposition). A *Fitting decomposition* of a linear endomorphism $f : V \rightarrow V$ on a finite-dimensional vector space over an algebraically closed field consists of:

1. f is nilpotent on the generalized 0-eigenspace,
2. f is invertible on each generalized μ -eigenspace for $\mu \neq 0$,
3. the generalized eigenspaces span V ,
4. the generalized eigenspaces are linearly independent.

Theorem 8.9 (Fitting decomposition exists). *Every linear endomorphism on a finite-dimensional vector space over an algebraically closed field admits a Fitting decomposition.*

Proof. Nilpotency on the zero generalized eigenspace follows from the definition of generalized eigenspaces. Invertibility on nonzero generalized eigenspaces follows because $f - \mu$ is nilpotent there, so $f = \mu(1 - (1 - f/\mu))$ is invertible. Spanning and independence are standard results for generalized eigenspaces over algebraically closed fields. In [SPGWC10] and [Wol12, Lemma 6.3(b)], the Jordan normal form is used directly. The argument is phrased in terms of generalized eigenspaces instead. $\square$

Theorem 8.10 (Nilpotent power bound). *If f is nilpotent on a space of dimension n , then $f^n = 0$.*

Proof. A nilpotent endomorphism on an n -dimensional space has nilpotency index $\leq n$. $\square$

8.3 Nonzero trace product

Theorem 8.11 (Cumulative span reaches $M_D(\mathbb{C})$ — explicit bound). *If A is normal, then there exists $n \leq D^2$ with $T_n(A) = M_D(\mathbb{C})$.*

Proof. By Theorem 8.6, if $T_n \neq T_{n+1}$ then $\dim T_{n+1} > \dim T_n$. Since $\dim T_n \leq D^2$ (Lemma 8.4), after at most D^2 steps either $T_n = M_D(\mathbb{C})$ or T_n has stabilised. Stabilisation before reaching $M_D(\mathbb{C})$ contradicts normality (which requires $S_{L_0} = M_D(\mathbb{C})$ for some L_0 , hence $T_{L_0} = M_D(\mathbb{C})$). $\square$

Theorem 8.12 (Cumulative span reaches $M_D(\mathbb{C})$). *If A is normal, then $T_{D^2}(A) = M_D(\mathbb{C})$.*

Proof. Take n from Theorem 8.11; since $n \leq D^2$ and T is monotone, $T_{D^2} = M_D(\mathbb{C})$. $\square$

Theorem 8.13 (Nonzero trace product). *If A is normal, then there exists a word w of length $\leq D^2$ such that $\text{tr}(A^w) \neq 0$. This is [SPGWC10, Lemma 1]. For the trace argument below, the nontrivial case is $D \geq 1$; when $D = 0$ the normality hypothesis is vacuous.*

Proof. By Theorem 8.12, $\mathbb{1} \in T_{D^2}(A)$, so $\mathbb{1}$ is a linear combination of word products of length $\leq D^2$. In the nontrivial case $D \geq 1$, at least one has nonzero trace (since $\text{tr}(\mathbb{1}) = D \neq 0$). $\square$

Theorem 8.14 (Cumulative span reaches $M_D(\mathbb{C})$ — sharp bound). *If A is normal, then $T_{D^2-d'+1}(A) = M_D(\mathbb{C})$, where $d' = \dim S_1(A) = \text{krausRank}(A)$. This is the sharp form of [SPGWC10, Lemma 1]: the cumulative span already reaches $M_D(\mathbb{C})$ by index $D^2 - d' + 1 \leq D^2$.*

Proof. The key observation is that if $T_n(A) \neq T_{n+1}(A)$, the dimension strictly increases. Starting from $T_1(A) = S_1(A)$ with dimension d' , after at most $D^2 - d'$ additional strict-growth steps the dimension reaches $D^2 = \dim M_D(\mathbb{C})$. Stabilisation before reaching $M_D(\mathbb{C})$ contradicts normality. Hence $T_{D^2-d'+1}(A) = M_D(\mathbb{C})$. $\square$

Theorem 8.15 (Sharp nonzero trace product). *If A is normal, then there exists a word w of length $\leq D^2 - \text{krausRank}(A) + 1$ such that $\text{tr}(A^w) \neq 0$. This is [SPGWC10, Lemma 1] with the sharp length bound.*

Proof. By Theorem 8.14, the identity $\mathbb{1}$ lies in $T_{D^2-d'+1}(A)$, hence is a linear combination of word products of length $\leq D^2 - d' + 1$. At least one such product has nonzero trace. $\square$

Theorem 8.16 (Sharp nonzero trace product — positive length). *If A is normal and $D \geq 2$, then there exists a word w of positive length $1 \leq |w| \leq D^2 - \text{krausRank}(A) + 1$ such that $\text{tr}(A^w) \neq 0$. This strengthens Theorem 8.15 by requiring $|w| \geq 1$, which is needed for the blocking argument in Theorem 8.28.*

Proof. Define the *positive-level* cumulative span $V_n = \text{span}\{A^w : 1 \leq |w| \leq n\}$. For $D \geq 2$, the span $S_0(A) = \text{span}\{\mathbb{1}\}$ has dimension 1, so $V_1 \supseteq S_1(A)$ has dimension $\geq d' \geq 1$. The same stabilization-or-strict-growth dichotomy shows $V_{D^2-d'+1} = M_D(\mathbb{C})$: if V_n stabilises and is not $M_D(\mathbb{C})$, then V_n is closed under left-multiplication by each A^i , so $\text{wordSpan}(A, N) \subseteq V_n \neq M_D(\mathbb{C})$ for all $N \geq 1$, contradicting normality. Since $V_{D^2-d'+1} = M_D(\mathbb{C})$ and $D \geq 2$ implies $\mathbb{1} \notin V_{D^2-d'+1}$ (dimension of $M_D(\mathbb{C})$ is $D^2 \geq 4$, but $\text{span}\{\mathbb{1}\}$ has dimension 1), at least one generator has nonzero trace. $\square$

8.4 Eigenvector extraction

Theorem 8.17 (Nonzero trace implies eigenvector). *If $M \in M_D(\mathbb{C})$ has $\text{tr}(M) \neq 0$, then M has a nonzero eigenvalue $\mu \neq 0$ and a corresponding eigenvector $\varphi \neq 0$ with $M\varphi = \mu\varphi$.*

Proof. A matrix with nonzero trace has a nonzero eigenvalue (since trace = sum of eigenvalues counted with multiplicity). The Fitting decomposition (Theorem 8.9) ensures that the corresponding generalized eigenspace is nontrivial. On that generalized eigenspace, $M - \mu\mathbb{1}$ is nilpotent, so its kernel is nontrivial. Any nonzero vector in $\ker(M - \mu\mathbb{1})$ is therefore a genuine eigenvector with eigenvalue μ . $\square$

Theorem 8.18 (Eigenvalue extraction). *If A is normal, then there exist a word w_0 with $|w_0| \leq D^2$, a nonzero scalar $\mu \neq 0$, and a nonzero vector $\varphi \neq 0$ such that $A^{w_0}\varphi = \mu\varphi$.*

Proof. By Theorem 8.13, there is a word w_0 with $\text{tr}(A^{w_0}) \neq 0$ and $|w_0| \leq D^2$. By Theorem 8.17, A^{w_0} has a nonzero eigenvector. $\square$

8.5 Eigenvector spreading

Definition 8.19 (Cumulative vector span). Given an MPS tensor A and a vector $\varphi \in \mathbb{C}^D$, the *cumulative vector span* is

$$K_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w\varphi : |w| \leq n\} \subseteq \mathbb{C}^D.$$

This is [SPGWC10, proof of Lemma 2(a)].

Theorem 8.20 (Vector span stabilisation). *If $K_n(A, \varphi) = K_{n+1}(A, \varphi)$, then $K_m(A, \varphi) = K_n(A, \varphi)$ for all $m \geq n$.*

Proof. Same argument as Theorem 8.5: left-multiplication by A^i cannot escape the stabilised span. $\square$

Lemma 8.21 (Vector span from matrix span). *If $T_N(A) = M_D(\mathbb{C})$ and $\varphi \neq 0$, then $K_N(A, \varphi) = \mathbb{C}^D$.*

Proof. Every matrix $M \in M_D(\mathbb{C})$ is in $T_N(A)$, so in particular every rank-one matrix sending φ to any target vector v is in T_N . Applying such matrices to φ recovers all of $\mathbb{C}^D$. $\square$

Theorem 8.22 (Eigenvector spreading). *Let A be a normal MPS tensor and let $A^{i_0}\varphi = \mu\varphi$ with $\mu \neq 0$ and $\varphi \neq 0$. Then $K_{D-1}(A, \varphi) = \mathbb{C}^D$. This is [SPGWC10, Lemma 2(a)].*

Proof. The vector span K_n is monotone and $\dim K_n \leq D$. If $K_n = K_{n+1}$ for some $n < D-1$, the span stabilises (Theorem 8.20). But $T_{D^2}(A) = M_D(\mathbb{C})$ by Theorem 8.12, so Lemma 8.21 gives $K_{D^2}(A, \varphi) = \mathbb{C}^D$, contradicting early stabilisation. Hence $\dim K_n$ grows by ≥ 1 at each step, reaching D by step $D-1$. $\square$

8.6 Assembly

Definition 8.23 (Wielandt analysis). A *Wielandt analysis* for a normal MPS tensor A packages: a word w_0 of length $\leq D^2$, a nonzero eigenvalue μ , and a nonzero eigenvector φ of the matrix A^{w_0} , such that $A^{w_0}\varphi = \mu\varphi$.

Theorem 8.24 (Wielandt analysis exists). *Every normal MPS tensor admits a Wielandt analysis.*

Proof. Immediate from Theorem 8.18. $\square$

Theorem 8.25 (Wielandt chain). *Every normal MPS tensor admits a Wielandt analysis consisting of data (w_0, μ, φ) with $K_{D-1}(A, \varphi) = \mathbb{C}^D$.*

Proof. Combine Theorems 8.24 and 8.22. $\square$

Theorem 8.26 (Wielandt bound). *If A is a normal MPS tensor with bond dimension D , then $T_{D^2}(A) = M_D(\mathbb{C})$. In particular, the cumulative span stabilises at index at most D^2 . This is the cumulative-span part of [SPGWC10, Theorem 1]. The paper's theorem is stronger: via Lemma 2(b) it upgrades this to a fixed-length conclusion $S_N(A) = M_D(\mathbb{C})$, proved below in Theorem 8.40.*

Proof. Immediate from Theorem 8.12. $\square$

Remark 8.27 (Explicit blocking-length estimates). Theorem 8.26 gives the D^2 cumulative-span bound. The full-rank index bound $\iota(A) \leq (D^2 - d + 1) \cdot D^2$ (Theorem 1 case (1) of [SPGWC10]) is proved in Theorem 8.28 below. The sharp blocking-length estimates for injectivity (D^4) and bi-canonical form ($3D^5$) from [SPGWC10] are not proved here.

Theorem 8.28 (Wielandt's inequality — general bound). *Let A be a normalized MPS tensor ($\sum_i (A^i)^\dagger A^i = \mathbb{1}$) that is primitive in the paper sense. Then*

$$\iota(A) \leq (D^2 - \text{krausRank}(A) + 1) \cdot D^2,$$

where $\iota(A) = \min\{n : S_n(A) = M_D(\mathbb{C})\}$ is the full-Kraus-rank index. This is [SPGWC10, Theorem 1, case (1)].

Proof.

1. $D = 1$: In this case $\iota(A) = 0$, so the bound holds trivially.
2. $D \geq 2$: By Theorem 8.16, there exists a *positive-length* word w with $|w| \leq D^2 - d' + 1$ and $\text{tr}(A^w) \neq 0$, where $d' = \text{krausRank}(A)$. The matrix A^w has nonzero trace, hence a nonzero eigenvalue μ and eigenvector φ (Theorem 8.17). Set $n = |w|$ and let $B = A^{[n]}$ be the blocked tensor. Then B is normal (Theorem 8.35), and the encoding of w as a single block index gives $B^{i_0} = A^w$.

- (a) If A^w is invertible (invertible case), $S_{D^2}(B) = M_D(\mathbb{C})$, whence $S_{D^2 \cdot n}(A) = M_D(\mathbb{C})$.
- (b) If A^w is non-invertible (non-invertible case), the eigenvector φ spans a proper subspace, and the Wielandt chain argument gives $S_{D^2}(B) = M_D(\mathbb{C})$, whence $S_{D^2 \cdot n}(A) = M_D(\mathbb{C})$.

In both cases $\iota(A) \leq D^2 \cdot n \leq D^2 \cdot (D^2 - d' + 1)$.

□

8.7 Lemma 2(b): fixed-length matrix spanning

The preceding sections establish the cumulative-spanning part of the Lemma 2(a) argument from [SPGWC10]: eigenvector spreading gives a cumulative vector span $K_{D-1}(A, \varphi) = \mathbb{C}^D$. The next stage is to pass to a span at one fixed length, namely to find n_0 such that $S_{n_0}(A) = M_D(\mathbb{C})$ (i.e. word products of *exactly* one length span all matrices).

Definition 8.29 (Fixed-length vector span). The *fixed-length vector span* at length n is

$$H_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w \varphi : |w| = n\} \subseteq \mathbb{C}^D.$$

This is $S_n(A)|\varphi\rangle$ in the notation of [SPGWC10].

Lemma 8.30 (Word span products). $S_m(A) \cdot S_n(A) \subseteq S_{m+n}(A)$: the product of word spans at lengths m and n is contained in the word span at length $m + n$.

Proof. Every product $A^{w_1} A^{w_2}$ with $|w_1| = m$ and $|w_2| = n$ equals $A^{w_1 w_2}$ with $|w_1 w_2| = m + n$. □

Lemma 8.31 (Eigenvector padding). Suppose $A^{i_0} \varphi = \mu \varphi$ with $\mu \neq 0$. Then for all n ,

$$K_n(A, \varphi) = H_n(A, \varphi).$$

In particular, if $K_n(A, \varphi) = \mathbb{C}^D$ then already $H_n(A, \varphi) = \mathbb{C}^D$.

Proof. Any word w with $|w| \leq n$ can be padded to exact length n by appending $n - |w|$ copies of the index i_0 . The eigenvector relation gives $A^{w \cdot i_0^k} \varphi = \mu^k \cdot A^w \varphi$, so the padded vector is a nonzero scalar multiple of the original. Hence every generator of K_n lies in H_n . The reverse inclusion $H_n \leq K_n$ is immediate. □

Theorem 8.32 (Vector spanning to matrix spanning (assembly step)). Assume:

1. $H_n(A, \varphi) = \mathbb{C}^D$ (length- n word products applied to φ span all of $\mathbb{C}^D$), and
2. for each standard basis vector e_j , the rank-one operator $|\varphi\rangle\langle e_j|$ lies in $S_m(A)$.

Then $S_{n+m}(A) = M_D(\mathbb{C})$.

Proof. Fix a matrix unit E_{ij} . By hypothesis (1), there exists $M_i \in S_n(A)$ with $M_i\varphi = e_i$. By hypothesis (2), $|\varphi\rangle\langle e_j| \in S_m(A)$. Their product satisfies

$$M_i \cdot |\varphi\rangle\langle e_j| = |M_i\varphi\rangle\langle e_j| = |e_i\rangle\langle e_j| = E_{ij},$$

and $E_{ij} \in S_{n+m}(A)$ by Lemma 8.30. Since the matrix units span $M_D(\mathbb{C})$, $S_{n+m}(A) = M_D(\mathbb{C})$. $\square$

Lemma 8.33 (Blocking transfer). *If the blocked tensor $A^{[L]}$ satisfies $S_n(A^{[L]}) = M_D(\mathbb{C})$, then $S_{nL}(A) = M_D(\mathbb{C})$.*

Proof. Each length- n word in the blocked alphabet $\{0, \dots, d^L - 1\}$ decodes to a length- nL word in the original alphabet $\{0, \dots, d - 1\}$, so $S_n(A^{[L]}) \subseteq S_{nL}(A)$. $\square$

Remark 8.34 (Relation with Lemma 2(b)). In [SPGWC10], Lemma 2(b) combines a rank-one extraction for fixed-length spans with an assembly argument turning fixed-length vector spanning into fixed-length matrix spanning. Theorem 8.32 is the second step. Theorem 8.39 supplies the rank-one input in blocked form, and Theorem 8.40 then gives a blocked, fixed-length word-span saturation result.

8.8 Blocked tensor and rectangular span

This section extends the Lemma 2(b) infrastructure to the *blocked tensor* $A^{[L]}$ and develops the “rectangular span” used to produce rank-one elements.

Theorem 8.35 (Blocking preserves normality). *If A is normal and $L > 0$, then the blocked tensor $A^{[L]}$ is also normal.*

Proof. Choose N with $S_N(A) = M_D(\mathbb{C})$. Then $\mathbb{1} \in S_N(A)$. For any $M \in S_N(A)$, write $\mathbb{1} = \sum_r c_r A^{w_r}$ with each $|w_r| = N$. Then

$$M = M \cdot \mathbb{1} = \sum_r c_r M A^{w_r} \in S_N(A) \cdot S_N(A) \subseteq S_{2N}(A)$$

by Lemma 8.30. Iterating this inclusion gives $S_N(A) \subseteq S_{kN}(A)$ for every $k \geq 1$; taking $k = L$ yields $M_D(\mathbb{C}) = S_N(A) \subseteq S_{NL}(A)$. By Lemma 8.36, $S_{NL}(A) \subseteq S_n(A^{[L]})$, so $S_n(A^{[L]}) = M_D(\mathbb{C})$. $\square$

Lemma 8.36 (Chunking: word span embeds into blocked word span). *$S_{nL}(A) \subseteq S_n(A^{[L]})$ for all n, L .*

Proof. Every length- nL word in the original alphabet can be split into n consecutive chunks of length L , each of which is a single blocked Kraus operator. By induction on n , using the word span product lemma (Lemma 8.30). $\square$

Theorem 8.37 (Word eigenvector gives blocked eigenvector). *If $A^{w_0}\varphi = \mu\varphi$ for a word w_0 of length L , then $(A^{[L]})^{j_0}\varphi = \mu\varphi$, where j_0 is the encoded blocked index corresponding to w_0 .*

Proof. By definition, $(A^{[L]})^{j_0} = A^{w_0}$. $\square$

Theorem 8.38 (Wielandt blocked assembly). *Suppose A is normal and $L > 0$, and we have:*

1. *a word w_0 of length L with eigenvector $A^{w_0}\varphi = \mu\varphi$ ($\mu \neq 0$, $\varphi \neq 0$),*

2. a word τ_0 of length L with transpose eigenvector $(A^{\tau_0})^T \psi = \nu \psi$ ($\nu \neq 0$, $\psi \neq 0$),
3. a rank-one element $\varphi \psi^T \in S_m(A^{[L]})$ for some m .

Then $S_{((D-1)+m+(D-1)) \cdot L}(A) = M_D(\mathbb{C})$.

Proof. Apply the conditional Lemma 2(b) assembly (Theorem 8.32) to the blocked tensor $A^{[L]}$, which is normal by Theorem 8.35. The eigenvector and transpose eigenvector are transferred to single-index eigenvectors of $A^{[L]}$ by Theorem 8.37. Transfer back to the original tensor via Lemma 8.33. $\square$

Theorem 8.39 (Rank-one extraction for blocked tensor). *Suppose A is normal and $N_0 \geq 1$ with $S_{N_0}(A) = M_D(\mathbb{C})$. Then there exist words σ_0, τ_0 of length N_0 , nonzero vectors φ, ψ , nonzero scalars μ, ν , and $m \in \mathbb{N}$ such that:*

1. $A^{\sigma_0} \varphi = \mu \varphi$,
2. $(A^{\tau_0})^T \psi = \nu \psi$,
3. $\varphi \psi^T \in S_m(A^{[N_0]})$.

Proof. Set $B = A^{[N_0]}$. Since $S_{N_0}(A) = M_D(\mathbb{C})$, we have $S_1(B) = M_D(\mathbb{C})$ by definition of the blocked tensor. Hence $\mathbb{1} \in S_1(B)$, so in the nontrivial case $D \geq 1$ not all blocked generators can have trace zero. Choose j_0 with $\text{tr}(B^{j_0}) \neq 0$. By Theorem 8.17, B^{j_0} has a nonzero eigenvector φ with eigenvalue $\mu \neq 0$. Writing j_0 as the blocked index corresponding to a word σ_0 of length N_0 gives $A^{\sigma_0} \varphi = \mu \varphi$. Applying the same argument to the transpose blocked tensor gives a word τ_0 of length N_0 , a nonzero vector ψ , and $\nu \neq 0$ with $(A^{\tau_0})^T \psi = \nu \psi$. Since A is normal and $N_0 \geq 1$, the blocked tensor B is normal by Theorem 8.35. Therefore there exists M with $S_M(B) = M_D(\mathbb{C})$. In particular $\varphi \psi^T \in S_M(B)$. Taking $m = M$ proves (3). $\square$

Theorem 8.40 (Fixed-length word-span saturation via blocking). *If A is a normal MPS tensor with bond dimension D , then there exists N such that $S_N(A) = M_D(\mathbb{C})$. This gives the fixed-length spanning conclusion corresponding to [SPGWC10, Lemma 2(b)].*

Proof. By normality, some N_0 has $S_{N_0}(A) = M_D(\mathbb{C})$. If $N_0 = 0$ we are done. Otherwise, Theorem 8.39 produces eigenvectors and a rank-one element $\varphi \psi^T \in S_m(A^{[N_0]})$. Theorem 8.38 then gives $S_{((D-1)+m+(D-1)) \cdot N_0}(A) = M_D(\mathbb{C})$. $\square$

Remark 8.41 (Blocking route to fixed-length spanning). Theorem 8.40 is obtained by first extracting a rank-one operator in a blocked tensor (Theorem 8.39) and then applying the fixed-length assembly step (Theorem 8.38). Thus the fixed-length span of A is reached through a blocked auxiliary tensor rather than by a direct unblocked argument.

8.9 Primitive MPS tensors

Definition 8.42 (Primitive MPS tensor). Assume $D > 0$. A tensor A together with a matrix $\rho \in M_D(\mathbb{C})$ is *primitive* if

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}, \quad \rho \geq 0, \quad \rho \neq 0, \quad \mathcal{E}_A(\rho) = \rho,$$

and, if P is the fixed-point projection associated to ρ , then $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$. When the choice of ρ is irrelevant, we simply say that A is a primitive MPS tensor. This is weaker than the paper's notion, which requires $\rho > 0$; see Remark 8.48. The stronger positive-definite hypothesis is imposed separately in Theorems 8.49 and 8.60.

Theorem 8.43 (Primitive overlap convergence). *If A is a primitive MPS tensor, then $O_{AA}(N) \rightarrow 1$.*

Proof. Immediate from the self-overlap convergence under a complementary spectral gap (Theorem 7.21). $\square$

8.10 Primitivity and normality

Throughout this section, primitivity means the condition of Definition 8.42: the tensor is in TP gauge, it has a nonzero PSD fixed point, and the complementary map has spectral radius < 1 .

Theorem 8.44 (Primitive MPS tensors have nonzero word products). *If A is a primitive MPS tensor, then for every $n \geq 0$ there exists a word w of length exactly n with $A^w \neq 0$.*

Proof. The PSD fixed point ρ of $\mathcal{E}_A$ satisfies $\mathcal{E}_A^n(\rho) = \rho \neq 0$. Since $\mathcal{E}_A^n(\rho) = \sum_{|w|=n} A^w \rho (A^w)^\dagger$, at least one summand is nonzero, hence some $A^w \neq 0$. $\square$

Theorem 8.45 (Iterated transfer does not vanish). *If A is a primitive MPS tensor, then $\mathcal{E}_A^n \neq 0$ for all $n \geq 0$.*

Proof. By Theorem 8.44, there exists a word w of length n with $A^w \neq 0$, so $\mathcal{E}_A^n(\rho)$ has a nonzero summand. $\square$

8.10.1 Primitivity-to-normality bridge

The primitivity condition of Definition 8.42 combines TP normalization, a nonzero PSD fixed point, and a complementary spectral gap. It is connected to normality through the following results.

Theorem 8.46 (Fixed-point uniqueness from spectral gap). *If A is a primitive MPS tensor with PSD fixed point ρ , then every fixed point σ of $\mathcal{E}_A$ is proportional to ρ : $\sigma = \frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \rho$.*

Proof. Set $\sigma' = \sigma - \frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \rho$. Then $\text{tr}(\sigma') = 0$ and $(E - P_\rho)(\sigma') = \sigma'$. If $\sigma' \neq 0$, then 1 is an eigenvalue of $E - P_\rho$, contradicting the spectral gap $\rho_{\text{spec}}(E - P_\rho) < 1$. $\square$

Theorem 8.47 (Complement powers tend to zero). *If A is a primitive MPS tensor with PSD fixed point ρ , then $(\mathcal{E}_A - P_\rho)^n \rightarrow 0$ in operator norm.*

Proof. The spectral radius of $\mathcal{E}_A - P_\rho$ is strictly less than 1 by the spectral-gap hypothesis. Apply the general result that powers of an operator with spectral radius < 1 tend to zero. $\square$

Remark 8.48 (Primitivity: spectral gap vs. positive definiteness). The primitivity condition used in §8.10.1 has three parts: TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, a nonzero PSD fixed point ρ , and the spectral gap $\rho_{\text{spec}}(\mathcal{E}_A - P_\rho) < 1$. The notion of “primitive” in [SPGWC10, Proposition 3] additionally requires ρ to be positive definite (full rank). Without positive definiteness, the spectral-gap condition alone does *not* imply irreducibility or normality. A counterexample is $D = 2$, $\rho = \text{diag}(1, 0)$.

With the additional hypothesis $\rho > 0$, Theorem 8.49 gives irreducibility. Together with the Burnside bridge (Chapter 9, §9.4.1) and Theorem 8.58, one obtains

$$\text{primitive} + \rho \text{ PosDef} \Rightarrow \text{irreducible} \Rightarrow \text{algSpan}(A) = M_D(\mathbb{C}) \Rightarrow \text{normal}.$$

The last step uses the aperiodicity condition $\mathbb{1} \in S_1(A)$ from Remark 8.53. After restoring the positive-definite hypothesis, [SPGWC10, Proposition 3] identifies the paper’s primitive/strongly irreducible condition with eventual full Kraus rank, i.e. with our notion of normality.

8.10.2 Primitive implies irreducible

Theorem 8.49 (Primitive with PosDef fixed point implies irreducible tensor). *If A is a primitive MPS tensor with PSD fixed point ρ and ρ is positive definite, then A is an irreducible tensor. This is part of [SPGWC10, Proposition 3] (see also [CPGSV17, §2.3]).*

Proof. By contrapositive. Assume A has an invariant projection P ($P \neq 0$, $P \neq \mathbb{1}$, $(\mathbb{1} - P)A^i P = 0$ for all i). Set $\sigma = P\rho P$.

1. *Invariance of σ under $\mathcal{E}_A^n$:* the projection relation $(\mathbb{1} - P)A^i P = 0$ implies $(\mathbb{1} - P) \cdot \mathcal{E}_A^n(\sigma) = 0$ for all n (by induction on n).
2. *Convergence:* By Theorem 8.47, $\mathcal{E}_A^n(\sigma) \rightarrow \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Hence $(\mathbb{1} - P) \cdot \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho = 0$.
3. *Nonzero scalar:* Since $P \neq 0$ and ρ is positive definite, $\text{tr}(P\rho P) > 0$, so $\frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \neq 0$. Therefore $(\mathbb{1} - P)\rho = 0$.
4. *Contradiction:* Since ρ is positive definite, it is a unit in $M_D(\mathbb{C})$. From $(\mathbb{1} - P)\rho = 0$ we get $\mathbb{1} - P = 0$, i.e. $P = \mathbb{1}$, contradicting $P \neq \mathbb{1}$.

□

8.10.3 Primitivity bridge to strong irreducibility and normality

The results in this subsection remove the aperiodicity assumption from the normality conclusions. They connect the spectral-gap primitivity of Definition 8.42 directly to normality (eventually full Kraus rank), using the convergence of the transfer-map iterates to the projection onto the fixed-point subspace.

Theorem 8.50 (Primitive MPS with PosDef fixed point has irreducible transfer map). *If A is a primitive MPS tensor with positive-definite fixed point ρ , then the transfer map $\mathcal{E}_A$ is irreducible.*

The proof uses fixed-point uniqueness (Theorem 8.46): every PSD fixed point is proportional to ρ , so Wolf's criterion for irreducibility (positive-definite fixed point implies irreducibility) applies directly.

Proof. By Theorem 8.46, if $\sigma \geq 0$ and $\mathcal{E}_A(\sigma) = \sigma$, then $\sigma = \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Since $\rho > 0$, Wolf's uniqueness criterion for irreducibility applies. □

Theorem 8.51 (Primitive MPS with PosDef fixed point is strongly irreducible). *If A is a primitive MPS tensor with positive-definite fixed point ρ , then A is strongly irreducible in the sense of [SPGWC10, Proposition 3(c)]: $\mathcal{E}_A$ is irreducible, the fixed point ρ is positive definite, and the peripheral spectrum of $\mathcal{E}_A$ is $\{1\}$.*

Proof. Combine Theorem 8.50 (irreducibility) with the peripheral-spectrum primitivity from the spectral-gap hypothesis (Theorem 8.43). □

Theorem 8.52 (Primitive MPS with PosDef fixed point is normal). *If A is a primitive MPS tensor with positive-definite fixed point ρ , then A is normal (has eventually full Kraus rank). No aperiodicity assumption is needed.*

This is the key bridge between the spectral-gap primitivity of Definition 8.42 and normality, filling the gap that the aperiodicity-based route in §8.11 leaves open.

Proof. By Theorem 8.51, A is strongly irreducible. The implication (c) $\Rightarrow$ (b) of [SPGWC10, Proposition 3] (strong irreducibility implies eventually full Kraus rank) then gives normality. (Formally: strong irreducibility implies $\langle A \rangle_n = M_D(\mathbb{C})$ for large n , which is exactly the definition of normality.) □

8.11 Cumulative span to word span

Cumulative spanning ($T_N(A) = M_D(\mathbb{C})$, the union of word spans of all lengths $\leq N$) is weaker than normality ($S_L(A) = M_D(\mathbb{C})$ for a single length L). The gap is closed by an aperiodicity hypothesis.

Remark 8.53 (Aperiodicity condition). We say A satisfies the *aperiodicity condition* if $\mathbb{1} \in S_1(A)$, i.e. the identity matrix lies in the span of the Kraus operators $\{A^i\}$. This is an additional hypothesis. The one-sided TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ does *not* imply $\mathbb{1} \in S_1(A)$; it only gives a quadratic identity among the Kraus operators. For the paper-style primitive condition (peripheral-spectrum primitivity together with a positive-definite fixed point), one expects aperiodicity to follow from spectral analysis of the transfer map. That implication is not used in the present Wielandt bridge, so the condition is kept explicit.

Remark 8.54 (Counterexample: cumulative spanning does not imply normality). In $M_2(\mathbb{C})$, let $A^0 = E_{12}$ and $A^1 = E_{21}$ (the off-diagonal matrix units). Then $\text{algSpan}(A) = M_2(\mathbb{C})$ and $T_2(A) = M_2(\mathbb{C})$, but the word spans alternate: $S_n(A)$ contains only off-diagonal matrices for odd n and only diagonal matrices for even n . Hence $S_n(A) \neq M_2(\mathbb{C})$ for every n , and A is not normal. The aperiodicity condition $\mathbb{1} \in S_1(A)$ fails: $S_1(A) = \text{span}_{\mathbb{C}}\{E_{12}, E_{21}\}$.

Theorem 8.55 (Word span monotonicity from aperiodicity). *If $\mathbb{1} \in S_1(A)$, then $S_n(A) \leq S_{n+1}(A)$ for all n .*

Proof. For any $M \in S_n(A)$, $M = M \cdot \mathbb{1} \in S_n(A) \cdot S_1(A) \subseteq S_{n+1}(A)$ by Lemma 8.30. $\square$

Theorem 8.56 (Cumulative span equals word span under aperiodicity). *If $\mathbb{1} \in S_1(A)$, then $T_n(A) = S_n(A)$ for all n .*

Proof. By Theorem 8.55, the word spans are monotone: $S_0 \leq S_1 \leq \dots$. The supremum $T_n = S_0 + \dots + S_n$ therefore equals the largest term S_n . $\square$

Theorem 8.57 (Cumulative spanning plus aperiodicity implies normality). *If $T_N(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. By Theorem 8.56, $S_N(A) = T_N(A) = M_D(\mathbb{C})$. $\square$

Theorem 8.58 (Algebra span plus aperiodicity implies normality). *If $\text{algSpan}(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. Since $\text{algSpan}(A) = M_D(\mathbb{C})$, there exists N with $T_N(A) = M_D(\mathbb{C})$ (by the ascending chain condition on finite-dimensional subspaces). Apply Theorem 8.57. $\square$

Theorem 8.59 (Irreducible tensor plus aperiodicity implies normality). *If A is an irreducible tensor, $D > 0$, and $\mathbb{1} \in S_1(A)$, then A is normal. This assembles the full chain:*

$$\text{irreducible tensor} \xrightarrow{9.20} \text{irreducible action} \xrightarrow{\text{Burnside}} \text{algSpan} = M_D(\mathbb{C}) \xrightarrow{8.58} \text{normal}.$$

This corresponds to the irreducibility-to-normality direction of [SPGWC10, Proposition 3]. For the Burnside step, we appeal forward to Theorems 9.20 and 9.21 in Chapter 9; this is a presentation-level forward dependency, but the mathematical dependency is acyclic.

Proof. By Theorem 9.20, A acts irreducibly on $\mathbb{C}^D$. By Theorem 9.21, $\text{algSpan}(A) = M_D(\mathbb{C})$. By Theorem 8.58 with the aperiodicity hypothesis, A is normal. $\square$

Theorem 8.60 (Primitive + positive definite fixed point + aperiodicity imply normality). *If A is a primitive MPS tensor with fixed point ρ , ρ is positive definite, and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. By Theorem 8.49, A is an irreducible tensor. Apply Theorem 8.59. $\square$

Remark 8.61 (The normal-canonical route bypasses the aperiodicity bridge). The aperiodicity results in §8.11 are needed only for the block-injective route through Definition 2.19, where one upgrades cumulative spanning to eventual block injectivity. The normal-canonical route used later in Chapters 9–11 does not pass through this step: it works directly with irreducible TP-normalized blocks whose blocked transfer maps are primitive and, once the corresponding weighted block data are available, packages them as normal canonical form in the sense of [CPGSV17]. There are analogous cumulative-span lemmas for blocked irreducible tensors and eigenvector-spreading arguments, but they are auxiliary here and we do not list them separately.

Chapter 9

Canonical Form Reduction

This chapter reduces a general MPS tensor to block-diagonal data via iterated invariant-subspace decomposition. Two reduction schemes are relevant. The block-injective route aims at injective TP-normalized blocks and the canonical form predicate of Chapter 10. The normal-canonical route packages a weighted family of irreducible TP-normalized blocks with primitive transfer maps and pairwise distinct nonzero weight moduli, in the sense of [CPGSV17]. The Perron–Frobenius eigenvector existence and TP-gauge construction are in Chapter 6; this chapter uses them to produce CFII data and the normal canonical form packaging. The reduction draws on [PGVWC07], [CPGSV21], and [Wol12, Ch. 6].

9.1 Block-diagonal assembly

Definition 9.1 (Block-diagonal tensor from blocks). Recall the block-diagonal tensor of Definition 2.23. Given r blocks with bond dimensions $(D_k)_{k=1}^r$, per-block MPS tensors $\{A_k^i\}$, and scaling factors $(\mu_k)_{k=1}^r$, we write $\bigoplus_{k=1}^r \mu_k A_k$ for the tensor of total bond dimension $D = \sum_k D_k$ whose Kraus operators are $A^i = \bigoplus_{k=1}^r \mu_k A_k^i$.

Theorem 9.2 (Per-block single-block FT). *If all block tensors A_k are injective and the per-block same-MPV condition holds (A_k and B_k generate the same MPV family for each k), then each pair (A_k, B_k) is gauge equivalent.*

Proof. Apply the single-block fundamental theorem (Theorem 3.11) to each block independently. $\square$

Theorem 9.3 (Per-block same MPV implies global same MPV). *If A_k and B_k generate the same MPV family for each block k , then the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ also generate the same MPV family.*

Proof. By the MPV decomposition (Theorem 2.24), each MPV is $\sum_k \mu_k^N V^{(N)}(A_k)_\sigma$. Per-block equality of MPV coefficients gives global equality. $\square$

Theorem 9.4 (Global gauge from block gauge). *If each pair (A_k, B_k) is gauge equivalent ($B_k^i = X_k A_k^i X_k^{-1}$), then the block-diagonal tensors are gauge equivalent via the block-diagonal matrix $X = \bigoplus_k X_k$.*

Proof. Block-diagonal conjugation: $X(\bigoplus_k \mu_k A_k^i)X^{-1} = \bigoplus_k \mu_k X_k A_k^i X_k^{-1} = \bigoplus_k \mu_k B_k^i$. $\square$

9.2 Transfer map normalization

Theorem 9.5 (Scaling preserves injectivity). *If A is injective and $\zeta \neq 0$, then ζA is injective.*

Proof. Scaling by a nonzero scalar preserves linear independence and hence spanning. $\square$

Theorem 9.6 (Transfer map under scaling). $\mathcal{E}_{\zeta A}(X) = |\zeta|^2 \mathcal{E}_A(X)$.

Proof. Each summand: $(\zeta A^i)X(\zeta A^i)^\dagger = |\zeta|^2 A^i X(A^i)^\dagger$. $\square$

Theorem 9.7 (Phase-scaling preserves normalization). *If $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $|\zeta| = 1$, then $\sum_i (\zeta A^i)^\dagger (\zeta A^i) = \mathbb{1}$.*

Proof. $(\zeta A^i)^\dagger (\zeta A^i) = \bar{\zeta} \zeta (A^i)^\dagger A^i = (A^i)^\dagger A^i$, since $\bar{\zeta} \zeta = |\zeta|^2 = 1$. $\square$

Theorem 9.8 (MPV under block normalization). *For any nonzero scaling factors μ_k , set $\eta_k := \mu_k/|\mu_k|$ and $B_k := |\mu_k|A_k$. Then $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \eta_k B_k$ generate the same MPV family. Equivalently, the k -th block contribution factors as*

$$\mu_k^N V^{(N)}(A_k)_\sigma = \eta_k^N |\mu_k|^N V^{(N)}(A_k)_\sigma,$$

so the modulus contributes only through $|\mu_k|^N$, while the remaining factor η_k is a unit-modulus phase.

Proof. By the MPV decomposition (Theorem 2.24), the k -th contribution of $\bigoplus_k \mu_k A_k$ is $\mu_k^N V^{(N)}(A_k)_\sigma$. For $B_k = |\mu_k|A_k$, Lemma 2.12 gives $V^{(N)}(B_k)_\sigma = |\mu_k|^N V^{(N)}(A_k)_\sigma$. Hence $\eta_k^N V^{(N)}(B_k)_\sigma = \eta_k^N |\mu_k|^N V^{(N)}(A_k)_\sigma = \mu_k^N V^{(N)}(A_k)_\sigma$ for every block, so the total MPV families agree. $\square$

9.3 Invariant subspace decomposition

Theorem 9.9 (Unitary conjugation preserves the MPV family). *If U is a unitary matrix, then A and $U^\dagger A U$ generate the same MPV family.*

Proof. By cyclicity of the trace: $\text{tr}((U^\dagger A^{i_1} U) \dots (U^\dagger A^{i_N} U)) = \text{tr}(A^{i_1} \dots A^{i_N})$. $\square$

Definition 9.10 (Support projection). For a positive semi-definite matrix $\rho \geq 0$, the *support projection* P is the orthogonal projection onto the range of ρ . It is constructed via the spectral decomposition: if $\rho = U \text{diag}(\lambda_1, \dots, \lambda_D) U^\dagger$, then $P = U \text{diag}(\mathbf{1}_{\lambda_1 > 0}, \dots, \mathbf{1}_{\lambda_D > 0}) U^\dagger$, where $\mathbf{1}_{\lambda_j > 0}$ is 1 if $\lambda_j > 0$ and 0 otherwise.

Definition 9.11 (Invariant projection predicate). An MPS tensor A has an *invariant projection* if there exists an orthogonal projection P with $P \neq 0$, $P \neq \mathbb{1}$, and $(\mathbb{1} - P)A^i P = 0$ for all i . The negation of this predicate defines an *irreducible tensor* (Definition 9.14).

Theorem 9.12 (PSD fixed point gives invariant projection). *Let $\rho \geq 0$ be a fixed point of $\mathcal{E}_A$, and let P be the support projection of ρ . Then for all i , $(\mathbb{1} - P)A^i P = 0$: the range of P is invariant under all A^i .*

Proof. From $\mathcal{E}_A(\rho) = \rho$ and positivity of each summand $A^i \rho (A^i)^\dagger$, $(\mathbb{1} - P)A^i \rho (A^i)^\dagger (\mathbb{1} - P) = 0$ for each i . Since ρ is supported on the range of P , this forces $(\mathbb{1} - P)A^i P = 0$.

This is the finite-dimensional version of [Wol12, Prop. 6.10] (“Stationary subspaces”: the support projection of any stationary state is sub-harmonic, i.e. invariant under the channel). The equivalence between irreducibility and absence of nontrivial invariant projections is [Wol12, Thm. 6.2]; see also [CPGSV21, §IV.A.1]. $\square$

Theorem 9.13 (Two-block decomposition). *If the transfer map has a PSD fixed point ρ that is not positive definite (i.e. ρ has a nontrivial kernel), then A is unitarily equivalent to a two-block upper-triangular tensor. Moreover, the block-diagonal tensor obtained by discarding the off-diagonal blocks generates the same MPV family as A .*

Proof. The support projection P of ρ is neither 0 nor $\mathbb{1}$ (since $\rho \neq 0$ and ρ is not PD). Theorem 9.12 gives $(\mathbb{1} - P)A^i P = 0$ for all i . A unitary change of basis that block-diagonalizes P puts A^i in upper-triangular block form (Theorem 9.9 ensures the MPV is preserved by the conjugation). The trace of the upper-triangular product equals the sum of the diagonal-block traces, so the block-diagonal tensor obtained by discarding the off-diagonal blocks has the same MPV coefficients as the conjugated tensor, and hence as A . $\square$

9.4 Iterated reduction

Definition 9.14 (Irreducible tensor). An MPS tensor A is *irreducible* if its transfer map $\mathcal{E}_A$ is irreducible in the sense of Definition 4.12.

Theorem 9.15 (Irreducible block decomposition). *Every MPS tensor A can be decomposed (up to unitary equivalence) into a block-diagonal tensor $\bigoplus_{k=1}^r A_k$ where each A_k is irreducible. The block-diagonal tensor generates the same MPV as A .*

Proof. By strong induction on the bond dimension D . If A is already irreducible, take $r = 1$. Otherwise, by Definition 9.14, A has a nontrivial invariant projection P . Diagonalizing P yields the same two-block upper-triangular splitting used in Theorem 9.13; the block-diagonal tensor obtained by discarding the off-diagonal part has the same MPV family as A , and the two diagonal blocks have strictly smaller bond dimensions. Apply the induction hypothesis to each block and concatenate. (If one prefers a positive-eigenvector route without assuming TP normalization, the correct replacement for the channel-specific Cesàro theorem is Theorem 6.15; the induction itself needs only the invariant projection.)

Both [PGVWC07] (Theorem 4, “TI canonical form”) and [CPGSV21] (§IV.A.1) obtain the block decomposition by the same invariant-projection splitting used here. An alternative route, developed in [Wol12, Thm. 6.14], characterises the fixed-point set of a quantum channel as a $*$ -algebra and then applies the Wedderburn–Artin decomposition ([Wol12, Eq. (1.39)]) to read off the block structure directly. The block decomposition is obtained by iterating the invariant-projection splitting and using strong induction on D to ensure termination. $\square$

Theorem 9.16 (Unitary diagonal positive fixed point in TP gauge). *Let A be an irreducible MPS tensor satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and assume $D > 0$. Then there exist a unitary U and a diagonal positive definite matrix Λ such that, for $B^i := U^\dagger A^i U$,*

$$\sum_i (B^i)^\dagger B^i = \mathbb{1} \quad \text{and} \quad \mathcal{E}_B(\Lambda) = \Lambda.$$

Proof. The TP hypothesis makes $\mathcal{E}_A$ a channel, so it has a nonzero PSD fixed point. The irreducible tensor condition implies that $\mathcal{E}_A$ is irreducible as a CP map, and Theorem 6.3 upgrades the fixed point to a positive definite one. The spectral theorem diagonalizes this matrix by a unitary conjugation, and unitary conjugation preserves the TP normalization. $\square$

9.4.1 Burnside bridge: irreducibility and spanning

The irreducible-tensor predicate (Definition 9.14) is formulated in terms of invariant *projections*. For Burnside-style arguments it is more natural to work with invariant *subspaces* of $\mathbb{C}^D$ under the Kraus operators.

Definition 9.17 (Algebra span). For an MPS tensor A , let $\text{algSpan}(A)$ be the unital $\mathbb{C}$ -subalgebra of $M_D(\mathbb{C})$ generated by the matrices $\{A^i\}$:

$$\text{algSpan}(A) = \mathbb{C}\langle A^i : i = 0, \dots, d-1 \rangle \subseteq M_D(\mathbb{C}).$$

Definition 9.18 (Invariant submodule). A subspace $W \leq \mathbb{C}^D$ is *A-invariant* if $A^i W \subseteq W$ for every i .

Definition 9.19 (Irreducible action). The Kraus operators of A act *irreducibly* on $\mathbb{C}^D$ if every A -invariant subspace is trivial, i.e. equal to 0 or to $\mathbb{C}^D$.

Theorem 9.20 (Irreducible tensor implies irreducible action). *If A is irreducible as a tensor (Definition 9.14), then the Kraus operators act irreducibly on $\mathbb{C}^D$ (Definition 9.19).*

Proof. Let $W \leq \mathbb{C}^D$ be an A -invariant subspace. In finite dimension the orthogonal projection P onto W exists. Invariance of W implies $(\mathbb{1} - P)A^i P = 0$ for all i . Thus P is a nontrivial invariant projection unless $W = 0$ or $W = \mathbb{C}^D$, contradicting irreducibility of the tensor. $\square$

Theorem 9.21 (Burnside's theorem for Kraus algebras). *Assume $D > 0$. If A acts irreducibly on $\mathbb{C}^D$, then the generated unital subalgebra is the full matrix algebra:*

$$\text{algSpan}(A) = M_D(\mathbb{C}).$$

This is the complex finite-dimensional case of Burnside's theorem (equivalently, Jacobson's density theorem); see [Jac09].

Proof. Transport $\text{algSpan}(A)$ along the algebra equivalence $M_D(\mathbb{C}) \simeq_{\mathbb{C}} \text{End}_{\mathbb{C}}(\mathbb{C}^D)$. The irreducible-action hypothesis makes $\mathbb{C}^D$ a simple module for this algebra. Over the algebraically closed field $\mathbb{C}$, Schur's lemma identifies the endomorphisms commuting with the action with scalars, and Jacobson density then forces the action map to be surjective onto all of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$. $\square$

Remark 9.22. This Burnside/Jacobson-density step is a substantial external algebraic dependency for the formalization. In the current project it is formalized directly as the theorem tagged above, rather than imported from Mathlib; if Mathlib later acquires a suitable finite-dimensional Burnside theorem, this chapter could be retargeted to that library result.

Theorem 9.23 (Irreducible action implies eventual cumulative spanning). *Assume $D > 0$. If A acts irreducibly on $\mathbb{C}^D$, then there exists N such that $T_N(A) = M_D(\mathbb{C})$.*

Proof. By Theorem 9.21, $\text{algSpan}(A) = M_D(\mathbb{C})$. Every element of the algebra span is a linear combination of word products, hence belongs to $T_n(A)$ for some n . Since $M_D(\mathbb{C})$ is finite-dimensional, the ascending chain $T_0(A) \leq T_1(A) \leq \dots$ stabilizes, so one can choose a uniform N with $T_N(A) = M_D(\mathbb{C})$. $\square$

9.5 Proportional single-block theorem

Theorem 9.24 (Proportional MPV implies gauge-phase equivalence). *Let A and B be injective MPS tensors satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. If A and B generate proportional MPV families $(V^{(N)}(A))_\sigma = c_N V^{(N)}(B)_\sigma$ for some $c_N \in \mathbb{C}$ and $O_{AA}(N) \rightarrow 1$, $O_{BB}(N) \rightarrow 1$, then A and B are gauge-phase equivalent.*

Proof. By contradiction. If A and B are not gauge-phase equivalent, then $O_{AB}(N) \rightarrow 0$ (Theorem 7.17). From the proportionality hypothesis, $O_{AB}(N) = c_N O_{BB}(N)$. Since $O_{BB}(N) \rightarrow 1$, it follows that $c_N \rightarrow 0$. But $O_{AA}(N) = |c_N|^2 O_{BB}(N) \rightarrow 0$, contradicting $O_{AA}(N) \rightarrow 1$.

In [PGVWC07], the proportional case is handled as part of the canonical form argument. Thus the proportional case is obtained directly from the spectral gap in Chapter 7. $\square$

9.6 Canonical form from peripheral primitivity

The canonical form predicate (Definition 10.9) includes an explicit hypothesis that the self-overlap $O_{A_k A_k}(N) \rightarrow 1$ for each block. The following theorem derives this from per-block peripheral-spectrum primitivity, removing it as an independent assumption.

Theorem 9.25 (Canonical form from peripheral primitivity). *Let $(\mu_k, A_k)_{k=1}^r$ be a collection of block tensors and scaling factors satisfying:*

1. *each A_k is injective,*
2. *each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,*
3. *$|\mu_1| > \dots > |\mu_r|$ (strictly decreasing),*
4. *all $\mu_k \neq 0$,*
5. *each transfer map $\mathcal{E}_{A_k}$ is primitive, i.e. has peripheral spectrum $\{1\}$.*

Then (μ_k, A_k) satisfies the canonical form predicate (Definition 10.9).

Proof. Fix a block k and write $E := \mathcal{E}_{A_k}$. By injectivity and TP normalization, Theorem 6.9 provides a nonzero positive definite fixed point for E . To apply the spectral-gap criterion (Theorem 4.39), we check its three hypotheses:

1. E is primitive, since the peripheral spectrum is $\{1\}$ by assumption;
2. every eigenvalue of E has modulus at most 1, because $E = F_{A_k A_k}$ is a normalized mixed transfer operator and Theorem 7.10 applies;
3. any trace-zero fixed point of E is zero; this is the trace-zero uniqueness consequence extracted in Chapter 6 from the uniqueness of the PSD fixed point (Theorem 6.5).

Hence E has a spectral gap on the complement of its fixed-point projection. Theorem 7.21 then gives $O_{A_k A_k}(N) \rightarrow 1$. Since this holds for every block, the overlap clause required in Definition 10.9 follows from the other hypotheses. $\square$

Remark 9.26. Theorem 9.25 shows that the overlap convergence hypothesis in the canonical form predicate is redundant once peripheral-spectrum primitivity is known. The argument factors through the spectral-gap formulation of Chapter 7: peripheral-spectrum primitivity is first converted to that spectral-gap condition, and the overlap limit is then deduced from it. In the

paper [CPGSV21], primitivity is obtained from irreducibility plus periodicity removal by blocking to period one. Theorem 9.27 is precisely this periodicity-removal step under irreducibility and TP normalization.

Theorem 9.27 (Blocking yields a peripheral-spectrum primitive transfer map). *Let A be an irreducible MPS tensor with $D > 0$. Assume that A satisfies the TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then there exists a blocking length $p > 0$ such that the transfer map $\mathcal{E}_{A^{[p]}}$ of the blocked tensor $A^{[p]}$ is primitive, i.e. its peripheral spectrum is $\{1\}$.*

Proof. Pass to the conjugate-transposed Kraus family $K_i := (A^i)^\dagger$. The TP normalization of A makes K unital, and irreducibility of A gives irreducibility of the associated Kraus map. The TP fixed-point theorem for $\mathcal{E}_A$ provides a positive definite fixed point for the adjoint map of K , so Theorem 4.34 shows that every peripheral eigenvalue is a root of unity. Since $\text{peripheral}(\mathcal{E}_A)$ is finite (Theorem 4.26), Lemma 4.30 gives a common exponent p such that $\mu^p = 1$ for all $\mu \in \text{peripheral}(\mathcal{E}_A)$. Lemma 4.31 then implies $\text{peripheral}(\mathcal{E}_A^p) = \{1\}$. Finally, blocking corresponds to taking a power of the transfer map, so $\mathcal{E}_{A^{[p]}}$ is primitive. $\square$

Remark 9.28. Theorem 9.27 is the “periodicity removal by blocking” step in [CPGSV17, Appendix A] for an irreducible block already placed in TP gauge. The later existence theorems simply reuse this step.

9.7 Normal canonical form

Definition 9.29 (Normal canonical form predicate). A collection of scaling factors $(\mu_k)_{k=1}^r$ and block tensors $(A_k)_{k=1}^r$ satisfies the *normal canonical form predicate* if:

1. each A_k is irreducible,
2. each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. each transfer map $\mathcal{E}_{A_k}$ is primitive (equivalently, has peripheral spectrum $\{1\}$),
4. the moduli are strictly decreasing: $|\mu_1| > |\mu_2| > \dots > |\mu_r|$,
5. all $\mu_k \neq 0$,
6. all bond dimensions are positive.

Here “normal” is used in the spectral sense of [CPGSV17]: irreducible blocks whose TP-normalized transfer maps are primitive. As noted in the remark after Definition 2.19, this is equivalent to the eventual block-injectivity formulation of Definition 2.19; see [SPGWC10, Proposition 3].

Theorem 9.30 (Self-overlap is derived in normal canonical form). *If (μ_k, A_k) satisfies the normal canonical form predicate, then $O_{A_k A_k}(N) \rightarrow 1$ for every block k .*

Proof. Peripheral-spectrum primitivity is part of the data. Together with irreducibility and TP normalization, it yields the spectral-gap primitive theorem needed for overlap convergence, so the self-overlap condition follows from the other hypotheses rather than appearing as a separate assumption. $\square$

Theorem 9.31 (Modulus-one eigenvalue rigidity for irreducible TP blocks). *Let A and B be irreducible MPS tensors satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. If the mixed transfer map has spectral radius at least 1 (equivalently, it has a peripheral eigenvalue of modulus 1), then A and B are gauge-phase equivalent.*

Proof. The argument is the same as in Theorem 7.12, but the fixed-point input comes from irreducibility rather than injectivity. Take a modulus-one eigenvector of the mixed transfer map. Because A and B are TP-normalized, each transfer map has a nonzero PSD fixed point by Theorem 6.8. Irreducibility upgrades these fixed points to positive definite ones by Theorem 6.3. Theorem 6.10 therefore gauges A and B separately to unital Kraus families. In this unital gauge, the embedded modulus-one eigenvector saturates the Kadison–Schwarz inequality, and the same equality / multiplicative-domain argument as in Theorem 7.12 produces an invertible intertwiner between the two Kraus families. That intertwiner is exactly the required gauge-phase equivalence. $\square$

9.8 Steps toward canonical form existence

This section assembles the components of the canonical-form construction following [CPG17, §2.3 and Appendix A]. The approach here requires blocking to remove non-trivial periodicity. An alternative framework that handles periodic blocks without blocking is developed in [DCCSP17]; see Chapter 12 for a discussion of that extension.

Theorem 9.32 (Irreducible block decomposition). *Every MPS tensor A generates the same MPV family as a block-diagonal tensor $\bigoplus_{k=1}^r A_k$ with each block irreducible.*

Proof. Apply Theorem 9.15. $\square$

Theorem 9.33 (CFII data for irreducible TP blocks). *Let A be an irreducible MPS tensor already in TP gauge ($\sum_i (A^i)^\dagger A^i = \mathbb{1}$) with $D > 0$. Then there exist a unitary U and a diagonal positive definite matrix Λ such that:*

1. $B^i := U^\dagger A^i U$ is TP,
2. $\mathcal{E}_B(\Lambda) = \Lambda$,
3. Λ is diagonal and positive definite,
4. A and B generate the same MPV family.

This is the “Canonical Form II” (CFII) step of [CPG17, Appendix A]. The unitary conjugation occurs only after TP normalization has already been achieved; it is a second step inside TP gauge, not a unitary normalization of an arbitrary irreducible block.

Proof. Apply Theorem 9.16 to obtain the unitary and the diagonal PD fixed point. Unitary conjugation preserves trace-preservation and the MPV family (Theorem 9.9). $\square$

Theorem 9.34 (CFII continuation after irreducible block decomposition). *For any MPS tensor A , there exist irreducible blocks $(A_k)_{k=1}^r$ that generate the same MPV family as A . Moreover, for each block k , if that block has already been placed in TP gauge and $D_k > 0$, one obtains CFII data (unitary, diagonal PD fixed point, MPV preservation). The conclusion is CFII data; the normal-canonical-form packaging requires additional downstream steps.*

Proof. Combine Theorems 9.32 and 9.33. $\square$

Theorem 9.35 (Normal canonical form from a primitive block decomposition). *Suppose A generates the same MPV family as a weighted block-diagonal tensor $\bigoplus_k \mu_k A_k$, and assume that:*

1. each A_k is irreducible,

2. each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. each transfer map $\mathcal{E}_{A_k}$ is primitive (equivalently, has peripheral spectrum $\{1\}$),
4. the moduli $|\mu_k|$ are pairwise distinct,
5. all $\mu_k \neq 0$,
6. all bond dimensions are positive.

Then there exist a blocking length $p > 0$, weights (ν_k) , and blocked tensors (B_k) such that $A^{[p]}$ generates the same MPV family as $\bigoplus_k \nu_k B_k$, and (ν_k, B_k) satisfies the normal canonical form predicate.

Proof. No new blocking is needed here: the input blocks are already primitive, so one may take $p = 1$. The theorem is stated with a blocking length because earlier reduction steps may require a common blocking period; under the present hypotheses that period can be chosen to be 1. Since the moduli $|\mu_k|$ are pairwise distinct, there is a reordering of the blocks for which they become strictly decreasing. After this reindexing, set $\nu_k := \mu_k$ and $B_k := A_k$ in the new order. Irreducibility, TP normalization, primitivity, nonzero weights, and positive bond dimensions are unchanged by the reordering, so the reordered family satisfies Definition 9.29 and generates the same MPV family as A . $\square$

9.9 Zero-block separation and honest block decomposition

In the irreducible block decomposition, some blocks may have all-zero Kraus operators. Such blocks contribute to the MPV only at word length $N = 0$ (where their contribution equals their bond dimension). The following results separate the zero blocks from the “live” (nonzero) blocks.

Definition 9.36 (Zero MPS tensor). The *zero MPS tensor* of physical dimension d and bond dimension D , denoted $\mathbf{0}_{d,D}$, is the tensor whose Kraus operators are all zero.

Theorem 9.37 (MPV of zero tensor). For any spin configuration σ of length N ,

$$V^{(N)}(\mathbf{0}_{d,D})_\sigma = \begin{cases} D & \text{if } N = 0, \\ 0 & \text{if } N \geq 1. \end{cases}$$

Proof. For $N = 0$, $\text{evalWord}(\mathbf{0}, []) = \mathbb{1}_D$, so the MPV equals $\text{tr}(\mathbb{1}_D) = D$. For $N \geq 1$, every word product of a zero tensor is zero. $\square$

Theorem 9.38 (Zero-block separation). Every MPS tensor A admits a block decomposition into:

1. a zero tail of bond dimension D_0 (accumulating all all-zero irreducible blocks), and
2. a finite family of live blocks $(B_k)_{k=1}^r$, each irreducible, each with at least one nonzero Kraus operator, and each with positive bond dimension,

such that

$$V^{(N)}(A)_\sigma = V^{(N)}(\mathbf{0}_{d,D_0})_\sigma + V^{(N)}(\bigoplus_{k=1}^r B_k)_\sigma$$

for every spin configuration σ of length N . At $N = 0$ this reads $D = D_0 + \sum_{k=1}^r D_k$. For $N \geq 1$ the zero-tail term vanishes.

Proof. Apply the irreducible block decomposition (Theorem 9.15) and partition the blocks by whether each has a nonzero Kraus operator. The all-zero blocks satisfy a dimension bound via irreducibility; their total bond dimension is D_0 . The MPV identity follows from the block structure. $\square$

9.10 Arbitrary-input TP-gauge pipeline

Theorem 9.39 (Arbitrary-input TP-gauge pipeline with zero-block separation). *For any MPS tensor A , there exist:*

1. a zero-tail bond dimension D_0 ,
2. live blocks $(B_k)_{k=1}^r$ with nonzero weights $(\mu_k)_{k=1}^r$, each block irreducible and left-canonical $(\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1})$, with positive bond dimension,

such that

$$V^{(N)}(A)_\sigma = V^{(N)}(\mathbf{0}_{d,D_0})_\sigma + V^{(N)}(\bigoplus_{k=1}^r \mu_k B_k)_\sigma.$$

This is the furthest honest arbitrary-input pipeline step available before periodicity removal.

Proof. Apply Theorem 9.38 to separate the zero tail and live blocks. Then apply the blockwise TP-gauge theorem to the live blocks. The MPV identity chains through both steps. $\square$

9.11 Blocking infrastructure

Theorem 9.40 (Blocking distributes over MPV equivalence). *If A generates the same MPV family as $\bigoplus_k \mu_k B_k$, then for any $p \geq 0$,*

$$A^{[p]} \sim \bigoplus_k \mu_k^p B_k^{[p]}.$$

Proof. The identity $V^{(N)}(A^{[p]})_\sigma = V^{(Np)}(A)_{\sigma_{\text{flat}}}$ holds for any tensor by definition of blocking. The MPV equivalence of A and $\bigoplus_k \mu_k B_k$ at length Np then gives the equivalence at length N with weights μ_k^p . $\square$

Theorem 9.41 (Primitive channels remain primitive under powers). *If $\mathcal{E}$ has peripheral spectrum $\{1\}$ (is primitive) and $m \geq 1$, then $\mathcal{E}^m$ is also primitive.*

Proof. The peripheral spectrum of $\mathcal{E}^m$ is the set of m -th powers of that of $\mathcal{E}$. Since $\{1\}^m = \{1\}$, the result follows. $\square$

Theorem 9.42 (Blocking-period divisibility preserves primitivity). *If the transfer map of $A^{[p]}$ is primitive, $p \mid q$, and $q \geq 1$, then the transfer map of $A^{[q]}$ is also primitive.*

Proof. Write $q = p \cdot m$. Then $\mathcal{E}_{A^{[q]}} = \mathcal{E}_{A^{[p]}}^m$. Apply Theorem 9.41 with $\mathcal{E} = \mathcal{E}_{A^{[p]}}$. $\square$

Theorem 9.43 (Common blocking period for a block family). *Let $(B_k)_{k=1}^r$ be a finite family of MPS tensors with positive bond dimensions, each admitting a blocking period p_k making its transfer map primitive. Then $p = \text{lcm}(p_1, \dots, p_r)$ is a common period: $\mathcal{E}_{B_k^{[p]}}$ is primitive for every k .*

Proof. Each $p_k \mid p$, so Theorem 9.42 applies. $\square$

9.12 Cyclic sector decomposition

Theorem 9.44 (Compression to a supported projection sector). *Let A be a left-canonical MPS tensor with $\sum_i (A^i)^\dagger A^i = P$ for an orthogonal projection P , and suppose $PA^iP = A^i$ for every i . Then there exists a left-canonical tensor C of bond dimension $n = \text{tr}(P)$ with*

$$V^{(N)}(C)_\sigma = \text{tr}(P \cdot A^\sigma)$$

for every σ .

Proof. Diagonalize P via a unitary U (spectral theorem). The eigenvalues are in $\{0, 1\}$. Set $n = \text{tr} P$; let C^i be the top-left $n \times n$ block of $U^\dagger A^i U$ (restricted to eigenvalue-1 indices). TP follows from the projection normalization; the MPV identity follows from trace cyclicity. $\square$

Theorem 9.45 (Block decomposition from commuting projections). *Let A be a left-canonical MPS tensor, and let $(P_k)_{k=1}^m$ be orthogonal projections summing to $\mathbb{1}$ that commute with every Kraus operator: $P_k A^i = A^i P_k$ for all i, k . Then there exist left-canonical blocks $(C_k)_{k=1}^m$ with unit weights such that A generates the same MPV family as $\bigoplus_{k=1}^m C_k$.*

Proof. For each k , the tensor $P_k A^i$ is supported on P_k (using commutativity and $P_k^2 = P_k$). The TP condition $\sum_i (P_k A^i)^\dagger (P_k A^i) = P_k$ follows from $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Apply Theorem 9.44 to each sector. The MPV identity uses $\sum_k P_k = \mathbb{1}$ and linearity of trace. $\square$

Theorem 9.46 (Block decomposition from adjoint-fixed projections). *Let A be a left-canonical MPS tensor, and let $(P_k)_{k=1}^m$ be orthogonal projections summing to $\mathbb{1}$ that are fixed by the adjoint transfer map: $\mathcal{E}_A^\dagger(P_k) = P_k$ for every k . Then the same block decomposition conclusion holds as in Theorem 9.45.*

Proof. The Kadison–Schwarz equality condition for a TP map implies that a projection fixed by $\mathcal{E}_A^\dagger$ commutes with every Kraus operator. Apply Theorem 9.45. $\square$

9.13 Assembly: arbitrary tensor to primitive block decomposition

Theorem 9.47 (Arbitrary tensor to primitive block decomposition). *For any MPS tensor A , there exist a period $p \geq 1$, a zero-tail bond dimension D_0 , nonzero weights $(\mu_k)_{k=1}^r$, and left-canonical blocks $(B_k)_{k=1}^r$ with primitive transfer maps and positive bond dimensions, such that*

$$V^{(N)}(A^{[p]})_\sigma = V^{(N)}(\mathbf{0}_{d^{[p]}, D_0})_\sigma + V^{(N)}(\bigoplus_k \mu_k B_k)_\sigma.$$

Proof. Apply Theorem 9.39 to get the zero tail and left-canonical live blocks. Apply Theorem 9.43 to find a common blocking period p making all transfer maps primitive. Thread through via Theorem 9.40. $\square$

Theorem 9.48 (TP-primitive irreducible tensor is normal). *Let A be a left-canonical MPS tensor with $D > 0$, irreducible transfer map, and peripheral spectrum $\{1\}$. Then A is normal.*

Proof. Peripheral primitivity and irreducibility yield the spectral-gap condition. The PSD fixed point is positive definite by Theorem 6.3. Apply Theorem 8.52. $\square$

Theorem 9.49 (Blocking preserves normality). *If A is normal and $p \geq 1$, then $A^{[p]}$ is normal.*

Proof. Since A is normal, there exists N with $S_N(A) = M_D(\mathbb{C})$. Then $S_{Np}(A) = M_D(\mathbb{C})$ (word products at multiples still span), and $S_N(A^{[p]}) \supseteq S_{Np}(A) = M_D(\mathbb{C})$. $\square$

9.14 Open directions

Remark 9.50 (Canonical form pipeline: current status). The core pipeline steps are in place. The arbitrary-input pipeline (Theorem 9.39) produces a zero tail plus left-canonical live blocks. After a common blocking step (Theorem 9.47), the blocked blocks have primitive transfer maps.

The remaining steps to full normal canonical form require: establishing tensor irreducibility for each blocked block (blocking does not in general preserve tensor irreducibility), and establishing pairwise distinct weight norms. The cyclic-sector decomposition (Theorems 9.45 and 9.46) provides the infrastructure for the periodic-to-aperiodic splitting.

Chapter 10

Block Permutation and Separation

This chapter establishes the algebraic structure of automorphisms of products of matrix algebras and uses it to prove that the same-MPV condition on block-diagonal tensors descends to per-block gauge equivalence. The results correspond to the canonical form uniqueness of [PGVWC07] and the block separation results of [CPGSV21]. The proof technique used here—induction on blocks via the spectral gap—is specific to this formalization.

10.1 Block permutation algebra

Theorem 10.1 (Ring isomorphisms permute block ideals). *Let $\{R_k\}_{k=1}^r$ be simple rings. Every ring isomorphism $T : \prod_k R_k \rightarrow \prod_k R_k$ permutes the block ideals: there exists a permutation π such that T maps the k -th block ideal to the $\pi(k)$ -th block ideal.*

Proof. Each block ideal $I_k = \{x : x_j = 0 \text{ for } j \neq k\}$ is an atom in the lattice of two-sided ideals (since R_k is simple). A ring isomorphism preserves the ideal lattice and hence permutes the atoms. Injectivity of the induced map on atoms gives the permutation π . $\square$

Remark 10.2 (Ring level versus algebra level). Theorem 10.1 is a ring-level statement: it only uses the ideal structure of $\prod_k R_k$. Theorems 10.3 and 10.4 require an upgrade to $\mathbb{C}$ -algebra automorphisms. The reason is threefold: dimension preservation compares the blocks as $\mathbb{C}$ -vector spaces; Skolem–Noether applies to the resulting central simple $\mathbb{C}$ -algebras; and the per-block maps extracted from T must commute with scalar multiplication. Thus the formalization separates the ring-level permutation step from the later algebra-level decomposition.

Theorem 10.3 (Dimension preservation). *If T is an algebra automorphism of $\prod_{k=1}^r M_{D_k}(\mathbb{C})$, and π is the induced permutation, then $D_{\pi(k)} = D_k$ for all k .*

Proof. The restriction of T to the k -th block ideal induces a $\mathbb{C}$ -algebra isomorphism $M_{D_k}(\mathbb{C}) \cong M_{D_{\pi(k)}}(\mathbb{C})$, which forces $D_k = D_{\pi(k)}$ (by comparing dimensions as $\mathbb{C}$ -vector spaces). $\square$

Theorem 10.4 (Decomposition of automorphisms of $\prod M_{D_k}(\mathbb{C})$). *Every $\mathbb{C}$ -algebra automorphism T of $\prod_{k=1}^r M_{D_k}(\mathbb{C})$ decomposes as a block permutation π followed by per-block inner automorphisms: there exist $\pi \in S_r$ with $D_{\pi(k)} = D_k$ and invertible matrices $X_k \in \text{GL}_{D_k}(\mathbb{C})$ such*

that, for every tuple $M = (M_1, \dots, M_r)$, $T(M)_k = X_k M_{\pi(k)} X_k^{-1}$, where $T(M)_k$ denotes the k -th block component of $T(M)$.

Proof. Theorem 10.1 gives the permutation π . Theorem 10.3 gives $D_{\pi(k)} = D_k$. The restriction of T to the k -th block ideal yields a $\mathbb{C}$ -algebra isomorphism $M_{D_k}(\mathbb{C}) \cong M_{D_{\pi(k)}}(\mathbb{C})$; after Theorem 10.3 identifies the two matrix sizes, Skolem–Noether (Theorem 3.9) makes this restriction inner.

This is a standard result in the theory of semisimple algebras; see [CPGSV21, §IV.A] for the MPS context. $\square$

10.2 Product algebra linear extension

Definition 10.5 (Per-block linear extension). Given injective tensors A_k, B_k of bond dimension D_k with the same MPV for each block k , the *per-block linear extension* T_k is the unique linear map $M_{D_k}(\mathbb{C}) \rightarrow M_{D_k}(\mathbb{C})$ satisfying $T_k(A_k^i) = B_k^i$ for all i . It exists (and is unique) because $\{A_k^i\}$ spans $M_{D_k}(\mathbb{C})$ by injectivity.

Definition 10.6 (Product algebra automorphism). The *product algebra automorphism* is the blockwise assembled map $T : \prod_k M_{D_k}(\mathbb{C}) \rightarrow \prod_k M_{D_k}(\mathbb{C})$ defined by $T(M)_k := T_k(M_k)$. Theorem 10.7 shows that each T_k is a $\mathbb{C}$ -algebra homomorphism and that this assembled map is indeed a $\mathbb{C}$ -algebra automorphism.

Theorem 10.7 (Per-block linear extensions assemble). *Let A_k and B_k be injective tensors of bond dimension D_k for $k = 1, \dots, r$, and suppose the per-block same-MPV condition holds. Then the per-block linear extensions $T_k : A_k^i \mapsto B_k^i$ assemble into an algebra automorphism of $\prod_k M_{D_k}(\mathbb{C})$, which decomposes as a permutation π with $D_{\pi(k)} = D_k$ and per-block conjugation matrices X_k .*

Proof. Each T_k exists uniquely and is multiplicative by Theorems 3.6 and 3.7. Promotion to algebra homomorphisms (Lemma 3.10) and assembly into the product gives an algebra automorphism T . Theorem 10.4 decomposes T into a permutation plus per-block inner automorphisms. Because T is assembled blockwise, it preserves each block ideal, so in this theorem the permutation is actually $\pi = \text{id}$. A nontrivial block permutation only appears after block separation identifies which blocks match globally. $\square$

Theorem 10.8 (Per-block same MPV gives global gauge equivalence). *If all block tensors A_k are injective and per-block same-MPV holds, then each pair (A_k, B_k) is gauge equivalent, and the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ are globally gauge equivalent.*

Proof. Per-block gauge equivalence follows from the single-block FT (Theorem 3.11) applied to each block. Global gauge equivalence follows by assembling the per-block gauge matrices into a block-diagonal matrix (Theorem 9.4). $\square$

10.3 Block separation

Recall from Definition 2.31 and Theorem 7.7 that the bilinear overlap is

$$O_{AB}(N) = \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma} = \text{Tr}(F_{AB}^N).$$

We use this notation throughout the block-separation argument.

Definition 10.9 (Canonical form predicate). A collection of scaling factors $(\mu_k)_{k=1}^r$ and block tensors $(A_k)_{k=1}^r$ satisfies the *canonical form predicate* if:

1. each A_k is injective,
2. each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. the moduli are strictly decreasing: $|\mu_1| > |\mu_2| > \dots > |\mu_r|$,
4. all $\mu_k \neq 0$,
5. the self-overlap converges: $O_{A_k A_k}(N) \rightarrow 1$ for each k .

Only this one-sided normalization is part of the predicate; no unital condition is assumed here. Condition (5) is a primitivity hypothesis: for a primitive channel ([Wol12, Thm. 6.7, item 3]), $T^n(\rho) \rightarrow \rho_\infty$ for every initial state, which forces the self-overlap to converge to 1. See also [Wol12, Thm. 6.8] for the completely-positive characterisation and Chapter 7 for the spectral-gap proof.

Remark 10.10 (Normalization convention). The convention here differs from [PGVWC07, Thm. 4], which uses the unital gauge $\sum_i A_k^i (A_k^i)^\dagger = \mathbb{1}$. This chapter instead works in the dual TP gauge $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$. A positive-definite fixed point of the adjoint transfer map provides a gauge transformation between these two normalizations (Theorem 6.11).

Remark 10.11. In the literature [CPGSV21], condition (5) is derived from primitivity of the transfer map (peripheral spectrum = $\{1\}$). Theorem 9.25 gives this implication: per-block primitivity implies overlap convergence, so condition (5) is redundant when primitivity holds. The canonical form predicate retains condition (5) explicitly because it can be verified independently of primitivity.

Remark 10.12 (Normal-canonical companion). Definition 10.9 is the block-injective predicate. The normal-canonical route instead uses Definition 9.29: injectivity and the explicit self-overlap hypothesis are replaced by irreducibility, TP normalization, and primitive transfer maps. Theorem 9.30 then supplies the self-overlap limit automatically.

Theorem 10.13 (Vandermonde separation). *If $\mu_1, \dots, \mu_r$ are pairwise distinct complex numbers and $\sum_{k=1}^r c_k \mu_k^N = 0$ for $N = 0, 1, \dots, r-1$, then $c_k = 0$ for all k .*

Proof. The coefficient vector c satisfies $Vc = 0$ where V is the Vandermonde matrix of the μ_k . Injectivity of μ makes V invertible. $\square$

Lemma 10.14 (Block separation core). *Let $((\mu_k), (A_k))$ satisfy the canonical form predicate, and let (B_k) be injective tensors satisfying the same TP normalization $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$. If $\sum_k \mu_k^N (V^{(N)}(A_k)_\sigma - V^{(N)}(B_k)_\sigma) = 0$ for all N, σ , then each pair (A_k, B_k) generates the same MPV family.*

Proof. By induction on the number of blocks r . In the inductive step, take overlaps with the leading block A_0 ; after division by μ_0^N , the contributions from $k \geq 1$ decay exponentially by Theorems 7.15 and 7.17, leaving $O_{A_0 A_0}(N) - O_{A_0 B_0}(N) \rightarrow 0$. Condition (5) gives $O_{A_0 A_0}(N) \rightarrow 1$, hence $O_{A_0 B_0}(N) \rightarrow 1$. Since this limit is nonzero, the contrapositive of Theorem 7.17 forces A_0 and B_0 to be gauge-phase equivalent; the limit $O_{A_0 B_0}(N) \rightarrow 1$ then makes the phase trivial, so A_0 and B_0 generate the same MPV family. Peeling off the leading block and dividing by μ_0^N , the induction hypothesis applies. $\square$

Theorem 10.15 (Block separation from canonical form). *Let $((\mu_k), (A_k))$ satisfy the canonical form predicate, and let (B_k) be injective tensors satisfying the same TP normalization $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$. If the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family, then each pair (A_k, B_k) generates the same MPV family.*

Proof. Apply Lemma 10.14 to the identity obtained from the equality of the two block-diagonal MPV families. The canonical form predicate supplies injectivity, TP normalization, strict nonzero weights, and self-overlap convergence for (A_k) , while the hypotheses on (B_k) supply the comparison injectivity and TP normalization. $\square$

Theorem 10.16 (Block separation from normal canonical form). *Let $((\mu_k), (A_k))$ satisfy the normal canonical form predicate, and let (B_k) be irreducible tensors of the same bond dimensions, satisfying the same TP normalization $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$. If the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family, then each pair (A_k, B_k) generates the same MPV family.*

Proof. The proof follows the same peeling argument as Theorem 10.15. Theorem 9.30 replaces the explicit self-overlap hypothesis on the leading block, and Theorem 9.31 identifies the leading comparison block from the resulting nondecaying mixed overlap. Since both block families already use the same dimension function, the conclusion is direct per-block same MPV. $\square$

Theorem 10.17 (Canonical form fundamental theorem, same-structure version). *Under the hypotheses of Theorem 10.15, each pair (A_k, B_k) is gauge equivalent. Hence the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ are gauge equivalent.*

Proof. Theorem 10.15 gives per-block same MPV. Theorem 10.8 converts this to per-block gauge equivalence and then to global gauge equivalence of the block-diagonal tensors. $\square$

Remark 10.18 (Relation to the literature). The results of this chapter correspond to the canonical-form uniqueness of [PGVWC07] and the block-separation statements of [CPGSV21]. The proof technique used here—an induction on the number of blocks driven by the spectral gap of the mixed transfer operators—is specific to this formalization.

Chapter 11

Bases of Normal Tensors

This chapter introduces *bases of normal tensors* (BNT), the canonical-form variants with BNT separation, the cross-overlap decay theorem, and the permutation-rigidity arguments used in the final assembly. This chapter also contains the auxiliary results on coefficient convergence and the Newton–Girard identities. The main references are [PGVWC07] and [CPGSV21].

11.1 Bases of normal tensors

Definition 11.1 (Basis of Normal Tensors (BNT)). A *basis of normal tensors* (BNT) for a total tensor A_{tot} consists of g block tensors $(A_1, \dots, A_g)$ with bond dimensions $(D_1, \dots, D_g)$ such that:

1. each A_j is normal (Definition 2.19),
2. at each system size N , the MPV of A_{tot} lies in the span of the block MPVs $\{|V^{(N)}(A_j)\}_{j=1}^g$,
3. for sufficiently large N , the block MPV vectors $\{|V^{(N)}(A_j)\}_{j=1}^g$ are linearly independent.

This is [CPGSV21, Definition 4.2].

Definition 11.2 (Canonical form with BNT separation). A family is in *canonical form with BNT separation* if it satisfies Definition 10.9 and, in addition, distinct blocks of the same bond dimension are *not* gauge-phase equivalent.

Definition 11.3 (Normal canonical form with BNT separation). A family is in *normal canonical form with BNT separation* if it satisfies Definition 9.29 and, in addition, distinct blocks of the same bond dimension are not gauge-phase equivalent.

Theorem 11.4 (Cross-overlap decay for distinct blocks in canonical form with BNT separation). *If $((\mu_k), (A_k))$ is in canonical form with BNT separation and $j \neq k$, then $O_{A_j A_k}(N) \rightarrow 0$.*

Proof. Definition 11.2 supplies injectivity and TP normalization for each block, which are the hypotheses required by the spectral-gap theorems. If $D_j \neq D_k$, the rectangular overlap-decay theorem (Theorem 7.15) gives $O_{A_j A_k}(N) \rightarrow 0$. If $D_j = D_k$, the separation condition says that A_j and A_k are not gauge-phase equivalent, so the same-dimension overlap-decay theorem (Theorem 7.17) again gives decay to 0. $\square$

Theorem 11.5 (Canonical form with BNT separation yields a basis of normal tensors). *If $((\mu_k), (A_k))$ is in canonical form with BNT separation, then the block tensors form a basis of normal tensors for the block-diagonal tensor $\bigoplus_k \mu_k A_k$.*

Proof. Each block is injective, so it is normal by taking blocking length 1. The MPV decomposition (Theorem 2.24) gives

$$|V^{(N)}(A_{\text{tot}})\rangle = \sum_k \mu_k^N |V^{(N)}(A_k)\rangle,$$

so the MPV of A_{tot} lies in the span of the block MPVs at each system size. For eventual linear independence, consider the Gram matrix

$$G_{jk}(N) = \langle V^{(N)}(A_j) | V^{(N)}(A_k) \rangle.$$

By Lemma 2.32, $G_{jk}(N) = \overline{O_{A_j A_k}(N)}$, so the same convergence statements hold after conjugation: the diagonal entries tend to 1 by the self-overlap part of the canonical-form hypothesis, and the off-diagonal entries tend to 0 by Theorem 11.4. Thus $G(N) \rightarrow I$, hence $G(N)$ is eventually invertible. Therefore the vectors $\{|V^{(N)}(A_j)\rangle\}_{j=1}^g$ are linearly independent for all sufficiently large N . $\square$

Remark 11.6 (The normality gap for normal canonical form with BNT separation). Definition 11.3 uses “normal” in the normal-canonical sense, not the eventual block-injective predicate of Definition 2.19. Therefore data satisfying Definition 11.3 do not by themselves yield a basis of normal tensors in the sense of Definition 11.1; one still needs eventual block injectivity for each block. This gap is closable: primitivity of the transfer map implies eventual full Kraus rank by [SPGWC10, Proposition 3], and the bridge from a primitive transfer map to eventual block injectivity is available via the Wielandt theory of Chapter 8. It is simply not built into Definition 11.3 itself.

11.2 Permutation rigidity for bases of normal tensors

Theorem 11.7 (Permutation rigidity for proportional MPV decompositions). *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be injective families satisfying the TP normalization $\sum_i (A_j^i)^\dagger A_j^i = \mathbb{1}$ and $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$, whose self-overlaps converge to 1 and whose cross-overlaps within each family converge to 0. Assume moreover that there exist MPS tensors A_{tot} and B_{tot} , coefficient arrays $a_{N,j}$ and $b_{N,k}$, and a sequence c_N with nonzero limits a_j^∞ , b_k^∞ , c^∞ such that for every N ,*

$$|V^{(N)}(A_{\text{tot}})\rangle = \sum_{j=1}^{g_A} a_{N,j} |V^{(N)}(A_j)\rangle, \quad |V^{(N)}(B_{\text{tot}})\rangle = \sum_{k=1}^{g_B} b_{N,k} |V^{(N)}(B_k)\rangle,$$

with $a_{N,j} \rightarrow a_j^\infty \neq 0$ and $b_{N,k} \rightarrow b_k^\infty \neq 0$, and also

$$|V^{(N)}(A_{\text{tot}})\rangle = c_N |V^{(N)}(B_{\text{tot}})\rangle$$

with $c_N \rightarrow c^\infty \neq 0$. Then $g_A = g_B$, and there is a permutation π of the common block index set such that $A_{\pi(k)}$ and B_k have the same bond dimension and are gauge-phase equivalent for each k .

Proof. Substitute the two decomposition formulas into the proportionality relation and take overlaps with the block MPVs. Passing to the limit using the convergence of $a_{N,j}$, $b_{N,k}$, and c_N produces a limiting mixed-overlap matrix. The asymptotically orthonormal overlap hypotheses within each family force every row and column of that matrix to contain at most one nonzero entry, hence determine a permutation π . If one of the matched mixed overlaps came from blocks of different bond dimensions, Theorem 7.15 would force it to decay to 0; if the bond dimensions

agree but the tensors are not gauge-phase equivalent, Theorem 7.17 does the same. Therefore every nonzero matched entry comes from blocks of the same bond dimension that are gauge-phase equivalent.

This is the overlap-and-decomposition form of the permutation step corresponding to [CPGSV21, Theorem 4.1]. $\square$

Remark 11.8. In the canonical-form application, the abstract data above come from block-diagonal assemblies

$$A_{\text{tot}} = \bigoplus_{j=1}^{g_A} \mu_j A_j, \quad B_{\text{tot}} = \bigoplus_{k=1}^{g_B} \nu_k B_k,$$

and Theorem 2.24 gives the coefficient arrays $a_{N,j} = \mu_j^N$ and $b_{N,k} = \nu_k^N$. Definition 11.2 supplies injectivity, TP normalization, and the within-family overlap limits, so the theorem isolates the abstract permutation argument from this canonical-form packaging.

Theorem 11.9 (Permutation rigidity with asymptotically orthonormal overlaps). *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be two families of injective tensors satisfying the TP normalization $\sum_i (A_j^i)^\dagger A_j^i = \mathbb{1}$ and $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$, whose self-overlaps converge to 1 and whose cross-overlaps within each family converge to 0. If for every system size N ,*

$$\text{span}_{\mathbb{C}}\{|V^{(N)}(A_j)\rangle : 1 \leq j \leq g_A\} = \text{span}_{\mathbb{C}}\{|V^{(N)}(B_k)\rangle : 1 \leq k \leq g_B\},$$

then $g_A = g_B$, and there is a permutation π of the common block index set such that A_j and $B_{\pi(j)}$ have the same bond dimension and are gauge-phase equivalent for each j .

Proof. For each family, the Gram matrix converges to the identity by Lemma 2.32 together with the self-overlap and off-diagonal decay hypotheses. Hence for all sufficiently large N both families of MPV vectors are linearly independent. At such an N , the equality of spans in the statement forces the two common subspaces to have the same dimension, so $g_A = g_B$. The span equality then gives change-of-basis matrices between the two families, and the same mixed-overlap matching argument shows that the mixed-overlap matrix has exactly one nonzero entry in each row and column. This produces the permutation π , while Theorems 7.15 and 7.17 upgrade each nonzero match to equality of bond dimensions and gauge-phase equivalence. $\square$

11.3 Coefficient convergence

Theorem 11.10 (Coefficient ratio decay). *Under the canonical form predicate of Definition 10.9, indexed so that μ_0 is the dominant weight, one has $(\mu_k/\mu_0)^N \rightarrow 0$ for every $k \geq 1$ as $N \rightarrow \infty$.*

Proof. Since $|\mu_k| < |\mu_0|$ (strict ordering of moduli), $|\mu_k/\mu_0| < 1$, so the geometric sequence converges to 0. $\square$

Remark 11.11 (Strict-dominance regime). Theorem 11.10 applies when one modulus is strictly dominant: $|\mu_0| > |\mu_k|$ for $k \geq 1$. In the more general periodic setting of [CPGSV21, Eq. (72a)], the coefficient attached to block j can take the form $\sum_q \mu_{j,q}^N$, where the $\mu_{j,q}$ are the individual weights contributed by that block. The oscillatory case with $|\mu_{j,q}| = 1$ is treated separately in the full paper.

11.4 Newton–Girard identities

Theorem 11.12 (Newton–Girard trace recursion). *If $A, B \in M_n(\mathbb{C})$ satisfy $\text{tr}(A^k) = \text{tr}(B^k)$ for all $k \geq 1$, then A and B have the same characteristic polynomial.*

Proof. Let e_m denote the m th elementary symmetric polynomial in the eigenvalues, with $e_0 = 1$. The Newton–Girard recursion

$$me_m = \sum_{i=1}^m (-1)^{i-1} e_{m-i} \text{tr}(A^i)$$

expresses the coefficients of the characteristic polynomial in terms of the power-sum traces $\text{tr}(A^i)$. Equal power sums for A and B therefore give equal coefficients, hence equal characteristic polynomials. $\square$

Theorem 11.13 (Equal power sums imply equal multisets). *Let $\alpha, \beta : \{1, \dots, n\} \rightarrow \mathbb{C}$. If $\sum_i \alpha_i^k = \sum_i \beta_i^k$ for all $k \geq 1$, then the multisets $\{\alpha_i\}$ and $\{\beta_i\}$ are equal.*

Proof. Construct diagonal matrices $\text{diag}(\alpha)$ and $\text{diag}(\beta)$. The power-sum hypothesis gives $\text{tr}(\text{diag}(\alpha)^k) = \text{tr}(\text{diag}(\beta)^k)$. By Theorem 11.12, the characteristic polynomials agree: $\prod_i (X - \alpha_i) = \prod_i (X - \beta_i)$. Extracting roots gives multiset equality. $\square$

Remark 11.14. Theorems 11.12 and 11.13 are auxiliary results retained for potential future use. The current route to Theorem 12.5 uses linear independence of the block MPV family supplied by a basis of normal tensors, rather than Newton–Girard identities, to match the weight multiset.

Chapter 12

Full Assembly

This chapter proves the Fundamental Theorem of Matrix Product States. We state the Fundamental Theorem in two versions: one for block-injective tensors in canonical form with basis of normal tensors (BNT) separation (CF-BNT), and one for the normal-canonical variant. We then treat the equal-MPV case, including the common-block-structure specialization that deduces gauge equivalence. The results combine the canonical form reduction (Chapter 9), block separation (Chapter 10), spectral gap (Chapter 7), and the BNT permutation arguments of Chapter 11.

The theorems in Sections 12.1–12.3 follow the approach of [CPGSV17]: they first block the tensor to remove non-trivial periodicity, then apply the aperiodic fundamental theorem. Section 12.4 outlines the stronger “irreducible form” framework of [DICCSPG17], which gives a fundamental theorem valid for periodic tensors directly, with an extra Z -gauge degree of freedom.

12.1 Weak Fundamental Theorem: pipeline end-to-end

The following theorem packages the full upstream pipeline (zero-block separation, TP gauge, blocking, primitivity) with the downstream Fundamental Theorem for normal canonical form data. It is conditional on the extra hypotheses (tensor irreducibility and distinct weight norms) that the automatic pipeline does not yet produce.

Theorem 12.1 (Weak Fundamental Theorem (conditional)). *Let $(\mu_j^A, A_j)_{j=1}^{g_A}$ and $(\mu_k^B, B_k)_{k=1}^{g_B}$ be two families in normal canonical form, each satisfying the non-degeneracy condition that no two distinct blocks in the same family are gauge-phase equivalent. Let A_{tot} and B_{tot} be tensors with MPV decompositions*

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_j a_{N,j} V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_k b_{N,k} V^{(N)}(B_k)_\sigma,$$

with $a_{N,j} \rightarrow a_j \neq 0$ and $b_{N,k} \rightarrow b_k \neq 0$. Suppose moreover that $V^{(N)}(A_{\text{tot}})_\sigma = c_N V^{(N)}(B_{\text{tot}})_\sigma$ with $c_N \rightarrow c \neq 0$. Then $g_A = g_B$ and there is a permutation π such that $D_j^A = D_{\pi(j)}^B$ and A_j is gauge-phase equivalent to $B_{\pi(j)}$ for every j .

Proof. This packages the normal-canonical-form Fundamental Theorem (Theorem 12.4). The normal canonical form data provide the BNT separation and overlap convergence hypotheses; the non-degeneracy condition ensures the block matching is well-defined. $\square$

12.2 Proportional Fundamental Theorem

Theorem 12.2 (Fundamental Theorem — proportional MPV case). *Let A_{tot} and B_{tot} be MPS tensors in canonical form with BNT separation, with associated families $(\mu_j^A, A_j)_{j=1}^{g_A}$ and $(\mu_k^B, B_k)_{k=1}^{g_B}$. Suppose there exist coefficient arrays $a_{N,j}$, $b_{N,k}$ and nonzero limits a_j^∞ , b_k^∞ , c^∞ such that*

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^{g_A} a_{N,j} V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_{k=1}^{g_B} b_{N,k} V^{(N)}(B_k)_\sigma,$$

with $a_{N,j} \rightarrow a_j^\infty \neq 0$ and $b_{N,k} \rightarrow b_k^\infty \neq 0$, and suppose also that

$$V^{(N)}(A_{\text{tot}})_\sigma = c_N V^{(N)}(B_{\text{tot}})_\sigma$$

for a sequence $c_N \rightarrow c^\infty \neq 0$. Then $g_A = g_B$, and there is a permutation π such that $D_j^A = D_{\pi(j)}^B$ and A_j is gauge-phase equivalent to $B_{\pi(j)}$ for each j .

Proof. Apply Theorem 11.7 to the two CF-BNT decompositions. The CF-BNT hypotheses provide injectivity, TP normalization, diagonal overlap limits, and off-diagonal decay via Theorem 11.4. With the supplied decomposition identities and convergence hypotheses, Theorem 11.7 yields the equality of block counts, the permutation, and the per-block gauge-phase equivalence. $\square$

Remark 12.3 (Coefficient arrays in the CF-BNT setting). In applications the coefficient arrays in Theorem 12.2 come from the block-diagonal decomposition itself. If

$$A_{\text{tot}} = \bigoplus_{j=1}^{g_A} \mu_j^A A_j, \quad B_{\text{tot}} = \bigoplus_{k=1}^{g_B} \mu_k^B B_k,$$

then Theorem 2.24 gives

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^{g_A} (\mu_j^A)^N V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_{k=1}^{g_B} (\mu_k^B)^N V^{(N)}(B_k)_\sigma.$$

Thus one takes $a_{N,j} = (\mu_j^A)^N$ and $b_{N,k} = (\mu_k^B)^N$, while the sequence c_N is the global proportionality factor from the hypothesis $V^{(N)}(A_{\text{tot}}) = c_N V^{(N)}(B_{\text{tot}})$. Theorem 12.2 keeps these arrays explicit because Theorem 11.7 is formulated in that abstract form.

Theorem 12.4 (Fundamental Theorem — proportional MPV case, NT route). *Let A_{tot} and B_{tot} be MPS tensors in normal canonical form with BNT separation, with associated families $(\mu_j^A, A_j)_{j=1}^{g_A}$ and $(\mu_k^B, B_k)_{k=1}^{g_B}$. Suppose there exist coefficient arrays $a_{N,j}$, $b_{N,k}$ and nonzero limits a_j^∞ , b_k^∞ , c^∞ such that*

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^{g_A} a_{N,j} V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_{k=1}^{g_B} b_{N,k} V^{(N)}(B_k)_\sigma,$$

with $a_{N,j} \rightarrow a_j^\infty \neq 0$ and $b_{N,k} \rightarrow b_k^\infty \neq 0$, and suppose also that

$$V^{(N)}(A_{\text{tot}})_\sigma = c_N V^{(N)}(B_{\text{tot}})_\sigma$$

for a sequence $c_N \rightarrow c^\infty \neq 0$. Then $g_A = g_B$, and there is a permutation π such that $D_j^A = D_{\pi(j)}^B$ and A_j is gauge-phase equivalent to $B_{\pi(j)}$ for each j .

Proof. Normal canonical form with BNT separation provides irreducibility, TP normalization, and primitive self-overlap convergence via Theorem 9.30. Theorem 9.31 supplies the leading-block identification step: a nondecaying mixed overlap forces the corresponding comparison block to be gauge-phase equivalent. The same peeling argument as in the injective CF-BNT case therefore yields the block permutation and gauge-phase equivalence. Remark 11.6 explains why the argument stops at this separated normal-canonical level. $\square$

12.3 Equal-MPV Fundamental Theorem

Theorem 12.5 (Fundamental Theorem — equal MPV case). *Let A_{tot} and B_{tot} be as in Theorem 12.2, with associated families $(\mu_j^A, A_j)_{j=1}^{g_A}$ and $(\mu_k^B, B_k)_{k=1}^{g_B}$. If*

$$V^{(N)}(A_{\text{tot}})_\sigma = V^{(N)}(B_{\text{tot}})_\sigma$$

for every N and σ , then the block-diagonal tensors $\bigoplus_{j=1}^{g_A} \mu_j^A A_j$ and $\bigoplus_{k=1}^{g_B} \mu_k^B B_k$ are gauge equivalent.

Proof. Apply Theorem 12.2 with $c_N = 1$. Then $g_A = g_B$, and after reindexing the B -blocks by a permutation π there are phases $e^{i\phi_j}$ and invertible matrices X_j such that

$$B_{\pi(j)}^i = e^{i\phi_j} X_j A_j^i X_j^{-1}$$

for every block j and physical index i . By Theorem 2.24,

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^{g_A} (\mu_j^A)^N V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_{j=1}^{g_A} (\mu_{\pi(j)}^B)^N V^{(N)}(B_{\pi(j)})_\sigma.$$

The gauge-phase identity gives

$$V^{(N)}(B_{\pi(j)})_\sigma = e^{iN\phi_j} V^{(N)}(A_j)_\sigma,$$

so the equality of MPVs becomes

$$\sum_{j=1}^{g_A} \left((\mu_j^A)^N - (\mu_{\pi(j)}^B e^{i\phi_j})^N \right) V^{(N)}(A_j)_\sigma = 0.$$

By Theorem 11.5 and Definition 11.1, the block MPVs $\{V^{(N)}(A_j)\}_{j=1}^{g_A}$ are linearly independent for all sufficiently large N . Hence, for each j and all large N ,

$$(\mu_j^A)^N = (\mu_{\pi(j)}^B e^{i\phi_j})^N.$$

Taking two consecutive large values of N and using that the weights are nonzero gives

$$\mu_j^A = \mu_{\pi(j)}^B e^{i\phi_j}.$$

Therefore

$$\mu_{\pi(j)}^B B_{\pi(j)}^i = \mu_j^A X_j A_j^i X_j^{-1}$$

for every j and i . With $X = \bigoplus_j X_j$, the reindexed block-diagonal tensors satisfy

$$\bigoplus_j \mu_{\pi(j)}^B B_{\pi(j)}^i = X \left(\bigoplus_j \mu_j^A A_j^i \right) X^{-1}.$$

Composing with the permutation matrix of π gives gauge equivalence of the original block-diagonal tensors. $\square$

Theorem 12.6 (Fundamental Theorem — equal MPV case, common block structure). *Fix a common block structure: the same number of blocks g , the same bond dimensions $(D_k)_{k=1}^g$, and the same weights $(\mu_k)_{k=1}^g$. Let $(\mu_k, A_k)_{k=1}^g$ and $(\mu_k, B_k)_{k=1}^g$ be two families in canonical form with BNT separation on this common data. If the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family, then each pair (A_k, B_k) is gauge equivalent, and hence the assembled block-diagonal tensors are gauge equivalent.*

Proof. Since both families share the canonical form data, the BNT separation condition is not needed. The canonical-form fundamental theorem (Theorem 10.17) then applies directly to this common (g, D_k, μ_k) structure, yielding per-block gauge equivalence and hence global gauge equivalence. $\square$

Remark 12.7 (Recovery of the literature statement). Theorem 12.6 is the common-block-structure specialization. The proof of Theorem 12.5 combines Theorem 12.2 with the BNT linear independence furnished by Theorem 11.5 and Definition 11.1, and therefore recovers the full [CPGSV21, Corollary IV.5] without assuming common block structure. The weight matching is obtained directly from the BNT expansion, rather than from Newton–Girard identities.

12.4 Periodic Fundamental Theorem (de las Cuevas–Cirac–Schuch–Pérez-García)

The theorems in the preceding sections all require that each block has a *primitive* transfer map, i.e. peripheral spectrum exactly $\{1\}$. This is achieved by blocking the original tensor by a common period. The paper [DICCSPG17] develops a framework that avoids this blocking step, giving a fundamental theorem that applies directly to periodic tensors.

12.4.1 Irreducible form

Definition 12.8 (Periodic tensor). An irreducible MPS tensor A of bond dimension D is *m-periodic* (for some $m \geq 1$) if the peripheral spectrum of its transfer map $\mathcal{E}_A$ is exactly the m -th roots of unity $\{e^{2\pi i k/m} : 0 \leq k < m\}$. A 1-periodic tensor is exactly an irreducible tensor with primitive transfer map (Definition 4.37).

Definition 12.9 (Irreducible form). A tensor A is in *irreducible form* if it is block-diagonal

$$A^i = \bigoplus_{j \in J} \mu_j A_j^i,$$

where each block A_j is an m_j -periodic tensor, each weight $\mu_j \neq 0$, and the blocks are pairwise non-repeated (no two distinct j, j' have A_j gauge-equivalent to $A_{j'}$). This is the standard form introduced in [DICCSPG17, §II]. It extends the normal canonical form (Definition 9.29), which requires the stronger condition that every block is 1-periodic.

Remark 12.10 (Relation to normal canonical form). Every tensor in normal canonical form (Definition 9.29) is in irreducible form with all periods equal to 1. The irreducible form is strictly more general: it accommodates blocks such as the antiferromagnet, which are 2-periodic.

The current formalization (Chapters 9–12) works exclusively with period-1 blocks, obtaining them by a blocking step. Extending the formalization to the irreducible form of [DICCSPG17] would require:

1. a definition of m -periodic irreducible tensors and their cyclic-sector decomposition (partial infrastructure exists in Section 9.12 of Chapter 9),
2. the off-diagonal structure of a periodic block's transfer matrix (the cyclic permutation of sectors), and
3. the notion of equivalence between periodic blocks that accounts for the Z -gauge freedom described below.

12.4.2 The Z -gauge degree of freedom

Definition 12.11 (Z -gauge equivalence). Let A be an m -periodic irreducible tensor of bond dimension D . A Z -gauge transformation of A is a diagonal unitary matrix $Z \in M_D(\mathbb{C})$ satisfying:

1. Z commutes with every Kraus operator: $[A^i, Z] = 0$ for all physical indices i , and
2. $Z^m = \mathbb{1}_D$ (all diagonal entries are m -th roots of unity).

Replacing A by ZA (i.e. $A^i \mapsto ZA^i$) leaves the MPV family invariant for lengths N that are multiples of m , because the factor $\text{tr}(Z^N) = \text{tr}(\mathbb{1}) = D$ for $N \equiv 0 \pmod{m}$. Two tensors A, B are Z -gauge equivalent if there exists an invertible Y and a Z -gauge transformation Z for A such that $ZA^i = YB^iY^{-1}$ for all i .

Remark 12.12 (Z -gauge is trivial for period-1 blocks). When $m = 1$ the condition $Z^1 = \mathbb{1}$ forces $Z = \mathbb{1}$, so Z -gauge equivalence reduces to ordinary gauge equivalence (Definition 2.7). The Z -gauge is therefore a genuine new degree of freedom that appears only for $m > 1$.

12.4.3 Statement of the periodic Fundamental Theorem

The following is Theorem 3.8 (“equal case”) of [DICCSPG17].

Theorem 12.13 (Periodic Fundamental Theorem). *Let A and B be MPS tensors in irreducible form, with bases of periodic blocks $\{A_j\}_{j \in J}$ and $\{B_k\}_{k \in K}$ respectively. Suppose that $\mathcal{V}(A) = \mathcal{V}(B)$ (equal MPV families for all lengths N).*

Then $|J| = |K|$ and there is a bijection $\pi : J \rightarrow K$ such that A_j and $B_{\pi(j)}$ are m_j -periodic for the same period m_j . Moreover, for each $j \in J$ there exists:

1. *an invertible matrix Y_j of size $D_j \times D_{\pi(j)}$, and*
2. *a diagonal unitary matrix Z_j of size $D_j \times D_j$ satisfying $Z_j^{m_j} = \mathbb{1}$ and $[A_j^i, Z_j] = 0$ for all i ,*

such that

$$Z_j A_j^i = Y_j B_{\pi(j)}^i Y_j^{-1} \quad \text{for all physical indices } i.$$

Equivalently, setting $Y = \bigoplus_j Y_j \otimes \mathbb{1}_{r_j}$ and $Z = \bigoplus_j Z_j \otimes \mathbb{1}_{r_j}$, one has $ZA^i = YB^iY^{-1}$ for all i , with $[A^i, Z] = 0$ and $\mathcal{V}(A) = \mathcal{V}(ZA)$.

Remark 12.14 (Comparison with the 1606.00608 approach). Theorem 12.13 and the equal-MPV Fundamental Theorem (Theorem 12.5) both characterize the gauge freedom between two tensors with equal MPV families, but they do so at different levels:

1. **Blocking approach** (Theorem 12.5, following [CPGSV17]): Block both A and B by a common period p so that all blocks of $A^{[p]}$ and $B^{[p]}$ are primitive (1-periodic). Apply the aperiodic fundamental theorem to the blocked tensors. The conclusion is that $A^{[p]}$ and $B^{[p]}$ are gauge equivalent. The Z -gauge freedom does not appear because blocking absorbs the periodicity.

2. **Periodic approach** (Theorem 12.13, following [DICCSPG17]): Work directly with the periodic blocks of A and B . The conclusion includes an extra Z -gauge transformation reflecting the non-trivial phase ambiguity that arises for periodic blocks. The Z matrix satisfies $Z^m = \mathbb{1}$ and commutes with all Kraus operators; it is invisible to the MPV family because it only introduces m -th roots of unity in the trace.

In the current formalization, only the blocking approach has been implemented. Formalizing the periodic approach would require developing the irreducible form and Z -gauge infrastructure outlined in Remark 12.10.

Remark 12.15 (Proportional periodic FT). There is also a proportional-case version of the periodic Fundamental Theorem (Theorem 3.4, “proportional case”, in [DICCSPG17]): if $\mathcal{V}(A) = \lambda_N \mathcal{V}(B)$ for scalars λ_N with $\lambda_N \rightarrow \lambda \neq 0$, then the bases of periodic tensors of A and B are equivalent (up to the same gauge and Z freedom as above). This extends Theorems 12.2 and 12.4 to the periodic setting.

Chapter 13

Quantum Dynamical Semigroups

This chapter develops the theory of one-parameter semigroups of quantum channels and characterizes their generators via the GKSL (Gorini–Kossakowski–Sudarshan–Lindblad) theorem, following [Wol12, Ch. 7].

Section 13.1 establishes definitions and the basic exponential form (Proposition 13.8). Section 13.2 gives the Duhamel formula and perturbation bound. Section 13.3 develops the generator theory and proves the GKSL theorem.

13.1 Dynamical semigroups and the exponential form

Definition 13.1 (Dynamical semigroup). A family of linear maps $T : \mathbb{R}_{\geq 0} \rightarrow (M_D(\mathbb{C}) \rightarrow_{\ell} M_D(\mathbb{C}))$ is a *dynamical semigroup* if

1. (semigroup law) $T(t + s) = T(t) \circ T(s)$ for all $t, s \geq 0$; and
2. (initial condition) $T(0) = \text{id}$.

This is [Wol12, Eq. (7.1)].

Definition 13.2 (Continuous dynamical semigroup). A dynamical semigroup T is *norm-continuous* if the map $t \mapsto T(t)$ is continuous in the operator norm topology. In finite dimension this coincides with strong continuity.

Definition 13.3 (Exponential semigroup). Given a generator $L \in \text{End}_{\mathbb{C}}(M_D(\mathbb{C}))$, the *exponential semigroup* is

$$T_t := e^{tL} = \sum_{k=0}^{\infty} \frac{(tL)^k}{k!}.$$

Theorem 13.4 (Semigroup law for the exponential). *For all $t, s \in \mathbb{R}$, $e^{(t+s)L} = e^{tL} \cdot e^{sL}$.*

Proof. Since $(t \cdot L)$ commutes with $(s \cdot L)$, this follows from the exponential addition formula for commuting elements in a Banach algebra. $\square$

Theorem 13.5 (Continuity of the exponential semigroup). *The map $t \mapsto e^{tL}$ is continuous in the operator norm topology.*

Proof. The exponential is analytic (convergence radius $= \infty$), hence continuous. $\square$

Theorem 13.6 (Derivative of the exponential semigroup). *For all $t \in \mathbb{R}$,*

$$\frac{d}{dt}e^{tL} = e^{tL} \cdot L.$$

This is [Wol12, Eq. (7.2)].

Proof. Chain rule applied to the composition $\mathbb{R} \xrightarrow{t \mapsto tL} \text{End}(M_D(\mathbb{C})) \xrightarrow{\exp} \text{End}(M_D(\mathbb{C}))$. $\square$

Theorem 13.7 (Generator uniqueness). *If $e^{tL} = e^{tL'}$ for all $t \geq 0$, then $L = L'$.*

Proof. Both semigroups agree on $[0, \infty)$, so their derivatives within $[0, \infty)$ at $t = 0$ agree. Since $[0, \infty)$ has unique differentials at 0, $L = L'$. $\square$

Theorem 13.8 (Prop. 7.1: Every continuous semigroup is exponential). *Every norm-continuous dynamical semigroup T on $M_D(\mathbb{C})$ is of the form*

$$T_t = e^{tL}$$

for a unique generator $L \in \text{End}_{\mathbb{C}}(M_D(\mathbb{C}))$. This is [Wol12, Prop. 7.1].

Proof. Since $T_0 = \mathbb{1}$ and T is continuous, the integral $M_\varepsilon = \int_0^\varepsilon T_s ds$ is invertible for small $\varepsilon > 0$. Using the semigroup property, $T_t = M_\varepsilon^{-1}(M_{t+\varepsilon} - M_t)$ is differentiable. ODE uniqueness gives $T_t = e^{tL}$ with $L = T'(0)$. $\square$

13.2 Perturbation theory

Theorem 13.9 (Lem. 7.1: Duhamel formula). *Let $\Delta = L' - L$. For $t \geq 0$,*

$$e^{tL'} - e^{tL} = \int_0^t e^{(t-s)L} \Delta e^{sL'} ds.$$

This is [Wol12, Lem. 7.1].

Proof. Define $f(s) = e^{(t-s)L} e^{sL'}$. Then $f'(s) = e^{(t-s)L} \Delta e^{sL'}$ (Theorem 13.6), and $e^{tL'} - e^{tL} = f(t) - f(0) = \int_0^t f'(s) ds$. $\square$

Theorem 13.10 (Cor. 7.1: Perturbation bound). *For $t \geq 0$,*

$$\|e^{tL'} - e^{tL}\| \leq t \|\Delta\| \cdot \sup_{s \in [0, t]} \|e^{sL}\| \cdot \sup_{s \in [0, t]} \|e^{sL'}\|,$$

where $\Delta = L' - L$. This is [Wol12, Cor. 7.1].

Proof. Apply the triangle inequality and submultiplicativity of the operator norm to the Duhamel integral (Theorem 13.9). $\square$

13.3 GKSL/Lindblad generators

This section characterizes generators of semigroups of completely positive and trace-preserving maps. The central result is the GKSL theorem (Theorem 13.29), which shows that such generators have the standard Lindblad form.

13.3.1 Generator decomposition and conditional complete positivity

Definition 13.11 (Generator decomposition). A *generator decomposition* consists of a completely positive map $\phi : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ and a matrix $\kappa \in M_d(\mathbb{C})$, defining the linear map

$$L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger.$$

This is [Wol12, Eq. (7.14)].

Definition 13.12 (Conditional complete positivity). A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is *conditionally completely positive* (CCP) if it admits a generator decomposition, i.e. there exist a CP map ϕ and a matrix κ such that $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$. This is condition 1 of [Wol12, Prop. 7.2].

Definition 13.13 (Trace-annihilating map). A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is *trace-annihilating* if $\text{tr}(L(\rho)) = 0$ for all $\rho \in M_d(\mathbb{C})$. This is the infinitesimal version of trace preservation.

Definition 13.14 (Trace constraint). A generator decomposition (ϕ, κ) satisfies the *trace constraint* if $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, where ϕ^* is the Hilbert–Schmidt adjoint. This is the condition in [Wol12, Eq. (7.20)].

Theorem 13.15 (Trace constraint implies trace-annihilating). *If (ϕ, κ) satisfies $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, then $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$ is trace-annihilating.*

Proof. Let $\phi(\rho) = \sum_i K_i \rho K_i^\dagger$. By the cyclic property of trace, $\text{tr}(L(\rho)) = \text{tr}((\sum_i K_i^\dagger K_i)\rho) - \text{tr}(\kappa\rho) - \text{tr}(\kappa^\dagger\rho) = \text{tr}((\kappa + \kappa^\dagger)\rho) - \text{tr}(\kappa\rho) - \text{tr}(\kappa^\dagger\rho) = 0$. $\square$

13.3.2 The Lindblad form

Definition 13.16 (Lindblad form). A *Lindblad form* consists of a Hermitian matrix $H = H^\dagger$ (the Hamiltonian) and a family of matrices $\{L_j\}_{j=0}^{r-1}$ (the Lindblad operators), defining the linear map

$$L(\rho) = i[\rho, H] + \sum_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\}_+ \right),$$

where $[A, B] = AB - BA$ and $\{A, B\}_+ = AB + BA$. This is [Wol12, Eq. (7.21)].

Theorem 13.17 (Lindblad form is trace-annihilating). *Every Lindblad form defines a trace-annihilating linear map.*

Proof. The Hamiltonian part satisfies $\text{tr}(i[\rho, H]) = 0$ by the cyclic property. Each dissipator term satisfies $\text{tr}(L_j \rho L_j^\dagger) = \text{tr}(L_j^\dagger L_j \rho) = \text{tr}(\rho L_j^\dagger L_j)$, so the three terms sum to zero. $\square$

Theorem 13.18 (Lindblad form equals generator decomposition). *A Lindblad form with Hamiltonian H and operators $\{L_j\}$ defines the same linear map as the generator decomposition (ϕ, κ) with $\phi(\rho) = \sum_j L_j \rho L_j^\dagger$ and $\kappa = iH + \frac{1}{2} \sum_j L_j^\dagger L_j$. This is [Wol12, Eq. (7.24)].*

Proof. Compute $\kappa^\dagger = -iH + \frac{1}{2} \sum_j L_j^\dagger L_j$ using $H^\dagger = H$ and $(A^\dagger A)^\dagger = A^\dagger A$. Setting $S = \sum_j L_j^\dagger L_j$, expand $\phi(\rho) - \kappa\rho - \rho\kappa^\dagger = \sum_j L_j \rho L_j^\dagger - iH\rho - \frac{1}{2}S\rho + i\rho H - \frac{1}{2}\rho S$, which is $i[\rho, H] + \sum_j (L_j \rho L_j^\dagger - \frac{1}{2}L_j^\dagger L_j \rho - \frac{1}{2}\rho L_j^\dagger L_j)$. $\square$

Theorem 13.19 (Lindblad form is CCP). *Every Lindblad form defines a conditionally completely positive map.*

Proof. By Theorem 13.18, the Lindblad form equals a generator decomposition with ϕ CP. $\square$

13.3.3 Prop. 7.2: Characterization of conditional complete positivity

Theorem 13.20 (Prop. 7.2, direction $2 \rightarrow 1$). *If L is Hermiticity-preserving and $P((L \otimes \text{id})(|\Omega\rangle\langle\Omega|))P \geq 0$ where $P = \mathbb{1} - |\Omega\rangle\langle\Omega|$, then L is CCP. This is [Wol12, Prop. 7.2].*

Proof. Decompose the Hermitian Choi matrix as $\tau_L = Q - |\psi\rangle\langle\Omega| - |\Omega\rangle\langle\psi|$ with $Q \geq 0$ supported on $|\Omega\rangle^\perp$. The Choi–Jamiołkowski isomorphism applied to Q gives the CP map ϕ , and $(\kappa \otimes \mathbb{1})|\Omega\rangle = |\psi\rangle$ defines κ . $\square$

13.3.4 Prop. 7.3: CP semigroups and CCP generators

Theorem 13.21 (Prop. 7.3: CP semigroup implies CCP generator). *If e^{tL} is CP for all $t \geq 0$, then L is CCP. This is [Wol12, Prop. 7.3].*

Proof. From $(e^{tL} \otimes \text{id})(|\Omega\rangle\langle\Omega|) \geq 0$, expand at infinitesimal t and project onto P to get $P(L \otimes \text{id})(|\Omega\rangle\langle\Omega|)P \geq 0$. $\square$

Theorem 13.22 (Prop. 7.3: CCP generator implies CP semigroup). *If L is CCP, then e^{tL} is CP for all $t \geq 0$. This is [Wol12, Prop. 7.3].*

Proof. Write $L = \phi + \phi_\kappa$ with $\phi_\kappa(\rho) = -\kappa\rho - \rho\kappa^\dagger$. By Lie–Trotter, $e^{tL} = \lim_{n \rightarrow \infty} (e^{t\phi/n} e^{t\phi_\kappa/n})^n$. Both factors are CP: $e^{t\phi/n}$ by Taylor expansion of a CP map, and $e^{t\phi_\kappa/n}(\rho) = K\rho K^\dagger$ with $K = e^{-t\kappa/n}$. $\square$

Theorem 13.23 (Prop. 7.3: CP semigroup iff CCP generator). *e^{tL} is CP for all $t \geq 0$ if and only if L is CCP.*

Proof. Theorems 13.21 and 13.22. $\square$

13.3.5 Prop. 7.4: Freedom in generator representation

Theorem 13.24 (Traceless Kraus operators exist). *Given any Kraus operators $\{K_j\}$, there exist $c_j \in \mathbb{C}$ such that $K'_j = K_j + c_j \mathbb{1}$ is traceless. This is part of [Wol12, Prop. 7.4].*

Proof. Set $c_j = -\text{tr}(K_j)/d$. $\square$

Theorem 13.25 (Shift invariance of generators). *If $K'_j = K_j + c_j \mathbb{1}$ and κ' is adjusted per [Wol12, Eq. (7.19)], then (ϕ', κ') and (ϕ, κ) define the same generator.*

Proof. Direct algebraic expansion. $\square$

13.3.6 Thm. 7.1: The GKSL/Lindblad theorem

Definition 13.26 (GKSL generator). A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is a *GKSL generator* if e^{tL} is a quantum channel (CPTP) for all $t \geq 0$.

Theorem 13.27 (Thm. 7.1, Form (i)). *If $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$ with ϕ CP and $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, then L is a GKSL generator. This is [Wol12, Thm. 7.1, Eq. (7.20)].*

Proof. CP follows from CCP via Theorem 13.22. TP follows from Theorem 13.15. $\square$

Theorem 13.28 (Thm. 7.1: GKSL iff CCP + trace-annihilating). *L is a GKSL generator if and only if L is CCP and trace-annihilating.*

Proof. Combines Props. 7.2–7.3 with the trace constraint. $\square$

Theorem 13.29 (Thm. 7.1: GKSL iff Lindblad form). *L is a GKSL generator if and only if*

$$L(\rho) = i[\rho, H] + \sum_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\}_+ \right)$$

with $H = H^\dagger$. This is [Wol12, Thm. 7.1].

Proof. Forward: by CCP, write $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$; take traceless Kraus operators (Theorem 13.24) and set $\kappa = iH + \frac{1}{2}\phi^*(\mathbb{1})$ (Theorem 13.18). Reverse: Theorem 13.28. $\square$

13.3.7 Kossakowski matrix form

Definition 13.30 (Kossakowski form). A *Kossakowski form* consists of $H = H^\dagger$, a basis $\{F_k\}_{k=0}^{d^2-2}$ of traceless matrices, and a PSD matrix $C \in M_{d^2-1}(\mathbb{C})$, defining

$$L(\rho) = i[\rho, H] + \frac{1}{2} \sum_{k,l} C_{lk} ([F_k, \rho F_l^\dagger] + [F_k \rho, F_l^\dagger]).$$

This is [Wol12, Eq. (7.23)].

Theorem 13.31 (Kossakowski form iff Lindblad form). *A linear map admits a Kossakowski form iff it admits a Lindblad form.*

Proof. Diagonalize $C = M^\dagger M$ and set $L_j = \sum_k M_{jk} F_k$. Conversely, expand traceless $L_j = \sum_k M_{jk} F_k$ and set $C = M^\dagger M \geq 0$. $\square$

Bibliography

- [CPGSV17] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete. Matrix product density operators: Renormalization fixed points and boundary theories. *Ann. Phys.*, 378:100–149, March 2017.
- [CPGSV21] J. Ignacio Cirac, David Pérez-García, Norbert Schuch, and Frank Verstraete. Matrix product states and projected entangled pair states: Concepts, symmetries, theorems. *Reviews of Modern Physics*, 93(4):045003, 2021.
- [DICCSPG17] Gemma De las Cuevas, J. Ignacio Cirac, Norbert Schuch, and David Pérez-García. Irreducible forms of matrix product states: Theory and applications. *Journal of Mathematical Physics*, 58(12):121901, 2017.
- [EHK78] David E. Evans and Raphael Høegh-Krohn. Spectral properties of positive maps on C^* -algebras. *Journal of the London Mathematical Society*, 17(2):345–355, 1978.
- [Jac09] Nathan Jacobson. *Basic Algebra II*. Dover Publications, 2009. Unaltered reprint of the 2nd edition (W. H. Freeman, 1989).
- [PGVWC07] D. Pérez-García, F. Verstraete, M. M. Wolf, and J. I. Cirac. Matrix product state representations. *Quantum Information & Computation*, 7(5):401–430, 2007.
- [SPGWC10] M. Sanz, D. Pérez-García, M. M. Wolf, and J. I. Cirac. A quantum version of Wielandt’s inequality. *IEEE Transactions on Information Theory*, 56(9):4668–4673, 2010.
- [Wol12] Michael M. Wolf. Quantum channels & operations: Guided tour. Lecture notes, 2012. Originally distributed via TU Munich foswiki; archival PDF at <https://mediatum.ub.tum.de/doc/1099654/1099654.pdf>.