

Tensor Network Theory:

A formalization blueprint

Sirui Lu

Erickson Tjoa

J. Ignacio Cirac

Last update: July 15, 2026

Contents

1	Introduction	7
2	Matrix Product Vectors	9
2.1	Basic definitions	9
2.2	Site-periodic tensor families	11
2.3	Gauge equivalence and same MPV	11
2.4	Injectivity and normality	13
2.5	Canonical form	14
2.6	Blocking	16
2.7	MPV overlap	17
3	The Single-Block Fundamental Theorem	18
3.1	The multiplicative linear extension	18
3.2	Inner automorphism and the single-block theorem	19
4	Quantum Channels and Positive Maps	21
4.1	Positive maps and channels	21
4.2	Irreducibility	27
4.3	Cesàro fixed-point theorem	28
4.4	Fixed-point projection	28
4.5	Peripheral spectrum and primitivity	29
4.5.1	Periodicity removal by powering	29
4.5.2	The Kadison–Schwarz inequality and the multiplicative domain	30
4.5.3	Peripheral closure via adjoint fixed point	31
4.6	Fixed-point algebra	33
5	Schwarz Inequalities and Multiplicative Domains	36
5.1	Kadison–Schwarz inequality	36
5.1.1	Two-positive maps and the generalized Schwarz inequality	37
5.1.2	The map T_η and strictness of the chain	48
5.1.3	The positivity hierarchy and Kadison–Schwarz from two-positivity	50
5.1.4	Douglas-type factorization lemmas	50
5.2	Multiplicative domain	51
5.2.1	Full multiplicative-domain structure	52
5.2.2	Schwarz inequality for normal operators	53
5.2.3	Schwarz inequality for subnormal and commuting-dominant operators	53
5.2.4	Positive maps preserve order and spectral intervals	54
5.2.5	A positive Schwarz map that is not completely positive	55

6	Perron–Frobenius Theory for Channels and Transfer Maps	56
6.1	Positive definiteness	56
6.2	Uniqueness	57
6.3	Existence and the Perron–Frobenius theorem	58
6.4	Right- and left-canonical gauges	58
6.5	Similarity preserves irreducibility	59
6.6	Perron–Frobenius eigenvector existence	60
6.7	Exponential positivity for irreducible CP maps	61
6.8	Ergodicity of irreducible channels	62
6.9	Spectral radius at the Perron eigenvalue	62
6.10	Spectral characterization of irreducibility	63
7	Transfer-Operator Gaps and Block Separation	64
7.1	Mixed transfer operator	64
7.2	MPV overlap as transfer trace	65
7.3	Frobenius norm estimates	66
7.4	Eigenvalue bound and transfer-operator gap	66
7.5	Transfer-operator gap — rectangular case	68
7.6	MPV overlap decay	69
7.7	Block separation	69
7.8	Transfer-operator gap under irreducible-TP hypotheses	70
7.9	Conditional expectation from a faithful fixed point	71
7.10	Stationary support	72
7.10.1	Faithful compression onto the support sector	73
7.11	Wedderburn decomposition of the fixed-point algebra	74
7.12	Further irreducibility and primitivity equivalences	91
7.13	Multi-cycle block-permutation structure	92
7.14	Peripheral eigenvalue group structure	93
7.14.1	Cyclic decomposition of irreducible Schwarz maps	94
7.14.2	Group structure of the peripheral spectrum	96
7.15	Primitive overlap convergence (complementary transfer-map gap form)	96
8	Wielandt Bound	97
8.1	Cumulative span	97
8.2	Fitting decomposition	99
8.3	Nonzero trace product	99
8.4	Eigenvector extraction	100
8.5	Eigenvector spreading	100
8.6	Proof of the cumulative Wielandt bound	101
8.7	Fixed-length matrix spanning	105
8.8	Blocked tensor and rectangular span	107
8.9	Primitive MPS tensors	108
8.10	Primitivity and normality	109
8.10.1	From primitivity to normality	109
8.10.2	Primitive implies irreducible	110
8.10.3	From irreducibility to full spanning via Burnside’s theorem	110
8.10.4	From primitivity to strong irreducibility and normality	111
8.11	Cumulative span to word span	112
8.12	Block injectivity from TP/primitive/irreducible blocks	113

9	Canonical Form Reduction	114
9.1	Normal tensors and canonical forms	115
9.1.1	Normalization conventions	116
9.2	Invariant-subspace decomposition and irreducible blocks	116
9.2.1	Iterated reduction to irreducible blocks	121
9.3	Normal canonical form	123
9.3.1	Canonical form from primitivity	124
9.4	Left-canonical and dual diagonalization of irreducible blocks	125
9.5	Zero-block separation and the trace-preserving gauge	126
9.5.1	Trace-preserving gauge for arbitrary tensors	127
9.6	Blocking, period removal, and primitive blocks	127
9.6.1	Period removal by cyclic sectors	128
9.6.2	Reduction to primitive blocks	131
9.7	Blocking and weighted direct sums	133
10	Basis of Normal Tensors	136
10.1	Bases of normal tensors	136
10.2	Permutation rigidity for bases of normal tensors	141
10.3	Newton–Girard identities and power-sum recovery	143
10.4	The BNT canonical form on a sector decomposition	145
10.4.1	Sector decompositions and phase matching	145
10.4.2	Prepared-block construction of BNT canonical forms	148
10.5	Coefficient comparison and copy-weight recovery	151
10.5.1	Matched-sector weight multiset equality	151
10.6	Strong existential matching by exact linear independence	152
10.6.1	Bijective matching by symmetry	153
11	Proof of the Fundamental Theorem	154
11.1	Coefficient identity, copy weights, and the block-diagonal gauge	154
11.2	The equal and proportional cases	162
12	Symmetries and String Order	165
12.1	Physical Symmetries Give Virtual Gauges	165
12.2	Virtual Representation Theorem	165
12.3	Cohomology Classes of Virtual Symmetry Cocycles	168
12.3.1	Non-triviality via the commutator phase	169
12.4	Permutation-Based Physical Symmetry	169
12.5	String Order and Local Symmetry Equivalence	170
12.6	SPT Phase Labels and String-Order Universality	176
13	Parent Hamiltonians	178
13.1	The N -site Hilbert space	178
13.2	Local ground space $G_L(A)$	179
13.3	Parent interaction and chain Hamiltonian	180
13.4	Intersection property and unique ground state	182
13.4.1	Restriction maps	183
13.4.2	Forward direction: ground space restricts	183
13.4.3	Injectivity of the ground-space map	184
13.4.4	The intersection property	184
13.4.5	Unique ground state	185

13.5	Periodic-boundary comparison for injective tensors	189
13.6	Periodic unique ground state	194
13.7	Injectivity-length reduction for the closure property	195
13.8	Periodic-boundary comparison for normal tensors	198
13.8.1	Reverse comparison for boundary-crossing restrictions	209
13.8.2	Normal unique-ground-state consequences	213
13.9	Commuting Local Terms for Parent Hamiltonians	214
13.10	Decorrelation and the idempotent-product parent-commuting condition	235
13.11	Spectral gap	239
13.11.1	Martingale condition for a finite family of projections	243
13.11.2	Cyclic-window overlap estimates	246
13.11.3	Martingale gap consequences	250
13.12	Block injectivity and degenerate ground spaces	254
13.13	Block separation and intersection	257
13.13.1	Trace identities for block intersections	264
13.13.2	Eventual block-intersection consequences	273
13.14	Block-diagonal boundary conditions for periodic ground spaces	276
13.14.1	Cyclic windows at the boundary cut	280
13.14.2	Periodic-boundary ground space for a block-diagonal tensor	295
13.15	Periodic-boundary ground space generated by a basis of normal tensors	302
14	Exponential Decay of Correlations	308
14.1	Connected correlators from the transfer map	308
14.2	Spectral expansion and exponential decay	309
14.3	Quantitative transfer-map gap bounds	310
14.4	Relation to parent Hamiltonians	311
15	Concrete Examples	312
15.1	The GHZ state	312
15.2	The AKLT state	314
15.2.1	Correlation length	316
15.3	The Majumdar–Ghosh state	317
15.4	The W state	318
15.5	The cluster state	319
15.6	Parent Hamiltonian statements for the examples	321
16	Channel Representations and Normal Forms	324
16.1	Representations	324
16.2	Choi–Jamiolkowski isomorphism	325
16.3	Representation corollaries for channel decompositions	326
16.4	Kraus representation theorem	327
16.5	Stinespring dilation	330
16.6	Ordered CP maps, Radon–Nikodym, and open-system representation	331
16.7	POVMs and Naimark dilation	334
16.8	Trace-pairing expansion in transfer-matrix form	336
16.9	SVD normal form (existence)	337
16.10	Lorentz normal form	338
16.11	Determinant of a quantum channel	340

17 Quantum Dynamical Semigroups	342
17.1 Dynamical semigroups and the exponential form	342
17.2 Perturbation theory	344
17.3 GKSL/Lindblad generators	346
17.3.1 Generator decomposition and conditional complete positivity	346
17.3.2 The Lindblad form	346
17.3.3 Characterization of conditional complete positivity	347
17.3.4 Completely positive semigroups and conditionally completely positive generators	348
17.3.5 Freedom in generator representation	348
17.3.6 Uniqueness of the traceless Lindblad form	348
17.3.7 Trace-annihilation and trace preservation	349
17.3.8 The GKSL/Lindblad theorem	349
17.3.9 Kossakowski matrix form	350
17.4 Primitivity and irreducibility of QDS	350
17.4.1 Auxiliary spectral and semigroup lemmas	350
17.5 Kernel of the adjoint Liouvillian	354
17.6 Reducibility of quantum dynamical semigroups	355
17.6.1 Sufficient conditions for non-reducibility	357
18 Operator Convexity and Jensen Inequalities	359
18.1 Diagonal Jensen inequality	359
18.2 Trace concavity and convexity of matrix powers	359
18.3 Operator convexity of real powers	360
18.4 Operator Jensen inequality for positive maps	360
18.5 Lieb concavity theorem	364
18.6 Resolvent monotonicity toward the Lieb concavity theorem	367
19 Quantum Entropy	369
19.1 Von Neumann entropy	369
19.2 Tripartite partial traces	378
19.3 Strong subadditivity	378
19.4 Mutual information	381
19.5 Entropy formulations	381
19.6 Mutual information: monotonicity and area-law bound	382
19.7 Trivial-factor corollaries	383
20 Positive but Not Completely Positive Maps	384
20.1 The reduction criterion	384
20.2 The Breuer–Hall map	386
20.3 Choi-type maps	387
20.4 Decomposable positive maps	390
20.5 The partial transpose and the PPT property	391
20.6 Separable states and the PPT criterion	393
20.7 The Schmidt number and the full reduction criterion	395

21	Matrix Product Operators and Density Operators	400
21.1	MPO tensors	400
21.2	Transfer maps	402
21.3	MPDOs and LPDOs	403
21.4	Horizontal and Vertical Canonical Forms	404
22	Pure States: Renormalization of Matrix Product States	434
22.1	Physical blocking and transfer idempotence	434
22.2	Zero-correlation-length conditions	436
22.3	Normal tensors and isometry canonical forms	438
22.4	Direct sums and residual isometries	442
22.5	Renormalization flow and physical correlations	447
23	Mixed States: Renormalization of Matrix Product Operators	450
23.1	Renormalization fixed points and pure-state recovery	450
23.2	Zero correlation length	456
23.3	Purification fixed points	457
23.4	Saturation of the area law	462
23.5	Simple local structure from SAL and ZCL	469
23.5.1	Markov decomposition and inverse-map sector factorization	469
23.5.2	Neighboring operators and commuting bond products	476
23.5.3	Closed-sector contractions and coherent rephasing	487
23.5.4	Normalized preparations and controlled partial traces	498
23.5.5	Refinement and coarse-graining channels	505
23.5.6	Primitivity and rank-one trace matrices	513
23.6	Commuting form and GSNNCH	521
24	Mixed States: General Case and Algebraic Structure	524
24.1	Local structure and two-site blocking	524
24.2	Algebra structure	526
24.3	Diagonal χ -matrices and the trace-power formula	531
24.4	Fusion isometries	541

Chapter 1

Introduction

The central result is the Fundamental Theorem for translation-invariant matrix product vectors. The basic object is a family of matrices $A = \{A^i\}_{i=0}^{d-1}$, which assigns to each chain length N the vector $|V^{(N)}(A)\rangle$ with coefficients indexed by $\sigma = (i_1, \dots, i_N)$,

$$V^{(N)}(A)_\sigma = \text{tr}(A^{i_1} \dots A^{i_N}).$$

The main argument, for translation-invariant tensors, follows the canonical-form and basis-of-normal-tensors argument of [CPGSV16, CPGSV21], with the single-block injective case anchored in the earlier representation theorem of [PGVWC07].

Notation

$\mathbb{C}, \mathbb{N}, \mathbb{Z}$	complex numbers, natural numbers, integers
$M_D(\mathbb{C}), \text{GL}_D(\mathbb{C})$	complex matrices and invertible complex matrices
$\text{tr}(X), \text{span}_{\mathbb{C}}(S)$	trace and complex linear span
$A = \{A^i\}_{i=0}^{d-1}$	a translation-invariant MPS tensor
$w = (i_1, \dots, i_L), A^w$	a word and the product $A^{i_1} \dots A^{i_L}$
$ V^{(N)}(A)\rangle$	the length- N matrix product vector
$V^{(N)}(A)_\sigma$	the coefficient of $ V^{(N)}(A)\rangle$ at σ
$O_{AB}(N)$	the bilinear overlap $\sum_\sigma V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma}$
$\mathcal{E}_A(X)$	the transfer map $\sum_i A^i X (A^i)^\dagger$
$A^{[L]}$	the length- L blocked tensor

In canonical form one writes

$$A^i = \bigoplus_{j=1}^g \mu_j A_j^i, \quad B^i = \bigoplus_{k=1}^h \nu_k B_k^i, \quad (1.1)$$

where the displayed blocks are normal tensors. If $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every positive chain length N , then the bases of normal tensors have the same size. After relabelling the blocks of B , each matched pair satisfies

$$B_j^i = \zeta_j X_j A_j^i X_j^{-1}, \quad (1.2)$$

for some invertible matrix X_j and nonzero scalar ζ_j , and the equal case compares the coefficient power sums attached to the relabelled bases. The block gauges (1.2) and the weight comparison

then combine into a single gauge conjugacy of the two weighted block-diagonal tensors (1.1). For a single injective block this reduces to the familiar conclusion $B^i = X A^i X^{-1}$ for all physical indices i , the single-block specialization of (1.2).

Chapters 2–11 carry out this argument. Chapter 2 fixes the matrix product vector notation, word evaluations, transfer maps, gauge equivalence, injectivity, blocking, and site-periodic tensor families. Chapter 3 proves the single-block theorem by trace pairings, linear extension, simplicity of matrix algebras, and Skolem–Noether. Chapters 4, 5, 6, and 7 develop the positive-map, multiplicative-domain, Perron–Frobenius, and overlap-decay tools used to control normal blocks. Chapters 8 and 9 supply the blocking, primitivity, irreducibility, and canonical-form reductions. Chapter 9 includes the invariant-projection splitting. Chapter 10 supplies bases of normal tensors and the power-sum comparison.

Chapter 12 uses the Fundamental Theorem to obtain virtual symmetry gauges, projective bond representations, and string-order criteria; its Kraus-mixing step appeals only to the later channel-representation chapter, not to any additional input for the Fundamental Theorem.

Beyond Chapter 12, the later chapters develop parent-Hamiltonian consequences, correlation estimates, examples, channel representations, normal forms, quantum dynamical semigroups, operator convexity, and entropy. They also include the standard examples of positive maps that fail to be completely positive. The final three chapters introduce matrix product operators and density operators, develop the renormalization fixed points of pure matrix product states, and then treat the zero-correlation-length and fixed-point conditions for mixed states.

Chapter 2

Matrix Product Vectors


Fix a physical dimension d and a bond dimension D . This chapter studies the tensors that generate matrix product vector (MPV) families. All tensors here are translation-invariant with periodic boundary conditions (PBC).

2.1 Basic definitions

Definition 2.1.1 (MPS tensor). [Lean](#) [✓ Stmt](#) A (translation-invariant, PBC) *MPS tensor* with physical dimension d and bond dimension D is a collection of matrices

$$\{A^i\}_{i=0}^{d-1}, \quad A^i \in M_D(\mathbb{C}),$$

indexed by a physical index $i \in \{0, \dots, d-1\}$. Such a tensor defines an MPV family. Diagrammatically,




The black node denotes the tensor A , the horizontal legs are virtual, and the upper leg is the physical index i .

Definition 2.1.2 (Word evaluation). [Lean](#) [✓ Stmt](#) Given a word $w = (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$, the *word evaluation* is the matrix product

$$A^w := A^{i_1} A^{i_2} \dots A^{i_L} \in M_D(\mathbb{C}),$$

with the convention that the empty word evaluates to the identity: $A^\emptyset = \mathbb{1}_D$. In tensor-network notation,



in which each black node denotes the same local tensor A , the virtual legs remain open, and the physical legs are labelled by the word $(i_1, \dots, i_L)$.

Lemma 2.1.3 (Word evaluation along a listed configuration). [Lean](#) [✓ Stmt](#) For a configuration $\sigma = (\sigma_0, \dots, \sigma_{N-1})$, let $w_\sigma = [\sigma_0, \dots, \sigma_{N-1}]$. Then

$$A^{w_\sigma} = A^{\sigma_0} A^{\sigma_1} \dots A^{\sigma_{N-1}}.$$

Proof. [✓ Proof](#) This follows by induction on N from the recursive definition of word evaluation. $\square$

Lemma 2.1.4 (Multiplicativity of word evaluation). [Lean](#) [✓ Stmt](#) For any words w_1, w_2 , $A^{w_1 \cdot w_2} = A^{w_1} A^{w_2}$.

Proof. [✓ Proof](#) The claim follows from the definition of A^w . $\square$

Lemma 2.1.5 (Centre of $M_D(\mathbb{C})$). [Lean](#) [✓ Stmt](#) If a matrix $Z \in M_D(\mathbb{C})$ commutes with every element of $M_D(\mathbb{C})$, then there exists $\lambda \in \mathbb{C}$ such that $Z = \lambda \mathbb{1}_D$.

Proof. [✓ Proof](#) Since Z commutes with every matrix unit, all off-diagonal entries of Z vanish and all diagonal entries are equal. $\square$

Lemma 2.1.6 (Rectangular amplified commutant). [Lean](#) [✓ Stmt](#) Let $(A_i)_{i \in I}$ span $M_D(\mathbb{C})$, and let

$$C : \mathbb{C}^{m_2} \otimes \mathbb{C}^D \longrightarrow \mathbb{C}^{m_1} \otimes \mathbb{C}^D$$

satisfy

$$(\mathbb{1}_{m_1} \otimes A_i)C = C(\mathbb{1}_{m_2} \otimes A_i) \quad (i \in I).$$

Then there is a matrix $F : \mathbb{C}^{m_2} \rightarrow \mathbb{C}^{m_1}$ such that

$$C = F \otimes \mathbb{1}_D.$$

Proof. [✓ Proof](#) Each (a, b) multiplicity block of C commutes with every A_i . Since the A_i span the full matrix algebra, Lemma 2.1.5 makes this block a scalar multiple of the identity. These scalars are the entries of F . $\square$

Definition 2.1.7 (Matrix product vector). [Lean](#) [✓ Stmt](#) The *matrix product vector* (MPV) of a tensor A at system size N is the vector

$$|V^{(N)}(A)\rangle = \sum_{i_1, \dots, i_N=0}^{d-1} \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}) |i_1, \dots, i_N\rangle \in (\mathbb{C}^d)^{\otimes N}.$$


Equivalently, for a configuration $\sigma = (i_1, \dots, i_N) \in \{0, \dots, d-1\}^N$, we write

$$V^{(N)}(A)_\sigma := \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}).$$

The coefficient function $\sigma \mapsto V^{(N)}(A)_\sigma$ gives the components; the displayed ket is the corresponding vector in $(\mathbb{C}^d)^{\otimes N}$. For a general word w , we also write

$$c_w(A) := \text{tr}(A^w).$$

The *MPV family* generated by A is the collection $\mathcal{V}(A) = \{|V^{(N)}(A)\rangle\}_{N \geq 1}$. The coefficient $V^{(N)}(A)_\sigma$ is the periodic contraction




of N copies of the local tensor A , with the outer virtual legs closed by the trace.

Definition 2.1.8 (Transfer map). [Lean](#) [✓ Stmt](#) The *transfer map* associated to a tensor A is the linear map $\mathcal{E}_A : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$\mathcal{E}_A(X) = \sum_{i=0}^{d-1} A^i X (A^i)^\dagger.$$

Diagrammatically, the transfer map is the double-layer contraction



in which the upper node denotes A , the lower node denotes $A^\dagger$, and the physical index is summed over between them.

Remark 2.1.9. [✓ Stmt](#) The transfer map is automatically positive: if $X \geq 0$ then $\mathcal{E}_A(X) \geq 0$, because each summand $A^i X (A^i)^\dagger$ is positive semidefinite. The abstract positive-map framework appears in Chapter 4.

2.2 Site-periodic tensor families

Definition 2.2.1 (Site-periodic tensor family). [Lean](#) [✓ Stmt](#) For a fixed period m , a *site-periodic tensor family* is a family of local tensors indexed by $\{0, \dots, m-1\}$, with local physical dimension d and bond dimension D .

Definition 2.2.2 (Periodic same-state and gauge-equivalence relations). [Lean](#) [Lean](#) [✓ Stmt](#) For site-periodic tensors A, B at fixed period m :

- equality of the induced site-periodic states;
- cyclic gauge equivalence.


Lemma 2.2.3 (Equivalence structures for periodic relations). [Lean](#) [Lean](#) [✓ Stmt](#) Both *periodic relations* are *equivalence relations* (*reflexive, symmetric, transitive*).

2.3 Gauge equivalence and same MPV

Definition 2.3.1 (Gauge equivalence). [Lean](#) [✓ Stmt](#) Two tensors A and B of the same bond dimension D are *gauge equivalent* if there exists an invertible matrix $X \in \text{GL}_D(\mathbb{C})$ such that, for every $i \in \{0, \dots, d-1\}$,

$$B^i = X A^i X^{-1}.$$

In tensor-network notation,



the black node denotes the tensor named in the surrounding formula, the upper leg is the physical index i , and the two red side nodes denote the gauge matrices acting on the virtual legs.

Definition 2.3.2 (Same MPV). [Lean](#) [✓ Stmt](#) Two tensors A and B (of the same bond dimension) generate the same MPV family if $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every system size N .

Definition 2.3.3 (Same MPV - different bond dimensions). [Lean](#) [✓ Stmt](#) For tensors A and B of possibly different bond dimensions D_1 and D_2 , we write $\mathcal{V}(A) = \mathcal{V}(B)$ if $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every N .

Definition 2.3.4 (Positive-length same MPV - different bond dimensions). [Lean](#) [✓ Stmt](#) For tensors A and B of possibly different bond dimensions, we write $\mathcal{V}^+(A) = \mathcal{V}^+(B)$ if $|V^{(N)}(A)\rangle = |V^{(N)}(B)\rangle$ for every $N > 0$.

Lemma 2.3.5 (Length-zero MPV coefficient). [Lean](#) [✓ Stmt](#) For any MPS tensor A with bond dimension D , the length-zero MPV coefficient is D , the trace of the identity on the bond space.

Lemma 2.3.6 (Positive-length equality with equal bond dimensions). [Lean](#) [✓ Stmt](#) If two tensors have the same positive-length MPV family and the same bond dimension, then they have the same MPV family at every length.

Remark 2.3.7. [✓ Stmt](#) Definition 2.3.2 is the same-bond-dimension specialization of Definition 2.3.3. We keep both because later results use both the same-dimension and different-dimension forms.

Definition 2.3.8 (Proportional MPV). [Lean](#) [✓ Stmt](#) Two tensors A and B (of possibly different bond dimensions) generate weakly proportional MPV families if for every N there exists $c_N \in \mathbb{C}$ such that $|V^{(N)}(A)\rangle = c_N |V^{(N)}(B)\rangle$.

Definition 2.3.9 (Nonzero proportional MPV). [Lean](#) [✓ Stmt](#) Two tensors A and B (of possibly different bond dimensions) generate nonzero proportional MPV families if for every N there is a nonzero scalar $c_N \in \mathbb{C}$ such that $|V^{(N)}(A)\rangle = c_N |V^{(N)}(B)\rangle$. This is the projective reading of the proportionality hypothesis in [CPGSV16, Theorem 2.10].

Definition 2.3.10 (Eventually nonzero proportional MPV). [Lean](#) [✓ Stmt](#) Two tensors A and B are eventually nonzero proportional if, for all sufficiently large N , there is a nonzero scalar $c_N \in \mathbb{C}$ such that $|V^{(N)}(A)\rangle = c_N |V^{(N)}(B)\rangle$. This auxiliary form is used with the eventual linear-independence statement Lemma 10.1.14.

Lemma 2.3.11 (Elementary nonzero proportionality rules). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Nonzero proportionality implies weak proportionality and eventual nonzero proportionality, is symmetric, and equality of MPV families is the special case with scalar 1.

Proof. [✓ Proof](#) Forgetting the nonvanishing condition gives weak proportionality. The eventual form follows by viewing an all-length statement as an eventual one. Symmetry follows by inverting the scalar c_N at each length, and equality is the case $c_N = 1$. $\square$

Lemma 2.3.12 (Scaling of word evaluation). [Lean](#) [✓ Stmt](#) Let ζA denote the tensor with matrices $\{\zeta A^i\}_{i=0}^{d-1}$. Then for any word w ,

$$(\zeta A)^w = \zeta^{|w|} A^w,$$

where $|w|$ is the length of w .

Proof. ✓ Proof Induction on w : the base case gives $\zeta^0 = 1$, and each step picks up one factor of ζ . □

Lemma 2.3.13 (Scaling of MPV components). Lean ✓ Stmt Let ζA denote the tensor with matrices $\{\zeta A^i\}_{i=0}^{d-1}$. Then for every N and every configuration $\sigma = (i_1, \dots, i_N)$,

$$V^{(N)}(\zeta A)_\sigma = \zeta^N V^{(N)}(A)_\sigma.$$

Proof. ✓ Proof Apply Lemma 2.3.12 to the word $w = (i_1, \dots, i_N)$ of length N , then take the trace; the scalar ζ^N pulls out by linearity. □

Definition 2.3.14 (Gauge-phase equivalence). Lean Lean ✓ Stmt Two tensors A and B are *gauge-phase equivalent* if there exist $X \in \text{GL}_D(\mathbb{C})$ and $\zeta \in \mathbb{C} \setminus \{0\}$ such that, for every physical index i ,

$$B^i = \zeta X A^i X^{-1}.$$

Proof. ✓ Proof Gauge equivalence is the special case $\zeta = 1$. □

Remark 2.3.15. ✓ Stmt After spectral-radius normalization one may restrict ζ to a unit-modulus phase $e^{i\phi}$. We allow an arbitrary nonzero scalar because later canonical-form statements keep the block-scaling factors explicit.

Lemma 2.3.16 (Gauge covariance of word evaluation). Lean ✓ Stmt If $B^i = X A^i X^{-1}$ for all i , then for every word w ,

$$B^w = X A^w X^{-1}.$$

Proof. ✓ Proof Induction on the word length, using $XX^{-1} = \mathbb{1}$. □

Theorem 2.3.17 (Gauge invariance of MPVs). Lean ✓ Stmt If A and B are gauge equivalent, then they generate the same MPV family.

Proof. ✓ Proof Let $B^i = X A^i X^{-1}$. By Lemma 2.3.16, $B^w = X A^w X^{-1}$ for every word w . Then $\text{tr}(B^w) = \text{tr}(X A^w X^{-1}) = \text{tr}(A^w)$ by cyclicity of the trace. □

Theorem 2.3.18 (Gauge-phase scaling of MPVs). Lean ✓ Stmt If, for every physical index i ,

$$B^i = \zeta X A^i X^{-1},$$

then for every system size N and configuration σ one has

$$V^{(N)}(B)_\sigma = \zeta^N V^{(N)}(A)_\sigma.$$

Proof. ✓ Proof Combine gauge covariance of word evaluation with scalar scaling, then take traces. □

2.4 Injectivity and normality

Definition 2.4.1 (Injectivity). Lean ✓ Stmt A tensor A is *injective* if its matrices span the full matrix algebra: $\text{span}_{\mathbb{C}}\{A^i : i = 0, \dots, d-1\} = M_D(\mathbb{C})$.

The inverse-tensor characterization of injectivity says that A is injective exactly when there is a tensor A^{-1} such that $\sum_i (A^i)_{\alpha, \beta} (A^{-1})_{\alpha', \beta'}^i = \delta_{\alpha, \alpha'} \delta_{\beta, \beta'}$. Thus contracting A^{-1} with A through their common physical index leaves two disconnected virtual pairings:

$$A^{-1}A = \mathbb{1}$$

This is the identity used in [CPGSV16, Section 2.3, lines 326–331].

Definition 2.4.2 (*L*-block injectivity). Lean ✓ Stmt A tensor A is *L*-block injective if $\text{span}_{\mathbb{C}}\{A^{i_1} \cdots A^{i_L} : (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L\} = M_D(\mathbb{C})$. An injective tensor is 1-block injective.

Lemma 2.4.3 (One-site injectivity gives one-block injectivity). Lean ✓ Stmt If the one-site matrices $\{A^i\}_{i=0}^{d-1}$ span $M_D(\mathbb{C})$, then the length-one word products span $M_D(\mathbb{C})$, so A is 1-block injective.

Proof. ✓ Proof The length-one words are exactly the letters $i \in \{0, \dots, d-1\}$, and evaluating a length-one word gives the matrix A^i . Thus the length-one word span is the same span as in one-site injectivity. $\square$

Definition 2.4.4 (Normal tensor). Lean ✓ Stmt A tensor A is *normal* [CPGSV21] if there exists a finite L_0 such that A is L_0 -block injective.

Lemma 2.4.5 (Gauge invariance of block injectivity). Lean ✓ Stmt If A and B are gauge equivalent and A is *L*-block injective, then B is *L*-block injective.

Proof. ✓ Proof Conjugation by an invertible matrix maps the span of the length- L words of A onto the span of the length- L words of B . Since conjugation is an automorphism of the full matrix algebra, fullness of the first span implies fullness of the second. $\square$

Theorem 2.4.6 (Gauge transport of normality). Lean ✓ Stmt If A and B are gauge equivalent and A is normal, then B is normal.

Proof. ✓ Proof Apply Lemma 2.4.5 at a blocking length for which A is block-injective. $\square$

Remark 2.4.7. ✓ Stmt The algebraic characterization of normality in Definition 2.4.4 — eventual full matrix span, equivalently eventual block injectivity — is equivalent to the spectral characterization used in [CPGSV21, Definition IV.1]: after TP normalization, the transfer map $\mathcal{E}_A$ is a primitive channel, equivalently irreducible with trivial peripheral spectrum. This equivalence is proved in [SPGWC10, Proposition 3]. In what follows, the directions of the equivalence are supplied by Lemma 8.10.2.1, Theorem 8.10.4.3 for primitive normalized tensors with positive-definite fixed point, the Burnside route through Theorem 8.11.7 for the aperiodic algebraic criterion, and Theorem 9.3.1.1. In particular, every subsequent use of “normal” here refers to this single notion; Definition 9.3.3 employs the spectral formulation, but it describes the same class of tensors.

2.5 Canonical form

Definition 2.5.1 (Block-injective canonical form). Lean ✓ Stmt Here, a *block-injective canonical form* for a tensor consists of:

1. a number of blocks $r \geq 1$,
2. bond dimensions $D_1, \dots, D_r$,
3. injective tensors $\{A_k^i\}_{i=0}^{d-1}$ with $A_k^i \in M_{D_k}(\mathbb{C})$ for each block $k \in \{1, \dots, r\}$,
4. scaling factors $\mu_1, \dots, \mu_r \in \mathbb{C}$.

The associated tensor is the block-diagonal tensor

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

This block-injective formulation of the canonical form is used in [PGVWC07]; see also [CPGSV21] for the normal canonical form.

Remark 2.5.2. ✓ Stmt Here we use the block-injective canonical form: each block is injective in the sense of Definition 2.4.1. This is stronger than the NT-block canonical form of [CPGSV21], where the blocks are only normal in the spectral sense. Chapter 9 treats the normal canonical form needed for the blocking-based multi-block theory, while the periodic irreducible-form extension is kept separate. This section isolates the stronger block-injective form used in the injective and block-injective arguments.

Definition 2.5.3 (Block projection). Lean ✓ Stmt For a finite direct sum $\bigoplus_k \mathbb{C}^{D_k}$, the block projection P_j is the matrix that is the identity on the j -th summand and zero on all other summands.

Definition 2.5.4 (Block-diagonal matrix). Lean ✓ Stmt A matrix on $\bigoplus_k \mathbb{C}^{D_k}$ is block diagonal if it is a direct sum of square matrices, one on each summand.

Lemma 2.5.5 (Off-block characterization). Lean ✓ Stmt A matrix on $\bigoplus_k \mathbb{C}^{D_k}$ is block diagonal if and only if every entry from one summand to a distinct summand is zero.

Proof. ✓ Proof The forward direction follows from the definition of a direct sum of matrices. Conversely, if all off-block entries vanish, the diagonal blocks reconstruct the original matrix. $\square$

Lemma 2.5.6 (Commutation with block projections). Lean ✓ Stmt If a matrix on $\bigoplus_k \mathbb{C}^{D_k}$ commutes with every block projection P_j , then it is block diagonal.

Proof. ✓ Proof Compare the (i, j) block of the identity $XP_j = P_jX$ for $i \neq j$. The right multiplication by P_j keeps that block, while the left multiplication by P_j kills it. Hence every off-block entry is zero, and Lemma 2.5.5 applies. $\square$

Definition 2.5.7 (Block-diagonal tensor from blocks). Lean ✓ Stmt Given r blocks with bond dimensions $(D_k)_{k=1}^r$, per-block tensors $\{A_k^i\}$, and scaling factors $(\mu_k)_{k=1}^r$, the associated *block-diagonal tensor* is the tensor of total bond dimension $D = \sum_{k=1}^r D_k$ defined by

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

Lemma 2.5.8 (Word evaluation of a block-diagonal tensor). Lean ✓ Stmt For a word w , the word evaluation of a block-diagonal tensor remains block diagonal:

$$A^w = \bigoplus_k \mu_k^{w|} A_k^w.$$

Proof. ✓ Proof Induct on the word and multiply block-diagonal matrices block by block; every letter contributes one scalar factor μ_k on the k -th block. $\square$

Theorem 2.5.9 (MPV decomposition). [Lean](#) [✓ Stmt](#) The MPV of a block-diagonal tensor decomposes as

$$|V^{(N)}(A)\rangle = \sum_{k=1}^r \mu_k^N |V^{(N)}(A_k)\rangle.$$

Equivalently, for each configuration σ one has

$$V^{(N)}(A)_\sigma = \sum_{k=1}^r \mu_k^N V^{(N)}(A_k)_\sigma.$$


Proof. [✓ Proof](#) By Lemma 2.5.8, the full word evaluation is block diagonal with blocks $\mu_k^N A_k^w$ for a word w of length N . Taking the trace of a block-diagonal matrix gives the sum of the block traces: $\text{tr}(A^w) = \sum_k \mu_k^N \text{tr}(A_k^w)$. $\square$

2.6 Blocking

Definition 2.6.1 (Blocked tensor). [Lean](#) [✓ Stmt](#) Given a tensor A and a blocking length L , the L -blocked tensor is the tensor with physical index set $\{0, \dots, d-1\}^L$ (hence physical dimension d^L) and the same bond dimension D , whose matrices are indexed by words $(i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$:

$$(A^{[L]})^{(i_1, \dots, i_L)} = A^{i_1} A^{i_2} \dots A^{i_L}.$$

Blocking coarse-grains L neighbouring sites into one tensor: the inner virtual bonds are contracted, the L physical legs merge into a single composite index, and the two outer bonds remain as the bond of $A^{[L]}$.



Lemma 2.6.2 (Blocked word evaluation). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) If $w = (J_1, \dots, J_m)$ is a word in the blocked alphabet, where each J_k is an L -tuple in $\{0, \dots, d-1\}^L$, then concatenating the block words gives a word $\tilde{w}$ of length mL in the original alphabet and one has

$$(A^{[L]})^w = A^{\tilde{w}}.$$

Proof. [✓ Proof](#) Induct on the blocked word w and split off the first block word. Lemma 2.1.4 turns concatenation of blocks into multiplication of the corresponding evaluations. $\square$

Lemma 2.6.3 (Blocking preserves MPV coefficients). [Lean](#) [✓ Stmt](#) For each basis state $\sigma = (\sigma_1, \dots, \sigma_N)$ of the blocked system, where each σ_k is an L -tuple in $\{0, \dots, d-1\}^L$, there exists a basis state $\tilde{\sigma} = (\tilde{\sigma}_1, \dots, \tilde{\sigma}_{NL})$ of the original system such that

$$V^{(N)}(A^{[L]})_\sigma = V^{(NL)}(A)_{\tilde{\sigma}}.$$

Proof. [✓ Proof](#) Concatenate the block words and apply the blocked word evaluation lemma (Lemma 2.6.2). $\square$

Theorem 2.6.4 (Blocking preserves same MPV). [Lean](#) [✓ Stmt](#) If A and B generate the same MPV family, then for any blocking length L , the blocked tensors $A^{[L]}$ and $B^{[L]}$ also generate the same MPV family.

Proof. ✓ Proof By Lemma 2.6.3, each MPV coefficient of the blocked tensor $A^{[L]}$ equals a coefficient of A at the corresponding concatenated configuration. The same holds for $B^{[L]}$ (with the same concatenated configuration). Since A and B agree on all configurations, so do $A^{[L]}$ and $B^{[L]}$. $\square$

2.7 MPV overlap

Definition 2.7.1 (MPV as Hilbert-space vector). Lean ✓ Stmt We regard $|V^{(N)}(A)\rangle$ as an element of $\mathbb{C}^{d^N}$ indexed by configurations $\sigma \in \{0, \dots, d-1\}^N$, with σ -component $V^{(N)}(A)_\sigma$ as in Definition 2.1.7. This allows us to use the standard Euclidean inner product on $\mathbb{C}^{d^N}$.

Definition 2.7.2 (MPV inner product). Lean ✓ Stmt The *MPV inner product* between two tensors A and B at system size N is


$$\langle V^{(N)}(A) | V^{(N)}(B) \rangle := \sum_{\sigma \in \{0, \dots, d-1\}^N} \overline{V^{(N)}(A)_\sigma} V^{(N)}(B)_\sigma,$$

i.e. the Euclidean inner product on $\mathbb{C}^{d^N}$ (conjugate-linear in the first argument).

Definition 2.7.3 (MPV overlap). Lean ✓ Stmt The *MPV overlap* between two tensors A (of bond dimension D_1) and B (of bond dimension D_2) at system size N is

$$O_{AB}(N) := \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma}.$$

Diagrammatically, the overlap contracts the MPV ring of A against the conjugate ring of B along their shared physical indices, both virtual rings closed by the trace:



This is the bilinear overlap (without complex conjugation) used later. By Lemma 2.7.4, it is the complex conjugate of the Hilbert-space inner product from Definition 2.7.2.

Lemma 2.7.4 (Overlap equals conjugate inner product). Lean ✓ Stmt $O_{AB}(N) = \overline{\langle V^{(N)}(A) | V^{(N)}(B) \rangle}$.

Proof. ✓ Proof The overlap sums $V_\sigma \overline{W_\sigma}$ while the inner product sums $\overline{V_\sigma} W_\sigma$. Complex conjugation swaps these. $\square$

Lemma 2.7.5 (Overlap determined by positive-length MPV). Lean ✓ Stmt If $V^{(N)}(A)_\sigma = V^{(N)}(A')_\sigma$ and $V^{(N)}(B)_\sigma = V^{(N)}(B')_\sigma$ for all $N > 0$ and all σ , then $O_{AB}(N) = O_{A'B'}(N)$ for all positive N .

Proof. ✓ Proof Direct pointwise rewriting: if the MPV coefficients agree for all positive chain lengths, the overlap sums are identical. $\square$

Chapter 3

The Single-Block Fundamental Theorem

An injective MPS tensor that generates the same matrix-product vectors as a second tensor of equal bond dimension is conjugate to it [CPGSV21, Corollary IV.5]: there is an invertible X with $B^i = XA^iX^{-1}$ for every physical index i . The argument is purely algebraic. Equality of the matrix-product vectors gives $\text{tr}(A^w) = \text{tr}(B^w)$ for every word w ; the length-two and length-three traces produce a linear map T with $T(A^i) = B^i$ that is multiplicative and nonzero; a nonzero multiplicative endomorphism of the simple algebra $M_D(\mathbb{C})$ is a unital automorphism; and by Skolem–Noether every automorphism of $M_D(\mathbb{C})$ is conjugation by an invertible X . Evaluating at A^i closes the proof. The same conjugation, one normal block at a time, is the central step of the multi-block theorem in Chapter 11.

3.1 The multiplicative linear extension

Theorem 3.1.1 (Nondegeneracy of the trace pairing). Lean ✓ Stmt For $M \in M_D(\mathbb{C})$, $\text{tr}(MN) = 0$ for all $N \in M_D(\mathbb{C})$ if and only if $M = 0$.

Proof. ✓ Proof Testing against the matrix unit E_{ji} gives $\text{tr}(E_{ji}M) = M_{ij}$, so $\text{tr}(MN) = 0$ for all N forces every entry of M to vanish. □

Lemma 3.1.2 (Same matrix-product vectors give trace agreement). Lean ✓ Stmt If A and B generate the same matrix-product vectors, then $\text{tr}(A^w) = \text{tr}(B^w)$ for every word w .

Proof. ✓ Proof The configuration $\sigma = w$ gives $V^{(|w|)}(A)_w = \text{tr}(A^w)$. Equality of the matrix-product vectors gives $V^{(N)}(A) = V^{(N)}(B)$ for all N , so $\text{tr}(A^w) = \text{tr}(B^w)$. □

The trace conditions constrain every product of the matrices A^i . To bring the algebra structure of $M_D(\mathbb{C})$ to bear, record how a matrix pairs against the family $\{A^i\}$ under the trace.

Definition 3.1.3 (Trace pairing map). Lean ✓ Stmt For a tensor A , the trace pairing map $\Phi_A : M_D(\mathbb{C}) \rightarrow \mathbb{C}^d$ is

$$\Phi_A(M)_i = \text{tr}(MA^i).$$

Theorem 3.1.4 (The trace pairing map is injective for injective tensors). Lean ✓ Stmt If A is injective, then $\ker \Phi_A = \{0\}$.

Proof. ✓ Proof If $\Phi_A(M) = 0$, then $\text{tr}(MA^i) = 0$ for every i . The $\{A^i\}$ span $M_D(\mathbb{C})$, so $\text{tr}(MN) = 0$ for all N , and $M = 0$ by Theorem 3.1.1. $\square$

Lemma 3.1.5 (Injectivity transfers across a range inclusion). Lean ✓ Stmt *Let $\Phi, \Psi : V \rightarrow W$ be $\mathbb{C}$ -linear maps with V finite-dimensional. If $\ker \Phi = \{0\}$ and $\text{range } \Phi \subseteq \text{range } \Psi$, then $\ker \Psi = \{0\}$.*

Proof. ✓ Proof By rank–nullity, $\ker \Phi = \{0\}$ gives $\dim \text{range } \Phi = \dim V$, and the range inclusion gives

$$\dim V = \dim \text{range } \Phi \leq \dim \text{range } \Psi \leq \dim V,$$

so $\dim \text{range } \Psi = \dim V$ and $\ker \Psi = \{0\}$. $\square$

Theorem 3.1.6 (The linear extension exists and is unique). Lean ✓ Stmt *If A is injective and A, B generate the same matrix-product vectors, then there is a unique linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ with $T(A^i) = B^i$ for all i .*

Proof. ✓ Proof Length-two trace agreement gives $\Phi_A(A^i) = \Phi_B(B^i)$ for every i . The $\{A^i\}$ span $M_D(\mathbb{C})$, so $\text{range } \Phi_A \subseteq \text{range } \Phi_B$. With $\ker \Phi_A = \{0\}$ (Theorem 3.1.4), Lemma 3.1.5 gives $\ker \Phi_B = \{0\}$. Let g be a left inverse of Φ_B and set $T = g \circ \Phi_A$; then

$$T(A^i) = g(\Phi_A(A^i)) = g(\Phi_B(B^i)) = B^i.$$

Uniqueness holds because the $\{A^i\}$ span $M_D(\mathbb{C})$. $\square$

Theorem 3.1.7 (The linear extension is multiplicative). Lean ✓ Stmt *Under the hypotheses of Theorem 3.1.6, the map T satisfies $T(MN) = T(M)T(N)$ for all $M, N \in M_D(\mathbb{C})$.*

Proof. ✓ Proof Length-two trace agreement gives $\Phi_B(T(A^i)) = \Phi_B(B^i) = \Phi_A(A^i)$; since the $\{A^i\}$ span $M_D(\mathbb{C})$ this extends to $\Phi_B \circ T = \Phi_A$, so $\text{range } \Phi_A = \text{range } (\Phi_B \circ T) \subseteq \text{range } \Phi_B$. With $\ker \Phi_A = \{0\}$ (Theorem 3.1.4), Lemma 3.1.5 makes Φ_B injective. Length-three trace agreement gives, for all i, j, k ,

$$\Phi_B(T(A^i A^j))_k = \Phi_A(A^i A^j)_k = \text{tr}(A^i A^j A^k) = \text{tr}(B^i B^j B^k) = \Phi_B(B^i B^j)_k,$$

so injectivity of Φ_B yields $T(A^i A^j) = B^i B^j$. Extend in two stages: for fixed j , the maps $M \mapsto T(MA^j)$ and $M \mapsto T(M)B^j$ agree on the spanning $\{A^i\}$, hence $T(MA^j) = T(M)B^j$ for all M ; then for fixed M , the maps $N \mapsto T(MN)$ and $N \mapsto T(M)T(N)$ agree on the spanning $\{A^j\}$, hence $T(MN) = T(M)T(N)$ for all N . $\square$

3.2 Inner automorphism and the single-block theorem

The rigidity of the full matrix algebra now finishes the proof. A nonzero multiplicative map on $M_D(\mathbb{C})$ is forced to be bijective (simplicity), unit-preserving (surjectivity), and conjugation by a single invertible matrix (Skolem–Noether). The next four results establish these steps and the nonvanishing of T , which the final theorem combines into the gauge X .

Theorem 3.2.1 (A nonzero multiplicative endomorphism of $M_D(\mathbb{C})$ is bijective). Lean ✓ Stmt *A nonzero multiplicative $\mathbb{C}$ -linear endomorphism of $M_D(\mathbb{C})$ is bijective.*

Proof. ✓ Proof The kernel of a multiplicative linear map is a two-sided ideal. The algebra $M_D(\mathbb{C})$ is simple, so the kernel is $\{0\}$ or all of $M_D(\mathbb{C})$; the latter forces the map to be zero. Hence the map is injective, and surjective by finite-dimensionality. $\square$

Lemma 3.2.2 (A surjective multiplicative map preserves the unit). [Lean](#) [✓ Stmt](#) *If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a surjective multiplicative $\mathbb{C}$ -linear map, then $T(\mathbb{1}) = \mathbb{1}$, so T is a $\mathbb{C}$ -algebra homomorphism.*

Proof. [✓ Proof](#) Choose X with $T(X) = \mathbb{1}$. Then $T(\mathbb{1}) = T(X)T(\mathbb{1}) = T(X\mathbb{1}) = T(X) = \mathbb{1}$. $\square$


Lemma 3.2.3 (An injective tensor forces a nonzero partner). [Lean](#) [✓ Stmt](#) *Let $D \geq 1$. If A is injective and A, B generate the same matrix-product vectors, then the B^i are not all zero.*

Proof. [✓ Proof](#) If $B^i = 0$ for every i , then Lemma 3.1.2 on length-one words gives $\text{tr}(A^i) = 0$ for all i . The $\{A^i\}$ span $M_D(\mathbb{C})$, so the trace functional vanishes identically, contradicting $\text{tr}(\mathbb{1}) = D \neq 0$. $\square$

Theorem 3.2.4 (Skolem–Noether). [Lean](#) [✓ Stmt](#) *Every $\mathbb{C}$ -algebra automorphism f of $M_D(\mathbb{C})$ is inner: there is $X \in \text{GL}_D(\mathbb{C})$ with $f(M) = XMX^{-1}$ for all M .*

Proof. [✓ Proof](#) Under the identification $M_D(\mathbb{C}) \cong \text{End}_{\mathbb{C}}(\mathbb{C}^D)$, every algebra automorphism of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$ is conjugation by an invertible linear map. Translating back gives the invertible X . $\square$

Theorem 3.2.5 (Single-block Fundamental Theorem). [Lean](#) [✓ Stmt](#) *Let A be an injective MPS tensor and B a tensor of the same bond dimension. If A and B generate the same matrix-product vectors, then there is $X \in \text{GL}_D(\mathbb{C})$ with $B^i = XA^iX^{-1}$ for all i . In tensor-network notation the gauge conjugates the local tensor on its virtual legs, with the physical index i on the upper leg:*



Proof. [✓ Proof](#) Theorem 3.1.6 gives the unique linear T with $T(A^i) = B^i$, and Theorem 3.1.7 makes it multiplicative. It is nonzero: otherwise $B^i = 0$ for all i , against Lemma 3.2.3. By Theorem 3.2.1, T is bijective, and by Lemma 3.2.2 it fixes $\mathbb{1}$, so T is a $\mathbb{C}$ -algebra automorphism. Theorem 3.2.4 gives $X \in \text{GL}_D(\mathbb{C})$ with $T(M) = XMX^{-1}$; evaluating at $M = A^i$ gives $B^i = XA^iX^{-1}$. $\square$

Chapter 4

Quantum Channels and Positive Maps

This chapter develops the theory of positive maps and quantum channels on matrix algebras, following [Wol12].

4.1 Positive maps and channels

Definition 4.1.1 (Positive map). [Lean](#) [✓ Stmt](#) A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *positive* if $E(X) \geq 0$ whenever $X \geq 0$, where $X \geq 0$ means that X is positive semidefinite.

Remark 4.1.2. [✓ Stmt](#) We use the Loewner order on matrices: $X \leq Y$ means $Y - X \geq 0$.

Definition 4.1.3 (Associated positive linear map). [Lean](#) [✓ Stmt](#) A positive matrix map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ determines the same linear map regarded as a positive linear map: if $A \leq B$, then $E(A) \leq E(B)$.

Definition 4.1.4 (Trace-preserving map). [Lean](#) [✓ Stmt](#) A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *trace-preserving* if $\text{tr}(E(X)) = \text{tr}(X)$ for all X .

Theorem 4.1.5 (Transposition is positive and trace-preserving). [Lean](#) [Lean](#) [✓ Stmt](#) The matrix transposition map $\theta(X) = X^T$ on $M_D(\mathbb{C})$ is positive and trace-preserving.

Proof. [✓ Proof](#) If $X \geq 0$, write $X = C^\dagger C$. Then $X^T = C^T \overline{C} = \overline{C}^\dagger \overline{C}$, so $X^T \geq 0$. Trace preservation is the identity $\text{tr}(X^T) = \text{tr}(X)$. $\square$

Definition 4.1.6 (Reduction map). [Lean](#) [✓ Stmt](#) The reduction map on $M_D(\mathbb{C})$, for $k \in \mathbb{N}$, is

$$T_k(X) = \text{tr}(X)\mathbb{1} - k^{-1}X.$$

Theorem 4.1.7 (The first reduction map is positive). [Lean](#) [✓ Stmt](#) For $D \geq 1$, the map $T_1(X) = \text{tr}(X)\mathbb{1} - X$ on $M_D(\mathbb{C})$ is positive.

Proof. [✓ Proof](#) If $X \geq 0$, diagonalize X . Its eigenvalues $\lambda_1, \dots, \lambda_D$ are nonnegative, and for each j ,

$$\lambda_j \leq \sum_{i=1}^D \lambda_i = \text{tr}(X).$$

Therefore all eigenvalues of $\text{tr}(X)\mathbb{1} - X$ are nonnegative. $\square$

Theorem 4.1.8 (Self-duality of the reduction map). [Lean](#) [✓ Stmt](#) *The reduction map is self-dual for the trace pairing:*

$$\text{tr}(T_k(\rho)X) = \text{tr}(\rho T_k(X)).$$

Proof. [✓ Proof](#) Expand both sides using $T_k(Y) = \text{tr}(Y)\mathbb{1} - k^{-1}Y$ and bilinearity of the trace. Both sides are

$$\text{tr}(\rho) \text{tr}(X) - k^{-1} \text{tr}(\rho X).$$

□

Theorem 4.1.9 (Choi matrix of the reduction map). [Lean](#) [✓ Stmt](#) *For $D \geq 1$, the Choi matrix of $T_k(X) = \text{tr}(X)\mathbb{1} - k^{-1}X$ is*

$$\tau(T_k) = D^{-1}\mathbb{1} - k^{-1}|\Omega\rangle\langle\Omega|.$$

Proof. [✓ Proof](#) The trace term sends the (a, b) slice of $|\Omega\rangle\langle\Omega|$ to its trace times the identity, giving the contribution $D^{-1}\mathbb{1}$. The second term is the Choi matrix of k^{-1} times the identity map, namely $k^{-1}|\Omega\rangle\langle\Omega|$. □

Theorem 4.1.10 (Conjugation filters preserve positivity). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *For every $X \in M_D(\mathbb{C})$, the conjugation filter*

$$\rho \mapsto X\rho X^\dagger$$

is completely positive, hence positive. Therefore, if $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive, then $\rho \mapsto T(X\rho X^\dagger)$ is positive. If in addition $X^\dagger X = \mathbb{1}$, then the conjugation filter is trace-preserving and hence is a quantum channel.

Proof. [✓ Proof](#) Complete positivity follows from the single Kraus operator X . Positivity follows immediately. If $X^\dagger X = \mathbb{1}$, cyclicity of trace gives

$$\text{tr}(X\rho X^\dagger) = \text{tr}(X^\dagger X\rho) = \text{tr}(\rho).$$

□

Theorem 4.1.11 (Filtered trace-normalization criterion). [Lean](#) [✓ Stmt](#) *Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive map, and let $X \in M_D(\mathbb{C})$ satisfy*

$$X^\dagger T^*(\mathbb{1})X = \mathbb{1},$$

where T^ is the trace-pairing adjoint. Then*

$$\rho \mapsto T(X\rho X^\dagger)$$

is positive and trace-preserving.

This is the algebraic trace-normalization step in the proof of [Wol12, Chapter 3, Lemma “Making positive maps trace preserving”]. The inverse-square-root choice is recorded in the next theorem.

Proof. [✓ Proof](#) Positivity follows because $X\rho X^\dagger$ is positive whenever ρ is positive, and T is positive. For trace preservation, use the trace-pairing adjoint identity and cyclicity of trace:

$$\text{tr}(T(X\rho X^\dagger)) = \text{tr}(T^*(\mathbb{1})X\rho X^\dagger) = \text{tr}(X^\dagger T^*(\mathbb{1})X\rho) = \text{tr}(\rho).$$

□

Lemma 4.1.12 (Normalizing conjugation by inverse square root). [Lean](#) [✓ Stmt](#) Let $A \in M_D(\mathbb{C})$ be positive definite and set $S = A^{1/2}$. Then

$$(S^{-1})^\dagger A S^{-1} = \mathbb{1}.$$

Proof. [✓ Proof](#) Since A is positive definite, S is invertible and self-adjoint. Moreover $S^2 = A$. Therefore

$$(S^{-1})^\dagger A S^{-1} = S^{-1} S^2 S^{-1} = \mathbb{1}.$$

□

Theorem 4.1.13 (Trace normalization by inverse square root). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive map such that $T^*(\mathbb{1})$ is positive definite. Put

$$X = (T^*(\mathbb{1}))^{-1/2}.$$

Then

$$\rho \mapsto T(X\rho X^\dagger)$$

is positive and trace-preserving.

Proof. [✓ Proof](#) Let $S = (T^*(\mathbb{1}))^{1/2}$. Since $T^*(\mathbb{1})$ is positive definite, S is invertible and self-adjoint, and $S^2 = T^*(\mathbb{1})$. For $X = S^{-1}$,

$$X^\dagger T^*(\mathbb{1}) X = S^{-1} S^2 S^{-1} = \mathbb{1}.$$

The preceding trace-normalization criterion applies.

□

Theorem 4.1.14 (Forward Lorentz-cone trace inequality). [Lean](#) [✓ Stmt](#) If $A \in M_D(\mathbb{C})$ is positive semidefinite, then

$$\operatorname{Re} \operatorname{tr}(A^2) \leq (\operatorname{Re} \operatorname{tr}(A))^2.$$

Proof. [✓ Proof](#) Diagonalize A with nonnegative eigenvalues λ_i . Then $\operatorname{tr}(A^2) = \sum_i \lambda_i^2$ and $\operatorname{tr}(A) = \sum_i \lambda_i$, so the desired inequality is

$$\sum_i \lambda_i^2 \leq \left(\sum_i \lambda_i \right)^2,$$

which follows because all mixed products are nonnegative.

□

Theorem 4.1.15 (Trace-nonnegative Lorentz-cone converse). [Lean](#) [✓ Stmt](#) Let $A \in M_D(\mathbb{C})$ be Hermitian and suppose that $\operatorname{Re} \operatorname{tr}(A) \geq 0$. If

$$(D-1) \operatorname{Re} \operatorname{tr}(A^2) \leq (\operatorname{Re} \operatorname{tr}(A))^2,$$

then A is positive semidefinite.

This is the trace-nonnegative form of the converse in [Wol12, Proposition 3.9]. The unrestricted squared implication printed there is false without a trace sign condition; this is recorded in `docs/paper-gaps/wolf_ch3_lorentz_cone_trace_sign.tex`.

Proof. ✓ Proof Diagonalize A with real eigenvalues λ_i . Suppose that some eigenvalue λ_0 is negative. Let

$$T = \sum_i \lambda_i, \quad R = \sum_{i \neq 0} \lambda_i.$$

Since $T \geq 0$ and $T = \lambda_0 + R$, one has $T < R$, and hence $T^2 < R^2$. Cauchy's inequality gives

$$R^2 \leq (D-1) \sum_{i \neq 0} \lambda_i^2 \leq (D-1) \sum_i \lambda_i^2,$$

contradicting the assumed Lorentz inequality. Thus all eigenvalues are nonnegative. $\square$

Definition 4.1.16 (Sum of the largest eigenvalues). Lean ✓ Stmt Let $A \in M_D(\mathbb{C})$ be Hermitian with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_D$ listed in decreasing order. For $0 \leq k \leq D$ set

$$S_k(A) = \sum_{i=1}^k \lambda_i,$$

the sum of the k largest (signed) eigenvalues. Indices beyond D contribute zero, so the value stabilizes at $\text{Re tr}(A)$ once k reaches D .

The *Ky-Fan k -norm* $\|A\|_{(k)}$ is by definition the sum of the k largest *singular values* of A . It agrees with $S_k(A)$ exactly when A is positive semidefinite, where eigenvalues and singular values coincide; for an indefinite Hermitian A the two differ (for $A = \text{diag}(1, -2)$ one has $S_1(A) = 1$ but $\|A\|_{(1)} = 2$). The results below concern $S_k(A)$ for arbitrary Hermitian A , so on the positive-semidefinite cone they are statements about the Ky-Fan norm.

Theorem 4.1.17 (Achievability of the largest-eigenvalue sum). Lean ✓ Stmt For every Hermitian $A \in M_D(\mathbb{C})$ and every $k \geq 0$ there is an orthogonal projection P of rank $\min(k, D)$ with

$$\text{Re tr}(PA) = S_k(A).$$

Proof. ✓ Proof Diagonalize $A = U \text{diag}(\lambda_i) U^\dagger$ and take P to be the projection onto the span of the eigenvectors with the $\min(k, D)$ largest eigenvalues. Then $\text{Re tr}(PA) = \sum_{i \leq k} \lambda_i = S_k(A)$. $\square$

Theorem 4.1.18 (Upper bound for the largest-eigenvalue sum). Lean ✓ Stmt For every Hermitian $A \in M_D(\mathbb{C})$ and every orthogonal projection P of rank $k < D$,

$$\text{Re tr}(PA) \leq S_k(A).$$

Proof. ✓ Proof Conjugating P by U gives an orthogonal projection Q of the same rank, and $\text{Re tr}(PA) = \sum_i w_i \lambda_i$ where w_i are the diagonal entries of Q . Each w_i lies in $[0, 1]$ and the weights sum to $\text{tr } P = k$. With the threshold $c = \lambda_{k+1}$,

$$\sum_{i \leq k} \lambda_i - \sum_i w_i \lambda_i = \sum_{i \leq k} (\lambda_i - c)(1 - w_i) + \sum_{i > k} (c - \lambda_i) w_i \geq 0,$$

since for $i \leq k$ the eigenvalues dominate c while $1 - w_i \geq 0$, and for $i > k$ they are dominated by c while $w_i \geq 0$. $\square$

Theorem 4.1.19 (Ky Fan's maximum principle). Lean ✓ Stmt Let $A \in M_D(\mathbb{C})$ be Hermitian and $0 \leq k < D$. Then the sum of its k largest eigenvalues is the largest value of $\text{Re tr}(PA)$ attained by an orthogonal projection P of rank k :

$$S_k(A) = \max_{P^2 = P = P^\dagger, \text{tr } P = k} \text{Re tr}(PA).$$

This is Ky Fan's maximum principle [Fan49]; see also [Bha97, Problem I.6.15, Exercise II.1.13]. For a positive semidefinite A , where $S_k(A)$ coincides with the Ky-Fan norm $\|A\|_{(k)}$, it gives the extremal overlap of [Wol12, Lemma 3.1].

Proof. ✓ Proof The achievability half (Theorem 4.1.17) provides a rank- k projection whose trace against A equals $S_k(A)$, so the value is attained. The upper bound (Theorem 4.1.18) shows that no rank- k projection exceeds it. Hence $S_k(A)$ is the maximum. $\square$

Definition 4.1.20 (Maps preserving the positive semidefinite cone). Lean ✓ Stmt A linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ maps the positive semidefinite cone onto itself if T is positive and every positive semidefinite matrix is the image under T of a positive semidefinite matrix. This is condition (1) of [Wol12, Proposition 3.8].

Theorem 4.1.21 (Conjugations preserve the positive semidefinite cone). Lean Lean ✓ Stmt Let $Y \in M_D(\mathbb{C})$ be invertible. Then the map $X \mapsto YXY^\dagger$ and the map $X \mapsto YX^TY^\dagger$ each map the positive semidefinite cone onto itself. This is the implication (3) $\Rightarrow$ (1) of [Wol12, Proposition 3.8].

Proof. ✓ Proof Conjugation by any matrix preserves positive semidefiniteness, and the transpose of a positive semidefinite matrix is positive semidefinite, so both maps send the cone into itself. For surjectivity onto the cone, given a positive semidefinite A set $W = Y^{-1}A(Y^{-1})^\dagger$, which is again positive semidefinite. Then $YWY^\dagger = A$ proves the conjugation case, and $Y(W^T)^TY^\dagger = A$ proves the transpose case, with W and W^T positive semidefinite. $\square$

Definition 4.1.22 (Completely positive map). Lean ✓ Stmt A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *completely positive* (CP) if it admits a *Kraus representation*: there exist operators $\{K_i\}_{i=0}^{r-1}$ with $K_i \in M_D(\mathbb{C})$ such that, for every $X \in M_D(\mathbb{C})$,

$$E(X) = \sum_{i=0}^{r-1} K_i X K_i^\dagger.$$

By Choi's theorem, this is equivalent to $E \otimes \text{id}_n$ being positive for all n .

Theorem 4.1.23 (Transposition is not completely positive). Lean ✓ Stmt If $D \geq 2$, the matrix transposition map $\theta(X) = X^T$ on $M_D(\mathbb{C})$ is not completely positive.

Proof. ✓ Proof If θ were completely positive, its Choi matrix would be positive semidefinite. By Theorem 16.2.3, this Choi matrix is $D^{-1}F$. For the antisymmetric vector $v = |01\rangle - |10\rangle$,

$$\langle v | D^{-1}F | v \rangle = -\frac{2}{D} < 0,$$

contradicting positive semidefiniteness. $\square$

Theorem 4.1.24 (Entrywise complete positivity from a Kraus representation). Lean ✓ Stmt Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be completely positive in the Kraus sense. For every k and every positive block matrix $M \in M_k(M_D(\mathbb{C}))$, the entrywise image

$$(E(M_{ab}))_{a,b}$$

is again positive in $M_k(M_D(\mathbb{C}))$.

Proof. [✓ Proof](#) Write $E(X) = \sum_i K_i X K_i^\dagger$. For each i , let d_i be the block-diagonal matrix with K_i on every diagonal block. Then

$$(E(M_{ab}))_{a,b} = \sum_i d_i M d_i^\dagger.$$

Each term on the right is positive because conjugation preserves positivity, and a finite sum of positive matrices is positive. $\square$

Theorem 4.1.25 (Kraus maps as abstract completely positive maps). [Lean](#) [✓ Stmt](#) *Every Kraus-represented completely positive map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ determines a completely positive map between the matrix C^* -algebras in the entrywise positivity sense.*

Proof. [✓ Proof](#) The defining entrywise positivity condition is exactly Theorem 4.1.24. $\square$

Theorem 4.1.26 (CP maps are positive). [Lean](#) [✓ Stmt](#) *Every completely positive map is positive.*

Proof. [✓ Proof](#) If $X \geq 0$, then each summand $K_i X K_i^\dagger \geq 0$ (conjugation preserves positive semidefiniteness), so the sum is PSD. $\square$

Theorem 4.1.27 (Channels are positive). [Lean](#) [✓ Stmt](#) *Every quantum channel is positive.*

Proof. [✓ Proof](#) A quantum channel is completely positive by definition, so Theorem 4.1.26 applies. $\square$

Theorem 4.1.28 (Positive maps preserve Hermiticity). [Lean](#) [✓ Stmt](#) *If E is positive and $X = X^\dagger$, then $E(X) = E(X)^\dagger$.*

Proof. [✓ Proof](#) Regard E as the associated positive linear map (Definition 4.1.3). For positive linear maps between matrix C^* -algebras, $E(Y^*) = E(Y)^*$. Since $X = X^\dagger$ means $X = X^*$, one obtains $E(X)^\dagger = E(X)^* = E(X^*) = E(X)$. $\square$

Definition 4.1.29 (Quantum channel). [Lean](#) [✓ Stmt](#) A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a *quantum channel* if it is both completely positive and trace-preserving (CPTP), following [Wol12].

Definition 4.1.30 (Density matrices). [Lean](#) [Lean](#) [✓ Stmt](#) The set of *density matrices* in $M_D(\mathbb{C})$ is

$$\mathcal{D}_D = \{\rho \in M_D(\mathbb{C}) : \rho \geq 0, \text{tr}(\rho) = 1\}.$$

Theorem 4.1.31 (Density matrices are compact). [Lean](#) [✓ Stmt](#) $\mathcal{D}_D$ is compact (in the entry-wise topology on $M_D(\mathbb{C})$).

Proof. [✓ Proof](#) The PSD cone is closed (preimage of the closed non-negative cone under continuous quadratic forms), and if $\rho \geq 0$ with $\text{tr}(\rho) = 1$ then each entry satisfies $|\rho_{ij}|^2 \leq \rho_{ii}\rho_{jj} \leq 1$: the diagonal entries are non-negative and sum to 1, and the off-diagonal bound is the PSD Cauchy–Schwarz inequality. Hence the matrix entries are uniformly bounded, and Heine–Borel gives compactness. $\square$

Theorem 4.1.32 (Density matrices are convex). [Lean](#) [✓ Stmt](#) $\mathcal{D}_D$ is convex.

Proof. [✓ Proof](#) Convex combination of PSD matrices is PSD, and trace is linear. $\square$

Theorem 4.1.33 (Density matrices are nonempty). [Lean](#) [✓ Stmt](#) For $D \geq 1$, $\mathcal{D}_D \neq \emptyset$.

Proof. [✓ Proof](#) $\frac{1}{D}\mathbb{1}_D$ is a density matrix. $\square$

Theorem 4.1.34 (Channels preserve density matrices). [Lean](#) [✓ Stmt](#) If E is a quantum channel, then $E(\mathcal{D}_D) \subseteq \mathcal{D}_D$.

Proof. [✓ Proof](#) Complete positivity implies positivity (Theorem 4.1.26), so $E(\rho) \geq 0$; trace preservation gives $\text{tr}(E(\rho)) = \text{tr}(\rho) = 1$. $\square$

4.2 Irreducibility

Definition 4.2.1 (Orthogonal projection). [Lean](#) [✓ Stmt](#) A matrix $P \in M_D(\mathbb{C})$ is an *orthogonal projection* if $P = P^\dagger$ and $P^2 = P$.

Theorem 4.2.2 (Orthogonal projections and involutive-algebra projections). [Lean](#) [Lean](#) [✓ Stmt](#) For a complex square matrix, the two descriptions $P = P^\dagger$, $P^2 = P$ and $P^* = P$, $P^2 = P$ are equivalent.

Proof. [✓ Proof](#) Since the involution on $M_D(\mathbb{C})$ is conjugate transpose, $P^* = P^\dagger$. Therefore

$$P^\dagger = P, P^2 = P \iff P^* = P, P^2 = P.$$

$\square$

Theorem 4.2.3 (Complement of an orthogonal projection). [Lean](#) [✓ Stmt](#) If $P \in M_D(\mathbb{C})$ is an orthogonal projection, then $\mathbb{1} - P$ is an orthogonal projection.

Proof. [✓ Proof](#) In the involutive algebra $M_D(\mathbb{C})$, P satisfies $P^\dagger = P$ and $P^2 = P$. Hence

$$(\mathbb{1} - P)^\dagger = \mathbb{1} - P, \quad (\mathbb{1} - P)^2 = \mathbb{1} - 2P + P^2 = \mathbb{1} - P.$$

Thus $\mathbb{1} - P$ is an orthogonal projection. $\square$

Definition 4.2.4 (Irreducible map). [Lean](#) [✓ Stmt](#) A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *irreducible* if whenever P is an orthogonal projection satisfying $E(PM_D(\mathbb{C})P) \subseteq PM_D(\mathbb{C})P$, then $P = 0$ or $P = \mathbb{1}$. This is [Wol12, Theorem 6.2(1)]. The definition applies to any linear map; complete positivity is not required.

Theorem 4.2.5 (Injectivity implies irreducibility). [Lean](#) [✓ Stmt](#) If A is an injective MPS tensor, then its transfer map $\mathcal{E}_A$ is irreducible.

Proof. [✓ Proof](#) If P is an invariant projection for $\mathcal{E}_A$, then $(\mathbb{1} - P)A^iP = 0$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, $(\mathbb{1} - P)MP = 0$ for all M , so $P = 0$ or $P = \mathbb{1}$. $\square$

Theorem 4.2.6 (Transfer map is CP). [Lean](#) [✓ Stmt](#) The transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is completely positive.

Proof. [✓ Proof](#) It is already written in Kraus form, with Kraus operators $\{A^i\}$. $\square$

Theorem 4.2.7 (Transfer map is a channel). [Lean](#) [✓ Stmt](#) If $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, then the transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is a quantum channel (CPTP).

Proof. [✓ Proof](#) Complete positivity is the observation of Theorem 4.2.6. Trace preservation: $\text{tr}(\mathcal{E}_A(X)) = \sum_i \text{tr}(A^i X (A^i)^\dagger) = \text{tr}((\sum_i (A^i)^\dagger A^i)X) = \text{tr}(X)$ by cyclicity and the normalization hypothesis. $\square$

4.3 Cesàro fixed-point theorem

Definition 4.3.1 (Cesàro mean). [Lean](#) [✓ Stmt](#) The *Cesàro mean* of a linear map E at order N , applied to a starting matrix ρ_0 , is

$$\sigma_N = \frac{1}{N} \sum_{n=0}^{N-1} E^n(\rho_0).$$

Theorem 4.3.2 (Cesàro telescope). [Lean](#) [✓ Stmt](#) $E(\sigma_N) - \sigma_N = \frac{1}{N}(E^N(\rho_0) - \rho_0)$.

Proof. [✓ Proof](#) Telescoping: $E(\sigma_N) = \frac{1}{N} \sum_{n=1}^N E^n(\rho_0)$ and subtracting σ_N cancels all but the first and last terms. $\square$

Theorem 4.3.3 (Hermitian fixed-point decomposition). [Lean](#) [✓ Stmt](#) Let E be a quantum channel and $X = E(X)$ a Hermitian fixed point with decomposition $X = P_+ - P_-$ into orthogonal positive and negative parts ($P_{\pm} \geq 0$, $\text{tr}(P_+ P_-) = 0$). Then $E(P_+) = P_+$ and $E(P_-) = P_-$. This is [Wol12, Proposition 6.8].

Proof. [✓ Proof](#) Let Q be the support projection of P_+ . Trace preservation and positivity give $\text{tr}(QE(P_-)) = 0$ and $E(P_-) \geq 0$, so $QE(P_-) = 0$. The fixed-point equation $E(P_+) - E(P_-) = P_+ - P_-$ then identifies $E(P_{\pm}) = P_{\pm}$. $\square$

Theorem 4.3.4 (Cesàro fixed-point theorem). [Lean](#) [✓ Stmt](#) Every quantum channel $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ (with $D \geq 1$) has a nonzero positive semidefinite fixed point: there exists $\rho \geq 0$, $\rho \neq 0$, with $E(\rho) = \rho$.

Proof. [✓ Proof](#) Start with any $\rho_0 \in \mathcal{D}_D$ (Theorem 4.1.33). Each Cesàro mean σ_N lies in $\mathcal{D}_D$ by applying channel preservation to every iterate, adding positive semidefinite matrices, rescaling by the positive average factor, and using trace linearity. By compactness (Theorem 4.1.31), extract a convergent subsequence $\sigma_{N_k} \rightarrow \rho$. The telescope identity (Theorem 4.3.2) gives $\|E(\sigma_N) - \sigma_N\| = \frac{1}{N} \|E^N(\rho_0) - \rho_0\| \rightarrow 0$, so $E(\rho) = \rho$ by continuity. In [Wol12, Theorem 6.11], existence is proved via Brouwer's fixed-point theorem; we use the Cesàro argument instead. $\square$

4.4 Fixed-point projection

Definition 4.4.1 (Fixed-point projection). [Lean](#) [✓ Stmt](#) Given a linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ and a fixed point ρ with $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$, the *fixed-point projection* is the rank-one map

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho.$$

This projects onto the span of ρ along the kernel of the trace functional. When the fixed point is unique up to scaling, this span is the full fixed-point space.

Theorem 4.4.2 (Power decomposition). [Lean](#) [✓ Stmt](#) If E is trace-preserving and $E(\rho) = \rho$, then for $n \geq 1$,

$$E^n = P + (E - P)^n,$$

where P is the fixed-point projection.

Proof. [✓ Proof](#) P is idempotent ($P^2 = P$), the fixed-point relation $E(\rho) = \rho$ gives $E \circ P = P$, and trace preservation gives $P \circ E = P$. These imply $(E - P) \circ P = 0$ and $P \circ (E - P) = 0$, so in the binomial expansion of $(P + (E - P))^n$ all cross terms vanish, leaving $E^n = P^n + (E - P)^n = P + (E - P)^n$. $\square$

4.5 Peripheral spectrum and primitivity

Definition 4.5.1 (Peripheral spectrum). [Lean](#) [✓ Stmt](#) The *peripheral spectrum* of a bounded linear operator T is the set of spectral values μ with $|\mu|$ equal to the spectral radius of T .

Definition 4.5.2 (Peripheral eigenvalues). [Lean](#) [✓ Stmt](#) The *peripheral eigenvalues* of a linear map E on a finite-dimensional space are the eigenvalues λ on the unit circle: $|\lambda| = 1$. For a channel, the spectral radius is 1 [Wol12, Proposition 6.1], so these coincide with the eigenvalues of maximal modulus.

The condition $|\lambda| = 1$ is appropriate for channels, since the spectral radius of a channel is 1; for a general linear map with spectral radius $r \neq 1$ the condition would generalise to $|\lambda| = r$.

Remark 4.5.3 (Peripheral conventions for channels). [✓ Stmt](#) The spectral-radius convention of Definition 4.5.1 is the usual bounded-operator notion. For channels the spectral radius is 1, so the channel arguments below use the unit-circle set of Definition 4.5.2.

Theorem 4.5.4 (1 is a peripheral eigenvalue). [Lean](#) [✓ Stmt](#) If E has a nonzero fixed point $\rho \neq 0$ with $E(\rho) = \rho$, then 1 is a peripheral eigenvalue of E .

Proof. [✓ Proof](#) ρ is an eigenvector with eigenvalue 1, and $|1| = 1$. □

Theorem 4.5.5 (Finiteness of peripheral eigenvalues). [Lean](#) [✓ Stmt](#) On a finite-dimensional space, the set of peripheral eigenvalues is finite.

Proof. [✓ Proof](#) Peripheral eigenvalues are a subset of all eigenvalues, which is a finite set in finite dimensions. □

Theorem 4.5.6 (Power-stable peripheral eigenvalues are roots of unity). [Lean](#) [✓ Stmt](#) Let E be a linear endomorphism on a finite-dimensional space. If μ is a peripheral eigenvalue of E and μ^n is an eigenvalue of E for every $n \geq 1$, then μ is a root of unity.

Proof. [✓ Proof](#) Since μ^n is an eigenvalue for every positive n , the pigeonhole principle on the finite eigenvalue set gives $\mu^a = \mu^b$ for some $a \neq b$, hence $\mu^{|a-b|} = 1$. □

Theorem 4.5.7 (Peripheral powers imply root of unity). [Lean](#) [✓ Stmt](#) If the set of peripheral eigenvalues of a linear endomorphism E on a finite-dimensional space is closed under powers ($\mu \in \text{peripheral}(E)$ and $n \geq 1$ imply $\mu^n \in \text{peripheral}(E)$), then every peripheral eigenvalue is a root of unity.

Proof. [✓ Proof](#) The closure hypothesis provides $\mu^n \in \text{peripheral}(E)$ for all positive n , which in particular means μ^n is an eigenvalue. Theorem 4.5.6 then gives $\mu^p = 1$ for some $p > 0$. □

4.5.1 Periodicity removal by powering

Remark 4.5.1.1. [✓ Stmt](#) The results in this subsection are purely spectral: they use only finiteness of a set of complex numbers and the spectral mapping theorem for powers. In particular, they do not rely on complete positivity. For transfer maps, replacing a tensor by its p -blocked tensor replaces $\mathcal{E}_A$ by $\mathcal{E}_A^p$, so this powering step is the operator-theoretic form of blocking.

Lemma 4.5.1.2 (Common exponent for a finite set of roots of unity). [Lean](#) [✓ Stmt](#) Let $s \subseteq \mathbb{C}$ be a finite set. Assume that for every $\mu \in s$ there exists an exponent $p_\mu > 0$ with $\mu^{p_\mu} = 1$. Then there exists $p > 0$ such that $\mu^p = 1$ for all $\mu \in s$.

Proof. [✓ Proof](#) Take p to be the least common multiple of the exponents p_μ . □

Lemma 4.5.1.3 (Powering collapses peripheral eigenvalues). [Lean](#) [✓ Stmt](#) Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a linear map, and assume that E has a nonzero fixed point $\rho \neq 0$. Let $p > 0$. If every peripheral eigenvalue μ of E satisfies $\mu^p = 1$, then $\text{peripheral}(E^p) = \{1\}$.

Proof. [✓ Proof](#) The fixed point ρ gives $1 \in \text{peripheral}(E^p)$. For the reverse inclusion, let $\nu \in \text{peripheral}(E^p)$. Spectral mapping gives $\nu = \mu^p$ for some $\mu \in \sigma(E)$. From $|\nu| = 1$ and $p > 0$ we obtain $|\mu| = 1$, hence $\mu \in \text{peripheral}(E)$. The hypothesis $\mu^p = 1$ then forces $\nu = 1$. $\square$

For a normalized MPS tensor, the transfer map is naturally trace-preserving but need not be unital. In general one should not expect a single gauge choice to make it both unital and trace-preserving.

Remark 4.5.1.4 (Bi-canonical special case). [✓ Stmt](#) When a Kraus map happens to be both unital and trace-preserving (the “bi-canonical” case), the Kadison–Schwarz equality at peripheral eigenvectors (Theorem 4.5.2.6) shows directly that the peripheral eigenvalues are closed under powers, and hence are roots of unity by Theorem 4.5.7. It requires both normalizations simultaneously. The more general adjoint-fixed-point formulation below covers the case where only unitality and a positive definite adjoint fixed point are available.

4.5.2 The Kadison–Schwarz inequality and the multiplicative domain

Definition 4.5.2.1 (Kraus maps, adjoints, and normalizations). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For operators $\{K_i\}_{i=0}^{d-1} \subseteq M_D(\mathbb{C})$, the associated *Kraus map* and *adjoint Kraus map* are

$$E(X) = \sum_i K_i X K_i^\dagger, \quad E^*(X) = \sum_i K_i^\dagger X K_i.$$

The Kraus family is *unital* when $\sum_i K_i K_i^\dagger = \mathbb{1}$ and *trace-preserving* when $\sum_i K_i^\dagger K_i = \mathbb{1}$.

Theorem 4.5.2.2 (Kadison–Schwarz inequality). [Lean](#) [✓ Stmt](#) Let $E(X) = \sum_i K_i X K_i^\dagger$ be a unital Kraus map. Then for every $X \in M_D(\mathbb{C})$,

$$E(X^\dagger X) \geq E(X)^\dagger E(X).$$

This is [Wol12, Equation (5.2)].

Proof. [✓ Proof](#) Apply the unital completely positive map entrywise to the positive 2×2 block matrix $\begin{pmatrix} X^\dagger X & X^\dagger \\ X & \mathbb{1} \end{pmatrix}$ and take the Schur complement of the lower-right block. $\square$

Theorem 4.5.2.3 (Hilbert–Schmidt contraction). [Lean](#) [✓ Stmt](#) If a Kraus map is both unital and trace-preserving, then

$$\text{tr}(E(X)^\dagger E(X)) \leq \text{tr}(X^\dagger X)$$

for all $X \in M_D(\mathbb{C})$.

Proof. [✓ Proof](#) The Kadison–Schwarz gap is positive semidefinite by Theorem 4.5.2.2. Trace preservation then identifies the traces of its two terms. $\square$

Theorem 4.5.2.4 (KS gap decomposition). [Lean](#) [✓ Stmt](#) For a unital Kraus map,

$$E(X^\dagger X) - E(X)^\dagger E(X) = \sum_i R_i^\dagger R_i, \quad R_i := X K_i^\dagger - K_i^\dagger E(X).$$

Proof. [✓ Proof](#) Expand the right-hand side, distribute the products, and use $\sum_i K_i K_i^\dagger = \mathbb{1}$ to recover the Kadison–Schwarz gap. $\square$

Theorem 4.5.2.5 (KS equality implies Kraus commutation). [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then, for every i ,

$$XK_i^\dagger = K_i^\dagger E(X).$$

Proof. [✓ Proof](#) The Kadison–Schwarz gap is a sum of positive semidefinite terms $\sum_i R_i^\dagger R_i$. If the gap vanishes, each R_i must vanish. $\square$

Theorem 4.5.2.6 (KS equality for peripheral eigenvectors). [Lean](#) [✓ Stmt](#) Let E be a Kraus map that is both unital and trace-preserving. If $E(X) = \mu X$ with $|\mu| = 1$, then

$$E(X^\dagger X) = E(X)^\dagger E(X).$$

Proof. [✓ Proof](#) The Kadison–Schwarz gap is positive semidefinite by Theorem 4.5.2.2. Trace preservation and the relation $E(X) = \mu X$ with $|\mu| = 1$ show that its trace is zero, hence the gap vanishes. $\square$

Theorem 4.5.2.7 (Left multiplicative identity from KS equality). [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then for every $Y \in M_D(\mathbb{C})$,

$$E(X^\dagger Y) = E(X)^\dagger E(Y).$$

Proof. [✓ Proof](#) The commutation relation of Theorem 4.5.2.5 lets one move $X^\dagger$ past each Kraus operator inside the sum. $\square$

Definition 4.5.2.8 (Multiplicative domain). [Lean](#) [✓ Stmt](#) The *multiplicative domain* of a Kraus map E is the set of matrices that are simultaneously left- and right-multiplicative for E .

Theorem 4.5.2.9 (Multiplicative domain is a *-subalgebra). [Lean](#) [✓ Stmt](#) For a unital Kraus map, the multiplicative domain is a *-subalgebra of $M_D(\mathbb{C})$.

Proof. [✓ Proof](#) The multiplicative identities are stable under addition, multiplication, and adjoint, and they contain the identity matrix. $\square$

4.5.3 Peripheral closure via adjoint fixed point

The following results replace the trace-preservation hypothesis by the existence of a positive definite fixed point for the *adjoint* Kraus map, matching the weighted-trace argument in [CPGSV16, Appendix A].

Lemma 4.5.3.1 (Peripheral closure via adjoint fixed point). [Lean](#) [✓ Stmt](#) Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital ($\sum_i K_i K_i^\dagger = \mathbb{1}$), and assume:

1. the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point $\rho > 0$, and
2. E is irreducible.

Then $\text{peripheral}(E)$ is closed under powers: if $\mu \in \text{peripheral}(E)$ then $\mu^n \in \text{peripheral}(E)$ for every $n \in \mathbb{N}$.

Proof. ✓ Proof Let $E(X) = \mu X$ with $|\mu| = 1$, and set $G := E(X^\dagger X) - E(X)^\dagger E(X)$. By Kadison–Schwarz, $G \geq 0$. Since $E^*(\rho) = \rho$,

$$\text{tr}(\rho G) = \text{tr}(\rho E(X^\dagger X)) - \text{tr}(\rho E(X)^\dagger E(X)) = \text{tr}(E^*(\rho) X^\dagger X) - \text{tr}(\rho E(X)^\dagger E(X)) = \text{tr}(\rho X^\dagger X) - |\mu|^2 \text{tr}(\rho X^\dagger X) = 0.$$

Thus the trace of the Kadison–Schwarz gap against the adjoint fixed point vanishes. Because $\rho > 0$ and $G \geq 0$, this forces $G = 0$, so $E(X^\dagger X) = E(X)^\dagger E(X)$. Hence $X^\dagger X$ is a PSD fixed point, which is positive definite by irreducibility (Theorem 6.1.3). So X is invertible. By Theorem 4.5.2.5 (KS equality implies Kraus commutation), X commutes with each Kraus operator; iterating via the left multiplicative identity (Theorem 4.5.2.7) gives $E(X^n) = \mu^n X^n$, so μ^n is peripheral. $\square$

Theorem 4.5.3.2 (Roots of unity via adjoint fixed point). Lean ✓ Stmt Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital, and assume that the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point and that E is irreducible. Then every peripheral eigenvalue of E is a root of unity.

Proof. ✓ Proof Combine the closure-under-powers result (Lemma 4.5.3.1) with Theorem 4.5.7. $\square$

Remark 4.5.3.3. ✓ Stmt Lemma 4.5.3.1 and Theorem 4.5.3.2 cover the bi-canonical (unital + TP) case as well: trace preservation gives $E^*(1) = 1$, which is a positive definite fixed point of the adjoint. The adjoint-fixed-point formulation matches the argument in [CPGSV16, Appendix A], which uses a faithful invariant state rather than bi-canonicity. Theorem 4.5.3.2 establishes the root-of-unity conclusion in this adjoint-fixed-point formulation. In the same adjoint-fixed-point formulation, Chapter 7 proves the corresponding cyclic structure: the cyclic group of peripheral eigenvalues is Theorem 7.14.3, and the cyclic projection decomposition is Theorem 7.14.1.13.

Definition 4.5.3.4 (Channel period). Lean ✓ Stmt The *period* of a channel E is the number of peripheral eigenvalues (counted without multiplicity): $p(E) := |\text{peripheral}(E)|$.

Definition 4.5.3.5 (Primitive map). Lean ✓ Stmt A linear map is *primitive* if its only peripheral eigenvalue is $\lambda = 1$, i.e. $\text{peripheral}(E) = \{1\}$. For channels this corresponds to [Wol12, Theorem 6.7]. This definition applies to any linear endomorphism, not only to channels.

Theorem 4.5.3.6 (Primitivity equals period one). Lean ✓ Stmt A linear map with a nonzero fixed point is primitive if and only if its period is 1.

Proof. ✓ Proof If $\text{peripheral}(E) = \{1\}$, then $|\text{peripheral}(E)| = 1$. Conversely, if $|\text{peripheral}(E)| = 1$, then since $1 \in \text{peripheral}(E)$ (Theorem 4.5.4), the only element is 1. $\square$

Remark 4.5.3.7 (Period-one characterization). ✓ Stmt The period $p(E)$ counts peripheral eigenvalues without multiplicity. Theorem 4.5.3.6 is therefore the period-one characterization of primitivity. The spectral estimates below are stated directly with Definition 4.5.3.5, since they use the absence of peripheral eigenvalues other than 1, not a separate counting hypothesis.

Theorem 4.5.3.8 (Complementary transfer-map gap for primitive maps). Lean ✓ Stmt Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be trace-preserving, and let $\rho \neq 0$ satisfy $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$. Let

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$$

be the fixed-point projection associated to ρ . Assume that:

1. E is primitive;
2. every eigenvalue μ of E satisfies $|\mu| \leq 1$;
3. whenever $E(X) = X$ and $\text{tr}(X) = 0$, one has $X = 0$.

Then every eigenvalue ν of $E - P$ satisfies $|\nu| < 1$.

Proof. ✓ **Proof** Let $(E - P)(X) = \nu X$ with $X \neq 0$. Then

$$E(X) = \nu X + P(X).$$

If $\text{tr}(X) = 0$, then $P(X) = 0$, so $E(X) = \nu X$ and hence ν is an eigenvalue of E with $|\nu| \leq 1$. If $|\nu| = 1$, primitivity forces $\nu = 1$, so X is a trace-zero fixed point of E ; by assumption this implies $X = 0$, a contradiction. Thus $|\nu| < 1$.

If instead $\text{tr}(X) \neq 0$, trace preservation and $\text{tr}(P(X)) = \text{tr}(X)$ give

$$\text{tr}(X) = \text{tr}(E(X)) = \nu \text{tr}(X) + \text{tr}(P(X)) = \nu \text{tr}(X) + \text{tr}(X),$$

so $\nu \text{tr}(X) = 0$ and therefore $\nu = 0$. In either case, $|\nu| < 1$. □

4.6 Fixed-point algebra

Definition 4.6.1 (Kraus fixed points). Lean ✓ **Stmt** The *fixed-point set* of a Kraus map $E(X) = \sum_i K_i X K_i^\dagger$ is $\text{Fix}(E) = \{X \in M_D(\mathbb{C}) : E(X) = X\}$.

Theorem 4.6.2 (Fixed points lie in the multiplicative domain). Lean ✓ **Stmt** Let E be a trace-preserving Kraus map whose Schrödinger-picture map has a positive definite fixed point $\rho > 0$. Then every adjoint fixed point belongs to the multiplicative domain of the adjoint Kraus family.

Proof. ✓ **Proof** The Kadison–Schwarz equality for fixed points forces both one-sided multiplicative identities, so the fixed point lies in the full multiplicative domain. □

Theorem 4.6.3 (Fixed points form a $*$ -subalgebra). Lean ✓ **Stmt** Let E be a unital Kraus map whose adjoint map E^* has a positive definite fixed point $\rho > 0$. Then $\text{Fix}(E)$ is a $*$ -subalgebra of $M_D(\mathbb{C})$. This is [Wol12, Theorem 6.12].

Proof. ✓ **Proof** Closure under addition and scalar multiplication follows from linearity. Closure under $\dagger$: if $E(X) = X$ then $E(X^\dagger) = X^\dagger$ because fixed points are stable under adjoint. Closure under multiplication: if $E(X) = X$ and $E(Y) = Y$, the weighted Kadison–Schwarz argument gives $E(X^\dagger X) = E(X)^\dagger E(X)$ and, applied to $X^\dagger$, also $E(X X^\dagger) = E(X) E(X)^\dagger$. Thus X lies in the multiplicative domain, and Theorem 4.5.2.7 yields $E(XY) = E(X)E(Y) = XY$. □

Remark 4.6.4. ✓ **Stmt** Theorem 4.6.2 is the trace-preserving / Heisenberg-picture counterpart of the same multiplicative-domain argument.

Theorem 4.6.5 (Projected elements lie in the multiplicative domain). Lean ✓ **Stmt** Let E be a unital Kraus map whose adjoint map has a positive definite fixed point, and suppose $E^2 = E$. Then $E(X)$ lies in the multiplicative domain of E for every $X \in M_D(\mathbb{C})$.

Proof. [✓ Proof](#) Idempotence gives $E(E(X)) = E(X)$, so $E(X)$ is a fixed point. By the argument in Theorem 4.6.3, the Kadison–Schwarz gap for the fixed point $E(X)$ vanishes, hence

$$E(E(X)^\dagger E(X)) = E(X)^\dagger E(X).$$

Fixed points are closed under $\dagger$, so the same argument applied to $E(X)^\dagger$ gives

$$E(E(X)E(X)^\dagger) = E(X)E(X)^\dagger.$$

The vanishing of both Kadison–Schwarz gaps, again by the argument of Theorem 4.6.3, places $E(X)$ in the multiplicative domain. $\square$

Theorem 4.6.6 (Choi–Effros left absorption). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, for all $X, Y \in M_D(\mathbb{C})$,*

$$E(E(X)Y) = E(E(X)E(Y)).$$

Proof. [✓ Proof](#) The previous theorem places $E(X)$ in the multiplicative domain, so $E(E(X)Y) = E(E(X))E(Y) = E(X)E(Y)$. Applying the same multiplicativity to $E(X)$ and $E(Y)$ gives $E(E(X)E(Y)) = E(E(X))E(E(Y))$, and idempotence rewrites this as $E(X)E(Y)$. $\square$

Theorem 4.6.7 (Choi–Effros right absorption). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, for all $X, Y \in M_D(\mathbb{C})$,*

$$E(XE(Y)) = E(E(X)E(Y)).$$

Proof. [✓ Proof](#) The previous theorem places $E(Y)$ in the multiplicative domain, so $E(XE(Y)) = E(X)E(E(Y)) = E(X)E(Y)$. Applying the same multiplicativity to $E(X)$ and $E(Y)$ gives $E(E(X)E(Y)) = E(E(X))E(E(Y))$, and idempotence rewrites this as $E(X)E(Y)$. $\square$

Theorem 4.6.8 (Choi–Effros symmetric absorption). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, for all $X, Y \in M_D(\mathbb{C})$,*

$$E(E(X)Y) = E(XE(Y)).$$

Proof. [✓ Proof](#) Both sides equal $E(E(X)E(Y))$ by the two preceding theorems. $\square$

Theorem 4.6.9 (Choi–Effros projected product). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, for all $X, Y \in M_D(\mathbb{C})$,*

$$E(E(X)E(Y)) = E(X)E(Y).$$

Proof. [✓ Proof](#) The previous theorem places $E(X)$ and $E(Y)$ in the multiplicative domain. Therefore multiplicativity gives

$$E(E(X)E(Y)) = E(E(X))E(E(Y)).$$

Since E is idempotent under the standing hypotheses, $E(E(X)) = E(X)$ and $E(E(Y)) = E(Y)$, so

$$E(E(X)E(Y)) = E(X)E(Y).$$

$\square$

Definition 4.6.10 (Adjoint fixed points). [Lean](#) [✓ Stmt](#) The *adjoint fixed-point set* $\text{Fix}(E^*) = \{X \in M_D(\mathbb{C}) : E^*(X) = X\}$, where $E^*(X) = \sum_i K_i^\dagger X K_i$.

Theorem 4.6.11 (Fixed points form a $*$ -subalgebra in the Heisenberg picture). [Lean](#) [Lean](#) [✓ Stmt](#) *If the Kraus map is trace-preserving and has a positive definite fixed point $\rho > 0$ with $E(\rho) = \rho$, then $\text{Fix}(E^*)$ is a $*$ -subalgebra of $M_D(\mathbb{C})$.*

Proof. ✓ Proof Apply Theorem 4.6.3 to the adjoint family $\{K_i^\dagger\}$. □

Definition 4.6.12 (Kraus commutant). Lean Lean ✓ Stmt The *Kraus commutant* is $\{K_i, K_i^\dagger\}' = \{X \in M_D(\mathbb{C}) : [X, K_i] = [X, K_i^\dagger] = 0 \forall i\}$. It is a $*$ -subalgebra of $M_D(\mathbb{C})$.

Theorem 4.6.13 (Adjoint fixed points equal the Kraus commutant). Lean ✓ Stmt Under the hypotheses of Theorem 4.6.11, the adjoint fixed-point $*$ -subalgebra equals the Kraus commutant:

$$\text{Fix}(E^*) = \{K_i, K_i^\dagger\}'.$$

This is [Wol12, Theorem 6.13].

Proof. ✓ Proof ($\supseteq$): if X commutes with all K_i and $K_i^\dagger$, then $E^*(X) = \sum_i K_i^\dagger X K_i = X \sum_i K_i^\dagger K_i = X$ by trace preservation. ($\subseteq$): any $*$ -subalgebra inside $\text{Fix}(E^*)$ lies in the Kraus commutant, because for $X \in \text{Fix}(E^*)$ with $X^\dagger X \in \text{Fix}(E^*)$ the Kadison–Schwarz gap vanishes, which forces $XK_i = K_iX$ and $XK_i^\dagger = K_i^\dagger X$. □

Theorem 4.6.14 (Kraus commutant inside the adjoint fixed points). Lean ✓ Stmt Under the hypotheses above, the Kraus commutant is the largest $*$ -subalgebra contained in the adjoint fixed-point set.

Proof. ✓ Proof By Theorem 4.6.13, every $*$ -subalgebra inside the adjoint fixed points is automatically contained in the Kraus commutant, while the Kraus commutant itself is such a $*$ -subalgebra. □

Chapter 5

Schwarz Inequalities and Multiplicative Domains

This chapter develops the Schwarz-inequality theory of [Wol12, Chapter 5]: the Kadison–Schwarz inequality for Kraus maps, the multiplicative domain as a $*$ -subalgebra on which the map restricts to a $*$ -homomorphism, the extension to normal, subnormal, and commuting-dominant inputs for positive subunital maps, order and spectral-interval contractivity, and the transpose-trace map as a Schwarz map that is not completely positive. The peripheral-eigenvector and normal-block consequences feed the Fundamental Theorem of Chapter 11. The operator-convexity and Jensen material from [Wol12, Chapter 5] is treated separately in Chapter 18.

5.1 Kadison–Schwarz inequality

A linear map is completely positive (Definition 4.1.22) if and only if it has a Kraus form. The Kadison–Schwarz inequality and the multiplicative-domain characterisation are proved first for Kraus maps. The transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is a Kraus map (Theorem 4.2.6), so these results apply directly to transfer maps.

Definition 5.1.1 (Kraus map). Lean ✓ Stmt Given operators $\{K_i\}_{i=0}^{d-1}$ with $K_i \in M_D(\mathbb{C})$, the *Kraus map* is $E(X) = \sum_{i=0}^{d-1} K_i X K_i^\dagger$. A linear map is completely positive (Definition 4.1.22) if and only if it can be written in this form.

Definition 5.1.2 (Adjoint Kraus map). Lean ✓ Stmt The *adjoint Kraus map* is $E^*(X) = \sum_{i=0}^{d-1} K_i^\dagger X K_i$. When the $\{K_i\}$ are the matrices of an MPS tensor A , this is the transfer map of the conjugate-transposed family $i \mapsto (A^i)^\dagger$.

Definition 5.1.3 (Unital Kraus map). Lean ✓ Stmt A Kraus map is *unital* if $\sum_{i=0}^{d-1} K_i K_i^\dagger = \mathbb{1}$. Equivalently, $E(\mathbb{1}) = \mathbb{1}$.

Definition 5.1.4 (Trace-preserving Kraus map). Lean ✓ Stmt A Kraus map is *trace-preserving* if $\sum_{i=0}^{d-1} K_i^\dagger K_i = \mathbb{1}$. Equivalently, the adjoint Kraus map is unital. This is the standard MPS normalization condition. In the later gauge language it is the left-canonical condition, so Kadison–Schwarz arguments are often applied to the adjoint map.

Theorem 5.1.5 (Kadison–Schwarz inequality). [Lean](#) [✓ Stmt](#) Let $E(X) = \sum_i K_i X K_i^\dagger$ be a unital Kraus map (i.e. $\sum_i K_i K_i^\dagger = \mathbb{1}$). Then for all $X \in M_D(\mathbb{C})$,

$$E(X^\dagger X) \geq E(X)^\dagger E(X).$$

This is [Wol12, Equation (5.2)].

Proof. [✓ Proof](#) The 2×2 block matrix $\begin{bmatrix} X^\dagger X & X^\dagger \\ X & \mathbb{1} \end{bmatrix}$ is PSD ($= vv^\dagger$ with $v = [X^\dagger, \mathbb{1}]^T$). Applying the unital CP map blockwise preserves positive semidefiniteness (each block $K_i(\cdot)K_i^\dagger$ preserves PSD, and unitality ensures the $(2,2)$ -block maps to $\mathbb{1}$). The Schur complement of the $(2,2)$ -block $\mathbb{1}$ gives the result. $\square$

Theorem 5.1.6 (Kadison–Schwarz for the adjoint channel). [Lean](#) [✓ Stmt](#) If $\sum_i K_i^\dagger K_i = \mathbb{1}$ (trace-preserving), then the adjoint Kraus map satisfies $E^*(X^\dagger X) \geq E^*(X)^\dagger E^*(X)$.

Proof. [✓ Proof](#) The adjoint operators $\{K_i^\dagger\}$ satisfy $\sum_i K_i^\dagger (K_i^\dagger)^\dagger = \sum_i K_i^\dagger K_i = \mathbb{1}$, so they form a unital family. Apply Theorem 5.1.5 to $\{K_i^\dagger\}$. $\square$

5.1.1 Two-positive maps and the generalized Schwarz inequality

Definition 5.1.1.1 (k -positive map). [Lean](#) [✓ Stmt](#) A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is k -positive if $E \otimes \text{id}_k$ is positive on $M_D(\mathbb{C}) \otimes M_k(\mathbb{C})$.

Definition 5.1.1.2 (Two-positive map). [Lean](#) [✓ Stmt](#) A linear map is *two-positive* if it is 2-positive.

Definition 5.1.1.3 (Blockwise ampliation). [Lean](#) [✓ Stmt](#) For a linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, its k -fold ampliation acts on block matrices by

$$E^{(k)}(X)_{(i,p),(j,q)} = E((X_{(a,p),(b,q)})_{a,b})_{i,j}.$$

Theorem 5.1.1.4 (k -positivity as positivity of the ampliation). [Lean](#) [✓ Stmt](#) A linear map is k -positive if and only if its blockwise ampliation is positive on $M_D(\mathbb{C}) \otimes M_k(\mathbb{C})$.

Proof. [✓ Proof](#) This is precisely the preceding definition of the ampliation. $\square$

Definition 5.1.1.5 (Schmidt rank). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A vector $\psi \in \mathbb{C}^D \otimes \mathbb{C}^k$ is identified with its coefficient matrix $C_\psi \in M_{D,k}(\mathbb{C})$. Its Schmidt rank is

$$\text{SR}(\psi) = \text{rank}(C_\psi).$$

We write $\text{SR}(\psi) \leq r$ for the corresponding bounded-rank condition.

Theorem 5.1.1.6 (Basic bounds for Schmidt rank). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The zero vector has Schmidt rank zero, every vector satisfies

$$\text{SR}(\psi) \leq \min(D, k),$$

and every product vector $u \otimes v$ has Schmidt rank at most one.

Proof. [✓ Proof](#) The zero-vector claim is $\text{rank}(0) = 0$. For the dimension bounds, $\text{SR}(\psi) = \text{rank}(C_\psi) \leq D$ and $\text{SR}(\psi) = \text{rank}(C_\psi) \leq k$ follow from the row-rank and column-rank bounds on a $D \times k$ matrix, hence $\text{SR}(\psi) \leq \min(D, k)$. For a product vector, each column of $C_{u \otimes v}$ is a scalar multiple of u , so the column span has dimension at most one. $\square$

Definition 5.1.1.7 (Schmidt singular values). [Lean](#) [✓ Stmt](#) The Schmidt singular values of a vector $\psi \in \mathbb{C}^D \otimes \mathbb{C}^k$ are the singular values of the coefficient matrix C_ψ , viewed as a linear map $\mathbb{C}^k \rightarrow \mathbb{C}^D$.

Theorem 5.1.1.8 (Schmidt singular values and Schmidt rank). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The Schmidt singular values are indexed from zero. The nonzero Schmidt singular values of ψ are exactly those with index below $\text{SR}(\psi)$. Equivalently, for zero-based indexing,

$$s_k(\psi) = 0 \iff \text{SR}(\psi) \leq k.$$

Proof. [✓ Proof](#) For a finite-dimensional linear map, the support of the singular-value sequence is the initial interval whose length is the dimension of the range. The coefficient matrix C_ψ represents the corresponding map $\mathbb{C}^k \rightarrow \mathbb{C}^D$, and this range dimension is $\text{rank}(C_\psi)$. $\square$

Lemma 5.1.1.9 (Monotonicity of bounded Schmidt rank). [Lean](#) [✓ Stmt](#) If $\text{SR}(\psi) \leq k$ and $k \leq l$, then $\text{SR}(\psi) \leq l$.

Lemma 5.1.1.10 (Rank factorization through a coordinate space). [Lean](#) [✓ Stmt](#) If $A \in M_{m,n}(\mathbb{C})(\mathbb{C})$ has rank at most k , then there are matrices $B \in M_{m,k}(\mathbb{C})(\mathbb{C})$ and $C \in M_{k,n}(\mathbb{C})(\mathbb{C})$ such that $BC = A$.

Proof. [✓ Proof](#) The range of the linear map represented by A has dimension at most k . Embed this range into $\mathbb{C}^k$, compose with the range map, and then map back to the ambient codomain. The corresponding matrices give the desired factorization. $\square$

Theorem 5.1.1.11 (Maximal overlap with a fixed Schmidt rank). [Lean](#) [✓ Stmt](#) Let $\phi \in \mathbb{C}^D \otimes \mathbb{C}^k$ be a normalized vector with reduced density matrix $\rho = \text{tr}_2 |\phi\rangle\langle\phi|$ on the first factor, and let $1 \leq n < D$. The largest squared overlap $|\langle\phi|\psi\rangle|^2$ attained by a normalized vector ψ of Schmidt rank at most n is the Ky-Fan n -norm of ρ :

$$\max_{\text{SR}(\psi) \leq n, \|\psi\|=1} |\langle\phi|\psi\rangle|^2 = \|\rho\|_{(n)}.$$

Because ρ is positive semidefinite, this norm is the sum of its n largest eigenvalues, which coincide there with its singular values. This is [Wol12, Lemma 3.1] for $1 \leq n < D$; the top index $n = D$, where the value is $\text{tr} \rho = 1$, is Theorem 5.1.1.12.

Proof. [✓ Proof](#) Write C for the coefficient matrix of ϕ , so that $\rho = CC^\dagger$ and the overlap with a vector ψ of coefficient matrix B is the Frobenius pairing $\langle\phi|\psi\rangle = \text{tr}(C^\dagger B)$; normalization reads $\text{tr}(B^\dagger B) = 1$ and the Schmidt-rank constraint reads $\text{rank}(B) \leq n$.

For the upper bound, let P be the orthogonal projection onto the column space of B , so $\text{rank}(P) \leq n$ and $PB = B$. The Cauchy–Schwarz inequality for the Frobenius pairing gives

$$|\text{tr}(C^\dagger B)|^2 = |\text{tr}((PC)^\dagger B)|^2 \leq \text{tr}((PC)^\dagger (PC)) \text{tr}(B^\dagger B) = \text{tr}(P\rho).$$

Since P has rank at most n and ρ is positive semidefinite, the positive-semidefinite form of Ky Fan’s maximum principle (Theorem 4.1.19) gives

$$\text{tr}(P\rho) \leq \|\rho\|_{(n)},$$

so $|\langle\phi|\psi\rangle|^2 \leq \|\rho\|_{(n)}$ for every admissible ψ .

For the converse, let P_n be the eigenprojection of ρ onto its n largest eigenvalues, so $\text{tr}(P_n \rho) = \|\rho\|_{(n)}$, and set

$$B = \|\rho\|_{(n)}^{-1/2} P_n C.$$

Then $\text{rank}(B) \leq \text{rank}(P_n) = n$ and $\text{tr}(B^\dagger B) = \|\rho\|_{(n)}^{-1} \text{tr}(P_n \rho) = 1$, while

$$|\text{tr}(C^\dagger B)|^2 = \|\rho\|_{(n)}^{-1} |\text{tr}(P_n \rho)|^2 = \|\rho\|_{(n)},$$

so the bound is attained. (For a normalized ϕ with $1 \leq n$ one has $\|\rho\|_{(n)} \geq \|\rho\|_{(1)} > 0$, since $\text{tr} \rho = 1$, which justifies the normalization.) $\square$

Theorem 5.1.1.12 (Maximal overlap at the top Schmidt rank). Lean ✓ Stmt *At the top index $n = D$ the Schmidt-rank constraint $\text{SR}(\psi) \leq D$ is vacuous, so the largest squared overlap is the Ky-Fan D -norm of ρ ,*

$$\max_{\|\psi\|=1} |\langle \phi | \psi \rangle|^2 = \|\rho\|_{(D)},$$

attained at $\psi = \phi$. The Ky-Fan D -norm sums all eigenvalues, so for a normalized ϕ it collapses to $\|\rho\|_{(D)} = \text{tr} \rho = 1$. Together with Theorem 5.1.1.11 this covers the full range $1 \leq n \leq D$ of [Wol12, Lemma 3.1].

Proof. ✓ Proof The Ky-Fan D -norm sums all eigenvalues of ρ , so $\|\rho\|_{(D)} = \text{tr} \rho = \|\phi\|^2 = 1$. The bound 1 is attained at $\psi = \phi$, whose Schmidt rank is at most D ; and for any normalized ψ the Cauchy–Schwarz inequality gives $|\langle \phi | \psi \rangle|^2 \leq \|\phi\|^2 \|\psi\|^2 = 1$. $\square$

Definition 5.1.1.13 (Reduced density of an eigenvector). Lean ✓ Stmt Let τ be a Hermitian operator on $\mathbb{C}^{D'} \otimes \mathbb{C}^D$ with normalized eigenvectors ϕ_i . The reduced density operator of the i -th eigenvector on the first factor is

$$\rho_i = \text{tr}_2 |\phi_i\rangle\langle\phi_i|.$$

Lemma 5.1.1.14 (Coefficient-matrix form of the reduced density). Lean ✓ Stmt *Writing C_i for the coefficient matrix of ϕ_i , the reduced density operator factors as $\rho_i = C_i C_i^\dagger$.*

Proof. ✓ Proof The partial trace of $|\phi_i\rangle\langle\phi_i|$ over the second factor has entries $\sum_b (C_i)_{a,b} \overline{(C_i)_{a',b}}$, which is the (a, a') entry of $C_i C_i^\dagger$. $\square$

Lemma 5.1.1.15 (Reduced density is positive semidefinite). Lean ✓ Stmt *The reduced density operator ρ_i is positive semidefinite.*

Proof. ✓ Proof By the coefficient-matrix form $\rho_i = C_i C_i^\dagger$, which is positive semidefinite for any matrix C_i . $\square$

Lemma 5.1.1.16 (Eigenbasis Rayleigh expansion). Lean ✓ Stmt *For a Hermitian operator τ with eigenvalues ν_i and normalized eigenvectors ϕ_i , the expectation in a vector ψ expands as*

$$\langle \psi | \tau | \psi \rangle = \sum_i \nu_i |\langle \phi_i | \psi \rangle|^2.$$

Proof. ✓ Proof Diagonalizing $\tau = U \text{diag}(\nu_i) U^\dagger$ with U the unitary of eigenvectors, the change of variable $y = U^\dagger \psi$ gives $\langle \psi | \tau | \psi \rangle = \sum_i \nu_i |y_i|^2$, and $y_i = \langle \phi_i | \psi \rangle$ is the i -th eigenvector overlap. $\square$

Lemma 5.1.1.17 (Eigenbasis Parseval identity). Lean ✓ Stmt *For a Hermitian operator τ with normalized eigenvectors ϕ_i , the eigenvector overlaps recover the squared norm of ψ :*

$$\sum_i |\langle \phi_i | \psi \rangle|^2 = \|\psi\|^2.$$

Proof. ✓ Proof The eigenvectors form an orthonormal basis, so with $y = U^\dagger \psi$ one has $\sum_i |\langle \phi_i | \psi \rangle|^2 = \sum_i |y_i|^2 = \|y\|^2 = \|\psi\|^2$, the last equality because U is unitary. $\square$

Theorem 5.1.1.18 (Spectral lower bound for the Schmidt-rank expectation). [Lean](#) [✓ Stmt](#) Let τ be a Hermitian operator on $\mathbb{C}^{D'} \otimes \mathbb{C}^D$ with eigenvalues ν_i and normalized eigenvectors ϕ_i , and write $\rho_i = \text{tr}_2 |\phi_i\rangle\langle\phi_i|$ for the reduced density operator of ϕ_i on the first factor. Let $1 \leq n < D'$ and let $\nu_0 \geq 0$ be a lower bound for the positive eigenvalues (the smallest positive eigenvalue). Then every normalized vector ψ of Schmidt rank at most n satisfies

$$\nu_0 + \sum_{i:\nu_i \leq 0} (\nu_i - \nu_0) \|\rho_i\|_{(n)} \leq \langle \psi | \tau | \psi \rangle,$$

and hence the same bound holds for the infimum over such vectors. This is the lower bound of [Wol12, Chapter 3, Proposition 3.2] for $n < D'$; the top index $n = D'$ is Theorem 5.1.1.20.

Proof. [✓ Proof](#) In the eigenbasis the expectation expands as $\langle \psi | \tau | \psi \rangle = \sum_i \nu_i |\langle \phi_i | \psi \rangle|^2$, while Parseval's identity gives $\sum_i |\langle \phi_i | \psi \rangle|^2 = \|\psi\|^2 = 1$. Subtracting ν_0 times the latter from the former,

$$\langle \psi | \tau | \psi \rangle = \nu_0 + \sum_i (\nu_i - \nu_0) |\langle \phi_i | \psi \rangle|^2.$$

For an eigenvalue $\nu_i > 0$ the factor $\nu_i - \nu_0$ is nonnegative, so those terms only increase the sum and may be dropped. For an eigenvalue $\nu_i \leq 0$ the factor $\nu_i - \nu_0$ is nonpositive, and the maximal-overlap lemma (Theorem 5.1.1.11) bounds $|\langle \phi_i | \psi \rangle|^2 \leq \|\rho_i\|_{(n)}$; multiplying by the nonpositive factor reverses the inequality, which gives the claimed lower bound. $\square$

Theorem 5.1.1.19 (Spectral upper bound for the Schmidt-rank expectation). [Lean](#) [✓ Stmt](#) With τ , ϕ_i and ρ_i as above, let $1 \leq n < D'$ and suppose every eigenvalue of τ is at most ν . For any eigenvector index j there is a normalized vector ψ of Schmidt rank at most n with

$$\langle \psi | \tau | \psi \rangle \leq \nu + (\nu_j - \nu) \|\rho_j\|_{(n)},$$

and hence the infimum over such vectors lies below that value. Taking ν the largest positive eigenvalue and j the index of the unique non-positive eigenvalue ν_- recovers the upper bound $\nu + (\nu_- - \nu) \|\rho_- \|_{(n)}$ of [Wol12, Chapter 3, Proposition 3.2]. The statement is for $n < D'$; the top index $n = D'$ is Theorem 5.1.1.20.

Proof. [✓ Proof](#) Replacing ν_0 by ν in the eigenbasis expansion of the previous proof,

$$\langle \psi | \tau | \psi \rangle = \nu + \sum_i (\nu_i - \nu) |\langle \phi_i | \psi \rangle|^2,$$

every factor $\nu_i - \nu$ is nonpositive, so all terms are nonpositive and keeping only the $i = j$ term gives $\langle \psi | \tau | \psi \rangle \leq \nu + (\nu_j - \nu) |\langle \phi_j | \psi \rangle|^2$. The maximal-overlap lemma (Theorem 5.1.1.11) supplies a normalized ψ of Schmidt rank at most n attaining $|\langle \phi_j | \psi \rangle|^2 = \|\rho_j\|_{(n)}$; at that vector the right-hand side equals $\nu + (\nu_j - \nu) \|\rho_j\|_{(n)}$. $\square$

Theorem 5.1.1.20 (Spectral criterion at the top Schmidt rank). [Lean](#) [Lean](#) [✓ Stmt](#) At the top index $n = D'$ the Schmidt-rank constraint is vacuous and each Ky-Fan D' -norm collapses to $\|\rho_i\|_{(D')} = \text{tr} \rho_i = 1$, so both bounds (3.7) and (3.8) reduce to the Rayleigh characterization of the least eigenvalue: the lower bound (3.7) becomes $\nu_{\min} \leq \langle \psi | \tau | \psi \rangle$ for every normalized ψ , and the upper bound (3.8) on the infimum becomes the existence of a normalized eigenvector with $\langle \psi | \tau | \psi \rangle = \nu_{\min}$. Together they give

$$\inf_{\|\psi\|=1} \langle \psi | \tau | \psi \rangle = \nu_{\min},$$

with no Schmidt-rank restriction. Together with Theorems 5.1.1.18 and 5.1.1.19 this covers the full range $1 \leq n \leq D'$ of [Wol12, Chapter 3, Proposition 3.2].

Proof. ✓ Proof The eigenbasis expansion $\langle \psi | \tau | \psi \rangle = \sum_i \nu_i |\langle \phi_i | \psi \rangle|^2$ together with Parseval's identity $\sum_i |\langle \phi_i | \psi \rangle|^2 = 1$ bounds each weight ν_i below by $\nu_{\min}$, giving $\nu_{\min} \leq \langle \psi | \tau | \psi \rangle$. Evaluating at the eigenvector ϕ_j for a least-eigenvalue index j returns $\langle \phi_j | \tau | \phi_j \rangle = \nu_j = \nu_{\min}$, so the lower bound is attained and the infimum equals $\nu_{\min}$. $\square$

Theorem 5.1.1.21 (Rank-one test for k -positivity). Lean ✓ Stmt A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is k -positive if and only if, for every vector $\phi \in \mathbb{C}^D \otimes \mathbb{C}^k$,

$$E^{(k)}(|\phi\rangle\langle\phi|) \geq 0.$$

This is the pure-state reduction used in [Wol12, Chapter 3, Proposition 3.1].

Proof. ✓ Proof The forward implication is the definition of k -positivity, since $|\phi\rangle\langle\phi|$ is positive semidefinite. Conversely, every positive semidefinite matrix on $\mathbb{C}^D \otimes \mathbb{C}^k$ is a finite sum $\sum_i |\phi_i\rangle\langle\phi_i|$, and

$$E^{(k)}\left(\sum_i |\phi_i\rangle\langle\phi_i|\right) = \sum_i E^{(k)}(|\phi_i\rangle\langle\phi_i|) \geq 0$$

by linearity of $E^{(k)}$, the hypothesis, and closure of the positive semidefinite cone under finite sums. $\square$

Definition 5.1.1.22 (Right compression of the Choi matrix). Lean Lean Lean ✓ Stmt For $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and a Choi matrix τ , the right-factor compression in the blockwise ampliation index convention has entries

$$C_{(i,p),(j,q)} = \sum_{a,b} X_{a,p} \tau_{(i,a),(j,b)} \overline{X_{b,q}}.$$

The associated vector is the normalized vector with coefficients $D^{-1/2} X_{i,p}$.

Definition 5.1.1.23 (Right tensor factor for Choi compression). Lean ✓ Stmt For $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$, let R_X be the matrix from $\mathbb{C}^D \otimes \mathbb{C}^D$ to $\mathbb{C}^D \otimes \mathbb{C}^k$ with entries

$$(R_X)_{(i,p),(j,a)} = \delta_{i,j} X_{a,p}.$$

Theorem 5.1.1.24 (Right tensor sandwich formula). Lean ✓ Stmt If τ is the Choi matrix of T , then

$$R_X \tau R_X^\dagger = C_T(X),$$

where $C_T(X)$ is the right-factor Choi compression by X .

Proof. ✓ Proof The $(i,p), (j,q)$ entry of $R_X \tau R_X^\dagger$ is

$$\sum_{r,s,a,b} \delta_{i,r} X_{a,p} \tau_{(r,a),(s,b)} \delta_{j,s} \overline{X_{b,q}} = \sum_{a,b} X_{a,p} \tau_{(i,a),(j,b)} \overline{X_{b,q}},$$

which is the defining entry of $C_T(X)$. $\square$

Lemma 5.1.1.25 (Compressed maximally entangled vectors have bounded Schmidt rank). ✓ Stmt Lean For $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$, the vector

$$\psi_X = (D^{-1/2} X_{i,p})_{(i,p)} \in \mathbb{C}^D \otimes \mathbb{C}^k$$

has Schmidt rank at most k .

Proof. ✓ Proof The coefficient matrix of ψ_X is a scalar multiple of X . Therefore its rank is bounded by the number of columns, which is k . $\square$

Theorem 5.1.1.26 (Rank of a compressed maximally entangled vector). Lean ✓ Stmt Assume $D > 0$. For $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$, the vector

$$\psi_X = (D^{-1/2}X_{i,p})_{(i,p)} \in \mathbb{C}^D \otimes \mathbb{C}^k$$

satisfies

$$\text{SR}(\psi_X) = \text{rank } X.$$

Proof. ✓ Proof The coefficient matrix of ψ_X is $D^{-1/2}X$. Since $D > 0$, the scalar $D^{-1/2}$ is nonzero, and multiplication by this scalar preserves rank. $\square$

Theorem 5.1.1.27 (Square representative with exact Schmidt rank). Lean Lean ✓ Stmt Assume $D > 0$. For every $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ there is $X \in M_D(\mathbb{C})(\mathbb{C})$ such that

$$\psi_{(i,p)} = D^{-1/2}X_{i,p} \quad \text{and} \quad \text{rank } X = \text{SR}(\psi).$$

In particular, if $\text{SR}(\psi) \leq r$, then X may be chosen with $\text{rank } X \leq r$.

Proof. ✓ Proof Take $X_{i,p} = D^{1/2}\psi_{(i,p)}$. Since $D > 0$, this gives $D^{-1/2}X_{i,p} = \psi_{(i,p)}$. The rank identity follows from Theorem 5.1.1.26. The bounded-rank statement follows by comparison with the assumed Schmidt-rank bound. $\square$

Lemma 5.1.1.28 (Square parametrization of bounded Schmidt rank). Lean ✓ Stmt Assume $D > 0$. A vector $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ has Schmidt rank at most k if and only if there is a matrix $X \in M_D(\mathbb{C})(\mathbb{C})$ of rank at most k such that

$$\psi_{(i,j)} = D^{-1/2}X_{i,j}.$$

Proof. ✓ Proof The forward direction is the bounded-rank part of Theorem 5.1.1.27. Conversely, if $\psi_{(i,j)} = D^{-1/2}X_{i,j}$, then the coefficient matrix of ψ is

$$C_\psi = D^{-1/2}X,$$

so $\text{rank}(C_\psi) = \text{rank}(X)$, since multiplication by a nonzero scalar preserves rank. $\square$

Lemma 5.1.1.29 (Schmidt rank under the adjoint right tensor factor). Lean Lean ✓ Stmt Let $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and let $\eta \in \mathbb{C}^D \otimes \mathbb{C}^k$. Then $R_X^\dagger \eta \in \mathbb{C}^D \otimes \mathbb{C}^D$ has Schmidt rank at most k .

Proof. ✓ Proof If η is read as a $D \times k$ coefficient matrix E , then the coefficient matrix of $R_X^\dagger \eta$ is $EX^\dagger$. Therefore

$$\text{rank}(EX^\dagger) \leq \text{rank}(X^\dagger) = \text{rank}(X) \leq k.$$

$\square$

Lemma 5.1.1.30 (Right-tensor representatives of bounded Schmidt rank). Lean ✓ Stmt If $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ has Schmidt rank at most k , then there exist $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and $\eta \in \mathbb{C}^D \otimes \mathbb{C}^k$ such that

$$\psi = R_X^\dagger \eta.$$

Proof. ✓ Proof Let C_ψ be the coefficient matrix of ψ . The rank assumption gives a factorisation $C_\psi = BC$ through $\mathbb{C}^k$. Taking $X = C^\dagger$ and reading B as the coefficient matrix of η gives the required identity, because the coefficient matrix of $R_X^\dagger \eta$ is $BX^\dagger = BC$. $\square$

Theorem 5.1.1.31 (Quadratic form of a right Choi compression). Lean Lean ✓ Stmt For $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and $\eta \in \mathbb{C}^D \otimes \mathbb{C}^k$,

$$\langle \eta, C_T(X)\eta \rangle = \langle R_X^\dagger \eta, \tau_T R_X^\dagger \eta \rangle.$$

Proof. ✓ Proof The sandwich identity gives $C_T(X) = R_X \tau_T R_X^\dagger$. Hence

$$\langle \eta, R_X \tau_T R_X^\dagger \eta \rangle = \langle R_X^\dagger \eta, \tau_T R_X^\dagger \eta \rangle.$$

$\square$

Theorem 5.1.1.32 (Schmidt-rank Choi expectation from k -positivity). Lean ✓ Stmt Assume $D > 0$. If T is k -positive, then

$$\langle \psi, \tau_T \psi \rangle \geq 0$$

for every $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ with $\text{SR}(\psi) \leq k$.

Proof. ✓ Proof Write $\psi = R_X^\dagger \eta$ using Lemma 5.1.1.30. By Theorem 5.1.1.36, the compression $C_T(X)$ is positive semidefinite. Its quadratic form at η is therefore nonnegative, and Theorem 5.1.1.31 identifies this number with $\langle \psi, \tau_T \psi \rangle$. $\square$

Theorem 5.1.1.33 (Converse Schmidt-rank Choi test). Lean ✓ Stmt Assume $D > 0$. If

$$\langle \psi, \tau_T \psi \rangle \geq 0$$

for every $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ with $\text{SR}(\psi) \leq k$, then T is k -positive.

Proof. ✓ Proof By Theorem 5.1.1.36, it is enough to prove that $C_T(X)$ is positive semidefinite for each $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$. For any η , the quadratic-form identity identifies $\langle \eta, C_T(X)\eta \rangle$ with $\langle R_X^\dagger \eta, \tau_T R_X^\dagger \eta \rangle$, and Lemma 5.1.1.29 gives $\text{SR}(R_X^\dagger \eta) \leq k$. Thus every quadratic form of $C_T(X)$ is nonnegative. Over a complex finite-dimensional space, polarization recovers Hermiticity from this real nonnegative quadratic form condition, so $C_T(X)$ is positive semidefinite. $\square$

Theorem 5.1.1.34 (Schmidt-rank Choi criterion). Lean ✓ Stmt Assume $D > 0$. Then T is k -positive if and only if

$$\langle \psi, \tau_T \psi \rangle \geq 0$$

for every $\psi \in \mathbb{C}^D \otimes \mathbb{C}^D$ with $\text{SR}(\psi) \leq k$.

Proof. ✓ Proof Combine Theorem 5.1.1.32 with Theorem 5.1.1.33. $\square$

Lemma 5.1.1.35 (Rank-one ampliation as Choi compression). Lean ✓ Stmt Applying the k -fold ampliation of T to the rank-one matrix $|\psi\rangle\langle\psi|$, where $\psi_{(a,p)} = D^{-1/2}X_{a,p}$, is exactly the right-factor compression of the Choi matrix of T by X .

Proof. ✓ Proof In entries, both sides are

$$\sum_{a,b} X_{a,p} T(D^{-1}E_{a,b})_{i,j} \overline{X_{b,q}}.$$

If $\psi_{(a,p)} = D^{-1/2}X_{a,p}$, then for fixed p, q the corresponding block of $|\psi\rangle\langle\psi|$ is

$$(|\psi\rangle\langle\psi|)_{(\cdot,p),(\cdot,q)} = D^{-1} \sum_{a,b} X_{a,p} \overline{X_{b,q}} E_{a,b}.$$

Hence

$$T\left(D^{-1} \sum_{a,b} X_{a,p} \overline{X_{b,q}} E_{a,b}\right)_{i,j} = D^{-1} \sum_{a,b} X_{a,p} \overline{X_{b,q}} (T(E_{a,b}))_{i,j} = \sum_{a,b} X_{a,p} \tau_{(i,a),(j,b)} \overline{X_{b,q}},$$

where the second equality uses $\tau_{(i,a),(j,b)} = D^{-1}(T(E_{a,b}))_{i,j}$. □

Theorem 5.1.1.36 (Rectangular Choi-compression test for k -positivity). Lean ✓ Stmt Assume $D > 0$. A linear map $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ is k -positive if and only if, for every $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$, the right-factor compression of the Choi matrix of T by X is positive semidefinite.

Proof. ✓ Proof The forward direction applies the rank-one k -positivity test to the normalized vector ψ_X with coefficients $(\psi_X)_{(i,p)} = D^{-1/2}X_{i,p}$, then uses Lemma 5.1.1.35 in the form

$$(T \otimes \text{id}_k)(|\psi_X\rangle\langle\psi_X|) = C_T(X).$$

Conversely, given a vector ψ on the amplified space, take $X_{i,p} = D^{1/2}\psi_{(i,p)}$. Since $D > 0$, this choice satisfies $D^{-1/2}X_{i,p} = \psi_{(i,p)}$. The same compression identity gives

$$(T \otimes \text{id}_k)(|\psi\rangle\langle\psi|) = C_T(X),$$

where $C_T(X)$ is the right-factor Choi compression. Positivity of $C_T(X)$ is therefore positivity of the ampliation on the rank-one matrix $|\psi\rangle\langle\psi|$, and the rank-one test gives k -positivity. □

Theorem 5.1.1.37 (Right tensor Choi sandwiches of a k -positive map). Lean ✓ Stmt Assume $D > 0$. If $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ is k -positive, then for every $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$,

$$R_X \tau R_X^\dagger \geq 0,$$

where τ is the Choi matrix of T .

Proof. ✓ Proof By Theorem 5.1.1.36, $C_T(X) \geq 0$ for every $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$. The sandwich formula Theorem 5.1.1.24 identifies $R_X \tau R_X^\dagger$ with $C_T(X)$. □

Lemma 5.1.1.38 (Right-tensor multiplication). Lean ✓ Stmt Let $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$. Then the right-tensor factors satisfy

$$R_X R_P = R_{PX}.$$

Proof. ✓ Proof This is an entrywise calculation: the Kronecker delta in the physical index forces the two right-tensor factors to have the same physical coordinate, and the remaining sum is the matrix product PX . □

Lemma 5.1.1.39 (Square sandwich implies fixed rectangular compression). [Lean](#) [✓ Stmt](#) Let $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ have Choi matrix τ . Suppose $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ and $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ satisfy $PX = X$. If

$$R_P \tau R_P^\dagger \geq 0,$$

then the rectangular right compression of τ by X is positive semidefinite.

Proof. [✓ Proof](#) Since $PX = X$, Lemma 5.1.1.38 gives $R_X R_P = R_X$. Therefore

$$R_X \tau R_X^\dagger = R_X (R_P \tau R_P^\dagger) R_X^\dagger,$$

which is positive because positive semidefiniteness is preserved under compression. The right-hand side is the rectangular compression by Theorem 5.1.1.24. $\square$

Lemma 5.1.1.40 (Rank- k projection fixing a rectangular right factor). [Lean](#) [✓ Stmt](#) Let $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$ and suppose $k \leq D$. Then there exists a Hermitian projection $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ of rank k such that

$$PX = X.$$

Proof. [✓ Proof](#) The column space of X has dimension at most k . Since $k \leq D$, extend this column space to a k -dimensional subspace $U \subseteq \mathbb{C}^D$. Let P be the orthogonal projection onto U . Then P is Hermitian and idempotent, its rank is $\dim U = k$, and it fixes each column of X . $\square$

Lemma 5.1.1.41 (Right-tensor factorization for $VV^\dagger$). [Lean](#) [✓ Stmt](#) For every $V \in M_{D,k}(\mathbb{C})(\mathbb{C})$, the right-tensor factor satisfies

$$R_{VV^\dagger} = R_V^\dagger R_V.$$

Proof. [✓ Proof](#) This is the entrywise calculation obtained by expanding the definition of R_V and summing over the middle physical index. $\square$

Lemma 5.1.1.42 (Factorized right-tensor Choi sandwich). [Lean](#) [✓ Stmt](#) Let $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ have Choi matrix τ . For every $V \in M_{D,k}(\mathbb{C})(\mathbb{C})$,

$$R_{VV^\dagger} \tau R_{VV^\dagger}^\dagger = R_V^\dagger (R_V \tau R_V^\dagger) R_V.$$

Proof. [✓ Proof](#) Substitute Lemma 5.1.1.41 on the left and reassociate the three matrix products. $\square$

Theorem 5.1.1.43 (Factorized right projections in the Choi criterion). [Lean](#) [✓ Stmt](#) Assume $D > 0$. Let $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ be k -positive, with Choi matrix τ . If a right-factor matrix has the form $P = VV^\dagger$ with $V \in M_{D,k}(\mathbb{C})(\mathbb{C})$, then

$$R_P \tau R_P^\dagger \geq 0.$$

Proof. [✓ Proof](#) In the right-tensor convention one has

$$R_{VV^\dagger} = R_V^\dagger R_V.$$

Therefore

$$R_{VV^\dagger} \tau R_{VV^\dagger}^\dagger = R_V^\dagger (R_V \tau R_V^\dagger) R_V.$$

The middle factor is positive by Theorem 5.1.1.37, and positive semidefiniteness is preserved under compression by R_V . $\square$

Lemma 5.1.1.44 (Rank- k Hermitian projection factorization). [Lean](#) [✓ Stmt](#) Let $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ be Hermitian, idempotent, and of rank k . Then there exists $V \in M_{D,k}(\mathbb{C})(\mathbb{C})$ such that

$$P = VV^\dagger.$$

Proof. [✓ Proof](#) Regard P as a linear operator on $\mathbb{C}^D$. Hermiticity and idempotence say that this operator is the orthogonal projection onto its range. Choose an orthonormal basis of the range, indexed by $\{0, \dots, k-1\}$ using the rank hypothesis, and let V be the matrix whose columns are these basis vectors. The standard rank-one expansion of an orthogonal projection gives $P = VV^\dagger$. $\square$

Theorem 5.1.1.45 (Rank- k right projections in the Choi criterion). [Lean](#) [✓ Stmt](#) Assume $D > 0$. Let $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ be k -positive, with Choi matrix τ . If $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ is Hermitian, satisfies $P^2 = P$, and has rank k , then

$$R_P \tau R_P^\dagger \geq 0.$$

This is the forward rank- k projection-compression direction of [Wol12, Chapter 3, Proposition 3.1].

Proof. [✓ Proof](#) By Lemma 5.1.1.44, write $P = VV^\dagger$ with $V \in M_{D,k}(\mathbb{C})(\mathbb{C})$. The conclusion is then exactly Theorem 5.1.1.43. $\square$

Theorem 5.1.1.46 (Projection-compression converse for the Choi criterion). [Lean](#) [✓ Stmt](#) Assume $D > 0$ and $k \leq D$. Let $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ have Choi matrix τ . Suppose that

$$R_P \tau R_P^\dagger \geq 0$$

for every Hermitian projection $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ of rank k . Then T is k -positive.

Proof. [✓ Proof](#) By the rectangular compression criterion, it is enough to prove positivity of the right-factor compression by an arbitrary $X \in M_{D,k}(\mathbb{C})(\mathbb{C})$. Choose a rank- k Hermitian projection P with $PX = X$ using Lemma 5.1.1.40. The assumed positivity of $R_P \tau R_P^\dagger$, together with Lemma 5.1.1.39, gives positivity of the rectangular compression by X . $\square$

Theorem 5.1.1.47 (Rank- k projection form of the Choi criterion). [Lean](#) [✓ Stmt](#) Assume $D > 0$ and $k \leq D$. A linear map $T : M_{D,D}(\mathbb{C})(\mathbb{C}) \rightarrow M_{D,D}(\mathbb{C})(\mathbb{C})$ is k -positive if and only if, for every Hermitian projection $P \in M_{D,D}(\mathbb{C})(\mathbb{C})$ of rank k ,

$$R_P \tau R_P^\dagger \geq 0,$$

where τ is the Choi matrix of T . This is the rank- k projection formulation of [Wol12, Chapter 3, Proposition 3.1].

Proof. [✓ Proof](#) The forward direction is Theorem 5.1.1.45. The reverse direction is Theorem 5.1.1.46. $\square$

Definition 5.1.1.48 (Trace-pairing adjoint). [Lean](#) [✓ Stmt](#) The trace-pairing adjoint E^* of a linear map on matrices is the adjoint for the bilinear pairing $(A, B) \mapsto \text{tr}(AB)$.

Theorem 5.1.1.49 (Trace-pairing adjoint identity). [Lean](#) [✓ Stmt](#) For every linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ and all matrices ρ and X , one has

$$\text{tr}(E^*(\rho)X) = \text{tr}(\rho E(X)).$$

Proof. ✓ Proof Write $X = \sum_{i,j} X_{ij} e_{ij}$ in the standard matrix units and use linearity in X . For $X = e_{ij}$ one has $\text{tr}(E^*(\rho) e_{ij}) = (E^*(\rho))_{ji}$, while the definition of E^* gives $(E^*(\rho))_{ji} = \text{tr}(\rho E(e_{ij}))$. $\square$

Theorem 5.1.1.50 (Trace-pairing adjoint is involutive). Lean ✓ Stmt *For every linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the trace-pairing adjoint of E^* is E .*

Proof. ✓ Proof Fix an arbitrary matrix $N \in M_D(\mathbb{C})$. The adjoint identity $\text{tr}(E^*(\rho)X) = \text{tr}(\rho E(X))$, applied first to E^* and then to E , gives

$$\text{tr}((E^*)^*(X)N) = \text{tr}(XE^*(N)) = \text{tr}(E^*(N)X) = \text{tr}(NE(X)) = \text{tr}(E(X)N).$$

Since this holds for every N and the trace pairing is nondegenerate, $(E^*)^* = E$. $\square$

Theorem 5.1.1.51 (Self-duality of the positive semidefinite cone). Lean ✓ Stmt *Let $A \in M_D(\mathbb{C})$ be Hermitian. If $\text{tr}(AB) \geq 0$ for every positive semidefinite matrix B , then $A \geq 0$.*

Proof. ✓ Proof Test the hypothesis on rank-one positive semidefinite matrices $B = xx^\dagger$. The identity $\text{tr}(Axx^\dagger) = x^\dagger Ax$ and the hypothesis give $x^\dagger Ax \geq 0$ for every vector x , so the Hermitian matrix A is positive semidefinite. $\square$

Theorem 5.1.1.52 (Trace-pairing adjoint of a positive map). Lean ✓ Stmt *If $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive, then its trace-pairing adjoint E^* is positive.*

Proof. ✓ Proof Let $\rho \geq 0$. First $E^*(\rho)$ is Hermitian. Indeed, positivity of E gives $E(A^\dagger) = E(A)^\dagger$, and hence for matrix units e_{ij} ,

$$\overline{\text{tr}(\rho E(e_{ij}))} = \text{tr}(E(e_{ij})^\dagger \rho) = \text{tr}(E(e_{ji}) \rho) = \text{tr}(\rho E(e_{ji})).$$

For every $B \geq 0$, the adjoint identity gives

$$\text{tr}(E^*(\rho)B) = \text{tr}(\rho E(B)).$$

Since $E(B) \geq 0$, the right side is nonnegative by positivity of the trace product of two positive semidefinite matrices. Self-duality of the positive semidefinite cone therefore gives $E^*(\rho) \geq 0$. $\square$

Theorem 5.1.1.53 (Trace-pairing adjoints preserve positivity exactly). Lean ✓ Stmt *A linear map is positive if and only if its trace-pairing adjoint is positive.*

Proof. ✓ Proof One direction is the preceding theorem. The other follows by applying the same theorem to E^* and using $(E^*)^* = E$. $\square$

Theorem 5.1.1.54 (Trace-pairing adjoint of an ampliation). Lean ✓ Stmt *For every linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the trace-pairing adjoint commutes with ampliation:*

$$(E^{(k)})^* = (E^*)^{(k)}.$$

Proof. ✓ Proof For block matrices write ρ_{pq} and X_{pq} for their $M_D(\mathbb{C})$ -valued blocks. Expanding the trace by blocks gives

$$\text{tr}((E^*)^{(k)}(\rho)X) = \sum_{p,q} \text{tr}(E^*(\rho_{pq})X_{qp}).$$

The trace-adjoint identity changes each summand to $\text{tr}(\rho_{pq}E(X_{qp}))$, and summing the blocks back gives $\text{tr}(\rho E^{(k)}(X))$. Nondegeneracy of the trace pairing identifies the two adjoints. $\square$

Theorem 5.1.1.55 (Trace-pairing adjoint of a k -positive map). [Lean](#) [✓ Stmt](#) *If E is k -positive, then E^* is k -positive.*

Proof. [✓ Proof](#) Since E is k -positive, its ampliation $E^{(k)}$ is positive. The trace-pairing adjoint of this positive map is positive, and the preceding theorem identifies that adjoint with $(E^*)^{(k)}$. $\square$

Theorem 5.1.1.56 (k -positivity is trace-adjoint invariant). [Lean](#) [✓ Stmt](#) *A linear map is k -positive if and only if its trace-pairing adjoint is k -positive.*

Proof. [✓ Proof](#) One direction is the preceding theorem. The other follows by applying the same theorem to E^* and using $(E^*)^* = E$. $\square$

Theorem 5.1.1.57 (Completely positive maps are k -positive). [Lean](#) [✓ Stmt](#) *Every completely positive map is k -positive for all $k \geq 1$.*

Theorem 5.1.1.58 (One-positive maps are positive). [Lean](#) [✓ Stmt](#) *A linear map is 1-positive if and only if it is positive.*

Proof. [✓ Proof](#) The ampliation by $M_1(\mathbb{C})$ is identified with the original matrix algebra: all block indices have the unique value, so positivity of $E \otimes \text{id}_1$ is exactly positivity of E . $\square$

Theorem 5.1.1.59 (D -positive maps on nonzero $M_D(\mathbb{C})$ are completely positive). [Lean](#) [✓ Stmt](#) *For maps $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ with $D \geq 1$, complete positivity is equivalent to D -positivity.*

Proof. [✓ Proof](#) Complete positivity implies k -positivity for every k . Conversely, assume $D \geq 1$. If E is D -positive, apply $E \otimes \text{id}_D$ to the maximally entangled projector $|\Omega\rangle\langle\Omega|$. The result is the Choi matrix of E , hence it is positive semidefinite. Choi's theorem then gives complete positivity. $\square$

Theorem 5.1.1.60 ($(k+1)$ -positive implies k -positive). [Lean](#) [✓ Stmt](#) *Positivity under amplification is monotone in k .*

Theorem 5.1.1.61 (k -positive implies m -positive for $m \leq k$). [Lean](#) [✓ Stmt](#) *If a linear map is k -positive and $m \leq k$, then it is m -positive.*

Theorem 5.1.1.62 (Zero map is k -positive). [Lean](#) [✓ Stmt](#) *The zero map is k -positive.*

Theorem 5.1.1.63 (Sums of k -positive maps). [Lean](#) [✓ Stmt](#) *If E and F are k -positive, then $E + F$ is k -positive.*

Theorem 5.1.1.64 (Nonnegative multiples of k -positive maps). [Lean](#) [✓ Stmt](#) *If E is k -positive and $c \geq 0$, then cE is k -positive.*

Theorem 5.1.1.65 (Closedness of k -positive maps). [Lean](#) [✓ Stmt](#) *For each k , the set of k -positive linear maps is closed in the finite-dimensional space of linear maps.*

5.1.2 The map T_η and strictness of the chain

The cones of k -positive maps form a decreasing chain from the completely positive maps, at the top amplification dimension, down to the merely positive maps. This subsection shows that every inclusion between consecutive dimensions, below the top one, is strict. The witnesses come from a one-parameter family of maps whose positivity threshold is the parameter itself, so that tuning the parameter between two integers separates the two adjacent cones.

Definition 5.1.2.1 (The family T_η). [Lean](#) [✓ Stmt](#) For a real parameter η , the map T_η on $M_D(\mathbb{C})$ is

$$T_\eta(\rho) = \text{tr}(\rho) \mathbb{1} - \eta^{-1} \rho.$$

The map is defined for every real η ; at $\eta = 0$ it reduces to $\rho \mapsto \text{tr}(\rho) \mathbb{1}$, and the positivity threshold below assumes $\eta > 0$.

Theorem 5.1.2.2 (Choi matrix of T_η). [Lean](#) [✓ Stmt](#) For $D \geq 1$, the Choi matrix of T_η is

$$\tau_\eta = D^{-1} \mathbb{1} - \eta^{-1} |\Omega\rangle\langle\Omega|.$$

Proof. [✓ Proof](#) The Choi matrix is linear in the map, so it splits over the two terms of T_η . The trace term contributes

$$\tau(\rho \mapsto \text{tr}(\rho) \mathbb{1}) = D^{-1} \mathbb{1},$$

since each elementary slice of $|\Omega\rangle\langle\Omega|$ contributes its trace times the identity. The identity map contributes

$$\tau(\text{id}) = |\Omega\rangle\langle\Omega|.$$

Subtracting η^{-1} times the second from the first gives

$$\tau_\eta = D^{-1} \mathbb{1} - \eta^{-1} |\Omega\rangle\langle\Omega|.$$

□

Theorem 5.1.2.3 (Positivity threshold of T_η). [Lean](#) [✓ Stmt](#) Let $\eta > 0$ and $1 \leq k < D$. Then T_η is k -positive if and only if $\eta \geq k$. This is [Wol12, Equation (3.11)] for $1 \leq k < D$; the top index $k = D$, where k -positivity is complete positivity, is Theorem 5.1.2.4.

Proof. [✓ Proof](#) By the Schmidt-rank Choi criterion, k -positivity is the nonnegativity of the Choi quadratic form on all vectors of Schmidt rank at most k . For a normalized such vector ψ , the Choi matrix $\tau_\eta = D^{-1} \mathbb{1} - \eta^{-1} |\Omega\rangle\langle\Omega|$ gives

$$\langle\psi|\tau_\eta|\psi\rangle = D^{-1} - \eta^{-1} |\langle\Omega|\psi\rangle|^2.$$

The reduced density of $|\Omega\rangle$ on the first factor is $\mathbb{1}/D$, so by the maximal-overlap principle the supremum of $|\langle\Omega|\psi\rangle|^2$ over Schmidt-rank- k normalized vectors is $\|\mathbb{1}/D\|_{(k)} = k/D$. The infimum of the quadratic form is therefore $D^{-1} - \eta^{-1}(k/D) = D^{-1}(1 - k/\eta)$, which is nonnegative precisely when $\eta \geq k$. Degree-two homogeneity extends the bound from normalized vectors to all vectors of Schmidt rank at most k . □

Theorem 5.1.2.4 (Positivity threshold of T_η at the top index). [Lean](#) [✓ Stmt](#) Let $\eta > 0$ and $D \geq 1$. Then T_η is D -positive if and only if $\eta \geq D$. Together with Theorem 5.1.2.3 this is [Wol12, Equation (3.11)] over the full range $1 \leq k \leq D$, where the top index $k = D$ is complete positivity.

Proof. [✓ Proof](#) On $M_D(\mathbb{C})$, D -positivity is complete positivity, which is positive semidefiniteness of the Choi operator $\tau_\eta = D^{-1} \mathbb{1} - \eta^{-1} |\Omega\rangle\langle\Omega|$. Its quadratic form on a vector ψ is $D^{-1} \|\psi\|^2 - \eta^{-1} |\langle\Omega|\psi\rangle|^2$; since $|\Omega\rangle$ is normalized, Cauchy-Schwarz gives $|\langle\Omega|\psi\rangle|^2 \leq \|\psi\|^2$ with equality at $\psi = \Omega$, so the form is nonnegative for all ψ precisely when $\eta \geq D$. □

Corollary 5.1.2.5 (Strictness of the positivity chain). [Lean](#) [✓ Stmt](#) Let $1 \leq k$ with $k + 1 < D$. Then there is a map on $M_D(\mathbb{C})$ that is k -positive but not $(k + 1)$ -positive; the cone of k -positive maps strictly contains the cone of $(k + 1)$ -positive maps. Every inclusion between consecutive amplification dimensions below the top one is therefore strict.

Proof. [✓ Proof](#) Take T_η at $\eta = k$. By the threshold equivalence it is k -positive, since $\eta = k \geq k$, but not $(k + 1)$ -positive, since $\eta = k < k + 1$. □

5.1.3 The positivity hierarchy and Kadison–Schwarz from two-positivity

Theorem 5.1.3.1 (Completely positive implies two-positive). [Lean](#) [✓ Stmt](#) *Every completely positive map is two-positive.*

Theorem 5.1.3.2 (Two-positive implies positive). [Lean](#) [✓ Stmt](#) *Every two-positive map is positive.*

Definition 5.1.3.3 (Unital linear map). [Lean](#) [✓ Stmt](#) A linear map E is unital if $E(\mathbb{1}) = \mathbb{1}$.

Theorem 5.1.3.4 (Kadison–Schwarz for unital two-positive maps). [Lean](#) [✓ Stmt](#) *If E is unital and two-positive, then for all $X \in M_D(\mathbb{C})$,*

$$E(X^\dagger X) \geq E(X)^\dagger E(X).$$

Proof. Form the 2×2 block matrix $\begin{pmatrix} \mathbb{1} & X \\ X^\dagger & X^\dagger X \end{pmatrix} \geq 0$. Apply $E \otimes \text{id}_2$ (positive by 2-positivity) and read off the $(2, 2)$ -block Schur complement: $E(X^\dagger X) - E(X)^\dagger E(X) \geq 0$. $\square$

Theorem 5.1.3.5 (Kraus-form Kadison–Schwarz as a special case). [Lean](#) [✓ Stmt](#) *Let $K = (K_i)_{i=0}^{d-1}$ be a unital Kraus family, that is, $\sum_i K_i K_i^\dagger = \mathbb{1}$. Then for all $X \in M_D(\mathbb{C})$,*

$$\mathcal{K}(X^\dagger X) \geq \mathcal{K}(X)^\dagger \mathcal{K}(X),$$

where $\mathcal{K}(Y) = \sum_i K_i Y K_i^\dagger$.

5.1.4 Douglas-type factorization lemmas

Theorem 5.1.4.1 (Range inclusion implies right factorization). [Lean](#) [✓ Stmt](#) *If $\text{ran}(A) \subseteq \text{ran}(B)$, then there exists C such that $A = BC$.*

Proof. In finite dimensions, $\text{ran}(A) \subseteq \text{ran}(B)$ implies $A = B(B^\dagger B)^\dagger B^\dagger A$, where $(B^\dagger B)^\dagger$ denotes the Moore–Penrose pseudoinverse. Set $C = (B^\dagger B)^\dagger B^\dagger A$. $\square$

Theorem 5.1.4.2 (Vectorwise range criterion implies factorization). [Lean](#) [✓ Stmt](#) *If $Av \in \text{ran}(B)$ for every vector v , then there exists C with $A = BC$.*

Theorem 5.1.4.3 (Hilbert–Schmidt contraction). [Lean](#) [✓ Stmt](#) *If a Kraus map is both unital and trace-preserving, then $\text{tr}(E(X)^\dagger E(X)) \leq \text{tr}(X^\dagger X)$ for all $X \in M_D(\mathbb{C})$.*

Proof. [✓ Proof](#) By Theorem 5.1.5, $E(X^\dagger X) - E(X)^\dagger E(X) \geq 0$, so $\text{tr}(E(X^\dagger X)) \geq \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. $\square$

Lemma 5.1.4.4 (Trace pairing for Kraus map and adjoint map). [Lean](#) [✓ Stmt](#) *For all $X, Y \in M_D(\mathbb{C})$,*

$$\text{tr}(X E(Y)) = \text{tr}(E^*(X) Y).$$

Lemma 5.1.4.5 (Positive-definite trace test). [Lean](#) [✓ Stmt](#) *If $A > 0$, $X \geq 0$, and $\text{tr}(AX) = 0$, then $X = 0$.*

Theorem 5.1.4.6 (Fixed-point peripheral eigenvectors satisfy KS equality). [Lean](#) [✓ Stmt](#) *Let E be unital and let E^* be trace-preserving with a positive definite fixed point $\rho > 0$. If $E(X) = \mu X$ with $|\mu| = 1$, then*

$$E(X^\dagger X) = E(X)^\dagger E(X).$$

5.2 Multiplicative domain

Theorem 5.2.1 (KS gap decomposition). [Lean](#) [✓ Stmt](#) For a unital Kraus map E , the Kadison–Schwarz gap decomposes as

$$E(X^\dagger X) - E(X)^\dagger E(X) = \sum_{i=0}^{d-1} R_i^\dagger R_i,$$

where $R_i := XK_i^\dagger - K_i^\dagger E(X)$.

Proof. [✓ Proof](#) Expand the right-hand side $\sum_i (XK_i^\dagger - K_i^\dagger E(X))^\dagger (XK_i^\dagger - K_i^\dagger E(X))$, distribute, and use unitality $\sum_i K_i K_i^\dagger = \mathbb{1}$ to identify the result with the KS gap. $\square$

Theorem 5.2.2 (KS equality implies Kraus commutation). [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$ for some X , then $XK_i^\dagger = K_i^\dagger E(X)$ for all i .

Proof. [✓ Proof](#) By Theorem 5.2.1, the KS gap is $\sum_i R_i^\dagger R_i$ with $R_i = XK_i^\dagger - K_i^\dagger E(X)$. If the gap vanishes, $\sum_i R_i^\dagger R_i = 0$. Each summand $R_i^\dagger R_i$ is PSD, so each must be zero, giving $R_i = 0$. $\square$

Theorem 5.2.3 (KS equality for peripheral eigenvectors). [Lean](#) [✓ Stmt](#) Let E be a Kraus map that is both unital and trace-preserving. If $E(X) = \mu X$ with $|\mu| = 1$, then the KS gap vanishes: $E(X^\dagger X) = E(X)^\dagger E(X)$.

Proof. [✓ Proof](#) The gap $G := E(X^\dagger X) - E(X)^\dagger E(X)$ is PSD by Theorem 5.1.5. Its trace satisfies $\text{tr}(G) = \text{tr}(E(X^\dagger X)) - \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. Using $E(X) = \mu X$ and $|\mu| = 1$, $\text{tr}(E(X)^\dagger E(X)) = |\mu|^2 \text{tr}(X^\dagger X) = \text{tr}(X^\dagger X)$. So $\text{tr}(G) = 0$, and a PSD matrix with zero trace must be zero. $\square$

Theorem 5.2.4 (Left multiplicative identity from KS equality). [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then $E(X^\dagger Y) = E(X)^\dagger E(Y)$ for all $Y \in M_D(\mathbb{C})$.

Proof. [✓ Proof](#) From Theorem 5.2.2, $XK_i^\dagger = K_i^\dagger E(X)$ for all i . Taking conjugate transposes: $K_i X^\dagger = E(X)^\dagger K_i$. Then

$$\begin{aligned} E(X^\dagger Y) &= \sum_i K_i X^\dagger Y K_i^\dagger \\ &= \sum_i E(X)^\dagger K_i Y K_i^\dagger \\ &= E(X)^\dagger E(Y). \end{aligned}$$

$\square$

Theorem 5.2.5 (Right multiplicative identity from KS equality). [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then $E(YX) = E(Y)E(X)$ for all $Y \in M_D(\mathbb{C})$.

Proof. [✓ Proof](#) From Theorem 5.2.2, $XK_i^\dagger = K_i^\dagger E(X)$ for all i . Then

$$\begin{aligned} E(YX) &= \sum_i K_i Y X K_i^\dagger \\ &= \sum_i K_i Y K_i^\dagger E(X) \\ &= E(Y)E(X). \end{aligned}$$

$\square$

5.2.1 Full multiplicative-domain structure

Definition 5.2.1.1 (Right and left multiplicative domains). [Lean](#) [Lean](#) [✓ Stmt](#) For a Kraus map E , define

$$\mathcal{A}_R(E) := \{X \in M_D(\mathbb{C}) : \forall Y \in M_D(\mathbb{C}), E(XY) = E(X)E(Y)\},$$

and

$$\mathcal{A}_L(E) := \{X \in M_D(\mathbb{C}) : \forall Y \in M_D(\mathbb{C}), E(YX) = E(Y)E(X)\}.$$

We follow the convention that $\mathcal{A}_R(E)$ controls right multiplication and $\mathcal{A}_L(E)$ controls left multiplication. These are the two one-sided multiplicative domains from [Wol12, Equations (5.11)–(5.12)].

Definition 5.2.1.2 (Full multiplicative domain). [Lean](#) [✓ Stmt](#) The *multiplicative domain* of E is

$$\mathcal{A}(E) := \mathcal{A}_R(E) \cap \mathcal{A}_L(E).$$

Theorem 5.2.1.3 (Multiplicative-domain characterization). [Lean](#) [Lean](#) [✓ Stmt](#) Let E be a unital Kraus map. Then

$$X \in \mathcal{A}_R(E) \iff E(XX^\dagger) = E(X)E(X)^\dagger,$$

and

$$X \in \mathcal{A}_L(E) \iff E(X^\dagger X) = E(X)^\dagger E(X).$$

Together these are the two one-sided characterizations from [Wol12, Theorem 5.7].

Proof. [✓ Proof](#) If X lies in a one-sided multiplicative domain, evaluate the defining identity at $Y = X^\dagger$. Conversely, the left-domain implication is exactly Theorem 5.2.4, and the right-domain implication follows by applying Theorem 5.2.4 to $X^\dagger$. $\square$

Theorem 5.2.1.4 (One-sided multiplicative domains are subalgebras). [Lean](#) [Lean](#) [✓ Stmt](#) For a unital Kraus map E , both $\mathcal{A}_R(E)$ and $\mathcal{A}_L(E)$ are unital subalgebras of $M_D(\mathbb{C})$.

Proof. [✓ Proof](#) Closure under addition follows from linearity of E . Closure under multiplication: if $X, Y \in \mathcal{A}_R(E)$, then $E((XY)Z) = E(X)E(YZ) = E(X)E(Y)E(Z)$ for all Z , using Theorem 5.2.1.3 iteratively, and the $\mathcal{A}_L(E)$ case is analogous. Containment of $\mathbb{1}$ follows from unitality of E . $\square$

Theorem 5.2.1.5 (Multiplicative domain is a *-subalgebra). [Lean](#) [✓ Stmt](#) For a unital Kraus map E , the full multiplicative domain $\mathcal{A}(E)$ is a *-subalgebra of $M_D(\mathbb{C})$. This is the algebraic content of [Wol12, Theorem 5.7].

Proof. [✓ Proof](#) If $X \in \mathcal{A}(E) = \mathcal{A}_R(E) \cap \mathcal{A}_L(E)$, then $E(X^\dagger X) = E(X)^\dagger E(X)$ and $E(XX^\dagger) = E(X)E(X)^\dagger$. The characterization of Theorem 5.2.1.3 shows $X^\dagger \in \mathcal{A}(E)$. Combined with Theorem 5.2.1.4, this gives a *-subalgebra. $\square$

Theorem 5.2.1.6 (Restriction to multiplicative domain is a *-homomorphism). [Lean](#) [✓ Stmt](#) If E is unital, then the restricted map $E|_{\mathcal{A}(E)} : \mathcal{A}(E) \rightarrow M_D(\mathbb{C})$ is a *-algebra homomorphism. This is the homomorphism conclusion of [Wol12, Theorem 5.7].

Proof. [✓ Proof](#) Multiplicativity on $\mathcal{A}(E)$ follows from the definition. For $X \in \mathcal{A}(E)$, the Kraus commutation relation from Theorem 5.2.2 gives $E(X^\dagger) = E(X)^\dagger$, so the restriction preserves adjoints. $\square$

5.2.2 Schwarz inequality for normal operators

Theorem 5.2.2.1 (CP Schwarz inequality for normal operators). [Lean](#) [Lean](#) [✓ Stmt](#) Let $E^*(X) = \sum_i K_i^\dagger X K_i$ be the adjoint Kraus map of a trace-preserving Kraus family, and let $A \in M_D(\mathbb{C})$ be normal. Then the Schwarz gap

$$E^*(A^\dagger A) - E^*(A^\dagger)E^*(A) \quad (5.1)$$

is positive semidefinite. Equivalently,

$$E^*(A^\dagger)E^*(A) \leq E^*(A^\dagger A). \quad (5.2)$$

This is the completely positive case of [Wol12, Proposition 5.1].

Proof. [✓ Proof](#) The positivity statement is exactly Theorem 5.1.6. The second formulation (5.2) is the Loewner-order reformulation of the first (5.1). $\square$

Theorem 5.2.2.2 (Schwarz inequality for normal operators). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $T(\mathbb{1}) \leq \mathbb{1}$. If $A \in M_D(\mathbb{C})$ is normal, then

$$T(A^\dagger A) - T(A^\dagger)T(A) \geq 0.$$

This is [Wol12, Proposition 5.1].

Proof. [✓ Proof](#) Since A is normal, diagonalize $A = UDU^\dagger$ with D diagonal and U unitary. Express $A^\dagger A = U|D|^2U^\dagger$ as a sum $\sum_{i,j} \bar{d}_i d_j P_i P_j$ where $P_i = U e_i e_i^\dagger U^\dagger$. Each P_i is rank-one PSD, and the images $T(P_i)$ pairwise commute because they are images of mutually orthogonal projectors under a positive map. The Schwarz gap then reduces to a block quadratic form in the images, which is shown non-negative by a simultaneous-diagonalization argument: positivity of T ensures the scalar PSD matrix $(\langle v_k, T(P_i P_j) v_k \rangle)_{i,j}$ dominates the product $(\langle v_k, T(P_i) v_k \rangle \langle v_k, T(P_j) v_k \rangle)_{i,j}$, and combining over a common eigenbasis of the $T(P_i)$ yields the result. $\square$

Remark 5.2.2.3. The proof follows the positivity-on-abelian argument of [Wol12, Proposition 1.6]: a positive map restricted to the commutative $*$ -subalgebra generated by the spectral projectors of A satisfies the Schwarz inequality directly, without invoking the Arveson extension theorem.

5.2.3 Schwarz inequality for subnormal and commuting-dominant operators

Theorem 5.2.3.1 (Schwarz inequality for subnormal operators). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $T(\mathbb{1}) \leq \mathbb{1}$. If $A \in M_D(\mathbb{C})$ is subnormal (i.e. there exists a normal operator on a larger space whose compression to $\mathbb{C}^D$ is A), then

$$T(A^\dagger A) - T(A^\dagger)T(A) \geq 0.$$

This is [Wol12, Theorem 5.5].

Proof. [✓ Proof](#) Embed A as the northwest corner of a normal operator N on $\mathbb{C}^D \oplus \mathbb{C}^E$. Extend T to a positive subunitary map $\tilde{T}$ on the larger space (acting as T on the northwest block, zero elsewhere). Apply Theorem 5.2.2.2 to $\tilde{T}$ and N , then compress the resulting inequality back to $\mathbb{C}^D$. $\square$

Theorem 5.2.3.2 (Schwarz inequality for commuting-dominant operators). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $T(\mathbb{1}) \leq \mathbb{1}$. If $A \in M_D(\mathbb{C})$ admits a positive semidefinite dominant $B \geq 0$ with $A^\dagger A \leq B$ and B commuting with A ($BA = AB$), then

$$T(A^\dagger)T(A) \leq T(B) \quad \text{and} \quad T(A)T(A^\dagger) \leq T(B).$$

This is [Wol12, Theorem 5.6].

Proof. [✓ Proof](#) First take $B > 0$. Since B commutes with A and $A^\dagger A \leq B$, also $AA^\dagger \leq B$, so $D_R = (B - A^\dagger A)^{1/2}$ and $D_L = (B - AA^\dagger)^{1/2}$ are positive semidefinite. The block matrix $N = \begin{pmatrix} A & D_L \\ D_R & -A^\dagger \end{pmatrix}$ is normal, with $N^\dagger N = NN^\dagger = B \oplus B$. As in the subnormal case, extend T to a positive subunital map on $M_{2D}(\mathbb{C})$, apply the normal-operator Schwarz inequality (Theorem 5.2.2.2) to that extension and N , and compress back to $\mathbb{C}^D$. Since N is normal, both $T(A^\dagger)T(A) \leq T(B)$ and $T(A)T(A^\dagger) \leq T(B)$ follow. For general $B \geq 0$, apply this to $B_\varepsilon = B + \varepsilon \mathbb{1} > 0$ and let $\varepsilon \rightarrow 0$. $\square$

Theorem 5.2.3.3 (CP variant: Kadison–Schwarz for commuting-dominant inputs). [Lean](#) [✓ Stmt](#) Let E^* be the adjoint of a trace-preserving Kraus map. If $A \in M_D(\mathbb{C})$ admits a positive semidefinite dominant B commuting with A and satisfying $A^\dagger A \leq B$, then $E^*(A^\dagger)E^*(A) \leq E^*(B)$ and $E^*(A)E^*(A^\dagger) \leq E^*(B)$.

Proof. [✓ Proof](#) Apply Theorem 5.2.3.2 to the adjoint Kraus map (which is unital under the trace-preservation hypothesis). $\square$

5.2.4 Positive maps preserve order and spectral intervals

Theorem 5.2.4.1 (Positive maps are monotone). [Lean](#) [✓ Stmt](#) If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive and $A \leq B$, then $T(A) \leq T(B)$.

Proof. [✓ Proof](#) Since $B - A \geq 0$, positivity gives $T(B - A) \geq 0$. By linearity, this is $T(B) - T(A) \geq 0$. $\square$

Theorem 5.2.4.2 (Positive maps preserve adjoints). [Lean](#) [✓ Stmt](#) If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is positive, then $T(A^\dagger) = T(A)^\dagger$ for all $A \in M_D(\mathbb{C})$.

Proof. [✓ Proof](#) Regard T as the associated positive linear map (Definition 4.1.3). For positive linear maps between matrix C^* -algebras, $T(A^*) = T(A)^*$. Since the star on $M_D(\mathbb{C})$ is the conjugate transpose, $T(A^\dagger) = T(A^*) = T(A)^* = T(A)^\dagger$. $\square$

Theorem 5.2.4.3 (Order intervals are preserved). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be positive with $T(\mathbb{1}) \leq \mathbb{1}$. If $a \leq 0 \leq b$ and $a\mathbb{1} \leq A \leq b\mathbb{1}$, then

$$a\mathbb{1} \leq T(A) \leq b\mathbb{1}.$$

This is the matrix form of [Wol12, Equation (5.21)].

Proof. [✓ Proof](#) By monotonicity, $T(a\mathbb{1}) \leq T(A) \leq T(b\mathbb{1})$. For the lower bound, $T(a\mathbb{1}) = aT(\mathbb{1})$. Since $a \leq 0$ and $T(\mathbb{1}) \leq \mathbb{1}$, we have $aT(\mathbb{1}) \geq a\mathbb{1}$. For the upper bound, $T(b\mathbb{1}) = bT(\mathbb{1}) \leq b\mathbb{1}$ since $b \geq 0$. Therefore $a\mathbb{1} \leq T(A) \leq b\mathbb{1}$. $\square$

Theorem 5.2.4.4 (Spectral interval contractivity). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be positive with $T(\mathbb{1}) \leq \mathbb{1}$, and let $A \in M_D(\mathbb{C})$ be Hermitian. If $\text{spec}(A) \subseteq [a, b]$ with $a \leq 0 \leq b$, then

$$\text{spec}(T(A)) \subseteq [a, b].$$

This is [Wol12, Equation (5.21)].

Proof.  **Proof** For Hermitian matrices, the inclusion $\text{spec}(A) \subseteq [a, b]$ is equivalent to the order bounds $a\mathbb{1} \leq A \leq b\mathbb{1}$. Apply Theorem 5.2.4.3 and translate the resulting inequalities back into spectral bounds. $\square$

5.2.5 A positive Schwarz map that is not completely positive

Example 5.2.5.1 (Transpose-trace map on $M_2(\mathbb{C})$).   **Stmt** Consider the linear map $T : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$ defined by

$$T(A) = \frac{1}{2}A^T + \frac{1}{4}\text{tr}(A)\mathbb{1}.$$

This is the map from [Wol12, Example 5.3].

Theorem 5.2.5.2 (The transpose-trace map is positive).   **Stmt** The map from Example 5.2.5.1 is positive.

Proof.  **Proof** It is a non-negative linear combination of the transpose map and the map $A \mapsto \text{tr}(A)\mathbb{1}$, and both of these maps send positive semidefinite matrices to positive semidefinite matrices. $\square$

Theorem 5.2.5.3 (The transpose-trace map satisfies the Schwarz inequality).   **Stmt** For every $A \in M_2(\mathbb{C})$,

$$T(A^\dagger A) - T(A^\dagger)T(A) \geq 0,$$

where T is the map from Example 5.2.5.1.

Proof.  **Proof** Write the Schwarz gap as $F(A)$. One checks that $F(A + \lambda\mathbb{1}) = F(A)$ for every scalar λ , so it is enough to treat the case $\text{tr}(A) = 0$. In that case a direct 2×2 computation shows that $F(A)$ dominates $\frac{1}{2}(A^\dagger A)^T$, hence is positive semidefinite. $\square$

Theorem 5.2.5.4 (The transpose-trace map is not completely positive).   **Stmt** The map from Example 5.2.5.1 is not completely positive.

Proof.  **Proof** Compute the Choi matrix of T and test it on the antisymmetric vector $|01\rangle - |10\rangle$. The resulting expectation value is negative, so the Choi matrix is not positive semidefinite. $\square$

Chapter 6

Perron–Frobenius Theory for Channels and Transfer Maps

This chapter establishes the Perron–Frobenius theory: existence, positive definiteness, and uniqueness of fixed points for injective and irreducible transfer maps, canonical gauge constructions, ergodicity of irreducible channels, the spectral-radius identification of Perron eigenvalues, and Wolf’s spectral characterization of irreducibility. The conceptual source is Wolf’s treatment of irreducible positive maps [Wol12, Chapter 6, especially Theorems 6.2–6.5 and Corollary 6.3] and [EHK78].

Theorem 6.1 (Brouwer fixed point on density matrices). Lean Lean Lean Lean Lean ✓ Stmt
Let $D > 0$. Every continuous map from the compact convex set of density matrices in $M_D(\mathbb{C})$ to itself has a fixed point.

Proof. ✓ Proof The proof starts from Brouwer’s theorem for products of simplices, transfers it to closed cubes, and then to compact retracts of finite-dimensional real normed spaces. The density-matrix set is such a compact retract, using the explicit density retraction. $\square$

6.1 Positive definiteness

Remark 6.1.1. ✓ Stmt Wolf’s general theory treats irreducible positive maps first ([Wol12, Theorem 6.2]), then derives non-degeneracy and positive definiteness ([Wol12, Theorem 6.3]). The injective-tensor version (Theorem 6.1.2) uses a shorter direct kernel argument; the irreducible CP-map version (Theorem 6.1.3) follows via the support-projection argument of [Wol12, Theorem 6.2].

Theorem 6.1.2 (PSD fixed point is positive definite). Lean ✓ Stmt *Let A be an injective MPS tensor and let $\rho \geq 0$, $\rho \neq 0$, be a fixed point of the transfer map $\mathcal{E}_A(\rho) = \rho$. Then ρ is positive definite.*

Proof. ✓ Proof Suppose $v^\dagger \rho v = 0$ for some $v \neq 0$. Since $\rho \geq 0$, this implies $\rho v = 0$. The fixed-point equation gives $v^\dagger \rho v = \sum_i ((A^i)^\dagger v)^\dagger \rho ((A^i)^\dagger v)$. Each term is non-negative; if the sum is zero, then $\rho (A^i)^\dagger v = 0$ for all i : the kernel of ρ is invariant under all $(A^i)^\dagger$. Since the $\{A^i\}$ span $M_D(\mathbb{C})$, we have $\rho M v = 0$ for every matrix M . Given any vector w , choose M with $M v = w$. Then $\rho w = 0$ for every w , so $\rho = 0$, contradicting $\rho \neq 0$. In [Wol12, Theorem 6.3(2)], the same conclusion is reached via the growth condition $(\mathbb{1} + \mathcal{E}_A)^{D-1}(\rho) > 0$ of [Wol12, Theorem 6.2, condition (2)]; here the direct kernel argument is shorter under injectivity. $\square$

Theorem 6.1.3 (PSD fixed point is positive definite — irreducible version). Lean ✓ Stmt If $\mathcal{E}_A$ is irreducible and $\rho \geq 0$, $\rho \neq 0$ is a fixed point, then ρ is positive definite.

Proof. ✓ Proof Suppose ρ is not positive definite. Then ρ has a nontrivial kernel; let Q be the orthogonal projection onto the support of ρ (the complement of the kernel). A direct computation using the fixed-point equation $\mathcal{E}_A(\rho) = \rho$ shows that Q is an invariant projection for $\mathcal{E}_A$: it satisfies $Q\mathcal{E}_A(QXQ)Q = \mathcal{E}_A(QXQ)$ for all X . By irreducibility of $\mathcal{E}_A$ (Definition 4.2.4), every invariant projection is 0 or $\mathbb{1}$. If $Q = 0$ then $\rho = 0$, contradicting $\rho \neq 0$. If $Q = \mathbb{1}$ then ρ is positive definite, contradicting our assumption. Hence ρ must be positive definite. This is the support-projection direction of [Wol12, Theorem 6.2(1)]. $\square$

Theorem 6.1.4 (Orthogonal trace condition). Lean ✓ Stmt Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, and let $A, B \geq 0$ be nonzero positive semidefinite matrices with $\text{tr}(BA) = 0$. Then there exists t with $1 \leq t \leq D - 1$ such that

$$\text{tr}(B \cdot E^t(A)) > 0.$$

Proof. ✓ Proof The growth theorem for irreducible CP maps ([Wol12, Theorem 6.2(2)]) gives $(\mathbb{1} + E)^{D-1}(A) > 0$ (positive definite). Since $B \geq 0$ is nonzero, $\text{tr}(B \cdot (\mathbb{1} + E)^{D-1}(A)) > 0$. Expanding by the binomial theorem:

$$\text{tr} \left(B \cdot \sum_{k=0}^{D-1} \binom{D-1}{k} E^k(A) \right) > 0.$$

Each term $\text{tr}(B \cdot E^k(A)) \geq 0$ since both factors are PSD. The $k = 0$ term vanishes by hypothesis ($\text{tr}(BA) = 0$). Hence at least one $k \in \{1, \dots, D - 1\}$ contributes a strictly positive term. This is [Wol12, Theorem 6.2(4)]: item (2) (growth condition) implies item (4) by a binomial expansion and PSD positivity argument. $\square$

6.2 Uniqueness

Theorem 6.2.1 (Uniqueness of PSD fixed point). Lean ✓ Stmt Let A be an injective MPS tensor. If $\rho, \sigma \geq 0$ are both nonzero fixed points of $\mathcal{E}_A$, then $\sigma = c\rho$ for some $c \in \mathbb{C}$.

Proof. ✓ Proof Both ρ and σ are positive definite by Theorem 6.1.2. Write $\rho = SS^\dagger$ via a spectral square root and set $H = (S^\dagger)^{-1}\sigma S^{-1}$, which is Hermitian and positive definite. Let c_0 be the minimal eigenvalue of H . Then $\tau := \sigma - c_0\rho$ is a PSD fixed point which is *not* positive definite (it has a zero eigenvalue by construction of c_0). By Theorem 6.1.2, τ must be zero, giving $\sigma = c_0\rho$. In [Wol12, Theorem 6.3, item 2] uniqueness is proved via the growth condition of [Wol12, Theorem 6.2]; the same minimal-eigenvalue argument applies here, using the injective-tensor positive-definiteness (Theorem 6.1.2) in place of the growth condition. $\square$

Remark 6.2.2. ✓ Stmt The proof produces a positive real scalar $c_0 > 0$: it is the minimal eigenvalue of the positive definite Hermitian matrix $H = (S^\dagger)^{-1}\sigma S^{-1}$.

Theorem 6.2.3 (Irreducible fixed-point uniqueness). Lean ✓ Stmt Let A be an MPS tensor whose transfer map is irreducible. If $\rho, \sigma \geq 0$ are fixed points of $\mathcal{E}_A$, and if ρ is nonzero, then $\sigma = c\rho$ for some scalar c .

Proof. ✓ Proof The irreducible positive-definiteness theorem (Theorem 6.1.3) upgrades every nonzero positive semidefinite fixed point to a positive definite one. The same minimal eigenvalue argument used in Theorem 6.2.1 then applies. $\square$

Theorem 6.2.4 (Uniqueness of positive eigenvalue for irreducible CP maps). [Lean](#) [✓ Stmt](#) Let $D \geq 1$ and let E be an irreducible completely positive map on $M_D(\mathbb{C})$. Suppose $\rho, \sigma \geq 0$ are nonzero positive semidefinite matrices satisfying $E(\rho) = r_1 \rho$ and $E(\sigma) = r_2 \sigma$ for real scalars $r_1, r_2 > 0$. Then $r_1 = r_2$.

Proof. [✓ Proof](#) Extract Kraus operators K for E . Irreducibility of E passes to the adjoint transfer map $E^\dagger(\cdot) = \sum_i K_i^\dagger(\cdot) K_i$, which is a positive CP map. The Perron–Frobenius existence theorem (Theorem 6.6.1) together with the irreducible positive-definiteness upgrade (Theorem 6.1.3) gives a positive definite eigenvector $\tau > 0$ with eigenvalue $t > 0$: $E^\dagger(\tau) = t \tau$. For any nonzero PSD X with $E(X) = sX$, the trace pairing identity yields

$$s \cdot \text{tr}(\tau X) = \text{tr}(\tau \cdot E(X)) = \text{tr}(E^\dagger(\tau) \cdot X) = t \cdot \text{tr}(\tau X).$$

Since $\tau > 0$ and $X \geq 0$, $X \neq 0$, the scalar $\text{tr}(\tau X) > 0$, so $s = t$. Applying this to both (ρ, r_1) and (σ, r_2) gives $r_1 = t = r_2$. This is [Wol12, Theorem 6.3, item 3], the non-degeneracy of the spectral-radius eigenvalue for irreducible CP maps; the proof follows Wolf’s dual-map trace argument. $\square$

6.3 Existence and the Perron–Frobenius theorem

Theorem 6.3.1 (Existence of PSD fixed point). [Lean](#) [✓ Stmt](#) Assume $D \geq 1$. Let A be an MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ (so that $\mathcal{E}_A$ is trace-preserving). Then there exists $\rho \geq 0$, $\rho \neq 0$, with $\mathcal{E}_A(\rho) = \rho$.

Proof. [✓ Proof](#) Under the normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, the transfer map is a quantum channel (Theorem 4.2.7). The Cesàro fixed-point theorem (Theorem 4.3.4) then provides the fixed point; injectivity is not needed for this step. $\square$

Theorem 6.3.2 (Quantum Perron–Frobenius). [Lean](#) [✓ Stmt](#) Assume $D \geq 1$. Let A be an injective MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then the transfer map $\mathcal{E}_A$ has a unique PSD fixed point ρ (up to scaling), and ρ is positive definite.

Proof. [✓ Proof](#) Combine Theorems 6.3.1, 6.1.2, and 6.2.1. $\square$

Theorem 6.3.3 (Quantum Perron–Frobenius for irreducible transfer maps). [Lean](#) [✓ Stmt](#) Assume $D \geq 1$. Let A be an MPS tensor such that its transfer map is irreducible and $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ (so that $\mathcal{E}_A$ is trace-preserving). Then $\mathcal{E}_A$ has a unique positive definite fixed point, up to scalar multiple.

Proof. [✓ Proof](#) Existence is the channel fixed-point theorem (Theorem 6.3.1). Irreducibility upgrades the fixed point to positive definite, and Theorem 6.2.3 gives uniqueness up to scalar multiple. $\square$

6.4 Right- and left-canonical gauges

Theorem 6.4.1 (Right-canonical (unital) gauge). [Lean](#) [✓ Stmt](#) Let $\mathcal{E}_A(\rho) = \rho$ and let S be an invertible matrix with $SS^\dagger = \rho$. Define the gauged operators $A'^i = S^{-1}A^iS$. Then $\sum_i A'^i(A'^i)^\dagger = \mathbb{1}$, i.e. the gauged Kraus map is unital. This is the right-canonical normalization.

Proof. [✓ Proof](#) Expand each summand: $(S^{-1}A^iS)(S^{-1}A^iS)^\dagger = S^{-1}A^i \underbrace{SS^\dagger}_\rho (A^i)^\dagger (S^\dagger)^{-1}$. Summing over i and using $\sum_i A^i \rho (A^i)^\dagger = \rho$ gives $S^{-1} \rho (S^\dagger)^{-1} = S^{-1} S S^\dagger (S^\dagger)^{-1} = \mathbb{1}$. $\square$

Theorem 6.4.2 (Left-canonical (trace-preserving) gauge). [Lean](#) [✓ Stmt](#) Let σ be a fixed point of the adjoint transfer map $\sum_i (A^i)^\dagger \sigma A^i = \sigma$, and let S be invertible with $S^\dagger S = \sigma$. Define $A'^i = S A^i S^{-1}$. Then $\sum_i (A'^i)^\dagger A'^i = \mathbb{1}$, i.e. the gauged Kraus map is trace-preserving. This is the left-canonical normalization.

Proof. [✓ Proof](#) Expand each summand: $(S A^i S^{-1})^\dagger (S A^i S^{-1}) = (S^\dagger)^{-1} (A^i)^\dagger \underbrace{S^\dagger S}_\sigma A^i S^{-1}$. Summing over i and using $\sum_i (A^i)^\dagger \sigma A^i = \sigma$ gives $(S^\dagger)^{-1} \sigma S^{-1} = (S^\dagger)^{-1} S^\dagger S S^{-1} = \mathbb{1}$. $\square$

Theorem 6.4.3 (Square-root trace-preserving gauge from an adjoint fixed point). [Lean](#) [✓ Stmt](#) Let σ be positive definite and satisfy $\sum_i (A^i)^\dagger \sigma A^i = \sigma$. Define

$$B^i = \sigma^{1/2} A^i \sigma^{-1/2}.$$

Then B is trace-preserving: $\sum_i (B^i)^\dagger B^i = \mathbb{1}$.

Proof. [✓ Proof](#) Since $\sigma^{1/2}$ is self-adjoint and $\sigma^{1/2} \sigma^{1/2} = \sigma$, each summand becomes $(\sigma^{1/2})^{-1} (A^i)^\dagger \sigma A^i \sigma^{-1/2}$. Summing over i and using the fixed-point equation leaves $(\sigma^{1/2})^{-1} \sigma \sigma^{-1/2} = \mathbb{1}$. $\square$

Lemma 6.4.4 (Square-root trace-preserving gauge is a gauge equivalence). [Lean](#) [✓ Stmt](#) If σ is positive definite and $B^i = \sigma^{1/2} A^i \sigma^{-1/2}$, then A and B are gauge-equivalent.

Proof. [✓ Proof](#) The gauge matrix is $\sigma^{1/2}$, which is invertible because σ is positive definite. $\square$

Remark 6.4.5. [✓ Stmt](#) The right-canonical gauge of Theorem 6.4.1 and the left-canonical gauge of Theorem 6.4.2 are generally different similarity transforms: one is built from a fixed point of $\mathcal{E}_A$, the other from a fixed point of the adjoint transfer map. This section does not claim that a single gauge makes the transfer map simultaneously unital and trace-preserving.

6.5 Similarity preserves irreducibility

Lemma 6.5.1 (Similarity transform preserves irreducibility). [Lean](#) [✓ Stmt](#) Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $C \in M_D(\mathbb{C})$ be invertible ($\det C \neq 0$), and let $c > 0$. Define the similarity-transformed map

$$E'(X) = c \cdot C^{-1} E(C X C^\dagger) (C^\dagger)^{-1}.$$

Then E' is also irreducible.

Proof. [✓ Proof](#) Suppose Q is an invariant projection for E' : $Q \neq 0$, $Q \neq \mathbb{1}$, and $(\mathbb{1}-Q)E'(QXQ)Q = 0$ for all X . Setting $R = CQC^\dagger$, the support projection P of R is an invariant projection for E . Irreducibility of E forces $P = 0$ or $P = \mathbb{1}$. The invertibility of C then forces $Q = 0$ or $Q = \mathbb{1}$, a contradiction. This is the full similarity version of [Wol12, Proposition 6.6]; the scalar case ($C = \mathbb{1}$) gives the scaling result stated next. $\square$

Lemma 6.5.2 (Similarity is an involution). [Lean](#) [✓ Stmt](#) For any invertible $C \in M_D(\mathbb{C})$ and any linear map E on $M_D(\mathbb{C})$, the similarity transforms by C and C^{-1} compose to the identity:

$$X \mapsto C(C^{-1} E(C X C^\dagger) (C^\dagger)^{-1}) C^\dagger = E(X).$$

Proof. [✓ Proof](#) Direct computation: the inner conjugation $C \cdot (C^{-1} X (C^\dagger)^{-1}) \cdot C^\dagger$ collapses to X using $CC^{-1} = \mathbb{1}$ and $(C^\dagger)^{-1} C^\dagger = \mathbb{1}$, and the outer conjugation by $C(\cdot)C^\dagger$ then cancels the inner $C^{-1}(\cdot)(C^\dagger)^{-1}$ on $E(X)$ for the same reason. $\square$

Lemma 6.5.3 (Irreducibility is invariant under similarity). Lean ✓ Stmt For any invertible $C \in M_D(\mathbb{C})$ and any linear map E on $M_D(\mathbb{C})$, the similarity transform $X \mapsto C^{-1} E(C X C^\dagger) (C^\dagger)^{-1}$ is irreducible if and only if E is.

Proof. ✓ Proof The forward direction follows from Lemma 6.5.1 applied with the inverse C^{-1} (whose determinant is also nonzero): if the similarity transform of E by C is irreducible, then so is its similarity transform by C^{-1} , which equals E by Lemma 6.5.2. The reverse direction is Lemma 6.5.1 itself (with $c = 1$). $\square$

Theorem 6.5.4 (Scaling preserves irreducibility). Lean ✓ Stmt If E is an irreducible map and $c \neq 0$, then cE is also irreducible. This is the scalar special case ($C = \mathbb{1}$) of [Wol12, Proposition 6.6].

Proof. ✓ Proof An invariant projection for cE is an invariant projection for E (scale out the nonzero constant from the commutation relation). $\square$

6.6 Perron–Frobenius eigenvector existence

This section establishes the existence of a positive definite eigenvector for the adjoint transfer map of an irreducible MPS tensor, and uses it to construct a TP-gauge normalization. The PSD-eigenvector step corresponds to [Wol12, Theorem 6.5] (general positive maps) and [EHK78]; the upgrade to positive definiteness uses the irreducible-map theory of [Wol12, Theorem 6.3]. The TP-gauge application follows [CPGSV16, Appendix A].

Theorem 6.6.1 (PSD eigenvector for positive maps). Lean ✓ Stmt Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $D > 0$, and assume that $E(\sigma) \neq 0$ for every nonzero PSD matrix σ . Then there exist a nonzero PSD matrix ρ and a positive real $r > 0$ such that $E(\rho) = r\rho$.

This is the finite-dimensional Perron–Frobenius theorem for positive maps on matrix algebras ([Wol12, Theorem 6.5]; see also [EHK78]). The proof applies Brouwer’s fixed-point theorem to the normalization map $\rho \mapsto E(\rho)/\text{tr}(E(\rho))$ on the compact convex set of density matrices.

Theorem 6.6.2 (Adjoint transfer map is nonzero). Lean ✓ Stmt If some Kraus operator $A^i \neq 0$, then the adjoint transfer map $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is nonzero.

Proof. ✓ Proof If $\mathcal{E}_A^\dagger = 0$ then $\mathcal{E}_A^\dagger(\mathbb{1}) = 0$, so $\sum_i (A^i)^\dagger A^i = 0$. Since each summand $(A^i)^\dagger A^i$ is PSD, each $(A^i)^\dagger = 0$ and hence each $A^i = 0$. $\square$

Theorem 6.6.3 (Positive-definite adjoint eigenvector for irreducible tensors). Lean ✓ Stmt Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive definite matrix σ and a positive real $r > 0$ such that $\mathcal{E}_A^\dagger(\sigma) = r\sigma$.

This combines the general PF existence ([Wol12, Theorem 6.5]) with the irreducible upgrade to positive definiteness ([Wol12, Theorem 6.3(2)]); the application to the adjoint transfer map follows [CPGSV16, Appendix A].

Proof. ✓ Proof By Theorem 6.6.2, $\mathcal{E}_A^\dagger \neq 0$. Since $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is a Kraus map, it is positive. Irreducibility of A transfers to the adjoint transfer map. More precisely: if P is invariant for $\mathcal{E}_A$ (i.e. $(\mathbb{1} - P)K_i P = 0$ for all Kraus operators K_i), then taking adjoints gives $P K_i^\dagger (\mathbb{1} - P) = 0$, which means $\mathbb{1} - P$ is invariant for $\mathcal{E}_A^\dagger$. Hence $\mathcal{E}_A$ is irreducible if and only if $\mathcal{E}_A^\dagger$ is irreducible, and together with $\mathcal{E}_A^\dagger \neq 0$ this implies $\mathcal{E}_A^\dagger(\tau) \neq 0$ for every nonzero PSD matrix τ . Hence Theorem 6.6.1 applies to $\mathcal{E}_A^\dagger$, giving $\sigma_0 \geq 0$ (PSD, nonzero) and $r > 0$ with

$\mathcal{E}_A^\dagger(\sigma_0) = r\sigma_0$. Rescale: set $T^i = r^{-1/2}(A^i)^\dagger$. Then $\mathcal{E}_T(\sigma_0) = \sigma_0$, and $\mathcal{E}_T$ is irreducible (by Theorem 6.5.4, since irreducibility of $\mathcal{E}_A$ transfers to the adjoint and is preserved under scaling). Since $\mathcal{E}_T$ is irreducible with a PSD fixed point, the positive-definiteness upgrade (Theorem 6.1.3) gives σ_0 positive definite. $\square$

Theorem 6.6.4 (TP-gauge existence for irreducible tensors). [Lean](#) [✓ Stmt](#) *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive real r , a positive definite matrix σ , and a gauge-equivalent tensor B such that*

$$B^i = \sigma^{1/2} \cdot (r^{-1/2} A^i) \cdot \sigma^{-1/2}$$

is trace-preserving: $\sum_i (B^i)^\dagger B^i = \mathbb{1}$.

This is the “spectral rescaling + TP gauge” step of [CPGSV16, Appendix A]. The similarity is generally non-unitary.

Proof. [✓ Proof](#) By Theorem 6.6.3, obtain σ positive definite and $r > 0$ with $\mathcal{E}_A^\dagger(\sigma) = r\sigma$. Set $c = r^{-1/2}$ and $A'^i = cA^i$. Then $\mathcal{E}_{A'}^\dagger(\sigma) = \sigma$. By the square-root trace-preserving gauge construction (Theorem 6.4.3), $B^i = \sigma^{1/2} A'^i \sigma^{-1/2}$ satisfies $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. Lemma 6.4.4 gives the required gauge equivalence. $\square$

Theorem 6.6.5 (Unital-gauge existence for irreducible tensors). [Lean](#) [✓ Stmt](#) *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive real r , a positive definite matrix ρ , and a gauge-equivalent tensor B such that*

$$B^i = r^{-1/2} \rho^{-1/2} A^i \rho^{1/2}$$

is unital: $\sum_i B^i (B^i)^\dagger = \mathbb{1}$.

This is the Perron–Frobenius unital-gauge orientation used in [PGVWC07, Theorem 4, lines 765–770], stated for one irreducible nonzero block.

Proof. [✓ Proof](#) Apply Theorem 6.6.3 to the conjugate-transposed Kraus family. Irreducibility transfers to that family, and the nonzero Kraus hypothesis is preserved by taking adjoints. Translating the resulting adjoint eigenvector equation back gives $\mathcal{E}_A(\rho) = r\rho$ with ρ positive definite and $r > 0$. The unital gauge theorem then gives the displayed unital representative, and the same square-root similarity gives gauge equivalence to $r^{-1/2} A$. $\square$

6.7 Exponential positivity for irreducible CP maps

Theorem 6.7.1 (Exponential truncation is positive definite). [Lean](#) [✓ Stmt](#) *Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $A \geq 0$ be nonzero, and let $t > 0$. Then the finite exponential truncation*

$$\sum_{k=0}^{D-1} \frac{t^k}{k!} E^k(A)$$

is positive definite. This is the finite-sum core of [Wol12, Theorem 6.2(3)].

Proof. [✓ Proof](#) If a nonzero vector v annihilated this quadratic form, then every term $E^k(A)$ with $0 \leq k \leq D-1$ would annihilate v . The growth theorem for irreducible CP maps (that is, $(\mathbb{1} + E)^{D-1}(A) > 0$ for irreducible E and nonzero PSD A , as established in the proof of Theorem 6.1.4) then forces $(\mathbb{1} + E)^{D-1}(A)$ to annihilate v , contradicting its positive definiteness. $\square$

Theorem 6.7.2 (Exponential positivity for irreducible CP maps). [Lean](#) [✓ Stmt](#) Let E be an irreducible completely positive map on $M_D(\mathbb{C})$, let $A \geq 0$ be nonzero, and let $t > 0$. Then

$$\exp(tE)(A) = \sum_{k=0}^{\infty} \frac{t^k}{k!} E^k(A)$$

is positive definite. This is [\[Wol12, Theorem 6.2\(3\)\]](#).

Proof. [✓ Proof](#) By Theorem 6.7.1, the first D terms already form a positive definite matrix. Every remaining term is positive semidefinite, so the full exponential series is the sum of a positive definite matrix and a positive semidefinite tail. $\square$

6.8 Ergodicity of irreducible channels

Theorem 6.8.1 (Unique density-matrix fixed point for an irreducible channel). [Lean](#) [✓ Stmt](#) Let E be an irreducible quantum channel on $M_D(\mathbb{C})$ with $D > 0$. Then there exists a unique density matrix σ such that $E(\sigma) = \sigma$. Moreover σ is positive definite. This is the qualitative form of [\[Wol12, Corollary 6.3\]](#).

Proof. [✓ Proof](#) Every Cesàro mean of a density matrix stays inside the compact convex set of density matrices, so a subsequence converges. Its limit is a fixed point by the telescoping argument, positive definiteness follows from irreducibility, and uniqueness comes from the uniqueness of positive semidefinite Perron eigenvectors at eigenvalue 1. $\square$

Theorem 6.8.2 (Cesàro convergence for irreducible channels). [Lean](#) [✓ Stmt](#) Let E be an irreducible quantum channel on $M_D(\mathbb{C})$ with $D > 0$, and let ρ be any density matrix. Then the Cesàro means

$$\frac{1}{N} \sum_{n=0}^{N-1} E^n(\rho)$$

converge to the unique positive definite density-matrix fixed point of E . This is [\[Wol12, Corollary 6.3\]](#).

Proof. [✓ Proof](#) Every subsequential limit of the Cesàro means is again a density-matrix fixed point. By Theorem 6.8.1, there is only one such fixed point, so every convergent subsequence has the same limit. Compactness then forces the whole sequence to converge to that limit. $\square$

6.9 Spectral radius at the Perron eigenvalue

Theorem 6.9.1 (Spectral radius is the Perron eigenvalue). [Lean](#) [✓ Stmt](#) Let E be an irreducible completely positive map on $M_D(\mathbb{C})$. Suppose $\rho > 0$ and $E(\rho) = r\rho$ with $r > 0$. Then the spectral radius of E is r . This is [\[Wol12, Theorem 6.3\(4\)\]](#).

Proof. [✓ Proof](#) Choose a positive definite eigenvector for the adjoint map with the same eigenvalue, gauge by its square root, and rescale by r^{-1} . The resulting map is trace-preserving and has a positive definite fixed point, so 1 is an eigenvalue. The spectral radius of a trace-preserving map F satisfies $\rho(F) \leq 1$, since $\text{tr}(F^n(X)) = \text{tr}(X)$ for all n bounds the growth of iterates. Hence its spectral radius is 1. Undoing the similarity and scalar rescaling recovers the spectral radius of E . $\square$

Theorem 6.9.2 (Real spectral-radius identity). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 6.9.1, the real-valued spectral radius of E is also equal to r .

Proof. [✓ Proof](#) This is the real-valued reformulation of Theorem 6.9.1. □

6.10 Spectral characterization of irreducibility

Definition 6.10.1 (Spectral properties). [Lean](#) [✓ Stmt](#) A completely positive map E on $M_D(\mathbb{C})$ has the *spectral properties* of [Wol12, Theorem 6.4] if there exist a positive real number r , a positive definite right eigenvector ρ , and a positive definite left eigenvector σ for the adjoint map such that $E(\rho) = r\rho$, $E^\dagger(\sigma) = r\sigma$, every positive semidefinite right eigenvector for eigenvalue r is a scalar multiple of ρ , and the spectral radius of E is equal to r .

Theorem 6.10.2 (Irreducibility gives the spectral properties). [Lean](#) [✓ Stmt](#) Every nonzero irreducible completely positive map on $M_D(\mathbb{C})$ has the spectral properties of Definition 6.10.1.

Proof. [✓ Proof](#) Combine Perron–Frobenius existence for the map and its adjoint with the uniqueness theorem for positive semidefinite Perron eigenvectors and the spectral-radius identity of Theorem 6.9.1. □

Theorem 6.10.3 (Spectral properties imply irreducibility). [Lean](#) [✓ Stmt](#) If a completely positive map on $M_D(\mathbb{C})$ has the spectral properties of Definition 6.10.1, then it is irreducible.

Proof. [✓ Proof](#) Gauge by the positive definite left eigenvector and rescale by the Perron eigenvalue to obtain a trace-preserving map with a positive definite fixed point. A nontrivial invariant projection would then produce, by Cesàro averaging in its corner algebra, a second positive semidefinite fixed point not proportional to the Perron vector, contradicting the uniqueness clause in Definition 6.10.1. □

Theorem 6.10.4 (Irreducibility iff the spectral properties hold). [Lean](#) [✓ Stmt](#) Let E be a nonzero completely positive map on $M_D(\mathbb{C})$. Then E is irreducible if and only if it has the spectral properties of Definition 6.10.1. This is [Wol12, Theorem 6.4].

Proof. [✓ Proof](#) Combine Theorems 6.10.2 and 6.10.3. □

Chapter 7

Transfer-Operator Gaps and Block Separation

This chapter proves that the MPV overlaps between distinct (non-gauge-phase-equivalent) MPS blocks decay to zero as system size grows. It also proves the corresponding self-overlap convergence to 1 under the complementary transfer-map gap hypothesis that encodes primitivity in the cases of interest. The argument is based on a spectral analysis of the *mixed transfer operator*, following [PGVWC07].

This chapter concerns gaps in the spectrum of transfer operators, writing $\rho_{\text{spec}}(\cdot)$ for the spectral radius: $\rho_{\text{spec}}(F_{AB}) < 1$ for mixed transfer operators, and $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$ after removing the fixed-point projection. The unqualified phrase *spectral gap* is reserved for the energy gap of a many-body Hamiltonian in Chapter 13.

The algebraic mixed-transfer identities in the first two sections do not use normalization. Starting with the spectral-radius estimates, we assume the trace-preserving normalization

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}. \quad (7.1)$$

7.1 Mixed transfer operator

Definition 7.1.1 (Mixed transfer operator). [Lean](#) [✓ Stmt](#) The *mixed transfer operator* (or *cross transfer operator*) for two MPS tensors A and B of the same bond dimension D is the linear map $F_{AB} : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$F_{AB}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Definition 7.1.2 (Rectangular mixed transfer operator). [Lean](#) [✓ Stmt](#) For MPS tensors A of bond dimension D_1 and B of bond dimension D_2 (possibly $D_1 \neq D_2$), the *rectangular mixed transfer operator* is $F_{AB}^{\text{rect}} : M_{D_1 \times D_2}(\mathbb{C}) \rightarrow M_{D_1 \times D_2}(\mathbb{C})$ defined by

$$F_{AB}^{\text{rect}}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Theorem 7.1.3 (Self-transfer). [Lean](#) [✓ Stmt](#) Setting $B = A$ recovers the ordinary transfer map: $F_{AA} = \mathcal{E}_A$.

Proof. ✓ Proof Substituting $B = A$ into the defining sum $F_{AB}(X) = \sum_i A^i X(B^i)^\dagger$ gives $\sum_i A^i X(A^i)^\dagger = \mathcal{E}_A(X)$. $\square$

Lemma 7.1.4 (Iterated mixed transfer). Lean ✓ Stmt For all $X \in M_D(\mathbb{C})$ and $N \geq 0$,

$$F_{AB}^N(X) = \sum_{\sigma \in \{0, \dots, d-1\}^N} A^\sigma X(B^\sigma)^\dagger,$$

where $A^\sigma = A^{\sigma_1} \dots A^{\sigma_N}$ denotes the word evaluation (Definition 2.1.2).

Proof. ✓ Proof Induction on N : the base case is $F^0(X) = X$; the inductive step applies F_{AB} to each summand and re-indexes via $\{0, \dots, d-1\}^{N+1} \cong \{0, \dots, d-1\} \times \{0, \dots, d-1\}^N$. $\square$

7.2 MPV overlap as transfer trace

Recall from Definition 2.7.3 that

$$O_{AB}(N) = \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma}.$$

This section rewrites that scalar as the operator trace of the mixed transfer power.

Lemma 7.2.1 (Trace expansion over matrix units). Lean ✓ Stmt For any linear endomorphism T on $M_{D_1 \times D_2}(\mathbb{C})$, the operator trace can be expanded as

$$\text{Tr}(T) = \sum_{p=0}^{D_1-1} \sum_{q=0}^{D_2-1} (T(E_{pq}))_{pq},$$

where $E_{pq} \in M_{D_1 \times D_2}(\mathbb{C})$ is the matrix with 1 in position (p, q) and 0 elsewhere.

Proof. ✓ Proof The operator trace $\text{Tr}(T)$ equals the sum of $\text{tr}(T(E_{pq}) \cdot E_{qp})$ over all (p, q) , and $\text{tr}(M \cdot E_{qp}) = M_{pq}$. $\square$

Lemma 7.2.2 (Overlap equals transfer trace). Lean ✓ Stmt For tensors A, B of bond dimension $D \geq 1$,

$$O_{AB}(N) = \text{Tr}(F_{AB}^N).$$

Proof. ✓ Proof Expand the operator trace using Lemma 7.2.1 as a sum over matrix units E_{pq} . Apply Lemma 7.1.4 to evaluate $F_{AB}^N(E_{pq})$, extract the (p, q) -entry, and reassemble. The resulting double sum over (p, q) and σ equals $\sum_\sigma V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma} = O_{AB}(N)$. $\square$

Lemma 7.2.3 (Rectangular overlap as transfer trace). Lean ✓ Stmt For tensors A of bond dimension $D_1 \geq 1$ and B of bond dimension $D_2 \geq 1$,

$$O_{AB}(N) = \text{Tr}((F_{AB}^{\text{rect}})^N).$$

Proof. ✓ Proof Expand the operator trace over the rectangular matrix-unit basis. Then apply the rectangular iterated-transfer formula and sum over the matrix-unit indices exactly as in Lemma 7.2.2. $\square$

7.3 Frobenius norm estimates

The transfer-operator gap proofs use the Frobenius (Hilbert–Schmidt) norm as the natural norm for the finite-dimensional matrix algebra. We write $\|\cdot\|_F$ for this norm; it coincides with, and we use it interchangeably with, the Hilbert–Schmidt norm $\|\cdot\|_{\text{HS}}$ on finite-dimensional spaces. We use the squared version and an isometric embedding into Euclidean space.

Definition 7.3.1 (Frobenius norm squared). [Lean](#) [✓ Stmt](#) For a (possibly rectangular) matrix $X \in M_{m \times n}(\mathbb{C})$, the *Frobenius norm squared* is

$$\|X\|_F^2 := \sum_{i=0}^{m-1} \sum_{j=0}^{n-1} |X_{ij}|^2.$$

Lemma 7.3.2 (Trace formula for Frobenius norm). [Lean](#) [Lean](#) [✓ Stmt](#) $\|X\|_F^2 = \text{Re tr}(X^\dagger X)$.

Proof. [✓ Proof](#) Expand the matrix product and trace entry by entry. $\square$

Lemma 7.3.3 (Trace product of positive semidefinite matrices). [Lean](#) [✓ Stmt](#) If $A, B \geq 0$, then $\text{tr}(AB) \geq 0$.

Proof. [✓ Proof](#) Write $B = U\Lambda U^\dagger$ with U unitary and Λ diagonal. Then

$$\text{tr}(AB) = \text{tr}((U^\dagger AU)\Lambda) = \sum_i (U^\dagger AU)_{ii} \Lambda_{ii} \geq 0,$$

since $U^\dagger AU \geq 0$ and $\Lambda_{ii} \geq 0$ for every i . $\square$

Definition 7.3.4 (Euclidean-space embedding). [Lean](#) [✓ Stmt](#) The map $\text{vec} : M_{m \times n}(\mathbb{C}) \rightarrow \mathbb{C}^{mn}$ that flattens matrix entries is an isometric embedding with respect to the Frobenius norm: $\|\text{vec}(X)\|^2 = \|X\|_F^2$.

Lemma 7.3.5 (Norm of embedded matrix). [Lean](#) [✓ Stmt](#) $\|\text{vec}(X)\|^2 = \|X\|_F^2$.

Proof. [✓ Proof](#) Both sides equal $\sum_{i,j} \|X_{ij}\|^2$: the left side by the Euclidean entry-norm formula, and the right side by the definition of the Frobenius norm. $\square$

7.4 Eigenvalue bound and transfer-operator gap

The eigenvalue bound comes from a uniform Frobenius-norm estimate obtained directly from the trace-preserving normalization (cf. [Wol12, Proposition 6.1] for the general operator-norm version via Russo–Dye). For the rigidity theorem, one gauges A and B separately using positive fixed points of their transfer maps to obtain unital Kraus families, and then applies the Kadison–Schwarz equality argument ([Wol12, Theorem 5.3]) to a block embedding of a mixed-transfer eigenvector.

Theorem 7.4.1 (Eigenvalue bound). [Lean](#) [✓ Stmt](#) Let A, B be normalized MPS tensors. Every eigenvalue μ of F_{AB} satisfies $|\mu| \leq 1$.

Proof. [✓ Proof](#) Expand $F_{AB}^n(X)$ as a sum over words and use Cauchy–Schwarz, together with $\sum_i (A^i)^\dagger A^i = \sum_i (B^i)^\dagger B^i = \mathbb{1}$ (Equation (7.1), applied to both tensors), to obtain a uniform Frobenius-norm bound $\|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for all n , where C_D depends only on D . If $F_{AB}(X) = \mu X$ and $X \neq 0$, then $|\mu|^n \|X\|_{\text{HS}} = \|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for every n . Hence $|\mu| \leq 1$.

The argument uses only the trace-preserving normalization (7.1) and does not appeal to Perron–Frobenius theory. $\square$

Theorem 7.4.2 (Rectangular eigenvalue bound). [Lean](#) [✓ Stmt](#) Let A and B be normalized MPS tensors of bond dimensions $D_1 \geq 1$ and $D_2 \geq 1$. Every eigenvalue μ of F_{AB}^{rect} satisfies $|\mu| \leq 1$.

Proof. [✓ Proof](#) The word-expansion and Frobenius-norm estimate used in Theorem 7.4.1 apply equally to rectangular matrices: if $F_{AB}^{\text{rect}}(X) = \mu X$ and $X \neq 0$, then $|\mu|^n \|X\|_{\text{HS}} \leq D_1 \|X\|_{\text{HS}}$ for every n . Hence $|\mu| \leq 1$. $\square$

Lemma 7.4.3 (Spectral radius bound). [Lean](#) [✓ Stmt](#) $\rho_{\text{spec}}(F_{AB}) \leq 1$ for normalized tensors.

Proof. [✓ Proof](#) The spectral radius is the supremum of $|\mu|$ over all eigenvalues, and each satisfies $|\mu| \leq 1$ by Theorem 7.4.1. $\square$

Theorem 7.4.4 (Spectral radius ≥ 1 implies gauge-phase equivalence). [Lean](#) [✓ Stmt](#) Let A, B be injective normalized MPS tensors. If $\rho_{\text{spec}}(F_{AB}) \geq 1$, then A and B are gauge-phase equivalent.

Proof. [✓ Proof](#) Since $\rho_{\text{spec}}(F_{AB}) \leq 1$ (Lemma 7.4.3) and $\rho_{\text{spec}}(F_{AB}) \geq 1$ by hypothesis, there exists an eigenvalue μ with $|\mu| = 1$ and eigenvector $X \neq 0$: $F_{AB}(X) = \mu X$. The proof proceeds in five steps.

Step 1 (Separate gauging). Choose positive definite fixed points ρ_A, ρ_B of the transfer maps $\mathcal{E}_A, \mathcal{E}_B$ (Theorem 6.3.2) and gauge each tensor separately to a unital Kraus family (Theorem 6.4.1): set $A'^i = \rho_A^{-1/2} A^i \rho_A^{1/2}$ and similarly for B . Let $X' = \rho_A^{-1/2} X \rho_B^{-1/2}$ denote the correspondingly gauged eigenvector.

Step 2 (Block Kraus family and off-diagonal embedding). Form the block Kraus operators $C^i = \begin{pmatrix} A'^i & 0 \\ 0 & B'^i \end{pmatrix}$. These constitute a unital Kraus family on $M_{2D}(\mathbb{C})$. The off-diagonal matrix $Y = \begin{pmatrix} 0 & X' \\ 0 & 0 \end{pmatrix}$ satisfies $E_C(Y) = \mu Y$.

Step 3 (KS equality and Kraus commutation). Because $|\mu| = 1$ and E_C is unital and trace-preserving, the Kadison–Schwarz equality for peripheral eigenvectors (Theorem 5.2.3) gives $E_C(Y^\dagger Y) = E_C(Y)^\dagger E_C(Y)$. By the Kraus commutation relation (Theorem 5.2.2), Y commutes with every block Kraus operator: $C^i Y = Y C^i$ for all i .

Step 4 (Intertwining identity and invertibility). Expanding the commutation relation block by block gives $A'^i X' = X' B'^i$ for every i . Since $\{A'^i\}$ span $M_D(\mathbb{C})$ (injectivity of A), X' is a nonzero intertwiner from $M_D(\mathbb{C})$ to $M_D(\mathbb{C})$. The left multiplicative identity (Theorem 5.2.4) gives $E_C(Y^\dagger Y) = E_C(Y^\dagger) E_C(Y)$, so $X'^\dagger X'$ is a positive semi-definite fixed point of $\mathcal{E}_{B'}$, positive definite by injectivity. Hence X' is invertible.

Step 5 (Skolem–Noether conclusion). The map $M \mapsto X' M (X')^{-1}$ is a $\mathbb{C}$ -algebra automorphism of $M_D(\mathbb{C})$ sending B'^i to A'^i . By Skolem–Noether (Theorem 3.2.4) this is an inner automorphism. Undoing the separate gauges gives $B^i = \bar{\mu} X A^i X^{-1}$ for all i , establishing gauge-phase equivalence (Definition 2.3.14).

The gauging is applied to the individual transfer maps $\mathcal{E}_A, \mathcal{E}_B$ separately; the mixed transfer map F_{AB} need not be simultaneously unital and trace-preserving. $\square$

Lemma 7.4.5 (Gauged peripheral eigenvector yields Kraus intertwining). [Lean](#) [✓ Stmt](#) Let A and B be normalized tensors, and let $X \neq 0$ satisfy $F_{AB}(X) = \mu X$ with $|\mu| = 1$. Choose positive-definite fixed points ρ_A, ρ_B of the transfer maps and gauge to unital families $A'^i = S_A^{-1} A^i S_A$ and $B'^i = S_B^{-1} B^i S_B$ with $S_A S_A^\dagger = \rho_A$ and $S_B S_B^\dagger = \rho_B$. Then the transported matrix $X' = S_A^{-1} X (S_B^\dagger)^{-1}$ is nonzero, the families A' and B' are unital, and for every i one has

$$X' (B'^i)^\dagger = \mu (A'^i)^\dagger X', \quad A'^i X' = \mu X' B'^i.$$

Proof. [✓ Proof](#) Block-embed the gauged families and transported eigenvector into a single unital Kraus map, apply Kadison–Schwarz equality for the modulus-one eigenvalue, then extract the intertwining identities from the block structure. $\square$

Lemma 7.4.6 (Intertwining yields gauge-phase equivalence). [Lean](#) [Stmnt](#) *Let A and B be tensors of the same bond dimension. Suppose there are invertible gauges taking them to families A' and B' , an invertible matrix X' , and a scalar μ with $|\mu| = 1$ such that, for every physical index i ,*

$$A'^i X' = \mu X' B'^i.$$

Then A and B are gauge-phase equivalent.

Proof. [Proof](#) Compose the two gauge matrices with the intertwiner to obtain a single invertible matrix conjugating A to a scalar multiple of B . $\square$

Remark 7.4.7 (Two gauge conventions). [Stmnt](#) The right-canonical gauge used here, $A'^i = \rho^{-1/2} A^i \rho^{1/2}$, makes $\{A'^i\}$ a *unital* Kraus family ($\sum_i A'^i (A'^i)^\dagger = \mathbb{1}$). A different convention, $\tilde{A}^i = \rho^{1/2} A^i \rho^{-1/2}$, makes the family *trace-preserving* ($\sum_i (\tilde{A}^i)^\dagger \tilde{A}^i = \mathbb{1}$). Both conventions appear in practice: the unital version via direct matrix square-root inversion (used in the transfer-operator gap proof above), and the trace-preserving version (used in the canonical-form reduction of Chapter 9).

Theorem 7.4.8 (Strict transfer-operator gap). [Lean](#) [Stmnt](#) *Let A, B be injective normalized MPS tensors that are not gauge-phase equivalent. Then $\rho_{\text{spec}}(F_{AB}) < 1$.*

Proof. [Proof](#) $\rho_{\text{spec}}(F_{AB}) \leq 1$ by Lemma 7.4.3. If $\rho_{\text{spec}}(F_{AB}) = 1$, Theorem 7.4.4 gives gauge-phase equivalence, contradicting the hypothesis. $\square$

7.5 Transfer-operator gap — rectangular case

Lemma 7.5.1 (Rectangular intertwining forces equal bond dimension). [Lean](#) [Stmnt](#) *Let A and B be injective tensors of bond dimensions D_1 and D_2 . If there exist $X \in M_{D_1 \times D_2}(\mathbb{C}) \setminus \{0\}$ and $\mu \in \mathbb{C}$ such that, for every physical index i ,*

$$X(B^i)^\dagger = \mu(A^i)^\dagger X, \quad A^i X = \mu X B^i,$$

then $D_1 = D_2$.

Proof. [Proof](#) The intertwining relation shows that $\ker(X)$ is invariant under all generators of B . Since B is injective, its generators span the full matrix algebra, so $\ker(X)$ is trivial and $D_2 \leq D_1$. Applying the same argument to $X^\dagger$ gives $D_1 \leq D_2$. $\square$

Theorem 7.5.2 (Rectangular transfer-operator gap). [Lean](#) [Stmnt](#) *Let A be an injective normalized tensor of bond dimension D_1 and B an injective normalized tensor of bond dimension D_2 , with $D_1 \neq D_2$. Then $\rho_{\text{spec}}(F_{AB}^{\text{rect}}) < 1$.*

Proof. [Proof](#) The same word-expansion argument as in Theorem 7.4.2 gives $|\mu| \leq 1$ for every eigenvalue of F_{AB}^{rect} . If $\rho_{\text{spec}}(F_{AB}^{\text{rect}}) = 1$, choose a modulus-one eigenvector $X \in M_{D_1 \times D_2}(\mathbb{C}) \setminus \{0\}$.

Gauge A, B to unital families A', B' and transport X to a nonzero rectangular intertwiner X' with $A'^i X' = \mu X' B'^i$ for every i . This is the hypothesis of Lemma 7.5.1, so $D_1 = D_2$, contradicting the hypothesis. $\square$

Theorem 7.5.3 (Rectangular overlap decay). [Lean](#) [Stmnt](#) *If $D_1 \neq D_2$ and both tensors are injective and normalized, then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. [Proof](#) The transfer-operator gap $\rho_{\text{spec}}(F_{AB}^{\text{rect}}) < 1$ (Theorem 7.5.2) implies $(F_{AB}^{\text{rect}})^N \rightarrow 0$, so the operator trace converges to 0. By Lemma 7.2.3, $O_{AB}(N) = \text{Tr}((F_{AB}^{\text{rect}})^N) \rightarrow 0$. $\square$

7.6 MPV overlap decay

Two lemmas about a complex Banach algebra translate the transfer-operator gap into convergence statements.

Lemma 7.6.1 (Powers vanish below unit spectral radius). [Lean](#) [✓ Stmt](#) *Let a be an element of a complex Banach algebra with $\rho_{\text{spec}}(a) < 1$. Then $a^n \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. [✓ Proof](#) Choose r with $\rho_{\text{spec}}(a) < r < 1$. The Gelfand formula $\|a^n\|^{1/n} \rightarrow \rho_{\text{spec}}(a)$ gives $\|a^n\| \leq r^n$ for all sufficiently large n , and $r^n \rightarrow 0$. $\square$

Lemma 7.6.2 (An idempotent below unit spectral radius vanishes). [Lean](#) [✓ Stmt](#) *Let a be an element of a complex Banach algebra with $a^2 = a$ and $\rho_{\text{spec}}(a) < 1$. Then $a = 0$.*

Proof. [✓ Proof](#) By Lemma 7.6.1, $a^n \rightarrow 0$. Idempotence gives $a^{n+1} = a$ for every n , so the shifted power sequence is constant with value a ; uniqueness of limits gives $a = 0$. $\square$

Theorem 7.6.3 (Transfer powers converge to zero). [Lean](#) [✓ Stmt](#) *Let A, B be injective normalized tensors that are not gauge-phase equivalent. Then $F_{AB}^n(X) \rightarrow 0$ for every $X \in M_D(\mathbb{C})$.*

Proof. [✓ Proof](#) The transfer-operator gap $\rho_{\text{spec}}(F_{AB}) < 1$ (Theorem 7.4.8) implies $\|F_{AB}^n\|_{\text{op}} \rightarrow 0$ by Lemma 7.6.1, so $F_{AB}^n(X) \rightarrow 0$ for every X . $\square$

Theorem 7.6.4 (Overlap decay). [Lean](#) [✓ Stmt](#) *Let A, B be injective normalized tensors of the same bond dimension that are not gauge-phase equivalent. Then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. [✓ Proof](#) By Lemma 7.2.2, $O_{AB}(N) = \text{Tr}(F_{AB}^N)$. Expand the operator trace over matrix units (Lemma 7.2.1): $\text{Tr}(F_{AB}^N) = \sum_{p,q} (F_{AB}^N(E_{pq}))_{pq}$. By Theorem 7.6.3, each $F_{AB}^N(E_{pq}) \rightarrow 0$ entry-wise. Since the sum is finite, $O_{AB}(N) \rightarrow 0$. $\square$

7.7 Block separation

The block-separation results below record the overlap asymptotics used in the canonical-form separation of Chapter 9.

Theorem 7.7.1 (Cross-correlation decay). [Lean](#) [✓ Stmt](#) *If A and B are injective normalized tensors that are not gauge-phase equivalent, then for every $X \in M_D(\mathbb{C})$ one has $\text{tr}(F_{AB}^N(X)) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. [✓ Proof](#) By Theorem 7.6.3, $F_{AB}^N(X) \rightarrow 0$ in operator norm, so $\text{tr}(F_{AB}^N(X)) \rightarrow 0$. $\square$

Theorem 7.7.2 (Self-correlation persists). [Lean](#) [✓ Stmt](#) *If ρ is a fixed point of $\mathcal{E}_A$, then $\text{tr}(\mathcal{E}_A^N(\rho)) = \text{tr}(\rho)$ for all $N \geq 0$. In particular this applies to the distinguished QPF fixed point from Theorem 6.3.2.*

Proof. [✓ Proof](#) Since ρ is a fixed point, $\mathcal{E}_A^N(\rho) = \rho$ for every N , so $\text{tr}(\mathcal{E}_A^N(\rho)) = \text{tr}(\rho)$. $\square$

7.8 Transfer-operator gap under irreducible-TP hypotheses

The preceding transfer-operator gap results require *injectivity* of both tensors. The following variants weaken this to *irreducibility* combined with the trace-preserving normalization (7.1), following Cirac et al. [CPGSV17] and the irreducibility theory of [Wol12, Section 6.2]. The argument replaces the Kraus-commutation step with a Cauchy–Schwarz rigidity argument for the Perron–Frobenius fixed point.

Theorem 7.8.1 (Gauge-equivalence from irreducible-TP spectral radius). Lean ✓ Stmt *Let A, B be irreducible trace-preserving normalized MPS tensors of bond dimension D . If $\rho_{\text{spec}}(F_{AB}) \geq 1$, then A and B are gauge-phase equivalent.*

Proof. ✓ Proof Choose a modulus-one eigenvector as in Theorem 7.4.4. The proof follows the five-step structure of Theorem 7.4.4, with one key adaptation. Steps 1–3 (separate gauging, block Kraus family, KS equality and Kraus commutation) proceed identically: the gauging uses the unique positive definite fixed points guaranteed by irreducibility (Theorem 6.3.2) and the right-canonical gauge (Theorem 6.4.1). The change is in Step 4: the Kadison–Schwarz equality gives $X'^\dagger X'$ as a nonzero positive semidefinite fixed point of $\mathcal{E}_{B'}$. Under injectivity this is immediately positive definite (the fixed point is unique up to scaling). Under the weaker hypothesis of irreducibility, Theorem 6.1.3 upgrades it to positive definite, giving invertibility of X' . Formally, Lemma 7.4.5 combines the separately gauged eigenvector into the required intertwining relations. Once X' is invertible, Lemma 7.4.6 upgrades the intertwining to gauge-phase equivalence. $\square$

Theorem 7.8.2 (Strict transfer-operator gap (irreducible-TP)). Lean ✓ Stmt *Let A, B be irreducible trace-preserving normalized MPS tensors of the same bond dimension that are not gauge-phase equivalent. Then $\rho_{\text{spec}}(F_{AB}) < 1$.*

Proof. ✓ Proof Contrapositive of Theorem 7.8.1, combined with $\rho_{\text{spec}}(F_{AB}) \leq 1$ (Lemma 7.4.3). $\square$

Theorem 7.8.3 (Overlap decay (irreducible-TP)). Lean ✓ Stmt *Let A, B be irreducible trace-preserving normalized tensors of the same bond dimension that are not gauge-phase equivalent. Then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. ✓ Proof Combine the transfer-operator gap (Theorem 7.8.2) with the overlap–trace identity (Lemma 7.2.2). $\square$

Theorem 7.8.4 (Rectangular transfer-operator gap (irreducible-TP)). Lean ✓ Stmt *Let A (bond dimension D_1) and B (bond dimension D_2) be irreducible trace-preserving normalized tensors with $D_1 \neq D_2$. Then $\rho_{\text{spec}}(F_{AB}^{\text{rect}}) < 1$.*

Proof. ✓ Proof Assume for contradiction that $\rho_{\text{spec}}(F_{AB}^{\text{rect}}) = 1$. Choose a modulus-one eigenvector $X \in M_{D_1 \times D_2}(\mathbb{C})$. Gauge A, B separately to unital families A', B' and transport X to $X' \neq 0$. The matrix $X'X'^\dagger$ ($D_1 \times D_1$) is a nonzero positive-semidefinite fixed point of $\mathcal{E}_{A'}$, so by Theorem 6.1.3 it is positive definite, hence invertible; thus X' has full row rank and $D_1 \leq D_2$. Symmetrically $X'^\dagger X'$ ($D_2 \times D_2$) is positive definite, giving full column rank and $D_2 \leq D_1$. Hence $D_1 = D_2$, contradicting the hypothesis. $\square$

Theorem 7.8.5 (Rectangular overlap decay (irreducible-TP)). Lean ✓ Stmt *If $D_1 \neq D_2$ and both tensors are irreducible and trace-preserving normalized, then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. ✓ Proof Combine Theorem 7.8.4 with Lemma 7.2.3. $\square$

7.9 Conditional expectation from a faithful fixed point

This section follows [Wol12, Section 6.4]; the conditional expectation is built from the fixed-point algebra of [Wol12, Theorem 6.14] via [Wol12, (1.40)]. The scalar conditional expectation E_σ is the special case where the fixed-point $*$ -subalgebra is the scalar algebra $\mathbb{C} \cdot \mathbb{1}$ (the primitive-channel case). The general irreducible case with nontrivial period requires the Wedderburn block decomposition of [Wol12, Theorem 6.14].

Definition 7.9.1 (Conditional expectation). Lean ✓ Stmt Given a $*$ -subalgebra $S \subseteq M_D(\mathbb{C})$, a linear map $P : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a conditional expectation onto S if it is idempotent, unital, has range contained in S , and satisfies $P(X) = X$ for all $X \in S$.

Definition 7.9.2 (Scalar conditional expectation). Lean ✓ Stmt For $\sigma \geq 0$ with $\text{tr}(\sigma) \neq 0$, define

$$E_\sigma(X) := \frac{\text{tr}(\sigma X)}{\text{tr}(\sigma)} \mathbb{1}.$$

Lemma 7.9.3 (Evaluation formula). Lean ✓ Stmt $E_\sigma(X) = \frac{\text{tr}(\sigma X)}{\text{tr}(\sigma)} \mathbb{1}$.

Lemma 7.9.4 (Unitality). Lean ✓ Stmt $E_\sigma(\mathbb{1}) = \mathbb{1}$.

Lemma 7.9.5 (Idempotence). Lean ✓ Stmt $E_\sigma(E_\sigma(X)) = E_\sigma(X)$ for all X .

Lemma 7.9.6 (Range is scalar). Lean ✓ Stmt For every X , there exists $c \in \mathbb{C}$ with $E_\sigma(X) = c \mathbb{1}$.

Lemma 7.9.7 (Fixes scalar operators). Lean ✓ Stmt For $c \in \mathbb{C}$, $E_\sigma(c \mathbb{1}) = c \mathbb{1}$.

Lemma 7.9.8 (Absorbs the adjoint map). Lean ✓ Stmt If $T(\sigma) = \sigma$, then $E_\sigma \circ T^* = E_\sigma$.

Lemma 7.9.9 (Adjoint map absorbs scalar projection). Lean ✓ Stmt If T is trace-preserving, then $T^* \circ E_\sigma = E_\sigma$.

Lemma 7.9.10 (Image lies in adjoint fixed points). Lean ✓ Stmt For all X , $E_\sigma(X)$ is fixed by T^* .

Theorem 7.9.11 (Scalar map is a conditional expectation). Lean ✓ Stmt Let K be trace-preserving, let $\rho > 0$ be a positive-definite fixed point of the associated channel $T(\rho) = \sum_i K_i \rho K_i^\dagger$ (so $T(\rho) = \rho$), and assume every adjoint fixed point of T^* is a scalar multiple of $\mathbb{1}$. Then E_ρ is a conditional expectation onto the adjoint fixed-point $*$ -subalgebra.

Lemma 7.9.12 (Complete positivity of the scalar conditional expectation). Lean ✓ Stmt For any positive semidefinite σ , the scalar conditional expectation E_σ is a completely positive map.

Proof. ✓ Proof Write $S = \sqrt{\sigma}$ and define the Kraus family $K_{j,i} = |j\rangle\langle i| S$ indexed by $(j,i) \in \{1, \dots, D\}^2$. Using $|j\rangle\langle i| M |i\rangle\langle j| = M_{ii} |j\rangle\langle j|$ and summing first over i and then over j ,

$$\sum_{j,i} K_{j,i} X K_{j,i}^\dagger = \sum_j |j\rangle\langle j| \text{tr}(SX S) = \text{tr}(\sigma X) \mathbb{1},$$

where we used cyclicity of the trace and $S^2 = \sigma$. Hence $E_\sigma = (\text{tr}(\sigma))^{-1} \cdot (X \mapsto \sum_{j,i} K_{j,i} X K_{j,i}^\dagger)$ is the Kraus map scaled by the non-negative real number $(\text{tr}(\sigma))^{-1}$, and therefore completely positive. $\square$

Lemma 7.9.13 (σ -trace preservation of the scalar conditional expectation). Lean ✓ Stmt If $\text{tr}(\sigma) \neq 0$, then for every X

$$\text{tr}(\sigma E_\sigma(X)) = \text{tr}(\sigma X).$$

Proof. ✓ Proof Direct computation: $\text{tr}(\sigma \cdot \frac{\text{tr}(\sigma X)}{\text{tr}(\sigma)} \mathbb{1}) = \frac{\text{tr}(\sigma X)}{\text{tr}(\sigma)} \text{tr}(\sigma) = \text{tr}(\sigma X)$. $\square$

7.10 Stationary support

This section develops the support-projection theory behind stationary states, following [Wol12, Section 6.4, Propositions 6.10–6.11].

Lemma 7.10.1 (Lower-triangular Kraus condition implies corner invariance). Lean ✓ Stmt *Let A be an MPS tensor and let P be an orthogonal projection. If, for every physical index i ,*

$$(\mathbb{1} - P)A^iP = 0$$

then

$$P\mathcal{E}_A(PXP)P = \mathcal{E}_A(PXP)$$

for all $X \in M_D(\mathbb{C})$.

Proof. ✓ Proof Expand the transfer map and use $(\mathbb{1} - P)A^iP = 0$ to reduce each Kraus term to the P -corner. □

Lemma 7.10.2 (Support projection invariance). Lean ✓ Stmt *Let E be a quantum channel and $\rho \geq 0$ a positive semidefinite fixed point ($E(\rho) = \rho$). Let P denote the support projection of ρ . Then*

$$PE(PXP)P = E(PXP)$$

for all $X \in M_D(\mathbb{C})$.

Proof. ✓ Proof Write E in Kraus form $E(X) = \sum_i K_i X K_i^\dagger$. From $E(\rho) = \rho$ and $\rho \geq 0$ one deduces $(\mathbb{1} - P)K_iP = 0$ for all i . Apply Lemma 7.10.1. □

Definition 7.10.3 (Stationary state). Lean ✓ Stmt For an irreducible channel E on $M_D(\mathbb{C})$ (with $D \geq 1$), the *stationary state* is the unique density-matrix fixed point ρ_∞ satisfying $E(\rho_\infty) = \rho_\infty$, $\rho_\infty > 0$, and $\text{tr}(\rho_\infty) = 1$. Uniqueness follows from the quantum Perron–Frobenius theorem (Theorem 6.3.2).

Definition 7.10.4 (Stationary support). Lean ✓ Stmt The *stationary support* of an irreducible channel E is the support projection of the stationary state (Definition 7.10.3).

Theorem 7.10.5 (Stationary support is full). Lean ✓ Stmt *For an irreducible channel E , the stationary support equals the identity: $\text{supp}(\rho_\infty) = \mathbb{1}$.*

Proof. ✓ Proof The support projection P of ρ_∞ is an orthogonal projection satisfying $PE(PXP)P = E(PXP)$ for all X (Lemma 7.10.2). By irreducibility, $P = 0$ or $P = \mathbb{1}$. Since $\rho_\infty \neq 0$, we have $P \neq 0$, so $P = \mathbb{1}$. □

Remark 7.10.6 (Stationary-support formulations). The non-trivial formulations needed here are the following results from Wolf:

- the converse direction of [Wol12, Proposition 6.10]: if the support projection of every positive semidefinite fixed point is full, then the channel is irreducible;
- [Wol12, Proposition 6.11]: minimality of the stationary support among invariant projections, without assuming irreducibility.

These statements are not used elsewhere in this chapter.

7.10.1 Faithful compression onto the support sector

This subsection establishes the results on the corner algebra, $*$ -structure, and compression of a PSD fixed point onto its support sector needed for [Wol12, Corollary 6.6]: the corner carries a $\mathbb{C}$ -algebra and $*$ -structure, the two standard presentations of the corner are identified, and the compression of a PSD fixed point onto its support sector is positive definite.

Lemma 7.10.1.1 (Support projection annihilates the kernel). [Lean](#) [✓ Stmt](#) Let $\rho \succeq 0$ on $\mathbb{C}^D$ with support projection $P = \text{supp}(\rho)$. For every $v \in \mathbb{C}^D$, if $\rho v = 0$, then $Pv = 0$.

Proof. [✓ Proof](#) Diagonalize $\rho = U \text{diag}(\lambda) U^\dagger$ with $\lambda_j \geq 0$. Writing $w := U^\dagger v$, the hypothesis $\rho v = 0$ gives $\lambda_j w_j = 0$ for every j ; hence $w_j = 0$ whenever $\lambda_j > 0$. The support projection acts as $\text{diag}(\chi_{\lambda_j > 0})$ in the eigenbasis, so $Pv = U \text{diag}(\chi_{\lambda_j > 0}) w = 0$. $\square$

Definition 7.10.1.2 (Corner submodule / idempotent-corner identification). [Lean](#) [✓ Stmt](#) For an idempotent $P \in M_D(\mathbb{C})$, the subspace $\{X \mid PXP = X\}$ and the corner $\{PXP \mid X \in M_D(\mathbb{C})\}$ are identified as $\mathbb{C}$ -linear spaces by the identity on underlying matrices.

Remark 7.10.1.3 ($\mathbb{C}$ -algebra and $*$ -structure on the corner). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The corner algebra associated to an idempotent already carries a ring structure with unit P . In the matrix case it also has $\mathbb{C}$ -module and $\mathbb{C}$ -algebra structures and, under the additional assumption $P^\dagger = P$, star, star ring, and star module structures over $\mathbb{C}$.

Theorem 7.10.1.4 (Faithful compression onto the support). [Lean](#) [✓ Stmt](#) Let $\rho \succeq 0$ on $\mathbb{C}^D$ with support projection $P = \text{supp}(\rho)$, and let $V : \mathbb{C}^D \rightarrow \mathbb{C}^k$ be a compression isometry satisfying $VV^\dagger = \mathbb{1}_k$ and $V^\dagger V = P$. Then the compression

$$V \rho V^\dagger \in M_k(\mathbb{C})$$

is positive definite.

Proof. [✓ Proof](#) Hermiticity follows from Hermiticity of ρ and the product-star identity. For strict positivity, let $w \neq 0$, set $u := V^\dagger w$, and note that $VV^\dagger = \mathbb{1}_k$ forces $u \neq 0$, while $V^\dagger V = P$ forces $Pu = u$. The quadratic form computes as $w^\dagger (V \rho V^\dagger) w = u^\dagger \rho u$, which is non-negative by positive semidefiniteness. If it vanishes, then $\rho u = 0$, and by Lemma 7.10.1.1 also $Pu = 0$, contradicting $Pu = u \neq 0$. $\square$

Theorem 7.10.1.5 (Corner-restricted fixed points form a $*$ -algebra). [Lean](#) [✓ Stmt](#) Let $T^*(Y) = \sum_i K_i^\dagger Y K_i$ be a trace-preserving Schwarz map, let $\rho \succeq 0$ satisfy $T(\rho) = \rho$ for the Schrödinger map $T(X) = \sum_i K_i X K_i^\dagger$, and let $Q = \text{supp}(\rho)$ be the support projection of ρ . Then

$$\{Y \in Q M_D(\mathbb{C}) Q \mid Q T^*(Y) Q = Y\}$$

is a $*$ -subalgebra of the corner algebra $Q M_D(\mathbb{C}) Q$.

Proof. [✓ Proof](#) The stated set is the fixed-point set of the compressed map $\tilde{T}^*$ on the support sector. Since ρ is a fixed point of T , the support projection satisfies $(\mathbb{1} - Q)K_i Q = 0$ (Theorem 7.10.2), so the compressed family $QK_i Q$ is supported on the corner and is trace-preserving there, $\sum_i (QK_i Q)^\dagger (QK_i Q) = Q$. Compress this family to the support sector: the compressed map is unital, and the compression of ρ onto the sector is positive definite (Theorem 7.10.1.4) and is a fixed point of the compressed Schrödinger map. The structure theorem for fixed points of unital Schwarz maps with a positive-definite fixed point (Theorem 4.6.11) makes the compressed fixed points a $*$ -algebra. The compression isomorphism is multiplicative and star-preserving, intertwines the compressed adjoint map with the corner-restricted map, and transports this $*$ -algebra structure back to the corner. $\square$

7.11 Wedderburn decomposition of the fixed-point algebra

This section proves the structure theorem for the fixed-point algebra of a trace-preserving Kraus map, following [Wol12, Theorems 6.12–6.14]. The proof uses the Jacobson radical characterization of semisimplicity together with the Wedderburn–Artin structure theorem for finite-dimensional semisimple algebras over an algebraically closed field.

The algebraic core is the semisimplicity of an arbitrary $*$ -subalgebra of a complex matrix algebra, from which the abstract Wedderburn–Artin decomposition follows. The fixed-point algebra of a trace-preserving Kraus map is then the special case used in this section.

Lemma 7.11.1 (A $*$ -subalgebra of a matrix algebra is semisimple). Lean ✓ Stmt *Every $*$ -subalgebra of $M_D(\mathbb{C})$ is a semisimple ring.*

Proof. ✓ Proof The subalgebra is finite-dimensional, hence Artinian, so its Jacobson radical J is nilpotent. If $x \in J$ then $x^*x \in J$, since the radical is a two-sided ideal. In the matrix algebra x^*x is positive semidefinite, and a positive semidefinite nilpotent matrix vanishes: its trace is a nilpotent complex number, hence zero, and the trace of x^*x vanishes only when $x = 0$. Thus $J = 0$ and the subalgebra is semisimple. $\square$

Theorem 7.11.2 (Abstract Wedderburn–Artin decomposition for $*$ -subalgebras). Lean ✓ Stmt *There exist $n \in \mathbb{N}$ and dimensions $d_1, \dots, d_n \geq 1$ such that every $*$ -subalgebra of $M_D(\mathbb{C})$ is $\mathbb{C}$ -algebra isomorphic to*

$$\prod_{k=1}^n M_{d_k}(\mathbb{C}).$$

Proof. ✓ Proof Combine semisimplicity (Lemma 7.11.1) with the Wedderburn–Artin structure theorem for finite-dimensional semisimple algebras over the algebraically closed field $\mathbb{C}$. $\square$

The abstract decomposition records only the ring structure. To realize it by a single unitary change of basis, as in [Wol12, Theorem 6.14], one uses the spatial feature of a $*$ -subalgebra: it reduces orthogonally. Whenever a subspace is invariant, so is its orthogonal complement, so the representation space splits into mutually orthogonal invariant pieces rather than merely into a direct sum of them.

Lemma 7.11.3 (Orthogonal reducibility of a $*$ -subalgebra). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, and let $W \subseteq \mathbb{C}^D$ be a subspace invariant under every member of S , meaning $AW \subseteq W$ for all $A \in S$. Then the orthogonal complement $W^\perp$ is also invariant under every member of S : for each $A \in S$,*

$$AW^\perp \subseteq W^\perp.$$

Proof. ✓ Proof Fix $A \in S$. Since S is closed under the adjoint, $A^\dagger \in S$, and therefore $A^\dagger W \subseteq W$. For $v \in W^\perp$ and $w \in W$,

$$\langle w, Av \rangle = \langle A^\dagger w, v \rangle = 0,$$

because $A^\dagger w \in W$ while v is orthogonal to W . Hence $Av \in W^\perp$. $\square$

Lemma 7.11.4 (Invariant subspaces split off orthogonally). Lean ✓ Stmt *With S and W as in Lemma 7.11.3, the representation space decomposes as the orthogonal direct sum*

$$\mathbb{C}^D = W \oplus W^\perp,$$

in which the orthogonal complement $W^\perp$ is invariant under every member of S .

Proof. ✓ Proof In a finite-dimensional inner product space the orthogonal complement of a subspace is a vector-space complement, which gives the orthogonal direct sum. Invariance of $W^\perp$ is Lemma 7.11.3. □

Orthogonal reducibility is the inductive step behind a full orthogonal decomposition. A subspace with no proper nonzero invariant subspace is irreducible; a subspace that has one splits off its orthogonal complement, and both pieces are again invariant and strictly smaller. Iterating on dimension exhausts the space into irreducible pieces, mutually orthogonal because at each step the two pieces are.

Definition 7.11.5 (Irreducible invariant subspace). Lean ✓ Stmt Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. A subspace $W \subseteq \mathbb{C}^D$ is irreducible under S when it is nonzero, invariant under every member of S , and its only invariant subspaces are $\{0\}$ and W itself.

Lemma 7.11.6 (Orthogonal decomposition of an invariant subspace). Lean ✓ Stmt Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W \subseteq \mathbb{C}^D$ be invariant under every member of S . Then W is the sum of a finite family of pairwise orthogonal subspaces contained in W , each irreducible under S .

Proof. ✓ Proof Induct on the dimension of W . If W is zero the empty family works, and if W is irreducible the family $\{W\}$ works. Otherwise W has a proper nonzero invariant subspace U . By Lemma 7.11.4 the orthogonal complement of U inside W is invariant, and both U and that complement are strictly smaller than W and span it. Their inductive decompositions consist of subspaces lying inside U and inside its complement respectively, hence are orthogonal to one another; the two families together form the required family. □

Theorem 7.11.7 (Orthogonal irreducible decomposition). Lean ✓ Stmt Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. Then $\mathbb{C}^D$ is the sum of a finite family of pairwise orthogonal subspaces, each irreducible under S ; that is, $\mathbb{C}^D$ is an orthogonal direct sum of irreducible invariant subspaces.

Proof. ✓ Proof Apply Lemma 7.11.6 to the whole space $W = \mathbb{C}^D$, which is invariant under every member of S . □

The irreducibility notion just used is simplicity in the language of modules. The members of S act on $\mathbb{C}^D$ as the operators their matrices represent, making $\mathbb{C}^D$ a module over the algebra S .

Lemma 7.11.8 (Irreducible invariant subspaces are the simple submodules). Lean Lean ✓ Stmt Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, acting on $\mathbb{C}^D$. A complex subspace $W \subseteq \mathbb{C}^D$ is invariant under every member of S if and only if W is the underlying complex subspace of a submodule of $\mathbb{C}^D$ over S . Under this identification an invariant subspace is irreducible under S precisely when, as a module over S , it is simple: nonzero, with no submodule other than $\{0\}$ and itself.

Proof. ✓ Proof The underlying set of a submodule of $\mathbb{C}^D$ over S is closed under addition and under the action of every member of S ; since the complex scalars lie in S , it is a complex subspace, and it is invariant. Conversely an invariant complex subspace W is closed under addition and under the action of every member of S , so W underlies a submodule q of $\mathbb{C}^D$ over S with the same members. Under this correspondence the submodules below q are exactly the invariant subspaces below W :

$$\{q' \subseteq q : q' \text{ a submodule over } S\} = \{W' \subseteq W : W' \text{ invariant under } S\}.$$

Hence q has a submodule other than $\{0\}$ and q exactly when W has an invariant subspace other than $\{0\}$ and W : simplicity of q as a module over S is irreducibility of W under S . □

Lemma 7.11.9 (Euclidean space is a semisimple module over the subalgebra). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. As a module over S , the space $\mathbb{C}^D$ is semisimple: every submodule admits a complementary submodule.*

Proof. [✓ Proof](#) The ring S is semisimple (Lemma 7.11.1), and every module over a semisimple ring is semisimple. $\square$

The next step towards the structure theorem groups the irreducible pieces by isomorphism type. The grouping is governed by Schur's lemma: a map commuting with S between two pieces is either zero or an isomorphism, and over the complex numbers the commutant of S on a single piece is the scalars.

Lemma 7.11.10 (Schur's lemma: an intertwiner of irreducible pieces is zero or an isomorphism). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be irreducible under S . Let $f: \mathbb{C}^D \rightarrow \mathbb{C}^D$ be a linear operator that carries W into W' and commutes on W with every member of S , that is,*

$$f(Ax) = A(fx)$$

for every $A \in S$ and every $x \in W$. Then either f is zero on W , or f is injective on W and maps W onto W' .

Proof. [✓ Proof](#) The vectors of W annihilated by f form a subspace of W that is invariant under S , because f commutes with S on W . By irreducibility of W this kernel is either all of W , in which case f is zero on W , or trivial, in which case f is injective on W . Likewise the image of W under f is a subspace of W' invariant under S , so by irreducibility of W' it is either trivial, again forcing f to be zero on W , or all of W' . Hence f is zero on W , or it is injective on W with image W' . $\square$

Lemma 7.11.11 (Scalar commutant on an irreducible piece). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W \subseteq \mathbb{C}^D$ be irreducible under S . Let $f: \mathbb{C}^D \rightarrow \mathbb{C}^D$ be a linear operator that maps W into itself and commutes on W with every member of S , that is,*

$$f(Ax) = A(fx)$$

for every $A \in S$ and every $x \in W$. Then f acts on W as a scalar: there is a complex number c with

$$fx = cx$$

for every $x \in W$.

Proof. [✓ Proof](#) Restricting f to W gives an operator on a nonzero finite-dimensional complex vector space, which over the complex numbers has an eigenvalue c together with an eigenvector inside W . The set of vectors of W on which f acts as multiplication by c is a subspace contained in W ; it is nonzero because it contains that eigenvector, and it is invariant under S because f commutes with every member of S on W while the scalar c commutes with everything. By irreducibility this subspace is all of W , so f acts as multiplication by c throughout W . $\square$

Schur's lemma turns the qualitative notion of two pieces "looking the same" into a usable relation. Call a linear operator on $\mathbb{C}^D$ an intertwiner from W into W' when it carries W into W' and commutes on W with every member of S . Two pieces are of the same type when some intertwiner between them does not vanish on the source; on irreducible pieces such an intertwiner is automatically an isomorphism.

Definition 7.11.12 (Intertwiner between subspaces). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be subspaces. A linear operator $f: \mathbb{C}^D \rightarrow \mathbb{C}^D$ is an intertwiner from W into W' when it carries W into W' and, for every $A \in S$ and every $x \in W$,

$$f(Ax) = A(fx).$$

An intertwiner whose source is irreducible scales the inner product by one nonnegative real factor; when the intertwiner does not vanish on the source, that factor is positive, so after rescaling it is an isometry. This is the metric ingredient that turns each same-type isomorphism into a partial isometry.

Lemma 7.11.13 (Intertwiner scales the inner product). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, let $W \subseteq \mathbb{C}^D$ be irreducible under S , and let f be an intertwiner from W into a subspace W' . Then there is a real number $c \geq 0$ such that for all $x, y \in W$,

$$\langle f(x), f(y) \rangle = c \langle x, y \rangle.$$

Proof. [✓ Proof](#) Following f with its adjoint and the orthogonal projection P_W onto W gives an operator $g = P_W f^* f$ that maps W into itself, since the projection lands in W . On W it commutes with every member of S : for $A \in S$ the adjoint A^* is again a member, and combining the intertwining identity of f with the pairing of the projected adjoint against W gives $g(Ax) = A(gx)$ for $x \in W$. By Schur's lemma (Lemma 7.11.11) the operator g acts on W as a scalar, $gx = \lambda x$. Pairing with a vector $y \in W$ and using that P_W is invisible against W gives $\langle f(x), f(y) \rangle = \langle g(x), y \rangle = \lambda \langle x, y \rangle$. Pairing a nonzero $x \in W$ with itself shows that $c = \lambda = \|f(x)\|^2 / \|x\|^2$, a nonnegative real. $\square$

Lemma 7.11.14 (Nonzero intertwiner scales by a positive factor). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, let $W \subseteq \mathbb{C}^D$ be irreducible under S , and let f be an intertwiner from W into a subspace W' that does not vanish on W . Then there is a real number $c > 0$ such that for all $x, y \in W$,

$$\langle f(x), f(y) \rangle = c \langle x, y \rangle.$$

Proof. [✓ Proof](#) The nonnegative factor from Lemma 7.11.13 satisfies $\|f(x)\|^2 = c \|x\|^2$. Choosing $x \in W$ with $f(x) \neq 0$ forces both squared norms to be positive, so $c > 0$. $\square$

Definition 7.11.15 (Same type for invariant subspaces). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be invariant under S . The pieces W and W' are of the same type when there is an intertwiner from W into W' that is not zero on W .

Lemma 7.11.16 (Same type is the existence of an isomorphism). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be irreducible under S . Then W and W' are of the same type if and only if there is an intertwiner from W into W' that is injective on W and maps W onto W' .

Proof. [✓ Proof](#) If W and W' are of the same type, then an intertwiner that is nonzero on W is, by Schur's lemma (Lemma 7.11.10), injective on W with image W' . Conversely, an intertwiner injective on W with image W' cannot vanish on W , since W' is nonzero, so the two pieces are of the same type. $\square$

Lemma 7.11.17 (Same-type pieces are unitarily identified). [Lean](#) [✓ Stmt](#) Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be irreducible under S . If W is of the same type as W' , then there is an intertwiner u from W into W' that maps W onto W' and preserves the inner product on W : for all $x, y \in W$,

$$\langle u(x), u(y) \rangle = \langle x, y \rangle.$$

Proof. ✓ Proof Same type gives, by Lemma 7.11.16, an intertwiner f that is injective on W with image W' . Since W' is nonzero, f does not vanish on W , so by Lemma 7.11.14 there is a real number $c > 0$ with $\langle f(x), f(y) \rangle = c \langle x, y \rangle$ for all $x, y \in W$. Set $u = f/\sqrt{c}$. Scaling by the nonzero factor $1/\sqrt{c}$ keeps the commuting identity and the image W' , and turns the inner-product factor into one,

$$\langle u(x), u(y) \rangle = \frac{1}{c} \langle f(x), f(y) \rangle = \langle x, y \rangle,$$

so u maps W onto W' and is an isometry on W . □

Lemma 7.11.18 (Reflexivity of the same-type relation). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W \subseteq \mathbb{C}^D$ be irreducible under S . Then W is of the same type as itself.*

Proof. ✓ Proof The identity operator carries W into itself, commutes with every member of S , and is nonzero on W because W is nonzero. □

Lemma 7.11.19 (Symmetry of the same-type relation). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be irreducible under S . If W is of the same type as W' , then W' is of the same type as W .*

Proof. ✓ Proof An isomorphism f of W onto W' has an inverse g , obtained by projecting onto W' , inverting on the subspace, and including back into W . For every $A \in S$ and $w \in W'$, since f is an intertwiner,

$$g(Aw) = f^{-1}(Aw) = f^{-1}(A f(f^{-1}w)) = f^{-1}(f(A f^{-1}w)) = A g(w),$$

so g is an intertwiner from W' into W that is injective on W' with image W , and the two pieces are of the same type in the reverse order. □

Lemma 7.11.20 (Transitivity of the same-type relation). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W', W'' \subseteq \mathbb{C}^D$ be irreducible under S . If W is of the same type as W' and W' is of the same type as W'' , then W is of the same type as W'' .*

Proof. ✓ Proof Composing an isomorphism f of W onto W' with an isomorphism g of W' onto W'' gives a map $g \circ f$ for which, for every $A \in S$ and $x \in W$,

$$(g \circ f)(Ax) = g(f(Ax)) = g(A f(x)) = A g(f(x)) = A (g \circ f)(x),$$

so it is an intertwiner from W into W'' that is injective on W with image W'' , which is nonzero on W because W'' is nonzero. □

Lemma 7.11.21 (The orthogonal projection onto an invariant subspace commutes with the subalgebra). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W \subseteq \mathbb{C}^D$ be invariant under S , with orthogonal projection P onto W . Then for every $A \in S$,*

$$P(Ax) = A(Px)$$

holds for every $x \in \mathbb{C}^D$.

Proof. ✓ Proof Decompose $x = Px + P^\perp x$ into its projection onto W and its orthogonal part, where $P^\perp$ is the projection onto $W^\perp$. Each member of S sends Px into W (invariance) and $P^\perp x$ into $W^\perp$ (orthogonal reducibility, Lemma 7.11.3), so

$$P(Ax) = P(APx + AP^\perp x) = APx + P(AP^\perp x) = APx,$$

because $APx \in W$ is fixed by P and $AP^\perp x \in W^\perp$ is annihilated by P . □

Lemma 7.11.22 (Pieces of different type are orthogonal). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $W, W' \subseteq \mathbb{C}^D$ be irreducible under S . If W and W' are not of the same type, then W and W' are orthogonal.*

Proof. [✓ Proof](#) The orthogonal projection onto W' commutes with every member of S (Lemma 7.11.21) and carries W into W' , so it is an intertwiner from W into W' . By Schur's lemma (Lemma 7.11.10) it is either an isomorphism, which would make the two pieces of the same type, or zero on W . Since the pieces are not of the same type, the projection $P_{W'}$ onto W' vanishes on W . For every $x \in W$ and $y \in W'$ we have $P_{W'}y = y$ and $P_{W'}x = 0$, so self-adjointness of $P_{W'}$ gives

$$\langle x, y \rangle = \langle x, P_{W'}y \rangle = \langle P_{W'}x, y \rangle = \langle 0, y \rangle = 0,$$

that is, W lies in the orthogonal complement of W' . □

Each isotypic component is the supremum of one same-type class; the suprema of two different classes are orthogonal, and the classes cover the whole family, so the components span $\mathbb{C}^D$.

Lemma 7.11.23 (Suprema of cross-type classes are orthogonal). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, and let $\mathcal{D}_1, \mathcal{D}_2$ be two sets of subspaces irreducible under S . If no piece of $\mathcal{D}_1$ is of the same type as any piece of $\mathcal{D}_2$, then*

$$\bigvee_{W \in \mathcal{D}_1} W \perp \bigvee_{W' \in \mathcal{D}_2} W'.$$

Proof. [✓ Proof](#) Each pair $W \in \mathcal{D}_1, W' \in \mathcal{D}_2$ is of different type, hence orthogonal by Lemma 7.11.22. Orthogonality to a fixed subspace is preserved by suprema on each side, so the supremum of $\mathcal{D}_1$ is orthogonal to the supremum of $\mathcal{D}_2$. □

An irreducible subspace of the supremum of a same-type class need not be one of its pieces, but its type is determined by the class.

Lemma 7.11.24 (Irreducible subspaces of a supremum inherit a type). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$, let $\mathcal{D}$ be a family of subspaces irreducible under S , and let W be a subspace irreducible under S with $W \leq \bigvee_{W' \in \mathcal{D}} W'$. Then W is of the same type as some piece of $\mathcal{D}$.*

Proof. [✓ Proof](#) Suppose W is of a different type than every piece of $\mathcal{D}$. Pieces of different type are orthogonal (Lemma 7.11.22), so W is orthogonal to every piece of $\mathcal{D}$, hence to their supremum, hence to itself. An irreducible subspace is nonzero, a contradiction. □

Theorem 7.11.25 (Isotypic-component decomposition). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. Then $\mathbb{C}^D$ is the supremum of a finite family $\mathcal{C}$ of pairwise orthogonal subspaces, the isotypic components, and each component $C \in \mathcal{C}$ is the supremum of a finite, nonempty class $\mathcal{D}_C$ of pairwise orthogonal subspaces irreducible under S that are pairwise of the same type:*

$$C = \bigvee_{W \in \mathcal{D}_C} W.$$

Moreover, for distinct components $C \neq C'$, no subspace irreducible under S contained in C is of the same type as any subspace irreducible under S contained in C' .

Proof. ✓ **Proof** The orthogonal irreducible decomposition (Theorem 7.11.7) supplies a finite, pairwise orthogonal family $\mathcal{D}$ of irreducible pieces with $\bigvee_{W \in \mathcal{D}} W = \mathbb{C}^D$. For each piece W let $\mathcal{D}_W$ be the class of pieces of $\mathcal{D}$ of the same type as W , and let C_W be its supremum. Each piece lies in its own class by reflexivity (Lemma 7.11.18), so each class is nonempty, the classes cover $\mathcal{D}$, and the components span $\mathbb{C}^D$. Each class is contained in $\mathcal{D}$, so its pieces are pairwise orthogonal. Two pieces of one class are of the same type with each other through the representative, by symmetry and transitivity (Lemmas 7.11.19 and 7.11.20). If two representatives are of the same type, transitivity makes their classes equal, hence their components equal; so two distinct components come from representatives W, W' that are not of the same type. Write $\sim$ for the same-type relation. For $a \in \mathcal{D}_W$ and $b \in \mathcal{D}_{W'}$ one has $W \sim a$ and $W' \sim b$ by definition of the classes; were $a \sim b$, symmetry and transitivity would give

$$W \sim a \sim b \sim W',$$

hence $W \sim W'$, contradicting that W and W' are not of the same type. So no piece of $\mathcal{D}_W$ is of the same type as any piece of $\mathcal{D}_{W'}$, and the components are orthogonal (Lemma 7.11.23).

Finally, let $V \leq C_W$ and $V' \leq C_{W'}$ be irreducible under S with $C_W \neq C_{W'}$. By Lemma 7.11.24 there are pieces $a \in \mathcal{D}_W$ and $b \in \mathcal{D}_{W'}$ with $V \sim a$ and $V' \sim b$. Were $V \sim V'$, symmetry and transitivity would give

$$W \sim a \sim V \sim V' \sim b \sim W',$$

contradicting again that W and W' are not of the same type. □

The pieces of one class are unitarily identified copies of a single irreducible piece. An orthonormal basis of one piece therefore spreads, through the inner-product-preserving intertwiners, to an adapted orthonormal basis of the whole component, and in this basis every member of S acts by one matrix on the irreducible index, identically across the multiplicity index. This is the tensor factorization of the isotypic components behind the block form $M_{d_k}(\mathbb{C}) \otimes \mathbb{1}_{m_k}$ of [Wol12, Theorem 6.14].

Theorem 7.11.26 (Adapted orthonormal basis of an isotypic component). Lean ✓ **Stmt** *Let S be a *-subalgebra of $M_D(\mathbb{C})$ and let $\mathcal{D}$ be a finite, nonempty family of pairwise orthogonal subspaces of $\mathbb{C}^D$, irreducible under S and pairwise of the same type. Write m for the number of pieces of $\mathcal{D}$; all pieces share one dimension d . Then the component $C = \bigvee_{W \in \mathcal{D}} W$ has an orthonormal basis $(f_{i,j})_{0 \leq i < m, 0 \leq j < d}$ such that for every $A \in S$ there is a matrix $B \in M_d(\mathbb{C})$ with, for all $i < m$ and $j < d$,*

$$A f_{i,j} = \sum_{j'=0}^{d-1} B_{j'j} f_{i,j'}.$$

In the adapted basis the member A acts on C by the matrix B on the irreducible index, identically across the multiplicity index; after reordering the indices this is the block $M_d(\mathbb{C}) \otimes \mathbb{1}_m$ of [Wol12, Theorem 6.14]. The vectors $(f_{i,j})_{0 \leq j < d}$ of the i -th copy span one of the pieces of $\mathcal{D}$.

Proof. ✓ **Proof** Fix a piece $W \in \mathcal{D}$, of dimension d , with an orthonormal basis $e_0, \dots, e_{d-1}$, and enumerate $\mathcal{D} = \{W'_0, \dots, W'_{m-1}\}$. Every piece of $\mathcal{D}$ is of the same type as W , by hypothesis for the other pieces and by reflexivity (Lemma 7.11.18) for W itself, so Lemma 7.11.17 provides intertwiners u_i mapping W onto W'_i and preserving the inner product on W . Set $f_{i,j} = u_i(e_j)$. Within one copy the vectors $f_{i,0}, \dots, f_{i,d-1}$ are orthonormal because u_i preserves inner products; across copies they are orthogonal because distinct pieces of $\mathcal{D}$ are. Since u_i maps W onto W'_i , the vectors of the i -th copy span W'_i ; in particular each W'_i has dimension d , and all the vectors

together span C . For $A \in S$ the piece W is invariant, so $Ae_j = \sum_{j'} B_{j'j} e_{j'}$ with $B_{j'j} = \langle e_{j'}, Ae_j \rangle$, and the intertwining identity gives

$$Af_{i,j} = Au_i(e_j) = u_i(Ae_j) = \sum_{j'} B_{j'j} u_i(e_{j'}) = \sum_{j'} B_{j'j} f_{i,j'},$$

the same matrix B for every copy i . $\square$

Theorem 7.11.27 (Spatial tensor factorization of the isotypic components). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. Then $\mathbb{C}^D$ is the supremum of a finite family of pairwise orthogonal isotypic components, and each component carries an orthonormal basis $(f_{i,j})$, indexed by a multiplicity index $i < m$ and an irreducible index $j < d$, such that every $A \in S$ acts by*

$$Af_{i,j} = \sum_{j'=0}^{d-1} B_{j'j} f_{i,j'}$$

for a matrix $B \in M_d(\mathbb{C})$ depending only on A and the component. In the adapted bases the action of S on each component is by matrix-times-identity blocks, the form $M_{d_k}(\mathbb{C}) \otimes \mathbb{1}_{m_k}$ of [Wol12, Theorem 6.14].

Proof. [✓ Proof](#) The isotypic-component decomposition (Theorem 7.11.25) writes $\mathbb{C}^D$ as the supremum of finitely many pairwise orthogonal components, each the supremum of a finite, nonempty class of pairwise orthogonal irreducible pieces of one type. Applying Theorem 7.11.26 to each class equips each component with its adapted orthonormal basis. $\square$

The adapted orthonormal bases of the pairwise orthogonal isotypic components concatenate into a single orthonormal basis of $\mathbb{C}^D$, indexed by a component index k , a multiplicity index i , and an irreducible index j . In this basis every member of S acts block-diagonally, one matrix-times-identity block per component. A $*$ -subalgebra contains the identity matrix, so every component carries irreducible pieces of S : this is the unital case of the block representation invoked by [Wol12, Theorem 6.14], without the zero block.

Theorem 7.11.28 (Adapted orthonormal basis of the whole space). [Lean](#) [✓ Stmt](#) *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. There are $K \in \mathbb{N}$, positive dimensions $d_0, \dots, d_{K-1}$ and multiplicities $m_0, \dots, m_{K-1}$, and an orthonormal basis $(f_{k,i,j})_{k < K, i < m_k, j < d_k}$ of $\mathbb{C}^D$ such that for every $A \in S$ there are matrices $B_k \in M_{d_k}(\mathbb{C})$ with, for all $k < K$, $i < m_k$, and $j < d_k$,*

$$Af_{k,i,j} = \sum_{j'=0}^{d_k-1} (B_k)_{j'j} f_{k,i,j'}.$$

In this basis the member A acts on the k -th component as $\mathbb{1}_{m_k} \otimes B_k$. Each row $(f_{k,i,j})_{0 \leq j < d_k}$ spans a subspace irreducible under S , and rows drawn from distinct components span subspaces that are never of the same type. Only the containment direction is asserted: every member of S takes this form. The equality with the block algebra, including the reverse inclusion, is Theorem 7.11.32.

Proof. [✓ Proof](#) The isotypic-component decomposition (Theorem 7.11.25) writes $\mathbb{C}^D$ as the supremum of finitely many pairwise orthogonal isotypic components, each the supremum of a finite, nonempty same-type class of pairwise orthogonal irreducible pieces. Theorem 7.11.26 equips the k -th component with an adapted orthonormal family $(f_{k,i,j})_{i < m_k, j < d_k}$ spanning it, where m_k is the number of pieces of the class, positive because the class is nonempty, and d_k is their common dimension, positive because irreducible pieces are nonzero. Concatenating the families over k

yields an orthonormal family: two vectors of one component are orthonormal by the adapted family, and two vectors of distinct components are orthogonal because the components are. The concatenated family spans $\mathbb{C}^D$ because each component is spanned by its family and the components have supremum $\mathbb{C}^D$; a spanning orthonormal family is an orthonormal basis. The action property holds componentwise by the adapted families. Each row of the concatenated family spans one of the irreducible pieces of its component's class (Theorem 7.11.26), and irreducible subspaces of distinct components are never of the same type (Theorem 7.11.25), so neither are the row spans of distinct components. $\square$

The change of basis to an orthonormal basis indexed by such triples is implemented by a unitary matrix, and it carries every matrix acting blockwise on the basis to a block-diagonal matrix.

Lemma 7.11.29 (Unitary change of basis to block-diagonal form). Lean ✓ Stmt *Let $(f_{k,i,j})_{k < K, i < m_k, j < d_k}$ be an orthonormal basis of $\mathbb{C}^D$. There are an identification of the index set of triples (k, i, j) with $\{0, \dots, D-1\}$, realizing $\sum_k d_k m_k = D$, and a unitary $U \in M_D(\mathbb{C})$, whose columns are the basis vectors, such that every matrix $A \in M_D(\mathbb{C})$ acting on the basis by*

$$A f_{k,i,j} = \sum_{j'=0}^{d_k-1} (B_k)_{j'j} f_{k,i,j'}$$

for matrices $B_k \in M_{d_k}(\mathbb{C})$ satisfies

$$U^\dagger A U = \bigoplus_{k=0}^{K-1} \mathbb{1}_{m_k} \otimes B_k.$$

Proof. ✓ Proof Let U be the change-of-basis matrix from the standard orthonormal basis of $\mathbb{C}^D$ to $(f_{k,i,j})$, so the columns of U are the basis vectors; a change of basis between orthonormal bases is unitary. Both bases have D elements, which identifies the index set of triples (k, i, j) with $\{0, \dots, D-1\}$ and gives $\sum_k d_k m_k = D$. For A acting as displayed, the matrix $U^\dagger A U = U^{-1} A U$ is the matrix of the operator $x \mapsto Ax$ in the basis $(f_{k,i,j})$, with entries

$$\langle f_{k',i',j'}, A f_{k,i,j} \rangle = \sum_{j''} (B_k)_{j''j} \langle f_{k',i',j'}, f_{k,i,j''} \rangle = \delta_{k'k} \delta_{i'i} (B_k)_{j'j}$$

by the action property and orthonormality. This is the block-diagonal matrix with blocks $\mathbb{1}_{m_k} \otimes B_k$. $\square$

Theorem 7.11.30 (Global unitary block-diagonal form). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. There are $K \in \mathbb{N}$, positive dimensions $d_0, \dots, d_{K-1}$ and multiplicities $m_0, \dots, m_{K-1}$ with $\sum_k d_k m_k = D$, realized by an explicit identification of the index sets, and a unitary $U \in M_D(\mathbb{C})$ such that every $A \in S$ satisfies*

$$U^\dagger A U = \bigoplus_{k=0}^{K-1} \mathbb{1}_{m_k} \otimes B_k$$

for matrices $B_k \in M_{d_k}(\mathbb{C})$ depending on A . Up to reordering the two tensor factors of each block this is the containment direction of the unital case of the block representation of [Wol12, Theorem 6.14]: S is carried by U into the block algebra $\bigoplus_k \mathbb{1}_{m_k} \otimes M_{d_k}(\mathbb{C})$. The equality, including the reverse inclusion, is Theorem 7.11.32.

Proof. ✓ Proof Let $(f_{k,i,j})$ be the adapted orthonormal basis of Theorem 7.11.28 and let U be the unitary supplied by Lemma 7.11.29 for this basis. Every $A \in S$ acts on the basis by matrices B_k on the irreducible index, identically across the multiplicity index, so the lemma gives $U^\dagger A U = \bigoplus_k \mathbb{1}_{m_k} \otimes B_k$. $\square$

The reverse inclusion of the block representation rests on the double-commutant characterization: in finite dimensions a $*$ -subalgebra contains every matrix that commutes with its commutant. The proof is the Jacobson density theorem for $\mathbb{C}^D$ as a semisimple module over the subalgebra.

Lemma 7.11.31 (Membership through the double commutant). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$ and let $T \in M_D(\mathbb{C})$. If T commutes with every linear operator on $\mathbb{C}^D$ that commutes with all members of S , then $T \in S$.*

Proof. ✓ Proof The space $\mathbb{C}^D$ is a semisimple module over S (Lemma 7.11.9), and the linear operators commuting with all members of S are exactly the endomorphisms of this module. The hypothesis therefore makes T linear over the endomorphism ring of the module. By the Jacobson density theorem, for every finite set of vectors there is a member of S acting on them as T does. Applied to a basis of $\mathbb{C}^D$, this produces a member of S whose action agrees with T on a basis, hence equals T . $\square$

Combining the containment direction with the double-commutant criterion yields the full block representation of a finite-dimensional $*$ -subalgebra, the unital case of Eq. (1.39) of [Wol12] invoked by [Wol12, Theorem 6.14]: up to a unitary change of basis, S equals the block algebra $\bigoplus_k \mathbb{1}_{m_k} \otimes M_{d_k}(\mathbb{C})$.

Theorem 7.11.32 (Full block form of a $*$ -subalgebra). Lean ✓ Stmt *Let S be a $*$ -subalgebra of $M_D(\mathbb{C})$. There are $K \in \mathbb{N}$, positive dimensions $d_0, \dots, d_{K-1}$ and multiplicities $m_0, \dots, m_{K-1}$ with $\sum_k d_k m_k = D$, realized by an explicit identification of the index sets, and a unitary $U \in M_D(\mathbb{C})$ such that a matrix $A \in M_D(\mathbb{C})$ belongs to S exactly when*

$$U^\dagger A U = \bigoplus_{k=0}^{K-1} \mathbb{1}_{m_k} \otimes B_k$$

for some matrices $B_k \in M_{d_k}(\mathbb{C})$; that is, up to reordering the two tensor factors of each block,

$$S = U \left(\bigoplus_{k=0}^{K-1} \mathbb{1}_{m_k} \otimes M_{d_k}(\mathbb{C}) \right) U^\dagger.$$

This is the block representation of a finite-dimensional $$ -algebra in Eq. (1.39) of [Wol12], invoked by [Wol12, Theorem 6.14], in the unital case: a $*$ -subalgebra contains the identity matrix, so there is no zero block.*

Proof. ✓ Proof Let $(f_{k,i,j})$ be the adapted orthonormal basis of Theorem 7.11.28, write $W_{k,i}$ for the span of the row $(f_{k,i,j})_{j < d_k}$, and let U be the unitary supplied by Lemma 7.11.29 for this basis: any matrix A whose action on the adapted basis is $A f_{k,i,j} = \sum_{j'} (B_k)_{j'j} f_{k,i,j'}$ satisfies the displayed conjugation identity; in particular every member of S does, which is the containment direction.

For the reverse inclusion, fix blocks $(B_k)_k$ and let T be the matrix acting on the adapted basis by $T f_{k,i,j} = \sum_{j'} (B_k)_{j'j} f_{k,i,j'}$. By Lemma 7.11.31 it suffices to show that T commutes with

every operator g that commutes with all members of S ; we compute the matrix of such a g in the adapted basis. For a component index k and multiplicity indices i, i' , consider the operator

$$\tau : v \mapsto \sum_j \langle f_{k,i',j}, v \rangle f_{k,i,j},$$

which reads off the coefficients of a vector along the row (k, i') and rebuilds them along the row (k, i) . Because every member of S acts on the rows (k, i') and (k, i) by the same matrix, τ commutes with all members of S , and so does $\tau \circ g$. The composite maps $\mathbb{C}^D$ into $W_{k,i}$, and $W_{k,i}$ is irreducible, so by Schur's lemma (Lemma 7.11.11) there is $\gamma_{i'i}^{(k)} \in \mathbb{C}$ with $\tau(gx) = \gamma_{i'i}^{(k)} x$ for all $x \in W_{k,i}$. Evaluating at $x = f_{k,i,j}$ and comparing coefficients gives

$$\langle f_{k,i',j'}, g f_{k,i,j} \rangle = \gamma_{i'i}^{(k)} \delta_{j'j}.$$

Across components the matrix of g vanishes: for $k' \neq k$,

$$\langle f_{k',i',j'}, g f_{k,i,j} \rangle = 0 :$$

with τ built from the rows (k', i') of another component, the composite $\tau \circ g$ carries $W_{k,i}$ into $W_{k',i'}$ and commutes with S on $W_{k,i}$, so a nonzero such entry would make it an intertwiner that does not vanish on $W_{k,i}$, giving $W_{k,i}$ and $W_{k',i'}$ the same type and contradicting Theorem 7.11.28. Expanding $g f_{k,i,j}$ in the adapted basis therefore gives

$$g f_{k,i,j} = \sum_{i'} \gamma_{i'i}^{(k)} f_{k,i',j} :$$

in the adapted basis g is block diagonal with blocks $C_k \otimes \mathbb{1}_{d_k}$, opposite to the blocks of S . Since T acts by one matrix B_k on the irreducible index of every row and g acts by one coefficient family $\gamma^{(k)}$ on the multiplicity index of every column, the two actions commute on every basis vector, so $Tg = gT$. By Lemma 7.11.31, $T \in S$.

Finally, if a matrix A satisfies $U^\dagger A U = \bigoplus_k \mathbb{1}_{m_k} \otimes B_k$, then $U^\dagger A U = U^\dagger T U$ for the member $T \in S$ built from the blocks $(B_k)_k$, so $A = T \in S$. $\square$

Theorem 7.11.33 (Fixed-point algebra is semisimple). [Lean](#) [✓ Stmt](#) *The adjoint-fixed-point *-subalgebra of a trace-preserving Kraus map on $M_D(\mathbb{C})$ is a semisimple ring.*

Proof. [✓ Proof](#) The subalgebra is finite-dimensional (as a subalgebra of $M_D(\mathbb{C})$), hence Artinian. Following [Wol12, Theorem 6.14], it suffices to show the Jacobson radical J vanishes. If $x \in J$, then $x^*x \in J$ (left-ideal closure). The radical of an Artinian ring is nilpotent, so x^*x is nilpotent. A self-adjoint nilpotent matrix satisfies $\|x^*x\|^{2^k} = \|(x^*x)^{2^k}\| = 0$, hence $x^*x = 0$, and the C^* -identity gives $x = 0$. $\square$

Theorem 7.11.34 (Abstract Wedderburn–Artin decomposition). [Lean](#) [✓ Stmt](#) *There exist $n \in \mathbb{N}$ and dimensions $d_1, \dots, d_n \geq 1$ such that the adjoint-fixed-point *-subalgebra is $\mathbb{C}$ -algebra isomorphic to*

$$\prod_{k=1}^n M_{d_k}(\mathbb{C}).$$

Proof. [✓ Proof](#) Combine semisimplicity (Theorem 7.11.33) with the Wedderburn–Artin structure theorem for finite-dimensional semisimple algebras over an algebraically closed field. $\square$

Theorem 7.11.35 (Wedderburn–Artin decomposition of the unital fixed-point algebra). [Lean](#)

[✓ Stmt](#) Let E be a unital Kraus map on $M_D(\mathbb{C})$ whose adjoint map E^* has a positive definite fixed point $\rho > 0$. There exist $n \in \mathbb{N}$ and dimensions $d_1, \dots, d_n \geq 1$ such that the fixed-point $*$ -subalgebra $\text{Fix}(E)$ is $\mathbb{C}$ -algebra isomorphic to

$$\text{Fix}(E) \cong \prod_{k=1}^n M_{d_k}(\mathbb{C}).$$

Proof. [✓ Proof](#) By Theorem 4.6.3 the fixed-point set $\text{Fix}(E)$ is a $*$ -subalgebra of $M_D(\mathbb{C})$. Applying the abstract Wedderburn–Artin decomposition for $*$ -subalgebras (Theorem 7.11.2) gives

$$\text{Fix}(E) \cong \prod_{k=1}^n M_{d_k}(\mathbb{C})$$

with each block size $d_k \geq 1$. □

Theorem 7.11.36 (Fixed-point algebra decomposition with multiplicities). [Lean](#) [✓ Stmt](#) There exist integers n , block sizes $d_1, \dots, d_n \geq 1$, multiplicities $m_1, \dots, m_n \geq 1$, and a $\mathbb{C}$ -algebra isomorphism from the adjoint-fixed-point $*$ -subalgebra to

$$\prod_{k=1}^n M_{d_k}(\mathbb{C})$$

such that

$$\sum_{k=1}^n d_k m_k \leq D.$$

Proof. [✓ Proof](#) Apply the abstract Wedderburn decomposition (Theorem 7.11.34). Since the adjoint-fixed-point algebra is realized as a subalgebra of the ambient matrix algebra $M_D(\mathbb{C})$, this realization also yields multiplicities m_k and the bound $\sum_k d_k m_k \leq D$. □

Remark 7.11.37 (Block form of the fixed-point algebra). Wolf’s block form identifies the adjoint-fixed-point algebra, up to a unitary change of basis, with

$$0 \oplus \bigoplus_k M_{d_k}(\mathbb{C}) \otimes \mathbb{1}_{m_k}$$

together with the density-block form $M_{d_k}(\mathbb{C}) \otimes \rho_k$. The theorem above gives the product decomposition, multiplicities, and dimension bound from this block form; the unitary conjugation identity itself, in the unital case, is Theorem 7.11.38 below.

Theorem 7.11.38 (Unitary block form of the Heisenberg-picture fixed points). [Lean](#) [✓ Stmt](#)

Let E be a trace-preserving Kraus map on $M_D(\mathbb{C})$ with a positive definite fixed point $\rho > 0$, $E(\rho) = \rho$. There are $n \in \mathbb{N}$, positive dimensions $d_0, \dots, d_{n-1}$ and multiplicities $m_0, \dots, m_{n-1}$ with $\sum_k d_k m_k = D$, realized by an explicit identification of the index sets, and a unitary $U \in M_D(\mathbb{C})$ such that a matrix $X \in M_D(\mathbb{C})$ satisfies $E^*(X) = X$ exactly when

$$U^\dagger X U = \bigoplus_{k=0}^{n-1} \mathbb{1}_{m_k} \otimes B_k$$

for some matrices $B_k \in M_{d_k}(\mathbb{C})$; that is, up to reordering the two tensor factors of each block,

$$\text{Fix}(E^*) = U \left(\bigoplus_{k=0}^{n-1} \mathbb{1}_{m_k} \otimes M_{d_k}(\mathbb{C}) \right) U^\dagger.$$

The positive definite fixed point removes the zero block: this is the unital case of the block representation in Equation (1.39) of [Wol12], invoked by [Wol12, Theorem 6.14]. The general form of [Wol12, Theorem 6.14], with a zero block and density weights ρ_k on the Schrödinger-picture fixed points, is not asserted here.

Proof. ✓ Proof The fixed points of E^* form a $*$ -subalgebra of $M_D(\mathbb{C})$ because E is trace-preserving with a positive definite fixed point (Theorem 4.6.11). The full block form of a $*$ -subalgebra (Theorem 7.11.32) supplies the unitary U , the dimensions and multiplicities, and the equivalence between membership and the block-diagonal conjugation identity. $\square$

Theorem 7.11.39 (Unitary block form of the fixed points of a unital Kraus map). Lean ✓ Stmt Let E be a unital Kraus map on $M_D(\mathbb{C})$ whose adjoint map E^* has a positive definite fixed point $\rho > 0$. There are $n \in \mathbb{N}$, positive dimensions d_k and multiplicities m_k with $\sum_k d_k m_k = D$, and a unitary $U \in M_D(\mathbb{C})$ such that a matrix $X \in M_D(\mathbb{C})$ satisfies $E(X) = X$ exactly when

$$U^\dagger X U = \bigoplus_{k=0}^{n-1} \mathbb{1}_{m_k} \otimes B_k$$

for some matrices $B_k \in M_{d_k}(\mathbb{C})$. This is Theorem 7.11.38 with the roles of the map and its adjoint exchanged.

Proof. ✓ Proof The fixed points of the unital map E form a $*$ -subalgebra of $M_D(\mathbb{C})$ (Theorem 4.6.3), and the full block form of a $*$ -subalgebra (Theorem 7.11.32) supplies the unitary and the equivalence. $\square$

Without a full-rank hypothesis on the fixed point, the corner-restricted fixed-point $*$ -algebra of Theorem 7.10.1.5 still carries the block structure through the compression onto the support sector; this is the $\sum_k d_k m_k \leq D$ support-sector form of the block representation in Equation (1.39) of [Wol12] invoked by [Wol12, Theorem 6.14]. A corner-restricted fixed point extended by zero on the complement of the support sector need not be a fixed point of the ambient map, so the corner block form does not by itself give the ambient fixed-point space a zero-block representation.

Theorem 7.11.40 (Block form of the corner-restricted fixed points). Lean ✓ Stmt Let $E(X) = \sum_i K_i X K_i^\dagger$ be a trace-preserving Kraus map on $M_D(\mathbb{C})$, with Heisenberg-picture adjoint $E^*(Y) = \sum_i K_i^\dagger Y K_i$, let $\rho \succeq 0$ satisfy $E(\rho) = \rho$, and let Q be the support projection of ρ . There are a sector dimension $r \leq D$, a number n of blocks, positive dimensions $d_0, \dots, d_{n-1}$ and multiplicities $m_0, \dots, m_{n-1}$ with $\sum_k d_k m_k = r$, realized by an explicit identification of the index sets, and an isometry $W : \mathbb{C}^r \rightarrow \mathbb{C}^D$ with $W^\dagger W = \mathbb{1}_r$ and $W W^\dagger = Q$, such that a matrix $Y \in M_D(\mathbb{C})$ satisfies $Q Y Q = Y$ and $Q E^*(Y) Q = Y$ exactly when

$$Y = W \left(\bigoplus_{k=0}^{n-1} \mathbb{1}_{m_k} \otimes B_k \right) W^\dagger$$

for some matrices $B_k \in M_{d_k}(\mathbb{C})$; that is, up to reordering the two tensor factors of each block,

$$\{Y \in Q M_D(\mathbb{C}) Q \mid Q E^*(Y) Q = Y\} = W \left(\bigoplus_{k=0}^{n-1} \mathbb{1}_{m_k} \otimes M_{d_k}(\mathbb{C}) \right) W^\dagger.$$

The right-hand side is the support-sector block representation in Equation (1.39) of [Wol12]; the equivalence characterizes the corner-restricted set, not the ambient fixed-point space.

Proof. ✓ Proof Since ρ is a fixed point of E , the support projection satisfies $(\mathbb{1} - Q)K_i Q = 0$ (Theorem 7.10.2), so the compressed family $QK_i Q$ is supported on the corner and trace-preserving there. Compress it to the support sector along the isometry $V : \mathbb{C}^r \rightarrow \mathbb{C}^D$ of Theorem 9.6.1.4, which satisfies $V^\dagger V = \mathbb{1}_r$ and $VV^\dagger = Q$: the compressed family C is trace-preserving on $\mathbb{C}^r$, and the compression $\sigma = V^\dagger \rho V$ of ρ is a positive definite fixed point of the compressed Schrödinger map (Theorem 7.10.1.4). The block form of the Heisenberg-picture fixed points of C (Theorem 7.11.38) yields a unitary $U \in M_r(\mathbb{C})$, dimensions d_k and multiplicities m_k with $\sum_k d_k m_k = r$, and the equivalence between the fixed-point equation on the sector and the block-diagonal conjugation identity. Writing $\Phi(X) := VXV^\dagger$ for the compression isomorphism onto the corner and C^* for the Heisenberg-picture adjoint of the compressed family, the corner-restricted fixed-point condition corresponds to the fixed-point equation of the compressed map,

$$QE^*(\Phi(X))Q = \Phi(X) \iff C^*(X) = X,$$

so, with $W := VU$, which satisfies $W^\dagger W = \mathbb{1}_r$ and $WW^\dagger = Q$, a corner element Y is a corner-restricted fixed point exactly when $Y = W(\bigoplus_k \mathbb{1}_{m_k} \otimes B_k)W^\dagger$ for some blocks B_k . Finally $r \leq D$ because the isometry V embeds $\mathbb{C}^r$ into $\mathbb{C}^D$. $\square$

Theorem 7.11.41 (Weighted fixed points form a $*$ -subalgebra). Lean ✓ Stmt Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving, and let $\rho > 0$ satisfy $T(\rho) = \rho$. If

$$F_T := \{X \in M_D(\mathbb{C}) \mid T(X) = X\},$$

then $\rho^{-1/2} F_T \rho^{-1/2}$ is a $*$ -subalgebra of $M_D(\mathbb{C})$.

Proof. ✓ Proof Set $B_i = \rho^{-1/2} K_i \rho^{1/2}$. The equation $T(\rho) = \rho$ implies that the family $\{B_i\}$ is unital, and trace preservation of T implies that ρ is a positive-definite fixed point of the adjoint map of the gauged family. Apply Theorem 4.6.3 to B . The identity

$$\sum_i B_i X B_i^\dagger = X \iff T(\rho^{1/2} X \rho^{1/2}) = \rho^{1/2} X \rho^{1/2}$$

identifies the fixed points of the gauged map with $\rho^{-1/2} F_T \rho^{-1/2}$. $\square$

For a singular fixed point the conjugation is taken with the inverse square root on the support of ρ , and the conjugated set lives in the corner algebra of the support projection. The corollary in [Wol12] takes a maximum-rank fixed point, for which every fixed point of T is supported on the support of ρ ; the following two statements take an arbitrary positive semidefinite fixed point and restrict to the fixed points supported on the support of ρ . Theorem 7.11.46 below supplies a fixed point whose support carries every fixed point, and at such a fixed point the restriction disappears (Theorem 7.11.47).

Theorem 7.11.42 (Weighted corner fixed points form a $*$ -subalgebra). Lean ✓ Stmt Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving, let $\rho \succeq 0$ satisfy $T(\rho) = \rho$, and let Q be the support projection of ρ . The corner elements $Y \in Q M_D(\mathbb{C}) Q$ such that $\sqrt{\rho} Y \sqrt{\rho}$ is a fixed point of T form a $*$ -subalgebra of the corner algebra $Q M_D(\mathbb{C}) Q$.

Proof. ✓ Proof Closure under addition, scalars, and conjugation is linearity together with the stability of fixed points under conjugate transposition; the corner unit Q belongs to the set because $\sqrt{\rho} Q \sqrt{\rho} = \rho$ is a fixed point, the support projection absorbing the square root: $Q \sqrt{\rho} = \sqrt{\rho} = \sqrt{\rho} Q$, since $(\mathbb{1} - Q) \sqrt{\rho} ((\mathbb{1} - Q) \sqrt{\rho})^\dagger = (\mathbb{1} - Q) \rho (\mathbb{1} - Q) = 0$.

For closure under multiplication, compress to the support sector: since ρ is a fixed point of T , the support projection satisfies $(\mathbb{1} - Q)K_i Q = 0$ (Theorem 7.10.2), the compressed family $QK_i Q$ is trace-preserving on the corner, and the compression along the isometry $V : \mathbb{C}^r \rightarrow \mathbb{C}^D$ of Theorem 9.6.1.4, with $V^\dagger V = \mathbb{1}_r$ and $VV^\dagger = Q$, produces a trace-preserving family C on $\mathbb{C}^r$ whose Schrödinger map has the positive definite fixed point $\sigma = V^\dagger \rho V$ (Theorem 7.10.1.4). The square root compresses along the isometry,

$$\sqrt{\sigma} = V^\dagger \sqrt{\rho} V,$$

by uniqueness of the positive square root, since $(V^\dagger \sqrt{\rho} V)(V^\dagger \sqrt{\rho} V) = V^\dagger \sqrt{\rho} Q \sqrt{\rho} V = \sigma$. Hence for corner-supported Y ,

$$\sqrt{\sigma} (V^\dagger Y V) \sqrt{\sigma} = V^\dagger (\sqrt{\rho} Y \sqrt{\rho}) V,$$

and the fixed-point equations correspond: for corner-supported W the compressed map evaluates to

$$C(V^\dagger W V) = V^\dagger Q T(W) Q V = V^\dagger T(W) V,$$

since $T(W)$ is again corner-supported, so $T(W) = W$ exactly when the compressed map fixes $V^\dagger W V$. The set therefore corresponds, element by element, to the weighted fixed points of the compressed family with respect to the positive definite fixed point σ , which form a $*$ -subalgebra by Theorem 7.11.41. Closure under multiplication transports back through the correspondence: a corner-supported Y_j satisfies $Y_j Q = Y_j$ (right-multiply $Q Y_j Q = Y_j$ by Q and use $Q^2 = Q$), so

$$(V^\dagger Y_1 V)(V^\dagger Y_2 V) = V^\dagger Y_1 (V V^\dagger) Y_2 V = V^\dagger (Y_1 Q) Y_2 V = V^\dagger (Y_1 Y_2) V,$$

and the product of two members corresponds to the product of their compressed images, which lies in the compressed $*$ -subalgebra. $\square$

Combined with the conjugation correspondence of Theorem 7.11.43, the carrier of this $*$ -subalgebra realizes

$$\rho^{-1/2} \{X \in M_D(\mathbb{C}) \mid T(X) = X, QXQ = X\} \rho^{-1/2},$$

with the inverse square root taken on the support of ρ : the singular case of Corollary 6.7 of [Wol12], restricted to the fixed points supported on the support of ρ .

Theorem 7.11.43 (Conjugation by the square root is onto the corner-supported fixed points).

Lean **Stmt** In the setting of Theorem 7.11.42, every fixed point X of T with $QXQ = X$ arises as $X = \sqrt{\rho} Y \sqrt{\rho}$ for a corner-supported Y with $\sqrt{\rho} Y \sqrt{\rho}$ fixed by T .

Proof. **Proof** Compress to the support sector as in the proof of Theorem 7.11.42 and set

$$Y = V (\sqrt{\sigma}^{-1} (V^\dagger X V) \sqrt{\sigma}^{-1}) V^\dagger,$$

where $\sigma = V^\dagger \rho V$ is positive definite, so $\sqrt{\sigma}$ is invertible. Then Y is corner-supported, and, writing $Z = \sqrt{\sigma}^{-1} (V^\dagger X V) \sqrt{\sigma}^{-1}$ for the compressed middle factor, the transport identity displayed above gives

$$V^\dagger (\sqrt{\rho} Y \sqrt{\rho}) V = \sqrt{\sigma} (V^\dagger Y V) \sqrt{\sigma} = \sqrt{\sigma} Z \sqrt{\sigma} = V^\dagger X V,$$

hence $\sqrt{\rho} Y \sqrt{\rho} = QXQ = X$, both sides being corner-supported. In particular $\sqrt{\rho} Y \sqrt{\rho}$ is fixed by T . $\square$

The support hypothesis on the fixed points is discharged by exhibiting a fixed point whose support carries every fixed point. The first ingredient is that Loewner domination transfers the support.

Lemma 7.11.44 (The transfer map of a trace-preserving family is a channel). [Lean](#) [✓ Stmt](#) Let $K_0, \dots, K_{d-1}$ be a trace-preserving Kraus family. The map $E(X) = \sum_i K_i X K_i^\dagger$ is a quantum channel.

Proof. [✓ Proof](#) The trace-preservation hypothesis is the normalization $\sum_i K_i^\dagger K_i = \mathbf{1}$ of the channel form of the transfer map (Theorem 4.2.7). $\square$

Lemma 7.11.45 (Loewner domination transfers the support). [Lean](#) [✓ Stmt](#) Let $\rho, P \succeq 0$ with $P \preceq \rho$, and let Q be the support projection of ρ . Then

$$QP = P, \quad PQ = P.$$

Proof. [✓ Proof](#) The support projection absorbs ρ , so $(\mathbf{1} - Q)\rho = 0$ and

$$0 \preceq (\mathbf{1} - Q)P(\mathbf{1} - Q) \preceq (\mathbf{1} - Q)\rho(\mathbf{1} - Q) = 0,$$

the middle inequality because conjugation by $\mathbf{1} - Q$ preserves the Loewner order. Hence

$$((\mathbf{1} - Q)\sqrt{P})(\mathbf{1} - Q)\sqrt{P}^\dagger = (\mathbf{1} - Q)P(\mathbf{1} - Q) = 0,$$

so $(\mathbf{1} - Q)\sqrt{P} = 0$, and

$$(\mathbf{1} - Q)P = ((\mathbf{1} - Q)\sqrt{P})\sqrt{P} = 0,$$

that is $QP = P$; taking conjugate transposes gives $PQ = P$ since P and Q are Hermitian. $\square$

Theorem 7.11.46 (A fixed point of maximal support). [Lean](#) [✓ Stmt](#) Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving. There is a positive semidefinite fixed point ρ_0 of T such that, with Q_0 the support projection of ρ_0 , every fixed point X of T satisfies

$$Q_0 X Q_0 = X.$$

This is the maximal-support property of fixed points in Section 6.4 of [Wol12]; there the witness is the image of the identity under the projection onto the fixed-point space, while here it is a sum of positive fixed points.

Proof. [✓ Proof](#) Let $X_0, \dots, X_{m-1}$ be a basis of the fixed-point space of T . Each X_j splits into fixed Hermitian components: with

$$H_{1,j} = X_j + X_j^\dagger, \quad H_{2,j} = i(X_j - X_j^\dagger),$$

both are Hermitian fixed points (the conjugate transpose of a fixed point is fixed), and

$$X_j = \frac{1}{2} H_{1,j} - \frac{i}{2} H_{2,j}.$$

By the decomposition of Hermitian fixed points into positive parts (Theorem 4.3.3) there are positive semidefinite fixed points with

$$H_{1,j} = P_{1,j}^+ - P_{1,j}^-, \quad H_{2,j} = P_{2,j}^+ - P_{2,j}^-.$$

Set

$$\rho_0 = \sum_{j=0}^{m-1} (P_{1,j}^+ + P_{1,j}^- + P_{2,j}^+ + P_{2,j}^-),$$

a positive semidefinite fixed point. Every summand is dominated by ρ_0 in the Loewner order, because the remaining summands are positive semidefinite, so the support projection Q_0 of ρ_0 absorbs each of them (Lemma 7.11.45):

$$Q_0 P_{k,j}^\pm Q_0 = P_{k,j}^\pm.$$

Substituting into the two displayed decompositions gives

$$Q_0 H_{k,j} Q_0 = Q_0 (P_{k,j}^+ - P_{k,j}^-) Q_0 = H_{k,j}, \quad Q_0 X_j Q_0 = \frac{1}{2} Q_0 H_{1,j} Q_0 - \frac{i}{2} Q_0 H_{2,j} Q_0 = X_j.$$

A fixed point X of T is a linear combination of the basis $X_0, \dots, X_{m-1}$, so $Q_0 X Q_0 = X$. $\square$

Theorem 7.11.47 (Conjugation by the square root at a fixed point of maximal support). [Lean](#)

[✓ Stmt](#) Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving. There is a positive semidefinite fixed point ρ_0 of T such that every fixed point X of T arises as

$$X = \sqrt{\rho_0} Y \sqrt{\rho_0}$$

for a corner-supported Y with $\sqrt{\rho_0} Y \sqrt{\rho_0}$ fixed by T .

Proof. [✓ Proof](#) Take the fixed point ρ_0 of Theorem 7.11.46. Every fixed point X of T satisfies $Q_0 X Q_0 = X$ for the support projection Q_0 of ρ_0 , so Theorem 7.11.43 applies to X and produces the corner-supported Y . $\square$

At the fixed point ρ_0 , with the inverse square root taken on the support of ρ_0 , the carrier of the $*$ -subalgebra of Theorem 7.11.42 therefore realizes

$$\rho_0^{-1/2} \{X \in M_D(\mathbb{C}) \mid T(X) = X\} \rho_0^{-1/2} :$$

the conjugated fixed-point set of Corollary 6.7 of [Wol12] at a fixed point of maximal support, without a support restriction on the fixed points.

The corollary in [Wol12] quantifies over an arbitrary maximum-rank fixed-point density matrix. The transfer from the constructed witness to every fixed point of maximum rank goes through the rank of the support projection.

Lemma 7.11.48 (The support projection has the rank of the matrix). [Lean](#) [✓ Stmt](#) For $\rho \succeq 0$ on $\mathbb{C}^D$, the support projection of ρ has the rank of ρ .

Proof. [✓ Proof](#) In the spectral decomposition $\rho = U \text{diag}(\lambda) U^\dagger$ the support projection is $U \text{diag}(\mathbf{1}_{\lambda>0}) U^\dagger$. Multiplication by the invertible U and $U^\dagger$ preserves the rank, and

$$\text{rank } \text{diag}(\mathbf{1}_{\lambda>0}) = \#\{i \mid \lambda_i > 0\} = \#\{i \mid \lambda_i \neq 0\} = \text{rank } \text{diag}(\lambda),$$

the middle equality because the eigenvalues of $\rho \succeq 0$ are nonnegative. $\square$

Lemma 7.11.49 (The trace of the support projection is the rank). [Lean](#) [✓ Stmt](#) For $\rho \succeq 0$ on $\mathbb{C}^D$, the trace of the support projection of ρ is the rank of ρ .

Proof. [✓ Proof](#) With $\rho = U \text{diag}(\lambda) U^\dagger$ as above,

$$\text{tr}(U \text{diag}(\mathbf{1}_{\lambda>0}) U^\dagger) = \text{tr } \text{diag}(\mathbf{1}_{\lambda>0}) = \#\{i \mid \lambda_i > 0\} = \text{rank } \rho,$$

the last equality because the rank of a positive semidefinite matrix is the number of its positive eigenvalues. $\square$

Theorem 7.11.50 (The support of a fixed point of maximum rank carries every fixed point).

Lean **✓ Stmt** Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving and let $\rho \succeq 0$ be a fixed point of T whose rank bounds the rank of every fixed-point density matrix: $\text{rank } \sigma \leq \text{rank } \rho$ for every $\sigma \succeq 0$ with $\text{tr } \sigma = 1$ and $T(\sigma) = \sigma$. Then the support projection Q of ρ satisfies

$$Q X Q = X$$

for every fixed point X of T .

Proof. **✓ Proof** Let ρ_0 be the fixed point of maximal support of Theorem 7.11.46, with support projection Q_0 , and let P be the support projection of ρ . If $\rho_0 = 0$, then every fixed point vanishes, because $Q_0 = 0$ annihilates it, and the claim is trivial; so let $\rho_0 \neq 0$. Its trace is then a nonzero nonnegative real number, and $\sigma = (\text{tr } \rho_0)^{-1} \rho_0$ is a fixed-point density matrix with $\text{rank } \sigma = \text{rank } \rho_0$, so the rank hypothesis gives $\text{rank } \rho_0 \leq \text{rank } \rho$. Conversely $Q_0 \rho Q_0 = \rho$, so

$$\text{rank } \rho = \text{rank}(Q_0 \rho Q_0) \leq \text{rank } Q_0 = \text{rank } \rho_0,$$

the last equality by Lemma 7.11.48; hence the ranks agree. From $\rho Q_0 = \rho$ one has $\rho(1 - Q_0) = 0$, so the range of $1 - Q_0$ lies in the kernel of ρ ; the support projection P of ρ vanishes on that kernel, so

$$P(1 - Q_0) = 0, \quad P Q_0 = P,$$

and taking conjugate transposes also $Q_0 P = P$. The difference $R = Q_0 - P$ is then Hermitian and idempotent,

$$R^2 = Q_0 - P - P + P = R,$$

and its trace vanishes by Lemma 7.11.49 and the equality of the ranks, so

$$\text{tr}(R^\dagger R) = \text{tr } R = 0,$$

hence $R = 0$ and $P = Q_0$. The claim now follows from the maximal-support property of ρ_0 . $\square$

Theorem 7.11.51 (Conjugation by the square root at every fixed point of maximum rank).

Lean **✓ Stmt** Let $T(X) = \sum_i K_i X K_i^\dagger$ be trace-preserving and let $\rho \succeq 0$ be a fixed point of T whose rank bounds the rank of every fixed-point density matrix. Then every fixed point X of T arises as

$$X = \sqrt{\rho} Y \sqrt{\rho}$$

for a corner-supported Y with $\sqrt{\rho} Y \sqrt{\rho}$ fixed by T .

Proof. **✓ Proof** By Theorem 7.11.50 every fixed point X of T satisfies $Q X Q = X$ for the support projection Q of ρ , so Theorem 7.11.43 applies to X and produces the corner-supported Y . $\square$

At every fixed point ρ of maximum rank, with the inverse square root taken on the support of ρ , the carrier of the $*$ -subalgebra of Theorem 7.11.42 therefore realizes the conjugated fixed-point set $\rho^{-1/2} \{X \in M_D(\mathbb{C}) \mid T(X) = X\} \rho^{-1/2}$ of Corollary 6.7 of [Wol12].

7.12 Further irreducibility and primitivity equivalences

This section collects several criteria and equivalences from [Wol12, Theorems 6.2, 6.7, 6.8] that are used later.

Theorem 7.12.1 (Exponential positivity condition). [Lean](#) [Lean](#) [✓ Stmt](#) Let E be an irreducible completely positive map, $A \geq 0$ with $A \neq 0$, and $t > 0$. Then $\exp(tE)(A)$ is positive definite. Conversely, if $\exp(tE)(A)$ is positive definite for every such t, A , then E is irreducible (full equivalence [Lean](#)).

Theorem 7.12.2 (Kraus-span equivalence). [Lean](#) [✓ Stmt](#) Under normalization, primitivity in Wolf’s sense is equivalent to eventual full Kraus rank.

Theorem 7.12.3 (Combined primitivity equivalence). [Lean](#) [✓ Stmt](#) Under normalization, primitivity in Wolf’s sense is equivalent to the conjunction of eventual full Kraus rank, normality, and strong irreducibility.

Remark 7.12.4 (Pairwise primitivity equivalences). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) These are the pairwise equivalences in [Wol12, Theorem 6.8].

Theorem 7.12.5 (Scalar fixed-point algebra case). [Lean](#) [✓ Stmt](#) In the scalar fixed-point algebra setting, the weighted map $X \mapsto \frac{\text{tr}(\rho X)}{\text{tr}(\rho)} \mathbb{1}$ is a conditional expectation onto the adjoint fixed-point $*$ -subalgebra.

7.13 Multi-cycle block-permutation structure

This section develops the block-permutation structure underlying the cycle decomposition of [Wol12, Theorem 6.16]. The multi-cycle decomposition refines the single-cycle case to handle arbitrary peripheral periods.

Definition 7.13.1 (Block-permutation structure). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$. A *block-permutation structure* for T consists of a finite index set ι , a family of orthogonal projections $P_k \in M_D(\mathbb{C})$ for $k \in \iota$, and a permutation $\sigma \in \text{Sym}(\iota)$ (not necessarily a single cycle—in general σ may have multiple disjoint cycles), such that

$$T(P_{\sigma(k)}) = P_k, \quad T(P_k X) = T(P_k) T(X), \quad T(X P_k) = T(X) T(P_k)$$

for every $k \in \iota$ and $X \in M_D(\mathbb{C})$.

Definition 7.13.2 (Multi-cycle block-permutation structure). [Lean](#) [✓ Stmt](#) A *multi-cycle decomposition* of $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ consists of a finite cycle index ι , a per-cycle period $m : \iota \rightarrow \mathbb{N}_{>0}$, a projection family $P_{c,k} \in M_D(\mathbb{C})$ for $c \in \iota$ and $k \in \{0, \dots, m(c)-1\}$ consisting of orthogonal projections. For every $c \in \iota$ and $k \in \{0, \dots, m(c)-1\}$, the per-cycle cyclic action is

$$T(P_{c,k+1}) = P_{c,k}.$$

The multiplicative-domain factorisations are $T(P_{c,k} X) = T(P_{c,k}) T(X)$ and $T(X P_{c,k}) = T(X) T(P_{c,k})$.

Lemma 7.13.3 (Per-cycle corner preservation). [Lean](#) [✓ Stmt](#) For every cycle $c \in \iota$ and index $k \in \{0, \dots, m(c)-1\}$, the iterate $T^{m(c)}$ preserves the corner $P_{c,k} M_D(\mathbb{C}) P_{c,k}$.

Proof. [✓ Proof](#) By induction on n , the n -th iterate T^n sends $P_{c,k+n} X P_{c,k+n}$ to $P_{c,k} T^n(X) P_{c,k}$, using the multiplicative-domain factorisations and the cyclic action $T(P_{c,j+1}) = P_{c,j}$ at each induction step. Specialising to $n = m(c)$ and using that the indices $k+m(c) = k$ in $\{0, \dots, m(c)-1\}$ yields corner preservation of $T^{m(c)}$. $\square$

Lemma 7.13.4 (Common-period corner preservation). [Lean](#) [✓ Stmt](#) If $N \in \mathbb{N}$ is divisible by every per-cycle period $m(c)$, then T^N preserves every corner $P_{c,k} M_D(\mathbb{C}) P_{c,k}$ of the decomposition.

Proof. ✓ Proof Write $N = m(c)q$. Corner preservation is stable under taking powers, so $T^N = (T^{m(c)})^q$ preserves each corner by Lemma 7.13.3. $\square$

Definition 7.13.5 (Flattening to a single-index block-permutation structure). Lean ✓ Stmt A multi-cycle decomposition gives a single-index block-permutation structure on the sigma index $\Sigma_{c \in \iota} \{0, \dots, m(c)-1\}$: the permutation is the sigma-product of per-cycle cyclic shifts $k \mapsto k+1$ (whose cycle decomposition has one cycle per $c \in \iota$), the projections are $(c, k) \mapsto P_{c,k}$, and the multiplicative-domain factorisations descend componentwise. The resulting map forgets the explicit cycle-indexing.

7.14 Peripheral eigenvalue group structure

This section proves the peripheral spectrum structure of [Wol12, Theorem 6.6] for irreducible Schwarz maps: the peripheral eigenvalues form a finite cyclic group. The trace-preserving refinement that the order divides the bond dimension is proved in Theorem 7.14.4.

Lemma 7.14.1 (Peripheral eigenvectors are invertible). Lean ✓ Stmt *Let E be an irreducible unital Kraus map on $M_D(\mathbb{C})$ with a positive-definite adjoint fixed point. If $X \neq 0$ satisfies $E(X) = \mu X$ with $|\mu| = 1$, then X is invertible.*

Proof. ✓ Proof The KS equality gives $E(X^\dagger X) = X^\dagger X$, so $X^\dagger X$ is a nonzero PSD fixed point. Irreducibility upgrades it to positive definite, hence invertible, so $\det(X) \neq 0$. $\square$

Lemma 7.14.2 (Peripheral eigenvalues: product closure). Lean ✓ Stmt *Let E be an irreducible unital Kraus map on $M_D(\mathbb{C})$ with a positive-definite adjoint fixed point. If μ, ν are peripheral eigenvalues of E , then so is $\mu\nu$.*

Proof. ✓ Proof Take eigenvectors X, Y for μ, ν . The KS equality at X puts X in the multiplicative domain, so $E(YX) = E(Y)E(X) = (\nu\mu)(YX)$. Since X, Y are units (Lemma 7.14.1), $YX \neq 0$ and $|\mu\nu| = 1$. $\square$

Theorem 7.14.3 (Peripheral eigenvalues: cyclic structure). Lean ✓ Stmt *Let E be an irreducible unital Schwarz map on $M_D(\mathbb{C})$ with a positive-definite adjoint fixed point (trace-preservation is not required). Then there exist $m \geq 1$ and a primitive m -th root of unity γ such that the peripheral eigenvalues of E are exactly $\{1, \gamma, \gamma^2, \dots, \gamma^{m-1}\}$.*

Proof. ✓ Proof The peripheral eigenvalues form a finite subset of $\mathbb{C}$, contain 1, and are closed under multiplication and inversion, so they form a finite subgroup of $\mathbb{C}^\times$. Any finite subgroup of $\mathbb{C}^\times$ is cyclic. The generator γ has order $m = |\text{peripheral spectrum}|$. $\square$

Theorem 7.14.4 (Peripheral eigenvalues form a cyclic group). Lean ✓ Stmt *Let E be an irreducible unital trace-preserving Schwarz map on $M_D(\mathbb{C})$ with a positive-definite adjoint fixed point. Then there exist $m \geq 1$ and a primitive m -th root of unity γ such that $m \mid D$ and the peripheral eigenvalues of E are exactly $\{1, \gamma, \gamma^2, \dots, \gamma^{m-1}\}$.*

Proof. ✓ Proof Apply Theorem 7.14.3 for the cyclic structure. Divisibility $m \mid D$ follows from the cyclic decomposition: m mutually orthogonal projections summing to $\mathbb{1}$ each have equal trace (by trace preservation), so $m \cdot \text{tr}(P_0) = D$ with $\text{tr}(P_0) \in \mathbb{N}$. $\square$

Theorem 7.14.5 (Period divides bond dimension). Lean ✓ Stmt *Let E be an irreducible unital trace-preserving Schwarz map on $M_D(\mathbb{C})$ with a positive-definite adjoint fixed point. Let γ be a primitive m -th root of unity such that the peripheral eigenvalues are exactly $\{\gamma^k : k \in \{0, \dots, m-1\}\}$. Then $m \mid D$.*

Proof. ✓ Proof The cyclic decomposition gives orthogonal projections $P_0, \dots, P_{m-1}$ with $\sum_k P_k = \mathbb{1}$ and $E(P_{k+1}) = P_k$. Trace preservation gives $\text{tr}(P_{k+1}) = \text{tr}(P_k)$ for every k , hence

$$D = \text{tr}(\mathbb{1}) = \sum_{k=0}^{m-1} \text{tr}(P_k) = m \text{tr}(P_0).$$

Since the trace of an orthogonal projection is an integer, $m \mid D$. □

7.14.1 Cyclic decomposition of irreducible Schwarz maps

Definition 7.14.1.1 (Corner preservation). Lean ✓ Stmt For an orthogonal projection P and linear map E , the corner $PM_D(\mathbb{C})P$ is preserved if $E(PXP) = PE(PXP)P$ for all X .

Definition 7.14.1.2 (Corner subspace). Lean ✓ Stmt The corner subspace associated with P is $\{PXP : X \in M_D(\mathbb{C})\}$.

Definition 7.14.1.3 (Restricted corner map). Lean ✓ Stmt The restriction of E to an invariant corner $PM_D(\mathbb{C})P$.

Definition 7.14.1.4 (Irreducible restriction on a corner). Lean ✓ Stmt Irreducibility of the corner-restricted map.

Definition 7.14.1.5 (Rank of an orthogonal projection). Lean ✓ Stmt The rank of an orthogonal projection $P \in M_D(\mathbb{C})$, i.e. the dimension of the corner algebra $PM_D(\mathbb{C})P$.

Lemma 7.14.1.6 (Compression isometry for an orthogonal projection). Lean ✓ Stmt Let $P \in M_D(\mathbb{C})$ be an orthogonal projection of rank n . The corner algebra $PM_D(\mathbb{C})P$ is linearly isomorphic to the full matrix algebra $M_n(\mathbb{C})$. The isomorphism is built from the spectral diagonalisation of P by conjugating the top-left $n \times n$ block by the unitary diagonalising P .

Lemma 7.14.1.7 (Rank equals trace for an orthogonal projection). Lean ✓ Stmt The rank of an orthogonal projection P equals its trace: $n = \text{tr } P$.

Lemma 7.14.1.8 (Scalar fixed points for irreducible unital maps). Lean ✓ Stmt If E is irreducible and unital, then every fixed point of E is a scalar multiple of $\mathbb{1}$.

Proof. ✓ Proof If $E(X) = X$, then the Hermitian matrices

$$X + X^\dagger, \quad i(X - X^\dagger)$$

are also fixed by E . The positive-semidefinite fixed-point uniqueness theorem gives $X + X^\dagger = a\mathbb{1}$ and $i(X - X^\dagger) = b\mathbb{1}$, hence

$$X = \frac{1}{2}(a - ib)\mathbb{1}.$$

□

Lemma 7.14.1.9 (Peripheral unitary eigenvector). Lean ✓ Stmt For an irreducible unital Schwarz map with faithful adjoint fixed point, each peripheral eigenvalue admits a unitary eigenvector.

Lemma 7.14.1.10 (Powers on a peripheral-unitary orbit). Lean ✓ Stmt If $E(U) = \mu U$ with $|\mu| = 1$ and U unitary, then $E(U^n) = \mu^n U^n$ for all n .

Lemma 7.14.1.11 (Normalized peripheral unitary). [Lean](#) [✓ Stmt](#) *Peripheral unitary eigenvectors can be normalized compatibly with the faithful fixed-point state.*

Lemma 7.14.1.12 (Cyclic projections from a peripheral unitary). [Lean](#) [✓ Stmt](#) *A primitive m -th peripheral root yields m orthogonal projections summing to $\mathbb{1}$ and cyclically permuted by the channel.*

Theorem 7.14.1.13 (Cyclic decomposition for irreducible Schwarz maps). [Lean](#) [✓ Stmt](#) *Under irreducibility and faithful fixed point hypotheses, there is a full cyclic projection decomposition realizing the peripheral period.*

Lemma 7.14.1.14 (Cyclic shift of the Kraus operators). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Let $P_0, \dots, P_{m-1}$ be orthogonal projections summing to the identity and cyclically permuted by a Kraus map, $\sum_v K_v P_{k+1} K_v^\dagger = P_k$. Then the projections are mutually orthogonal, none of them vanishes, and each Kraus operator intertwines consecutive projections,*

$$K_v P_{k+1} = P_k K_v.$$

Consequently a product of ℓ Kraus operators moves each projection back by ℓ steps; in particular a product of m Kraus operators commutes with every projection.

Proof. [✓ Proof](#) Compressing $\sum_v K_v P_{k+1} K_v^\dagger = P_k$ by $\mathbb{1} - P_k$ exhibits a vanishing sum of positive semidefinite matrices, so each summand vanishes: $(\mathbb{1} - P_k) K_v P_{k+1} = 0$. Mutual orthogonality follows by compressing the resolution of the identity by a fixed P_k , another vanishing sum of positive semidefinite matrices. Expanding $K_v P_{k+1}$ and $P_k K_v$ through the resolution of the identity, the off-diagonal compressions vanish by orthogonality, leaving $K_v P_{k+1} = P_k K_v P_{k+1} = P_k K_v$. If some projection vanished, the cyclic action would propagate the vanishing around the cycle, contradicting the resolution of the identity. The word statement is induction on the length. $\square$

Lemma 7.14.1.15 (Power maps preserve each cyclic sector). [Lean](#) [✓ Stmt](#) *Appropriate powers of the channel preserve each sector corner.*

Lemma 7.14.1.16 (Primitivity passes to a fixed corner). [Lean](#) [✓ Stmt](#) *Let $P \in M_D(\mathbb{C})$ be a nonzero idempotent and let E be a linear map on $M_D(\mathbb{C})$ that preserves the corner $PM_D(\mathbb{C})P$, is primitive, and fixes the corner projection, $E(P) = P$. Then the restriction of E to the corner $PM_D(\mathbb{C})P$ is again primitive.*

Proof. [✓ Proof](#) The corner projection P lies in its own corner and is fixed by the restriction, so 1 is a peripheral eigenvalue of the restriction. Conversely, every eigenvalue of the restriction of unit modulus arises from a nonzero corner element X with $E(X) = \mu X$; viewing X as an element of $M_D(\mathbb{C})$ exhibits μ as a peripheral eigenvalue of E . Primitivity of E forces $\mu = 1$, so the peripheral spectrum of the restriction is exactly $\{1\}$ and the restriction is primitive. $\square$

Lemma 7.14.1.17 (Irreducibility of restricted sector dynamics). [Lean](#) [✓ Stmt](#) *Each sector restriction is irreducible.*

Lemma 7.14.1.18 (Primitivity of restricted sector dynamics). [Lean](#) [✓ Stmt](#) *Each sector restriction is primitive.*

Lemma 7.14.1.19 (Corner invariance at the period). [Lean](#) [✓ Stmt](#) *The channel raised to the period preserves every cyclic corner.*

7.14.2 Group structure of the peripheral spectrum

Lemma 7.14.2.1 (Peripheral product closure). [Lean](#) [✓ Stmt](#) *The product of two peripheral eigenvalues is peripheral.*

Lemma 7.14.2.2 (Peripheral inverse closure). [Lean](#) [✓ Stmt](#) *The inverse of a peripheral eigenvalue is peripheral.*

Theorem 7.14.2.3 (Peripheral eigenvalue multiplicity one). [Lean](#) [✓ Stmt](#) *Each peripheral eigenvalue has algebraic multiplicity one.*

Proof. [✓ Proof](#) Given a peripheral unitary eigenvector U for γ , the multiplicative-domain identity gives $E(XU^\dagger) = XU^\dagger$ for any γ -eigenvector X . By the scalar fixed-point lemma for irreducible unital maps, $XU^\dagger = c \cdot \mathbb{1}$, hence $X = c \cdot U$ and the eigenspace is one-dimensional. $\square$

7.15 Primitive overlap convergence (complementary transfer-map gap form)

The theorems in this section are stated directly in terms of the complementary map $E - P$. This is the analytic form needed for the later overlap argument, corresponding to the primitive-case specialization of [Wol12, Theorem 6.7] where T_ϕ is a rank-one projection. For primitive normalized transfer maps, Chapter 4 supplies this complementary transfer-map input under explicit hypotheses.

Theorem 7.15.1 (Complementary transfer-map gap implies trace convergence to 1). [Lean](#) [✓ Stmt](#) *Let E be trace-preserving with fixed point ρ ($\text{tr}(\rho) \neq 0$) and let P be the fixed-point projection (Definition 4.4.1). If $\rho_{\text{spec}}(E - P) < 1$, then $\text{Tr}(E^n) \rightarrow 1$ as $n \rightarrow \infty$.*

Proof. [✓ Proof](#) By the power decomposition (Theorem 4.4.2), $E^n = P + (E - P)^n$ for $n \geq 1$. Since $\rho_{\text{spec}}(E - P) < 1$, the Gelfand formula gives $(E - P)^n \rightarrow 0$, so $\text{Tr}(E^n) \rightarrow \text{Tr}(P)$. The operator trace of P equals 1: since $P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)}\rho$, we have $\text{Tr}(P) = \frac{\text{tr}(\rho)}{\text{tr}(\rho)} = 1$. $\square$

Theorem 7.15.2 (Self-overlap convergence from a complementary transfer-map gap). [Lean](#) [✓ Stmt](#) *Let A be a normalized MPS tensor with a nonzero PSD fixed point ρ of the transfer map. If $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$ (where P is the fixed-point projection), then $O_{AA}(N) \rightarrow 1$ as $N \rightarrow \infty$.*

Proof. [✓ Proof](#) By Lemma 7.2.2 (with $A = B$), $O_{AA}(N) = \text{Tr}(\mathcal{E}_A^N)$. By Theorem 7.15.1, $\text{Tr}(\mathcal{E}_A^N) \rightarrow 1$. $\square$

Chapter 8

Wielandt Bound

This chapter proves a quantum analog of Wielandt's inequality: for a normal MPS tensor with bond dimension D , the cumulative span of matrix products stabilises after at most D^2 steps. The results follow [SPGWC10].

8.1 Cumulative span

Definition 8.1.1 (Word span). [Lean](#) [✓ Stmt](#) The *word span* at length n is the subspace

$$S_n(A) = \text{span}_{\mathbb{C}}\{A^w : |w| = n\} \subseteq M_D(\mathbb{C}).$$

This is [SPGWC10, Section II].

Definition 8.1.2 (Cumulative span). [Lean](#) [✓ Stmt](#) The *cumulative span* at level n is

$$T_n(A) = S_0(A) + S_1(A) + \dots + S_n(A).$$

Lemma 8.1.3 (Monotonicity of cumulative span). [Lean](#) [✓ Stmt](#) $T_n(A) \leq T_{n+1}(A)$ for all n .

Proof. [✓ Proof](#) T_{n+1} includes all words of length $\leq n+1$, which contains all words of length $\leq n$. $\square$

Lemma 8.1.4 (Dimension bound). [Lean](#) [✓ Stmt](#) $\dim T_n(A) \leq D^2$ for all n .

Proof. [✓ Proof](#) $T_n(A) \subseteq M_D(\mathbb{C})$ and $\dim M_D(\mathbb{C}) = D^2$. $\square$

Theorem 8.1.5 (Cumulative span stabilisation). [Lean](#) [✓ Stmt](#) If $T_n(A) = T_{n+1}(A)$, then $T_m(A) = T_n(A)$ for all $m \geq n$.

Proof. [✓ Proof](#) If $T_n = T_{n+1}$, then $S_{n+1} \subseteq T_n$. Left-multiplying any element of S_{n+1} by A^i gives an element of S_{n+2} , so $S_{n+2} \subseteq T_{n+1} = T_n$. By induction, $S_m \subseteq T_n$ for all $m > n$, hence $T_m = T_n$. $\square$

Theorem 8.1.6 (Dimension growth). [Lean](#) [✓ Stmt](#) If $T_n(A) \neq T_{n+1}(A)$, then $\dim T_{n+1}(A) > \dim T_n(A)$.

Proof. [✓ Proof](#) $T_n \leq T_{n+1}$ (Lemma 8.1.3) and $T_n \neq T_{n+1}$ imply strict submodule inclusion, hence strict rank increase. $\square$

Lemma 8.1.7 (Word span characterises block injectivity). [Lean](#) [✓ Stmt](#) $S_L(A) = M_D(\mathbb{C})$ if and only if A is L -block injective.

Proof. [✓ Proof](#) Both conditions assert that the span of all products of length L equals $M_D(\mathbb{C})$. $\square$

Theorem 8.1.8 (Positive-length propagation of block injectivity). [Lean](#) [Lean](#) [✓ Stmt](#) For every n ,

$$S_{n+1}(A) = \text{span}_{\mathbb{C}}\{A_i\}_i S_n(A).$$

Consequently, if $L > 0$ and A is L -block injective, then A is $(L + 1)$ -block injective. This is an auxiliary length-shift statement used to combine the block-injectivity discussion in [CPGSV16, lines 318–345] with Appendix C.3, Lemma L, lines 1835–1858; it is not stated there as a separate lemma.

Proof. [✓ Proof](#) Every word of length $n + 1$ factors into its first letter and a word of length n , which proves the displayed equality by bilinearity. Write $L = n + 1$. Since $S_L(A) = M_D(\mathbb{C})$, one has $S_n(A) \leq S_L(A)$. Hence

$$M_D(\mathbb{C}) = S_L(A) = \text{span}_{\mathbb{C}}\{A_i\}_i S_n(A) \leq \text{span}_{\mathbb{C}}\{A_i\}_i S_L(A) = S_{L+1}(A).$$

$\square$

Lemma 8.1.9 (Zero-length word span). [Lean](#) [✓ Stmt](#) If $D \geq 2$, then $S_0(A) \neq M_D(\mathbb{C})$.

Proof. [✓ Proof](#) The zero-length products consist only of the identity, so $S_0(A) = \text{span}_{\mathbb{C}}\{1\}$ has dimension 1. Since $\dim M_D(\mathbb{C}) = D^2 \geq 4$, this span cannot be all of $M_D(\mathbb{C})$. $\square$

Corollary 8.1.10 (0-block injectivity). [Lean](#) [✓ Stmt](#) If $D \geq 1$ and A is 0-block injective, then $D = 1$.

Proof. [✓ Proof](#) By Lemma 8.1.7, 0-block injectivity gives $S_0(A) = M_D(\mathbb{C})$. Lemma 8.1.9 excludes $D \geq 2$, hence the nonzero bond dimension must be 1. $\square$

Lemma 8.1.11 (Two-sided span from one nonzero matrix). [Lean](#) [✓ Stmt](#) If $C \in M_D(\mathbb{C})$ is nonzero, then

$$\text{span}_{\mathbb{C}}\{RCS : R, S \in M_D(\mathbb{C})\} = M_D(\mathbb{C}). \quad (8.1)$$

This is the matrix-factorization input used in [PGVWC07, Lemma 3].

Proof. [✓ Proof](#) Choose a nonzero entry C_{pq} . Then every matrix unit E_{ij} is a scalar multiple of $E_{ip}CE_{qj}$, so the span in (8.1) contains the standard matrix basis. $\square$

Lemma 8.1.12 (Exact words around a nonzero middle matrix). [Lean](#) [✓ Stmt](#) If A is L -block injective and $X \neq 0$, then

$$\text{span}_{\mathbb{C}}\{A^u X A^v : |u| = |v| = L\} = M_D(\mathbb{C}).$$

Proof. [✓ Proof](#) By L -block injectivity, the length- L word products span $M_D(\mathbb{C})$. Bilinearity of $(R, S) \mapsto RXS$ reduces the preceding lemma to products with R and S chosen among those word products. $\square$

8.2 Fitting decomposition

Definition 8.2.1 (Fitting decomposition). [Lean](#) [✓ Stmt](#) A *Fitting decomposition* of a linear endomorphism $f : V \rightarrow V$ on a finite-dimensional vector space over an algebraically closed field consists of:

1. f is nilpotent on the generalized 0-eigenspace,
2. f is invertible on each generalized μ -eigenspace for $\mu \neq 0$,
3. the generalized eigenspaces span V ,
4. the generalized eigenspaces are linearly independent.

Theorem 8.2.2 (Fitting decomposition exists). [Lean](#) [✓ Stmt](#) Every linear endomorphism on a finite-dimensional vector space over an algebraically closed field admits a Fitting decomposition.

Proof. [✓ Proof](#) Nilpotency on the zero generalized eigenspace follows from the definition of generalized eigenspaces. Invertibility on nonzero generalized eigenspaces follows because $f - \mu$ is nilpotent there, so $f = \mu(1 - (1 - f/\mu))$ is invertible. Spanning and independence are standard results for generalized eigenspaces over algebraically closed fields. In [SPGWC10] and [Wol12, Lemma 6.3(b)], the Jordan normal form is used directly. The argument is phrased in terms of generalized eigenspaces instead. $\square$

Theorem 8.2.3 (Nilpotent power bound). [Lean](#) [✓ Stmt](#) If f is nilpotent on a space of dimension n , then $f^n = 0$.

Proof. [✓ Proof](#) A nilpotent endomorphism on an n -dimensional space has nilpotency index $\leq n$. $\square$

8.3 Nonzero trace product

Theorem 8.3.1 (Cumulative span reaches $M_D(\mathbb{C})$ — explicit bound). [Lean](#) [✓ Stmt](#) If A is normal, then there exists $n \leq D^2$ with $T_n(A) = M_D(\mathbb{C})$.

Proof. [✓ Proof](#) By Theorem 8.1.6, if $T_n \neq T_{n+1}$ then $\dim T_{n+1} > \dim T_n$. Since $\dim T_n \leq D^2$ (Lemma 8.1.4), after at most D^2 steps either $T_n = M_D(\mathbb{C})$ or T_n has stabilised. Stabilisation before reaching $M_D(\mathbb{C})$ contradicts normality (which requires $S_{L_0} = M_D(\mathbb{C})$ for some L_0 , hence $T_{L_0} = M_D(\mathbb{C})$). $\square$

Theorem 8.3.2 (Cumulative span reaches $M_D(\mathbb{C})$). [Lean](#) [✓ Stmt](#) If A is normal, then $T_{D^2}(A) = M_D(\mathbb{C})$.

Proof. [✓ Proof](#) Take n from Theorem 8.3.1; since $n \leq D^2$ and T is monotone, $T_{D^2} = M_D(\mathbb{C})$. $\square$

Theorem 8.3.3 (Nonzero trace product). [Lean](#) [✓ Stmt](#) If A is normal, then there exists a word w of length $\leq D^2$ such that $\text{tr}(A^w) \neq 0$. This is [SPGWC10, Lemma 1]. For the trace argument below, the nontrivial case is $D \geq 1$; when $D = 0$ the normality hypothesis is vacuous.

Proof. [✓ Proof](#) By Theorem 8.3.2, $\mathbb{1} \in T_{D^2}(A)$, so $\mathbb{1}$ is a linear combination of word products of length $\leq D^2$. In the nontrivial case $D \geq 1$, at least one has nonzero trace (since $\text{tr}(\mathbb{1}) = D \neq 0$). $\square$

Theorem 8.3.4 (Cumulative span reaches $M_D(\mathbb{C})$ — sharp bound). [Lean](#) [✓ Stmt](#) *If A is normal, then $T_{D^2-d'+1}(A) = M_D(\mathbb{C})$, where $d' = \dim S_1(A) = \text{kr}(A)$. This is the sharp form of [SPGWC10, Lemma 1]: the cumulative span already reaches $M_D(\mathbb{C})$ by index $D^2 - d' + 1 \leq D^2$.*

Proof. [✓ Proof](#) The key observation is that if $T_n(A) \neq T_{n+1}(A)$, the dimension strictly increases. Since $S_1(A) \subseteq T_1(A)$ and $\dim S_1(A) = d'$, we have $\dim T_1(A) \geq d'$. After at most $D^2 - d'$ additional strict-growth steps the dimension reaches $D^2 = \dim M_D(\mathbb{C})$. Stabilisation before reaching $M_D(\mathbb{C})$ contradicts normality. Hence $T_{D^2-d'+1}(A) = M_D(\mathbb{C})$. $\square$

Theorem 8.3.5 (Sharp nonzero trace product). [Lean](#) [✓ Stmt](#) *If A is normal, then there exists a word w of length $\leq D^2 - \text{kr}(A) + 1$ such that $\text{tr}(A^w) \neq 0$. This is [SPGWC10, Lemma 1] with the sharp length bound.*

Proof. [✓ Proof](#) By Theorem 8.3.4, the identity $\mathbb{1}$ lies in $T_{D^2-d'+1}(A)$, hence is a linear combination of word products of length $\leq D^2 - d' + 1$. At least one such product has nonzero trace. $\square$

Theorem 8.3.6 (Sharp nonzero trace product — positive length). [Lean](#) [✓ Stmt](#) *If A is normal and $D \geq 2$, then there exists a word w of positive length $1 \leq |w| \leq D^2 - \text{kr}(A) + 1$ such that $\text{tr}(A^w) \neq 0$. This strengthens Theorem 8.3.5 by requiring $|w| \geq 1$, a feature needed for the blocking argument in the general Wielandt bound below.*

Proof. [✓ Proof](#) Define the *positive-level* cumulative span $V_n = \text{span}\{A^w : 1 \leq |w| \leq n\}$. For $D \geq 2$, the span $S_0(A) = \text{span}\{\mathbb{1}\}$ has dimension 1, so $V_1 \supseteq S_1(A)$ has dimension $\geq d' \geq 1$. The same stabilization-or-strict-growth dichotomy shows $V_{D^2-d'+1} = M_D(\mathbb{C})$: if V_n stabilises and is not $M_D(\mathbb{C})$, then V_n is closed under left-multiplication by each A^i , so $S_N(A) \subseteq V_n \neq M_D(\mathbb{C})$ for all $N \geq 1$, contradicting normality. Since $V_{D^2-d'+1} = M_D(\mathbb{C})$, the identity $\mathbb{1}$ lies in $V_{D^2-d'+1}$ and is therefore a linear combination of word products of positive length. At least one such product has nonzero trace (since $\text{tr}(\mathbb{1}) = D \geq 2 \neq 0$). $\square$

8.4 Eigenvector extraction

Theorem 8.4.1 (Nonzero trace implies eigenvector). [Lean](#) [✓ Stmt](#) *If $M \in M_D(\mathbb{C})$ has $\text{tr}(M) \neq 0$, then M has a nonzero eigenvalue $\mu \neq 0$ and a corresponding eigenvector $\varphi \neq 0$ with $M\varphi = \mu\varphi$.*

Proof. [✓ Proof](#) The trace is the sum of the eigenvalues of M , counted with multiplicity, so a nonzero trace gives a nonzero eigenvalue μ . Choosing a nonzero vector $\varphi \in \ker(M - \mu\mathbb{1})$ gives $M\varphi = \mu\varphi$. $\square$

Theorem 8.4.2 (Eigenvalue extraction). [Lean](#) [✓ Stmt](#) *If A is normal, then there exist a word w_0 with $|w_0| \leq D^2$, a nonzero scalar $\mu \neq 0$, and a nonzero vector $\varphi \neq 0$ such that $A^{w_0}\varphi = \mu\varphi$.*

Proof. [✓ Proof](#) By Theorem 8.3.3, there is a word w_0 with $\text{tr}(A^{w_0}) \neq 0$ and $|w_0| \leq D^2$. By Theorem 8.4.1, A^{w_0} has a nonzero eigenvector. $\square$

8.5 Eigenvector spreading

Definition 8.5.1 (Cumulative vector span). [Lean](#) [✓ Stmt](#) Given an MPS tensor A and a vector $\varphi \in \mathbb{C}^D$, the *cumulative vector span* is

$$K_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w\varphi : |w| \leq n\} \subseteq \mathbb{C}^D.$$

This is [SPGWC10, proof of Lemma 2(a)].

Theorem 8.5.2 (Vector span stabilisation). [Lean](#) [✓ Stmt](#) If $K_n(A, \varphi) = K_{n+1}(A, \varphi)$, then $K_m(A, \varphi) = K_n(A, \varphi)$ for all $m \geq n$.

Proof. [✓ Proof](#) Same argument as Theorem 8.1.5: left-multiplication by A^i cannot escape the stabilised span. $\square$

Lemma 8.5.3 (Vector span from matrix span). [Lean](#) [✓ Stmt](#) If $T_N(A) = M_D(\mathbb{C})$ and $\varphi \neq 0$, then $K_N(A, \varphi) = \mathbb{C}^D$.

Proof. [✓ Proof](#) Every matrix $M \in M_D(\mathbb{C})$ is in $T_N(A)$, so in particular every rank-one matrix sending φ to any target vector v is in T_N . Applying such matrices to φ recovers all of $\mathbb{C}^D$. $\square$

Theorem 8.5.4 (Eigenvector spreading). [Lean](#) [✓ Stmt](#) Let A be a normal MPS tensor and let $A^{i_0} \varphi = \mu \varphi$ with $\mu \neq 0$ and $\varphi \neq 0$. Then $K_{D-1}(A, \varphi) = \mathbb{C}^D$. This is [SPGWC10, Lemma 2(a)].

Proof. [✓ Proof](#) The vector span K_n is monotone and $\dim K_n \leq D$. If $K_n = K_{n+1}$ for some $n < D - 1$, the span stabilises (Theorem 8.5.2). But $T_{D^2}(A) = M_D(\mathbb{C})$ by Theorem 8.3.2, so Lemma 8.5.3 gives $K_{D^2}(A, \varphi) = \mathbb{C}^D$, contradicting early stabilisation. Hence $\dim K_n$ grows by ≥ 1 at each step, reaching D by step $D - 1$. $\square$

8.6 Proof of the cumulative Wielandt bound

Theorem 8.6.1 (Wielandt bound). [Lean](#) [✓ Stmt](#) If A is a normal MPS tensor with bond dimension D , then $T_{D^2}(A) = M_D(\mathbb{C})$. In particular, the cumulative span stabilises at index at most D^2 . This is the cumulative-span part of [SPGWC10, Theorem 1]. The cited theorem is stronger: via Lemma 2(b) it upgrades this to a fixed-length conclusion $S_N(A) = M_D(\mathbb{C})$, stated below as a separate statement.

Proof. [✓ Proof](#) The claim follows from Theorem 8.3.2. $\square$

Remark 8.6.2 (Explicit blocking-length estimates). [✓ Stmt](#) Theorem 8.6.1 gives the D^2 cumulative-span bound. The full-rank index bound $\iota(A) \leq (D^2 - d + 1) \cdot D^2$ ([SPGWC10, Theorem 1, case (1)]) is proved in Theorem 8.6.15 below. The left-canonical normal blocked-injectivity estimate D^4 is proved below in Theorem 8.7.6. The sharper bi-canonical blocking estimate $3D^5$ from [SPGWC10] is not proved here.

Definition 8.6.3 (One-step augmentation). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor with Kraus family $\{A^i\}_{i=0}^{d-1}$, and let $X \in M_D(\mathbb{C})$. The *one-step augmentation* of A by X is the tensor $\tilde{A}$ with physical dimension $d + 1$ whose first Kraus operator is X and whose remaining Kraus operators are those of A :

$$\tilde{A}^0 = X, \quad \tilde{A}^{i+1} = A^i.$$

Lemma 8.6.4 (Kraus operators lie in the one-step span). [Lean](#) [✓ Stmt](#) Every Kraus operator A^i belongs to the one-step span $S_1(A)$.

Proof. [✓ Proof](#) This is the length-one word with letter i . $\square$

Lemma 8.6.5 (Redundant one-step augmentation). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $X \in S_1(A)$, and let $\tilde{A}$ be the one-step augmentation of A by X . Then

$$S_n(\tilde{A}) = S_n(A)$$

for every $n \geq 0$.

Proof. [✓ Proof](#) The new generator X is already in $S_1(A)$, and each original generator remains in the augmented family. Hence the one-step spans agree. Multiplying word spans and arguing by induction on the word length gives equality at every length. $\square$

Lemma 8.6.6 (Normality and Kraus rank under one-step augmentation). [Lean](#) [Lean](#) [✓ Stmt](#) Let $X \in S_1(A)$, and let $\tilde{A}$ be the one-step augmentation of A by X . If A is normal, then $\tilde{A}$ is normal, and

$$\text{kr}(\tilde{A}) = \text{kr}(A).$$

Proof. [✓ Proof](#) Since all exact word spans are unchanged by Lemma 8.6.5, eventual full word span for A gives eventual full word span for $\tilde{A}$. The equality of Kraus ranks is the equality of the dimensions of the common one-step span. $\square$

Theorem 8.6.7 (Wielandt case (2), invertible generator). [Lean](#) [Lean](#) [✓ Stmt](#) Let A be a normal MPS tensor with bond dimension D , and suppose that some Kraus operator A^{i_0} is invertible. Then

$$S_{D^2 - \text{kr}(A) + 1}(A) = M_D(\mathbb{C}), \quad \iota(A) \leq D^2 - \text{kr}(A) + 1.$$

Proof. [✓ Proof](#) The dimensions of the exact word spans cannot decrease, because left multiplication by the invertible Kraus operator is injective. If this dimension growth stopped before the full matrix algebra was reached, the same argument applied at all later lengths would contradict normality. Since $\dim S_1(A) = \text{kr}(A)$ and $\dim M_D(\mathbb{C}) = D^2$, the full algebra is reached by length $D^2 - \text{kr}(A) + 1$. The index bound follows from the definition of $\iota(A)$. $\square$

Theorem 8.6.8 (Wielandt case (2), one-step invertible element). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor with bond dimension $D \geq 1$, satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and suppose that A is primitive. If $S_1(A)$ contains an invertible matrix X , then

$$S_{D^2 - \text{kr}(A) + 1}(A) = M_D(\mathbb{C}). \tag{8.2}$$

In the terminology of [SPGWC10], this is the case where the first application space $S_1(A)$ contains an invertible operator. This is the exact word-span form of [SPGWC10, Theorem 1, case (2)].

Proof. [✓ Proof](#) Adjoin X as an extra first Kraus operator, obtaining $\tilde{A}$. Primitivity and normalization give normality of A , hence normality of $\tilde{A}$; the Kraus rank and all exact word spans are unchanged. The first Kraus operator of $\tilde{A}$ is invertible, so Theorem 8.6.7 gives $S_{D^2 - \text{kr}(\tilde{A}) + 1}(\tilde{A}) = M_D(\mathbb{C})$. Transfer this equality back to A . $\square$

Corollary 8.6.9 (Wielandt case (2), index bound). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 8.6.8,

$$\iota(A) \leq D^2 - \text{kr}(A) + 1,$$

where $\iota(A) = \min\{n : S_n(A) = M_D(\mathbb{C})\}$.

Proof. [✓ Proof](#) The word-span equality (8.2) gives an admissible value in the defining minimum for $\iota(A)$. $\square$

Corollary 8.6.10 (Wielandt case (2), invertible Kraus operator). [Lean](#) [Lean](#) [✓ Stmt](#) Let A be an MPS tensor with bond dimension $D \geq 1$, satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and suppose that A is primitive. If some Kraus operator A^{i_0} is invertible, then

$$S_{D^2 - \text{kr}(A) + 1}(A) = M_D(\mathbb{C}), \quad \iota(A) \leq D^2 - \text{kr}(A) + 1.$$

Proof. ✓ **Proof** Apply the one-step case to $X = A^{i_0}$, which belongs to $S_1(A)$ by Lemma 8.6.4. $\square$

Lemma 8.6.11 (Rank-one extraction from a non-invertible eigenvector). Lean ✓ **Stmt** *Let A be a normal MPS tensor with bond dimension $D \geq 1$. Suppose that a Kraus operator A^{i_0} is non-invertible and that there exists a vector $\varphi \in \mathbb{C}^D$ and a nonzero scalar μ such that $A^{i_0}\varphi = \mu\varphi$. Then, for every $\psi \in \mathbb{C}^D$,*

$$|\varphi\rangle\langle\psi| \in S_{D^2-D+1}(A).$$

Proof. ✓ **Proof** Let r be the nilpotent index of left multiplication by A^{i_0} . The dimension-growth argument gives a level n_0 for which

$$(A^{i_0})^r S_{n_0}(A) = \{(A^{i_0})^r M : M \in M_D(\mathbb{C})\}.$$

Since $A^{i_0}\varphi = \mu\varphi$ and $\mu \neq 0$, iterating gives $(A^{i_0})^r\varphi = \mu^r\varphi$, hence

$$\varphi = \mu^{-r}(A^{i_0})^r\varphi \in \text{range}((A^{i_0})^r).$$

Hence, for every $\psi \in \mathbb{C}^D$,

$$|\varphi\rangle\langle\psi| \in S_{r+n_0}(A).$$

The nilpotent-index estimate gives

$$r + n_0 \leq D^2 - D + 1.$$

Since

$$A^{i_0}|\varphi\rangle\langle\psi| = |A^{i_0}\varphi\rangle\langle\psi| = \mu|\varphi\rangle\langle\psi|$$

and $\mu \neq 0$, the equality

$$|\varphi\rangle\langle\psi| = \mu^{-1}A^{i_0}|\varphi\rangle\langle\psi|$$

implies

$$|\varphi\rangle\langle\psi| \in S_k(A) \implies |\varphi\rangle\langle\psi| \in S_{k+1}(A)$$

because left multiplication by A^{i_0} maps $S_k(A)$ into $S_{k+1}(A)$. Iterating from $k = r + n_0$ and using $r + n_0 \leq D^2 - D + 1$ gives

$$|\varphi\rangle\langle\psi| \in S_{D^2-D+1}(A).$$

$\square$

Theorem 8.6.12 (Wielandt case (3), one-step non-invertible element). Lean ✓ **Stmt** *Let A be an MPS tensor with bond dimension $D \geq 1$, satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and suppose that A is primitive. If $S_1(A)$ contains a non-invertible matrix X with a nonzero eigenvalue, then*

$$S_{D^2}(A) = M_D(\mathbb{C}).$$

More explicitly, it is enough to have $X\varphi = \mu\varphi$ with $\varphi \neq 0$ and $\mu \neq 0$. In the terminology of [SPGWC10], this is the case where the first application space $S_1(A)$ contains a non-invertible operator with a nonzero eigenvalue. This is the exact word-span form of [SPGWC10, Theorem 1, case (3)].

Proof. ✓ **Proof** Adjoin X as an extra first Kraus operator. The augmented tensor is normal and has the same exact word spans as A . The eigenvector relation for X is now a one-generator eigenvector relation. The eigenvector-spreading argument gives full fixed-length vector span by length $D - 1$, and Lemma 8.6.11 gives the rank-one operators $|\varphi\rangle\langle\psi|$ in length $D^2 - D + 1$. Multiplying these two fixed-length spans gives S_{D^2} for the augmented tensor, hence for A . $\square$

Corollary 8.6.13 (Wielandt case (3), index bound). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 8.6.12,

$$\iota(A) \leq D^2.$$

Proof. [✓ Proof](#) The equality $S_{D^2}(A) = M_D(\mathbb{C})$ gives an admissible value in the defining minimum for $\iota(A)$. $\square$

Corollary 8.6.14 (Wielandt case (3), non-invertible Kraus operator). [Lean](#) [Lean](#) [✓ Stmt](#) Let A be an MPS tensor with bond dimension $D \geq 1$, satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and suppose that A is primitive. If some Kraus operator A^{i_0} is non-invertible and has an eigenvector $\varphi \neq 0$ with nonzero eigenvalue μ , then

$$S_{D^2}(A) = M_D(\mathbb{C}), \quad \iota(A) \leq D^2.$$

Proof. [✓ Proof](#) Apply the one-step case to $X = A^{i_0}$, which belongs to $S_1(A)$ by Lemma 8.6.4. $\square$

Theorem 8.6.15 (Wielandt's inequality — general bound). [Lean](#) [✓ Stmt](#) Let A be a normalized MPS tensor ($\sum_i (A^i)^\dagger A^i = \mathbb{1}$) that is primitive. Then

$$\iota(A) \leq (D^2 - \text{kr}(A) + 1) \cdot D^2,$$

where $\iota(A) = \min\{n : S_n(A) = M_D(\mathbb{C})\}$ is the full-Kraus-rank index. This is [SPGWC10, Theorem 1, case (1)].

Proof. [✓ Proof](#)

1. $D = 1$: In this case $\iota(A) = 0$, so the bound holds trivially.
2. $D \geq 2$: By Theorem 8.3.6, there exists a *positive-length* word w with $|w| \leq D^2 - d' + 1$ and $\text{tr}(A^w) \neq 0$, where $d' = \text{kr}(A)$. The matrix A^w has nonzero trace, hence a nonzero eigenvalue μ and eigenvector φ (Theorem 8.4.1). Set $n = |w|$ and let $B = A^{[n]}$ be the blocked tensor. Then B is normal (Theorem 8.8.1), and the encoding of w as a single block index gives $B^{i_0} = A^w$.
 - (a) If A^w is invertible (invertible case), $S_{D^2}(B) = M_D(\mathbb{C})$, whence $S_{D^2 \cdot n}(A) = M_D(\mathbb{C})$.
 - (b) If A^w is non-invertible (non-invertible case), the eigenvector φ spans a proper subspace, and the rank-one extraction argument gives $S_{D^2}(B) = M_D(\mathbb{C})$, whence $S_{D^2 \cdot n}(A) = M_D(\mathbb{C})$.

In the invertible case, the invertible Kraus operator B^{i_0} allows direct dimension growth, giving $S_{D^2 - k_B + 1}(B) = M_D(\mathbb{C})$ where $k_B = \text{kr}(B)$. In the non-invertible case, the bound $S_{D^2}(B) = M_D(\mathbb{C})$ follows from tracking indices through the rank-one extraction argument of [SPGWC10, Lemma 2(b)]; the separate qualitative fixed-length conclusion stated below does not supply this explicit bound. In both cases $\iota(A) \leq D^2 \cdot n \leq D^2 \cdot (D^2 - d' + 1)$. $\square$

Remark 8.6.16 (Quantitative and qualitative Wielandt bounds). [✓ Stmt](#) The quantitative bound $\iota(A) \leq (D^2 - d' + 1) \cdot D^2$ is cited from [SPGWC10, Theorem 1, case (1)]. The blueprint's own lemmas prove the qualitative conclusion used on the fundamental-theorem route, namely Theorem 8.8.6: there exists N with $S_N(A) = M_D(\mathbb{C})$. The quantitative bound is not on the fundamental-theorem critical path.

8.7 Fixed-length matrix spanning

The preceding sections establish the cumulative-spanning part of the Lemma 2(a) argument from [SPGWC10]: eigenvector spreading gives a cumulative vector span $K_{D-1}(A, \varphi) = \mathbb{C}^D$. The next stage is to pass to a span at one fixed length, namely to find n_0 such that $S_{n_0}(A) = M_D(\mathbb{C})$ (i.e. word products of *exactly* one length span all matrices).

Definition 8.7.1 (Fixed-length vector span). Lean ✓ Stmt The *fixed-length vector span* at length n is

$$H_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w \varphi : |w| = n\} \subseteq \mathbb{C}^D.$$

This is $S_n(A)|\varphi\rangle$ in the notation of [SPGWC10].

Lemma 8.7.2 (Word span products). Lean ✓ Stmt $S_m(A) \cdot S_n(A) \subseteq S_{m+n}(A)$: the product of word spans at lengths m and n is contained in the word span at length $m + n$.

Proof. ✓ Proof Every product $A^{w_1} A^{w_2}$ with $|w_1| = m$ and $|w_2| = n$ equals $A^{w_1 w_2}$ with $|w_1 w_2| = m + n$. □

Lemma 8.7.3 (Eigenvector padding). Lean ✓ Stmt Suppose $A^{i_0} \varphi = \mu \varphi$ with $\mu \neq 0$. Then for all n ,

$$K_n(A, \varphi) = H_n(A, \varphi).$$

In particular, if $K_n(A, \varphi) = \mathbb{C}^D$ then already $H_n(A, \varphi) = \mathbb{C}^D$.

Proof. ✓ Proof Any word w with $|w| \leq n$ can be padded to exact length n by appending $n - |w|$ copies of the index i_0 . The eigenvector relation gives $A^{w \cdot i_0^k} \varphi = \mu^k \cdot A^w \varphi$, so the padded vector is a nonzero scalar multiple of the original. Hence every generator of K_n lies in H_n . The reverse inclusion $H_n \leq K_n$ is immediate. □

Theorem 8.7.4 (Fixed-length vector spanning implies matrix spanning). Lean ✓ Stmt Assume:

1. $H_n(A, \varphi) = \mathbb{C}^D$ (length- n word products applied to φ span all of $\mathbb{C}^D$), and
2. for each standard basis vector e_j , the rank-one operator $|\varphi\rangle\langle e_j|$ lies in $S_m(A)$.

Then $S_{n+m}(A) = M_D(\mathbb{C})$.

Proof. ✓ Proof Fix a matrix unit E_{ij} . By hypothesis (1), there exists $M_i \in S_n(A)$ with $M_i \varphi = e_i$. By hypothesis (2), $|\varphi\rangle\langle e_j| \in S_m(A)$. Their product satisfies

$$M_i \cdot |\varphi\rangle\langle e_j| = |M_i \varphi\rangle\langle e_j| = |e_i\rangle\langle e_j| = E_{ij},$$

and $E_{ij} \in S_{n+m}(A)$ by Lemma 8.7.2. Since the matrix units span $M_D(\mathbb{C})$, $S_{n+m}(A) = M_D(\mathbb{C})$. □

Lemma 8.7.5 (Blocking transfer). Lean ✓ Stmt If the blocked tensor $A^{[L]}$ satisfies $S_n(A^{[L]}) = M_D(\mathbb{C})$, then $S_{nL}(A) = M_D(\mathbb{C})$.

Proof. ✓ Proof Each length- n word in the blocked alphabet $\{0, \dots, d^L - 1\}$ decodes to a length- nL word in the original alphabet $\{0, \dots, d - 1\}$, so $S_n(A^{[L]}) \subseteq S_{nL}(A)$. □

Theorem 8.7.6 (Bounded injective blocking for left-canonical normal tensors). [Lean](#) [✓ Stmt](#) Let A be a left-canonical normal MPS tensor with bond dimension $D > 0$. Then there is a positive blocking length $L \leq D^4$ such that the blocked tensor $A^{[L]}$ is one-site injective.

This is the blocked-injectivity form of the quantum Wielandt input used in the Fundamental Theorem for translation-invariant tensors. It corresponds to the statement in [CPGSV16, Section II, line 332] that every normal tensor becomes injective after at most D^4 blockings, using the index estimate of [SPGWC10, Theorem 1]. The Lean statement includes the left-canonical/trace-preserving hypothesis because that is the canonical-form context in which the blocked-injectivity input is consumed.

Proof. [✓ Proof](#) The proof uses the full-Kraus-rank index bound from the quantum Wielandt inequality. Since A is normal, the index is attained; the one-step Kraus rank is nonzero in the left-canonical case, so the bound $(D^2 - \text{rank } S_1(A) + 1)D^2$ is at most D^4 . The passage from exact word-span fullness to injectivity of the blocked tensor is Lemma 8.7.5. $\square$

Theorem 8.7.7 (Common exact block-injectivity length). [Lean](#) [✓ Stmt](#) Let $(A_j)_{j=1}^g$ be a finite family of positive-bond-dimension, left-canonical normal tensors. There is a positive integer L such that every A_j is L -block injective.

Proof. [✓ Proof](#) Apply Theorem 8.7.6 to each block. The product of the finitely many positive lengths is a common positive multiple, and exact block injectivity persists under positive multiples. $\square$

Corollary 8.7.8 (Uniform injective blocking for separated normal-canonical families). [Lean](#) [Lean](#) [✓ Stmt](#) Let $(A_x)_{x \in X}$ be a finite separated normal-canonical family whose blocks have positive virtual bond dimension. Then there is a positive integer L such that every blocked tensor $A_x^{[L]}$ is one-site injective. The same conclusion holds simultaneously for two finite separated normal-canonical families whose blocks have positive virtual bond dimension.

Here the family is a basis of normal-canonical representatives in the sense of Definition 10.1.19; the blocks are distinct and the weight moduli are non-increasing but need not be strictly ordered. It assumes that the representative family has already been selected; it does not reconstruct the full CPSV sector decomposition with repeated copies inside one sector. The restriction is recorded in `docs/paper-gaps/ft_one_copy_scope_restriction.tex`.

Proof. [✓ Proof](#) Apply Theorem 8.7.6 to each block. Taking the product of the finitely many resulting positive blocking lengths gives a common positive multiple, and injectivity persists under positive further blocking. $\square$

Remark 8.7.9 (Relation with the rank-one extraction lemma). [✓ Stmt](#) In [SPGWC10], Lemma 2(b) combines a rank-one extraction for fixed-length spans with a product argument turning fixed-length vector spanning into fixed-length matrix spanning. Theorem 8.7.4 is the second step. Theorem 8.8.5 supplies the rank-one hypothesis in blocked form, and Theorem 8.8.6 then gives a blocked, fixed-length word-span saturation result.

Remark 8.7.10 (Use of the quantum Wielandt theorem in the MPS route). The source papers use the quantum Wielandt theorem as an input converting normality into injectivity after blocking. In the notation of this chapter, the input has two distinct conclusions. First, Theorem 8.6.1 gives cumulative saturation $T_{D^2}(A) = M_D(\mathbb{C})$. This alone is not the injectivity statement used in canonical form, because injectivity requires products of one fixed length. Second, Lemma 2(b) of [SPGWC10], in the form of Theorem 8.8.6, supplies a length N for which $S_N(A) = M_D(\mathbb{C})$. This is the statement that matches the injectivity of Γ_N and can be transported through blocking by Lemma 8.7.5. Thus any later use of Wielandt in the Fundamental Theorem must specify

whether it is using cumulative span, exact-length span, or the blocked version of the exact-length conclusion.

8.8 Blocked tensor and rectangular span

This section extends the Lemma 2(b) argument to the *blocked tensor* $A^{[L]}$ and develops the “rectangular span” used to produce rank-one elements.

Theorem 8.8.1 (Blocking preserves normality). [Lean](#) [✓ Stmt](#) *If A is normal and $L > 0$, then the blocked tensor $A^{[L]}$ is also normal.*

Proof. [✓ Proof](#) Choose N with $S_N(A) = M_D(\mathbb{C})$. Then $\mathbb{1} \in S_N(A)$. For any $M \in S_N(A)$, write $\mathbb{1} = \sum_r c_r A^{w_r}$ with each $|w_r| = N$. Then

$$M = M \cdot \mathbb{1} = \sum_r c_r M A^{w_r} \in S_N(A) \cdot S_N(A) \subseteq S_{2N}(A) \quad (8.3)$$

by Lemma 8.7.2. Iterating the inclusion (8.3) gives $S_N(A) \subseteq S_{kN}(A)$ for every $k \geq 1$; taking $k = L$ yields $M_D(\mathbb{C}) = S_N(A) \subseteq S_{NL}(A)$. By Lemma 8.8.2, $S_{NL}(A) \subseteq S_N(A^{[L]})$, so $S_N(A^{[L]}) = M_D(\mathbb{C})$. $\square$

Lemma 8.8.2 (Chunking: word span embeds into blocked word span). [Lean](#) [✓ Stmt](#) $S_{nL}(A) \subseteq S_n(A^{[L]})$ for all n, L .

Proof. [✓ Proof](#) Every length- nL word in the original alphabet can be split into n consecutive chunks of length L , each of which is a single blocked Kraus operator. By induction on n , using the word span product lemma (Lemma 8.7.2). $\square$

Theorem 8.8.3 (Word eigenvector gives blocked eigenvector). [Lean](#) [✓ Stmt](#) *If $A^{w_0} \varphi = \mu \varphi$ for a word w_0 of length L , then $(A^{[L]})^{j_0} \varphi = \mu \varphi$, where j_0 is the encoded blocked index corresponding to w_0 .*

Proof. [✓ Proof](#) By definition, $(A^{[L]})^{j_0} = A^{w_0}$. $\square$

Theorem 8.8.4 (Blocked fixed-length matrix spanning). [Lean](#) [✓ Stmt](#) *Suppose A is normal and $L > 0$, and we have:*

1. a word w_0 of length L with eigenvector $A^{w_0} \varphi = \mu \varphi$ ($\mu \neq 0, \varphi \neq 0$),
2. a word τ_0 of length L with transpose eigenvector $(A^{\tau_0})^T \psi = \nu \psi$ ($\nu \neq 0, \psi \neq 0$),
3. a rank-one element $\varphi \psi^T \in S_m(A^{[L]})$ for some m .

Then there exists N such that $S_N(A) = M_D(\mathbb{C})$.

Remark. The proof gives the explicit bound $N = ((D-1) + m + (D-1)) \cdot L$; the statement above gives only the existential conclusion.

Proof. [✓ Proof](#) Apply Theorem 8.7.4 to the blocked tensor $A^{[L]}$, which is normal by Theorem 8.8.1. The eigenvector and transpose eigenvector are transferred to single-index eigenvectors of $A^{[L]}$ by Theorem 8.8.3. Transfer back to the original tensor via Lemma 8.7.5. $\square$

Theorem 8.8.5 (Rank-one extraction for blocked tensor). [Lean](#) [✓ Stmt](#) *Suppose A is normal and $N_0 \geq 1$ with $S_{N_0}(A) = M_D(\mathbb{C})$. Then there exist words σ_0, τ_0 of length N_0 , nonzero vectors φ, ψ , nonzero scalars μ, ν , and $m \in \mathbb{N}$ such that:*

1. $A^{\sigma_0} \varphi = \mu \varphi$,
2. $(A^{\tau_0})^T \psi = \nu \psi$,
3. $\varphi \psi^T \in S_m(A^{[N_0]})$.

Proof. ✓ Proof Set $B = A^{[N_0]}$. Since $S_{N_0}(A) = M_D(\mathbb{C})$, we have $S_1(B) = M_D(\mathbb{C})$ by definition of the blocked tensor. Hence $\mathbb{1} \in S_1(B)$, so in the nontrivial case $D \geq 1$ not all blocked generators can have trace zero. Choose j_0 with $\text{tr}(B^{j_0}) \neq 0$. By Theorem 8.4.1, B^{j_0} has a nonzero eigenvector φ with eigenvalue $\mu \neq 0$. Writing j_0 as the blocked index corresponding to a word σ_0 of length N_0 gives $A^{\sigma_0} \varphi = \mu \varphi$. Applying the same argument to the transpose blocked tensor gives a word τ_0 of length N_0 , a nonzero vector ψ , and $\nu \neq 0$ with $(A^{\tau_0})^T \psi = \nu \psi$. Since A is normal and $N_0 \geq 1$, the blocked tensor B is normal by Theorem 8.8.1. Therefore there exists M with $S_M(B) = M_D(\mathbb{C})$. In particular $\varphi \psi^T \in S_M(B)$. Taking $m = M$ proves (3). $\square$

Theorem 8.8.6 (Fixed-length word-span saturation via blocking). Lean ✓ Stmt *If A is a normal MPS tensor with bond dimension D , then there exists N such that $S_N(A) = M_D(\mathbb{C})$. This gives the fixed-length spanning conclusion corresponding to [SPGWC10, Lemma 2(b)].*

Proof. ✓ Proof By normality, some N_0 has $S_{N_0}(A) = M_D(\mathbb{C})$. If $N_0 = 0$ we are done. Otherwise, Theorem 8.8.5 produces eigenvectors and a rank-one element $\varphi \psi^T \in S_m(A^{[N_0]})$. Theorem 8.8.4 then gives $S_{((D-1)+m+(D-1)) \cdot N_0}(A) = M_D(\mathbb{C})$. $\square$

Remark 8.8.7 (Blocking argument for fixed-length spanning). ✓ Stmt Theorem 8.8.6 is obtained by first extracting a rank-one operator in a blocked tensor (Theorem 8.8.5) and then applying the fixed-length matrix-spanning theorem (Theorem 8.8.4). Thus the fixed-length span of A is reached through a blocked auxiliary tensor rather than by a direct unblocked argument.

8.9 Primitive MPS tensors

Definition 8.9.1 (Primitive MPS tensor). Lean ✓ Stmt Assume $D > 0$. A tensor A together with a matrix $\rho \in M_D(\mathbb{C})$ is *primitive* if

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}, \quad \rho \geq 0, \quad \rho \neq 0, \quad \mathcal{E}_A(\rho) = \rho,$$

and, if P is the fixed-point projection associated to ρ , then $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$. When the choice of ρ is irrelevant, we simply say that A is a primitive MPS tensor. This is weaker than the primitive condition of [SPGWC10], which requires $\rho > 0$; see Remark 8.10.1.3. The stronger positive-definite hypothesis is imposed separately in Lemma 8.10.2.1 and Theorem 8.10.4.3.

Theorem 8.9.2 (Primitive overlap convergence). Lean ✓ Stmt *If A is a primitive MPS tensor, then $O_{AA}(N) \rightarrow 1$.*

Proof. ✓ Proof The claim follows from the self-overlap convergence under a complementary transfer-map gap (Theorem 7.15.2). $\square$

Theorem 8.9.3 (Peripheral primitivity and irreducibility give normalized overlap). Lean ✓ Stmt *Let A have positive bond dimension and satisfy*

$$\sum_i (A^i)^\dagger A^i = I.$$

If A is tensor-irreducible and the transfer map is primitive, then $O_{AA}(N) \rightarrow 1$.

Proof. ✓ Proof Irreducible Perron–Frobenius theory supplies a nonzero positive fixed point. Peripheral primitivity gives a strict spectral-radius bound on the complementary transfer map. Thus A is a primitive MPS tensor, and Theorem 8.9.2 applies. $\square$

8.10 Primitivity and normality

Throughout this section, primitivity means the condition of Definition 8.9.1: the tensor is in TP gauge, it has a nonzero PSD fixed point, and the complementary transfer map has spectral radius < 1 .

Lemma 8.10.1 (Primitive MPS tensors have nonzero word products). Lean ✓ Stmt *If A is a primitive MPS tensor, then for every $n \geq 0$ there exists a word w of length exactly n with $A^w \neq 0$.*

Proof. ✓ Proof The PSD fixed point ρ of $\mathcal{E}_A$ satisfies $\mathcal{E}_A^n(\rho) = \rho \neq 0$. Since $\mathcal{E}_A^n(\rho) = \sum_{|w|=n} A^w \rho (A^w)^\dagger$, at least one summand is nonzero, hence some $A^w \neq 0$. $\square$

Lemma 8.10.2 (Iterated transfer does not vanish). Lean ✓ Stmt *If A is a primitive MPS tensor, then $\mathcal{E}_A^n \neq 0$ for all $n \geq 0$.*

Proof. ✓ Proof By Lemma 8.10.1, there exists a word w of length n with $A^w \neq 0$, so $\mathcal{E}_A^n(\rho)$ has a nonzero summand. $\square$

8.10.1 From primitivity to normality

The primitivity condition of Definition 8.9.1 combines TP normalization, a nonzero PSD fixed point, and a complementary transfer-map gap. It is connected to normality through the following results.

Lemma 8.10.1.1 (Fixed-point uniqueness from a complementary transfer-map gap). Lean ✓ Stmt *If A is a primitive MPS tensor with PSD fixed point ρ , then every fixed point σ of $\mathcal{E}_A$ is proportional to ρ : $\sigma = \frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \rho$.*

Proof. ✓ Proof Set $\sigma' = \sigma - \frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \rho$. Then $\text{tr}(\sigma') = 0$ and $(\mathcal{E}_A - P_\rho)(\sigma') = \sigma'$. If $\sigma' \neq 0$, then 1 is an eigenvalue of $\mathcal{E}_A - P_\rho$, contradicting the complementary transfer-map gap $\rho_{\text{spec}}(\mathcal{E}_A - P_\rho) < 1$. $\square$

Lemma 8.10.1.2 (Complement powers tend to zero). Lean ✓ Stmt *If A is a primitive MPS tensor with PSD fixed point ρ , then $(\mathcal{E}_A - P_\rho)^n \rightarrow 0$ in operator norm.*

Proof. ✓ Proof The spectral radius of $\mathcal{E}_A - P_\rho$ is strictly less than 1 by the complementary transfer-map gap hypothesis. Apply the general result that powers of an operator with spectral radius < 1 tend to zero. $\square$

Remark 8.10.1.3 (Primitivity: transfer-map gap vs. positive definiteness). ✓ Stmt The primitivity condition used in Section 8.10.1 has three parts: TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, a nonzero PSD fixed point ρ , and the complementary transfer-map gap $\rho_{\text{spec}}(\mathcal{E}_A - P_\rho) < 1$. The notion of “primitive” in [SPGWC10, Proposition 3] additionally requires ρ to be positive definite (full rank). Without positive definiteness, the transfer-map gap condition alone does *not* imply irreducibility or normality. A counterexample is $D = 2$, $\rho = \text{diag}(1, 0)$.

With the additional hypothesis $\rho > 0$, Lemma 8.10.2.1 gives irreducibility. Together with the Burnside argument below (Section 8.10.3) and Theorem 8.11.7, one obtains

$$\text{primitive} + \rho \text{ positive definite} \Rightarrow \text{irreducible} \Rightarrow \text{alg}(A) = M_D(\mathbb{C}) \Rightarrow \text{normal}.$$

The last step uses the aperiodicity condition $\mathbb{1} \in S_1(A)$ from Remark 8.11.1. After restoring the positive-definite hypothesis, [SPGWC10, Proposition 3] identifies that primitive/strongly irreducible condition with eventual full Kraus rank, i.e. with our notion of normality.

8.10.2 Primitive implies irreducible

Lemma 8.10.2.1 (Primitive tensors with positive-definite fixed point are irreducible). [Lean](#) [Stmnt](#) If A is a primitive MPS tensor with PSD fixed point ρ and ρ is positive definite, then A is an irreducible tensor. This is part of [SPGWC10, Proposition 3] (see also [CPGSV16, Section 2.3]).

Proof. [Lean](#) [Proof](#) By contrapositive. Assume A has an invariant projection P ($P \neq 0$, $P \neq \mathbb{1}$, $(\mathbb{1} - P)A^iP = 0$ for all i). Set $\sigma = P\rho P$.

1. *Invariance of σ under $\mathcal{E}_A^n$:* the projection relation $(\mathbb{1} - P)A^iP = 0$ implies $(\mathbb{1} - P) \cdot \mathcal{E}_A^n(\sigma) = 0$ for all n (by induction on n).
2. *Convergence:* By Lemma 8.10.1.2, $\mathcal{E}_A^n(\sigma) \rightarrow \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Hence $(\mathbb{1} - P) \cdot \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho = 0$.
3. *Nonzero scalar:* Since $P \neq 0$ and ρ is positive definite, $\text{tr}(P\rho P) > 0$, so $\frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \neq 0$. Therefore $(\mathbb{1} - P)\rho = 0$.
4. *Contradiction:* Since ρ is positive definite, it is a unit in $M_D(\mathbb{C})$. From $(\mathbb{1} - P)\rho = 0$ we get $\mathbb{1} - P = 0$, i.e. $P = \mathbb{1}$, contradicting $P \neq \mathbb{1}$.

□

8.10.3 From irreducibility to full spanning via Burnside's theorem

The irreducibility condition (Definition 9.2.1.1) is formulated in terms of invariant *projections*. For Burnside-style arguments it is more natural to work with invariant *subspaces* of $\mathbb{C}^D$ under the Kraus operators.

Definition 8.10.3.1 (Algebra span). [Lean](#) [Stmnt](#) For an MPS tensor A , let $\text{alg}(A)$ be the unital $\mathbb{C}$ -subalgebra of $M_D(\mathbb{C})$ generated by the matrices $\{A^i\}_{i=0}^{d-1}$:

$$\text{alg}(A) = \mathbb{C}\langle A^i : i = 0, \dots, d-1 \rangle \subseteq M_D(\mathbb{C}).$$

Definition 8.10.3.2 (Invariant submodule). [Lean](#) [Stmnt](#) A subspace $W \leq \mathbb{C}^D$ is *A-invariant* if $A^iW \subseteq W$ for every i .

Definition 8.10.3.3 (Irreducible action). [Lean](#) [Stmnt](#) The Kraus operators of A act *irreducibly* on $\mathbb{C}^D$ if every A -invariant subspace is trivial, i.e. equal to 0 or to $\mathbb{C}^D$.

Theorem 8.10.3.4 (Irreducible tensor implies irreducible action). [Lean](#) [Stmnt](#) If A is irreducible as a tensor (Definition 9.2.1.1), then the Kraus operators act irreducibly on $\mathbb{C}^D$ (Definition 8.10.3.3).

Proof. [Lean](#) [Proof](#) Let $W \leq \mathbb{C}^D$ be an A -invariant subspace. In finite dimension the orthogonal projection P onto W exists. Invariance of W implies $(\mathbb{1} - P)A^iP = 0$ for all i . Thus P is a nontrivial invariant projection unless $W = 0$ or $W = \mathbb{C}^D$, contradicting irreducibility of the tensor. □

Theorem 8.10.3.5 (Burnside’s theorem for Kraus algebras). [Lean] ✓ Stmt Assume $D > 0$. If A acts irreducibly on $\mathbb{C}^D$, then the generated unital subalgebra is the full matrix algebra:

$$\text{alg}(A) = M_D(\mathbb{C}).$$

This is the complex finite-dimensional case of Burnside’s theorem (equivalently, Jacobson’s density theorem); see [Jac09].

Proof. ✓ Proof Transport $\text{alg}(A)$ along the algebra equivalence $M_D(\mathbb{C}) \simeq_{\mathbb{C}} \text{End}_{\mathbb{C}}(\mathbb{C}^D)$. The irreducible-action hypothesis makes $\mathbb{C}^D$ a simple module for this algebra. Over the algebraically closed field $\mathbb{C}$, Schur’s lemma identifies the endomorphisms commuting with the action with scalars, and Jacobson density then forces the action map to be surjective onto all of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$. □

Remark 8.10.3.6. ✓ Stmt This Burnside/Jacobson-density step is a substantial external algebraic ingredient in the chapter. We use it here as the finite-dimensional complex Burnside theorem, namely the statement that an irreducible matrix algebra action already generates the full endomorphism algebra.

8.10.4 From primitivity to strong irreducibility and normality

The results in this subsection remove the aperiodicity assumption from the normality conclusions. They connect primitivity in Definition 8.9.1, formulated via a complementary transfer-map gap, directly to normality (eventually full Kraus rank), using the convergence of the transfer-map iterates to the projection onto the fixed-point subspace.

Lemma 8.10.4.1 (Primitive tensors with positive-definite fixed point have irreducible transfer map). [Lean] ✓ Stmt If A is a primitive MPS tensor with positive-definite fixed point ρ , then the transfer map $\mathcal{E}_A$ is irreducible.

The proof uses fixed-point uniqueness (Lemma 8.10.1.1): every PSD fixed point is proportional to ρ , so Wolf’s criterion for irreducibility (positive-definite fixed point implies irreducibility) applies directly.

Proof. ✓ Proof By Lemma 8.10.1.1, if $\sigma \geq 0$ and $\mathcal{E}_A(\sigma) = \sigma$, then $\sigma = \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Since $\rho > 0$, Wolf’s uniqueness criterion for irreducibility applies. □

Lemma 8.10.4.2 (Primitive tensors with positive-definite fixed point are strongly irreducible). [Lean] ✓ Stmt If A is a primitive MPS tensor with positive-definite fixed point ρ , then A is strongly irreducible in the sense of [SPGWC10, Proposition 3(c)]: $\mathcal{E}_A$ is irreducible, the fixed point ρ is positive definite, and the peripheral spectrum of $\mathcal{E}_A$ is $\{1\}$.

Proof. ✓ Proof Combine Lemma 8.10.4.1 (irreducibility) with the peripheral-spectrum primitivity from the transfer-map gap hypothesis (Theorem 8.9.2). □

Theorem 8.10.4.3 (Primitive MPS with positive-definite fixed point is normal). [Lean] ✓ Stmt If A is a primitive MPS tensor with positive-definite fixed point ρ , then A is normal (has eventually full Kraus rank). No aperiodicity assumption is needed.

This is the key connection between primitivity in Definition 8.9.1, formulated via a complementary transfer-map gap, and normality; it removes the need for the aperiodicity-based argument.

Proof. ✓ Proof By Lemma 8.10.4.2, A is strongly irreducible, i.e. primitive in Wolf’s sense. Under normalization, Theorem 7.12.3 identifies this primitivity condition directly with normality (equivalently, eventual full Kraus rank). □

8.11 Cumulative span to word span

Cumulative spanning ($T_N(A) = M_D(\mathbb{C})$, the union of word spans of all lengths $\leq N$) is weaker than normality ($S_L(A) = M_D(\mathbb{C})$ for a single length L). An aperiodicity hypothesis upgrades cumulative spanning to a single-length word span.

Remark 8.11.1 (Aperiodicity condition). ✓ Stmt We say A satisfies the *aperiodicity condition* if $\mathbb{1} \in S_1(A)$, i.e. the identity matrix lies in the span of the Kraus operators $\{A^i\}_{i=0}^{d-1}$. This is an additional hypothesis. The one-sided TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ does *not* imply $\mathbb{1} \in S_1(A)$; it only gives a quadratic identity among the Kraus operators. The aperiodicity condition is kept explicit in this section for generality. For the primitive condition with positive-definite fixed point (peripheral-spectrum primitivity together with a positive-definite fixed point), normality follows directly from Wolf's primitivity equivalence (Theorem 7.12.3) without passing through aperiodicity.

Remark 8.11.2 (Counterexample: cumulative spanning does not imply normality). ✓ Stmt In $M_2(\mathbb{C})$, let $A^0 = E_{12}$ and $A^1 = E_{21}$ (the off-diagonal matrix units). Then $\text{alg}(A) = M_2(\mathbb{C})$ and $T_2(A) = M_2(\mathbb{C})$, but the word spans alternate: $S_n(A)$ contains only off-diagonal matrices for odd n and only diagonal matrices for even n . Hence $S_n(A) \neq M_2(\mathbb{C})$ for every n , and A is not normal. The aperiodicity condition $\mathbb{1} \in S_1(A)$ fails: $S_1(A) = \text{span}_{\mathbb{C}}\{E_{12}, E_{21}\}$.

Theorem 8.11.3 (Word span monotonicity from aperiodicity). Lean ✓ Stmt If $\mathbb{1} \in S_1(A)$, then $S_n(A) \leq S_{n+1}(A)$ for all n .

Proof. ✓ Proof For any $M \in S_n(A)$, $M = M \cdot \mathbb{1} \in S_n(A) \cdot S_1(A) \subseteq S_{n+1}(A)$ by Lemma 8.7.2. □

Theorem 8.11.4 (Homogeneous word-span propagation under normalization). Lean Lean

Lean ✓ Stmt Suppose

$$\sum_a A_a A_a^\dagger = \mathbb{1} \quad \text{and} \quad S_L(A) = M_D(\mathbb{C}).$$

Then, for every $m \geq L$,

$$S_m(A) = M_D(\mathbb{C}).$$

Equivalently, L -block injectivity propagates to every larger homogeneous length.

Proof. ✓ Proof If $S_n(A) = M_D(\mathbb{C})$, then for any $X \in M_D(\mathbb{C})$,

$$X = \left(\sum_a A_a A_a^\dagger \right) X = \sum_a A_a (A_a^\dagger X).$$

Since $A_a^\dagger X \in S_n(A)$, each summand lies in $S_1(A)S_n(A) \subseteq S_{n+1}(A)$. Thus $S_{n+1}(A) = M_D(\mathbb{C})$. Iterating this one-step implication gives the result for all $m \geq L$. □

Theorem 8.11.5 (Cumulative span equals word span under aperiodicity). Lean ✓ Stmt If $\mathbb{1} \in S_1(A)$, then $T_n(A) = S_n(A)$ for all n .

Proof. ✓ Proof By Theorem 8.11.3, the word spans are monotone: $S_0 \leq S_1 \leq \dots$. The supremum $T_n = S_0 + \dots + S_n$ therefore equals the largest term S_n . □

Theorem 8.11.6 (Cumulative spanning plus aperiodicity implies normality). Lean ✓ Stmt If $T_N(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.

Proof. ✓ Proof By Theorem 8.11.5, $S_N(A) = T_N(A) = M_D(\mathbb{C})$. □

Theorem 8.11.7 (Algebra span plus aperiodicity implies normality). [Lean](#) [✓ Stmt](#) *If $\text{alg}(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. [✓ Proof](#) Since $\text{alg}(A) = M_D(\mathbb{C})$, there exists N with $T_N(A) = M_D(\mathbb{C})$ (by the ascending chain condition on finite-dimensional subspaces). Apply Theorem 8.11.6. $\square$

Remark 8.11.8 (Normal canonical form avoids the aperiodicity step). The aperiodicity results in Section 8.11 are needed only for the block-injective argument through Definition 2.4.4, where one upgrades cumulative spanning to eventual block injectivity. The normal-canonical theory used later in Chapters 9–10 does not pass through this step: it works directly with irreducible TP-normalized blocks whose blocked transfer maps are primitive and, once the corresponding weighted block data are available, places them in normal canonical form in the sense of [CPGSV16]. There are analogous cumulative-span lemmas for blocked irreducible tensors and eigenvector-spreading arguments, but they are not needed for the implication stated below.

8.12 Block injectivity from TP/primitive/irreducible blocks

Theorem 8.12.1 (TP-primitive irreducible tensor is injective after blocking). [Lean](#) [✓ Stmt](#) *Let A be a left-canonical MPS tensor with $D > 0$, irreducible tensor, and primitive transfer map. Then some positive blocking $A^{[L]}$ is one-site injective.*

Proof. [✓ Proof](#) If $D = 1$, trace preservation gives a nonzero one-site Kraus matrix, and every nonzero scalar matrix spans $M_1(\mathbb{C})$; hence A is one-site injective, so Lemma 2.4.3 gives the positive witness $L = 1$. Suppose now that $D \geq 2$. The normality theorem (Theorem 9.6.2.1) gives a block-injectivity witness L . If $L = 0$, Corollary 8.1.10 would force $D = 1$, a contradiction. Thus $L > 0$, and the blocked-chain equivalence identifies this L -block injectivity with one-site injectivity of the blocked tensor $A^{[L]}$. $\square$

Remark 8.12.2. Blocking preserves normality by Theorem 8.8.1.

Chapter 9

Canonical Form Reduction

This is the first of three chapters that together prove the Fundamental Theorem of matrix product states. Here an arbitrary translation-invariant MPS tensor is reduced to a weighted family of primitive blocks; Chapter 10 constructs the basis of normal tensors carried by those blocks and compares their coefficients, and Chapter 11 proves the theorem itself. The reduction follows [PGVWC07], [CPGSV16], [CPGSV21], and [Wol12, Chapter 6]. Its goal is to expose the primitive blocks inside the tensor. The transfer operator of an arbitrary tensor can be reducible and can have a period greater than one; the reduction splits it along its invariant subspaces and brings the period of each block down to one, so that each block is primitive, with peripheral spectrum $\{1\}$. The blocks are returned in a fixed gauge, and the sections below carry out the reduction. The periodic decomposition theorem of [PGVWC07, Theorem 5] also gives an explicit decomposition of the state vector into p -periodic summands; here we use only its blocked cyclic-sector consequence (Theorem 9.6.1.1), and that explicit state-vector decomposition lies outside this chapter.

Two normalizations appear. The unital orientation $\sum_i A_k^i (A_k^i)^\dagger = \mathbb{1}$ is used in [PGVWC07, Theorem 4]; the trace-preserving (left-canonical) orientation $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$ is used in the TP-gauge and after-blocking reductions below. They exchange under the conjugate-transposed Kraus family $K_i := (A^i)^\dagger$, whose transfer map is the Frobenius adjoint of $\mathcal{E}_A$.

Theorem 9.1 (Translation-invariant canonical form). Lean Stmt *Let A be a translation-invariant MPS representation of bond dimension D . There is a positive scalar s and a finite family of blocks A_k , possibly empty, such that the globally rescaled tensor $s^{-1}A$ has the same MPV coefficients on every positive-length ring as*

$$\bigoplus_k \mu_k A_k^i.$$

The weights are positive and satisfy $1 \geq |\mu_k|$; if the family is nonempty, then some weight has norm 1. For each block,

$$\sum_i A_k^i (A_k^i)^\dagger = \mathbb{1}, \quad \sum_i (A_k^i)^\dagger \Lambda_k A_k^i = \Lambda_k,$$

where Λ_k is diagonal, positive, and full-rank. Each block has positive bond dimension. Moreover the only fixed points of $\mathcal{E}_{A_k}(X) = \sum_i A_k^i X (A_k^i)^\dagger$ are the scalar multiples of $\mathbb{1}$: for every X ,

$$\mathcal{E}_{A_k}(X) = X \implies \exists c \in \mathbb{C}, X = c \mathbb{1}.$$

The total bond dimension of the decomposition is at most D .

This is the positive-length form of [PGVWC07, Theorem 4], with the global spectral-radius normalization step made explicit.

Remark 9.2 (Source proof order). The proof follows [PGVWC07, Theorem 4] in order: normalize the spectral radius and use a full-rank positive fixed point to obtain the unital gauge; derive the invariant support projection from a singular positive fixed point by the positivity contradiction; split the finite-ring trace into the supported and orthogonal summands; iterate, using a non-scalar fixed point to split further until each block has only scalar fixed points; and finally repeat the fixed-point argument for the dual map and diagonalize the resulting full-rank positive fixed point by a unitary gauge. The primitive-block results below are used only after their hypotheses have been obtained from this argument.

Remark 9.3 (CPSV blocked canonical form (source statement)). The blocked canonical-form reduction of [CPGSV16, Section 2.3] and [CPGSV21, Section IV.A] states that an arbitrary translation-invariant MPS tensor becomes, after blocking by some period $p \geq 1$, the direct sum of an all-zero summand and a weighted family of normal tensors $\bigoplus_k \mu_k B_k$ with $|\mu_k| \leq 1$ and at least one $|\mu_k| = 1$. The closest proved results here are Theorem 9.6.2.2 and Theorem 10.4.2.9; the source upper normalization $|\mu_k| \leq 1$ with a unit-modulus witness is the one remaining input, recorded in Remark 10.4.2.10.

9.1 Normal tensors and canonical forms

This section records the CPSV definitions of *normal tensor* and *canonical form* from [CPGSV16]. The basis of normal tensors is stated in Chapter 10. Write

$$\sigma_{\partial}(\mathcal{E}) = \{\lambda \in \sigma(\mathcal{E}) : |\lambda| = r(\mathcal{E})\}$$

for the peripheral spectrum of a transfer map $\mathcal{E}$, where $r(\mathcal{E})$ is its spectral radius. The stronger predicates used later (Definition 9.3.3 and the canonical-form predicate of Chapter 10) add left-canonical normalization and spectral primitivity. The weight moduli are taken non-increasing, not strictly decreasing; equal-modulus and repeated-copy sectors are retained and compared by the coefficient argument of Chapter 10.

Definition 9.1.1 (CPSV normal tensor (NT)). Lean ✓ Stmt A tensor A of physical dimension d and bond dimension D is a *normal tensor* if, after the spectral-radius normalization of [CPGSV16], (i) A admits no nontrivial invariant orthogonal projection and (ii) the associated CPM $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ has spectral radius $r(\mathcal{E}_A) = 1$ and peripheral spectrum $\sigma_{\partial}(\mathcal{E}_A) = \{1\}$.

Remark 9.1.2. Condition (i) cannot be dropped. A reducible tensor may have peripheral spectrum $\{1\}$ while still admitting a nontrivial invariant projection, so condition (ii) alone does not characterize normality.

The CPSV *canonical form* (CF) of a tensor A is the normal-block decomposition

$$A^i \cong \bigoplus_{k=1}^r \mu_k A_k^i$$

with weights normalized by $|\mu_k| \leq 1$ and with at least one $|\mu_k| = 1$; see [CPGSV16, Section 2.3]. In the statements below the structural part is supplied by the stronger normal canonical form of Definition 9.3.3, and the global weight normalization is invoked as a separate source hypothesis where needed.

Remark 9.1.3 (Relation to the strong predicates). A tensor A which is irreducible (Definition 9.2.1.1), has transfer map of spectral radius one, and has primitive transfer map is normal in the sense of Definition 9.1.1. The reverse implications from the stronger predicates of Definition 9.3.3 and of Chapter 10 require the fundamental-theorem step or the algebraic-to-spectral normality equivalence, established later.

9.1.1 Normalization conventions

Remark 9.1.1.1 (Scalar normalization conventions). The reductions use four scalar facts: scaling preserves injectivity; the transfer map under a scalar ζ rescales by $|\zeta|^2$; unit-modulus phase rescaling preserves $\sum_i (A^i)^\dagger A^i = \mathbb{1}$; and the MPV of a weighted block-diagonal tensor factorizes as $\mu_k^N V^{(N)}(A_k)_\sigma = \eta_k^N |\mu_k|^N V^{(N)}(A_k)_\sigma$ with $\eta_k := \mu_k / |\mu_k|$.

Left-canonical normalization is required before the blocks can be compared. Without it the implication $\mathcal{V}(\bigoplus_k \mu_k A_k) = \mathcal{V}(\bigoplus_k \mu_k B_k) \Rightarrow \forall k, \mathcal{V}(A_k) = \mathcal{V}(B_k)$ already fails for two scalar blocks: for $\mu = (1, 2)$, $A = (1, 3/2)$, $B = (3, 1/2)$ the two direct sums generate identical MPV families but the first blocks differ at $N = 1$. Left-canonical normalization removes this particular ambiguity; the sectors are then matched, up to a permutation, by the exact fixed-length linear independence of Chapter 10, which requires no ordering of the weight moduli.

9.2 Invariant-subspace decomposition and irreducible blocks

The block decomposition rests on one principle: a positive semidefinite fixed point of $\mathcal{E}_A$ that is not full rank has a proper invariant support, which block-diagonalizes the Kraus operators without changing the MPV family. Iterating the resulting splitting decomposes any tensor into irreducible blocks.

Theorem 9.2.1 (Unitary conjugation preserves the MPV family). Lean ✓ Stmt *If U is a unitary matrix, then A and $U^\dagger A U$ generate the same MPV family.*

Proof. ✓ Proof By cyclicity of the trace, $\text{tr}((U^\dagger A^{i_1} U) \dots (U^\dagger A^{i_N} U)) = \text{tr}(A^{i_1} \dots A^{i_N})$. □

Theorem 9.2.2 (Invariant projection gives a two-block decomposition). Lean ✓ Stmt *Suppose P is an orthogonal projection satisfying, for every physical index i ,*

$$(\mathbb{1} - P)A^i P = 0.$$

Then there exist n, m with $n + m = D$ and MPS tensors A_1, A_2 of bond dimensions n and m such that A and the two-block tensor $A_1 \oplus A_2$ generate the same MPV family in the sense of Definition 2.3.3.

Proof. ✓ Proof Diagonalize P by a unitary U . Since $P^2 = P$, the diagonal form of P has only 0- and 1-eigenvalues, so the basis splits as $n + m = D$. Conjugating A by U preserves the MPV family by Theorem 9.2.1. In this basis the relation $(\mathbb{1} - P)A^i P = 0$ makes each letter upper block triangular. Write

$$B^i = \begin{pmatrix} A_1^i & * \\ 0 & A_2^i \end{pmatrix}$$

for the conjugated letters in this split basis. Since $B^{i_1} \dots B^{i_N}$ is again upper block triangular, its trace satisfies

$$\text{tr}(B^{i_1} \dots B^{i_N}) = \text{tr}(A_1^{i_1} \dots A_1^{i_N}) + \text{tr}(A_2^{i_1} \dots A_2^{i_N}).$$

Hence the block-diagonal tensor $A_1 \oplus A_2$ has the same MPV coefficients as the conjugated tensor. □

Theorem 9.2.3 (Nontrivial invariant projection gives strict two-block reduction). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 9.2.2, assume in addition that $P \neq 0$ and $P \neq \mathbb{1}$. Then there exist n, m with $n + m = D$, $n < D$, and $m < D$, together with block tensors A_1, A_2 , such that A and $A_1 \oplus A_2$ generate the same MPV family in the sense of Definition 2.3.3.

Proof. [✓ Proof](#) Apply Theorem 9.2.2. In the diagonal form of P , the 1-eigenspace is nonempty because $P \neq 0$, and the 0-eigenspace is nonempty because $P \neq \mathbb{1}$. Hence both block sizes are positive. Since $n + m = D$, this implies $n < D$ and $m < D$. $\square$

Definition 9.2.4 (Support projection). [Lean](#) [✓ Stmt](#) For a positive semidefinite matrix $\rho \geq 0$, the *support projection* P is the orthogonal projection onto the range of ρ . Via the spectral decomposition $\rho = U \text{diag}(\lambda_1, \dots, \lambda_D) U^\dagger$,

$$P = U \text{diag}(\mathbf{1}_{\lambda_1 > 0}, \dots, \mathbf{1}_{\lambda_D > 0}) U^\dagger,$$

where $\mathbf{1}_{\lambda_j > 0}$ is 1 if $\lambda_j > 0$ and 0 otherwise.

Definition 9.2.5 (Invariant projection predicate). [Lean](#) [✓ Stmt](#) An MPS tensor A has an *invariant projection* if there exists an orthogonal projection P with $P \neq 0$, $P \neq \mathbb{1}$, and $(\mathbb{1} - P)A^i P = 0$ for all i . A tensor is *irreducible* if it has no nontrivial invariant projection (Definition 9.2.1.1).

Definition 9.2.6 (Invariant-projector closure). [Lean](#) [✓ Stmt](#) An MPS tensor A has *invariant-projector closure* if every orthogonal projection P satisfying $PA^i = PA^i P$ for all i also satisfies $A^i P = PA^i P$ for all i . This is the projector hypothesis in the sufficient condition for canonical form of [CPGSV16, Section 2].

Theorem 9.2.7 (Invariant projections are reducing under projector closure). [Lean](#) [✓ Stmt](#) Suppose that A has invariant-projector closure and that P is an orthogonal projection satisfying $(\mathbb{1} - P)A^i P = 0$ for all i . Then P reduces every tensor matrix: $PA^i = A^i P$ for all i .

Proof. [✓ Proof](#) The lower-corner equation gives

$$(\mathbb{1} - P)A^i = (\mathbb{1} - P)A^i(\mathbb{1} - P).$$

Applying projector closure to $\mathbb{1} - P$ gives $A^i(\mathbb{1} - P) = (\mathbb{1} - P)A^i(\mathbb{1} - P)$, hence the upper corner $PA^i(\mathbb{1} - P)$ also vanishes. Therefore

$$PA^i = PA^i P = A^i P.$$

$\square$

Theorem 9.2.8 (Invariant-projector closure gives a reducing projection). [Lean](#) [✓ Stmt](#) If A has invariant-projector closure and admits a nontrivial invariant projection, then it admits a nontrivial orthogonal projection commuting with every A^i .

Proof. [✓ Proof](#) Apply Theorem 9.2.7 to the projection supplied by Definition 9.2.5. $\square$

Lemma 9.2.9 (Support isometry of an orthogonal projection). [Lean](#) [✓ Stmt](#) Every orthogonal projection P on $\mathbb{C}^D$ factors through an isometry: there are $n \in \mathbb{N}$ and $V \in \mathbb{C}^{D \times n}$ with $V^\dagger V = \mathbb{1}$, $VV^\dagger = P$, and $n = \text{tr}(P)$.

Proof. [✓ Proof](#) Since P is Hermitian and idempotent, its eigenvalues are 0 and 1, and unitary diagonalization gives

$$P = U \text{diag}(\underbrace{1, \dots, 1}_n, 0, \dots, 0) U^\dagger$$

with U unitary and $n = \text{tr}(P)$. The matrix $V \in \mathbb{C}^{D \times n}$ formed by the first n columns of U satisfies $V^\dagger V = \mathbb{1}$ and $VV^\dagger = P$. $\square$

Lemma 9.2.10 (Corner tensors inherit projector closure). [Lean](#) [✓ Stmt](#) Let A have invariant-projector closure, and let V be an isometry, $V^\dagger V = \mathbb{1}$, whose range projection $VV^\dagger$ commutes with every A^i . Then the corner tensor $B^i = V^\dagger A^i V$ again has invariant-projector closure.

Proof. [✓ Proof](#) Let Q be an orthogonal projection with $QB^i = QB^i Q$ for all i . Then $VQV^\dagger$ is an orthogonal projection. Commutation of $VV^\dagger$ with each A^i gives $V^\dagger A^i = B^i V^\dagger$, so left-invariance of Q yields

$$(VQV^\dagger)A^i = VQB^iV^\dagger = VQB^iQV^\dagger = (VQV^\dagger)A^i(VQV^\dagger).$$

Projector closure for A yields $A^i(VQV^\dagger) = (VQV^\dagger)A^i(VQV^\dagger)$, and compressing both sides between $V^\dagger$ and V returns $B^iQ = QB^iQ$. $\square$

Lemma 9.2.11 (Matrix product vectors split along an isometric partition). [Lean](#) [✓ Stmt](#) Let $V_1, \dots, V_r$ be isometries, $V_k^\dagger V_k = \mathbb{1}$, with $\sum_k V_k V_k^\dagger = \mathbb{1}$, and suppose $A^i V_k = V_k B_k^i$ for some tensors $B_1, \dots, B_r$. Then $\mathcal{V}(A) = \mathcal{V}(\bigoplus_{k=1}^r B_k)$.

Proof. [✓ Proof](#) The intertwining relation propagates from letters to words, so $A^w V_k = V_k B_k^w$ for every word w . Inserting the partition of unity $\sum_k V_k V_k^\dagger = \mathbb{1}$ into the trace and cycling gives

$$\mathrm{tr}(A^w) = \sum_{k=1}^r \mathrm{tr}(V_k^\dagger A^w V_k) = \sum_{k=1}^r \mathrm{tr}(B_k^w),$$

which is the matrix product vector of the direct sum. $\square$

Theorem 9.2.12 (Decomposition into irreducible corners under projector closure). [Lean](#) [✓ Stmt](#) Suppose A has invariant-projector closure. Then there are isometries $V_1, \dots, V_r$ with $V_k^\dagger V_k = \mathbb{1}$, pairwise orthogonal ranges, and $\sum_k V_k V_k^\dagger = \mathbb{1}$, such that $A^i V_k = V_k A_k^i$ for the corner tensors $A_k^i = V_k^\dagger A^i V_k$, every corner tensor A_k is irreducible, and $\mathcal{V}(A) = \mathcal{V}(\bigoplus_{k=1}^r A_k)$.

Proof. [✓ Proof](#) Strong induction on the bond dimension. If A is irreducible, the single corner $V = \mathbb{1}$ suffices. Otherwise A has a nontrivial invariant projection P , which commutes with every A^i by Theorem 9.2.7. Factoring $P = V_1 V_1^\dagger$ and $\mathbb{1} - P = V_2 V_2^\dagger$ through support isometries (Lemma 9.2.9) splits A into two corner tensors of strictly smaller bond dimension: for $j = 1, 2$, commutation gives

$$A^i V_j = V_j (V_j^\dagger A^i V_j),$$

and each corner inherits invariant-projector closure by Lemma 9.2.10. The two recursively obtained families are combined; ranges from different corners are orthogonal since $V_1^\dagger P = V_1^\dagger$ and $(\mathbb{1} - P)V_2 = V_2$ give

$$V_1^\dagger V_2 = V_1^\dagger P(\mathbb{1} - P)V_2 = 0.$$

The equality of matrix product vectors is Lemma 9.2.11. $\square$

Definition 9.2.13 (No nontrivial periodic vectors). [Lean](#) [✓ Stmt](#) An MPS tensor A has *no nontrivial p -periodic vectors* if, for every isometry V with $V^\dagger V = \mathbb{1}$ and $A^i V = V B^i$ for all i with B irreducible (equivalently, for every restriction B of A to a minimal invariant subspace), and for every $\rho > 0$ and $r > 0$ with $\mathcal{E}_B(\rho) = r\rho$, every eigenvalue of $\mathcal{E}_B$ of modulus r equals r . The eigenvalue r is the spectral radius of $\mathcal{E}_B$ (Theorem 6.9.1), so the condition states that no block of the invariant-subspace decomposition of [CPGSV16, Section 2], rescaled to spectral radius one, has a peripheral eigenvalue $e^{2\pi i q/p} \neq 1$. Eigenvectors with such eigenvalues are the p -periodic vectors of [CPGSV16, Section 2].

Lemma 9.2.14 (Transfer map of an intertwined corner). [Lean](#) [Lean](#) [✓ Stmt](#) Let V satisfy $A^i V = V B^i$ for all i . Then for every X ,

$$\mathcal{E}_A(VXV^\dagger) = V\mathcal{E}_B(X)V^\dagger.$$

If moreover $V^\dagger V = \mathbb{1}$, then every eigenvalue of $\mathcal{E}_B$ is an eigenvalue of $\mathcal{E}_A$.

Proof. [✓ Proof](#) Expanding $\mathcal{E}_A(VXV^\dagger)$ termwise, the intertwining $A^i V = V B^i$ and its adjoint $V^\dagger (A^i)^\dagger = (B^i)^\dagger V^\dagger$ move both factors past V and $V^\dagger$. For the eigenvalue statement, if $\mathcal{E}_B(X) = \mu X$ with $X \neq 0$, then $\mathcal{E}_A(VXV^\dagger) = \mu VXV^\dagger$, and $VXV^\dagger \neq 0$ because compressing with the isometry returns $V^\dagger(VXV^\dagger)V = X$. $\square$

Lemma 9.2.15 (Perron eigenpair of a nonzero irreducible tensor). [Lean](#) [✓ Stmt](#) Let B be an irreducible MPS tensor of positive bond dimension, not all of whose matrices vanish. Then there are $\rho > 0$ and $r > 0$ with $\mathcal{E}_B(\rho) = r\rho$.

Proof. [✓ Proof](#) Irreducibility of the tensor makes the transfer map an irreducible completely positive map, and a nonzero matrix makes it nonzero, so it does not annihilate any nonzero PSD matrix. The Perron–Frobenius theorem for positive maps (Theorem 6.6.1) gives a nonzero PSD eigenvector with eigenvalue $r > 0$, and irreducibility upgrades it to a positive-definite one ([Wol12, Theorem 6.3(2)]). $\square$

Lemma 9.2.16 (Spectral normalization of an irreducible block). [Lean](#) [✓ Stmt](#) Let B be an irreducible MPS tensor of positive bond dimension with $\mathcal{E}_B(\rho) = r\rho$ for some $\rho > 0$ and $r > 0$, and suppose every eigenvalue of $\mathcal{E}_B$ of modulus r equals r . Then $r^{-1/2}B$ is a normal tensor.

Proof. [✓ Proof](#) The rescaling identity

$$\mathcal{E}_{r^{-1/2}B}(X) = r^{-1} \mathcal{E}_B(X)$$

gives $\mathcal{E}_{r^{-1/2}B}(\rho) = r^{-1} \cdot r\rho = \rho$, so ρ is a positive-definite fixed point of $\mathcal{E}_{r^{-1/2}B}$. The rescaled map remains irreducible and completely positive, so Theorem 6.9.1 identifies its spectral radius with the eigenvalue one. The eigenvalues of modulus one of $\mathcal{E}_{r^{-1/2}B}$ are the eigenvalues of $\mathcal{E}_B$ of modulus r , divided by r ; by hypothesis the only such eigenvalue is one, so the peripheral eigenvalue set is $\{1\}$. Irreducibility is unchanged by a nonzero rescaling. $\square$

Theorem 9.2.17 (Sufficient condition for canonical form). [Lean](#) [✓ Stmt](#) Suppose A has invariant-projector closure and no nontrivial p -periodic vectors. Then there are normal tensors $A_1, \dots, A_r$ of positive bond dimensions $D_1, \dots, D_r$ with $\sum_k D_k \leq D$, and nonzero weights $\mu_1, \dots, \mu_r$, such that

$$\mathcal{V}^+(A) = \mathcal{V}^+\left(\bigoplus_{k=1}^r \mu_k A_k\right).$$

This is the sufficient condition for canonical form of [CPGSV16, Section 2], with that paper’s convention $\sum_k D_k \leq D$ allowing zero blocks: a corner of A carrying the zero tensor cannot be rescaled to spectral radius one, and contributes its bond dimension only to the length-zero coefficient.

Proof. [✓ Proof](#) Decompose A into irreducible corners B_k along isometries V_k (Theorem 9.2.12); every matrix product vector coefficient of A is the sum of those of the corners. A corner whose matrices all vanish contributes nothing at positive length and is dropped. Every remaining corner has a Perron eigenpair $\mathcal{E}_{B_k}(\rho_k) = r_k \rho_k$ (Lemma 9.2.15). The absence of p -periodic vectors, applied to the corner B_k , gives that every eigenvalue of $\mathcal{E}_{B_k}$ of modulus r_k equals r_k , so

$A_k := r_k^{-1/2} B_k$ is normal (Lemma 9.2.16). Setting $\mu_k := r_k^{1/2}$ gives $\mu_k A_k = B_k$ at the level of matrices, so for every word w of positive length N the weight cancels,

$$\mu_k^N \operatorname{tr}(A_k^w) = (r_k^{1/2})^N (r_k^{-1/2})^N \operatorname{tr}(B_k^w) = \operatorname{tr}(B_k^w),$$

and summing over k identifies the positive-length coefficients of $\bigoplus_k \mu_k A_k$ with those of A . The kept dimensions satisfy $\sum_k D_k \leq \sum_k \operatorname{tr}(V_k V_k^\dagger) = \operatorname{tr}(\mathbb{1}) = D$. $\square$

Theorem 9.2.18 (Canonical form at every length without zero corners). [Lean] [✓ Stmt] *Suppose A has invariant-projector closure, has no nontrivial p -periodic vectors, and every restriction of A to a minimal invariant subspace of positive dimension has a nonzero matrix. Then there are normal tensors $A_1, \dots, A_r$ and nonzero weights $\mu_1, \dots, \mu_r$ with*

$$\mathcal{V}(A) = \mathcal{V}\left(\bigoplus_{k=1}^r \mu_k A_k\right).$$

The no-zero-corner hypothesis is not part of the source statement; it excludes exactly the zero blocks of [CPGSV16, Section 2], which change only the length-zero coefficient.

Proof. [✓ Proof] As for Theorem 9.2.17, with no corner dropped: every corner is nonzero by hypothesis, so $\bigoplus_k \mu_k A_k$ has the same coefficients as the unit-weight direct sum of the corners at every length, including zero. $\square$

Theorem 9.2.19 (PSD fixed point gives invariant projection). [Lean] [✓ Stmt] *Let $\rho \geq 0$ be a fixed point of $\mathcal{E}_A$, and let P be its support projection. Then $(\mathbb{1} - P)A^i P = 0$ for all i : the range of P is invariant under every A^i .*

Proof. [✓ Proof] From $\mathcal{E}_A(\rho) = \rho$ and positivity of each summand $A^i \rho (A^i)^\dagger$, one gets $(\mathbb{1} - P)A^i \rho (A^i)^\dagger (\mathbb{1} - P) = 0$ for each i . Since ρ is supported on the range of P , this forces $(\mathbb{1} - P)A^i P = 0$.

This is the finite-dimensional case of [Wol12, Proposition 6.10]: the support projection of any stationary state is sub-harmonic. The equivalence between irreducibility and absence of nontrivial invariant projections is [Wol12, Theorem 6.2]; see also [CPGSV21, Section IV.A.1]. $\square$

Theorem 9.2.20 (Two-block decomposition). [Lean] [✓ Stmt] *If $\mathcal{E}_A$ has a nonzero PSD fixed point ρ that is not positive definite, then there exist MPS tensors A_1, A_2 of smaller bond dimensions with $\mathcal{V}(A) = \mathcal{V}(A_1 \oplus A_2)$.*

Proof. [✓ Proof] The support projection P of ρ is neither 0 nor $\mathbb{1}$ (since $\rho \neq 0$ and ρ is not positive definite). Theorem 9.2.19 gives $(\mathbb{1} - P)A^i P = 0$, so a unitary change of basis block-diagonalizing P puts A^i in upper-triangular block form with diagonal blocks B_{11}^i, B_{22}^i . Theorem 9.2.1 preserves the MPV under the conjugation, and the trace of an upper-triangular product equals the sum of the diagonal-block traces, so deleting the off-diagonal blocks preserves all MPV coefficients. $\square$

Theorem 9.2.21 (Non-scalar fixed point gives a further split). [Lean] [Lean] [✓ Stmt] *Suppose A is unital, $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. If X is a Hermitian fixed point of $\mathcal{E}_A$ which is not a scalar multiple of $\mathbb{1}$, then there is a nonzero PSD fixed point ρ which is not positive definite, and consequently A admits a two-block MPV-equivalent decomposition with both block dimensions strictly smaller than D . This is the fixed-point shift and further decomposition step in [PGVWC07, Theorem 4].*

Proof. [✓ Proof] Let $\lambda_{\max}$ be the largest eigenvalue of X . Then $\rho := \lambda_{\max} \mathbb{1} - X$ is positive semidefinite, singular, and nonzero (since X is not scalar), and unitality with the fixed-point equation for X gives $\mathcal{E}_A(\rho) = \rho$. Apply Theorem 9.2.20. The non-scalar hypothesis is needed because scalar multiples of $\mathbb{1}$ are fixed by every unital map and yield no nontrivial support projection. $\square$

9.2.1 Iterated reduction to irreducible blocks

Iterating the two-block splitting on bond dimension decomposes any tensor into irreducible blocks; the remaining lemmas record the spectral facts about irreducible unital blocks and the algebra-spanning input used later.

Definition 9.2.1.1 (Irreducible tensor). [Lean](#) [✓ Stmt](#) An MPS tensor A is *irreducible* if it has no nontrivial invariant projection.

Theorem 9.2.1.2 (Irreducible block decomposition). [Lean](#) [✓ Stmt](#) Every MPS tensor A admits a block-diagonal tensor $\bigoplus_{k=1}^r A_k$ with each A_k irreducible and $\mathcal{V}(A) = \mathcal{V}(\bigoplus_k A_k)$.

Proof. [✓ Proof](#) By strong induction on D . If A is irreducible, take $r = 1$. Otherwise a nontrivial invariant projection P gives the upper-triangular splitting of Theorem 9.2.20; deleting the off-diagonal part preserves the MPV family and yields two blocks of strictly smaller bond dimension. Apply the induction hypothesis to each block.

Both [PGVWC07, Theorem 4] and [CPGSV21, Section IV.A.1] obtain the block decomposition by the same invariant-projection splitting. $\square$

Theorem 9.2.1.3 (Scalar fixed points in an irreducible unital block). [Lean](#) [✓ Stmt](#) If A is irreducible and unital, $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, and $\mathcal{E}_A(X) = X$, then X is a scalar multiple of $\mathbb{1}$. This is the final fixed-point assertion in the proof of [PGVWC07, Theorem 4].

Proof. [✓ Proof](#) Tensor irreducibility gives irreducibility of the completely positive transfer map. By Wolf’s Theorem 6.6 for irreducible unital Kraus maps, every fixed point is scalar. $\square$

Theorem 9.2.1.4 (Unital gauge from a full-rank Perron–Frobenius eigenvector). [Lean](#) [Lean](#) [✓ Stmt](#) Let ρ be positive definite with $\mathcal{E}_A(\rho) = r\rho$ for some $r > 0$. For $B^i := r^{-1/2}\rho^{-1/2}A^i\rho^{1/2}$,

$$\sum_i B^i (B^i)^\dagger = \mathbb{1}. \quad (9.1)$$

If $r = 1$, then the unscaled gauge $B^i := \rho^{-1/2}A^i\rho^{1/2}$ is unital and generates the same MPV family as A . This is the spectral-radius normalization and full-rank fixed-point gauge of [PGVWC07, Theorem 4].

Proof. [✓ Proof](#) The positive square root of ρ is invertible. Substituting $B^i = r^{-1/2}\rho^{-1/2}A^i\rho^{1/2}$ into $\sum_i B^i (B^i)^\dagger$ and using $\mathcal{E}_A(\rho) = r\rho$ gives (9.1). For $r = 1$ the change from A to B is a similarity, so cyclicity of the trace gives equality of all MPV coefficients. $\square$

Lemma 9.2.1.5 (Range factorization of the support projection). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The support projection π of a positive semidefinite matrix ρ factors as $\pi = \rho W$ for a spectral pseudo-inverse W , witnessing that the range of π is the range of ρ . For a rectangular matrix Y , the support projection π of $YY^\dagger$ fixes Y , $\pi Y = Y$; and whenever $AY = YG$ for some matrix G — that is, A maps the range of Y into itself — the one-sided invariance $(\mathbb{1} - \pi)A\pi = 0$ holds.

Proof. [✓ Proof](#) In the spectral decomposition $\rho = U \text{diag}(\lambda) U^\dagger$ take $W = U \text{diag}(\lambda^+) U^\dagger$, where λ^+ inverts the positive eigenvalues and vanishes on the kernel; then $\rho W = U \text{diag}(\mathbf{1}_{\lambda>0}) U^\dagger = \pi$. From $\pi(YY^\dagger) = YY^\dagger$ the matrix $((\mathbb{1} - \pi)Y)((\mathbb{1} - \pi)Y)^\dagger$ vanishes, hence $\pi Y = Y$. Finally $(\mathbb{1} - \pi)A\pi = (\mathbb{1} - \pi)AY(Y^\dagger W) = (\mathbb{1} - \pi)Y G(Y^\dagger W) = 0$. $\square$

Theorem 9.2.1.6 (Irreducibility under rescaled conjugation). [Lean](#) [✓ Stmt](#) Let B be an irreducible tensor, let X be invertible, and let $c \neq 0$. Then the rescaled conjugated tensor with letters $cX^{-1}B^vX$ is irreducible. In particular the rescaled unital gauge of an irreducible tensor is irreducible; this is the normalization of [CPGSV16, lines 224–225] applied to an irreducible corner.

Proof. ✓ Proof An invariant orthogonal projection $P \notin \{0, \mathbb{1}\}$ of the conjugated tensor makes the columns of $Y = XP$ span a subspace invariant under every letter of B : $B^v Y = Y G_v$ with $G_v = X^{-1} B^v X P$. The support projection of $Y Y^\dagger$ is then a nontrivial invariant orthogonal projection of B by Lemma 9.2.1.5: it is nonzero because $Y \neq 0$, and it is not the identity because any nonzero vector of the kernel of P transports through $(X^\dagger)^{-1}$ to the kernel of $Y Y^\dagger$. This contradicts irreducibility of B . $\square$

Theorem 9.2.1.7 (Unital Schwarz data for an irreducible corner). Lean ✓ Stmt *Let B be an irreducible tensor with a positive definite transfer eigenvector, $\mathcal{E}_B(\rho) = r\rho$ with $\rho > 0$ and $r > 0$. Then the rescaled unital gauge $K^v = r^{-1/2} \rho^{-1/2} B^v \rho^{1/2}$ is a unital Kraus family, is an irreducible tensor with an irreducible transfer map, and its adjoint map has a positive definite fixed point. These are the hypotheses of the cyclic decomposition of the peripheral spectrum, [Wol12, Theorem 6.6].*

Proof. ✓ Proof Unitality is Theorem 9.2.1.4; irreducibility is Theorem 9.2.1.6. The adjoint of a unital family is trace preserving, so its transfer map is a channel and has a positive semidefinite fixed point by the Cesàro argument; irreducibility of the adjoint map upgrades the fixed point to a positive definite one. $\square$

Lemma 9.2.1.8 (Eigenvalue transport to the unital gauge). Lean ✓ Stmt *In the setting of Theorem 9.2.1.7, every transfer eigenvalue μ of B becomes the transfer eigenvalue μ/r of the rescaled unital gauge: the gauge conjugates the transfer map by the congruence with $\rho^{1/2}$ and divides it by r . In particular the peripheral eigenvalues $e^{2\pi i q/p}$ of [CPGSV16, lines 225–230] arise from the transfer eigenvalues of modulus r .*

Proof. ✓ Proof If $\mathcal{E}_B(Z) = \mu Z$ then $Z' = \rho^{-1/2} Z \rho^{-1/2}$ is nonzero and satisfies $\mathcal{E}_K(Z') = (\mu/r) Z'$ by direct computation. $\square$

Theorem 9.2.1.9 (Unitary diagonal positive fixed point in TP gauge). Lean ✓ Stmt *Let A be an irreducible MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $D > 0$. Then there exist a unitary U and a diagonal positive definite Λ such that, for $B^i := U^\dagger A^i U$,*

$$\sum_i (B^i)^\dagger B^i = \mathbb{1}, \quad \mathcal{E}_B(\Lambda) = \Lambda.$$

Proof. ✓ Proof The TP hypothesis makes $\mathcal{E}_A$ a channel with a nonzero PSD fixed point. Irreducibility makes $\mathcal{E}_A$ an irreducible CP map, and Theorem 6.1.3 upgrades the fixed point to positive definite. The spectral theorem diagonalizes it by a unitary conjugation, which preserves the TP normalization. $\square$

Remark 9.2.1.10 (Burnside input for cumulative spanning). ✓ Stmt The passage from irreducible tensors to full algebra spanning first passes through the irreducible action on $\mathbb{C}^D$. Burnside's theorem then says that the resulting irreducible subalgebra of $M_D(\mathbb{C})$ is all of $M_D(\mathbb{C})$. Once the algebra span is full, finite-dimensionality gives a single cumulative span that already equals $M_D(\mathbb{C})$.

Lemma 9.2.1.11 (The algebra span reaches a finite cumulative span). Lean ✓ Stmt *If the generated unital algebra satisfies $\text{alg}(A) = M_D(\mathbb{C})$, then there exists N such that $T_N(A) = M_D(\mathbb{C})$.*

Proof. ✓ Proof Every element of $\text{alg}(A)$ lies in some cumulative span $T_n(A)$: the generators lie in $T_1(A)$, scalar multiples of the identity lie in $T_0(A)$, and the family $\{T_n(A)\}_{n \geq 0}$ is closed under addition and under multiplication with lengths adding. Since $M_D(\mathbb{C})$ is finite-dimensional, the ascending chain $T_0(A) \subseteq T_1(A) \subseteq \dots$ stabilizes, so one can choose a single N with $T_N(A) = M_D(\mathbb{C})$. $\square$

Lemma 9.2.1.12 (Irreducible action gives an irreducible tensor). [Lean](#) [✓ Stmt](#) *If the letters of A act irreducibly on $\mathbb{C}^D$, then A has no nontrivial invariant orthogonal projection.*

Proof. [✓ Proof](#) If P were a nontrivial invariant orthogonal projection, then its range would be invariant under every letter of A . Irreducibility of the action forces this range to be either 0 or all of $\mathbb{C}^D$, so P must be 0 or $\mathbb{1}$, a contradiction. $\square$

9.3 Normal canonical form

This section fixes the two canonical-form predicates used downstream: the block-injective form of Definition 9.3.1 and the spectral *normal* canonical form of Definition 9.3.3. It then records how the self-overlap clause and gauge-phase equivalence follow from primitivity, and closes with the periodicity-removal step that produces primitive transfer maps.

Recall from Definition 2.7.3 and Lemma 7.2.2 that the bilinear overlap is

$$O_{AB}(N) = \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma} = \text{Tr}(F_{AB}^N).$$

This notation enters the self-overlap clause of the canonical-form predicate below.

Definition 9.3.1 (Canonical form conditions). [Lean](#) [✓ Stmt](#) A collection of scaling factors $(\mu_k)_{k=1}^r$ and block tensors $(A_k)_{k=1}^r$ satisfies the *canonical form conditions* if:

1. each A_k is injective,
2. each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. the moduli are non-increasing: $|\mu_1| \geq |\mu_2| \geq \dots \geq |\mu_r|$,
4. all $\mu_k \neq 0$,
5. the self-overlap converges: $O_{A_k A_k}(N) \rightarrow 1$ for each k .

Only this one-sided normalization is assumed; no unital condition is assumed here. Condition (5) is a primitivity hypothesis: for a primitive channel ([Wol12, Theorem 6.7(3)]), $T^n(\rho) \rightarrow \rho_\infty$ for every initial state, which forces the self-overlap to converge to 1. See also [Wol12, Theorem 6.8] for the completely-positive characterisation and Chapter 7 for the transfer-operator gap proof.

Remark 9.3.2 (Normalization convention). [PGVWC07, Theorem 4] uses the unital gauge $\sum_i A_k^i (A_k^i)^\dagger = \mathbb{1}$; here the blocks are in the dual TP gauge $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$. A positive-definite fixed point of the adjoint transfer map provides the change of gauge between the two (Theorem 6.4.2).

Definition 9.3.3 (Normal canonical form conditions). [Lean](#) [✓ Stmt](#) Scaling factors $(\mu_k)_{k=1}^r$ and block tensors $(A_k)_{k=1}^r$ satisfy the *normal canonical form conditions* if:

1. each A_k is irreducible,
2. each A_k satisfies $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. each $\mathcal{E}_{A_k}$ is primitive (equivalently, $\sigma_\partial(\mathcal{E}_{A_k}) = \{1\}$),
4. $|\mu_1| \geq |\mu_2| \geq \dots \geq |\mu_r|$,
5. all $\mu_k \neq 0$,

6. all bond dimensions are positive.

Here “normal” is the spectral sense of [CPGSV16], which by [SPGWC10, Proposition 3] is equivalent to the eventual block-injectivity formulation of Definition 2.4.4.

Theorem 9.3.4 (Self-overlap is derived in normal canonical form). Lean ✓ Stmt *If (μ_k, A_k) satisfies the normal canonical form conditions, then $O_{A_k A_k}(N) \rightarrow 1$ for every block k .*

Proof. ✓ Proof Fix a block k . Irreducibility and the TP normalization give a nonzero positive semidefinite fixed point ρ of $\mathcal{E}_{A_k}$ with fixed-point projector P ; primitivity gives the complementary gap $\rho_{\text{spec}}(\mathcal{E}_{A_k} - P) < 1$ (Theorem 4.5.3.8). Theorem 7.15.2 then yields $O_{A_k A_k}(N) \rightarrow 1$. $\square$

Remark 9.3.5. See Theorem 7.8.1 for the modulus-one eigenvalue rigidity statement for irreducible trace-preserving blocks.

Lemma 9.3.6 (Mixed-transfer spectral radius from unit-modulus overlap). Lean ✓ Stmt *Let A and B have the same bond dimension. If $\|V^{(N)}(A)|V^{(N)}(B)\rangle\| \rightarrow 1$, then the mixed transfer map has spectral radius at least 1.*

Proof. ✓ Proof If the mixed transfer spectral radius were < 1 , the iterated mixed transfer map would tend to zero in operator norm, hence so would its trace; via $\text{Tr}(F_{AB}^N) = \langle V^{(N)}(A)|V^{(N)}(B)\rangle$ the overlap would tend to 0, contradicting the hypothesis. $\square$

Theorem 9.3.7 (Gauge-phase equivalence from unit-modulus overlap). Lean ✓ Stmt *Let A and B be irreducible MPS tensors of the same bond dimension with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. If $\|V^{(N)}(A)|V^{(N)}(B)\rangle\| \rightarrow 1$, then A and B are gauge-phase equivalent.*

Proof. ✓ Proof By Lemma 9.3.6 the mixed transfer spectral radius is at least 1, and Theorem 7.8.1 then gives gauge-phase equivalence. $\square$

9.3.1 Canonical form from primitivity

A normal block needs a primitive transfer map; an irreducible TP block may still have peripheral spectrum larger than $\{1\}$, namely the roots of unity coming from its period. Blocking by the least common multiple of those periods raises the transfer map to a power whose peripheral spectrum is $\{1\}$, removing the periodicity. In [CPGSV21] this is the passage from irreducibility to primitivity by blocking to period one.

Theorem 9.3.1.1 (Blocking yields a peripheral-spectrum primitive transfer map). Lean ✓ Stmt *Let A be an irreducible MPS tensor with $D > 0$ and $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then there exists a blocking length $p > 0$ such that $\sigma_\partial(\mathcal{E}_{A^{[p]}}) = \{1\}$.*

Proof. ✓ Proof Pass to the conjugate-transposed Kraus family $K_i := (A^i)^\dagger$, which is unital with irreducible Kraus map. The TP fixed-point theorem for $\mathcal{E}_A$ provides a positive definite fixed point for the adjoint of K , so Theorem 4.5.3.2 shows every peripheral eigenvalue is a root of unity. Since $\sigma_\partial(\mathcal{E}_A)$ is finite (Theorem 4.5.5), Lemma 4.5.1.2 gives a common exponent p with $\mu^p = 1$ for all $\mu \in \sigma_\partial(\mathcal{E}_A)$, and Lemma 4.5.1.3 gives $\sigma_\partial(\mathcal{E}_A^p) = \{1\}$. Blocking by p takes the p th power of the transfer map, so $\mathcal{E}_{A^{[p]}}$ is primitive. $\square$

Remark 9.3.1.2. Theorem 9.3.1.1 is the periodicity-removal-by-blocking step of [CPGSV16, Appendix A] for an irreducible block already in TP gauge.

9.4 Left-canonical and dual diagonalization of irreducible blocks

Each irreducible block in trace-preserving gauge admits the PGVWC07 unital orientation together with a diagonal positive-definite dual fixed point (Theorem 9.4.1, the Perron–Frobenius gauge of Chapter 6, and Theorem 9.2.1.3); the full PGVWC07 canonical form is Theorem 9.1.

Theorem 9.4.1 (Left-canonical normalization for irreducible TP blocks). Lean ✓ Stmt *Let A be an irreducible MPS tensor in TP gauge ($\sum_i (A^i)^\dagger A^i = \mathbb{1}$) with $D > 0$. Then there exist a unitary U and a diagonal positive definite Λ such that:*

1. $B^i := U^\dagger A^i U$ is TP,
2. $\mathcal{E}_B(\Lambda) = \Lambda$,
3. Λ is diagonal and positive definite,
4. A and B generate the same MPV family.

This is the left-canonical normalization step (Canonical Form II) of [CPGSV16, Appendix A]. The unitary conjugation is a second step inside TP gauge, performed only after TP normalization.

Proof. ✓ Proof Theorem 9.2.1.9 provides the unitary and the diagonal PD fixed point, and unitary conjugation preserves trace-preservation and the MPV family (Theorem 9.2.1). □

Theorem 9.4.2 (Dual diagonalization for irreducible unital blocks). Lean ✓ Stmt *Let A be an irreducible MPS tensor in unital gauge $\sum_i A^i (A^i)^\dagger = \mathbb{1}$ with $D > 0$. Then there exist a unitary U and a diagonal positive definite Λ such that, for $B^i := U^\dagger A^i U$,*

$$\sum_i B^i (B^i)^\dagger = \mathbb{1}, \quad \sum_i (B^i)^\dagger \Lambda B^i = \Lambda, \quad (9.2)$$

and A , B generate the same MPV family. This is the dual-map fixed-point and unitary-diagonalization step of [PGVWC07, Theorem 4].

Proof. ✓ Proof Apply Theorem 9.4.1 to the conjugate-transposed family $i \mapsto (A^i)^\dagger$: unitality of A is trace preservation for this family, whose transfer map is the dual of $\mathcal{E}_A$. Translating back gives the equations (9.2). □

Lemma 9.4.3 (Common scalar rescaling of block weights). Lean ✓ Stmt *Multiplying every block weight in a block-diagonal MPS tensor by a scalar c multiplies every length- N MPV coefficient by c^N .*

Proof. ✓ Proof Expand the block-diagonal MPV coefficient as a finite sum over blocks and factor c^N out of the sum. □

Theorem 9.4.4 (Normal canonical form from a primitive block decomposition). Lean ✓ Stmt *Suppose A generates the same MPV family as $\bigoplus_k \mu_k A_k$, and that each A_k is irreducible, satisfies $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$, and has primitive transfer map, all $\mu_k \neq 0$, and all bond dimensions are positive. Then there exist a blocking length $p > 0$, weights (ν_k) , and blocked tensors (B_k) such that $A^{[p]}$ generates the same MPV family as $\bigoplus_k \nu_k B_k$, and (ν_k, B_k) satisfies the normal canonical form conditions. The weight moduli need not be distinct; equal-modulus blocks are allowed.*

Proof. ✓ Proof The given blocks are already primitive, so take $p = 1$. Reorder the blocks by non-increasing weight modulus, with ties allowed, and set $\nu_k := \mu_k$ and $B_k := A_k$ in that order. Irreducibility, TP normalization, primitivity, nonzero weights, and positive bond dimensions are preserved, so the reordered family satisfies Definition 9.3.3 and generates the same MPV family as A . $\square$

Remark 9.4.5 (Scope of Theorem 9.4.4). Theorem 9.4.4 assumes all six block hypotheses in advance and produces the normal canonical form by reordering. It is not the arbitrary-input existence theorem of [CPGSV16, Section 2.3]; that primitive-block reduction, including the zero-tail count, is Theorem 9.6.2.2 in Section 9.6.2.

9.5 Zero-block separation and the trace-preserving gauge

In an irreducible block decomposition some blocks may have all-zero Kraus operators. An all-zero block contributes $V^{(N)} = 0$ for every $N \geq 1$, so it adds nothing to the positive-length MPV family; at $N = 0$ the MPV coefficient is the bond dimension. The all-zero summand therefore enters the output not as a tensor but as a single count D_0 in the length-zero identity $D = D_0 + \sum_k D_k$, and all positive-length content is an equality of MPV families. This is the positive-length convention used throughout the reduction: comparisons of weighted block sums are stated for $N \geq 1$, with the zero block carried only by D_0 . After separating the zero summand, each nonzero block is placed in the trace-preserving gauge, the furthest unconditional reduction available before periodicity removal.

Theorem 9.5.1 (Zero-block separation). Lean ✓ Stmt *Every MPS tensor A admits a block decomposition into a zero-block bond dimension D_0 and a finite family of nonzero blocks $(B_k)_{k=1}^r$, each irreducible, with at least one nonzero Kraus operator, and of positive bond dimension, recorded by two facts. The positive-length equality of MPV families: for every $N \geq 1$ and every σ ,*

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_{k=1}^r B_k)_\sigma.$$

The length-zero dimension identity

$$D = D_0 + \sum_{k=1}^r D_k,$$

where D_0 is the total bond dimension of the all-zero blocks.

Proof. ✓ Proof Apply the irreducible block decomposition (Theorem 9.2.1.2) and partition the blocks by whether each has a nonzero Kraus operator; D_0 accumulates the bond dimensions of the all-zero blocks. For every all-zero irreducible block C_j and every $N \geq 1$, $V^{(N)}(C_j)_\sigma = 0$, so

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_{k=1}^r B_k)_\sigma.$$

At $N = 0$ the MPV coefficient is the bond dimension, giving $D = D_0 + \sum_{k=1}^r D_k$. $\square$

Theorem 9.5.2 (Blockwise TP-gauge normalization of irreducible blocks). Lean ✓ Stmt *Suppose A generates the same MPV family as $\bigoplus_{k=1}^r B_k$, where each B_k is irreducible with at least one nonzero Kraus operator. Then there are nonzero weights $(\mu_k)_{k=1}^r$ and blocks $(C_k)_{k=1}^r$ such that, for every N and every σ ,*

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_{k=1}^r \mu_k C_k)_\sigma = \sum_{k=1}^r \mu_k^N V^{(N)}(C_k)_\sigma,$$

each C_k irreducible and left-canonical, $\sum_i (C_k^i)^\dagger C_k^i = \mathbb{1}$, with positive bond dimension. This is the blockwise form used in the arbitrary-tensor trace-preserving gauge below.

Proof. ✓ Proof The Perron–Frobenius TP-gauge construction is applied to each irreducible block. Gauge equivalence preserves irreducibility and the MPV family, and the positive scaling factors are absorbed into the weights μ_k . □

9.5.1 Trace-preserving gauge for arbitrary tensors

Theorem 9.5.1.1 (TP-gauge reduction for arbitrary tensors with zero-block separation). Lean

✓ Stmt For any MPS tensor A , there exist a zero-block bond dimension D_0 and nontrivial blocks $(B_k)_{k=1}^r$ with nonzero weights $(\mu_k)_{k=1}^r$, each block irreducible, left-canonical ($\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$), and of positive bond dimension, such that for every $N \geq 1$ and every σ ,

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_{k=1}^r \mu_k B_k)_\sigma,$$

and the zero-block contribution is recorded by the length-zero identity

$$D = D_0 + \sum_{k=1}^r D_k.$$

Proof. ✓ Proof Theorem 9.5.1 supplies D_0 and the nonzero irreducible blocks $(C_k)_{k=1}^r$. For every $N \geq 1$ and every σ ,

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_{k=1}^r C_k)_\sigma.$$

The blockwise TP-gauge theorem then replaces each C_k by a left-canonical block B_k with weight μ_k . Again for every $N \geq 1$ and every σ ,

$$V^{(N)}(\bigoplus_{k=1}^r C_k)_\sigma = V^{(N)}(\bigoplus_{k=1}^r \mu_k B_k)_\sigma.$$

The length-zero identity $D = D_0 + \sum_{k=1}^r D_k$ is preserved. □

9.6 Blocking, period removal, and primitive blocks

Blocking by an integer p carries MPV equalities, weights, transfer maps, and identifications of the blocked physical alphabet. The first consequence is the existence of a common blocking period under which every transfer map in a finite family becomes primitive; the period-removal and primitive-block reductions then proceed from that common period.

Theorem 9.6.1 (Common blocking period for a block family). Lean ✓ Stmt Let $(B_k)_{k=1}^r$ be a finite family of positive-bond-dimension MPS tensors, each admitting a blocking period p_k making its transfer map primitive. Then $p = \text{lcm}(p_1, \dots, p_r)$ is a common period: $\mathcal{E}_{B_k^{[p]}}$ is primitive for every k .

Proof. ✓ Proof Each p_k divides p , so primitivity of $\mathcal{E}_{B_k^{[p_k]}}$ carries over to $\mathcal{E}_{B_k^{[p]}}$ via the divisibility identity for transfer-map powers. □

The following stability statements are used when a primitive irreducible block is blocked again for a common-refinement or injectivity length. This later blocking is distinct from the period-removal length that produced the sector block.

Theorem 9.6.2 (Blocking preserves irreducibility for primitive irreducible blocks). [Lean](#) [✓ Stmt](#) Let A be left-canonical, irreducible, primitive, of positive bond dimension. For every $P > 0$, the blocked tensor $A^{[P]}$ is irreducible: there is no orthogonal projection Q with $Q \neq 0$, $Q \neq \mathbb{1}$ such that, for every P -blocked index α ,

$$(\mathbb{1} - Q)(A^{[P]})^\alpha Q = 0.$$

Proof. [✓ Proof](#) The blocked transfer map is the corresponding positive power of the original one:

$$\mathcal{E}_{A^{[P]}}(X) = \mathcal{E}_A^P(X).$$

Hence a positive-definite fixed point of $\mathcal{E}_A$ remains fixed by $\mathcal{E}_{A^{[P]}}$. Primitivity gives uniqueness of positive semidefinite fixed points for the blocked map, hence its transfer map is irreducible, which is the irreducibility criterion for $A^{[P]}$. $\square$

Theorem 9.6.3 (Extra blocking of primitive irreducible sector blocks). [Lean](#) [✓ Stmt](#) Let A be trace-preserving, primitive, and tensor-irreducible. For every $k > 0$,

$$\sum_{\alpha} ((A^{[k]})^\alpha)^\dagger (A^{[k]})^\alpha = \mathbb{1}, \quad \sigma_{\partial}(\mathcal{E}_{A^{[k]}}) = \{1\},$$

and $A^{[k]}$ is tensor-irreducible. This k is a later common-refinement or injectivity length, distinct from the period-removal length that produced the sector block.

Proof. [✓ Proof](#) Trace preservation follows from blocking a trace-preserving tensor, primitivity from a positive power of a primitive transfer map, and irreducibility from Theorem 9.6.2. $\square$

9.6.1 Period removal by cyclic sectors

The step needed below is the period-removing blocking described in [CPGSV16, Section 2.3]: after blocking each irreducible TP block by the least common multiple of its peripheral periods, the result has no nontrivial p -periodic vectors. The cyclic-sector decomposition comes from the peripheral spectrum of the adjoint transfer map [Wol12, Theorem 6.6].

Theorem 9.6.1.1 (Period removal by blocking). [Lean](#) [✓ Stmt](#) Let A be a left-canonical irreducible tensor. Then there is a period-removing length $m \geq 1$ such that $A^{[m]}$ admits:

1. left-canonical sector tensors $(C_k)_{k=1}^m$ of nonzero bond dimension,
2. a unit-weight direct-sum MPV decomposition $A^{[m]} \sim \bigoplus_{k=1}^m C_k$,
3. orthogonal projections $(P_k)_{k=1}^m$ summing to $\mathbb{1}$ with $\mathcal{E}_{A^\dagger}(P_{k+1}) = P_k$,
4. commutation relations $P_k A_i^{[m]} = A_i^{[m]} P_k$ for every blocked letter i ,
5. the sector trace formula $f_{C_k}(\sigma) = \text{tr}(P_k A_{\sigma_1}^{[m]} \cdots A_{\sigma_N}^{[m]})$,
6. linear isomorphisms onto the compression corners $\varphi_k : M_{\dim C_k}(\mathbb{C}) \xrightarrow{\sim} P_k M_D(\mathbb{C}) P_k$,
7. transfer-intertwining identities $\varphi_k(\mathcal{E}_{C_k^\dagger}(X)) = \mathcal{E}_{(P_k A^{[m]})^\dagger}(\varphi_k(X))$,
8. multiplicativity and adjoint compatibility $\varphi_k(XY) = \varphi_k(X)\varphi_k(Y)$ and $\varphi_k(X^\dagger) = \varphi_k(X)^\dagger$.

This is the cyclic-sector part of the period-removal step of [CPGSV16, Section 2.3]; the projections P_k come from the peripheral spectrum of the adjoint transfer map ([Wol12, Theorem 6.6]).

Proof. ✓ Proof The cyclic decomposition of the adjoint transfer map gives projections (P_k) with $\mathcal{E}_{A^\dagger}(P_{k+1}) = P_k$. Iterating the cyclic relation m times gives $(\mathcal{E}_{A^\dagger})^m(P_k) = P_k$, and the blocked-transfer identity $\mathcal{E}_{(A^{[m]})^\dagger} = (\mathcal{E}_{A^\dagger})^m$ identifies these with the adjoint transfer map of $A^{[m]}$. Apply the block decomposition from adjoint-fixed projections. $\square$

Theorem 9.6.1.2 (Unconditional primitive blocked sectors). Lean ✓ Stmt *Let A be a left-canonical irreducible tensor whose adjoint transfer map has peripheral spectrum $\{\gamma^j : 0 \leq j < m\}$ for a primitive m th root of unity γ , with blocked cyclic-sector decomposition $(C_u)_{u=1}^m, (P_u)_{u=1}^m$, and linear isomorphisms $\varphi_u : M_{\dim C_u}(\mathbb{C}) \xrightarrow{\sim} P_u M_D(\mathbb{C}) P_u$. Assume explicitly that the P_u are orthogonal projections summing to $\mathbb{1}$ and satisfying $\mathcal{E}_{A^\dagger}(P_{u+1}) = P_u$, that each sector has nonzero bond dimension, and that the compression maps satisfy*

$$\varphi_u(\mathcal{E}_{C_u^\dagger}(X)) = \mathcal{E}_{(P_u A^{[m]})^\dagger}(\varphi_u(X)), \quad \varphi_u(XY) = \varphi_u(X)\varphi_u(Y), \quad \varphi_u(X^\dagger) = \varphi_u(X)^\dagger.$$

Here $P_u A^{[m]}$ denotes the blocked tensor with letters $P_u(A^{[m]})^\alpha$. Then every sector tensor satisfies $\sigma_\partial(\mathcal{E}_{C_u}) = \{1\}$ and is tensor-irreducible.

Proof. ✓ Proof Theorem 9.7.10 gives irreducibility of $(\mathcal{E}_A^\dagger)^m$ on each corner P_u . The compression equivalences φ_u intertwine the adjoint transfer map of C_u with the corresponding corner restriction of $(\mathcal{E}_A^\dagger)^m$. The peripheral spectrum of the blocked adjoint map is

$$\sigma_\partial((\mathcal{E}_A^\dagger)^m) = \{(\gamma^j)^m : 0 \leq j < m\} = \{1\},$$

because γ is a primitive m th root of unity. Hence each nonzero irreducible corner restriction is primitive. Primitivity and irreducibility are preserved across φ_u ; then use Theorem 9.7.2 to pass from the adjoint blocked map to the primal transfer map of C_u . $\square$

Theorem 9.6.1.3 (Period removal with primitive irreducible sectors). Lean ✓ Stmt *Let A be a left-canonical irreducible tensor. Then there is a period $m \geq 1$ and a unit-weight decomposition of $A^{[m]}$ into sector blocks $(C_k)_{k=1}^m$, each of which is left-canonical, has primitive transfer map, is tensor-irreducible, and has nonzero bond dimension. Equivalently, the cyclic-sector decomposition of $A^{[m]}$ already satisfies the trace-preservation, primitivity, and irreducibility hypotheses needed for the common-blocking step (Lemma 9.7.9).*

Proof. ✓ Proof Left-canonicity makes $K_i := (A^i)^\dagger$ unital, irreducibility passes to the adjoint map, and the quantum Perron–Frobenius theorem provides a positive-definite fixed point; the cyclic peripheral-spectrum theorem and Theorem 9.6.1.1 then produce the cyclic-sector decomposition. Keep the projections and compression equivalences, and apply Theorem 9.6.1.2 to the sector blocks. $\square$

Theorem 9.6.1.4 (Supported corner compression with a support isometry). Lean ✓ Stmt *Let A be a tensor normalized on the support of an orthogonal projection P , so that $PA^iP = A^i$ and $\sum_i (A^i)^\dagger A^i = P$. Then there are a corner tensor C , a linear isomorphism*

$$\varphi : M_{\dim C}(\mathbb{C}) \xrightarrow{\sim} PM_D(\mathbb{C})P,$$

and a rectangular isometry $V : \mathbb{C}^{\dim C} \rightarrow \mathbb{C}^D$ such that

$$V^\dagger V = \mathbb{1}, \quad VV^\dagger = P, \quad \varphi(X) = VXV^\dagger.$$

Moreover $\varphi(C^i) = A^i$ for every physical letter, and for every word $\sigma = (\sigma_1, \dots, \sigma_N)$,

$$V^{(N)}(C)_\sigma = \text{tr}(PA^{\sigma_1} \dots A^{\sigma_N}).$$

Proof. ✓ **Proof** Diagonalize the projection P by a unitary change of basis. The inclusion of the support coordinates gives a rectangular isometry V with $V^\dagger V = \mathbb{1}$ and $VV^\dagger = P$. Compress the letters of A to this support and let φ be the corresponding corner isomorphism. The support relation $PA^iP = A^i$ gives the corner-letter identity. For each word $\sigma = (\sigma_1, \dots, \sigma_N)$ the MPV component is obtained from the trace computation

$$V^{(N)}(C)_\sigma = \text{tr}(C^{\sigma_1} \dots C^{\sigma_N}) = \text{tr}(V^\dagger A^{\sigma_1} \dots A^{\sigma_N} V) = \text{tr}(PA^{\sigma_1} \dots A^{\sigma_N}).$$

The final equality uses cyclicity of the trace and $VV^\dagger = P$. □

Theorem 9.6.1.5 (Commuting projections give sector isometries). Lean ✓ **Stmt** *Let a left-canonical tensor A commute with a family of orthogonal projections (P_k) summing to $\mathbb{1}$. Then A decomposes into compressed sector tensors (C_k) , with a unit-weight direct-sum MPV decomposition, corner isomorphisms $\varphi_k : M_{\dim C_k}(\mathbb{C}) \rightarrow P_k M_D(\mathbb{C}) P_k$, and rectangular isometries V_k satisfying*

$$V_k^\dagger V_k = \mathbb{1}, \quad V_k V_k^\dagger = P_k, \quad \varphi_k(X) = V_k X V_k^\dagger.$$

The sector letters obey $\varphi_k(C_k^i) = P_k A^i P_k$.

Proof. ✓ **Proof** Apply Theorem 9.6.1.4 to the compression of A on each sector P_k . Orthogonality and $\sum_k P_k = \mathbb{1}$ give the direct-sum decomposition. For every word $\sigma = (\sigma_1, \dots, \sigma_N)$ the partition of unity gives

$$V^{(N)}(A)_\sigma = \text{tr}\left(\left(\sum_k P_k\right) A^{\sigma_1} \dots A^{\sigma_N}\right) = \sum_k \text{tr}(P_k A^{\sigma_1} \dots A^{\sigma_N}) = \sum_k V^{(N)}(C_k)_\sigma,$$

which is the unit-weight matrix-product-vector identity. □

Theorem 9.6.1.6 (Adjoint-fixed projections give sector isometries). Lean ✓ **Stmt** *Let A be left-canonical and let (P_k) be orthogonal projections summing to $\mathbb{1}$ and fixed by the adjoint transfer map. Then the same sector decomposition and support isometries of Theorem 9.6.1.5 exist. In particular, the adjoint fixed-point condition $\mathcal{E}_A^\dagger(P_k) = P_k$ gives*

$$P_k A^i = A^i P_k,$$

which gives the corner-letter identity

$$\varphi_k(C_k^i) = P_k A^i P_k.$$

Proof. ✓ **Proof** The fixed-point equation $\mathcal{E}_A^\dagger(P_k) = P_k$, together with left-canonicity $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, places P_k in the multiplicative domain of $\mathcal{E}_A^\dagger$, forcing

$$P_k A^i = A^i P_k$$

for every letter A^i . The result then follows from Theorem 9.6.1.5. □

Theorem 9.6.1.7 (Spectral cyclic sectors with support isometries). Lean ✓ **Stmt** *Let A be a left-canonical irreducible tensor, and suppose the adjoint transfer map $\mathcal{E}_{A^\dagger}$ has a positive-definite fixed point, is irreducible as a positive map, and has peripheral spectrum*

$$\sigma_\partial(\mathcal{E}_{A^\dagger}) = \{\gamma^j : j \in \mathbb{Z}/m\mathbb{Z}\}$$

for a primitive m th root of unity γ , with $m \geq 1$. Then the spectral cyclic-sector decomposition of $A^{[m]}$ has compression maps which may be chosen with rectangular support isometries V_k satisfying

$$V_k^\dagger V_k = \mathbb{1}, \quad V_k V_k^\dagger = P_k, \quad \varphi_k(X) = V_k X V_k^\dagger.$$

These isometries also realize the blocked corner letters $\varphi_k(C_k^\alpha) = P_k (A^{[m]})^\alpha P_k$.

Proof. ✓ Proof The cyclic peripheral-spectrum decomposition applied to $\mathcal{E}_{A^\dagger}$ provides projections P_k satisfying $\mathcal{E}_{A^\dagger}(P_{k+1}) = P_k$. Iterating m times gives $(\mathcal{E}_{A^\dagger})^m(P_k) = P_k$, and the identity $(\mathcal{E}_{A^\dagger})^m = \mathcal{E}_{(A^{[m]})^\dagger}$ shows that each P_k is fixed by the adjoint transfer map of $A^{[m]}$. Applying Theorem 9.6.1.6 to that blocked tensor gives the compressed sector tensors and the support isometries. $\square$

9.6.2 Reduction to primitive blocks

This subsection completes the reduction after period removal. A TP-primitive-irreducible block is normal, and an arbitrary tensor becomes, after a positive blocking length, a weighted family of left-canonical primitive blocks with the zero-block dimension recorded separately. For two tensors with the same MPV family, the cyclic sectors can be expressed over one common blocked alphabet, which is the form used by the Fundamental Theorem.

Theorem 9.6.2.1 (TP-primitive irreducible tensor is normal). Lean ✓ Stmt *Let A be a left-canonical MPS tensor with $D > 0$, irreducible transfer map, and peripheral spectrum $\{1\}$. Then A is normal.*

Proof. ✓ Proof Irreducibility upgrades the PSD fixed point to positive definite (Theorem 6.1.3), and Theorem 8.10.4.3 then gives normality. $\square$

An arbitrary translation-invariant tensor reduces, after blocking, to a weighted direct sum of primitive irreducible blocks.

Theorem 9.6.2.2 (Arbitrary tensor to primitive block decomposition). Lean ✓ Stmt *For any MPS tensor A , there exist $p \geq 1$, a zero-tail bond dimension D_0 , nonzero weights $(\mu_k)_{k=1}^r$, and blocks $(B_k)_{k=1}^r$ such that, for every blocked σ of positive length N ,*

$$V^{(N)}(A^{[p]})_\sigma = V^{(N)}(\bigoplus_{k=1}^r \mu_k B_k)_\sigma.$$

The zero-block contribution is recorded separately by the length-zero identity $D_0 + \sum_{k=1}^r D_k = D$. Each block is left-canonical and primitive,

$$\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}, \quad \sigma_\partial(\mathcal{E}_{B_k}) = \{1\},$$

and each bond dimension is positive.

Proof. ✓ Proof Theorem 9.5.1.1 supplies the zero-block dimension, the left-canonical nontrivial blocks, the positive-length equality, and the identity $D = D_0 + \sum_k D_k$. Theorem 9.6.1 provides a common blocking period p making all transfer maps primitive, Lemma 9.7.8 keeps the blocked weights nonzero, and Lemma 9.7.6 carries the positive-length equality through blocking. $\square$

Theorem 9.6.2.3 (Cyclic sectors after the zero-block split). Lean ✓ Stmt *If A and B generate the same MPV family, then after the zero-block split and the trace-preserving gauge there are weighted nonzero-block tensors $\bigoplus_j \mu_j^A A_j$ and $\bigoplus_k \mu_k^B B_k$ with, for every $N \geq 1$ and σ ,*

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_j \mu_j^A A_j)_\sigma, \quad V^{(N)}(B)_\sigma = V^{(N)}(\bigoplus_k \mu_k^B B_k)_\sigma,$$

and the two nonzero parts agree:

$$V^{(N)}(\bigoplus_j \mu_j^A A_j)_\sigma = V^{(N)}(\bigoplus_k \mu_k^B B_k)_\sigma.$$

Each nonzero block has primitive irreducible cyclic sectors: for each j , there is $m_j \geq 1$ with a unit-weight sector decomposition $A_j^{[m_j]} \sim \bigoplus_{u=1}^{m_j} C_{j,u}^A$, and similarly for B_k .

Proof. ✓ Proof Apply Theorem 9.5.1.1 separately to A and B . Its positive-length conclusion gives, for $N \geq 1$,

$$V^{(N)}(A)_\sigma = V^{(N)}(\bigoplus_j \mu_j^A A_j)_\sigma, \quad V^{(N)}(B)_\sigma = V^{(N)}(\bigoplus_k \mu_k^B B_k)_\sigma.$$

Chaining these through $V(A) = V(B)$ identifies the two nonzero parts. Finally apply Theorem 9.6.1.3 to every trace-preserving irreducible nonzero-weight block. $\square$

Theorem 9.6.2.4 (Common blocking length for cyclic sectors). Lean ✓ Stmt *If two tensors generate the same MPV family, then the cyclic-sector decompositions on the two sides can be expressed at one common positive blocking length p . For every $N \geq 1$ and every blocked σ , each blocked tensor equals its powered nonzero part,*

$$V^{(N)}(A^{[p]})_\sigma = V^{(N)}(\bigoplus_j (\mu_j^A)^p (A_j)^{[p]})_\sigma, \quad V^{(N)}(B^{[p]})_\sigma = V^{(N)}(\bigoplus_k (\mu_k^B)^p (B_k)^{[p]})_\sigma,$$

and the two powered nonzero parts agree,

$$V^{(N)}(\bigoplus_j (\mu_j^A)^p (A_j)^{[p]})_\sigma = V^{(N)}(\bigoplus_k (\mu_k^B)^p (B_k)^{[p]})_\sigma.$$

Proof. ✓ Proof Start from Theorem 9.6.2.3. With p_A, p_B the least common multiples of the period-removal lengths on the two sides, use $p = \text{lcm}(p_A, p_B)$. The prescribed-common-multiple lemma constructs both one-sided sector families at this length, and the positive-length blocking lemmas carry the one-sided tensor/nonzero-part equalities over to the common alphabet; for every $N \geq 1$,

$$V^{(N)}(A^{[p]})_\sigma = V^{(N)}(\bigoplus_j (\mu_j^A)^p (A_j)^{[p]})_\sigma.$$

The same applies on the B side, and the two-sided blocking lemma gives equality of the two powered nonzero parts. $\square$

Theorem 9.6.2.5 (Unconditional common primitive block decompositions). Lean ✓ Stmt *Let A and B have the same MPV family. Then there is a positive blocking length at which both blocked tensors have the same positive-length MPV family as weighted direct sums of trace-preserving, primitive, tensor-irreducible blocks with positive bond dimensions and nonzero weights. The two weighted nonzero-sector families have the same positive-length MPV family.*

Proof. ✓ Proof Apply Theorem 9.6.2.4 to choose the common blocking length and the two common cyclic-sector families. For either one of these families, write G_k for its cyclic-sector blocks, and write (ν_x, C_x) for the weights and blocks of the flattened common-sector family from Lemma 9.7.16. The common-alphabet nonzero-part identity gives, for every length N and blocked word σ ,

$$V^{(N)}(\bigoplus_k \mu_k^p G_k^{[p]})_\sigma = V^{(N)}(\bigoplus_x \nu_x C_x)_\sigma,$$

Applying this on the two sides of the comparison yields a positive length p and weighted primitive block families such that, for every $N \geq 1$ and blocked word σ ,

$$V^{(N)}(A^{[p]})_\sigma = V^{(N)}(\bigoplus_x \mu_x^A A_x)_\sigma, \quad V^{(N)}(B^{[p]})_\sigma = V^{(N)}(\bigoplus_y \mu_y^B B_y)_\sigma,$$

with the two weighted nonzero parts equal at every positive length. The structural properties and the nonvanishing of the weights come from the common-sector families. $\square$

A weighted block sum with distinct sectors is the simplest sector decomposition: one block per sector, each of multiplicity one. Equal-modulus and repeated-copy sectors are not single scalar powers, already for $A = C \oplus (-C)$ with $V^{(N)}(A)_\sigma = (1 + (-1)^N) V^{(N)}(C)_\sigma$; they are compared by the basis-of-normal-tensors construction of Chapter 10.

Definition 9.6.2.6 (Single-copy sector decomposition). [Lean](#) [✓ Stmt](#) Given nonzero weights $(\mu_k)_{k=1}^r$ and blocks $(B_k)_{k=1}^r$, the *single-copy sector decomposition* has r sectors, one basis block B_k per sector, multiplicity 1, and sector weight μ_k . As a finite sum,

$$\sum_{k=1}^r \mu_k^N V^{(N)}(B_k)_\sigma = V^{(N)}(\bigoplus_k \mu_k B_k)_\sigma.$$

Lemma 9.6.2.7 (Single-copy sector decomposition preserves the represented family). [Lean](#) [✓ Stmt](#) The single-copy sector decomposition of $(\mu_k, B_k)_k$ represents the same MPS family as $\bigoplus_k \mu_k B_k$.

Proof. [✓ Proof](#) Expanding the sector decomposition gives $\sum_k \mu_k^N V^{(N)}(B_k)_\sigma$, the finite block-sum expression for $\bigoplus_k \mu_k B_k$. $\square$

9.7 Blocking and weighted direct sums

Blocking by a length p commutes with weighted direct sums up to the p th power of each weight: for nonzero weights (μ_k) and blocks (B_k) ,

$$(\bigoplus_k \mu_k B_k)^{[p]} \sim \bigoplus_k \mu_k^p B_k^{[p]},$$

where $\sim$ is equality of MPV families. When two block families are reblocked to a common length $p = m_k e_k$, the p -blocked alphabet is identified with the e_k -fold tensor power of the m_k -blocked alphabet, so each sector reblocked through its own period agrees with the directly p -blocked block.

Lemma 9.7.1 (Canonical form from peripheral primitivity). [Lean](#) [✓ Stmt](#) Let $(\mu_k, A_k)_{k=1}^r$ be nonzero weights and injective left-canonical blocks with positive bond dimensions, non-increasing moduli $|\mu_1| \geq \dots \geq |\mu_r| > 0$, and $\sigma_\partial(\mathcal{E}_{A_k}) = \{1\}$. Then $(\mu_k, A_k)_{k=1}^r$ satisfies the canonical-form conditions of Definition 9.3.1, and $O_{A_k A_k}(N) \rightarrow 1$ for every k .

Proof. [✓ Proof](#) The injectivity, left-canonical equation, weight ordering, and nonzero-weight clauses are exactly the hypotheses. For each block, peripheral primitivity gives the complementary spectral gap $\rho_{\text{spec}}(\mathcal{E}_{A_k} - P) < 1$ by Theorem 4.5.3.8. Theorem 7.15.2 then yields $O_{A_k A_k}(N) \rightarrow 1$. $\square$

Lemma 9.7.2 (Adjoint primitivity equivalence). [Lean](#) [✓ Stmt](#) Let $\mathcal{E}$ be a finite-dimensional completely positive map. Then $\mathcal{E}$ is primitive if and only if its Frobenius adjoint $\mathcal{E}^\dagger$ is primitive; equivalently, $\sigma_\partial(\mathcal{E}) = \{1\}$ iff $\sigma_\partial(\mathcal{E}^\dagger) = \{1\}$, since the peripheral spectrum of $\mathcal{E}^\dagger$ is the complex conjugate of that of $\mathcal{E}$.

Lemma 9.7.3 (Primitivity under blocking by a multiple). [Lean](#) [✓ Stmt](#) Let A have positive bond dimension and let $p_0, p \geq 1$ with $p_0 \mid p$. If $\mathcal{E}_{A[p_0]}$ is primitive, then $\mathcal{E}_{A[p]}$ is primitive.

Lemma 9.7.4 (Blocking the weighted block sum). [Lean](#) [✓ Stmt](#) For nonzero weights $(\mu_k)_{k=1}^r$, blocks $(B_k)_{k=1}^r$, and any $p \geq 1$,

$$(\bigoplus_k \mu_k B_k)^{[p]} \sim \bigoplus_k \mu_k^p B_k^{[p]},$$

where $\sim$ denotes generation of the same MPV family at every length.

Lemma 9.7.5 (Positive-length equality survives blocking). [Lean](#) [✓ Stmt](#) If two tensors have the same MPV coefficients at every positive length, then so do their blocked tensors after any positive blocking length.

Lemma 9.7.6 (Positive-length blocking of a weighted block sum). [Lean](#) [✓ Stmt](#) If A and $\bigoplus_k \mu_k B_k$ have the same MPV coefficients at every positive length, then after any positive blocking length p , $A^{[p]}$ and $\bigoplus_k \mu_k^p B_k^{[p]}$ have the same MPV coefficients at every positive length.

Lemma 9.7.7 (Positive-length equality of powered block sums). [Lean](#) [✓ Stmt](#) If $\bigoplus_j \mu_j^A A_j$ and $\bigoplus_k \mu_k^B B_k$ have the same MPV coefficients at every positive length, then after any positive common blocking length p , $\bigoplus_j (\mu_j^A)^p A_j^{[p]}$ and $\bigoplus_k (\mu_k^B)^p B_k^{[p]}$ have the same MPV coefficients at every positive length.

Lemma 9.7.8 (Nonzero weights survive blocking). [Lean](#) [✓ Stmt](#) If $\mu \in \mathbb{C}$ is nonzero and $p \geq 1$, then $\mu^p \neq 0$; blocking preserves the nonzero-weight hypothesis of a weighted block decomposition.

Lemma 9.7.9 (Common reblocking of cyclic-sector families). [Lean](#) [✓ Stmt](#) Given a finite family $(B_k)_{k=1}^r$, each with primitive irreducible cyclic sectors, there is a common blocking length p such that each $B_k^{[p]}$ admits a unit-weight direct-sum decomposition over a single common blocked alphabet $d^{[p]}$ into left-canonical blocks with primitive transfer maps, tensor irreducibility, and positive bond dimensions.

Lemma 9.7.10 (Sector irreducibility for the adjoint blocked map). [Lean](#) [✓ Stmt](#) Let A be a left-canonical irreducible MPS tensor with $D > 0$, let m be the period of its cyclic-sector decomposition, and let $(P_u)_{u=1}^m$ be the cyclic projections of the adjoint transfer map with $\mathcal{E}_{A^\dagger}(P_{u+1}) = P_u$ and $\sum_u P_u = \mathbb{1}$. Then the restriction of $(\mathcal{E}_{A^\dagger})^m$ to each corner $P_u M_D(\mathbb{C}) P_u$ is irreducible.

Lemma 9.7.11 (Iterated blocking at the transfer-map level). [Lean](#) [✓ Stmt](#) For any MPS tensor A and $m, n \geq 0$,

$$\mathcal{E}_{(A^{[m]})^{[n]}} = \mathcal{E}_{A^{[mn]}}.$$

Definition 9.7.12 (Primitive irreducible cyclic-sector decomposition). [Lean](#) [✓ Stmt](#) A has primitive irreducible cyclic sectors of period $m \geq 1$ if there is a family $(C_k)_{k=1}^m$ with $A^{[m]} \sim \bigoplus_{k=1}^m C_k$ (unit weights), each C_k left-canonical, with primitive transfer map, tensor-irreducible, and of positive bond dimension.

Lemma 9.7.13 (Common reblocking at a prescribed multiple). [Lean](#) [✓ Stmt](#) With hypotheses as in Lemma 9.7.9, and given any common multiple p of the per-block period-removal lengths (m_k) , the common reblocking can be performed at the prescribed length p .

Lemma 9.7.14 (Structural properties of the common reblocked sectors). [Lean](#) [✓ Stmt](#) Each sector tensor in the common reblocked cyclic-sector family of Lemma 9.7.9 is trace-preserving ($\sum_i (B^i)^\dagger B^i = \mathbb{1}$), has primitive transfer map ($\sigma_\partial(\mathcal{E}_B) = \{1\}$), is tensor-irreducible, and has positive bond dimension.

Lemma 9.7.15 (Common-alphabet flattening). [Lean](#) [✓ Stmt](#) With $p = m_k e_k$ for each k , the iterated blocked tensor $(B_k^{[m_k]})^{[e_k]}$ agrees with $B_k^{[p]}$ in MPV family, after the canonical identification of the p -blocked alphabet $d^{[p]}$ with the e_k -fold tensor power of the m_k -blocked alphabet $d^{[m_k]}$.

Lemma 9.7.16 (Reindexed nonzero-part identity). [Lean](#) [✓ Stmt](#) The powered nonzero-part decomposition $\bigoplus_k \mu_k^p B_k^{[p]}$ carries over under the common-alphabet identification of Lemma 9.7.15: the common-alphabet sectors have the same nonzero-part decomposition as the direct p -blocked tensors.

Lemma 9.7.17 (Direct blocking in the common alphabet). [Lean](#) [✓ Stmt](#) *For every original block B_k , direct blocking at the common length $p = m_k e_k$ agrees, as an MPV family, with the same block transported to the common blocked alphabet. Write $\widetilde{B}_k$ for this common-alphabet image. Then*

$$V^{(N)}(B_k^{[p]})_\sigma = V^{(N)}(\widetilde{B}_k)_\sigma.$$

Chapter 10

Basis of Normal Tensors

A *basis of normal tensors* (BNT) for an MPS tensor A is a finite family of normal blocks $\{A_j\}$ whose MPV states $\{|V^{(N)}(A_j)\rangle\}$ span $|V^{(N)}(A)\rangle$ at every length and are eventually linearly independent. Two payoffs follow. First, equal MPV families force a bijective gauge-phase matching of the basis sectors $B_k = e^{i\phi_k} X_k A_j X_k^{-1}$. Second, the sector coefficients $c_N^{(j)} = \sum_q \mu_{j,q}^N$ recover the per-copy weight multisets by Newton–Girard. The references are [PGVWC07], [CPGSV16], and [CPGSV21].

10.1 Bases of normal tensors

Definition 10.1.1 (Basis of Normal Tensors (BNT)). Lean ✓ Stmt A *basis of normal tensors* (BNT) for a total tensor A_{tot} consists of g block tensors $(A_1, \dots, A_g)$ with bond dimensions $(D_1, \dots, D_g)$ such that each A_j is normal (Definition 2.4.4); at each system size N the MPV of A_{tot} lies in the span of the block MPVs $\{|V^{(N)}(A_j)\rangle\}_{j=1}^g$; and for sufficiently large N those block MPVs are linearly independent. This is [CPGSV21, Definition IV.2]. In the canonical-form characterization of [CPGSV16, Section 2.3], a BNT is a minimal family of representatives for $\tilde{A}_k^i = e^{i\phi_k} X_k A_j^i X_k^{-1}$ among the normal blocks.

Definition 10.1.2 (Basis of normal tensors, coefficient form). Lean ✓ Stmt The coefficient form of a basis of normal tensors: each block A_j is a CPSV normal tensor (Definition 9.1.1), the MPV of A is given at every length by explicit coefficients $c_j^{(N)} \in \mathbb{C}$ with $|V^{(N)}(A)\rangle = \sum_{j=1}^g c_j^{(N)} |V^{(N)}(A_j)\rangle$, and the block MPVs $\{|V^{(N)}(A_j)\rangle\}$ are eventually linearly independent.

Lemma 10.1.3 (Perron gauges preserve primitivity). Lean ✓ Stmt Let σ be positive definite. The transfer map of the Perron gauge determined by σ is primitive if and only if the original transfer map is primitive.

Proof. ✓ Proof The two transfer maps are similar through the invertible congruence $X \mapsto \sigma^{-1/2} X \sigma^{-1/2}$. Similar linear maps have the same spectrum, so the condition that 1 is the only eigenvalue on the unit circle is unchanged. $\square$

Theorem 10.1.4 (Spectral normality gives a pure trace-preserving gauge). Lean ✓ Stmt Let A be a normal tensor of positive bond dimension in the sense of [CPGSV16, Section 2.2]. There is a positive definite matrix σ such that the gauge obtained from σ is trace-preserving, primitive, and irreducible. No scalar rescaling is needed.

Proof. ✓ Proof The irreducible Perron–Frobenius theorem gives positive definite right and left eigenvectors with positive eigenvalues r and t . The spectral normalization gives $r = 1$. Adjointness with respect to the trace pairing gives

$$r \operatorname{Tr}(\sigma\rho) = t \operatorname{Tr}(\sigma\rho).$$

Since σ and ρ are positive definite, their pairing is nonzero, hence $t = 1$. Thus the Perron gauge is a pure similarity. Irreducibility and primitivity are invariant under this similarity. $\square$

Theorem 10.1.5 (Spectral normality gives eventual block injectivity). Lean ✓ Stmt *Every positive-bond-dimension normal tensor in the sense of [CPGSV16, Section 2.2] is injective after blocking by some finite length.*

Proof. ✓ Proof Apply Theorem 10.1.4. The trace-preserving, primitive, irreducible representative is eventually block-injective, and gauge invariance transports this conclusion back to the original tensor. $\square$

Lemma 10.1.6 (Exact MPV phase equivalence determines the bond dimension). Lean ✓ Stmt *Suppose that, for two tensors A and B and some nonzero $\zeta \in \mathbb{C}$,*

$$|V^{(N)}(B)\rangle = \zeta^N |V^{(N)}(A)\rangle$$

at every nonnegative length N . Then A and B have the same bond dimension.

Proof. ✓ Proof At length zero the two coefficients are the traces of the corresponding identity matrices, while $\zeta^0 = 1$. Their equality is therefore equality of the two bond dimensions. $\square$

Theorem 10.1.7 (Non-decaying rectangular overlap determines the bond dimension). Lean ✓ Stmt *Let A and B be irreducible trace-preserving tensors of positive, possibly different, bond dimensions. If*

$$|O_{AB}(N)| \rightarrow 1,$$

then their bond dimensions are equal.

Proof. ✓ Proof If the bond dimensions differed, the rectangular overlap-decay theorem would give $O_{AB}(N) \rightarrow 0$, and hence $|O_{AB}(N)| \rightarrow 0$, contradicting the assumed limit. $\square$

Theorem 10.1.8 (Phase-equivalent normal tensors are gauge-phase equivalent). Lean Lean Lean ✓ Stmt *Let A and B be normal tensors of positive, possibly different, bond dimensions. Suppose that, for some nonzero $\zeta \in \mathbb{C}$,*

$$|V^{(N)}(B)\rangle = \zeta^N |V^{(N)}(A)\rangle \quad (N \geq 0).$$

Then the bond dimensions are equal and there is an invertible matrix X and a scalar η of unit modulus such that

$$B^i = \eta X A^i X^{-1}$$

for every physical letter i . This is the gauge recovery used in [CPGSV16, Appendix A].

Proof. ✓ Proof Apply Theorem 10.1.4 to both tensors. Gauge invariance of matrix product vectors preserves the phase relation, whose self-overlap identities imply unit modulus for the phase. The resulting rectangular cross-overlap has norm tending to one. If the bond dimensions differed, Theorem 10.1.7 would give a contradiction. Thus the bond dimensions agree, and the unit-overlap rigidity theorem gives a gauge-phase equivalence between the trace-preserving representatives. Conjugating by the two Perron gauges gives the asserted relation for A and B . $\square$

Theorem 10.1.9 (Normal-tensor overlap dichotomy). [Lean](#) [✓ Stmt](#) Let A and B be normal tensors of positive, possibly different, bond dimensions. Their self-overlaps satisfy

$$O_{AA}(N) \longrightarrow 1, \quad O_{BB}(N) \longrightarrow 1.$$

Moreover, either

$$O_{AB}(N) \longrightarrow 0,$$

or their bond dimensions are equal; call the common value D . Then there are $X \in \text{GL}_D(\mathbb{C})$ and $\zeta \in \mathbb{C}$, with $|\zeta| = 1$, such that

$$B^i = \zeta X A^i X^{-1}$$

for every physical letter i . In the latter case, $|O_{AB}(N)| \rightarrow 1$. This is [\[CPGSV16, Lemma A.2\]](#).

Proof. [✓ Proof](#) Put both tensors in their pure trace-preserving gauges. If their bond dimensions differ, rectangular overlap decay gives $O_{AB}(N) \rightarrow 0$. If their dimensions agree but they are not gauge-phase equivalent, the equal-dimension overlap-decay theorem gives the same conclusion. In the remaining case, $B^i = \zeta X A^i X^{-1}$. By Theorem [10.1.8](#), $|\zeta| = 1$. Hence

$$O_{AB}(N) = \bar{\zeta}^N O_{AA}(N), \quad |O_{AB}(N)| = |O_{AA}(N)| \longrightarrow 1.$$

□

Theorem 10.1.10 (Exact phase alternative for normal tensors). [Lean](#) [✓ Stmt](#) Let A and B be normal tensors of positive, possibly different, bond dimensions. Then either $O_{AB}(N) \rightarrow 0$, or there is a phase $\zeta \in \mathbb{C}$, with $|\zeta| = 1$, such that for every chain length N ,

$$|V^{(N)}(B)\rangle = \zeta^N |V^{(N)}(A)\rangle.$$

This is [\[CPGSV16, Corollary A.3\]](#).

Proof. [✓ Proof](#) Apply Theorem [10.1.9](#). In its second alternative, gauge-phase covariance gives $V^{(N)}(B)_\sigma = \zeta^N V^{(N)}(A)_\sigma$ for every configuration σ , and hence the asserted equality of vectors. □

Theorem 10.1.11 (Separated normal tensors are eventually independent). [Lean](#) [✓ Stmt](#) Let $(A_j)_{j=1}^g$ be a finite family of positive-bond-dimension normal tensors, no two of which are gauge-phase equivalent after identifying equal bond dimensions. Then the matrix product vectors $\{|V^{(N)}(A_j)\rangle\}_{j=1}^g$ are linearly independent for all sufficiently large N , as in [\[CPGSV16, Appendix A\]](#).

Proof. [✓ Proof](#) Put every tensor in the pure trace-preserving gauge of Theorem [10.1.4](#). Distinct gauge-phase classes have overlaps tending to zero, while each self-overlap tends to one. The finite Gram matrix therefore tends to the identity and is invertible for all sufficiently large lengths. Gauge invariance of matrix product vectors gives the conclusion for the original tensors. □

Theorem 10.1.12 (Simultaneous one-letter span after blocking a BNT). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $(A_j)_{j=1}^g$ be a basis of normal tensors, with bond dimensions D_j . There is a positive integer L such that the one-letter evaluations of the blocked tensors span the full product algebra:

$$\text{span}_{\mathbb{C}} \left\{ ((A_1^{[L]})^i, \dots, (A_g^{[L]})^i) : 0 \leq i < d^L \right\} = \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

This is the existence form of simultaneous block injectivity associated with [\[CPGSV16, lines 317–345\]](#); no numerical bound on L is asserted here.

Proof. [✓ Proof](#) Eventual linear independence implies $D_j > 0$ for every j and excludes gauge-phase equivalence between two distinct blocks. Put every block in a trace-preserving gauge. Normality gives a common positive length p at which every gauged block is p -block injective; this injectivity persists at all larger lengths. Pairwise direct-sum separation at lengths p , $p + 1$, and $3(p + 1)$ gives, for each ordered pair of distinct blocks, a length- $3(p + 1)$ polynomial equal to the identity on the first block and zero on the second. Multiplying the $g - 1$ pairwise polynomials gives a selector for each block. Combining these selectors with the common length- p word spans yields the full product-algebra span at

$$L = p + (g - 1)3(p + 1).$$

Conjugation by the Perron gauges preserves this span. Finally, a one-letter evaluation of $A_j^{[L]}$ is precisely a length- L word evaluation of A_j . $\square$




Figure 10.1: The decomposition behind the coefficient form: the tensor is $A^i = X(\bigoplus_j \bigoplus_q \mu_{j,q} A_j^i)X^{-1}$ with A_j a basis of normal tensors, and the direct sums are drawn as the block-index lines j and q threading the gauge X , the basis tensor A_j , the coefficient tensor μ , and X^{-1} . This is [CPGSV16, Section 2.3, lines 304–313].

The eventual linear-independence clause is produced from the asymptotic overlaps: diagonal overlaps tending to one and off-diagonal overlaps tending to zero make the Gram matrix converge to the identity.

Lemma 10.1.13 (Eventual linear independence from finite-index overlap orthonormality). [Lean](#)

[✓ Stmt](#) Let I be a finite index set and let $(A_i)_{i \in I}$ be a finite family of tensors such that $O_{A_i A_i}(N) \rightarrow 1$ for each i , and $O_{A_i A_j}(N) \rightarrow 0$ whenever $i \neq j$. Then the MPV states $\{|V^{(N)}(A_i)\rangle\}_{i \in I}$ are linearly independent for all sufficiently large N .

Proof. [✓ Proof](#) The Gram matrix of the family $\{|V^{(N)}(A_i)\rangle\}_{i \in I}$ converges entrywise to the identity matrix. Hence for all sufficiently large N it is invertible, so the MPV states are linearly independent. $\square$

Lemma 10.1.14 (Eventual linear independence from overlap orthonormality). [Lean](#) [✓ Stmt](#)

Let $(A_j)_{j=1}^g$ be a finite family of tensors such that $O_{A_j A_j}(N) \rightarrow 1$ for each j and $O_{A_i A_j}(N) \rightarrow 0$ for $i \neq j$. Then the MPV states $\{|V^{(N)}(A_j)\rangle\}_{j=1}^g$ are linearly independent for all sufficiently large N .

Proof. [✓ Proof](#) Apply Lemma 10.1.13 with $I = \{1, \dots, g\}$. $\square$

The same conclusion is often more convenient with an explicit threshold rather than the eventual quantifier.

Theorem 10.1.15 (Existential BNT independence from overlap limits). [Lean](#) [✓ Stmt](#) If the self-overlaps of a finite block family tend to 1 and the cross-overlaps of distinct blocks tend to 0, then the MPV states are linearly independent for all sufficiently large system sizes.

Proof. ✓ Proof This is Lemma 10.1.14, restated with an explicit threshold N_0 . □

Eventual linear independence is also what converts an equality of two finite linear combinations into equality of their coefficients; this is the mechanism behind the coefficient identities of Chapter 11.

Lemma 10.1.16 (Eventual coefficient extraction from linear independence). Lean ✓ Stmt *If a finite family of vectors is linearly independent for all sufficiently large N , and two finite linear combinations of that family are equal for all sufficiently large N , then the corresponding coefficients are equal for all sufficiently large N .*

Proof. ✓ Proof Move the two linear combinations to one side. Linear independence forces each coefficient difference to vanish at every sufficiently large length. □

When two BNT families must be compared against each other, the same Gram-matrix criterion applies to their disjoint union, provided the mixed overlaps also decay. This is the form used in the exact sector matching of Section 10.6.

Lemma 10.1.17 (Eventual linear independence for two families). Lean ✓ Stmt *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be finite families of tensors. Suppose the overlaps inside each family are asymptotically orthonormal, and suppose every mixed overlap $O_{A_j B_k}(N)$ tends to 0. Then the combined MPV family*

$$\{|V^{(N)}(A_j)\rangle\}_{j=1}^{g_A} \cup \{|V^{(N)}(B_k)\rangle\}_{k=1}^{g_B}$$

is linearly independent for all sufficiently large N .

Proof. ✓ Proof Apply the finite-index Gram-matrix criterion to the disjoint union of the two index sets. The within-family hypotheses give the diagonal and off-diagonal limits inside each block, and the mixed overlap hypothesis gives the off-diagonal limits between the two blocks. □

We now isolate the separation condition that supplies the off-diagonal decay and combine it with the canonical-form hypotheses.

Definition 10.1.18 (No gauge-phase-equivalent distinct blocks). Lean ✓ Stmt A finite block family has no gauge-phase-equivalent distinct blocks when, for any two distinct indices with the same bond dimension, the corresponding blocks are not gauge-phase equivalent.

Definition 10.1.19 (Normal canonical form with BNT separation). Lean ✓ Stmt A family is in *normal canonical form with BNT separation* if it satisfies Definition 9.3.3 and distinct blocks of the same bond dimension are not gauge-phase equivalent. The weight moduli are non-increasing but need not be strictly decreasing, so equal-modulus blocks are allowed; the grouping into a BNT is governed by gauge-phase equivalence, not by distinctness of moduli. Repeated equal-modulus copies of one basis tensor are not separate blocks here; they appear as the weights $\mu_{j,q}$ in $c_N^{(j)} = \sum_q \mu_{j,q}^N$ in the sector decomposition below.

The off-diagonal decay needed for the BNT criterion follows from the separation condition together with the overlap-decay theorems of the preceding chapter.

Theorem 10.1.20 (Cross-overlap decay from explicit BNT hypotheses). Lean ✓ Stmt *Let $(A_j)_{j=1}^r$ be a family in which each A_j is injective and left-canonical, $\sum_i (A_j^i)^\dagger A_j^i = \mathbb{1}$, and distinct equal-dimension blocks are not gauge-phase equivalent. Then $O_{A_j A_k}(N) \rightarrow 0$ for every pair $j \neq k$.*

Proof. ✓ **Proof** If the bond dimensions differ, Theorem 7.5.3 gives the decay. If the bond dimensions agree, the non-equivalence hypothesis excludes gauge-phase equivalence, so Theorem 7.6.4 gives the decay. $\square$

Lemma 10.1.21 (BNT from explicit blockwise hypotheses). Lean ✓ **Stmt** *Let $(\mu_k, A_k)_{k=1}^r$ be a weighted block family. Assume each A_k is injective and left-canonical, $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$, that $O_{A_k A_k}(N) \rightarrow 1$ for each k , and that distinct equal-dimension blocks are not gauge-phase equivalent. Then the blocks form a basis of normal tensors for the block-diagonal tensor $\bigoplus_k \mu_k A_k$.*

Proof. ✓ **Proof** Injective blocks are normal. The block-diagonal decomposition theorem gives the MPV expansion of $\bigoplus_k \mu_k A_k$ in terms of the block MPVs. The self-overlap limits are part of the hypotheses, and Theorem 10.1.20 supplies the off-diagonal decay. Therefore Lemma 10.1.14 gives the eventual linear independence required in Definition 10.1.1. $\square$

The normal-canonical formulation encodes normality spectrally (primitive transfer map), the BNT definition algebraically (eventual block injectivity).

Remark 10.1.22 (Comparing the two normality conditions). The spectral normality of Definition 9.3.3 and the algebraic normality of Definition 2.4.4 agree: the Wielandt theory of Chapter 8 supplies the implication from primitivity to eventual block injectivity used here.

Lemma 10.1.23 (Restricted normal-CF-BNT hypotheses give a BNT). Lean ✓ **Stmt** *If a family is in normal canonical form with BNT separation and each block is also normal in the algebraic sense, then the blocks form a basis of normal tensors for the associated block-diagonal tensor. The blocks here are distinct representatives.*

Proof. ✓ **Proof** The normal-canonical hypotheses already give the block-diagonal MPV expansion, the self-overlap normalization, and the eventual linear independence coming from irreducible trace-preserving overlap decay. Supplying algebraic normality for each block adds the remaining clause in Definition 10.1.1. $\square$

10.2 Permutation rigidity for bases of normal tensors

When two bases of normal tensors generate the same MPV subspace at every length, the blocks must agree up to a permutation and per-block gauge phases. The argument turns the equality of spans into an invertible change-of-basis matrix, then analyses its asymptotics through the overlaps.

Lemma 10.2.1 (Invertible change of basis). Lean ✓ **Stmt** *Let $v_1, \dots, v_g$ and $w_1, \dots, w_g$ be two linearly independent families in a complex vector space with the same span. Then there exists an invertible matrix $U \in \text{GL}_g(\mathbb{C})$ such that*

$$w_j = \sum_{i=1}^g U_{ij} v_i$$

for every j .

Proof. ✓ **Proof** Because the two families span the same subspace, each w_j has a unique expansion in the basis $\{v_i\}_{i=1}^g$; these coefficients form the columns of U . If $Ua = 0$, then the corresponding linear relation among the w_j is trivial because the w_j are linearly independent. Thus U is injective, hence invertible. $\square$

Lemma 10.2.2 (Eventual change of basis from equal spans). [Lean](#) [✓ Stmt](#) *If two finite families of blocks have asymptotically orthonormal overlaps within each family and their MPV spans agree for every system size N , then for all sufficiently large N there exists an invertible matrix U_N expressing the MPV states of one family as linear combinations of the MPV states of the other.*

Proof. [✓ Proof](#) By Lemma 10.1.14, both families are linearly independent for all sufficiently large N . At such an N , the span equality lets us apply Lemma 10.2.1. $\square$

The change-of-basis matrix is now used to match the blocks. We first treat equal block counts, then remove that restriction.

Theorem 10.2.3 (Permutation rigidity for equal block counts). [Lean](#) [✓ Stmt](#) *Let $(A_j)_{j=1}^g$ and $(B_j)_{j=1}^g$ be injective families of positive bond dimension satisfying trace-preserving normalization, with self-overlaps tending to 1 and off-diagonal overlaps tending to 0 within each family. If the spans of the MPV states agree for every system size N , then there is a permutation π of $\{1, \dots, g\}$ such that A_j and $B_{\pi(j)}$ have the same bond dimension and are gauge-phase equivalent.*

Proof. [✓ Proof](#) For large N , Lemma 10.2.2 gives an invertible matrix U_N relating the two MPV families. Fix j and write $|V^{(N)}(B_j)\rangle = \sum_i u_{ij}^{(N)} |V^{(N)}(A_i)\rangle$. With $G_A(N) = (O_{A_i A_\ell}(N))_{i,\ell}$ and $b_j(N) = (O_{A_i B_j}(N))_i$, this gives

$$G_A(N) \overline{u_j(N)} = b_j(N), \quad O_{B_j B_j}(N) = u_j(N)^\top G_A(N) \overline{u_j(N)}.$$

If all mixed overlaps $O_{A_i B_j}(N)$ tended to 0, then $b_j(N) \rightarrow 0$; since $G_A(N) \rightarrow I$, one obtains $\overline{u_j(N)} \rightarrow 0$ and hence $u_j(N) \rightarrow 0$. Thus $O_{B_j B_j}(N) \rightarrow 0$, contradicting $O_{B_j B_j}(N) \rightarrow 1$. Thus each B_j has a matched block $A_{f(j)}$ with non-decaying mixed overlap. Theorems 7.5.3 and 7.6.4 then force equal bond dimension and gauge-phase equivalence for each match. Off-diagonal decay within the B -family makes f injective, hence bijective, and its inverse is the required permutation. $\square$

Lemma 10.2.4 (Permutation rigidity with asymptotically orthonormal overlaps). [Lean](#) [✓ Stmt](#) *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be two families of injective tensors of positive bond dimension satisfying the TP normalization $\sum_i (A_j^i)^\dagger A_j^i = \mathbb{1}$ and $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$, whose self-overlaps converge to 1 and whose cross-overlaps within each family converge to 0. If for every system size N ,*

$$\text{span}_{\mathbb{C}}\{|V^{(N)}(A_j)\rangle : 1 \leq j \leq g_A\} = \text{span}_{\mathbb{C}}\{|V^{(N)}(B_k)\rangle : 1 \leq k \leq g_B\},$$

then $g_A = g_B$, and there is a permutation π of the common block index set such that A_j and $B_{\pi(j)}$ have the same bond dimension and are gauge-phase equivalent for each j .

Proof. [✓ Proof](#) By Lemma 10.1.14, both families are linearly independent for all sufficiently large N . At such an N , the common span has the same dimension when computed from the A -family or the B -family, so $g_A = g_B$. After identifying the two index sets, Theorem 10.2.3 yields the permutation conclusion. $\square$

The two preceding results obtain the block matching from the asymptotic orthonormality of the overlaps. The same matching is obtained by the exact sector comparison of Section 10.6, which pairs the sectors of two canonical forms through a bijection fixed by their bond dimensions and gauge-phase equivalences. The proof of the Fundamental Theorem in Chapter 11 proceeds through the sector comparison.

The same overlap dichotomy also shows that the MPV states of distinct normal blocks are non-proportional at arbitrarily large lengths; this is an input to the exact sector matching later in the chapter.

Theorem 10.2.5 (Non-equivalent irreducible TP blocks are eventually non-proportional). [Lean](#)

[Lean](#) [✓ Stmt](#) *Let A and B be irreducible trace-preserving blocks whose self-overlaps converge to 1. If they are not gauge-phase equivalent, then for every lower bound N_0 there is $N \geq N_0$ such that $|V^{(N)}(A)\rangle$ is not proportional to $|V^{(N)}(B)\rangle$. Consequently, in an already separated same-dimensional BNT block family, any two distinct blocks have such non-proportional MPV states at arbitrarily large lengths.*

Proof. [✓ Proof](#) Suppose that for all sufficiently large N one had $|V^{(N)}(A)\rangle = c_N |V^{(N)}(B)\rangle$. Then

$$O_{AA}(N) = |c_N|^2 O_{BB}(N), \quad O_{AB}(N) = c_N O_{BB}(N). \quad (10.1)$$

Since $O_{AA}(N) \rightarrow 1$ and $O_{BB}(N) \rightarrow 1$, the first identity in (10.1) gives $|c_N| \rightarrow 1$, and hence $|O_{AB}(N)| \rightarrow 1$. The irreducible trace-preserving overlap decay theorem gives $O_{AB}(N) \rightarrow 0$ when $A \not\sim_{\text{gp}} B$, a contradiction. $\square$

10.3 Newton–Girard identities and power-sum recovery

The coefficients $c_N^{(j)} = \sum_q \mu_{j,q}^N$ that appear in the two-layer BNT expansion are power sums of the copy weights. Recovering the individual weights from these sums is a Newton–Girard problem: equal power sums determine equal characteristic polynomials, hence equal multisets of roots.

Theorem 10.3.1 (Newton–Girard trace recursion). [Lean](#) [Lean](#) [✓ Stmt](#) *If $A, B \in M_n(\mathbb{C})$ satisfy $\text{tr}(A^k) = \text{tr}(B^k)$ for all $k \geq 1$, then A and B have the same characteristic polynomial. It is enough to assume this equality for $1 \leq k \leq n$.*

Proof. [✓ Proof](#) Let e_m denote the m th elementary symmetric polynomial in the eigenvalues, with $e_0 = 1$. The Newton–Girard recursion

$$m e_m = \sum_{i=1}^m (-1)^{i-1} e_{m-i} \text{tr}(A^i)$$

expresses the coefficients of the characteristic polynomial in terms of the power-sum traces $\text{tr}(A^i)$. Equal power sums for A and B therefore give equal coefficients, hence equal characteristic polynomials. $\square$

The matched-sector comparison below applies the power-sum identities to a single matched pair, after the BNT permutation and phase gauges have identified which basis tensors are compared: the coefficient $\sum_q \mu_{j,q}^N$ then equals $\sum_q (\nu_{j,q} e^{i\phi_j})^N$, and Newton–Girard recovers the weight multiset with multiplicity.

Lemma 10.3.2 (Eventual vanishing of a finite geometric sum). [Lean](#) [✓ Stmt](#) *Let $w_1, \dots, w_n$ be nonzero complex numbers. If*

$$\sum_i c_i w_i^N = 0$$

for every sufficiently large N , then the same equality holds for every nonnegative exponent.

Proof. [✓ Proof](#) Induct on the number of terms. Subtracting w_1 times the equality at exponent N from the equality at exponent $N+1$ gives

$$\sum_{i=2}^n c_i (w_i - w_1) w_i^N = 0.$$

The induction hypothesis yields this equality at every exponent, so the original sum $S_N = \sum_i c_i w_i^N$ satisfies $S_{N+1} = w_1 S_N$ for every N . Since S_N eventually vanishes and $w_1 \neq 0$, it follows that $S_0 = 0$ and hence $S_N = 0$ for every N . $\square$

Lemma 10.3.3 (Eventual power-sum equality holds at every exponent). [Lean](#) [✓ Stmt](#) Let $a_1, \dots, a_m$ and $b_1, \dots, b_n$ be nonzero complex numbers. If their power sums agree for every sufficiently large exponent, then

$$\sum_i a_i^k = \sum_j b_j^k$$

for every nonnegative exponent k .

Proof. [✓ Proof](#) Set $w_l = a_l$ for $l \leq m$ and $w_{m+l} = b_l$ for $l \leq n$, with coefficients $c_l = 1$ for $l \leq m$ and $c_l = -1$ for $l > m$. Then

$$\sum_{l=1}^{m+n} c_l w_l^k = \sum_i a_i^k - \sum_j b_j^k.$$

The hypothesis makes this expression zero for every sufficiently large k . Lemma 10.3.2 makes it zero for every k , which is the required power-sum equality. $\square$

Theorem 10.3.4 (Finite-range equal power sums imply equal multisets). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $\alpha, \beta : \{0, \dots, n-1\} \rightarrow \mathbb{C}$. If $\sum_i \alpha_i^k = \sum_i \beta_i^k$ for $1 \leq k \leq n$, then the multisets $\{\alpha_i\}$ and $\{\beta_i\}$ are equal.

Two variants relax the equal-cardinality assumption. If $\alpha : \{0, \dots, m-1\} \rightarrow \mathbb{C}$ and $\beta : \{0, \dots, n-1\} \rightarrow \mathbb{C}$ have no zero entries and their power sums agree for $1 \leq k \leq \max(m, n)$, then $m = n$ and the multisets are equal. Without a nonzero hypothesis, the list form $\sum_{k=1}^{x_a} \lambda_{a,k}^N = \sum_{k=1}^{x_b} \lambda_{b,k}^N$ for $1 \leq N \leq \max(x_a, x_b)$ gives equality of the nonzero multisets; if it also holds at $N = 0$ then $x_a = x_b$ and the full lists agree.

Proof. [✓ Proof](#) Apply the finite-range Newton–Girard theorem to $\text{diag}(\alpha)$ and $\text{diag}(\beta)$ and extract the roots of $\prod_i (X - \alpha_i)$ and $\prod_i (X - \beta_i)$. For unequal cardinalities, pad both lists with zeros to length $\max(m, n)$; the nonzero hypothesis makes the padded-zero counts $\max(m, n) - m$ and $\max(m, n) - n$, forcing $m = n$. Without that hypothesis, delete the zero entries first (unchanged for positive N); the equality at $N = 0$ gives $x_a = x_b$. $\square$

Corollary 10.3.5 (A nonzero finite family has a nonvanishing power sum). [Lean](#) [✓ Stmt](#) Let $m \geq 1$ and let $\lambda_0, \dots, \lambda_{m-1} \in \mathbb{C}$ be nonzero. Then there is an exponent N with $1 \leq N \leq m$ such that

$$\sum_{q=0}^{m-1} \lambda_q^N \neq 0.$$

This is the consequence of the Appendix power-sum lemma used in the proof of Lemma L [CPGSV16, Appendix C.3, lines 1851–1858]; the finite-range comparison is [CPGSV16, Appendix A, lines 1155–1163].

Proof. [✓ Proof](#) Suppose that every power sum with $1 \leq N \leq m$ vanishes. Comparing $\lambda_0, \dots, \lambda_{m-1}$ with the empty family in Theorem 10.3.4 gives $m = 0$, contradicting $m \geq 1$. $\square$

Theorem 10.3.6 (Equal power sums imply equal multisets). [Lean](#) [✓ Stmt](#) Let $\alpha, \beta : \{0, \dots, n-1\} \rightarrow \mathbb{C}$. If $\sum_i \alpha_i^k = \sum_i \beta_i^k$ for all $k \geq 1$, then the multisets $\{\alpha_i\}$ and $\{\beta_i\}$ are equal.

Proof. [✓ Proof](#) Construct diagonal matrices $\text{diag}(\alpha)$ and $\text{diag}(\beta)$. The power-sum hypothesis gives $\text{tr}(\text{diag}(\alpha)^k) = \text{tr}(\text{diag}(\beta)^k)$. By Theorem 10.3.1, the characteristic polynomials agree: $\prod_i (X - \alpha_i) = \prod_i (X - \beta_i)$. Extracting roots gives multiset equality. $\square$

10.4 The BNT canonical form on a sector decomposition

This section introduces the BNT canonical-form predicate of [CPGSV16, Section 2], defined on a sector decomposition with raw sector weights $\mu_{j,q}$ and coefficients $c_N^{(j)} = \sum_q \mu_{j,q}^N$; no equal-modulus factorization is assumed.

10.4.1 Sector decompositions and phase matching

The sector decomposition and the MPV phase-equivalence predicates introduced here feed the BNT canonical-form predicate below.

Definition 10.4.1.1 (Sector weights and sector decomposition). [Lean](#) [Lean](#) [✓ Stmt](#) A sector-weight tuple over g basis blocks consists of multiplicities $r_j > 0$, nonzero weights $\mu_{j,q}$ for $q \in \{1, \dots, r_j\}$, and the sector coefficients

$$c_{S,j}^{(N)} := \sum_{q=1}^{r_j} \mu_{j,q}^N.$$

A *sector decomposition* consists of the basis blocks $(A_j)_{j=1}^g$ together with these multiplicities and weights.

Definition 10.4.1.2 (Physical blocking of a sector decomposition). [Lean](#) [✓ Stmt](#) Let P be a sector decomposition with representatives A_j , multiplicities r_j , and copy weights $\mu_{j,q}$. Its physical p -site blocking has the same representative and copy indices, with

$$A_j \mapsto A_j^{[p]}, \quad \mu_{j,q} \mapsto \mu_{j,q}^p.$$

The two-layer decomposition is [CPGSV16, lines 283–301]. Its blocked form is an auxiliary construction for the planned formal proof of Appendix C.3, Lemma L, lines 1835–1858; it is not stated there as a separate construction.

Theorem 10.4.1.3 (Indices, dimensions, weights, and representatives under blocking). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Physical blocking preserves the copy sets and bond dimensions. On the flattened copy index, the weight is raised to the p -th power and the representative is replaced by its p -site blocking.*

Proof. [✓ Proof](#) These identities follow directly from the definition. □

Theorem 10.4.1.4 (Blocked sector coefficients and assembled tensor). [Lean](#) [Lean](#) [✓ Stmt](#) *The coefficient of the blocked decomposition satisfies*

$$c_{P^{[p]},j}^{(N)} = \sum_q (\mu_{j,q}^p)^N = c_{P,j}^{(Np)}.$$

Moreover, blocking the tensor assembled from P gives the same MPV family as the tensor assembled from the blocked sector decomposition.

Proof. [✓ Proof](#) The coefficient identity is $(\mu^p)^N = \mu^{pN}$. The tensor identity is Theorem 9.7.4 applied to the flattened copy family. □

Lemma 10.4.1.5 (Sector decomposition formula). [Lean](#) [✓ Stmt](#) *For a sector decomposition S with associated tensor A_{tot} , the MPV satisfies*

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^g c_{S,j}^{(N)} V^{(N)}(A_j)_\sigma.$$

Two blocks are identified for canonical-form purposes when their MPV families differ by a scalar power. The next three definitions formalize this relation, the common cover it generates between two families, and the phase classes of a single family.

Definition 10.4.1.6 (MPV phase equivalence for nonzero-weight blocks). [Lean](#) [✓ Stmt](#) Two blocks A and B are MPV-phase equivalent if there is a nonzero scalar ζ such that, for every system size N and every physical word,

$$V^{(N)}(B)_\sigma = \zeta^N V^{(N)}(A)_\sigma.$$

The two bond dimensions are allowed to differ.

Definition 10.4.1.7 (Common MPV phase cover). [Lean](#) [✓ Stmt](#) For two finite nonzero-weight block families, a common MPV phase cover consists of a third finite family of blocks, a map from each nonzero-weight block family onto that common family, and MPV-phase equivalences between each block and its common representative. Both maps are required to be surjective, so the common family contains no unused representative.

Definition 10.4.1.8 (MPV phase classes). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For a finite family of blocks, put $j \sim k$ when there is a nonzero scalar factor ζ such that

$$|V^{(N)}(B_k)\rangle = \zeta^N |V^{(N)}(B_j)\rangle$$

for every length N . The associated tuple enumerates the finite quotient, chooses one representative per class, enumerates that representative first in its class, records the scalar relating the representative to each member, and proves that distinct representatives are not MPV-phase equivalent. The pairs (j, q) , with j a phase class and q a copy in that class, are in bijection with the original block indices.

Lemma 10.4.1.9 (Existence of an MPV phase class). [Lean](#) [✓ Stmt](#) A nonempty finite family of blocks has at least one MPV phase class.

Proof. [✓ Proof](#) Choose one original block index. Its occurrence in the phase-class enumeration supplies a phase class containing it. $\square$

Lemma 10.4.1.10 (Regrouping a matrix sum by MPV phase classes). [Lean](#) [✓ Stmt](#) Let F_k be a finite family of square matrices indexed by the original blocks. If $k = (j, q)$ enumerates the copies in the MPV phase classes, then

$$\sum_j \sum_q F_{j,q} = \sum_k F_k.$$

Proof. [✓ Proof](#) Evaluate both sides at a pair of matrix indices and apply the bijective enumeration of the finite phase-class quotient. $\square$

Definition 10.4.1.11 (BNT canonical form). [Lean](#) [✓ Stmt](#) Given a sector decomposition P , the BNT canonical-form predicate records the minimal BNT hypotheses of [CPGSV16, CPGSV21] without any equal-modulus or strict-ordering condition on the raw sector weights. It asserts that each basis block has positive bond dimension, is irreducible, and left-canonical; the normalized self-overlap of every basis block converges to 1; the basis MPV family is eventually linearly independent; distinct same-dimension basis blocks are not gauge-phase equivalent after identifying equal bond dimensions; and the [CPGSV16, Section 2.3] canonical-form normalization holds,

$$\forall (j, q), |\mu_{j,q}| \leq 1, \quad \exists (j_*, q_*) : |\mu_{j_*, q_*}| = 1.$$

The MPV expansion takes the raw two-layer form

$$V^{(N)}(P)_\sigma = \sum_{j=1}^g \left(\sum_{q=1}^{r_j} \mu_{j,q}^N \right) V^{(N)}(P_j)_\sigma,$$

matching [CPGSV16, Section 2.3] and [CPGSV21, Definition IV.2]. The canonical-form normalization admits every example of the source paper (in particular $C \oplus D$, $C \oplus (-C)$, $C \oplus (1/2)C$, and $C \oplus (-C) \oplus (1/2)C$) and is not derivable from the per-block normality hypotheses alone.

The matching arguments below read off three facts from a BNT canonical form: the raw coefficient identity, the cross-overlap decay between distinct basis blocks, and the combined-family linear independence.

Lemma 10.4.1.12 (Raw two-layer sector coefficient identity). [Lean](#) [✓ Stmt](#) *For every sector decomposition P , length N , and basis index j ,*

$$c_N^{(j)} = \sum_{q=1}^{r_j} \mu_{j,q}^N.$$

This is [CPGSV16, Section 2.3].

Proof. [✓ Proof](#) The claim follows from the definition of the sector coefficient. $\square$

Lemma 10.4.1.13 (Cross-overlap decay between distinct basis blocks). [Lean](#) [✓ Stmt](#) *If P is a BNT canonical form and $j \neq k$, then $O_{P_j P_k}(N) \rightarrow 0$.*

Proof. [✓ Proof](#) Case on the bond dimensions. If they differ, the dimension-mismatch overlap-decay theorem applies. If they agree, the basis-distinctness clause rules out gauge-phase equivalence, so the equal-dimension overlap-decay theorem applies. $\square$

Lemma 10.4.1.14 (Combined-family eventual linear independence). [Lean](#) [✓ Stmt](#) *Let P and Q each be a BNT canonical form, and suppose every mixed overlap $O_{P_j Q_k}(N)$ tends to 0. Then the union family*

$$\{|V^{(N)}(P_j)\rangle\}_{j=1}^{g_P} \cup \{|V^{(N)}(Q_k)\rangle\}_{k=1}^{g_Q}$$

is linearly independent for all sufficiently large N . This is [CPGSV16, Appendix A].

Proof. [✓ Proof](#) Feed the per-family normalized self-overlaps, the within-family cross-overlap decay (Lemma 10.4.1.13), and the inter-family decay hypothesis into the two-family overlap-orthonormality criterion (Lemma 10.1.17). $\square$

Lemma 10.4.1.15 (Unit-modulus weight witness from the canonical-form normalization). [Lean](#) [✓ Stmt](#) *Under the BNT canonical form, the [CPGSV16, Section 2.3] unit-modulus existential provides indices (j_*, q_*) with $|\mu_{j_*, q_*}| = 1$.*

Proof. [✓ Proof](#) The canonical-form normalization includes a unit-modulus weight $|\mu_{j_*, q_*}| = 1$. $\square$

Example 10.4.1.16 ($C \oplus (1/2)C$, unequal-modulus admissibility). [Lean](#) [✓ Stmt](#) The single-sector two-copy decomposition with raw weights $(1, 1/2)$ is a BNT canonical form, so unequal-modulus copies are admissible: $|1/2| \leq 1$ and the unit witness is the first copy.

10.4.2 Prepared-block construction of BNT canonical forms

The construction below separates the after-blocking preparation of suitable blocks from the normalized-weight BNT construction: the prepared blocks are trace-preserving, primitive, irreducible, and injective; normalized weights then quotient gauge-phase-equivalent blocks into sectors to produce the canonical form of Definition 10.4.1.11.

Lemma 10.4.2.1 (Common injective blocking for primitive irreducible families). Lean ✓ Stmt *Let $(B_k)_{k=1}^r$ be a finite family of positive-bond-dimension MPS tensors, each trace-preserving, primitive, and tensor-irreducible. Then there is a single $L \geq 1$ such that each $B_k^{[L]}$ is one-site injective, and this blocking preserves trace preservation, primitivity, and irreducibility for every block.*

Proof. ✓ Proof For each block, Theorem 8.12.1 gives a positive length at which it becomes one-site injective; the product of these lengths is a common length. Injectivity persists under positive further blocking, and Theorem 9.6.3 preserves the other hypotheses. $\square$

Theorem 10.4.2.2 (Prepared BNT blocks after blocking). Lean ✓ Stmt *Let A be an MPS tensor. Then there is a blocking length $p \geq 1$, nonzero weights μ_k , and blocks B_k over the p -blocked alphabet such that:*

1. *each block has positive bond dimension;*
2. *each block is trace-preserving, primitive, tensor-irreducible, and one-site injective;*
3. *$A^{[p]}$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family at every positive length.*

No weight-modulus normalization is asserted at this stage.

Proof. ✓ Proof Apply the common primitive irreducible block decomposition to (A, A) , giving a positive blocking length and a nonzero weighted family of trace-preserving primitive irreducible blocks with the positive-length MPV identity. Lemma 10.4.2.1 supplies one common further blocking making every block one-site injective while preserving the other hypotheses; finally identify the iterated blocked alphabet with the direct one. $\square$

With the prepared blocks in hand, normalized weights produce a BNT canonical form by quotienting gauge-phase-equivalent blocks into sectors.

Theorem 10.4.2.3 (BNT from normal blocks with normalized weights). Lean ✓ Stmt *Let $r \in \mathbb{N}$, and for each $k = 1, \dots, r$ let $D_k \in \mathbb{N}$, $\mu_k \in \mathbb{C}$, and let B_k be an MPS tensor of physical dimension d and bond dimension D_k . Assume:*

1. **positive bond dimensions:** $D_k > 0$ for all k ,
2. **TP normalization:** $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$,
3. **primitivity:** $\mathcal{E}_{B_k}$ is primitive for each k ,
4. **irreducibility:** each B_k is irreducible,
5. **nonzero weights:** $\mu_k \neq 0$ for all k ,
6. **bounded weights:** $|\mu_k| \leq 1$ for all k ,
7. **global unit witness:** $|\mu_k| = 1$ for some k .

Then there exists a sector decomposition P such that P and $\bigoplus_{k=1}^r \mu_k B_k$ generate the same MPV family at every length, and P satisfies the basis-of-normal-tensors conditions (Definition 10.1.2) together with the multiplicity, span, and weight-bound conditions of Definition 10.4.1.11.

Proof. ✓ Proof The phase-class quotient groups gauge-phase-equivalent blocks into sectors with unit-modulus per-copy scalars. Under the TP, primitive, and irreducible hypotheses the phase-class construction applies, and for distinct representatives $B_j, B_{j'}$ ($j \neq j'$) the cross-overlap decays,

$$\lim_{N \rightarrow \infty} O_{B_j B_{j'}}(N) = 0, \quad \lim_{N \rightarrow \infty} O_{B_j B_j}(N) = 1,$$

which gives the eventual linear-independence condition with no injectivity required. The resulting sector decomposition inherits the span and multiplicity conditions, and the bounded-weight and global-unit hypotheses give the weight-norm conditions. $\square$

The BNT construction preserves total bond dimension because CF summands assigned to the same BNT representative have equal bond dimension. Indeed, suppose trace-preserving primitive irreducible NTs X and Y of positive bond dimension have MPV families related by $|V^{(N)}(Y)\rangle = \zeta^N |V^{(N)}(X)\rangle$ with $\zeta \neq 0$. The normalized self-overlaps give $|O_{XX}(N)| \rightarrow 1$. The phase relation gives the cross-overlap identity

$$O_{YX}(N) = \zeta^N O_{XX}(N), \quad |O_{YX}(N)| = |\zeta|^N |O_{XX}(N)|.$$

Since the same normalization forces $|\zeta| = 1$, this gives $|O_{YX}(N)| \rightarrow 1$. If their bond dimensions differed, the rectangular overlap-decay theorem would give

$$O_{YX}(N) \rightarrow 0,$$

a contradiction.

Lemma 10.4.2.4 (Total dimension after grouping by BNT representatives). Lean ✓ Stmt
Suppose CF summands are grouped by BNT representative, every original copy weight is nonzero, and every summand assigned to a representative has that representative's bond dimension. Then the total bond dimension of the resulting BNT sector decomposition is the sum of the original block dimensions.

Proof. ✓ Proof The collapsed decomposition counts one representative dimension for each assigned copy. Let $e(j, q)$ denote the original CF summand assigned to the q th copy of representative P_j . Since $(j, q) \mapsto e(j, q)$ reindexes the original summands and each assigned summand has the representative's bond dimension,

$$\sum_{j=1}^g \sum_{q=1}^{r_j} \dim(P_j) = \sum_{j=1}^g \sum_{q=1}^{r_j} D_{e(j, q)} = \sum_{k=1}^r D_k.$$

This is the equality between the collapsed total dimension and the original direct-sum dimension. $\square$

Lemma 10.4.2.5 (Prepared phase-equivalent NTs have the same bond dimension). Lean ✓ Stmt
Let X and Y be trace-preserving, primitive, irreducible NTs of positive bond dimension. If their MPV families differ by a nonzero scalar power, then X and Y have the same bond dimension.

Proof. [✓ Proof](#) Choose $\zeta \neq 0$ with $|V^{(N)}(Y)\rangle = \zeta^N |V^{(N)}(X)\rangle$ for every length N . The normalized self-overlaps give $|O_{XX}(N)| \rightarrow 1$, and the phase relation gives

$$O_{YX}(N) = \zeta^N O_{XX}(N), \quad |O_{YX}(N)| = |\zeta|^N |O_{XX}(N)|.$$

Since the same normalization forces $|\zeta| = 1$, this gives $|O_{YX}(N)| \rightarrow 1$. If the two bond dimensions differed, the rectangular overlap-decay theorem would force

$$O_{YX}(N) \rightarrow 0,$$

contradicting the preceding limit. □

Theorem 10.4.2.6 (Gauge realization of prepared phase-equivalent NTs). [Lean](#) [✓ Stmt](#) *Let X and Y be trace-preserving, primitive, irreducible NTs of positive bond dimensions D_X and D_Y . If their MPV families differ by a nonzero scalar power, then $D_X = D_Y$, and after identifying the bond spaces there exist $G \in \text{GL}_{D_Y}(\mathbb{C})$ and $\zeta \in \mathbb{C} \setminus \{0\}$ such that, for every physical index i ,*

$$Y^i = \zeta G X^i G^{-1}.$$

The scalar is not asserted to be positive, and the gauge is not asserted to be unitary.

Proof. [✓ Proof](#) Lemma 10.4.2.5 identifies the bond dimensions. If no gauge-phase relation existed after this identification, Theorem 10.2.5 would give a length N at which the two MPV states are not proportional. This contradicts the assumed scalar-power identity at every length. □

Lemma 10.4.2.7 (Prepared BNT representative classes preserve bond dimension). [Lean](#) [✓ Stmt](#) *In a prepared family of positive-bond-dimension trace-preserving primitive irreducible NTs, every CF summand assigned to a BNT representative has that representative's bond dimension.*

Proof. [✓ Proof](#) Apply Lemma 10.4.2.5 to each representative and each assigned summand. □

Lemma 10.4.2.8 (Prepared collapsed BNT total dimension). [Lean](#) [✓ Stmt](#) *For prepared trace-preserving primitive irreducible NTs of positive bond dimension with all copy weights nonzero, grouping CF summands by BNT representative preserves the total bond dimension: the collapsed sector decomposition has total bond dimension equal to the sum of the original block dimensions.*

Proof. [✓ Proof](#) Every assigned summand shares the bond dimension of its representative, by the equal-dimension fact above. The displayed regrouping identity in Lemma 10.4.2.4 then identifies the collapsed sum over pairs (j, q) with $\sum_{k=1}^r D_k$, the original direct-sum dimension. □

Combining the after-blocking preparation with the normalized-weight construction gives the arbitrary-input statement, conditional on the weight normalization.

Theorem 10.4.2.9 (Conditional BNT form after blocking for an arbitrary tensor). [Lean](#) [Lean](#) [✓ Stmt](#) *Let A be an MPS tensor. Then there exists a blocking length $p \geq 1$, nonzero weights μ_k , and blocks B_k over the blocked alphabet, such that:*

1. every block has positive bond dimension;
2. every block is TP-normalized, primitive, irreducible, and injective;
3. $A^{[p]}$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family at every positive length.

Moreover, under $|\mu_k| \leq 1$ for every k and $|\mu_k| = 1$ for at least one k , there exists a sector decomposition P over the blocked alphabet in the BNT canonical form of Definition 10.4.1.11 such that $A^{[p]}$ and P generate the same MPV family at every positive length; the identity upgrades to all lengths once the total bond dimension of P equals that of $A^{[p]}$.

Proof. ✓ Proof The after-blocking reduction supplies the TP-normalized primitive irreducible injective blocks and the positive-length MPV identity. The normalized-weight hypotheses are the choice that [CPGSV16, Section 2.3] makes; Theorem 10.4.2.3 then produces the sector decomposition, following the BNT selection of [CPGSV16, Appendix A]. At length zero the MPV coefficient is the bond dimension, so equal total bond dimensions supply the missing length-zero coefficient. The remaining rescaling step is recorded in Remark 10.4.2.10. $\square$

Remark 10.4.2.10 (Arbitrary-input reduction). The reduction is conditional on the two weight-normalization hypotheses of Theorem 10.4.2.9.

10.5 Coefficient comparison and copy-weight recovery

The exact matching of Section 10.6 produces, on each matched pair, a coefficient identity. This section turns that identity into equality of the per-copy weight multisets [CPGSV16, Appendix A]. No analytic non-decay estimate enters: the fixed-length linear independence isolates each sector coefficient exactly.

10.5.1 Matched-sector weight multiset equality

Given a matched pair $(j_0, \tilde{k}_0)$ and a nonzero scalar ζ with $c_N^{(j_0)}(P) = \zeta^N c_N^{(\tilde{k}_0)}(Q)$ past a threshold N_0 , this section records two conclusions: the multiset equality of the per-copy weights from power-sum recovery (Theorem 10.3.6), and its upgrade to an explicit per-copy permutation. Together they give the $\mu_{j,q} = \nu_{j,q} e^{i\varphi_j}$ recovery of [CPGSV16, Appendix A]. The coefficient identity is supplied, in the equal-MPV case, by Lemma 11.1.3 of Chapter 11; in the proportional case it is the remaining hypothesis of the conditional proportional global gauge (Theorem 11.1.13).

Lemma 10.5.1.1 (Matched-sector weight multiset equality). Lean ✓ Stmt *Let P, Q be sector decompositions with weights $\mu_{j,q}$ and $\nu_{k,q'}$, and fix indices $j_0, \tilde{k}_0$, and a nonzero scalar $\zeta \in \mathbb{C} \setminus \{0\}$. Suppose that for every $N > N_0$,*

$$c_N^{(j_0)}(P) = \zeta^N \cdot c_N^{(\tilde{k}_0)}(Q). \quad (10.2)$$

Then the per-sector multiplicities agree, $r_{j_0}(P) = r_{\tilde{k}_0}(Q)$, and the rescaled per-copy weight multisets coincide. This is [CPGSV16, Appendix A] read on a single matched sector.

Proof. ✓ Proof Expand both sides of the coefficient identity (10.2) into per-copy power sums $\sum_q \mu_{j_0,q}^N$ and $\sum_{q'} \nu_{\tilde{k}_0,q'}^N$, and distribute ζ^N through the right-hand sum to obtain, for every $N > N_0$,

$$\sum_q \mu_{j_0,q}^N = \sum_{q'} (\zeta \cdot \nu_{\tilde{k}_0,q'})^N.$$

Extend to all positive exponents and apply the unequal-cardinality multiset-recovery lemma (Theorem 10.3.4), using $\zeta \neq 0$ together with the weight non-vanishing assumption to ensure the input weights are nonzero. This yields the multiplicity match and the multiset equality. $\square$

Lemma 10.5.1.2 (Matched-sector weight-permutation extractor). [Lean](#) [✓ Stmt](#) Under the hypotheses of Lemma 10.5.1.1, the per-sector multiplicities agree and there is a permutation τ of the per-copy index set identifying the individual weights up to the gauge phase: for every q ,

$$\nu_{k_0, \tau(q)} = \zeta^{-1} \cdot \mu_{j_0, q}.$$

This is the per-copy realisation of [CPGSV16, Appendix A]: the multiset equality from Lemma 10.5.1.1 is upgraded to a concrete indexing between the matched P - and Q -copies.

Proof. [✓ Proof](#) Invoke Lemma 10.5.1.1 to obtain the cardinality match and the multiset equality. From this multiset-map equality extract a permutation matching the elements pointwise (proved by induction on the cardinality). Compose with the cardinality identification and cancel ζ via $\zeta \neq 0$ to convert the gauge factor on the right of the pointwise identity to ζ^{-1} on the left. $\square$

10.6 Strong existential matching by exact linear independence

The exact matching reads [CPGSV16, Appendix A] as a statement at fixed length: equal MPV families force each basis sector of one form to pair with a basis sector of the other. The non-vanishing of the sector coefficients isolates the matched sector exactly, with no limit of a single-sector projection.

Lemma 10.6.1 (Sector coefficient is not eventually zero). [Lean](#) [✓ Stmt](#) For a BNT canonical form P and any sector j , the sector coefficient $c_{P,j}^{(N)} = \sum_q \mu_{j,q}^N$ is not eventually zero in N . Only nonvanishing of the copy weights is used, not any modulus condition.

Proof. [✓ Proof](#) If $c_{P,j}^{(N)} = 0$ for all $N \geq M$, the signed geometric-sequence vanishing lemma forces $\sum_q \mu_{j,q}^k = 0$ at every exponent k , in particular $\sum_q 1 = r_j = 0$ at $k = 0$, contradicting $r_j > 0$. Repeated ratios are handled by the same extrapolation, so distinct weights are not needed. $\square$

Theorem 10.6.2 (Exact single-sector matching). [Lean](#) [✓ Stmt](#) Let P, Q be two BNT canonical forms with the same MPV family (at all positive lengths). For every sector j of P there exist a sector k of Q with matched bond dimension $\dim_P(j) = \dim_Q(k)$, a gauge-phase equivalence after identifying the matched bond dimensions, and a non-decaying cross-overlap. No per-sector unit-modulus hypothesis is required.

Proof. [✓ Proof](#) Suppose j decays against every sector of Q , and let T be the set of P -sectors with this property. For $j' \notin T$ the overlap dichotomy turns a non-decaying overlap with some $Q_{k(j')}$ into a gauge-phase equivalence, hence a scalar $\omega_{j'} \in \mathbb{C}^\times$ with $V^{(N)}(P_{j'}) = \omega_{j'}^N V^{(N)}(Q_{k(j')})$, so $V^{(N)}(P_{j'})$ lies in the span of $\{V^{(N)}(Q_k)\}_k$. The family $\{P_{j'} : j' \in T\} \cup \{Q_k\}_k$ then has all pairwise cross-overlaps decaying, hence is eventually linearly independent (Lemma 10.4.1.14). Rewriting the common-MPV identity over this family,

$$\sum_{j' \in T} c_N^{(j')} V^{(N)}(P_{j'}) = \sum_k \left(c_N^{(k)} - \sum_{j' \notin T, k(j')=k} c_N^{(j')} \omega_{j'}^N \right) V^{(N)}(Q_k),$$

exact coefficient extraction forces $c_N^{(j)} = 0$ at each fixed large N , contradicting Lemma 10.6.1. Hence a matched sector k exists; the gauge-phase equivalence is recovered from the non-decaying overlap by the irreducible trace-preserving overlap dichotomy. $\square$

Theorem 10.6.3 (Strong existential matching, full-basis form). Lean ✓ Stmt *Let P, Q be two BNT canonical forms with the same MPV family. Then for every sector k of Q there exist a sector j of P with matched bond dimension $\dim_P(j) = \dim_Q(k)$, a gauge-phase equivalence after identifying the matched bond dimensions, and a non-decaying cross-overlap.*

Proof. ✓ Proof Apply Theorem 10.6.2 with the roles of P and Q swapped and the same-MPV hypothesis flipped by symmetry of gauge-phase equivalence and overlap convergence. $\square$

10.6.1 Bijective matching by symmetry

Applying the exact matching once more with P and Q swapped gives the reverse direction; the basis-distinctness clause forces injectivity both ways, so $g_a = g_b$ and the source relabelling $B_k = e^{i\phi_k} X_k A_{j_k} X_k^{-1}$ holds on the full basis [CPGSV16, Appendix A].

Theorem 10.6.1.1 (Bijective matching by symmetry). Lean ✓ Stmt *Let P, Q be two BNT canonical forms with the same MPV family. Then there is a bijection*

$$\beta : J(Q) \simeq J(P)$$

such that for every $k \in J(Q)$ the bond dimensions match, the basis tensors $P_{\beta(k)}$ and Q_k are gauge-phase equivalent after identifying the matched bond dimensions, and the basis cross-overlap does not tend to zero.

Proof. ✓ Proof Theorem 10.6.2 applied to (Q, P) gives a matching map $m : J(Q) \rightarrow J(P)$, and applied to (P, Q) a matching map $J(P) \rightarrow J(Q)$.

Injectivity of m . If $m(k_1) = m(k_2) = j$, the two witnesses give a common centre P_j gauge-phase equivalent, after identifying bond dimensions, to both Q_{k_1} and Q_{k_2} ; transitivity yields a gauge-phase equivalence $Q_{k_1} \sim Q_{k_2}$ under $\dim_Q(k_1) = \dim_Q(k_2)$. The basis-distinctness clause on Q then forces $k_1 = k_2$; the reverse map is injective by symmetry.

Bijection. The two injections give equal finite sector counts, so m is a bijection $\beta : J(Q) \simeq J(P)$. $\square$

Chapter 11

Proof of the Fundamental Theorem

For BNT canonical forms P and Q , write the weighted direct sums over normal-tensor sectors $P^i = \bigoplus_j \mu_j P_j^i$ and $Q^i = \bigoplus_k \nu_k Q_k^i$. The proportional theorem gives a bijection β of the normal-tensor sectors with unit-modulus phases ζ_k , invertible matrices X_k , and $Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$; the equal-MPV corollary collapses the per-block phases into a single X with $Q_{\text{tot}}^i = X P_{\text{tot}}^i X^{-1}$. Given the sector matching of Chapter 10, extract the per-sector coefficient identities $c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$ from eventual linear independence, recover the copy weights by finite power-sum comparison, and combine the block gauges into $X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k)$.

Throughout, “eventually” abbreviates “for all sufficiently large N ”: there is N_0 such that the stated relation holds for every $N > N_0$. The comparison is made at a fixed such N , where BNT linear independence detects a vanishing coefficient even when $|\mu| < 1$ lets it decay.

11.1 Coefficient identity, copy weights, and the block-diagonal gauge

The bijective matching theorem of Chapter 10 (Theorem 10.6.1.1) gives the sector phases ζ_k and block gauges X_k . The coefficient comparison of [CPGSV16, Appendix MPV proof, lines 1187–1188] then turns the equality of total MPV families into the per-sector coefficient identity $c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$. Neither step uses a per-sector unit-modulus hypothesis.

Lemma 11.1.1 (Unit phase from matched normalized BNT blocks). [Lean] [✓ Stmt] *Let P and Q be BNT canonical forms. If, for every length N and word σ , a sector P_j and a sector Q_k have MPV families related by*

$$V^{(N)}(Q_k)_\sigma = \zeta^N V^{(N)}(P_j)_\sigma \quad (11.1)$$

then $|\zeta| = 1$.

Proof. [✓ Proof] The BNT normalization gives self-overlap limits of modulus 1 for both sectors. The MPV identity (11.1) gives, for every N ,

$$O_{Q_k Q_k}(N) = |\zeta|^{2N} O_{P_j P_j}(N).$$

Since $O_{Q_k Q_k}(N) \rightarrow 1$ and $O_{P_j P_j}(N) \rightarrow 1$, taking limits forces $|\zeta|^{2N} \rightarrow 1$ and hence $|\zeta| = 1$. $\square$

Theorem 11.1.2 (Proportional sector matching with unit phases). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms whose total tensors generate eventually nonzero proportional MPV families. Then there are a bijection $\beta : J(Q) \simeq J(P)$, bond-dimension equalities $\dim P_{\beta(k)} = \dim Q_k$, scalars $\zeta_k \in \mathbb{C}$, and block gauges X_k such that $|\zeta_k| = 1$ and, for every sector k and physical index i ,*

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1} \quad (11.2)$$

Equivalently, for every length N and word σ ,

$$V^{(N)}(Q_k)_\sigma = \zeta_k^N V^{(N)}(P_{\beta(k)})_\sigma.$$

This is [CPGSV16, Appendix MPV theorem statement, lines 1167–1170, and proof line 1182].

Proof. [✓ Proof](#) Apply the proportional sector-matching theorem to obtain the bijection and dimension-compatible gauge-phase equivalences. Choosing the gauges and scalars gives $Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$, and therefore

$$O_{Q_k Q_k}(N) = |\zeta_k|^{2N} O_{P_{\beta(k)} P_{\beta(k)}}(N).$$

The normalized self-overlap limits give $O_{Q_k Q_k}(N) \rightarrow 1$ and $O_{P_{\beta(k)} P_{\beta(k)}}(N) \rightarrow 1$, so $|\zeta_k|^{2N} \rightarrow 1$ and hence $|\zeta_k| = 1$. The MPV scalar-power identity follows from the same gauge equation (11.2): $V^{(N)}(Q_k)_\sigma = \zeta_k^N V^{(N)}(P_{\beta(k)})_\sigma$. $\square$

Lemma 11.1.3 (Full-basis coefficient identity from matched sectors). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms with the same MPV family, matched by a bijection $\beta : J(Q) \simeq J(P)$ with bond-dimension equalities and a gauge-phase equivalence between Q_k and $P_{\beta(k)}$ for every k . Then each sector carries a unit-modulus phase ζ_k , and eventually*

$$c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q). \quad (11.3)$$

This is the equal-MPV coefficient comparison of [CPGSV16, Appendix MPV proof, lines 1187–1188].

Proof. [✓ Proof](#) For each matched sector, the gauge-phase equivalence gives $V^{(N)}(Q_k)_\sigma = \zeta_k^N V^{(N)}(P_{\beta(k)})_\sigma$. Lemma 11.1.1 gives $|\zeta_k| = 1$. Applying Lemma 11.1.4 to the same phases gives $c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$ for all sufficiently large N . $\square$

Lemma 11.1.4 (Full-basis coefficient identity from matched phases). [Lean](#) [✓ Stmt](#) *Let P be a BNT canonical form, let Q be a sector decomposition, and assume P_{tot} and Q_{tot} have the same MPV family. Suppose that there is a bijection*

$$\beta : J(Q) \simeq J(P)$$

and scalars $\zeta_k \in \mathbb{C}$ such that, for every $k \in J(Q)$, every length N , and every spin configuration σ of length N ,

$$V^{(N)}(Q_k)_\sigma = \zeta_k^N V^{(N)}(P_{\beta(k)})_\sigma. \quad (11.4)$$

Then for every $k \in J(Q)$ there is N_0 such that, for all $N > N_0$,

$$c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q).$$

This is the coefficient-comparison step of [CPGSV16, Appendix MPV proof, lines 1187–1188] in full-basis form.

Proof. ✓ Proof For every N , expand the equality of total MPVs as

$$\sum_{j \in J(P)} c_N^{(j)}(P) |V^{(N)}(P_j)\rangle = \sum_{k \in J(Q)} c_N^{(k)}(Q) |V^{(N)}(Q_k)\rangle.$$

The phase relation (11.4) gives $|V^{(N)}(Q_k)\rangle = \zeta_k^N |V^{(N)}(P_{\beta(k)})\rangle$, so reindexing the Q -sum by β onto the P -basis yields

$$\sum_{j \in J(P)} c_N^{(j)}(P) |V^{(N)}(P_j)\rangle = \sum_{j \in J(P)} \zeta_{\beta^{-1}(j)}^N c_N^{(\beta^{-1}(j))}(Q) |V^{(N)}(P_j)\rangle.$$

For all sufficiently large N the P -basis MPV family $\{|V^{(N)}(P_j)\rangle\}_{j \in J(P)}$ is linearly independent, by the BNT eventual linear independence of P . Lemma 10.1.16 then extracts the coefficient equalities $c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$ from this eventual linear independence, for all sufficiently large N . $\square$

The coefficient identity (11.3) is now converted, sector by sector, into a matching of the individual copy weights.

Definition 11.1.5 (Copy-weight matching for matched BNT sectors). Lean ✓ Stmt Let the sectors of two decompositions P and Q be matched by $\beta : J(Q) \simeq J(P)$, with sector phases ζ_k . A copy-weight matching consists of copy permutations τ_k such that, for all sectors k and copies q ,

$$\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}.$$

This is the copy-level conclusion of [CPGSV16, Appendix MPV proof, line 1188].

Lemma 11.1.6 (Copy-weight matching from matched-sector coefficient identities). Lean ✓ Stmt Let the sectors of P and Q be matched by $\beta : J(Q) \simeq J(P)$, with nonzero sector phases ζ_k . If the matched-sector coefficient identities

$$c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q) \tag{11.5}$$

hold eventually, then the phases ζ_k determine a copy-weight matching $\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}$. This is, sector by sector, [CPGSV16, Appendix MPV proof, line 1188].

Proof. ✓ Proof Eventually the coefficient identity (11.5) in sector k reads

$$\sum_r \mu_{\beta(k), r}^N = \zeta_k^N \sum_q \nu_{k,q}^N = \sum_q (\zeta_k \nu_{k,q})^N.$$

Newton–Girard finite power-sum comparison (Lemma 10.5.1.2) applies sector by sector, producing a copy permutation τ_k with $\mu_{\beta(k), \tau_k(q)} = \zeta_k \nu_{k,q}$; since $\zeta_k \neq 0$ this is $\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}$. $\square$

The matched block gauges and copy weights now combine into a single block-diagonal gauge.

Definition 11.1.7 (Flattened block-diagonal gauge). Lean Lean ✓ Stmt Given per-block gauges $X_k \in \text{GL}_{D_k}(\mathbb{C})$, form the block-diagonal element $\bigoplus_k X_k$ and then reindex the dependent block-coordinate space with the standard bond index $\{0, \dots, \sum_k D_k - 1\}$ of the assembled tensor. The result is an element of $\text{GL}_{\sum_k D_k}(\mathbb{C})$.

Lemma 11.1.8 (Unitary block-diagonal gauges). [Lean](#) [Lean](#) [✓ Stmt](#) A unitary matrix U_k defines an invertible block gauge with inverse $U_k^\dagger$. If every block gauge is unitary, then the flattened block-diagonal gauge

$$U = \bigoplus_k U_k$$

is unitary.

Proof. [✓ Proof](#) The products are computed blockwise:

$$UU^\dagger = \bigoplus_k U_k U_k^\dagger = \bigoplus_k \mathbb{1} = \mathbb{1}.$$

□

Lemma 11.1.9 (Direct-sum conjugation by the flattened gauge). [Lean](#) [Lean](#) [✓ Stmt](#) If, for every block k and physical index i , $B_k^i = X_k A_k^i X_k^{-1}$, then the weighted direct sums satisfy

$$\bigoplus_k \mu_k B_k^i = X \left(\bigoplus_k \mu_k A_k^i \right) X^{-1},$$

where X is the flattened block-diagonal gauge from Definition 11.1.7. Hence the two assembled tensors are gauge equivalent.

Proof. [✓ Proof](#) Multiplication of block-diagonal matrices is computed blockwise. The scalar weight μ_k commutes with the conjugation on the k th block, and the reindexing from dependent block coordinates to the flattened bond index preserves products and inverses. □

Theorem 11.1.10 (Global gauge from matched sectors and copy weights). [Lean](#) [✓ Stmt](#) Suppose that the sectors of two sector decompositions P and Q are matched by a bijection $\beta : J(Q) \simeq J(P)$, with bond-dimension equalities, copy permutations τ_k , nonzero phases ζ_k , and block gauges X_k . If, for every sector k , copy q , and physical index i ,

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}, \quad \nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}, \quad (11.6)$$

then, in the flattened Q -coordinates,

$$Q_{\text{flat}}^i = X P_{\text{matched}}^i X^{-1}, \quad X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k).$$

This is the direct-sum gauge construction of [CPGSV16, Appendix MPV proof, lines 1189–1192], separated from the coefficient-comparison step that supplies the weight identities.

Proof. [✓ Proof](#) For each matched copy,

$$\nu_{k,q} Q_k^i = (\zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}) (\zeta_k X_k P_{\beta(k)}^i X_k^{-1}) = \mu_{\beta(k), \tau_k(q)} X_k P_{\beta(k)}^i X_k^{-1}.$$

Thus every flattened weighted block is conjugated by its sector gauge X_k . Lemma 11.1.9 combines these block conjugacies into the direct-sum gauge $X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k)$. □

Theorem 11.1.11 (Global gauge from a copy-weight matching). [Lean](#) [✓ Stmt](#) Suppose that the sectors of P and Q are matched, with bond-dimension equalities, nonzero phases, and block gauges satisfying

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}.$$

If the same phases also give a copy-weight matching, then the flattened Q -tensor is obtained from the matched-coordinate P -tensor by the direct-sum gauge

$$X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k).$$

This is only a reformulation of Theorem 11.1.10.

Proof. ✓ Proof The copy-weight matching supplies $\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}$. Together with $Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$, these are exactly the two displayed hypotheses (11.6) of Theorem 11.1.10. $\square$

We can now state the proportional global gauge, first from a copy-weight matching and then from the coefficient identities that produce one.

Theorem 11.1.12 (Conditional proportional global gauge from a copy-weight matching). Lean

✓ Stmt Let P and Q be BNT canonical forms whose total tensors generate eventually nonzero proportional MPV families. Then the proportional sector-matching theorem gives β , the bond-dimension equalities, the unit phases ζ_k , and the block gauges X_k . If the remaining proportional coefficient comparison supplies a copy-weight matching for these same phases, then the flattened Q -tensor is obtained from the matched-coordinate P -tensor by the direct-sum gauge

$$X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k).$$

Proof. ✓ Proof The proportional sector-matching theorem supplies β , X_k , and unit phases ζ_k satisfying $Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$. Since $|\zeta_k| = 1$, each ζ_k is nonzero. A copy-weight matching gives $\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}$, so Theorem 11.1.11 gives the direct-sum gauge. $\square$

Theorem 11.1.13 (Conditional proportional global gauge from coefficient identities). Lean

✓ Stmt Let P and Q be BNT canonical forms whose total tensors generate eventually nonzero proportional MPV families. Then there are a bijection $\beta : J(Q) \simeq J(P)$, bond-dimension equalities, unit-modulus phases ζ_k , and block gauges X_k such that

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$$

for every sector k and physical index i . If these same phases satisfy the matched-sector coefficient identities

$$c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$$

for all sufficiently large N , then they determine a copy-weight matching and hence the flattened Q -tensor is obtained from the matched-coordinate P -tensor by the direct-sum gauge

$$X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k).$$

Proof. ✓ Proof The sector-matching theorem gives β , X_k , and $|\zeta_k| = 1$ with $Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}$. Since $\zeta_k \neq 0$, the coefficient identities $c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q)$ produce $\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}$ by Lemma 11.1.6. Theorem 11.1.12 then applies to this copy-weight matching. $\square$

The coefficient-identity hypothesis of Theorem 11.1.13 is the scalar-free identity (11.3). From the proportional MPV hypothesis alone one obtains the same identity carrying the global length-scalar of the proportionality, recorded next.

Lemma 11.1.14 (Proportional matched-phase coefficient identity, scalar-threaded). [Lean](#) [✓ Stmt](#)

Let P be a BNT canonical form and Q a sector decomposition whose total tensors generate eventually nonzero proportional MPV families. Suppose a bijection $\beta : J(Q) \simeq J(P)$ and scalars ζ_k satisfy the matched MPV phase identities (11.4) for every sector k . Then there is a single eventually-nonzero scalar sequence c_N and a threshold N_0 such that, for all $N > N_0$ and every $k \in J(Q)$,

$$c_N^{(\beta(k))}(P) = c_N \zeta_k^N c_N^{(k)}(Q). \quad (11.7)$$

The scalar c_N is the same for every sector k : it is the single global proportionality scalar of [CPGSV16, Theorem `thm1`, lines 1167–1170] at length N .

Proof. [✓ Proof](#) For each N on the proportionality tail, expand both total MPVs in their sector bases and use the global proportionality scalar c_N :

$$\sum_{j \in J(P)} c_N^{(j)}(P) |V^{(N)}(P_j)\rangle = c_N \sum_{k \in J(Q)} c_N^{(k)}(Q) |V^{(N)}(Q_k)\rangle.$$

The matched phase identities (11.4) give $|V^{(N)}(Q_k)\rangle = \zeta_k^N |V^{(N)}(P_{\beta(k)})\rangle$; because every sector is matched, reindexing the Q -sum by β rewrites the right-hand side entirely in the P -basis. The BNT eventual linear independence of the P -basis and Lemma 10.1.16 then extract the coefficient identities (11.7) for all sufficiently large N . $\square$

Remark 11.1.15 (Residual length-scalar control). The identity (11.7) discharges the hypothesis of Theorem 11.1.13 — equivalently (11.3) — precisely when $c_N \equiv 1$, which holds if and only if $c_N = \xi^N$ for a single fixed phase ξ (the power ξ^N is then absorbed into the sector phases ζ_k). The gauge-phase conclusion of [CPGSV16, Theorem `thm1`] itself forces $c_N = \xi^N$, so this is the necessary form. Proving that the global proportionality scalar of two BNT canonical forms is a pure phase power is a length-scalar control statement requiring peripheral-spectrum / almost-periodicity analysis of the bounded power sums $c_N^{(j)}(\cdot)$; it is not derivable from (11.7) alone. Until this length-scalar control is supplied, the unconditional full-tensor proportional gauge equivalence is not stated; the unconditional per-normal-tensor proportional fundamental theorem is Theorem 11.2.1.

In the equal-MPV case all ingredients are unconditional.

Lemma 11.1.16 (Equal-MPV matched copy-weight witnesses). [Lean](#) [✓ Stmt](#) Let P and Q be BNT canonical forms whose total tensors have the same MPV family. Then there are a bijection $\beta : J(Q) \simeq J(P)$, bond-dimension equalities, unit-modulus phases ζ_k , and block gauges X_k such that

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}.$$

For the same phases there is a copy-weight matching: after a permutation of the copies inside each matched sector,

$$\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}.$$

Proof. [✓ Proof](#) The full-basis matching theorem gives the bijection and block gauges. The self-overlap comparison

$$\langle V_N(Q_k), V_N(Q_k) \rangle = |\zeta_k|^{2N} \langle V_N(P_{\beta(k)}), V_N(P_{\beta(k)}) \rangle$$

together with the BNT normalization limits on both sides forces $|\zeta_k| = 1$. The matched MPV phase identities also give the eventual coefficient identities

$$c_N^{(\beta(k))}(P) = \zeta_k^N c_N^{(k)}(Q).$$

Lemma 11.1.6 applies the finite power-sum comparison in each sector and gives the copy-weight identities. $\square$

Theorem 11.1.17 (Full-basis global gauge in matched coordinates). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms whose total tensors have the same MPV family. Then there exist:*

- a bijection $\beta : J(Q) \simeq J(P)$,
- bond-dimension equalities $\dim P_{\beta(k)} = \dim Q_k$,
- copy-number equalities $r_{\beta(k)}(P) = r_k(Q)$ and copy permutations τ_k ,
- unit-modulus phases ζ_k with $|\zeta_k| = 1$ and block gauges X_k ,

such that, for every sector k , copy q , and physical index i ,

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}, \quad \nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}.$$

Moreover, if

$$X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k)$$

is the flattened block-diagonal gauge, then, for every physical index i , the flattened Q -coordinates satisfy

$$Q_{\text{flat}}^i = X P_{\text{matched}}^i X^{-1}.$$

This is the explicit direct-sum gauge construction of [CPGSV16, Appendix MPV proof, lines 1189–1192].

Proof. [✓ Proof](#) Lemma 11.1.16 gives β , the bond-dimension identifications, the phases ζ_k , the block gauges X_k , and copy permutations τ_k satisfying

$$\nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}.$$

Theorem 11.1.11 combines the block conjugacies into the flattened direct-sum conjugation by $X = \bigoplus_k (\mathbf{1}_{r_k} \otimes X_k)$. $\square$

Theorem 11.1.18 (Literal equal-MPV gauge after sector-coordinate permutation). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms whose total tensors have the same MPV family. Then the total bond dimensions agree; write their common value as D . There is*

$$Y \in \text{GL}_D(\mathbb{C}) \tag{11.8}$$

such that, for every physical index i ,

$$Q_{\text{tot}}^i = Y P_{\text{tot}}^i Y^{-1},$$

where P_{tot}^i and Q_{tot}^i are both regarded as matrices in $M_D(\mathbb{C})$ using the total bond-dimension equality. This is the BNT form of [CPGSV16, Corollary II.2 and Appendix MPV proof, lines 1189–1192].

Proof. [✓ Proof](#) Apply Theorem 11.1.17 to obtain the flattened gauge after the sectors of P have been reindexed by the bijection between basis blocks of P and Q . The sector-coordinate permutation ρ , induced by this bijection, the copy permutations, and the bond-dimension equalities, identifies the reindexed P -side direct sum with the total tensor P_{tot} , transported along the total bond-dimension equality. Composing the permutation gauge for ρ with the sector-permuted gauge gives the single matrix Y of (11.8). $\square$

Lemma 11.1.19 (Unitary gauge in matched sector coordinates). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms whose total tensors generate the same MPV family at every positive length. There are a sector bijection β , copy bijections τ_k , phases ζ_k , sector unitaries U_k , and a unitary block-diagonal matrix X such that*

$$Q_k^i = \zeta_k U_k P_{\beta(k)}^i U_k^\dagger, \quad \nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}, \quad |\zeta_k| = 1.$$

The matched sectors and copies have equal bond dimensions and multiplicities. If P_{match} denotes the corresponding reordered block assembly, then

$$Q_{\text{flat}}^i = X P_{\text{match}}^i X^{-1}.$$

Proof. [✓ Proof](#) Apply Lemma 11.2.4 to each matched sector, retaining the phase in the copy-weight identity. Repeating U_k on every copy gives $X = \bigoplus_{k,q} U_k$, which is unitary by Lemma 11.1.8. For every matched copy (k, q) ,

$$\nu_{k,q} Q_k^i = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)} \zeta_k U_k P_{\beta(k)}^i U_k^\dagger = \mu_{\beta(k), \tau_k(q)} U_k P_{\beta(k)}^i U_k^\dagger.$$

These are precisely the blocks of $Q_{\text{flat}}^i = X P_{\text{match}}^i X^{-1}$. □

Corollary 11.1.20 (Canonical Form II unitary refinement in the equal case). [Lean](#) [✓ Stmt](#) *Let P and Q be BNT canonical forms whose total tensors generate the same MPV family at every positive length. Their total bond dimensions agree; writing the common value as D , there is a unitary matrix $U \in \text{U}(D)$ such that, for every physical index i ,*

$$Q_{\text{tot}}^i = U P_{\text{tot}}^i U^\dagger.$$

This is the equal case of [CPGSV16, Corollary C.5].

Proof. [✓ Proof](#) For each matched sector, retain the phase in both identities

$$Q_k^i = \zeta_k U_k P_{\beta(k)}^i U_k^\dagger, \quad \nu_{k,q} = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)}.$$

For every matched copy (k, q) ,

$$\nu_{k,q} \zeta_k = \zeta_k^{-1} \mu_{\beta(k), \tau_k(q)} \zeta_k = \mu_{\beta(k), \tau_k(q)}.$$

Hence $Q_{\text{flat}}^i = X P_{\text{match}}^i X^{-1}$. Repeating U_k on its matched copies gives

$$X = \bigoplus_{k,q} U_k, \quad X^\dagger X = \mathbb{1}.$$

If Π is the permutation matrix restoring the original sector and copy order, then $\Pi^\dagger \Pi = \mathbb{1}$. Hence

$$U = X \Pi, \quad U^\dagger U = \mathbb{1},$$

and the permutation of matched coordinates gives

$$Q_{\text{tot}}^i = U P_{\text{tot}}^i U^\dagger.$$

□

11.2 The equal and proportional cases

Theorem 11.2.1 (Multi-block fundamental theorem for normal tensors). Lean ✓ Stmt Let P and Q be BNT canonical forms (Definition 10.4.1.11), with bases of normal tensors $(P_j)_{j=1}^{g_a}$ and $(Q_k)_{k=1}^{g_b}$. If the total tensors $P_{\text{tot}}, Q_{\text{tot}}$ generate eventually nonzero proportional MPV families — for all sufficiently large N there is a nonzero scalar c_N with $V^{(N)}(P_{\text{tot}}) = c_N V^{(N)}(Q_{\text{tot}})$ — then $g_a = g_b$. After relabelling the basis of normal tensors of P by a bijection $\beta: J(Q) \simeq J(P)$, there are scalars $\zeta_k \in \mathbb{C}$ with $|\zeta_k| = 1$ and invertible matrices Y_k such that, for every k and every physical index i ,

$$Q_k^i = \zeta_k Y_k P_{\beta(k)}^i Y_k^{-1}.$$

Thus the paired normal tensors have the same bond dimension ($\dim P_{\beta(k)} = \dim Q_k$) and the same gauge-phase class.

Proof. ✓ Proof The proportional sector-matching Theorem 11.1.2, applied to the BNT canonical forms P and Q , supplies the bijection β , the bond-dimension equalities, the unit phases ζ_k , and the block gauges Y_k with $Q_k^i = \zeta_k Y_k P_{\beta(k)}^i Y_k^{-1}$; the matching needs no per-sector unit-modulus hypothesis. $\square$

This is [CPGSV16, Theorem 2.10]; see also [CPGSV21, Theorem IV.4 and Corollary IV.5]. The statement is given on the BNT canonical-form surface (Definition 10.4.1.11), which is the paper’s “ A and B in canonical form” with chosen bases of normal tensors; the reduction of an arbitrary proportional-MPV pair to a common primitive block decomposition is Theorem 9.6.2.4 of Chapter 9.

Definition 11.2.2 (Left-canonical normalization). Lean ✓ Stmt An MPS tensor $A = (A^i)_i$ is left-canonical when

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}.$$

Lemma 11.2.3 (Phase normalization for an irreducible gauge). Lean ✓ Stmt Let A and B be left-canonical tensors of common positive bond dimension, with B irreducible. If X is invertible, $\zeta \neq 0$, and

$$B^i = \zeta X A^i X^{-1}$$

for every i , then $|\zeta| = 1$.

Proof. ✓ Proof Let ρ be a nonzero positive fixed point of $\mathcal{E}_A$ and set $\sigma = X \rho X^\dagger$. Then σ is nonzero and positive, and

$$\mathcal{E}_B(\sigma) = |\zeta|^2 \sigma.$$

The map $\mathcal{E}_B$ also has a nonzero positive fixed point. Its irreducibility therefore forces $|\zeta|^2 = 1$. $\square$

Lemma 11.2.4 (Unitary replacement with the same phase). Lean ✓ Stmt Let A and B be left-canonical irreducible tensors of common positive bond dimension satisfying

$$B^i = \zeta X A^i X^{-1}$$

for an invertible X and a nonzero scalar ζ . There is a unitary matrix U , with the same scalar ζ , such that $|\zeta| = 1$ and

$$B^i = \zeta U A^i U^\dagger$$

for every i .

Proof. ✓ Proof Lemma 11.2.3 gives $|\zeta| = 1$. Substitution into the left-canonical identities gives

$$\sum_i A^{i\dagger}(X^\dagger X)A^i = X^\dagger X.$$

Thus $X^\dagger X$ is a positive-semidefinite fixed point of the adjoint transfer map of A . Irreducibility gives $X^\dagger X = c\mathbb{1}$ for some $c > 0$, so $X^{-1} = c^{-1}X^\dagger$ and $U = c^{-1/2}X$ is unitary. Moreover,

$$UA^iU^\dagger = c^{-1}XA^iX^\dagger = XA^iX^{-1}.$$

□

Lemma 11.2.5 (Unitary gauge for left-canonical irreducible tensors). Lean ✓ Stmt *Let A and B be left-canonical irreducible tensors of common positive bond dimension that are gauge-phase equivalent. Then the gauge can be taken unitary: there are a unitary matrix U and a scalar ζ with $|\zeta| = 1$ such that, for every physical index i ,*

$$B^i = \zeta UA^iU^\dagger.$$

Proof. ✓ Proof Unpack the gauge-phase equivalence as $B^i = \zeta XA^iX^{-1}$ with X invertible and $\zeta \neq 0$. Lemma 11.2.4 replaces X by a unitary U while retaining this same phase and proves $|\zeta| = 1$. □

Corollary 11.2.6 (Canonical Form II unitary refinement of proportional matching). Lean ✓ Stmt *Under the hypotheses of Theorem 11.2.1, there are a bijection $\beta : J(Q) \simeq J(P)$, unitary matrices $U_k \in \text{U}(\dim Q_k)$, and phases $\zeta_k \in \mathbb{C}$ with $|\zeta_k| = 1$, such that $\dim P_{\beta(k)} = \dim Q_k$ and, for every k and every physical index i ,*

$$Q_k^i = \zeta_k U_k P_{\beta(k)}^i U_k^\dagger.$$

This is the proportional part of [CPGSV16, Corollary C.5].

Proof. ✓ Proof The proportional sector-matching theorem gives a bijection β , invertible matrices X_k , and unit-modulus phases ζ_k with

$$Q_k^i = \zeta_k X_k P_{\beta(k)}^i X_k^{-1}.$$

The matched tensors are left-canonical and irreducible. Hence Lemma 11.2.5 gives unitary matrices U_k satisfying

$$X_k P_{\beta(k)}^i X_k^{-1} = U_k P_{\beta(k)}^i U_k^\dagger,$$

and therefore

$$Q_k^i = \zeta_k U_k P_{\beta(k)}^i U_k^\dagger.$$

□

Theorem 11.2.7 (Equal BNT canonical forms are globally conjugate). Lean ✓ Stmt *Let P and Q be BNT canonical forms (Definition 10.4.1.11) whose total tensors P_{tot} , Q_{tot} generate the same MPV family for every system size N . Then their total bond dimensions coincide; writing the common value as D , there is one invertible matrix $X \in \text{GL}_D(\mathbb{C})$ such that, for every physical index i ,*

$$Q_{\text{tot}}^i = X P_{\text{tot}}^i X^{-1}.$$

This is [CPGSV16, Corollary 2.11] on the BNT canonical-form surface; the canonical form of [CPGSV16, Section 2.3], a direct sum of normal tensors with scalar weights, is a BNT canonical form. The reduction of an arbitrary pair of same-MPV tensors to a common BNT canonical form is Chapter 9.

Proof.  **Proof** This is Theorem 11.1.18. The BNT comparison of Chapter 10 provides the basis relabelling, the per-block gauge-phase equivalences, and the copy-weight multiset matching, which combine into the block-diagonal global gauge. The exact linear-independence matching needs no per-sector unit-modulus hypothesis, and the BNT canonical form needs no one-site injectivity. $\square$

This is also [CPGSV21, Corollary IV.5].

Chapter 12

Symmetries and String Order

This chapter collects the symmetry-theoretic consequences of the MPS Fundamental Theorem [PGVWC07], following the symmetry analysis of [PGWS⁺08]. For injective tensors, physical on-site symmetries determine virtual gauges, those gauges define projective representations on the bond space, and virtual intertwiners enter the string-order criteria.

12.1 Physical Symmetries Give Virtual Gauges

Definition 12.1.1 (Physical-index rotation). [Lean](#) [✓ Stmt](#) Given a matrix $M \in M_d(\mathbb{C})$ and an MPS tensor A , the *physical-index rotation* is the tensor $(A_M)^i := \sum_j M_{ij} A^j$. When the matrix is a unitary u , the rotation is written A_u .

Corollary 12.1.2 (Periodic symmetry yields gauge equivalence up to a root of unity). [Lean](#) [✓ Stmt](#) Suppose that for any two tensors in irreducible form II with the same MPV family there is $m > 0$ making them gauge equivalent up to an m -th root of unity. Let A be in irreducible form II and let M be a matrix such that $B := A_M$ is also in irreducible form II and $\mathcal{V}(A) = \mathcal{V}(B)$. Then there is $m > 0$ such that A and B are gauge equivalent up to an m -th root of unity.

Proof. [✓ Proof](#) Apply the assumed equivalence to A and B , which are in irreducible form II with $\mathcal{V}(A) = \mathcal{V}(B)$. □

12.2 Virtual Representation Theorem

With a full group of on-site symmetries (rather than just a single unitary), the resulting gauges determine a projective representation on the bond space.

Definition 12.2.1 (Twisted tensor). [Lean](#) [✓ Stmt](#) Given a group representation $U : G \rightarrow \mathrm{GL}_d(\mathbb{C})$ on the physical index, the *g -twisted tensor* is defined, for each $i \in \{0, \dots, d-1\}$, by

$$\tilde{A}_g^i := \sum_{j=0}^{d-1} U(g)_{ij} A^j.$$

Definition 12.2.2 (On-site symmetry). [Lean](#) [✓ Stmt](#) A tensor A is *on-site symmetric* under a group representation $U : G \rightarrow \mathrm{GL}_d(\mathbb{C})$ if $\mathcal{V}(A) = \mathcal{V}(\tilde{A}_g)$ for every $g \in G$.

Lemma 12.2.3 (Functoriality of twisting). Lean ✓ Stmt *The twisted tensor satisfies the composition law*

$$\widetilde{A}_{gh} = \widetilde{(\widetilde{A}_h)}_g,$$

i.e. twisting by gh is the same as first twisting by h and then by g .

Proof. ✓ Proof Expanding both sides using $U(gh) = U(g)U(h)$, then swapping the order of summation. □

Lemma 12.2.4 (Identity twist). Lean ✓ Stmt *Twisting by the identity group element is trivial: $\widetilde{A}_1 = A$.*

Proof. ✓ Proof Since $U(1) = \mathbb{1}_d$, each component reduces to $\sum_j \delta_{ij} A^j = A^i$. □


Lemma 12.2.5 (Symmetric twists are gauge equivalent). Lean ✓ Stmt *If A is injective and on-site symmetric under U , then for every $g \in G$ the twisted tensor $\widetilde{A}_g$ is gauge equivalent to A .*

Proof. ✓ Proof On-site symmetry gives $\mathcal{V}(A) = \mathcal{V}(\widetilde{A}_g)$, and the single-block Fundamental Theorem turns this equality of MPV families into gauge equivalence. □

Remark 12.2.6 (Fundamental-Theorem route for injective symmetry). The injective virtual-representation theorem is not proved from string-order rigidity. The on-site symmetry condition first identifies A and $\widetilde{A}_g$ as the same MPV family; the single-block Fundamental Theorem gives the gauge in Lemma 12.2.5. Gauge uniqueness and the twist composition law then give Theorem 12.2.13. The string-order results later in this chapter use this virtual representation as an input.

Theorem 12.2.7 (Virtual symmetry equation). Lean ✓ Stmt *If A is injective and on-site symmetric under U , then for every $g \in G$ there exist an invertible matrix $X(g) \in \text{GL}_D(\mathbb{C})$ and a nonzero scalar $\varphi(g) \in \mathbb{C}^\times$ (in fact $\varphi = 1$ in the single-block case) such that $\widetilde{A}_g^i = \varphi(g) X(g) A^i X(g)^{-1}$ for all i .*

Proof. ✓ Proof The physical action is transferred to the virtual level by the same local move that replaces the twisted tensor by a gauge-conjugated one:



Apply Lemma 12.2.5 to obtain $X(g)$, then set $\varphi(g) = 1$. □

Lemma 12.2.8 (Gauge uniqueness up to scalar). Lean ✓ Stmt *Let A be an injective tensor. If $X, Y \in \text{GL}_D(\mathbb{C})$ both satisfy $B^i = X A^i X^{-1} = Y A^i Y^{-1}$ for all i , then there exists a nonzero scalar $u \in \mathbb{C}^\times$ such that $Y = u \cdot X$.*

Proof. ✓ Proof The matrix $Z = Y^{-1}X$ commutes with every A^i . Since A is injective, $\{A^i\}$ spans $M_D(\mathbb{C})$, so Z lies in the centre of $M_D(\mathbb{C})$. By Lemma 2.1.5, Z is a scalar matrix. Since Z is invertible, that scalar is nonzero, and $Y = u \cdot X$. □

Theorem 12.2.9 (Gauge uniqueness from equality of MPV families). [Lean](#) [✓ Stmt](#) Let A be an injective tensor and assume that A and B generate the same MPV family. If $X, Y \in \text{GL}_D(\mathbb{C})$ both satisfy $B^i = X A^i X^{-1} = Y A^i Y^{-1}$ for all i , then there is a nonzero scalar $u \in \mathbb{C}^\times$ such that $Y = u \cdot X$.

Proof. [✓ Proof](#) This is exactly Lemma 12.2.8; the extra MPV hypothesis specifies the setting in which the two gauges are usually produced. $\square$

Definition 12.2.10 (Scalar 2-cochain). [Lean](#) [✓ Stmt](#) A scalar 2-cochain on a group G is a function $\omega : G \times G \rightarrow \mathbb{C}^\times$.

Definition 12.2.11 (Projective representation). [Lean](#) [✓ Stmt](#) Given a scalar 2-cochain ω , a projective representation of G on $\mathbb{C}^D$ is a map $\rho : G \rightarrow \text{GL}_D(\mathbb{C})$ satisfying, for all $g, h \in G$,

$$\rho(g) \rho(h) = \omega(g, h) \rho(gh).$$

Lemma 12.2.12 (Associativity forces the cocycle condition). [Lean](#) [✓ Stmt](#) If ρ is a projective representation with factor system ω and $D > 0$, then ω satisfies the 2-cocycle identity:

$$\omega(g, h) \omega(gh, k) = \omega(g, hk) \omega(h, k).$$

Proof. [✓ Proof](#) Expanding $\rho(g) (\rho(h) \rho(k))$ and $(\rho(g) \rho(h)) \rho(k)$ using the projective multiplication law and equating via matrix associativity. The hypothesis $D > 0$ allows cancellation of the invertible matrix factor. $\square$

Theorem 12.2.13 (Virtual representation theorem for injective MPS). [Lean](#) [✓ Stmt](#) Let A be an injective MPS tensor that is on-site symmetric under a group representation $U : G \rightarrow \text{GL}_d(\mathbb{C})$. Then there exist:


1. a scalar 2-cochain $\omega : G \times G \rightarrow \mathbb{C}^\times$, and
2. a map $\rho : G \rightarrow \text{GL}_D(\mathbb{C})$ satisfying, for all $g, h \in G$,

$$\rho(g) \rho(h) = \omega(g, h) \rho(gh).$$

such that, defining the gauge matrices $X(g) := \rho(g^{-1})$, one has, for every $g \in G$ and $i \in \{0, \dots, d-1\}$,

$$\tilde{A}_g^i = X(g) A^i X(g)^{-1}.$$

In tensor-network notation, for each fixed g this is the local gauge relation



with the black node denoting the tensor named in the surrounding formula and the two red side nodes acting on the virtual legs. In particular, ρ is a projective representation of G on the bond space $\mathbb{C}^D$.

Proof. [✓ Proof](#) Since A is injective and on-site symmetric, each twisted tensor $\tilde{A}_g$ is gauge equivalent to A (Lemma 12.2.5). Choose $X(g) \in \text{GL}_D(\mathbb{C})$ with $\tilde{A}_g^i = X(g) A^i X(g)^{-1}$ for all i .

By functoriality (Lemma 12.2.3), the product $X(h) X(g)$ also intertwines A with $\tilde{A}_{gh}$. Since $X(gh)$ does the same, gauge uniqueness (Lemma 12.2.8) gives a nonzero scalar $\omega'(h, g)$ with $X(h) X(g) = \omega'(h, g) X(gh)$.

Defining $\rho(g) := X(g^{-1})$ converts this to the standard law: $\rho(g) \rho(h) = \omega(g, h) \rho(gh)$ where $\omega(g, h) := \omega'(h^{-1}, g^{-1})$. $\square$

Corollary 12.2.14 (2-cocycle identity for the factor system). [Lean](#) [✓ Stmt](#) If $D > 0$, the factor system ω from Theorem 12.2.13 satisfies the 2-cocycle identity for all $g, h, k \in G$:

$$\omega(g, h) \omega(gh, k) = \omega(g, hk) \omega(h, k).$$

Proof. [✓ Proof](#) Theorem 12.2.13 gives ρ and ω . By Lemma 12.2.12, the cocycle identity follows from associativity of matrix multiplication and invertibility of the representation matrices. $\square$

12.3 Cohomology Classes of Virtual Symmetry Cocycles

Two cocycles may differ by a coboundary yet represent the same projective-representation class. The following definitions and results make this precise and establish the quotient $H^2(G, \mathbb{C}^\times)$.

Definition 12.3.1 (Coboundary). [Lean](#) [✓ Stmt](#) A scalar 2-cocycle ω is a *coboundary* if there exists $\varphi: G \rightarrow \mathbb{C}^\times$ such that, for all $g, h \in G$,

$$\omega(g, h) = \varphi(g) \varphi(h) \varphi(gh)^{-1}.$$

Definition 12.3.2 (Cohomologous cocycles). [Lean](#) [✓ Stmt](#) Two cocycles ω_1, ω_2 are *cohomologous* (written $\omega_1 \sim \omega_2$) if there exists $\varphi: G \rightarrow \mathbb{C}^\times$ such that, for all $g, h \in G$,

$$\omega_1(g, h) = \varphi(g) \varphi(h) \varphi(gh)^{-1} \omega_2(g, h).$$

Theorem 12.3.3 (Cohomology relation is an equivalence relation). [Lean](#) [✓ Stmt](#) The relation “cohomologous to” is reflexive, symmetric, and transitive on the set of scalar 2-cocycles.


Proof. [✓ Proof](#) Reflexivity: take $\varphi = 1$. Symmetry: replace φ by φ^{-1} . Transitivity: compose the witnesses φ and ψ pointwise. $\square$

Lemma 12.3.4 (Coboundary iff cohomologous to the trivial cocycle). [Lean](#) [✓ Stmt](#) A cocycle ω is a coboundary if and only if $\omega \sim 1$, where $1(g, h) = 1$ for all g, h .

Proof. [✓ Proof](#) Both directions follow by absorbing or introducing the factor $1(g, h) = 1$ via $x \cdot 1 = x$. $\square$

Theorem 12.3.5 (Gauge independence of the cocycle class). [Lean](#) [✓ Stmt](#) If the bond dimension satisfies $D \geq 1$ and two projective representations ρ_1, ρ_2 (with cocycles ω_1, ω_2) both arise as virtual representations of the same injective MPS tensor A under an on-site symmetry U , then $\omega_1 \sim \omega_2$.

Proof. [✓ Proof](#) For each g , gauge uniqueness gives a nonzero scalar $f(g)$ with $X_2(g^{-1}) = f(g) X_1(g^{-1})$. Equivalently, for each fixed g the two local gauges



describe the same twisted tensor, so they differ by a scalar. Substituting into the projective law and cancelling $X_1(g \cdot h)$ yields $\omega_2(g, h) = f(g) f(h) f(gh)^{-1} \omega_1(g, h)$. $\square$

Definition 12.3.6 (Multiplicative 2-cocycle condition). [Lean](#) [✓ Stmt](#) A scalar 2-cochain $\omega: G \times G \rightarrow \mathbb{C}^\times$ is a *2-cocycle* if it satisfies, for all $g, h, k \in G$,

$$\omega(g, h) \omega(gh, k) = \omega(g, hk) \omega(h, k).$$

Definition 12.3.7 (Second cohomology quotient). [Lean](#) [✓ Stmt](#) The second cohomology set is the quotient

$$H^2(G, \mathbb{C}^\times) := Z^2(G, \mathbb{C}^\times) / \sim,$$

where $\sim$ is the coboundary-equivalence relation.

Definition 12.3.8 (Projective equivalence). [Lean](#) [✓ Stmt](#) Two projective representations are *projectively equivalent* when their factor systems are cohomologous.

Theorem 12.3.9 (Projective equivalence and cohomologous cocycles). [Lean](#) [✓ Stmt](#) Let ρ_1, ρ_2 be projective representations with factor systems ω_1, ω_2 . Then

$$\rho_1 \sim_{\text{proj}} \rho_2 \iff \omega_1 \sim \omega_2.$$

Proof. [✓ Proof](#) This follows from the definition of projective equivalence as cohomology-class equality for factor systems. $\square$

12.3.1 Non-triviality via the commutator phase

For an abelian symmetry group the cohomology class of a factor system is detected by an anti-symmetric pairing, which gives a computable test for non-triviality.

Definition 12.3.1.1 (Commutator phase). [Lean](#) [✓ Stmt](#) The *commutator phase* of a 2-cochain ω at a pair g, h is $\beta_\omega(g, h) := \omega(g, h)\omega(h, g)^{-1}$. It is defined for any $\mathbb{C}^\times$ -valued 2-cochain, the cocycle condition is not needed.

Theorem 12.3.1.2 (The commutator phase is a class invariant). [Lean](#) [✓ Stmt](#) If $\omega_1 \sim \omega_2$ and $gh = hg$, then $\beta_{\omega_1}(g, h) = \beta_{\omega_2}(g, h)$.

Proof. [✓ Proof](#) Write $\omega_1(g, h) = \varphi(g)\varphi(h)\varphi(gh)^{-1}\omega_2(g, h)$ for a coboundary witness φ . Substituting into the ratio and using $gh = hg$ and the commutativity of $\mathbb{C}^\times$,

$$\beta_{\omega_1}(g, h) = \frac{\varphi(g)\varphi(h)\varphi(gh)^{-1}\omega_2(g, h)}{\varphi(h)\varphi(g)\varphi(hg)^{-1}\omega_2(h, g)} = \frac{\omega_2(g, h)}{\omega_2(h, g)} = \beta_{\omega_2}(g, h),$$

the φ -factors cancelling since $\varphi(gh) = \varphi(hg)$. $\square$

Definition 12.3.1.3 (Non-trivial cohomology class). [Lean](#) [✓ Stmt](#) A factor system ω has a *non-trivial class* when it is not cohomologous to the constant cocycle 1.

Theorem 12.3.1.4 (Commutator-phase non-triviality test). [Lean](#) [✓ Stmt](#) If a commuting pair g, h has $\beta_\omega(g, h) \neq 1$, then ω has a non-trivial class.

Proof. [✓ Proof](#) The trivial cocycle has commutator phase 1 at every pair, so by Theorem 12.3.1.2 any cocycle cohomologous to it has $\beta = 1$ at every commuting pair. Contrapositively, $\beta_\omega(g, h) \neq 1$ for one commuting pair forces ω to be non-cohomologous to the trivial cocycle. $\square$

12.4 Permutation-Based Physical Symmetry


The linear-combination twist $\tilde{A}_g^i = \sum_j U(g)_{ij} A^j$ is the standard form used in the virtual representation theorem. The following alternative formulation replaces the matrix action by a permutation of the physical index; it is useful for symmetries that act by reordering basis states rather than by a general linear map.

Definition 12.4.1 (Permutation symmetry datum). [Lean](#) [✓ Stmt](#) An *on-site symmetry datum* for a group G on a physical space of dimension d consists of a matrix-valued map $u : G \rightarrow M_d(\mathbb{C})$ and a physical-index action $\sigma : G \rightarrow \text{End}(\{0, \dots, d-1\})$.

Definition 12.4.2 (Permutation-twisted tensor). [Lean](#) [✓ Stmt](#) Given an on-site symmetry datum (u, σ) , the *permutation-twisted tensor* at group element g is

$$(A^{\sigma_g})^i := A^{\sigma(g)(i)}.$$

In tensor-network notation the local tensor is unchanged and only the physical leg is relabelled:



Lemma 12.4.3 (Identity permutation twist). [Lean](#) [✓ Stmt](#) If $\sigma(1) = \text{id}$, then $(A^{\sigma_1})^i = A^i$ for all i .

Proof. [✓ Proof](#) The claim follows from $\sigma(1)(i) = i$ for all i . □

Composition order. The linear-combination twist (Lemma 12.2.3) composes as $\widetilde{(\widetilde{A}_h)}_g = \widetilde{A}_{gh}$, applying h first then g . The permutation twist below uses the same left-action convention $\sigma(gh) = \sigma(g) \circ \sigma(h)$, so $(A^{\sigma_g})^{\sigma_h} = A^{\sigma_{gh}}$; the inner permutation $\sigma(h)$ acts first on the index.

Lemma 12.4.4 (Composition of permutation twists). [Lean](#) [✓ Stmt](#) If $\sigma(gh) = \sigma(g) \circ \sigma(h)$ for all $g, h \in G$, then

$$A^{\sigma_{gh}} = (A^{\sigma_g})^{\sigma_h}.$$

Proof. [✓ Proof](#) Expanding both sides: $(A^{\sigma_{gh}})^i = A^{\sigma(g)(\sigma(h)(i))} = ((A^{\sigma_g})^{\sigma_h})^i$. Diagrammatically this is just two consecutive relabellings of the same local tensor:



□


12.5 String Order and Local Symmetry Equivalence

The string order parameter [PGWS⁺08] detects whether an on-site unitary u acts as a local symmetry on the state generated by an injective MPS tensor.

Definition 12.5.1 (Twisted transfer map). [Lean](#) [✓ Stmt](#) Given an MPS tensor A and a physical-index matrix u , the *twisted transfer map* is

$$\mathcal{E}_u(X) := \sum_{n, n'} \langle n' | u | n \rangle A_n X A_{n'}^\dagger.$$

Diagrammatically one inserts the physical operator u on the doubled local transfer cell:




The upper and lower black nodes denote A_n and $A_n^\dagger$, and the red node labelled u couples the physical legs.

Definition 12.5.2 (String order parameter). [Lean](#) [✓ Stmt](#) The *string order parameter* for twist u and stationary state Λ at length L is

$$R_L(u) := \text{tr}(\Lambda \cdot \mathcal{E}_u^L(\mathbb{1})).$$

In tensor-network notation this is the length- L doubled chain with a copy of u inserted on each physical rung:



The boundary matrix Λ is contracted into the virtual boundary, the red nodes labelled u sit on the physical rungs, and the right boundary is traced.

Definition 12.5.3 (Physical string correlator). [Lean](#) [✓ Stmt](#) For physical operators $x, y \in M_d(\mathbb{C})$, a twist $u \in M_d(\mathbb{C})$, and boundary state Λ , the transfer-form string correlator of [PGWS⁺08, display SOPMP, lines 176–181] is

$$S_N(x, y, u) := \text{tr}(\Lambda \mathcal{E}_x \mathcal{E}_u^N(\mathcal{E}_y(\mathbb{1}))).$$

This is the physical-endpoint expression corresponding to $\langle \Psi | x \otimes u^{\otimes N} \otimes y | \Psi \rangle$.

Definition 12.5.4 (Physical string order). [Lean](#) [Lean](#) [✓ Stmt](#) A tensor has physical string order, in the sense of [PGWS⁺08, display SOP, lines 112–122], if there exist a unitary $u \neq \mathbb{1}$ and physical endpoint operators x, y for which the absolute value of the correlator has a positive limit: for some $s > 0$,

$$|S_N(x, y, u)| \rightarrow s.$$

The comparison with virtual boundary matrices is a separate endpoint-realizability statement.

Definition 12.5.5 (Boundary string order parameter). [Lean](#) [✓ Stmt](#) For a boundary state Λ and boundary matrices $X, Y \in M_D(\mathbb{C})$, the *boundary string order parameter* is

$$R_L(u; X, Y) := \text{tr}(\Lambda X \mathcal{E}_u^L(Y)).$$

The ordinary string order parameter is the special case $X = Y = \mathbb{1}$. The matrices X, Y act on the virtual level. In the string correlator of [PGWS⁺08] the string of twists is instead flanked by local operators x, y acting on physical sites; expanding such operators in the transfer picture produces particular virtual matrices, and [PGWS⁺08] argues that for injective tensors, physical operators on sufficiently many sites produce every virtual matrix. The results below are stated directly in this virtual-boundary form.

Definition 12.5.6 (Local symmetry). [Lean](#) [✓ Stmt](#) An MPS tensor A has *local symmetry* under a unitary u with respect to a boundary state Λ if there exist a unitary matrix V and a phase μ with $|\mu| = 1$ such that $V^\dagger \Lambda V = \Lambda$ and, for every physical index i ,

$$\sum_j u_{ij} A^j = \mu V A^i V^\dagger.$$

The unitary V intertwines the physical action of u with a virtual conjugation, and the boundary state Λ is invariant under V .


Definition 12.5.7 (String order). [Lean](#) [✓ Stmt](#) String order *exists* for u with respect to a boundary state Λ if there exist boundary matrices X, Y and a positive constant $c > 0$ such that $c \leq \|R_L(u; X, Y)\|$ for all L , where $R_L(u; X, Y)$ is the boundary-refined string-order parameter. Thus some boundary-refined string-order sequence does not decay to zero.

String order is here an existence statement over the virtual matrices X, Y , for a fixed twist u ; no claim is made for any particular boundary choice. Definition 12.5.4 records the physical-endpoint source definition, which quantifies existentially over u, x, y and requires a positive limit rather than a lower bound uniform in the length; that paper also notes that the correlator for a particular endpoint choice can vanish even when the symmetry is present.


Lemma 12.5.8 (String order prevents decay to zero). [Lean](#) [✓ Stmt](#) If A has string order for u with respect to a boundary state Λ , then for some boundary matrices X, Y the sequence $R_L(u; X, Y)$ does not tend to 0 as $L \rightarrow \infty$.

Proof. [✓ Proof](#) A uniform lower bound $c \leq \|R_L(u; X, Y)\|$ is incompatible with convergence to 0. $\square$

Definition 12.5.9 (Local physical intertwining). [Lean](#) [✓ Stmt](#) The local physical intertwining relation is $\sum_j u_{ij} A^j = V A^i V^\dagger$. Locally, the physical action of u can therefore be pushed to the virtual legs:



Definition 12.5.10 (Transfer-map covariance). [Lean](#) [✓ Stmt](#) Transfer-map covariance is the identity that the transfer map commutes with virtual conjugation: $\mathcal{E}(V X V^\dagger) = V \mathcal{E}(X) V^\dagger$. In doubled-layer notation, the virtual operators slide through the local transfer cell:




Definition 12.5.11 (Commutation in the doubled transfer picture). [Lean](#) [✓ Stmt](#) In the transfer-map language, the commutation of the doubled transfer matrix $E = \sum_i A_i \otimes \bar{A}_i$ with $V \otimes \bar{V}$ is

$$V \mathcal{E}_1(X) V^\dagger = \mathcal{E}_1(V X V^\dagger).$$

Theorem 12.5.12 (Transfer-map covariance and doubled commutation are equivalent). [Lean](#) [✓ Stmt](#) Transfer-map covariance and commutation in the doubled transfer picture are equivalent for any matrix V .

Proof. [✓ Proof](#) Both sides express the same local picture with opposite orientation:



The two formulas are the same identity read from opposite sides. $\square$

Lemma 12.5.13 (Unitary Kraus mixing). [Lean](#) [✓ Stmt](#) If u is unitary, then replacing Kraus operators $\{A_i\}$ by their unitary mixtures $\{\sum_j u_{ij} A_j\}$ preserves the channel:

$$\sum_i \left(\sum_j u_{ij} A_j \right) Y \left(\sum_j u_{ij} A_j \right)^\dagger = \sum_i A_i Y A_i^\dagger.$$

Proof. [✓ Proof](#) This is exactly the usual invariance of a Kraus decomposition under a unitary change of Kraus operators; see Theorem 16.4.4. $\square$

Theorem 12.5.14 (Local physical intertwining implies transfer-map covariance). [Lean](#) [✓ Stmt](#) If V and u are unitary and the local physical intertwining relation holds, then the transfer map is covariant.

Proof. [✓ Proof](#) Starting from

one inserts $V^\dagger V = \mathbb{1}$ so that each local transfer cell is written in terms of conjugated Kraus operators, then replaces those operators using the intertwining relation. After that, the physical u -boxes disappear by the unitary Kraus-mixing identity. $\square$

Theorem 12.5.15 (Transfer-map covariance implies local physical intertwining). [Lean](#) [✓ Stmt](#) For an injective MPS, if V is unitary and the transfer map is covariant, then there exists a unitary u satisfying the local physical intertwining relation.

Proof. [✓ Proof](#) Set $B^i := V A^i V^\dagger$. The transfer-map covariance relation

implies that the two Kraus families B and A define the same transfer map. Kraus freedom for equal channels converts $\mathcal{E}_B = \mathcal{E}_A$ into a unitary Kraus mixing matrix u with $B^i = \sum_j u_{ij} A^j$. $\square$

Definition 12.5.16 (Twisted companion tensor). [Lean](#) [✓ Stmt](#) For an MPS tensor A and a physical-index matrix u , define the companion tensor B_u by

$$B_u^n := \sum_{n'=0}^{d-1} \overline{u_{n'n}} A^{n'}.$$

Lemma 12.5.17 (Twisted transfer as a mixed transfer operator). [Lean](#) [✓ Stmt](#) The twisted transfer map of A is the mixed transfer operator of the pair (A, B_u) :

$$\mathcal{E}_u = F_{A, B_u}.$$

Proof. [✓ Proof](#) Expand both definitions and rearrange the double sum. $\square$

Theorem 12.5.18 (Twisted-transfer eigenvalues have modulus at most one). [Lean](#) [✓ Stmt](#) Let A be an injective MPS tensor with $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, and let u satisfy $uu^\dagger = \mathbb{1}$. If

$$\mathcal{E}_u(X) = \lambda X$$

for some nonzero $X \in M_D(\mathbb{C})$, then $|\lambda| \leq 1$.

Proof. [✓ Proof](#) By Lemma 12.5.17, the twisted transfer map is the mixed transfer operator F_{A,B_u} . Because B_u is obtained from A by unitary mixing of the Kraus operators, the transfer maps of A and B_u coincide, so the two adjoint channels share a common positive definite fixed point. Gauging both families by that fixed point puts them in a common trace-preserving gauge. Theorem 7.4.1 then applies to the mixed transfer operator in that gauge, and similarity preserves the eigenvalue λ . $\square$

Theorem 12.5.19 (Modulus-one twisted eigenvalue implies gauge-phase equivalence). [Lean](#) [✓ Stmt](#) Let A be an injective MPS tensor with $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, and let u satisfy $uu^\dagger = \mathbb{1}$. If

$$\mathcal{E}_u(X) = \lambda X \tag{12.1}$$

for some nonzero $X \in M_D(\mathbb{C})$ with $|\lambda| = 1$, then A is gauge-phase equivalent to the companion tensor B_u .

Proof. [✓ Proof](#) By Lemma 12.5.17, the twisted transfer map is the mixed transfer operator F_{A,B_u} . Because B_u is obtained from A by unitary mixing of the Kraus operators, the transfer maps of A and B_u coincide, so the two adjoint channels share a common positive definite fixed point. Gauging both families by that fixed point puts them in a common trace-preserving gauge. In that gauge the eigenvalue equation (12.1) becomes a modulus-one peripheral eigenvalue for an irreducible mixed transfer operator, so Theorem 7.8.1 yields gauge-phase equivalence. Undoing the common trace-preserving gauge gives the claim for A and B_u . $\square$

Lemma 12.5.20 (Spectral radius < 1 forces boundary-string decay). [Lean](#) [✓ Stmt](#) If $\rho(\mathcal{E}_u) < 1$, then, as $L \rightarrow \infty$, for all boundary matrices $X, Y, \Lambda \in M_D(\mathbb{C})$ one has

$$R_L(u; X, Y) \rightarrow 0.$$

Proof. [✓ Proof](#) In finite dimension, $\rho(\mathcal{E}_u) < 1$ implies $\mathcal{E}_u^L \rightarrow 0$ in operator norm. Composing with the fixed trace pairing $Z \mapsto \text{tr}(\Lambda X Z)$ gives the stated scalar convergence. $\square$

Theorem 12.5.21 (String order gives gauge-phase equivalence with the companion tensor). [Lean](#) [✓ Stmt](#) Let A be injective, let $\Lambda \in M_D(\mathbb{C})$, and assume $uu^\dagger = \mathbb{1}$ and $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. If string order exists for u with boundary parameter Λ , then A is gauge-phase equivalent to the companion tensor B_u .

Proof. [✓ Proof](#) If $\rho(\mathcal{E}_u) < 1$, Lemma 12.5.20 forces every boundary string-order sequence to decay to zero, contradicting Definition 12.5.7. Hence $\rho(\mathcal{E}_u) \geq 1$. Theorem 12.5.18 shows that every eigenvalue has modulus at most 1, so some peripheral eigenvalue has modulus exactly 1. Applying Theorem 12.5.19 then gives the gauge-phase equivalence. $\square$

Theorem 12.5.22 (Gauge-phase equivalence with the companion tensor yields a virtual unitary). [Lean](#) [✓ Stmt](#) Let A be injective, assume $uu^\dagger = \mathbb{1}$ and $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. If A is gauge-phase equivalent to the companion tensor B_u , then there exist a unitary V and a phase μ with $|\mu| = 1$ such that, for every physical index i ,

$$\sum_j u_{ij} A^j = \mu V A^i V^\dagger. \tag{12.2}$$

Proof. ✓ Proof Start from an invertible intertwiner X between A and B_u . The positive matrix $Q := XX^\dagger$ is a positive fixed point of $\mathcal{E}_A$. By uniqueness of positive fixed points for an injective normalized tensor, Q is a positive scalar multiple of the identity. Rescaling X by the square root of that scalar produces a unitary V , and the original intertwining relation becomes the phased covariance equation (12.2), with $|\mu| = 1$. $\square$

Theorem 12.5.23 (Virtual unitary gives a twisted-transfer eigenmatrix). Lean ✓ Stmt Assume $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$ and that, for every physical index i ,

$$\sum_j u_{ij} A^j = \mu V A^i V^\dagger$$

with V unitary. Then V is an eigenmatrix of the twisted transfer map:

$$\mathcal{E}_u(V) = \mu V.$$

Proof. ✓ Proof Expand $\mathcal{E}_u(V)$, insert the intertwining relation into each local term, and use $V^\dagger V = \mathbb{1}$ together with $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. $\square$

Theorem 12.5.24 (Boundary-state invariance of the virtual unitary). Lean ✓ Stmt Let A be injective, assume $uu^\dagger = \mathbb{1}$, and let Λ be a positive definite boundary state with $\text{tr}(\Lambda) = 1$ and $\mathcal{E}_A^\dagger(\Lambda) = \Lambda$. If V is a unitary and μ a phase satisfying, for every physical index i ,

$$\sum_j u_{ij} A^j = \mu V A^i V^\dagger, \tag{12.3}$$

then

$$V^\dagger \Lambda V = \Lambda.$$

Proof. ✓ Proof Relation (12.3) shows that $V^\dagger \Lambda V$ is another positive fixed point of the adjoint transfer map. Since Λ is the normalized positive fixed point, uniqueness forces $V^\dagger \Lambda V = \Lambda$. $\square$

Lemma 12.5.25 (Local symmetry implies string order). Lean ✓ Stmt Assume $\text{tr}(\Lambda) = 1$ and $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. If u is a local symmetry of A with respect to Λ , then string order exists for u .

Proof. ✓ Proof Choose the boundary matrices $X = V^\dagger$ and $Y = V$, where V is the unitary from the local-symmetry datum. By Theorem 12.5.23, one has $\mathcal{E}_u^L(V) = \mu^L V$. Therefore

$$R_L(u; V^\dagger, V) = \mu^L \text{tr}(\Lambda) = \mu^L,$$

whose norm is constantly 1. $\square$

Theorem 12.5.26 (Local symmetry iff a unitary peripheral eigenmatrix exists). Lean ✓ Stmt Let A be injective, and assume $uu^\dagger = \mathbb{1}$, $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, Λ is positive definite, $\text{tr}(\Lambda) = 1$, and $\mathcal{E}_A^\dagger(\Lambda) = \Lambda$. Then u is a local symmetry if and only if there exist a unitary $V \in M_D(\mathbb{C})$ and a phase μ with $|\mu| = 1$ such that

$$\mathcal{E}_u(V) = \mu V.$$

By Theorem 12.5.18, this witness form is equivalent to the peripheral-spectrum condition $\rho(\mathcal{E}_u) = 1$.

Proof. ✓ Proof The forward direction is Theorem 12.5.23. For the reverse direction, start from a unitary peripheral eigenmatrix. Theorem 12.5.19 gives gauge-phase equivalence with the companion tensor; then Theorem 12.5.22 yields a virtual unitary and phase satisfying the local covariance relation, and Theorem 12.5.24 gives the boundary-state invariance needed for local symmetry. $\square$

Theorem 12.5.27 (String order $\Leftrightarrow$ local symmetry). [Lean](#) [✓ Stmt](#) For an injective MPS, assume $uu^\dagger = \mathbb{1}$, $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, Λ is positive definite, $\text{tr}(\Lambda) = 1$, and $\mathcal{E}_A^\dagger(\Lambda) = \Lambda$. Then string order for u exists iff u is a local symmetry.

Proof. [✓ Proof](#) If string order exists, Theorem 12.5.21 gives a gauge-phase equivalence with the companion tensor. Theorem 12.5.22 extracts a virtual unitary and phase from that equivalence, and Theorem 12.5.24 shows that the same unitary preserves the boundary state, so one obtains local symmetry. Conversely, Lemma 12.5.25 turns local symmetry into a non-decaying boundary string-order sequence. $\square$

Theorem 12.5.28 (Virtual unitary from string order). [Lean](#) [✓ Stmt](#) For an injective MPS, let $\Lambda \in M_D(\mathbb{C})$, and assume $uu^\dagger = \mathbb{1}$ and $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$. If string order exists for u with boundary parameter Λ , then there exists a virtual unitary V and a unit-modulus scalar μ such that $\sum_j u_{ij} A^j = \mu V A^i V^\dagger$ for all i . Diagrammatically this is the same local intertwining relation, but only up to an overall unit scalar on the virtual side:

Proof. [✓ Proof](#) String order first gives gauge-phase equivalence with the companion tensor by Theorem 12.5.21. Theorem 12.5.22 then normalizes the resulting intertwiner to a unitary and produces the phase μ . $\square$

12.6 SPT Phase Labels and String-Order Universality

Two injective symmetric tensors are defined to be in the same symmetry-protected topological (SPT) phase when their virtual representation cocycles are cohomologous. The identification of this cocycle class with phase equivalence under symmetric gapped paths is the classification theorem of Chen, Gu, and Wen [CGW11] and Schuch, Pérez-García, and Cirac [SPGC11, Section II.F] (Remark 12.6.2). The theorems of this section show that string order holds for every injective symmetric tensor with canonical normalisations, so existence of string order is a symmetry diagnostic and does not separate SPT phases (Remark 12.6.5).

Definition 12.6.1 (Same SPT phase). [Lean](#) [✓ Stmt](#) Injective tensors A and B , both symmetric under an on-site representation $U: G \rightarrow \text{GL}_d(\mathbb{C})$, are in the *same SPT phase* if there exist virtual cocycles ω_A, ω_B with corresponding projective representations that intertwine the respective twisted tensors and satisfy $\omega_A \sim \omega_B$ (cohomologous).

Remark 12.6.2 (Scope of the classification theorem). For injective tensors with a fixed on-site symmetry and no symmetry breaking, two tensors can be connected by a symmetric gapped path if and only if their virtual cocycle classes in $H^2(G, \text{U}(1))$ coincide ([CGW11]; [SPGC11, Section II.F]). The results below use only the virtual cocycle class and its gauge independence (Theorem 12.3.5), and Definition 12.6.1 takes cohomologous cocycles as the definition of phase equality.

Theorem 12.6.3 (String order holds for injective symmetric MPS). [Lean](#) [✓ Stmt](#) Let A be an injective tensor, symmetric under a unitary representation U . Assume $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, Λ is positive definite, $\text{tr}(\Lambda) = 1$, and $\mathcal{E}_A^\dagger(\Lambda) = \Lambda$. Then string order exists for every group element g .

Proof. ✓ **Proof** The virtual representation theorem produces $\rho(g^{-1}) \in \text{GL}_D(\mathbb{C})$ satisfying $\sum_j U(g)_{ij} A^j = \rho(g^{-1}) A^i \rho(g^{-1})^{-1}$. A direct computation using $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$ shows that $\mathcal{E}_{U(g)}(\rho(g^{-1})) = \rho(g^{-1})$, so $\rho(g^{-1})$ is a nonzero eigenvector with eigenvalue 1. Theorem 12.5.19 gives gauge-phase equivalence with the companion tensor, and Theorem 12.5.22 extracts a virtual unitary and phase. By Theorem 12.5.24, this unitary preserves the boundary state, so Lemma 12.5.25 gives string order. $\square$

Theorem 12.6.4 (String order existence agrees for any two injective symmetric tensors). Lean
✓ **Stmnt** *If A and B are injective, both on-site symmetric under a unitary representation U , and if $\mathcal{E}_A(\mathbb{1}) = \mathbb{1}$, $\mathcal{E}_B(\mathbb{1}) = \mathbb{1}$, Λ_A and Λ_B are positive definite, $\text{tr}(\Lambda_A) = 1$, $\text{tr}(\Lambda_B) = 1$, $\mathcal{E}_A^\dagger(\Lambda_A) = \Lambda_A$, and $\mathcal{E}_B^\dagger(\Lambda_B) = \Lambda_B$, then for every $g \in G$, string order exists for A with twist $U(g)$ if and only if it exists for B . In particular, since Definition 12.6.1 implies on-site symmetry for both tensors, this theorem applies to tensors in the same SPT phase; it applies equally to symmetric tensors in different SPT phases.*

Proof. ✓ **Proof** Both directions follow from Theorem 12.6.3, which shows that string order holds universally for injective symmetric tensors with canonical normalisations. $\square$

Remark 12.6.5 (String order detects symmetry, not the SPT phase). Theorem 12.6.3 applies to every injective symmetric tensor with canonical normalisations, irrespective of its cocycle class, so tensors in different SPT phases all have string order for every group element. Existence of string order is equivalent to the local symmetry itself (Theorem 12.5.27); the invariant that separates SPT phases is the cohomology class of the virtual representation cocycle (Definition 12.6.1).

Chapter 13

Parent Hamiltonians

For a translation-invariant matrix product state with tensor A , the *parent Hamiltonian* is the translation-invariant sum of local terms, each the orthogonal projector onto the complement of the length- L local space $G_L(A) = \text{im}(\Gamma_L)$. This is the space $\mathcal{G}_L^A$ of [PGVWC07]; we write $\Gamma_L(X)(\sigma) = \text{tr}(A^\sigma X)$, which is the same as the source convention $\text{tr}(XA^\sigma)$ by cyclicity of the trace. We follow the construction of [PGVWC07] and [CPGSV21], identifying the N -site space with functions on the computational basis,

$$\{0, \dots, d-1\}^N \rightarrow \mathbb{C},$$

rather than as an explicit tensor product.

The chapter establishes the ground-space structure in increasing generality. For an injective tensor, when $N \geq 2$ and $1 < L \leq N$, the parent Hamiltonian is frustration-free and the periodic MPS vector $V^{(N)}(A)$ is its unique ground state, obtained from the intersection property of the local ground spaces. For a normal tensor the corresponding conclusion follows, under the stated closure-property and length hypotheses, from the inverting-and-growing-back argument at the injectivity length. The nearest-neighbor section separates the translated parent-term commutation and zero-energy equations for $V^{(N)}(A)$ from the source ground-space spanning clause in the pure-state theorem relating renormalization fixed points, zero correlation length, and nearest-neighbor commuting parent Hamiltonians in [CPGSV16]. The spectral-gap section records the source anticommutator martingale condition for overlapping local terms, together with cyclic norm-compression estimates as sufficient stronger hypotheses. For a tensor in block-injective canonical form $A^i = \bigoplus_j \mu_j A_j^i$ with a basis of normal tensors $\{A_j\}$, [CPGSV21, Theorem IV.16] and [PGVWC07, Theorem 12] state that the periodic parent-Hamiltonian ground space is spanned by the vectors $\{V^{(N)}(A_j)\}$. The sections below record the open-segment recursion and the intersection and closure steps, together with separate conditional statements for the block-diagonal boundary-condition comparison in the periodic ring.

13.1 The N -site Hilbert space

Definition 13.1.1 (N -site Hilbert space). [Lean](#) [✓ Stmt](#) For local dimension d and chain length N , the N -site Hilbert space is

$$\mathcal{H}_N^{(d)} := \{0, \dots, d-1\}^N \rightarrow \mathbb{C}.$$

This is the computational-basis identification used throughout the chapter for periodic-chain states.

13.2 Local ground space $G_L(A)$

Fix an MPS tensor $A = \{A^i\}_{i=0}^{d-1}$ with $A^i \in M_D(\mathbb{C})$ and a block length $L \geq 1$. For a word $w = (i_1, \dots, i_L)$, write $A^w := A^{i_1} \dots A^{i_L}$. The boundary-condition parametrization is


$$\Gamma_L : M_D(\mathbb{C}) \rightarrow (\{0, \dots, d-1\}^L \rightarrow \mathbb{C}), \quad \Gamma_L(X)(\sigma) := \text{tr}(A^\sigma X),$$

where A^σ abbreviates $A^{\sigma(0)} \dots A^{\sigma(L-1)}$. By cyclicity of the trace this is the same as the convention of [PGVWC07], namely $\text{tr}(XA^\sigma)$ for the local space $\mathcal{G}_L^A$.

Definition 13.2.1 (Ground-space map). [Lean](#) [✓ Stmt](#) The map Γ_L above is a $\mathbb{C}$ -linear map

$$\Gamma_L : M_D(\mathbb{C}) \rightarrow (\{0, \dots, d-1\}^L \rightarrow \mathbb{C}).$$

In tensor-network notation,



the upper chain denotes the word tensor A^σ with open physical indices $\sigma_1, \dots, \sigma_L$, and the lower red node denotes the boundary matrix X inserted on the closing virtual bond before taking the trace. The chain length L is fixed by the surrounding formula, not by an extra label inside the diagram.

Lemma 13.2.2 (Evaluation formula). [Lean](#) [✓ Stmt](#) For every boundary matrix $X \in M_D(\mathbb{C})$ and every length- L word σ ,

$$\Gamma_L(X)(\sigma) = \text{tr}(A^\sigma X).$$

Definition 13.2.3 (Local ground space). [Lean](#) [✓ Stmt](#) The local ground space is the image

$$G_L(A) := \text{im}(\Gamma_L) \subseteq (\{0, \dots, d-1\}^L \rightarrow \mathbb{C}).$$

Lemma 13.2.4 (Dimension bound). [Lean](#) [✓ Stmt](#) For every L ,

$$\dim G_L(A) \leq D^2.$$

Proof. [✓ Proof](#) Since $G_L(A)$ is the range of a linear map from $M_D(\mathbb{C})$,

$$\dim G_L(A) \leq \dim M_D(\mathbb{C}) = D^2.$$

□

Lemma 13.2.5 (Ambient-space dimension). [Lean](#) [✓ Stmt](#) The above coefficient space satisfies

$$\dim(\{0, \dots, d-1\}^L \rightarrow \mathbb{C}) = d^L.$$

Lemma 13.2.6 (Nontriviality criterion). [Lean](#) [✓ Stmt](#) If $d^L > D^2$, then $G_L(A)$ is a proper subspace of the local Hilbert space:

$$G_L(A) \neq (\{0, \dots, d-1\}^L \rightarrow \mathbb{C}).$$

Proof. [✓ Proof](#) If $G_L(A)$ were the whole space, its dimension would be d^L by Lemma 13.2.5; Lemma 13.2.4 gives the contradiction $d^L \leq D^2$. □

13.3 Parent interaction and chain Hamiltonian

The canonical parent interaction at range L is

$$h_L(A) = 1 - \Pi_{G_L(A)} = \Pi_{G_L(A)^\perp}, \quad \ker h_L(A) = G_L(A).$$

The sources also allow any positive local interaction with this kernel; the orthogonal projector is the canonical representative used here.

Definition 13.3.1 (Local ground space under the canonical ℓ^2 identification). [Lean](#) [✓ Stmt](#) Let

$$\iota_L : (\{0, \dots, d-1\}^L \rightarrow \mathbb{C}) \simeq \ell^2(\{0, \dots, d-1\}^L)$$

be the canonical computational-basis identification. The local ground space under this ℓ^2 identification is

$$G_L^{\text{ES}}(A) := \iota_L(G_L(A)) \subseteq \ell^2(\{0, \dots, d-1\}^L).$$

Lemma 13.3.2 (Membership under the canonical ℓ^2 identification). [Lean](#) [✓ Stmt](#) For $v \in \ell^2(\{0, \dots, d-1\}^L)$,

$$v \in G_L^{\text{ES}}(A) \iff \iota_L^{-1}v \in G_L(A).$$

Proof. [✓ Proof](#) This is the defining equation $G_L^{\text{ES}}(A) = \iota_L(G_L(A))$. □

Definition 13.3.3 (Canonical parent interaction). [Lean](#) [✓ Stmt](#) The parent interaction at range L is the canonical orthogonal projector

$$h_L(A) := \Pi_{G_L(A)^\perp} = 1 - \Pi_{G_L(A)}, \quad \ker h_L(A) = G_L(A),$$

on the local L -site space.

Lemma 13.3.4 (Parent interaction is idempotent). [Lean](#) [✓ Stmt](#) The local parent interaction is a projector:

$$h_L(A)^2 = h_L(A).$$

Proof. [✓ Proof](#) This is the idempotency of the orthogonal projector onto $G_L(A)^\perp$, transported from the Hilbert-space realization of the local coefficient space. □

Lemma 13.3.5 (Parent interaction annihilates ground-space elements). [Lean](#) [✓ Stmt](#) For any $v \in G_L(A)$, we have $h_L(A)v = 0$.

Proof. [✓ Proof](#) Since $h_L(A)$ is the orthogonal projector onto $G_L(A)^\perp$, it annihilates every element of $G_L(A)$. □

Definition 13.3.6 (Restriction to a cyclic interval). [Lean](#) [✓ Stmt](#) For $\sigma \in \{0, \dots, d-1\}^N$ and a starting site i , write

$$\sigma|_{[i, i+L-1]} \in \{0, \dots, d-1\}^L$$

for the word defined by

$$(\sigma|_{[i, i+L-1]})(r) = \sigma(i + r \bmod N), \quad 0 \leq r < L.$$

Definition 13.3.7 (Prescribing a cyclic interval). [Lean](#) [✓ Stmt](#) If $L \leq N$ and $\tau \in \{0, \dots, d-1\}^L$, write

$$\sigma^{[i, i+L-1] \leftarrow \tau}$$

for the configuration defined by

$$\sigma^{[i, i+L-1] \leftarrow \tau}(k) = \begin{cases} \tau(r), & k \equiv i + r \pmod{N} \text{ for some } 0 \leq r < L, \\ \sigma(k), & \text{otherwise.} \end{cases}$$

The index r in the first case is unique because $L \leq N$.

Lemma 13.3.8 (Extracting a replaced window). [Lean](#) [✓ Stmt](#) If $L \leq N$, then restricting to the same cyclic interval after prescribing its values recovers the inserted word:

$$(\sigma^{[i, i+L-1] \leftarrow \tau})|_{[i, i+L-1]} = \tau.$$

Lemma 13.3.9 (Replacing an extracted window). [Lean](#) [✓ Stmt](#) If $L \leq N$, then prescribing on a cyclic interval the values already carried by σ leaves σ unchanged:

$$\sigma^{[i, i+L-1] \leftarrow \sigma|_{[i, i+L-1]}} = \sigma.$$

Proof. [✓ Proof](#) Case-split on whether the offset $(k - i + N) \bmod N$ lies in $[0, L)$: inside the cyclic interval the prescribed value is the original value of σ ; outside the interval both sides already agree with σ . $\square$

Lemma 13.3.10 (Replacing the same window twice). [Lean](#) [✓ Stmt](#) If $L \leq N$, then two prescriptions on the same cyclic interval reduce to the second prescription:

$$(\sigma^{[i, i+L-1] \leftarrow \tau})^{[i, i+L-1] \leftarrow v} = \sigma^{[i, i+L-1] \leftarrow v}.$$

Proof. [✓ Proof](#) On the cyclic interval both sides take the value prescribed by v . Away from that interval neither second prescription changes the value. $\square$

For a periodic chain of length N , the parent Hamiltonian is the sum of translated copies of the local interaction.

Definition 13.3.11 (Translated local term). [Lean](#) [✓ Stmt](#) For each site index $i \in \{0, \dots, N-1\}$, the translated local term is determined by

$$(h_i \psi)(\sigma) = \begin{cases} [h_L(A)(\tau \mapsto \psi(\sigma^{[i, i+L-1] \leftarrow \tau}))](\sigma|_{[i, i+L-1]}), & L \leq N, \\ 0, & L > N. \end{cases}$$

Lemma 13.3.12 (Pointwise form of a translated local term). [Lean](#) [✓ Stmt](#) If $L \leq N$, then the translated term is obtained by applying the local parent interaction on the chosen cyclic window:

$$(h_i(A, L) \psi)(\sigma) = [h_L(A)(\tau \mapsto \psi(\sigma^{[i, i+L-1] \leftarrow \tau}))](\sigma|_{[i, i+L-1]}).$$

Proof. [✓ Proof](#) This is the first case in the definition of the translated local term. $\square$

Lemma 13.3.13 (Translated local terms are idempotent). [Lean](#) [✓ Stmt](#) For every site i ,

$$h_i(A, L)^2 = h_i(A, L).$$

Proof. ✓ Proof If $L > N$, the translated term is zero. Suppose $L \leq N$ and put

$$f_{\sigma,i}(\tau) = \psi(\sigma^{[i,i+L-1] \leftarrow \tau}).$$

The pointwise formula for translated terms, together with Lemmas 13.3.8 and 13.3.10, gives

$$(h_i(A, L)^2 \psi)(\sigma) = (h_L(A)^2 f_{\sigma,i})(\sigma|_{[i,i+L-1]}).$$

Lemma 13.3.4 turns the right-hand side into $(h_L(A) f_{\sigma,i})(\sigma|_{[i,i+L-1]})$, which is $(h_i(A, L) \psi)(\sigma)$. $\square$

Definition 13.3.14 (Parent Hamiltonian). Lean ✓ Stmt The chain Hamiltonian is

$$H_N(A, L) := \sum_{i=0}^{N-1} h_i.$$

Definition 13.3.15 (Frustration-free condition). Lean ✓ Stmt A vector ψ is a frustration-free ground state of the parent Hamiltonian when each local interaction annihilates it:

$$\forall i \in \{0, \dots, N-1\}, \quad h_i \psi = 0.$$

Lemma 13.3.16 (Periodic MPS vector window membership). Lean ✓ Stmt For any window of L consecutive sites starting at position i in an N -site periodic chain ($L \leq N$), the restriction of the periodic MPS vector $V^{(N)}(A)$ to that window lies in $G_L(A)$.

Proof. ✓ Proof The boundary matrix is the product of A -matrices on the complement sites. Trace cyclicity rotates the full N -site product so that window indices come first, matching the ground-space map definition. $\square$

Lemma 13.3.17 (Local term annihilates the periodic MPS vector). Lean ✓ Stmt For each site i , $h_i V^{(N)}(A) = 0$.

Proof. ✓ Proof Combines periodic-MPS-vector window membership with the fact that the parent interaction annihilates ground-space elements. $\square$

Lemma 13.3.18 (Parent Hamiltonian annihilates the periodic MPS vector). Lean ✓ Stmt For every N with $L \leq N$, the periodic MPS vector satisfies

$$H_N(A, L) V^{(N)}(A) = 0.$$

Proof. ✓ Proof Sum of zeros, since each local term annihilates the periodic MPS vector. $\square$

Lemma 13.3.19 (Frustration-freeness of the periodic MPS vector). Lean ✓ Stmt The periodic MPS vector is a frustration-free ground state: for every site i ,

$$h_i V^{(N)}(A) = 0.$$

Proof. ✓ Proof The claim follows from Lemma 13.3.17. $\square$

13.4 Intersection property and unique ground state

This section develops the intersection property of local ground spaces for injective MPS and the resulting uniqueness of the ground state on the periodic chain.

Given a state ψ on $L + 1$ sites, we define two restriction maps.

Definition 13.4.1.2 (Left restriction). Lean ✓ Stmt Fix the last physical index j :

$$\sigma \longmapsto \psi(\sigma_1, \dots, \sigma_L, j).$$

$$\sigma \mapsto \psi(i, \sigma_1, \dots, \sigma_L).$$

Definition 13.4.1.5 (Right ground membership). [Lean] [✓ Stmt] A state ψ on $L + 1$ sites satisfies the right ground condition if for every i , the state obtained by fixing the first index to i lies in $G_L(A)$.

Lemma 13.4.2.1 (Left restriction). [Lean] [✓ Stmt] If $\psi \in G_{L+1}(A)$, i.e. $\psi(\sigma) = \text{tr}(A^\sigma X)$ for some X , then for each j , the state obtained by fixing the last index to j lies in $G_L(A)$ with witness $A^j X$.

Diagram illustrating a quantum circuit structure. The top line shows qubits labeled $\sigma_1, \dots, \sigma_L$. The bottom line shows a sequence of operations: $A^{A^j} X$, followed by a red dot, and then L . The entire sequence is enclosed in large square brackets.

☐

Diagram illustrating a quantum circuit with L qubits. The top wire has control dots labeled $\sigma_1, \dots, \sigma_L$. The bottom wire has a target dot. A gate labeled $A^X A^i$ is applied between the first and last qubits.

☐

13.4.3 Injectivity of the ground-space map

Theorem 13.4.3.1 (Ground-space map injectivity). [Lean](#) [✓ Stmt](#) *If A is injective and $L \geq 1$, then Γ_L is injective.*

Proof. [✓ Proof](#) If $\Gamma_L(X) = 0$ then $\text{tr}(A^\sigma X) = 0$ for every length- L word σ . Since A is injective, the products $\{A^\sigma\}_\sigma$ span $M_D(\mathbb{C})$, so the trace pairing gives $X = 0$. $\square$

Theorem 13.4.3.2 (Ground-space map injectivity from word span). [Lean](#) [✓ Stmt](#) *If the length- L products $\{A^\sigma : |\sigma| = L\}$ span $M_D(\mathbb{C})$, then Γ_L is injective.*

Proof. [✓ Proof](#) If $\Gamma_L(X) = 0$, then $\text{tr}(A^\sigma X) = 0$ for every word σ of length L . The spanning hypothesis turns this into $\text{tr}(MX) = 0$ for every $M \in M_D(\mathbb{C})$, and the trace pairing gives $X = 0$. $\square$

Lemma 13.4.3.3 (Ground-space map injectivity after blocking). [Lean](#) [✓ Stmt](#) *If the length- L_0 products span the full matrix algebra,*

$$\text{span}\{A^\omega : |\omega| = L_0\} = M_D(\mathbb{C}),$$

then Γ_{L_0} is injective.

Proof. [✓ Proof](#) The hypothesis is precisely the fixed-length spanning condition needed by Theorem 13.4.3.2. $\square$

Theorem 13.4.3.4 (Ground-space dimension). [Lean](#) [✓ Stmt](#) *If A is injective and $L \geq 1$, then $\dim G_L(A) = D^2$.*


Proof. [✓ Proof](#) The upper bound $\dim G_L(A) \leq D^2$ is Lemma 13.2.4. Injectivity of Γ_L gives $\dim G_L(A) = \dim M_D(\mathbb{C}) = D^2$. $\square$

13.4.4 The intersection property

Theorem 13.4.4.1 (One inclusion of the intersection property). [Lean](#) [✓ Stmt](#) *If A is injective and $L \geq 2$, then the nontrivial inclusion in the intersection property holds:*

$$G_L^{\text{left}} \cap G_L^{\text{right}} \subseteq G_{L+1}(A).$$

Proof. [✓ Proof](#) The “inverting and growing back” argument compares the left and right restrictions on their common $(L-1)$ -site overlap and then reconstructs the full $(L+1)$ -site word from a single boundary matrix:



1. From the right ground condition, for each physical index i there exists a unique $Y_i \in M_D(\mathbb{C})$ (by injectivity of Γ_L) such that $\psi(i, \sigma) = \text{tr}(A^\sigma Y_i)$.
2. Similarly, from the left ground condition, for each j there exists a unique Z_j with $\psi(\sigma, j) = \text{tr}(A^\sigma Z_j)$.
3. Matching on the $(L-1)$ -site overlap and using nondegeneracy of the trace pairing: $A^j Y_i = Z_j A^i$ for all i, j .

4. Since A is injective, the matrices $\{A^j\}$ span $M_D(\mathbb{C})$ (Definition 2.4.1), so $\mathbb{1} = \sum_j c_j A^j$ for some coefficients $c_j \in \mathbb{C}$. Then $Y_i = \mathbb{1} \cdot Y_i = \sum_j c_j A^j Y_i = \sum_j c_j Z_j A^i = X' A^i$, where $X' := \sum_j c_j Z_j$.
5. By trace cyclicity, $\psi(\sigma) = \text{tr}(A^\sigma X')$, so $\psi \in G_{L+1}(A)$.

Theorem 13.4.4.2 (Intersection property). Lean ✓ Stmt For injective A and $L \geq 2$:

$$G_L^{\text{left}} \cap G_L^{\text{right}} = G_{L+1}(A),$$

equivalently $\psi \in G_{L+1}(A) \iff \psi \in G_L^{\text{left}} \cap G_L^{\text{right}}$.

Proof. ✓ Proof The claim follows from the two forward-direction lemmas and Theorem 13.4.4.1.

13.4.5 Unique ground state

Contiguous and cyclic interval restrictions

Remark 13.4.5.1 (Contiguous interval restrictions). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#)

✓ Stmt For a non-wrapping interval $[s, s + M - 1] \subseteq \{0, \dots, N - 1\}$ and a full configuration ρ supplying the complementary values, the restriction map is

$$(\mathrm{Res}_{s,M}^\rho \psi)(\omega) = \psi(\rho^{[s,s+M-1] \leftarrow \omega}).$$

The listed identities are the equations for $s = 0$, $M = N$, for deleting the first or last site, and for first fixing K sites and then restricting to $[s + K, s + K + L - 1]$.

Remark 13.4.5.2 (Cyclic interval restrictions). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) On the periodic chain, for a full configuration ρ supplying the complementary values, put

$$\delta_i(k) := (k + N - i) \bmod N, \quad \rho_{i,L,\omega}^{\text{cyc}} := \begin{cases} \omega(\delta_i(k)), & \delta_i(k) < L, \\ \rho(k), & \delta_i(k) \geq L. \end{cases}$$

The restriction to the cyclic interval beginning at i is

$$(\mathrm{Res}_{i,L}^\rho \psi)(\omega) = \psi(\rho_{i,L,\omega}^{\mathrm{cyc}}),$$

with all site labels read modulo N . If $L \leq N$, this is the same configuration as $\rho^{[i, i+L-1] \leftarrow \omega}$. If two boundary conditions agree whenever $\delta_i(k) \geq L$, their restrictions are equal. In particular, filling the interval inside the outside configuration leaves the restriction unchanged:

$$\mathrm{Res}_{i,L}^{\rho_{i,L,\omega}^{\mathrm{cyc}}} \psi = \mathrm{Res}_{i,L}^{\rho} \psi.$$

For every $r < L$, the inserted word is recovered:

$$\rho_{i,L,\omega}^{\text{cyc}}(i+r) = \omega(r).$$

Here site labels are taken modulo N . The remaining listed identities say that deleting the final site of an $(L + 1)$ -interval gives the L -interval beginning at i ; if $L + 1 \leq N$, deleting the first site gives the L -interval beginning at $i + 1$; and the cyclic formula agrees with the contiguous formula when the interval does not wrap around the ring.

Lemma 13.4.5.3 (One-site restrictions of cyclic-window boundary conditions). [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *If a cyclic $(L+1)$ -window is represented by a boundary matrix Y , then deleting one endpoint gives*

$$\Gamma_{L+1}(Y)|_{last=b} = \Gamma_L(A^b Y), \quad \Gamma_{L+1}(Y)|_{first=a} = \Gamma_L(Y A^a).$$

Hence two adjacent cyclic windows whose restrictions agree on the common length- L overlap have boundary matrices satisfying

$$Y_1 A^a = A^b Y_2.$$

Consequently, for a finite chain of adjacent overlaps indexed by $0 \leq r < n$,

$$Y_r A^{a_r} = A^{b_r} Y_{r+1},$$

one has

$$Y_0 A^{a_0} \dots A^{a_{n-1}} = A^{b_0} \dots A^{b_{n-1}} Y_n.$$

In particular, if the two endpoint words are named by the equations

$$a_r = \rho(i_0 + r), \quad b_r = \rho(i_0 + r + L + 1),$$

with site labels read modulo N , then the same transport identity is written as

$$Y_0 A^a = A^b Y_n.$$

Proof. ✓ Proof The first two identities are the cyclic first- and last-site deletion identities, followed by the corresponding ground-space restriction formulas. For adjacent windows, equality of the completed outside configurations gives the common-overlap equality

$$\Gamma_{L_i}(Y_0 A^a) = \Gamma_{L_i}(A^b Y_n).$$

Injectivity of Γ_L gives $Y_0 A^a = A^b Y_n$. Iterating the displayed adjacent identity gives the word-product identity. $\square$

Lemma 13.4.5.4 (Iterated contiguous-window intersection). [Lean] ✓ Stmt *If an N -site state satisfies the L -site ground condition on every non-wrapping contiguous window (with $L \geq 2$), then it lies in $G_N(A)$.*

Proof. ✔ Proof The induction repeatedly merges two neighbouring admissible windows $[i, i+L-1]$ and $[i+1, i+L]$ into the larger window $[i, i+L]$ by applying Theorem 13.4.4.1 to their common overlap $[i+1, i+L-1]$. $\square$

Definition 13.4.5.5 (Suffix restriction). [Lean] [✓ Stmt] Fix a prefix $u \in \{0, \dots, d-1\}^K$. The suffix restriction of an $(K+L)$ -site state ψ is the L -site state given by

$$(R_u^{\text{tail}}\psi)(\sigma) = \psi(u, \sigma).$$

Definition 13.4.5.6 (Suffix ground membership). [Lean] [Stm] A state ψ on $K + L$ sites lies in the suffix ground condition if every fixed prefix u gives a suffix state in $G_L(A)$.

Lemma 13.4.5.7 (Ground-space suffix restriction). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) If $\psi \in G_{K+L}(A)$, then every fixed-prefix suffix restriction of ψ lies in $G_L(A)$. For a vector in $G_{L+1}(A)$, fixing either endpoint gives the expected left or right multiplication of its boundary matrix.

Proof. ✓ Proof Write $\psi(\rho) = \text{tr}(A^\rho X)$ for a boundary matrix X . Fixing the final site gives the boundary matrix $A^j X$, while fixing the first site gives $X A^i$ by cyclicity of the trace. Fixing a prefix u gives

$$\psi(u, \sigma) = \text{tr}(A^u A^\sigma X) = \text{tr}(A^\sigma X A^u),$$

so the suffix restriction again has the form of a ground-space vector. □

Lemma 13.4.5.8 (Left-tail compatibility). Lean ✓ Stmt Assume ψ on $K + L_0 + 1$ sites satisfies the left ground condition on the first $K + L_0$ sites and the suffix ground condition on every fixed prefix of length K . Then there exist matrices $(Z_j)_j$ and $(Y_u)_u$ such that

$$\psi(\sigma, j) = \text{tr}(A^\sigma Z_j), \quad \psi(u, \tau) = \text{tr}(A^\tau Y_u),$$

and for all j and u ,

$$Z_j A^u = A^j Y_u.$$

Proof. ✓ Proof Choose Z_j from the long left-window condition and Y_u from the suffix ground condition. The two restrictions give the same vector in $G_{L_0}(A)$:

$$\Gamma_{L_0}(Z_j A^u) = \Gamma_{L_0}(A^j Y_u).$$

Injectivity of Γ_{L_0} , obtained from L_0 -block injectivity, gives $Z_j A^u = A^j Y_u$. □

Theorem 13.4.5.9 (Inverting step for word compatibility). Lean ✓ Stmt Assume A is L_0 -block-injective with $L_0 > 0$. If matrices $(Z_j)_j$ satisfy

$$Z_j A^\sigma = A^j Y_\sigma$$

for every word σ of length K , then there exists a matrix X such that $Z_j = A^j X$ for all j .

Proof. ✓ Proof For $K = 1$, the hypothesis gives $Z_j A^i = A^j Y_i$ for every letter i . Every blocked word of length L_0 is nonempty, so this identity extends to the coefficients of the blocked tensor $A^{[L_0]}$. Since $A^{[L_0]}$ is injective, the identity matrix is a linear combination of those blocked coefficients, and substituting this decomposition yields $Z_j = A^j X$ for a common matrix X . For larger K , strip the first letter: for each i , the matrices $Z_j A^i$ satisfy the same compatibility hypothesis with words of length $K - 1$, so the induction hypothesis reduces the problem to the case $K = 1$. □

Theorem 13.4.5.10 (Growing-back step at the injectivity length). Lean ✓ Stmt Assume A is L_0 -block-injective with $L_0 > 0$. If an element ψ on $K + L_0 + 1$ sites satisfies the left ground condition on the first $K + L_0$ sites and the suffix ground condition on every prefix of length K , then $\psi \in G_{K+L_0+1}(A)$.

Proof. ✓ Proof Lemma 13.4.5.8 produces matrices $(Z_j)_j$ and $(Y_u)_u$ with $Z_j A^u = A^j Y_u$ for every prefix word u of length K . Theorem 13.4.5.9 gives a common right factor X such that $Z_j = A^j X$ for all j . Therefore each last-site restriction of ψ agrees with the corresponding last-site restriction of the ground-space vector determined by X . Since every configuration is obtained by adjoining its final letter to its initial $(K + L_0)$ -tuple, the two states coincide, so ψ lies in $G_{K+L_0+1}(A)$. □

Theorem 13.4.5.11 (Open-chain iteration of the intersection property). Lean ✓ Stmt Let A be L_0 -block-injective with $D \geq 1$ and $L_0 > 0$. If an N -site state satisfies the ground-space constraint on every non-wrapping contiguous interval of length $L_0 + 1$, for every fixed choice of the complementary sites, with $N \geq L_0 + 1$, then it lies in $G_N(A)$.

Proof. ✓ Proof Induct on the extra length beyond $L_0 + 1$. The induction hypothesis gives the long left-window ground condition after deleting the last site, while Definition 13.4.5.5 and the contiguous-interval restriction identities above identify every fixed-prefix suffix with one of the assumed length- $L_0 + 1$ intervals. Theorem 13.4.5.10 then grows back the rightmost site. $\square$

Lemma 13.4.5.12 (Open-chain iteration for a sequence of subspaces). Lean ✓ Stmt Let $S_m \subseteq (\mathbb{C}^d)^{\otimes m}$ be subspaces, and let $0 < L \leq N$. Suppose that every non-wrapping length- L interval of ψ , for every fixed choice of the complementary sites, lies in S_L . Suppose also that, for every $m \geq L$,

$$\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d = S_{m+1}.$$

Then $\psi \in S_N$.

Proof. ✓ Proof Induct on the length m of a non-wrapping interval. The case $m = L$ is the assumed local condition. If all length- m intervals lie in S_m , then the two one-site restrictions of a length- $m + 1$ interval are adjacent length- m intervals. Hence the length- $m + 1$ interval lies in $\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d$, and the displayed identity places it in S_{m+1} . Taking $m = N$ and the interval starting at the first site gives $\psi \in S_N$. $\square$

Lemma 13.4.5.13 (Periodic constraints and propagated subspaces). Lean ✓ Stmt Let $S_m \subseteq (\mathbb{C}^d)^{\otimes m}$ be subspaces, and let $0 < L \leq N$. Suppose that $G_L(A) \subseteq S_L$. Suppose also that, for every $m \geq L$,

$$\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d = S_{m+1}.$$

Then the periodic local constraints imply

$$\mathcal{G}_{N,L}(A) \subseteq S_N.$$

Proof. ✓ Proof If $\psi \in \mathcal{G}_{N,L}(A)$, then every length- L cyclic interval of ψ lies in $G_L(A)$, and hence in S_L . A non-wrapping interval is the cyclic interval starting at the same site. Therefore all non-wrapping length- L intervals lie in S_L , and Lemma 13.4.5.12 gives $\psi \in S_N$. $\square$

Lemma 13.4.5.14 (Cyclic-window range antitonicity). Lean Lean ✓ Stmt On a nonempty periodic chain, longer cyclic-window constraints imply all shorter cyclic-window constraints: if $L' \leq L \leq N$, then $\mathcal{G}_{N,L}(A) \subseteq \mathcal{G}_{N,L'}(A)$.

Proof. ✓ Proof For every cyclic window vector $v \in G_{L+1}(A)$ and every last letter b ,

$$v|_{\text{last}=b} \in G_L(A).$$

Thus $\mathcal{G}_{N,L+1}(A) \subseteq \mathcal{G}_{N,L}(A)$, and iteration gives the general antitonicity. $\square$

Theorem 13.4.5.15 (Open-chain containment from cyclic local constraints). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$. If an N -site periodic-chain state satisfies all cyclic ground-space constraints at some range $L_0 + 1 \leq L \leq N$, then it lies in the open-chain ground space $G_N(A)$.

Proof. ✓ Proof Lemma 13.4.5.14 reduces the cyclic hypotheses to range $L_0 + 1$. On every non-wrapping interval, the cyclic and contiguous restriction maps agree, so Theorem 13.4.5.11 applies. $\square$

13.5 Periodic-boundary comparison for injective tensors

On the periodic ring the two cyclic supports that cross the periodic-boundary cut expose the same complementary word, and matching their boundary matrices forces X to commute with every A^j . This section derives the boundary-crossing compatibilities $C_\tau^+ A^j X = Y_\tau A^j$ and $X A^j C_\tau^- = A^j Y_\tau$ and combines them into the commutation $X A^j = A^j X$.

Lemma 13.5.1 (Boundary-crossing compatibility at the last site). Lean ✓ Stmt Assume A is L_0 -block-injective with $L_0 > 0$ and $M \geq L_0$. If the reduced cyclic interval crossing the last site has boundary matrices Y_τ , then for every physical letter j one has

$$C_\tau^+ A^j X = Y_\tau A^j, \quad (13.1)$$

where C_τ^+ is the complementary word seen by that cyclic interval.

Proof. ✓ Proof The trace identity for this cyclic interval is tested against all words of length L_0 . Injectivity of Γ_{L_0} , which follows from L_0 -block injectivity, turns these test-word identities into the displayed matrix identity (13.1). $\square$

Lemma 13.5.2 (Boundary-crossing compatibility at the second boundary position). Lean ✓ Stmt Assume A is L_0 -block-injective with $L_0 > 0$ and $M \geq L_0$. If the second boundary-crossing reduced cyclic interval has boundary matrices Y_τ , then

$$X A^j C_\tau^- = A^j Y_\tau$$

for every physical letter j , where C_τ^- is the complementary word seen from the second cyclic position.

Proof. ✓ Proof Factor the cyclic word at the second cyclic position, rotate the trace, and use injectivity of Γ_{L_0} to remove the length- L_0 test word. $\square$

Theorem 13.5.3 (The two boundary-crossing compatibilities). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$. Suppose $\psi \in \mathcal{G}_{N,L}(A)$, $N \geq 2$, $L_0 < L \leq N$, and $\psi = \Gamma_N(X)$ for a boundary matrix X . Then there are two families of boundary matrices Y_τ^+ and Y_τ^- , indexed by fixed complement values τ , such that for every physical letter j ,

$$C_\tau^+ A^j X = Y_\tau^+ A^j, \quad X A^j C_\tau^- = A^j Y_\tau^-, \quad (13.2)$$

where C_τ^+ and C_τ^- are the complementary words exposed by the two boundary-crossing cyclic intervals, both of reduced length $L_0 + 1$.

Proof. ✓ Proof First use Lemma 13.4.5.14 to reduce the cyclic constraints from range L to range $L_0 + 1$. Then apply the two cyclic-interval compatibility lemmas at the two boundary positions to obtain the displayed one-sided identities (13.2). $\square$

Lemma 13.5.4 (Boundary-crossing equations for $\Gamma_{M+1}(X)$). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$, $D \geq 1$, and $L_0 \leq M$. Suppose $\psi = \Gamma_{M+1}(X)$, and suppose matrices $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Then, for every physical letter j and boundary condition τ ,

$$A^{\tau L_0} \dots A^{\tau M-1} A^j X = Y_M(\tau) A^j, \quad X A^j A^{\tau_1} \dots A^{\tau_{M-L_0}} = A^j Y_{M+1-L_0}(\tau). \quad (13.3)$$

Proof. [✓ Proof](#) Substitute $\psi = \Gamma_{M+1}(X)$ in the two cyclic restrictions at the last site and at the second boundary-crossing support. Lemmas [13.5.1](#) and [13.5.2](#) then give the two equations in [\(13.3\)](#). $\square$

Lemma 13.5.5 (One-sided products for a common complement word). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$, $D \geq 1$, and $L_0 \leq M$. Suppose $\psi = \Gamma_{M+1}(X)$, and suppose matrices $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions*

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

For every boundary letter η , every physical letter j , and every complementary word μ ,

$$Y_M(\tau_\eta^+(\mu))A^j = A^\mu A^j X, \quad XA^j A^\mu = A^j Y_{M+1-L_0}(\tau_\eta^-(\mu)).$$

Proof. [✓ Proof](#) Substitute $\tau_\eta^+(\mu)$ and $\tau_\eta^-(\mu)$ into Lemma [13.5.4](#). The exposed-complement identities of Lemma [13.5.6](#) identify both complementary products with A^μ . $\square$

Lemma 13.5.6 (Common complement word for the two boundary supports). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *On a chain of length $N = M + 1$, with $L_0 \leq M$, fix a physical letter η used outside the complementary sites and a word μ of length $M + 1 - (L_0 + 1)$. Write $\tau_\eta^+(\mu)$ for the boundary condition at the support crossing the last site and $\tau_\eta^-(\mu)$ for the boundary condition at the second boundary-crossing support. Their exposed complements are both exactly μ .*

Proof. [✓ Proof](#) Put the same letter η on the remaining sites. Fill the physical sites $L_0, \dots, M - 1$ with μ for the support crossing the last site, and fill the sites $1, \dots, M - L_0$ with μ for the second boundary-crossing support. The two complement-extraction identities follow from the index arithmetic. The same outside-site calculation gives, for any ρ with $\rho_{k+L_0} = \mu_k$,

$$\text{Res}_{M, L_0+1}^\rho \psi = \text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)} \psi,$$

and, for any ρ with $\rho_{k+1} = \mu_k$,

$$\text{Res}_{M+1-L_0, L_0+1}^\rho \psi = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)} \psi.$$

$\square$

Lemma 13.5.7 (Boundary equations from a common complement word). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective, with $L_0 > 0$, and let $L_0 \leq M$. Suppose two length- $(L_0 + 1)$ restrictions of a state ψ at the same cyclic support are represented by*

$$\text{Res}_{i, L_0+1}^\rho \psi = \Gamma_{L_0+1}(Y_\rho), \quad \text{Res}_{i, L_0+1}^\tau \psi = \Gamma_{L_0+1}(Y_\tau).$$

If these two restrictions are equal, then for every physical letter j ,

$$Y_\rho A^j = Y_\tau A^j.$$

In particular, the last-site boundary-crossing condition satisfies

$$\rho_{k+L_0} = \mu_k \implies Y_\rho A^j = Y_{\tau_\eta^+(\mu)} A^j,$$

and the second boundary-crossing condition satisfies

$$\rho_{k+1} = \mu_k \implies Y_\rho A^j = Y_{\tau_\eta^-(\mu)} A^j.$$

Consequently, for every word σ of length L_0 , the two displayed implications remain true after right multiplication by A^σ .

Proof. ✓ Proof Apply the first-letter restriction to both equal length- (L_0+1) restrictions. The two resulting length- L_0 vectors are represented by $\Gamma_{L_0}(Y_\rho A^j)$ and $\Gamma_{L_0}(Y_\tau A^j)$, and block injectivity makes Γ_{L_0} injective. The two displayed special cases follow from Lemma 13.5.6. The length- L_0 word forms are obtained by multiplying these equations on the right by A^σ . $\square$

Lemma 13.5.8 (Reduction to compatible boundary conditions). Lean ✓ Stmt Let A be L_0 -block-injective, with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$, and let the matrices $Y_i(\tau)$ represent its length- (L_0+1) cyclic restrictions:

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Fix a boundary letter η and a complementary word μ . Let ρ^+ and ρ^- be two boundary conditions satisfying

$$\rho_{k+L_0}^+ = \mu_k, \quad \rho_{k+1}^- = \mu_k.$$

If the adjacent-window comparison gives, for every physical letter j and every word σ of length L_0 ,

$$Y_M(\rho^+)A^jA^\sigma = Y_{M+1-L_0}(\rho^-)A^jA^\sigma,$$

then the same product equation holds for $\tau_\eta^+(\mu)$ and $\tau_\eta^-(\mu)$:

$$Y_M(\tau_\eta^+(\mu))A^jA^\sigma = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma.$$

Proof. ✓ Proof By Lemma 13.5.7,

$$Y_M(\rho^+)A^j = Y_M(\tau_\eta^+(\mu))A^j, \quad Y_{M+1-L_0}(\rho^-)A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Multiplying on the right by A^σ and applying the hypothesis gives the displayed product equation. $\square$

Theorem 13.5.9 (Pointwise reduction to compatible boundary conditions). Lean ✓ Stmt Let A be L_0 -block-injective, with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$, and let the matrices $Y_i(\tau)$ represent its length- (L_0+1) cyclic restrictions:

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Fix a boundary letter η and a complementary word μ . Suppose that, for each physical letter j and word σ of length L_0 , boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ satisfy, for every complementary position k ,

$$\rho_{j,\sigma}^+(k+L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k+1) = \mu_k.$$

If, for every j and σ ,

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma,$$

then, for every j and σ ,

$$Y_M(\tau_\eta^+(\mu))A^jA^\sigma = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma.$$

Proof. ✓ Proof Fix j and σ . Lemma 13.5.7 gives

$$Y_M(\rho_{j,\sigma}^+)A^j = Y_M(\tau_\eta^+(\mu))A^j, \quad Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Multiply the two identities on the right by A^σ and use the assumed product equation for the chosen pair j, σ . $\square$

Lemma 13.5.10 (Endpoint words for the periodic-boundary windows). [Lean](#) [Stmnt](#) Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$. Suppose that matrices $Y_r(\rho)$, for $0 \leq r \leq L_0 - 1$, represent the length- $(L_0 + 1)$ cyclic restrictions beginning at the sites $M+1-L_0+r$:

$$\text{Res}_{M+1-L_0+r, L_0+1}^\rho \psi = \Gamma_{L_0+1}(Y_r(\rho)).$$

Then

$$Y_0(\rho) A^{\rho_{M+1-L_0}} \dots A^{\rho_{M-1}} = A^{\rho_1} \dots A^{\rho_{L_0-1}} Y_{L_0-1}(\rho).$$

For $L_0 = 1$, both products are empty.

Proof. [Stmnt](#) Use Lemma 13.4.5.3 for the $L_0 - 1$ adjacent restrictions starting at $i_0 = M+1-L_0$. For $0 \leq r < L_0 - 1$, the two endpoint words are determined by

$$i_0 + r = M + 1 - L_0 + r, \quad i_0 + r + L_0 + 1 \equiv r + 1 \pmod{M + 1},$$

Substituting these two words in the iterated transport identity gives the displayed product equation. $\square$

Lemma 13.5.11 (Boundary-condition product for the periodic-boundary windows). [Lean](#) [Lean](#) [Stmnt](#) Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$, and suppose matrices $Y_i(\rho)$ represent the length- $(L_0 + 1)$ cyclic restrictions

$$\text{Res}_{i, L_0+1}^\rho(\psi) = \Gamma_{L_0+1}(Y_i(\rho)).$$

Then, for every boundary condition ρ ,

$$Y_{M+1-L_0}(\rho) A^{\rho_{M+1-L_0}} \dots A^{\rho_{M-1}} = A^{\rho_1} \dots A^{\rho_{L_0-1}} Y_M(\rho).$$

For $L_0 = 1$, both products are empty. The same equation remains true after multiplying both sides on the right by any matrix R .

Proof. [Stmnt](#) Apply Lemma 13.4.5.3 to the $L_0 - 1$ restrictions beginning at $M+1-L_0+r$, with

$$Y_r(\rho) = Y_{M+1-L_0+r}(\rho).$$

With $i_0 = M+1-L_0$, the site labels satisfy

$$i_0 + r = M + 1 - L_0 + r, \quad i_0 + r + L_0 + 1 \equiv r + 1 \pmod{M + 1}$$

hence, for $r = 0, \dots, L_0 - 2$,

$$Y_{M+1-L_0+r}(\rho) A^{\rho_{M+1-L_0+r}} = A^{\rho_{r+1}} Y_{M+2-L_0+r}(\rho).$$

Multiplying these $L_0 - 1$ equations from left to right gives the displayed product equation. The right-multiplied form follows by multiplying this equality by R . $\square$

Lemma 13.5.12 (Transport followed by the one-sided boundary equation). [Lean](#) [Stmnt](#) Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$, and let $L_0 \leq M$. Let $\psi = \Gamma_{M+1}(X)$, and suppose that matrices $Y_i(\rho)$ represent all length- $(L_0 + 1)$ cyclic restrictions of ψ . Then, for every boundary condition ρ and every physical letter j ,

$$Y_{M+1-L_0}(\rho) A^{\rho_{M+1-L_0}} \dots A^{\rho_{M-1}} A^j = A^{\rho_1} \dots A^{\rho_{L_0-1}} A^{\rho_{L_0}} \dots A^{\rho_{M-1}} A^j X.$$

For $\rho = \tau_\eta^-(\mu)$ this is the transport identity supplied by the adjacent-window argument, with the single factor A^j following $Y_{M+1-L_0}(\tau_\eta^-(\mu))$. The padded identity, whose corresponding factor is $A^j A^\sigma$, is supplied separately.

Proof. [✓ Proof](#) Lemma 13.5.11 gives

$$Y_{M+1-L_0}(\rho)A^{\rho_{M+1-L_0}} \dots A^{\rho_{M-1}}A^j = A^{\rho_1} \dots A^{\rho_{L_0-1}}Y_M(\rho)A^j.$$

The one-sided equation for the window beginning at M gives

$$Y_M(\rho)A^j = A^{\rho_{L_0}} \dots A^{\rho_{M-1}}A^jX.$$

Substitution gives the displayed formula. $\square$

Lemma 13.5.13 (Long boundary-condition product for the periodic-boundary windows). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M + 1$, and suppose matrices $Y_i(\rho)$ represent the length- $(L_0 + 1)$ cyclic restrictions*

$$\text{Res}_{i, L_0+1}^\rho(\psi) = \Gamma_{L_0+1}(Y_i(\rho)).$$

Then, for every boundary condition ρ ,

$$Y_M(\rho)A^{\rho_M}A^{\rho_0} \dots A^{\rho_{M-L_0}} = A^{\rho_{L_0}} \dots A^{\rho_M}A^{\rho_0}Y_{M+1-L_0}(\rho).$$

Each side has $M + 2 - L_0$ one-site factors, with site labels read modulo $M + 1$.

Proof. [✓ Proof](#) Apply Lemma 13.4.5.3 to the $M + 2 - L_0$ restrictions beginning at $M + r$ modulo $M + 1$. With $i_0 = M$, the site labels satisfy

$$i_0 + r \equiv M + r, \quad i_0 + r + L_0 + 1 \equiv L_0 + r \pmod{M + 1}.$$

The final window is

$$i_0 + M + 2 - L_0 \equiv M + 1 - L_0 \pmod{M + 1}.$$

Substituting these site labels in the iterated product identity gives the displayed equation. $\square$

Lemma 13.5.14 (One matrix Y_μ from compared boundary matrices). [Lean](#) [✓ Stmt](#) *Fix a physical letter η used outside the complementary sites. Suppose the two one-sided boundary matrix families Y^+ and Y^- have been extracted from the cyclic windows. If, for every word on the complementary sites μ , these boundary matrices agree on the boundary conditions built from η ,*

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-,$$

then there is a single family Y_μ satisfying

$$A^\mu A^b X = Y_\mu A^b, \quad X A^a A^\mu = A^a Y_\mu$$

for all letters a, b .

Proof. [✓ Proof](#) Define Y_μ to be the first boundary matrix evaluated on the boundary condition $\tau_\eta^+(\mu)$. Lemma 13.5.6 rewrites the first cyclic-window complement as μ , while the assumed comparison at $\tau_\eta^-(\mu)$ rewrites the second boundary matrix to the same Y_μ . $\square$

Lemma 13.5.15 (Commutation from two one-sided equations). [Lean](#) [✓ Stmt](#) *Suppose a single family of boundary matrices Y_μ , indexed by words μ on the complementary sites, satisfies the two one-sided identities*

$$A^\mu A^b X = Y_\mu A^b, \quad X A^a A^\mu = A^a Y_\mu \tag{13.4}$$

for every pair of physical letters a, b . Then X commutes with every word product $A^a A^\mu A^b$ obtained by adjoining one letter on each side of the word A^μ .

Proof. ✓ Proof Multiply the second identity in (13.4) on the right by A^b and the first identity on the left by A^a :

$$XA^a A^\mu A^b = A^a Y_\mu A^b = A^a A^\mu A^b X.$$

□

Theorem 13.5.16 (Boundary matrix commutation from periodic closure). Lean ✓ Stmt Let A be injective with $D \geq 1$, and let $\psi = \Gamma_N(X)$ for some boundary matrix $X \in M_D(\mathbb{C})$. If every cyclic restriction of length L of ψ lies in $G_L(A)$, with $N \geq 2$ and $1 < L \leq N$, then, for every $j = 0, \dots, d-1$,

$$XA^j = A^j X$$

Proof. ✓ Proof The periodic boundary condition produces matrix identities of the form $XMW = MY$ for arbitrary test matrices M and complementary words W . Spanning first over the length- $(L-1)$ tails and then over the complementary words, injectivity forces $(XM - MX)W = 0$ for every W , hence $XM = MX$ for every matrix M . In particular, X commutes with each letter A^j . □

13.6 Periodic unique ground state

This section gives three periodic ground-space facts: the span of the periodic MPS vector $V^{(N)}(A) = \Gamma_N(\mathbb{1})$, its membership in every cyclic window constraint $\mathcal{G}_{N,L}(A)$, and its nonvanishing for block-injective tensors. Together with the boundary-matrix commutation $XA^j = A^j X$ of the previous section, these establish that the periodic ground space is spanned by $V^{(N)}(A)$; the one-dimensionality conclusion is completed downstream.

Definition 13.6.1 (Span of the periodic MPS vector). Lean ✓ Stmt The subspace spanned by the MPS vector $V^{(N)}(A)(\sigma) = \text{tr}(A^{\sigma_0} \dots A^{\sigma_{N-1}})$.

Lemma 13.6.2 (The periodic MPS vector is the ground-space map at the identity). Lean ✓ Stmt For every chain length N ,

$$V^{(N)}(A) = \Gamma_N(\mathbb{1}).$$

Lemma 13.6.3 (The periodic MPS vector lies in the local ground space). Lean ✓ Stmt $V^{(N)}(A) \in G_N(A)$ for every N , with boundary matrix $X = I$.

Proof. ✓ Proof $V^{(N)}(A)(\sigma) = \text{tr}(A^\sigma) = \text{tr}(A^\sigma \cdot I) = \Gamma_N(I)(\sigma)$. □

Definition 13.6.4 (One-dimensional ground space). Lean ✓ Stmt A finite-dimensional ground space S is one-dimensional if $\dim S = 1$.

Theorem 13.6.5 (One-dimensionality criterion). Lean ✓ Stmt For finite-dimensional S , $\dim S = 1$ iff there exists a nonzero $\psi_0 \in S$ such that every $\psi \in S$ is of the form $c\psi_0$.

Proof. ✓ Proof Apply the standard finite-dimensional criterion that a nonzero space has dimension one exactly when all of its vectors are proportional to a single nonzero vector. □

Definition 13.6.6 (Periodic parent-Hamiltonian ground space). Lean ✓ Stmt For $N > 0$ and $L \leq N$, define the *periodic parent-Hamiltonian ground space*

$$\mathcal{G}_{N,L}(A) := \{\psi \in (\{0, \dots, d-1\}^N \rightarrow \mathbb{C}) : \forall i \in \{0, \dots, N-1\}, \psi|_{[i, i+L-1]} \in G_L(A)\},$$

where the condition $\psi|_{[i, i+L-1]} \in G_L(A)$ means that for every choice of physical indices outside the window, the restriction of ψ to that window lies in $G_L(A)$. The one-index notation $G_L(A)$

denotes the local ground space $\mathcal{G}_L$ of [CPGSV21, Section IV.C]. The two-index notation $\mathcal{G}_{N,L}(A)$ used here denotes the periodic intersection of those local constraints over all translated length- L windows on the N -site ring. When $N = 0$ or $L > N$, set $\mathcal{G}_{N,L}(A) := \top$ by convention.

Lemma 13.6.7 (Cyclic-window representation matrices inside the periodic ground space). [Lean](#)

[✓ Stmt](#) *If $0 < N$, $L \leq N$, and $\psi \in \mathcal{G}_{N,L}(A)$, then for every cyclic window and every outside configuration there is a boundary matrix Y such that the corresponding restricted L -site state is $\Gamma_L(Y)$.*

Proof. [✓ Proof](#) This is the defining cyclic-window condition for membership in $\mathcal{G}_{N,L}(A)$, together with the definition of the local ground space as the range of Γ_L . $\square$

Lemma 13.6.8 (The periodic MPS vector satisfies every cyclic window constraint). [Lean](#) [✓ Stmt](#)

If $0 < N$ and $L \leq N$, then

$$V^{(N)}(A) \in \mathcal{G}_{N,L}(A).$$

Theorem 13.6.9 (Periodic MPS vector nonvanishing for block-injective tensors). [Lean](#) [✓ Stmt](#)

If A is L_0 -block-injective with $L_0 > 0$, $D \geq 1$, and $N \geq L_0 + 1$, then $V^{(N)}(A) \neq 0$ in the N -site space.

Proof. [✓ Proof](#) Suppose for contradiction that $V^{(N)}(A) = 0$, i.e. $\text{tr}(A^\sigma) = 0$ for every word σ of length N . For any words w and u with $|w| = N - L_0$ and $|u| = L_0$, we have $\text{tr}(A^w A^u) = 0$. Since A is L_0 -block-injective, $\{A^u : |u| = L_0\}$ spans $M_D(\mathbb{C})$. By nondegeneracy of the trace pairing, $A^w = 0$ for every $|w| = N - L_0$. Repeating the argument with words of length $N - 2L_0$ (since $|w'| + L_0 = N - L_0$, the product $A^{w'} A^{u_1}$ has length $N - L_0$, so $A^{w'} A^{u_1} = 0$ by the previous step) and iterating, we eventually reach $\text{tr}(A^u) = 0$ for $|u| = L_0$, which gives $I = 0$, a contradiction. $\square$

13.7 Injectivity-length reduction for the closure property

This section implements the inverting-and-growing-back reduction from [CPGSV21, Section IV.C]: a commutation or annihilation relation for length- m word products is reduced to the injectivity length L_0 , then used in the periodic-boundary closure-property step. Throughout, $A^w := A^{i_1} \cdots A^{i_L}$ denotes the product along a word $w = (i_1, \dots, i_L)$, Γ_N sends a boundary matrix to its open-chain vector, and A being L_0 -block-injective means $\text{span}\{A^u : |u| = L_0\} = M_D(\mathbb{C})$, so a relation valid on every length- L_0 product extends by linearity to all of $M_D(\mathbb{C})$.

Theorem 13.7.1 (Common right factor from suffix equations). [Lean](#) [✓ Stmt](#) *Assume A is L_0 -block-injective with $L_0 > 0$. Let $\{Z_u\}_{|u|=L_0}$ be a family indexed by words of length L_0 . If for some $K \geq 0$ and every suffix word w of length K there exists $Y_w \in M_D(\mathbb{C})$ such that, for every word u of length L_0 ,*

$$Z_u A^w = A^u Y_w,$$

then there exists $X \in M_D(\mathbb{C})$ with $Z_u = A^u X$ for every word u of length L_0 .

Proof. [✓ Proof](#) Induct on K . For $K = 0$ the empty suffix gives $Z_u = A^u Y_\emptyset$. For the successor step, fix the first physical index i and apply the induction hypothesis to the shortened suffix. This gives a matrix X_i satisfying

$$Z_u A^i = A^u X_i$$

for every word u of length L_0 . Extending this letter equation to all length- L_0 words and decomposing $\mathbb{1}$ in the spanning family $\{A^u : |u| = L_0\}$ yields a common factor X . $\square$

Theorem 13.7.2 (Common boundary matrix from spanning word identities). [Lean](#) [✓ Stmt](#) Let A be a tensor whose length- K word products span $M_D(\mathbb{C})$. Let $\{F_b\}_{b \in B}$ and $\{Z_b\}_{b \in B}$ be two families of matrices. If, for every length- K word w , there is a matrix Y_w such that, for every $b \in B$,

$$Z_b A^w = F_b Y_w,$$

then there is a single matrix Y such that, for every $b \in B$,

$$Z_b = F_b Y.$$

Proof. [✓ Proof](#) Suppose $Z_b M_1 = F_b Y_1$ and $Z_b M_2 = F_b Y_2$ for every $b \in B$. Then

$$Z_b(M_1 + M_2) = Z_b M_1 + Z_b M_2 = F_b(Y_1 + Y_2).$$

For a scalar c ,

$$Z_b(cM_1) = cZ_b M_1 = F_b(cY_1).$$

Thus the displayed identity holds for every matrix in the span of the length- K word products. Since this span is all of $M_D(\mathbb{C})$, apply it to the identity matrix. $\square$

Theorem 13.7.3 (Injectivity-length commutation from length- m commutation). [Lean](#) [✓ Stmt](#) If A is L_0 -block-injective with $L_0 > 0$, $m \geq L_0$, and $X \in M_D(\mathbb{C})$ commutes with every length- m product A^w , then X already commutes with every length- L_0 product A^u .

Proof. [✓ Proof](#) Write each length- m word as a concatenation uw with $|u| = L_0$ and $|w| = m - L_0$. The hypothesis gives

$$X A^u A^w = A^u A^w X = A^u (A^w X).$$

Thus the family $Z_u := X A^u$ satisfies the hypothesis of Theorem 13.7.1. One obtains $X A^u = A^u X$ for all $|u| = L_0$. Decomposing $\mathbb{1}$ in the span of the length- L_0 words shows $R = X$, hence $X A^u = A^u X$. $\square$

Theorem 13.7.4 (Centrality from length- m commutation). [Lean](#) [✓ Stmt](#) If A is L_0 -block-injective with $L_0 > 0$, $m \geq L_0$, and $X \in M_D(\mathbb{C})$ commutes with every length- m word product, then X commutes with every matrix in $M_D(\mathbb{C})$.

Proof. [✓ Proof](#) By Theorem 13.7.3, X commutes with every length- L_0 word product. These products span $M_D(\mathbb{C})$ by block injectivity, so linearity gives $XM = MX$ for every $M \in M_D(\mathbb{C})$. $\square$

Lemma 13.7.5 (Amplifying fixed-length word commutation). [Lean](#) [✓ Stmt](#) If a boundary matrix X commutes with every word product of length m , then it commutes with every word product whose length is a multiple qm .

Proof. [✓ Proof](#) Proceed by induction on q . For $q = 0$, the claim is trivial. For the inductive step, let a word of length $(q+1)m$ be given and write it as a length- m prefix followed by a length- qm suffix. The prefix commutes with X by hypothesis, and the suffix commutes by the induction hypothesis. $\square$

Lemma 13.7.6 (Padded complement annihilation). [Lean](#) [Lean](#) [✓ Stmt](#) If A is L_0 -block-injective, $q \geq 1$, and $k \leq qL_0$, then the two one-sided annihilation statements are:

$$Z A^w = 0 \quad (|w| = k) \quad \implies \quad Z = 0.$$

The left-handed form is

$$A^w Z = 0 \quad (|w| = k) \quad \implies \quad Z = 0.$$

Proof. ✓ Proof The hypothesis $\text{span}\{A^u : |u| = L_0\} = M_D(\mathbb{C})$ implies the same spanning statement at every positive multiple qL_0 . Each length- qL_0 word factors as a length- k prefix followed by padding, so Z annihilates all generators of the full span and hence annihilates the identity. For the left-handed form, factor each length- qL_0 word as a prefix followed by a length- k suffix and use the same spanning argument. $\square$

Lemma 13.7.7 (One-sided boundary matrices are unique). Lean Lean ✓ Stmt *Let A be L_0 -block-injective with $L_0 > 0$. If two boundary matrices have the same product with every one-site tensor on one fixed side, then the two boundary matrices are equal.*

Proof. ✓ Proof If $Y_1 A^j = Y_2 A^j$ for every physical index j , then for every nonempty word w , induction on the word gives

$$Y_1 A^w = Y_2 A^w.$$

Since A is L_0 -block-injective,

$$\text{span}\{A^w : |w| = L_0\} = M_D(\mathbb{C}).$$

Thus the left multiplication maps $M \mapsto Y_1 M$ and $M \mapsto Y_2 M$ are equal on $M_D(\mathbb{C})$, and evaluating on the identity gives $Y_1 = Y_2$. The other one-sided statement is the same argument, with right multiplication in place of left multiplication. $\square$

Lemma 13.7.8 (Generator commutation from length- m commutation). Lean ✓ Stmt *Under the hypotheses of Theorem 13.7.4, one has $XA^i = A^i X$ for every physical index i .*

Proof. ✓ Proof Apply Theorem 13.7.4 with $M = A^i$. $\square$

Theorem 13.7.9 (MPS-line containment from length- m commutation). Lean ✓ Stmt *Let A be injective after blocking $L_0 > 0$ sites, and let $m \geq L_0$. If a boundary matrix X commutes with every length- m word product, then*

$$\Gamma_N(X) \in \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ Proof Theorem 13.7.4 makes X commute with the whole matrix algebra. The center of $M_D(\mathbb{C})$ consists of scalar matrices, so $X = c\mathbf{1}$, and therefore $\Gamma_N(X) = cV^{(N)}(A)$. $\square$

Theorem 13.7.10 (MPS-line containment from positive-length commutation). Lean ✓ Stmt *Let A be injective after blocking $L_0 > 0$ sites. If X commutes with every word product of some positive length m , then*

$$\Gamma_N(X) \in \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ Proof Lemma 13.7.5 amplifies the length- m commutation hypothesis to length $L_0 m \geq L_0$. Then Theorem 13.7.9 applies. $\square$

Theorem 13.7.11 (MPS-line containment from two one-sided equations). Lean ✓ Stmt *Let A be injective after blocking $L_0 > 0$ sites. Suppose there are matrices Y_μ such that*

$$A^\mu A^b X = Y_\mu A^b, \quad X A^a A^\mu = A^a Y_\mu,$$

for every pair of physical letters a, b . Then

$$\Gamma_N(X) \in \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ Proof The identities in (13.4) give commutation with all words of length $|\mu| + 2$, a positive length. The positive-length containment theorem then applies. $\square$

Theorem 13.7.12 (Boundary equality implies containment in the MPS line). Lean ✓ Stmt Let A be injective after blocking $L_0 > 0$ sites, and fix a physical letter η used outside the complementary sites. Suppose that, for every boundary condition τ and every physical letter j ,

$$C_\tau^+ A^j X = Y_\tau^+ A^j, \quad X A^j C_\tau^- = A^j Y_\tau^-,$$

where C_τ^+ and C_τ^- are the complementary words exposed by the two boundary-crossing cyclic windows. Suppose also that the two boundary matrices satisfy

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-,$$

for every word μ on the complementary sites, then

$$\Gamma_N(X) \in \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ Proof Lemma 13.5.14 gives a family Y_μ satisfying

$$A^\mu A^b X = Y_\mu A^b, \quad X A^a A^\mu = A^a Y_\mu.$$

Theorem 13.7.11 applies. □

Theorem 13.7.13 (Closure-property step for periodic chains). Lean ✓ Stmt Let A be injective after blocking $L_0 > 0$ sites and let $D \geq 1$. Assume $N \geq 2$ and $L_0 < L \leq N$. Suppose that, for every physical letter η used in the two boundary conditions and every boundary matrix obtained from an N -site chain ground state, the identities

$$A^\mu A^j X = Y_{\tau_\eta^+(\mu)}^+ A^j, \quad X A^j A^\mu = A^j Y_{\tau_\eta^-(\mu)}^-,$$

hold for every physical letter j and every complementary word μ , and are supplemented, for every μ , by

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-.$$

Then

$$\mathcal{G}_{N,L}(A) \subseteq \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ Proof The cyclic-to-open-chain reduction writes any chain ground state as $\Gamma_N(X)$. The two boundary-crossing cyclic windows give

$$A^\mu A^j X = Y_{\tau_\eta^+(\mu)}^+ A^j, \quad X A^j A^\mu = A^j Y_{\tau_\eta^-(\mu)}^-.$$

Together with

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-,$$

Theorem 13.7.12 finishes. □

13.8 Periodic-boundary comparison for normal tensors

This section isolates the closure-property part of the periodic-boundary argument for normal tensors. In the source proof, after the intersection property has grown the open interval, the same inverting-and-growing-back argument is applied when closing the boundaries [CPGSV21,

Section IV.C]. For a vector $\psi = \Gamma_{M+1}(X)$, the closure property asks that the two boundary-crossing restrictions $\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi)$ and $\text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi)$ agree. The symbols $\tau_\eta^\pm(\mu)$ index these restrictions, and the matrices $Y_i(\tau)$ below satisfy

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Once the one-sided boundary products are known, equality of the restrictions follows from the boundary-matrix commutation relation $XA^j = A^jX$ for every physical letter j . The block-window identity

$$XA^\alpha A^\nu = A^\alpha Y_\nu$$

is derived from the cyclic-window constraints crossing the periodic cut.

Lemma 13.8.1 (Block-to-letter translation of the boundary block matrix equation). [Lean](#) [Stmt](#) *Let A be an MPS tensor and let $L_0, K \geq 0$. Let X be a matrix, and let Y_{c_b} be a matrix for each iterated block index c_b of the complement. Suppose that, for every block letter b of the alphabet $\{0, \dots, d-1\}^{L_0}$ and every iterated block index c_b of the complement, with $w(b)$ the length- L_0 word of b and $\widetilde{w(c_b)}$ the length- $L_0 K$ complement word obtained by concatenating the blocks of c_b ,*

$$XA^{w(b)} A^{\widetilde{w(c_b)}} = A^{w(b)} Y_{c_b}.$$

Then there is a family Y'_c , indexed by the length- $L_0 K$ words c , with

$$XA^s A^c = A^s Y'_c$$

for every length- L_0 word s and every length- $L_0 K$ word c .

Proof. [Proof](#) Each length- L_0 word s is the word $w(b)$ of its block letter b , and each length- $L_0 K$ word c regroups into K blocks, recovering it as the concatenated complement word $\widetilde{w(c_b)}$ of an iterated block index c_b ; both identifications are bijective. Substituting these representations into the hypothesis and setting $Y'_c = Y_{c_b}$ gives the asserted equation for every s and c . $\square$

Lemma 13.8.2 (Boundary matrix commutation from block-window equations). [Lean](#) [Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$. Let X be a boundary matrix, and let Y_v be a matrix for each complementary word v of length K . Suppose that, for every word u of length L_0 and every word v of length K ,*

$$XA^u A^v = A^u Y_v.$$

Then, for every physical letter j ,

$$XA^j = A^j X.$$

This is the boundary-matrix commutation step obtained from the equations $XA^u A^v = A^u Y_v$, for all length- L_0 words u and all length- K words v , as in [CPGSV21, Section IV.C, lines 2049–2090].

Proof. [Proof](#) By Lemma 8.1.7,

$$\text{span}\{A^u : |u| = L_0\} = M_D(\mathbb{C}).$$

Hence, for every $M \in M_D(\mathbb{C})$ and every complementary word v of length K , the displayed hypothesis extends by linearity in the length- L_0 block to

$$XMA^v = MY_v$$

Taking $M = 1$ gives

$$Y_v = XA^v.$$

Taking $M = A^j$ and substituting the preceding identity gives, for every word v of length K ,

$$(XA^j - A^jX)A^v = 0$$

Since $K \leq (K+1)L_0$ and $K+1 \geq 1$, Lemma 13.7.6 gives $XA^j - A^jX = 0$. $\square$

Lemma 13.8.3 (Trace identity from the boundary-crossing cyclic window). [Lean](#) [✓ Stmt](#) *Let A be a tensor with virtual dimension at least one, and let $L_0 > 0$ and $L_0 < M$. Let*

$$\psi = \Gamma_{M+1}(X)$$

and assume that every cyclic restriction of length $L_0 + 1$ belongs to $G_{L_0+1}(A)$. Then there are boundary matrices Y_ν , indexed by nonempty complementary words ν of length $M+1 - (L_0+1)$, such that, for every physical letter j and every word α of length L_0 ,

$$\text{tr}(A^j X A^\alpha A^\nu) = \text{tr}(A^j A^\alpha Y_\nu).$$

Proof. [✓ Proof](#) Let Y_ν be the matrix representing the cyclic restriction whose window begins at the last site. The cyclic word is

$$\alpha \nu j$$

after deleting the first letter from the window and reading the complement, so the ground-space representation gives

$$\text{tr}(A^\alpha A^\nu A^j X) = \text{tr}(A^j A^\alpha Y_\nu).$$

Cyclicity of the trace gives

$$\text{tr}(A^j X A^\alpha A^\nu) = \text{tr}(A^\alpha A^\nu A^j X).$$

Therefore

$$\text{tr}(A^j X A^\alpha A^\nu) = \text{tr}(A^j A^\alpha Y_\nu).$$

$\square$

Lemma 13.8.4 (One-site-injective boundary block-window identity). [Lean](#) [✓ Stmt](#) *Let A be a tensor with virtual dimension at least one whose single-site matrices span the full matrix algebra. Let $L_0 > 0$ and $L_0 < M$, and let*

$$\psi = \Gamma_{M+1}(X).$$

If every cyclic restriction of length $L_0 + 1$ belongs to $G_{L_0+1}(A)$, then there are boundary matrices Y_ν , indexed by nonempty complementary words ν of length $M+1 - (L_0+1)$, such that, for every word α of length L_0 ,

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

Proof. [✓ Proof](#) For every physical letter j , Lemma 13.8.3 gives

$$\text{tr}(A^j X A^\alpha A^\nu) = \text{tr}(A^j A^\alpha Y_\nu)$$

Put

$$Z = XA^\alpha A^\nu - A^\alpha Y_\nu.$$

For every physical letter j , the preceding identity gives

$$\text{tr}(A^j Z) = 0.$$

Since the matrices A^j span the full matrix algebra, the trace pairing gives

$$(\forall j, \operatorname{tr}(A^j Z) = 0) \implies Z = 0.$$

Hence $XA^\alpha A^\nu = A^\alpha Y_\nu$. □

Theorem 13.8.5 (Trace identities with length- L_0 probes from cyclic-window constraints). Lean

✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

and assume that every cyclic restriction of length $L_0 + 1$ belongs to $G_{L_0+1}(A)$. Then there are boundary matrices Y_ν , indexed by nonempty complementary words ν of length $M + 1 - (L_0 + 1)$, such that, for every word α of length L_0 and every word β of length L_0 ,

$$\operatorname{tr}(A^\beta X A^\alpha A^\nu) = \operatorname{tr}(A^\beta A^\alpha Y_\nu).$$

Proof. ✓ Proof Choose representation matrices for every cyclic restriction of length $L_0 + 1$. Fix α and ν . The boundary-crossing equations move the matrix X through the first boundary letter. In local boundary notation this has the form

$$XA^{\rho_0} A^{\rho_1 \cdots \rho_{M-L_0}} = A^{\rho_0} Y_{M+1-L_0}(\rho).$$

The adjacent boundary-window product then transports this comparison through the remaining $L_0 - 1$ boundary letters:

$$Y_{M+1-L_0}(\rho) A^{\rho_{M+1-L_0} \cdots \rho_{M-1}} = A^{\rho_1 \cdots \rho_{L_0-1}} Y_M(\rho).$$

Finally, the outside-label uniqueness lemma identifies the final boundary matrix with the matrix Y_ν attached to the complementary word ν . Thus

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

Multiplying on the left by A^β and taking the trace gives the asserted identity for every length- L_0 word β . □

Theorem 13.8.6 (Length-word trace separation for the boundary block window). Lean ✓ Stmt

Let A be L_0 -block-injective. Let X be a boundary matrix, and let Y_ν be a matrix for each complementary word ν of length K . Suppose that, for every word α of length L_0 , every word β of length L_0 , and every complementary word ν ,

$$\operatorname{tr}(A^\beta X A^\alpha A^\nu) = \operatorname{tr}(A^\beta A^\alpha Y_\nu).$$

Then, for every word α of length L_0 and every complementary word ν ,

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

Proof. ✓ Proof The trace identities say exactly that

$$\Gamma_{L_0}(XA^\alpha A^\nu) = \Gamma_{L_0}(A^\alpha Y_\nu)$$

for each fixed pair (α, ν) . Since A is L_0 -block-injective, Γ_{L_0} is injective by Lemma 13.4.3.3. Hence

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

□

Lemma 13.8.7 (Matrix identity $XA^\alpha A^\nu = A^\alpha Y_\nu$ from cyclic-window constraints). [Lean](#) [✓ Stmt](#)
Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

and assume that every cyclic restriction of length $L_0 + 1$ belongs to $G_{L_0+1}(A)$. Then there are boundary matrices Y_ν , indexed by nonempty complementary words ν of length $M + 1 - (L_0 + 1)$, such that for every word α of length L_0 ,

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

Proof. [✓ Proof](#) Theorem 13.8.5 gives, for every word β of length L_0 ,

$$\text{tr}(A^\beta XA^\alpha A^\nu) = \text{tr}(A^\beta A^\alpha Y_\nu).$$

Theorem 13.8.6 then gives the displayed matrix equation. □

Lemma 13.8.8 (Boundary restrictions from commutation and one-sided products). [Lean](#) [✓ Stmt](#)
Let A be L_0 -block-injective with $L_0 > 0$. Let X be a boundary matrix, let W_η and V_η be matrix families indexed by a physical letter η , and let μ be a word. Suppose that, for every physical letter j ,

$$XA^j = A^j X$$

Suppose also that, for every η and every j ,

$$W_\eta A^j = A^\mu A^j X, \quad XA^j A^\mu = A^j V_\eta.$$

Then, for every η and every j ,

$$W_\eta A^j = V_\eta A^j.$$

Proof. [✓ Proof](#) The commutation relation $XA^j = A^j X$ extends by induction: for every word w ,

$$XA^w = A^w X$$

Hence the second one-sided equation gives, for every k ,

$$A^k V_\eta = XA^k A^\mu = A^k XA^\mu = A^k A^\mu X.$$

Lemma 13.7.7 gives $V_\eta = A^\mu X$. The first one-sided equation then gives

$$W_\eta A^j = A^\mu A^j X = A^\mu XA^j = V_\eta A^j.$$

□

Lemma 13.8.9 (Products with one-site tensors determine the boundary matrix). [Lean](#) [✓ Stmt](#)
Let A be L_0 -block-injective with $L_0 > 0$. Fix two matrix families Y^+ and Y^- and the two reindexed boundary conditions $\tau_\eta^+(\mu)$ and $\tau_\eta^-(\mu)$. If, for every physical letter j ,

$$Y_{\tau_\eta^+(\mu)}^+ A^j = Y_{\tau_\eta^-(\mu)}^- A^j,$$

then

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-.$$

Proof. [✓ Proof](#) Apply Lemma 13.7.7 to the two matrices $Y_{\tau_\eta^+(\mu)}^+$ and $Y_{\tau_\eta^-(\mu)}^-$. □

Lemma 13.8.10 (Outside labels determine boundary-restriction matrices). [Lean](#) [Lean](#) [Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$ and $L_0 \leq M$. If an outside configuration ρ has the same word μ as $\tau_\eta^+(\mu)$ on the sites outside the last-site cyclic window, then the two matrices representing that restriction agree:

$$Y_\rho = Y_{\tau_\eta^+(\mu)}.$$

The same assertion holds for the second boundary-crossing window: if ρ has the same outside word as $\tau_\eta^-(\mu)$, then

$$Y_\rho = Y_{\tau_\eta^-(\mu)}.$$

This is the boundary-matrix independence step used to choose one representative outside configuration for each outside word in the closure-property matrix comparison.

Proof. [Proof](#) Equality of the cyclic restrictions gives, for every physical letter j ,

$$Y_\rho A^j = Y_{\tau_\eta^\pm(\mu)} A^j.$$

Lemma 13.7.7 identifies the two matrices from these one-site products. □

Lemma 13.8.11 (First-letter restrictions determine a vector). [Lean](#) [Stmt](#) Let R_j be restriction to first letter j :

$$(R_j \phi)(\sigma_1, \dots, \sigma_L) = \phi(j, \sigma_1, \dots, \sigma_L).$$

For $\phi, \psi \in (\mathbb{C}^d)^{\otimes(L+1)}$,

$$(\forall j, R_j \phi = R_j \psi) \implies \phi = \psi.$$

Proof. [Proof](#) At $(\sigma_0, \sigma_1, \dots, \sigma_L)$ the required equality is

$$\phi(\sigma_0, \sigma_1, \dots, \sigma_L) = (R_{\sigma_0} \phi)(\sigma_1, \dots, \sigma_L) = (R_{\sigma_0} \psi)(\sigma_1, \dots, \sigma_L) = \psi(\sigma_0, \sigma_1, \dots, \sigma_L).$$

□

Lemma 13.8.12 (Auxiliary restriction equality for the closure property). [Lean](#) [Stmt](#) Let $L_0 > 0$ and $L_0 \leq M$, and set

$$B_{\eta,\mu}^+(\psi) = \text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi), \quad B_{\eta,\mu}^-(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

With R_j as in Lemma 13.8.11,

$$(\forall j, R_j B_{\eta,\mu}^+(\psi) = R_j B_{\eta,\mu}^-(\psi)) \implies B_{\eta,\mu}^+(\psi) = B_{\eta,\mu}^-(\psi).$$

Proof. [Proof](#) Apply Lemma 13.8.11 to the two length- $(L_0 + 1)$ restrictions, using the displayed family of first-letter equalities. □

Lemma 13.8.13 (Boundary-crossing restriction from right products). [Lean](#) [Stmt](#) Let A be an MPS tensor, let $L_0 > 0$, and let $L_0 \leq M$. Suppose the two boundary-crossing restrictions of length $L_0 + 1$ are represented by boundary matrices $Y_{\eta,\mu}^+$ and $Y_{\eta,\mu}^-$:

$$B_{\eta,\mu}^+(\psi) = \Gamma_{L_0+1}(Y_{\eta,\mu}^+), \quad B_{\eta,\mu}^-(\psi) = \Gamma_{L_0+1}(Y_{\eta,\mu}^-).$$

If, for every physical letter j ,

$$Y_{\eta,\mu}^+ A^j = Y_{\eta,\mu}^- A^j,$$

then

$$B_{\eta,\mu}^+(\psi) = B_{\eta,\mu}^-(\psi).$$

Proof. ✓ Proof Fix j and take the first-letter restriction of the two displayed length- $(L_0 + 1)$ restrictions. These are

$$\Gamma_{L_0}(Y_{\eta, \mu}^+ A^j), \quad \Gamma_{L_0}(Y_{\eta, \mu}^- A^j).$$

The one-site product equality gives, for every j ,

$$\Gamma_{L_0}(Y_{\eta, \mu}^+ A^j) = \Gamma_{L_0}(Y_{\eta, \mu}^- A^j).$$

Lemma 13.8.11 identifies the two original restrictions. □

Lemma 13.8.14 (Boundary restrictions from first-letter products). Lean ✓ Stmt Let $L_0 > 0$ and $L_0 \leq M$. Let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Fix a complementary word μ . If, for every boundary letter η and physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j,$$

then for every η ,

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Proof. ✓ Proof Apply Lemma 13.8.13 for each boundary letter η , using the corresponding first-letter product equations. □

Lemma 13.8.15 (Equality of the two restrictions crossing the periodic cut). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$, let $L_0 < M$, and let

$$\psi \in \mathcal{G}_{M+1, L_0+1}(A), \quad \psi = \Gamma_{M+1}(X).$$

For every boundary letter η and complementary word μ , the two restrictions crossing the periodic cut satisfy

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Proof. ✓ Proof The cyclic-window representation lemma gives matrices $Y_M(\tau_\eta^+(\mu))$ and $Y_{M+1-L_0}(\tau_\eta^-(\mu))$ such that

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \Gamma_{L_0+1}(Y_M(\tau_\eta^+(\mu))), \quad \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi) = \Gamma_{L_0+1}(Y_{M+1-L_0}(\tau_\eta^-(\mu))).$$

Theorem 13.8.1.8 gives, for every j ,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Lemma 13.8.13 identifies the two length- $(L_0 + 1)$ restrictions. □

Lemma 13.8.16 (Boundary-matrix product from boundary-crossing restrictions). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M + 1$, and let $Y_i(\tau)$ be matrices satisfying

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Suppose that, for every outside letter η ,

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Then there are boundary conditions $\rho_{j, \sigma}^+$ and $\rho_{j, \sigma}^-$ such that, for every complementary position k ,

$$\rho_{j, \sigma}^+(k + L_0) = \mu_k, \quad \rho_{j, \sigma}^-(k + 1) = \mu_k,$$

and, for every physical letter j and every word σ of length L_0 ,

$$Y_M(\rho_{j, \sigma}^+)A^j A^\sigma = Y_{M+1-L_0}(\rho_{j, \sigma}^-)A^j A^\sigma.$$

Proof. ✓ Proof For the product indexed by j and σ , specialize the preceding equality at the outside letter $\eta = j$. Take $\rho_{j,\sigma}^+ = \tau_j^+(\mu)$ and $\rho_{j,\sigma}^- = \tau_j^-(\mu)$. The complement equations are the two identities of Lemma 13.5.6. Fix j . Applying the first-letter restriction to the displayed equality gives

$$\Gamma_{L_0}(Y_M(\tau_j^+(\mu))A^j) = \Gamma_{L_0}(Y_{M+1-L_0}(\tau_j^-(\mu))A^j),$$

where the two sides are identified by Lemma 13.4.5.3. By Lemma 13.4.3.3,

$$Y_M(\tau_j^+(\mu))A^j = Y_{M+1-L_0}(\tau_j^-(\mu))A^j.$$

Right multiplication by A^σ gives the asserted product equation. □

Lemma 13.8.17 (Boundary-matrix product from right products). Lean ✓ Stmt *Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$, and let $Y_i(\tau)$ be matrices satisfying*

$$\text{Res}_{i,L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Fix a complementary word μ of length $M+1-(L_0+1)$. Suppose that, for every boundary letter η and every physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Then there are boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ such that, for every complementary position k ,

$$\rho_{j,\sigma}^+(k+L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k+1) = \mu_k,$$

and, for every physical letter j and every word σ of length L_0 ,

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma.$$

Proof. ✓ Proof For each boundary letter η , Lemma 13.8.13 gives

$$\text{Res}_{M,L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0,L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Applying these equalities, Lemma 13.8.16 gives the displayed boundary conditions and product equation. □

Lemma 13.8.18 (Boundary-crossing comparison after multiplication by words). Lean ✓ Stmt *Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Fix a complementary word μ , a matrix X , and a family of matrices $Y_i(\tau)$. Suppose that, for every boundary letter η , every physical letter j , and every word σ of length L_0 ,*

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma = A^\mu A^j X A^\sigma.$$

Then, for every η and j ,

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j = A^\mu A^j X.$$

Proof. ✓ Proof Fix η and j , and set

$$Z_{\eta,j} = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j - A^\mu A^j X.$$

The hypothesis gives, for every σ with $|\sigma| = L_0$,

$$Z_{\eta,j}A^\sigma = 0.$$

Since A is L_0 -block-injective, the products A^σ with $|\sigma| = L_0$ span the full matrix algebra. Applying Lemma 13.7.6 gives $Z_{\eta,j} = 0$. □

Lemma 13.8.19 (Boundary-crossing comparison after left multiplication by words). [Lean](#)
[✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Fix a complementary word μ , a matrix X , and a family of matrices $Y_i(\tau)$. Suppose that, for every boundary letter η , every physical letter j , and every pair of words σ, α of length L_0 ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma) = A^\alpha (A^\mu A^j X A^\sigma).$$

Then, for every η, j , and σ ,

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma = A^\mu A^j X A^\sigma.$$

Proof. [✓ Proof](#) Fix η, j , and σ , and set

$$Z_{\eta,j,\sigma} = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma - A^\mu A^j X A^\sigma.$$

The hypothesis says that $A^\alpha Z_{\eta,j,\sigma} = 0$ for every word α of length L_0 . The left-handed form of Lemma 13.7.6 gives $Z_{\eta,j,\sigma} = 0$. $\square$

Lemma 13.8.20 (Boundary-matrix product from the boundary-crossing comparison). [Lean](#)
[✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M+1$, and let $Y_i(\tau)$ be matrices satisfying

$$\text{Res}_{i,L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Fix a complementary word μ and a matrix X . Suppose that, for every boundary letter η and every physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = A^\mu A^j X.$$

Suppose also that, for every η, j , and every word σ of length L_0 ,

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma = A^\mu A^j X A^\sigma.$$

Then there are boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ such that, for every complementary position k ,

$$\rho_{j,\sigma}^+(k + L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k + 1) = \mu_k,$$

and, for every physical letter j and every word σ of length L_0 ,

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma.$$

Proof. [✓ Proof](#) Lemma 13.8.18 gives, for every η and j ,

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j = A^\mu A^j X.$$

Combining this equation with

$$Y_M(\tau_\eta^+(\mu))A^j = A^\mu A^j X$$

gives

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Lemma 13.8.17 then gives the displayed boundary conditions and product equation. $\square$

Lemma 13.8.21 (Boundary-matrix product from the left-multiplied boundary-crossing comparison). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, and let $L_0 \leq M$. Let ψ be a state on the chain of length $M + 1$, and let $Y_i(\tau)$ be matrices satisfying

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

Fix a complementary word μ and a matrix X . Suppose that, for every boundary letter η and every physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = A^\mu A^j X.$$

Suppose also that, for every η, j , and every pair of words σ, α of length L_0 ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j A^\sigma) = A^\alpha (A^\mu A^j X A^\sigma).$$

Then there are boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ satisfying the complementary-word equations

$$\rho_{j,\sigma}^+(k + L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k + 1) = \mu_k,$$

and the product equation

$$Y_M(\rho_{j,\sigma}^+)A^j A^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^j A^\sigma.$$

Proof. [✓ Proof](#) Lemma 13.8.19 gives

$$Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j A^\sigma = A^\mu A^j X A^\sigma.$$

Lemma 13.8.20 then gives the displayed boundary conditions and product equation. $\square$

Theorem 13.8.22 (Boundary-matrix product from left-multiplied equations). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

be an open-chain representation, and let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Fix a complementary word μ . Suppose that, for every boundary letter η , physical letter j , and length- L_0 words σ, α ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j A^\sigma) = A^\alpha (A^\mu A^j X A^\sigma).$$

Then there are boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ satisfying the complementary-word equations

$$\rho_{j,\sigma}^+(k + L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k + 1) = \mu_k,$$

and the product equation

$$Y_M(\rho_{j,\sigma}^+)A^j A^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^j A^\sigma.$$

Proof. [✓ Proof](#) Lemma 13.5.5 gives the equation for the boundary-crossing window beginning at M ,

$$Y_M(\tau_\eta^+(\mu))A^j = A^\mu A^j X.$$

The assumed left-multiplied coordinate equation and Lemma 13.8.21 then give the displayed boundary conditions and product equation. $\square$

Theorem 13.8.23 (Boundary restrictions from the left-multiplied comparison). [Lean](#) [✓ Stmt](#)
Let A be L_0 -block-injective with $L_0 > 0$, $L_0 \leq M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

be an open-chain representation, and let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Fix a complementary word μ . Suppose that, for every boundary letter η , physical letter j , and length- L_0 words α, σ ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma) = A^\alpha (A^\mu A^j X A^\sigma).$$

Then, for every boundary letter η ,

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Proof. [✓ Proof](#) The assumed equations and Lemma 13.8.19 give, for every η, j , and σ ,

$$Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma = A^\mu A^j X A^\sigma.$$

Lemma 13.8.18 removes the right word:

$$Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j = A^\mu A^j X.$$

The equation for the boundary-crossing window beginning at M , from Lemma 13.5.5, gives

$$Y_M(\tau_\eta^+(\mu)) A^j = A^\mu A^j X.$$

Hence the first-letter products agree, and Lemma 13.8.14 identifies the two restrictions. $\square$

Theorem 13.8.24 (Equality of the boundary-crossing restrictions). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

be an open-chain representation. Assume that for all i and τ ,

$$\text{Res}_{i, L_0+1}^\tau(\psi) \in G_{L_0+1}(A).$$

In local coordinate notation, fix a nonempty word μ on the sites outside the two cyclic windows. For every outside letter η , choose the two outside configurations so that, for each index k of this outside word,

$$\tau_\eta^+(\mu)_{k+L_0} = \mu_k, \quad \tau_\eta^-(\mu)_{k+1} = \mu_k,$$

and put the letter η on the remaining sites. The notation $\tau_\eta^\pm(\mu)$ is a coordinate parametrization for the comparison used at the periodic boundary in [CPGSV21, Section IV.C, lines 2078–2079]; it is not notation from the source. The corresponding boundary-crossing restriction equality is

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Proof. [✓ Proof](#) Choose matrices $Y_i(\tau)$ representing the length- $(L_0 + 1)$ restrictions. The one-sided boundary-product lemma gives, for every outside letter η and physical letter j ,

$$Y_M(\tau_\eta^+(\mu)) A^j = A^\mu A^j X, \quad X A^j A^\mu = A^j Y_{M+1-L_0}(\tau_\eta^-(\mu)).$$

Lemma 13.8.7 gives matrices Y_ν satisfying, for every word α of length L_0 and every complementary word ν ,

$$XA^\alpha A^\nu = A^\alpha Y_\nu.$$

Lemma 13.8.2 then gives, for every physical letter k , the commutation identity

$$XA^k = A^k X.$$

Lemma 13.8.8 gives

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Lemma 13.8.14 then identifies the two boundary-crossing restrictions, giving the periodic-boundary comparison of [CPGSV21, lines 2078–2079]. $\square$

13.8.1 Reverse comparison for boundary-crossing restrictions

This subsection runs the comparison in the reverse direction: from equality of the two cyclic restrictions it recovers the first-letter product equation $Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j$ and the left-multiplied boundary comparison.

Lemma 13.8.1.1 (First-letter products from equal boundary restrictions). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$ and $L_0 \leq M$. Let ψ be a vector on $M + 1$ sites, and suppose the two length- $(L_0 + 1)$ boundary restrictions are represented by*

$$\Gamma_{L_0+1}(Y_M(\tau_\eta^+(\mu))), \quad \Gamma_{L_0+1}(Y_{M+1-L_0}(\tau_\eta^-(\mu))).$$

If these two restrictions are equal, then for every physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Thus the first-letter restriction of the preceding equality is precisely the displayed product equation.

Proof. [✓ Proof](#) Restricting both sides to the first physical letter j gives

$$\Gamma_{L_0}(Y_M(\tau_\eta^+(\mu))A^j) = \Gamma_{L_0}(Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j).$$

Since A is L_0 -block-injective, Γ_{L_0} is injective, and the displayed product equality follows. $\square$

Lemma 13.8.1.2 (Left-multiplied comparison from equal boundary restrictions). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$ and $L_0 \leq M$. Let ψ be a vector on $M + 1$ sites, and let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Assume that for every boundary letter η ,*

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Then for every boundary letter η , physical letter j , and length- L_0 words α, σ ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j A^\sigma) = A^\alpha (Y_M(\tau_\eta^+(\mu))A^j A^\sigma).$$

Proof. [✓ Proof](#) The preceding lemma gives

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Multiplying on the left by A^α and on the right by A^σ gives the desired comparison. $\square$

Theorem 13.8.1.3 (Left-multiplied comparison of boundary matrices). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

be an open-chain representation, and let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Fix a nonempty complementary word μ . Then for every boundary letter η , physical letter j , and length- L_0 words α, σ ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma) = A^\alpha (Y_M(\tau_\eta^+(\mu)) A^j A^\sigma).$$

Proof. [✓ Proof](#) The matrices $Y_i(\tau)$ give the local ground-space representations needed in the preceding lemma. Once the preceding boundary-restriction equality is proved, it gives

$$\text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

Applying the left-multiplied comparison for equal boundary restrictions gives

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma) = A^\alpha (Y_M(\tau_\eta^+(\mu)) A^j A^\sigma).$$

□

Theorem 13.8.1.4 (Left-multiplied coordinate comparison at the periodic boundary). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$, $L_0 < M$, and $D \geq 1$. Let

$$\psi = \Gamma_{M+1}(X)$$

be an open-chain representation, and let $Y_i(\tau)$ represent the length- $(L_0 + 1)$ cyclic restrictions of ψ . Fix a nonempty complementary word μ . Then for every boundary letter η , physical letter j , and length- L_0 words α, σ ,

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma) = A^\alpha (A^\mu A^j X A^\sigma).$$

Proof. [✓ Proof](#) Once the comparison of the two cyclic restrictions is available, it gives

$$A^\alpha (Y_{M+1-L_0}(\tau_\eta^-(\mu)) A^j A^\sigma) = A^\alpha (Y_M(\tau_\eta^+(\mu)) A^j A^\sigma).$$

The one-sided equation for the boundary-crossing window beginning at M gives

$$Y_M(\tau_\eta^+(\mu)) A^j = A^\mu A^j X.$$

Substitution gives the displayed identity.

□

Lemma 13.8.1.5 (Letterwise comparison of boundary-crossing restrictions). [Lean](#) [✓ Stmt](#) Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$, let $L_0 < M$, and let

$$\psi \in \mathcal{G}_{M+1, L_0+1}(A), \quad \psi = \Gamma_{M+1}(X).$$

Let R_j denote restriction to first physical letter j . For every outside letter η , complementary word μ , and physical letter j , one has

$$R_j \text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) = R_j \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi).$$

This is the one-letter product form of the local comparison corresponding to [CPGSV21, lines 2078–2079].

Proof. ✓ Proof Let $Y_i(\tau)$ be the representation matrices supplied by Lemma 13.6.7. By Theorem 13.8.1.8,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Fixing the first physical letter in the two length- $(L_0 + 1)$ windows gives

$$\begin{aligned} R_j \text{Res}_{M, L_0+1}^{\tau_\eta^+(\mu)}(\psi) &= \Gamma_{L_0}(Y_M(\tau_\eta^+(\mu))A^j), \\ R_j \text{Res}_{M+1-L_0, L_0+1}^{\tau_\eta^-(\mu)}(\psi) &= \Gamma_{L_0}(Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j). \end{aligned}$$

The two restrictions are therefore equal. □

Theorem 13.8.1.6 (Boundary-matrix product for the two boundary-crossing restrictions).

Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$. Let ψ be a state on the chain of length $M + 1$ with open-chain representation $\psi = \Gamma_{M+1}(X)$. Let $Y_i(\tau)$ be matrices satisfying

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

If $L_0 < M$, then for every nonempty complementary word μ there are boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$, depending on the physical letter j and the word σ of length L_0 , such that, for every complementary position k ,

$$\rho_{j,\sigma}^+(k + L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k + 1) = \mu_k,$$

and, for every j and σ ,

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma.$$

Proof. ✓ Proof Theorem 13.8.1.4 gives, for every boundary letter η , physical letter j , and words α, σ of length L_0 ,

$$A^\alpha(Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma) = A^\alpha(A^\mu A^j X A^\sigma).$$

Theorem 13.8.22 applies this left-multiplied comparison together with the one-sided boundary products and gives the displayed boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$, including

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma.$$

□

Theorem 13.8.1.7 (Product equality at the periodic boundary). Lean ✓ Stmt Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$. Let $\psi \in \mathcal{G}_{M+1, L_0+1}(A)$ have an open-chain representation $\psi = \Gamma_{M+1}(X)$. Let $Y_i(\tau)$ be matrices satisfying

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

If $L_0 < M$, then for every boundary letter η , complementary word μ , physical letter j , and word σ of length L_0 ,

$$Y_M(\tau_\eta^+(\mu))A^jA^\sigma = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma.$$

Proof. ✓ Proof Theorem 13.8.1.6 gives boundary conditions $\rho_{j,\sigma}^+$ and $\rho_{j,\sigma}^-$ satisfying

$$\rho_{j,\sigma}^+(k + L_0) = \mu_k, \quad \rho_{j,\sigma}^-(k + 1) = \mu_k,$$

and

$$Y_M(\rho_{j,\sigma}^+)A^jA^\sigma = Y_{M+1-L_0}(\rho_{j,\sigma}^-)A^jA^\sigma.$$

Theorem 13.5.9 replaces these auxiliary conditions by the displayed boundary conditions:

$$Y_M(\tau_\eta^+(\mu))A^jA^\sigma = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^jA^\sigma,$$

which is the product equality needed for the periodic-boundary comparison. $\square$

Theorem 13.8.1.8 (Right-product equality at the periodic boundary). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$. Let $\psi \in \mathcal{G}_{M+1, L_0+1}(A)$ have an open-chain representation $\psi = \Gamma_{M+1}(X)$. Let $Y_i(\tau)$ be matrices satisfying*

$$\text{Res}_{i, L_0+1}^\tau(\psi) = \Gamma_{L_0+1}(Y_i(\tau)).$$

If $L_0 < M$, then for every letter η used in the two boundary conditions, every word μ on the complementary sites, and every physical letter j ,

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

This is the one-site product equality used to compare the two boundary matrices in the periodic-boundary comparison.

Proof. [✓ Proof](#) Fix j and set

$$Z_j = Y_M(\tau_\eta^+(\mu))A^j - Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

Theorem 13.8.1.7 gives, after subtracting the two sides, for every word σ of length L_0 ,

$$Z_j A^\sigma = 0.$$

Since A is L_0 -block-injective,

$$\text{span}\{A^\sigma : |\sigma| = L_0\} = M_D(\mathbb{C}).$$

Applying Lemma 13.7.6 gives $Z_j = 0$, hence

$$Y_M(\tau_\eta^+(\mu))A^j = Y_{M+1-L_0}(\tau_\eta^-(\mu))A^j.$$

$\square$

Theorem 13.8.1.9 (Equality of the two boundary-matrix families). [Lean](#) [✓ Stmt](#) *Let A be L_0 -block-injective with $L_0 > 0$ and $D \geq 1$, let $N \geq 2$, $L_0 + 1 < N$, and $L_0 < L \leq N$, and let $\psi \in \mathcal{G}_{N, L}(A)$ have an open-chain representation $\psi = \Gamma_N(X)$. Suppose Y^+ and Y^- are two families of matrices obtained from the two boundary-crossing cyclic-window constraints, and that they satisfy, for every physical letter j and every boundary condition τ ,*

$$A^{\tau_{L_0} \cdots \tau_{N-2}} A^j X = Y_\tau^+ A^j, \quad X A^j A^{\tau_1 \cdots \tau_{N-L_0-1}} = A^j Y_\tau^-.$$

For every physical letter $\eta \in \{0, \dots, d-1\}$ and every word μ on the complementary sites, define two boundary conditions by

$$\tau_\eta^+(\mu)_i = \begin{cases} \mu_{i-L_0}, & L_0 \leq i < N-1, \\ \eta, & \text{otherwise,} \end{cases} \quad \tau_\eta^-(\mu)_i = \begin{cases} \mu_{i-1}, & 1 \leq i < N-L_0, \\ \eta, & \text{otherwise.} \end{cases}$$

Then the two families agree on these two boundary conditions:

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-.$$

This is the boundary-matrix comparison required for equality of the two restrictions crossing the periodic cut.

Proof. ✓ **Proof** The reduced cyclic-window compatibilities at the boundary give, for every physical letter j ,

$$A^\mu A^j X = Y_{\tau_\eta^+(\mu)}^+ A^j, \quad X A^j A^\mu = A^j Y_{\tau_\eta^-(\mu)}^-.$$

By Lemma 13.8.9, it is enough to prove, for every j ,

$$Y_{\tau_\eta^+(\mu)}^+ A^j = Y_{\tau_\eta^-(\mu)}^- A^j.$$

Write $N = M + 1$. The two boundary-crossing supports start at M and $M + 1 - L_0$. Let $Y_i(\tau)$ denote the cyclic-window representation matrix given by Lemma 13.6.7 at the window beginning at i . The first step identifies the two abstract boundary-condition families with the two boundary-restriction matrices:

$$Y_{\tau_\eta^+(\mu)}^+ = Y_M(\tau_\eta^+(\mu)), \quad Y_{\tau_\eta^-(\mu)}^- = Y_{M+1-L_0}(\tau_\eta^-(\mu)).$$

Together with the preceding identifications, Theorem 13.8.1.8 gives, for every physical letter j ,

$$Y_{\tau_\eta^+(\mu)}^+ A^j = Y_{\tau_\eta^-(\mu)}^- A^j.$$

Lemma 13.8.9 then gives the desired equality of boundary matrices. □

13.8.2 Normal unique-ground-state consequences

In the injective case, for $N \geq 2$ and $1 < L \leq N$, the chain ground space equals $\text{span}\{V^{(N)}(A)\}$, so the parent Hamiltonian has a unique ground state in that length range. In the normal case the same conclusion is the payoff of the section, given the boundary-crossing comparison and the length hypotheses $N \geq 2$, $L_0 + 1 < N$, and $L_0 < L \leq N$.

Theorem 13.8.2.1 (Chain ground space is spanned by the MPS vector in the injective case).

Lean ✓ **Stmnt** If A is injective, $D \geq 1$, $N \geq 2$, and $1 < L \leq N$, then

$$\mathcal{G}_{N,L}(A) = \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ **Proof** Let $\psi \in \mathcal{G}_{N,L}(A)$. The non-wrapping window conditions and Lemma 13.4.5.4 imply $\psi \in G_N(A)$, so $\psi = \Gamma_N(X)$ for some boundary matrix X . The periodic boundary condition now applies Theorem 13.5.16 and forces X to commute with every letter A^j . Since A is injective, the letters generate the full matrix algebra, hence X is scalar. Therefore ψ is proportional to $V^{(N)}(A)$, while the reverse inclusion is Lemma 13.6.8. □

Theorem 13.8.2.2 (Normal tensor: $\mathcal{G}_{N,L}(A) \subseteq \text{span}\{V^{(N)}(A)\}$). Lean ✓ **Stmnt** If A is normal, $D \geq 1$, and injective after blocking $L_0 > 0$ sites, with $N \geq 2$, $L_0 + 1 < N$, and $L_0 < L \leq N$, then

$$\mathcal{G}_{N,L}(A) \subseteq \text{span}\{V^{(N)}(A)\}.$$

Proof. ✓ **Proof** For a chain ground state, the open-chain containment gives $\psi = \Gamma_N(X)$. The two boundary-crossing cyclic-window constraints give

$$A^\mu A^j X = Y_{\tau_\eta^+(\mu)}^+ A^j, \quad X A^j A^\mu = A^j Y_{\tau_\eta^-(\mu)}^-.$$

The boundary-crossing comparison is the equality

$$Y_{\tau_\eta^+(\mu)}^+ = Y_{\tau_\eta^-(\mu)}^-.$$

These equations imply

$$\psi \in \text{span}\{V^{(N)}(A)\}.$$

□

The algebraic part of the overlapping nearest-neighbor commutation condition is

$$Q_{AX}^2 = Q_{AX}, \quad Q_{XB}^2 = Q_{XB}, \quad Q_{AX}Q_{XB} = Q_{XB}Q_{AX}$$

after these two local actions are viewed on the common three-site space. This is the local-support part of the parent commuting Hamiltonian condition in [CPGSV16, Definition D.2]. The cyclic windows beginning at i and $i + 1$ realize the two supports AX and XB , and hence overlap at the middle site.

The source local condition also fixes a three-site subspace K_{AXB} and imposes the kernel-intersection equation

$$K_{AXB} = \ker((Q_{AX})_{AX}) \cap \ker((Q_{XB})_{XB}).$$

Forgetting this intersection equation gives the algebraic overlapping two-site commutation condition displayed above.

The two lift maps also distribute over complementary projectors:

$$(1 - Q_{AX})_{AX} = 1 - (Q_{AX})_{AX}, \quad (1 - Q_{XB})_{XB} = 1 - (Q_{XB})_{XB}.$$

Consequently, if the projectors Q_{AX} and Q_{XB} commute on the three-site space, then their complementary projectors

$$P_{AX} = 1 - Q_{AX}, \quad P_{XB} = 1 - Q_{XB}$$

commute after the same lifts. This is the elementary algebraic passage used when one moves between the Q -projectors of [CPGSV16, Appendix D.2] and the complementary ground-space projectors P . Since the original two projectors are idempotent, the complementary projectors are idempotent as well; hence the pair P_{AX}, P_{XB} satisfies the same idempotent-and-commutation condition on the three-site space.

Theorem 13.9.2 (Matrices of the two adjacent factor actions). [Lean](#) [Lean](#) [✓ Stmt](#) Let $\mathcal{H} = \mathbb{C}^d$, and let Q be an endomorphism of $\mathcal{H} \otimes \mathcal{H}$. Under the canonical identifications of two- and three-site configurations with pairs and right-associated triples, the matrices of its two adjacent actions are

$$[Q_{AX}] = [Q] \otimes I_d, \quad [Q_{XB}] = I_d \otimes [Q].$$

Equivalently, for all $a, x, b, a', x', b' \in \{0, \dots, d - 1\}$,

$$[Q_{AX}]_{(a,x,b),(a',x',b')} = [Q]_{(a,x),(a',x')} \delta_{b,b'},$$

and

$$[Q_{XB}]_{(a,x,b),(a',x',b')} = \delta_{a,a'} [Q]_{(x,b),(x',b')}.$$

This is the homogeneous three-single-site specialization of the factor actions on $\mathcal{H}_A \otimes \mathcal{H}_X \otimes \mathcal{H}_B$ in [CPGSV16, Definition D.2]; here $\mathcal{H}_A = \mathcal{H}_X = \mathcal{H}_B = \mathcal{H}$. No assertion is made for unequal factor spaces.

Proof. [✓ Proof](#) On the first pair, the final coordinate is unchanged, which contributes $\delta_{b,b'}$ to the matrix coefficient. On the final pair, the first coordinate is unchanged, which contributes $\delta_{a,a'}$. $\square$

Theorem 13.9.3 (Three-site translated local terms). [Lean](#) [Lean](#) [✓ Stmt](#) Put $q(A) = 1 - P_{G_2(A)}$ for the canonical two-site parent interaction. On the three-site space $\mathcal{H}_A \otimes \mathcal{H}_X \otimes \mathcal{H}_B$, with

sites 0, 1, 2 labelled A, X, B , the two adjacent translated length-two parent interactions are the two local actions

$$h_0^{(3)}(A, 2) = q(A)_{AX}, \quad h_1^{(3)}(A, 2) = q(A)_{XB}.$$

This identifies the two translated parent interactions with the two supports appearing in [CPGSV16, Definition D.2]; it does not assert that they commute.

Proof. ✓ Proof Let $\sigma = (a, x, b)$ be a three-site configuration. For the first translated term, the length-two window and the corresponding replacement satisfy

$$\sigma_{\{0,1\}} = (a, x) = \sigma_{AX}, \quad \sigma^{\{0,1\} \leftarrow (\alpha, \xi)} = (\alpha, \xi, b) = \sigma^{AX \leftarrow (\alpha, \xi)}.$$

For the second translated term,

$$\sigma_{\{1,2\}} = (x, b) = \sigma_{XB}, \quad \sigma^{\{1,2\} \leftarrow (\xi, \beta)} = (a, \xi, \beta) = \sigma^{XB \leftarrow (\xi, \beta)}.$$

Substituting these two pairs of identities into the definition of the translated parent term gives the two displayed equations. □

Theorem 13.9.4 (Three-site commutation from overlapping supports). Lean ✓ Stmt Suppose the canonical two-site parent interaction $q(A) = 1 - P_{G_2(A)}$ satisfies the overlapping AX/XB commutation condition of Definition 13.9.1. Then, on the three-site chain,

$$h_0^{(3)}(A, 2)h_1^{(3)}(A, 2) = h_1^{(3)}(A, 2)h_0^{(3)}(A, 2).$$

This only identifies the first two translated local terms with the lifted AX and XB actions; the construction of the [CPGSV16, Appendix D.2] projectors associated with the [CPGSV16, Appendix B] basic-vector form is a separate step.

Proof. ✓ Proof By Theorem 13.9.3, the two translated local terms are $q(A)_{AX}$ and $q(A)_{XB}$. The assumed overlapping support condition says exactly that these two lifted endomorphisms commute. □

Theorem 13.9.5 (Three-site commutation from two-site representatives). Lean ✓ Stmt Let Q_{AX} and Q_{XB} be two-site projectors whose lifted AX and XB actions commute. Suppose that

$$q(A) = Q_{AX}, \quad q(A) = Q_{XB}.$$

Then the first two translated length-two local terms on the three-site chain commute.

Proof. ✓ Proof The lift identities identify the translated terms with the lifted actions of the two projectors:

$$h_0^{(3)}(A, 2) = (Q_{AX})_{AX}, \quad h_1^{(3)}(A, 2) = (Q_{XB})_{XB}.$$

The overlapping-support hypothesis is the equation

$$(Q_{AX})_{AX}(Q_{XB})_{XB} = (Q_{XB})_{XB}(Q_{AX})_{AX}.$$

Substituting the two lifted identities gives the desired commutation of $h_0^{(3)}(A, 2)$ and $h_1^{(3)}(A, 2)$. □

Theorem 13.9.6 (Three-site commutation from source projectors). [Lean](#) [✓ Stmt](#) Suppose the projectors Q_{AX} and Q_{XB} satisfy the [CPGSV16, Definition D.2] local parent-commuting condition on AXB , and suppose that the canonical two-site parent interaction is identified, as a two-site operator, with the two source projectors:

$$q(A) = Q_{AX}, \quad q(A) = Q_{XB}.$$

After lifting these two equalities to the AX and XB faces, the first two translated length-two local terms on the three-site chain commute. Thus

$$h_0^{(3)}(A, 2)h_1^{(3)}(A, 2) = h_1^{(3)}(A, 2)h_0^{(3)}(A, 2).$$

Proof. [✓ Proof](#) The lift identities of Theorem 13.9.3 identify the translated terms with $(Q_{AX})_{AX}$ and $(Q_{XB})_{XB}$. After forgetting the kernel-intersection equation, the [CPGSV16, Definition D.2] condition says exactly that these two lifted projectors commute. $\square$

Theorem 13.9.7 (Three-site commutation from complementary Appendix projectors). [Lean](#)

[Lean](#) [✓ Stmt](#) Suppose the Appendix D.2 projectors Q_{AX} and Q_{XB} satisfy the local parent-commuting condition on AXB , and suppose that the canonical two-site parent interaction is their complement on both faces:

$$q(A) = 1 - Q_{AX} = 1 - Q_{XB}.$$

Then the first two translated length-two local terms on the three-site chain commute,

$$h_0^{(3)}(A, 2)h_1^{(3)}(A, 2) = h_1^{(3)}(A, 2)h_0^{(3)}(A, 2).$$

Proof. [✓ Proof](#) The lift identities of Theorem 13.9.3 identify the translated terms with $(1 - Q_{AX})_{AX}$ and $(1 - Q_{XB})_{XB}$. The Appendix D.2 condition implies the overlapping two-site commutation condition after forgetting the kernel-intersection equation. Definition 13.9.1 then gives commutation of the complementary lifted projectors, hence the displayed commutation relation. $\square$

Definition 13.9.8 (Commuting local terms for a parent Hamiltonian). [Lean](#) [✓ Stmt](#) The parent Hamiltonian with block length L on N sites has *commuting* local terms if $h_i h_j = h_j h_i$ for all $i, j \in \{0, \dots, N-1\}$. This records the translated commutation equation $[\tau_j(P_L), P_L] = 0$ from [CPGSV16, Definition 3.9]; the same source definition also requires the ground space to be spanned, for $N > L$, by the periodic MPS vectors of the BNT components.

Definition 13.9.9 (Nearest-neighbor commuting parent Hamiltonian). [Lean](#) [✓ Stmt](#) The nearest-neighbor commuting condition is the length-2 specialization of the preceding commutation equation. For every $i, j \in \{0, \dots, N-1\}$,

$$h_i h_j = h_j h_i.$$

The source definition [CPGSV16, Definition 3.9] also requires the corresponding parent Hamiltonian to have $L = 2$ and to have the prescribed ground-space spanning property for $N > 2$.

Lemma 13.9.10 (Overlap reduction for commuting local terms). [Lean](#) [Lean](#) [✓ Stmt](#) Let $L \leq N$. Suppose that $h_i h_j = h_j h_i$ whenever the length- L cyclic supports of the two local terms meet. Then the parent Hamiltonian has commuting local terms:

$$h_i h_j = h_j h_i \quad \text{for all } i, j.$$

In the case $L = 2$, this gives the nearest-neighbor commuting condition.

Proof. [✓ Proof](#) If the two cyclic supports meet, the hypothesis gives the commutation relation. If they are disjoint, the corresponding local terms commute by locality. $\square$

Lemma 13.9.11 (Length-two cyclic-overlap alternatives). [Lean](#) [✓ Stmt](#) For $i, j \in \mathbb{Z}/N\mathbb{Z}$, if the two length-two cyclic supports starting at i and j overlap, then

$$j = i \quad \text{or} \quad j \equiv i + 1 \pmod{N} \quad \text{or} \quad i \equiv j + 1 \pmod{N}.$$

Proof. [✓ Proof](#) Expanding each length-two cyclic support gives shifts $r_i, r_j \in \{0, 1\}$ and an equality of sites $i + r_i \equiv j + r_j \pmod{N}$. The four possible pairs (r_i, r_j) give exactly the three alternatives displayed above. $\square$

Lemma 13.9.12 (Adjacent two-site translates suffice for nearest-neighbor commutation). [Lean](#) [Lean](#) [✓ Stmt](#) Let $N \geq 2$. Suppose that, for every $i \in \mathbb{Z}/N\mathbb{Z}$, the translated length-two local term commutes with its cyclic right neighbor:

$$h_i(A, 2) h_{i+1}(A, 2) = h_{i+1}(A, 2) h_i(A, 2).$$

Then all translated length-two local terms commute, and the family $p_i = h_i(A, 2)$ satisfies the finite-chain local-projector hypotheses used in the conditional Appendix B extraction.

Proof. [✓ Proof](#) By Lemma 13.9.11, overlapping length-two cyclic supports force

$$j = i \quad \text{or} \quad j \equiv i + 1 \pmod{N} \quad \text{or} \quad i \equiv j + 1 \pmod{N}.$$

The equal-start case gives $h_i(A, 2) = h_j(A, 2)$. The two nearest-neighbor cases are the displayed hypothesis and its transpose. The overlap-reduction lemma supplies the remaining disjoint pairs by locality. $\square$

Lemma 13.9.13 (Local-projector hypotheses from overlapping two-site commutation). [Lean](#) [✓ Stmt](#) Let $N \geq 2$. Suppose that the translated length-two local terms commute for every pair of overlapping cyclic windows:

$$h_i(A, 2) h_j(A, 2) = h_j(A, 2) h_i(A, 2).$$

Then the family $p_i = h_i(A, 2)$ gives the finite-chain local-projector hypotheses used in the conditional Appendix B extraction.

Proof. [✓ Proof](#) Lemma 13.9.10 turns the overlapping-pair commutation equations into commutation for all pairs. The idempotency equation is Lemma 13.3.13. $\square$

Definition 13.9.14 (Commutation and zero-energy equations for nearest-neighbor local terms). [Lean](#) [✓ Stmt](#) This condition records the commutation equation and the zero-energy equation for the nearest-neighbor local terms:

$$h_i h_j = h_j h_i, \quad h_i V^{(N)}(A) = 0.$$

This is the zero-energy part of the condition appearing in [CPGSV16, Theorem 3.10(iii)]: the MPS vector has zero energy for the translated two-site local terms, and those terms commute. The missing source assertion is that the whole parent-Hamiltonian ground space is spanned by the corresponding periodic MPS vectors from the BNT components, as recorded in Definition 13.9.15.

Theorem 13.9.19 (Chain ground-space vectors are frustration-free). [Lean](#) [✓ Stmt](#) Assume $N > 0$ and $L \leq N$. Let ψ be a vector in the periodic chain ground space $G_L^{(N)}(A)$. Then ψ satisfies all local zero-energy equations: for every site i ,

$$h_i(A, L)\psi = 0.$$

Proof. [✓ Proof](#) Write $R_{i,\eta}\psi$ for the cyclic length- L restriction of ψ at site i , with the outside configuration fixed by η . Membership in $G_L^{(N)}(A)$ gives, for every i and η ,

$$R_{i,\eta}\psi \in G_L(A) = \ker h_L(A).$$

Hence

$$h_L(A)R_{i,\eta}\psi = 0.$$

For a full configuration σ , taking $\eta = \sigma|_{\{i, \dots, i+L-1\}^c}$ gives

$$(h_i(A, L)\psi)(\sigma) = (h_L(A)R_{i,\eta}\psi)(\sigma|_{\{i, \dots, i+L-1\}}) = 0.$$

Thus $h_i(A, L)\psi = 0$ for every i . □

Lemma 13.9.20 (Zero-energy BNT vectors lie in the parent-Hamiltonian kernel). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let B be a canonical-form tensor with BNT components $A_1, \dots, A_g$. If every component vector has zero energy for the parent Hamiltonian,

$$H_L^{(N)}(B)V^{(N)}(A_j) = 0 \quad (1 \leq j \leq g),$$

then

$$\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\} \subseteq \ker H_L^{(N)}(B).$$

It is enough to assume the local frustration-free equations $h_i V^{(N)}(A_j) = 0$ for every i and j . For $N > 0$ and $L \leq N$, in the block-diagonal case $B = \bigoplus_j \mu_j A_j$, with every $\mu_j \neq 0$, the same inclusion follows because each local space $G_L(A_j)$ is contained in $G_L(B)$. This is the easy inclusion in the source ground-space spanning equation of [CPGSV16, Definition 3.9]; the reverse inclusion is the remaining spanning direction.

Proof. [✓ Proof](#) The kernel is a linear subspace. Hence it is enough to check the generators $V^{(N)}(A_j)$. The first hypothesis gives this directly. Under the local frustration-free equations,

$$H_L^{(N)}(B)V^{(N)}(A_j) = \sum_i h_i V^{(N)}(A_j) = 0.$$

For a block-diagonal tensor, write $R_{i,\tau}$ for the cyclic length- L restriction at site i , with the outside configuration fixed by τ . Then

$$R_{i,\tau}V^{(N)}(A_j) \in G_L(A_j) \subseteq G_L(B).$$

The preceding theorem therefore gives $h_i(B, L)V^{(N)}(A_j) = 0$ for every i . □

Lemma 13.9.21 (Kernel containment from chain-space splitting). [Lean](#) [✓ Stmt](#) Let $B = \bigoplus_{j=0}^{r-1} \mu_j A_j$. Suppose $N > 0$, $L \leq N$, and the periodic chain constraints for B split into the block constraints:

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=0}^{r-1} \mathcal{G}_{N,L}(A_j).$$

If every block chain space is contained in its periodic MPS line,

$$\mathcal{G}_{N,L}(A_j) \subseteq \mathbb{C}V^{(N)}(A_j) \quad (0 \leq j < r),$$

then the zero-energy space of the parent Hamiltonian is contained in the BNT vector span:

$$\ker H_L^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}.$$

Proof. [✓ Proof](#) Theorem 13.11.12 gives

$$\ker H_L^{(N)}(B) \subseteq \mathcal{G}_{N,L}(B).$$

Together with the splitting hypothesis,

$$\ker H_L^{(N)}(B) \subseteq \mathcal{G}_{N,L}(B) \subseteq \bigvee_j \mathcal{G}_{N,L}(A_j).$$

The blockwise containments give

$$\bigvee_j \mathcal{G}_{N,L}(A_j) \subseteq \bigvee_j \mathbb{C}V^{(N)}(A_j),$$

and Lemma 13.12.4 gives

$$\bigvee_j \mathbb{C}V^{(N)}(A_j) = \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}.$$

□

Lemma 13.9.22 (Kernel containment from the block-diagonal boundary comparison). [Lean](#)

[✓ Stmt](#) Let $B = \bigoplus_{j=0}^{r-1} \mu_j A_j$, with every $\mu_j \neq 0$. Suppose the physical alphabet and all block dimensions are nonempty. Assume that the block family is irreducible, left-canonical, has normalized self-overlap, and has no two blocks gauge-phase equivalent. Suppose also that each A_j is L_0 -block-injective, that $L_0 > 0$, and that

$$\sum_a A_a^j (A_a^j)^\dagger = I \quad (0 \leq j < r).$$

Let $N > 0$, $L > 0$, $L \leq N$, $L + L_0 \leq N$, and

$$(L_0 + 1) + (r-1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))) + 1 \leq L.$$

Assume the opened-boundary C^j, D^j comparison: whenever $\psi \in \mathcal{G}_{N,L}(B)$ and $\psi = \Gamma_N^B(\bigoplus_j X_j)$, there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every ρ , and every β ,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

If every block chain space is contained in its periodic MPS line,

$$\mathcal{G}_{N,L}(A_j) \subseteq \mathbb{C}V^{(N)}(A_j) \quad (0 \leq j < r),$$

then

$$\ker H_L^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}.$$

Proof. [✓ Proof](#) Lemma 13.14.2.9 gives

$$\mathcal{G}_{N,L}(B) = \bigvee_{j=0}^{r-1} \mathcal{G}_{N,L}(A_j)$$

from the stated block hypotheses and the opened-boundary comparison. Lemma 13.9.21 then applies to this equality and to the blockwise containments. $\square$

Lemma 13.9.23 (Kernel containment from boundary-crossing comparisons). [Lean](#) [✓ Stmt](#) Let $B = \bigoplus_{j=0}^{r-1} \mu_j A_j$, with every $\mu_j \neq 0$. Suppose the physical alphabet and all block dimensions are nonempty. Assume that the block family is irreducible, left-canonical, has normalized self-overlap, and has no two blocks gauge-phase equivalent. Suppose also that each A_j is L_0 -block-injective, that $L_0 > 0$, and that

$$\sum_a A_a^j (A_a^j)^\dagger = I \quad (0 \leq j < r).$$

Let $N > 0$, $L > 0$, $L \leq N$, $L + L_0 \leq N$, and

$$(L_0 + 1) + (r - 1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))) + 1 \leq L.$$

Assume that, for every boundary-crossing interval beginning at i , the simultaneous products

$$w \mapsto (A_w^1, \dots, A_w^r), \quad |w| = N - i,$$

span the product algebra $\prod_j M_{D_j}(\mathbb{C})$. If every block chain space is contained in its periodic MPS line,

$$\mathcal{G}_{N,L}(A_j) \subseteq \text{CV}^{(N)}(A_j) \quad (0 \leq j < r),$$

then

$$\ker H_L^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r - 1\}.$$

Proof. [✓ Proof](#) Lemma 13.14.2.11 gives

$$\mathcal{G}_{N,L}(B) = \bigvee_{j=0}^{r-1} \mathcal{G}_{N,L}(A_j)$$

from the block hypotheses, the local boundary-crossing constraints, and the stated tail-word span. Lemma 13.9.21 then applies to this equality and to the blockwise containments. $\square$

Lemma 13.9.24 (Block-diagonal spanning is equivalent to the reverse inclusion). [Lean](#) [Lean](#) [✓ Stmt](#) Let $B = \bigoplus_{j=0}^{r-1} \mu_j A_j$, with every $\mu_j \neq 0$. Then, for every $N > L$, the source parent-Hamiltonian spanning condition

$$\ker H_L^{(N)}(B) = \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r - 1\}$$

is equivalent to the reverse containment, for the same N :

$$\ker H_L^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r - 1\}.$$

In the nearest-neighbor case $L = 2$, the all-chain source condition of Definition 13.9.17 is equivalent to the conjunction

$$\begin{aligned} & (\forall N > 2, \forall i, j \in \{0, \dots, N - 1\}, h_i(B, 2)h_j(B, 2) = h_j(B, 2)h_i(B, 2)) \\ & \wedge \left(\forall N > 2, \ker H_2^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r - 1\} \right). \end{aligned}$$

Proof. ✓ **Proof** The forward direction of the displayed equality gives the containment by rewriting the kernel. Conversely, for $N > L$ the easy inclusion above gives

$$\text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\} \subseteq \ker H_L^{(N)}(B),$$

and the assumed reverse containment gives equality. For $L = 2$, the all-chain nearest-neighbor condition of Definition 13.9.17 decomposes as the conjunction

$$\begin{aligned} & (\forall N > 2, \forall i, j \in \{0, \dots, N-1\}, h_i(B, 2)h_j(B, 2) = h_j(B, 2)h_i(B, 2)) \\ & \wedge (\forall N > 2, \ker H_2^{(N)}(B) \subseteq \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}). \end{aligned}$$

The second conjunct is the reverse containment just identified with the source spanning clause. □

Lemma 13.9.25 (Chain-space comparison gives the block-diagonal spanning equation). Lean

Lean ✓ **Stmt** Let

$$B = \bigoplus_{j=0}^{r-1} \mu_j A_j, \quad \mu_j \neq 0.$$

Suppose that, for every $N > L$,

$$G_L^{(N)}(B) = \bigvee_{j=0}^{r-1} \text{span}\{V^{(N)}(A_j)\}.$$

Then the source parent-Hamiltonian spanning equation holds:

$$\ker H_N(B, L) = \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}.$$

It is enough to prove the two equations

$$G_L^{(N)}(B) = \bigvee_{j=0}^{r-1} G_L^{(N)}(A_j), \quad G_L^{(N)}(A_j) = \text{span}\{V^{(N)}(A_j)\}.$$

Proof. ✓ **Proof** The finite-volume identity gives

$$\ker H_N(B, L) = G_L^{(N)}(B).$$

The assumed chain-space comparison therefore rewrites the kernel as

$$\bigvee_{j=0}^{r-1} \text{span}\{V^{(N)}(A_j)\}.$$

The latter supremum is the BNT vector span. Conversely, the easy inclusion of Lemma 13.9.20 gives

$$\text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\} \subseteq \ker H_N(B, L),$$

so the displayed equation is the source spanning clause. If the comparison is first stated as $G_L^{(N)}(B) = \bigvee_j G_L^{(N)}(A_j)$, the single-block equations $G_L^{(N)}(A_j) = \text{span}\{V^{(N)}(A_j)\}$ give the previous hypothesis. □

Theorem 13.9.29 (Cyclic trace expansion for a closed MPS chain). [Lean](#) [✓ Stmt](#) For a nonempty closed chain, the trace of the word $A^{\sigma_0} \dots A^{\sigma_{L-1}}$ is the sum over cyclic virtual-index configurations:

$$\mathrm{tr}(A^{\sigma_0} \dots A^{\sigma_{L-1}}) = \sum_{\alpha_0, \dots, \alpha_{L-1}} \prod_{t=0}^{L-1} (A^{\sigma_t})_{\alpha_t, \alpha_{t+1}}, \quad \alpha_L = \alpha_0.$$

Proof. [✓ Proof](#) The trace is the diagonal sum:

$$\mathrm{tr}(A^{\sigma_0} \dots A^{\sigma_{L-1}}) = \sum_{\alpha_0} (A^{\sigma_0} \dots A^{\sigma_{L-1}})_{\alpha_0, \alpha_0}.$$

For prescribed endpoints, the matrix product entry expands as

$$(A^{\sigma_0} \dots A^{\sigma_{L-1}})_{\alpha_0, \alpha_L} = \sum_{\alpha_1, \dots, \alpha_{L-1}} \prod_{t=0}^{L-1} (A^{\sigma_t})_{\alpha_t, \alpha_{t+1}}.$$

Substituting $\alpha_L = \alpha_0$ in the trace gives the cyclic formula. $\square$

Definition 13.9.30 (Physical isometry in the basic-vector expression). [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that A is equipped with Appendix B structural data (X, Λ, U) : the matrix X is invertible, the diagonal entries of Λ are strictly positive, $A^i = X \Lambda U^i X^{-1}$, and U satisfies the pair-index isometry equation below. Let $\mathcal{H}_a$ and $\mathcal{H}_b$ have bases indexed by $\alpha, \beta \in \{0, \dots, D-1\}$. The coefficients $U_{\alpha, \beta}^i$ define maps

$$U : \mathcal{H}_a \otimes \mathcal{H}_b \longrightarrow \mathcal{H}_{\mathrm{phys}}, \quad (Uv)_i = \sum_{\alpha, \beta} U_{\alpha, \beta}^i v_{\alpha, \beta},$$

and

$$(U^* \psi)_{\alpha, \beta} = \sum_i \overline{U_{\alpha, \beta}^i} \psi_i.$$

Theorem 13.9.31 (The physical map is an isometric embedding). [Lean](#) [Lean](#) [✓ Stmt](#) Let A be equipped with Appendix B structural data (X, Λ, U) as in Definition 13.9.30. Its pair-index isometry equation

$$\sum_i \overline{U_{\alpha, \beta}^i} U_{\alpha', \beta'}^i = \delta_{\alpha, \alpha'} \delta_{\beta, \beta'}$$

implies $U^* U = 1$. In particular, U is injective.

Proof. [✓ Proof](#) Expanding the coefficient of $U^* U v$ at (α, β) and interchanging the two finite sums gives

$$(U^* U v)_{\alpha, \beta} = \sum_{\alpha', \beta'} \left(\sum_i \overline{U_{\alpha, \beta}^i} U_{\alpha', \beta'}^i \right) v_{\alpha', \beta'} = v_{\alpha, \beta}.$$

A map with a left inverse is injective. $\square$

Definition 13.9.32 (Tensor powers of the physical isometry). [Lean](#) [Lean](#) [✓ Stmt](#) For every $N \geq 0$, the coefficients of U define

$$U^{\otimes N} : (\mathcal{H}_a \otimes \mathcal{H}_b)^{\otimes N} \longrightarrow \mathcal{H}_{\mathrm{phys}}^{\otimes N},$$

with

$$(U^{\otimes N} v)_{\sigma} = \sum_{p_0, \dots, p_{N-1}} \left(\prod_{t=0}^{N-1} U_{(p_t)_a, (p_t)_b}^{\sigma_t} \right) v_{p_0, \dots, p_{N-1}}.$$

The conjugate coefficient map is

$$(U^{*\otimes N}\psi)_p = \sum_{\sigma} \left(\prod_{t=0}^{N-1} \overline{U_{(p_t)_a, (p_t)_b}^{\sigma_t}} \right) \psi_{\sigma}.$$

This is the tensor power occurring in the basic-vector expression [CPGSV16, Equation (3.17)].

Theorem 13.9.33 (The tensor powers are isometric embeddings). [Lean](#) [Lean](#) [✓ Stmt](#) For every $N \geq 0$,

$$U^{*\otimes N} U^{\otimes N} = 1.$$

Hence $U^{\otimes N}$ is injective.

Proof. [✓ Proof](#) For virtual-pair configurations p and q , interchange the two finite sums and factor the sum over physical configurations site by site:

$$\sum_{\sigma} \prod_t \overline{U_{(p_t)_a, (p_t)_b}^{\sigma_t}} U_{(q_t)_a, (q_t)_b}^{\sigma_t} = \prod_t \sum_i \overline{U_{(p_t)_a, (p_t)_b}^i} U_{(q_t)_a, (q_t)_b}^i.$$

The pair-index isometry equation turns the last product into $\prod_t \delta_{p_t, q_t} = \delta_{p, q}$. Therefore the coefficient at p is v_p . A map with a left inverse is injective. $\square$

Theorem 13.9.34 (Appendix B basis change and the parent interaction). [Lean](#) [✓ Stmt](#) Let $A^i = X \Lambda U^i X^{-1}$ be the Appendix B structural form, and write $A_{\text{core}}^i = \Lambda U^i$. The canonical local parent interaction is unchanged by this basis change:

$$q_L(A) = q_L(A_{\text{core}}).$$

Proof. [✓ Proof](#) The basis change gives a gauge equivalence between A_{core} and A , hence equality of their length- L local ground spaces. The parent interaction is the orthogonal complement projector of this local ground space. $\square$

Definition 13.9.35 (Appendix B two-site basic support). [Lean](#) [✓ Stmt](#) For Appendix B structural data, the two-site basic support is

$$G_2(A_{\text{core}}) = G_2(\Lambda U).$$

Definition 13.9.36 (Two-site bond insertion and basic-vector embedding). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Put

$$|\varphi\rangle = \sum_b \lambda_b |b, b\rangle.$$

For outer-boundary coefficients $v_{a,c}$, insertion of this vector on the middle virtual bond is the map I_{φ} defined by

$$(I_{\varphi} v)_{(a,b), (b',c)} = \delta_{b,b'} \lambda_b v_{a,c}.$$

The two-site physical map is $U^{\otimes 2} I_{\varphi}$. A boundary matrix Y determines outer-boundary coefficients by

$$(v_Y)_{a,c} = \lambda_a Y_{c,a}.$$

Opening the outer virtual bond in the two-site contraction of [CPGSV16, Equations (3.17)–(3.18)] gives these coefficients.

Definition 13.9.37 (Virtual bond projector and its adjacent placements). [Lean](#) [Lean](#) [Lean](#)

[✓ Stmt](#) Write


$$s = \langle \varphi, \varphi \rangle = \sum_b \lambda_b^2.$$

The orthogonal projector onto $\mathbb{C}\varphi$ has pair-index coefficients

$$(P_\varphi v)_{b,b'} = \delta_{b,b'} \frac{\lambda_b}{s} \sum_k \lambda_k v_{k,k}.$$

On three virtual site pairs, let P_{01} place this projector on the bond joining the right index of site zero to the left index of site one, and let P_{12} place it on the next bond.

Theorem 13.9.38 (Commutation of the adjacent virtual bond projectors). [Lean](#) [✓ Stmt](#) *The two adjacent placements commute:*



Proof. [✓ Proof](#) The contracted bond coordinates are disjoint. Thus replacing the first bond by the diagonal basis vector $|k, k\rangle$ and the second by $|\ell, \ell\rangle$ is independent of the order. Expanding both rank-one projectors gives the same double sum over (k, ℓ) . $\square$

Theorem 13.9.39 (Surjectivity of the weighted boundary parametrization). [Lean](#) [✓ Stmt](#) *Strict positivity of the λ_a makes $Y \mapsto v_Y$ surjective.*

Proof. [✓ Proof](#) For prescribed coefficients $v_{a,c}$, take $Y_{c,a} = \lambda_a^{-1} v_{a,c}$. $\square$

Theorem 13.9.40 (The two-site basic-vector image). [Lean](#) [Lean](#) [✓ Stmt](#) *Opening the outer virtual bond in the two-site contraction of [CPGSV16, Equations (3.17)–(3.18)] gives the following local-support family. For every boundary matrix Y ,*

$$U^{\otimes 2} I_\varphi(v_Y) = \Gamma_2(\Lambda U)(Y).$$

Consequently,

$$\text{ran}(U^{\otimes 2} I_\varphi) = G_2(\Lambda U).$$

Proof. [✓ Proof](#) At a two-site physical configuration (i, j) , the bond insertion gives

$$\sum_{a,b,c} U_{a,b}^i U_{b,c}^j \lambda_b \lambda_a Y_{c,a}.$$

Expanding $\text{tr}((\Lambda U^i)(\Lambda U^j)Y)$ gives the same sum, so the two maps agree after the weighted boundary parametrization. Since that parametrization is surjective, their ranges are equal. $\square$

Definition 13.9.41 (Two-site basic support projector). [Lean](#) [✓ Stmt](#) Let P_2 be the orthogonal projector onto $G_2(\Lambda U)$, as in the local parent-space construction of [CPGSV16, Definition 3.9].

Theorem 13.9.42 (Range and complement of the two-site basic support projector). [Lean](#) [Lean](#)
[✓ Stmt](#) The projector P_2 satisfies

$$P_2^2 = P_2, \quad \text{ran}(P_2) = \text{ran}(U^{\otimes 2} I_\varphi) = G_2(\Lambda U), \quad P_2 = 1 - q_2(\Lambda U).$$

In particular, $q_2(\Lambda U)$ annihilates the image of $U^{\otimes 2} I_\varphi$, while P_2 fixes that image pointwise.

Proof. [✓ Proof](#) Orthogonal projection onto $G_2(\Lambda U)$ has that subspace as its range. The preceding theorem identifies this range with $\text{ran}(U^{\otimes 2} I_\varphi)$. Orthogonal projection onto the complementary subspace is $q_2(\Lambda U)$, hence $P_2 = 1 - q_2(\Lambda U)$. Since $q_2(\Lambda U)^2 = q_2(\Lambda U)$,

$$P_2^2 = (1 - q_2(\Lambda U))^2 = 1 - q_2(\Lambda U) = P_2.$$

If ψ lies in $\text{ran}(U^{\otimes 2} I_\varphi)$, then $\psi \in G_2(\Lambda U) = \ker q_2(\Lambda U)$, and therefore

$$q_2(\Lambda U)\psi = 0, \quad P_2\psi = (1 - q_2(\Lambda U))\psi = \psi.$$

□

Theorem 13.9.43 (Commutation of adjacent two-site basic support projectors). [Lean](#) [✓ Stmt](#)
Let $(P_2)_{01}$ and $(P_2)_{12}$ be the two adjacent placements of the two-site basic support projector on three physical sites. Then

$$(P_2)_{01}(P_2)_{12} = (P_2)_{12}(P_2)_{01}.$$

This is the local three-site consequence of the parent-space construction and the basic-vector form in [CPGSV16, Definition 3.9, Equations (3.16)–(3.18), and Appendix B]. It does not assert commutation for every chain length.


Proof. [✓ Proof](#) The one-site map U is an isometry. The preceding range identity therefore identifies the physical projector with the isometric transport of the virtual bond projector:

$$P_2 = U^{\otimes 2} P_\varphi (U^*)^{\otimes 2}.$$

On three sites, multiplication of the two adjacent transports cancels the middle factors by $U^*U = 1$. Consequently their products are

$$U^{\otimes 3} P_{01} P_{12} (U^*)^{\otimes 3} \quad \text{and} \quad U^{\otimes 3} P_{12} P_{01} (U^*)^{\otimes 3},$$

respectively. They agree by Theorem 13.9.38. Diagrammatically, the same argument transports the two disjoint virtual bond contractions through the three one-site isometries:



□

Definition 13.9.44 (Appendix B AX coefficient representative). [Lean](#) [✓ Stmt](#) The AX coefficient-space representative is, before the AX -lift, the canonical two-site parent interaction of the core tensor:

$$\widehat{Q}_{AX} = q_2(A_{\text{core}}).$$

The source projector Q_{AX} of Definition D.2 acts on $\mathcal{H}_A \otimes \mathcal{H}_X$. The displayed operator is the two-site coefficient representative that will be placed on the AX face; it is not by itself the construction of that source projector.

Definition 13.9.45 (Appendix B XB coefficient representative). [Lean](#) [✓ Stmt](#) The XB coefficient-space representative is, before the XB -lift, the same coefficient-space representative:

$$\widehat{Q}_{XB} = \widehat{Q}_{AX}.$$

The source projector Q_{XB} of Definition D.2 acts on $\mathcal{H}_X \otimes \mathcal{H}_B$. The displayed operator is the coefficient representative later placed on the XB face.

Theorem 13.9.46 (The (AX) factor action). [Lean](#) [✓ Stmt](#) On the homogeneous three-single-site coefficient space $(\mathbb{C}^d)^{\otimes 3}$, the hatted Appendix B (AX) representative acts as

$$[\widehat{Q}_{AX}]_{AX} = [\widehat{Q}_{AX}] \otimes I_d.$$

This equation gives the coefficient-factor specialization of the left local action in [CPGSV16, Definition D.2]. It does not construct a projector for an arbitrary factorization $\mathcal{H}_A \otimes \mathcal{H}_X \otimes \mathcal{H}_B$.

Proof. [✓ Proof](#) Apply the left identity in Theorem 13.9.2 to $\widehat{Q}_{AX}$. □

Theorem 13.9.47 (The (XB) factor action). [Lean](#) [✓ Stmt](#) On the homogeneous three-single-site coefficient space $(\mathbb{C}^d)^{\otimes 3}$, the hatted Appendix B (XB) representative acts as

$$[\widehat{Q}_{XB}]_{XB} = I_d \otimes [\widehat{Q}_{XB}].$$

This equation gives the coefficient-factor specialization of the right local action in [CPGSV16, Definition D.2]. It does not construct a projector for an arbitrary factorization $\mathcal{H}_A \otimes \mathcal{H}_X \otimes \mathcal{H}_B$.

Proof. [✓ Proof](#) Apply the right identity in Theorem 13.9.2 to $\widehat{Q}_{XB}$. □

Lemma 13.9.48 (Appendix B coefficient representatives and the parent interaction). [Lean](#) [Lean](#) [✓ Stmt](#) As coefficient-space representatives,

$$\widehat{Q}_{AX} = q_2(A), \quad \widehat{Q}_{XB} = q_2(A).$$

Proof. [✓ Proof](#) Both operators are $q_2(A_{\text{core}})$, and the Appendix B basis change leaves q_2 unchanged. □

Lemma 13.9.49 (Appendix B two-site representative idempotents). [Lean](#) [Lean](#) [✓ Stmt](#) The two Appendix B two-site coefficient representatives are idempotent:

$$\widehat{Q}_{AX}^2 = \widehat{Q}_{AX}, \quad \widehat{Q}_{XB}^2 = \widehat{Q}_{XB}.$$

Proof. [✓ Proof](#) By the preceding identification, both representatives are the canonical two-site parent interaction $q_2(A)$. The latter is an orthogonal projector, hence is idempotent. □

Theorem 13.9.50 (Appendix B overlapping two-site commutation). [Lean](#) [✓ Stmt](#) *The two Appendix B coefficient representatives satisfy*

$$\widehat{Q}_{AX}^2 = \widehat{Q}_{AX}, \quad \widehat{Q}_{XB}^2 = \widehat{Q}_{XB},$$

and their adjacent actions on the common three-site coefficient space commute:

$$(\widehat{Q}_{AX})_{AX}(\widehat{Q}_{XB})_{XB} = (\widehat{Q}_{XB})_{XB}(\widehat{Q}_{AX})_{AX}.$$

Thus they satisfy the algebraic overlapping two-site condition of Definition 13.9.1. This is the coefficient-space analogue of the projector-and-commutator part of [CPGSV16, Definition D.2]. The hatted maps have not yet been identified with the source projectors on the explicitly factored spaces. By itself the result does not prove the kernel-intersection equation in that definition and makes no assertion for every chain length. The coefficient-space kernel equation is established in the next two results.

Proof. [✓ Proof](#) Both coefficient representatives are the complement $1 - P_2$ of the two-site basic support projector. Their idempotency was established in Lemma 13.9.49. Since the two adjacent lifts of P_2 commute, their complementary lifts commute as well. $\square$

Theorem 13.9.51 (Three-site intersection of adjacent parent kernels). [Lean](#) [Lean](#) [✓ Stmt](#) *If A is injective, then*

$$G_3(A) = \ker(q_2(A)_{AX}) \cap \ker(q_2(A)_{XB}).$$

More generally, on $L + 1$ sites, simultaneous annihilation by the two adjacent length- L parent interactions is equivalent to membership in $G_{L+1}(A)$ whenever A is injective and $L > 1$.

Proof. [✓ Proof](#) By Theorem 13.4.4.1, a three-site vector belongs to $G_3(A)$ exactly when its restrictions to both adjacent pairs belong to $G_2(A)$. Since $\ker q_2(A) = G_2(A)$, this is equivalent to annihilation by both adjacent terms. Theorem 13.9.3 identifies these terms with the two displayed lifts. $\square$

Theorem 13.9.52 (Appendix B coefficient-space analogue of the Appendix D kernel datum).

[Lean](#) [Lean](#) [✓ Stmt](#) *Let A be injective and let an Appendix B structural datum be fixed. Put*

$$\widehat{K}_{AXB} = G_3(A), \quad \widehat{Q}_{AX} = q_2(A), \quad \widehat{Q}_{XB} = q_2(A),$$

with the two copies of $q_2(A)$ lifted to the indicated faces of the three-site coefficient space. Then

$$[(\widehat{Q}_{AX})_{AX}, (\widehat{Q}_{XB})_{XB}] = 0, \quad \widehat{K}_{AXB} = \ker((\widehat{Q}_{AX})_{AX}) \cap \ker((\widehat{Q}_{XB})_{XB}).$$

These are the coefficient-space analogues of the idempotence, commutation, and kernel-intersection equations in [CPGSV16, Definition D.2]. The hatted operators are not identified here with the source projectors on $\mathcal{H}_A \otimes \mathcal{H}_X$ and $\mathcal{H}_X \otimes \mathcal{H}_B$. For a normal left-canonical RFP tensor, injectivity follows from the RFP structural theorem and is not an additional hypothesis.

Proof. [✓ Proof](#) Idempotency and commutation are Theorem 13.9.50. The two coefficient representatives are $q_2(A)$, and Theorem 13.9.51 supplies the remaining kernel equation. $\square$

Theorem 13.9.53 (Transport of the three-site commutator). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Suppose the first two length-two parent interactions commute on the three-site chain:*

$$h_0^{(3)}h_1^{(3)} = h_1^{(3)}h_0^{(3)}.$$


Then, for every $N > 2$ and every $i \in \mathbb{Z}/N\mathbb{Z}$, the two adjacent translates commute:

$$h_i^{(N)} h_{i+1}^{(N)} = h_{i+1}^{(N)} h_i^{(N)}.$$

For each exterior configuration τ , let $R_{i,\tau}^{(3)}$ restrict to the cyclic three-site window beginning at i . It intertwines the two N -site terms with the two three-site terms:

$$R_{i,\tau}^{(3)} h_i^{(N)} = h_0^{(3)} R_{i,\tau}^{(3)}, \quad R_{i,\tau}^{(3)} h_{i+1}^{(N)} = h_1^{(3)} R_{i,\tau}^{(3)}.$$

Diagrammatically, the same local AXB calculation is placed at every cyclic translate:



The hypothesis $N > 2$ is essential to this transport: at $N = 2$ the two cyclic windows contain the same sites in opposite orders. This case is not part of [CPGSV16, Theorem 3.10].

Proof. ✓ Proof Restrict an N -site vector to the cyclic window $(i, i+1, i+2)$ while holding all other physical indices fixed. Window extraction and replacement identify the two restricted local terms with $h_0^{(3)}$ and $h_1^{(3)}$. Apply the three-site commutator to the restricted vector and then evaluate at the original three-site configuration. Replacing a window by the values extracted from that configuration recovers the original N -site configuration. $\square$

Theorem 13.9.54 (Appendix B adjacent commutation on periodic chains). Lean ✓ Stmt For every Appendix B structural datum and every periodic chain of length $N > 2$, the adjacent translated length-two parent interactions commute.

Proof. ✓ Proof Theorem 13.9.50 gives the three-site commutator, and Theorem 13.9.53 places it at every adjacent pair on the periodic chain. $\square$

Definition 13.9.55 (Appendix B local-projector family). Lean ✓ Stmt For an Appendix B structural datum and a periodic chain of length $N > 2$, set

$$p_i = h_i^{(N)}, \quad i \in \mathbb{Z}/N\mathbb{Z},$$

where $h_i^{(N)}$ is the translated length-two parent interaction. These maps are idempotent and commute pairwise: adjacent commutation follows from Theorem 13.9.54, while disjoint length-two interactions commute by locality.

Theorem 13.9.56 (Left-canonical RFP commutation and zero energy). Lean Lean Lean ✓ Stmt A normal left-canonical renormalization fixed-point tensor of nonzero virtual dimension has pairwise commuting length-two parent interactions on every periodic chain of length $N > 2$, and its periodic matrix product vector is annihilated by every such interaction. The removal of the left-canonical hypothesis and the ground-space spanning equation in [CPGSV16, Definition 3.9] remain separate.

Proof. ✓ Proof The Appendix B structural theorem supplies the datum used in Definition 13.9.55. Its commuting idempotents are precisely the translated parent interactions. The standard frustration-free identity shows that each interaction annihilates the periodic matrix product vector. $\square$

Lemma 13.9.57 (Appendix B overlapping condition from the lifted commutator). Lean ✓ Stmt
Suppose the lifted AX and XB representatives commute:

$$(\widehat{Q}_{AX})_{AX}(\widehat{Q}_{XB})_{XB} = (\widehat{Q}_{XB})_{XB}(\widehat{Q}_{AX})_{AX}.$$

Then $\widehat{Q}_{AX}$ and $\widehat{Q}_{XB}$ satisfy the algebraic overlapping two-site condition of Definition 13.9.1.

Proof. ✓ Proof The idempotency clauses are supplied by Lemma 13.9.49. The only remaining clause in Definition 13.9.1 is exactly the displayed lifted commutator. $\square$

Lemma 13.9.58 (Definition D.2 projectors identified with Appendix B representatives). Lean ✓ Stmt
Let Q_{AX} and Q_{XB} be the two-site coefficient-space transports of source projectors satisfying the Definition D.2 local condition on AXB . If these transported projectors agree with the Appendix B coefficient representatives,

$$Q_{AX} = \widehat{Q}_{AX}, \quad Q_{XB} = \widehat{Q}_{XB},$$

then $\widehat{Q}_{AX}$ and $\widehat{Q}_{XB}$ satisfy the algebraic overlapping two-site condition of Definition 13.9.1. Consequently,

$$h_0^{(3)}(A, 2)h_1^{(3)}(A, 2) = h_1^{(3)}(A, 2)h_0^{(3)}(A, 2).$$

Proof. ✓ Proof Rewriting the two projectors in the Definition D.2 condition as $\widehat{Q}_{AX}$ and $\widehat{Q}_{XB}$ gives the algebraic overlapping condition. The three-site commutation is then Lemma 13.9.60. $\square$

Lemma 13.9.59 (Three-site lifts of the Appendix B coefficient representatives). Lean Lean ✓ Stmt
On the three-site A, X, B window, the two adjacent length-two local terms are the two local lifts

$$h_0^{(3)}(A, 2) = (\widehat{Q}_{AX})_{AX}, \quad h_1^{(3)}(A, 2) = (\widehat{Q}_{XB})_{XB}.$$

Proof. ✓ Proof Apply the three-site local-term identities to $q_2(A)$, and replace $q_2(A)$ by $\widehat{Q}_{AX}$ or $\widehat{Q}_{XB}$ using the preceding two-site identifications. $\square$

Lemma 13.9.60 (Appendix B overlapping condition gives three-site local-term commutation). Lean ✓ Stmt
If the Appendix B AX and XB coefficient representatives satisfy the overlapping two-site commutation datum

$$\widehat{Q}_{AX}^2 = \widehat{Q}_{AX}, \quad \widehat{Q}_{XB}^2 = \widehat{Q}_{XB}, \quad (\widehat{Q}_{AX})_{AX}(\widehat{Q}_{XB})_{XB} = (\widehat{Q}_{XB})_{XB}(\widehat{Q}_{AX})_{AX},$$

then the corresponding first two translated length-two local terms on the three-site window commute:

$$h_0^{(3)}(A, 2)h_1^{(3)}(A, 2) = h_1^{(3)}(A, 2)h_0^{(3)}(A, 2).$$

Proof. ✓ Proof By Lemma 13.9.59, the two translated local terms are exactly the AX and XB lifts of $\widehat{Q}_{AX}$ and $\widehat{Q}_{XB}$. The displayed equality is therefore the commutation clause of the overlapping two-site datum. $\square$

[Lean](#) [Lean](#) [✓ Stmt](#) *If the Appendix B AX and XB coefficient representatives satisfy the lifted Definition D.2 commutator*

then the corresponding first two translated length-two local terms on the three-site window commute:

The same conclusion holds if one supplies a Definition D.2-shaped local condition whose two operators are these coefficient-space representatives, rather than the source projectors themselves.

$$K_{AXB} = \ker((\widehat{Q}_{AX})_{AX}) \cap \ker((\widehat{Q}_{XB})_{XB}).$$

[illegible]

$$A^i = X \Lambda U^i X^{-1}.$$

$$\sum_i \overline{(U^i)_{\alpha,\beta}} (U^i)_{\alpha',\beta'} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

$$\sum_i \overline{(\Lambda U^i)_{\alpha,\beta}} (\Lambda U^i)_{\alpha',\beta'} = \lambda_\alpha \lambda_{\alpha'} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

$$C_L(\sigma) = \sum_{\alpha_0, \dots, \alpha_{L-1}} \prod_{t=0}^{L-1} \Lambda_{\alpha_t}(U^{\sigma_t})_{\alpha_t, \alpha_{t+1}}, \quad \alpha_L = \alpha_0.$$

$$|V^{(L)}(A_j)\rangle = U^{\otimes L}|\varphi_j\rangle^{\otimes L}, \quad |\varphi_j\rangle = \sum_m \lambda_m |m, m\rangle,$$

$$C_2(\sigma) = \phi_2(\sigma).$$

The change of basis does not alter either the local spaces G_L or the periodic MPS coefficients:

$$G_L(A) = G_L(\Lambda U).$$

It therefore also leaves the canonical parent interaction unchanged:

$$q_L(A) = q_L(\Lambda U).$$

At length two, the Appendix B two-site basic support is $G_2(\Lambda U)$, and the coefficient-space representatives $\widehat{Q}_{AX}$ and $\widehat{Q}_{XB}$ are the two copies of $q_2(\Lambda U)$ before they are lifted to the AX and XB faces. Hence

$$\widehat{Q}_{AX} = q_2(A), \quad \widehat{Q}_{XB} = q_2(A),$$

as coefficient-space representatives. On the three-site window their two lifts give the local identifications

$$h_0^{(3)}(A, 2) = (\widehat{Q}_{AX})_{AX}, \quad h_1^{(3)}(A, 2) = (\widehat{Q}_{XB})_{XB}.$$

Consequently, for every chain length $L \geq 1$, the original tensor has the same cyclic coefficient formula:

$$V^{(L)}(A)_\sigma = C_L(\sigma).$$

In particular, the corresponding two-site amplitude is

$$\phi_2(a, b) = V^{(2)}(A)(a, b) = V^{(2)}(\Lambda U)(a, b).$$

Thus the one-pair case of the factorization is already verified for the core tensor:

$$V^{(2)}(\Lambda U)(\sigma) = \phi_2(\sigma_0, \sigma_1).$$

The even-chain physical-pair factorization is the following condition, for a positive number n of pairs,

$$V^{(2n)}(A)(\sigma) = \prod_{p=0}^{n-1} \phi_2(\sigma_{2p}, \sigma_{2p+1}).$$

The case $n = 0$ is not part of this condition: the empty-chain MPV coefficient is the virtual bond dimension, whereas the empty product on the right-hand side is 1. The first equality, $V^{(L)}(\Lambda U)_\sigma = C_L(\sigma)$, is the trace expansion of the closed matrix product and is the coefficient form of $U^{\otimes L} \varphi_j^{\otimes L}$. The remaining coefficient step is to relate this source expression, with the physical word σ fixed, to the even-chain physical-pair factorization above. On a three-site chain, the two adjacent length-two local terms are the AX and XB actions of the canonical two-site parent interaction. Theorem 13.9.50 constructs the adjacent physical support projectors from the basic-vector expression and proves the overlapping relation

$$(\widehat{Q}_{AX})_{AX}(\widehat{Q}_{XB})_{XB} = (\widehat{Q}_{XB})_{XB}(\widehat{Q}_{AX})_{AX}.$$

Theorem 13.9.53 transports this identity around a periodic chain. Consequently, for every $N > 2$ and every pair of sites $i, j \in \mathbb{Z}/N\mathbb{Z}$,

$$h_i(A, 2)h_j(A, 2) = h_j(A, 2)h_i(A, 2).$$

Definition 13.9.55 collects these maps into a commuting idempotent family; idempotency is Lemma 13.3.13. Together with Lemma 13.3.19, this gives the zero-energy equation for $V^{(N)}(A)$.

Neither conclusion requires the even-chain physical-pair factorization displayed above. That factorization remains a separate conditional coefficient statement. The full source implication additionally requires removal of the left-canonical hypothesis and the ground-space spanning equation. The coefficient-space analogue of the local kernel-intersection equation is Theorem 13.9.52; identifying its hatted maps with the source projectors remains separate and is not needed for the chain commutator.

Theorem 13.9.63 (Even-chain factorization and ground-space spanning give all-chain NNCPH).

[Lean](#) [Lean](#) [✓ Stmt](#) Suppose the conditional even-chain hypotheses give commuting two-site parent terms for a tensor B . If the nearest-neighbor parent Hamiltonian also satisfies the ground-space spanning equation of Definition 13.9.15, with BNT components $A_1, \dots, A_g$, then B satisfies the all-chain nearest-neighbor commuting parent-Hamiltonian ground-space condition of Definition 13.9.17.

In particular, if a normal left-canonical RFP tensor has nonzero bond dimension, the same conditional Appendix B hypotheses, and the same ground-space spanning equation, then it satisfies the full all-chain condition in [CPGSV16, Theorem 3.10(iii)].

Proof. [✓ Proof](#) The conditional hypotheses give $h_i h_j = h_j h_i$ for all translated two-site local terms. Lemma 13.3.19 gives $h_i V^{(N)}(B) = 0$. For every $N > 2$, the spanning hypothesis supplies

$$\ker H_2^{(N)}(B) = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$$

These are exactly the three clauses in Definition 13.9.16, uniformly in N . $\square$

Theorem 13.9.64 (All-chain nearest-neighbor commuting parent-Hamiltonian ground spaces imply a renormalization fixed point). [Lean](#) [Lean](#) [✓ Stmt](#) Let B be normal and in left-canonical form. Suppose that the family A_j is a BNT for B , and that the nearest-neighbor parent Hamiltonian of B has the all-chain ground-space property of Definition 13.9.17: for every $N > 2$ the length-two translated parent interactions commute,

$$h_i(B, 2)h_j(B, 2) = h_j(B, 2)h_i(B, 2),$$

the vector $V^{(N)}(B)$ has zero energy, and

$$\ker H_2^{(N)}(B) = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

Then B is a renormalization fixed point. At present this reverse implication remains an unproved hypothesis, representing Beigi's one-dimensional nearest-neighbor commuting-Hamiltonian ground-space theorem and the identification of the resulting locally orthogonal states with the BNT of B .

13.10 Decorrelation and the idempotent-product parent-commuting condition

This section develops the idempotent-product formulation used for the decorrelation implication from Appendix D. It is not the full source condition. In [CPGSV16, Appendix D.2], the parent commuting Hamiltonian condition is stated for local projectors Q_{AX} and Q_{XB} with

$$[Q_{AX}, Q_{XB}] = 0, \quad K_{AXB} = (K_{AX} \otimes H_B) \cap (H_A \otimes K_{XB}).$$

Writing $P_{AX} = 1 - Q_{AX}$ and $P_{XB} = 1 - Q_{XB}$ gives, for the idempotent P_K used below, the identity $P_{AX}P_{XB} = P_K$. The following definition records this idempotent-product consequence, not the tensor-product locality, orthogonal-projector, or ground-space-intersection hypotheses of Appendix D.2.

The following preparatory result is independent of the decorrelation condition. For a positive semidefinite operator on a bipartite space, the support projector of the retained left marginal, lifted to the full space, fixes the original operator. These are the support-projector identities used in [CPGSV16, Appendix D.2, lines 2225–2235].

Lemma 13.10.1 (Adjoint identity for the right partial trace). Lean ✓ Stmt *For every operator M on H_L and every operator ρ on $H_L \otimes H_R$,*

$$\text{tr}((M \otimes \mathbf{1}_R)\rho) = \text{tr}(M \text{tr}_R \rho).$$

Proof. ✓ Proof Expand both traces in product bases and sum first over the identity entry on H_R . □

Theorem 13.10.2 (Left absorption by the marginal support projector). Lean ✓ Stmt *Let ρ be positive semidefinite on $H_L \otimes H_R$, and let P_L be the orthogonal projector onto the support of $\text{tr}_R \rho$. Then*

$$(P_L \otimes \mathbf{1}_R)\rho = \rho.$$

Proof. ✓ Proof Set $Q_L = \mathbf{1}_L - P_L$. Since $Q_L \text{tr}_R \rho = 0$, the adjoint identity gives

$$\text{tr}((Q_L \otimes \mathbf{1}_R)\rho) = \text{tr}(Q_L \text{tr}_R \rho) = 0.$$

Positivity of ρ then implies $(Q_L \otimes \mathbf{1}_R)\rho = 0$. Finally, $\mathbf{1} - (Q_L \otimes \mathbf{1}_R) = P_L \otimes \mathbf{1}_R$ gives the claim. □

Theorem 13.10.3 (Right absorption by the marginal support projector). Lean ✓ Stmt *Under the same assumptions,*

$$\rho(P_L \otimes \mathbf{1}_R) = \rho.$$

For $H_L = H_A \otimes H_X$ and $H_R = H_B$, the two absorption identities are precisely the P_{AX} support-projector identities in [CPGSV16, Appendix D.2, lines 2228–2235].

Proof. ✓ Proof Apply $(\cdot)^\dagger$ to the left absorption identity. Since $\rho^\dagger = \rho$ and $(P_L \otimes \mathbf{1}_R)^\dagger = P_L \otimes \mathbf{1}_R$,

$$\rho(P_L \otimes \mathbf{1}_R) = [(P_L \otimes \mathbf{1}_R)\rho]^\dagger = \rho^\dagger = \rho.$$

□

Definition 13.10.4 (Decorrelation). Lean ✓ Stmt Given an endomorphism P_K and sets of observables $\mathcal{O}_A, \mathcal{O}_B$, regions A and B are *decorrelated* w.r.t. K when $P_K \circ \mathcal{O}_A \circ (1 - P_K) \circ \mathcal{O}_B \circ P_K = 0$ for all $\mathcal{O}_A \in \mathcal{O}_A, \mathcal{O}_B \in \mathcal{O}_B$.

Definition 13.10.5 (Idempotent-product parent-commuting structure). Lean ✓ Stmt The structure used below records only the idempotent-product consequence of the parent commuting Hamiltonian condition of [CPGSV16, Appendix D.2]. It consists of commuting idempotent endomorphisms P_{AX} and P_{XB} such that, for the idempotent P_K associated to K ,

$$P_{AX} \circ P_{XB} = P_K.$$

In the source, this is the equation obtained from local projectors Q_{AX} and Q_{XB} satisfying $[Q_{AX}, Q_{XB}] = 0$ and $K_{AXB} = (K_{AX} \otimes H_B) \cap (H_A \otimes K_{XB})$.

Theorem 13.10.6 (Idempotence gives tautological idempotent-product data). [Lean](#) [✓ Stmt](#) If

$$P_K \circ P_K = P_K,$$

then P_K has idempotent-product data.

Proof. [✓ Proof](#) Take $P_{AX} = P_{XB} = P_K$. Their commutativity is immediate, and the product identity is exactly the assumed idempotence of P_K . $\square$

Lemma 13.10.7 (Cancellation for commuting idempotents). [Lean](#) [✓ Stmt](#) If P and Q are commuting idempotents, then

$$Q \circ (1 - P \circ Q) \circ P = 0.$$

Proof. [✓ Proof](#) The cancellation is the computation

$$Q(1 - PQ)P = QP - QPQP = QP - QPPQ = QP - QP = 0,$$

using commutativity and idempotence. $\square$

Theorem 13.10.8 (Idempotence of the product projector). [Lean](#) [✓ Stmt](#) If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_K \circ P_K = P_K.$$

Proof. [✓ Proof](#) Since P_{AX} and P_{XB} are idempotent and commute, their product $P_{AX} \circ P_{XB}$ is idempotent. Substituting $P_K = P_{AX} \circ P_{XB}$ gives the claim. $\square$

Theorem 13.10.9 (Absorption by the left projector). [Lean](#) [✓ Stmt](#) If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_{AX} \circ P_K = P_K.$$

Proof. [✓ Proof](#) Using $P_K = P_{AX} \circ P_{XB}$ and the idempotence of P_{AX} ,

$$P_{AX} \circ P_K = P_{AX} \circ P_{AX} \circ P_{XB} = P_{AX} \circ P_{XB} = P_K.$$

$\square$

Theorem 13.10.10 (Absorption by the right projector). [Lean](#) [✓ Stmt](#) If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_K \circ P_{XB} = P_K.$$

Proof. [✓ Proof](#) Using $P_K = P_{AX} \circ P_{XB}$ and the idempotence of P_{XB} ,

$$P_K \circ P_{XB} = P_{AX} \circ P_{XB} \circ P_{XB} = P_{AX} \circ P_{XB} = P_K.$$

$\square$

Theorem 13.10.11 (Cross-absorption by the right projector). [Lean](#) [✓ Stmt](#) If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_{XB} \circ P_K = P_K.$$

Proof. [✓ Proof](#) By commutativity, $P_{XB} \circ P_K = P_{XB} \circ P_{AX} \circ P_{XB} = P_{AX} \circ P_{XB} \circ P_{XB}$, and idempotence of P_{XB} yields $P_{XB} \circ P_K = P_K$. $\square$

Theorem 13.10.12 (Cross-absorption by the left projector). [Lean](#) [✓ Stmt](#) If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_K \circ P_{AX} = P_K.$$

Proof. ✓ Proof By commutativity, $P_K \circ P_{AX} = P_{AX} \circ P_{XB} \circ P_{AX} = P_{AX} \circ P_{AX} \circ P_{XB}$, and idempotence of P_{AX} yields $P_K \circ P_{AX} = P_K$. $\square$

Theorem 13.10.13 (Reverse product identity). Lean ✓ Stmt If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_{XB} \circ P_{AX} = P_K.$$

Proof. ✓ Proof This follows from commutativity of P_{AX} and P_{XB} together with the identity $P_{AX} \circ P_{XB} = P_K$. $\square$

Theorem 13.10.14 (Commuting complementary projectors). Lean ✓ Stmt If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data and

$$Q_{AX} = 1 - P_{AX}, \quad Q_{XB} = 1 - P_{XB},$$

then

$$Q_{AX} \circ Q_{XB} = Q_{XB} \circ Q_{AX}.$$

Proof. ✓ Proof Expanding both products and using $P_{AX} \circ P_{XB} = P_{XB} \circ P_{AX}$ shows that the complementary projectors commute. $\square$

Theorem 13.10.15 (Idempotent-product data imply decorrelation). Lean ✓ Stmt If P_{AX} and P_{XB} are idempotent endomorphisms satisfying $P_{AX} \circ P_{XB} = P_K$ and $P_{AX} \circ P_{XB} = P_{XB} \circ P_{AX}$, and A -observables commute with P_{XB} while B -observables commute with P_{AX} , then regions A and B are decorrelated w.r.t. K . This is the idempotent-product form of the backward direction of [CPGSV16, Appendix D, Section D.2, Proposition D.3].

Proof. ✓ Proof The key identity is $P_{XB} \circ (1 - P_{AX} \circ P_{XB}) \circ P_{AX} = 0$. Substituting $P_K = P_{AX} \circ P_{XB}$ and using $O_A \circ P_{XB} = P_{XB} \circ O_A$ and $O_B \circ P_{AX} = P_{AX} \circ O_B$ gives

$$P_K \circ O_A \circ (1 - P_K) \circ O_B \circ P_K = P_{AX} \circ O_A \circ [P_{XB} \circ (1 - P_{AX} \circ P_{XB}) \circ P_{AX}] \circ O_B \circ P_{XB} = 0.$$

$\square$

Theorem 13.10.16 (Decorrelating idempotent-product data). Lean ✓ Stmt Corollary of Theorem 13.10.15: if P_K has idempotent-product data, the A -observables commute with P_{XB} , and the B -observables commute with P_{AX} , then regions A and B are decorrelated w.r.t. K .

Proof. ✓ Proof Apply Theorem 13.10.15 to the witnessing endomorphisms P_{AX} and P_{XB} supplied by the idempotent-product data. $\square$

Lemma 13.10.17 (Frustration-free identity). Lean ✓ Stmt For any endomorphisms P and Q ,

$$(1 - P) + (1 - Q) - (1 - P) \circ (1 - Q) = 1 - P \circ Q.$$

Proof. ✓ Proof Expanding the left-hand side and cancelling gives the result. $\square$

Theorem 13.10.18 (Idempotent-product Hamiltonian identity). Lean ✓ Stmt If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$Q_{AX} + Q_{XB} - Q_{AX} \circ Q_{XB} = 1 - P_K,$$

where $Q_{AX} = 1 - P_{AX}$ and $Q_{XB} = 1 - P_{XB}$ are the complementary projectors. This is the algebraic identity obtained from the complementary projectors in [CPGSV16, Appendix D, Section D.2]; it is not the full source parent commuting Hamiltonian condition.

Proof. ✓ Proof Applying Lemma 13.10.17 to P_{AX} and P_{XB} , and substituting $P_{AX} \circ P_{XB} = P_K$. $\square$

Theorem 13.10.19 (Ground-space membership). Lean ✓ Stmt If $P_K = P_{AX} \circ P_{XB}$ is idempotent-product data, then

$$P_K v = v \iff P_{AX} v = v \text{ and } P_{XB} v = v.$$

This is the algebraic consequence corresponding to equation (D.2) from [CPGSV16]: $K_{AXB} = (K_{AX} \otimes H_B) \cap (H_A \otimes K_{XB})$. The source equation itself also carries the tensor-product local-subspace hypotheses.

Proof. ✓ Proof $(\Rightarrow)$ If $P_K v = v$, then $P_{AX}(P_K v) = (P_{AX} \circ P_K)v = P_K v = v$ by left absorption, and similarly for P_{XB} .

$(\Leftarrow)$ If $P_{AX} v = v$ and $P_{XB} v = v$, then $P_K v = (P_{AX} \circ P_{XB})v = P_{AX}(P_{XB} v) = P_{AX} v = v$. $\square$

13.11 Spectral gap

A frustration-free Hamiltonian is *gapped* if the smallest nonzero eigenvalue, equivalently the gap above the ground space, is bounded away from zero uniformly in the chain length N . For MPS parent Hamiltonians, the martingale method of Fannes–Nachtergaele–Werner and Nachtergaele [FNW92, Nac96], as recalled by Cirac–Perez-Garcia–Schuch–Verstraete [CPGSV21, Section IV.C] and in the later formulation of Kastoryano and Lucia [KL18], reduces a uniform gap to an anticommutator estimate for overlapping local terms. Principal angles between the corresponding local ground spaces give the geometric interpretation of this estimate. The results below state the anticommutator condition for the cyclic-window local terms and also record cyclic norm-compression estimates as sufficient stronger hypotheses.

Definition 13.11.1 (Parent ground space under the canonical ℓ^2 identification). Lean ✓ Stmt Denote by $\mathcal{G}_{N,L}^{\text{ES}}(A)$ the parent ground space after identifying the coefficient space with the Hilbert space $\ell^2(\{0, \dots, d-1\}^N)$.

Definition 13.11.2 (Parent Hamiltonian under the canonical ℓ^2 identification). Lean ✓ Stmt Denote by $H_{N,L}^{\text{ES}}(A)$ the conjugate of $H_N(A, L)$ by the canonical identification between the coefficient space and $\ell^2(\{0, \dots, d-1\}^N)$.

Theorem 13.11.3 (Kernel under the canonical ℓ^2 identification). Lean ✓ Stmt Under the canonical ℓ^2 identification of the N -site state space, the parent ground space is exactly the kernel of the corresponding parent Hamiltonian.

Proof. ✓ Proof This is an immediate unraveling of the definitions: both constructions are obtained from the same parent Hamiltonian by conjugating with the canonical identification between the site space and the ℓ^2 space, so the ground space is precisely the kernel after this identification. $\square$

Lemma 13.11.4 (Membership criterion under the canonical ℓ^2 identification). Lean ✓ Stmt A vector belongs to $\mathcal{G}_{N,L}^{\text{ES}}(A)$ exactly when the parent Hamiltonian annihilates it.

Remark 13.11.5 (Local projectors under the canonical ℓ^2 identification). Lean Lean Lean Lean ✓ Stmt Let $P_L^{\text{ES}}(A)$ denote the orthogonal projector onto $(G_L^{\text{ES}}(A))^\perp$ on the canonical ℓ^2 L -site Hilbert space. For a cyclic window starting at i and an outside configuration τ , the map $R_{i,\tau}$ restricts an N -site vector to that length- L cyclic window with the outside configuration fixed by τ . The local term $h_{i,\text{ES}}$ is the corresponding N -site local projector, and $R_{i,\tau}^\dagger P_L^{\text{ES}}(A) R_{i,\tau}$ is the basic positive summand in its cyclic-averaging formula.

Lemma 13.11.6 (Positivity of the Hilbert-space local summands). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The Hilbert-space local projector $P_L^{\text{ES}}(A)$ is positive. Consequently, every summand $R_{i,\tau}^\dagger P_L^{\text{ES}}(A) R_{i,\tau}$ is positive, and so is the finite sum over τ .*

Proof. [✓ Proof](#) Orthogonal projectors are positive. Conjugation by $R_{i,\tau}$ preserves positivity, and finite sums of positive operators remain positive. $\square$

Lemma 13.11.7 (Cyclic averaging for the Hilbert-space local terms). [Lean](#) [Lean](#) [✓ Stmt](#) *The Hilbert-space parent Hamiltonian is the sum of the Hilbert-space local terms $H_{N,\text{ES}} = \sum_i h_{i,\text{ES}}$, and each local term admits the exact cyclic-averaging formula*

$$h_{i,\text{ES}} = d^{-L} \sum_{\tau} R_{i,\tau}^\dagger P_L^{\text{ES}}(A) R_{i,\tau}. \quad (13.5)$$

Proof. [✓ Proof](#) Unfold the Hilbert-space local term and compare it configuration-by-configuration with the cyclic restrictions. The d^{-L} factor compensates for the d^L choices of the window coordinates in the ambient configuration parameter τ . $\square$

Lemma 13.11.8 (Positivity of the Hilbert-space Hamiltonian). [Lean](#) [Lean](#) [✓ Stmt](#) *Each Hilbert-space local term $h_{i,\text{ES}}$ is positive, hence the full Hilbert-space parent Hamiltonian $H_{N,\text{ES}}$ is positive.*

Proof. [✓ Proof](#) Apply the averaging formula (13.5): $h_{i,\text{ES}}$ is a non-negative scalar multiple of a finite sum of positive conjugates of $P_L^{\text{ES}}(A)$, hence is itself positive. Summing over i yields positivity of $H_{N,\text{ES}}$. $\square$

Lemma 13.11.9 (Hilbert-space local terms are symmetric projections). [Lean](#) [Lean](#) [✓ Stmt](#) *The Hilbert-space local projector $P_L^{\text{ES}}(A)$ is a symmetric projection, and every Hilbert-space local term $h_{i,\text{ES}}$ on the N -site chain is a symmetric projection.*

Proof. [✓ Proof](#) The local operator $P_L^{\text{ES}}(A)$ is an orthogonal projection onto $(G_L^{\text{ES}}(A))^\perp$. For $L \leq N$, restricting $h_{i,\text{ES}} v$ to a cyclic window gives $P_L^{\text{ES}}(A)$ applied to the corresponding restriction of v ; applying $h_{i,\text{ES}}$ a second time therefore applies $(P_L^{\text{ES}}(A))^2 = P_L^{\text{ES}}(A)$ on that window. The positivity result above supplies symmetry, and the case $L > N$ is the zero projection by definition. $\square$

Lemma 13.11.10 (Kernel of the Hilbert-space parent interaction). [Lean](#) [✓ Stmt](#) *The Hilbert-space local interaction $P_L^{\text{ES}}(A)$ annihilates exactly the Hilbert-space local ground space:*

$$P_L^{\text{ES}}(A)v = 0 \iff v \in G_L^{\text{ES}}(A).$$

Proof. [✓ Proof](#) Since $P_L^{\text{ES}}(A)$ is the orthogonal projector onto $(G_L^{\text{ES}}(A))^\perp$, its kernel is the original subspace $G_L^{\text{ES}}(A)$. $\square$

Lemma 13.11.11 (Local-term kernel via cyclic restrictions). [Lean](#) [Lean](#) [✓ Stmt](#) *For $L \leq N$, a Hilbert-space local term $h_{i,\text{ES}}$ annihilates a vector v iff every restriction of v to the cyclic length- L window starting at i , with the outside configuration fixed, lies in $G_L^{\text{ES}}(A)$.*

Proof. [✓ Proof](#) Restricting $h_{i,\text{ES}} v$ to the same cyclic window gives $P_L^{\text{ES}}(A)$ applied to the corresponding restriction of v . Lemma 13.11.10 identifies the vanishing of this local projector with membership in $G_L^{\text{ES}}(A)$. $\square$

Theorem 13.11.12 (Zero energy gives the cyclic local constraints). [Lean](#) [✓ Stmt](#) Assume $N > 0$ and $L \leq N$. If a vector ψ has zero energy for the finite periodic parent Hamiltonian,

$$H_N(A, L)\psi = 0,$$

then every cyclic length- L restriction of ψ lies in the local MPS ground space. Equivalently,

$$\ker H_N(A, L) \subseteq G_L^{(N)}(A).$$

Proof. [✓ Proof](#) Let e_N be the canonical ℓ^2 identification of the N -site space. Since $H_N(A, L)$ is the e_N -conjugate of $H_{N,L}^{\text{ES}}(A)$, the hypothesis gives

$$H_{N,L}^{\text{ES}}(A)(e_N^{-1}\psi) = e_N^{-1}(H_N(A, L)\psi) = 0.$$

Put $P_i = h_{i,\text{ES}}$ and $H = \sum_i P_i$. If $Hv = 0$, then

$$0 = \text{Re}\langle Hv, v \rangle = \sum_i \text{Re}\langle P_i v, v \rangle = \sum_i \|P_i v\|^2.$$

Each summand is nonnegative; hence $\|P_i v\|^2 = 0$ and therefore $P_i v = 0$ for every i . Applying this to $H_{N,L}^{\text{ES}}(A) = \sum_i h_{i,\text{ES}}$ gives the vanishing of every Hilbert-space local term. Lemma 13.11.11 identifies these equations with the conditions: for all i and τ ,

$$R_{i,\tau}\psi \in G_L(A),$$

which are exactly the cyclic-window defining equations of $G_L^{(N)}(A)$. □

Lemma 13.11.13 (Cyclic constraints give zero energy). [Lean](#) [✓ Stmt](#) Assume $N > 0$ and $L \leq N$. Then

$$G_L^{(N)}(A) \subseteq \ker H_N(A, L).$$

Proof. [✓ Proof](#) If $\psi \in G_L^{(N)}(A)$, then every local parent interaction annihilates ψ , for every site i :

$$h_i(A, L)\psi = 0.$$

Summing over all sites gives

$$H_N(A, L)\psi = \sum_i h_i(A, L)\psi = 0.$$

□

Definition 13.11.14 (Hilbert-space cyclic constraint space). [Lean](#) [✓ Stmt](#) The Hilbert-space cyclic constraint space is the transport of $G_L^{(N)}(A)$ by the canonical ℓ^2 -identification:

$$G_{L,\text{ES}}^{(N)}(A) = e_N^{-1}G_L^{(N)}(A).$$

Theorem 13.11.15 (The finite-volume kernel is the cyclic constraint space). [Lean](#) [Lean](#) [✓ Stmt](#) Assume $N > 0$ and $L \leq N$. Then

$$\ker H_N(A, L) = G_L^{(N)}(A).$$

Equivalently, after the canonical Hilbert-space identification,

$$\mathcal{K}_N^{\text{ES}}(A, L) = e_N^{-1}G_L^{(N)}(A),$$

where $\mathcal{K}_N^{\text{ES}}(A, L)$ is the zero-energy space of $H_{N,L}^{\text{ES}}(A)$.

Proof. ✓ Proof The inclusion

$$\ker H_N(A, L) \subseteq G_L^{(N)}(A)$$

is Theorem 13.11.12. For the converse, a vector in $G_L^{(N)}(A)$ satisfies all local equations $h_i(A, L)\psi = 0$. Therefore

$$H_N(A, L)\psi = \sum_i h_i(A, L)\psi = 0.$$

The Hilbert-space statement is the same equality transported by e_N^{-1} . □

Theorem 13.11.16 (Intersection property for two adjacent local kernels). Lean Lean Lean

✓ Stmt *If A is injective and $L \geq 2$, then on an $(L + 1)$ -site chain the kernels of the two adjacent Hilbert-space L -site local terms have common kernel equal to the Hilbert-space $(L + 1)$ -site MPS ground space:*

$$h_{0, \text{ES}}v = 0 \quad \text{and} \quad h_{1, \text{ES}}v = 0 \quad \Longleftrightarrow \quad v \in G_{L+1}^{\text{ES}}(A).$$

Proof. ✓ Proof Lemma 13.11.11 translates the two local kernel assumptions into the left and right L -site ground conditions. The open-chain intersection property, Theorem 13.4.4.1, then grows back the shared $(L - 1)$ -site overlap to the $(L + 1)$ -site ground space. The converse uses the forward restriction lemmas for ground-space vectors and the same kernel–restriction characterization. □

Lemma 13.11.17 (Non-overlap positivity for Hilbert-space local terms). Lean Lean Lean

Lean Lean Lean ✓ Stmt *Let $L \leq N$ and let $h_{i, \text{ES}}, h_{j, \text{ES}}$ be Hilbert-space local terms whose cyclic L -windows are site-disjoint: no site of the periodic chain has cyclic offset $< L$ from both i and j . Then the terms commute pointwise: for every vector v ,*

$$h_{i, \text{ES}}(h_{j, \text{ES}}v) = h_{j, \text{ES}}(h_{i, \text{ES}}v),$$

The same commutation identity holds for the corresponding local terms before passing to the Hilbert-space model. Their ordered cross term is non-negative:

$$0 \leq \text{Re}\langle h_{i, \text{ES}}v, h_{j, \text{ES}}v \rangle.$$

Proof. ✓ Proof In coordinates, $h_{i, \text{ES}}$ applies $P_L^{\text{ES}}(A)$ only to the i -window variables and leaves the complementary variables fixed. Site disjointness gives two replacement identities: replacing the i -window does not change the extracted j -window, and the i - and j -window replacements commute. Therefore the two embedded local operators commute. The preceding projection-geometry identity for commuting symmetric projections gives the non-negative ordered cross term. □

Lemma 13.11.18 (Quadratic form to norm bound). Lean ✓ Stmt *Let H be a positive semidefinite self-adjoint operator on a finite-dimensional complex Hilbert space, and suppose that, for all v ,*

$$\gamma \langle v, Hv \rangle \leq \langle Hv, Hv \rangle$$

i.e. $H^2 \geq \gamma H$ as a quadratic form, for some $\gamma > 0$. Then for every v in the orthogonal complement of $\ker H$,

$$\gamma \|v\| \leq \|Hv\|.$$

In particular, every positive eigenvalue of H is at least γ . The positive-semidefiniteness hypothesis is essential: without it, $H = -\text{Id}$ with $\gamma = 2$ satisfies the quadratic-form inequality vacuously but violates the norm bound.

Proof. ✓ Proof By the finite-dimensional spectral theorem, H diagonalises in an orthonormal eigenbasis with real eigenvalues λ_k . For a unit eigenvector ψ_k with eigenvalue λ_k the hypothesis gives $\gamma \lambda_k \leq \lambda_k^2$, whence $\lambda_k \leq 0$ or $\lambda_k \geq \gamma$. Since H is positive semidefinite (its quadratic form is ≥ 0), the negative case is excluded and every nonzero eigenvalue satisfies $\lambda_k \geq \gamma$. Writing $v = \sum_k c_k \psi_k$ and projecting onto $(\ker H)^\perp$ keeps only terms with $\lambda_k \geq \gamma$, so $\|Hv\|^2 = \sum_k |c_k|^2 \lambda_k^2 \geq \gamma^2 \sum_k |c_k|^2 = \gamma^2 \|v\|^2$. $\square$

13.11.1 Martingale condition for a finite family of projections

Lemma 13.11.1.1 (Parent-Hamiltonian quadratic-form reduction). Lean Lean ✓ Stmt Let $H_{N,L}^{\text{ES}}(A)$ be the Hilbert-space representative of the parent Hamiltonian. If a constant $\gamma > 0$ satisfies the uniform quadratic-form estimate

$$\gamma \operatorname{Re}\langle H_{N,L}^{\text{ES}}(A)v, v \rangle \leq \operatorname{Re}\langle H_{N,L}^{\text{ES}}(A)v, H_{N,L}^{\text{ES}}(A)v \rangle \quad (13.6)$$

for every $N \geq 2L$ and every vector v , then

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|$$

for every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$. In particular, with $L > 1$ and $\gamma = 1/(4L)$, the explicit gap-bound conclusion follows once this same quadratic-form estimate (13.6) is supplied.

Proof. ✓ Proof The Hilbert-space Hamiltonian is positive by Lemma 13.11.8, and its Hilbert-space ground space is its kernel by Theorem 13.11.3. Therefore the abstract spectral-theorem conversion of Lemma 13.11.18 applies directly. The explicit constant is positive because $L > 1$ implies $4L > 0$. $\square$

Remark 13.11.1.2 (Projection-geometry identities for row-sum arguments). Lean Lean Lean Lean Lean Lean Lean Lean Lean ✓ Stmt The projection-geometry reduction uses the following elementary Hilbert-space identities: the diagonal identity $\operatorname{Re}\langle Pv, Pv \rangle = \operatorname{Re}\langle Pv, v \rangle$ for symmetric projections; nonnegativity of this quadratic form; nonnegativity of $\operatorname{Re}\langle Pv, Qv \rangle$ for commuting symmetric projections P, Q ; the Cauchy–Schwarz conversion from $\|PQv\| \leq c\|Pv\|$ to $\operatorname{Re}\langle Pv, Qv \rangle \geq -c\operatorname{Re}\langle Pv, v \rangle$; the symmetric variant which, for $c \geq 0$, converts the operator-product bound $\|PQv\| \leq c\|Qv\|$ to the anticommutator estimate $\operatorname{Re}\langle Pv, Qv \rangle + \operatorname{Re}\langle Qv, Pv \rangle \geq -c(\operatorname{Re}\langle Pv, v \rangle + \operatorname{Re}\langle Qv, v \rangle)$; finite-sum expansions of $\operatorname{Re}\langle (\sum_i P_i)v, v \rangle$ and $\operatorname{Re}\langle (\sum_i P_i)v, (\sum_i P_i)v \rangle$; the conclusion that a vector annihilated by a finite sum of symmetric projections is annihilated by each projection; the diagonal/off-diagonal split of the ordered double sum; the aggregate off-diagonal estimate obtained from row sums $\sum_{j \neq i} c_{ij} \leq 1$; and the corresponding aggregate estimate for the anticommutator martingale condition.

Lemma 13.11.1.3 (Ordered projection row-sum quadratic-form reduction). Lean ✓ Stmt Let P_i be a finite family of symmetric projections and let $H = \sum_i P_i$. If the ordered off-diagonal forms obey

$$\operatorname{Re}\langle P_i v, P_j v \rangle \geq -(1 - \gamma)c_{ij} \operatorname{Re}\langle P_i v, v \rangle \quad (i \neq j),$$

with row sums $\sum_{j \neq i} c_{ij} \leq 1$ and $\gamma \leq 1$, then $H^2 \geq \gamma H$ as a quadratic form.

Proof. ✓ Proof Expand $\operatorname{Re}\langle Hv, Hv \rangle$ as an ordered double sum. Symmetric idempotence identifies each diagonal term with $\operatorname{Re}\langle P_i v, v \rangle$, while the row-summable ordered cross-term bounds control the off-diagonal contribution by $-(1 - \gamma) \sum_i \operatorname{Re}\langle P_i v, v \rangle$. $\square$

Lemma 13.11.1.4 (Projection anticommutator row-sum reduction). [Lean](#) [✓ Stmt](#) Let P_i be a finite family of symmetric projections and let $H = \sum_i P_i$. Suppose the anticommutator forms obey

$$\operatorname{Re}\langle P_i v, P_j v \rangle + \operatorname{Re}\langle P_j v, P_i v \rangle \geq -(1 - \gamma) c_{ij} (\operatorname{Re}\langle P_i v, v \rangle + \operatorname{Re}\langle P_j v, v \rangle) \quad (i \neq j),$$

with row sums $\sum_{j \neq i} c_{ij} \leq 1$, column sums $\sum_{i \neq j} c_{ij} \leq 1$, and $\gamma \leq 1$. Then $H^2 \geq \gamma H$ as a quadratic form.

Proof. [✓ Proof](#) Expand $\operatorname{Re}\langle H v, H v \rangle$ as an ordered double sum. The anticommutator bound controls the sum of each ordered term with its reversed term. The row and column estimates bound the coefficient-weighted diagonal contribution by twice $\sum_i \operatorname{Re}\langle P_i v, v \rangle$; after dividing by two, the same quadratic-form estimate follows. $\square$

Lemma 13.11.1.5 (Finite-overlap projection anticommutator reduction). [Lean](#) [✓ Stmt](#) Let P_i be a finite family of symmetric projections and let $\gamma \leq 1$. Suppose an interaction predicate $i \sim j$ has at most m off-diagonal neighbours in each row and in each column, where $m > 0$. If noninteracting pairs have non-negative anticommutator form and interacting pairs satisfy

$$\operatorname{Re}\langle P_i v, P_j v \rangle + \operatorname{Re}\langle P_j v, P_i v \rangle \geq -(1 - \gamma) \frac{1}{m} (\operatorname{Re}\langle P_i v, v \rangle + \operatorname{Re}\langle P_j v, v \rangle),$$

then $H = \sum_i P_i$ satisfies $H^2 \geq \gamma H$ as a quadratic form.

Proof. [✓ Proof](#) Choose $c_{ij} = 1/m$ on interacting pairs and $c_{ij} = 0$ otherwise. The row and column cardinality bounds give the corresponding coefficient sums, and the noninteracting anticommutator non-negativity supplies the zero-coefficient estimates. Apply Lemma 13.11.1.4. $\square$

Lemma 13.11.1.6 (Finite-overlap projection row-sum reduction). [Lean](#) [✓ Stmt](#) Let P_i be a finite family of symmetric projections and let $\gamma \leq 1$. Suppose an interaction predicate $i \sim j$ has at most m off-diagonal neighbours in each row for some positive integer m . If noninteracting ordered cross terms are non-negative and, on each interacting pair,

$$\operatorname{Re}\langle P_i v, P_j v \rangle \geq -(1 - \gamma) \frac{1}{m} \operatorname{Re}\langle P_i v, v \rangle,$$

then $H = \sum_i P_i$ satisfies $H^2 \geq \gamma H$ as a quadratic form.

Proof. [✓ Proof](#) Choose $c_{ij} = 1/m$ on interacting pairs and $c_{ij} = 0$ otherwise. The row cardinality bound gives $\sum_{j \neq i} c_{ij} \leq 1$; noninteracting nonnegativity supplies the zero-coefficient cross-term estimates. Apply Lemma 13.11.1.3. $\square$

Lemma 13.11.1.7 (Finite-overlap projection norm-compression reduction). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let P_i be a finite family of symmetric projections and let $\gamma \leq 1$. Suppose an interaction predicate has at most $m > 0$ off-diagonal neighbours in each row, noninteracting ordered cross terms are non-negative, and interacting pairs satisfy the norm-compression estimate

$$\|P_i P_j v\| \leq (1 - \gamma) \frac{1}{m} \|P_i v\|. \quad (13.7)$$

Then $H = \sum_i P_i$ satisfies $H^2 \geq \gamma H$ as a quadratic form. The same conclusion holds from a separate compression coefficient η whenever $\eta \leq (1 - \gamma)/m$. In particular, if $0 \leq \eta$ and $\eta m < 1$, then

$$H^2 \geq (1 - \eta m) H$$

and $1 - \eta m > 0$.

Proof. ✓ Proof Cauchy–Schwarz and symmetric idempotence turn the norm-compression estimate into $\text{Re}\langle P_i v, P_j v \rangle \geq -(1 - \gamma)m^{-1} \text{Re}\langle P_i v, v \rangle$. If a separate coefficient η is used, the assumption $\eta \leq (1 - \gamma)/m$ first weakens that estimate to (13.7). The finite-overlap row-sum reduction then applies. For the strict form take $\gamma = 1 - \eta m$. $\square$

Lemma 13.11.1.8 (Quadratic-form estimate from the martingale condition). Lean ✓ Stmt For a fixed chain length N , suppose the Hilbert-space local terms $h_{i,\text{ES}}$ satisfy the ordered row-sum estimate

$$\text{Re}\langle h_{i,\text{ES}} v, h_{j,\text{ES}} v \rangle \geq -(1 - \gamma)c_{ij} \text{Re}\langle h_{i,\text{ES}} v, v \rangle \quad (i \neq j), \quad (13.8)$$

with $\sum_{j \neq i} c_{ij} \leq 1$ and $\gamma \leq 1$. Then the Hilbert-space parent Hamiltonian satisfies

$$\gamma \text{Re}\langle H_{N,L}^{\text{ES}} v, v \rangle \leq \text{Re}\langle H_{N,L}^{\text{ES}} v, H_{N,L}^{\text{ES}} v \rangle$$

for every vector v .

Proof. ✓ Proof Lemma 13.11.9 supplies the symmetric-projection hypothesis for the family $P_i = h_{i,\text{ES}}$. Apply Lemma 13.11.1.3 to this family and use $H_{N,L}^{\text{ES}} = \sum_i h_{i,\text{ES}}$. $\square$

Lemma 13.11.1.9 (Parent-Hamiltonian gap from ordered local row bounds). Lean ✓ Stmt Assume uniformly for every $N \geq 2L$ that the Hilbert-space local terms satisfy the ordered row-sum estimate (13.8) with $\gamma = 1/(4L)$. If $L > 1$, then $1/(4L) > 0$ and, for every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$,

$$\frac{1}{4L} \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. ✓ Proof The fixed-chain row-sum reduction supplies the quadratic-form estimate with $\gamma = 1/(4L)$ for every admissible N . The positivity and kernel identification of the Hilbert-space Hamiltonian then allow Lemma 13.11.1.1 to convert this quadratic-form estimate into the norm lower bound on the orthogonal complement of the ground space. $\square$

Lemma 13.11.1.10 (Parent-Hamiltonian gap from the anticommutator martingale condition). Lean Lean ✓ Stmt Assume uniformly for every $N \geq 2L$ that the Hilbert-space local terms satisfy the anticommutator martingale estimate

$$\text{Re}\langle h_{i,\text{ES}} v, h_{j,\text{ES}} v \rangle + \text{Re}\langle h_{j,\text{ES}} v, h_{i,\text{ES}} v \rangle \geq -c_{ij} \left(1 - \frac{1}{4L}\right) (\text{Re}\langle h_{i,\text{ES}} v, v \rangle + \text{Re}\langle h_{j,\text{ES}} v, v \rangle)$$

for $i \neq j$, with row and column coefficient sums bounded by one. If $L > 1$, then $1/(4L) > 0$ and, for every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$,

$$\frac{1}{4L} \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. ✓ Proof The fixed-chain anticommutator row-sum reduction supplies the quadratic-form estimate with $\gamma = 1/(4L)$ for every admissible N . The positivity and kernel identification of the Hilbert-space Hamiltonian then convert this quadratic-form estimate into the norm lower bound on the orthogonal complement of the ground space. $\square$

Lemma 13.11.1.11 (Parent-Hamiltonian finite-overlap anticommutator reduction). Lean Lean ✓ Stmt Fix $L > 1$ and set $m = 2(L - 1)$. Suppose uniformly for every $N \geq 2L$ that the overlap relation on local windows has at most m off-diagonal neighbours in each row and in each column. If non-overlapping local terms have non-negative anticommutator form, and overlapping local terms satisfy

$$\text{Re}\langle h_{i,\text{ES}} v, h_{j,\text{ES}} v \rangle + \text{Re}\langle h_{j,\text{ES}} v, h_{i,\text{ES}} v \rangle \geq -\left(1 - \frac{1}{4L}\right) \frac{1}{m} (\text{Re}\langle h_{i,\text{ES}} v, v \rangle + \text{Re}\langle h_{j,\text{ES}} v, v \rangle),$$

then the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. ✓ Proof Apply the finite-overlap anticommutator projection reduction to the Hilbert-space local terms. The resulting quadratic-form estimate has $\gamma = 1/(4L)$, and Lemma 13.11.1.1 converts it to the norm lower bound. $\square$

Lemma 13.11.1.12 (Parent-Hamiltonian finite-overlap martingale quadratic reduction). Lean

✓ Stmt For a fixed chain length N , let m be a positive integer and let $\gamma \leq 1$. Suppose an overlap predicate on local windows has at most m overlapping off-diagonal windows in each row. If non-overlapping local terms have non-negative ordered cross terms, and overlapping local terms satisfy the one-sided ordered cross-term estimate

$$\text{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle \geq -(1-\gamma) \frac{1}{m} \text{Re}\langle h_{i,\text{ES}}v, v \rangle,$$

and the Hilbert-space local terms are symmetric projections, then the Hilbert-space Hamiltonian satisfies $H_{N,L}^{\text{ES}^2} \geq \gamma H_{N,L}^{\text{ES}}$ as a quadratic form.

Proof. ✓ Proof Apply the finite-overlap projection reduction with $P_i = h_{i,\text{ES}}$. $\square$

Lemma 13.11.1.13 (Finite-overlap norm-compression reduction). Lean Lean ✓ Stmt For a fixed chain length N , let $m > 0$ and $\gamma \leq 1$. Suppose an overlap predicate on local windows has at most m overlapping off-diagonal windows in each row, non-overlapping local terms have non-negative ordered cross terms, and overlapping local terms satisfy

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq (1-\gamma) \frac{1}{m} \|h_{i,\text{ES}}v\|.$$

Then $H_{N,L}^{\text{ES}^2} \geq \gamma H_{N,L}^{\text{ES}}$ as a quadratic form. The same conclusion holds from a separate coefficient η satisfying $\eta \leq (1-\gamma)/m$.

Proof. ✓ Proof Apply the finite-overlap norm-compression projection reduction to the family $P_i = h_{i,\text{ES}}$. If a separate coefficient η is used, first weaken the compression estimate using $\eta \leq (1-\gamma)/m$. $\square$

13.11.2 Cyclic-window overlap estimates

Definition 13.11.2.1 (Cyclic-window support). Lean ✓ Stmt For a length- L cyclic window on $\{0, \dots, N-1\}$ starting at i , its support is

$$S_i = \{i, i+1, \dots, i+L-1\} \pmod{N},$$

with repeated visits counted once.

Lemma 13.11.2.2 (Cyclic-window support offsets). Lean ✓ Stmt If $L \leq N$, then a site k belongs to the cyclic support S_i exactly when its cyclic offset from i is below L :

$$k \in S_i \iff (k-i) \bmod N < L.$$

Proof. ✓ Proof Write $k = i + r \pmod{N}$ for a representative offset r . Membership in S_i gives an offset $r < L$, and conversely an offset below L gives the corresponding point of the support. $\square$

Definition 13.11.2.3 (Cyclic-window overlap). Lean ✓ Stmt Two length- L cyclic windows overlap, written $i \sim_L j$, when their supports meet: $S_i \cap S_j \neq \emptyset$.

Lemma 13.11.2.4 (Cyclic-window overlap symmetry). [Lean](#) [✓ Stmt](#) For length- L cyclic windows, $i \sim_L j$ if and only if $j \sim_L i$.

Proof. [✓ Proof](#) Both assertions say that the two cyclic supports have a common site. $\square$

Lemma 13.11.2.5 (Concrete non-overlap positivity for cyclic supports). [Lean](#) [Lean](#) [✓ Stmt](#) If $L \leq N$ and the cyclic supports S_i and S_j do not meet, then the Hilbert-space local terms have non-negative ordered cross term:

$$0 \leq \operatorname{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle.$$

Proof. [✓ Proof](#) The support-offset characterization converts failure of $S_i \cap S_j \neq \emptyset$ into site-disjointness of the two cyclic windows. The disjoint-window positivity lemma then applies. $\square$

Lemma 13.11.2.6 (Cyclic-window overlap row cardinality). [Lean](#) [✓ Stmt](#) If $2L \leq N$ and $L > 1$, then for every $i \in \{0, \dots, N-1\}$,

$$\#\{j \neq i : i \sim_L j\} \leq 2(L-1).$$

Proof. [✓ Proof](#) If S_i and S_j meet, choose offsets $a, b < L$ with $i + a \equiv j + b \pmod{N}$. Since $2L \leq N$, the offset from i to j is either the clockwise distance $a - b \in \{1, \dots, L-1\}$ or the counterclockwise distance $b - a \in \{1, \dots, L-1\}$; the zero distance is excluded by $j \neq i$. Thus every overlapping off-diagonal start lies among the $L-1$ clockwise starts or the $L-1$ counterclockwise starts. $\square$

Lemma 13.11.2.7 (Parent-Hamiltonian gap from a finite-overlap martingale estimate). [Lean](#) [✓ Stmt](#) Fix $L > 1$ and set $m = 2(L-1)$. Suppose uniformly for every $N \geq 2L$ that the hypotheses of the fixed-chain finite-overlap martingale reduction hold with $\gamma = 1/(4L)$ and this value of m . Then $1/(4L) > 0$, and the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. [✓ Proof](#) The fixed-chain finite-overlap reduction gives the required quadratic-form estimate for every admissible N . Lemma 13.11.1.1 converts that estimate to the norm lower bound. $\square$

Lemma 13.11.2.8 (Parent-Hamiltonian gap from a cyclic-window martingale estimate). [Lean](#) [✓ Stmt](#) Fix $L > 1$ and let $i \sim_L j$ mean that the cyclic supports of the length- L windows at i and j meet. Suppose uniformly for every $N \geq 2L$ that overlapping off-diagonal terms satisfy

$$\operatorname{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle \geq -(1 - \frac{1}{4L}) \frac{1}{2(L-1)} \operatorname{Re}\langle h_{i,\text{ES}}v, v \rangle. \quad (13.9)$$

Then $1/(4L) > 0$, and the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. [✓ Proof](#) Lemma 13.11.2.6 gives the row-cardinality bound $m = 2(L-1)$ for the overlap relation among length- L local windows, and Lemma 13.11.9 supplies the symmetric projection structure of each Hilbert-space local term. If two cyclic supports do not meet, the corresponding windows are site-disjoint, so Lemma 13.11.2.5 gives the required non-negative ordered cross term. Apply Lemma 13.11.2.7 with these facts and the overlapping-window estimate (13.9). $\square$

Lemma 13.11.2.9 (Parent-Hamiltonian gap from an overlapping cyclic martingale estimate).

[Lean](#) [✓ Stmt](#) Fix $L > 1$ and let $i \sim_L j$ mean that the cyclic supports of the length- L windows at i and j meet. Suppose uniformly for every $N \geq 2L$ that overlapping off-diagonal terms satisfy

$$\operatorname{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle \geq -\left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \operatorname{Re}\langle h_{i,\text{ES}}v, v \rangle. \quad (13.10)$$

Then the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. [✓ Proof](#) Apply Lemma 13.11.2.8. Its non-overlap positivity and finite-family hypotheses are already internal to the preceding reduction; the overlapping estimate (13.10) supplies its remaining ordered cross-term condition. $\square$

Lemma 13.11.2.10 (Parent-Hamiltonian gap from a cyclic anticommutator estimate). [Lean](#)

[✓ Stmt](#) Fix $L > 1$ and let $i \sim_L j$ mean that the cyclic supports of the length- L windows at i and j meet. Suppose uniformly for every $N \geq 2L$ that overlapping off-diagonal terms satisfy

$$\operatorname{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle + \operatorname{Re}\langle h_{j,\text{ES}}v, h_{i,\text{ES}}v \rangle \geq -\left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} (\operatorname{Re}\langle h_{i,\text{ES}}v, v \rangle + \operatorname{Re}\langle h_{j,\text{ES}}v, v \rangle).$$

Then the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. [✓ Proof](#) The cyclic overlap relation is symmetric, and Lemma 13.11.2.6 gives at most $2(L-1)$ overlapping off-diagonal windows in each row and column. If two cyclic supports do not meet, Lemma 13.11.2.5 gives non-negative ordered cross terms in both orders. Apply the finite-overlap anticommutator reduction. $\square$

Lemma 13.11.2.11 (Parent-Hamiltonian gap from sufficient cyclic norm compression). [Lean](#)

[✓ Stmt](#) Fix $L > 1$. Suppose uniformly for every $N \geq 2L$ and every overlapping off-diagonal pair that

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \|h_{i,\text{ES}}v\|.$$

Then the explicit norm lower bound with constant $1/(4L)$ follows for vectors in $(\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$.

Proof. [✓ Proof](#) For an overlapping pair, Cauchy–Schwarz and $h_{i,\text{ES}}^2 = h_{i,\text{ES}}$ give

$$\begin{aligned} \operatorname{Re}\langle h_{i,\text{ES}}v, h_{j,\text{ES}}v \rangle &= \operatorname{Re}\langle h_{i,\text{ES}}v, h_{i,\text{ES}}h_{j,\text{ES}}v \rangle \\ &\geq -\|h_{i,\text{ES}}v\| \|h_{i,\text{ES}}h_{j,\text{ES}}v\| \\ &\geq -\left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \|h_{i,\text{ES}}v\|^2 \\ &= -\left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \operatorname{Re}\langle h_{i,\text{ES}}v, v \rangle. \end{aligned}$$

This is the ordered lower bound used by the preceding overlapping-cyclic martingale reduction. $\square$

Lemma 13.11.2.12 (Parent-Hamiltonian gap from a cyclic overlap compression constant).

[Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Fix $L > 1$, and let $m = 2(L-1)$. Suppose uniformly for every $N \geq 2L$ and every overlapping off-diagonal pair that

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{i,\text{ES}}v\|.$$

If $\gamma > 0$, $\gamma \leq 1$, and $\eta \leq (1 - \gamma)/m$, then $0 < \gamma$ and, for every admissible N and every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^{\perp}$,

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

In particular, if $\eta \geq 0$ and $\eta m < 1$, then $0 < 1 - \eta m$ and, for every admissible N and every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^{\perp}$,

$$(1 - \eta m) \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

This strict compression form is a norm-compression sufficient condition, not the principal-angle estimate itself. It is a sufficient strengthening of the anticommutator martingale estimate in Cirac–Perez–Garcia–Schuch–Verstraete [CPGSV21, Section IV.C]. It does not prove the remaining MPS-specific estimate; it isolates the finite-row martingale argument once such a strengthening is supplied.

Proof. ✓ Proof The overlap-cardinality lemma gives at most $m = 2(L-1)$ overlapping off-diagonal neighbours in each row, while the non-overlap positivity lemma handles disjoint supports. The finite-overlap norm-compression reduction gives the quadratic-form estimate with the chosen γ , and the positive quadratic-form criterion converts it to the norm lower bound. The strict form is the same statement with $\gamma = 1 - \eta m$. $\square$

Lemma 13.11.2.13 (Parent-Hamiltonian gap from a cyclic operator-product norm constant).

Lean Lean ✓ Stmt Fix $L > 1$, and let $m = 2(L-1)$. Suppose uniformly for every $N \geq 2L$ and every overlapping off-diagonal pair that

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{j,\text{ES}}v\|,$$

that is, a bound $\|h_{i,\text{ES}}h_{j,\text{ES}}\| \leq \eta$ on the operator-product norm. If $\gamma > 0$, $\gamma \leq 1$, and $\eta \leq (1 - \gamma)/m$, then $0 < \gamma$ and, for every admissible N and every $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^{\perp}$,

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

In particular, if $\eta \geq 0$ and $\eta m < 1$, then $0 < 1 - \eta m$ and the same norm bound holds with $\gamma = 1 - \eta m$.

This is the symmetric (operator-product) counterpart of Lemma 13.11.2.12, obtained from the anticommutator conversion recorded in Remark 13.11.1.2. Like that lemma it is a conditional reduction: the excitation projections satisfy $\|h_{i,\text{ES}}h_{j,\text{ES}}\| = 1$ whenever the overlapping ranges share a common nonzero vector, so the hypothesis is unsatisfiable for $\eta < 1$ in that case. It isolates the finite-row martingale argument once such a bound is supplied; the gap with the cited principal-angle estimates is documented in the paper-gap note `docs/paper-gaps/cpgsv21_martingale_overlap.tex`.

Proof. ✓ Proof The overlap-cardinality lemma gives at most $m = 2(L-1)$ overlapping off-diagonal neighbours in each row, while the non-overlap positivity lemma handles disjoint supports. The symmetric anticommutator conversion of Remark 13.11.1.2 turns the operator-product bound into the per-pair anticommutator estimate with coefficient $(1 - \gamma)/m$, using the hypothesis $\eta \leq (1 - \gamma)/m$; the anticommutator finite-overlap reduction then gives the quadratic-form estimate with the chosen γ , and the positive quadratic-form criterion converts it to the norm lower bound. The strict form is the same statement with $\gamma = 1 - \eta m$, so the coefficient is then η . $\square$

13.11.3 Martingale gap consequences

Theorem 13.11.3.1 (Martingale criterion for spectral gap). Not ready Consider a frustration-free Hamiltonian

$$H = \sum_{i=1}^n h_i$$

with projector interaction terms ($h_i^2 = h_i$). Suppose that for every unordered pair $\{i, j\}$ of overlapping terms there exist constants $c_{\{i, j\}} \geq 0$ and $\gamma > 0$ satisfying

$$h_i h_j + h_j h_i \geq -c_{\{i, j\}}(1 - \gamma)(h_i + h_j), \quad (13.11)$$

and suppose further that the coefficients attached to the terms overlapping h_i satisfy

$$\sum_{\substack{j \neq i \\ \{i, j\} \text{ overlapping}}} c_{\{i, j\}} \leq 1$$

for every i . Then the smallest nonzero eigenvalue of H satisfies $\lambda_{\min}^+(H) \geq \gamma$.

Proof. Not ready The relevant geometry is a family of translated length- L projector windows on the ring; two terms h_i and h_j interact only when their supports overlap, equivalently when the cyclic distance $d_N(i, j)$ is smaller than L . From $h_i^2 = h_i$,

$$H^2 = \sum_i h_i + \sum_{\substack{i < j \\ \text{overlapping}}} (h_i h_j + h_j h_i) + \sum_{\substack{i < j \\ \text{non-overlapping}}} (h_i h_j + h_j h_i).$$

Non-overlapping projectors commute ($h_i h_j = h_j h_i$), so their product $h_i h_j$ is itself a projector and hence PSD; thus the last sum is ≥ 0 . For the overlapping sum, the martingale condition (13.11) gives $h_i h_j + h_j h_i \geq -c_{\{i, j\}}(1 - \gamma)(h_i + h_j)$. Summing over unordered pairs $\{i, j\}$:

$$\sum_{\{i, j\}, \text{overlap}} (h_i h_j + h_j h_i) \geq -(1 - \gamma) \sum_{\{i, j\}, \text{overlap}} c_{\{i, j\}} (h_i + h_j).$$

Collecting terms for each h_i :

$$\sum_{\{i, j\}, \text{overlap}} c_{\{i, j\}} (h_i + h_j) = \sum_i \left(\sum_{\substack{j \neq i \\ \text{overlap}}} c_{\{i, j\}} \right) h_i \leq H,$$

where the last inequality uses the row-sum hypothesis $\sum_{\substack{j \neq i \\ \{i, j\} \text{ overlapping}}} c_{\{i, j\}} \leq 1$. Combining,

$H^2 \geq H - (1 - \gamma)H = \gamma H$. For any eigenvector with $H\psi = \lambda\psi$ and $\lambda > 0$, this gives $\lambda^2 \geq \gamma\lambda$, hence $\lambda \geq \gamma$ [KL18]. $\square$

Lemma 13.11.3.2 (MPS anticommutator input for the martingale condition). Not ready Let A be injective and fix an interaction range $L \geq 2$ as in Definition 13.3.3. Let $h_L(A) := \Pi_{G_L(A)^\perp}$ be the L -site parent interaction, and for each chain length and site i let h_i denote its translate as in Definition 13.3.11. For every pair of overlapping terms h_i and h_j with $0 < d_N(i, j) < L$, where $d_N(i, j) := \min(|i - j|, N - |i - j|)$ is the cyclic distance on the ring (i.e. whose L -site supports share at least one site under periodic boundary conditions), the source martingale argument asks for constants $c_{\{i, j\}}$ and $\gamma > 0$, independent of N , for which the martingale condition of

Theorem 13.11.3.1 holds. Cirac–Perez-Garcia–Schuch–Verstraete [CPGSV21, Section IV.C] use the martingale estimate

$$h_i h_j + h_j h_i \geq -c_{ij}(1 - \gamma)(h_i + h_j)$$

and interpret it as a lower bound on the smallest non-zero principal angle between the local ground spaces $\ker h_i$ and $\ker h_j$. In the Hilbert-space formulation above, the source condition is the corresponding quadratic-form inequality

$$\operatorname{Re}\langle h_i v, h_j v \rangle + \operatorname{Re}\langle h_j v, h_i v \rangle \geq -c_{ij}(1 - \gamma)(\operatorname{Re}\langle h_i v, v \rangle + \operatorname{Re}\langle h_j v, v \rangle),$$

with row-summable coefficients. Lemma 13.11.1.10 gives the implication from this anticommutator estimate to the spectral gap. A directed norm-compression estimate, independent of N , such as

$$\|h_i h_j v\| \leq \eta \|h_i v\|, \quad \eta 2(L - 1) < 1,$$

for every overlapping off-diagonal pair is a sufficient stronger hypothesis; Theorem 13.11.3.5 records that consequence. (When $L = 1$ the interaction terms have disjoint supports, so there are no overlapping pairs and the spectral gap follows directly without the martingale argument.)

Proof. **Not ready** The cited martingale paragraph compares the anticommutator estimate with principal angles between the local ground spaces. If $q_i := I - h_i$ and $q_j := I - h_j$ denote the corresponding ground-space projectors, the Kastoryano–Lucia estimate cited there concerns

$$\delta(\ell) = \|q_i q_j - P\|,$$

where P projects onto $\ker(h_i + h_j) = \ker h_i \cap \ker h_j$ and ℓ is the overlap length. What remains is to derive the anticommutator estimate above, with its blocking convention and constants, for the cyclic-window local terms. The intersection property identifies $\ker h_i \cap \ker h_j$ with the parent ground space on the union of the two windows, but it does not by itself supply this numerical inequality. The directed excitation-projection estimate

$$\|h_i h_j v\| \leq \eta \|h_i v\|, \quad \eta 2(L - 1) < 1.$$

remains useful as a sufficient strengthening of the source condition.

Row-sum bound. Since $L \geq 2$, each site i overlaps with at most $2(L - 1)$ neighbours (those j with $0 < d_N(i, j) < L$). Set $c_{\{i, j\}} := \frac{1}{2(L-1)}$ for every unordered overlapping pair and $c_{\{i, j\}} := 0$ otherwise. These constants depend only on the cyclic distance $d_N(i, j)$. Then $\sum_{\substack{j \neq i \\ \{i, j\} \text{ overlapping}}} c_{\{i, j\}} \leq 2(L - 1) \cdot \frac{1}{2(L-1)} = 1$, satisfying the row-sum hypothesis of Theorem 13.11.3.1.

Anticommutator constant. It remains to compare the cited principal-angle estimates with the cyclic-window anticommutator statement

$$h_i h_j + h_j h_i \geq -c_{\{i, j\}}(1 - \gamma)(h_i + h_j).$$

If this estimate is obtained with the displayed row-summable coefficients, Lemma 13.11.1.10 gives the gap. A sufficient stronger route is the norm-compression statement

$$\|h_i h_j v\| \leq \eta \|h_i v\|, \quad \eta 2(L - 1) < 1.$$

If such an η is obtained, Theorem 13.11.3.5 gives the gap constant $1 - \eta 2(L - 1)$. The frequently displayed fixed-coefficient compression form is the specialization with

$$\eta = \left(1 - \frac{1}{4L}\right) \frac{1}{2(L - 1)}, \quad 1 - \eta 2(L - 1) = \frac{1}{4L}.$$

In the corresponding anticommutator form the martingale condition is

$$h_i h_j + h_j h_i \geq - \left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} (h_i + h_j)$$

for every overlapping off-diagonal pair, and a sufficient one-sided strengthening is

$$\|h_i h_j v\| \leq \left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \|h_i v\|.$$

A qualitative statement that the relevant smallest non-zero principal angles are positive is not by itself enough: the martingale reduction needs a numerical anticommutator bound with row-summable coefficients, or a stronger norm-compression bound with $\eta 2(L-1) < 1$. Thus the remaining comparison is whether the principal-angle estimates cited in [FNW92, Nac96, KL18] supply such a cyclic-window estimate, with constants independent of N , under the convention used here. $\square$

Theorem 13.11.3.3 (Gap bound from cyclic overlap compression). Lean ✓ Stmt *Let A be a tensor and let $L > 1$. Suppose that, for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,*

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)} \|h_{i,\text{ES}}v\|.$$

Then the parent Hamiltonian has the explicit gap constant

$$\gamma_L := \frac{1}{4L} > 0.$$

Concretely, if $N \geq 2L$ and $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^{\perp}$, then

$$\gamma_L \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. ✓ Proof The local averaging, positivity, and symmetric-projection statements are now established in Lemmas 13.11.7, 13.11.8, and 13.11.9. Lemma 13.11.1.1 reduces the present theorem to the remaining MPS-specific task: deriving the uniform quadratic-form estimate. Lemma 13.11.1.9 establishes the finite-sum algebra from ordered cross-term row bounds, while Lemma 13.11.1.10 records the source anticommutator martingale condition. Lemmas 13.11.1.12 and 13.11.2.7 state the common finite-overlap ordered-cross-term specialization with $m = 2(L-1)$. Lemma 13.11.1.17 supplies the non-overlap positivity statement for site-disjoint windows, and Lemma 13.11.2.5 translates this to the concrete support-disjoint cyclic windows. Lemma 13.11.2.6 gives the finite-overlap row-cardinality bound. Lemma 13.11.2.8 combines these results with the finite-overlap reduction, so the only additional input is the ordered anticommutator or cross-term estimate for overlapping cyclic windows. Lemma 13.11.2.9 states the same overlap-only formulation, and Lemma 13.11.2.11 states a sufficient norm-compression formulation. Equivalently, this is the constant-compression theorem below with

$$\eta = \left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)}.$$

Since

$$1 - \eta 2(L-1) = \frac{1}{4L},$$

the displayed hypothesis gives the stated lower bound with $\gamma_L = 1/(4L)$. $\square$

Theorem 13.11.3.4 (Gap bound from a cyclic anticommutator estimate). [Lean](#) [✓ Stmt](#) Let $L > 1$. Suppose that for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,

$$h_{i,\text{ES}}h_{j,\text{ES}} + h_{j,\text{ES}}h_{i,\text{ES}} \geq -\left(1 - \frac{1}{4L}\right) \frac{1}{2(L-1)}(h_{i,\text{ES}} + h_{j,\text{ES}})$$

in quadratic form. Then the parent Hamiltonian has the explicit gap constant

$$\gamma_L := \frac{1}{4L} > 0.$$

Concretely, if $N \geq 2L$ and $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$, then

$$\gamma_L \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. [✓ Proof](#) This is Lemma 13.11.2.10 stated as the public conditional gap theorem, with the anticommutator inequality retained as an explicit hypothesis. $\square$

Theorem 13.11.3.5 (Gap bound from a strict cyclic compression constant). [Lean](#) [✓ Stmt](#) Let $L > 1$. Put $m = 2(L-1)$. Suppose that there is a constant $\eta \geq 0$ such that $\eta m < 1$, and suppose that, for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{i,\text{ES}}v\|.$$

Then the parent Hamiltonian has the gap constant $1 - \eta m > 0$. Concretely, if $N \geq 2L$ and $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$, then

$$(1 - \eta m) \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. [✓ Proof](#) Lemma 13.11.2.12, with $\gamma = 1 - \eta m$, applies to the displayed hypothesis

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{i,\text{ES}}v\|.$$

It gives $0 < 1 - \eta m$ and the stated norm lower bound. The preceding fixed-coefficient theorem is the specialization $\eta = (1 - 1/(4L))/(2(L-1))$. $\square$

Theorem 13.11.3.6 (Gap bound from a cyclic operator-product norm constant). [Lean](#) [✓ Stmt](#) Let $L > 1$. Put $m = 2(L-1)$. Suppose there is a constant $\eta \geq 0$ with $\eta m < 1$ and that, for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{j,\text{ES}}v\|,$$

that is, a bound $\|h_{i,\text{ES}}h_{j,\text{ES}}\| \leq \eta$ on the operator-product norm rather than the directional bound of Theorem 13.11.3.5. Then the parent Hamiltonian has the gap constant $1 - \eta m > 0$. Concretely, if $N \geq 2L$ and $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$, then

$$(1 - \eta m) \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

This is a conditional reduction. The local terms $h_{i,\text{ES}} = p_i$ are the excitation projections. If their overlapping ranges have a common nonzero vector, then $\|h_{i,\text{ES}}h_{j,\text{ES}}\| = 1$, so the hypothesis is unsatisfiable for $\eta < 1$ in that case. In degenerate cases, such as a zero excitation projection, the operator-product norm can be smaller. In any case this is not the quantity controlled by the cited principal-angle estimates. The comparison is documented in `docs/paper-gaps/cpgsv21_martingale_overlap.tex`.

Proof. [✓ Proof](#) Lemma 13.11.2.13, with $\gamma = 1 - \eta m$, applies to the displayed operator-product hypothesis and gives $0 < 1 - \eta m$ together with the stated norm lower bound. $\square$

Theorem 13.11.3.7 (Conditional spectral gap for MPS parent Hamiltonians). [Lean](#) [Lean](#)
[✓ Stmt](#) For a tensor A and a range $L > 1$, assume either the cyclic-window anticommutator estimate of Theorem 13.11.3.4 or the stronger overlapping cyclic-window norm-compression estimate of Theorem 13.11.3.3. Then there exists $\gamma > 0$ such that for every $N \geq 2L$ and every vector $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$ one has

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

By Lemma 13.11.18, this implies the usual uniform lower bound on every nonzero eigenvalue of the parent Hamiltonian.

Proof. [✓ Proof](#) This follows immediately from the corresponding explicit theorem by taking γ to be the constant produced there. $\square$

Theorem 13.11.3.8 (Conditional spectral gap from a strict cyclic compression constant). [Lean](#) [✓ Stmt](#) For a tensor A and a range $L > 1$, suppose there is a constant $\eta \geq 0$ such that $\eta 2(L-1) < 1$ and, for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{i,\text{ES}}v\|.$$

Then there exists $\gamma > 0$ such that, for every $N \geq 2L$ and every vector $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$,

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

Proof. [✓ Proof](#) This follows from Theorem 13.11.3.5 by taking $\gamma = 1 - \eta 2(L-1)$. $\square$

Theorem 13.11.3.9 (Conditional spectral gap from an operator-product norm constant). [Lean](#)
[✓ Stmt](#) For a tensor A and a range $L > 1$, suppose there is a constant $\eta \geq 0$ such that $\eta 2(L-1) < 1$ and, for every $N \geq 2L$ and every overlapping off-diagonal pair of cyclic length- L windows,

$$\|h_{i,\text{ES}}(h_{j,\text{ES}}v)\| \leq \eta \|h_{j,\text{ES}}v\|.$$

Then there exists $\gamma > 0$ such that, for every $N \geq 2L$ and every vector $v \in (\mathcal{G}_{N,L}^{\text{ES}}(A))^\perp$,

$$\gamma \|v\| \leq \|H_{N,L}^{\text{ES}}(A)v\|.$$

As in Theorem 13.11.3.6 this is a conditional reduction: the operator-product hypothesis is unsatisfiable for any overlapping pair whose excitation ranges have a common nonzero vector, since then $\|h_{i,\text{ES}}h_{j,\text{ES}}\| = 1$. The statement does not assert this norm identity in degenerate cases.

Proof. [✓ Proof](#) This follows from Theorem 13.11.3.6 by taking $\gamma = 1 - \eta 2(L-1)$. $\square$

13.12 Block injectivity and degenerate ground spaces

For an MPS with more than one block in canonical form, after blocking to block-injective canonical form, the physical-virtual correspondence is restricted to block-diagonal boundary conditions. In the notation of [CPGSV21, lines 2113–2117], set

$$\tilde{A}^\omega := (A_1^\omega, \dots, A_g^\omega) \in \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

Then block injectivity is the statement

$$\forall X = (X_j)_{j=1}^g \in \bigoplus_{j=1}^g M_{D_j}(\mathbb{C}), \quad \exists (c_\omega(X))_\omega, \quad X = \sum_\omega c_\omega(X) \tilde{A}^\omega.$$

Equivalently, with the same coefficient family,

$$\forall j, \quad X_j = \sum_\omega c_\omega(X) A_j^\omega.$$

The strengthened parent-Hamiltonian theorem in [CPGSV21, Theorem IV.16, lines 2126–2128] is

$$N \geq L_0 + 1 \implies \ker H_N = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

The statements below record the corresponding periodic-boundary ground-space formulation; the passage to $\ker H_N$ is kept separate.

Definition 13.12.1 (Periodic-boundary ground space for the block-diagonal tensor). Not ready For a basis of normal tensors $A = (A_j)_{j=1}^g$ and nonzero weights μ_j , let

$$B := \bigoplus_{j=1}^g \mu_j A_j.$$

The ground space on the N -site ring considered in this section is

$$\mathcal{G}_{N,L}(B).$$

Lemma 13.12.2 (Block-diagonal periodic-boundary ground space). Not ready By definition, for $B = \bigoplus_{j=1}^g \mu_j A_j$ one has

$$\mathcal{G}_{N,L}(B) = \mathcal{G}_{N,L}\left(\bigoplus_{j=1}^g \mu_j A_j\right).$$

Definition 13.12.3 (Vector space generated by a basis of normal tensors). Lean ✓ Stmt For a basis of normal tensors $A_1, \dots, A_g$, the vector space generated by their length- N matrix product vectors is

$$\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

Lemma 13.12.4 (BNT span as the sum of one-vector spaces). Lean ✓ Stmt For a family $A_0, \dots, A_{r-1}$ and a chain length N , let

$$M_j = \text{span}\{V^{(N)}(A_j)\}.$$

Then

$$\bigvee_{j=0}^{r-1} M_j = \text{span}\{V^{(N)}(A_j) : j = 0, \dots, r-1\}.$$

Proof. ✓ Proof Both sides are the span of the same family of vectors. □

Definition 13.12.5 (Fixed-length block-injective canonical-form condition). Lean ✓ Stmt For a fixed blocked length m , the fixed-length form of the block-injective canonical-form condition is

$$\text{span}\{(A_1^\omega, \dots, A_g^\omega) : |\omega| = m\} = \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

Equivalently,

$$\forall X = (X_j)_j \in \bigoplus_{j=1}^g M_{D_j}(\mathbb{C}), \quad \exists (c_\omega(X))_{|\omega|=m}, \quad (X_j)_{j=1}^g = \sum_{|\omega|=m} c_\omega(X) (A_1^\omega, \dots, A_g^\omega).$$

This is the fixed-length form of the block-injective canonical-form condition of [CPGSV21, lines 2113–2117].

Lemma 13.12.6 (Each normal-tensor vector lies in the periodic-boundary ground space).

Not ready If $1 < L$ and $N \geq L + 1$, then for each normal tensor A_j in the basis the vector $V^{(N)}(A_j)$ lies in $\mathcal{G}_{N,L}(\bigoplus_k \mu_k A_k)$.

Proof. **Not ready** Write Γ_N^B for the length- N boundary-to-state map of the tensor B , and let $\iota_j(Z)$ be the block-diagonal matrix with Z in the j -th block and 0 elsewhere. The direct-sum evaluation gives

$$\Gamma_N^{\bigoplus_k \mu_k A_k}(\iota_j(\mu_j^{-N} X)) = \Gamma_N^{A_j}(X).$$

Taking $X = I$ gives the vector $V^{(N)}(A_j)$, and Lemma 13.6.8 applied to the individual block gives its membership in $\mathcal{G}_{N,L}(\bigoplus_k \mu_k A_k)$. $\square$

Lemma 13.12.7 (Inclusion of a block ground space). **Lean** **✓ Stmt** For each block index j with $\mu_j \neq 0$, and whenever $0 < N$ and $L \leq N$, the periodic-boundary ground space of the block A_j embeds into the corresponding ground space of the direct-sum tensor:

$$\mathcal{G}_{N,L}(A_j) \subseteq \mathcal{G}_{N,L}\left(\bigoplus_k \mu_k A_k\right).$$

Proof. **✓ Proof** The definition by length- L windows on the ring reduces the inclusion to the local block-embedding statement that each ground vector of A_j on the window lifts to the direct sum by the identity

$$\Gamma_L^{\bigoplus_k \mu_k A_k}(\iota_j(\mu_j^{-L} X)) = \Gamma_L^{A_j}(X).$$

Here ι_j inserts a matrix in the j -th diagonal block and has zero off that block. Thus $\iota_j(\mu_j^{-L} X)$ has exactly the local vector of the j -th block on that window. $\square$

Lemma 13.12.8 (Forward inclusion from the block ground spaces). **Lean** **✓ Stmt** Assume $\mu_j \neq 0$ for every block index j . Whenever $0 < N$ and $L \leq N$, the linear sum of the blockwise ground spaces with periodic boundary conditions is contained in the direct-sum tensor's ground space with periodic boundary conditions:

$$\sum_j \mathcal{G}_{N,L}(A_j) \subseteq \mathcal{G}_{N,L}\left(\bigoplus_k \mu_k A_k\right),$$

where j ranges over all block indices.

Proof. **✓ Proof** Take the linear sum of the inclusions

$$\mathcal{G}_{N,L}(A_j) \subseteq \mathcal{G}_{N,L}\left(\bigoplus_k \mu_k A_k\right)$$

over all block indices j . $\square$

Lemma 13.12.9 (Forward inclusion for the direct-sum tensor). Not ready Let $A^i = \bigoplus_j \mu_j A_j^i$ be in block-injective canonical form, where $A_1, \dots, A_g$ is a basis of normal tensors. The linear sum of the blockwise ground spaces with periodic boundary conditions is contained in the ground space of the block-diagonal tensor on the N -site ring:

$$\sum_j \mathcal{G}_{N,L}(A_j) \subseteq \mathcal{G}_{N,L}\left(\bigoplus_k \mu_k A_k\right).$$

Proof. Not ready The block-injective canonical-form hypothesis supplies $\mu_j \neq 0$ for every j , and the block-diagonal periodic-boundary ground space is $\mathcal{G}_{N,L}(\bigoplus_j \mu_j A_j)$ by definition. $\square$

13.13 Block separation and intersection

This section establishes that block-separating words make the simultaneous block-word tuples $\omega \mapsto (A_1^\omega, \dots, A_g^\omega)$ span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$ at a single length, and that this single-length span, together with the normalization $\sum_a A_a^j A_a^{j\dagger} = I$, yields the one-step intersection identity $\bigvee_j G_{n+2}(A^j)$ used for the degenerate parent-Hamiltonian ground space.

Lemma 13.13.1 (Block-diagonal commutant reduction from span extension). Not ready Let S be a family of virtual matrices. If each sector projection satisfies

$$P_j \in \text{span } S$$

and a boundary matrix X commutes with every M in S ,

$$XM = MX,$$

then, for every sector j ,

$$XP_j = P_j X,$$

and hence, for distinct sectors j and k ,

$$P_j X P_k = 0.$$

Thus X is block diagonal with respect to the sector decomposition.

Proof. Not ready The equations $XM = MX$ are linear in M , so they hold for every $M \in \text{span } S$. Substituting $M = P_j$ gives $XP_j = P_j X$. Multiplying this identity on the left by P_j and on the right by P_k with $j \neq k$ gives $P_j X P_k = 0$. $\square$

Lemma 13.13.2 (Block projections from product-algebra span). Not ready Let $T_a = (T_a(i))_i$ be a family of tuples spanning the full product algebra $\prod_i M_{n_i}(\mathbb{C})$, and let $c_i \neq 0$ be nonzero scalars. Then, for every sector k , the sector projection P_k lies in the span of the scaled dependent block-diagonal matrices

$$\bigoplus_i c_i T_a(i).$$

Proof. Not ready Apply the linear map $(M_i)_i \mapsto \bigoplus_i c_i M_i$ to the full-span product-algebra element with $(c_k)^{-1}I$ in the k -th component and zero in every other component. Its image is exactly P_k . $\square$

Lemma 13.13.3 (Sector projections from blockwise word span). Not ready If the simultaneous length- m block-word tuples

$$\omega \mapsto (A_j^\omega)_j$$

span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$ and every μ_j is nonzero, then the pulled-back length- m word products of $\bigoplus_j \mu_j A_j$ span every virtual sector projection P_j . If a virtual matrix X commutes with the pulled-back word product for every word ω of length m ,

$$X \left(\bigoplus_j \mu_j^m A_j^\omega \right) = \left(\bigoplus_j \mu_j^m A_j^\omega \right) X,$$

then, for distinct sectors j and k ,

$$P_j X P_k = 0.$$

Proof. Not ready First embed the product algebra linearly as dependent block-diagonal matrices, scaling the j -th block by μ_j^m . Since $\mu_j^m \neq 0$, the product-algebra span contains the tuple with $(\mu_j^m)^{-1}I$ in one component and zero elsewhere; its scaled diagonal image is exactly P_j . The word-evaluation formula for the direct-sum tensor identifies these scaled diagonal matrices with the pulled-back direct-sum word products, and the commutant conclusion follows from Lemma 13.13.1. □

Definition 13.13.4 (Pairwise block-separating equations). Lean Lean ✓ Stmt Fix a finite target set T of block indices. A length- S block-separating equation for k against T is a choice of coefficients c_ω such that

$$\sum_{|\omega|=S} c_\omega A_\ell^\omega = \begin{cases} I, & \ell = k, \\ 0, & \ell \in T. \end{cases}$$

The pairwise version requires this equation for each ordered pair of distinct blocks, with $T = \{j\}$.

Definition 13.13.5 (Trace-dual pair product-span condition). Lean Lean Lean Lean Lean Lean ✓ Stmt For two blocks A and B , the length- S pair word tuple is $w \mapsto (A^w, B^w)$. We also use the subalgebra of $M_{D_A}(\mathbb{C}) \times M_{D_B}(\mathbb{C})$ generated by the one-letter pairs (A_i, B_i) . The pair product-span condition says that the length- S tuples span $M_{D_A}(\mathbb{C}) \times M_{D_B}(\mathbb{C})$. Equivalently, in the trace-dual formulation of the same product-span condition, the only pair of test matrices (Δ_A, Δ_B) whose trace pairing with every length- S tuple vanishes is $(0, 0)$. The cumulative variant replaces one fixed length by all word lengths at most S , and the all-length variant allows every finite word.

Lemma 13.13.6 (Three-block trace reduction from a nonzero middle matrix). Lean Lean ✓ Stmt Let A and B be tensors with virtual bond dimensions D_A and D_B . Assume that A is L -block injective. Let $\Delta_A \in M_{D_A}(\mathbb{C})$ with $\Delta_A \neq 0$, and let $\Delta_B \in M_{D_B}(\mathbb{C})$. Suppose that, for all words u, v, t of length L ,

$$\text{tr}(\Delta_A A_v A_t A_u) + \text{tr}(\Delta_B B_v B_t B_u) = 0.$$

Then for every $Z \in M_{D_A}(\mathbb{C})$ there is a matrix $W \in M_{D_B}(\mathbb{C})$ such that, for all words t of length L ,

$$\text{tr}(Z A_t) + \text{tr}(W B_t) = 0.$$

This is the trace-dual algebraic step in the two-block case of the direct-sum lemma of [PGVWC07].

Proof. ✓ Proof Since $\Delta_A \neq 0$, Lemma 8.1.12 writes Z as a linear combination of matrices $A_u \Delta_A A_v$. For one such term, cyclicity of the trace rewrites the three-block relation as

$$\text{tr}(A_u \Delta_A A_v A_t) + \text{tr}(B_u \Delta_B B_v B_t) = 0.$$

Taking the same linear combination of the matrices $B_u \Delta_B B_v$ gives the required W , and linearity gives the general case. $\square$

Lemma 13.13.7 (Finite-chain image inclusion from a three-block relation). Lean Lean Lean Lean Lean ✓ Stmt Under the hypotheses of Lemma 13.13.6, the length- L spaces $\mathcal{G}_L^A$ and $\mathcal{G}_L^B$ satisfy

$$\mathcal{G}_L^A \subseteq \mathcal{G}_L^B.$$

If the corresponding nonzero middle matrix exists on both sides, then $\mathcal{G}_L^A = \mathcal{G}_L^B$.

Proof. ✓ Proof Write a vector in $\mathcal{G}_L^A$ as $t \mapsto \text{tr}(A_t Z)$. The preceding lemma supplies W with $\text{tr}(Z A_t) + \text{tr}(W B_t) = 0$ for every word t . Cyclicity of the trace gives

$$\text{tr}(A_t Z) = \text{tr}(Z A_t) = -\text{tr}(W B_t) = \text{tr}(B_t (-W)),$$

so the vector lies in $\mathcal{G}_L^B$. The equality statement follows by applying the same argument with the two blocks exchanged. $\square$

Lemma 13.13.8 (Dimension step for two-block local spaces). Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean Lean ✓ Stmt If A and B are length- L block-injective, the map $Z \mapsto (t \mapsto \text{tr}(A_t Z))$ has image dimension D_A^2 , and similarly for B . Therefore an inclusion $\mathcal{G}_L^A \subseteq \mathcal{G}_L^B$, together with $D_B \leq D_A$, forces $D_A = D_B$ and $\mathcal{G}_L^A = \mathcal{G}_L^B$. In particular, the three-block trace relation from Lemma 13.13.7 gives this equality whenever the ordering $D_B \leq D_A$ is available. If the inclusion comes from the strictly larger block, $D_B < D_A$, then the same dimension step already contradicts the existence of the three-block trace relation. More generally, let H be a source hypothesis which excludes the simultaneous conclusion

$$D_A = D_B, \quad \mathcal{G}_L^A = \mathcal{G}_L^B.$$

Then the three-block intersection vanishes:

$$H \implies \mathcal{G}_{3L}^A \cap \mathcal{G}_{3L}^B = \{0\}.$$

With injectivity at the three-block length, the strict-size branch gives

$$D_B < D_A, \quad \mathcal{G}_{3L}^A \cap \mathcal{G}_{3L}^B = \{0\} \implies \text{span}\{(A^\omega, B^\omega) : |\omega| = 3L\} = M_{D_A}(\mathbb{C}) \times M_{D_B}(\mathbb{C}).$$

One such source hypothesis is non-proportionality of the periodic MPS vectors at a sufficiently long chain length:

$$\mathbb{C} V^{(N)}(A) \neq \mathbb{C} V^{(N)}(B).$$

Indeed,

$$\mathcal{G}_L^A = \mathcal{G}_L^B \implies \mathcal{G}_{N,L}(A) = \mathcal{G}_{N,L}(B) \implies \mathbb{C} V^{(N)}(A) = \mathbb{C} V^{(N)}(B),$$

by equality of the periodic-boundary constraint spaces, followed by injective uniqueness. For same-dimensional normalized BNT representatives that are not gauge-phase equivalent, the non-equivalence condition supplies this non-proportionality at arbitrarily large chain lengths; the

direct-sum injectivity hypotheses remain explicit. Combining the equal-dimension and unequal-dimension branches over all ordered block pairs gives one common positive length S_0 with

$$\text{span}\{(A_1^\omega, \dots, A_g^\omega) : |\omega| = S_0\} = \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

For an ordered pair of blocks, the trace-separation conclusion has the form

$$[\forall |\omega| = S, \text{tr}(\Delta_A A^\omega) + \text{tr}(\Delta_B B^\omega) = 0] \implies \Delta_A = \Delta_B = 0.$$

Equivalently, the homogeneous pair span is full:

$$\text{span}\{(A^\omega, B^\omega) : |\omega| = S\} = M_{D_A}(\mathbb{C}) \times M_{D_B}(\mathbb{C}).$$

Concatenating words propagates this equality to positive multiples of S , and hence to a finite period window for every ordered pair. Combining that window with the conditional period-window homogenization theorem gives a single length S_0 such that

$$\text{span}\{(A_1^\omega, \dots, A_g^\omega) : |\omega| = S_0\} = \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

The same conclusion applies to the corresponding normal canonical-form hypotheses with a basis of normal tensors once one-site injectivity is supplied explicitly, since those hypotheses carry the irreducibility and normalization assumptions used in the direct-sum comparison.

Proof. ✓ Proof Block injectivity makes the boundary-to-chain map injective, so $\dim \mathcal{G}_L^A = D_A^2$ and $\dim \mathcal{G}_L^B = D_B^2$. Inclusion gives $D_A^2 \leq D_B^2$, while $D_B \leq D_A$ gives the reverse inequality. Hence $D_A = D_B$, and equality of dimensions inside the inclusion gives $\mathcal{G}_L^A = \mathcal{G}_L^B$. In the strict-size branch this contradicts $D_B < D_A$. For the conditional directness statement, suppose $\psi \in \mathcal{G}_{3L}^A \cap \mathcal{G}_{3L}^B$ has boundary representations

$$\psi = \Gamma_{3L}^A(X) = \Gamma_{3L}^B(Y).$$

If $X = 0$, injectivity of the A boundary-to-chain map gives $\psi = 0$. Otherwise equality of the coefficient formulas gives

$$\forall |t| = 3L, \quad \text{tr}(XA^t) - \text{tr}(YB^t) = 0,$$

equivalently

$$\forall |t| = 3L, \quad \text{tr}(XA^t) + \text{tr}((-Y)B^t) = 0.$$

The dimension step applied to this homogeneous three-block trace relation gives $D_A^2 \leq D_B^2$, contradicting $D_B < D_A$ in the strict-size branch. In the equal-size branch, equal local spaces impose the same length- L constraints on the N -site ring:

$$\mathcal{G}_L^A = \mathcal{G}_L^B \implies \mathcal{G}_{N,L}(A) = \mathcal{G}_{N,L}(B).$$

The injective uniqueness theorem identifies these periodic ground spaces with the one-dimensional MPS spans,

$$\mathcal{G}_{N,L}(A) = \mathbb{C} V^{(N)}(A), \quad \mathcal{G}_{N,L}(B) = \mathbb{C} V^{(N)}(B),$$

and hence

$$\mathbb{C} V^{(N)}(A) = \mathbb{C} V^{(N)}(B),$$

contradicting non-proportionality of the two MPS vectors. □

Lemma 13.13.11 (Conditional padding from exact homogeneous identity witnesses). [Lean](#)

[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#)

[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) ✓ Stmt

Homogeneous identity-padding witnesses multiply by concatenation of words. Therefore, if one has a positive period and a full residue window of homogeneous pair spans, then the simultaneous identity pair lies in all sufficiently large homogeneous pair spans. Combined with all-length pair trace separation, this period-window input gives one homogeneous pair trace-separation length. This does not prove the block-separation span condition of [PGVWC07] by itself: fullness of spans up to a bounded length is weaker than the exact homogeneous input used here.

Proof. ✓ **Proof** The length-zero word gives the identity only at length zero, so it is not a positive-length padding theorem. The useful padding facts are homogeneous: if the identity pair is in the span at lengths m and n , componentwise multiplication puts it in the span at length $m + n$. A period witness and a complete residue window therefore give homogeneous padding at every sufficiently large length, which is the input needed by the all-length-to-fixed-length implication. □

Lemma 13.13.12 (Pair trace separation gives pairwise block separation). [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *At a fixed length S , the pair trace-separation criterion is dual to full pair product-span. Therefore a pair trace-separation witness supplies a linear combination of word evaluations that is identity on the first block and zero on the second. A common witness for all ordered distinct pairs gives pairwise block-separating words at the same length.*

Proof. ✓ Proof If the pair span were proper, a nonzero linear functional would vanish on it. Nondegeneracy of the matrix trace pairing represents that functional as (Δ_A, Δ_B) , contradicting pair trace separation. Once the pair span is the whole product algebra, the tuple $(I, 0)$ is in the span. Using the same coefficients on the ambient block family gives the required pairwise block-separating equation. $\square$

Lemma 13.13.13 (Block-separating equations from pairwise separation). [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✔ Stmt](#) *If every ordered pair of distinct blocks admits a common-length pairwise block-separating equation, then for each k there are coefficients $c_\omega^{(k)}$ with*

$$\sum_{|\omega|=(g-1)S} c_{\omega}^{(k)} A_j^{\omega} = \begin{cases} I, & j = k, \\ 0, & j \neq k. \end{cases}$$

Hence common block injectivity plus pairwise separation implies the finite-length blockwise physical-virtual correspondence of the block-injectivity proposition in [CPGSV16, Section 2.3] and the block-injective canonical-form trace-separation condition.

Proof. ✓ Proof The empty word gives the identity tuple, corresponding to an empty target set. The span of word tuples is closed under pointwise multiplication because the product of a length- L word and a length- S word is the concatenated length- $(L+S)$ word. Inducting over the finite set of blocks different from k and multiplying the corresponding pairwise equations keeps the value I on k and introduces one additional zero block at each step. Extracting coefficients from the final tuple-span membership gives the coefficient form of the displayed block-separating equation. $\square$

Lemma 13.13.14 (Homogeneous padding for blockwise product spans). [\[Lean\]](#) [\[Lean\]](#) [\[StmT\]](#)
Suppose the length- L simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$, and suppose the simultaneous identity tuple $(I, \dots, I)$ lies in the span of the length- S simultaneous block-word tuples. Then the length- $L + S$ simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$. In particular, two full homogeneous spans at lengths L and S give a full homogeneous span at length $L + S$.

Proof. ✓ Proof Let $M = (M_j)_j$ be a target tuple. Write

$$M_j = \sum_{|u|=L} a_u A_u^j, \quad I = \sum_{|v|=S} b_v A_v^j$$

simultaneously for every block j . Multiplying the two expansions gives

$$M_j = M_j I = \sum_{|u|=L} \sum_{|v|=S} a_u b_v A_u^j A_v^j = \sum_{|u|=L} \sum_{|v|=S} a_u b_v A_{uv}^j.$$

The words uv have length $L + S$, so the target tuple lies in the length- $L + S$ span. □

Lemma 13.13.15 (Period-window propagation for blockwise product spans). Lean Lean ✓ Stmt
Let $p > 0$. Suppose the length- p simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$, and suppose that for every $0 \leq r < p$, the length- $s + r$ simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$. Then there is a length N such that, for every $n \geq N$, the length- n simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$. More concretely, one may take $N = s$: writing

$$n = s + r + qp, \quad 0 \leq r < p,$$

the span at length $s + r$ propagates to the span at length n by multiplying q times by the period- p span.

Proof. ✓ Proof The Euclidean division of $n - s$ by p gives $n = s + r + qp$ with $0 \leq r < p$. The assumed residue-window span gives the full product algebra at length $s + r$. Applying homogeneous padding with the length- p span once gives the full product algebra at length $s + r + p$; iterating gives the full product algebra at length $s + r + qp$. □

Lemma 13.13.16 (Blockwise word span from block-separating equations). Lean Lean ✓ Stmt
Let J be a finite set of r blocks, and let $(A_j)_{j \in J}$ be a finite family of tensors with common block injectivity at length L . Suppose there are length- S block-separating equations: for each block k ,

$$\sum_{|\omega|=S} c_\omega^{(k)} A_j^\omega = \begin{cases} I, & j = k, \\ 0, & j \neq k. \end{cases}$$

Then the simultaneous length- $(L + S)$ block-word evaluations span $\prod_j M_{D_j}(\mathbb{C})$. If the block-separating equations are first obtained from pairwise block-separating equations at length S , the same conclusion holds at length $L + (r - 1)S$.

Proof. ✓ Proof Block injectivity writes each target block matrix as a length- L linear combination of word evaluations. For a target tuple $(M_j)_j$, write M_k in this form in block k , multiply by the block-separating equation for k , and sum over k . Thus, if $M_k = \sum_{|u|=L} a_u^{(k)} A_k^u$ and $\sum_{|v|=S} b_v^{(k)} A_j^v = \delta_{j,k} I$, then for every block j ,

$$M_j = \sum_{k \in J} \sum_{|u|=L} \sum_{|v|=S} a_u^{(k)} b_v^{(k)} A_j^u A_j^v.$$

The right side is a linear combination of length- $(L + S)$ word tuples, so these tuples span the full product algebra. If one starts with pairwise separation instead, first use Lemma 13.13.13 to multiply the pairwise equations into the full equations. □

13.13.1 Trace identities for block intersections

Once the simultaneous block-word tuples span the full product algebra at a single length, non-degeneracy of the product trace pairing turns blockwise trace equalities into blockwise matrix identities $A_b^j C_a^j = A_b^j E^j A_a^j$. The lemmas here use those identities to place left-boundary trace decompositions in the block sum $\bigvee_j G_{n+2}(A^j)$ and to obtain the one-step block-intersection identity.

Lemma 13.13.1.1 (BNT block separation gives a finite blockwise product span). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Stmt](#) *Let $A^1, \dots, A^r$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent. Fix $L_0 > 0$. Assume that, for every block j , the map*

$$\Gamma_{L_0}^{A^j} : M_{D_j}(\mathbb{C}) \rightarrow (\mathbb{C}^d)^{\otimes L_0}, \quad X \mapsto \sum_{|u|=L_0} \text{tr}(X A_u^j) |u\rangle$$

is injective, and that the homogeneous word spans at the two source lengths are full:

$$\text{span}\{A_u^j : |u| = L_0 + 1\} = M_{D_j}(\mathbb{C}), \quad \text{span}\{A_v^j : |v| = 3(L_0 + 1)\} = M_{D_j}(\mathbb{C}).$$

Then for every ordered pair $k \neq j$ there are coefficients $c_\omega^{k,j}$ such that

$$\sum_{|\omega|=3(L_0+1)} c_\omega^{k,j} A_\omega^k = I, \quad \sum_{|\omega|=3(L_0+1)} c_\omega^{k,j} A_\omega^j = 0.$$

Consequently the simultaneous block-word tuples at length

$$(L_0 + 1) + (r - 1)3(L_0 + 1)$$

span the full product algebra

$$\prod_{j=1}^r M_{D_j}(\mathbb{C}).$$

Proof. [Proof](#) Lemma 13.13.8 gives homogeneous trace separation at length $3(L_0 + 1)$ for every ordered pair of distinct normalized BNT representatives. Lemma 13.13.12 converts these trace-separation equations into the displayed pairwise block-separating equations. Multiplying the equations for the $r - 1$ blocks different from k gives a block-separating equation for k , and Lemma 13.13.16 then supplies the displayed product span. $\square$

Lemma 13.13.1.2 (Blockwise matrix equality from trace pairings). [Lean](#) [Lean](#) [Stmt](#) *Let $A^1, \dots, A^g$ be block tensors whose length- m simultaneous word tuples span the product algebra. If, for every word w of length m , two blockwise matrix families X_j and Y_j satisfy*

$$\sum_j \text{tr}(X_j A_w^j) = \sum_j \text{tr}(Y_j A_w^j)$$

then $X_j = Y_j$ for every block j .

Proof. [Proof](#) For every word w of length m ,

$$\sum_j \text{tr}((X_j - Y_j) A_w^j) = \sum_j \text{tr}(X_j A_w^j) - \sum_j \text{tr}(Y_j A_w^j) = 0.$$

Nondegeneracy of the product trace pairing, together with the spanning hypothesis, gives $X_j - Y_j = 0$ for every block j . $\square$

Lemma 13.13.1.3 (Boundary identities from block-sum membership). Lean ✓ Stmt Let $A^1, \dots, A^g$ be block tensors whose length- m simultaneous word tuples span the product algebra. Let $C_a^j \in M_{D_j}(\mathbb{C})$ be matrices indexed by blocks j and physical letters a . Suppose that, for every pair of boundary letters a, b and every word w of length m , a vector ψ has the left-boundary trace form

$$\psi(a, w, b) = \sum_j \text{tr}(A_b^j C_a^j A_w^j)$$

and if

$$\psi \in \bigvee_j G_{m+2}(A^j),$$

then there are matrices E_j such that, for every block j and all a, b ,

$$A_b^j C_a^j = A_b^j E_j A_a^j$$

Proof. ✓ Proof Write $\psi = \sum_j \phi_j$ with $\phi_j \in G_{m+2}(A^j)$. For each j , choose E_j whose image under the length- $m+2$ ground-space map for A^j is ϕ_j . Comparing the coefficient of the word awb in the two expressions for ψ gives

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(A_b^j E_j A_a^j A_w^j).$$

Lemma 13.13.1.2 gives the displayed blockwise identities. □

Lemma 13.13.1.4 (Blockwise boundary compatibility from traces). Lean ✓ Stmt Let $A^1, \dots, A^g$ be block tensors with the following common word-span property: for a fixed m , the simultaneous tuples

$$w \mapsto (A_w^1, \dots, A_w^g), \quad |w| = m,$$

span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Suppose that matrices $C_a^j, D_b^j \in M_{D_j}(\mathbb{C})$ satisfy, for all physical indices a, b and all words w of length m ,

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

Then

$$A_b^j C_a^j = D_b^j A_a^j$$

for every block j and all a, b .

Proof. ✓ Proof For fixed a, b , set

$$\Delta_j = A_b^j C_a^j - D_b^j A_a^j.$$

The assumed trace equality says that

$$\sum_j \text{tr}(\Delta_j A_w^j) = 0$$

for every word w of length m . Since the tuples $(A_w^1, \dots, A_w^g)$ span the product algebra, the product trace pairing is nondegenerate and hence every Δ_j is zero. □

Lemma 13.13.1.5 (Word-valued boundary compatibility from traces). [Lean](#) [Lean](#) [✓ Stmt](#) Let $A^1, \dots, A^g$ be block tensors with the common word-span property at length m . Suppose that $C_\rho^j \in M_{D_j}(\mathbb{C})$ is indexed by a block j and a word ρ of some fixed length M , and that $D_\beta^j \in M_{D_j}(\mathbb{C})$ is indexed by a block j and a word β of some fixed length K . If, for every word ρ of length M , every word β of length K , and every middle word w of length m ,

$$\sum_j \operatorname{tr}(A_\beta^j C_\rho^j A_w^j) = \sum_j \operatorname{tr}(D_\beta^j A_\rho^j A_w^j),$$

then

$$A_\beta^j C_\rho^j = D_\beta^j A_\rho^j$$

for every block j and all words β, ρ . In particular, taking $D_\beta^j = X_j A_\beta^j$ gives

$$A_\beta^j C_\rho^j = (X_j A_\beta^j) A_\rho^j.$$

Proof. [✓ Proof](#) Fix β and ρ , and set

$$\Delta_j = A_\beta^j C_\rho^j - D_\beta^j A_\rho^j.$$

The trace equality says that

$$\sum_j \operatorname{tr}(\Delta_j A_w^j) = 0$$

for every word w of length m . The same nondegenerate product trace-pairing argument as in Lemma 13.13.1.4 gives $\Delta_j = 0$ for every j . The specialization $D_\beta^j = X_j A_\beta^j$ is immediate. $\square$

Lemma 13.13.1.6 (Fixed complementary-word compatibility from traces). [Lean](#) [✓ Stmt](#) Let $A^1, \dots, A^g$ be block tensors with the common word-span property at length m . Fix a complementary word ρ of length M and matrices $C_\rho^j, X_j \in M_{D_j}(\mathbb{C})$. If, for every wrapped word β of length K and every middle word w of length m ,

$$\sum_j \operatorname{tr}(A_\beta^j C_\rho^j A_w^j) = \sum_j \operatorname{tr}((X_j A_\beta^j) A_\rho^j A_w^j),$$

then

$$A_\beta^j C_\rho^j = (X_j A_\beta^j) A_\rho^j$$

for every block j and every wrapped word β .

Proof. [✓ Proof](#) Fix β and set

$$\Delta_j = A_\beta^j C_\rho^j - (X_j A_\beta^j) A_\rho^j.$$

The trace equality gives

$$\sum_j \operatorname{tr}(\Delta_j A_w^j) = 0$$

for every word w of length m . The common word-span property and the product trace pairing imply $\Delta_j = 0$ for every j . $\square$

Lemma 13.13.1.7 (Normalized boundary matrix identities). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $(A_a)_{a \in I}$, $(C_a)_{a \in I}$, and $(D_b)_{b \in I}$ be matrices in $M_D(\mathbb{C})$, indexed by a finite alphabet. If

$$\sum_a A_a A_a^\dagger = \mathbb{1}, \quad A_b C_a = D_b A_a$$

for all $a, b \in I$, and if

$$E = \sum_a C_a A_a^\dagger,$$

then

$$D_b = A_b E, \quad A_b C_a = A_b E A_a$$

for all $a, b \in I$.

In particular, the same calculation applies when the alphabet is the finite set of words of a fixed length and A_β denotes $A_{\beta_1} \cdots A_{\beta_K}$.

The same argument also applies when the normalized family is indexed by a finite set J , while $(A_b)_{b \in I}$ and $(D_b)_{b \in I}$ are indexed by a set I . If $(B_a)_{a \in J}$ and $(C_a)_{a \in J}$ are indexed by J , and if for all $a \in J$ and $b \in I$,

$$\sum_a B_a B_a^\dagger = \mathbb{1}, \quad A_b C_a = D_b B_a$$

then with

$$E = \sum_a C_a B_a^\dagger$$

one has

$$D_b = A_b E, \quad A_b C_a = A_b E B_a.$$

This includes the case where I and J are word sets of different lengths.

Proof.  **Proof** Multiply D_b by the normalization:

$$D_b = D_b \sum_a A_a A_a^\dagger = \sum_a D_b A_a A_a^\dagger = \sum_a A_b C_a A_a^\dagger = A_b E.$$

Substituting this in $A_b C_a = D_b A_a$ gives $A_b C_a = A_b E A_a$. For two alphabets, the corresponding step is

$$D_b = D_b \sum_a B_a B_a^\dagger = \sum_a D_b B_a B_a^\dagger = \sum_a A_b C_a B_a^\dagger = A_b E,$$

where $E = \sum_a C_a B_a^\dagger$. Substituting this in $A_b C_a = D_b B_a$ gives $A_b C_a = A_b E B_a$. □

Lemma 13.13.1.8 (Explicit C^j, D^j, E^j identities from trace decompositions).  **Lean**  **Stmt** Let $A^1, \dots, A^g$ be block tensors with the common word-span property at length m . Fix matrices X_j , and put

$$D_\beta^j = X_j A_\beta^j.$$

Suppose that C_ρ^j is indexed by a block j and a complementary word ρ of length M , and assume that, for every wrapped word β of length K , every complementary word ρ , and every middle word w of length m ,

$$\sum_j \text{tr}(A_\beta^j C_\rho^j A_w^j) = \sum_j \text{tr}(D_\beta^j A_\rho^j A_w^j).$$

If each block satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I,$$

then for every block j there is a matrix

$$E^j = \sum_\rho C_\rho^j (A_\rho^j)^\dagger$$

such that, for every wrapped word β and complementary word ρ ,

$$D_\beta^j = A_\beta^j E^j, \quad A_\beta^j C_\rho^j = A_\beta^j E^j A_\rho^j.$$

This is the calculation in [PGVWC07, Theorem 12, proof lines 1446–1451], with the adjoint on A_ρ^j required by the displayed normalization.

Proof. ✓ **Proof** Lemma 13.13.1.5 gives

$$A_\beta^j C_\rho^j = D_\beta^j A_\rho^j.$$

The single-letter normalization gives the word normalization

$$\sum_\rho A_\rho^j (A_\rho^j)^\dagger = I.$$

Lemma 13.13.1.7, applied to the two word alphabets, gives

$$E^j = \sum_\rho C_\rho^j (A_\rho^j)^\dagger, \quad D_\beta^j = A_\beta^j E^j, \quad A_\beta^j C_\rho^j = A_\beta^j E^j A_\rho^j.$$

□

Lemma 13.13.1.9 (Block-diagonal boundary C^j, D^j, E^j identities from trace decompositions).

Lean ✓ **Stmt** Let $A^1, \dots, A^g$ be block tensors with the common word-span property at length m . Fix $N \in \mathbb{N}$, complex numbers μ_j , and block-diagonal boundary matrices X_j , and put

$$D_\beta^j = (\mu_j^N X_j) A_\beta^j.$$

Suppose that, for every wrapped word β of length K , every complementary word ρ of length M , and every middle word w of length m ,

$$\sum_j \text{tr}(A_\beta^j C_\rho^j A_w^j) = \sum_j \text{tr}((\mu_j^N X_j) A_\beta^j A_\rho^j A_w^j).$$

If each block satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I,$$

then for every block j there is a matrix

$$E^j = \sum_\rho C_\rho^j (A_\rho^j)^\dagger$$

such that, for every wrapped word β and complementary word ρ ,

$$(\mu_j^N X_j) A_\beta^j = A_\beta^j E^j, \quad A_\beta^j C_\rho^j = A_\beta^j E^j A_\rho^j.$$

This is the boundary-matrix calculation in [PGVWC07, Theorem 12, proof lines 1446–1451], after substituting the block-diagonal boundary matrix $D_\beta^j = (\mu_j^N X_j) A_\beta^j$.

Proof. ✓ **Proof** Apply Lemma 13.13.1.8 with $D_\beta^j = (\mu_j^N X_j) A_\beta^j$. □

Lemma 13.13.1.10 (Complementary-word boundary identities from a compatibility hypothesis). Lean ✓ Stmt Let A_β be the products over wrapped words of length K , let A_ρ be the products over complementary words of length M , and let $X \in M_D(\mathbb{C})$ be a matrix. Suppose that the complementary-word products satisfy

$$\sum_{\rho} A_{\rho} A_{\rho}^{\dagger} = \mathbb{1}$$

and that there are matrices C_{ρ} , indexed by complementary words, such that for every wrapped word β and complementary word ρ ,

$$A_{\beta} C_{\rho} = (X A_{\beta}) A_{\rho}.$$

Then, for every complementary word ρ , with

$$E_{\rho} = \left(\sum_{\gamma} C_{\gamma} A_{\gamma}^{\dagger} \right) A_{\rho},$$

one has, for every wrapped word β ,

$$(X A_{\beta}) A_{\rho} = A_{\beta} E_{\rho}.$$

Proof. ✓ Proof Apply Lemma 13.13.1.7 with $D_{\beta} = X A_{\beta}$. It gives

$$A_{\beta} C_{\rho} = A_{\beta} E A_{\rho}$$

for $E = \sum_{\gamma} C_{\gamma} A_{\gamma}^{\dagger}$. With $E_{\rho} = E A_{\rho}$, the compatibility hypothesis gives

$$(X A_{\beta}) A_{\rho} = A_{\beta} C_{\rho} = A_{\beta} E A_{\rho} = A_{\beta} E_{\rho}.$$

□

Lemma 13.13.1.11 (Complementary-word boundary identities from trace decompositions). Lean ✓ Stmt Let $A^1, \dots, A^g$ be block tensors with the common word-span property at length m . Fix matrices X_j . Suppose that C_{ρ}^j is indexed by a block j and a complementary word ρ of length M . Assume that, for every wrapped word β of length K , every complementary word ρ , and every middle word w of length m ,

$$\sum_j \text{tr}(A_{\beta}^j C_{\rho}^j A_w^j) = \sum_j \text{tr}((X_j A_{\beta}^j) A_{\rho}^j A_w^j).$$

If each block satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I,$$

then, for every block j and every complementary word ρ , there is a matrix $E_{j,\rho}$ such that, for every wrapped word β ,

$$(X_j A_{\beta}^j) A_{\rho}^j = A_{\beta}^j E_{j,\rho}.$$

Proof. ✓ Proof Lemma 13.13.1.5 applied with $D_{\beta}^j = X_j A_{\beta}^j$ gives

$$A_{\beta}^j C_{\rho}^j = (X_j A_{\beta}^j) A_{\rho}^j.$$

The hypothesis $\sum_a A_a^j A_a^{j\dagger} = I$ extends to words of length M :

$$\sum_{\rho} A_{\rho}^j A_{\rho}^{j\dagger} = I$$

. Lemma 13.13.1.10 gives the matrices $E_{j,\rho}$. □

Lemma 13.13.1.12 (A left-boundary summand is a ground-space vector). [Lean](#) [✓ Stmt](#) Let A be a block tensor, and let $C_a, E \in M_D(\mathbb{C})$ satisfy

$$A_b C_a = A_b E A_a$$

for all physical indices a, b . For a word $\sigma = (\sigma_1, \dots, \sigma_{n+2})$, define

$$\alpha(\sigma) = \text{tr}(A_{\sigma_{n+2}} C_{\sigma_1} A_{\sigma_2} \cdots A_{\sigma_{n+1}}).$$

Then $\alpha \in G_{n+2}(A)$.

Proof. [✓ Proof](#) The boundary identity gives

$$A_{\sigma_{n+2}} C_{\sigma_1} = A_{\sigma_{n+2}} E A_{\sigma_1}.$$

Hence, by cyclicity of the trace,

$$\alpha(\sigma) = \text{tr}(A_{\sigma_1} A_{\sigma_2} \cdots A_{\sigma_{n+2}} E),$$

which is the ground-space parametrization of E . □

Lemma 13.13.1.13 (Left-boundary summands are ground-space vectors). [Lean](#) [✓ Stmt](#) Let $A^1, \dots, A^g$ be block tensors, and let $C_a^j, E^j \in M_{D_j}(\mathbb{C})$ satisfy

$$A_b^j C_a^j = A_b^j E^j A_a^j$$

for every block j and all physical indices a, b . For a word $\sigma = (\sigma_1, \dots, \sigma_{n+2})$, define

$$\alpha_j(\sigma) = \text{tr}(A_{\sigma_{n+2}}^j C_{\sigma_1}^j A_{\sigma_2}^j \cdots A_{\sigma_{n+1}}^j).$$

Then

$$\sum_j \alpha_j \in \bigvee_j G_{n+2}(A^j).$$

Proof. [✓ Proof](#) Fix j . By the assumed boundary identity and cyclicity of the trace,

$$\text{tr}(A_{\sigma_{n+2}}^j C_{\sigma_1}^j A_{\sigma_2}^j \cdots A_{\sigma_{n+1}}^j) = \text{tr}(A_{\sigma_1}^j A_{\sigma_2}^j \cdots A_{\sigma_{n+2}}^j E^j).$$

Thus α_j is the image of E^j under the parametrization of $G_{n+2}(A^j)$. Hence each α_j lies in $G_{n+2}(A^j)$, and the finite sum lies in the supremum of these subspaces. □

Lemma 13.13.1.14 (Left-boundary trace decompositions lie in the block ground spaces). [Lean](#) [✓ Stmt](#) Let $A^1, \dots, A^g$ be block tensors with common word-span separation at length n , and suppose that

$$\sum_a A_a^j A_a^{j\dagger} = \mathbb{1}$$

for every block j . Suppose a vector $\psi \in (\mathbb{C}^d)^{\otimes(n+2)}$ has the left-boundary trace decomposition

$$\psi = \sum_j \alpha_j, \quad \alpha_j(i_1, \dots, i_{n+2}) = \text{tr}(A_{i_{n+2}}^j C_{i_1}^j A_{i_2}^j \cdots A_{i_{n+1}}^j),$$

and suppose that the two trace decompositions agree for every middle word:

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

Then

$$\psi \in \bigvee_j G_{n+2}(A^j).$$

Proof. ✓ Proof The equality of the two boundary trace expressions and the common word-span hypothesis give

$$A_b^j C_a^j = D_b^j A_a^j$$

for every j, a, b . By the normalization,

$$D_b^j = D_b^j \sum_a A_a^j A_a^{j\dagger} = \sum_a A_b^j C_a^j A_a^{j\dagger} = A_b^j E^j, \quad E^j := \sum_a C_a^j A_a^{j\dagger}.$$

Hence $A_b^j C_a^j = A_b^j E^j A_a^j$. Lemma 13.13.1.13 then gives

$$\psi = \sum_j \alpha_j \in \bigvee_j G_{n+2}(A^j).$$

□

Lemma 13.13.1.15 (Trace decompositions from fixed-boundary restrictions). Lean ✓ Stmt Let $A^1, \dots, A^g$ be block tensors. If $\psi \in (\mathbb{C}^d)^{\otimes(n+2)}$ satisfies

$$\psi(a, -) \in \bigvee_j G_{n+1}(A^j), \quad \psi(-, b) \in \bigvee_j G_{n+1}(A^j)$$

for every physical index a, b , then there are matrices C_a^j and D_b^j such that

$$\psi = \sum_j \alpha_j, \quad \alpha_j(i_1, \dots, i_{n+2}) = \text{tr}(A_{i_{n+2}}^j C_{i_1}^j A_{i_2}^j \dots A_{i_{n+1}}^j),$$

and, for every middle word w of length n ,

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

Proof. ✓ Proof Choose finite decompositions of the fixed-boundary vectors:

$$\psi(a, i_2, \dots, i_{n+2}) = \sum_j \text{tr}(A_{i_2}^j \dots A_{i_{n+2}}^j C_a^j),$$

and

$$\psi(i_1, \dots, i_{n+1}, b) = \sum_j \text{tr}(A_{i_1}^j \dots A_{i_{n+1}}^j D_b^j).$$

For every middle word w ,

$$\psi(a, w, b) = \sum_j \text{tr}(A_w^j A_b^j C_a^j) = \sum_j \text{tr}(A_b^j C_a^j A_w^j),$$

and the second decomposition gives

$$\psi(a, w, b) = \sum_j \text{tr}(A_a^j A_w^j D_b^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

Hence

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

For an arbitrary word $(i_1, \dots, i_{n+2})$, the first decomposition gives

$$\psi(i_1, \dots, i_{n+2}) = \sum_j \text{tr}(A_{i_2}^j \dots A_{i_{n+2}}^j C_{i_1}^j) = \sum_j \text{tr}(A_{i_{n+2}}^j C_{i_1}^j A_{i_2}^j \dots A_{i_{n+1}}^j),$$

by cyclicity of the trace. □

Lemma 13.13.1.16 (One-step block intersection from block restrictions). [Lean](#) [Stmt](#) Let $A^1, \dots, A^g$ be block tensors with common word-span separation at length n , and suppose

$$\sum_a A_a^j A_a^{j\dagger} = \mathbb{1}$$

for every block j . If $\psi \in (\mathbb{C}^d)^{\otimes(n+2)}$ satisfies

$$\psi(a, -) \in \bigvee_j G_{n+1}(A^j), \quad \psi(-, b) \in \bigvee_j G_{n+1}(A^j)$$

for every physical index a, b , then

$$\psi \in \bigvee_j G_{n+2}(A^j).$$

Proof. [Proof](#) Choose finite decompositions of the fixed-boundary vectors:

$$\psi(a, i_2, \dots, i_{n+2}) = \sum_j \text{tr}(A_{i_2}^j \cdots A_{i_{n+2}}^j C_a^j),$$

and

$$\psi(i_1, \dots, i_{n+1}, b) = \sum_j \text{tr}(A_{i_1}^j \cdots A_{i_{n+1}}^j D_b^j).$$

Evaluating these two decompositions on (a, w, b) , where w has length n , gives

$$\sum_j \text{tr}(A_b^j C_a^j A_w^j) = \sum_j \text{tr}(D_b^j A_a^j A_w^j).$$

The first decomposition also gives

$$\psi = \sum_j \alpha_j, \quad \alpha_j(i_1, \dots, i_{n+2}) = \text{tr}(A_{i_{n+2}}^j C_{i_1}^j A_{i_2}^j \cdots A_{i_{n+1}}^j).$$

Lemma 13.13.1.14 therefore gives

$$\psi \in \bigvee_j G_{n+2}(A^j).$$

□

Lemma 13.13.1.17 (One-step block intersection characterization). [Lean](#) [Stmt](#) Under the hypotheses of Lemma 13.13.1.16, an $(n+2)$ -site vector satisfies

$$\psi \in \bigvee_j G_{n+2}(A^j)$$

if and only if

$$\psi(a, -) \in \bigvee_j G_{n+1}(A^j), \quad \psi(-, b) \in \bigvee_j G_{n+1}(A^j)$$

for every physical index a, b .

Proof. [Proof](#) The reverse implication is Lemma 13.13.1.16. For the forward implication, write

$$\psi = \sum_j \gamma_j, \quad \gamma_j \in G_{n+2}(A^j).$$

Each γ_j has fixed-boundary restrictions in $G_{n+1}(A^j)$:

$$\gamma_j(a, -) \in G_{n+1}(A^j), \quad \gamma_j(-, b) \in G_{n+1}(A^j).$$

Taking the finite sum over j gives the two restrictions of ψ in $\bigvee_j G_{n+1}(A^j)$. □

Lemma 13.13.1.18 (Subspace form of the one-step block intersection). [Lean](#) [✓ Stmt](#) Under the hypotheses of Lemma 13.13.1.16, set

$$S_n := \bigvee_j G_{n+1}(A^j).$$

Then

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. [✓ Proof](#) Lemma 13.13.1.17 gives the displayed equivalence for each vector ψ . Interpreting the two sides of that equivalence as subspaces gives the stated equality. $\square$

13.13.2 Eventual block-intersection consequences

Propagating the single-length product span through a period window gives the block-intersection identity for all sufficiently large n , both from a general period window and from the normalized BNT representative hypotheses. The full product span at length n also makes the sum of the local spaces $G_n(A^j)$ direct.

Lemma 13.13.2.1 (Period-window form of the block intersection identity). [Lean](#) [✓ Stmt](#) Let $p > 0$. Suppose the length- p simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$, and suppose that for every $0 \leq r < p$, the length- $s + r$ simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$. Assume also the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I$$

for every block j . Then there is a length N such that, for every $n \geq N$, if

$$S_n := \bigvee_j G_{n+1}(A^j),$$

then

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. [✓ Proof](#) Lemma 13.13.15 gives a length N such that the simultaneous block-word tuples span $\prod_j M_{D_j}(\mathbb{C})$ at every length $n \geq N$. For each such n , Lemma 13.13.1.18 applies and gives the displayed subspace equality. $\square$

Lemma 13.13.2.2 (BNT block separation gives one block-intersection step). [Lean](#) [✓ Stmt](#) Under the hypotheses of Lemma 13.13.1.1, assume in addition the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I$$

for every block j . Let

$$n = (L_0 + 1) + (r - 1)3(L_0 + 1), \quad S_n = \bigvee_j G_{n+1}(A^j).$$

Then

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. ✓ **Proof** Lemma 13.13.1.1 gives

$$\text{span}\{(A_w^1, \dots, A_w^r) : |w| = n\} = \prod_j M_{D_j}(\mathbb{C}).$$

Substituting this span into Lemma 13.13.1.18, together with the normalization, gives the stated subspace equality. $\square$

Lemma 13.13.2.3 (BNT block separation with a block-injectivity window). Lean ✓ **Stmt** *Let $A^1, \dots, A^r$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent. Assume $L > 1$, and assume that, for every block j ,*

$$\text{span}\{A_a^j : a = 0, \dots, d-1\} = M_{D_j}(\mathbb{C}), \quad \text{span}\{A_u^j : |u| = L\} = M_{D_j}(\mathbb{C}), \quad \text{span}\{A_v^j : |v| = 3L\} = M_{D_j}(\mathbb{C}).$$

Set

$$S = 3L, \quad q = (r-1)S.$$

Suppose that $p > 0$ and let s_0 be a starting length. Assume that, for every block j ,

$$\text{span}\{A_u^j : |u| = p\} = M_{D_j}(\mathbb{C}),$$

and that, for every $0 \leq s < p + q$ and every block j ,

$$\text{span}\{A_v^j : |v| = s_0 + s\} = M_{D_j}(\mathbb{C}).$$

Assume also the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I$$

for every block j . Then there is N such that, for every $n \geq N$, if

$$S_n := \bigvee_j G_{n+1}(A^j),$$

then

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. ✓ **Proof** The direct-sum dimension step gives, for each ordered pair of distinct blocks $k \neq j$, coefficients $c_\omega^{k,j}$ of length S such that

$$\sum_{|\omega|=S} c_\omega^{k,j} A_\omega^k = I, \quad \sum_{|\omega|=S} c_\omega^{k,j} A_\omega^j = 0.$$

Multiplying these equations over the $r-1$ blocks $j \neq k$ gives coefficients $c_\omega^{(k)}$ of total length q satisfying

$$\sum_{|\omega|=q} c_\omega^{(k)} A_\omega^j = \begin{cases} I, & j = k, \\ 0, & j \neq k. \end{cases}$$

Combining this block-separating equation with block injectivity at length p gives

$$\text{span}\{(A_w^1, \dots, A_w^r) : |w| = p + q\} = \prod_j M_{D_j}(\mathbb{C}).$$

Combining the same separating equations with the assumed block-injectivity window gives the full product span at lengths $s_0 + q + s$, for $0 \leq s < p + q$. The period-window form of Lemma 13.13.2.1 then gives the displayed equality for all sufficiently large n . $\square$

Lemma 13.13.2.4 (Normalized block-separating equations give all large product spans). [Lean](#)

[Lean](#) [✓ Stmt](#) Let $A^1, \dots, A^r$ be blocks satisfying the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I$$

and assume that every block is injective at length L :

$$\text{span}\{A_u^j : |u| = L\} = M_{D_j}(\mathbb{C}).$$

If there are block-separating equations of length S , then, for every $n \geq L + (r-1)S$,

$$\text{span}\{(A_w^1, \dots, A_w^r) : |w| = n\} = \prod_j M_{D_j}(\mathbb{C}).$$

In particular, for normalized BNT representatives satisfying the displayed unital normalization, if $L_0 > 0$ and $\Gamma_{L_0}^{A^j}$ is injective for every block j , the same conclusion holds with

$$S = 3(L_0 + 1), \quad n \geq (L_0 + 1) + (r-1)3(L_0 + 1).$$

Proof. [✓ Proof](#) By Theorem 8.11.4, the normalization propagates the length- L span equality to every larger homogeneous length:

$$m \geq L \implies \text{span}\{A_u^j : |u| = m\} = M_{D_j}(\mathbb{C}).$$

For $n \geq L + (r-1)S$, take $m = n - (r-1)S$. Combining the length- m block span with the $r-1$ block-separating equations gives the displayed product span at length n . In the BNT case, the injectivity of $\Gamma_{L_0}^{A^j}$ gives the full span at length L_0 , and the normalization propagates it to the lengths $L_0 + 1$ and $3(L_0 + 1)$. Lemma 13.13.1.1 then supplies the block-separating equations with $S = 3(L_0 + 1)$. $\square$

Lemma 13.13.2.5 (Product span makes the sum of local spaces direct). [Lean](#) [Lean](#) [✓ Stmt](#)
Suppose

$$\text{span}\{(A_w^1, \dots, A_w^r) : |w| = n\} = \prod_j M_{D_j}(\mathbb{C}).$$

Then the sum of the local spaces $G_n(A^j)$ is direct: if

$$\phi_j \in G_n(A^j), \quad \sum_j \phi_j = 0,$$

then $\phi_j = 0$ for every j . In particular, for the normalized BNT hypotheses above, this directness holds for every

$$n \geq (L_0 + 1) + (r-1)3(L_0 + 1).$$

Proof. [✓ Proof](#) Write

$$\phi_j(\sigma) = \text{tr}(A_\sigma^j X_j).$$

Evaluating $\sum_j \phi_j = 0$ on each word w of length n gives

$$\sum_j \text{tr}(A_w^j X_j) = 0.$$

Since $\text{tr}(A_w^j X_j) = \text{tr}(X_j A_w^j)$, the product-span equation gives, for every $(M_1, \dots, M_r) \in \prod_j M_{D_j}(\mathbb{C})$,

$$\sum_j \text{tr}(X_j M_j) = 0.$$

Taking only the j -th component nonzero and using nondegeneracy of the trace pairing yields $X_j = 0$, hence $\phi_j = 0$. The BNT statement follows by substituting Lemma 13.13.2.4. $\square$

13.14 Block-diagonal boundary conditions for periodic ground spaces

For a block-diagonal tensor $B = \bigoplus_{j=1}^g \mu_j A_j$ with $\mu_j \neq 0$, this section proves implications from explicit block-diagonal boundary identities to the component membership

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

These implications supply the boundary-condition step in the periodic ring used by [CPGSV21, Theorem IV.16].

Lemma 13.14.1 (Normalized BNT representatives give the large-length block intersection).

[Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Let $A^1, \dots, A^r$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent. Suppose $L_0 > 0$, and for every block j the map*

$$\Gamma_{L_0}^{A_j} : M_{D_j}(\mathbb{C}) \rightarrow (\mathbb{C}^d)^{\otimes L_0}$$

is injective. Assume also the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Set

$$N = (L_0 + 1) + (r - 1)3(L_0 + 1).$$

For every $n \geq N$, set

$$S_n := \bigvee_j G_{n+1}(A^j),$$

and the two consecutive sums are internal direct sums:

$$\bigvee_j G_{n+1}(A^j) = \bigoplus_j G_{n+1}(A^j), \quad \bigvee_j G_{n+2}(A^j) = \bigoplus_j G_{n+2}(A^j).$$

With this notation,

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. [✓ Proof](#) Lemma 13.13.2.4 gives

$$\text{span}\{(A_w^1, \dots, A_w^r) : |w| = n\} = \prod_j M_{D_j}(\mathbb{C})$$

for every $n \geq N$. Substituting this product-span equation into Lemma 13.13.1.18, together with the normalization equation, gives the displayed intersection identity. Lemma 13.13.2.5 at lengths $n + 1$ and $n + 2$ gives the two directness statements. $\square$

Lemma 13.14.2 (Normalized BNT representatives give an eventual block intersection). [Lean](#)

[✓ Stmt](#) *Let $A^1, \dots, A^r$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent. Suppose $L_0 > 0$, and for every block j the map*

$$\Gamma_{L_0}^{A_j} : M_{D_j}(\mathbb{C}) \rightarrow (\mathbb{C}^d)^{\otimes L_0}$$

is injective. Assume also the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Then there is a length N such that, for every $n \geq N$, with

$$S_n := \bigvee_j G_{n+1}(A^j),$$

one has

$$\{\psi : \forall b, \psi(-, b) \in S_n\} \cap \{\psi : \forall a, \psi(a, -) \in S_n\} = \bigvee_j G_{n+2}(A^j).$$

Proof. [✓ Proof](#) Apply Lemma 13.14.1 and take

$$N = (L_0 + 1) + (r - 1)3(L_0 + 1).$$

□

Lemma 13.14.3 (Block-diagonal boundary-condition formula). [Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{j=1}^g \mu_j A_j.$$

For block-diagonal boundary conditions X_j ,

$$\Gamma_L^B \left(\bigoplus_{j=1}^g X_j \right) = \sum_{j=1}^g \Gamma_L^{A_j}(\mu_j^L X_j).$$

Proof. [✓ Proof](#) This is the diagonal-block trace formula specialized to a block-diagonal boundary matrix. The j -th diagonal block of $\bigoplus_k X_k$ is X_j , giving the displayed sum. □

Lemma 13.14.4 (One block lies in the local parent space of the direct-sum tensor). [Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{k=1}^g \mu_k A_k.$$

If $\mu_j \neq 0$, then, for every length L ,

$$G_L(A_j) \subseteq G_L(B).$$

Proof. [✓ Proof](#) A vector in $G_L(A_j)$ has the form $\Gamma_L^{A_j}(X)$. Put $\mu_j^{-L}X$ in the j -th diagonal block of a boundary matrix for B and put zero in every other block. If $\iota_j(Z)$ denotes this block insertion, Lemma 13.14.3 gives

$$\Gamma_L^B(\iota_j(\mu_j^{-L}X)) = \sum_{k=1}^g \Gamma_L^{A_k}(\mu_k^L(\iota_j(\mu_j^{-L}X))_{kk}) = \Gamma_L^{A_j}(X).$$

Hence $\Gamma_L^{A_j}(X) \in G_L(B)$. □

Lemma 13.14.5 (Local ground space of a block-diagonal tensor). [Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{j=1}^g \mu_j A_j, \quad \mu_j \neq 0 \quad (1 \leq j \leq g).$$

Then, for every length L ,

$$G_L(B) = \bigvee_{j=1}^g G_L(A_j).$$

Proof. [✓ Proof](#) A boundary matrix X for B has diagonal blocks X_j , and block-diagonality gives

$$\Gamma_L^B(X) = \sum_{j=1}^g \Gamma_L^{A_j}(\mu_j^L X_j).$$

This proves $G_L(B) \subseteq \bigvee_j G_L(A_j)$. The reverse inclusion is Lemma 13.14.4 for each block j . $\square$

Lemma 13.14.6 (Periodic constraints for a block-diagonal tensor). [Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{j=1}^g \mu_j A_j, \quad \mu_j \neq 0 \quad (1 \leq j \leq g).$$

Set

$$S_M := \bigvee_{j=1}^g G_M(A_j).$$

If, for every $M \geq L$,

$$\mathbb{C}^d \otimes S_M \cap S_M \otimes \mathbb{C}^d = S_{M+1},$$

then

$$\mathcal{G}_{N,L}(B) \subseteq S_N.$$

Proof. [✓ Proof](#) Lemma 13.14.5 gives $G_L(B) = S_L$. Apply Lemma 13.4.5.13 to the sequence S_M . $\square$

Lemma 13.14.7 (Periodic constraints lie in the block local-space sum). [Lean](#) [✓ Stmt](#) Let $A_1, \dots, A_g$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent. Suppose $L_0 > 0$, and for every block j the map

$$\Gamma_{L_0}^{A_j} : M_{D_j}(\mathbb{C}) \rightarrow (\mathbb{C}^d)^{\otimes L_0}$$

is injective. Assume also the normalization

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

For nonzero scalars μ_j , set

$$B = \bigoplus_{j=1}^g \mu_j A_j, \quad S_M := \bigvee_{j=1}^g G_M(A_j).$$

If

$$L \geq (L_0 + 1) + (g - 1)3(L_0 + 1) + 1,$$

then, for every $N \geq L$,

$$\mathcal{G}_{N,L}(B) \subseteq S_N.$$

Proof. ✓ **Proof** Put

$$T = (L_0 + 1) + (g - 1)3(L_0 + 1).$$

For $M \geq L$, one has $M - 1 \geq T$. Applying Lemma 13.14.1 at $n = M - 1$ gives

$$\mathbb{C}^d \otimes S_M \cap S_M \otimes \mathbb{C}^d = S_{M+1}$$

for every $M \geq L$. Lemma 13.14.6 then propagates the local constraint from L to N . □

Lemma 13.14.8 (Periodic constraints and internality of the block local-space sum). Lean

✓ **Stmnt** With the hypotheses and notation of Lemma 13.14.7, set

$$S_N := \bigvee_{j=1}^g G_N(A_j).$$

Then, for every $N \geq L$,

$$\mathcal{G}_{N,L}(B) \subseteq S_N,$$

and the sum defining S_N is internal: if

$$\phi_j \in G_N(A_j), \quad \sum_{j=1}^g \phi_j = 0,$$

then $\phi_j = 0$ for every j .

Proof. ✓ **Proof** The inclusion is Lemma 13.14.7. Since $N \geq L$ and

$$L \geq (L_0 + 1) + (g - 1)3(L_0 + 1) + 1,$$

the product-span directness range of Lemma 13.13.2.5 applies at length N :

$$\text{span}\{(A_w^1, \dots, A_w^g) : |w| = N\} = \prod_j M_{D_j}(\mathbb{C}) \implies \ker \left(\bigoplus_{j=1}^g G_N(A_j) \longrightarrow (\mathbb{C}^d)^{\otimes N}, \quad (\phi_j)_j \mapsto \sum_j \phi_j \right) = 0.$$

□

Lemma 13.14.9 (Periodic parent-space vectors decompose in the local block spaces). Lean

✓ **Stmnt** With the hypotheses and notation of Lemma 13.14.7, every vector

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_{j=1}^g \mu_j A_j \right)$$

has a unique family $(\psi_j)_{j=1}^g$ such that

$$\psi = \sum_{j=1}^g \psi_j, \quad \psi_j \in G_N(A_j).$$

Proof. ✓ **Proof** Lemma 13.14.8 gives

$$\mathcal{G}_{N,L} \left(\bigoplus_{j=1}^g \mu_j A_j \right) \subseteq \bigvee_{j=1}^g G_N(A_j),$$

and the spaces $G_N(A_j)$ are independent: for every j and every $x_j \in G_N(A_j)$,

$$\sum_j x_j = 0 \implies x_j = 0.$$

Thus every $\psi \in \mathcal{G}_{N,L}(\bigoplus_j \mu_j A_j)$ has a unique decomposition $\psi = \sum_j \psi_j$ with $\psi_j \in G_N(A_j)$. □

Lemma 13.14.10 (Block-diagonal boundary conditions for periodic parent-space vectors). [Lean](#)

[✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.7, every vector

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_{j=1}^g \mu_j A_j \right)$$

admits block-diagonal boundary conditions X_j such that

$$\psi = \Gamma_N^B \left(\bigoplus_{j=1}^g X_j \right), \quad \Gamma_N^{A_j}(\mu_j^N X_j) \in G_N(A_j)$$

for every j .

Proof. [✓ Proof](#) Lemma 13.14.9 gives

$$\psi = \sum_{j=1}^g \phi_j, \quad \phi_j \in G_N(A_j).$$

Choose Y_j with $\phi_j = \Gamma_N^{A_j}(Y_j)$, and set $X_j = \mu_j^{-N} Y_j$. Since $\mu_j \neq 0$,

$$\Gamma_N^{A_j}(\mu_j^N X_j) = \phi_j.$$

The block-diagonal boundary formula then gives

$$\Gamma_N^B \left(\bigoplus_{j=1}^g X_j \right) = \sum_{j=1}^g \Gamma_N^{A_j}(\mu_j^N X_j) = \psi.$$

□

The lemma gives

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in G_N(A_j).$$

This is not enough for periodic-boundary membership when a cyclic window crosses the chosen cut. The boundary-cut comparison of Lemma 13.14.1.24 concerns the separate assertion

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

13.14.1 Cyclic windows at the boundary cut

A cyclic length- L window either stays inside the chosen linear interval ($i + L \leq N$) or crosses the boundary cut ($N < i + L$). The non-wrapping case follows from trace cyclicity, while the crossing case requires a boundary-matrix identity $\mu_j^N X_j A_\beta^j = A_\beta^j Y_j$. The lemmas below establish the single-block periodic membership $\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j)$ in both cases.

Lemma 13.14.1.1 (Cyclic restrictions of block-diagonal boundary vectors). [Lean](#) [✓ Stmt](#) Let $B = \bigoplus_{j=1}^g \mu_j A_j$, and fix block-diagonal boundary conditions X_j . For every cyclic restriction $R_{i,\tau}$ from N sites to L sites,

$$R_{i,\tau} \left(\Gamma_N^B \left(\bigoplus_{j=1}^g X_j \right) \right) = \sum_{j=1}^g R_{i,\tau} \left(\Gamma_N^{A_j}(\mu_j^N X_j) \right).$$

Proof. ✓ Proof Apply Lemma 13.14.3 at length N , then use linearity of the cyclic restriction $R_{i,\tau}$. $\square$

Lemma 13.14.1.2 (Local direct-sum constraint from block-diagonal boundary conditions). Lean ✓ Stmt Let $B = \bigoplus_{j=1}^g \mu_j A_j$, with $\mu_j \neq 0$, and suppose

$$\psi = \Gamma_N^B \left(\bigoplus_{j=1}^g X_j \right), \quad \psi \in \mathcal{G}_{N,L}(B).$$

Then every local constraint of ψ gives

$$\sum_{j=1}^g R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in \bigvee_{j=1}^g G_L(A_j).$$

Proof. ✓ Proof The hypothesis $\psi \in \mathcal{G}_{N,L}(B)$ says that the left-hand side before decomposition lies in $G_L(B)$. Lemma 13.14.1.1 rewrites this local vector as the displayed sum, and Lemma 13.14.5 identifies $G_L(B)$ with $\bigvee_j G_L(A_j)$. $\square$

Lemma 13.14.1.3 (Cyclic windows not crossing the cut). Lean ✓ Stmt Fix block-diagonal boundary conditions X_j . If the cyclic window beginning at i stays inside the chosen linear interval, $i + L \leq N$, then for every block j ,

$$R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in G_L(A_j).$$

Proof. ✓ Proof When the window stays inside the chosen linear interval, the boundary matrix remains outside the window. Write $A_{\text{left}} = A_{\tau_0}^j \cdots A_{\tau_{i-1}}^j$ and $A_{\text{right}} = A_{\tau_{i+L}}^j \cdots A_{\tau_{N-1}}^j$. Trace cyclicity gives

$$R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) = \Gamma_L^{A_j} (A_{\text{right}} \mu_j^N X_j A_{\text{left}}).$$

Hence the restricted vector lies in $G_L(A_j)$. $\square$

Lemma 13.14.1.4 (Boundary-crossing coefficient formula). Lean ✓ Stmt Assume $0 < N, L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Write $a = i + L - N$. For a local word σ of length L , set

$$\beta = \sigma_{N-i} \cdots \sigma_{L-1}, \quad \rho = \tau_a \cdots \tau_{i-1}, \quad \alpha = \sigma_0 \cdots \sigma_{N-i-1}.$$

Then the corresponding coefficient of the j -th block component is

$$R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) (\sigma) = \text{tr} \left(A_\beta^j A_\rho^j A_\alpha^j \mu_j^N X_j \right).$$

Proof. ✓ Proof Let ω be the N -site word obtained by inserting σ in the cyclic interval beginning at i and by using τ outside that interval. Since the interval crosses the cut,

$$\omega = \beta \rho \alpha.$$

Hence

$$\Gamma_N^{A_j} (\mu_j^N X_j) (\omega) = \text{tr} \left(A_\beta^j A_\rho^j A_\alpha^j \mu_j^N X_j \right).$$

This is exactly $R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) (\sigma)$. $\square$

Lemma 13.14.1.5 (Boundary-crossing trace decomposition from the local constraint). [Lean](#) [✓ Stmt](#) Assume $0 < N$, $L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Put $a = i + L - N$. For a local word σ of length L , write

$$\beta = \sigma_{N-i} \cdots \sigma_{L-1}, \quad \rho = \tau_a \cdots \tau_{i-1}, \quad \alpha = \sigma_0 \cdots \sigma_{N-i-1}.$$

These are the word coordinates obtained by opening the periodic boundary at the chosen cut, in the comparison with the source expressions involving $C_{i_1}^j$ and $D_{i_{m+1}}^j$. If

$$\sum_j R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in \bigvee_j G_L(A_j),$$

then there are matrices $C_{i,\tau}^j$ such that, for every σ ,

$$\sum_j \text{tr} \left(A_\beta^j C_{i,\tau}^j A_\alpha^j \right) = \sum_j \text{tr} \left(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_\alpha^j \right).$$

Proof. [✓ Proof](#) Decompose the local vector in $\bigvee_j G_L(A_j)$ as a sum of vectors $\Gamma_L^{A_j}(C_{i,\tau}^j)$. Evaluating at σ gives

$$\sum_j \text{tr} \left(A_{\alpha\beta}^j C_{i,\tau}^j \right).$$

Trace cyclicity rewrites this as $\sum_j \text{tr} (A_\beta^j C_{i,\tau}^j A_\alpha^j)$. On the other hand, Lemma 13.14.1.4 gives the second trace expression for the same local coefficient sum. $\square$

Lemma 13.14.1.6 (Boundary representation gives fixed-tail boundary traces). [Lean](#) [✓ Stmt](#) Assume that the physical dimension d is positive, $0 < N$, $L \leq N$, and $\mu_j \neq 0$ for every block j . Let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right), \quad \psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

For every outside word ρ of length $N - L$, there are matrices $C_{i,\rho}^j$ such that, for every word β of length $i + L - N$ before the cut and every wrapped tail word α of length $N - i$,

$$\sum_j \text{tr} \left(A_\beta^j C_{i,\rho}^j A_\alpha^j \right) = \sum_j \text{tr} \left(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_\alpha^j \right).$$

Proof. [✓ Proof](#) Since $d > 0$, extend the outside word ρ to a full outside configuration τ whose entries between the end of the interval and the cut are the word ρ . By the nonzero-weight hypothesis, Lemma 13.14.1.2 applies and puts the sum of the corresponding block restrictions in $\bigvee_j G_L(A_j)$. Applying Lemma 13.14.1.5 gives the displayed trace equality after rewriting the opened cyclic word as $\beta\rho\alpha$. $\square$

Lemma 13.14.1.7 (Boundary representation gives all fixed-tail boundary traces). [Lean](#) [✓ Stmt](#) Assume that the physical dimension d is positive, $0 < N$, $L \leq N$, and $\mu_j \neq 0$ for every block j . Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right), \quad \psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

Then there are matrices $C_{i,\rho}^j$, indexed by the boundary-crossing start i and by an outside word ρ of length $N - L$, such that whenever $N < i + L$,

$$\sum_j \operatorname{tr}(A_\beta^j C_{i,\rho}^j A_\alpha^j) = \sum_j \operatorname{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_\alpha^j)$$

for every word β of length $i + L - N$ before the cut and every wrapped tail word α of length $N - i$.

Proof. [✓ Proof](#) Apply Lemma 13.14.1.6 to each boundary-crossing start i and each outside word ρ , and choose the resulting matrices $C_{i,\rho}^j$. $\square$

Theorem 13.14.1.8 (Fixed boundary-crossing C^j, D^j comparison from the local block-sum constraint). [Lean](#) [✓ Stmt](#) Assume $0 < N$, $L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Put $a = i + L - N$ and $\rho = \tau_a \cdots \tau_{i-1}$. Suppose also that the block tensors have the common word-span property at length $N - i$. If

$$\sum_j R_{i,\tau}(\Gamma_N^A(\mu_j^N X_j)) \in \bigvee_j G_L(A_j),$$

then there are matrices $C_{i,\tau}^j$ such that, for every block j and every word β of length a ,

$$A_\beta^j C_{i,\tau}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Proof. [✓ Proof](#) Lemma 13.14.1.5 gives matrices $C_{i,\tau}^j$ with

$$\sum_j \operatorname{tr}(A_\beta^j C_{i,\tau}^j A_\alpha^j) = \sum_j \operatorname{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_\alpha^j)$$

for every word α of length $N - i$ and every word β of length a . The fixed complementary-word trace comparison, applied with X_j replaced by $\mu_j^N X_j$, gives the displayed matrix identity. $\square$

Theorem 13.14.1.9 (C^j, D^j comparison for a block-diagonal boundary representation). [Lean](#)

[✓ Stmt](#) Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right), \quad \psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

Assume that the physical alphabet is nonempty, that every μ_j is nonzero, that $0 < N$, and that $L \leq N$. For every boundary-crossing interval beginning at i , assume that the simultaneous products

$$w \mapsto (A_w^1, \dots, A_w^r), \quad |w| = N - i,$$

span the product algebra $\prod_j M_{D_j}(\mathbb{C})$. Then there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every block j , and every word β before the cut,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Proof. [✓ Proof](#) For a boundary-crossing interval beginning at i and an outside word ρ , choose a configuration whose outside segment is ρ . The local constraint on ψ gives a vector in $\bigvee_j G_L(A_j)$ by Lemma 13.14.1.2. Theorem 13.14.1.8 applied with the stated span at length $N - i$ gives the displayed C^j, D^j comparison for that interval and outside word. $\square$

Theorem 13.14.1.10 (Fixed boundary-crossing component in the local block space). [Lean](#)
[✓ Stmt](#) Assume $0 < N$, $L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Suppose that the block tensors have the common word-span property at length $N - i$ and that

$$\sum_j R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in \bigvee_j G_L(A_j).$$

Then each block component of the fixed boundary-crossing restriction lies in its local block space:

$$R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in G_L(A_j).$$

Proof. [✓ Proof](#) Theorem 13.14.1.8 gives matrices $C_{i,\tau}^j$ with

$$A_\beta^j C_{i,\tau}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Thus $E = C_{i,\tau}^j$ satisfies the boundary matrix identity required in Lemma 13.14.1.11, and the displayed membership follows for every block j . $\square$

Lemma 13.14.1.11 (Boundary-crossing cyclic intervals from the boundary matrix identity). [Lean](#) [✓ Stmt](#) Assume $0 < N$, $L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Put $a = i + L - N$. Suppose that there is a matrix E such that for every word β of length a ,

$$\mu_j^N X_j A_\beta^j A_{\tau_a}^j \cdots A_{\tau_{i-1}}^j = A_\beta^j E.$$

Then

$$R_{i,\tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in G_L(A_j).$$

Proof. [✓ Proof](#) Write a length- L word in the boundary-crossing interval as $\alpha\beta$, where α is the segment inside $\{i, \dots, N-1\}$ and β is the segment after the interval wraps past the cut, inside $\{0, \dots, a-1\}$. The full N -site word appearing in the restriction is then

$$\beta \tau_a \cdots \tau_{i-1} \alpha.$$

Trace cyclicity gives

$$\text{tr} \left(A_\beta^j A_{\tau_a}^j \cdots A_{\tau_{i-1}}^j A_\alpha^j \mu_j^N X_j \right) = \text{tr} \left(A_\alpha^j \mu_j^N X_j A_\beta^j A_{\tau_a}^j \cdots A_{\tau_{i-1}}^j \right).$$

The assumed boundary matrix identity changes the last expression to

$$\text{tr} \left(A_\alpha^j A_\beta^j E \right),$$

which is the coefficient of $\Gamma_L^{A_j}(E)$ at the word $\alpha\beta$. Hence the restricted vector lies in $G_L(A_j)$. $\square$

Lemma 13.14.1.12 (Last crossing-window component trace form). [Lean](#) [✓ Stmt](#) Write $N = M + 1$ and let the local length be $m + 2 \leq M + 1$. Consider the cyclic interval beginning at the last site M . For an outside configuration τ , put

$$\rho = \tau_{m+1} \cdots \tau_{M-1}, \quad C_a^j = A_\rho^j A_a^j (\mu_j^{M+1} X_j).$$

Then the j -th component of the last crossing-window restriction has the left-boundary trace form

$$R_{M,\tau} \left(\Gamma_{M+1}^{A_j} (\mu_j^{M+1} X_j) \right) = \Phi^j, \quad \Phi^j(a, w, b) = \text{tr} \left(A_b^j C_a^j A_w^j \right).$$

Proof. ✓ **Proof** The cyclic interval beginning at M reads the last site first and then wraps to the sites $0, \dots, m$. Thus, for the local word awb , the full word in the trace is

$$wb\rho a.$$

Hence the j -th coefficient is

$$\mathrm{tr}(A_w^j A_b^j A_\rho^j A_a^j (\mu_j^{M+1} X_j)).$$

Trace cyclicity rotates this to

$$\mathrm{tr}(A_b^j A_\rho^j A_a^j (\mu_j^{M+1} X_j) A_w^j),$$

which is the displayed left-boundary trace form. □

Lemma 13.14.1.13 (Last crossing-window trace form). Lean ✓ **Stmt** Write $N = M + 1$ and let the local length be $m + 2 \leq M + 1$. Consider the cyclic interval beginning at the last site M . For an outside configuration τ , put

$$\rho = \tau_{m+1} \cdots \tau_{M-1}, \quad C_a^j = A_\rho^j A_a^j (\mu_j^{M+1} X_j).$$

Then the block sum of the last crossing-window restrictions has the left-boundary trace form

$$\sum_j R_{M,\tau}(\Gamma_{M+1}^{A_j}(\mu_j^{M+1} X_j)) = \sum_j \Phi^j, \quad \Phi^j(a, w, b) = \mathrm{tr}(A_b^j C_a^j A_w^j).$$

Proof. ✓ **Proof** Sum Lemma 13.14.1.12 over the block index. □

Lemma 13.14.1.14 (Last crossing-window coefficient comparison). Lean ✓ **Stmt** In the setting of Lemma 13.14.1.13, assume that the length- m simultaneous word tuples span the product algebra and that

$$\sum_j R_{M,\tau}(\Gamma_{M+1}^{A_j}(\mu_j^{M+1} X_j)) \in \bigvee_j G_{m+2}(A^j).$$

Then there are matrices E_j such that, for every block j and all boundary letters a, b ,

$$A_b^j(A_\rho^j A_a^j (\mu_j^{M+1} X_j)) = A_b^j E_j A_a^j.$$

Proof. ✓ **Proof** Lemma 13.14.1.13 writes the local block sum in the form

$$\Phi(a, w, b) = \sum_j \mathrm{tr}(A_b^j C_a^j A_w^j), \quad C_a^j = A_\rho^j A_a^j (\mu_j^{M+1} X_j).$$

Applying Lemma 13.13.1.3 gives the displayed blockwise equations. □

Lemma 13.14.1.15 (Last crossing-window component membership). Lean ✓ **Stmt** In the setting of Lemma 13.14.1.13, assume that the length- m simultaneous word tuples span the product algebra and that

$$\sum_j R_{M,\tau}(\Gamma_{M+1}^{A_j}(\mu_j^{M+1} X_j)) \in \bigvee_j G_{m+2}(A^j).$$

Then, for every block j ,

$$R_{M,\tau}(\Gamma_{M+1}^{A_j}(\mu_j^{M+1} X_j)) \in G_{m+2}(A^j).$$

Proof. ✓ Proof Lemma 13.14.1.14 gives matrices E_j with

$$A_b^j C_a^j = A_b^j E_j A_a^j.$$

By Lemma 13.14.1.12, the j -th cyclic restriction is the left-boundary component determined by C_a^j . The displayed identity makes that component equal to $\Gamma_{m+2}^{A_j}(E_j)$, hence it lies in $G_{m+2}(A^j)$. □

Lemma 13.14.1.16 (Last crossing-window coefficients from a block-diagonal chain vector).

Lean ✓ Stmt Let $B = \bigoplus_j \mu_j A^j$, and assume $\mu_j \neq 0$ for every block j . Suppose

$$\psi = \Gamma_{M+1}^B \left(\bigoplus_j X_j \right), \quad \psi \in \mathcal{G}_{M+1, m+2}(B).$$

Assume also that the length- m simultaneous word tuples span the product algebra. Then, for every outside configuration τ , with $\rho = \tau_{m+1} \cdots \tau_{M-1}$, there are matrices E_j such that

$$A_b^j (A_\rho^j A_a^j (\mu_j^{M+1} X_j)) = A_b^j E_j A_a^j$$

for every block j and all boundary letters a, b .

Proof. ✓ Proof The local constraint of ψ at the cyclic interval beginning at M gives

$$\sum_j R_{M, \tau} \left(\Gamma_{M+1}^{A_j} (\mu_j^{M+1} X_j) \right) \in \bigvee_j G_{m+2}(A^j)$$

by Lemma 13.14.1.2. Applying Lemma 13.14.1.14 gives the displayed equations. □

Lemma 13.14.1.17 (Last crossing-window component membership from a chain vector). Lean

✓ Stmt Let $B = \bigoplus_j \mu_j A^j$, and assume $\mu_j \neq 0$ for every block j . Suppose

$$\psi = \Gamma_{M+1}^B \left(\bigoplus_j X_j \right), \quad \psi \in \mathcal{G}_{M+1, m+2}(B).$$

If the length- m simultaneous word tuples span the product algebra, then for every outside configuration τ and every block j ,

$$R_{M, \tau} \left(\Gamma_{M+1}^{A_j} (\mu_j^{M+1} X_j) \right) \in G_{m+2}(A^j).$$

Proof. ✓ Proof Lemma 13.14.1.2 applied to the window beginning at the last site gives

$$\sum_j R_{M, \tau} \left(\Gamma_{M+1}^{A_j} (\mu_j^{M+1} X_j) \right) \in \bigvee_j G_{m+2}(A^j).$$

Lemma 13.14.1.15 then applies to each block component. □

Lemma 13.14.1.18 (Boundary-crossing intervals from a right boundary-matrix identity). Lean

✓ Stmt Assume $0 < N$, $L \leq N$, and let the cyclic interval beginning at i cross the boundary cut, so $N < i + L$. Put $a = i + L - N$. Suppose that there is a matrix Y such that for every word β of length a ,

$$\mu_j^N X_j A_\beta^j = A_\beta^j Y.$$

Then

$$R_{i, \tau} \left(\Gamma_N^{A_j} (\mu_j^N X_j) \right) \in G_L(A_j).$$

Proof. ✓ Proof Let

$$E := Y A_{\tau_a}^j \cdots A_{\tau_{i-1}}^j.$$

Then, for every word β on the segment $0, \dots, a-1$,

$$\mu_j^N X_j A_\beta^j A_{\tau_a}^j \cdots A_{\tau_{i-1}}^j = A_\beta^j E.$$

Lemma 13.14.1.11 gives the stated local constraint. □

Lemma 13.14.1.19 (Boundary-crossing cyclic intervals suffice). Lean ✓ Stmt Assume $0 < N$ and $L \leq N$. Fix block-diagonal boundary conditions X_j . Suppose that, for every block j , every cyclic interval beginning at i with $N < i + L$, and every configuration τ ,

$$R_{i,\tau}(\Gamma_N^{A_j}(\mu_j^N X_j)) \in G_L(A_j).$$

Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof By definition, membership in $\mathcal{G}_{N,L}(A_j)$ is the collection of all cyclic length- L local constraints. For an interval beginning at i , either $i + L \leq N$ or $N < i + L$. In the first case use Lemma 13.14.1.3. In the second case use the displayed hypothesis. □

Lemma 13.14.1.20 (Right boundary-matrix identities give single-block periodic-boundary constraints). Lean ✓ Stmt Assume $0 < N$ and $L \leq N$. Fix block-diagonal boundary conditions X_j . Suppose that, for every block j , every boundary-crossing interval beginning at i , and every configuration τ , there is a matrix $Y_{j,i,\tau}$ such that, for every word β of length $a = i + L - N$,

$$\mu_j^N X_j A_\beta^j = A_\beta^j Y_{j,i,\tau}.$$

Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Lemma 13.14.1.18 turns the displayed boundary identity into the local constraint for each boundary-crossing interval. Lemma 13.14.1.19 then yields

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j),$$

handling both boundary-crossing intervals $N < i + L$ and non-crossing intervals $i + L \leq N$. □

Lemma 13.14.1.21 (Spanning matrix identities give single-block constraints). Lean ✓ Stmt Assume $0 < N$, $L \leq N$, and that for every block j , the length- $N - L$ word products of A_j span the full matrix algebra. Suppose that, for every block j , every boundary-crossing interval beginning at i , and every outside word ρ of length $N - L$, there is a matrix $E_{j,i,\rho}$ such that for every word β before the cut, of length $i + L - N$,

$$\mu_j^N X_j A_\beta^j A_\rho^j = A_\beta^j E_{j,i,\rho}.$$

Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof For fixed j, i , apply Theorem 13.7.2 to the family

$$Z_\beta = \mu_j^N X_j A_\beta^j, \quad F_\beta = A_\beta^j.$$

Since the length- $N - L$ outside-word products span the full matrix algebra, there is a single matrix $Y_{j,i}$ such that, for every β ,

$$\mu_j^N X_j A_\beta^j = A_\beta^j Y_{j,i}.$$

Lemma 13.14.1.20 then gives the periodic-boundary constraints for each block. □

Lemma 13.14.1.22 (Matrix identities under block injectivity give single-block constraints).

Lean ✓ Stmt Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies the normalization equation

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Suppose also that the boundary-crossing matrix identities of Lemma 13.14.1.21 hold for every outside word ρ of length $N - L$. Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Since $L + L_0 \leq N$, the complementary length satisfies $L_0 \leq N - L$. Theorem 8.11.4 therefore gives, for every block j ,

$$S_{N-L}(A_j) = M_{D_j}(\mathbb{C}).$$

Apply Lemma 13.14.1.21. □

Lemma 13.14.1.23 (Normalization for words). Lean ✓ Stmt If

$$\sum_a A_a A_a^\dagger = I,$$

then, for every length M ,

$$\sum_\rho A_\rho A_\rho^\dagger = I,$$

where ρ ranges over all words of length M .

Proof. ✓ Proof The case $M = 0$ is the identity matrix. For the induction step, write a word of length $M + 1$ as $a\rho$. Since $A_{a\rho} = A_a A_\rho$,

$$\sum_{a,\rho} A_a A_\rho A_\rho^\dagger A_a^\dagger = \sum_a A_a \left(\sum_\rho A_\rho A_\rho^\dagger \right) A_a^\dagger = \sum_a A_a A_a^\dagger = I.$$

□

Lemma 13.14.1.24 (Compatibility hypotheses under block injectivity give single-block constraints). Lean ✓ Stmt Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Fix block-diagonal boundary conditions X_j . For every boundary-crossing interval beginning at i , suppose there are matrices $C_{i,\rho}^j$, indexed by outside words ρ of length $N - L$, such that for every word β before the cut, of length $i + L - N$,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

After opening the periodic boundary at the chosen cut, this is the source comparison

$$A_{i_{m+1}}^j C_{i_1}^j = D_{i_{m+1}}^j A_{i_1}^j, \quad D_\beta^j = (\mu_j^N X_j) A_\beta^j.$$

Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof The normalization equation gives, by induction on word length,

$$\sum_\rho A_\rho^j A_\rho^{j\dagger} = I$$

for words ρ of length $N - L$. Applying Lemma 13.13.1.10 with $X = \mu_j^N X_j$ gives, for every outside word ρ , a matrix $E_{j,i,\rho}$ such that for every word β before the cut,

$$((\mu_j^N X_j) A_\beta^j) A_\rho^j = A_\beta^j E_{j,i,\rho}.$$

Lemma 13.14.1.22 then gives the periodic-boundary constraint for each block. □

Lemma 13.14.1.25 (Boundary representation and crossing constraints give periodic components). Lean ✓ Stmt Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right), \quad \psi = \Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

If every boundary-crossing interval has the corresponding tail-word spanning property, then there are block-diagonal boundary conditions X_j with the displayed representation of ψ and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Theorem 13.14.1.9 supplies matrices $C_{i,\rho}^j$ satisfying

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Lemma 13.14.1.24 then gives the single-block periodic constraints. □

Lemma 13.14.1.26 (Boundary comparison implies trace comparison). Lean ✓ Stmt Fix block-diagonal boundary matrices X_j . Suppose the opened-boundary matrices $C_{i,\rho}^j$ satisfy, for every boundary-crossing interval i , every outside word ρ , and every word β before the cut,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Then for every middle word w ,

$$\sum_j \text{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \text{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j).$$

Proof. ✓ **Proof** Substituting the displayed comparison identity in each summand gives

$$\mathrm{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \mathrm{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j).$$

Summing over j gives the conclusion. □

Lemma 13.14.1.27 (Boundary trace comparisons give single-block constraints). Lean ✓ **Stmt** Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Assume also that the simultaneous tuples $w \mapsto (A_w^1, \dots, A_w^g)$ at the middle-word length m span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Fix complex numbers μ_j and block-diagonal boundary conditions X_j . For every boundary-crossing interval beginning at i , suppose there are matrices $C_{i,\rho}^j$, indexed by outside words ρ of length $N - L$, such that for every word β before the cut, outside word ρ , and middle word w ,

$$\sum_j \mathrm{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \mathrm{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j).$$

Then, for every block j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ **Proof** Applying Lemma 13.13.1.9 with the same scalars μ_j and boundary matrices X_j , for every block j and every boundary-crossing interval i , this gives a matrix E_i^j such that, for every word β before the cut,

$$(\mu_j^N X_j) A_\beta^j = A_\beta^j E_i^j.$$

Hence for every outside word ρ , taking $E_{j,i,\rho} = E_i^j A_\rho^j$ gives

$$((\mu_j^N X_j) A_\beta^j) A_\rho^j = A_\beta^j E_{j,i,\rho}.$$

These are precisely the block-diagonal boundary-condition equations of Lemma 13.14.1.22, which gives the periodic-boundary constraint for each block. □

Lemma 13.14.1.28 (Boundary trace comparisons with a block-diagonal representation). Lean ✓ **Stmt** Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Assume also that the simultaneous tuples $w \mapsto (A_w^1, \dots, A_w^g)$ at the middle-word length m span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Let

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

for some block-diagonal boundary conditions X_j . Suppose that whenever ψ has such a representation, there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every word β of length $i + L - N$ before the cut, and every word w of length m ,

$$\sum_j \mathrm{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \mathrm{tr}(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j).$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Choose a block-diagonal boundary representation of ψ . For that representation, the boundary trace comparison supplies the matrices $C_{i,\rho}^j$. Applying Lemma 13.14.1.27 gives the desired containment $\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j)$ for every block j . □

Lemma 13.14.1.29 (C^j boundary comparison with a block-diagonal representation). Lean
✓ Stmt Assume $0 < N$, $L \leq N$, and $L + L_0 \leq N$. Suppose each block A_j is L_0 -block-injective and satisfies

$$\sum_a A_a^j A_a^{j\dagger} = I.$$

Assume also that the simultaneous tuples $w \mapsto (A_w^1, \dots, A_w^g)$ at the middle-word length m span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Let

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

for some block-diagonal boundary conditions X_j . Suppose that whenever ψ has such a representation, there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, and every word β of length $i + L - N$ before the cut,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Choose a block-diagonal boundary representation of ψ . For that representation, Lemma 13.14.1.26 gives, for every boundary-crossing interval i , outside word ρ , word β before the cut, and middle word w ,

$$\sum_j \text{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \text{tr}((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j.$$

Applying Lemma 13.14.1.28 to these trace comparisons gives, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

□

Lemma 13.14.1.30 (Boundary trace comparisons give periodic-boundary single-block states).

[Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.10, assume in addition that $L + L_0 \leq N$. Assume also that, for some middle-word length m , the simultaneous tuples $w \mapsto (A_w^1, \dots, A_w^g)$, where w ranges over words of length m in $\{0, \dots, d-1\}$, span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right).$$

Suppose that whenever

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every word β before the cut, and every word w of length m ,

$$\sum_j \text{tr} \left(A_\beta^j C_{i,\rho}^j A_w^j \right) = \sum_j \text{tr} \left(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j \right).$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. [✓ Proof](#) By Lemma 13.14.10 there are matrices X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

Lemma 13.14.1.28 applies to this representation and the assumed equality of boundary trace expressions. Thus, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

□

Lemma 13.14.1.31 (Basis-of-normal-tensors product span gives boundary-trace periodic components). [Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.10, assume in addition that $L + L_0 \leq N$ and

$$L \geq (L_0 + 1) + (g - 1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))) + 1.$$

Put

$$m = (L_0 + 1) + (g - 1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))).$$

Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right).$$

Suppose that whenever

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every word β before the cut, and every word w of length m ,

$$\sum_j \text{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \text{tr}((\mu_j^N X_j) A_\beta^j A_\rho^j A_w^j).$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Lemma 13.13.2.4 gives

$$\text{span}\{(A_w^1, \dots, A_w^g) : |w| = m\} = \prod_j M_{D_j}(\mathbb{C}).$$

Applying Lemma 13.14.1.30 with this middle-word span gives the stated block-diagonal boundary conditions and the periodic-boundary constraint in each block. $\square$

Lemma 13.14.1.32 (C^j boundary comparison gives periodic single-block states). Lean ✓ Stmt
 With the hypotheses and notation of Lemma 13.14.10, assume in addition that $L + L_0 \leq N$. Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right).$$

Suppose that whenever

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, and every word β before the cut,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ **Proof** By Lemma 13.14.10 there are matrices X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

For these boundary conditions, the assumed C^j boundary comparison supplies the matrices $C_{i,\rho}^j$. Lemma 13.14.1.24 gives, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

□

Lemma 13.14.1.33 (Matrix identities give periodic single-block states). Lean ✓ **Stmt** With the hypotheses and notation of Lemma 13.14.10, assume in addition that $L + L_0 \leq N$. Let

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right).$$

Suppose that whenever

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

the corresponding boundary conditions satisfy the block-diagonal boundary-condition equations: for every boundary-crossing interval i , every word β before the cut, and every outside word ρ of length $N - L$,

$$\mu_j^N X_j A_\beta^j A_\rho^j = A_\beta^j E_{j,i,\rho}.$$

Then there are block-diagonal boundary conditions X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ **Proof** By Lemma 13.14.10 there are matrices X_j with

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right).$$

These X_j satisfy the assumed identities, so Lemma 13.14.1.22 gives, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

□

13.14.2 Periodic-boundary ground space for a block-diagonal tensor

The periodic ground-space theorem states that, for sufficiently long periodic systems, the periodic-boundary ground space of a parent Hamiltonian is generated by a basis of normal tensors of the initial tensor A [CPGSV21, Theorem IV.16]. The lemmas in this subsection are conditional reductions toward that statement. Their hypotheses explicitly include a boundary inclusion, a block-diagonal boundary representation, or periodic-boundary membership for the single-block states; in the source theorem these conditions are obtained from the boundary-condition comparison in the periodic ring. The source proof [PGVWC07] writes this comparison with $C_{i_1}^j$, $D_{i_{m+1}}^j$, and the normalized matrix E^j ; the symbols β and ρ below are the words obtained after opening the periodic boundary at the chosen cut. The source theorem writes the number of blocks as b and assumes $b \geq 2$; here the same number is denoted by g . In these coordinates the source comparison input becomes

$$A_{\beta}^j C_{i,\rho}^j = (\mu_j^N X_j A_{\beta}^j) A_{\rho}^j,$$

which is the form obtained after opening the boundary from $A_{i_{m+1}}^j C_{i_1}^j = D_{i_{m+1}}^j A_{i_1}^j$ with $D_{\beta}^j = (\mu_j^N X_j) A_{\beta}^j$. Once this compatibility is known, the normalized E^j calculation is the matrix identity of Lemma 13.13.1.9; the local boundary form used after opening the periodic boundary is Lemma 13.13.1.10.

Lemma 13.14.2.1 (The block-diagonal periodic-boundary space lies between two block sums).

[Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.7, one has

$$\bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j) \subseteq \mathcal{G}_{N,L}\left(\bigoplus_{j=1}^g \mu_j A_j\right) \subseteq \bigvee_{j=1}^g G_N(A_j),$$

and the sum defining the right-hand side is internal.

Proof. [✓ Proof](#) The first inclusion is Lemma 13.12.8. The second inclusion and internal directness are Lemma 13.14.8. $\square$

Lemma 13.14.2.2 (Boundary inclusion gives the block-diagonal periodic-boundary equality).

[Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{j=1}^g \mu_j A_j, \quad \mu_j \neq 0 \quad (1 \leq j \leq g).$$

If the periodic-boundary comparison with block-diagonal boundary conditions gives

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j),$$

then

$$\mathcal{G}_{N,L}(B) = \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

Proof. [✓ Proof](#) The reverse inclusion is Lemma 13.12.8; combine the two inclusions. $\square$

Lemma 13.14.2.3 (Componentwise periodic-boundary decomposition gives the reverse inclusion). [Lean](#) [✓ Stmt](#) Let

$$B = \bigoplus_{j=1}^g \mu_j A_j.$$

Suppose that every vector $\psi \in \mathcal{G}_{N,L}(B)$ can be written as

$$\psi = \sum_{j=1}^g \psi_j, \quad \psi_j \in \mathcal{G}_{N,L}(A_j).$$

Then

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof For $\psi \in \mathcal{G}_{N,L}(B)$, choose the displayed decomposition. Each summand lies in the corresponding subspace $\mathcal{G}_{N,L}(A_j)$, hence their finite sum lies in $\bigvee_j \mathcal{G}_{N,L}(A_j)$. $\square$

Lemma 13.14.2.4 (Block-diagonal boundary vectors in the periodic-boundary block sum).

Lean ✓ Stmt Let $B = \bigoplus_{j=1}^g \mu_j A_j$. Fix block-diagonal boundary conditions X_j . Suppose that, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Then

$$\Gamma_N^B\left(\bigoplus_{j=1}^g X_j\right) \in \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof The block-diagonal boundary formula gives

$$\Gamma_N^B\left(\bigoplus_{j=1}^g X_j\right) = \sum_{j=1}^g \Gamma_N^{A_j}(\mu_j^N X_j).$$

Each summand belongs to the corresponding block periodic-boundary space by assumption, so the finite sum lies in the linear span of these spaces. $\square$

Lemma 13.14.2.5 (Block-diagonal boundary representation gives the reverse inclusion). Lean

✓ Stmt Let $B = \bigoplus_{j=1}^g \mu_j A_j$. Suppose every $\psi \in \mathcal{G}_{N,L}(B)$ admits block-diagonal boundary conditions X_j such that

$$\psi = \Gamma_N^B\left(\bigoplus_{j=1}^g X_j\right),$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Then

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

Proof. ✓ Proof Apply Lemma 13.14.2.4 to the block-diagonal boundary representation of each $\psi \in \mathcal{G}_{N,L}(B)$. $\square$

Lemma 13.14.2.6 (Block-diagonal boundary representation reduces to periodic-boundary equality).

Lean ✓ Stmt With the hypotheses and notation of Lemma 13.14.7, suppose every vector $\psi \in \mathcal{G}_{N,L}(B)$ has block-diagonal boundary conditions X_j such that

$$\psi = \Gamma_N^B\left(\bigoplus_{j=1}^g X_j\right),$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Then

$$\mathcal{G}_{N,L}(B) = \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j),$$

and the sum defining $\bigvee_j G_N(A_j)$ is internal.

Proof. ✓ Proof The block-diagonal boundary representation gives

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j)$$

by Lemma 13.14.2.5. The equality follows from Lemma 13.14.2.2; the internality statement is Lemma 13.14.8. □

Lemma 13.14.2.7 (Complementary-word matrix identities reduce to periodic-boundary equality). Lean ✓ Stmt *With the hypotheses and notation of Lemma 13.14.2.6, assume in addition that $L + L_0 \leq N$. Assume further that, for every*

$$\psi \in \mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right)$$

and every block-diagonal boundary representation

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j}\left(\bigoplus_j X_j\right),$$

the complementary-word matrix identities hold: for every block j , boundary-crossing interval i , word β before the cut, and outside word ρ of length $N - L$, there is a matrix $E_{j,i,\rho}$ such that

$$\mu_j^N X_j A_{\beta}^j A_{\rho}^j = A_{\beta}^j E_{j,i,\rho}.$$

Then

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the N -site ground spaces $G_N(A_j) = \mathcal{G}_{N,N}(A_j)$ have internal sum.

Proof. ✓ Proof For each ψ , Lemma 13.14.1.33 gives matrices X_j with

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j}\left(\bigoplus_j X_j\right)$$

and, for every j , the membership

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j)$$

holds. Lemma 13.14.2.6 then gives

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the internality of the sum of the $G_N(A_j)$. □

Lemma 13.14.2.8 (C^j, D^j boundary-condition comparisons reduce to periodic-boundary equality). [Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.2.6, assume in addition that $L + L_0 \leq N$. Assume further that, for every

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right)$$

and every block-diagonal boundary representation

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, and every word β before the cut,

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

This is the C^j, D^j compatibility input; the subsequent normalized E^j identities follow from Lemma 13.13.1.10. Then

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the N -site ground spaces $G_N(A_j) = \mathcal{G}_{N,N}(A_j)$ have internal sum.

Proof. [✓ Proof](#) For each ψ , Lemma 13.14.1.32 gives matrices X_j with

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Lemma 13.14.2.6 then gives

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the internality of the sum of the $G_N(A_j)$. □

Lemma 13.14.2.9 (C^j, D^j comparisons give the periodic block ground-space equality). [Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.2.8, one has

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j).$$

This records only the ground-space equality conclusion obtained after the C^j, D^j boundary-condition comparison input in [PGVWC07, Theorem 12, proof lines 1446–1456]; the local normalized E^j calculation is Lemma 13.13.1.10. The independence of the N -site spaces is a separate conclusion of Lemma 13.14.2.8. The statement is still in the length- L_0 injectivity range displayed in that lemma, not the shorter range in [PGVWC07, Theorem 12].

Proof. ✓ Proof The claim is the equality part of Lemma 13.14.2.8. □

Lemma 13.14.2.10 (C^j, D^j comparisons at boundary-crossing intervals reduce to periodic-boundary equality). Lean ✓ Stmt *With the hypotheses and notation of Lemma 13.14.2.6, assume in addition that $L + L_0 \leq N$. Assume also that, for every boundary-crossing interval beginning at i , the simultaneous products*

$$w \mapsto (A_w^1, \dots, A_w^g), \quad |w| = N - i,$$

span the product algebra $\prod_j M_{D_j}(\mathbb{C})$. Then

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the N -site ground spaces $G_N(A_j) = \mathcal{G}_{N,N}(A_j)$ have internal sum. The present statement is the span-dependent consequence, after opening the boundary, of the C^j, D^j compatibility input in [PGVWC07, Theorem 12, proof lines 1436–1451] in the length- L_0 injectivity range

$$L \geq (L_0 + 1) + 3(g - 1)(L_0 + 1) + 1.$$

The source theorem has the shorter bound

$$L \geq 3(g - 1)(L_0 + 1) + 1.$$

Proof. ✓ Proof Lemma 13.14.2.6 reduces the claim to producing, for each ψ , block-diagonal boundary conditions X_j whose single-block components lie in $\mathcal{G}_{N,L}(A_j)$. The crossing comparison theorem supplies this: it first uses the boundary-comparison-free BNT representation to obtain X_j , and then uses the stated tail-word spans and Theorem 13.14.1.9 to produce matrices $C_{i,\rho}^j$ satisfying

$$A_\beta^j C_{i,\rho}^j = ((\mu_j^N X_j) A_\beta^j) A_\rho^j.$$

Lemma 13.14.1.24 then gives

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j)$$

for every j . Applying Lemma 13.14.2.6 to these boundary conditions gives the displayed equality and internality. □

Lemma 13.14.2.11 (Boundary-crossing comparisons give the periodic block ground-space equality). Lean ✓ Stmt *With the hypotheses and notation of Lemma 13.14.2.10, one has*

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j).$$

This records the ground-space equality conclusion of the conditional crossing-span version above. It is the corresponding part of [PGVWC07, Theorem 12], but only under the visible tail-word span hypothesis and in the longer stated range. The independence of the N -site spaces is a separate conclusion of Lemma 13.14.2.10. Thus the statement remains conditional on the short tail-word spans; it is not the unconditional boundary-condition comparison in the source theorem.

Proof. ✓ Proof The claim is the equality part of Lemma 13.14.2.10. □

Lemma 13.14.2.12 (Boundary trace comparisons reduce to periodic-boundary equality). [Lean](#)
[✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.2.6, assume in addition that $L + L_0 \leq N$. Assume also that, for some middle-word length m , the simultaneous tuples $w \mapsto (A_w^1, \dots, A_w^g)$, where w ranges over words of length m in $\{0, \dots, d-1\}$, span the full product algebra $\prod_j M_{D_j}(\mathbb{C})$. Assume further that, for every

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right)$$

and every block-diagonal boundary representation

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every word β before the cut, and every word w of length m ,

$$\sum_j \text{tr} \left(A_\beta^j C_{i,\rho}^j A_w^j \right) = \sum_j \text{tr} \left(((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j \right).$$

Then

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the N -site ground spaces $G_N(A_j) = \mathcal{G}_{N,N}(A_j)$ have internal sum.

Proof. [✓ Proof](#) For each ψ , Lemma 13.14.1.30 gives matrices X_j with

$$\psi = \Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right)$$

and, for every j ,

$$\Gamma_N^{A_j}(\mu_j^N X_j) \in \mathcal{G}_{N,L}(A_j).$$

Lemma 13.14.2.6 then gives

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the internality of the sum of the $G_N(A_j)$. □

Lemma 13.14.2.13 (Basis-of-normal-tensors product span gives boundary-trace ground-space equality). [Lean](#) [✓ Stmt](#) With the hypotheses and notation of Lemma 13.14.2.6, assume in addition that $L + L_0 \leq N$ and

$$L \geq (L_0 + 1) + (g - 1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))) + 1.$$

Put

$$m = (L_0 + 1) + (g - 1)((L_0 + 1) + ((L_0 + 1) + (L_0 + 1))).$$

Assume that, for every

$$\psi \in \mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right)$$

and every block-diagonal boundary representation

$$\psi = \Gamma_N^{\oplus_j \mu_j A_j} \left(\bigoplus_j X_j \right),$$

there are matrices $C_{i,\rho}^j$ such that, for every boundary-crossing interval i , every outside word ρ of length $N - L$, every word β before the cut, and every word w of length m ,

$$\sum_j \text{tr}(A_\beta^j C_{i,\rho}^j A_w^j) = \sum_j \text{tr}((\mu_j^N X_j) A_\beta^j) A_\rho^j A_w^j.$$

Then

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) = \bigvee_j \mathcal{G}_{N,L}(A_j),$$

and the N -site ground spaces $G_N(A_j) = \mathcal{G}_{N,N}(A_j)$ have internal sum.

Proof. [✓ Proof](#) Lemma 13.13.2.4 gives

$$\text{span}\{(A_w^1, \dots, A_w^g) : |w| = m\} = \prod_j M_{D_j}(\mathbb{C}).$$

Applying Lemma 13.14.2.12 with this middle-word span gives the equality of periodic-boundary ground spaces and the internality of the sum of the N -site spaces. $\square$

Lemma 13.14.2.14 (Boundary decomposition reduces to periodic-boundary equality). [Lean](#)
[✓ Stmt](#) Let $A_1, \dots, A_g$ be normalized representatives of a basis of normal tensors: each representative is irreducible, the family is left-canonical, the self-overlaps are normalized, and no two distinct same-dimensional representatives are gauge-phase equivalent, and let

$$B = \bigoplus_{j=1}^g \mu_j A_j, \quad \mu_j \neq 0.$$

Assume each block is injective at length L_0 , assume $\sum_a A_a^j A_a^{j\dagger} = I$ for every j , and suppose

$$L \geq (L_0 + 1) + (g - 1)3(L_0 + 1) + 1, \quad N \geq L.$$

If every vector $\psi \in \mathcal{G}_{N,L}(B)$ can be written as

$$\psi = \sum_{j=1}^g \psi_j, \quad \psi_j \in \mathcal{G}_{N,L}(A_j),$$

then

$$\mathcal{G}_{N,L}(B) = \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j),$$

and the sum $\bigvee_j \mathcal{G}_{N,L}(A_j)$ is internal.

Proof. [✓ Proof](#) Lemma 13.14.2.3 gives

$$\mathcal{G}_{N,L}(B) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

The blockwise inclusion gives the reverse containment, hence equality. The internality of $\bigvee_j \mathcal{G}_{N,L}(A_j)$ is the directness conclusion of Lemma 13.14.8. $\square$

13.15 Periodic-boundary ground space generated by a basis of normal tensors

For a block-diagonal tensor $\bigoplus_j \mu_j A_j$ with $\mu_j \neq 0$, this section relates three ground-space statements. Each block A_j contributes the local space $G_m(A_j)$ on an open length- m segment, and we write $S_m := \sum_{j=1}^g G_m(A_j)$ for their linear sum; internal directness is proved only under the BNT separation hypotheses and the stated length ranges. On the N -site ring the parent Hamiltonian has the periodic-boundary ground space $\mathcal{G}_{N,L}(\bigoplus_j \mu_j A_j)$. The results below establish the open-segment recursion $\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d = S_{m+1}$ and, conditional on the block-diagonal boundary-condition comparison in the periodic ring, reduce the periodic-boundary ground space to the span of the vectors $V^{(N)}(A_j)$ associated to the chosen basis of normal tensors:

$$\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

Theorem 13.15.1 (Block-diagonal local recursion and boundary-condition comparison). Not ready
Let $A_1, \dots, A_g$ be the blocks in a block-injective canonical form, and assume that their MPV families are pairwise different:

$$j \neq k \implies \exists n \geq 1, V^{(n)}(A_j) \neq V^{(n)}(A_k).$$

Let L_0 be a common injectivity length. Assume the multi-block range used in [PGVWC07, Theorem 12]:

$$g \geq 2, \quad L_0 \geq 1, \quad L \geq 3(g-1)(L_0+1)+1.$$

Set

$$S_m := \bigoplus_{j=1}^g G_m(A_j).$$

The open-segment recursion in [PGVWC07, Theorem 12] says that, for every $m \geq L$,

$$\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d = S_{m+1}.$$

Let $H_{S_L, N}$ denote the periodic parent Hamiltonian whose local ground space is S_L . The periodic-boundary conclusion of the same theorem is

$$\ker H_{S_L, N} = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$$

for every $N \geq L + L_0$.

Proof. Not ready The source proof takes a vector ψ in the left-hand side and writes its two boundary descriptions as

$$\psi = \sum_{j=1}^g \alpha_j = \sum_{j=1}^g \beta_j, \quad \alpha_j \in \mathbb{C}^d \otimes G_m(A_j), \quad \beta_j \in G_m(A_j) \otimes \mathbb{C}^d.$$

In the source notation, the two components have boundary matrices $C_{i_1}^j$ and $D_{i_{m+1}}^j$: the first family leaves the initial physical index open, and the second family leaves the final physical index open.

$$\alpha_j = \sum_{i_1, \dots, i_{m+1}} \text{tr}(A_{i_{m+1}}^j C_{i_1}^j A_{i_2}^j \cdots A_{i_m}^j) |i_1 \cdots i_{m+1}\rangle,$$

$$\beta_j = \sum_{i_1, \dots, i_{m+1}} \text{tr}(D_{i_{m+1}}^j A_{i_1}^j A_{i_2}^j \cdots A_{i_m}^j) |i_1 \cdots i_{m+1}\rangle.$$

The direct-sum separation lemma used in [PGVWC07, Theorem 12], together with injectivity of each block, gives the blockwise matrix identity

$$A_{i_{m+1}}^j C_{i_1}^j = D_{i_{m+1}}^j A_{i_1}^j$$

for every j , i_1 , and i_{m+1} . For fixed j , set

$$E^j := \sum_a C_a^j A_a^{j\dagger}.$$

The printed formula in [PGVWC07, Theorem 12, proof lines 1449–1451] omits the adjoint on A_a^j ; the adjointed form is forced by the displayed normalization below. In the normalization used in the source proof, $\sum_a A_a^j A_a^{j\dagger} = \mathbb{1}$, so Lemma 13.13.1.7 gives

$$D_b^j = \sum_a D_b^j A_a^j A_a^{j\dagger} = \sum_a A_b^j C_a^j A_a^{j\dagger} = A_b^j E^j.$$

Hence

$$A_b^j C_a^j = A_b^j E^j A_a^j$$

for every a, b , and cyclicity of the trace gives

$$\alpha_j = \sum_{i_1, \dots, i_{m+1}} \text{tr}(A_{i_1}^j \cdots A_{i_m}^j A_{i_{m+1}}^j E^j) |i_1 \cdots i_{m+1}\rangle = \Gamma_{m+1}^{A_j}(E^j) \in G_{m+1}(A_j).$$

Since $\psi = \sum_j \alpha_j$, this gives

$$\psi \in \bigoplus_j G_{m+1}(A_j).$$

For one block the preceding implication is

$$\mathbb{C}^d \otimes G_m(A_j) \cap G_m(A_j) \otimes \mathbb{C}^d \subseteq G_{m+1}(A_j).$$

The reverse inclusion is blockwise:

$$G_{m+1}(A_j) \subseteq \mathbb{C}^d \otimes G_m(A_j) \cap G_m(A_j) \otimes \mathbb{C}^d.$$

Thus

$$\mathbb{C}^d \otimes G_m(A_j) \cap G_m(A_j) \otimes \mathbb{C}^d = G_{m+1}(A_j).$$

With

$$S_m := \bigoplus_{j=1}^g G_m(A_j),$$

the preceding calculation is the open-segment recursion

$$\mathbb{C}^d \otimes S_m \cap S_m \otimes \mathbb{C}^d = S_{m+1}$$

for every $m \geq L$. For $B = \bigoplus_j \mu_j A_j$ with every $\mu_j \neq 0$, Lemma 13.14.5 gives

$$G_L(B) = \bigvee_j G_L(A_j).$$

In the range where the sums $\bigvee_j G_m(A_j)$ are direct, this identifies $G_L(B)$ with the subspace denoted in the source by $S = \bigoplus_j \mathcal{G}_L^{A_j}$. The passage from the open-segment recursion to the N -site ring is the separate boundary-condition comparison in the periodic ring in [PGVWC07, Theorem 12]. The source periodic-boundary conclusion is

$$\ker H_{S_L, N} = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

□

Lemma 13.15.2 (Conditional reduction to block-diagonal boundary conditions). Not ready Let $B^i = \bigoplus_j \mu_j A_j^i$, and assume $1 < L$, $N \geq L + 1$, and $0 < m$. Assume that every virtual sector projection P_j lies in the pulled-back span of the length- m words B^ω , and assume that every vector $\psi \in \mathcal{G}_{N, L}(\bigoplus_j \mu_j A_j)$ has a boundary matrix X with

$$\psi = \Gamma_N^B(X), \quad XB^\omega = B^\omega X$$

for the boundary-crossing words ω . Then

$$\mathcal{G}_{N, L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \sum_j \mathcal{G}_{N, L}(A_j),$$

with equality after the forward inclusion. This is a conditional form of the block-diagonal boundary-condition step in [CPGSV21, lines 2126–2128].

Proof. Not ready The commutant reduction makes the pulled-back boundary matrix block diagonal. Restricting the same word-commutation identities to a diagonal block and using the injectivity of that block shows, by the scalar-commutant lemma, that each diagonal block is scalar. Write Γ_N^B for the length- N boundary-to-state map of the tensor B . For a block-diagonal boundary matrix,

$$\Gamma_N^{\bigoplus_j \mu_j A_j}\left(\bigoplus_j X_j\right) = \sum_j \mu_j^N \Gamma_N^{A_j}(X_j).$$

If $X_j = \lambda_j I$, this is a finite sum of scalar multiples of the vectors $V^{(N)}(A_j)$. The reverse inclusion and equality statements are reformulations of this calculation, combined with the forward inclusion and the blockwise uniqueness theorem. □

Lemma 13.15.3 (Block-diagonal boundary reduction from blockwise word span). Not ready Let $A^i = \bigoplus_j \mu_j A_j^i$ be in block-injective canonical form, and assume $1 < L$, $N \geq L + 1$, and $0 < m$. If

$$\text{span}\{(A_1^\omega, \dots, A_g^\omega) : |\omega| = m\} = \prod_j M_{D_j}(\mathbb{C})$$

and every vector in $\mathcal{G}_{N, L}(\bigoplus_j \mu_j A_j)$ comes with a boundary matrix representation commuting with all length- m direct-sum words, then

$$\mathcal{G}_{N, L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \sum_j \mathcal{G}_{N, L}(A_j).$$

Combined with the forward inclusion, this gives the corresponding conditional equality. In the parent-ground-space formulation, the same hypotheses imply containment in $\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$.

Proof. Not ready The blockwise word-span hypothesis gives

$$P_j \in \text{span} \left\{ \left(\bigoplus_r \mu_r A_r \right)^\omega : |\omega| = m \right\}.$$

If X commutes with all displayed word products, then $XP_j = P_jX$ for every j , hence $X = \bigoplus_j X_j$. The direct-sum boundary formula

$$\Gamma_N^{\bigoplus_j \mu_j A_j} \left(\bigoplus_j X_j \right) = \sum_j \mu_j^N \Gamma_N^{A_j}(X_j)$$

gives the reverse inclusion, and the blockwise uniqueness theorem gives the stated span containment. $\square$

Lemma 13.15.4 (Block-diagonal boundary reduction from block-separating equations). Not ready Let $A^i = \bigoplus_j \mu_j A_j^i$ be in block-injective canonical form, with $A_1, \dots, A_g$ a basis of normal tensors. Assume $1 < L$ and $N \geq L + 1$, and suppose that for each block k there are coefficients $c_\omega^{(k)}$ such that

$$\sum_{|\omega|=S} c_\omega^{(k)} A_j^\omega = \begin{cases} I, & j = k, \\ 0, & j \neq k. \end{cases}$$

If every vector in $\mathcal{G}_{N,L}(\bigoplus_j \mu_j A_j)$ has a boundary matrix representation commuting with all length- $(1 + S)$ direct-sum words, then

$$\mathcal{G}_{N,L} \left(\bigoplus_j \mu_j A_j \right) \subseteq \sum_j \mathcal{G}_{N,L}(A_j).$$

Together with the forward inclusion, this gives the corresponding conditional equality. In the parent-ground-space formulation, the same hypotheses imply containment in $\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$. This is the block-injective ground-space argument of [CPGSV21, lines 2120–2128], where the re-growing step is restricted to block-diagonal boundary conditions.

Proof. Not ready Finite-length injectivity gives a block length at which each block has the required physical–virtual correspondence. In the length-one specialization, Lemma 2.4.3 supplies the corresponding one-block injectivity. The displayed separation equations imply

$$\text{span}\{(A_1^\omega, \dots, A_g^\omega) : |\omega| = 1 + S\} = \prod_j M_{D_j}(\mathbb{C}).$$

This is Lemma 13.13.16 with $L = 1$. Therefore Lemma 13.15.3 applies with $m = 1 + S$. Its conclusion is the displayed containment, and the equality follows after adding the forward inclusion. $\square$

Theorem 13.15.5 (Periodic-boundary containment in the BNT span). Not ready If $A^i = \bigoplus_j \mu_j A_j^i$ is in block-injective canonical form, where $A_1, \dots, A_g$ is a basis of normal tensors, assume $g \geq 2$ and $\mu_j \neq 0$ for every j . Let $L_0 \geq 1$ be a common injectivity length for the blocks, and suppose

$$L_0 < L, \quad L \geq 3(g - 1)(L_0 + 1) + 1, \quad N \geq L + L_0,$$

and assume the block-diagonal boundary-condition comparison in the periodic ring,

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \bigvee_{j=1}^g \mathcal{G}_{N,L}(A_j).$$

then

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

Proof. **Not ready** The length hypotheses are the conservative range used in [PGVWC07, Theorem 12]. The shorter range stated in [CPGSV21, lines 2126–2128] requires the strengthened block-diagonal re-growing step. These hypotheses do not by themselves give the displayed boundary-condition comparison in the periodic ring. That comparison follows only after the block-diagonal boundary conditions have been compared across the windows crossing the chosen periodic-boundary cut. Since $g \geq 2$ and $L_0 \geq 1$,

$$L \geq 3(g-1)(L_0+1)+1 \geq 3L_0+4 > L_0, \quad N \geq L+L_0 \geq L+1,$$

Because $A_1, \dots, A_g$ is a basis of normal tensors, the block MPV families are non-redundant:

$$j \neq k \implies \exists n \geq 1, V^{(n)}(A_j) \neq V^{(n)}(A_k).$$

By the assumed block-diagonal boundary-condition comparison,

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \bigvee_j \mathcal{G}_{N,L}(A_j).$$

Theorem 13.8.2.3 gives

$$\mathcal{G}_{N,L}(A_j) = \text{span}\{V^{(N)}(A_j)\}$$

for every j . Substituting these one-block identities into the join gives the displayed containment. $\square$

Theorem 13.15.6 (Periodic-boundary equality with the BNT span). **Not ready** Under the hypotheses of Theorem 13.15.5, the corresponding periodic-boundary ground space is exactly the span of the BNT component vectors:

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) = \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}.$$

Proof. **Not ready** We prove the two inclusions separately. The inclusion

$$\mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right) \subseteq \text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$$

is exactly Theorem 13.15.5. For the reverse inclusion, the nonzero-weight hypothesis and Lemma 13.12.6 show that each vector $V^{(N)}(A_j)$ lies in $\mathcal{G}_{N,L}(\bigoplus_j \mu_j A_j)$. Since this ground space is linear, it contains every linear combination of such vectors, and hence contains $\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\}$. Thus

$$\text{span}\{V^{(N)}(A_j) : j = 1, \dots, g\} \subseteq \mathcal{G}_{N,L}\left(\bigoplus_j \mu_j A_j\right),$$

and the two spaces are equal. $\square$

Remark. A separate kernel-identification statement for these direct-sum block families would convert the periodic-boundary ground-space statement into the corresponding statement for $\ker(H_N)$.

Chapter 14

Exponential Decay of Correlations

This chapter derives the transfer-matrix expression for thermodynamic-limit correlators of a normal MPS and the resulting exponential decay estimate. The key mechanism is spectral: after removing the leading eigenvalue contribution, the connected part is governed by subleading eigenvalues of the transfer map.

When the MPS tensor generates a parent Hamiltonian (Chapter 13), the correlator quantifies spatial correlations in the ground state. The exponential decay rate is set by the spectral properties of the transfer map, i.e. by the modulus of its subleading eigenvalues.

The spectral analysis of the transfer map and the mixed transfer operator, including the eigenvalue bound $\rho(F_{AB}) \leq 1$, the strict transfer-operator gap $\rho(F_{AB}) < 1$ for non-equivalent blocks, and the resulting overlap decay, is developed in Chapter 7. This chapter focuses on the *correlator* side: one-point and two-point expectations, their spectral decomposition, and quantitative bounds on decay rates.

14.1 Connected correlators from the transfer map


Definition 14.1.1 (One-point expectation). [Lean](#) [✓ Stmt](#) Fix an MPS tensor and a right fixed point of its transfer map. Write the tensor as A and the fixed point as ρ^R . Under the hypotheses of Theorem 6.3.2, that fixed point exists and is unique up to scaling; we normalize it so that $\text{tr}(\rho^R) = 1$. For a local observable $X \in M_D(\mathbb{C})$ define

$$\langle X_0 \rangle := \text{tr}(X \rho^R).$$

Definition 14.1.2 (Two-point expectation at distance n). [Lean](#) [✓ Stmt](#) For observables $X, Y \in M_D(\mathbb{C})$ and $n \geq 0$,

$$\langle X_0 Y_n \rangle := \text{tr}(Y \mathcal{E}_A^n(X \rho^R)).$$

Equivalently: insert X at site 0, propagate by $\mathcal{E}_A^n$, then insert Y and take the trace.



Definition 14.1.3 (Connected correlator). [Lean](#) [✓ Stmt](#) The connected two-point function is

$$C(X, Y; n) := \langle X_0 Y_n \rangle - \langle X_0 \rangle \langle Y_0 \rangle.$$

14.2 Spectral expansion and exponential decay

Theorem 14.2.1 (Sum of exponentials (spectral expansion interface)). [Lean](#) [✓ Stmt](#) If coefficients $c_j \in \mathbb{C}$ and eigenvalues $\lambda_j \in \mathbb{C}$ ($j = 1, \dots, D^2 - 1$) satisfy, for all $n \geq 0$,

$$C(X, Y; n) = \sum_{j=1}^{D^2-1} c_j \lambda_j^n,$$

then the connected correlator equals the stated sum of exponentials. The source [\[CPGSV21, Section II.B.3\]](#) states that the connected correlator is a sum of $D^2 - 1$ pure exponentials $C(X, Y; n) = \sum_{j \geq 2}^{D^2} c_{XY}(j) \lambda_j^n$, which always exists for a normal MPS via the transfer-map spectral decomposition.

Proof. [✓ Proof](#) Given the expansion hypothesis, the equality is immediate. The source [\[CPGSV21, Section II.B.3\]](#) derives this expansion from the spectral decomposition of $\mathcal{E}_A^n$ restricted to the complement of the fixed-point eigenspace: the leading eigenvalue $\lambda_1 = 1$ contributes $\langle X_0 \rangle \langle Y_0 \rangle$, which is subtracted in the connected correlator, leaving the sum over subleading eigenvalues. $\square$

Theorem 14.2.2 (Exponential decay bound (interface)). [Lean](#) [✓ Stmt](#) If a constant $C_{XY} \in \mathbb{R}$ and $\lambda_2 \in \mathbb{C}$ satisfy, for all $n \geq 0$,

$$|C(X, Y; n)| \leq C_{XY} |\lambda_2|^n,$$

then the connected correlator satisfies that exponential decay bound. The source [\[CPGSV21, Section II.B.3\]](#) obtains this by combining the sum-of-exponentials expansion with the triangle inequality, and [Sec. 4](#) provides the transfer-map gap condition $|\lambda_j| < 1$ for the subleading eigenvalues.

Proof. [✓ Proof](#) Given the bound hypothesis, the estimate is immediate. The source [\[CPGSV21, Section II.B.3\]](#) obtains this bound by applying the triangle inequality to the sum-of-exponentials expansion and using the transfer-map gap condition $|\lambda_j| \leq |\lambda_2| < 1$ for all subleading eigenvalues. $\square$

Remark 14.2.3 (Transfer-map spectral hypothesis). The hypothesis “ $|\lambda_2| < 1$ ” in [Theorem 14.2.2](#) is a transfer-map gap condition: all non-leading eigenvalues of $\mathcal{E}_A$ lie strictly inside the unit disk. It is distinct from the Hamiltonian spectral gap of [Chapter 13](#). The complementary transfer-map gap $\rho(\mathcal{E}_A - P) < 1$ is established for primitive channels ([Theorem 4.5.3.8](#), [Chapter 4](#)). The mixed-transfer gap needed for block separation is supplied for irreducible trace-preserving tensors ([Theorem 7.8.2](#)), and quantitative single-tensor transfer-map bounds for injective tensors are recorded in [Theorem 14.3.3](#). Thus the analytical input needed to make [Theorems 14.2.1](#) and [14.2.2](#) unconditional is already present; what remains is extracting the spectral expansion coefficients c_j, λ_j from the eigendecomposition of $\mathcal{E}_A$ ([Issue #1447](#)).

Definition 14.2.4 (Correlation length). [Lean](#) [✓ Stmt](#) The correlation length associated with a subleading eigenvalue λ_2 of the transfer map ($|\lambda_2| < 1$) is

$$\xi := -\frac{1}{\log |\lambda_2|}.$$

Lemma 14.2.5 (Correlation length is positive). [Lean](#) [✓ Stmt](#) If $|\lambda_2| < 1$ and $\lambda_2 \neq 0$, then $\xi > 0$.

Proof. [✓ Proof](#) Since $0 < |\lambda_2| < 1$, we have $\log |\lambda_2| < 0$, so $\xi = -1/\log |\lambda_2| > 0$. $\square$

14.3 Quantitative transfer-map gap bounds

The transfer-map gap theorems in Chapter 4 establish $\rho(\mathcal{E}_A - P) < 1$ for primitive channels qualitatively. This section gives the *quantitative* refinements with explicit constants and rates for the single-tensor transfer map $\mathcal{E}_A$.

Theorem 14.3.1 (Exponential convergence of injective primitive channels). [Lean](#) [✓ Stmt](#) Fix an injective, trace-preserving MPS tensor A with positive definite fixed point ρ . Let

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$$

be the corresponding fixed-point projection. Then there exist $C > 0$ and $0 < \delta \leq 1$ such that for all $n \geq 0$ and $X \in M_D(\mathbb{C})$,

$$\|\mathcal{E}_A^n(X) - P(X)\| \leq C(1 - \delta)^n \|X\|.$$

The transfer-map gap δ exists by primitivity (Theorem 4.5.3.8, the complementary transfer-map gap).

Proof. [✓ Proof](#) Injectivity implies primitivity of the transfer map. The complementary spectral radius $\rho(\mathcal{E}_A - P) < 1$ then follows from the primitive transfer-map gap theorem for the complementary map. The Gelfand formula yields a geometric bound $\|(\mathcal{E}_A - P)^n\| \leq C(1 - \delta)^n$ for suitable constants, giving the claimed estimate. $\square$

Theorem 14.3.2 (Correlation length bound). [Lean](#) [✓ Stmt](#) For an injective trace-preserving MPS tensor, there exist $C > 0$ and $\xi > 0$ such that for all $n \geq 0$ and all traceless $X \in M_D(\mathbb{C})$ (i.e. $\text{tr}(X) = 0$),

$$\|\mathcal{E}_A^n(X)\| \leq C e^{-n/\xi} \|X\|.$$

Traceless matrices lie in $\ker P$, where P is the fixed-point projection. Since $\mathcal{E}_A - P$ has spectral radius strictly less than 1, the Gelfand formula gives exponential decay at rate $\xi = -1/\log \rho(\mathcal{E}_A - P)$.

Proof. [✓ Proof](#) For traceless X we have $P(X) = 0$, so $\mathcal{E}_A^n(X) = (\mathcal{E}_A - P)^n(X)$. The exponential convergence theorem gives a geometric bound $C(1 - \delta)^n \|X\|$. Rewriting $(1 - \delta)^n = e^{-n/\xi}$ with $\xi = -1/\log(1 - \delta)$ yields the claimed estimate. $\square$

Theorem 14.3.3 (Transfer-map gap from injectivity). [Lean](#) [✓ Stmt](#) For an injective trace-preserving MPS tensor, there exists $\delta > 0$ such that every eigenvalue $\mu \neq 1$ of the transfer map satisfies $|\mu| \leq 1 - \delta$.

Proof. [✓ Proof](#) Injectivity implies eventual full Kraus rank. This yields primitivity, hence a complementary transfer-map gap. Since the transfer map has only finitely many eigenvalues, one may choose a uniform $\delta > 0$ separating 1 from the remaining spectrum. $\square$

14.4 Relation to parent Hamiltonians

Remark 14.4.1. When the MPS tensor A generates a parent Hamiltonian $H_N(A, L)$ (Definition 13.3.14) whose ground state is the matrix product vector (Lemma 13.3.18), the exponential decay bound (Theorem 14.2.2) shows that ground-state correlations decay with a finite correlation length ξ .

Chapter 15

Concrete Examples

This chapter collects concrete MPS tensors from [CPGSV21, Section III]. Each example is specified by its physical dimension d , bond dimension D , and explicit matrix family $\{A^i\}$. Key structural properties (injectivity, RFP status, symmetry) are stated for each tensor.

15.1 The GHZ state

The Greenberger–Horne–Zeilinger (GHZ) state is the prototypical non-injective, renormalization-fixed-point MPS.

Definition 15.1.1 (GHZ tensor). [Lean](#) [✓ Stmt](#) Set $d = D = 2$. The GHZ tensor is the family $\{A^0, A^1\} \subset M_2(\mathbb{C})$ defined by

$$A^0 = |0\rangle\langle 0| = \text{diag}(1, 0), \quad A^1 = |1\rangle\langle 1| = \text{diag}(0, 1).$$

For a system of N sites the resulting MPV is the GHZ state

$$|\text{GHZ}_N\rangle = |0\rangle^{\otimes N} + |1\rangle^{\otimes N}.$$

Definition 15.1.2 (Two-site computational vector). [Lean](#) [✓ Stmt](#) For $a, b \in \{0, 1\}$, the vector

$$|ab\rangle \in (\{0, 1\}^2 \rightarrow \mathbb{C})$$

is the indicator of the configuration

$$(\sigma_1, \sigma_2) = (a, b).$$

Definition 15.1.3 (Constant product vector). [Lean](#) [✓ Stmt](#) For $a \in \{0, 1\}$ and $N \geq 0$,

$$|a\rangle^{\otimes N}(\sigma_1, \dots, \sigma_N) = \begin{cases} 1, & (\sigma_1, \dots, \sigma_N) = (a, \dots, a), \\ 0, & \text{otherwise.} \end{cases}$$

Theorem 15.1.4 (GHZ transfer map is idempotent). [Lean](#) [✓ Stmt](#) The transfer map of the GHZ tensor satisfies $\mathcal{E}_A^2 = \mathcal{E}_A$.

Proof. [✓ Proof](#) The transfer map acts by $\mathcal{E}_A(X)_{ij} = \delta_{ij}X_{ij}$ (extraction of the diagonal). This operation is idempotent. $\square$

Theorem 15.1.5 (GHZ tensor is an RFP). [Lean](#) [✓ Stmt](#) *The GHZ tensor is a renormalization fixed point.*

Proof. [✓ Proof](#) The claim follows from Theorem 15.1.4. $\square$

Theorem 15.1.6 (GHZ tensor is not injective). [Lean](#) [✓ Stmt](#) *The GHZ tensor is not injective: $\text{span}_{\mathbb{C}}\{A^0, A^1\}$ is the two-dimensional space of diagonal matrices in $M_2(\mathbb{C})$, which does not equal $M_2(\mathbb{C})$.*

Proof. [✓ Proof](#) Every element of $\text{span}_{\mathbb{C}}\{A^0, A^1\}$ has zero off-diagonal entries, so the matrix with all entries equal to 1 is not in the span. $\square$

Theorem 15.1.7 ($\mathbb{Z}_2$ on-site symmetry of the GHZ tensor). [Lean](#) [✓ Stmt](#) *Let $\mathbb{Z}_2 = \{1, \sigma_x\}$ act on $\mathbb{C}^2$ by the Pauli-X matrix σ_x . The GHZ tensor is on-site symmetric under this action: for each $g \in \mathbb{Z}_2$, the twisted tensor $\tilde{A}_g$ defines the same MPV family as A , i.e. $\mathcal{V}(A) = \mathcal{V}(\tilde{A}_g)$.*

Proof. [✓ Proof](#) The identity element is trivial. For the generator $g = \sigma_x$, the twisted tensor satisfies $\tilde{A}_g^i = A^{1-i}$, i.e. the two matrices are swapped. Conjugation by σ_x implements the same swap, giving gauge equivalence and hence the same MPV family. $\square$

Definition 15.1.8 (Virtual $\mathbb{Z}_2$ representation of the GHZ tensor). [Lean](#) [✓ Stmt](#) *The virtual $\mathbb{Z}_2$ representation on the bond space is the homomorphism $\mathbb{Z}_2 \rightarrow M_2(\mathbb{C})$ sending the identity to I and the generator to $\sigma_z = \text{diag}(1, -1)$.*

Theorem 15.1.9 (Virtual $\mathbb{Z}_2$ symmetry of the GHZ tensor). [Lean](#) [✓ Stmt](#) *The matrix σ_z commutes with every GHZ matrix A^i . A non-scalar matrix commuting with all A^i is the hallmark of a non-injective ($\mathbb{Z}_2$ -injective) tensor: the on-site symmetry of Theorem 15.1.7 swaps the two one-dimensional injective blocks $A^0 \leftrightarrow A^1$, while σ_z acts on each block by a phase.*

Proof. [✓ Proof](#) Both A^0 and A^1 are diagonal, so each commutes with the diagonal matrix σ_z . $\square$

Theorem 15.1.10 (GHZ order parameter is a transfer-map fixed point). [Lean](#) [✓ Stmt](#) *The order-parameter operator σ_z is a fixed point of the GHZ transfer map, $\mathcal{E}_A(\sigma_z) = \sigma_z$.*

Proof. [✓ Proof](#) Since $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ and both A^i are diagonal, the claim is the entrywise computation $\mathcal{E}_A(\sigma_z) = \sigma_z$. $\square$

Remark 15.1.11. A primitive (injective) tensor has a one-dimensional transfer-map fixed-point space, spanned by a positive matrix that equals the identity only in a normalized gauge. That the GHZ transfer map fixes the additional non-scalar operator σ_z is the transfer-matrix signature of the extra non-decaying mode behind its long-range ferromagnetic correlations.

Theorem 15.1.12 (GHZ zero correlation length). [Lean](#) [✓ Stmt](#) *The GHZ tensor has zero correlation length: its idempotent transfer map (Theorem 15.1.5) gives correlations independent of the separation. This contrasts with the AKLT state (Theorem 15.2.1.3), whose subleading eigenvalue $-\frac{1}{3}$ produces a finite correlation length $1/\log 3$.*

Proof. [✓ Proof](#) The claim follows from the idempotence of the GHZ transfer map and the equivalence of zero correlation length with transfer-map idempotence (Theorem 22.5.13). $\square$

15.2 The AKLT state

The Affleck–Kennedy–Lieb–Tasaki (AKLT) state is the canonical example of a normal (block-injective) MPS with non-trivial symmetry-protected topological (SPT) order.

Definition 15.2.1 (AKLT tensor). [Lean](#) [✓ Stmt](#) Set $d = 3$ (spin-1) and $D = 2$. The AKLT tensor is the family $\{A^0, A^1, A^2\} \subset M_2(\mathbb{C})$ defined via the Pauli matrices by

$$A^0 = \frac{1}{\sqrt{3}}\sigma_z, \quad A^1 = \frac{\sqrt{2}}{\sqrt{3}}|0\rangle\langle 1|, \quad A^2 = -\frac{\sqrt{2}}{\sqrt{3}}|1\rangle\langle 0|,$$

where $\sigma_z = |0\rangle\langle 0| - |1\rangle\langle 1|$, so equivalently $A^1 = \frac{1}{\sqrt{3}}\sigma_+$ and $A^2 = -\frac{1}{\sqrt{3}}\sigma_-$ for $\sigma_+ = \sqrt{2}|0\rangle\langle 1|$ and $\sigma_- = \sqrt{2}|1\rangle\langle 0|$. Equivalently, the matrices are the Clebsch–Gordan coefficients coupling spin-1 and spin- $\frac{1}{2}$ to spin- $\frac{1}{2}$.

Theorem 15.2.2 (AKLT tensor is not injective). [Lean](#) [✓ Stmt](#) *The AKLT tensor is not injective at a single site: $\text{span}_{\mathbb{C}}\{A^0, A^1, A^2\}$ is the three-dimensional space of traceless matrices $\mathfrak{sl}(2, \mathbb{C})$, which is a proper subspace of $M_2(\mathbb{C})$.*

Theorem 15.2.3 (AKLT tensor is normal). [Lean](#) [✓ Stmt](#) *The transfer map of the AKLT tensor has a unique eigenvalue of maximal modulus (a simple spectral-radius eigenvalue); in particular the AKLT tensor is normal. Concretely, the length-2 blocked tensor is injective.*

Theorem 15.2.4 ($\mathbb{Z}_2 \times \mathbb{Z}_2$ on-site symmetry of the AKLT tensor). [Lean](#) [✓ Stmt](#) *The AKLT tensor is on-site symmetric under the $\mathbb{Z}_2 \times \mathbb{Z}_2$ subgroup of $\text{SO}(3)$ generated by two commuting π -rotations about orthogonal axes, acting on the spin-1 physical index.*

Theorem 15.2.5 (Anticommuting virtual gauges of the AKLT symmetry). [Lean](#) [✓ Stmt](#) *The two virtual matrices that implement the symmetry of Theorem 15.2.4, namely $i\sigma_y$ and σ_z , anti-commute.*

Definition 15.2.6 (AKLT factor system). [Lean](#) [✓ Stmt](#) The AKLT factor system on $\mathbb{Z}_2 \times \mathbb{Z}_2$ is $\omega(g, h) = (-1)^{(g_1+g_2)h_1}$ (the value -1 when $g_1 + g_2 = 1$ and $h_1 = 1$, and 1 otherwise). It is the factor system of the explicit projective representation with virtual gauges $i\sigma_y$ and σ_z ; the diagonal term $g_1 h_1$ relative to the cluster case reflects $(i\sigma_y)^2 = -I$.

Lemma 15.2.7 (AKLT commutator phase). [Lean](#) [✓ Stmt](#) *The commutator phase of the AKLT factor system on the two generators is -1 .*

Theorem 15.2.8 (The AKLT factor system is a 2-cocycle). [Lean](#) [✓ Stmt](#) *The AKLT factor system is a genuine 2-cocycle, being the factor system of an explicit projective representation, so it represents an element of $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \text{U}(1))$.*

Theorem 15.2.9 (The AKLT $\mathbb{Z}_2 \times \mathbb{Z}_2$ phase is non-trivial). [Lean](#) [✓ Stmt](#) *The AKLT factor system (Theorem 15.2.8), built from the anticommuting gauges $i\sigma_y$ and σ_z (Theorem 15.2.5), is not a coboundary: its commutator phase on the two generators is -1 . Its class in $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \text{U}(1)) \cong \mathbb{Z}_2$ is therefore non-trivial, so the AKLT chain realizes a non-trivial SPT phase rather than a trivial product state.*

Remark 15.2.10. This $\mathbb{Z}_2 \times \mathbb{Z}_2$ is the smallest subgroup of $\text{SO}(3)$ that still carries a non-trivial 2-cocycle. That minimality, and the full continuous classification over $\text{SO}(3)$, are not used in this discrete-subgroup example (Remark 15.2.20).

Definition 15.2.11 (Blocked AKLT tensor). [Lean](#) [✓ Stmt](#) The length-2 blocked AKLT tensor groups two physical sites into one, with blocked physical dimension $9 = 3^2$ and the same bond dimension $D = 2$.

Definition 15.2.12 (Blocked $\mathbb{Z}_2 \times \mathbb{Z}_2$ on-site action). [Lean](#) [✓ Stmt](#) The length-2 blocked on-site action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ on the 9-dimensional blocked physical space is the Kronecker square

$$g \mapsto P_g \otimes P_g$$

of the single-site spin-1 π -rotation representation P_g that implements the symmetry of Theorem 15.2.4. Identifying the blocked physical index with the pair of single-site indices, the entry of $P_g \otimes P_g$ at $(i_1 i_2, j_1 j_2)$ is the product $(P_g)_{i_1 j_1} (P_g)_{i_2 j_2}$ over the two grouped sites. This is the on-site action that makes the blocked twist agree with blocking the single-site twist.

Theorem 15.2.13 (The blocked AKLT tensor is injective). [Lean](#) [✓ Stmt](#) *The length-2 blocked AKLT tensor is injective: its nine matrices span $M_2(\mathbb{C})$.*

Proof. [✓ Proof](#) This is the blocked-tensor form of the normality of the single-site tensor (Theorem 15.2.3): the single-site tensor is 2-block injective, which is exactly injectivity of its length-2 blocking. $\square$

Theorem 15.2.14 ($\mathbb{Z}_2 \times \mathbb{Z}_2$ on-site symmetry of the blocked AKLT tensor). [Lean](#) [✓ Stmt](#) *The length-2 blocked AKLT tensor is on-site symmetric under $\mathbb{Z}_2 \times \mathbb{Z}_2$, with each group element acting by the Kronecker square $P_g \otimes P_g$ of the corresponding single-site spin-1 π -rotation P_g (Definition 15.2.12).*

Proof. [✓ Proof](#) Writing A for the single-site tensor and $A^{[2]}$ for its length-2 blocking, twisting the blocked tensor by $P_g \otimes P_g$ equals blocking the twisted single-site tensor,

$$(A^{[2]})^{P_g \otimes P_g} = (A^{P_g})^{[2]},$$

where the superscript denotes the twist. Since the single-site tensor is symmetric, A^{P_g} has the same matrix product vector family as A (Theorem 15.2.4), and blocking preserves the matrix product vector family, so $(A^{[2]})^{P_g \otimes P_g}$ has the same matrix product vector family as $A^{[2]}$. $\square$

Theorem 15.2.15 (The single-site $\mathbb{Z}_2 \times \mathbb{Z}_2$ action is unitary). [Lean](#) [✓ Stmt](#) *Each single-site spin-1 π -rotation matrix P_g of the $\mathbb{Z}_2 \times \mathbb{Z}_2$ action is unitary: $P_g P_g^\dagger = \mathbb{1}$.*

Theorem 15.2.16 (The AKLT state has string order under its $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry). [Lean](#) [✓ Stmt](#) *For every element g of $\mathbb{Z}_2 \times \mathbb{Z}_2$, the length-2 blocked AKLT tensor has string order in the sense of Definition 12.5.7 for the twist by g , with the maximally mixed boundary state $\Lambda = \frac{1}{2}\mathbb{1}$ as its stationary boundary state.*

Proof. [✓ Proof](#) The blocked tensor is injective (Theorem 15.2.13) and on-site symmetric under $\mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 15.2.14), and each group element acts by a real symmetric involutive matrix, hence unitarily. The maximally mixed state $\Lambda = \frac{1}{2}\mathbb{1}$ is positive definite and has trace 1. The single-site AKLT matrices satisfy $\sum_i A^i (A^i)^\dagger = \mathbb{1}$ and $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. These identities propagate to the length-2 blocked tensor $A^{[2]}$ whose matrices are the products $A^{i_1} A^{i_2}$: the blocked transfer map is unital,

$$\mathcal{E}_{A^{[2]}}(\mathbb{1}) = \sum_{i_1, i_2} A^{i_1} A^{i_2} (A^{i_2})^\dagger (A^{i_1})^\dagger = \mathbb{1},$$

and $\Lambda = \frac{1}{2}\mathbb{1}$ is a fixed point of the adjoint transfer map,

$$\mathcal{E}_{A^{[2]}}^\dagger(\Lambda) = \sum_{i_1, i_2} (A^{i_2})^\dagger (A^{i_1})^\dagger \Lambda A^{i_1} A^{i_2} = \Lambda.$$

The first identity sums the right-canonical relation $\sum_i A^i (A^i)^\dagger = \mathbb{1}$ over the inner letter and then the outer; the second sums the left-canonical relation $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ over the outer letter and then the inner, using $\Lambda = \frac{1}{2}\mathbb{1}$. These are exactly the hypotheses of Theorem 12.6.3, which then yields string order for every g . Like the cluster state, the AKLT state is an explicit witness of that criterion, sharing the same virtual-boundary caveat on the string order parameter. $\square$

Definition 15.2.17 (Cartesian AKLT tensor). [Lean](#) [✓ Stmt](#) The Cartesian presentation of the AKLT tensor has $d = 3$, $D = 2$, and $A^x = \sigma_x$, $A^y = \sigma_y$, $A^z = \sigma_z$, the three Pauli matrices. This is the rotationally natural form used in [CPGSV21] (around line 1159); it is the AKLT state in the Cartesian basis of the spin-1 site, the counterpart of the $|m\rangle$ -basis form of Definition 15.2.1.

Theorem 15.2.18 (The spin- $\frac{1}{2}$ cover is surjective onto $\text{SO}(3)$). [Lean](#) [✓ Stmt](#) Every rotation $R \in \text{SO}(3)$ is realized by conjugating the Pauli vector by some $U \in \text{SU}(2)$: there is $U \in \text{SU}(2)$ with $(\frac{1}{2} \text{tr}(\sigma_i U \sigma_j U^{-1}))_{ij} = R$. Equivalently, the adjoint double cover $\text{SU}(2) \rightarrow \text{SO}(3)$ is surjective.

Proof. [✓ Proof](#) Factor $R = R_z(\alpha) R_x(\beta) R_z(\gamma)$ by its ZXZ Euler decomposition and map each coordinate rotation to an explicit $\text{SU}(2)$ generator: $\text{diag}(e^{-i\theta/2}, e^{i\theta/2}) \mapsto R_z(\theta)$ and $\begin{pmatrix} \cos \frac{\beta}{2} & -i \sin \frac{\beta}{2} \\ -i \sin \frac{\beta}{2} & \cos \frac{\beta}{2} \end{pmatrix} \mapsto R_x(\beta)$. $\square$

Theorem 15.2.19 ($\text{SO}(3)$ on-site symmetry of the AKLT tensor). [Lean](#) [✓ Stmt](#) The Cartesian AKLT tensor is on-site symmetric under the spin-1 (defining) representation of $\text{SO}(3)$ on the physical index.

Remark 15.2.20. The bond index carries the projective spin- $\frac{1}{2}$ representation: the gauge for a rotation is either of its two $\text{SU}(2)$ preimages, so the assignment is a representation only up to sign and factors through the double cover $\text{SU}(2) \rightarrow \text{SO}(3)$. Its class is the non-trivial element of $H^2(\text{SO}(3), \text{U}(1)) \cong \mathbb{Z}_2$, witnessing the SPT phase.

15.2.1 Correlation length

The AKLT state is the canonical example of a finite correlation length. Its transfer map has leading eigenvalue 1 and a single subleading eigenvalue $-\frac{1}{3}$ of multiplicity three; the three Pauli matrices are the subleading eigenvectors and the identity is the leading one. The subleading modulus $\frac{1}{3}$ gives the correlation length $\xi = 1/\log 3$, and connected correlations decay as $(\frac{1}{3})^n$.

Theorem 15.2.1.1 (Pauli eigenvectors of the AKLT transfer map). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Each of the three Pauli matrices σ_x , σ_y , σ_z is an eigenvector of the AKLT transfer map with eigenvalue $-\frac{1}{3}$:

$$\mathcal{E}_A(\sigma_k) = -\frac{1}{3} \sigma_k, \quad k \in \{x, y, z\}.$$

Proof. [✓ Proof](#) Each identity is an entrywise computation from the AKLT matrices, using $(\sqrt{3})^2 = 3$ and $(\sqrt{2})^2 = 2$ to clear the prefactors. $\square$

Theorem 15.2.1.2 (Subleading eigenvalue of the AKLT transfer map). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The value 1 is the leading eigenvalue of the AKLT transfer map, with the identity $\mathbb{1}$ as eigenvector, and $-\frac{1}{3}$ is a subleading eigenvalue, with σ_z as eigenvector. The subleading modulus is therefore $\frac{1}{3}$.

Theorem 15.2.1.3 (AKLT correlation length). [Lean](#) [Lean](#) [✓ Stmt](#) *The AKLT correlation length, computed from the subleading eigenvalue $\lambda_2 = -\frac{1}{3}$, equals*

$$\xi = -\frac{1}{\log |\lambda_2|} = \frac{1}{\log 3}.$$

It is finite and positive.

Proof. [✓ Proof](#) The subleading modulus is $|\lambda_2| = \frac{1}{3}$, so rewriting the correlation length at $\lambda_2 = -\frac{1}{3}$ gives $\xi = -1/\log \frac{1}{3} = 1/\log 3$. Since $0 < \frac{1}{3} < 1$, positivity is the specialization of Lemma 14.2.5 to $\lambda_2 = -\frac{1}{3}$. $\square$

15.3 The Majumdar–Ghosh state

The Majumdar–Ghosh state is the exact dimer ground state of the spin- $\frac{1}{2}$ chain $H = \sum S_i \cdot S_{i+1} + \frac{1}{2} \sum S_i \cdot S_{i+2}$. On an even periodic chain it is the symmetric superposition of the two nearest-neighbour singlet coverings $(1, 2)(3, 4) \dots$ and $(2, 3)(4, 5) \dots (N, 1)$, the one-dimensional analogue of a resonating valence-bond state.

Definition 15.3.1 (Majumdar–Ghosh tensor). [Lean](#) [✓ Stmt](#) Set $d = 2$ (spin- $\frac{1}{2}$) and $D = 3$. The Majumdar–Ghosh tensor is the family $\{A^0, A^1\} \subset M_3(\mathbb{C})$ defined by

$$A^0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 \end{pmatrix}, \quad A^1 = \begin{pmatrix} 0 & 0 & 1 \\ -\frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The three bond levels are the two spin- $\frac{1}{2}$ values of an open singlet partner together with the closed inter-dimer link; the relative sign $-\frac{1}{\sqrt{2}}$ is the antisymmetry of the singlet. The equality between the matrix product vectors of these matrices and the even-ring dimer superposition, with normalization and cyclic convention fixed, is a separate verification, not part of this definition.

Theorem 15.3.2 (Majumdar–Ghosh tensor is in left-canonical form). [Lean](#) [✓ Stmt](#) *The Majumdar–Ghosh tensor is isometric: $(A^0)^\dagger A^0 + (A^1)^\dagger A^1 = \mathbb{1}$.*

Proof. [✓ Proof](#) Direct multiplication of the two displayed matrices gives $(A^0)^\dagger A^0 = \text{diag}(\frac{1}{2}, 1, 0)$ and $(A^1)^\dagger A^1 = \text{diag}(\frac{1}{2}, 0, 1)$. $\square$

Theorem 15.3.3 (Transfer map of the Majumdar–Ghosh tensor). [Lean](#) [✓ Stmt](#) *The transfer map of the Majumdar–Ghosh tensor sends the identity to the diagonal matrix $\mathcal{E}_A(\mathbb{1}) = \text{diag}(2, \frac{1}{2}, \frac{1}{2})$.*

Proof. [✓ Proof](#) Substitute the two matrices in $\mathcal{E}_A(\mathbb{1}) = A^0(A^0)^\dagger + A^1(A^1)^\dagger$ and multiply. $\square$

Theorem 15.3.4 (Majumdar–Ghosh tensor is not injective). [Lean](#) [✓ Stmt](#) *The Majumdar–Ghosh tensor is not injective: every element of $\text{span}_{\mathbb{C}}\{A^0, A^1\}$ has vanishing $(2, 2)$ entry, so the span is a proper subspace of $M_3(\mathbb{C})$.*

Proof. [✓ Proof](#) The matrix unit E_{22} has nonzero $(2, 2)$ entry, while every linear combination of A^0 and A^1 has zero $(2, 2)$ entry. $\square$

Theorem 15.3.5 (Majumdar–Ghosh tensor is not normal). [Lean](#) [✓ Stmt](#) *The Majumdar–Ghosh tensor is not normal: no blocking length makes the products of that length span the full matrix algebra $M_3(\mathbb{C})$. This non-normality is the algebraic signature expected from the two-periodic dimer-covering structure.*

Proof. ✓ Proof The bond carries a two-periodic grading with complementary support sets $\{(0, 1), (0, 2), (1, 0), (2, 0)\}$ and $\{(0, 0), (1, 1), (1, 2), (2, 1), (2, 2)\}$ that are exchanged by left multiplication with a generator. Let E_{ij} denote the (i, j) matrix unit. Then, for every blocking length N ,

$$\begin{aligned} N \text{ odd} &\Rightarrow (\forall w, |w| = N \Rightarrow (A^w)_{22} = 0) \Rightarrow E_{22} \notin \text{span}_{\mathbb{C}}\{A^w : |w| = N\}, \\ N \text{ even} &\Rightarrow (\forall w, |w| = N \Rightarrow (A^w)_{10} = 0) \Rightarrow E_{10} \notin \text{span}_{\mathbb{C}}\{A^w : |w| = N\}. \end{aligned}$$

Hence no fixed-length word span is all of $M_3(\mathbb{C})$. □

Remark 15.3.6. The review writes the tensor in the valence-bond shorthand $A = [|0\rangle\langle 02| + (20| + |1\rangle\langle 12| + (21|)] \otimes Y$ with the singlet $Y = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Read literally as $(ab) = |a\rangle\langle b|$ on a separate three-level factor tensored with the two-level singlet space, that shorthand would describe a six-dimensional bond. Its two-site contractions do not have the singlet-pair support pattern: same-letter traces can be nonzero while opposite-letter traces vanish. The shorthand is therefore treated here as a schematic of the valence-bond construction: contracting the singlet Y leaves a matrix product operator on a three-dimensional bond, with the antisymmetry of Y supplying the relative sign. The contracted tensor is the $D = 3$ representative in Definition 15.3.1. The source-notation gap and the missing matrix-product-vector identification are recorded in docs/paper-gaps/rmp_majumdar_ghosh_tensor_gap.tex.

The remaining target for the Majumdar–Ghosh example is the ground-space description of its parent Hamiltonian. Because the tensor is non-normal (Theorem 15.3.5) with a two-periodic bond grading, its periodic ground space is governed by the block-diagonal boundary-condition machinery for non-normal tensors (Section 13.14) rather than by the injective single-state result. Thus the relevant target is the following range-three periodic ground-space statement.

Theorem 15.3.7 (Two-fold dimer degeneracy of the Majumdar–Ghosh ground space). Not ready

On an even periodic chain of length $N = 2m \geq 4$, let $H_{N, \text{MG}}^{(3)}$ be the range-three parent Hamiltonian of the Majumdar–Ghosh tensor, the periodic sum of translates of the local projector with kernel $\mathcal{G}_3(A_{\text{MG}})$. Then $H_{N, \text{MG}}^{(3)}$ has a two-dimensional ground space, spanned by the two nearest-neighbour singlet coverings $(1, 2)(3, 4) \cdots (N-1, N)$ and $(2, 3)(4, 5) \cdots (N, 1)$. This two-fold degeneracy reflects the spontaneous breaking of the one-site translation symmetry of the spin- $\frac{1}{2}$ chain $H = \sum S_i \cdot S_{i+1} + \frac{1}{2} \sum S_i \cdot S_{i+2}$ of [CPGSV21].

15.4 The W state

The W state on N sites is the single-excitation symmetric state

$$|W_N\rangle = |10 \cdots 0\rangle + |010 \cdots 0\rangle + \cdots + |0 \cdots 01\rangle,$$

the equal-weight superposition of all configurations with exactly one excitation. Unlike the preceding examples it has no translation-invariant representation with bond dimension bounded independently of N ; it is presented with open boundary conditions, where the matrix product is closed by a left and a right boundary vector rather than by a trace.

Definition 15.4.1 (Open-boundary contraction). Lean ✓ Stmt For a tensor A with bond dimension D , a left boundary covector $v_L \in \mathbb{C}^D$, a right boundary vector $v_R \in \mathbb{C}^D$, and a word $w = (i_1, \dots, i_k)$, the open-boundary contraction is the bilinear pairing

$$(v_L | A^{i_1} A^{i_2} \cdots A^{i_k} | v_R) = v_L^\top (A^{i_1} \cdots A^{i_k}) v_R.$$

Definition 15.4.2 (Open-boundary state). [Lean](#) [✓ Stmt](#) The open-boundary state on N sites assigns to each configuration $\sigma = (\sigma_1, \dots, \sigma_N)$ the amplitude $(v_L | A^{\sigma_1} \dots A^{\sigma_N} | v_R)$.

Definition 15.4.3 (W tensor and boundary vectors). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Set $d = D = 2$. The W tensor is the site-independent family

$$A^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A^1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},$$

with left boundary covector $\langle l | = \langle 0 |$ and right boundary vector $|r\rangle = |1\rangle$. Here A^1 is the single raising operator: it sends $|1\rangle \mapsto |0\rangle$ and annihilates $|0\rangle$, and squares to zero.

Definition 15.4.4 (W-state indicator). [Lean](#) [✓ Stmt](#) For $N \geq 0$, the W-state amplitude is the indicator of the single-excitation configurations:

$$|W_N\rangle(\sigma_1, \dots, \sigma_N) = \begin{cases} 1, & \text{exactly one } \sigma_j = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Theorem 15.4.5 (The open-boundary contraction is the W state). [Lean](#) [✓ Stmt](#) For every N , the open-boundary state of the W tensor with boundary vectors $\langle 0 |$ and $|1\rangle$ equals the W state:

$$|W_N\rangle = \sum_{\sigma} \langle 0 | A^{\sigma_1} \dots A^{\sigma_N} | 1 \rangle |\sigma\rangle.$$

Proof. [✓ Proof](#) Because A^0 is the identity and A^1 is the raising operator with $A^1 A^1 = 0$, the ordered product $A^{\sigma_1} \dots A^{\sigma_N}$ depends only on the number of excitations of σ : it is the identity with no excitation, the raising operator with one excitation, and zero with two or more. Reading off the $(0, 1)$ matrix element against the boundary vectors therefore gives 1 on the single-excitation configurations and 0 elsewhere, which is the W-state amplitude. $\square$

Theorem 15.4.6 (Three-site single excitations). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) On three sites, each of the three single-excitation configurations $|100\rangle$, $|010\rangle$, $|001\rangle$ receives W-state amplitude 1, while the vacuum $|000\rangle$ and the doubly-excited $|110\rangle$ receive amplitude 0.

Proof. [✓ Proof](#) Direct instances of Theorem 15.4.5: the excitation count is one in the first three cases and zero or two in the others. $\square$

Remark 15.4.7 (No fixed-bond translation-invariant family). The open-boundary representation above is not translation invariant: the chain is closed by boundary vectors, not by a trace. The review observes that there is no translation-invariant representation whose bond dimension is bounded independently of N : any translation-invariant matrix product state equal to $|W_N\rangle$ must have bond dimension growing with N , satisfying a bound of the form $D^3 \log D = \Omega(N)$. The review obtains this by combining the matrix product state representation theory of [PGVWC07] with the quantitative quantum Wielandt inequality. That asymptotic lower bound is not used in the present example.

15.5 The cluster state

The cluster state is a universal resource for measurement-based quantum computation and provides another example of a non-trivial SPT phase.

Definition 15.5.1 (Cluster-state tensor). [Lean](#) [✓ Stmt](#) Set $d = D = 2$. The cluster-state tensor is the family $\{A^0, A^1\} \subset M_2(\mathbb{C})$ defined by

$$A^0 = |+\rangle\langle 0| = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}, \quad A^1 = |-\rangle\langle 1| = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix},$$

where $|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$. The review [\[CPGSV21\]](#) writes the reflected convention $A^0 = |0\rangle\langle +|$, $A^1 = |1\rangle\langle -|$; the transpose taken here is the same state read in the opposite direction and carries the same SPT order.

Theorem 15.5.2 (Cluster-state tensor is not injective). [Lean](#) [✓ Stmt](#) *The cluster-state tensor is not injective: $\text{span}_{\mathbb{C}}\{A^0, A^1\} = \text{span}_{\mathbb{C}}\{|+\rangle\langle 0|, |-\rangle\langle 1|\}$ has dimension 2, not $\dim M_2(\mathbb{C}) = 4$.*

Proof. [✓ Proof](#) Both A^0 and A^1 are rank-one matrices with linearly independent column spaces, so they are linearly independent. Their span is two-dimensional, which is strictly less than $\dim M_2(\mathbb{C}) = 4$. $\square$

Theorem 15.5.3 (Cluster-state tensor is normal). [Lean](#) [✓ Stmt](#) *Although the cluster-state tensor is not injective at a single site, its length-2 blocking is injective: the four length-2 products span $M_2(\mathbb{C})$. In particular the cluster-state tensor is normal.*

Remark 15.5.4. Although the cluster-state tensor is not injective at length 1, the length-2 blocked tensor is injective. Using $\langle 0|\pm\rangle = \frac{1}{\sqrt{2}}$, $\langle 1|+\rangle = \frac{1}{\sqrt{2}}$, and $\langle 1|-\rangle = -\frac{1}{\sqrt{2}}$, the four products $A^{ij} = A^i A^j$ are

$$A^{00} = \frac{1}{\sqrt{2}}|+\rangle\langle 0|, \quad A^{01} = \frac{1}{\sqrt{2}}|+\rangle\langle 1|, \quad A^{10} = \frac{1}{\sqrt{2}}|-\rangle\langle 0|, \quad A^{11} = -\frac{1}{\sqrt{2}}|-\rangle\langle 1|. \quad (15.1)$$

Since $\{|+\rangle, |-\rangle\}$ and $\{|0\rangle, |1\rangle\}$ are bases, these four rank-one matrices are linearly independent and span $M_2(\mathbb{C})$.

Theorem 15.5.5 ($\mathbb{Z}_2 \times \mathbb{Z}_2$ on-site symmetry of the blocked cluster-state tensor). [Lean](#) [✓ Stmt](#) *The length-2 blocked cluster-state tensor is on-site symmetric under $\mathbb{Z}_2 \times \mathbb{Z}_2$. The two generators act on the 4-dimensional blocked physical space $(\mathbb{C}^2)^{\otimes 2}$ by $\sigma_x \otimes I$ and $I \otimes \sigma_x$, which commute and each square to the identity, defining a linear representation of $\mathbb{Z}_2 \times \mathbb{Z}_2$.*

Theorem 15.5.6 (Anticommuting virtual gauges of the cluster-state symmetry). [Lean](#) [✓ Stmt](#) *The virtual matrices implementing the two generators of Theorem 15.5.5, namely $V_1 = \sigma_z$ and $V_2 = \sigma_x$, anticommute ($V_1 V_2 = -V_2 V_1$).*

Definition 15.5.7 (Cluster-state factor system). [Lean](#) [✓ Stmt](#) The cluster-state factor system on $\mathbb{Z}_2 \times \mathbb{Z}_2$ is $\omega(g, h) = (-1)^{g_2 h_1}$ (the value -1 when $g_2 = 1$ and $h_1 = 1$, and 1 otherwise). It is the factor system of the explicit projective representation with virtual gauges σ_z and σ_x , both squaring to the identity, so there is no diagonal term.

Lemma 15.5.8 (Cluster-state commutator phase). [Lean](#) [✓ Stmt](#) *The commutator phase of the cluster-state factor system on the two generators is -1 .*

Theorem 15.5.9 (The cluster-state factor system is a 2-cocycle). [Lean](#) [✓ Stmt](#) *The cluster-state factor system is a genuine 2-cocycle, being the factor system of an explicit projective representation, so it represents an element of $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \text{U}(1))$.*

Theorem 15.5.10 (The cluster-state $\mathbb{Z}_2 \times \mathbb{Z}_2$ phase is non-trivial). [Lean](#) [✓ Stmt](#) *The cluster-state factor system (Theorem 15.5.9), built from the anticommuting gauges σ_z and σ_x (Theorem 15.5.6), is not a coboundary: its commutator phase on the two generators is -1 . Its class in $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \text{U}(1)) \cong \mathbb{Z}_2$ is therefore non-trivial, so the cluster state realizes a non-trivial SPT phase.*

Theorem 15.5.11 (The cluster state has string order under its $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry). [Lean](#) [✓ Stmt](#) For every element g of $\mathbb{Z}_2 \times \mathbb{Z}_2$, the length-2 blocked cluster-state tensor has string order in the sense of Definition 12.5.7 for the twist by g , with the maximally mixed boundary state $\Lambda = \frac{1}{2}\mathbb{1}$ as its stationary boundary state.

Proof. [✓ Proof](#) The blocked tensor is injective (Remark 15.5.4) and on-site symmetric under $\mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 15.5.5), and each group element acts by a real symmetric involutive permutation matrix, hence unitarily. The maximally mixed state $\Lambda = \frac{1}{2}\mathbb{1}$ is positive definite and has trace 1. The four blocked matrices in (15.1) satisfy $\sum_i A^i (A^i)^\dagger = \mathbb{1}$ and $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, so the transfer map is unital and Λ is a fixed point of the adjoint transfer map. These are exactly the hypotheses of Theorem 12.6.3, which then yields string order for every g . The cluster state is thus the first explicit witness of that criterion. $\square$

Theorem 15.5.12 (Blocked cluster transfer map is idempotent). [Lean](#) [Lean](#) [✓ Stmt](#) The length-2 blocked cluster transfer map satisfies $\mathcal{E}_A^2 = \mathcal{E}_A$. In closed form it sends every X to the scalar $(X_{00} + X_{11})/2$ times the identity, so its image is one-dimensional.

Proof. [✓ Proof](#) The single-site cluster transfer map is not idempotent, but the four blocked matrices give the closed form $\mathcal{E}_A(X) = \frac{1}{2}(X_{00} + X_{11})\mathbb{1}$, which is manifestly idempotent. $\square$

Theorem 15.5.13 (Cluster zero correlation length). [Lean](#) [✓ Stmt](#) The length-2 blocked cluster tensor has zero correlation length: its idempotent transfer map (Theorem 15.5.12) gives correlations independent of the separation. Like the GHZ state, and unlike the AKLT state, the cluster state is a renormalization fixed point with zero correlation length.

Proof. [✓ Proof](#) The claim follows from the idempotence of the blocked cluster transfer map and the equivalence of zero correlation length with transfer-map idempotence (Theorem 22.5.13). $\square$

15.6 Parent Hamiltonian statements for the examples

The parent-Hamiltonian construction of [CPGSV21, lines 1988–2000] is recorded here in the form

$$\mathcal{G}_L(A) = \left\{ \sum_{i_1, \dots, i_L} \text{tr}(A^{i_1} \dots A^{i_L} X) |i_1 \dots i_L\rangle : X \in M_D(\mathbb{C}) \right\}.$$

Thus

$$\ker h_L = \mathcal{G}_L(A), \quad H_N^{(L)} = \sum_i (h_L)_i.$$

We write

$$\text{GS}(H) = \ker(H - E_0(H)I)$$

for the ground eigenspace.

Theorem 15.6.1 (GHZ two-site parent space). [Lean](#) [✓ Stmt](#) For the GHZ tensor A of Definition 15.1.1,

$$\mathcal{G}_2(A) = \text{span}_{\mathbb{C}}\{|00\rangle, |11\rangle\}.$$

Proof. [✓ Proof](#) By definition,

$$\Gamma_2(X)(\sigma_1, \sigma_2) = \text{tr}(A^{\sigma_1} A^{\sigma_2} X).$$

For the GHZ matrices $A^0 = \text{diag}(1, 0)$ and $A^1 = \text{diag}(0, 1)$, whenever $\sigma_1 \neq \sigma_2$,

$$A^{\sigma_1} A^{\sigma_2} = 0.$$

Hence

$$\Gamma_2(X) = X_{00}|00\rangle + X_{11}|11\rangle,$$

and the image of Γ_2 is $\text{span}_{\mathbb{C}}\{|00\rangle, |11\rangle\}$. □

Theorem 15.6.2 (Constant GHZ product vectors satisfy the cyclic constraints). Lean ✓ Stmt
For $N \geq 2$ and $a \in \{0, 1\}$,

$$|a\rangle^{\otimes N} \in \mathcal{G}_{N,2}(A_{\text{GHZ}}).$$

Proof. ✓ Proof For a cyclic nearest-neighbour window and outside assignment τ , set

$$R_{i,\tau} := |a\rangle^{\otimes N} \Big|_{[i,i+1],\tau}.$$

For $b, c \in \{0, 1\}$ with $b \neq c$, the completed configuration is not constant, hence

$$R_{i,\tau}(b, c) = 0.$$

Therefore

$$R_{i,\tau} \in \{v : v(0, 1) = v(1, 0) = 0\} = \text{span}_{\mathbb{C}}\{|00\rangle, |11\rangle\} = \mathcal{G}_2(A_{\text{GHZ}}),$$

so $|a\rangle^{\otimes N} \in \mathcal{G}_{N,2}(A_{\text{GHZ}})$. □

Theorem 15.6.3 (GHZ periodic nearest-neighbour ground space). Lean ✓ Stmt For $N \geq 2$, the common nearest-neighbour cyclic ground space of the GHZ tensor satisfies

$$\mathcal{G}_{N,2}(A_{\text{GHZ}}) = \text{span}_{\mathbb{C}}\{|0\rangle^{\otimes N}, |1\rangle^{\otimes N}\}.$$

This is the cyclic-window form of the two-fold degeneracy in [CPGSV21, line 2205].

Proof. ✓ Proof Let $\psi \in \mathcal{G}_{N,2}(A_{\text{GHZ}})$. For every cyclic nearest-neighbour window and outside assignment τ ,

$$\psi|_{[i,i+1],\tau} \in \mathcal{G}_2(A_{\text{GHZ}}) = \text{span}_{\mathbb{C}}\{|00\rangle, |11\rangle\}.$$

Taking the outside assignment from the word σ gives

$$\psi|_{[i,i+1],\sigma}(\sigma_i, \sigma_{i+1}) = \psi(\sigma).$$

Hence a mixed nearest-neighbour word has zero coefficient:

$$\sigma_i \neq \sigma_{i+1} \implies \psi(\sigma) = 0.$$

A nonconstant binary cyclic word has a mixed edge,

$$\sigma \notin \{(0, \dots, 0), (1, \dots, 1)\} \implies \exists i, \sigma_i \neq \sigma_{i+1}.$$

Therefore

$$\psi = \psi(0, \dots, 0)|0\rangle^{\otimes N} + \psi(1, \dots, 1)|1\rangle^{\otimes N}.$$

Conversely, Theorem 15.6.2 gives

$$|0\rangle^{\otimes N}, |1\rangle^{\otimes N} \in \mathcal{G}_{N,2}(A_{\text{GHZ}}),$$

and hence their span lies in $\mathcal{G}_{N,2}(A_{\text{GHZ}})$. □

Remark 15.6.4 (GHZ parent interaction). For the GHZ tensor of Definition 15.1.1,

$$\mathcal{G}_2(A) = \text{span}\{|00\rangle, |11\rangle\}, \quad h_{\text{GHZ}} = |01\rangle\langle 01| + |10\rangle\langle 10| = I - |00\rangle\langle 00| - |11\rangle\langle 11|.$$

The periodic nearest-neighbour parent Hamiltonian therefore satisfies

$$H_{\text{GHZ},N} = \sum_i (h_{\text{GHZ}})_i, \quad \ker H_{\text{GHZ},N} = \text{span}\{|0\rangle^{\otimes N}, |1\rangle^{\otimes N}\},$$

using the cyclic-window equality

$$\mathcal{G}_{N,2}(A_{\text{GHZ}}) = \text{span}\{|0\rangle^{\otimes N}, |1\rangle^{\otimes N}\}$$

from Theorem 15.6.3. This is the ferromagnetic Ising parent Hamiltonian [CPGSV21, lines 1194–1196 and line 2205]; see also [FGSW⁺15, Section 3] for the displayed two-site space and interaction.

Remark 15.6.5 (Cluster-state stabilizer convention). The one-dimensional cluster tensor in Definition 15.5.1 is the reflected convention of the tensor displayed in [CPGSV21, lines 2364–2369]. The sourced stabilizer is

$$K_i^{ZXZ} = \sigma_i^z \sigma_{i+1}^x \sigma_{i+2}^z.$$

The corresponding positive parent interaction is

$$h_i^{ZXZ} = \frac{1}{2}(I - K_i^{ZXZ}), \quad H_{\text{cl},N}^{ZXZ} = \sum_i h_i^{ZXZ}, \quad \dim \ker H_{\text{cl},N}^{ZXZ} = 1$$

[PGVWC07, local TeX lines 374–387]. Thus an XZX formula should be used for this tensor only after establishing an equivalence such as

$$UH_{\text{cl},N}^{ZXZ}U^\dagger = H_{\text{cl},N}^{XZX} \quad \text{or} \quad TH_{\text{cl},N}^{ZXZ}T^{-1} = H_{\text{cl},N}^{XZX}.$$

Remark 15.6.6 (AKLT parent Hamiltonian and edge modes). For the AKLT tensor of Definition 15.2.1, with $\mathcal{G}_{[a,b]}$ denoting the local ground space on sites $a, \dots, b$,

$$L_0 = 2, \quad \mathcal{G}_{[1,2]} \cap \mathcal{G}_{[2,3]} = \mathcal{G}_{[1,3]}$$

[CPGSV21, line 2095]. The displayed local spin-chain Hamiltonian is

$$H_{\text{AKLT},N} = \sum_i \left(\vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3}(\vec{S}_i \cdot \vec{S}_{i+1})^2 \right),$$

up to an additive constant [PGVWC07, local TeX lines 340–349]; [SPGC11, lines 2126–2130]. The spin-2 projector form is related to the polynomial Hamiltonian by the spin-addition identity

$$P_{i,i+1}^{(2)} = \frac{1}{2} \left(\vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3}(\vec{S}_i \cdot \vec{S}_{i+1})^2 + \frac{2}{3}I \right)$$

for two spin-1 particles. For open boundary conditions,

$$\dim \text{GS}(H_{\text{AKLT},N}^{\text{open}}) = 4, \quad \dim(\text{GS}(H_{\text{AKLT},N}^{\text{open}}) \cap \mathcal{H}_{S=0}) = 1$$

[CPGSV21, lines 1169–1174].

Chapter 16

Channel Representations and Normal Forms

This chapter collects the representation material for finite-dimensional quantum channels from [Wol12, Chapter 2]: the Choi–Jamiołkowski isomorphism, Kraus representation theorems, Stinespring and Naimark dilations, ordered completely positive maps, Radon–Nikodym and open-system representations, trace-pairing expansions, SVD and Lorentz normal forms, and the channel determinant. These results are important for the later channel theory, but they are not prerequisites for the matrix-product-state Fundamental Theorem.

16.1 Representations

The following definitions support the Choi–Jamiołkowski isomorphism (Theorem 16.2.4) and the representation theorems of [Wol12, Chapter 2].

Definition 16.1.1 (Partial traces). [Lean](#) [Lean](#) [✓ Stmt](#) Let $X \in M_{d,d'}(\mathbb{C})$ be a bipartite matrix indexed by

$$(\{0, \dots, d-1\} \times \{0, \dots, d'-1\})^2.$$

The *left partial trace* $\text{tr}_A(X)$ and *right partial trace* $\text{tr}_B(X)$ are the $d' \times d'$ and $d \times d$ matrices defined by

$$(\text{tr}_A(X))_{ij} = \sum_k X_{(k,i),(k,j)}, \quad (\text{tr}_B(X))_{ij} = \sum_k X_{(i,k),(j,k)}.$$

Definition 16.1.2 (Maximally entangled state). [Lean](#) [✓ Stmt](#) The *maximally entangled state* on $\mathbb{C}^d \otimes \mathbb{C}^d$ is the rank-one projector

$$|\Omega\rangle\langle\Omega|, \quad |\Omega\rangle = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |j, j\rangle.$$

Concretely, $(|\Omega\rangle\langle\Omega|)_{(i_1,i_2),(j_1,j_2)} = \frac{1}{d} \delta_{i_1 j_1} \delta_{i_2 j_2}$ and $\text{tr}(|\Omega\rangle\langle\Omega|) = 1$.

Definition 16.1.3 (Flip operator). [Lean](#) [✓ Stmt](#) The *flip operator* F on $\mathbb{C}^d \otimes \mathbb{C}^d$ is

$$F = \sum_{i,j=0}^{d-1} |ij\rangle\langle ji|.$$

Its matrix entries are

$$F_{(i_1, i_2), (j_1, j_2)} = \begin{cases} 1, & i_1 = j_2 \text{ and } i_2 = j_1, \\ 0, & \text{otherwise.} \end{cases}$$

Definition 16.1.4 (Two-dimensional antisymmetric vector). [Lean](#) [✓ Stmt](#) In $\mathbb{C}^2 \otimes \mathbb{C}^2$, let

$$v = |01\rangle - |10\rangle.$$

Equivalently, $v_{(0,1)} = 1$, $v_{(1,0)} = -1$, and all other coefficients are zero.


Definition 16.1.5 (Tensor product of a map with identity). [Lean](#) [✓ Stmt](#) Given a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the *tensor extension* $T \otimes \text{id}$ acts on bipartite matrices $X \in M_{D \times D}(\mathbb{C})$ by applying T to each “slice”:

$$(T \otimes \text{id})(X)_{(i_1, i_2), (j_1, j_2)} = (T(X^{(i_2, j_2)}))_{i_1 j_1},$$

where $X_{ab}^{(i_2, j_2)} = X_{(a, i_2), (b, j_2)}$ is the bipartite slice.

16.2 Choi–Jamiołkowski isomorphism

The Choi–Jamiołkowski isomorphism bends the input legs of the channel T around the maximally entangled pair $|\Omega\rangle$, presenting the map as the matrix $\tau = (T \otimes \text{id})|\Omega\rangle\langle\Omega|$ on the doubled space:



Definition 16.2.1 (Choi matrix). [Lean](#) [✓ Stmt](#) The *Choi matrix* of a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is

$$\tau = (T \otimes \text{id})(|\Omega\rangle\langle\Omega|) \in M_{D \times D}(\mathbb{C}).$$

Concretely, $\tau_{(i_1, i_2), (j_1, j_2)} = \frac{1}{D} (T(E_{i_2 j_2}))_{i_1 j_1}$ where $E_{i_2 j_2}$ is the matrix unit. This is the Choi–Jamiołkowski convention of [Wol12, Proposition 2.1].

Theorem 16.2.2 (Choi matrix of the identity map). [Lean](#) [✓ Stmt](#) The identity channel has Choi matrix $|\Omega\rangle\langle\Omega|$.

Proof. [✓ Proof](#) Apply the identity map in the definition of the Choi matrix. □

Theorem 16.2.3 (Choi matrix of transposition). [Lean](#) [✓ Stmt](#) Let $\theta(X) = X^T$ be matrix transposition on $M_D(\mathbb{C})$, with $D \geq 1$. Then

$$(\theta \otimes \text{id})(|\Omega\rangle\langle\Omega|) = \frac{1}{D} F.$$

This is [Wol12, Equation 3.1].

Proof. [✓ Proof](#) The (i_2, j_2) slice of $|\Omega\rangle\langle\Omega|$ is $D^{-1} E_{i_2 j_2}$. Transposition sends this matrix unit to $D^{-1} E_{j_2 i_2}$, whose (i_1, j_1) entry is D^{-1} exactly when $i_1 = j_2$ and $i_2 = j_1$. □

Theorem 16.2.4 (Complete positivity iff the Choi matrix is positive semidefinite). [Lean](#) [✓ Stmt](#) A linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is completely positive (in the Kraus sense) if and only if its Choi matrix $\tau \geq 0$.

Proof. ✓ Proof ($\Rightarrow$) If $T(X) = \sum_i K_i X K_i^\dagger$, then $\tau = \sum_i |v_i\rangle\langle v_i|$ where $(v_i)_{(a,b)} = c K_i(a, b)$ and $c = 1/\sqrt{D}$. This is a sum of rank-one PSD matrices.

($\Leftarrow$) If $\tau \geq 0$, write $\tau = \sum_m |v_m\rangle\langle v_m|$ by spectral decomposition, define $K_m(a, b) = v_m(a, b)/c$, and recover $T(X) = \sum_m K_m X K_m^\dagger$ by linearity on matrix units. $\square$

Theorem 16.2.5 (Trace preservation implies the partial-trace condition). Lean ✓ Stmt If T is trace-preserving, then $\text{tr}_A(\tau) = \frac{1}{D} \mathbb{1}_D$.

Proof. ✓ Proof $(\text{tr}_A(\tau))_{ij} = \text{tr}(T(\Omega^{(ij)})) = \text{tr}(\Omega^{(ij)})$ by trace preservation, where $\Omega^{(ij)} = \frac{1}{D} E_{ij}$ is the (i, j) -slice of $|\Omega\rangle\langle\Omega|$. Taking the trace gives $\frac{1}{D} \delta_{ij}$. $\square$

Theorem 16.2.6 (Hermiticity preservation and the Choi matrix). Lean ✓ Stmt The Choi matrix τ is Hermitian if and only if $T(B^\dagger) = T(B)^\dagger$ for all $B \in M_D(\mathbb{C})$.

Proof. ✓ Proof The Choi matrix entries satisfy $\tau_{(a,i),(b,j)} = \frac{1}{D} (T(E_{ij}))_{ab}$. Hermiticity of τ says $\overline{\tau_{(b,j),(a,i)}} = \tau_{(a,i),(b,j)}$, which unravels to $T(E_{ji})_{ba} = \overline{T(E_{ij})_{ab}}$, i.e., $T(E_{ij}^\dagger) = T(E_{ij})^\dagger$. By linearity this extends to all B . $\square$

Theorem 16.2.7 (Trace of the Choi matrix of a trace-preserving map). Lean ✓ Stmt If T is trace-preserving, then $\text{tr}(\tau) = 1$.

Proof. ✓ Proof $\text{tr}(\tau) = \text{tr}(\text{tr}_A(\tau)) = \text{tr}(\frac{1}{D} \mathbb{1}_D) = 1$. $\square$

16.3 Representation corollaries for channel decompositions

The next three corollaries follow from the Choi/Kraus/Stinespring representations already developed above. They correspond to [Wol12, Propositions 2.2–2.4].

Theorem 16.3.1 (Polarization of sesquilinear sandwiches). Lean ✓ Stmt For any $A, B, X \in M_D(\mathbb{C})$,

$$4AXB^\dagger = (A+B)X(A+B)^\dagger - (A-B)X(A-B)^\dagger + i(A+iB)X(A+iB)^\dagger - i(A-iB)X(A-iB)^\dagger.$$

Each of the four summands on the right is a single-Kraus CP map, so every sesquilinear sandwich decomposes into a signed complex linear combination of CP maps.

Proof. ✓ Proof Working entrywise, reduces to the scalar polarization identity $4\alpha\bar{\delta} = \sum_{k=0}^3 i^k (\alpha + i^k \beta)(\gamma + i^k \delta)$ in $\mathbb{C}$, which holds after substituting $i^2 = -1$. $\square$

Theorem 16.3.2 (CP decomposition of sandwich sums). Lean ✓ Stmt For any finite Kraus-like families $\{A_i\}, \{B_i\}$, the map $T(X) = \sum_i A_i X B_i^\dagger$ satisfies $4T = T_1 - T_2 + iT_3 - iT_4$ with each T_k a CP map given explicitly by a single-side Kraus sum of the combined families.

Proof. ✓ Proof Apply the polarization identity summand-by-summand. $\square$

Theorem 16.3.3 (No information without disturbance). Lean ✓ Stmt Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be linear. If $T(|v\rangle\langle v|) = |v\rangle\langle v|$ for every $v \in \mathbb{C}^D$, then $T = \text{id}$. In particular, a quantum channel preserving every pure state equals the identity.

Proof. ✓ Proof Rank-one outer products $|u\rangle\langle v|$ polarize into rank-one self-outer-products:

$$4|u\rangle\langle v| = \sum_{k=0}^3 i^k |u + i^k v\rangle\langle u + i^k v|,$$

so T preserves every $|u\rangle\langle v|$. Since the matrix units $E_{ij} = |e_i\rangle\langle e_j|$ span $M_D(\mathbb{C})$, T equals the identity on a spanning set, hence everywhere. $\square$

Definition 16.3.4 (Pure-state ensemble density). Lean ✓ Stmt Given a finite family $\{\psi_i\}_{i \in \iota}$ of (unnormalized) vectors in $\mathbb{C}^D$, its *pure-ensemble density* is $\rho = \sum_i |\psi_i\rangle\langle\psi_i|$. Weights $p_i \geq 0$ with $\sum p_i = 1$ can be absorbed by replacing $\psi_i \mapsto \sqrt{p_i}\psi_i$.

Theorem 16.3.5 (Isometric mixing preserves the density). Lean ✓ Stmt If two pure-state ensembles $\{\psi_i\}_{i \in \iota_1}$ and $\{\phi_j\}_{j \in \iota_2}$ are related by an isometric mixing matrix $V \in \mathbb{C}^{\iota_1 \times \iota_2}$ with $V^\dagger V = \mathbb{1}$ and $\psi_i = \sum_j V_{ij} \phi_j$, then they induce the same pure-ensemble density operator. This is the sufficient direction of the Hughston–Jozsa–Wootters theorem; the converse is Theorem 16.3.6.

Proof. ✓ Proof Expanding and applying the orthogonality relation $\sum_i V_{ij} \overline{V_{ij'}} = \delta_{jj'}$ from $V^\dagger V = \mathbb{1}$:

$$\sum_i |\psi_i\rangle\langle\psi_i| = \sum_{j,j'} \left(\sum_i V_{ij} \overline{V_{ij'}} \right) |\phi_j\rangle\langle\phi_{j'}| = \sum_j |\phi_j\rangle\langle\phi_j|.$$

$\square$

Theorem 16.3.6 (Hughston–Jozsa–Wootters converse). Lean ✓ Stmt If two pure-state ensembles $\{\psi_i\}_{i \in \iota_1}$ and $\{\phi_j\}_{j \in \iota_2}$ induce the same pure-ensemble density operator and $|\iota_2| \leq |\iota_1|$, then there exists a tall isometric mixing matrix $V \in \mathbb{C}^{\iota_1 \times \iota_2}$ with $V^\dagger V = \mathbb{1}$ and $\psi_i = \sum_j V_{ij} \phi_j$. The cardinality hypothesis is what makes V a tall isometry; the symmetric case $|\iota_1| \leq |\iota_2|$ follows by swapping the roles of the ensembles.

Proof. ✓ Proof Embed each vector ψ_i, ϕ_j as the 0-th column of a $D \times D$ matrix with zeros elsewhere; call these $K_i, L_j \in M_D(\mathbb{C})$. A direct entry-wise computation shows that for any $X \in M_D(\mathbb{C})$ both $\sum_i K_i X K_i^\dagger$ and $\sum_j L_j X L_j^\dagger$ collapse to $X_{00} \cdot \rho$, so the density equality $\rho_\psi = \rho_\phi$ forces the two Kraus families to define the same CP map. Theorem 16.4.10 then supplies an isometry V with $V^\dagger V = \mathbb{1}$ and $K_i = \sum_j V_{ij} L_j$; reading the equation off at column 0 recovers the vector relation $\psi_i = \sum_j V_{ij} \phi_j$. $\square$

Theorem 16.3.7 (Hughston–Jozsa–Wootters equivalence). Lean ✓ Stmt Under the cardinality hypothesis $|\iota_2| \leq |\iota_1|$, two pure-state ensembles induce the same density operator iff they are related by a tall isometric mixing matrix.

Proof. ✓ Proof Combine the two directions above. $\square$

16.4 Kraus representation theorem

A Kraus representation writes the channel as $T(\rho) = \sum_i K_i \rho K_i^\dagger$. In tensor-network form the Kraus index i is summed between the K_i layer and the $K_i^\dagger$ layer:

Theorem 16.4.1 (Trace preservation implies Kraus normalization). [Lean](#) [✓ Stmt](#) If $T(X) = \sum_i K_i X K_i^\dagger$ is trace-preserving, then $\sum_i K_i^\dagger K_i = \mathbb{1}$.

Proof. [✓ Proof](#) For any N , $\text{tr}((\sum_i K_i^\dagger K_i) N) = \sum_i \text{tr}(K_i^\dagger K_i N) = \sum_i \text{tr}(K_i N K_i^\dagger) = \text{tr}(T(N)) = \text{tr}(N)$. Non-degeneracy of the trace pairing forces $\sum_i K_i^\dagger K_i = \mathbb{1}$. $\square$

Theorem 16.4.2 (Kraus normalization implies trace preservation). [Lean](#) [✓ Stmt](#) If $\sum_i K_i^\dagger K_i = \mathbb{1}$, then $T(X) = \sum_i K_i X K_i^\dagger$ is trace-preserving.

Proof. [✓ Proof](#) $\text{tr}(T(X)) = \sum_i \text{tr}(K_i X K_i^\dagger) = \sum_i \text{tr}(K_i^\dagger K_i X) = \text{tr}((\sum_i K_i^\dagger K_i) X) = \text{tr}(X)$. $\square$

Theorem 16.4.3 (Unitality implies Kraus unital normalization). [Lean](#) [✓ Stmt](#) If $T(X) = \sum_i K_i X K_i^\dagger$ satisfies $T(\mathbb{1}) = \mathbb{1}$, then

$$\sum_i K_i K_i^\dagger = \mathbb{1}.$$

Proof. [✓ Proof](#) Evaluate the Kraus formula at the identity matrix. $\square$

Theorem 16.4.4 (Unitary freedom in Kraus operators). [Lean](#) [✓ Stmt](#) Let U be a unitary $r \times r$ matrix ($U^\dagger U = \mathbb{1}$) and suppose $K_j = \sum_\ell U_{j\ell} \tilde{K}_\ell$. Then $\{K_j\}$ and $\{\tilde{K}_\ell\}$ define the same Kraus map: for every $X \in M_D(\mathbb{C})$,

$$\sum_j K_j X K_j^\dagger = \sum_\ell \tilde{K}_\ell X \tilde{K}_\ell^\dagger.$$

Proof. [✓ Proof](#) Substitute the combination formula, expand the triple sum over j, ℓ, ℓ' , swap summation order, and use the entry-wise form of $U^\dagger U = \mathbb{1}$ to collapse to $\ell = \ell'$ diagonal terms. $\square$

Theorem 16.4.5 (Isometric mixing preserves the Kraus map). [Lean](#) [✓ Stmt](#) Let W be an isometry between two Kraus index spaces, so $W^\dagger W = \mathbb{1}$. If

$$K_j = \sum_\ell W_{j\ell} \tilde{K}_\ell,$$

then the Kraus families $\{K_j\}$ and $\{\tilde{K}_\ell\}$ define the same completely positive map.

Proof. [✓ Proof](#) Expand the Kraus sums, interchange the finite summations, and use $W^\dagger W = \mathbb{1}$ to collapse the coefficient matrix. $\square$

Theorem 16.4.6 (The transition matrix of orthonormal Kraus families is unitary). [Lean](#) [✓ Stmt](#) Let $\{K_j\}_{j=0}^{r-1}$ and $\{K'_j\}_{j=0}^{r-1}$ be two Hilbert–Schmidt orthonormal Kraus families, and suppose

$$K_j = \sum_{\ell=0}^{r-1} U_{j\ell} K'_\ell.$$

Then the transition matrix U is unitary: $U^\dagger U = \mathbb{1}_r$.

Proof. [✓ Proof](#) Expand $\text{tr}(K_j^\dagger K_i)$ using the change of basis. Hilbert–Schmidt orthonormality of both Kraus families identifies the Gram matrix with the identity, yielding the matrix identity $U U^\dagger = \mathbb{1}_r$, hence also $U^\dagger U = \mathbb{1}_r$. $\square$

Conversely, diagonalizing the positive semidefinite Choi matrix and keeping only the nonzero eigenvalue summands produces a Kraus family with exactly $\text{rank}(\tau_E)$ operators. Hence the Choi rank is precisely the minimal Kraus cardinality.

Proof. [✓ Proof](#) For the forward implication, expand the Choi matrix using the Kraus formula and write $\tau_E = \sum_j v_j v_j^\dagger$ with one vector v_j for each Kraus operator. The column space of this sum lies in the span of the r vectors $\{v_j\}$, so the rank is at most r . For the converse, use the spectral decomposition of the positive semidefinite Choi matrix, discard the zero-eigenvalue terms, and rescale the remaining eigenvectors into Kraus operators. $\square$

16.5 Stinespring dilation

The Stinespring dilation realizes the channel through an isometry V into a larger space, $T(\rho) = \text{tr}_E[V\rho V^\dagger]$; tracing out the environment E returns the channel:

$$\begin{array}{c} \rho \\ \hline \bullet \\ \hline \bullet \\ \hline V^\dagger \end{array} \begin{array}{c} \\ Y \\ E \\ \\ \end{array} \begin{array}{c} \\ \text{tr}_E[V\rho V^\dagger] \\ \\ \end{array}$$

Definition 16.5.1 (Stinespring isometry). [Lean](#) [✓ Stmt](#) Given Kraus operators $\{K_j\}_{j=0}^{r-1}$ with $K_j \in M_D(\mathbb{C})$, the *Stinespring isometry* $V : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^r$ is the $(D \cdot r) \times D$ matrix

$$V_{(i,j),k} = (K_j)_{ik}.$$

This implements $V = \sum_j K_j \otimes |j\rangle$.

Lemma 16.5.2 (Entry formula for the Stinespring isometry). [Lean](#) [✓ Stmt](#) For indices $i, k \in \{0, \dots, D-1\}$ and $j \in \{0, \dots, r-1\}$,

$$V_{(i,j),k} = (K_j)_{ik}.$$

Theorem 16.5.3 (Stinespring Gram identity). [Lean](#) [✓ Stmt](#) The Stinespring isometry satisfies

$$V^\dagger V = \sum_{j=0}^{r-1} K_j^\dagger K_j.$$

Theorem 16.5.4 (Stinespring dual representation). [Lean](#) [✓ Stmt](#) For Kraus operators $\{K_j\}$ and $A \in M_D(\mathbb{C})$,

$$V^\dagger (A \otimes \mathbb{1}_r) V = \sum_j K_j^\dagger A K_j = T^*(A).$$

This is the Stinespring representation of the dual (Heisenberg picture) map T^* .

Proof. [✓ Proof](#) Compute $(V^\dagger (A \otimes \mathbb{1}_r) V)_{ab}$ by summing over the product index (i, j) ; the δ_{jl} from $\mathbb{1}_r$ collapses the sum over l , giving $\sum_j K_j^\dagger A K_j$. $\square$

Theorem 16.5.5 (The Stinespring isometry condition iff trace preservation). [Lean](#) [✓ Stmt](#) $V^\dagger V = \mathbb{1}_D$ if and only if $\sum_j K_j^\dagger K_j = \mathbb{1}_D$. In particular, V is an isometry precisely when the Kraus map is trace-preserving.

Proof. [✓ Proof](#) $(V^\dagger V)_{ab} = \sum_{(i,j)} \overline{V_{(i,j),a}} V_{(i,j),b} = \sum_j \sum_i \overline{(K_j)_{ia}} (K_j)_{ib} = (\sum_j K_j^\dagger K_j)_{ab}$. $\square$

Theorem 16.5.6 (Stinespring Schrödinger representation). [Lean](#) [✓ Stmt](#) *The Kraus map $T(\rho) = \sum_j K_j \rho K_j^\dagger$ equals the partial trace over the dilation space:*

$$T(\rho)_{ij} = \sum_{k=0}^{r-1} (V \rho V^\dagger)_{(i,k),(j,k)} = \text{tr}_r(V \rho V^\dagger).$$

This is the Schrödinger-picture Stinespring representation.

Proof. [✓ Proof](#) Expand $(V \rho V^\dagger)_{(i,k),(j,k)}$ and sum over k to recover $\sum_k K_k \rho K_k^\dagger$. □

Theorem 16.5.7 (Existence of a Stinespring dilation). [Lean](#) [✓ Stmt](#) *Every completely positive map admits an ancilla dimension r , Kraus operators $\{K_j\}_{j=0}^{r-1}$, and the concrete representation $\pi(A) = A \otimes \mathbb{1}_r$ such that*

$$E(A) = V^\dagger \pi(A) V.$$

Proof. [✓ Proof](#) Choose a Kraus representation of E and apply the explicit Heisenberg-picture Stinespring formula. □

Theorem 16.5.8 (CPTP maps admit isometric Stinespring dilations). [Lean](#) [✓ Stmt](#) *Every quantum channel admits a Stinespring dilation whose matrix V is an isometry, equivalently $V^\dagger V = \mathbb{1}$.*

Proof. [✓ Proof](#) Choose Kraus operators for the channel and use trace preservation to obtain the Stinespring isometry condition $V^\dagger V = \mathbb{1}$. □

16.6 Ordered CP maps, Radon–Nikodym, and open-system representation

The next three results form the structural part of [Wol12, Theorems 2.3–2.5]: the ordered CP-map theorem, the Radon–Nikodym theorem for completely positive maps, and the open-system representation. They refine the Stinespring dilation by tracking how two CP maps related by domination or decomposition sit inside a common dilation space.

Definition 16.6.1 (CP partial order). [Lean](#) [✓ Stmt](#) For linear maps $S, T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, write $T \leq S$ (in the CP order) when $S - T$ is completely positive.

Definition 16.6.2 (Block-top contraction). [Lean](#) [✓ Stmt](#) For natural numbers r, s , let $C = C_{r,s} \in \mathbb{C}^{r \times (r+s)}$ be the rectangular $r \times (r+s)$ matrix whose rows are the first r rows of the identity on $\mathbb{C}^{r+s}$: $C_{ij} = 1$ if $j = i < r$ and 0 otherwise.

Lemma 16.6.3 (C is a co-isometry). [Lean](#) [✓ Stmt](#) *The block-top matrix satisfies $CC^\dagger = \mathbb{1}_r$.*

Proof. [✓ Proof](#) Direct entrywise computation: $(CC^\dagger)_{ii'} = \sum_j C_{ij} \overline{C_{i'j}}$ collapses to $\delta_{ii'}$. □

Lemma 16.6.4 (C is a contraction). [Lean](#) [✓ Stmt](#) *The block-top matrix satisfies $C^\dagger C \leq \mathbb{1}_{r+s}$.*

Proof. [✓ Proof](#) The matrix $C^\dagger C$ is diagonal with a 1 on the first r coordinates and a 0 on the last s , so $\mathbb{1} - C^\dagger C$ is positive semidefinite. □

Lemma 16.6.5 (Intertwining of Stinespring isometries). [Lean](#) [✓ Stmt](#) *Let $K : \{0, \dots, r-1\} \rightarrow M_D(\mathbb{C})$ and $L : \{0, \dots, s-1\} \rightarrow M_D(\mathbb{C})$ be two Kraus families. Write $K ++ L$ for their concatenation as a family indexed by $\{0, \dots, r+s-1\}$. Then*

$$V_K = (\mathbb{1}_D \otimes C_{r,s}) V_{K++L}.$$

Proof. ✓ Proof Entrywise expansion of both sides: the Kronecker product with $C_{r,s}$ picks out the first r coordinates of the dilation space, matching the action of V_{K++L} on those coordinates, which is precisely V_K . $\square$

Theorem 16.6.6 (Ordered CP maps via Stinespring contractions). Lean ✓ Stmt *Let $T_1, T_2 : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be CP maps with $T_1 \leq T_2$. Then there exist an ancilla dimension m , Heisenberg-form Stinespring matrices V_1, V_2 realizing T_1, T_2 via $T_i(A) = V_i^\dagger(A \otimes \mathbb{1})V_i$, and a contraction $\tilde{C}$ on the dilation space with $\tilde{C}^\dagger \tilde{C} \leq \mathbb{1}$ such that*

$$V_1 = (\mathbb{1}_D \otimes \tilde{C}) V_2.$$

This is [Wol12, Theorem 2.3].

Proof. ✓ Proof Choose a Heisenberg-form Kraus family K_1 of T_1 with Stinespring matrix $V_1 = V_{K_1}$, and Kraus operators L of the CP map $T_2 - T_1$ in Schrödinger orientation. Let $L^\dagger$ denote the conjugate-transposed family. Set $m = r_1 + s$ and let $K_2 = K_1 ++ L^\dagger$. Then $T_2(A) = T_1(A) + (T_2 - T_1)(A) = \sum_i (K_1^i)^\dagger A K_1^i + \sum_j L_j A L_j^\dagger$, and the latter equals $\sum_i (K_1^i)^\dagger A K_1^i + \sum_j (L_j^\dagger)^\dagger A L_j^\dagger$ which is the Heisenberg form for K_2 , yielding $T_2(A) = V_{K_2}^\dagger(A \otimes \mathbb{1})V_{K_2}$. Taking $\tilde{C} = C_{r_1,s}$ the intertwining $V_{K_1} = (\mathbb{1}_D \otimes \tilde{C})V_{K_2}$ is Lemma 16.6.5, and $\tilde{C}^\dagger \tilde{C} \leq \mathbb{1}$ is Lemma 16.6.4. $\square$

Definition 16.6.7 (Block-diagonal projectors on the dilation space). Lean Lean ✓ Stmt For r, s , let $P_{\text{top}} = C_{r,s}^\dagger C_{r,s}$ and $P_{\text{bot}} = \mathbb{1} - P_{\text{top}}$. Both are PSD (for P_{bot} , by Lemma 16.6.4) and $P_{\text{top}} + P_{\text{bot}} = \mathbb{1}$.

Theorem 16.6.8 (Radon–Nikodym theorem for finite quantum instruments). Lean ✓ Stmt *Let $\{T_i\}_{i \in I}$ be a nonempty finite family of completely positive maps, let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ satisfy $\sum_{i \in I} T_i = T$, and suppose that T has a supplied Stinespring representation*

$$T(A) = V^\dagger(A \otimes \mathbb{1}_r)V.$$

Then there are positive semidefinite operators $P_i \in M_r(\mathbb{C})$ with

$$\sum_{i \in I} P_i = \mathbb{1}_r.$$

For every $i \in I$ and $A \in M_D(\mathbb{C})$,

$$T_i(A) = V^\dagger(A \otimes P_i)V$$

This is [Wol12, Theorem 2.4], with the nonempty-family condition made explicit; see docs/paper-gaps/wolf_radon_nikodym_nonempty_family.tex.

Proof. ✓ Proof Choose Kraus families for the component maps T_i and group all their Kraus operators into one finite family. Add zero Kraus operators to one component so that this grouped family has at least as many rows as the supplied Stinespring dilation. Write the grouped Kraus operators as B_α , with a label map $\ell(\alpha) \in I$, and write the Kraus operators obtained from the blocks of V as A_j . Rectangular Kraus freedom gives a matrix W with

$$W^\dagger W = \mathbb{1}_r, \quad B_\alpha = \sum_{j=0}^{r-1} W_{\alpha j} A_j.$$

For the fibre over i , set

$$(C_i)_{\alpha j} = \begin{cases} \overline{W_{\alpha j}}, & \ell(\alpha) = i, \\ 0, & \ell(\alpha) \neq i, \end{cases} \quad P_i = C_i^\dagger C_i.$$

Then $P_i \geq 0$ and $\sum_i P_i = \mathbb{1}_r$ follows from $W^\dagger W = \mathbb{1}_r$. Finally,

$$V^\dagger(A \otimes P_i)V = \sum_{\ell(\alpha)=i} B_\alpha A B_\alpha^\dagger = T_i(A),$$

which proves the claim. $\square$

Theorem 16.6.9 (Binary Radon–Nikodym theorem for CP maps). [Lean](#) [✓ Stmt](#) *Let $T_1, T_2 : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be completely positive. There exist an ancilla dimension m , a Kraus family K with Stinespring matrix $V = V_K$, and positive semidefinite operators $P_1, P_2 : \mathbb{C}^m \rightarrow \mathbb{C}^m$ with $P_1 + P_2 = \mathbb{1}_m$ such that, for $i = 1, 2$ and every $A \in M_D(\mathbb{C})$,*

$$T_i(A) = V^\dagger(A \otimes P_i)V.$$

Remark 16.6.10 (Relation with Wolf’s Radon–Nikodym theorem). Theorem 16.6.8 is the source-faithful finite-family statement relative to a supplied Stinespring dilation. Theorem 16.6.9 is the earlier binary block-diagonal corollary in which the common dilation is constructed from the Kraus families of T_1 and T_2 .

Proof. [✓ Proof](#) Take Heisenberg-form Kraus families K_1 for T_1 and Schrödinger-form L for T_2 , and form $K = K_1 ++ L^\dagger$ on $\mathbb{C}^{r_1+s}$. Choose $P_1 = P_{\text{top}}$, $P_2 = P_{\text{bot}}$ as in Definition 16.6.7. The Kronecker identity $A \otimes (C^\dagger C) = (\mathbb{1} \otimes C)^\dagger (A \otimes \mathbb{1})(\mathbb{1} \otimes C)$ rewrites $V^\dagger(A \otimes P_{\text{top}})V$ as $((\mathbb{1} \otimes C)V)^\dagger (A \otimes \mathbb{1})(\mathbb{1} \otimes C)V$, which equals $V_{K_1}^\dagger (A \otimes \mathbb{1}) V_{K_1} = T_1(A)$ by the intertwining Lemma 16.6.5. For the complementary block, $A \otimes P_{\text{bot}} = A \otimes \mathbb{1} - A \otimes P_{\text{top}}$ and sandwiching by V yields $(T_1 + T_2)(A) - T_1(A) = T_2(A)$. $\square$

Theorem 16.6.11 (Open-system representation of quantum channels). [Lean](#) [✓ Stmt](#) *Every quantum channel $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ admits a Stinespring isometry V on an ancilla space $\mathbb{C}^r$ such that*

$$T(\rho)_{ij} = \sum_{k=0}^{r-1} (V \rho V^\dagger)_{(i,k),(j,k)} = \text{tr}_r(V \rho V^\dagger).$$

Proof. [✓ Proof](#) Apply the existence of an isometric Stinespring dilation (Theorem 16.5.8) and the Schrödinger-picture identity (Theorem 16.5.6). $\square$

Theorem 16.6.12 (Open-system representation, partial-trace form). [Lean](#) [✓ Stmt](#) *Every quantum channel T admits an isometric Stinespring dilation V such that $T(\rho) = \text{tr}_E(V \rho V^\dagger)$.*

Proof. [✓ Proof](#) Rewrite the componentwise identity of Theorem 16.6.11 as a partial trace over the ancilla factor. $\square$

Lemma 16.6.13 (Rectangular Gram equality gives a unitary). [Lean](#) [✓ Stmt](#) *Let $A, B : \mathbb{C}^n \rightarrow \mathbb{C}^m$ be linear maps with $B^\dagger B = A^\dagger A$. Then there is a unitary U on $\mathbb{C}^m$ such that $B = UA$.*

Proof. [✓ Proof](#) $\square$

Definition 16.6.14 (First environment embedding). [Lean](#) [✓ Stmt](#) For $r \geq 1$, let $W_0 : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^r$ be the isometry $W_0 x = x \otimes e_0$.

Lemma 16.6.15 (First environment embedding is an isometry). [Lean](#) [✓ Stmt](#) *The first-environment embedding satisfies $W_0^\dagger W_0 = \mathbb{1}$.*

Proof. [✓ Proof](#) □

Theorem 16.6.16 (Open-system representation, unitary form). [Lean](#) [✓ Stmt](#) *For $D \geq 1$, every quantum channel T on $\mathbb{C}^D$ admits an environment dimension $r \geq 1$ and a unitary U on $\mathbb{C}^D \otimes \mathbb{C}^r$ such that, for every matrix ρ on $\mathbb{C}^D$,*

$$T(\rho) = \text{tr}_E(UW_0\rho W_0^\dagger U^\dagger).$$

Proof. [✓ Proof](#) Take the isometric open-system representation $T(\rho) = \text{tr}_E(V\rho V^\dagger)$. Since $V^\dagger V = \mathbb{1} = W_0^\dagger W_0$, Lemma 16.6.13 gives a unitary U with $V = UW_0$. Substituting this identity into the partial trace formula gives the displayed expression. □

16.7 POVMs and Naimark dilation

Definition 16.7.1 (Positive operator-valued measure). [Lean](#) [✓ Stmt](#) *A positive operator-valued measure with n outcomes on $\mathbb{C}^D$ is a family $\{E_i\}_{i=0}^{n-1}$ of positive semidefinite operators on $\mathbb{C}^D$ satisfying the resolution of identity*

$$\sum_{i=0}^{n-1} E_i = \mathbb{1}_D.$$

Definition 16.7.2 (Naimark Kraus square roots). [Lean](#) [Lean](#) [✓ Stmt](#) *For a POVM $\{E_i\}$, each effect $E_i \geq 0$ admits a square-root factorisation $E_i = M_i^\dagger M_i$ with $M_i \in M_D(\mathbb{C})$. The operators M_i are the *Naimark Kraus square roots*.*

Theorem 16.7.3 (Naimark square roots satisfy the Kraus normalization). [Lean](#) [✓ Stmt](#) *For a POVM $\{E_i\}$ with square roots $E_i = M_i^\dagger M_i$, one has*

$$\sum_i M_i^\dagger M_i = \mathbb{1}.$$

Proof. [✓ Proof](#) Sum the identities $E_i = M_i^\dagger M_i$ and use the defining resolution of the identity for the POVM. □

Definition 16.7.4 (Naimark isometry). [Lean](#) [✓ Stmt](#) *The *Naimark isometry* $V : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^n$ of a POVM is the Stinespring-type construction*

$$V = \sum_{i=0}^{n-1} M_i \otimes |i\rangle, \quad V_{(a,i),k} = (M_i)_{a,k}.$$

Definition 16.7.5 (Naimark projective measurement). [Lean](#) [Lean](#) [✓ Stmt](#) *The *Naimark projectors* on $\mathbb{C}^D \otimes \mathbb{C}^n$ are*

$$P_i = \mathbb{1}_D \otimes |i\rangle\langle i|, \quad (P_i)_{(a,b),(c,d)} = \delta_{a,c} \delta_{b,i} \delta_{d,i}.$$

Theorem 16.7.6 (Naimark isometry condition). [Lean](#) [✓ Stmt](#) *The Naimark isometry satisfies $V^\dagger V = \mathbb{1}_D$.*

Proof. [✓ Proof](#) The Stinespring Gram identity gives $V^\dagger V = \sum_i M_i^\dagger M_i = \sum_i E_i = \mathbb{1}_D$ by the resolution of identity. $\square$

Theorem 16.7.7 (Naimark projectors form a projective measurement). [Lean](#) [Lean](#) [Lean](#) [Lean](#)

[✓ Stmt](#) The Naimark projectors satisfy $P_i^2 = P_i$, $P_i^\dagger = P_i$, $P_i P_j = 0$ for $i \neq j$, and $\sum_i P_i = \mathbb{1}_{\mathbb{C}^D \otimes \mathbb{C}^n}$.

Proof. [✓ Proof](#) Each identity follows from direct entrywise computation using the delta-function form of $(P_i)_{(a,b),(c,d)}$. $\square$

Theorem 16.7.8 (Naimark dilation). [Lean](#) [✓ Stmt](#) Every POVM $\{E_i\}_{i=0}^{n-1}$ arises as a projective measurement on a dilation: for the isometry V and projectors P_i above,

$$E_i = V^\dagger P_i V.$$

Proof. [✓ Proof](#) Computing entrywise, $(V^\dagger P_i V)_{k,\ell} = \sum_a \overline{(M_i)_{a,k}} (M_i)_{a,\ell} = (M_i^\dagger M_i)_{k,\ell} = (E_i)_{k,\ell}$. $\square$

Theorem 16.7.9 (Existential Naimark dilation). [Lean](#) [✓ Stmt](#) For every POVM $\{E_i\}$ on $\mathbb{C}^D$ there exist a dilation dimension r , an isometry $V : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^r$, and a projective measurement $\{P_i\}$ on the dilation satisfying $E_i = V^\dagger P_i V$.

Proof. [✓ Proof](#) Take $r = n$ and the explicit witnesses $V = \sum_i M_i \otimes |i\rangle$ and $P_i = \mathbb{1}_D \otimes |i\rangle\langle i|$. $\square$

Definition 16.7.10 (Naimark dilation as a structure). [Lean](#) [✓ Stmt](#) A Naimark dilation of a POVM $\{E_i\}$ consists of an isometry V into a larger Hilbert space together with a projective measurement $\{P_i\}$ there such that $E_i = V^\dagger P_i V$ for all i .

Theorem 16.7.11 (Canonical dilation satisfies the Naimark dilation axioms). [Lean](#) [✓ Stmt](#) The explicit isometry $V = \sum_i M_i \otimes |i\rangle$ and projectors $P_i = \mathbb{1}_D \otimes |i\rangle\langle i|$ satisfy the defining axioms of a Naimark dilation.

Proof. [✓ Proof](#) Combine the previously proved identities $V^\dagger V = \mathbb{1}_D$, $P_i^2 = P_i$, $P_i^\dagger = P_i$, $P_i P_j = 0$ for $i \neq j$, $\sum_i P_i = \mathbb{1}$, and $V^\dagger P_i V = E_i$. $\square$

Theorem 16.7.12 (Concrete uniqueness for canonical projectors). [Lean](#) [✓ Stmt](#) Let $V : \mathbb{C}^D \rightarrow \mathbb{C}^D \otimes \mathbb{C}^n$ satisfy $V^\dagger P_i V = E_i$ for the canonical projectors $P_i = \mathbb{1}_D \otimes |i\rangle\langle i|$. Then there exists an isometry W on the dilated space such that $V = W V_0$, where V_0 is the canonical Naimark isometry of $\{E_i\}$.

Proof. [✓ Proof](#) For each outcome i , the i -th block of V has Gram matrix E_i . Comparing with the canonical square root M_i gives $V_i^\dagger V_i = M_i^\dagger M_i$, hence $V_i = U_i M_i$ for a unitary U_i on $\mathbb{C}^D$. Assemble the U_i block-diagonally to obtain an isometry $W = \bigoplus_i U_i$ on $\mathbb{C}^D \otimes \mathbb{C}^n$, and then $V = W \sum_i M_i \otimes |i\rangle = W V_0$. $\square$

Definition 16.7.13 (POVM from a PSD resolution of identity on a dilation). [Lean](#) [✓ Stmt](#) Given an isometry $V : \mathbb{C}^D \rightarrow \mathbb{C}^{d'}$ with $V^\dagger V = \mathbb{1}_D$ and a family $\{P_i\}_{i=0}^{n-1}$ of positive semidefinite operators on $\mathbb{C}^{d'}$ summing to the identity, the pulled-back operators

$$E_i := V^\dagger P_i V$$

form a POVM. In particular, any projective measurement on the dilation (a special case of a PSD resolution of identity) pulls back to a POVM.

Definition 16.7.14 (Quantum instrument). [Lean](#) [✓ Stmt](#) A *quantum instrument* with n outcomes is a family $\{\Phi_i\}_{i=0}^{n-1}$ of completely positive maps on $M_D(\mathbb{C})$ whose sum $\sum_i \Phi_i$ is trace-preserving.

Definition 16.7.15 (Instrument operations). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Associated to an instrument are the total channel $\sum_i \Phi_i$, the unnormalized update $\rho \mapsto \Phi_i(\rho)$ for each outcome i , the outcome probability $p_i(\rho) = \text{tr}(\Phi_i(\rho))$, and the normalized posterior state $\Phi_i(\rho)/p_i(\rho)$ whenever $p_i(\rho) \neq 0$.

Theorem 16.7.16 (Instrument total map is a channel). [Lean](#) [✓ Stmt](#) The total map $\sum_i \Phi_i$ of an instrument is a quantum channel.

Proof. [✓ Proof](#) Complete positivity of the sum follows from closure of CP maps under finite addition; trace preservation is built into the definition. $\square$

Theorem 16.7.17 (Conservation of probability for instruments). [Lean](#) [✓ Stmt](#) For every state $\rho \in M_D(\mathbb{C})$, the outcome probabilities $p_i(\rho) := \text{tr}(\Phi_i(\rho))$ sum to $\text{tr}(\rho)$.

Proof. [✓ Proof](#) Linearity of trace and trace preservation of $\sum_i \Phi_i$ give $\sum_i \text{tr}(\Phi_i(\rho)) = \text{tr}((\sum_i \Phi_i)(\rho)) = \text{tr}(\rho)$. $\square$

Theorem 16.7.18 (Non-negativity of instrument probabilities). [Lean](#) [✓ Stmt](#) If $\rho \in M_D(\mathbb{C})$ is positive semidefinite, then each outcome probability $p_i(\rho) = \text{tr}(\Phi_i(\rho))$ is a non-negative real number.

Proof. [✓ Proof](#) Each Φ_i is completely positive, hence positive, so $\Phi_i(\rho)$ is positive semidefinite and its trace is a non-negative real. $\square$

16.8 Trace-pairing expansion in transfer-matrix form

The trace-pairing expansion separates a linear map into the scalar input pairing, the transfer coefficients, and the output basis element:

$$T(\rho) = \sum_i \sum_j t_{ij} \text{tr}(\sigma_j \rho) \sigma_i.$$

The diagram records the same factorization by closing the input matrix with σ_j , weighting the resulting scalar by t_{ij} , and producing the output matrix through σ_i :

$$\begin{array}{c} \rho \\ \text{---} \bullet \sigma_j \\ \text{---} \text{tr} \end{array} \cdot \boxed{t_{ij}} \cdot \boxed{\sigma_i} = T(\rho)$$

Definition 16.8.1 (Trace-self-dual basis). [Lean](#) [✓ Stmt](#) A basis $\{\sigma_i\}_i$ of $M_D(\mathbb{C})$ is *trace-self-dual* when its coordinate functionals are given by trace pairing:

$$X = \sum_i \text{tr}(\sigma_i X) \sigma_i.$$

Definition 16.8.2 (Trace-pairing coefficients). [Lean](#) [✓ Stmt](#) For a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ and a trace-self-dual basis $\{\sigma_i\}$, define coefficients

$$t_{ij} := \text{tr}(\sigma_i T(\sigma_j)).$$

Theorem 16.8.3 (Trace-pairing expansion of a linear map). [Lean](#) [✓ Stmt](#) If $\{\sigma_i\}$ is trace-self-dual, then every linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ admits the expansion

$$T(\rho) = \sum_i \sum_j t_{ij} \operatorname{tr}(\sigma_j \rho) \sigma_i, \quad t_{ij} = \operatorname{tr}(\sigma_i T(\sigma_j)).$$

Proof. [✓ Proof](#) Expand ρ and each $T(\sigma_j)$ in the chosen basis and substitute. The trace-self-dual identity replaces coordinates by traces, giving the stated double-sum formula. $\square$

16.9 SVD normal form (existence)

[Wol12, Section 2.3] discusses two families of normal forms for the transfer-matrix representation of a quantum channel: the *singular value decomposition* (SVD) and, for qubit channels, the *Lorentz normal form* obtained by $\operatorname{SL}(2, \mathbb{C})$ filtering operations. The SVD normal form is proved below. The Lorentz normal form theorem (Theorem 16.10.13) requires the compactness and minimisation argument for $\operatorname{SL}(2, \mathbb{C})$ filterings in [Wol12, Propositions 2.9 and 2.11]; see Section 16.10.

Theorem 16.9.1 (SVD for positive semidefinite matrices). [Lean](#) [✓ Stmt](#) Every positive semidefinite matrix $M \in M_D(\mathbb{C})$ admits a decomposition

$$M = U \operatorname{diag}(\sigma) U^\dagger,$$

where $U \in \mathcal{U}(D)$ is unitary and $\sigma : \{0, \dots, D-1\} \rightarrow \mathbb{R}_{\geq 0}$ has non-negative entries.

Proof. [✓ Proof](#) The spectral theorem for Hermitian matrices provides an orthonormal eigenbasis with real eigenvalues. Positive semidefiniteness forces those eigenvalues to be non-negative, and the decomposition is written in SVD form with U the eigenvector unitary and σ the eigenvalue sequence. $\square$

Theorem 16.9.2 (SVD existence for invertible matrices). [Lean](#) [✓ Stmt](#) Every invertible complex square matrix $M \in M_D(\mathbb{C})$ admits a singular value decomposition

$$M = U \operatorname{diag}(\sigma) V^\dagger,$$

with $U, V \in \mathcal{U}(D)$ unitary and $\sigma_i > 0$ for all i .

Proof. [✓ Proof](#) Apply the spectral theorem to the positive definite matrix $M^\dagger M$ to obtain $M^\dagger M = V \operatorname{diag}(\lambda) V^\dagger$ with $\lambda_i > 0$. Set $\sigma_i := \sqrt{\lambda_i}$ and $\Sigma := \operatorname{diag}(\sigma)$; since Σ is a real positive diagonal, it is self-adjoint and invertible. Define $U := MV\Sigma^{-1}$. Then $U^\dagger U = \Sigma^{-1} V^\dagger (M^\dagger M) V \Sigma^{-1} = \Sigma^{-1} \operatorname{diag}(\lambda) \Sigma^{-1} = \mathbb{1}$, so U is unitary, and $U \Sigma V^\dagger = MV(\Sigma^{-1} \Sigma) V^\dagger = MVV^\dagger = M$. $\square$

Theorem 16.9.3 (SVD representation of a transfer matrix). [Lean](#) [✓ Stmt](#) Every invertible transfer matrix $\hat{T}$ of a linear super-operator $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ admits a singular value decomposition $\hat{T} = U \operatorname{diag}(\sigma) V^\dagger$ with U, V unitary on $\mathbb{C}^D \otimes \mathbb{C}^D$ and $\sigma_i > 0$.

Proof. [✓ Proof](#) Direct specialisation of Theorem 16.9.2 to the index type $\{0, \dots, D-1\} \times \{0, \dots, D-1\}$ used by the transfer-matrix representation. $\square$

Remark 16.9.4 (Sorted / unique singular values). Theorem 16.9.2 produces the singular values as an unordered family, without the usual convention that σ_i are sorted in non-increasing order or the statement that they are uniquely determined by M . Downstream applications in [Wol12, Section 2.3], such as trace-norm identities, polar decomposition, and the Lorentz normal form, will typically require the sorted / unique variant; that refinement is future work.

16.10 Lorentz normal form

This section states the existence theorems from [Wol12, Section 2.3].

Definition 16.10.1 (SL-filtering operation). An *SL-filtering* for $D \times D$ matrices is a completely positive map of the form $\Phi(X) = SXS^\dagger$ where $S \in M_D(\mathbb{C})$ satisfies $\det S = 1$. Such maps are invertible and have Kraus rank 1.

Definition 16.10.2 (Doubly-stochastic map). A linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *doubly-stochastic* if $T(\mathbb{1}) \propto \mathbb{1}$ and the reduced density matrix $\text{tr}_1[\tau]$ of its Choi matrix $\tau = (T \otimes \text{id})(|\Omega\rangle\langle\Omega|)$ is proportional to the identity. This is the normal form in [Wol12, Proposition 2.9].

Lemma 16.10.3 (Positive-definite trace lower bound). Lean Lean ✓ Stmt Let $D \geq 1$, let $M \in M_D(\mathbb{C})$ be positive-definite, and let $\lambda_{\min}(M)$ be its smallest Hermitian eigenvalue. For every $X \in M_D(\mathbb{C})$,

$$\lambda_{\min}(M) \text{tr}(X^\dagger X) \leq \text{tr}(XMX^\dagger).$$

Proof. ✓ Proof The matrix $M - \lambda_{\min}(M)\mathbb{1}$ is positive semidefinite by the spectral theorem. Since $X^\dagger X$ is also positive semidefinite, the trace product of these two matrices is non-negative. Expanding

$$\text{tr}(X^\dagger X(M - \lambda_{\min}(M)\mathbb{1}))$$

and cycling the trace gives the displayed inequality. □

Lemma 16.10.4 (Determinant AM–GM Hilbert–Schmidt bound). Lean ✓ Stmt Let $A \in M_D(\mathbb{C})$ with $D \geq 1$. If $|\det A| = 1$, then

$$D \leq \text{tr}(A^\dagger A).$$

Proof. ✓ Proof The matrix $A^\dagger A$ is positive semidefinite and has determinant $|\det A|^2 = 1$. Thus the product of its Hermitian eigenvalues is one. Denoting the eigenvalues of $A^\dagger A$ by $\lambda_1, \dots, \lambda_D \geq 0$, the arithmetic-geometric mean inequality gives

$$\frac{\lambda_1 + \dots + \lambda_D}{D} \geq (\lambda_1 \dots \lambda_D)^{1/D} = 1,$$

so $\text{tr}(A^\dagger A) = \lambda_1 + \dots + \lambda_D \geq D$. □

Lemma 16.10.5 (Kronecker Hilbert–Schmidt trace multiplicativity). Lean ✓ Stmt For complex matrices A and B of compatible rectangular sizes,

$$\text{tr}((A \otimes_k B)^\dagger (A \otimes_k B)) = \text{tr}(A^\dagger A) \text{tr}(B^\dagger B).$$

Proof. ✓ Proof Use $(A \otimes_k B)^\dagger = A^\dagger \otimes_k B^\dagger$, the mixed product identity for Kronecker products, and $\text{tr}(C \otimes_k D) = \text{tr}(C) \text{tr}(D)$. With $C = A^\dagger A$ and $D = B^\dagger B$, these identities give the equality. □

Lemma 16.10.6 (Infimum attainment for SL-filterings). Lean ✓ Stmt For a positive-definite Choi matrix τ , the infimum of $\text{tr}[(S_2 \otimes_k S_1)^\dagger \tau (S_2 \otimes_k S_1)]$ over $S_1, S_2 \in M_D(\mathbb{C})$ with $\det S_1 = \det S_2 = 1$ is attained.

Proof. ✓ Proof Write $X = S_2 \otimes_k S_1$. By the positive-definite trace lower bound (Lemma 16.10.3), $\text{tr}[X\tau X^\dagger] \geq \lambda_{\min}(\tau) \text{tr}[X^\dagger X]$, and by Hilbert–Schmidt multiplicativity (Lemma 16.10.5), $\text{tr}[X^\dagger X] = \text{tr}[S_2^\dagger S_2] \text{tr}[S_1^\dagger S_1]$. The determinant AM–GM estimate (Lemma 16.10.4) gives $\text{tr}[S_i^\dagger S_i] = \|S_i\|^2 \geq D$ whenever $\det S_i = 1$. Combining, the value dominates $\lambda_{\min}(\tau) D \|S_i\|^2$ for each factor, so any point whose value is at most the value at the identity lies in a fixed Frobenius ball $\{\|S\| \leq C\}$.

Intersecting with the closed set $\det S = 1$ produces a compact set, on which the continuous trace functional attains its minimum by the extreme-value theorem. A point outside the sublevel set has value larger than the value at the identity, so the minimiser on the compact set is a global minimiser. $\square$

Lemma 16.10.7 (Arithmetic–geometric-mean inequality, product/sum form). [Lean](#) [✓ Stmt](#) For a nonnegative family $f_0, \dots, f_{D-1}$ of real numbers,

$$D^D \prod_i f_i \leq \left(\sum_i f_i \right)^D.$$

Proof. [✓ Proof](#) Apply the weighted arithmetic–geometric-mean inequality with uniform weights $1/D$:

$$\left(\prod_i f_i \right)^{1/D} \leq \frac{1}{D} \sum_i f_i.$$

Raising both sides to the D -th power gives the stated bound. $\square$

Lemma 16.10.8 (Trace–determinant AM–GM for positive-semidefinite matrices). [Lean](#) [Lean](#) [✓ Stmt](#) For a positive-semidefinite $D \times D$ matrix M ,

$$D^D \det M \leq (\operatorname{tr} M)^D,$$

and equality holds if and only if $M = (\operatorname{tr} M/D) \mathbb{1}$.

Proof. [✓ Proof](#) Both $\det M = \prod_i \lambda_i$ and $\operatorname{tr} M = \sum_i \lambda_i$ are expressed through the nonnegative eigenvalues λ_i of M . The bound is the arithmetic–geometric-mean inequality with uniform weights $1/D$, raised to the D -th power; equality holds exactly when all eigenvalues coincide, that is, when M is a scalar matrix. $\square$

Theorem 16.10.9 (Generic normal form for CP maps). [Lean](#) [✓ Stmt](#) Let $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a completely positive map whose Choi matrix is positive-definite (equivalently, T has full Kraus rank). Then there exist SL -filterings Φ_1, Φ_2 such that $\Phi_2 \circ T \circ \Phi_1$ is doubly-stochastic.

This is the equal-dimension (square) case $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$; the source [\[Wol12, Proposition 2.9\]](#) states the proposition for a general $T : \mathcal{M}_{d_1} \rightarrow \mathcal{M}_{d_2}$ with separate filterings on each factor.

Proof. [✓ Proof](#) Let τ be the Choi matrix of T and take the minimiser (S_1, S_2) of $\operatorname{tr}[(S_2 \otimes_k S_1) \tau (S_2 \otimes_k S_1)^\dagger]$ over $\det S_1 = \det S_2 = 1$ from Lemma [16.10.6](#). Set Φ_1, Φ_2 to be the filterings with matrices $S_1^\dagger$ and S_2 , so that the Choi matrix of $\Phi_2 \circ T \circ \Phi_1$ equals $(S_2 \otimes_k S_1) \tau (S_2 \otimes_k S_1)^\dagger$. Holding one filtering factor fixed, the minimiser is optimal in the other coordinate, so each partial trace minimises a functional $\operatorname{tr}[SMS^\dagger]$ over $\det S = 1$ for a positive-semidefinite M , where

$$\operatorname{tr}[SMS^\dagger] \geq D(\det M)^{1/D},$$

with equality at the minimiser; by the equality case of Lemma [16.10.8](#) this forces $SMS^\dagger \propto \mathbb{1}$, hence both partial traces are proportional to the identity, which is exactly the doubly-stochastic condition. $\square$

Definition 16.10.10 (Diagonal Lorentz normal form). For a completely positive, trace-preserving qubit channel $T' : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$, write $\widehat{T}'$ for its matrix in the normalized Pauli basis $\{\sigma_0/\sqrt{2}, \sigma_1/\sqrt{2}, \sigma_2/\sqrt{2}, \sigma_3/\sqrt{2}\}$. The channel is in *diagonal Lorentz normal form* when it is unital and every off-diagonal entry of $\widehat{T}'$ is zero.

Definition 16.10.11 (Non-diagonal Lorentz normal form). For a completely positive, trace-preserving qubit channel $T' : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$, write $\widehat{T}'$ in the normalized Pauli basis. The channel is in *non-diagonal Lorentz normal form* when, for some $x \in [0, 1]$,

$$\widehat{T}' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & x/\sqrt{3} & 0 & 0 \\ 0 & 0 & x/\sqrt{3} & 0 \\ 2/3 & 0 & 0 & 1/3 \end{pmatrix}.$$

Trace preservation supplies the first row of the displayed matrix.

Definition 16.10.12 (Singular Lorentz normal form). For a completely positive, trace-preserving qubit channel $T' : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$, write $\widehat{T}'$ in the normalized Pauli basis. The channel is in *singular Lorentz normal form* when

$$\widehat{T}' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

equivalently, every input state is mapped to $(1 + \sigma_3)/2$. Trace preservation supplies the first row of the displayed matrix.

Theorem 16.10.13 (Lorentz normal form for qubit channels). *For every qubit channel $T : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$, there exist $\text{SL}(2, \mathbb{C})$ -filterings Φ_1, Φ_2 such that the filtered channel $T' = \Phi_2 \circ T \circ \Phi_1$ is in one of the three Lorentz normal forms: diagonal, non-diagonal, or singular. **Not yet proved**—proof still needs the generic normal form and the Lorentz group classification of $\text{SL}(2, \mathbb{C})$ orbits.*

16.11 Determinant of a quantum channel

Definition 16.11.1 (Channel determinant). [Lean](#) [✓ Stmt](#) For a linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the *channel determinant* $\det T$ is the determinant of the matrix of T with respect to the standard matrix-unit basis of $M_D(\mathbb{C})$. This is the quantity studied in [\[Wol12, Section 6.1\]](#).

Remark 16.11.2 (Channel determinant as an operator determinant). [Lean](#) [✓ Stmt](#) The channel determinant agrees with the ordinary determinant of the linear endomorphism $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$.

Definition 16.11.3 (Unitary channel). [Lean](#) [Lean](#) [✓ Stmt](#) For a unitary matrix $U \in \mathcal{U}(D)$, the *unitary channel* is $T(\rho) = U\rho U^\dagger$. It is automatically a quantum channel.

Theorem 16.11.4 (Unitary channels have determinant one). [Lean](#) [Lean](#) [✓ Stmt](#) *If $T(\rho) = U\rho U^\dagger$ is a unitary channel, then $\det T = 1$ and hence $|\det T| = 1$.*

Proof. [✓ Proof](#) Vectorization identifies T with a Kronecker product $\overline{U} \otimes U$, whose determinant is $\overline{\det U}^D (\det U)^D = 1$. □

Theorem 16.11.5 (Determinant bound for positive trace-preserving maps). [Lean](#) [Lean](#) [✓ Stmt](#) *For any positive trace-preserving map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, $|\det T| \leq 1$. This is [\[Wol12, Theorem 6.1\(1\)\]](#).*

Proof. ✓ **Proof** Every eigenvalue μ of T satisfies $|\mu| \leq 1$ (positivity and trace preservation force spectral radius ≤ 1). Since $\det T$ is the product of the eigenvalues (counted with algebraic multiplicity), the bound $|\det T| = \prod_i |\mu_i| \leq 1$ follows by induction. $\square$

Theorem 16.11.6 (Determinant one iff the channel is unitary). Lean ✓ **Stmt** For a CPTP map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$,

$$|\det T| = 1 \iff \exists U \in \mathcal{U}(D), \quad T(\rho) = U\rho U^\dagger.$$

This is [Wol12, Theorem 6.1(2)].

Proof. ✓ **Proof** Reverse direction is Theorem 16.11.4. Forward direction: $|\det T| = 1$ together with $|\mu| \leq 1$ for every eigenvalue μ (from trace preservation and positivity) forces every eigenvalue to satisfy $|\mu| = 1$. Hence T is bijective. A bijective CPTP map is a $*$ -automorphism: the Heisenberg dual $T^*(Y) = \sum_i K_i^\dagger Y K_i$ is unital (since T is TP), so the Kadison–Schwarz inequality $T^*(Y^\dagger Y) \geq T^*(Y)^\dagger T^*(Y)$ becomes an equality for every Y . Hence T^* , and therefore T , preserves products. By the Skolem–Noether theorem, every $*$ -automorphism of $M_D(\mathbb{C})$ has the form $T(\rho) = U\rho U^\dagger$ for some unitary U . Kraus freedom then gives $K_i = c_i U$ with $\sum_i |c_i|^2 = 1$. $\square$

Chapter 17

Quantum Dynamical Semigroups

This chapter develops the theory of one-parameter semigroups of quantum channels and characterizes their generators via the GKSL (Gorini–Kossakowski–Sudarshan–Lindblad) theorem, following [Wol12, Chapter 7].

Section 17.1 establishes definitions and the basic exponential form (Proposition 17.1.14). Section 17.2 gives the Duhamel formula and perturbation bound. Section 17.3 develops the generator theory and proves the GKSL theorem.

17.1 Dynamical semigroups and the exponential form

Definition 17.1.1 (Dynamical semigroup). [Lean](#) [✓ Stmt](#) A family of linear maps $T : \mathbb{R} \rightarrow (M_D(\mathbb{C}) \rightarrow_{\ell} M_D(\mathbb{C}))$ is a *dynamical semigroup* if

1. (semigroup law) $T_{t+s} = T_t \circ T_s$ for all $t, s \geq 0$; and
2. (initial condition) $T_0 = \text{id}$.

This is [Wol12, Equation (7.1)].

Definition 17.1.2 (Continuous dynamical semigroup). [Lean](#) [✓ Stmt](#) A dynamical semigroup T is *norm-continuous* if the map $t \mapsto T_t$ is continuous in the operator norm topology. In finite dimension this coincides with strong continuity.

Definition 17.1.3 (Exponential semigroup). [Lean](#) [Lean](#) [✓ Stmt](#) Given a generator $L \in \text{End}_{\mathbb{C}}(M_D(\mathbb{C}))$, the *exponential semigroup* is

$$T_t := e^{tL} = \sum_{k=0}^{\infty} \frac{(tL)^k}{k!}.$$

Theorem 17.1.4 (Semigroup law for the exponential). [Lean](#) [Lean](#) [✓ Stmt](#) For all $t, s \in \mathbb{R}$, $e^{(t+s)L} = e^{tL} \cdot e^{sL}$.

Proof. [✓ Proof](#) Since $(t \cdot L)$ commutes with $(s \cdot L)$, this follows from the exponential addition formula for commuting elements in a Banach algebra. $\square$

Theorem 17.1.5 (Exponential semigroup at time 0). [Lean](#) [✓ Stmt](#) The continuous-linear-map exponential satisfies $e^{0L} = \mathbb{1}$.

Proof. [✓ Proof](#) The exponential of the zero endomorphism is the identity. $\square$

Theorem 17.1.6 (Continuity of the exponential semigroup). [Lean](#) [Lean](#) [✓ Stmt](#) *The map $t \mapsto e^{tL}$ is continuous in the operator norm topology.*

Proof. [✓ Proof](#) The exponential is analytic (convergence radius = ∞), hence continuous. $\square$

Theorem 17.1.7 (Derivative of the exponential semigroup). [Lean](#) [✓ Stmt](#) *For all $t \in \mathbb{R}$,*

$$\frac{d}{dt}e^{tL} = e^{tL} \cdot L.$$

This is [Wol12, Equation (7.2)].

Proof. [✓ Proof](#) Chain rule applied to the composition $\mathbb{R} \xrightarrow{t \mapsto t \cdot L} \text{End}(M_D(\mathbb{C})) \xrightarrow{\exp} \text{End}(M_D(\mathbb{C}))$. $\square$

Theorem 17.1.8 (Derivative at the origin). [Lean](#) [✓ Stmt](#) *One has*

$$\left. \frac{d}{dt}e^{tL} \right|_{t=0} = L.$$

Proof. [✓ Proof](#) This is Theorem 17.1.7 at $t = 0$ together with Theorem 17.1.5. $\square$

Theorem 17.1.9 (Generator uniqueness). [Lean](#) [✓ Stmt](#) *If $e^{tL} = e^{tL'}$ for all $t \geq 0$, then $L = L'$.*

Proof. [✓ Proof](#) Both semigroups agree on $[0, \infty)$, so their derivatives within $[0, \infty)$ at $t = 0$ agree. Since $[0, \infty)$ has unique differentials at 0, $L = L'$. $\square$

Theorem 17.1.10 (CLM and linear-map forms coincide). [Lean](#) [✓ Stmt](#) *The linear-map exponential semigroup and the continuous-linear-map exponential semigroup agree under the canonical identification of endomorphisms with continuous linear endomorphisms.*

Proof. [✓ Proof](#) This is a direct unpacking of the definition of the linear-map exponential semigroup. $\square$

Theorem 17.1.11 (Pointwise derivative formula). [Lean](#) [✓ Stmt](#) *For every $X \in M_D(\mathbb{C})$ and every $t \in \mathbb{R}$,*

$$\frac{d}{dt}(e^{tL}(X)) = e^{tL}(L(X)).$$

Proof. [✓ Proof](#) Evaluate the derivative of the continuous-linear-map semigroup at X and transport the statement through Theorem 17.1.10. $\square$

Theorem 17.1.12 (Linear-map exponential semigroup at time 0). [Lean](#) [✓ Stmt](#) *The linear-map exponential semigroup satisfies $e^{0L} = \mathbb{1}$.*

Proof. [✓ Proof](#) Transport Theorem 17.1.5 through Theorem 17.1.10. $\square$

Theorem 17.1.13 (Exponential semigroup as a dynamical semigroup). [Lean](#) [✓ Stmt](#) *The family $t \mapsto e^{tL}$ satisfies the semigroup law and the initial condition, hence is a dynamical semigroup.*

Proof. [✓ Proof](#) Transport the continuous-linear-map semigroup law from Theorem 17.1.4 through Theorem 17.1.10; the initial condition is Theorem 17.1.12. $\square$

Theorem 17.1.14 (Every continuous semigroup is exponential). [Lean](#) [✓ Stmt](#) Every norm-continuous dynamical semigroup T on $M_D(\mathbb{C})$ is of the form

$$T_t = e^{tL}$$

for some generator $L \in \text{End}_{\mathbb{C}}(M_D(\mathbb{C}))$ and every $t \geq 0$. This is [Wol12, Proposition 7.1].

Proof. [✓ Proof](#) Since $T_0 = \mathbb{1}$ and T is continuous, the integral $M_\varepsilon = \int_0^\varepsilon T_s ds$ is invertible for small $\varepsilon > 0$. Using the semigroup property, $T_t = M_\varepsilon^{-1}(M_{t+\varepsilon} - M_t)$ is differentiable for $t \geq 0$. ODE uniqueness then gives $T_t = e^{tL}$ for all $t \geq 0$. $\square$

17.2 Perturbation theory

Theorem 17.2.1 (Duhamel formula). [Lean](#) [✓ Stmt](#) Let $\Delta = L' - L$. For $t \geq 0$,

$$e^{tL'} - e^{tL} = \int_0^t e^{(t-s)L} \Delta e^{sL'} ds.$$

This is [Wol12, Lemma 7.1].

Proof. [✓ Proof](#) Define $f(s) = e^{(t-s)L} e^{sL'}$. Then $f'(s) = e^{(t-s)L} \Delta e^{sL'}$ (Theorem 17.1.7), and $e^{tL'} - e^{tL} = f(t) - f(0) = \int_0^t f'(s) ds$. $\square$

Theorem 17.2.2 (Perturbation bound). [Lean](#) [✓ Stmt](#) For $t \geq 0$,

$$\|e^{tL'} - e^{tL}\| \leq t \|\Delta\| \cdot \sup_{s \in [0, t]} \|e^{sL}\| \cdot \sup_{s \in [0, t]} \|e^{sL'}\|,$$

where $\Delta = L' - L$. This is [Wol12, Corollary 7.1].

Proof. [✓ Proof](#) Apply the triangle inequality and submultiplicativity of the operator norm to the Duhamel integral (Theorem 17.2.1). $\square$

Corollary 17.2.3 (Perturbation bound under unit-norm control). [Lean](#) [✓ Stmt](#) For $t \geq 0$, if $\|e^{sL}\| \leq 1$ and $\|e^{sL'}\| \leq 1$ for every $s \in [0, t]$, then

$$\|e^{tL'} - e^{tL}\| \leq t \|L' - L\|.$$

Proof. [✓ Proof](#) Apply Theorem 17.2.2 and bound both suprema by 1. $\square$

Lemma 17.2.4 (Dyson-series summability from factorial majorant). [Lean](#) [✓ Stmt](#) Assume a factorial majorant of Dyson–Phillips iterates:

$$\|D_n(t)\| \leq M \frac{(t \|L' - L\| M)^n}{n!}.$$

Then the Dyson series is summable in operator norm.

Proof. [✓ Proof](#) This is the Weierstrass M-test step, using the scalar convergence of $\sum_n a^n/n!$ and lifting via norm-bounded summability. $\square$

Theorem 17.2.5 (Factorial norm bound for Dyson iterates). [Lean](#) [✓ Stmt](#) For $s \in [0, t]$ and $M = \sup_{u \in [0, t]} \|e^{uL}\|$,

$$\|\tilde{T}_s^{(n)}\| \leq M \frac{(s \|\Delta\| M)^n}{n!}.$$

Proof. ✓ Proof Induction on n . The base case uses the semigroup supremum bound. The inductive step bounds the integrand pointwise by $M\|\Delta\| \cdot M(u\|\Delta\|M)^n/n!$ and integrates $\int_0^s u^n du = s^{n+1}/(n+1)$, recovering the factorial. $\square$

Corollary 17.2.6 (Dyson series summability). Lean ✓ Stmt *The Dyson–Phillips series $\sum_n \tilde{T}_t^{(n)}$ converges in operator norm for every $t \geq 0$.*

Proof. ✓ Proof Compose the factorial norm bound (Theorem 17.2.5) with the Weierstrass M-test step (Lemma 17.2.4). $\square$

Corollary 17.2.7 (Factorial norm bound at the final time). Lean ✓ Stmt *For $t \geq 0$, setting $s = t$ in Theorem 17.2.5 gives*

$$\|\tilde{T}_t^{(n)}\| \leq M \frac{(t\|\Delta\|M)^n}{n!},$$

where $M = \sup_{u \in [0, t]} \|e^{uL}\|$.

Proof. ✓ Proof This is the special case $s = t$ of Theorem 17.2.5. $\square$

Theorem 17.2.8 (Continuity of Dyson iterates). Lean ✓ Stmt *Each Dyson iterate $t \mapsto \tilde{T}_t^{(n)}$ is continuous in the time parameter.*

Proof. ✓ Proof By induction on n . The base case is continuity of the matrix exponential. For the inductive step, the parametric integral $t \mapsto \int_0^t e^{(t-s)L} \Delta \tilde{T}_s^{(n)} ds$ is continuous by the continuous-parameter primitive theorem; the pasting lemma extends this to all of $\mathbb{R}$ since $\tilde{T}_t^{(n+1)} = 0$ for $t \leq 0$. $\square$

Theorem 17.2.9 (Factorial bound on the Dyson remainder). Lean ✓ Stmt *For $s \in [0, t]$, $M = \sup_{u \in [0, t]} \|e^{uL}\|$, and $M' = \sup_{u \in [0, t]} \|e^{uL'}\|$:*

$$\|T'_s - \sum_{n < N} \tilde{T}_s^{(n)}\| \leq M' \frac{(s\|\Delta\|M)^N}{N!}.$$

Proof. ✓ Proof Induction on N . The base case is the semigroup supremum bound. The inductive step uses the telescoping remainder identity $R_{N+1}(s) = \int_0^s e^{(s-u)L} \Delta R_N(u) du$ (derived from the Duhamel formula), bounds the integrand pointwise, and integrates to recover the factorial. $\square$

Theorem 17.2.10 (Dyson series identity [Wol12, Equation (7.13)]). Lean ✓ Stmt *For $t \geq 0$, the Dyson–Phillips series equals the perturbed semigroup:*

$$\sum_{n=0}^{\infty} \tilde{T}_t^{(n)} = e^{tL'}.$$

Proof. ✓ Proof The partial-sum remainder $\|e^{tL'} - \sum_{n < N} \tilde{T}_t^{(n)}\| \leq M' (t\|\Delta\|M)^N/N!$ tends to zero as $N \rightarrow \infty$, since $c^N/N! \rightarrow 0$ for any constant c . Together with the summability of the series (Corollary 17.2.6), this identifies the sum via uniqueness of limits. $\square$

Corollary 17.2.11 (Tsum form of the Dyson series). Lean ✓ Stmt $\sum_{n=0}^{\infty} \tilde{T}_t^{(n)} = e^{tL'}$ (tsum form).

Proof. ✓ Proof The claim follows from Theorem 17.2.10. $\square$

17.3 GKSL/Lindblad generators

This section characterizes generators of semigroups of completely positive and trace-preserving maps. The central result is the GKSL theorem (Theorem 17.3.8.5), which shows that such generators have the standard Lindblad form.

17.3.1 Generator decomposition and conditional complete positivity

Definition 17.3.1.1 (Generator decomposition). [Lean](#) [✓ Stmt](#) A *generator decomposition* consists of a completely positive map $\phi : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ and a matrix $\kappa \in M_d(\mathbb{C})$, defining the linear map

$$L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger.$$

This is [Wol12, Equation (7.14)].

Definition 17.3.1.2 (Conditional complete positivity). [Lean](#) [✓ Stmt](#) A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is *conditionally completely positive* (CCP) if it admits a generator decomposition, i.e. there exist a CP map ϕ and a matrix κ such that $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$. This is condition 1 of [Wol12, Proposition 7.2].

Definition 17.3.1.3 (Trace-annihilating map). [Lean](#) [✓ Stmt](#) A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is *trace-annihilating* if $\text{tr}(L(\rho)) = 0$ for all $\rho \in M_d(\mathbb{C})$. This is the infinitesimal version of trace preservation.

Definition 17.3.1.4 (Trace constraint). [Lean](#) [✓ Stmt](#) A generator decomposition (ϕ, κ) satisfies the *trace constraint* if $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, where ϕ^* is the Hilbert–Schmidt adjoint. This is the condition in [Wol12, Equation (7.20)].

Theorem 17.3.1.5 (Trace constraint implies trace-annihilating). [Lean](#) [✓ Stmt](#) If (ϕ, κ) satisfies $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, then $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$ is trace-annihilating.

Proof. [✓ Proof](#) Let $\phi(\rho) = \sum_i K_i \rho K_i^\dagger$. By the cyclic property of trace, $\text{tr}(L(\rho)) = \text{tr}((\sum_i K_i^\dagger K_i)\rho) - \text{tr}(\kappa\rho) - \text{tr}(\kappa^\dagger\rho) = \text{tr}((\kappa + \kappa^\dagger)\rho) - \text{tr}(\kappa\rho) - \text{tr}(\kappa^\dagger\rho) = 0$. $\square$

17.3.2 The Lindblad form

Definition 17.3.2.1 (Lindblad form). [Lean](#) [✓ Stmt](#) A *Lindblad form* consists of a Hermitian matrix $H = H^\dagger$ (the Hamiltonian) and a family of matrices $\{L_j\}_{j=0}^{r-1}$ (the Lindblad operators), defining the linear map

$$L(\rho) = i[\rho, H] + \sum_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\}_+ \right),$$

where $[A, B] = AB - BA$ and $\{A, B\}_+ = AB + BA$. This is [Wol12, Equation (7.21)].

Theorem 17.3.2.2 (Lindblad form is trace-annihilating). [Lean](#) [✓ Stmt](#) Every Lindblad form defines a trace-annihilating linear map.

Proof. [✓ Proof](#) The Hamiltonian part satisfies $\text{tr}(i[\rho, H]) = 0$ by the cyclic property. Each dissipator term satisfies $\text{tr}(L_j \rho L_j^\dagger) = \text{tr}(L_j^\dagger L_j \rho) = \text{tr}(\rho L_j^\dagger L_j)$, so the three terms sum to zero. $\square$

Theorem 17.3.2.3 (Lindblad form equals generator decomposition). [Lean](#) [✓ Stmt](#) A Lindblad form with Hamiltonian H and operators $\{L_j\}$ defines the same linear map as the generator decomposition (ϕ, κ) with $\phi(\rho) = \sum_j L_j \rho L_j^\dagger$ and $\kappa = iH + \frac{1}{2} \sum_j L_j^\dagger L_j$. This is [Wol12, Equation (7.24)].

Proof. [✓ Proof](#) Compute $\kappa^\dagger = -iH + \frac{1}{2} \sum_j L_j^\dagger L_j$ using $H^\dagger = H$ and $(A^\dagger A)^\dagger = A^\dagger A$. Setting $S = \sum_j L_j^\dagger L_j$, expand $\phi(\rho) - \kappa\rho - \rho\kappa^\dagger = \sum_j L_j \rho L_j^\dagger - iH\rho - \frac{1}{2}S\rho + i\rho H - \frac{1}{2}\rho S$, which is $i[\rho, H] + \sum_j (L_j \rho L_j^\dagger - \frac{1}{2}L_j^\dagger L_j \rho - \frac{1}{2}\rho L_j^\dagger L_j)$. $\square$

Theorem 17.3.2.4 (Lindblad form is CCP). [Lean](#) [✓ Stmt](#) Every Lindblad form defines a conditionally completely positive map.

Proof. [✓ Proof](#) By Theorem 17.3.2.3, the Lindblad form equals a generator decomposition with ϕ CP. $\square$

Definition 17.3.2.5 (Commutator form of Lindblad equation). [Lean](#) [✓ Stmt](#) The commutator form of the Lindblad equation writes the generator as

$$L(\rho) = i[\rho, H] + \frac{1}{2} \sum_j ([L_j, \rho L_j^\dagger] + [L_j \rho, L_j^\dagger]),$$

where $[A, B] = AB - BA$. This is [Wol12, Equation (7.22)].

Theorem 17.3.2.6 (Commutator form equals standard Lindblad form). [Lean](#) [✓ Stmt](#) The commutator form (Definition 17.3.2.5) defines the same linear map as the standard Lindblad form (Definition 17.3.2.1).

Proof. [✓ Proof](#) Expanding the commutator brackets $[L_j, \rho L_j^\dagger]$ and $[L_j \rho, L_j^\dagger]$ yields the standard dissipator terms. $\square$

17.3.3 Characterization of conditional complete positivity

Theorem 17.3.3.1 (Conditional complete positivity implies projected Choi positivity). [Lean](#) [✓ Stmt](#) If L is CCP and $P = \mathbb{1} - |\Omega\rangle\langle\Omega|$, then

$$P((L \otimes \text{id})(|\Omega\rangle\langle\Omega|))P \geq 0.$$

This is [Wol12, Proposition 7.2].

Proof. [✓ Proof](#) Write $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$. The left- and right-multiplication terms are annihilated by P , so the projected Choi matrix reduces to the projected Choi matrix of ϕ , which is positive. $\square$

Theorem 17.3.3.2 (Projected Choi positivity implies conditional complete positivity). [Lean](#) [✓ Stmt](#) If L is Hermiticity-preserving and $P((L \otimes \text{id})(|\Omega\rangle\langle\Omega|))P \geq 0$ where $P = \mathbb{1} - |\Omega\rangle\langle\Omega|$, then L is CCP. This is [Wol12, Proposition 7.2].

Proof. [✓ Proof](#) Decompose the Hermitian Choi matrix as $\tau_L = Q - |\psi\rangle\langle\Omega| - |\Omega\rangle\langle\psi|$ with $Q \geq 0$ supported on $|\Omega\rangle^\perp$. The Choi–Jamiołkowski isomorphism applied to Q gives the CP map ϕ , and $(\kappa \otimes \mathbb{1})|\Omega\rangle = |\psi\rangle$ defines κ . $\square$

17.3.4 Completely positive semigroups and conditionally completely positive generators

Theorem 17.3.4.1 (CP semigroups have CCP generators). [Lean](#) [✓ Stmt](#) If e^{tL} is CP for all $t \geq 0$, then L is CCP. This is [Wol12, Proposition 7.3].

Proof. [✓ Proof](#) From $(e^{tL} \otimes \text{id})(|\Omega\rangle\langle\Omega|) \geq 0$, expand at infinitesimal t and project onto P to get $P(L \otimes \text{id})(|\Omega\rangle\langle\Omega|)P \geq 0$. Then Theorem 17.3.3.2 gives CCP. $\square$

Theorem 17.3.4.2 (CCP generators yield CP semigroups). [Lean](#) [✓ Stmt](#) If L is CCP, then e^{tL} is CP for all $t \geq 0$. This is [Wol12, Proposition 7.3].

Proof. [✓ Proof](#) Approximate e^{tL} by the completely positive Euler steps

$$\rho \mapsto (\mathbb{1} - h\kappa)\rho(\mathbb{1} - h\kappa)^\dagger + h\phi(\rho), \quad h = \frac{t}{n+1}.$$

Each step is CP, hence so are its powers. These powers converge in operator norm to e^{tL} , and the cone of CP maps is closed. $\square$

Theorem 17.3.4.3 (CP semigroups iff CCP generators). [Lean](#) [✓ Stmt](#) e^{tL} is CP for all $t \geq 0$ if and only if L is CCP.

Proof. [✓ Proof](#) Combine Theorems 17.3.4.1 and 17.3.4.2. $\square$

17.3.5 Freedom in generator representation

Theorem 17.3.5.1 (Traceless Kraus operators exist). [Lean](#) [✓ Stmt](#) Assume $d > 0$. Given any Kraus operators $\{K_j\}$, there exist $c_j \in \mathbb{C}$ such that $K'_j = K_j + c_j\mathbb{1}$ is traceless. This is part of [Wol12, Proposition 7.4].

Proof. [✓ Proof](#) Set $c_j = -\text{tr}(K_j)/d$. $\square$

Theorem 17.3.5.2 (Shift invariance of generators). [Lean](#) [✓ Stmt](#) If $K'_j = K_j + c_j\mathbb{1}$ and κ' is adjusted per [Wol12, Equation (7.19)], then (ϕ', κ') and (ϕ, κ) define the same generator.

Proof. [✓ Proof](#) Direct algebraic expansion. $\square$

17.3.6 Uniqueness of the traceless Lindblad form

Definition 17.3.6.1 (Traceless Kraus operators). [Lean](#) [✓ Stmt](#) A Lindblad form has *traceless Kraus operators* if $\text{tr}(L_j) = 0$ for every j .

Theorem 17.3.6.2 (Uniqueness of the completely positive part). [Lean](#) [✓ Stmt](#) If two Lindblad forms F, F' induce the same generator and both have traceless Kraus operators, then their CP parts agree: $\phi = \phi'$. This is part of [Wol12, Proposition 7.4(2)].

Proof. [✓ Proof](#) Compare projected Choi matrices and use Choi injectivity. $\square$

Theorem 17.3.6.3 (Uniqueness of the drift term modulo phase). [Lean](#) [✓ Stmt](#) If two Lindblad forms F, F' induce the same generator and both have traceless Kraus operators, then their drift matrices differ by an imaginary scalar: $\kappa' = \kappa + i\lambda\mathbb{1}$ for some $\lambda \in \mathbb{R}$. This is part of [Wol12, Proposition 7.4(2)].

Proof. [✓ Proof](#) After identifying $\phi = \phi'$ (Theorem 17.3.6.2), the generator equality forces $\Delta\kappa \cdot \rho + \rho \cdot (\Delta\kappa)^\dagger = 0$ for all ρ . Setting $\rho = \mathbb{1}$ gives $\Delta\kappa + (\Delta\kappa)^\dagger = 0$ (skew-Hermiticity), which converts the equation to $[\Delta\kappa, \rho] = 0$ for all ρ . The scalar commutant lemma gives $\Delta\kappa = c \cdot \mathbb{1}$, and skew-Hermiticity forces $c = i\lambda$ for some $\lambda \in \mathbb{R}$. $\square$

Theorem 17.3.6.4 (Combined uniqueness of the traceless Lindblad form). [Lean](#) [✓ Stmt](#) *If two Lindblad forms F, F' induce the same generator and both have traceless Kraus operators, then $\phi = \phi'$ and $\kappa' = \kappa + i\lambda \mathbb{1}$ for some $\lambda \in \mathbb{R}$.*

Proof. [✓ Proof](#) Combine Theorems 17.3.6.2 and 17.3.6.3. $\square$

17.3.7 Trace-annihilation and trace preservation

Theorem 17.3.7.1 (Trace-annihilating generators yield trace-preserving semigroups). [Lean](#) [✓ Stmt](#) *If L is trace-annihilating, then e^{tL} is trace-preserving for every $t \in \mathbb{R}$.*

Proof. [✓ Proof](#) For fixed ρ , the function $f(t) = \text{tr}(e^{tL}(\rho))$ has derivative $f'(t) = \text{tr}(L(e^{tL}(\rho))) = 0$, so f is constant and $\text{tr}(e^{tL}(\rho)) = f(t) = f(0) = \text{tr}(\rho)$. $\square$

Theorem 17.3.7.2 (Trace-preserving semigroups have trace-annihilating generators). [Lean](#) [✓ Stmt](#) *If e^{tL} is trace-preserving for every $t \geq 0$, then L is trace-annihilating.*

Proof. [✓ Proof](#) Differentiate the identity $\text{tr}(e^{tL}(\rho)) = \text{tr}(\rho)$ at $t = 0$ from the right. $\square$

17.3.8 The GKSL/Lindblad theorem

Definition 17.3.8.1 (GKSL generator). [Lean](#) [✓ Stmt](#) A linear map $L : M_d(\mathbb{C}) \rightarrow M_d(\mathbb{C})$ is a *GKSL generator* if e^{tL} is a quantum channel (CPTP) for all $t \geq 0$.

Theorem 17.3.8.2 (A trace-constrained generator decomposition gives a GKSL generator). [Lean](#) [✓ Stmt](#) *If $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$ with ϕ CP and $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$, then L is a GKSL generator. This is [Wol12, Theorem 7.1, Equation (7.20)].*

Proof. [✓ Proof](#) CP follows from Theorem 17.3.4.2. The trace constraint implies trace-annihilation by Theorem 17.3.1.5, so Theorem 17.3.7.1 gives trace preservation. $\square$

Theorem 17.3.8.3 (GKSL generators admit a trace-constrained generator decomposition). [Lean](#) [✓ Stmt](#) *If L is a GKSL generator, then there exist a CP map ϕ and a matrix κ such that*

$$L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$$

and $\phi^(\mathbb{1}) = \kappa + \kappa^\dagger$.*

Proof. [✓ Proof](#) Apply Theorem 17.3.4.1 to obtain a CCP decomposition. Then apply Theorem 17.3.7.2 to the trace-preserving part of the semigroup and rewrite the resulting trace-annihilation condition as $\phi^*(\mathbb{1}) = \kappa + \kappa^\dagger$. $\square$

Theorem 17.3.8.4 (GKSL iff conditional complete positivity and trace annihilation). [Lean](#) [✓ Stmt](#) *L is a GKSL generator if and only if L is CCP and trace-annihilating.*

Proof. [✓ Proof](#) Forward: apply Theorem 17.3.4.1 to the CP part and Theorem 17.3.7.2 to the TP part. Reverse: combine Theorem 17.3.4.2 with Theorem 17.3.7.1. $\square$

Theorem 17.3.8.5 (GKSL iff Lindblad form). [Lean](#) [✓ Stmt](#) L is a GKSL generator if and only if

$$L(\rho) = i[\rho, H] + \sum_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\}_+ \right)$$

with $H = H^\dagger$. This is [Wol12, Theorem 7.1].

Proof. [✓ Proof](#) Forward: apply Theorem 17.3.8.3 to write $L(\rho) = \phi(\rho) - \kappa\rho - \rho\kappa^\dagger$ with trace constraint. Then choose traceless Kraus operators by Theorem 17.3.5.1 and rewrite κ as $iH + \frac{1}{2}\phi^*(\mathbb{1})$; Theorem 17.3.2.3 gives the Lindblad formula. Reverse: Theorems 17.3.2.4 and 17.3.2.2 show that every Lindblad form satisfies the right-hand side of Theorem 17.3.8.4. $\square$

17.3.9 Kossakowski matrix form

Definition 17.3.9.1 (Kossakowski form). [Lean](#) [✓ Stmt](#) A Kossakowski form consists of $H = H^\dagger$, a finite family of matrices $\{F_k\}_{k=0}^{n-1}$, and a positive semidefinite matrix $C \in M_n(\mathbb{C})$, defining

$$L(\rho) = i[\rho, H] + \frac{1}{2} \sum_{k,l} C_{lk} ([F_k, \rho F_l^\dagger] + [F_k \rho, F_l^\dagger]).$$

In Wolf's formula one takes $\{F_k\}$ to be a basis of traceless matrices; here we keep only the algebraic data needed for the conversion to Lindblad form.

Theorem 17.3.9.2 (Kossakowski form iff Lindblad form). [Lean](#) [✓ Stmt](#) A linear map admits a Kossakowski form iff it admits a Lindblad form.

Proof. [✓ Proof](#) Forward: diagonalize $C = B^\dagger B$ and set $L_j = \sum_k B_{jk} F_k$. Reverse: keep the same Hamiltonian and family $F_k = L_k$, and take $C = \mathbb{1}$. $\square$

17.4 Primitivity and irreducibility of QDS

17.4.1 Auxiliary spectral and semigroup lemmas

Definition 17.4.1.1 (Quantum dynamical semigroup). [Lean](#) [✓ Stmt](#) A quantum dynamical semigroup is a norm-continuous dynamical semigroup whose time slices are quantum channels for every $t \geq 0$.

Theorem 17.4.1.2 (Semigroup power identity). [Lean](#) [✓ Stmt](#) For $t \geq 0$ and $n \in \mathbb{N}$, one has $T_{nt} = (T_t)^n$.

Lemma 17.4.1.3 (Density matrices are nonzero). [Lean](#) [✓ Stmt](#) Every density matrix is nonzero.

Theorem 17.4.1.4 (Compression invariance is stable under powers). [Lean](#) [✓ Stmt](#) If a corner compression is invariant under E , it remains invariant under every power E^n .

Theorem 17.4.1.5 (Generator eigenvectors along the exponential semigroup). [Lean](#) [✓ Stmt](#) If $L(X) = \mu X$, then $e^{tL}(X) = e^{t\mu} X$ for all $t \in \mathbb{R}$.

Theorem 17.4.1.6 (Spectral mapping for semigroup generators). [Lean](#) [✓ Stmt](#) If μ is in the spectrum of L , then $e^{t\mu}$ is in the spectrum of e^{tL} .

Theorem 17.4.1.7 (Root-of-unity rigidity for phases). [Lean](#) [✓ Stmt](#) If $e^{it\theta}$ is a root of unity for every $t > 0$, then $\theta = 0$.

Theorem 17.4.1.8 (Peripheral generator eigenvalues are purely imaginary). [Lean](#) [✓ Stmt](#) *Peripheral spectral points of the generator have vanishing real part.*

Theorem 17.4.1.9 (Peripheral cardinality bound). [Lean](#) [✓ Stmt](#) *The number of peripheral eigenvalues is bounded by $\dim(M_D(\mathbb{C}))$.*

Theorem 17.4.1.10 (Bounded order under peripheral power-closure). [Lean](#) [✓ Stmt](#) *If all powers of a peripheral eigenvalue remain peripheral, then some positive power not exceeding $\dim(M_D(\mathbb{C}))$ equals 1.*

Theorem 17.4.1.11 (Peripheral powers stay peripheral under irreducibility). [Lean](#) [✓ Stmt](#) *For an irreducible channel with faithful fixed point, every power of a peripheral eigenvalue is again peripheral.*

Theorem 17.4.1.12 (Spectral-radius criterion from eigenvalue bounds). [Lean](#) [✓ Stmt](#) *If all eigenvalues satisfy $|\lambda| < 1$, then the spectral radius is < 1 .*

Theorem 17.4.1.13 (Primitive powers annihilate trace-zero matrices). [Lean](#) [✓ Stmt](#) *If E is an irreducible primitive channel with faithful fixed density σ , then $E^n(X) \rightarrow 0$ for every matrix X with $\text{tr}(X) = 0$.*

Proof. [✓ Proof](#) Decompose E^n into the fixed-point projection and its complementary part. On the complement, every eigenvalue has modulus < 1 , so the complementary powers tend to 0; the trace-zero hypothesis removes the fixed-point component. $\square$

Theorem 17.4.1.14 (Fixed points of an irreducible channel are trace multiples). [Lean](#) [✓ Stmt](#) *If E is an irreducible channel with faithful fixed density σ , then every nonzero fixed point V of E satisfies $V = \text{tr}(V)\sigma$.*

Proof. [✓ Proof](#) Subtract the trace multiple $\text{tr}(V)\sigma$ and apply the vanishing theorem for trace-zero fixed points of an irreducible channel. $\square$

Corollary 17.4.1.15 (Nonzero fixed points have nonzero trace). [Lean](#) [✓ Stmt](#) *Every nonzero fixed point of an irreducible channel has nonzero trace.*

Proof. [✓ Proof](#) Otherwise the preceding theorem would force the fixed point to vanish. $\square$

Theorem 17.4.1.16 (Peripheral eigenvectors become periodic fixed points). [Lean](#) [✓ Stmt](#) *For every positive time t , a quantum dynamical semigroup irreducible at every positive time has the following property: every peripheral eigenvalue μ of T_t admits a nonzero eigenvector V and a positive integer p such that $T_t(V) = \mu V$ and $T_{pt}(V) = V$.*

Proof. [✓ Proof](#) Peripheral powers remain peripheral for an irreducible channel with faithful fixed point. The bounded-order lemma then gives $\mu^p = 1$, and the semigroup power identity turns the eigenvector equation for T_t into a fixed-point equation for T_{pt} . $\square$

Theorem 17.4.1.17 (Peripheral eigenvectors with nonzero trace). [Lean](#) [✓ Stmt](#) *For every positive time t , under the same hypotheses every peripheral eigenvalue of T_t has a nonzero eigenvector with nonzero trace.*

Proof. [✓ Proof](#) Apply Theorem 17.4.1.16 and then use the previous corollary at the irreducible time pt . $\square$

Theorem 17.4.1.18 (Trace-preserving eigenvectors force eigenvalue 1). [Lean](#) [✓ Stmt](#) *If E is trace-preserving, $E(V) = \mu V$, and $\text{tr}(V) \neq 0$, then $\mu = 1$.*

Proof. ✓ Proof Take traces in $E(V) = \mu V$ and use trace preservation. □

Theorem 17.4.1.19 (Peripheral collapse under irreducibility at all times). Lean ✓ Stmt *If a quantum dynamical semigroup is irreducible at every positive time, then every peripheral eigenvalue of every positive-time slice equals 1.*

Proof. ✓ Proof Combine Theorem 17.4.1.17 with Theorem 17.4.1.18. □

Theorem 17.4.1.20 (Primitivity from irreducibility at all times). Lean ✓ Stmt *If a quantum dynamical semigroup is irreducible at every positive time, then every positive-time slice is primitive.*

Proof. ✓ Proof The unique faithful fixed density gives the one-dimensional peripheral spectrum criterion, and Theorem 17.4.1.19 collapses the peripheral set to $\{1\}$. □

Remark 17.4.1.21 (Semigroup irreducible/primitive equivalence). ✓ Stmt The semigroup analog of the irreducible/primitive equivalence starts from one irreducible time slice, constructs a primitive smaller time step, and lifts the conclusion to the full semigroup (Theorems 17.4.1.37 and 17.4.1.38).

Remark 17.4.1.22 (Reduction to a smaller time step for semigroups). ✓ Stmt Starting from one irreducible time t_0 , we construct a positive time step $u = t_0/N$ with $T_{t_0} = (T_u)^N$, prove peripheral eigenvalue collapse at time u , and deduce primitivity of T_u .

Theorem 17.4.1.23 (The stationary density is fixed for all times). Lean ✓ Stmt *If $t_0 > 0$, T_{t_0} is irreducible, and σ is its unique fixed density matrix, then $T_u(\sigma) = \sigma$ for every $u \geq 0$.*

Proof. ✓ Proof By semigroup commutativity, $T_{t_0}(T_u(\sigma)) = T_u(T_{t_0}(\sigma)) = T_u(\sigma)$, and uniqueness of the fixed density at time t_0 identifies $T_u(\sigma)$ with σ . □

Theorem 17.4.1.24 (Fixed points at the irreducible time are trace multiples). Lean ✓ Stmt *If T_{t_0} is irreducible with faithful fixed density σ , then every fixed point V of T_{t_0} satisfies $V = \text{tr}(V)\sigma$.*

Proof. ✓ Proof This is the fixed-point trace-scalar theorem for the irreducible channel T_{t_0} . □

Definition 17.4.1.25 (Residual slice index). Lean ✓ Stmt For $u \geq 0$ and $s > 0$, the residual slice index is

$$m_n = \left\lfloor \frac{nu}{s} \right\rfloor.$$

Definition 17.4.1.26 (Residual slice time). Lean ✓ Stmt For $u \geq 0$ and $s > 0$, the residual slice time is the remainder

$$r_n = s \text{ fract}\left(\frac{nu}{s}\right).$$

Theorem 17.4.1.27 (Residual slice decomposition). Lean Lean ✓ Stmt *For $u \geq 0$ and $s > 0$, one has $r_n \in [0, s]$ and*

$$nu = m_n s + r_n.$$

Proof. ✓ Proof This is the floor-plus-fraction decomposition of nu/s , multiplied by s . □

Theorem 17.4.1.28 (A residual slice can vanish). Lean ✓ Stmt *Suppose $u \geq 0$ and $s > 0$, and that $T_s(\delta) = \delta$ and $T_{nu}(\delta) \rightarrow 0$ as $n \rightarrow \infty$. Then there exists $a \in [0, s]$ such that $T_a(\delta) = 0$.*

Proof. ✓ Proof The residual slice times lie in the compact interval $[0, s]$, so a subsequence converges to some $a \in [0, s]$. The residual decomposition rewrites $T_{nu}(\delta)$ in terms of $T_{r_n}(\delta)$, and continuity of the semigroup identifies the limit with $T_a(\delta)$. □

Theorem 17.4.1.29 (Reduced-time eigenvector has nonzero trace). [Lean](#) [✓ Stmt](#) Under the reduced-time hypotheses $T_{t_0} = (T_u)^{(\dim M_D(\mathbb{C}))!}$, if μ is a peripheral eigenvalue of T_u , then T_u admits a μ -eigenvector with nonzero trace.

Theorem 17.4.1.30 (Peripheral collapse at the reduced time step). [Lean](#) [✓ Stmt](#) Under the same hypotheses, every peripheral eigenvalue of T_u equals 1.

Theorem 17.4.1.31 (Primitivity at the reduced time step). [Lean](#) [✓ Stmt](#) Under the same hypotheses, T_u is primitive.

Theorem 17.4.1.32 (Irreducibility descends to a reduced time step). [Lean](#) [✓ Stmt](#) If $T_{t_0} = (T_u)^N$ and T_{t_0} is irreducible, then T_u is irreducible.

Proof. [✓ Proof](#) Any invariant compression for T_u remains invariant under powers, hence also for T_{t_0} ; irreducibility of T_{t_0} rules this out. $\square$

Theorem 17.4.1.33 (Existence of an irreducible reduced time step). [Lean](#) [✓ Stmt](#) If $t_0 > 0$, T_{t_0} is irreducible, and σ is fixed for all times, then there exists a positive time step u such that

$$T_{t_0} = (T_u)^{(\dim M_D(\mathbb{C}))!},$$

the slice T_u is a channel, T_u is irreducible, and $T_u(\sigma) = \sigma$.

Proof. [✓ Proof](#) Take $u = t_0/N$ with $N = (\dim M_D(\mathbb{C}))!$. The semigroup power identity gives the factorization of T_{t_0} , channelity comes from the semigroup hypotheses, and Theorem 17.4.1.32 gives irreducibility of T_u . $\square$

Theorem 17.4.1.34 (Fixed points of the primitive reduced time step are trace-scalars). [Lean](#)

[✓ Stmt](#) If T_u is primitive and σ is the common fixed density, then every fixed point of any positive-time slice T_s has the form $\tau = \text{tr}(\tau)\sigma$.

Theorem 17.4.1.35 (Existence of a primitive reduced time step). [Lean](#) [✓ Stmt](#) If T_{t_0} is irreducible for some $t_0 > 0$, then there exists a positive reduced time step u such that T_u is primitive.

Theorem 17.4.1.36 (Irreducibility propagates to all positive times). [Lean](#) [✓ Stmt](#) If T_{t_0} is irreducible for some $t_0 > 0$, then every positive-time slice T_s is irreducible.

Proof. [✓ Proof](#) Let σ be the unique faithful fixed density of T_{t_0} . By Theorem 17.4.1.23, it is fixed for all times, and Theorem 17.4.1.24 makes the fixed-point space of T_{t_0} one-dimensional. Choose a primitive reduced time step by Theorem 17.4.1.35; then the trace-scalar description of fixed points from Theorem 17.4.1.34 gives the uniqueness criterion for irreducibility at every positive time. $\square$

Theorem 17.4.1.37 (An irreducible semigroup is primitive). [Lean](#) [✓ Stmt](#) Let $T_t = e^{tL}$ be a quantum dynamical semigroup. If T_{t_0} is irreducible for some $t_0 > 0$, then T_t is primitive (and irreducible) for every $t > 0$. This is [Wol12, Proposition 7.5].

Proof. [✓ Proof](#) First apply Theorem 17.4.1.36 to propagate irreducibility from the single time t_0 to every positive time. Then apply Theorem 17.4.1.20 to conclude primitivity of every positive-time slice. $\square$

Theorem 17.4.1.38 (Irreducibility iff primitivity for a quantum dynamical semigroup). [Lean](#)

[✓ Stmt](#) For a quantum dynamical semigroup $T_t = e^{tL}$:

$$(\exists t_0 > 0, T_{t_0} \text{ irreducible}) \iff (\forall t > 0, T_t \text{ primitive and irreducible}).$$

This is [Wol12, Proposition 7.5].

Proof. [✓ Proof](#) Forward: apply Theorem [17.4.1.37](#). Reverse: take $t_0 = 1$ (or any positive time); the right-hand side already includes irreducibility of T_t for every $t > 0$. $\square$

17.5 Kernel of the adjoint Liouvillian

Definition 17.5.1 (Faithful stationary state). [Lean](#) [✓ Stmt](#) A generator L has a faithful stationary state if there exists a positive definite density matrix ρ_0 with $L(\rho_0) = 0$.

Definition 17.5.2 (Adjoint dissipator). [Lean](#) [✓ Stmt](#) For a Lindblad operator L_j , the adjoint dissipator on observables is

$$\mathcal{D}_{L_j}^*(A) = L_j^\dagger A L_j - \frac{1}{2} L_j^\dagger L_j A - \frac{1}{2} A L_j^\dagger L_j.$$

Definition 17.5.3 (Adjoint generator). [Lean](#) [✓ Stmt](#) The adjoint generator of a Lindblad form is

$$L^*(A) = i[H, A] + \sum_j \mathcal{D}_{L_j}^*(A).$$

Theorem 17.5.4 (Trace-pairing characterization of the adjoint generator). [Lean](#) [✓ Stmt](#) For every density matrix ρ and every observable A ,

$$\text{tr}(\rho L^*(A)) = \text{tr}(L(\rho) A).$$

Proof. [✓ Proof](#) Expand the Schrödinger and Heisenberg formulas and apply cyclicity of the trace term by term. $\square$

Definition 17.5.5 (Commutant of the Lindblad data). [Lean](#) [✓ Stmt](#) The commutant of a Lindblad form is the set of matrices commuting with H , with every Lindblad operator L_j , and with every adjoint $L_j^\dagger$.

Definition 17.5.6 (Adjoint kernel). [Lean](#) [✓ Stmt](#) The adjoint kernel is the set of observables A satisfying $L^*(A) = 0$.

Theorem 17.5.7 (Commutant lies in the adjoint kernel). [Lean](#) [Lean](#) [✓ Stmt](#) If an observable commutes with H , with every L_j , and with every $L_j^\dagger$, then it lies in $\ker(L^*)$.

Proof. [✓ Proof](#) The Hamiltonian commutator term vanishes, and each adjoint dissipator $\mathcal{D}_{L_j}^*(A)$ is zero under the commutation hypotheses. $\square$

Theorem 17.5.8 (Adjoint kernel lies in the commutant). [Lean](#) [✓ Stmt](#) Let L be a GKSL generator in Lindblad form with Hamiltonian H and Lindblad operators $\{L_j\}$. Define the adjoint generator L^* by $\text{tr}(\rho L^*(A)) = \text{tr}(L(\rho) A)$. If L has a faithful stationary state $\rho_0 > 0$ with $L(\rho_0) = 0$, then for any A :

$$L^*(A) = 0 \implies [A, H] = [A, L_j] = [A, L_j^\dagger] = 0 \quad \forall j.$$

That is, $\ker(L^*) \subseteq \{H, L_j, L_j^\dagger\}'$. This is [[Wol12](#), Theorem 7.2].

Proof. [✓ Proof](#) From $L^*(A) = 0$ and $L^*(A^\dagger) = 0$, compute $L^*(A^\dagger A) = \sum_j [A, L_j]^\dagger [A, L_j]$. Faithfulness of ρ_0 and $\text{tr}(\rho_0 L^*(A^\dagger A)) = 0$ force $[A, L_j] = 0$ for all j . Similarly $[A, L_j^\dagger] = 0$, and then $L^*(A) = 0$ directly gives $[A, H] = 0$. $\square$

Theorem 17.5.9 (Adjoint kernel equals the commutant). [Lean](#) [✓ Stmt](#) Under the same hypotheses as Theorem 17.5.8,

$$\ker(L^*) = \{H, L_j, L_j^\dagger\}',$$

i.e. the adjoint kernel equals the commutant of $\{H, L_j, L_j^\dagger\}$. This is [Wol12, Theorem 7.2].

Proof. [✓ Proof](#) The inclusion $\supseteq$ is the easy direction (commutant elements are in the kernel by a direct calculation). The reverse $\subseteq$ is Theorem 17.5.8. $\square$

17.6 Reducibility of quantum dynamical semigroups

The four equivalent reducibility conditions of [Wol12, Proposition 7.6] are as follows.

Definition 17.6.1 (Nontrivial projection). [Lean](#) [✓ Stmt](#) A projection is nontrivial if it is orthogonal and neither 0 nor $\mathbb{1}$.

Definition 17.6.2 (Rank-deficient fixed density). [Lean](#) [✓ Stmt](#) Condition (1): there exists a density matrix ρ_0 with nontrivial kernel such that $e^{tL}(\rho_0) = \rho_0$ for all $t \geq 0$.

Definition 17.6.3 (Rank-deficient kernel element). [Lean](#) [✓ Stmt](#) Condition (2): there exists a density matrix ρ_0 with nontrivial kernel satisfying $L(\rho_0) = 0$.

Definition 17.6.4 (Invariant compression). [Lean](#) [✓ Stmt](#) Condition (3): there exists a nontrivial orthogonal projector P such that $e^{tL}(PM_D(\mathbb{C})P) \subseteq PM_D(\mathbb{C})P$ for all $t \geq 0$.

Definition 17.6.5 (Block-upper-triangular Lindblad form). [Lean](#) [✓ Stmt](#) Condition (4): there exists a nontrivial projector P and a Lindblad form $(H, \{L_j\})$ for L such that $(1 - P)L_jP = 0$ and $(1 - P)\kappa P = 0$ for all j , where $\kappa = iH + \frac{1}{2} \sum_j L_j^\dagger L_j$.

Definition 17.6.6 (Generator-preserved compression). [Lean](#) [✓ Stmt](#) For a projection P , the generator preserves the compression $PM_d(\mathbb{C})P$ if

$$P L(PXP) P = L(PXP)$$

for every $X \in M_d(\mathbb{C})$.

Definition 17.6.7 (Reducible quantum dynamical semigroup). [Lean](#) [✓ Stmt](#) A quantum dynamical semigroup is reducible if it has a nontrivial invariant compression.

Theorem 17.6.8 (Semigroup invariance implies generator invariance). [Lean](#) [✓ Stmt](#) If the semigroup preserves $PM_d(\mathbb{C})P$ for all non-negative times, then the generator preserves the same compression.

Proof. [✓ Proof](#) Differentiate the identity $Pe^{tL}(PXP)P = e^{tL}(PXP)$ at $t = 0$. $\square$

Theorem 17.6.9 (Generator invariance implies semigroup invariance). [Lean](#) [✓ Stmt](#) If the generator preserves $PM_d(\mathbb{C})P$, then so does the exponential semigroup e^{tL} for every $t \geq 0$.

Proof. [✓ Proof](#) Apply the compression map termwise to the exponential series. Every iterate of L preserves the compression, so the whole series does as well. $\square$

Theorem 17.6.10 (Reducibility as a generator-level criterion). [Lean](#) [✓ Stmt](#) A quantum dynamical semigroup is reducible if and only if its generator preserves some nontrivial compression.

Proof. [✓ Proof](#) Combine Theorems 17.6.8 and 17.6.9. □

Theorem 17.6.11 (Rank-deficient fixed densities and kernel elements). [Lean](#) [✓ Stmt](#) *A rank-deficient density matrix is a fixed point of e^{tL} for all $t \geq 0$ if and only if it lies in $\ker L$. This is [Wol12, Proposition 7.6, (1) $\Leftrightarrow$ (2)].*

Proof. [✓ Proof](#) The equivalence $L(\rho) = 0 \iff e^{tL}(\rho) = \rho$ for all $t \geq 0$ follows from the generator-semigroup correspondence: differentiate at $t = 0$ for the forward direction, and exponentiate for the reverse. □

Theorem 17.6.12 (A rank-deficient fixed density yields an invariant compression). [Lean](#) [✓ Stmt](#) *If L is a GKSL generator and there exists a rank-deficient fixed density matrix, then the semigroup preserves a nontrivial compression. This is [Wol12, Proposition 7.6, (1) $\Rightarrow$ (3)].*

Proof. [✓ Proof](#) Let ρ_0 be a rank-deficient fixed density. Let P be the support projection of ρ_0 . Since ρ_0 is rank-deficient, $P \neq \mathbb{1}$; since $\rho_0 \neq 0$, $P \neq 0$. Channel positivity and trace preservation force e^{tL} to preserve $PM_D(\mathbb{C})P$. □

Theorem 17.6.13 (Invariant compression yields block-upper-triangular Lindblad data). [Lean](#) [✓ Stmt](#) *If a GKSL generator preserves a nontrivial compression, then its Lindblad data can be chosen block-upper-triangular with respect to that compression.*

Proof. [✓ Proof](#) First pass from semigroup invariance to generator invariance by Theorem 17.6.8. The generator-level vanishing condition forces $(1 - P)L_j P = 0$ and $(1 - P)\kappa P = 0$ through the sum-of-squares identity. □

Theorem 17.6.14 (Block-upper-triangular Lindblad data yield invariant compression). [Lean](#) [✓ Stmt](#) *If the Lindblad data are block-upper-triangular with respect to a nontrivial projection, then the associated semigroup preserves that compression.*

Proof. [✓ Proof](#) The block-upper-triangular relations imply generator-level compression invariance; then apply Theorem 17.6.9. □

Theorem 17.6.15 (Invariant compression and triangular Lindblad data). [Lean](#) [Lean](#) [✓ Stmt](#) *For a GKSL generator L , the semigroup preserves a nontrivial compression if and only if the Lindblad data can be chosen block-upper-triangular. This is [Wol12, Proposition 7.6, (3) $\Leftrightarrow$ (4)].*

Proof. [✓ Proof](#) Combine Theorems 17.6.14 and 17.6.13. □

Theorem 17.6.16 (Triangular Lindblad data give deficient kernels). [Lean](#) [✓ Stmt](#) *If the Lindblad data of a GKSL generator are block-upper-triangular with respect to some nontrivial projector P , then there exists a density matrix ρ_0 with nontrivial kernel satisfying $L(\rho_0) = 0$. This is [Wol12, Proposition 7.6, (4) $\Rightarrow$ (2)].*

Proof. [✓ Proof](#) The block-upper-triangular structure implies L preserves the compression $PM_d(\mathbb{C})P$. Therefore the semigroup e^{tL} also preserves $PM_d(\mathbb{C})P$ for all $t \geq 0$. By the Brouwer fixed-point theorem, $e^{(1/m)L}$ has a fixed density matrix ρ_m supported in $PM_d(\mathbb{C})P$ for each m . Passing to a subsequential limit gives a density matrix ρ_0 with $P\rho_0 P = \rho_0$ and $L(\rho_0) = 0$. Since $P \neq \mathbb{1}$, the matrix ρ_0 has nontrivial kernel. □

Theorem 17.6.17 (Equivalent formulations of reducibility). [Lean](#) [✓ Stmt](#) *For a GKSL generator $L : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$, the four conditions*

1. rank-deficient fixed density,

2. rank-deficient kernel element,
3. invariant compression, and
4. block-upper-triangular Lindblad form

are equivalent. This is [Wol12, Proposition 7.6].

Proof. ✓ Proof The cycle $(1) \leftrightarrow (2)$, $(1) \rightarrow (3)$, $(3) \leftrightarrow (4)$, and $(4) \rightarrow (2)$ closes the equivalence chain. □

17.6.1 Sufficient conditions for non-reducibility

Definition 17.6.1.1 (Lindblad span). Lean ✓ Stmt The $\mathbb{C}$ -linear span of the Lindblad operators $\{L_j\}$.

Definition 17.6.1.2 (Hermitian-closed Lindblad span). Lean ✓ Stmt The span of the Lindblad operators is closed under Hermitian conjugation, i.e. it forms a $*$ -subspace of $M_d(\mathbb{C})$.

Definition 17.6.1.3 (Trivial Lindblad commutant). Lean ✓ Stmt The commutant of the Lindblad family $\{L_j\}'$ equals $\mathbb{C} \cdot \mathbb{1}$, i.e. the Lindblad operators act irreducibly on $M_d(\mathbb{C})$.

Definition 17.6.1.4 (Kossakowski rank). Lean ✓ Stmt The minimal number of Lindblad operators across all GKSL representations of L , which equals the rank of the Kossakowski matrix in the orthonormal-basis representation.

Theorem 17.6.1.5 (κ block vanishing from generator compression). Lean ✓ Stmt Let $(H, \{L_j\})$ be a Lindblad form with $\kappa = iH + \frac{1}{2} \sum_j L_j^\dagger L_j$, and let P be an orthogonal projection such that the generator preserves the compression $PM_d(\mathbb{C})P$. If $(1 - P)L_jP = 0$ for every j , then $(1 - P)\kappa P = 0$.

Proof. ✓ Proof From $L = \varphi - \kappa(\cdot) - (\cdot)\kappa^\dagger$ and $(1 - P)L(P) = 0$: the $(1 - P)\varphi(P)$ term vanishes because each $(1 - P)L_jP = 0$, and the $(1 - P)(P\kappa^\dagger)$ term vanishes because $(1 - P)P = 0$. What remains is $(1 - P)\kappa P = 0$. □

Theorem 17.6.1.6 (Full algebra generation forbids block-upper-triangular form). Lean ✓ Stmt Let $(H, \{L_j\})$ be a Lindblad form. If the algebra generated by $\{L_j\}$ and κ is the entire matrix algebra $M_d(\mathbb{C})$, then the Lindblad data do not admit a block-upper-triangular decomposition.

Proof. ✓ Proof Assume for contradiction that a block-upper-triangular decomposition exists with nontrivial projection P . By Theorem 17.6.1.5, $(1 - P)\kappa P = 0$. Induction on elements of the subalgebra shows $(1 - P)AP = 0$ for every generator. Since the generated algebra is all of $M_d(\mathbb{C})$, every matrix satisfies $(1 - P)AP = 0$, forcing $P = 0$ or $P = \mathbb{1}$ — a contradiction. □

Theorem 17.6.1.7 (Hermitian span with trivial commutant forbids triangular form). Lean ✓ Stmt If the Lindblad span is closed under Hermitian conjugation and its commutant is $\mathbb{C} \cdot \mathbb{1}$, then the Lindblad data do not admit a block-upper-triangular decomposition.

Proof. ✓ Proof Assume for contradiction that a block-upper-triangular decomposition exists with nontrivial projection P . The block vanishing $(1 - P)L_jP = 0$ and Hermitian closure give $PL_j(1 - P) = 0$ as well, so $[P, L_j] = 0$ for all j . But then P lies in the commutant of the Lindblad span, contradicting triviality. □

Theorem 17.6.1.8 (Large Kossakowski rank forbids block-upper-triangular form). [Lean](#) [✓ Stmt](#)
If the Kossakowski rank satisfies

$$\text{rank}(C) + d \geq d^2 + 1,$$

then no block-upper-triangular Lindblad form exists.

Proof. [✓ Proof](#) A block-upper-triangular traceless Lindblad family lies in a proper subspace whose dimension is at most $d^2 - d$. The formal rank minimization argument then contradicts the large-rank bound. $\square$

Theorem 17.6.1.9 (No block-upper-triangular Lindblad form implies non-reducibility). [Lean](#) [✓ Stmt](#)
If a GKSL generator admits no block-upper-triangular Lindblad form, then the associated quantum dynamical semigroup is not reducible.

Proof. [✓ Proof](#) A reducible semigroup would have an invariant compression, and Theorem 17.6.13 would then produce a block-upper-triangular Lindblad form. $\square$

Theorem 17.6.1.10 (Full algebra generation implies non-reducibility). [Lean](#) [✓ Stmt](#) *If the algebra generated by $\{L_j\}$ and κ is the entire matrix algebra $M_d(\mathbb{C})$, then the QDS is not reducible. This is [Wol12, Corollary 7.2(1)].*

Proof. [✓ Proof](#) By Theorem 17.6.1.6, no block-upper-triangular decomposition exists. Then apply Theorem 17.6.1.9. $\square$

Theorem 17.6.1.11 (Hermitian span and trivial commutant imply non-reducibility). [Lean](#) [✓ Stmt](#)
If the Lindblad span is Hermitian-closed and its commutant is trivial, then the QDS is not reducible. This is [Wol12, Corollary 7.2(2)].

Proof. [✓ Proof](#) By Theorem 17.6.1.7, no block-upper-triangular decomposition exists. Then apply Theorem 17.6.1.9. $\square$

Theorem 17.6.1.12 (Large Kossakowski rank implies non-reducibility). [Lean](#) [✓ Stmt](#) *If $\text{rank}(C) > d^2 - d$, then the QDS is not reducible. This is [Wol12, Corollary 7.2(3)].*

Proof. [✓ Proof](#) The rank hypothesis rules out block-upper-triangular Lindblad data by Theorem 17.6.1.8. Then apply Theorem 17.6.1.9. $\square$

Chapter 18

Operator Convexity and Jensen Inequalities

This chapter collects the operator convexity/concavity results, trace inequalities, the operator Jensen inequality for positive maps, and the Lieb concavity theorem. These are standard results in matrix analysis ([Bha97, Chapter V]; [Wol12, Theorems 5.12 and 5.13]; Hansen–Pedersen’s Jensen inequality [HP82]; and Lieb’s concavity theorem [Lie73]). The operator Jensen inequalities for real powers $x \mapsto x^p$ on the Loewner order are obtained from the Loewner integral representation of the power function. Lieb’s concavity theorem is derived from the operator integral representation of the fractional product together with the vectorization isometry, which turns the trace functional into a quadratic form of a positive-semidefinite matrix.

18.1 Diagonal Jensen inequality

Lemma 18.1.1 (Diagonal Jensen inequality). Lean ✓ Stmt Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be convex on $[0, \infty)$, let A be a positive semidefinite matrix on a finite-dimensional space indexed by n , and let $v \in \mathbb{C}^n$ satisfy $\langle v, v \rangle = 1$. Then

$$f(\Re \langle v, Av \rangle) \leq \Re \langle v, f(A)v \rangle,$$

where $f(A)$ is defined by the Hermitian continuous functional calculus.

Proof. ✓ Proof Write $A = U \operatorname{diag}(\mu) U^*$ by the spectral theorem and set $w = U^*v$. Since U is unitary, $p_i := |w_i|^2$ satisfies $\sum_i p_i = \|v\|^2 = 1$, so (p_i) is a probability distribution on n . The eigenvalues μ_i lie in $[0, \infty)$ because A is positive semidefinite. A direct computation yields $\Re \langle v, Av \rangle = \sum_i p_i \mu_i$ and $\Re \langle v, f(A)v \rangle = \sum_i p_i f(\mu_i)$, so the scalar Jensen inequality applied to the weights (p_i) and points (μ_i) gives the conclusion. See [Bha97, Chapter V]. $\square$

18.2 Trace concavity and convexity of matrix powers

Theorem 18.2.1 (Trace concavity of real powers). Lean Lean ✓ Stmt For $p \in [0, 1]$, PSD matrices A_1, A_2 , and $t \in [0, 1]$:

$$t \Re \operatorname{tr}(A_1^p) + (1 - t) \Re \operatorname{tr}(A_2^p) \leq \Re \operatorname{tr}((tA_1 + (1 - t)A_2)^p).$$

Follows from operator concavity of $x \mapsto x^p$ composed with trace monotonicity on the Loewner order.

Proof. ✓ Proof Let $A = tA_1 + (1-t)A_2$ and let $\{\psi_j\}$ with eigenvalues $\mu_j \geq 0$ be an orthonormal eigenbasis of A . A direct computation gives $\Re \operatorname{tr}(A^p) = \sum_j \mu_j^p$ and $\mu_j = t a_j + (1-t) b_j$ where $a_j = \Re \langle \psi_j, A_1 \psi_j \rangle$, $b_j = \Re \langle \psi_j, A_2 \psi_j \rangle$. Scalar concavity of $x \mapsto x^p$ on $[0, \infty)$ yields $\mu_j^p \geq t a_j^p + (1-t) b_j^p$, and the diagonal Jensen inequality (Lemma 18.1.1, concave form) gives $a_j^p \leq \Re \langle \psi_j, A_1^p \psi_j \rangle$ (and similarly for b_j). Summing over j and using $\sum_j \Re \langle \psi_j, M \psi_j \rangle = \Re \operatorname{tr} M$ yields the conclusion. See [Bha97, Chapter V]. $\square$

Theorem 18.2.2 (Trace convexity of real powers). Lean Lean ✓ Stmt For $p \in [1, 2]$, PSD matrices A_1, A_2 , and $t \in [0, 1]$:

$$\Re \operatorname{tr}((tA_1 + (1-t)A_2)^p) \leq t \Re \operatorname{tr}(A_1^p) + (1-t) \Re \operatorname{tr}(A_2^p).$$

Follows from operator convexity of $x \mapsto x^p$ composed with trace monotonicity on the Loewner order.

Proof. ✓ Proof Identical eigenbasis reduction as in Theorem 18.2.1, with scalar convexity (rather than concavity) of $x \mapsto x^p$ on $[0, \infty)$ for $p \geq 1$ and the convex form of the diagonal Jensen inequality. $\square$

18.3 Operator convexity of real powers

Theorem 18.3.1 (Operator convexity of real powers between one and two). Lean ✓ Stmt For elements a, b of a unital C^* -algebra with $0 \leq a, b$, an exponent $p \in [1, 2]$, and $t \in [0, 1]$, the map $a \mapsto a^p$ is convex for the Loewner order on the positive cone:

$$(ta + (1-t)b)^p \leq t a^p + (1-t) b^p.$$

This is the convex counterpart of the operator concavity of $x \mapsto x^p$ on $[0, 1]$.

Proof. ✓ Proof For $p \in (1, 2)$ one uses the integral representation $a^p = \int_0^\infty f_{p,s}(a) d\mu(s)$ of Carlen [Car10, Lemma 2.8], whose integrand $f_{p,s}(x) = s^{p-1}(s^{-1}x + s(s+x)^{-1} - 1)$ is operator convex: it decomposes into a linear term, a nonnegative multiple of the operator convex resolvent $x \mapsto (s \cdot 1 + x)^{-1}$, and a constant. Since $f_{p,s}(ta + (1-t)b) \leq t f_{p,s}(a) + (1-t) f_{p,s}(b)$ for every $s > 0$, integrating this pointwise Loewner inequality against μ gives

$$(ta + (1-t)b)^p = \int_0^\infty f_{p,s}(ta + (1-t)b) d\mu(s) \leq t \int_0^\infty f_{p,s}(a) d\mu(s) + (1-t) \int_0^\infty f_{p,s}(b) d\mu(s) = t a^p + (1-t) b^p.$$

The endpoint $p = 1$ is the identity, and $p = 2$ is $a \mapsto a^2$, operator convex because $t a^2 + (1-t) b^2 - (ta + (1-t)b)^2 = t(1-t)(a-b)^2 \geq 0$. See [Bha97, Chapter V]. $\square$

18.4 Operator Jensen inequality for positive maps

Definition 18.4.1 (Finite-POVM dilation construction). Lean Lean ✓ Stmt For a finite family of matrices $\{C_i\}_{i \in \iota}$ and a defect block S , one forms an isometric dilation whose rows are the adjoints of the matrices C_i together with the adjoint of S . One also forms the scalar block-diagonal matrix whose ι -blocks carry prescribed weights and whose defect block carries a single scalar.

Lemma 18.4.2 (Compression auxiliary for Jensen). [Lean](#) [✓ Stmt](#) Let Y be positive definite and let W be an isometry. Then

$$(W^*YW)^{-1} \leq W^*Y^{-1}W. \quad (18.1)$$

Proof. [✓ Proof](#) Apply the Schur-complement criterion to the block matrix $\begin{pmatrix} Y^{-1} & W \\ W^* & W^*YW \end{pmatrix}$. The lower-right Schur complement is zero, hence positive semidefinite, and compressing the resulting upper-left Schur complement by W gives the inequality (18.1). $\square$

Lemma 18.4.3 (Finite-POVM compression identities). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) If the defect block satisfies

$$SS^* = \mathbb{1} - \sum_{i \in \iota} C_i C_i^*, \quad (18.2)$$

then the dilation is an isometry, compressing the weighted scalar block-diagonal matrix gives

$$\sum_{i \in \iota} w_i C_i C_i^* + t SS^*,$$

the defect relation rewrites as

$$\sum_{i \in \iota} C_i C_i^* + SS^* = \mathbb{1},$$

and strict positivity of all weights together with $t > 0$ implies positive definiteness of the scalar block-diagonal matrix. If all weights are nonzero and the defect scalar is nonzero, the inverse diagonal has reciprocal weights, and compressing this inverse gives

$$\sum_{i \in \iota} w_i^{-1} C_i C_i^* + t^{-1} SS^*.$$

Proof. [✓ Proof](#) Expand the block matrix multiplication entrywise. The defect identity (18.2) gives the isometry relation after summing the ι -blocks and the defect block, and the weighted compression identity follows from the same expansion with the scalar diagonal inserted. Positive definiteness uses strict positivity of every diagonal entry, while the inverse formula uses the nonvanishing of every diagonal entry. $\square$

Lemma 18.4.4 (Finite-POVM resolvent inequality). [Lean](#) [✓ Stmt](#) Let $w_i \geq 0$ and $t > 0$. If the defect block satisfies

$$SS^* = \mathbb{1} - \sum_{i \in \iota} C_i C_i^*,$$

then

$$\left(\sum_{i \in \iota} w_i C_i C_i^* + t \mathbb{1} \right)^{-1} \leq \sum_{i \in \iota} (w_i + t)^{-1} C_i C_i^* + t^{-1} SS^*. \quad (18.3)$$

Proof. [✓ Proof](#) Apply the compression inequality to the positive definite scalar block-diagonal matrix with entries $w_i + t$ and defect entry t . The compression identities rewrite the compressed matrix as $\sum_i w_i C_i C_i^* + t \mathbb{1}$ and its inverse compression as the upper bound (18.3). $\square$

Lemma 18.4.5 (Spectral finite-POVM resolution). [Lean](#) [✓ Stmt](#) Let $A \geq 0$ be a finite matrix. There are positive semidefinite matrices P_i and nonnegative real numbers λ_i such that

$$\sum_i P_i = \mathbb{1}, \quad A = \sum_i \lambda_i P_i.$$

Proof. ✓ Proof Diagonalize the Hermitian matrix A by a unitary matrix. The matrices P_i are the conjugates of the rank-one diagonal matrix units, their sum is the identity, and positive semidefiniteness of A gives $\lambda_i \geq 0$. $\square$

Lemma 18.4.6 (Positive-map resolvent inequality). Lean ✓ Stmt Let T be a positive subunital map, let $A \geq 0$, and let $t > 0$. Then

$$(TA + t\mathbb{1})^{-1} \leq T((A + t\mathbb{1})^{-1}) + t^{-1}(\mathbb{1} - T\mathbb{1}).$$

Proof. ✓ Proof Use the spectral resolution $A = \sum_i \lambda_i P_i$. The matrices $T(P_i)$ are positive semidefinite and sum to $T\mathbb{1} \leq \mathbb{1}$. Write each $T(P_i)$ as $C_i C_i^*$ using the positive square root, and write the defect $\mathbb{1} - \sum_i T(P_i)$ as SS^* . The finite-POVM resolvent inequality gives the desired estimate after substituting the spectral formula for $(A + t\mathbb{1})^{-1}$. $\square$

Lemma 18.4.7 (Positive-map real-power integrand inequality). Lean ✓ Stmt Let T be a positive subunital map, let $A \geq 0$, let $p \in (0, 1)$, and let $t > 0$. Then

$$T(f_{p,t}(A)) \leq f_{p,t}(TA), \quad f_{p,t}(x) = t^p (t^{-1} - (t + x)^{-1}).$$

Proof. ✓ Proof Rewrite the continuous functional calculus for $f_{p,t}$ in resolvent form,

$$f_{p,t}(X) = t^{p-1}\mathbb{1} - t^p(t\mathbb{1} + X)^{-1}.$$

After applying the linear map T , the difference $f_{p,t}(TA) - T(f_{p,t}(A))$ is exactly t^p times the positive matrix supplied by the positive-map resolvent inequality. $\square$

Lemma 18.4.8 (Positive-map convex real-power integrand inequality). Lean ✓ Stmt Let T be a positive subunital map, let $A \geq 0$, let $p \in (1, 2)$, and let $t > 0$. Then

$$g_{p,t}(TA) \leq T(g_{p,t}(A)), \quad g_{p,t}(x) = t^{p-2}x + t^p(t + x)^{-1} - t^{p-1}.$$

The inequality is reversed relative to the concave integrand, since $g_{p,t}$ is operator convex.

Proof. ✓ Proof Rewrite the continuous functional calculus for $g_{p,t}$ in resolvent form,

$$g_{p,t}(X) = t^{p-2}X + t^p(t\mathbb{1} + X)^{-1} - t^{p-1}\mathbb{1}.$$

Under the linear map T the leading t^{p-2} terms cancel and the constant t^{p-1} terms combine, so the difference $T(g_{p,t}(A)) - g_{p,t}(TA)$ is exactly t^p times the positive matrix supplied by the positive-map resolvent inequality (using $t^p t^{-1} = t^{p-1}$). $\square$

Lemma 18.4.9 (Positive matrix-valued integrals). Lean ✓ Stmt Let f be a function from a measure space to a finite matrix algebra. If $f(x) \geq 0$ for almost every x , then $\int f(x) d\mu(x) \geq 0$.

Proof. ✓ Proof The Loewner order has closed positive cone in finite dimension. Hence the ordered Bochner integral preserves almost-everywhere nonnegativity. The corresponding almost-everywhere monotonicity statement is the general ordered Bochner integral theorem. $\square$

Theorem 18.4.10 (Jensen for concave real powers). Lean ✓ Stmt For a positive subunital map T , a positive semidefinite matrix A , and $p \in [0, 1]$:

$$T(A^p) \leq (TA)^p.$$

This is [Wol12, Theorem 5.13] for concave $f(x) = x^p$.

Proof. ✓ Proof For $p = 0$ the assertion is the subunitarity condition $T(\mathbb{1}) \leq \mathbb{1}$. For $p = 1$ it is an equality. For $0 < p < 1$, use Carlen's integral representation of A^p by the functions $f_{p,t}(x) = t^p(t^{-1} - (t+x)^{-1})$. The pointwise inequality $T(f_{p,t}(A)) \leq f_{p,t}(TA)$ is Lemma 18.4.7. Integrating this inequality with respect to the representing measure gives the result by Lemma 18.4.9 and linearity of the Bochner integral under T . □

Theorem 18.4.11 (Map form of Jensen for concave real powers). Lean ✓ Stmt For a positive subunital map T , a positive semidefinite matrix A , and $p \in [0, 1]$, the inequality $T(A^p) \leq (TA)^p$ holds.

Proof. ✓ Proof Direct application of Theorem 18.4.10. □

Theorem 18.4.12 (Jensen for convex real powers). Lean ✓ Stmt For a positive subunital map T , a positive semidefinite matrix A , and $p \in [1, 2]$:

$$(TA)^p \leq T(A^p).$$

This is [Wol12, Theorem 5.11] for the subunital operator-convex case $f(x) = x^p$.

Proof. ✓ Proof For $p = 1$ the assertion is an equality. For $1 < p < 2$, use Carlen's integral representation of A^p by the convex integrands $g_{p,t}(x) = t^{p-2}x + t^p(t+x)^{-1} - t^{p-1}$. The pointwise inequality $g_{p,t}(TA) \leq T(g_{p,t}(A))$ is Lemma 18.4.8. Integrating this inequality with respect to the representing measure and using Lemma 18.4.9 together with linearity of the Bochner integral under T gives the inequality for $1 < p < 2$. The endpoint $p = 2$ follows by taking the limit $q \rightarrow 2^-$ in $(TA)^q \leq T(A^q)$: since $M^q \rightarrow M^2$ in the operator norm as $q \rightarrow 2^-$ for any positive semidefinite M , and the Loewner order is closed, this gives $(TA)^2 \leq T(A^2)$. □

Theorem 18.4.13 (Map form of Jensen for convex real powers). Lean ✓ Stmt For a positive subunital map T , a positive semidefinite matrix A , and $p \in [1, 2]$, the inequality $(TA)^p \leq T(A^p)$ holds.

Proof. ✓ Proof Direct application of the theorem. □

Theorem 18.4.14 (Jensen for concave log). Lean ✓ Stmt For a positive unital map T and positive-definite A :

$$T(\log A) \leq \log(TA).$$

This is [Wol12, Theorem 5.13] for $f = \log$. Requires unitality ($T(\mathbb{1}) = \mathbb{1}$), not merely subunitarity.

Proof. ✓ Proof Apply Theorem 18.4.10 to $0 < p < 1$ and rewrite it as an inequality for $(A^p - \mathbb{1})/p$. The continuous functional calculus gives $(A^p - \mathbb{1})/p \rightarrow \log A$ as $p \rightarrow 0^+$, and the same limit holds for $T(A)$. Since the Loewner order graph is closed in finite dimension, the limiting inequality is $T(\log A) \leq \log(TA)$. □

Theorem 18.4.15 (Map form of Jensen for the concave logarithm). Lean ✓ Stmt For a positive unital map T and a positive-definite matrix A , the inequality $T(\log A) \leq \log(TA)$ holds.

Proof. ✓ Proof Direct application of Theorem 18.4.14. □

18.5 Lieb concavity theorem

The Lieb concavity theorem rests on a scalar integral identity that expresses the geometric interpolation $a^s b^{1-s}$ as a positive average of the elementary fractions $ab/(a+tb)$. This is the analytic core of the integral representation of $A^s B^{1-s}$.

Lemma 18.5.1 (Reflection integral). [Lean](#) [✓ Stmt](#) For $s \in (0, 1)$,

$$\int_0^\infty \frac{u^{s-1}}{1+u} du = \frac{\pi}{\sin(\pi s)}.$$

Proof. [✓ Proof](#) This is the value $B(s, 1-s)$ of the Beta function, which equals $\Gamma(s)\Gamma(1-s) = \pi/\sin(\pi s)$ by Euler's reflection formula. The substitution $x = u/(1+u)$ carries the half-line onto the unit interval and turns the integrand into the Beta integrand $x^{s-1}(1-x)^{-s}$. $\square$

Lemma 18.5.2 (Integrability of the Lieb integrand). [Lean](#) [✓ Stmt](#) For $a, b > 0$ and $s \in (0, 1)$, the function $t \mapsto t^{s-1} \frac{ab}{a+tb}$ is integrable on $(0, \infty)$.

Proof. [✓ Proof](#) The integrand is $O(t^{s-1})$ as $t \rightarrow 0^+$ and $O(t^{s-2})$ as $t \rightarrow \infty$, so it has integrable comparison functions on each tail. $\square$

Lemma 18.5.3 (Value of the Lieb integral). [Lean](#) [✓ Stmt](#) For $a, b > 0$ and $s \in (0, 1)$,

$$\int_0^\infty t^{s-1} \frac{ab}{a+tb} dt = a^s b^{1-s} \frac{\pi}{\sin(\pi s)}.$$

Proof. [✓ Proof](#) The scaling substitution $t = (a/b)u$ turns the integrand into $a^s b^{1-s} u^{s-1}/(1+u)$, whose half-line integral is the reflection integral $\pi/\sin(\pi s)$ of Lemma 18.5.1. $\square$

Lemma 18.5.4 (Scalar Lieb integral identity). [Lean](#) [✓ Stmt](#) For $a, b > 0$ and $s \in (0, 1)$,

$$a^s b^{1-s} = \frac{\sin(\pi s)}{\pi} \int_0^\infty t^{s-1} \frac{ab}{a+tb} dt.$$

Proof. [✓ Proof](#) By Lemma 18.5.3 the integral equals $a^s b^{1-s} \pi/\sin(\pi s)$, so

$$\frac{\sin(\pi s)}{\pi} \int_0^\infty t^{s-1} \frac{ab}{a+tb} dt = \frac{\sin(\pi s)}{\pi} \cdot a^s b^{1-s} \frac{\pi}{\sin(\pi s)} = a^s b^{1-s}.$$

$\square$

Lemma 18.5.5 (Diagonal Lieb integral identity). [Lean](#) [✓ Stmt](#) For positive diagonal matrices M and N and $s \in (0, 1)$,

$$M^s N^{1-s} = \frac{\sin(\pi s)}{\pi} \int_0^\infty t^{s-1} M (M + tN)^{-1} N dt.$$

Proof. [✓ Proof](#) Both sides are diagonal, so the identity holds entry by entry; on each diagonal entry it is the scalar Lieb integral identity (Lemma 18.5.4). $\square$

Lemma 18.5.6 (Operator integral representation). [Lean](#) [Stmnt](#) For positive-definite matrices A, B and $s \in (0, 1)$,

$$(A \otimes I)^s (I \otimes B^\top)^{1-s} = \frac{\sin(\pi s)}{\pi} \int_0^\infty t^{s-1} (A \otimes I)((A \otimes I) + t(I \otimes B^\top))^{-1} (I \otimes B^\top) dt,$$

where $A \otimes I$ and $I \otimes B^\top$ are regarded as operators on the Kronecker model space $\mathbb{C}^{D \times D}$.

Proof. [Proof](#) The pair $(A \otimes I, I \otimes B^\top)$ is simultaneously diagonalized by $V = U_A \otimes U_{B^\top}$ (eigenbases of A and $B^\top$). In that basis the pair becomes a commuting pair of positive diagonal matrices, and the identity follows from the diagonal case (Lemma 18.5.5). $\square$

Lemma 18.5.7 (Concavity of the parallel sum). [Lean](#) [Stmnt](#) For positive-definite matrices X_1, X_2, Y_1, Y_2 and $\theta \in [0, 1]$, the parallel sum $(X, Y) \mapsto X(X + Y)^{-1}Y$ is jointly concave in the Loewner order:

$$\theta X_1(X_1 + Y_1)^{-1}Y_1 + (1 - \theta) X_2(X_2 + Y_2)^{-1}Y_2 \leq X_\theta(X_\theta + Y_\theta)^{-1}Y_\theta,$$

where $X_\theta = \theta X_1 + (1 - \theta)X_2$ and $Y_\theta = \theta Y_1 + (1 - \theta)Y_2$.

Proof. [Proof](#) Write $Z = X(X + Y)^{-1}Y$. The identity $X(X + Y)^{-1}(X + Y) = X$ gives

$$(X - Z) - X(X + Y)^{-1}X = 0,$$

so the block matrix $\begin{pmatrix} X & Z \\ Z^\top & X + Y \end{pmatrix}$ is positive semidefinite (its Schur complement against $X + Y$ vanishes). Setting $Z_i = X_i(X_i + Y_i)^{-1}Y_i$ and $Z_\theta = \theta Z_1 + (1 - \theta)Z_2$, the convex combination of the two block matrices is $\begin{pmatrix} X_\theta & Z_\theta \\ Z_\theta^\top & X_\theta + Y_\theta \end{pmatrix}$, which is positive semidefinite, and its Schur complement against $X_\theta + Y_\theta$ yields

$$X_\theta(X_\theta + Y_\theta)^{-1}Y_\theta - Z_\theta \geq 0,$$

which is the claimed inequality (Ando, 1979). $\square$

Lemma 18.5.8 (Concavity of the resolvent integrand). [Lean](#) [Stmnt](#) For $t > 0$, positive-definite A_1, A_2, B_1, B_2 , and $\theta \in [0, 1]$, with $\hat{A} = A \otimes I$ and $\hat{B} = I \otimes B^\top$, setting $A_\theta = \theta A_1 + (1 - \theta)A_2$ and $B_\theta = \theta B_1 + (1 - \theta)B_2$:

$$\theta \hat{A}_1(\hat{A}_1 + t\hat{B}_1)^{-1}\hat{B}_1 + (1 - \theta) \hat{A}_2(\hat{A}_2 + t\hat{B}_2)^{-1}\hat{B}_2 \leq \hat{A}_\theta(\hat{A}_\theta + t\hat{B}_\theta)^{-1}\hat{B}_\theta.$$

Proof. [Proof](#) The integrand equals $t^{-1} \hat{A}(\hat{A} + t\hat{B})^{-1}(t\hat{B})$, a rescaled parallel sum of $\hat{A}$ and $t\hat{B}$, so concavity follows from Lemma 18.5.7 and the linearity of $A \mapsto \hat{A}$, $B \mapsto \hat{B}$. $\square$

Lemma 18.5.9 (Concavity of the fractional product). [Lean](#) [Stmnt](#) For positive-definite matrices A_1, A_2, B_1, B_2 , $s \in (0, 1)$, and $\theta \in [0, 1]$, with $\hat{A} = A \otimes I$ and $\hat{B} = I \otimes B^\top$, setting $A_\theta = \theta A_1 + (1 - \theta)A_2$ and $B_\theta = \theta B_1 + (1 - \theta)B_2$, the fractional product $\hat{A}^s \hat{B}^{1-s}$ is jointly concave in the Loewner order:

$$\theta \hat{A}_1^s \hat{B}_1^{1-s} + (1 - \theta) \hat{A}_2^s \hat{B}_2^{1-s} \leq \hat{A}_\theta^s \hat{B}_\theta^{1-s}.$$

Proof. [Proof](#) By the integral representation 18.5.6 each fractional product equals the positive multiple $\frac{\sin(\pi s)}{\pi}$ of the integral of the resolvent integrand against the nonnegative weight t^{s-1} on $(0, \infty)$. For each $t > 0$ the integrand is jointly concave by Lemma 18.5.8, so the difference of the averaged point and the convex combination is, almost everywhere, the nonnegative scalar t^{s-1} times a positive-semidefinite matrix. Integrating a positive-semidefinite integrand yields a positive-semidefinite matrix, and scaling by the positive constant $\frac{\sin(\pi s)}{\pi}$ preserves the Loewner order, which gives the inequality. $\square$

Theorem 18.5.10 (Lieb concavity). Lean ✓ Stmt For $s \in [0, 1]$, any matrix K , and positive-definite matrices A_1, A_2, B_1, B_2 , the map $(A, B) \mapsto \Re \operatorname{tr}(K^\dagger A^s K B^{1-s})$ is jointly concave:

$$t \Re \operatorname{tr}(K^\dagger A_1^s K B_1^{1-s}) + (1-t) \Re \operatorname{tr}(K^\dagger A_2^s K B_2^{1-s}) \leq \Re \operatorname{tr}(K^\dagger (tA_1 + (1-t)A_2)^s K (tB_1 + (1-t)B_2)^{1-s}). \quad (18.4)$$

See [Lie73, And79]. This statement is the positive-definite, boundary-exponent ($x + y = 1$) case of the Ando–Lieb theorem [Wol12, Theorem 5.15], which holds for positive semidefinite A, B and all exponents $x, y \geq 0$ with $x + y \leq 1$.

Proof. ✓ Proof For $s \in (0, 1)$, the concavity of the fractional product of the commuting left and right multiplication operators (Lemma 18.5.9) gives a Loewner-order inequality between positive-semidefinite matrices on the model space $\mathbb{C}^D \otimes \mathbb{C}^D$. Writing that product as $A^s \otimes (B^\top)^{1-s}$ and pairing it with the vectorization of $K^\top$,

$$\langle \operatorname{vec} K^\top, (A^s \otimes (B^\top)^{1-s}) \operatorname{vec} K^\top \rangle = \Re \operatorname{tr}(K^\dagger A^s K B^{1-s}),$$

expresses the trace functional as the quadratic form of the Kronecker product at that vector. Applying this positive quadratic form to the Loewner inequality transfers the concavity to the real part of the trace. The endpoints $s = 0$ and $s = 1$ reduce the varying side to a linear function of the convex-combination argument, hence hold with equality. $\square$

Theorem 18.5.11 (Lieb concavity, positive-semidefinite inputs). Lean ✓ Stmt For $s \in [0, 1]$, any matrix K , and positive-semidefinite matrices A_1, A_2, B_1, B_2 , the map $(A, B) \mapsto \Re \operatorname{tr}(K^\dagger A^s K B^{1-s})$ is jointly concave, satisfying the inequality (18.4) of Theorem 18.5.10. This is the full Ando–Lieb theorem [Wol12, Theorem 5.15] on the boundary line $x + y = 1$ (with $x = s, y = 1 - s$): the positive-definiteness restriction of Theorem 18.5.10 is lifted. The general two-exponent region is obtained in Theorem 18.5.12.

Proof. ✓ Proof For the interior exponents $s \in (0, 1)$, replace each input A_i by the positive-definite matrix $A_i + \varepsilon \mathbb{1}$, and likewise for the B_i . Theorem 18.5.10 applies to these positive-definite matrices for every $\varepsilon > 0$, and the average of the regularized inputs is the regularization of the average. Letting $\varepsilon \rightarrow 0^+$, each regularized power $(A_i + \varepsilon \mathbb{1})^s$ converges to A_i^s , since $x \mapsto x^s$ is continuous on the nonnegative reals for $0 < s < 1$, including at the origin; the trace functional is continuous, so the regularized inequality passes to the limit. The endpoints $s = 0$ and $s = 1$ are linear in the convex-combination argument and hold with equality. $\square$

Theorem 18.5.12 (Lieb concavity, sub-boundary exponents). Lean ✓ Stmt Let $x, y \geq 0$ with $x + y \leq 1$. For any matrix K and positive-semidefinite matrices A_1, A_2, B_1, B_2 , the map $(A, B) \mapsto \Re \operatorname{tr}(K^\dagger A^x K B^y)$ is jointly concave:

$$t \Re \operatorname{tr}(K^\dagger A_1^x K B_1^y) + (1-t) \Re \operatorname{tr}(K^\dagger A_2^x K B_2^y) \leq \Re \operatorname{tr}(K^\dagger (tA_1 + (1-t)A_2)^x K (tB_1 + (1-t)B_2)^y)$$

for every $t \in [0, 1]$. This is Wolf’s Theorem 5.15 in the finite-dimensional matrix setting.

Proof. ✓ Proof Put $r = x + y$. If $r = 0$, both exponents vanish and the functional is constant. If $0 < r \leq 1$, set $s = x/r$. Then $x = rs$ and $y = r(1 - s)$, with $s \in [0, 1]$. The functional becomes the boundary Lieb functional applied to the powered inputs A^r and B^r . Operator concavity of $C \mapsto C^r$ gives

$$tA_1^r + (1-t)A_2^r \leq (tA_1 + (1-t)A_2)^r$$

and the analogous inequality for B . The boundary positive-semidefinite theorem gives concavity at the left-hand powered inputs, and monotonicity of $(P, Q) \mapsto \Re \operatorname{tr}(K^\dagger P^s K Q^{1-s})$ on positive-semidefinite arguments transports the result to the powered convex combinations. $\square$

Theorem 18.5.13 (Map form of Lieb concavity). [Lean](#) [✓ Stmt](#) For $s \in [0, 1]$, any matrix K , and positive-definite matrices A_1, A_2, B_1, B_2 , the function $(A, B) \mapsto \Re \operatorname{tr}(K^\dagger A^s K B^{1-s})$ is jointly concave, satisfying the inequality (18.4) of Theorem 18.5.10.

Proof. [✓ Proof](#) Follows directly from Theorem 18.5.10. $\square$

Corollary 18.5.14 (Lieb concavity, $K = \mathbb{1}$ case). [Lean](#) [✓ Stmt](#) For $s \in [0, 1]$, $t \in [0, 1]$, and positive-definite matrices A_1, A_2, B_1, B_2 , the map $(A, B) \mapsto \Re \operatorname{tr}(A^s B^{1-s})$ is jointly concave:

$$t \Re \operatorname{tr}(A_1^s B_1^{1-s}) + (1-t) \Re \operatorname{tr}(A_2^s B_2^{1-s}) \leq \Re \operatorname{tr}((tA_1 + (1-t)A_2)^s (tB_1 + (1-t)B_2)^{1-s}).$$

Obtained from Theorem 18.5.13 by taking $K = \mathbb{1}$.

Proof. [✓ Proof](#) Specialize Theorem 18.5.13 at $K = \mathbb{1}$ and simplify $\mathbb{1}^\dagger = \mathbb{1}$ together with $\mathbb{1} \cdot X = X \cdot \mathbb{1} = X$. $\square$

Corollary 18.5.15 (Concavity of Lieb's map in the first argument). [Lean](#) [✓ Stmt](#) For $s \in [0, 1]$, $t \in [0, 1]$, matrix K , and positive-definite matrices A_1, A_2, B , the map $A \mapsto \Re \operatorname{tr}(K^\dagger A^s K B^{1-s})$ is concave:

$$t \Re \operatorname{tr}(K^\dagger A_1^s K B^{1-s}) + (1-t) \Re \operatorname{tr}(K^\dagger A_2^s K B^{1-s}) \leq \Re \operatorname{tr}(K^\dagger (tA_1 + (1-t)A_2)^s K B^{1-s}).$$

Proof. [✓ Proof](#) Specialize Theorem 18.5.13 at $B_1 = B_2 = B$, so that the convex combination collapses: $tB + (1-t)B = B$. $\square$

Corollary 18.5.16 (Concavity of Lieb's map in the second argument). [Lean](#) [✓ Stmt](#) For $s \in [0, 1]$, $t \in [0, 1]$, matrix K , and positive-definite matrices A, B_1, B_2 , the map $B \mapsto \Re \operatorname{tr}(K^\dagger A^s K B^{1-s})$ is concave:

$$t \Re \operatorname{tr}(K^\dagger A^s K B_1^{1-s}) + (1-t) \Re \operatorname{tr}(K^\dagger A^s K B_2^{1-s}) \leq \Re \operatorname{tr}(K^\dagger A^s K (tB_1 + (1-t)B_2)^{1-s}).$$

Proof. [✓ Proof](#) Specialize Theorem 18.5.13 at $A_1 = A_2 = A$, so that the convex combination collapses: $tA + (1-t)A = A$. $\square$

18.6 Resolvent monotonicity toward the Lieb concavity theorem

The integral-representation route toward the Lieb concavity theorem rests on the antitonicity of the resolvent of the commuting left- and right-multiplication superoperators. Its matrix-order foundation is the inverse-antitonicity of the Loewner order.

Lemma 18.6.1 (Loewner inverse-antitonicity). [Lean](#) [✓ Stmt](#) For positive-definite matrices A and B with $A \leq B$ in the Loewner order, the inverses satisfy $B^{-1} \leq A^{-1}$.

Proof. [✓ Proof](#) Compare the two Schur complements of the block matrix $\begin{pmatrix} A^{-1} & \mathbb{1} \\ \mathbb{1} & B \end{pmatrix}$. Complementing its (1, 1) block gives $B - (A^{-1})^{-1} = B - A \geq 0$, so the block matrix is positive semidefinite; complementing the (2, 2) block of the same matrix gives $A^{-1} - B^{-1}$, which is therefore positive semidefinite as well. $\square$

Lemma 18.6.2 (Joint Loewner-antitonicity of the commuting-multiplication resolvent). [Lean](#) [✓ Stmt](#) Fix $t > 0$ and positive-definite matrices with $A_1 \leq A_2$ and $B_1 \leq B_2$. The resolvent of the Kronecker model $A \otimes \mathbb{1} + t(\mathbb{1} \otimes B^\top)$ of the commuting left- and right-multiplication superoperators is antitone:

$$(A_2 \otimes \mathbb{1} + t(\mathbb{1} \otimes B_2^\top))^{-1} \leq (A_1 \otimes \mathbb{1} + t(\mathbb{1} \otimes B_1^\top))^{-1}.$$

Proof.  **Proof** Each operator $A \otimes \mathbb{1} + t(\mathbb{1} \otimes B^\top)$ is positive definite, being the sum of the positive-definite $A \otimes \mathbb{1}$ and the positive-semidefinite $t(\mathbb{1} \otimes B^\top)$. From $A_1 \leq A_2$ and $B_1 \leq B_2$ one gets $A_1 \otimes \mathbb{1} \leq A_2 \otimes \mathbb{1}$ and $\mathbb{1} \otimes B_1^\top \leq \mathbb{1} \otimes B_2^\top$; since $t > 0$, adding these yields

$$A_1 \otimes \mathbb{1} + t(\mathbb{1} \otimes B_1^\top) \leq A_2 \otimes \mathbb{1} + t(\mathbb{1} \otimes B_2^\top).$$

Inverse-antitonicity (Lemma [18.6.1](#)) reverses this inequality on passing to the resolvents. □

Chapter 19

Quantum Entropy

This chapter develops the von Neumann entropy and its basic properties for finite-dimensional quantum systems.

19.1 Von Neumann entropy

Definition 19.1.1 (Von Neumann entropy). [Lean](#) [✓ Stmt](#) For a Hermitian matrix $\rho \in M_D(\mathbb{C})$ with eigenvalues $\lambda_0, \dots, \lambda_{D-1}$, the *von Neumann entropy* is

$$S(\rho) = - \sum_{i=0}^{D-1} \lambda_i \log \lambda_i,$$

where $0 \log 0 := 0$.

Lemma 19.1.2 (Entropy respects equality). [Lean](#) [✓ Stmt](#) If two Hermitian matrices are equal, then their von Neumann entropies are equal.

Proof. [✓ Proof](#) Substitute the equality of the matrices in the defining eigenvalue sum. □

Lemma 19.1.3 (Entropy of the zero matrix). [Lean](#) [✓ Stmt](#) The zero matrix has zero von Neumann entropy:

$$S(0) = 0.$$

Proof. [✓ Proof](#) All eigenvalues of the zero matrix are zero, and $0 \log 0 = 0$. □

Theorem 19.1.4 (Entropy is non-negative). [Lean](#) [✓ Stmt](#) For any density matrix ρ , $S(\rho) \geq 0$.

Proof. [✓ Proof](#) Each eigenvalue λ_i of a density matrix satisfies $0 \leq \lambda_i \leq 1$, and $-x \log x \geq 0$ on $[0, 1]$. □

Lemma 19.1.5 (Eigenvalue sum). [Lean](#) [✓ Stmt](#) The eigenvalues of a density matrix sum to 1.

Proof. [✓ Proof](#) Follows from $\text{tr}(\rho) = \sum_i \lambda_i = 1$. □

Theorem 19.1.6 (Eigenvalue bound). [Lean](#) [✓ Stmt](#) Each eigenvalue of a density matrix lies in $[0, 1]$.

Proof. [✓ Proof](#) Non-negativity comes from positive semidefiniteness. The upper bound follows because the eigenvalues are non-negative and sum to 1. □

Theorem 19.1.7 (Entropy upper bound). [Lean](#) [✓ Stmt](#) For a density matrix $\rho \in M_D(\mathbb{C})$ with $D \geq 1$,

$$S(\rho) \leq \log D.$$

Proof. [✓ Proof](#) By Jensen's inequality applied to the concave function $-x \log x$, the entropy is maximized when all eigenvalues equal $1/D$. $\square$

Theorem 19.1.8 (Entropy bounded by log of rank). [Lean](#) [✓ Stmt](#) For a density matrix ρ of rank r ,

$$S(\rho) \leq \log r.$$

This refines the dimension bound: only the nonzero eigenvalues contribute to the entropy, and there are exactly r of them.

Proof. [✓ Proof](#) The entropy is the $-x \log x$ sum over all eigenvalues; the $D - r$ zero eigenvalues contribute nothing. Jensen's inequality applied to $-x \log x$ over the r nonzero eigenvalues, with uniform weights $1/r$, gives $S(\rho) \leq \log r$, the maximum attained when each nonzero eigenvalue equals $1/r$. The rank equals the number of nonzero eigenvalues of the Hermitian matrix ρ . $\square$

Lemma 19.1.9 (Entropy from characteristic polynomial roots). [Lean](#) [✓ Stmt](#) The von Neumann entropy of a Hermitian matrix is the $-x \log x$ sum over the real parts of the roots of its characteristic polynomial χ_ρ :

$$S(\rho) = \sum_{\lambda \in \text{roots}(\chi_\rho)} (-\text{Re}(\lambda) \log \text{Re}(\lambda)).$$

Proof. [✓ Proof](#) The eigenvalues of a Hermitian matrix are exactly the roots of its characteristic polynomial, counted with multiplicity, so the eigenvalue sum defining $S(\rho)$ equals the displayed sum over roots. $\square$

Theorem 19.1.10 (Trace-logarithm form of the entropy). [Lean](#) [✓ Stmt](#) Let ρ be a Hermitian matrix and let $\log \rho$ be its logarithm defined through the functional calculus. Then

$$S(\rho) = -\text{Re}(\text{tr}(\rho \log \rho)).$$

Proof. [✓ Proof](#) A Hermitian matrix and its logarithm are simultaneously diagonalized by a unitary U with $\rho = U \text{diag}(\lambda_i) U^*$, so

$$\text{tr}(\rho \log \rho) = \text{tr}(U \text{diag}(\lambda_i \log \lambda_i) U^*) = \sum_i \lambda_i \log \lambda_i.$$

Its negative is $\sum_i (-\lambda_i \log \lambda_i) = S(\rho)$. Zero eigenvalues contribute nothing under the convention $0 \log 0 = 0$, so no full-support assumption is required. $\square$

Remark 19.1.11. The logarithm is the totalized real logarithm, with $\log x = \log|x|$ and $\log 0 = 0$; both sides of the identity use it on every eigenvalue, so the equality holds for an arbitrary Hermitian matrix. It coincides with the physical entropy $-\text{tr}(\rho \log \rho)$ precisely when ρ is positive semidefinite. In that case (a density matrix ρ , positive semidefinite with unit trace) the eigenvalues λ_i are nonnegative, $\rho \log \rho$ is Hermitian with real trace, and the real-part extraction is superfluous: the identity reduces to the standard expression $S(\rho) = -\text{tr}(\rho \log \rho)$.

Definition 19.1.12 (Quantum relative entropy). [Lean](#) [✓ Stmt](#) For matrices $\rho, \sigma \in M_D(\mathbb{C})$, define the trace-log expression

$$D(\rho\|\sigma) = \operatorname{Re} \operatorname{tr}(\rho(\log \rho - \log \sigma)).$$

On the physical domain where ρ is a density matrix and σ is positive definite, this is the Umegaki relative entropy.

Lemma 19.1.13 (Trace-log splitting for relative entropy). [Lean](#) [✓ Stmt](#) For matrices $\rho, \sigma \in M_D(\mathbb{C})$,

$$D(\rho\|\sigma) = \operatorname{Re} \operatorname{tr}(\rho \log \rho) - \operatorname{Re} \operatorname{tr}(\rho \log \sigma).$$

Proof. [✓ Proof](#) Expand the matrix product over the difference $\log \rho - \log \sigma$ and use linearity of the trace and of the real part. $\square$

Lemma 19.1.14 (Relative entropy with identical arguments). [Lean](#) [✓ Stmt](#) For every matrix $\rho \in M_D(\mathbb{C})$,

$$D(\rho\|\rho) = 0.$$

Proof. [✓ Proof](#) The logarithmic difference $\log \rho - \log \rho$ vanishes. $\square$

Lemma 19.1.15 (Relative entropy with zero first argument). [Lean](#) [✓ Stmt](#) For every matrix $\sigma \in M_D(\mathbb{C})$,

$$D(0\|\sigma) = 0.$$

Proof. [✓ Proof](#) The trace-log expression is multiplied on the left by the zero matrix. $\square$

Theorem 19.1.16 (Relative entropy in entropy form). [Lean](#) [✓ Stmt](#) If ρ is Hermitian, then

$$D(\rho\|\sigma) = -S(\rho) - \operatorname{Re} \operatorname{tr}(\rho \log \sigma).$$

Proof. [✓ Proof](#) Apply Lemma 19.1.13 to write

$$D(\rho\|\sigma) = \operatorname{Re} \operatorname{tr}(\rho \log \rho) - \operatorname{Re} \operatorname{tr}(\rho \log \sigma).$$

The trace-logarithm identity

$$\operatorname{Re} \operatorname{tr}(\rho \log \rho) = -S(\rho)$$

then gives the result. $\square$

Theorem 19.1.17 (Klein's inequality). [Lean](#) [✓ Stmt](#) Let $\rho, \sigma \in M_D(\mathbb{C})$ be density matrices with σ of full rank. Then the relative entropy is nonnegative,

$$D(\rho\|\sigma) \geq 0.$$

Proof. [✓ Proof](#) Diagonalize $\rho = \sum_i p_i |e_i\rangle\langle e_i|$ and $\sigma = \sum_j q_j |f_j\rangle\langle f_j|$ in their eigenbases, with all $q_j > 0$ since σ has full rank. The overlap numbers $P_{ij} = |\langle e_i | f_j \rangle|^2$ are nonnegative with row sums and column sums equal to 1, because the two eigenbases are orthonormal. A trace computation gives

$$\operatorname{Re} \operatorname{tr}(\rho \log \rho) = \sum_i p_i \log p_i, \quad \operatorname{Re} \operatorname{tr}(\rho \log \sigma) = \sum_{i,j} p_i P_{ij} \log q_j,$$

so that

$$D(\rho\|\sigma) = \sum_i p_i \log p_i - \sum_{i,j} p_i P_{ij} \log q_j.$$

The row sums let the first term be written over the index pair, and the scalar tangent bound $\log x \leq x - 1$ applied to q_j/p_i gives, after weighting by $p_i P_{ij}$ and summing, the lower bound $\sum_{i,j} P_{ij} (q_j - p_i)$. By the column sums and the trace-one normalizations of p and q this telescopes to $\sum_j q_j - \sum_i p_i = 0$, which is the claim. $\square$

Theorem 19.1.18 (Klein’s inequality, support form). [Lean](#) [✓ Stmt](#) Let $\rho, \sigma \in M_D(\mathbb{C})$ be density matrices satisfying the support condition $\ker \sigma \subseteq \ker \rho$, that is, every vector annihilated by σ is annihilated by ρ . Then the relative entropy is nonnegative,

$$D(\rho\|\sigma) \geq 0.$$

Proof. [✓ Proof](#) Regularize σ by the trace-one perturbation $\sigma'_\varepsilon = (1 + \varepsilon D)^{-1}(\sigma + \varepsilon \mathbf{1})$, which is positive definite for every $\varepsilon > 0$, hence of full rank. The full-rank Klein inequality (Theorem 19.1.17) gives $D(\rho\|\sigma'_\varepsilon) \geq 0$. The perturbation shares the eigenbasis $\{|f_j\rangle\}$ of σ , so the cross term reduces to a scalar sum over the eigenvalues q_j of σ ,

$$\operatorname{Re} \operatorname{tr}(\rho \log \sigma'_\varepsilon) = \sum_j \langle f_j | \rho | f_j \rangle \log((1 + \varepsilon D)^{-1}(q_j + \varepsilon)).$$

For $q_j > 0$ the scalar logarithm converges to $\log q_j$; for $q_j = 0$ the eigenvector $|f_j\rangle$ lies in $\ker \sigma \subseteq \ker \rho$, so its diagonal weight $\langle f_j | \rho | f_j \rangle$ vanishes and the summand is identically zero. No eigenvalue-continuity input is needed, so $D(\rho\|\sigma'_\varepsilon) \rightarrow D(\rho\|\sigma)$ as $\varepsilon \rightarrow 0^+$ and the inequality passes to the limit. $\square$

Theorem 19.1.19 (Equality case of Klein’s inequality). [Lean](#) [✓ Stmt](#) Let $\rho, \sigma \in M_D(\mathbb{C})$ be density matrices with σ of full rank. Then the relative entropy vanishes exactly when the states coincide,

$$D(\rho\|\sigma) = 0 \iff \rho = \sigma.$$

Together with nonnegativity, this is the order property that makes D a divergence.

Proof. [✓ Proof](#) If $\rho = \sigma$ then $\log \rho - \log \sigma = 0$ and $D(\rho\|\sigma) = 0$. For the converse, diagonalize $\rho = \sum_i p_i |e_i\rangle\langle e_i|$ and $\sigma = \sum_j q_j |f_j\rangle\langle f_j|$, with all $q_j > 0$ by full rank, and set $P_{ij} = |\langle e_i | f_j \rangle|^2$. As in Theorem 19.1.17,

$$D(\rho\|\sigma) = \sum_{i,j} P_{ij} [p_i \log p_i - p_i \log q_j - (p_i - q_j)],$$

a sum of nonnegative terms: each is bounded below by the tangent inequality $\log x \leq x - 1$ at $x = q_j/p_i$, while the linear remainder telescopes through the doubly stochastic row and column sums,

$$\sum_{i,j} P_{ij} (p_i - q_j) = \sum_i p_i - \sum_j q_j = 0.$$

If the total vanishes then every term vanishes. On a row with $p_i = 0$ the term reads $P_{ij} q_j$, so $P_{ij} = 0$; on a row with $p_i > 0$ the term forces the tangent inequality to be tight,

$$\log \frac{q_j}{p_i} = \frac{q_j}{p_i} - 1,$$

and strict concavity of the logarithm ($\log x < x - 1$ for $x \neq 1$) gives $q_j = p_i$. Hence $q_j = p_i$ whenever $\langle e_i | f_j \rangle \neq 0$. Writing $W = U_\rho^\dagger U_\sigma$ for the overlap of the eigenvector unitaries, this matching says W intertwines the eigenvalue diagonals,

$$W \operatorname{diag}(q) = \operatorname{diag}(p) W,$$

so conjugating σ into the eigenbasis of ρ and using the unitarity $WW^\dagger = 1$,

$$U_\rho^\dagger \sigma U_\rho = W \operatorname{diag}(q) W^\dagger = \operatorname{diag}(p) WW^\dagger = \operatorname{diag}(p),$$

the spectral diagonal of ρ . Therefore $\sigma = U_\rho \operatorname{diag}(p) U_\rho^\dagger = \rho$. $\square$

Theorem 19.1.20 (Joint convexity of the relative entropy). [Lean](#) [✓ Stmt](#) *On pairs of positive definite matrices in $M_D(\mathbb{C})$, the map $(\rho, \sigma) \mapsto D(\rho\|\sigma)$ is jointly convex.*

Proof. [✓ Proof](#) For $s \in [0, 1)$ consider the approximant

$$g_s(\rho, \sigma) = \frac{\operatorname{tr} \rho - \operatorname{Re} \operatorname{tr}(\rho^s \sigma^{1-s})}{1-s}.$$

The trace $\operatorname{tr} \rho$ is real-affine in the pair, hence convex, while $(\rho, \sigma) \mapsto \operatorname{Re} \operatorname{tr}(\rho^s \sigma^{1-s})$ is jointly concave by the $K = \mathbb{1}$ case of the Lieb concavity theorem (Corollary 18.5.14). Their difference, scaled by the nonnegative factor $(1-s)^{-1}$, is therefore jointly convex, so each g_s is convex.

Writing ρ and σ in their eigenbases with eigenvalues $p_i, q_j > 0$ and overlap weights $P_{ij} = |\langle e_i | f_j \rangle|^2$, the approximant becomes the double sum $g_s(\rho, \sigma) = \sum_{i,j} (1-s)^{-1} (p_i - p_i^s q_j^{1-s}) P_{ij}$. As $s \rightarrow 1^-$ each per-pair term converges, via the scalar limit $(c^u - 1)/u \rightarrow \log c$, to $p_i (\log p_i - \log q_j) P_{ij}$, whose sum equals $D(\rho\|\sigma)$. Thus $g_s \rightarrow D$ pointwise on positive definite pairs, and since the pointwise limit of convex functions is convex, D is jointly convex. $\square$

Theorem 19.1.21 (Joint convexity on the support domain). [Lean](#) [✓ Stmt](#) *On pairs of density matrices (ρ, σ) in $M_D(\mathbb{C})$ satisfying the support condition $\ker \sigma \subseteq \ker \rho$, the map $(\rho, \sigma) \mapsto D(\rho\|\sigma)$ is jointly convex.*

Proof. [✓ Proof](#) The domain is convex: for a strict convex combination of two such pairs the kernel of $a\sigma_1 + b\sigma_2$ is $\ker \sigma_1 \cap \ker \sigma_2$, since the quadratic forms of the positive semidefinite summands are nonnegative and their positively weighted sum vanishes only when each does, and a positive semidefinite matrix annihilates exactly the vectors of zero quadratic form; the two pointwise support inclusions then give $\ker \sigma_1 \cap \ker \sigma_2 \subseteq \ker \rho_1 \cap \ker \rho_2$.

Regularize both arguments through the affine trace-one perturbation $M_\varepsilon = (1 + \varepsilon N)^{-1}(M + \varepsilon \mathbb{1})$, where N is the matrix dimension, which is positive definite for every $\varepsilon > 0$. Because the perturbation is affine, it commutes with the convex combination, so the four regularized endpoints and the regularized mixture all lie in the positive definite domain and the positive definite joint convexity (Theorem 19.1.20) gives the two-point inequality for the regularized pairs. As $\varepsilon \rightarrow 0^+$ the regularized relative entropy converges on each pair of the support domain,

$$D(\rho_\varepsilon\|\sigma_\varepsilon) \rightarrow D(\rho\|\sigma).$$

Indeed the perturbation shares the eigenbasis of its argument, so each trace-logarithm term is the diagonal sum $\sum_j w_j(\varepsilon) \log((1 + \varepsilon N)^{-1}(q_j + \varepsilon))$, where q_j runs over the eigenvalues of σ and $w_j(\varepsilon)$ is the corresponding diagonal weight. For a positive eigenvalue the scalar factor has the limit

$$\log((1 + \varepsilon N)^{-1}(q_j + \varepsilon)) \rightarrow \log q_j \quad (q_j > 0),$$

while at a zero eigenvalue the support condition makes the weight vanish, with $w_j(\varepsilon) = (1 + \varepsilon N)^{-1}\varepsilon$, so the summand is $(1 + \varepsilon N)^{-1}\varepsilon \log((1 + \varepsilon N)^{-1}\varepsilon) \rightarrow 0$ by $x \log x \rightarrow 0$ as $x \rightarrow 0^+$. The two-point inequality therefore passes to the limit, giving joint convexity on the support domain. $\square$

Lemma 19.1.22 (Functional calculus under unitary conjugation). [Lean](#) [✓ Stmt](#) *For a Hermitian matrix A , a real function f , and a unitary U , the continuous functional calculus satisfies*

$$f(UAU^\dagger) = U f(A) U^\dagger.$$

Proof. ✓ Proof The conjugation $\varphi : x \mapsto UxU^\dagger$ is a star-algebra automorphism of the matrix algebra, and the continuous functional calculus commutes with such automorphisms, $\varphi(f(A)) = f(\varphi(A))$; since $\varphi(A) = UAU^\dagger$, this is the claim. $\square$

Lemma 19.1.23 (Matrix logarithm under unitary conjugation). Lean ✓ Stmt For a Hermitian matrix A and a unitary U ,

$$\log(UAU^\dagger) = U (\log A) U^\dagger.$$

Proof. ✓ Proof This is the $f = \log$ case of Lemma 19.1.22: with $f(UAU^\dagger) = U f(A) U^\dagger$ specialized to the real logarithm,

$$\log(UAU^\dagger) = U (\log A) U^\dagger.$$

$\square$

Theorem 19.1.24 (Unitary invariance of the quantum relative entropy). Lean ✓ Stmt For Hermitian matrices ρ, σ and a unitary U ,

$$D(U\rho U^\dagger \| U\sigma U^\dagger) = D(\rho \| \sigma).$$

Proof. ✓ Proof Carry the two logarithms through the conjugation by Lemma 19.1.23, so that

$$D(U\rho U^\dagger \| U\sigma U^\dagger) = \operatorname{Re} \operatorname{tr}(U\rho(\log \rho - \log \sigma)U^\dagger) = \operatorname{Re} \operatorname{tr}(\rho(\log \rho - \log \sigma)) = D(\rho \| \sigma),$$

where the middle equality is trace cyclicity together with $U^\dagger U = 1$. $\square$

Lemma 19.1.25 (Logarithm of a tensor product of positive definite matrices). Lean ✓ Stmt For positive definite matrices ρ and τ ,

$$\log(\rho \otimes \tau) = \log \rho \otimes \mathbf{1} + \mathbf{1} \otimes \log \tau.$$

Proof. ✓ Proof The tensor product factors as $\rho \otimes \tau = (\rho \otimes \mathbf{1})(\mathbf{1} \otimes \tau)$ into a pair of commuting positive definite matrices, so the logarithm of the product is the sum of the logarithms of the factors. Each unital embedding $A \mapsto A \otimes \mathbf{1}$ and $B \mapsto \mathbf{1} \otimes B$ is a continuous star-algebra homomorphism, hence commutes with the functional calculus, which carries each logarithm onto its factor: $\log(\rho \otimes \mathbf{1}) = \log \rho \otimes \mathbf{1}$ and $\log(\mathbf{1} \otimes \tau) = \mathbf{1} \otimes \log \tau$. $\square$

Theorem 19.1.26 (Ancilla additivity of the quantum relative entropy). Lean ✓ Stmt For positive definite matrices ρ, σ and a positive definite matrix τ of unit trace,

$$D(\rho \otimes \tau \| \sigma \otimes \tau) = D(\rho \| \sigma).$$

Proof. ✓ Proof Splitting each tensor logarithm by Lemma 19.1.25, the common $\mathbf{1} \otimes \log \tau$ terms cancel in the difference, leaving $\log(\rho \otimes \tau) - \log(\sigma \otimes \tau) = (\log \rho - \log \sigma) \otimes \mathbf{1}$. Hence

$$D(\rho \otimes \tau \| \sigma \otimes \tau) = \operatorname{Re} \operatorname{tr}((\rho(\log \rho - \log \sigma)) \otimes \tau) = \operatorname{Re}(\operatorname{tr}(\rho(\log \rho - \log \sigma)) \cdot \operatorname{tr} \tau),$$

where the trace of a tensor product factors as a product of traces. As $\operatorname{tr} \tau = 1$, the right-hand side is $D(\rho \| \sigma)$. $\square$

Lemma 19.1.27 (Root-of-unity character-sum orthogonality). Lean ✓ Stmt Let ζ be a primitive d -th root of unity and let i, j range over the residues modulo d . Then

$$\sum_{b=0}^{d-1} \zeta^{bi} \overline{\zeta^{bj}} = \begin{cases} d & i = j, \\ 0 & i \neq j. \end{cases}$$

Proof. [✓ Proof](#) Each summand equals ξ^b with $\xi = \zeta^i \bar{\zeta}^j$, and $\xi^d = 1$. When $i = j$ the base ξ equals 1 and the sum is d . When $i \neq j$ the base is a root of unity different from 1, so the geometric sum $(\xi - 1) \sum_b \xi^b = \xi^d - 1 = 0$ forces the sum to vanish; the equivalence $\xi = 1 \iff i = j$ uses that $\bar{\zeta} = \zeta^{-1}$ and the injectivity of $b \mapsto \zeta^b$ on residues. $\square$

Theorem 19.1.28 (Weyl-operator unitary 1-design twirl). [Lean](#) [✓ Stmt](#) Fix a dimension $d \geq 1$ and a primitive d -th root of unity ζ , and let X be the cyclic shift $|i\rangle \mapsto |i+1\rangle$ and $Z = \text{diag}(\zeta^0, \dots, \zeta^{d-1})$ the clock operator. For every matrix M on $\mathbb{C}^d$, the uniform average of the conjugations by the d^2 Weyl operators $W(a, b) = X^a Z^b$ is the completely depolarizing channel:

$$\frac{1}{d^2} \sum_{a,b=0}^{d-1} W(a, b) M W(a, b)^\dagger = \frac{\text{tr } M}{d} \mathbf{1}.$$

Proof. [✓ Proof](#) The double average factors into a clock average followed by a shift average. The clock average $\sum_b Z^b M (Z^b)^\dagger$ multiplies the entry M_{ij} by $\sum_b \zeta^{bi} \bar{\zeta}^{bj}$, which by Lemma 19.1.27 is d when $i = j$ and 0 otherwise; the result is d times the diagonal part of M . The shift average $\sum_a X^a (\text{diagonal } v) (X^a)^\dagger$ cyclically permutes the diagonal entries, so each diagonal position receives the full sum $\sum_k v_k = \text{tr } M$, giving $(\text{tr } M) \mathbf{1}$. Combining the two factors of d with the prefactor d^{-2} leaves $(\text{tr } M/d) \mathbf{1}$. $\square$

Lemma 19.1.29 (Twirl as partial trace tensored with the maximally mixed ancilla). [Lean](#) [✓ Stmt](#) Fix a dimension $d_C \geq 1$ and a primitive d_C -th root of unity ζ . For every matrix M on $\mathcal{H}_S \otimes \mathbb{C}^{d_C}$, the uniform average of the conjugations by the d_C^2 unitaries $\mathbf{1}_S \otimes W(a, b)$ on the second factor is the partial trace over that factor tensored with the maximally mixed state $\mathbf{1}_C/d_C$:

$$\frac{1}{d_C^2} \sum_{a,b} (\mathbf{1}_S \otimes W(a, b)) M (\mathbf{1}_S \otimes W(a, b))^\dagger = (\text{tr}_C M) \otimes (\mathbf{1}_C/d_C).$$

Proof. [✓ Proof](#) On each pair of blocks indexed by the first factor, the conjugation by $\mathbf{1}_S \otimes W(a, b)$ acts as the Weyl conjugation $W(a, b) (\cdot) W(a, b)^\dagger$ of the corresponding block of M . Averaging over the d_C^2 Weyl operators sends each block to the depolarizing channel by Theorem 19.1.28, replacing it by its trace times $\mathbf{1}_C/d_C$. The block trace is exactly the corresponding entry of the partial trace $\text{tr}_C M$, so the average is $(\text{tr}_C M) \otimes (\mathbf{1}_C/d_C)$. $\square$

Lemma 19.1.30 (Reindexing invariance of the quantum relative entropy). [Lean](#) [✓ Stmt](#) For Hermitian matrices ρ, σ on a finite index set and any bijection e from that set onto another finite set,

$$D(\rho_{e^{-1}, e^{-1}} \parallel \sigma_{e^{-1}, e^{-1}}) = D(\rho \parallel \sigma).$$

Proof. [✓ Proof](#) Write $D(\rho \parallel \sigma) = \text{tr}(\rho(\log \rho - \log \sigma))$. The matrix logarithm is covariant under reindexing, $\log(\rho_{e^{-1}, e^{-1}}) = (\log \rho)_{e^{-1}, e^{-1}}$, because reindexing is a star-algebra isomorphism and so commutes with the continuous functional calculus. The same isomorphism preserves products and the trace,

$$(\rho M)_{e^{-1}, e^{-1}} = \rho_{e^{-1}, e^{-1}} M_{e^{-1}, e^{-1}}, \quad \text{tr}(M_{e^{-1}, e^{-1}}) = \text{tr}(M).$$

Taking $M = \log \rho - \log \sigma$ and applying these three identities gives $D(\rho_{e^{-1}, e^{-1}} \parallel \sigma_{e^{-1}, e^{-1}}) = D(\rho \parallel \sigma)$. $\square$

Theorem 19.1.31 (Data-processing inequality under the partial trace). [Lean](#) [✓ Stmt](#) For positive definite matrices ρ, σ on a tensor product of a system factor and an ancilla factor,

$$D(\text{tr}_C \rho \parallel \text{tr}_C \sigma) \leq D(\rho \parallel \sigma),$$

where tr_C is the partial trace over the ancilla factor. This is the positive-definite base case; the source inequality on the support domain $\ker \sigma \subseteq \ker \rho$ is Theorem 19.1.32.

Proof. ✓ Proof By ancilla additivity (Theorem 19.1.26) the reduced-state relative entropy equals $D((\text{tr}_C \rho) \otimes (\mathbf{1}_C/d_C) \| (\text{tr}_C \sigma) \otimes (\mathbf{1}_C/d_C))$. By Lemma 19.1.29 each tensored reduced state is the uniform average of the conjugations by the d_C^2 unitaries $U_{ab} = \mathbf{1}_S \otimes W(a, b)$, so this is the relative entropy of a convex combination of the conjugated pairs $(U_{ab} \rho U_{ab}^\dagger, U_{ab} \sigma U_{ab}^\dagger)$ with equal weights d_C^{-2} . Joint convexity bounds it above by the same convex combination of the per-term relative entropies, each of which equals $D(\rho \| \sigma)$ by unitary invariance (Theorem 19.1.24). As the weights sum to one, the bound is $D(\rho \| \sigma)$. $\square$

Theorem 19.1.32 (Data-processing inequality on the support domain). Lean ✓ Stmt For positive semidefinite ρ, σ on a tensor product of a system factor and an ancilla factor of dimension d_C , with the support condition $\ker \sigma \subseteq \ker \rho$,

$$D(\text{tr}_C \rho \| \text{tr}_C \sigma) \leq D(\rho \| \sigma),$$

where tr_C is the partial trace over the ancilla factor.

Proof. ✓ Proof Let N be the dimension of the joint space. Regularize both arguments through the affine trace-shrinking perturbation $M_\varepsilon = (1 + N\varepsilon)^{-1}(M + \varepsilon \mathbf{1})$, which is positive definite for every $\varepsilon > 0$, so the positive-definite data-processing inequality (Theorem 19.1.31) gives

$$D(\text{tr}_C \rho_\varepsilon \| \text{tr}_C \sigma_\varepsilon) \leq D(\rho_\varepsilon \| \sigma_\varepsilon), \quad \varepsilon > 0.$$

The right-hand side converges to $D(\rho \| \sigma)$ as $\varepsilon \rightarrow 0^+$, by the same shared-eigenbasis scalar-limit argument as the joint convexity on the support domain (Theorem 19.1.21).

For the left-hand side, the partial trace of the regularization is a differently-scaled regularization of the marginal: because the partial trace is linear and $\text{tr}_C \mathbf{1} = d_C \mathbf{1}$,

$$\text{tr}_C((1 + N\varepsilon)^{-1}(M + \varepsilon \mathbf{1})) = (1 + N\varepsilon)^{-1}(\text{tr}_C M + d_C \varepsilon \mathbf{1}),$$

the affine regularization of $\text{tr}_C M$ with the same scaling rate N but shift rate d_C and the smaller identity on the system factor. The support condition transfers to the marginals, $\ker(\text{tr}_C \sigma) \subseteq \ker(\text{tr}_C \rho)$: a vector annihilated by $\text{tr}_C \sigma$ has vanishing marginal quadratic form, which splits into the nonnegative joint quadratic forms of its single-ancilla lifts, so each lift lies in $\ker \sigma$, hence in $\ker \rho$, and reassembling the ancilla sum shows the vector lies in $\ker(\text{tr}_C \rho)$. With this support condition the arbitrary-rate affine regularization has the same both-arguments limit, so

$$D(\text{tr}_C \rho_\varepsilon \| \text{tr}_C \sigma_\varepsilon) \rightarrow D(\text{tr}_C \rho \| \text{tr}_C \sigma), \quad \varepsilon \rightarrow 0^+.$$

Passing the inequality through the two limits gives the support-domain bound. $\square$

Theorem 19.1.33 (Entropy is invariant under reindexing). Lean ✓ Stmt Let ρ be a Hermitian matrix indexed by a finite set J , and let $e : I \rightarrow J$ be a bijection from a finite set I . The reindexed matrix on I with entries $\rho_{e(i)e(j)}$ has the same von Neumann entropy as ρ .

Proof. ✓ Proof By Lemma 19.1.9 the entropy depends only on the characteristic polynomial. Reindexing conjugates ρ by a permutation matrix, so

$$\chi_{(\rho_{e(i)e(j)})} = \chi_\rho,$$

and the entropies coincide. $\square$

Lemma 19.1.34 (Cyclic invariance of the charpoly-root entropy sum). [Lean](#) [✓ Stmt](#) Let $A \in M_{m \times n}(\mathbb{C})$ and $B \in M_{n \times m}(\mathbb{C})$. Then the charpoly-root entropy sum is invariant under the cyclic swap $AB \mapsto BA$:

$$\sum_{\lambda \in \text{roots}(\chi_{AB})} (-\text{Re}(\lambda) \log \text{Re}(\lambda)) = \sum_{\mu \in \text{roots}(\chi_{BA})} (-\text{Re}(\mu) \log \text{Re}(\mu)),$$

with roots counted with algebraic multiplicity.

Proof. [✓ Proof](#) The rectangular characteristic-polynomial identity gives

$$X^n \chi_{AB} = X^m \chi_{BA}.$$

Thus the two characteristic polynomials have the same nonzero roots, with multiplicity; the additional zero roots contribute nothing because $0 \log 0 = 0$. $\square$

Theorem 19.1.35 (Entropy of AB equals entropy of BA). [Lean](#) [✓ Stmt](#) For matrices $A \in M_{m \times n}(\mathbb{C})$ and $B \in M_{n \times m}(\mathbb{C})$ such that AB and BA are Hermitian,

$$S(AB) = S(BA).$$

Proof. [✓ Proof](#) By Lemma 19.1.9, the entropy of a Hermitian matrix is its charpoly-root entropy sum. Lemma 19.1.34 identifies these two sums for AB and BA . $\square$

Theorem 19.1.36 (Entropy is additive over tensor products). [Lean](#) [✓ Stmt](#) For density matrices ω and τ ,

$$S(\omega \otimes \tau) = S(\omega) + S(\tau).$$

Proof. [✓ Proof](#) The tensor product is unitarily conjugate to the diagonal of eigenvalue products $\lambda_i \mu_j$. Using $-\lambda_i \mu_j \log(\lambda_i \mu_j) = \mu_j (-\lambda_i \log \lambda_i) + \lambda_i (-\mu_j \log \mu_j)$ and the unit eigenvalue sums collapses the double sum to $S(\omega) + S(\tau)$. $\square$

Theorem 19.1.37 (Entropy of a scaled density matrix). [Lean](#) [✓ Stmt](#) For a density matrix ω and a scalar c ,

$$S(c\omega) = c S(\omega) - c \log c.$$

Proof. [✓ Proof](#) The eigenvalues of $c\omega$ are $c\lambda_i$, so the entropy is $\sum_i -(c\lambda_i) \log(c\lambda_i)$. The splitting identity

$$-(c\lambda) \log(c\lambda) = \lambda (-c \log c) + c (-\lambda \log \lambda)$$

together with the unit eigenvalue sum $\sum_i \lambda_i = 1$ gives $\sum_i -(c\lambda_i) \log(c\lambda_i) = (-c \log c) \sum_i \lambda_i + c \sum_i (-\lambda_i \log \lambda_i) = -c \log c + c S(\omega)$, the stated form. $\square$

Theorem 19.1.38 (Entropy is additive over an orthogonal direct sum of Hermitian blocks).

[Lean](#) [✓ Stmt](#) For a family of Hermitian matrices M_j , the block-diagonal direct sum satisfies

$$S\left(\bigoplus_j M_j\right) = \sum_j S(M_j).$$

Proof. [✓ Proof](#) Each block diagonalizes by a unitary congruence, and the block-diagonal assembly of the block unitaries diagonalizes the direct sum, whose eigenvalue multiset is the disjoint union of the block eigenvalue multisets. Hence

$$S\left(\bigoplus_j M_j\right) = \sum_{\lambda \in \biguplus_j \text{spec}(M_j)} -\lambda \log \lambda = \sum_j \sum_{\lambda \in \text{spec}(M_j)} -\lambda \log \lambda = \sum_j S(M_j).$$

$\square$

19.2 Tripartite partial traces

Definition 19.2.1 (Partial trace over A). [Lean](#) [✓ Stmt](#) For a tripartite matrix ρ_{ABC} on $\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B} \otimes \mathbb{C}^{d_C}$, the partial trace over A is

$$(\mathrm{tr}_A \rho_{ABC})_{(b_1, c_1)(b_2, c_2)} = \sum_{a=0}^{d_A-1} (\rho_{ABC})_{(a, b_1, c_1)(a, b_2, c_2)}.$$

Definition 19.2.2 (Partial trace over C). [Lean](#) [✓ Stmt](#) The partial trace over C :

$$(\mathrm{tr}_C \rho_{ABC})_{(a_1, b_1)(a_2, b_2)} = \sum_{c=0}^{d_C-1} (\rho_{ABC})_{(a_1, b_1, c)(a_2, b_2, c)}.$$

Definition 19.2.3 (Partial trace over AC). [Lean](#) [✓ Stmt](#) The partial trace over A and C :

$$(\mathrm{tr}_{AC} \rho_{ABC})_{b_1 b_2} = \sum_{a=0}^{d_A-1} \sum_{c=0}^{d_C-1} (\rho_{ABC})_{(a, b_1, c)(a, b_2, c)}.$$

Lemma 19.2.4 (Partial trace preserves Hermiticity). [Lean](#) [✓ Stmt](#) If ρ_{ABC} is Hermitian, then $\mathrm{tr}_A(\rho_{ABC})$, $\mathrm{tr}_C(\rho_{ABC})$, and $\mathrm{tr}_{AC}(\rho_{ABC})$ are all Hermitian. The same holds for bipartite partial traces $\mathrm{tr}_A(\rho_{AB})$ and $\mathrm{tr}_B(\rho_{AB})$.

Proof. [✓ Proof](#) Follows from $\overline{\rho_{ji}} = \rho_{ij}$ applied entry-wise inside the summation defining each partial trace. $\square$

19.3 Strong subadditivity

Lemma 19.3.1 (Relative entropy against a maximally mixed reference). [Lean](#) [✓ Stmt](#) Let ρ be a density matrix on $A \otimes R$ with reduced state $\rho_R = \mathrm{tr}_A \rho$, and suppose the support condition $\ker((\mathbf{1}_A/d_A) \otimes \rho_R) \subseteq \ker \rho$ holds. Then

$$D(\rho \| (\mathbf{1}_A/d_A) \otimes \rho_R) = \log d_A + S(\rho_R) - S(\rho).$$

Proof. [✓ Proof](#) When ρ_R is singular the tensor logarithm of the reference does not split. Regularize the reduced state by the affine perturbation $\rho_{R, \varepsilon} = (1 + d_R \varepsilon)^{-1}(\rho_R + \varepsilon \mathbf{1})$, positive definite for $\varepsilon > 0$. The reference $(\mathbf{1}_A/d_A) \otimes \rho_{R, \varepsilon}$ is then a positive definite tensor product, whose logarithm splits, so the cross trace term evaluates to $-\log d_A + \mathrm{Re} \mathrm{tr}(\rho_R \log \rho_{R, \varepsilon})$. As $\varepsilon \rightarrow 0^+$ the support condition makes the zero eigenvalues of ρ_R contribute nothing, so

$$\mathrm{Re} \mathrm{tr}(\rho \log((\mathbf{1}_A/d_A) \otimes \rho_{R, \varepsilon})) \rightarrow -\log d_A - S(\rho_R).$$

Since $D(\rho \| \sigma) = -S(\rho) - \mathrm{Re} \mathrm{tr}(\rho \log \sigma)$, the evaluation follows. $\square$

Lemma 19.3.2 (The maximally mixed reference lies in the support domain). [Lean](#) [✓ Stmt](#) Let ρ be a positive semidefinite operator on $A \otimes R$ with reduced state $\rho_R = \mathrm{tr}_A \rho$. Then the singular reference $(\mathbf{1}_A/d_A) \otimes \rho_R$ satisfies the support condition $\ker((\mathbf{1}_A/d_A) \otimes \rho_R) \subseteq \ker \rho$.

Proof. [✓ Proof](#) A vector v annihilated by $(\mathbf{1}_A/d_A) \otimes \rho_R$ is annihilated by $\mathbf{1}_A \otimes \rho_R$, since $\mathbf{1}_A/d_A$ is invertible. Let P be the orthogonal projection onto the range of ρ_R . The complementary lift $\mathbf{1}_A \otimes (\mathbf{1} - P)$ then fixes v , while it annihilates ρ on the left because the reduced state of ρ on R is supported on the range of P . Hence $\rho v = 0$. $\square$

Theorem 19.3.3 (Strong subadditivity). [Lean](#) [✓ Stmt](#) For a tripartite density matrix ρ_{ABC} ,

$$S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC}).$$

Proof. [✓ Proof](#) Read the inequality as one instance of data processing under the partial trace over C , with the singular reference state $\sigma_{ABC} = (\mathbf{1}_A/d_A) \otimes \rho_{BC}$. Being a density operator is not enough to place the pair $(\rho_{ABC}, \sigma_{ABC})$ in the relative-entropy domain, which is the kernel inclusion $\ker \sigma_{ABC} \subseteq \ker \rho_{ABC}$; the marginal support lemma supplies it. Against this reference the relative entropy of each pair evaluates to an entropy difference,

$$D(\rho_{ABC} \| (\mathbf{1}_A/d_A) \otimes \rho_{BC}) = \log d_A + S(\rho_{BC}) - S(\rho_{ABC}),$$

$$D(\rho_{AB} \| (\mathbf{1}_A/d_A) \otimes \rho_B) = \log d_A + S(\rho_B) - S(\rho_{AB}).$$

For a singular reference the tensor logarithm does not split, so each evaluation regularizes the reduced state through the affine perturbation $\rho_{R,\varepsilon} = (1 + d_R\varepsilon)^{-1}(\rho_R + \varepsilon\mathbf{1})$, which is positive definite, and passes to the limit

$$\text{Retr}(\rho \log((\mathbf{1}_A/d_A) \otimes \rho_{R,\varepsilon})) \rightarrow -\log d_A - S(\rho_R) \quad (\varepsilon \rightarrow 0^+),$$

where the zero eigenvalues of ρ_R contribute nothing because the kernel inclusion makes the corresponding diagonal weights vanish. Data processing on the singular support domain under the partial trace over C , which sends $\rho_{ABC} \mapsto \rho_{AB}$ and $(\mathbf{1}_A/d_A) \otimes \rho_{BC} \mapsto (\mathbf{1}_A/d_A) \otimes \rho_B$, reads

$$D(\rho_{AB} \| (\mathbf{1}_A/d_A) \otimes \rho_B) \leq D(\rho_{ABC} \| (\mathbf{1}_A/d_A) \otimes \rho_{BC}).$$

Substituting the two evaluations cancels the common $\log d_A$ and rearranges to the claimed inequality. $\square$

Theorem 19.3.4 (Strong subadditivity, positive definite case). [Lean](#) [✓ Stmt](#) For a positive definite tripartite density matrix ρ_{ABC} ,

$$S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC}).$$

This is the positive definite case of Theorem 19.3.3.

Proof. [✓ Proof](#) Read the inequality as one instance of data processing under the partial trace over C , with reference state $\sigma_{ABC} = (\mathbf{1}_A/d_A) \otimes \rho_{BC}$, which is positive definite because ρ_{BC} is. Against this reference the relative entropy of each pair evaluates to an entropy difference,

$$D(\rho_{ABC} \| (\mathbf{1}_A/d_A) \otimes \rho_{BC}) = \log d_A + S(\rho_{BC}) - S(\rho_{ABC}),$$

$$D(\rho_{AB} \| (\mathbf{1}_A/d_A) \otimes \rho_B) = \log d_A + S(\rho_B) - S(\rho_{AB}),$$

each by the tensor logarithm split and the adjoint of the partial trace. Data processing under the partial trace over C , which sends $\rho_{ABC} \mapsto \rho_{AB}$ and $(\mathbf{1}_A/d_A) \otimes \rho_{BC} \mapsto (\mathbf{1}_A/d_A) \otimes \rho_B$, reads

$$D(\rho_{AB} \| (\mathbf{1}_A/d_A) \otimes \rho_B) \leq D(\rho_{ABC} \| (\mathbf{1}_A/d_A) \otimes \rho_{BC}).$$

Substituting the two evaluations cancels the common $\log d_A$ and rearranges to the claimed inequality. $\square$

Definition 19.3.5 (SSA equality). [Lean](#) [✓ Stmt](#) A tripartite density matrix ρ_{ABC} satisfies *SSA equality* if

$$S(\rho_{ABC}) + S(\rho_B) = S(\rho_{AB}) + S(\rho_{BC}).$$

Theorem 19.3.6 (SSA equality under factor relabeling). [Lean](#) [✓ Stmt](#) Let ρ be a Hermitian operator on $A \otimes B \otimes C$, and let $e_A : A' \rightarrow A$, $e_B : B' \rightarrow B$, and $e_C : C' \rightarrow C$ be bijections. Define ρ' by

$$\rho'_{(a',b',c'),(\tilde{a}',\tilde{b}',\tilde{c}')} = \rho_{(e_A(a'),e_B(b'),e_C(c')),(e_A(\tilde{a}'),e_B(\tilde{b}'),e_C(\tilde{c}'))}.$$

If ρ satisfies equality in strong subadditivity, then so does ρ' .

Proof. [✓ Proof](#) The four operators entering the equality are related by the induced bijections:

$$\rho' = \rho_{e_A \times e_B \times e_C}, \quad \rho'_{AB} = (\rho_{AB})_{e_A \times e_B}, \quad \rho'_{BC} = (\rho_{BC})_{e_B \times e_C}, \quad \rho'_B = (\rho_B)_{e_B}.$$

These identities follow by changing variables in the three finite sums defining the partial traces. Entropy invariance under reindexing makes the corresponding four entropy terms equal, so the strong-subadditivity equality for ρ gives the equality for ρ' . $\square$

Definition 19.3.7 (Quantum Markov decomposition). [Lean](#) [✓ Stmt](#) A Hayashi Markov decomposition of a tripartite state ρ_{ABC} consists of a finite direct-sum decomposition

$$H_B \cong \bigoplus_j H_{B_j^L} \otimes H_{B_j^R},$$

together with a unitary change of basis on B , a probability vector $(p_j)_j$, and density matrices $\rho_{AB_j^L}$ and $\rho_{B_j^R C}$ such that, in the adapted basis, the state becomes

$$\bigoplus_j p_j \rho_{AB_j^L} \otimes \rho_{B_j^R C}.$$

The terminology follows Hayashi's presentation of quantum Markov structure [Hay06]; the block decomposition used by the MPDO argument is the structure theorem of [HJPW04].

Theorem 19.3.8 (Hayashi SSA-equality characterization). [Lean](#) [✓ Stmt](#) For a tripartite density matrix ρ_{ABC} ,

$$S(\rho_{ABC}) + S(\rho_B) = S(\rho_{AB}) + S(\rho_{BC})$$

holds if and only if ρ_{ABC} admits a quantum Markov decomposition on the middle subsystem B .

The reverse implication, that a quantum Markov decomposition forces the equality, is proved as Theorem 19.3.9. The forward implication, that the equality forces the block-diagonal structure, is the deep part: it rests on recovery-map and Petz theory not yet available, so it is the only part that remains axiomatic. The biconditional combines the two.

This equality criterion is cited here as an input for the later MPDO arguments. It is the equality criterion needed by Appendix C of arXiv:1606.00608. The equality conditions are reviewed in [Hay06, Rus02]; the structural block-decomposition formulation used for MPDOs appears in [HJPW04].

Theorem 19.3.9 (Quantum Markov structure attains SSA equality). [Lean](#) [✓ Stmt](#) A tripartite density matrix ρ_{ABC} that admits a quantum Markov decomposition on the middle subsystem B satisfies

$$S(\rho_{ABC}) + S(\rho_B) = S(\rho_{AB}) + S(\rho_{BC}).$$

Proof. [✓ Proof](#) Write the state in the adapted basis as the block-diagonal direct sum $\bigoplus_j p_j \rho_{AB_j^L} \otimes \rho_{B_j^R C}$. The von Neumann entropy of a weighted orthogonal direct sum is $S(\bigoplus_j p_j \omega_j) = -\sum_j p_j \log p_j + \sum_j p_j S(\omega_j)$ by Theorems 19.1.38 and 19.1.37, and the entropy of a tensor product is additive,

$S(\omega \otimes \tau) = S(\omega) + S(\tau)$, by Theorem 19.1.36. Tracing out one tensor factor within each block, the three reduced states factor as block-diagonal direct sums,

$$\rho_{AB} \cong \bigoplus_j p_j \rho_{AB_j^L} \otimes \rho_{B_j^R}, \quad \rho_{BC} \cong \bigoplus_j p_j \rho_{B_j^L} \otimes \rho_{B_j^R C}, \quad \rho_B \cong \bigoplus_j p_j \rho_{B_j^L} \otimes \rho_{B_j^R},$$

with $\rho_{ABC} \cong \bigoplus_j p_j \rho_{AB_j^L} \otimes \rho_{B_j^R C}$ itself. Writing $H = -\sum_j p_j \log p_j$, the four entropies expand as

$$\begin{aligned} S(\rho_{ABC}) &= H + \sum_j p_j [S(\rho_{AB_j^L}) + S(\rho_{B_j^R C})], & S(\rho_{AB}) &= H + \sum_j p_j [S(\rho_{AB_j^L}) + S(\rho_{B_j^R})], \\ S(\rho_B) &= H + \sum_j p_j [S(\rho_{B_j^L}) + S(\rho_{B_j^R})], & S(\rho_{BC}) &= H + \sum_j p_j [S(\rho_{B_j^L}) + S(\rho_{B_j^R C})]. \end{aligned}$$

Both sides of the claimed identity equal $2H + \sum_j p_j [S(\rho_{AB_j^L}) + S(\rho_{B_j^R C}) + S(\rho_{B_j^L}) + S(\rho_{B_j^R})]$, so $S(\rho_{ABC}) + S(\rho_B) = S(\rho_{AB}) + S(\rho_{BC})$. The basis change on B and the direct-sum reindexing leave every entropy term unchanged. $\square$

19.4 Mutual information

Definition 19.4.1 (Mutual information). [Lean](#) [✓ Stmt](#) The *quantum mutual information* of a bipartite state ρ_{AB} is

$$I(A:B) = S(\rho_A) + S(\rho_B) - S(\rho_{AB}).$$

Theorem 19.4.2 (Mutual information is non-negative). [Lean](#) [✓ Stmt](#) For any bipartite density matrix ρ_{AB} , $I(A:B) \geq 0$.

Proof. [✓ Proof](#) Apply strong subadditivity (Theorem 19.3.3) with trivial B (one-dimensional middle system). The SSA inequality $S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC})$ reduces to subadditivity $S(\rho_{AC}) \leq S(\rho_A) + S(\rho_C)$, which gives $I(A:C) \geq 0$. $\square$

19.5 Entropy formulations

This section states the entropy formulations used later in the development: von Neumann entropy, strong subadditivity, quantum Markov decomposition, and mutual information. These statements are cited from the standard entropy literature and supply the entropy-theoretic input for the later MPDO arguments.

Definition 19.5.1 (Entropy formulation of von Neumann entropy). [Lean](#) [✓ Stmt](#) This formulation has the same value $S(\rho) = -\sum_i \lambda_i \log \lambda_i$ as in Definition 19.1.1.

Theorem 19.5.2 (Entropy formulation of strong subadditivity). [Lean](#) [✓ Stmt](#) For any tripartite density matrix ρ_{ABC} on $A \otimes B \otimes C$,

$$S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC}).$$

This formulation introduces no new axiom: it is the same strong-subadditivity statement as Theorem 19.3.3, which is proved there from Lieb concavity along the relative-entropy route [LR73].

Definition 19.5.3 (Entropy formulation of quantum Markov decomposition). [Lean](#) [✓ Stmt](#) This is the entropy formulation of the Hayashi Markov decomposition from Definition 19.3.7.

Theorem 19.5.4 (SSA equality iff quantum Markov decomposition). [Lean](#) [✓ Stmt](#) For any tripartite density matrix ρ_{ABC} , equality in strong subadditivity holds if and only if ρ_{ABC} admits a quantum Markov decomposition on the middle subsystem B .

This formulation introduces no new axiom: it is the same equality criterion as Theorem 19.3.8.

Definition 19.5.5 (Entropy formulation of mutual information). [Lean](#) [✓ Stmt](#) This formulation has the same value $I(A:B) = S(\rho_A) + S(\rho_B) - S(\rho_{AB})$ as in Definition 19.4.1.

Theorem 19.5.6 (Subadditivity from trivial-middle SSA). [Lean](#) [✓ Stmt](#) For a tripartite density matrix ρ_{ABC} with $\dim B = 1$,

$$S(\rho_{ABC}) \leq S(\rho_{AB}) + S(\rho_{BC}).$$

Proof. [✓ Proof](#) Apply Theorem 19.5.2 with one-dimensional middle subsystem. The reduced middle state has trace 1 by Theorem 19.7.3, hence its entropy vanishes by Theorem 19.7.4. $\square$

19.6 Mutual information: monotonicity and area-law bound

The two inequalities below are the downstream MPDO-facing consequences of the entropy inequalities in this chapter. The monotonicity inequality is the strong-subadditivity content underlying the MPDO mutual-information monotonicity $I_L \leq I_{L+1}$ (arXiv:1606.00608, Proposition C.1); the elementary area-law bound is the single-site entropy bound underlying the MPDO area-law bound $I_L \leq 4 \log D$ (arXiv:1606.00608, cited from the Wolf area-law bound). Both inequalities follow directly from strong subadditivity (Theorem 19.3.3) and the single-system $S(\rho) \leq \log D$ bound; neither introduces a new axiom.

Theorem 19.6.1 (Entropy nonnegativity for finite index sets). [Lean](#) [✓ Stmt](#) For any PSD Hermitian matrix ρ with $\text{tr}(\rho) = 1$ on an arbitrary finite index set, $S(\rho) \geq 0$. This is the analog of Theorem 19.1.4 for arbitrary finite index sets: it does not require the index set to be $\{0, \dots, D-1\}$, so it applies to bipartite density matrices on $\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B}$.

Proof. [✓ Proof](#) The eigenvalues of a PSD matrix are non-negative, and eigenvalues of a trace-1 Hermitian matrix with non-negative eigenvalues are bounded above by 1 (each single eigenvalue is at most the total sum). Apply $x \log x \leq 0$ on $[0, 1]$ pointwise and sum. $\square$

Theorem 19.6.2 (Mutual information is monotone under enlargement). [Lean](#) [✓ Stmt](#) For any tripartite density matrix ρ_{ABC} on $A \otimes B \otimes C$,

$$I(A:B)_{\text{tr}_C \rho_{ABC}} \leq I(A:BC)_{\rho_{ABC}},$$

where the left-hand side is the bipartite mutual information of the reduced state $\rho_{AB} = \text{tr}_C(\rho_{ABC})$ and the right-hand side is evaluated by expanding $I(A:BC) = S(\rho_A) + S(\rho_{BC}) - S(\rho_{ABC})$.

Proof. [✓ Proof](#) Expanding both sides in entropy form and cancelling the common $S(\rho_A)$ term, the inequality reduces to strong subadditivity $S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC})$. The bipartite B -reduced state of ρ_{AB} matches the tripartite B -reduced state $\text{tr}_{AC}(\rho_{ABC})$, ensuring the $S(\rho_B)$ term is the one supplied by Theorem 19.5.2. $\square$

Theorem 19.6.3 (Elementary area-law bound on mutual information). [Lean](#) [✓ Stmt](#) For a bipartite density matrix ρ_{AB} on $\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B}$ with $d_A, d_B \geq 1$ whose single-system reduced states are obtained by partial trace,

$$I(A:B) \leq \log d_A + \log d_B.$$

Proof. [✓ Proof](#) The reduced states are density matrices because partial trace preserves positivity and trace. Combine $S(\rho_A) \leq \log d_A$ and $S(\rho_B) \leq \log d_B$ (Theorem 19.1.7) with $S(\rho_{AB}) \geq 0$ (Theorem 19.6.1). $\square$

19.7 Trivial-factor corollaries

The partial traces and von Neumann entropy are introduced in Chapter 19. This supplement collects elementary trace-preservation identities and the dimension-1 entropy bound that follow directly from those definitions. They are listed as separate results because each proof requires only unfolding definitions and re-indexing finite sums.

Theorem 19.7.1 (Partial trace over A preserves the full trace). [Lean](#) [✓ Stmt](#) *For any tripartite matrix ρ_{ABC} ,*

$$\mathrm{tr}(\rho_{ABC}) = \mathrm{tr}(\mathrm{tr}_A(\rho_{ABC})).$$

Proof. [✓ Proof](#) Unfolding the definitions,

$$\mathrm{tr}(\mathrm{tr}_A(\rho_{ABC})) = \sum_{b,c} \sum_a (\rho_{ABC})_{(a,b,c)(a,b,c)} = \mathrm{tr}(\rho_{ABC}).$$

□

Theorem 19.7.2 (Partial trace over C preserves the full trace). [Lean](#) [✓ Stmt](#) *For any tripartite matrix ρ_{ABC} ,*

$$\mathrm{tr}(\rho_{ABC}) = \mathrm{tr}(\mathrm{tr}_C(\rho_{ABC})).$$

Proof. [✓ Proof](#) Unfolding the definitions,

$$\mathrm{tr}(\mathrm{tr}_C(\rho_{ABC})) = \sum_{a,b} \sum_c (\rho_{ABC})_{(a,b,c)(a,b,c)} = \mathrm{tr}(\rho_{ABC}).$$

□

Theorem 19.7.3 (Partial trace over AC preserves the full trace). [Lean](#) [✓ Stmt](#) *For any tripartite matrix ρ_{ABC} ,*

$$\mathrm{tr}(\rho_{ABC}) = \mathrm{tr}(\mathrm{tr}_{AC}(\rho_{ABC})).$$

Proof. [✓ Proof](#) Unfolding the definitions and interchanging the outer sums,

$$\mathrm{tr}(\mathrm{tr}_{AC}(\rho_{ABC})) = \sum_b \sum_{a,c} (\rho_{ABC})_{(a,b,c)(a,b,c)} = \mathrm{tr}(\rho_{ABC}).$$

□

Theorem 19.7.4 (Entropy vanishes in dimension 1). [Lean](#) [✓ Stmt](#) *If $\rho \in M_1(\mathbb{C})$ is Hermitian and $\mathrm{tr}(\rho) = 1$, then*

$$S(\rho) = 0.$$

Proof. [✓ Proof](#) The unique eigenvalue equals the trace, hence equals 1, and $-1 \cdot \log 1 = 0$:

$$\lambda_0 = \mathrm{tr}(\rho) = 1, \quad S(\rho) = -\lambda_0 \log \lambda_0 = -1 \cdot \log 1 = 0.$$

□

Chapter 20

Positive but Not Completely Positive Maps

Wolf, *Quantum Channels & Operations*, Chapter 3 studies positive maps that fail to be completely positive and their role in entanglement detection. This material sits outside the fundamental-theorem chain of the preceding chapters: the reduction criterion, the partial transpose and the positive-partial-transpose (PPT) property, separable states, and the Schmidt number with the full reduction criterion. The threshold n -positivity of the family T_η and the two-positivity content these results use live in the Kadison–Schwarz chapter (Chapter 5).

20.1 The reduction criterion

The reduction map and the family T_η are the same map under two parametrizations: the natural-number parameter k and the real parameter η agree at $\eta = k$. Identifying them lets the positivity threshold and the Choi formula carry over to the reduction map without restating them. Wolf attaches to this map the entanglement witness $W_n = D^{-1}\mathbb{1} - n^{-1}|\Omega\rangle\langle\Omega|$ and the operator inequality [Wol12, Equation (3.18)] that constrains the reduced densities of a bipartite state whose Schmidt number is at most n .

Theorem 20.1.1 (The reduction map is the family T_η at integer parameter). Lean ✓ Stmt For every $k \in \mathbb{N}$ the reduction map T_k equals the family map T_η at $\eta = k$:

$$T_k = T_\eta|_{\eta=k}.$$

Proof. ✓ Proof The two maps differ only in the scalar coefficient of X , namely k^{-1} against η^{-1} at $\eta = k$. The natural-number and real casts of k into the complex scalars coincide, so the coefficients agree and the maps are equal. □

Theorem 20.1.2 (The reduction map T_k is k -positive). Lean ✓ Stmt For $1 \leq k \leq D$ the reduction map T_k on $M_D(\mathbb{C})$ is k -positive.

Proof. ✓ Proof The reduction map equals the family map T_η at $\eta = k$, and T_η is k -positive exactly when $\eta \geq k$; the threshold is met with equality at $\eta = k$. For $k < D$ this is the Schmidt-rank threshold, and the top index $k = D$ is the completely positive endpoint supplied by the dimension-indexed threshold. □

Theorem 20.1.3 (The reduction map T_k is not $(k+1)$ -positive). [Lean](#) [✓ Stmt](#) For $1 \leq k < D$ the reduction map T_k on $M_D(\mathbb{C})$ is not $(k+1)$ -positive.

Proof. [✓ Proof](#) The reduction map equals T_η at $\eta = k$. For $k+1 < D$ the threshold gives that T_η is $(k+1)$ -positive iff $\eta \geq k+1$, so T_k would require $k \geq k+1$, which is false. For $k+1 = D$ the dimension-indexed threshold gives that T_η is D -positive iff $\eta \geq D$, so T_k would require $k \geq D$, contradicting $k < D$. $\square$

Remark 20.1.4. Together with the k -positivity of T_k (20.1.2), this shows that T_k is *exactly* k -positive for $1 \leq k < D$: every inclusion in Wolf's chain $T_{\text{cp}} = T_D \subsetneq \cdots \subsetneq T_1$ is strict.

Definition 20.1.5 (Reduction witness). [Lean](#) [✓ Stmt](#) The reduction witness on $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$, for $n \in \mathbb{N}$, is

$$W_n = D^{-1}\mathbb{1} - n^{-1}|\Omega\rangle\langle\Omega|.$$

Theorem 20.1.6 (The Choi matrix of the reduction map is the witness). [Lean](#) [✓ Stmt](#) For $D \geq 1$ the Choi matrix of T_n is the reduction witness:

$$\tau(T_n) = W_n = D^{-1}\mathbb{1} - n^{-1}|\Omega\rangle\langle\Omega|.$$

Proof. [✓ Proof](#) The reduction map is T_η at $\eta = n$, whose Choi matrix is $D^{-1}\mathbb{1} - n^{-1}|\Omega\rangle\langle\Omega|$, which is exactly W_n . $\square$

Theorem 20.1.7 (Reduction-criterion identity). [Lean](#) [✓ Stmt](#) Applying T_η to the first factor of a bipartite matrix ρ gives

$$(T_\eta \otimes \text{id})(\rho) = \mathbb{1} \otimes \rho_2 - \eta^{-1}\rho,$$

where ρ_2 is the reduced density of ρ on the second factor.

Proof. [✓ Proof](#) Fix block indices i_2, j_2 on the second factor and write $\rho^{(i_2, j_2)}$ for the $D \times D$ slice $\rho_{i_1 j_1}^{(i_2, j_2)} = \rho_{(i_1, i_2), (j_1, j_2)}$ on the first factor. Its trace is the corresponding entry of ρ_2 :

$$\text{tr}(\rho^{(i_2, j_2)}) = \sum_{i_1} \rho_{(i_1, i_2), (i_1, j_2)} = (\rho_2)_{i_2 j_2}.$$

Applying T_η entry by entry gives

$$((T_\eta \otimes \text{id})(\rho))_{(i_1, i_2), (j_1, j_2)} = \text{tr}(\rho^{(i_2, j_2)}) \delta_{i_1 j_1} - \eta^{-1} \rho_{(i_1, i_2), (j_1, j_2)} = (\rho_2)_{i_2 j_2} \delta_{i_1 j_1} - \eta^{-1} \rho_{(i_1, i_2), (j_1, j_2)},$$

and the first term is exactly the $(i_1, i_2), (j_1, j_2)$ entry of $\mathbb{1} \otimes \rho_2$. $\square$

Theorem 20.1.8 (Operator-implication step of the reduction criterion). [Lean](#) [Lean](#) [✓ Stmt](#) Let $n \geq 1$ and let ρ be a bipartite matrix, with ρ^F its image under the factor swap $\rho_{(i_2, i_1), (j_2, j_1)}^F = \rho_{(i_1, i_2), (j_1, j_2)}$. If $(T_n \otimes \text{id})(\rho) \geq 0$ then

$$n \mathbb{1} \otimes \rho_2 \geq \rho,$$

and if $(T_n \otimes \text{id})(\rho^F) \geq 0$ then

$$n \rho_1 \otimes \mathbb{1} \geq \rho,$$

where ρ_1 and ρ_2 are the reduced densities of ρ on the first and second factors. Because the factor swap is a unitary reindexing,

$$(\text{id} \otimes T_n)(\rho) \geq 0 \iff (T_n \otimes \text{id})(\rho^F) \geq 0,$$

so the second hypothesis is Wolf's symmetric condition $(\text{id} \otimes T_n)(\rho) \geq 0$ expressed through the swap.

This is the second step of Wolf's two-step argument for [Wol12, Equation (3.18)]. The source states the inequalities under the premise that ρ has Schmidt number at most n , and reaches $(T_n \otimes \text{id})(\rho) \geq 0$ from that premise by the n -positivity of T_n ; that first step is taken here as the hypothesis rather than derived. The first step and the composed full criterion with the Schmidt-number premise are Theorem 20.7.11.

Proof. ✓ Proof By the reduction-criterion identity, $(T_n \otimes \text{id})(\rho)$ equals $\mathbb{1} \otimes \rho_2 - n^{-1}\rho$. Scaling a positive semidefinite operator by $n > 0$ preserves positivity, so $n\mathbb{1} \otimes \rho_2 - \rho \geq 0$, which is the first inequality. For the second, the factor swap F satisfies $F^2 = \text{id}$ and is a unitary reindexing, so it preserves positive semidefiniteness and the order, and it exchanges the reduced densities and the two tensor slots:

$$(\rho^F)_2 = \rho_1, \quad (\mathbb{1} \otimes (\rho^F)_2 - n^{-1}\rho^F)^F = \rho_1 \otimes \mathbb{1} - n^{-1}\rho.$$

Applying the first form to ρ^F gives $n\mathbb{1} \otimes (\rho^F)_2 - \rho^F \geq 0$; conjugating by F turns this into $n\rho_1 \otimes \mathbb{1} - \rho \geq 0$. □

20.2 The Breuer–Hall map

A second concrete positive map of [Wol12, Chapter 3, Example 3.1] is built from an antisymmetric contraction U on $\mathbb{C}^D$, a matrix with $U^\top = -U$ and $U^\dagger U \leq \mathbb{1}$. The associated Breuer–Hall map [Wol12, Equation (3.19)] is

$$T_{\text{BH}}(X) = \text{tr}(X)\mathbb{1} - X - U X^\top U^\dagger.$$

It is positive; whether it is decomposable, and the threshold beyond which it fails to be completely positive, are not treated here.

Definition 20.2.1 (Breuer–Hall map). Lean ✓ Stmt For a matrix U on $M_D(\mathbb{C})$, the Breuer–Hall map on $M_D(\mathbb{C})$ is

$$T_{\text{BH}}(X) = \text{tr}(X)\mathbb{1} - X - U X^\top U^\dagger.$$

Lemma 20.2.2 (The Breuer–Hall map on rank-one operators). Lean ✓ Stmt If U is antisymmetric, $U^\top = -U$, and a contraction, $U^\dagger U \leq \mathbb{1}$, then for every vector v the image $T_{\text{BH}}(|v\rangle\langle v|)$ of the rank-one operator $|v\rangle\langle v|$ is positive semidefinite.

Proof. ✓ Proof On $|v\rangle\langle v|$ the image is

$$T_{\text{BH}}(|v\rangle\langle v|) = \|v\|^2 \mathbb{1} - |v\rangle\langle v| - |U\bar{v}\rangle\langle U\bar{v}|,$$

with $\bar{v}$ the entrywise conjugate of v . The two vectors v and $U\bar{v}$ are orthogonal: antisymmetry of U gives

$$\langle v|U\bar{v}\rangle = \bar{v}^\top U \bar{v} = \bar{v}^\top (-U^\top) \bar{v} = -\langle v|U\bar{v}\rangle,$$

hence $\langle v|U\bar{v}\rangle = 0$. The contraction bound gives $\|U\bar{v}\| \leq \|v\|$. For any w the quadratic form is

$$\|v\|^2 \|w\|^2 - |\langle v|w\rangle|^2 - |\langle U\bar{v}|w\rangle|^2,$$

which is nonnegative because the two-vector Bessel inequality for the orthogonal pair $v, U\bar{v}$ with $\|U\bar{v}\| \leq \|v\|$ gives

$$|\langle v|w\rangle|^2 + |\langle U\bar{v}|w\rangle|^2 \leq \|v\|^2 \|w\|^2.$$

□

Theorem 20.2.3 (The Breuer–Hall map is positive). [Lean](#) [✓ Stmt](#) If U is antisymmetric, $U^\top = -U$, and a contraction, $U^\dagger U \leq \mathbb{1}$, then the Breuer–Hall map T_{BH} sends every positive semidefinite matrix to a positive semidefinite matrix.

Proof. [✓ Proof](#) A positive semidefinite matrix is a sum of rank-one operators $|v\rangle\langle v|$. By Lemma 20.2.2 the Breuer–Hall map sends each such operator to a positive semidefinite matrix, and a sum of positive semidefinite matrices is positive semidefinite, so T_{BH} is positive. $\square$

20.3 Choi-type maps

The Choi-type maps in Wolf’s Example 3.1 are built from the diagonal projection $D(X)$ and cyclic shifts U_{k0} :

$$T_C(X) = (d - n)D(X) - X + \sum_{k=1}^n D(U_{k0} X U_{k0}^\dagger).$$

The formal statements below record the map and the rank-one reduction used in the standard positivity argument. Positivity is proved for the case $d = 3$, $n = 1$ and, for every $d \geq 3$, at the top of the range, $n = d - 2$; the scalar input for the latter is a reciprocal inequality valid for an arbitrary permutation in place of the cyclic shift. The positivity assertion for the remaining range $1 \leq n \leq d - 3$ and the indecomposability assertion remain open. The missing general positivity step is a cyclic reciprocal estimate for the weights in the rank-one reduction; it does not follow merely from equality of the total numerator and denominator weights. The general estimate is classical: the case $n = d - 2$ is due to Ando, the case $n = 1$ to Tanahashi and Tomiyama [TT88], and the whole range to Yamagami [Yam93], whose proof is a variational argument on the boundary of a projective simplex.

Definition 20.3.1 (Choi-type map). [Lean](#) [✓ Stmt](#) On the cyclic index set $\mathbb{Z}/d\mathbb{Z}$, the Choi-type map is

$$T_C(X) = (d - n)D(X) - X + \sum_{k=1}^n D(U_{k0} X U_{k0}^\dagger),$$

where D keeps the diagonal of a matrix and U_{k0} is the cyclic shift by k .

Lemma 20.3.2 (The Choi-type map on rank-one projectors). [Lean](#) [✓ Stmt](#) For a vector v on $\mathbb{Z}/d\mathbb{Z}$,

$$T_C(|v\rangle\langle v|) = \text{diag} \left((d - n)|v_i|^2 + \sum_{k=1}^n |v_{i-k}|^2 \right) - |v\rangle\langle v|.$$

Proof. [✓ Proof](#) The diagonal projection of a shifted rank-one projector records the shifted squared moduli:

$$D(U_{k0}|v\rangle\langle v|U_{k0}^\dagger)_{ii} = |v_{i-k}|^2.$$

Substituting this into the definition of T_C gives the displayed diagonal matrix minus the original rank-one projector. $\square$

Definition 20.3.3 (The Choi rank-one diagonal weight). [Lean](#) [✓ Stmt](#) For a vector v on $\mathbb{Z}/d\mathbb{Z}$, set

$$a_i = (d - n)|v_i|^2 + \sum_{k=1}^n |v_{i-k}|^2.$$

Lemma 20.3.4 (Rank-one complement from a unit bound). [Lean](#) [✓ Stmt](#) If p is a vector with

$$\sum_i |p_i|^2 \leq 1,$$

then

$$I - |p\rangle\langle p| \geq 0.$$

Proof. [✓ Proof](#) For every vector w ,

$$\langle w|(I - |p\rangle\langle p|)|w\rangle = \|w\|^2 - |\langle p|w\rangle|^2.$$

Since $\|p\|^2 = \sum_i |p_i|^2 \leq 1$, Cauchy's inequality gives

$$|\langle p|w\rangle|^2 \leq \|p\|^2 \|w\|^2 \leq \|w\|^2,$$

so the quadratic form is nonnegative. □

Lemma 20.3.5 (Diagonal rank-one Schur-complement criterion). [Lean](#) [✓ Stmt](#) Let $a_i \geq 0$ and suppose $v_i = 0$ whenever $a_i = 0$. If, with the convention that $|v_i|^2/a_i$ is read as 0 when $a_i = 0$,

$$\sum_i \frac{|v_i|^2}{a_i} \leq 1,$$

then

$$\text{diag}(a_i) - |v\rangle\langle v| \geq 0.$$

Proof. [✓ Proof](#) Put $p_i = 0$ when $a_i = 0$ and $p_i = v_i/\sqrt{a_i}$ otherwise. Then

$$\|p\|^2 = \sum_i |p_i|^2 = \sum_i \frac{|v_i|^2}{a_i} \leq 1$$

under the same zero-term convention. Lemma 20.3.4 gives $I - |p\rangle\langle p| \geq 0$. If $S = \text{diag}(\sqrt{a_i})$, then the support condition gives

$$S(I - |p\rangle\langle p|)S^\dagger = \text{diag}(a_i) - |v\rangle\langle v|.$$

Conjugation by S preserves positive semidefiniteness, which proves the claim. □

Lemma 20.3.6 (Rank-one positivity from the cyclic reciprocal bound). [Lean](#) [✓ Stmt](#) Assume $n \leq d - 2$. If, with the convention that the summand is 0 at indices where $a_i = 0$,

$$\sum_i \frac{|v_i|^2}{a_i} \leq 1,$$

where a_i is the weight of Definition 20.3.3, then

$$T_C(|v\rangle\langle v|) \geq 0.$$

Proof. [✓ Proof](#) The matrix in Lemma 20.3.2 has the form $\text{diag}(a_i) - |v\rangle\langle v|$. Each a_i is nonnegative. Since $n \leq d - 2$, the coefficient $d - n$ is positive; hence $a_i = 0$ forces $v_i = 0$. The hypotheses of Lemma 20.3.5 therefore apply and give the stated positivity. □

Lemma 20.3.7 (The first Choi rank-one positivity subcase). [Lean](#) [✓ Stmt](#) For $d = 3$ and $n = 1$, every vector $v = (v_0, v_1, v_2)$ satisfies

$$T_C(|v\rangle\langle v|) \geq 0.$$

Equivalently, with $x = |v_0|^2$, $y = |v_1|^2$, and $z = |v_2|^2$, the needed cyclic estimate is

$$\frac{x}{2x+z} + \frac{y}{2y+x} + \frac{z}{2z+y} \leq 1.$$

The zero-denominator terms are read as 0, as in the preceding Schur-complement criterion. This is only the 3×3 rank-one subcase; the cyclic estimate for all $1 \leq n \leq d-2$ remains open.

Proof. [✓ Proof](#) First assume that the denominators are nonzero. After multiplying by their product, the inequality is equivalent to

$$x^2y + y^2z + z^2x \geq 3xyz.$$

This follows from AM–GM applied to the three nonnegative numbers x^2y , y^2z , and z^2x . If one of the denominators vanishes, the corresponding two variables vanish. For example, if $2x+z=0$, then $x=z=0$, and the remaining term is

$$\frac{y}{2y} \leq \frac{1}{2} \leq 1$$

when $y > 0$, while all terms vanish when $y = 0$. The rank-one positivity then follows from Lemma 20.3.6. $\square$

Lemma 20.3.8 (Positivity from rank-one positivity). [Lean](#) [✓ Stmt](#) A linear map on matrices that sends every rank-one projector $|w\rangle\langle w|$ to a positive semidefinite matrix sends every positive semidefinite matrix to a positive semidefinite matrix.

Proof. [✓ Proof](#) Let Φ be the map and let $X \geq 0$. Scaling the eigenvectors of X by the square roots of its nonnegative eigenvalues produces vectors w_α with

$$X = \sum_{\alpha} |w_{\alpha}\rangle\langle w_{\alpha}|.$$

Linearity gives

$$\Phi(X) = \sum_{\alpha} \Phi(|w_{\alpha}\rangle\langle w_{\alpha}|),$$

each summand is positive semidefinite by hypothesis, and a sum of positive semidefinite matrices is positive semidefinite. $\square$

Theorem 20.3.9 (The first Choi map is positive). [Lean](#) [✓ Stmt](#) For $d = 3$ and $n = 1$, the Choi-type map T_C sends every positive semidefinite matrix to a positive semidefinite matrix.

Proof. [✓ Proof](#) Combine Lemma 20.3.7 with Lemma 20.3.8. $\square$

Lemma 20.3.10 (Permutation reciprocal inequality). [Lean](#) [✓ Stmt](#) Let $x_i \geq 0$ be indexed by a finite set, let $T = \sum_j x_j$, and let σ be a permutation of the index set. Then, with zero-denominator summands read as 0,

$$\sum_i \frac{x_i}{T + x_i - x_{\sigma(i)}} \leq 1.$$

Proof. ✓ Proof If $T = 0$, every summand vanishes. Otherwise write $\delta_i = x_i - x_{\sigma(i)}$. Each summand obeys

$$\frac{x_i}{T + \delta_i} \leq \frac{x_i T - x_i \delta_i + \delta_i^2/2}{T^2}.$$

for $\sigma(i) = i$ both sides equal x_i/T , and for $\sigma(i) \neq i$ the difference of the two sides is

$$\frac{\delta_i^2 (T - x_i - x_{\sigma(i)})}{2T^2 (T + \delta_i)} \geq 0,$$

by the pair bound $x_i + x_{\sigma(i)} \leq T$ and $T + \delta_i \geq 2x_i \geq 0$; a vanishing denominator forces $x_i = 0$ and leaves a nonnegative right-hand side. Because σ permutes the entries, $\sum_i x_{\sigma(i)}^2 = \sum_i x_i^2$, which gives the identity $\sum_i x_i \delta_i = \frac{1}{2} \sum_i \delta_i^2$. Summing the term bounds therefore yields

$$\sum_i \frac{x_i}{T + \delta_i} \leq \frac{T \cdot T - \sum_i x_i \delta_i + \frac{1}{2} \sum_i \delta_i^2}{T^2} = 1.$$

□

Lemma 20.3.11 (The Choi weight at the top of the range). Lean Lean ✓ Stmt Let $d \geq 3$ and $n = d - 2$. With $x_i = |v_i|^2$ and $T = \sum_j x_j$, the Choi rank-one diagonal weight equals

$$a_i = (d - n)x_i + \sum_{k=1}^n x_{i-k} = T + x_i - x_{i+1}.$$

Proof. ✓ Proof For $n = d - 2$ the backward shifts $i - 1, \dots, i - (d - 2)$ run over every index of $\mathbb{Z}/d\mathbb{Z}$ except i and $i + 1$, so the shifted sum equals $T - x_i - x_{i+1}$ and $a_i = 2x_i + T - x_i - x_{i+1}$. □

Lemma 20.3.12 (Rank-one positivity at the top of the range). Lean ✓ Stmt For $d \geq 3$ and $n = d - 2$, every vector v satisfies $T_C(|v\rangle\langle v|) \geq 0$.

Proof. ✓ Proof By Lemma 20.3.11, the reciprocal weight sum is

$$\sum_i \frac{x_i}{T + x_i - x_{i+1}}$$

with $x_i = |v_i|^2$. Lemma 20.3.10, applied to the cyclic shift $\sigma(i) = i + 1$, bounds this sum by 1, and Lemma 20.3.6 turns the bound into rank-one positivity. □

Theorem 20.3.13 (Choi-type maps at the top of the range are positive). Lean ✓ Stmt For every $d \geq 3$ and $n = d - 2$, the Choi-type map T_C sends every positive semidefinite matrix to a positive semidefinite matrix. This is the case $n = d - 2$ of the positivity assertion of Wolf's Example 3.1 and subsumes the case $d = 3, n = 1$.

Proof. ✓ Proof Combine Lemma 20.3.12 with Lemma 20.3.8. □

20.4 Decomposable positive maps

The Choi-type maps are stated in Wolf's Example 3.1 as positive and indecomposable. The formal language for the second adjective is the following standard one: a positive map is decomposable when it is the sum of a completely positive map and a completely copositive map.

Definition 20.4.1 (Decomposable and indecomposable positive maps). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#)
A map $\Phi: M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is completely copositive if $\Phi \circ \theta$ is completely positive, where θ is transposition. It is decomposable if

$$\Phi = \Phi_{\text{cp}} + \Phi_{\text{ccp}}$$

with Φ_{cp} completely positive and Φ_{ccp} completely copositive. It is indecomposable if it is positive and not decomposable.

Lemma 20.4.2 (Completely copositive maps are positive). [Lean](#) [✓ Stmt](#) *If Φ is completely copositive, then Φ is positive.*

Proof. [✓ Proof](#) Let $X \geq 0$. Transposition preserves positivity, so $\theta(X) \geq 0$. Since $\Phi \circ \theta$ is completely positive,

$$(\Phi \circ \theta)(\theta(X)) \geq 0.$$

Because $\theta^2 = \text{id}$, this is exactly $\Phi(X) \geq 0$. □

Lemma 20.4.3 (Decomposable maps are positive). [Lean](#) [✓ Stmt](#) *If Φ is decomposable, then Φ is positive.*

Proof. [✓ Proof](#) Write

$$\Phi = \Phi_{\text{cp}} + \Phi_{\text{ccp}},$$

with Φ_{cp} completely positive and Φ_{ccp} completely copositive. For $X \geq 0$, the two preceding positivity results give

$$\Phi_{\text{cp}}(X) \geq 0, \quad \Phi_{\text{ccp}}(X) \geq 0.$$

Their sum is positive, hence $\Phi(X) \geq 0$. □

20.5 The partial transpose and the PPT property

Transposition is a positive map that is not completely positive, so applying it to one factor of a bipartite operator can break positivity. The resulting operation, the partial transpose [Wol12, Equation (3.16)], sends $\rho \in M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$ to the operator ρ^{T_1} obtained by transposing the first tensor factor while leaving the second untouched, and the property $\rho^{T_1} \geq 0$ — positive partial transpose, or PPT — is the criterion [Wol12, Equation (3.17)] that separates many entangled states from the separable ones.

Definition 20.5.1 (Partial transpose over the first factor). [Lean](#) [✓ Stmt](#) For a bipartite matrix ρ on $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$, the partial transpose over the first factor is the matrix ρ^{T_1} with entries

$$(\rho^{T_1})_{(i,k),(j,l)} = \rho_{(j,k),(i,l)},$$

transposing the first index pair i, j and fixing the second pair k, l .

Definition 20.5.2 (Partial transpose over the second factor). [Lean](#) [✓ Stmt](#) For a bipartite matrix ρ on $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$, the partial transpose over the second factor is the matrix ρ^{T_2} with entries

$$(\rho^{T_2})_{(i,k),(j,l)} = \rho_{(i,l),(j,k)},$$

transposing the second index pair k, l and fixing the first pair i, j .

Theorem 20.5.3 (The partial transpose is an involution). [Lean](#) [✓ Stmt](#) *Applying the first-factor partial transpose twice returns the original matrix: $(\rho^{T_1})^{T_1} = \rho$.*

Proof. ✓ Proof Both entries equal $\rho_{(i,k),(j,l)}$: the first transposition exchanges the first index pair, the second restores it. $\square$

Theorem 20.5.4 (The partial transpose preserves the trace). Lean ✓ Stmt *The first-factor partial transpose preserves the trace: $\text{tr}(\rho^{T_1}) = \text{tr}(\rho)$.*

Proof. ✓ Proof On the diagonal the row and column indices coincide, so the index exchange in the definition is the identity and each diagonal entry is unchanged. $\square$

Theorem 20.5.5 (The partial transpose preserves Hermiticity). Lean ✓ Stmt *If ρ is Hermitian then so is its first-factor partial transpose ρ^{T_1} .*

Proof. ✓ Proof The entry $(\rho^{T_1})_{(i,k),(j,l)} = \rho_{(j,k),(i,l)}$ has complex conjugate $\overline{\rho_{(j,k),(i,l)}} = \rho_{(i,l),(j,k)}$ by Hermiticity of ρ , which is the $((j,l), (i,k))$ entry of ρ^{T_1} . $\square$

Theorem 20.5.6 (The two partial transposes differ by a global transposition). Lean ✓ Stmt *The second-factor partial transpose is the full transpose of the first-factor one:*

$$\rho^{T_2} = (\rho^{T_1})^T.$$

Proof. ✓ Proof Transposing both factors is the full transpose, so transposing the first factor and then the whole operator transposes only the second factor. $\square$

Definition 20.5.7 (Positive partial transpose property). Lean ✓ Stmt A bipartite matrix ρ has the positive-partial-transpose (PPT) property when its first-factor partial transpose is positive semidefinite, $\rho^{T_1} \geq 0$.

Theorem 20.5.8 (The PPT property is symmetric in the two factors). Lean ✓ Stmt *A bipartite matrix has the PPT property if and only if its second-factor partial transpose is positive semidefinite:*

$$\rho^{T_1} \geq 0 \iff \rho^{T_2} \geq 0.$$

Proof. ✓ Proof The two partial transposes differ by a global transposition, and transposition preserves positive semidefiniteness, so one is positive exactly when the other is. $\square$

Theorem 20.5.9 (Basis-change law for the partial transpose). Lean ✓ Stmt *For matrices U, V on the first factor and a bipartite matrix ρ , conjugating inside by V and U , taking the first-factor partial transpose, and conjugating outside by U and V produces the original partial transpose conjugated by UU^T and $V^T V$:*

$$(U \otimes \mathbb{1}) \left[((V \otimes \mathbb{1}) \rho (U \otimes \mathbb{1}))^{T_1} \right] (V \otimes \mathbb{1}) = ((UU^T) \otimes \mathbb{1}) \rho^{T_1} ((V^T V) \otimes \mathbb{1}).$$

Proof. ✓ Proof Expanding the conjugated operator entry by entry as a double sum over the first factor, the partial transpose exchanges the two summation indices and reverses the roles of the left and right conjugating matrices, leaving them transposed; collecting the Kronecker factors gives the stated form. $\square$

Theorem 20.5.10 (Conjugation invariance of the partial transpose). Lean ✓ Stmt *Let U be any matrix on the first factor. If ρ has the PPT property then*

$$(U \otimes \mathbb{1}) \left[((U^\dagger \otimes \mathbb{1}) \rho (U \otimes \mathbb{1}))^{T_1} \right] (U^\dagger \otimes \mathbb{1})$$

is positive semidefinite; no invertibility or unitarity of U is required. Specializing to a unitary U this is the partial transpose taken in the basis changed by U , so the positivity of the partial transpose is independent of the local basis, recovering Wolf's basis-independence statement [Wol12, Equation (3.17)].

Proof. ✓ Proof By the basis-change law with $V = U^\dagger$, the new-basis partial transpose equals $((UU^\dagger) \otimes \mathbb{1}) \rho^{T_1} ((UU^\dagger) \otimes \mathbb{1})^\dagger$, a conjugation of ρ^{T_1} that preserves positive semidefiniteness. $\square$

Definition 20.5.11 (Transposition witness). Lean ✓ Stmt For matrices A, B on $M_d(\mathbb{C})$ and the SWAP operator F on $M_d(\mathbb{C}) \otimes M_d(\mathbb{C})$, the transposition witness is

$$W = (A \otimes B) F (A \otimes B)^\dagger.$$

These are the entanglement witnesses associated with the transposition map.

Theorem 20.5.12 (Transposition witnesses are Hermitian). Lean ✓ Stmt Every transposition witness $W = (A \otimes B) F (A \otimes B)^\dagger$ is Hermitian.

Proof. ✓ Proof The SWAP operator is self-adjoint, $F^\dagger = F$, so taking the adjoint of $(A \otimes B) F (A \otimes B)^\dagger$ returns the same operator. $\square$

20.6 Separable states and the PPT criterion

A bipartite state is separable when it is a convex combination of pure product states $\sum_i p_i |a_i\rangle\langle a_i| \otimes |b_i\rangle\langle b_i|$. Each pure product state is a Kronecker product of two rank-one positive semidefinite matrices, the nonnegative weight p_i is absorbed into one factor, and conversely a positive semidefinite matrix is a nonnegative combination of rank-one projectors. Up to normalization this gives the equivalent description of separable states as finite sums of Kronecker products of positive semidefinite matrices, the cone generated by the product positive operators, which is the form used below.

Definition 20.6.1 (Separable bipartite matrix). Lean ✓ Stmt A bipartite matrix ρ on $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$ is separable when it is a finite sum of Kronecker products of positive semidefinite matrices,

$$\rho = \sum_i A_i \otimes B_i, \quad A_i \geq 0, \quad B_i \geq 0.$$

Theorem 20.6.2 (A Kronecker product of positive operators is separable). Lean ✓ Stmt For positive semidefinite A on $M_d(\mathbb{C})$ and B on $M_{d'}(\mathbb{C})$, the Kronecker product $A \otimes B$ is separable.

Proof. ✓ Proof It is the one-term sum $\sum_i A_i \otimes B_i$ with the single index taking the value $A \otimes B$. $\square$

Theorem 20.6.3 (Separable matrices are closed under addition). Lean ✓ Stmt If ρ and σ are separable then so is $\rho + \sigma$.

Proof. ✓ Proof Concatenating the two index families realizes $\rho + \sigma$ as a single sum of Kronecker products of positive semidefinite matrices. $\square$

Theorem 20.6.4 (Separable matrices are closed under nonnegative scaling). Lean ✓ Stmt If ρ is separable and $c \geq 0$ is real then $c\rho$ is separable.

Proof. ✓ Proof Absorbing c into the first factor of each Kronecker product keeps every factor positive semidefinite, since a nonnegative multiple of a positive semidefinite matrix is positive semidefinite. $\square$

Theorem 20.6.5 (A separable matrix is positive semidefinite). [Lean](#) [✓ Stmt](#) *Every separable matrix is positive semidefinite.*

Proof. [✓ Proof](#) Each summand $A_i \otimes B_i$ is a Kronecker product of positive semidefinite matrices, hence positive semidefinite, and a finite sum of positive semidefinite matrices is positive semidefinite. $\square$

Theorem 20.6.6 (The partial transpose distributes over a finite sum). [Lean](#) [✓ Stmt](#) *For a finite family of bipartite matrices,*

$$\left(\sum_i f_i\right)^{T_1} = \sum_i f_i^{T_1}.$$

Proof. [✓ Proof](#) The first-factor partial transpose is an entrywise reindexing of the matrix, so it commutes with the entrywise finite sum. $\square$

Theorem 20.6.7 (PPT criterion, forward direction). [Lean](#) [✓ Stmt](#) *Every separable matrix has the positive-partial-transpose property: if ρ is separable then $\rho^{T_1} \geq 0$. A negative partial transpose therefore certifies that a state is entangled.*

Proof. [✓ Proof](#) The first-factor partial transpose is linear and acts on a Kronecker product by transposing the first factor, $(A \otimes B)^{T_1} = A^T \otimes B$. Writing $\rho = \sum_i A_i \otimes B_i$ with each factor positive semidefinite, the partial transpose is $\sum_i A_i^T \otimes B_i$. The transpose of a positive semidefinite matrix is positive semidefinite, so each summand is a Kronecker product of positive semidefinite matrices, and the sum is positive semidefinite. $\square$

Theorem 20.6.8 (Applying a map to one factor distributes over a finite sum). [Lean](#) [✓ Stmt](#) *For a linear map T on the first factor and a finite family of bipartite matrices,*

$$(T \otimes \text{id})\left(\sum_i f_i\right) = \sum_i (T \otimes \text{id})(f_i).$$

Proof. [✓ Proof](#) Applying a map to the first factor is linear in the bipartite matrix, so it commutes with the finite sum. $\square$

Theorem 20.6.9 (Reduction map at the first level on a separable state). [Lean](#) [✓ Stmt](#) *For $d \geq 1$ and a separable matrix ρ on $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$, applying the reduction map $T_1(X) = \text{tr}(X)\mathbb{1} - X$ to the first factor keeps the result positive semidefinite:*

$$(T_1 \otimes \text{id})(\rho) \geq 0.$$

This is the separability instance of the operator inequality underlying the reduction criterion at the first level.

Proof. [✓ Proof](#) The reduction map T_1 is positive, so it maps each positive semidefinite first factor A_i of $\rho = \sum_i A_i \otimes B_i$ to a positive semidefinite matrix. The ampliation acts by $(T_1 \otimes \text{id})(A_i \otimes B_i) = T_1(A_i) \otimes B_i$, each summand is a Kronecker product of positive semidefinite matrices, and the sum is positive semidefinite. $\square$

20.7 The Schmidt number and the full reduction criterion

The Schmidt number of a bipartite state is the smallest n for which the state has a convex decomposition into pure states each of Schmidt rank at most n . Separable states are precisely those of Schmidt number one, and a state of Schmidt number at most n satisfies the reduction criterion

$$n \rho_1 \otimes \mathbb{1} \geq \rho \quad \text{and} \quad n \mathbb{1} \otimes \rho_2 \geq \rho,$$

which is [Wol12, Equation (3.18)]. The bound on the Schmidt number is recorded as a finite sum of pure-state projectors of Schmidt rank at most n , the unnormalized form of the convex decomposition; rescaling a vector by the square root of its weight scales its coefficient matrix and so leaves the Schmidt rank unchanged.

Definition 20.7.1 (Bounded Schmidt number). [Lean](#) [✓ Stmt](#) A bipartite matrix ρ on $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$ has Schmidt number at most n when it is a finite sum of pure-state projectors of Schmidt rank at most n ,

$$\rho = \sum_i |\psi_i\rangle\langle\psi_i|, \quad \text{SR}(\psi_i) \leq n.$$

Theorem 20.7.2 (Basic properties of bounded Schmidt number). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A matrix of Schmidt number at most n is positive semidefinite. States of Schmidt number at most n are closed under addition and under multiplication by a nonnegative real scalar, and a bound on the Schmidt number relaxes to any larger bound.

Proof. [✓ Proof](#) Each summand $|\psi_i\rangle\langle\psi_i|$ is a rank-one positive semidefinite matrix, and a finite sum of positive semidefinite matrices is positive semidefinite. Concatenating two pure-state families realizes a sum as a single finite sum of pure-state projectors of Schmidt rank at most n , and a pure summand of Schmidt rank at most n also has Schmidt rank at most any larger bound. Multiplication by a nonnegative real a rescales each pure summand ψ_i to $\sqrt{a}\psi_i$, which leaves its Schmidt rank unchanged and reproduces a times the original state. $\square$

Theorem 20.7.3 (Schmidt number one is separability). [Lean](#) [✓ Stmt](#) A bipartite matrix has Schmidt number at most one if and only if it is separable.

Proof. [✓ Proof](#) A pure state of Schmidt rank at most one has a coefficient matrix of rank at most one, which factors as $\psi(i, j) = u_i v_j$; its projector is the Kronecker product $|u\rangle\langle u| \otimes |v\rangle\langle v|$ of two rank-one positive semidefinite matrices, hence separable, and separable matrices are closed under addition. Conversely a Kronecker product $A \otimes B$ of positive semidefinite matrices, after expanding each factor into rank-one projectors, is a double sum of product-vector projectors, each of Schmidt rank at most one, so it has Schmidt number at most one; a separable matrix is a finite sum of such products. $\square$

Theorem 20.7.4 (Convexity of the Schmidt-number set). [Lean](#) [✓ Stmt](#) For each n the set S_n of bipartite states of Schmidt number at most n is convex.

Proof. [✓ Proof](#) A convex combination $ax + by$ with $a, b \geq 0$ is a sum of two nonnegatively scaled states of Schmidt number at most n , and that bound is preserved under nonnegative scaling and under addition; rescaling a vector by a nonnegative scalar scales its coefficient matrix and so does not increase the Schmidt rank. $\square$

Theorem 20.7.5 (Compactness of the Schmidt-number state set). [Lean](#) [✓ Stmt](#) For each n the set S_n of trace-one bipartite states of Schmidt number at most n is compact.

Proof. ✓ Proof A trace-one state of Schmidt number at most n is a convex combination of pure states $|\psi\rangle\langle\psi|$ whose vector ψ has unit norm and Schmidt rank at most n , so S_n is the convex hull of the set P_n of those pure states. The set P_n is the image, under the continuous projector map $\psi \mapsto |\psi\rangle\langle\psi|$, of the unit vectors of Schmidt rank at most n . In the Euclidean norm those unit vectors form the unit sphere, which is compact in finite dimension, intersected with the closed set on which the Schmidt rank is at most n (the coefficient matrix depends continuously on the vector and the matrices of rank at most n form a closed set); hence P_n is compact. The trace of $|\psi\rangle\langle\psi|$ is the squared Euclidean norm of ψ , so the trace-one constraint is exactly the unit-norm constraint and the weights of the convex decomposition are the squared norms, summing to the trace. The convex hull of a compact set is compact in a finite-dimensional real normed space. $\square$

Theorem 20.7.6 (Pure-state step of the reduction criterion). Lean ✓ Stmt For a pure state $|\psi\rangle\langle\psi|$ on the square bipartite system $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$ with $\text{SR}(\psi) \leq n$ and $1 \leq n < D$, applying T_n to the first factor keeps the result positive semidefinite:

$$(T_n \otimes \text{id})(|\psi\rangle\langle\psi|) \geq 0.$$

Proof. ✓ Proof The map T_n is n -positive for $1 \leq n < D$. Write ψ through the maximally entangled vector as $\psi = \psi_X$ for a square matrix X on the right factor with $\text{rank}(X) = \text{SR}(\psi) \leq n$. Then $(T_n \otimes \text{id})(|\psi\rangle\langle\psi|)$ is the right-factor Choi compression of T_n by X . Its quadratic form on any vector η is the Choi quadratic form of T_n on $R_X^\dagger \eta$, whose Schmidt rank is at most $\text{rank}(X) \leq n$. The Schmidt-rank Choi criterion makes that form nonnegative, so the compression is positive semidefinite. $\square$

Theorem 20.7.7 (Pure-state step for a general positive map). Lean ✓ Stmt For a pure state $|\psi\rangle\langle\psi|$ on the square bipartite system $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$ with $\text{SR}(\psi) \leq n$ and any n -positive map T , applying T to the first factor keeps the result positive semidefinite:

$$(T \otimes \text{id})(|\psi\rangle\langle\psi|) \geq 0.$$

Proof. ✓ Proof Write ψ through the maximally entangled vector as $\psi = \psi_X$ for a square matrix X on the right factor with $\text{rank}(X) = \text{SR}(\psi) \leq n$. Then $(T \otimes \text{id})(|\psi\rangle\langle\psi|)$ is the right-factor Choi compression of T by X , and its quadratic form on a vector η equals the Choi quadratic form C_T of T evaluated on $R_X^\dagger \eta$, a vector of Schmidt rank at most $\text{rank}(X) \leq n$. Because T is n -positive,

$$\langle R_X^\dagger \eta, C_T R_X^\dagger \eta \rangle \geq 0,$$

so the compression is positive semidefinite. $\square$

Theorem 20.7.8 (Positive maps and entanglement, only-if direction). Lean ✓ Stmt A bipartite state on the square system $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$ of Schmidt number at most n satisfies

$$(T \otimes \text{id})(\rho) \geq 0$$

for every n -positive map T . This is the square case $d = d' = D$ of the only-if direction of Wolf's Proposition 3.4 for a general bipartite system $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$ [Wol12, Chapter 3, Proposition 3.4].

Proof. ✓ Proof The state is a finite sum of pure-state projectors of Schmidt rank at most n . Applying T to the first factor is linear, so it distributes over the sum; the pure-state step makes each summand positive semidefinite, and a finite sum of positive semidefinite matrices is positive semidefinite. $\square$

Theorem 20.7.9 (Only-if direction for a general bipartite system). [Lean](#) [✓ Stmt](#) *On a general bipartite system $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$, with no relation imposed between the two factors, a state of Schmidt number at most n satisfies*

$$(T \otimes \text{id})(\rho) \geq 0$$

for every n -positive map T on $M_d(\mathbb{C})$. This is the only-if direction of Wolf's Proposition 3.4, the first step of eq. (3.18) [Wol12, Chapter 3, Proposition 3.4], with no restriction on the bipartite dimensions.

Proof. [✓ Proof](#) Pad the bipartite state to the square system $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$ with $D = \max(d, d')$ by zero rows and columns: when $d' \leq d$ the second factor is padded up to d ; when $d \leq d'$ the first factor is padded up to d' and T is extended to a map on $M_{d'}(\mathbb{C})$ that preserves n -positivity. Padding preserves the Schmidt-number bound, the square result makes the padded ampliation positive semidefinite, and the original ampliation is its principal submatrix, hence positive semidefinite. $\square$

Theorem 20.7.10 (Reduction step from the Schmidt-number premise). [Lean](#) [✓ Stmt](#) *For $1 \leq n < D$, a bipartite state on the square system $M_D(\mathbb{C}) \otimes M_D(\mathbb{C})$ of Schmidt number at most n satisfies*

$$(T_n \otimes \text{id})(\rho) \geq 0.$$

Proof. [✓ Proof](#) The state is a finite sum of pure-state projectors of Schmidt rank at most n . Applying T_n to the first factor is linear, so it distributes over the sum; the pure-state step makes each summand positive semidefinite, and a finite sum of positive semidefinite matrices is positive semidefinite. $\square$

Theorem 20.7.11 (Reduction criterion with the Schmidt-number premise). [Lean](#) [Lean](#) [✓ Stmt](#) *On a general bipartite system $M_d(\mathbb{C}) \otimes M_{d'}(\mathbb{C})$, a state of Schmidt number at most n satisfies the reduction criterion*

$$n \mathbb{1} \otimes \rho_2 \geq \rho \quad (1 \leq n < d) \quad \text{and} \quad n \rho_1 \otimes \mathbb{1} \geq \rho \quad (1 \leq n < d'),$$

with ρ_1 and ρ_2 the reduced densities on the first and second factors. No relation is imposed between d and d' ; each inequality carries only the n -positivity threshold of T_n on its acting factor.

Proof. [✓ Proof](#) The Schmidt-number premise gives $(T_n \otimes \text{id})(\rho) \geq 0$ on the general system (first step), and the dimension-general operator implication then yields $n \mathbb{1} \otimes \rho_2 \geq \rho$. The factor swap of a state of Schmidt number at most n again has Schmidt number at most n , since it sends each pure summand ψ to $\psi \circ \text{swap}$, whose Schmidt coefficient matrix is the transpose $M_{\psi \circ \text{swap}} = M_\psi^T$ of that of ψ and so has the same rank; applying the first step to the swapped state (whose first factor is d') and the operator implication in its symmetric form gives $n \rho_1 \otimes \mathbb{1} \geq \rho$. $\square$

Theorem 20.7.12 (Trace-form representation of a real-linear functional). [Lean](#) [✓ Stmt](#) *Every real-linear functional g on the space of square complex matrices is the trace form of a Hermitian matrix: there is a Hermitian H_0 with*

$$g(X) = \text{Re tr}(XH_0) \quad \text{for every Hermitian } X.$$

Proof. [✓ Proof](#) The real bilinear pairing $(X, H) \mapsto \text{Re tr}(XH)$ is nondegenerate, because on the diagonal pair $(H, H^\dagger)$ it equals the squared Frobenius norm $\text{Re tr}(HH^\dagger) = \sum_{i,j} |H_{ij}|^2$, which is positive unless $H = 0$. Hence the map sending H to the functional $X \mapsto \text{Re tr}(XH)$ is an injective real-linear map from the matrix space to its real dual. These two spaces have the same finite real dimension, so the map is also surjective, and every g is represented by some H . Replacing H by its Hermitian part $\frac{1}{2}(H + H^\dagger)$ keeps the value on Hermitian inputs, since $\text{tr}(XH^\dagger) = \text{tr}(XH)$ when X is Hermitian, and produces a Hermitian representing matrix. $\square$

Theorem 20.7.13 (Entanglement witness from a separating hyperplane). [Lean](#) [✓ Stmt](#) Let ρ be a trace-one Hermitian bipartite state whose Schmidt number exceeds n . Then there is a Hermitian operator W with

$$\operatorname{Re} \operatorname{tr}(W\rho) < 0, \quad \operatorname{Re} \langle \psi | W | \psi \rangle \geq 0 \quad \text{for every } \psi \text{ of Schmidt rank at most } n.$$

This is the only-if direction of Wolf’s Proposition 3.3 [Wol12, Chapter 3, Proposition 3.3]: a state of Schmidt number larger than n is detected by an entanglement witness for S_n .

Proof. [✓ Proof](#) The set S_n of trace-one states of Schmidt number at most n is convex and compact, and $\rho \notin S_n$. The geometric Hahn–Banach theorem separates the closed convex set $\{\rho\}$ from the compact convex set S_n by a continuous real-linear functional f and a constant c with $f(\rho) < c < f(\sigma)$ for every $\sigma \in S_n$. Representing f as the trace form of a Hermitian H_0 and setting $W = H_0 - c\mathbb{1}$ gives $\operatorname{Re} \operatorname{tr}(W\rho) = f(\rho) - c < 0$. For a vector ψ of Schmidt rank at most n the projector $|\psi\rangle\langle\psi|$, after normalization by $\|\psi\|^2$, is a trace-one state of S_n , so its value under f is at least c ; multiplying back by the squared norm gives $\operatorname{Re} \langle \psi | W | \psi \rangle \geq 0$, with the zero vector giving the value 0 directly. $\square$

Theorem 20.7.14 (Witness criterion for Schmidt number). [Lean](#) [✓ Stmt](#) Let ρ be a trace-one Hermitian bipartite state. Then the Schmidt number of ρ exceeds n if and only if there is a Hermitian operator W with

$$\operatorname{Re} \operatorname{tr}(W\rho) < 0, \quad \operatorname{Re} \langle \psi | W | \psi \rangle \geq 0 \quad \text{for every } \psi \text{ of Schmidt rank at most } n.$$

This is Wolf’s Proposition 3.3 [Wol12, Chapter 3, Proposition 3.3]: a state has Schmidt number larger than n exactly when it is detected by an entanglement witness for S_n .

Proof. [✓ Proof](#) The forward implication is the separating-hyperplane construction (20.7.13). For the converse, suppose ρ had Schmidt number at most n , so $\rho = \sum_i |\psi_i\rangle\langle\psi_i|$ with each ψ_i of Schmidt rank at most n . By linearity of the trace, $\operatorname{Re} \operatorname{tr}(W\rho) = \sum_i \operatorname{Re} \operatorname{tr}(W|\psi_i\rangle\langle\psi_i|)$, a sum of nonnegative terms, contradicting $\operatorname{Re} \operatorname{tr}(W\rho) < 0$. This converse uses no density-matrix hypotheses on ρ . $\square$

Theorem 20.7.15 (Surjectivity of the Choi correspondence). [Lean](#) [✓ Stmt](#) Assume $D > 0$. Every bipartite matrix W on $\mathbb{C}^D \otimes \mathbb{C}^D$ is the Choi matrix $\tau_T = (T \otimes \mathbb{1})(|\Omega\rangle\langle\Omega|)$ of some linear map $T: M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$.

Proof. [✓ Proof](#) The Choi correspondence $T \mapsto \tau_T$ is complex-linear and injective. Its domain, the space of linear maps on $M_D(\mathbb{C})$, and its codomain, the bipartite matrix space on $\mathbb{C}^D \otimes \mathbb{C}^D$, both have complex dimension D^4 , so the injective map is also surjective. Hence every W is a Choi matrix. $\square$

Theorem 20.7.16 (A witness is the Choi matrix of an n -positive map). [Lean](#) [✓ Stmt](#) Assume $D > 0$. Let W be a Hermitian operator on $\mathbb{C}^D \otimes \mathbb{C}^D$ with

$$\operatorname{Re} \langle \psi | W | \psi \rangle \geq 0 \quad \text{for every } \psi \text{ of Schmidt rank at most } n.$$

Then W is the Choi matrix of an n -positive map T . This is the Choi–Jamiołkowski translation of the entanglement witness of Wolf’s Proposition 3.3 into the n -positive map of Wolf’s Proposition 3.4 [Wol12, Chapter 3, Proposition 3.4].

Proof. ✓ Proof By the surjectivity of the Choi correspondence (20.7.15) there is a linear map T with $\tau_T = W$. Because W is Hermitian, the Choi quadratic form $\langle \psi | W | \psi \rangle$ is real, and it equals the witness expectation $\text{tr}(W |\psi\rangle\langle\psi|)$; the witness condition makes it nonnegative on every vector of Schmidt rank at most n . This is exactly the Schmidt-rank Choi criterion (5.1.1.34) for n -positivity of T . $\square$

Theorem 20.7.17 (Choi trace pairing through the trace-pairing adjoint). Lean ✓ Stmt For a linear map $T: M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ with Choi matrix τ_T and any bipartite matrix ρ on $\mathbb{C}^D \otimes \mathbb{C}^D$,

$$\text{tr}(\tau_T \rho) = \langle \Omega | (T^* \otimes \mathbb{1})(\rho) | \Omega \rangle,$$

where T^* is the trace-pairing adjoint of T .

Proof. ✓ Proof Since $\tau_T = (T \otimes \mathbb{1})(|\Omega\rangle\langle\Omega|)$, cyclicity of the trace gives $\text{tr}(\tau_T \rho) = \text{tr}(\rho (T \otimes \mathbb{1})(|\Omega\rangle\langle\Omega|))$. The $\mathbb{1}$ -ampliation of the trace-pairing adjoint is the trace-pairing adjoint of the ampliation, so moving T across the trace pairing replaces $(T \otimes \mathbb{1})$ by $(T^* \otimes \mathbb{1})$ acting on ρ and turns the pairing against $|\Omega\rangle\langle\Omega|$ into the quadratic form $\langle \Omega | \cdot | \Omega \rangle$. $\square$

Theorem 20.7.18 (Positive maps detect high Schmidt number). Lean ✓ Stmt Let ρ be a trace-one Hermitian bipartite state on $\mathbb{C}^D \otimes \mathbb{C}^D$ whose Schmidt number exceeds n . Then there is an n -positive map $T: M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ such that $(T \otimes \mathbb{1})(\rho)$ is not positive semidefinite. This is the if direction of Wolf's Proposition 3.4 [Wol12, Chapter 3, Proposition 3.4].

Proof. ✓ Proof The entanglement witness W for ρ (20.7.13) is the Choi matrix of an n -positive map P (20.7.16). Its trace-pairing adjoint $T = P^*$ is again n -positive (5.1.1.52). By the Choi trace pairing through the trace-pairing adjoint (20.7.17),

$$\langle \Omega | (T \otimes \mathbb{1})(\rho) | \Omega \rangle = \text{tr}(W \rho),$$

whose real part is negative. A positive semidefinite matrix has nonnegative quadratic forms, so $(T \otimes \mathbb{1})(\rho)$ is not positive semidefinite. $\square$

Chapter 21

Matrix Product Operators and Density Operators

Fix a physical dimension d and a bond dimension D . An MPO tensor carries two physical indices, one ket index and one bra index, together with the virtual left and right indices. The renormalization fixed-point conditions and their algebraic consequences are developed in Chapters 23 and 24.

21.1 MPO tensors

Definition 21.1.1 (MPO tensor). [Lean](#) [✓ Stmt](#) An *MPO tensor* is a family of matrices

$$\{M^{ij}\}_{i,j=0}^{d-1}, \quad M^{ij} \in M_D(\mathbb{C}),$$

indexed by a ket index i and a bra index j . Diagrammatically,



The black node denotes the tensor M , the horizontal legs are virtual, the upper leg is the ket index i , and the lower leg is the bra index j .

Definition 21.1.2 (Doubled-index MPS view). [Lean](#) [✓ Stmt](#) By combining the pair (i, j) into a single physical index in $\{0, \dots, d^2 - 1\}$, one obtains an MPS tensor with physical dimension d^2 and bond dimension D :

$$M_{(i,j)}^{\text{MPS}} = M^{ij},$$

identifying $\{0, \dots, d-1\} \times \{0, \dots, d-1\}$ with $\{0, \dots, d^2-1\}$ via the product encoding.

Definition 21.1.3 (Word evaluation). [Lean](#) [✓ Stmt](#) For words $u = (i_1, \dots, i_N)$ and $v = (j_1, \dots, j_N)$ in $\{0, \dots, d-1\}^N$, the *word evaluation* is

$$M^{u,v} := M^{i_1 j_1} M^{i_2 j_2} \dots M^{i_N j_N} \in M_D(\mathbb{C}).$$

The empty pair of words evaluates to the identity, $M^{\emptyset, \emptyset} = \mathbb{1}_D$, and word pairs of different lengths evaluate to 0.

Lemma 21.1.4 (Word-evaluation factorization). [Lean](#) [✓ Stmt](#) For words u_1, v_1 of equal length and arbitrary words u_2, v_2 , word evaluation factors as

$$M^{u_1 u_2, v_1 v_2} = M^{u_1, v_1} M^{u_2, v_2}.$$

Proof. [✓ Proof](#) We argue by induction on the common length of u_1 and v_1 . For the empty prefix,

$$M^{u_2, v_2} = \mathbb{1}_D M^{u_2, v_2}.$$

If $u_1 = iu$ and $v_1 = jv$, then

$$M^{iuu_2, jvv_2} = M^{ij} M^{uu_2, vv_2} = M^{ij} M^{u, v} M^{u_2, v_2} = M^{iu, jv} M^{u_2, v_2},$$

using associativity of matrix multiplication. $\square$

Lemma 21.1.5 (Trace cyclicity of the closed MPO word). [Lean](#) [✓ Stmt](#) Let $a, b \in \{0, \dots, d-1\}$, and let u, v be words of equal length. Writing au, bv for adding the letters at the beginning and ua, vb for adding them at the end,

$$\text{tr}(M^{au, bv}) = \text{tr}(M^{ua, vb}).$$

Proof. [✓ Proof](#) The factorization Lemma 21.1.4 gives

$$M^{ua, vb} = M^{u, v} M^{ab}, \quad M^{au, bv} = M^{ab} M^{u, v}.$$

Hence


$$\text{tr}(M^{ua, vb}) = \text{tr}(M^{u, v} M^{ab}) = \text{tr}(M^{ab} M^{u, v}) = \text{tr}(M^{au, bv}),$$

by cyclicity of the trace. $\square$

Definition 21.1.6 (MPO operator family). [Lean](#) [✓ Stmt](#) The operator generated by M on N sites is the matrix $\rho^{(N)}(M)$ indexed by configurations $\sigma, \tau \in \{0, \dots, d-1\}^N$ and given by

$$\rho^{(N)}(M)_{\sigma, \tau} = \text{tr}(M^{\sigma, \tau}).$$

Diagrammatically,



each black node denotes the same local tensor M , and the matrix entry $\rho^{(N)}(M)_{\sigma, \tau}$ is obtained by closing the outer virtual legs cyclically and taking the resulting trace.

Definition 21.1.7 (Cyclic-shift reindexing). [Lean](#) [✓ Stmt](#) The *cyclic shift* of configurations on N sites is the bijection that relabels each site by one position, $\sigma \mapsto \sigma \circ r$, where r is the cyclic rotation $i \mapsto i + 1$ of $\{0, \dots, N-1\}$.

Theorem 21.1.8 (Translation invariance of the periodic MPDO). [Lean](#) [✓ Stmt](#) The operator $\rho^{(N)}(M)$ is invariant under the simultaneous cyclic shift of its bra and ket configurations:

$$\rho^{(N)}(M)_{r\sigma, r\tau} = \rho^{(N)}(M)_{\sigma, \tau}.$$

Proof. [✓ Proof](#) The entry is a closed bond trace, $\rho^{(N)}(M)_{\sigma,\tau} = \text{tr}(M^{\sigma_1\tau_1} \dots M^{\sigma_N\tau_N})$. Shifting both configurations by r rotates the matrix product by one factor, and the trace is invariant under that cyclic rotation,

$$\text{tr}(M^{\sigma_2\tau_2} \dots M^{\sigma_N\tau_N} M^{\sigma_1\tau_1}) = \text{tr}(M^{\sigma_1\tau_1} \dots M^{\sigma_N\tau_N}),$$

so $\rho^{(N)}(M)_{r\sigma, r\tau} = \rho^{(N)}(M)_{\sigma,\tau}$. □

Definition 21.1.9 (Hermitian MPO tensor). [Lean](#) [✓ Stmt](#) An MPO tensor is *Hermitian* if, for all $i, j \in \{0, \dots, d-1\}$,

$$M^{ij} = (M^{ji})^\dagger$$

Definition 21.1.10 (Adjoint MPO tensor). [Lean](#) [Lean](#) [✓ Stmt](#) The *adjoint tensor* $M^\dagger$ of an MPO tensor M exchanges the ket and bra physical indices and conjugate-transposes each virtual matrix:

$$(M^\dagger)^{ij} = (M^{ji})^\dagger.$$

A tensor equals its adjoint tensor, $M^\dagger = M$, precisely when it is Hermitian.

Theorem 21.1.11 (Reflection identity for the adjoint tensor). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The adjoint tensor generates the adjoint operator family up to spatial reflection: entrywise,

$$\rho^{(N)}(M^\dagger)_{\sigma\tau} = \overline{\rho^{(N)}(M)_{\tau \circ \text{rev}, \sigma \circ \text{rev}}},$$

where *rev* reverses the site order. For an MPDO the conjugation disappears, $\rho^{(N)}(M^\dagger)_{\sigma\tau} = \rho^{(N)}(M)_{\sigma \circ \text{rev}, \tau \circ \text{rev}}$, and the adjoint tensor is again an MPDO.

Proof. [✓ Proof](#) At the level of word evaluations, the conjugate transpose of a product reverses the factors, so

$$(M^\dagger)^{\sigma,\tau} = (M^{\tau \circ \text{rev}, \sigma \circ \text{rev}})^\dagger.$$

Taking traces gives the entrywise identity

$$\rho^{(N)}(M^\dagger)_{\sigma\tau} = \text{tr}(M^{\tau \circ \text{rev}, \sigma \circ \text{rev}})^\dagger = \overline{\rho^{(N)}(M)_{\tau \circ \text{rev}, \sigma \circ \text{rev}}}.$$

For an MPDO, $\rho^{(N)}(M)$ is Hermitian, so $\overline{\rho^{(N)}(M)_{\tau \circ \text{rev}, \sigma \circ \text{rev}}} = \rho^{(N)}(M)_{\sigma \circ \text{rev}, \tau \circ \text{rev}}$; positive semidefiniteness is preserved under the simultaneous relabeling of rows and columns by the reflection. □

21.2 Transfer maps

Definition 21.2.1 (MPO transfer map). [Lean](#) [✓ Stmt](#) The *transfer map* associated to M is the linear map $\mathcal{E}_M : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$\mathcal{E}_M(X) = \sum_{i,j=0}^{d-1} M^{ij} X (M^{ij})^\dagger.$$

Lemma 21.2.2 (Identification with the doubled-index MPS transfer map). [Lean](#) [✓ Stmt](#) The transfer map of an MPO tensor agrees with the transfer map of its doubled-index MPS view:

$$\mathcal{E}_M = \mathcal{E}_{M^{\text{MPS}}}.$$

Proof. [✓ Proof](#) After identifying the single physical index of the doubled tensor with a pair (i, j) , for every $X \in M_D(\mathbb{C})$,

$$\mathcal{E}_{M^{\text{MPS}}}(X) = \sum_{(i,j)} M_{(i,j)}^{\text{MPS}} X (M_{(i,j)}^{\text{MPS}})^\dagger = \sum_{i,j=0}^{d-1} M^{ij} X (M^{ij})^\dagger = \mathcal{E}_M(X).$$

□

Theorem 21.2.3 (Positivity of the MPO transfer map). [Lean](#) [✓ Stmt](#) If $X \geq 0$, then $\mathcal{E}_M(X) \geq 0$.

Proof. [✓ Proof](#) By Lemma 21.2.2, $\mathcal{E}_M$ is the transfer map of the doubled-index MPS tensor. The corresponding MPS transfer map is positive, so $\mathcal{E}_M(X) \geq 0$ whenever $X \geq 0$. □

21.3 MPDOs and LPDOs

Definition 21.3.1 (MPDO). [Lean](#) [✓ Stmt](#) An MPO tensor is an *MPDO* if, for every $N \in \mathbb{N}$,

$$\rho^{(N)}(M) \geq 0$$

This is the global positivity condition of [CPGSV16, Section 4].

Definition 21.3.2 (LPDO). [Lean](#) [✓ Stmt](#) An MPO tensor is an *LPDO* if there exist a Kraus dimension d_K , an inner bond dimension D' , a bond-space identification $e : \{0, \dots, D-1\} \simeq \{0, \dots, D'-1\} \times \{0, \dots, D'-1\}$ (equivalently, D and $(D')^2$ have the same cardinality), and matrices $A^{(i,k)} \in M_{D'}(\mathbb{C})$ such that, for all physical indices i, j ,

$$M^{ij} = \left(\sum_{k=0}^{d_K-1} A^{(i,k)} \otimes \overline{A^{(j,k)}} \right)_{e,e}$$

where $\otimes$ denotes the Kronecker product, $\overline{(\cdot)}$ denotes entrywise complex conjugation, and $(\cdot)_{e,e}$ denotes reindexing along e back to $\{0, \dots, D-1\}$. See [CPGSV16, Section 4.3]. The local contraction is

where the straight internal line is the ancillary index k , while the ket and bra physical legs remain external.

Theorem 21.3.3 (LPDOs are MPDOs). [Lean](#) [✓ Stmt](#) Every LPDO tensor is an MPDO.

Proof. [✓ Proof](#) By the mixed-product property $(A \otimes B)(C \otimes D) = (AC) \otimes (BD)$, the N -site product decomposes as a sum of Kronecker products after transporting along the bond-space identification e :

$$M^{\sigma_1 \tau_1} \dots M^{\sigma_N \tau_N} = \left(\sum_{\kappa} P_{\kappa} \otimes \overline{Q_{\kappa}} \right)_e,$$

where $P_{\kappa} = A^{(\sigma_1, \kappa_1)} \dots A^{(\sigma_N, \kappa_N)}$ and $Q_{\kappa} = A^{(\tau_1, \kappa_1)} \dots A^{(\tau_N, \kappa_N)}$. Since the trace is invariant under the reindexing by e , we apply $\text{tr}(X \otimes Y) = \text{tr}(X) \text{tr}(Y)$ to obtain

$$\rho^{(N)}(M)_{\sigma, \tau} = \sum_{\kappa} \text{tr}(P_{\kappa}) \overline{\text{tr}(Q_{\kappa})} = \sum_{\kappa} \psi_{\kappa}(\sigma) \overline{\psi_{\kappa}(\tau)},$$

where $\psi_{\kappa}(\sigma) = \text{tr}(P_{\kappa})$. Hence $\rho^{(N)}(M) = \sum_{\kappa} |\psi_{\kappa}\rangle \langle \psi_{\kappa}| \geq 0$. □

21.4 Horizontal and Vertical Canonical Forms

Definition 21.4.1 (Horizontal canonical form for an MPO). [Lean](#) [✓ Stmt](#) An MPO tensor is in *horizontal canonical form* if its doubled-index MPS tensor is a BNT sector decomposition with an independent gauge on every repeated copy:

$$M^{ab} = \bigoplus_{j=0}^{g-1} \bigoplus_{q=0}^{r_j-1} \mu_{j,q} X_{j,q} A_j^{ab} X_{j,q}^{-1}.$$

Here the A_j are the distinct minimal representatives, $r_j > 0$, and every weight $\mu_{j,q}$ is nonzero, $|\mu_{j,q}| \leq 1$, and at least one weight has modulus one. The BNT hypotheses include irreducibility, left-canonicity, normalized self-overlap, eventual linear independence of the representative MPV families, and the exclusion of similarity up to phase between distinct representatives. Repeated gauge-equivalent copies are retained over the same representative rather than separated as distinct blocks. The total bond dimension of the displayed direct sum is exactly D . Equivalently, with the block-diagonal matrix $X = \bigoplus_{j,q} X_{j,q}$,

$$M^{ab} = X \left[\bigoplus_{j,q} \mu_{j,q} A_j^{ab} \right] X^{-1}, \quad X = \bigoplus_{j,q} X_{j,q}.$$

This is the canonical form and its BNT decomposition in [\[CPGSV16, lines 237–246 and 281–301\]](#).

Definition 21.4.2 (MPV representation associated with horizontal canonical form). [Lean](#) [✓ Stmt](#) An MPO tensor has a *horizontal-canonical-form MPV representation* if its doubled-index MPS tensor generates the same complete MPV family as a BNT canonical form

$$\bigoplus_{j=0}^{g-1} \bigoplus_{q=0}^{r_j-1} \mu_{j,q} A_j.$$

Its length- N matrix product vector is

$$\sum_{j=0}^{g-1} \left(\sum_{q=0}^{r_j-1} \mu_{j,q}^N \right) V^{(N)}(A_j).$$

This is the MPV-level decomposition in [\[CPGSV16, Section 2.3, lines 298–301\]](#).

Theorem 21.4.3 (Horizontal canonical form gives the BNT MPV representation). [Lean](#) [✓ Stmt](#) *Every MPO tensor in horizontal canonical form has the associated complete BNT matrix-product-vector representation.*

Proof. [✓ Proof](#) Each copy similarity cancels inside its closed matrix-product trace. Since the two bond dimensions agree, this also identifies the coefficient at length zero. $\square$

Definition 21.4.4 (Vertically viewed tensor of an MPO). [Lean](#) [Lean](#) [✓ Stmt](#) The *vertical direction* of an MPO tensor exchanges the notion of physical and virtual indices [\[CPGSV16, line 943\]](#): the vertically viewed tensor $\widetilde{M}$ has its letters indexed by the horizontal bond pair (a, b) , and each letter is the operator on the physical space with matrix elements

$$(\widetilde{M}_{ab})_{ij} = M_{ab}^{ij}.$$

Concatenating $\widetilde{M}$ vertically generates the matrix product vectors of the vertical direction.

$$K = X \left(\bigoplus_j \mathcal{K}_j \right) X^{-1}$$

Figure 21.1: Horizontal canonical form of the doubled tensor: the gauge X , one basis tensor $\mathcal{K}$ carrying the ket and bra legs, and X^{-1} , threaded by the block-index lines j and q . In the source specialized simple-biCF/ZCL display the weights μ_j are independent of the copy index and absorbed into the basis tensors, so the basis tensor taps only the line j while the gauge factors $X_{j,q}$ tap both lines. This is [CPGSV16, Appendix C.2, lines 1660–1665].

Lemma 21.4.5 (Vertical reflection identity for the adjoint tensor). [Lean](#) [Lean](#) [Lean](#) [Lean](#)

✓ Stmt Adjunction exchanges the two oriented horizontal bond indices in the vertical view:

$$\widetilde{M}^\dagger_{a,b} = (\widetilde{M}_{b,a})^\dagger.$$

Proof. ✓ Proof This follows entrywise from the exchange of the ket and bra indices and conjugate transposition of the virtual matrix. □

Definition 21.4.6 (Vertical canonical form for an MPO). [\[Lean\]](#) [\[Stm\]](#) An MPO tensor is in *vertical canonical form* if there are a basis of normal tensors $\{\tilde{M}_\alpha\}_\alpha$ for its vertically viewed tensor $\tilde{M}$, positive diagonal matrices $\mu_\alpha = \text{diag}(\omega_{\alpha,0}, \dots, \omega_{\alpha,r_\alpha-1})$ with $r_\alpha \geq 1$, and a matrix U from the physical space onto the retained nonzero sector space such that

$$U\widetilde{M}U^\dagger = B, \quad \widetilde{M} = U^\dagger B U, \quad B = \bigoplus_{\alpha} \mu_{\alpha} \otimes M_{\alpha}.$$

This is the conclusion of [CPGSV16, Proposition 4.13, lines 1863–1870]. In this orientation $UU^\dagger = I$ on the retained space, while $U^\dagger U$ is its support projection in the original physical space. The reconstruction identity says that every letter of $\widehat{M}$ is supported on this projection, so the discarded orthogonal complement and the off-diagonal corners vanish. The orthogonal complement may be a zero sector: the source’s general canonical-form construction explicitly permits the sum of the nonzero block dimensions to be strictly smaller than the original dimension [CPGSV16, lines 214–225]. Since each μ_α is diagonal, the summand $\mu_\alpha \otimes M_\alpha$ is the direct sum $\bigoplus_{k=0}^{r_\alpha-1} \omega_{\alpha,k} M_\alpha$ of weighted copies of M_α , so the right-hand side is a block-diagonal tensor over the pairs (α, k) in which the multiplicities r_α and the diagonal entries $\omega_{\alpha,k}$ stay explicit; the coefficients $m_\alpha = \text{tr}(\mu_\alpha) = \sum_{k=0}^{r_\alpha-1} \omega_{\alpha,k}$ appearing in the renormalization fixed-point characterization are recovered from them.

$$U\mathcal{K}U^\dagger = \bigoplus_k \mu_k \otimes A_k$$

Figure 21.2: Vertical direct-sum form of the tensor: after the physical isometry U , each block factorizes as the positive diagonal weight μ_α on the multiplicity factor times the basis tensor M_α , and only M_α carries the virtual legs. This is the decomposition $U\tilde{M}U^\dagger = \bigoplus_\alpha \mu_\alpha \otimes M_\alpha$ of [CPGSV16, Proposition 4.13, lines 1864–1870].

Remark 21.4.7 (Finite witnesses for block-injective canonical form). [Lean](#) [Lean](#) [Lean](#) [Lean](#)

[Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Learn](#) [Stm](#) Let $(A_k)_k$ be a finite family of block tensors of physical dimension d , and fix a length L ; write A_k^w for

the length- L word evaluation of A_k along a word $w \in \{0, \dots, d-1\}^L$. The finite-length spanning hypothesis is that the tuple-valued evaluations span the full product algebra of block matrices,

$$\text{span}_{\mathbb{C}} \{ (A_k^w)_k : w \in \{0, \dots, d-1\}^L \} = \prod_k M_{D_k}(\mathbb{C}).$$

The corresponding finite-length separation property is that some length L makes the trace pairing against block words nondegenerate: there is a length L such that, for every tuple $(\Delta_k)_k$ of block matrices,

$$\left(\forall w \in \{0, \dots, d-1\}^L, \sum_k \text{tr}(\Delta_k A_k^w) = 0 \right) \implies \forall k, \Delta_k = 0.$$

Spanning at length L gives separation at the same length: a trace functional that vanishes on every tuple $(A_k^w)_k$ vanishes on their span, hence on the whole product algebra, so nondegeneracy of the product trace pairing forces every $\Delta_k = 0$. Combined with injectivity, left-canonicity, and nonzero block weights, this finite-length separation supplies the older per-copy trace-separation hypotheses for the block-diagonal tensor.

The spanning hypothesis has an equivalent entrywise form. For a block index k and matrix entry (a, b) , let

$$w \mapsto (A_k^w)_{a,b}$$

be the corresponding scalar word family, indexed jointly by the triple (k, a, b) . Linear independence of this scalar family already implies the spanning condition, hence the separation property, and hence the older per-copy trace-separation hypotheses directly.

The duplicate-scalar counterexample shows that the separation property is strictly stronger than blockwise injectivity, left-canonicity, and nonzero weights alone. Take $d = 1$ and two blocks of bond dimension 1 with $A_0^0 = A_1^0 = 1$ and weights $\mu_0 = 1, \mu_1 = 2$: both blocks are injective and left-canonical, $(A_k^0)^\dagger A_k^0 = 1$, and the weights are nonzero, but at every length L there is a unique word w and $A_0^w = A_1^w = 1$, so $\Delta_0 = 1, \Delta_1 = -1$ satisfies $\sum_k \text{tr}(\Delta_k A_k^w) = 0$ while $\Delta \neq 0$. Thus the separation property fails, and no length makes the entrywise scalar family linearly independent.

The block-injectivity proposition of [CPGSV16, lines 340–345] gives a concrete sufficient criterion for the spanning condition: a common length L with every block L -block-injective, together with a length S at which, for every k , there is a coefficient family $(c_w)_{w \in \{0, \dots, d-1\}^S}$ with

$$\sum_w c_w A_k^w = \mathbb{1}, \quad \forall j \neq k, \sum_w c_w A_j^w = 0.$$

Concatenating the injective prefix with this finite selector family produces the spanning condition at length $L+S$, hence the separation property and the older per-copy trace-separation hypotheses.

The inverse tensor of a block-injective canonical form satisfies

$$\sum_p (\mathcal{K}_j^p)_{\alpha, \beta} (K_{j'}^{-1, p})_{\alpha', \beta'} = \delta_{\alpha, \alpha'} \delta_{\beta, \beta'} \delta_{j, j'}.$$

Contracting K^{-1} with the doubled physical index passes the virtual legs of X and X^{-1} to K^{-1} and identifies its block index with the line j :

$$K^{-1}K = \bigoplus_j \mathbb{1}_j$$

This is the inverse identity in [CPGSV16, Appendix C.2, lines 1666–1670].

Theorem 21.4.8 (Finite block-separation criterion). Lean ✓ Stmt *If a block family admits the finite-length data described in the block-injectivity proposition of [CPGSV16, lines 340–345], then it is in block-injective canonical form.*

Proof. ✓ Proof This is the direct implication from the finite-length hypotheses to the finite-length trace-separation statement. General BNT canonical-form hypotheses now yield a finite representative-indexed witness. What remains open in the sharper block-injectivity proposition of [CPGSV16, lines 340–345] is the stated numerical bound $3D^5$ and its particular selector construction. $\square$

Definition 21.4.9 (Representative word-tuple separation). Lean Lean ✓ Stmt Let $(A_j)_{j=0}^{g-1}$ be the basis representatives of a sector decomposition. They have *representative word-tuple separation at length L* when

$$\text{span}_{\mathbb{C}}\{(A_j^w)_{j=0}^{g-1} : w \in \{0, \dots, d-1\}^L\} = \prod_{j=1}^g M_{D_j}(\mathbb{C}).$$

They have *eventual representative word-tuple separation* when this equality holds at every sufficiently large length. Notice that the family is indexed by the minimal representatives j , and not by the repeated copies lying over each j .

Theorem 21.4.10 (Representative-grouped Lemma L). Lean Lean Lean ✓ Stmt Let A_{tot} be the tensor of a sector decomposition with basis representatives $(A_j)_{j=0}^{g-1}$, multiplicities $r_j > 0$, nonzero weights $\mu_{j,q}$, and grouped coefficients

$$c_j^{(N)} = \sum_{q=0}^{r_j-1} \mu_{j,q}^N.$$

Suppose that the first-site actions of $Y, Z \in M_d(\mathbb{C})$ agree on the MPV family of A_{tot} . If representative word-tuple separation holds at length L and $c_{j_0}^{(L+1)} \neq 0$, then

$$YA_{j_0} = ZA_{j_0}.$$

Consequently, under eventual representative word-tuple separation, $YA_j = ZA_j$ for every representative j . The same conclusion holds if the first-site equality is initially given for any tensor with the same complete MPV family as A_{tot} .

The representative grouping is the argument of [CPGSV16, Appendix C.3, lines 1835–1858].

In the first auxiliary assertion, eventual representative word-tuple separation is an explicit assumption. The later post-blocked theorem derives the needed finite representative separation from the full BNT canonical-form hypotheses. The source’s sharp $3D^5$ bound remains separate, as recorded in `docs/paper-gaps/cpgsv17_bicf_block_separation.tex`.

Proof. ✓ Proof At a fixed tail length L , expand the first-site equality at system size $L + 1$ and group all copies lying over the same representative. For every length- L word w this gives

$$\sum_{j=0}^{g-1} \text{tr}(c_j^{(L+1)}(YA_j - ZA_j)A_j^w) = 0.$$

Representative word-tuple separation therefore forces each matrix $c_j^{(L+1)}(YA_j - ZA_j)$ to vanish. A nonzero coefficient may then be cancelled.

For the eventual assertion, choose a threshold L_0 and put $M = L_0 + 1$. For a fixed representative j , apply Corollary 10.3.5 to the nonzero numbers $\mu_{j,q}^M$. It yields $r \geq 1$ such that

$$c_j^{(Mr)} = \sum_q (\mu_{j,q}^M)^r \neq 0.$$

Since $Mr - 1 \geq L_0$, the fixed-length argument applies with $L = Mr - 1$. Finally, equality of complete MPV families transports the first-site hypothesis to the sector-decomposition tensor. $\square$

Definition 21.4.11 (Direct sum with one marked letter). [Lean](#) [✓ Stmt](#) Let

$$A_{\text{tot}}^i = \bigoplus_{j=0}^{g-1} \bigoplus_{q=0}^{r_j-1} \mu_{j,q} A_j^i$$

be the tensor of a sector decomposition. Let s range over a second finite alphabet, and for each representative A_j let C_j^s be a matrix on its bond space. The corresponding direct sum with one marked letter is

$$C_{\text{tot}}^s = \bigoplus_{j=0}^{g-1} \bigoplus_{q=0}^{r_j-1} \mu_{j,q} C_j^s.$$

The marked alphabet need not coincide with the alphabet of the unmarked tensor. This is the algebraic form of the marked matrices in the two diagrams of [CPGSV16, Proposition 4.13, lines 1909–1919].

Lemma 21.4.12 (Trace expansion with one marked letter). [Lean](#) [✓ Stmt](#) For every marked index s and every unmarked word w ,

$$\text{tr}(C_{\text{tot}}^s A_{\text{tot}}^w) = \sum_{j=0}^{g-1} c_j^{(|w|+1)} \text{tr}(C_j^s A_j^w), \quad c_j^{(N)} = \sum_{q=0}^{r_j-1} \mu_{j,q}^N.$$

The exponent $|w| + 1$ includes the weight at the marked site.

Proof. [✓ Proof](#) Expand both matrices as direct sums. Their product is the direct sum of

$$\mu_{j,q}^{|w|+1} C_j^s A_j^w.$$

Taking the trace and summing first over q gives the stated coefficient $c_j^{(|w|+1)}$. $\square$

Theorem 21.4.13 (Algebraic marked-letter extension of representative separation). [Lean](#) [Lean](#) [✓ Stmt](#) Let (C_j^s) and (E_j^s) be two marked families over the same sector decomposition. Suppose that, for every marked index s and every unmarked word w ,

$$\text{tr}(C_{\text{tot}}^s A_{\text{tot}}^w) = \text{tr}(E_{\text{tot}}^s A_{\text{tot}}^w).$$

If representative word-tuple separation holds at length L and $c_{j_0}^{(L+1)} \neq 0$, then $C_{j_0} = E_{j_0}$. Under eventual representative word-tuple separation, one has $C_j = E_j$ for every j .

Lemma L is stated for marks induced by physical operators on the first site. After its closed-chain identity has been expanded, the separation calculation uses only the matrices in the marked slot. The theorem is the resulting algebraic extension of [CPGSV16, Appendix C.3, lines 1848–1858]; it is not attributed to the paper as a separately stated arbitrary-mark theorem.

Proof. [✓ Proof](#) Fix s and a tail length L , and put

$$\Delta_j = c_j^{(L+1)} (C_j^s - E_j^s).$$

Lemma 21.4.12 gives, for every length- L word w ,

$$\sum_{j=0}^{g-1} \text{tr}(\Delta_j A_j^w) = 0.$$

Representative word-tuple separation implies $\Delta_j = 0$ for every j . If $c_{j_0}^{(L+1)} \neq 0$, cancellation gives $C_{j_0}^s = E_{j_0}^s$. Since s was arbitrary, $C_{j_0} = E_{j_0}$.

Under eventual separation, apply Corollary 10.3.5 to the nonzero weights above each fixed representative, exactly as in Theorem 21.4.10. This selects a sufficiently large L with $c_j^{(L+1)} \neq 0$ and proves $C_j = E_j$. $\square$

Definition 21.4.14 (First-site action on a physical block). [Lean](#) [✓ Stmt](#) Let $Y \in M_d(\mathbb{C})$, and write physical indices of an $(L+1)$ -site block as $I = (i_0, \dots, i_L)$ and $J = (j_0, \dots, j_L)$. The induced matrix on the blocked physical space is

$$Y_{I,J}^{[L+1]} = Y_{i_0, j_0} \prod_{k=1}^L \delta_{i_k, j_k}.$$

Lemma 21.4.15 (First-site actions under physical blocking). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) This auxiliary statement supplies the transport needed to combine the block-injectivity argument [CPGSV16, lines 318–345] with Appendix C.3, Lemma L [CPGSV16, lines 1835–1858].

Let A be a tensor, and let $Y^{[L+1]}$ be the physical-block action induced by $Y \in M_d(\mathbb{C})$. For $I = (i_0, \dots, i_L)$, its insertion into the blocked tensor satisfies

$$(Y^{[L+1]} A^{[L+1]})^I = (YA)^{i_0} A^{i_1} \dots A^{i_L}.$$

Hence equality of the first-site actions of Y, Z on the MPV family of A implies equality of the first-block actions of $Y^{[L+1]}, Z^{[L+1]}$ on the MPV family of $A^{[L+1]}$.

The first-site actions of Y, Z agree at every length if and only if

$$\text{tr}((YA)^i A^w) = \text{tr}((ZA)^i A^w)$$

for every physical index i and every finite word w .

Conversely, suppose that A is L -block injective. If the two induced insertion tensors on $A^{[L+1]}$ are equal, then $YA = ZA$.

Proof. [✓ Proof](#) For the forward implication, split the first block into its first letter and its remaining L letters. The displayed insertion identity reduces the blocked MPV equality to the original first-site equality, with the remaining letters of the first block prepended to the flattened tail.

For the converse, put $\Delta^i = (YA)^i - (ZA)^i$. Equality after blocking gives, for every word w of length L ,

$$\Delta^i A^w = 0.$$

Since the length- L words span $M_D(\mathbb{C})$, the same equality holds with A^w replaced by the identity, and therefore $\Delta^i = 0$ for every physical index i . $\square$

Theorem 21.4.16 (Representative separation after injective blocking). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let P be a BNT canonical form with

basis representatives $(A_j)_{j=0}^{g-1}$, and suppose that the representatives have already been blocked so that every A_j is injective. Then representative word-tuple separation holds at every length

$$L \geq 6(g-1) + 1.$$

Consequently, if the first-site actions of $Y, Z \in M_d(\mathbb{C})$ agree on the complete MPV family of the tensor represented by P , then

$$YA_j = ZA_j \quad (0 \leq j < g).$$

The same conclusion holds when the first-site equality is given for any tensor with the same complete MPV family as the tensor represented by P .

More generally, let P be a BNT canonical form before physical blocking, let $p > 0$, and suppose that every p -blocked representative $A_j^{[p]}$ is injective. Then the blocked sector decomposition $P^{[p]}$ has representative word-tuple separation at every blocked length $L \geq 6(g-1) + 1$.

Without changing the physical alphabet, a BNT canonical form also has eventual representative word-tuple separation. If $p > 0$ is a common block-injectivity length, then one may take pairwise separating words of length $6p$, form selectors of length $6p(g-1)$, and append homogeneous words of any length at least p .

Finally, the BNT canonical-form hypotheses imply that there is a common $L > 0$ for which every unblocked representative A_j is L -block injective. Indeed, irreducibility and left-canonicity give a finite peripheral period m_j . The self-overlap tends to m_j along multiples of m_j , but by normalization it also tends to 1; hence $m_j = 1$. Thus every representative is normal, and finiteness gives a common positive block-injectivity length. If the first-site actions of Y, Z agree on the tensor represented by P , then

$$YA_j = ZA_j \quad (0 \leq j < g).$$

The same conclusion holds when the first-site equality is given for any tensor with the same complete MPV family. This composes the blocking conclusion of [CPGSV16, lines 318–345] with Appendix C.3, Lemma L [CPGSV16, lines 1835–1858].

No additional common-length hypothesis is required.

Proof. ✓ Proof Injectivity of A_j implies that its homogeneous words span $M_{D_j}(\mathbb{C})$ at every positive length. Apply the three-block separation argument with source length 1. For each ordered pair of distinct representatives it gives a pairwise block-separating equation at length 6. Multiplying the equations for the $g-1$ representatives different from A_j gives coefficients $c_w^{(j)}$ satisfying

$$\sum_{|w|=6(g-1)} c_w^{(j)} A_k^w = \begin{cases} I, & k = j, \\ 0, & k \neq j. \end{cases}$$

If $L \geq 6(g-1) + 1$, put $m = L - 6(g-1) > 0$. The length- m words of each A_j span $M_{D_j}(\mathbb{C})$. Concatenating these words with the displayed selectors shows that the simultaneous length- L word tuples span $\prod_j M_{D_j}(\mathbb{C})$. This proves eventual representative word-tuple separation. The representative-grouped Lemma L gives the two first-site conclusions.

For the unblocked canonical form, propagate p -block injectivity to the lengths $2p-1$, $2p$, and $6p$. Applying the direct-sum argument to the resulting injective tensors gives pair trace separation at original length $6p$. The bijection between blocked words of length 6 and original words of length $6p$ transports this to pair trace separation for the p -blocked representatives. The preceding selector argument then applies verbatim to $P^{[p]}$.

The same pairwise equations at original length $6p$ give selectors for the unblocked representatives directly. For every $L \geq 6p(g-1) + p$, the remaining prefix has length at least p and

hence spans each full representative matrix algebra. Concatenating the prefix with the selectors proves eventual representative word-tuple separation before blocking.

For the final assertion, first consider one representative A_j . Irreducibility and left-canonicity give a positive peripheral period m_j . The periodic self-overlap formula and the normalized self-overlap hypothesis give, along the same subsequence,

$$\lim_{k \rightarrow \infty} O_{A_j A_j}(m_j k) = m_j, \quad \lim_{k \rightarrow \infty} O_{A_j A_j}(m_j k) = 1.$$

Hence $m_j = 1$. Primitivity then implies normality. Taking a common multiple of the finitely many normality lengths gives a positive length L at which every representative is block injective. Block $L + 1$ sites. Injectivity propagates from length L to length $L + 1$, so the blocked representatives are injective. Act on the first letter of each physical block and apply the blocked representative separation followed by the representative-grouped Lemma L. This gives equality of the two induced insertions on every blocked representative. The remaining L letters of the first block form a spanning family, so Lemma 21.4.15 recovers $Y A_j = Z A_j$ before blocking. $\square$

Corollary 21.4.17 (Marked-letter separation for a BNT canonical form). [Lean] [Stmt] [Lean] Let (C_j^s) and (E_j^s) be two marked families over a BNT canonical form. If their closed marked chains agree for every marked index s and every unmarked word w , then

$$C_j^s = E_j^s$$

for every representative j and every marked index s .

Proof. [Proof] The BNT canonical-form hypotheses give eventual representative word-tuple separation. Apply Theorem 21.4.13. $\square$

Corollary 21.4.18 (Physical-letter marked separation in horizontal canonical form). [Lean] [Lean] [Lean] [Lean] [Lean] [Stmt] Let M be an MPO tensor in literal horizontal canonical form. For two coefficient families $f_{u,z}$ and $g_{u,z}$, put

$$C^u = \sum_z f_{u,z} M^z, \quad E^u = \sum_z g_{u,z} M^z.$$

If, for every marked index u and every word w ,

$$\text{tr}(C^u M^w) = \text{tr}(E^u M^w),$$

then $C^u = E^u$ for every u .

The marks are required to lie in the span of the physical letters. No assertion is made for arbitrary bond-space matrices, since off-diagonal matrices between repeated sectors are invisible to closed chains. The corollary is the physical-insertion consequence of [CPGSV16, Appendix C.3, lines 1835–1858] used in the contractions of [CPGSV16, Proposition 4.13, lines 1909–1919].

Proof. [Proof] Pass through the block-diagonal similarity in the literal horizontal canonical form. Linear combinations of physical letters are covariant under this similarity, and cyclicity of the trace removes it from each closed marked chain. On the representative-indexed sector decomposition one has

$$C_{\text{tot}}^u = \bigoplus_{j,q} \mu_{j,q} \sum_z f_{u,z} A_j^z, \quad E_{\text{tot}}^u = \bigoplus_{j,q} \mu_{j,q} \sum_z g_{u,z} A_j^z.$$

Corollary 21.4.17 identifies the two marks on every minimal representative. Taking their weighted direct sums and restoring the common block-diagonal similarity gives $C^u = E^u$. $\square$

of one representative before the trace-separation step; the block-injective hypothesis stated here instead demands trace separation simultaneously across every block, including gauge-equivalent copies, and so excludes decompositions with repeated normal tensors. Documented in `docs/paper-gaps/cpgsv17_bicf_block_separation.tex`.

$$YH^{(N)}(\widetilde{M}) = ZH^{(N)}(\widetilde{M})$$

$$YA_k = ZA_k$$

Figure 21.3: Blockwise equality from agreement on the MPV family. Top: the hypothesis $Y_1 V^{(N)}(A) = Z_1 V^{(N)}(A)$ for every length N , with the operator inserted on the physical leg of the first site. Bottom: the conclusion that the two insertions agree on every block A_k of the canonical form. These are the two displays of Lemma L, [CPGSV16, Appendix C.3, lines 1835–1846].

Proof. ✓ Proof This is the per-block-separated form of Lemma L of [CPGSV16, Appendix C.3, lines 1835–1846], using the finite criterion from Theorem 21.4.8 applied directly to the full block family rather than to a minimal basis of representatives. Expanding the hypothesis over a word of length $L + 1$ beginning with the fixed site and folding each block contribution into a trace pairing against the corresponding first-site insertion tensor yields, upon taking the Y - Z difference, a tuple $(\Delta_k)_k$ of block matrices such that

$$\sum_k \text{tr}(\mu_k^{L+1} \Delta_k A_k^w) = 0$$

for every word w of length L . The finite block-separation hypothesis applied to the tuple $(\mu_k^{L+1} \Delta_k)_k$ forces each weighted block difference to be zero, and the nonzero weights give the blockwise equality of the two first-site insertions. $\square$

The trace-pairing step used above is the following local identity:

$$\text{tr}(WYA_k) = \text{tr}(WZA_k)$$

For every W in the span of the length- N word evaluations of the canonical form, it states $\text{tr}(WA^Y) = \text{tr}(WA^Z)$, where A^Y is obtained by applying Y to the physical index. Compare the two displays in the proof of Lemma L, [CPGSV16, Appendix C.3, lines 1848–1858].

Theorem 21.4.21 (One-sided invariant corners of positive operators). Lean Lean ✓ Stmt Let $H \geq 0$ and let $P = P^\dagger$. If

$$PH = PHP,$$

then

$$(\mathbb{1} - P)HP = 0.$$

In particular, this conclusion holds for every density operator generated by an MPDO. This is the operator identity used in the proof of [CPGSV16, Proposition 4.13].

Proof. ✓ Proof Since H and P are Hermitian, taking adjoints gives

$$HP = (PH)^\dagger = (PHP)^\dagger = PHP.$$

Therefore $(\mathbb{1} - P)HP = HP - PHP = 0$. $\square$

Theorem 21.4.22 (First-site contraction reduction). [Lean](#) [✓ Stmt](#) Let $\bigoplus_{j,q} \mu_{j,q} A_j$ be a BNT canonical form and let $Y_L, Y_C, Y_R \in M_d(\mathbb{C})$, with $Y_1 V^{(N)}(A)$ denoting Y acting on the first physical index of the MPV. If, for every $N \geq 1$,

$$(Y_L)_1 V^{(N)}(A) = (Y_C)_1 V^{(N)}(A), \quad (Y_L)_1 V^{(N)}(A) = (Y_R)_1 V^{(N)}(A),$$

then $Y_R A_j = Y_C A_j$ for every minimal representative A_j . This is the representative-grouped application of Lemma L and anticipates the concrete instantiation with Y_L, Y_C, Y_R the physical-index insertions of P , PHP , HP in Theorem 21.4.26.

Proof. [✓ Proof](#) Apply the representative-grouped Lemma L first to the actions Y_L, Y_R and then to Y_L, Y_C . The BNT canonical-form hypotheses supply the required common block-injectivity length. Transitivity gives the stated equality on every minimal representative; the repeated copies have already been combined through $\sum_q \mu_{j,q}^N$. $\square$

Definition 21.4.23 (Doubled-index forms of the three first-site contractions). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $P \in M_d(\mathbb{C})$ and write doubled indices as $p = (a, b)$ and $q = (c, e)$. The first-site matrices corresponding respectively to $PH^{(N)}$, $H^{(N)}P$, and $PH^{(N)}P$ are

$$(Y_L)_{p,q} = P_{a,c} \delta_{b,e}, \quad (Y_R)_{p,q} = \delta_{a,c} P_{e,b}, \quad (Y_C)_{p,q} = P_{a,c} P_{e,b}.$$

The horizontal operator $H^{(N)}(\widetilde{M})$ is the cyclic word

$$H^{(N)}(\widetilde{M}) = \text{tr}_v(\widetilde{M}_1 \cdots \widetilde{M}_N)$$

Its virtual indices are contracted around the row, while every ket and bra index remains external. Reading each ket–bra pair as one doubled index identifies its entries with the matrix product vector of the doubled-index tensor. This is the contraction in [CPGSV16, Proposition 4.13, lines 1876–1880].

Lemma 21.4.24 (Coefficient form of the three first-site contractions). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For ket and bra configurations σ, τ ,

$$V^{(N)}(\widehat{M})_{(\sigma, \tau)} = H^{(N)}(M)_{\sigma, \tau}.$$

If $p_0 = (a, b)$ is the first doubled index and σ', τ' are the ket and bra configurations on the remaining N sites, contraction with Y_L , Y_R , and Y_C gives, respectively,

$$\begin{aligned} \sum_c P_{a,c} H_{(c, \sigma'), (b, \tau')}^{(N+1)}, \quad \sum_e H_{(a, \sigma'), (e, \tau')}^{(N+1)} P_{e,b}, \\ \sum_{c,e} P_{a,c} H_{(c, \sigma'), (e, \tau')}^{(N+1)} P_{e,b}. \end{aligned}$$

Proof. [✓ Proof](#) Decode the doubled index at every site as a ket–bra pair. The defining trace of the resulting matrix word is exactly the corresponding entry of $H^{(N)}(M)$. Summing the first doubled index against Y_L , Y_R , or Y_C leaves the displayed single or double sums. $\square$

Lemma 21.4.25 (Doubled-index insertions are vertical multiplications). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $\widehat{M}$ be the doubled-index tensor associated with M . Then insertion of Y_L , Y_R , and Y_C gives, respectively, the doubled-index tensors associated with $P\widehat{M}$, $\widehat{M}P$, and $P\widehat{M}P$.

$$PH^{(N+1)}, \quad H^{(N+1)}P, \quad PH^{(N+1)}P$$

Figure 21.4: The three first-site contractions of the horizontal operator on $N+1$ sites: contracting the first doubled leg with Y_L , Y_R , or Y_C places P on the ket leg, on the bra leg, or on both legs of the first site, and represents $PH^{(N+1)}$, $H^{(N+1)}P$, and $PH^{(N+1)}P$. Compare the first display in the proof of [CPGSV16, Proposition 4.13, lines 1881–1887].

Proof. ✓ Proof Write the output and summed doubled indices as (a, b) and (c, e) . The Kronecker deltas in Y_L and Y_R remove, respectively, the sums over e and c ; Y_C leaves both multiplications. The resulting coefficients are $\sum_c P_{a,c} M^{cb}$, $\sum_e P_{e,b} M^{ae}$, and $\sum_{c,e} P_{a,c} P_{e,b} M^{ce}$. $\square$

Theorem 21.4.26 (Representative reduction for the MPO contractions). Lean ✓ Stmt *Let the doubled-index tensor $\widehat{M}$ generate the same complete MPV family as a BNT canonical form $\bigoplus_{j,q} \mu_{j,q} A_j$. If, entrywise at every positive length,*

$$PH^{(N)} = PH^{(N)}P, \quad PH^{(N)} = H^{(N)}P,$$

then for every minimal representative A_j the insertions associated with $H^{(N)}P$ and $PH^{(N)}P$ agree. Repeated copies over A_j are grouped through the coefficient $\sum_q \mu_{j,q}^N$.

Proof. ✓ Proof Lemma 21.4.24 turns the two operator identities into equality of the corresponding first-site contractions. Equality of matrix product vectors transports these equalities to the chosen canonical-form decomposition. Apply Theorem 21.4.22 with (Y_L, Y_C, Y_R) . $\square$

Definition 21.4.27 (One-site actions of a matrix on the vertically viewed tensor). Lean Lean Lean Lean ✓ Stmt View the MPO tensor M vertically as the family of physical-space operators M_{ab} of Definition 21.4.4, indexed by the virtual indices, with matrix elements $(M_{ab})_{ij} = M_{ab}^{ij}$. A matrix $P \in M_d(\mathbb{C})$ then acts on the tensor by

$$(PM)^{ij} = \sum_k P_{ik} M^{kj}, \quad (MP)^{ij} = \sum_k P_{kj} M^{ik},$$

and on an $(N+1)$ -site chain by the first-spin operator $P \otimes \mathbb{1}^{\otimes N}$. On the letters of the vertically viewed tensor these actions are the left and right matrix products, $(PM)_{ab} = PM_{ab}$ and $(MP)_{ab} = M_{ab}P$. The first-spin operators compose site by site, $P_1 Q_1 = (PQ)_1$.

Theorem 21.4.28 (One-sided invariance and density-operator commutation). Lean Lean ✓ Stmt *Let M generate an MPDO and let $P = P^\dagger$ satisfy $PM = PMP$ on the vertically viewed tensor. For every total length $L \geq 1$, write $P_1 = P \otimes \mathbb{1}^{\otimes (L-1)}$. Then*

$$P_1 H^{(L)} = P_1 H^{(L)} P_1 = H^{(L)} P_1.$$

This is the displayed chain in the proof of [CPGSV16, Proposition 4.13]; the proposition assumes P is an orthogonal projector, but only self-adjointness of P is used.

Proof. ✓ Proof Put $L = N+1$. Splitting off the first site of a density-operator entry gives

$$H_{(i,\sigma),(j,\tau)}^{(N+1)} = \text{tr}(M^{ij} M^{\sigma_0 \tau_0} \dots M^{\sigma_{N-1} \tau_{N-1}}).$$

Summing the first ket index against P therefore pairs the operator $(PM)^{ij}$ against the remaining word; the hypothesis $PM = PMP$ replaces it by $(PMP)^{ij}$, which unfolds to the entry of

$P_1 H^{(N+1)} P_1$. This proves $P_1 H^{(N+1)} = P_1 H^{(N+1)} P_1$. Since $H^{(N+1)}$ is positive semidefinite and P is self-adjoint, Theorem 21.4.21 gives $(\mathbb{1} - P_1) H^{(N+1)} P_1 = 0$, hence $H^{(N+1)} P_1 = P_1 H^{(N+1)} P_1 = P_1 H^{(N+1)}$. $\square$

Theorem 21.4.29 (One-sided invariance on BNT representatives). [Lean](#) [✓ Stmt](#) *Let M generate an MPDO whose doubled-index tensor $\widehat{M}$ generates the same complete MPV family as a BNT canonical form $\bigoplus_{j,q} \mu_{j,q} A_j$, and let $P = P^\dagger$ satisfy $PM = PMP$ on the vertically viewed tensor. Then on every minimal representative A_j the insertions of MP and PMP agree; equivalently, $(\mathbb{1} - P)MP = 0$ on every representative.*

Proof. [✓ Proof](#) For every tail length N , Theorem 21.4.28 turns $PM = PMP$ into the entrywise identities $P_1 H^{(N+1)} = P_1 H^{(N+1)} P_1$ and $P_1 H^{(N+1)} = H^{(N+1)} P_1$. These are exactly the hypotheses of Theorem 21.4.26, whose conclusion is the equality of the insertions of MP and PMP on every minimal representative. $\square$

Theorem 21.4.30 (Commutation with a power of a positive semidefinite matrix). [Lean](#) [Lean](#) [✓ Stmt](#) *Let $H \geq 0$, let $p \geq 1$, and let Q satisfy $QH^p = H^p Q$. Then $QH = HQ$. In particular, a matrix commuting with a nonzero power of a density operator generated by an MPDO commutes with the density operator itself. This is the final operator implication in the periodic-sector contradiction in the proof of [CPGSV16, Proposition 4.13].*

Proof. [✓ Proof](#) Diagonalize $H = U\Lambda U^\dagger$ with U unitary and Λ the diagonal matrix of eigenvalues $\lambda_i \geq 0$. Conjugating the hypothesis by U gives, entrywise for $Q' = U^\dagger Q U$,

$$Q'_{ij} \lambda_j^p = \lambda_i^p Q'_{ij}.$$

Since $x \mapsto x^p$ is injective on the nonnegative reals, every entry Q'_{ij} with $\lambda_i \neq \lambda_j$ vanishes, so $Q'\Lambda = \Lambda Q'$. Conjugating back gives $QH = HQ$. $\square$

Definition 21.4.31 (Periodic-sector projector). [Lean](#) [✓ Stmt](#) *Let M be an MPO tensor. A periodic-sector projector for M with exponent $p \geq 1$ is an orthogonal projector Q on the physical space such that for every $N \geq 0$, with $Q_1 = Q \otimes \mathbb{1}^{\otimes N}$ acting on the first site,*

$$Q_1 H^{(N+1)} \neq H^{(N+1)} Q_1 \quad \text{and} \quad Q_1 [H^{(N+1)}]^p = [H^{(N+1)}]^p Q_1.$$

These are the commutation identities that the proof of [CPGSV16, Proposition 4.13], lines 1889–1891, attaches to a nontrivial vertical period; the second family is the displayed identity at lines 1890–1891.

Theorem 21.4.32 (No periodic-sector projector for a matrix product density operator). [Lean](#) [✓ Stmt](#) *Let M generate matrix product density operators. Then M admits no periodic-sector projector.*

Proof. [✓ Proof](#) Suppose Q were such a projector. Every $H^{(N)}$ is positive semidefinite, so commutation of Q_1 with $[H^{(N)}]^p$ implies commutation with $H^{(N)}$ (Theorem 21.4.30), contradicting $Q_1 H^{(N)} \neq H^{(N)} Q_1$. This is the contradiction concluding the periodic-sector step in the proof of [CPGSV16, Proposition 4.13], lines 1892–1893. $\square$

Definition 21.4.33 (Periodic vectors supply projectors). [Lean](#) [✓ Stmt](#) *Say that periodic vectors of M supply projectors if for every nontrivial p -periodic vector of the vertically viewed tensor $\widetilde{M}$ — an irreducible restriction B of $\widetilde{M}$ to a minimal invariant subspace whose transfer map has a peripheral eigenvalue different from its spectral radius, as in Definition 9.2.13 — there is an*

exponent $q \geq 1$ for which M admits a periodic-sector projector. The proof of [CPGSV16, Proposition 4.13], lines 1889–1891, asserts this implication from the general theory of the peripheral spectrum of completely positive maps, with q equal to the period p of the vector; only the existence of some exponent enters the exclusion argument. The projector, its word invariance, and its single-letter displacement are constructed from the cyclic decomposition of the peripheral spectrum (Theorem 21.4.41); the passage from the displacement to the all-length non-commutation family is not needed for the periodic-sector contradiction below.

Theorem 21.4.34 (No periodic vectors in the vertical direction). [Lean](#) [✓ Stmt](#) *Let M generate matrix product density operators and suppose that periodic vectors of M supply projectors. Then the vertically viewed tensor $\tilde{M}$ has no nontrivial p -periodic vectors. This is the periodic-sector step in the proof of [CPGSV16, Proposition 4.13], lines 1888–1893, granted the projector that the source draws from the general theory of the peripheral spectrum.*

Proof. [✓ Proof](#) A nontrivial p -periodic vector of $\tilde{M}$ supplies a periodic-sector projector for M , and Theorem 21.4.32 shows that no such projector exists. $\square$

Definition 21.4.35 (Vertically stacked layers). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Placing one MPO layer on top of another and contracting the shared physical index gives the *layer product* of two MPO tensors M and M' with the same physical dimension,

$$(MM')^{ij} = \sum_k M^{ik} \otimes M'^{kj},$$

an MPO tensor on the tensor product of the virtual spaces. The p -fold vertical stack $M^{[p]}$ of M is the iterated layer product of p copies of M ; the empty stack $M^{[0]}$ is the trivial-bond tensor with letters δ_{ij} , which generates the identity operator at every system size.

Theorem 21.4.36 (Stacked layers generate powers of the density operator). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *For every system size N , the density operator generated by a layer product is the product of the density operators generated by the layers,*

$$H^{(N)}(MM') = H^{(N)}(M) H^{(N)}(M').$$

Consequently the p -fold vertical stack generates $H^{(N)}(M^{[p]}) = [H^{(N)}(M)]^p$ for all N and p ; this is the reading of the powers in the displayed identity of [CPGSV16, Proposition 4.13, lines 1890–1891] as density operators of stacked layers. In particular, the vertical stack of a tensor generating matrix product density operators again generates matrix product density operators.

Proof. [✓ Proof](#) The word product of layer-product letters along a ket–bra configuration expands, by induction on the length and the mixed-product property of the Kronecker product, as a sum over configurations κ of the contracted physical index of Kronecker products of the layers' word products. Taking traces, each summand factors into the product of the two traces, so the entry of $H^{(N)}(MM')$ at (σ, τ) equals $\sum_{\kappa} H^{(N)}(M)_{\sigma\kappa} H^{(N)}(M')_{\kappa\tau}$, the entry of the matrix product. The power identity follows by induction on p , and positive semidefiniteness of $[H^{(N)}]^p$ follows from that of $H^{(N)}$. $\square$

Theorem 21.4.37 (Vertical letters of the stacked tensor). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The vertically viewed letters of a layer product are the products of the layers' vertically viewed letters, $(MM')_{(a,a'),(b,b')} = M_{ab} M'_{a'b'}$. Hence every vertically viewed letter of the p -fold stack $M^{[p]}$ is the evaluation of a length- p word of $\tilde{M}$. In particular, if a matrix Q satisfies the one-sided invariance $QA_w = QA_wQ$ for every product A_w of p letters of $\tilde{M}$, then $QM^{[p]} = QM^{[p]}Q$ in the sense of Definition 21.4.27.*

Proof. ✓ Proof The letter identity is a direct computation from the layer product: contracting the shared physical index composes the physical-space operators. Induction on p writes each vertical letter of $M^{[p]}$ as a length- p word. Thus, for every bond index v , there is a word w of length p such that $\widetilde{M^{[p]}}_v = A_w$, and

$$QA_w = QA_wQ \implies Q\widetilde{M^{[p]}}_v = Q\widetilde{M^{[p]}}_vQ.$$

Applying this identity to the two one-site actions gives $QM^{[p]} = QM^{[p]}Q$. □

Definition 21.4.38 (Periodic-sector projector from cyclic data). Lean ✓ Stmt Let M generate matrix product density operators, let $p \geq 1$, and let Q be an orthogonal projector on the physical space satisfying $QA_w = QA_wQ$ for every product A_w of p letters of $\widetilde{M}$ and $Q_1H^{(N)} \neq H^{(N)}Q_1$ for every length. Then Q is a periodic-sector projector for M with exponent p .

Proof. ✓ Proof By Theorem 21.4.37 the word invariance gives $QM^{[p]} = QM^{[p]}Q$ on the stacked tensor, which generates matrix product density operators by Theorem 21.4.36. Theorem 21.4.28 applied to $M^{[p]}$ yields $Q_1H^{(N)}(M^{[p]}) = H^{(N)}(M^{[p]})Q_1$ for every length, and $H^{(N)}(M^{[p]}) = [H^{(N)}]^p$ by the stacked-layers identity. Together with the assumed non-commutation family this is exactly the periodic-sector projector of Definition 21.4.31. □

Definition 21.4.39 (Periodic vectors supply cyclic projectors). Lean ✓ Stmt Say that *periodic vectors of M supply cyclic projectors* if for every nontrivial p -periodic vector of the vertically viewed tensor $\widetilde{M}$ — presented as in Definition 9.2.13 — there are an exponent $q \geq 1$ and an orthogonal projector Q on the physical space satisfying the one-sided invariance $QA_w = QA_wQ$ for every product A_w of q letters of $\widetilde{M}$, together with $Q_1H^{(N)} \neq H^{(N)}Q_1$ for every length. This is the shape of the cyclic decomposition of the peripheral spectrum [Wol12, Theorem 6.6] applied to the vertical transfer map: each cyclic projection of a period- p sector returns to itself after p applications of the map, while its nontrivial cyclic displacement under one application prevents commutation with $H^{(N)}$. Compared with Definition 21.4.33, the commutation of Q with $[H^{(N)}]^p$ is no longer assumed; it is derived through the stacked-layers identity.

Theorem 21.4.40 (Reduction of the periodic-sector step to the cyclic decomposition). Lean ✓ Stmt Let M generate matrix product density operators. If periodic vectors of M supply cyclic projectors, then they supply periodic-sector projectors, and the vertically viewed tensor $\widetilde{M}$ has no nontrivial p -periodic vectors. This reduces the hypothesis of the periodic-sector step in the proof of [CPGSV16, Proposition 4.13, lines 1888–1893] to the cyclic decomposition of the peripheral spectrum of the vertical transfer map.

Proof. ✓ Proof A nontrivial p -periodic vector supplies an invariant non-commuting projector, which is a periodic-sector projector by Definition 21.4.38. The exclusion of periodic vectors is then Theorem 21.4.34. □

Theorem 21.4.41 (The displaced invariant projector of a nontrivial periodic vector). Lean ✓ Stmt Let M be an MPO tensor and let $\widetilde{M}$ carry a nontrivial p -periodic vector, presented as in Definition 9.2.13: an irreducible restriction B of $\widetilde{M}$ to a minimal invariant subspace, reached through an isometry V , with a positive definite transfer eigenvector ρ of eigenvalue $r > 0$ and a transfer eigenvalue $\mu \neq r$ of modulus r . Then there are an exponent $q \geq 1$ and an orthogonal projector Q on the physical space such that $QA_w = QA_wQ$ for every product A_w of q letters of $\widetilde{M}$, while $Q\widetilde{M} \neq Q\widetilde{M}Q$ in the sense of Definition 21.4.27. This realizes the projector that the proof of [CPGSV16, Proposition 4.13, lines 1889–1891] draws from the general theory of the peripheral spectrum.

Proof. [✓ Proof](#) The rescaled unital gauge K of the corner satisfies the hypotheses of the cyclic decomposition of the peripheral spectrum (Theorem 9.2.1.7), and the eigenvalue $\mu/r \neq 1$ of modulus one (Lemma 9.2.1.8) forces the peripheral cyclic group to have order $m \geq 2$. Theorem 7.14.1.13 produces orthogonal projections $P_0, \dots, P_{m-1}$ summing to the identity with $\mathcal{E}_K(P_{k+1}) = P_k$, and the letter-level shift $K_v P_{k+1} = P_k K_v$ holds (Lemma 7.14.1.14). Set $q = m$ and let Q be the complement of the support projection π of $YY^\dagger$ for $Y = V\rho^{1/2}P_0$. A word A_w of m letters of $\widetilde{M}$ satisfies $A_w Y = Y G_w$, where G_w is a scalar multiple of the corresponding word in the letters of K , which commutes with P_0 by the full-period shift; so the range of Y is invariant and Lemma 9.2.1.5 gives $Q A_w = Q A_w Q$. If a single letter also satisfied the invariance, then π would fix $A_v Y$, whose columns lie in the sector $V\rho^{1/2}P_{m-1}$ by the cyclic shift; cancelling the injective factor $V\rho^{1/2}$ and using orthogonality of P_{m-1} and P_0 would force $P_{m-1} K_v = 0$ for every letter, hence $P_{m-1} = \mathcal{E}_K(P_0) = 0$, contradicting the nonvanishing of the cyclic projections. $\square$

Theorem 21.4.42 (Horizontal canonical form excludes vertical periodic sectors). [Lean](#) [Lean](#)
[✓ Stmt](#) *Let M be in horizontal canonical form. If an idempotent Q satisfies $Q\widetilde{M} \neq Q\widetilde{M}Q$, then for some $N \geq 0$,*

$$Q_1 H^{(N+1)} \neq H^{(N+1)} Q_1.$$

Consequently, if M generates matrix product density operators, then $\widetilde{M}$ has no nontrivial periodic vectors.

Proof. [✓ Proof](#) Suppose first that Q_1 commutes with $H^{(N+1)}$ for every N . Since $Q^2 = Q$, this gives

$$Q_1 H^{(N+1)} = Q_1 H^{(N+1)} Q_1$$

at every positive length. Theorem 21.4.19 and Lemma 21.4.25 then imply $Q\widetilde{M} = Q\widetilde{M}Q$, a contradiction. Thus a displaced idempotent has a noncommuting length $N + 1$.

A nontrivial periodic vector supplies the orthogonal projector Q of Theorem 21.4.41, invariant under every word of some full period $p \geq 1$. At the noncommuting length just obtained, the stacked-layer identity and the invariant-projection calculation give

$$Q_1 [H^{(N+1)}]^p = [H^{(N+1)}]^p Q_1.$$

Positive semidefiniteness and Theorem 21.4.30 imply that Q_1 commutes with $H^{(N+1)}$, which is the required contradiction. This uses the pointwise contradiction in [CPGSV16, Proposition 4.13, lines 1888–1893]; the stronger all-length inequality printed there is unnecessary. $\square$

Definition 21.4.43 (Displaced projectors never commute). [Lean](#) [✓ Stmt](#) Say that *displaced projectors of M never commute* if every orthogonal projector Q on the physical space with $Q\widetilde{M} \neq Q\widetilde{M}Q$ satisfies $Q_1 H^{(N)} \neq H^{(N)} Q_1$ at every length. This is the canonical-form step of the periodic-sector argument: the proof of [CPGSV16, Proposition 4.13] assumes the tensor is in canonical form in the horizontal direction, and the argument of lines 1874–1887 turns all-length commutation of Q_1 with $H^{(N)}$ into the letter-level invariance $Q\widetilde{M} = Q\widetilde{M}Q$. The literal transport through a horizontal canonical form is proved. This all-length condition is a stronger auxiliary assumption than is needed for Theorem 21.4.42, which uses the single noncommuting length supplied by the contrapositive of Lemma L.

Theorem 21.4.44 (Reduction of the periodic-sector step to the non-commutation family). [Lean](#)
[Lean](#) [✓ Stmt](#) *Let M generate matrix product density operators. If displaced projectors of M never commute, then periodic vectors of M supply cyclic projectors, and the vertically viewed tensor $\widetilde{M}$ has no nontrivial p -periodic vectors. This narrows the hypothesis of the periodic-sector step in the proof of [CPGSV16, Proposition 4.13, lines 1888–1893] to the canonical-form implication of lines 1874–1887: every other ingredient of the step is proved.*

P at the first site is $P_1 H^{(N+1)} P_1$ with $P_1 = P \otimes \mathbb{1}^{\otimes N}$, and for a matrix $E_\alpha \in M_D(\mathbb{C})$ the chain trace with one marked loop is

$$d_\alpha^{(N+1)} = \text{tr}(E_\alpha E^N).$$

For a representative tensor M_α whose letters are indexed by horizontal bond pairs, its closed vertical loop is defined entrywise by

$$(E_\alpha)_{ab} = \text{tr}(M_\alpha^{(a,b)}).$$

In this case the chain trace is the scalar d_α displayed in the proof of [CPGSV16, Proposition 4.13], resolved by chain length: one loop of M_α followed by loops of M , closed by the horizontal trace.

Lemma 21.4.50 (The trace of the density operator is a chain of loops). [Lean](#) [Lean](#) [✓ Stmt](#) *Summing the diagonal word evaluations of an MPO tensor over all length- N configurations closes the vertical loops site by site,*

$$\sum_{\sigma} M^{\sigma_0 \sigma_0} \dots M^{\sigma_{N-1} \sigma_{N-1}} = E^N,$$

and hence $\text{tr} H^{(N)} = \text{tr}(E^N)$.

Proof. [✓ Proof](#) By induction on N : splitting off the first site factors the sum as $(\sum_a M^{aa})$ times the sum over the remaining N sites. Taking horizontal traces and summing the diagonal entries of $H^{(N)}$ gives the trace identity. $\square$

Theorem 21.4.51 (Trace of a first-site compression). [Lean](#) [Lean](#) [✓ Stmt](#) *For every matrix $P \in M_d(\mathbb{C})$ and every chain length $N + 1$,*

$$\text{tr}(P_1 H^{(N+1)}) = \text{tr}(E_P E^N) :$$

the marked site closes to the loop with P inserted, and the remaining N sites close to plain loops. If P is idempotent, then also $\text{tr}(P_1 H^{(N+1)} P_1) = \text{tr}(E_P E^N)$.

Proof. [✓ Proof](#) For a length- N configuration ρ , write $E_\rho^{(N)} = M^{\rho_0 \rho_0} \dots M^{\rho_{N-1} \rho_{N-1}}$. Expanding at the first site gives

$$(P_1 H^{(N+1)})_{(a,\rho)(a,\rho)} = \sum_i P_{ai} \text{tr}(M^{ia} E_\rho^{(N)}).$$

Summing over a and ρ and collecting closes the remaining loops by Lemma 21.4.50. For the two-sided compression, first-site actions compose site by site, $P_1 Q_1 = (PQ)_1$, so cyclicity of the trace and idempotency reduce $\text{tr}(P_1 H^{(N+1)} P_1)$ to $\text{tr}(P_1 H^{(N+1)})$. $\square$

Definition 21.4.52 (Sector projectors of the vertical decomposition). [Lean](#) [Lean](#) [✓ Stmt](#) A matrix $P \in M_d(\mathbb{C})$ is a sector projector for the weight $\mu \in \mathbb{C}$ and the matrix $E_\alpha \in M_D(\mathbb{C})$ if P is an orthogonal projector and inserting P into the single-site vertical loop selects the sector contribution,

$$E_P = \mu E_\alpha.$$

In the vertical decomposition $\tilde{M} = \bigoplus_{\alpha,k} \mu_{\alpha,k} X_{\alpha,k} M_\alpha X_{\alpha,k}^{-1}$ of the vertically viewed tensor, carried by the physical space in the proof of [CPGSV16, Proposition 4.13], the projector $P_{\alpha,k}$ selecting the sector (α, k) has this property with $\mu = \mu_{\alpha,k}$ and $(E_\alpha)_{ab} = \text{tr}(M_\alpha^{(a,b)})$, independently of k . Indeed, cyclicity of trace cancels the internal similarity $X_{\alpha,k} M_\alpha^{(a,b)} X_{\alpha,k}^{-1}$. Applied to the physical isometry of a grouped normal sector, the gauge-corner identity constructs the projector and proves its loop identity.

Theorem 21.4.53 (A nonzero vertical corner has a nonzero sector compression). [Lean](#) [✓ Stmt](#) Let M be horizontally canonical and let P be a matrix for which $P\widehat{M}_vP$ is nonzero for some vertical letter v . Then the first-site compression $P_1H^{(N+1)}P_1$ is nonzero for some N . This is the separation step used in [CPGSV16, Proposition 4.13, line 1898].

Proof. [✓ Proof](#) If every compression vanished, its matrix entries would say that the first-site insertion by P has the same matrix product vectors as zero. Lemma L for the horizontal canonical form would then force the inserted tensor itself to vanish, contrary to the assumed nonzero corner. $\square$

Theorem 21.4.54 (The sector trace identity). [Lean](#) [✓ Stmt](#) Let P be a sector projector for the weight μ and the matrix E_α . Then for every chain length $N + 1$,

$$\text{tr}(P_1H^{(N+1)}P_1) = \mu d_\alpha^{(N+1)}.$$

Once spectral normalization and grouping identify the reducing projectors with sector projectors $P_{\alpha,k}$ satisfying Definition 21.4.52, this specializes to the trace identification $\text{tr}(P_{\alpha,k}H^{(N)}P_{\alpha,k}) = \mu_{\alpha,k}d_\alpha^{(N)}$ used in the proof of [CPGSV16, Proposition 4.13]; the source states it at the particular length where the compression does not vanish.

Proof. [✓ Proof](#) By Theorem 21.4.51 the trace of the compression is $\text{tr}(E_P E^N)$, and the sector-projector property replaces E_P by μE_α . $\square$

Theorem 21.4.55 (Positivity of the vertical weights). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let M generate an MPDO, let Q and P be sector projectors for the weights $\mu_{\alpha,1}$ and $\mu_{\alpha,k}$ and a common matrix E_α , and suppose $\mu_{\alpha,1} > 0$. If the compression of $H^{(N+1)}$ by P at the first site is nonzero, then

$$\mu_{\alpha,k} d_\alpha^{(N+1)} > 0, \quad d_\alpha^{(N+1)} > 0, \quad \mu_{\alpha,k} > 0.$$

Granted such sector projectors, this yields the conclusion of the trace argument in the proof of [CPGSV16, Proposition 4.13]: since w.l.o.g. $\mu_{\alpha,1} > 0$, the positive trace forces $d_\alpha > 0$ and hence $\mu_{\alpha,k} > 0$.

Proof. [✓ Proof](#) By Theorem 21.4.48 the nonzero compression has positive trace, which Theorem 21.4.54 identifies with $\mu_{\alpha,k} d_\alpha^{(N+1)}$. In particular $d_\alpha^{(N+1)} \neq 0$, so the trace of the compression by the sector projector of $(\alpha, 1)$ at the same length equals $\mu_{\alpha,1} d_\alpha^{(N+1)} \neq 0$; that compression is therefore also nonzero, and its trace is a positive real. Dividing by $\mu_{\alpha,1} > 0$ gives $d_\alpha^{(N+1)} > 0$, and dividing the first display by $d_\alpha^{(N+1)}$ gives $\mu_{\alpha,k} > 0$. $\square$

Theorem 21.4.56 (Gram rigidity of normal tensors). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let M_α be a normal tensor. Any matrix commuting with every matrix of M_α is a scalar multiple of the identity. In particular, if $X \neq 0$ and $X^\dagger X$ commutes with every matrix of M_α , then

$$X^\dagger X = \omega \mathbb{1}$$

for a necessarily positive constant ω , and $\omega^{-1/2}X$ is unitary. This is the normalized form $X_{\alpha,k}^\dagger X_{\alpha,k} = \omega_{\alpha,k} \mathbb{1}$ of the Gram-matrix proportionality in the proof of [CPGSV16, Proposition 4.13], with $X_{\alpha,1} = \mathbb{1}$ and $U_{\alpha,k} = \omega_{\alpha,k}^{-1/2} X_{\alpha,k}$. Deriving the commutation from self-adjointness of the sector compressions is a separate step.

Proof. [✓ Proof](#) Commutation with the matrices of M_α extends to all word evaluations, and after blocking these span the full matrix algebra, so the commuting matrix is central and hence scalar. The Gram matrix $X^\dagger X$ is positive semidefinite and nonzero, so its scalar value is a positive real ω ; dividing X by $\sqrt{\omega}$ makes the Gram matrix the identity. $\square$

Theorem 21.4.57 (Reflected marked-chain identity). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let M generate positive semidefinite operators at every length, let A be a vertical tensor, and suppose that a matrix V selects the vertical corner

$$V^\dagger \widetilde{M}_v V = cA^v, \quad c > 0.$$

For vertical indices r, s , write $(\mathcal{R}(A)^{rs})_{ab} = (A^{(a,b)})_{rs}$, and set $(A^\sharp)^v = (A^{v^{\text{op}}})^\dagger$. Then, for every pair of equal-length tail words σ, τ ,

$$\text{tr}(\mathcal{R}(A)^{rs} M^{\sigma_1 \tau_1} \dots M^{\sigma_L \tau_L}) = \text{tr}(\mathcal{R}(A^\sharp)^{rs} (M^\dagger)^{\sigma_L \tau_L} \dots (M^\dagger)^{\sigma_1 \tau_1}).$$

Thus the adjoint exchanges the oriented horizontal bond pair and reverses the unmarked tail. In particular, this is not a same-tail identity and does not identify A with $A^\sharp$ letterwise.

Proof. [✓ Proof](#) Compress the first site of the positive operator by V . The resulting matrix is Hermitian. Expanding one matrix entry gives the marked chain on the left and the complex conjugate of the entry with r, s and the ket–bra words exchanged. Taking the conjugate transpose reverses the tail product, cyclicity of trace returns the marked factor to the first position, and positivity of c permits cancellation without a phase. $\square$

Theorem 21.4.58 (Pairwise Gram conjugation of grouped vertical corners). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let M be in horizontal canonical form and generate positive semidefinite operators at every length. Suppose two vertical corners are positive scalar multiples of similarities of the same tensor A :

$$V_X^\dagger \widetilde{M}^v V_X = c_X X A^v X^{-1}, \quad V_Y^\dagger \widetilde{M}^v V_Y = c_Y Y A^v Y^{-1}, \quad c_X, c_Y > 0.$$

Then the two Gram conjugations agree for every vertical letter,

$$(X^\dagger X) A^v (X^\dagger X)^{-1} = (Y^\dagger Y) A^v (Y^\dagger Y)^{-1}.$$

Normality of A is not required for this equality; it enters only in the subsequent proportionality of the Gram matrices. This is the pairwise conclusion of Figures 7–8 in [CPGSV16, Proposition 4.13, lines 1909–1919].

Proof. [✓ Proof](#) Conjugate the open indices in the reflected marked-chain identity for the first corner by $X^\dagger$. Its left side becomes the marked chain of $(X^\dagger X) A^v (X^\dagger X)^{-1}$, while the gauge factors on the reflected side cancel and leave the marked chain of $A^\sharp$. The same calculation with Y gives the identical reflected chain. Thus the two Gram-dressed marks have equal closed chains for every tail word.

These marks lie in the physical-letter span. Indeed, with

$$L_X = V_X X, \quad R_X = V_X (X^{-1})^\dagger,$$

their coefficients are

$$f_{(r,s),(i,j)}^X = c_X^{-1} \overline{(L_X)_{ir}} (R_X)_{js},$$

and similarly for Y . The corner identities show that these linear combinations are precisely the horizontal slices of the two Gram-dressed tensors. Physical-letter marked separation therefore identifies every slice, hence every vertical letter. $\square$

Theorem 21.4.59 (Gram conjugation for an actual grouped sector). [Lean](#) [✓ Stmt](#) Consider the grouped vertical decomposition furnished by Theorem 21.4.76. For a phase class α and a copy q , transport its representative tensor to the bond space of that copy. If $X_{\alpha,q}$ is the corresponding gauge, then

$$(X_{\alpha,q}^\dagger X_{\alpha,q}) A_{\alpha,q}^v (X_{\alpha,q}^\dagger X_{\alpha,q})^{-1} = A_{\alpha,q}^v$$

for every vertical letter v . This is Figure 8 for the actual witnesses of the grouped decomposition.

Proof. [✓ Proof](#) Transport the distinguished corner of the phase class along the equality of its bond dimension with that of the copy q . Its distinguished gauge is the identity. Apply Theorem 21.4.58 to this corner and the corner indexed by q . The Gram conjugation belonging to the distinguished corner is the identity, which gives the stated equality. $\square$

Theorem 21.4.60 (Gram conjugation from the adjoint dressing). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let X be invertible and suppose that B and a target C satisfy

$$XBX^{-1} = (X^{-1})^\dagger CX^\dagger.$$

Then conjugation by the Gram matrix carries B to C ,

$$(X^\dagger X) B (X^\dagger X)^{-1} = C.$$

Hence any two gauges satisfying this conditional identity with the same target have equal Gram conjugations, and the ratio of their Gram matrices commutes with B . This is a useful algebraic route from an abstract common target to the relative identity in [CPGSV16, Proposition 4.13, lines 1909–1919]. The reflected marked-chain argument in the source does not identify the target separately for either sector. The same-letter choice $C = B^\dagger$ and the further specialization $B^\dagger = B$ are useful corollaries, but the latter is not assumed by the source.

Proof. [✓ Proof](#) Multiply the dressed-target identity by $X^\dagger$ on the left and by $(X^{-1})^\dagger$ on the right. The left side becomes $(X^\dagger X)B(X^\dagger X)^{-1}$ and the right side collapses to C . Eliminating the common target for two gauges and multiplying by the inverse Gram matrix gives the relative commutation relation. $\square$

$$X_\alpha^\dagger X_\alpha = Y_\alpha^\dagger Y_\alpha$$

Figure 21.5: Equality of the Gram conjugations for two copies of the same vertical normal tensor: $(X_{\alpha,k}^\dagger X_{\alpha,k})M_\alpha(X_{\alpha,k}^\dagger X_{\alpha,k})^{-1} = (X_{\alpha,1}^\dagger X_{\alpha,1})M_\alpha(X_{\alpha,1}^\dagger X_{\alpha,1})^{-1}$. Theorem 21.4.58 gives this equality from self-adjointness of the corresponding sector compressions and horizontal marked separation. Normality is used only in the subsequent proportionality argument. This is [CPGSV16, Proposition 4.13, lines 1915–1921].

Theorem 21.4.61 (Gram-matrix proportionality from relative Gram conjugations). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let M_α be a normal tensor, and let X and Y be invertible gauges whose Gram matrices induce the same conjugation on each letter M_α^v . Then there is a positive real number ω such that

$$X^\dagger X = \omega Y^\dagger Y,$$

and $\omega^{-1/2}XY^{-1}$ is unitary. Theorem 21.4.58 gives the Figure 8 comparison in [CPGSV16, Proposition 4.13, lines 1909–1921]. This comparison supplies precisely the relative hypothesis for a sector and the distinguished sector. In the normalization $Y = \mathbb{1}$, the resulting unitary is $\omega^{-1/2}X$. The common-target declarations attached to this entry are conditional algebraic corollaries, not the reflected marked-chain statement itself.

Proof. ✓ Proof By Theorem 21.4.60, the ratio $(Y^\dagger Y)^{-1}(X^\dagger X)$ commutes with every letter of M_α . Theorem 21.4.56 makes this ratio scalar. When the matrix dimension is positive, positive definiteness of the two Gram matrices makes the scalar a positive real number ω . In dimension zero the equality is vacuous and one may choose $\omega = 1$. Conjugating the relative Gram identity by Y^{-1} shows that XY^{-1} has Gram matrix $\omega \mathbb{1}$; division by $\sqrt{\omega}$ makes it unitary. $\square$

Theorem 21.4.62 (Normalization of the grouped vertical gauges). Lean Lean ✓ Stmt *In the grouped vertical decomposition of a horizontally canonical matrix-product density operator, let $X_{\alpha,k}$ be the gauge of a copy of the normal representative M_α , with the distinguished gauge fixed to $X_{\alpha,1} = \mathbb{1}$. Then there is a positive real number $\omega_{\alpha,k}$ such that*

$$X_{\alpha,k}^\dagger X_{\alpha,k} = \omega_{\alpha,k} \mathbb{1},$$

and

$$U_{\alpha,k} = \omega_{\alpha,k}^{-1/2} X_{\alpha,k}$$

is unitary. This is the normalization in [CPGSV16, Proposition 4.13, lines 1903–1921].

Proof. ✓ Proof Transport the normal representative to the bond space of the chosen copy. Theorem 21.4.59 states that conjugation by $X_{\alpha,k}^\dagger X_{\alpha,k}$ fixes every letter of the transported tensor. Apply Theorem 21.4.61 with the second gauge equal to the identity. The resulting Gram identity has a positive real scalar, and division by its positive square root gives the stated unitary. $\square$

Theorem 21.4.63 (Normalized physical maps of the grouped vertical sectors). Lean ✓ Stmt *In the grouped vertical decomposition, put*

$$Q_{\alpha,q} = \omega_{\alpha,q}^{-1/2} X_{\alpha,q}, \quad W_{\alpha,q} = V_{\alpha,q} Q_{\alpha,q},$$

and identify the bond space of each copy with that of its chosen representative. Then

$$W_{\alpha,q}^\dagger W_{\alpha,q} = \mathbb{1}, \quad W_{\alpha,q}^\dagger W_{\beta,p} = 0 \quad \text{if } (\alpha, q) \neq (\beta, p).$$

Writing $c_{\alpha,q} = \nu_{\alpha,q} \zeta_{\alpha,q} > 0$, the normalized maps satisfy

$$\widetilde{M}_v W_{\alpha,q} = W_{\alpha,q} (c_{\alpha,q} M_\alpha^v)$$

and retain the literal reconstruction

$$\widetilde{M}_v = \sum_\alpha \sum_q W_{\alpha,q} (c_{\alpha,q} M_\alpha^v) W_{\alpha,q}^\dagger.$$

These are the normalized sector identities in [CPGSV16, Proposition 4.13, lines 1895–1921].

Proof. ✓ Proof Each $Q_{\alpha,q}$ is unitary, so right multiplication by it preserves the isometry equation and the pairwise orthogonality of the physical sector maps. In the intertwining relation, the scalar normalization commutes with every letter and the adjacent factors $X_{\alpha,q}^{-1} X_{\alpha,q}$ cancel. The Gram identity gives $X_{\alpha,q}^\dagger = \omega_{\alpha,q} X_{\alpha,q}^{-1}$; hence

$$Q_{\alpha,q} M_\alpha^v Q_{\alpha,q}^\dagger = X_{\alpha,q} M_\alpha^v X_{\alpha,q}^{-1}.$$

Substitution into every summand of the grouped reconstruction proves the final identity. Transport along the equality of bond dimensions changes only the coordinate labels. $\square$

Lemma 21.4.64 (Matrix product vector of the diagonal tensor). [Lean](#) [✓ Stmt](#) For an MPO tensor M and a configuration σ , the matrix product vector of the diagonal tensor $i \mapsto M^{ii}$ equals the matrix product vector of the doubled-index tensor $\widehat{M}$ evaluated on the diagonal-paired configuration $k \mapsto (\sigma_k, \sigma_k)$:

$$V^{(N)}(i \mapsto M^{ii})(\sigma) = V^{(N)}(\widehat{M})(k \mapsto (\sigma_k, \sigma_k)).$$

Proof. [✓ Proof](#) At each site value i the diagonal tensor agrees with the doubled-index tensor on the diagonal pair,

$$M^{ii} = \widehat{M}^{(i,i)}.$$

The matrix product vector is multiplicative over sites, so replacing each factor $M^{\sigma_k \sigma_k}$ by $\widehat{M}^{(\sigma_k, \sigma_k)}$ reindexes the configuration from σ to $k \mapsto (\sigma_k, \sigma_k)$, which is the stated equality. $\square$

Lemma 21.4.65 (Diagonal tensor as a block decomposition under horizontal form). [Lean](#) [✓ Stmt](#) Suppose the doubled-index tensor $\widehat{M}$ shares its matrix product vector family with a block-diagonal assembly $\bigoplus_k \mu_k A_k$ of blocks A_k on the doubled index, as in a horizontal canonical form. Then the diagonal tensor of M generates the same matrix product vector family as the block-diagonal assembly $\bigoplus_k \mu_k A_k^{\text{diag}}$ on the single-spin index, where $A_k^{\text{diag}}(i) = A_k(i, i)$ is the diagonal restriction of each block:

$$V^{(N)}(i \mapsto M^{ii})(\sigma) = V^{(N)}\left(\bigoplus_k \mu_k A_k^{\text{diag}}\right)(\sigma).$$

Proof. [✓ Proof](#) By Lemma 21.4.64 the diagonal matrix product vector equals that of $\widehat{M}$ on the diagonal-paired configuration, which by hypothesis equals that of the block-diagonal assembly there. Expanding the assembly as the weighted sum $\sum_k \mu_k^N V^{(N)}(A_k)$ and restricting each block to the diagonal pair gives $\sum_k \mu_k^N V^{(N)}(A_k^{\text{diag}})$, the matrix product vector of the single-spin block-diagonal assembly. $\square$

Theorem 21.4.66 (Per-copy horizontal hypotheses do not control diagonal normality). [Lean](#)

[✓ Stmt](#) There is a left-canonical, injective doubled-index tensor with a nonzero block weight satisfying the older per-copy finite-length separation hypothesis whose restriction to the diagonal physical pairs is not normal. On bond space $\mathbb{C}^2$, take

$$A^{(a,b)} = c_a E_{ab}, \quad c_0 = \frac{3}{5}, \quad c_1 = \frac{4}{5}.$$

Then the four matrices $A^{(a,b)}$ span $M_2(\mathbb{C})$ and

$$\sum_{a,b} (A^{(a,b)})^* A^{(a,b)} = I.$$

For a single block, injectivity also gives finite-length block separation. The diagonal restriction, however, consists only of

$$A^{\text{diag}}(0) = \frac{3}{5} E_{00}, \quad A^{\text{diag}}(1) = \frac{4}{5} E_{11}.$$

Every product of these two matrices is diagonal, so no blocking length spans $M_2(\mathbb{C})$.

Proof. ✓ Proof The four nonzero scalar multiples of the matrix units span the full matrix algebra. The identity $(3/5)^2 + (4/5)^2 = 1$ gives left-canonicity. After diagonal restriction,

$$E_{ii}E_{jj} = \delta_{ij}E_{ii}.$$

Hence every word is either zero or a scalar multiple of some E_{ii} , and therefore has vanishing $(0, 1)$ entry. Since E_{01} does not, it lies outside the span at every blocking length. $\square$

Lemma 21.4.67 (Diagonal tensor word evaluation). Lean ✓ Stmt *For any word w , the matrix product of the diagonal tensor along w equals the operator word evaluation with equal ket and bra words, since both multiply the matrices $M^{w_k w_k}$ site by site.*

Proof. ✓ Proof By induction on the word. $\square$

Lemma 21.4.68 (Diagonal tensor evaluates the density-operator diagonal). Lean ✓ Stmt *For an MPO tensor M and a configuration σ , the matrix product vector of the diagonal tensor equals the diagonal entry of the generated operator,*

$$V^{(N)}(i \mapsto M^{ii})(\sigma) = \langle \sigma | \rho^{(N)}(M) | \sigma \rangle,$$

since both equal $\text{tr}(M^{\sigma_1 \sigma_1} \dots M^{\sigma_N \sigma_N})$.

Proof. ✓ Proof Word evaluation of the diagonal tensor multiplies the matrices $M^{\sigma_k \sigma_k}$, which is exactly the operator word evaluation with equal ket and bra indices; taking the trace gives the diagonal entry. $\square$

Lemma 21.4.69 (Nonnegativity of the diagonal matrix product vector). Lean ✓ Stmt *If $\rho^{(N)}(M)$ is positive semidefinite, then $V^{(N)}(i \mapsto M^{ii})(\sigma) \geq 0$ for every configuration σ .*

Proof. ✓ Proof By Lemma 21.4.68, the value is a diagonal entry of $\rho^{(N)}(M)$, and the diagonal entries of a positive semidefinite matrix are nonnegative. $\square$

Lemma 21.4.70 (Matrix product vector of the vertical-assembled tensor). Lean Lean Lean ✓ Stmt *Let g be a block count, $\text{dim}, \text{mult} : \{0, \dots, g-1\} \rightarrow \mathbb{N}$, weights $\omega_{\alpha j} \in \mathbb{C}$ for $0 \leq j < \text{mult}_{\alpha}$, and blocks A_{α} of bond dimension dim_{α} . The vertical-assembled tensor is the block-diagonal assembly over the flattened index (α, j) that repeats A_{α} with weight $\omega_{\alpha j}$, written $\bigoplus_{\alpha} \bigoplus_j \omega_{\alpha j} A_{\alpha}$. Its matrix product vector decomposes as*

$$V^{(N)}\left(\bigoplus_{\alpha} \bigoplus_j \omega_{\alpha j} A_{\alpha}\right)(\sigma) = \sum_{\alpha} \sum_j (\omega_{\alpha j})^N V^{(N)}(A_{\alpha})(\sigma).$$

Proof. ✓ Proof The vertical-assembled tensor is the block-diagonal assembly over the single flattened index q , with flattened weights $\tilde{\omega}_q$ and blocks $\tilde{A}_q$. By the block matrix-product-vector decomposition (Theorem 2.5.9),

$$V^{(N)}\left(\bigoplus_q \tilde{\omega}_q \tilde{A}_q\right)(\sigma) = \sum_q (\tilde{\omega}_q)^N V^{(N)}(\tilde{A}_q)(\sigma).$$

The flattening identifies q with the pair (α, j) via the bijection

$$\{(\alpha, j) : 0 \leq \alpha < g, 0 \leq j < \text{mult}_{\alpha}\} \simeq \{0, \dots, \sum_{\alpha} \text{mult}_{\alpha} - 1\},$$

under which $\tilde{\omega}_q = \omega_{\alpha j}$ and $\tilde{A}_q = A_{\alpha}$. Reindexing the sum along this bijection yields the double sum over (α, j) . $\square$

Lemma 21.4.71 (Regrouping a block family by repeated representatives). [Lean](#) [✓ Stmt](#) Let $(B_k)_{k \in I}$ be a finite family of blocks with weights ν_k , and let $(A_\alpha)_{\alpha \in J}$ be a finite family of representative blocks. Suppose there is a bijection

$$e : I \simeq \{(\alpha, j) : \alpha \in J, 0 \leq j < m_\alpha\}$$

such that, writing $e(k) = (\alpha, j)$, the paired blocks have equal bond dimensions, $\nu_k = \omega_{\alpha j}$, and for every positive length N ,

$$V^{(N)}(B_k) = V^{(N)}(A_\alpha)$$

Then

$$V^{(N)}\left(\bigoplus_{k \in I} \nu_k B_k\right) = V^{(N)}\left(\bigoplus_{\alpha \in J} \bigoplus_{j=0}^{m_\alpha-1} \omega_{\alpha j} A_\alpha\right)$$

for every length N , including $N = 0$. This is the finite-sum reindexing after the vertical sectors have been grouped in [\[CPGSV16, Proposition 4.13\]](#).

Proof. [✓ Proof](#) For $N > 0$ and every configuration σ , the two sides expand as

$$\sum_{k \in I} \nu_k^N V^{(N)}(B_k)(\sigma), \quad \sum_{\alpha \in J} \sum_{j=0}^{m_\alpha-1} \omega_{\alpha j}^N V^{(N)}(A_\alpha)(\sigma).$$

The summand indexed by k in the first sum equals the summand indexed by $e(k)$ in the second, so reindexing by e proves the equality. At length zero the pointwise dimension equality gives equality of the paired bond dimensions; the same reindexing gives equality of the total dimensions. $\square$

Theorem 21.4.72 (Invariant vertical projections are reducing). [Lean](#) [✓ Stmt](#) Let M be in horizontal canonical form and generate matrix product density operators. If an orthogonal projection P satisfies, for every vertical letter v ,

$$P\widetilde{M}_v = P\widetilde{M}_v P,$$

then, for every vertical letter v , it also satisfies

$$\widetilde{M}_v P = P\widetilde{M}_v P.$$

Thus the vertically viewed tensor has invariant-projector closure.

Proof. [✓ Proof](#) Translate the left-invariance identity into the one-site ket and bra actions of the original tensor. The horizontal canonical-form version of Lemma L gives the opposite one-sided identity. Translating back to the vertically viewed tensor proves the claim. $\square$

Theorem 21.4.73 (Irreducible reducing sectors of the vertically viewed tensor). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 21.4.72, there are positive integers d_k , tensors B_k of bond dimension d_k , and isometries $V_k : \mathbb{C}^{d_k} \rightarrow \mathbb{C}^d$ such that

$$V_k^\dagger V_k = \mathbb{1}, \quad \sum_k V_k V_k^\dagger = \mathbb{1}, \quad V_k^\dagger V_l = 0 \quad (k \neq l).$$

Every B_k is irreducible, and for every vertical letter v ,

$$\widetilde{M}_v V_k = V_k B_k^v, \quad V_k^\dagger \widetilde{M}_v = B_k^v V_k^\dagger, \quad B_k^v = V_k^\dagger \widetilde{M}_v V_k.$$

Moreover, $\widetilde{M}$ and the unit-weight direct sum $\bigoplus_k B_k$ have the same matrix product vectors at every length.

Proof. [✓ Proof](#) Apply the irreducible-corner decomposition under invariant-projector closure to $\widetilde{M}$. $\square$

Theorem 21.4.74 (Complete vertical reducing projectors). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, there are orthogonal projections P_k such that*

$$\sum_k P_k = \mathbb{1}, \quad P_k P_l = 0 \quad (k \neq l), \quad \widetilde{M}_v P_k = P_k \widetilde{M}_v$$

for every k and every vertical letter v .

Proof. [✓ Proof](#) For the isometries of Theorem 21.4.73, set $P_k = V_k V_k^\dagger$. The isometry, orthogonality, completeness, and intertwining identities give the four assertions directly. $\square$

$$U\mathcal{K}U^\dagger = \bigoplus_\alpha M_\alpha$$

Figure 21.6: The vertically viewed tensor decomposes through complete orthogonal physical sectors. Each V_k embeds the irreducible tensor B_k , while $P_k = V_k V_k^\dagger$ is its reducing range projection. The open-leg identity is equivalent, for every vertical letter v , to $\widetilde{M}_v = \sum_k V_k B_{k,v} V_k^\dagger$. This is the iterated invariant-subspace construction in [CPGSV16, lines 200–219], applied to the reduction in [CPGSV16, Proposition 4.13, lines 1873–1887].

Theorem 21.4.75 (Spectrally normalized vertical sectors with physical isometries). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, there are positive integers d_k , positive scalars ν_k , normal tensors B_k of bond dimension d_k , and isometries $V_k : \mathbb{C}^{d_k} \rightarrow \mathbb{C}^d$ with pairwise orthogonal ranges. For every sector k and every vertical letter v ,*

$$\widetilde{M}_v V_k = V_k (\nu_k B_k^v), \quad V_k^\dagger \widetilde{M}_v = (\nu_k B_k^v) V_k^\dagger, \quad \nu_k B_k^v = V_k^\dagger \widetilde{M}_v V_k.$$

Moreover, every vertical letter is reconstructed literally by

$$\widetilde{M}_v = \sum_k V_k (\nu_k B_k^v) V_k^\dagger.$$

Proof. [✓ Proof](#) Retain exactly the irreducible corners containing a nonzero letter. The Perron–Frobenius eigenvalue of each retained corner is positive, and the absence of nontrivial periodic vectors makes the inverse-square-root rescaling normal. Put the removed square root into ν_k and reindex the retained corners. The original complete reducing decomposition expresses each vertical letter as the sum of all compressed corners; every omitted summand vanishes, which gives the displayed reconstruction. Thus the retained range projections need not sum to the identity, but their orthogonal complement is a zero sector. $\square$

Theorem 21.4.76 (Vertical normal sectors grouped into BNT representatives). [Lean](#) [✓ Stmt](#) *Under the same hypotheses, partition the normalized vertical sectors by equality of their matrix product vector families up to a fixed nonzero scalar per site. There are normal representatives M_α , an enumeration $k = (\alpha, q)$ of the original sectors, invertible matrices $X_{\alpha,q}$, and complex numbers $\zeta_{\alpha,q}$ of unit modulus, with $X_{\alpha,1} = \mathbb{1}$ and $\zeta_{\alpha,1} = 1$, such that*

$$B_{\alpha,q}^v = \zeta_{\alpha,q} X_{\alpha,q} M_\alpha^v X_{\alpha,q}^{-1}.$$

The representatives are pairwise gauge-phase distinct and form a basis of normal tensors for the weighted block tensor $\bigoplus_k \nu_k B_k$. Every grouped coefficient $\nu_{\alpha,q} \zeta_{\alpha,q}$ is positive. With the physical isometries from Theorem 21.4.75 retained unchanged, one has both reducing identities, exact compression, and the literal reconstruction

$$\widetilde{M}_v = \sum_{\alpha} \sum_q V_{\alpha,q} (\nu_{\alpha,q} \zeta_{\alpha,q} X_{\alpha,q} M_{\alpha}^v X_{\alpha,q}^{-1}) V_{\alpha,q}^{\dagger}.$$

For each pair (α, q) , the range projector of $V_{\alpha,q}$ obeys the representative-loop identity and gives a nonzero compression at some finite length. This is the decomposition and positivity argument of [CPGSV16, Proposition 4.13, lines 1898–1902]. The relative Gram identity and proportional-unitary normalization of lines 1903–1908 are not asserted here.

Proof. ✓ Proof Take one representative from each phase class. Theorem 10.1.8 gives equality of bond dimensions and the displayed letterwise gauge-phase relation for every copy. Distinct representatives are gauge-phase separated by construction, so Theorem 10.1.11 gives eventual linear independence. The phase-class regrouping of the finite sum gives the BNT expansion. The representative copy is assigned the identity gauge and phase one, while the self-overlap normalization of normal tensors shows that every phase has unit modulus. Substitution in the two intertwining identities, the compression identity, and the literal reconstruction from Theorem 21.4.75 proves all remaining algebraic assertions without altering any physical isometry. Cyclicity of trace gives the sector-loop identities. The nonzero-corner separation theorem supplies a nonzero compression for every sector, and the sector trace identity and MPDO positivity then make every grouped coefficient positive. □

Definition 21.4.77 (Block-row map of sector isometries). Lean ✓ Stmt Let $W_k : \mathbb{C}^{D_k} \rightarrow \mathbb{C}^d$ be a finite family of rectangular matrices. Their block-row map $U : \mathbb{C}^d \rightarrow \bigoplus_k \mathbb{C}^{D_k}$ is defined by

$$Ux = (W_k^{\dagger} x)_k.$$

This realizes the coisometry in the conclusion of [CPGSV16, Proposition 4.13, lines 1863–1870] once the sector isometries have been normalized.

Theorem 21.4.78 (Conjugation and reconstruction from orthogonal sectors). Lean Lean Lean ✓ Stmt Suppose that

$$W_k^{\dagger} W_k = \mathbb{1}, \quad W_k^{\dagger} W_l = 0 \quad (k \neq l),$$

and, for every letter v and every sector k ,

$$T_v W_k = W_k B_k^v.$$

Then the block-row map U is a coisometry and conjugates T to the direct sum of the sector tensors:

$$UU^{\dagger} = \mathbb{1}, \quad UT_v U^{\dagger} = \bigoplus_k B_k^v.$$

If the letters also satisfy the literal sector decomposition

$$T_v = \sum_k W_k B_k^v W_k^{\dagger},$$

then

$$T_v = U^{\dagger} \left(\bigoplus_k B_k^v \right) U.$$

This last equality is a conditional consequence of the displayed literal sector decomposition; no additional assertion from the source is used.

Proof. ✓ Proof The (k, l) block of $UU^\dagger$ is $W_k^\dagger W_l$, hence the first identity follows from orthonormality. The corresponding block of $UT_v U^\dagger$ is

$$W_k^\dagger T_v W_l = W_k^\dagger W_l B_l^v,$$

which is B_k^v for $k = l$ and zero otherwise. Finally, multiplication of the block diagonal matrix by the block row gives the sum $\sum_k W_k B_k^v W_k^\dagger$, which is T_v by hypothesis. The row indices of U form the dependent disjoint union $\coprod_k \{0, \dots, D_k - 1\}$. Its canonical order-preserving bijection with $\{0, \dots, \sum_k D_k - 1\}$ gives the coordinate reindexing required by the vertical canonical-form definition. $\square$

Theorem 21.4.79 (Vertical canonical form from orthogonal grouped sectors). Lean ✓ Stmt Let $\{A_\alpha\}_{\alpha=0}^{g-1}$ be a BNT for the vertically viewed tensor $\widetilde{M}$. For each α , let $r_\alpha > 0$, let $\omega_{\alpha,q} > 0$ for $0 \leq q < r_\alpha$, and let $W_{\alpha,q} : \mathbb{C}^{D_\alpha} \rightarrow \mathbb{C}^d$ satisfy

$$W_{\alpha,q}^\dagger W_{\alpha,q} = \mathbb{1}, \quad W_{\alpha,q}^\dagger W_{\beta,p} = 0 \quad \text{if } (\alpha, q) \neq (\beta, p).$$

Suppose that, for every vertical letter v and every pair (α, q) ,

$$\widetilde{M}_v W_{\alpha,q} = W_{\alpha,q} (\omega_{\alpha,q} A_\alpha^v),$$

and that the letters have the exact sector decomposition

$$\widetilde{M}_v = \sum_{\alpha=0}^{g-1} \sum_{q=0}^{r_\alpha-1} W_{\alpha,q} (\omega_{\alpha,q} A_\alpha^v) W_{\alpha,q}^\dagger.$$

Then M is in vertical canonical form. This is the final finite-dimensional implication in [CPGSV16, Proposition 4.13, lines 1863–1870].

Proof. ✓ Proof Form the block-row map U from the adjoints $W_{\alpha,q}^\dagger$. Theorem 21.4.78 gives

$$UU^\dagger = \mathbb{1}, \quad U \widetilde{M}_v U^\dagger = \bigoplus_{\alpha,q} \omega_{\alpha,q} A_\alpha^v,$$

together with

$$\widetilde{M}_v = U^\dagger \left(\bigoplus_{\alpha,q} \omega_{\alpha,q} A_\alpha^v \right) U.$$

Identifying the dependent pairs (α, q) and their bond indices with the standard coordinates of the two finite direct sums gives the required matrix dimensions. Positivity of the multiplicities and weights, together with the BNT hypothesis, supplies the remaining conditions of vertical canonical form. $\square$

Lemma 21.4.80 (A retained vertical phase class exists). Lean ✓ Stmt Under the hypotheses of Theorem 21.4.76, the retained vertical decomposition has at least one sector and hence at least one phase class.

Proof. ✓ Proof The unit-modulus weight witness in the horizontal canonical form, together with left canonicity of its representative, makes the doubled-index tensor nonzero. Hence its vertical reshaping $\widetilde{M}$ is nonzero. If the retained decomposition had no sector, its literal reconstruction formula would make every letter of $\widetilde{M}$ zero, a contradiction. Passing from the nonempty finite set of retained sectors to its phase partition gives a retained representative. $\square$

Theorem 21.4.81 (Nonvanishing of the vertical reshaping). [Lean](#) [✓ Stmt](#) *If an MPO tensor M is in horizontal canonical form, then its vertical reshaping is nonzero:*

$$\widetilde{M} \neq 0.$$

This is the nonvanishing observation used in [CPGSV16, Proposition 4.13, lines 1895–1902].

Proof. [✓ Proof](#) Write the doubled-index tensor in horizontal canonical form as $\widehat{M} = GSG^{-1}$, where S is the weighted direct sum of its basis tensors. Choose a copy whose weight has unit modulus. Its basis tensor cannot have every letter equal to zero, since left canonicity gives

$$\sum_i (A^i)^\dagger A^i = \mathbb{1}.$$

The corresponding nonzero weight and nonzero letter give $S \neq 0$, hence $\widehat{M} \neq 0$. If $\widetilde{M} = 0$, the bijection between its vertical index and the two horizontal bond indices would give $\widehat{M} = 0$, a contradiction. $\square$

Theorem 21.4.82 (Algebraic BNT from a grouped vertical decomposition). [Lean](#) [✓ Stmt](#) *Let M be in horizontal canonical form and put $T = \widetilde{M}$. Suppose that B_k has positive bond dimension d_k , that $\mu_k \in \mathbb{C}$, and that $V_k : \mathbb{C}^{d_k} \rightarrow \mathbb{C}^d$ is an isometry for $0 \leq k < r$. Assume, for every vertical letter v , that*

$$V_k^\dagger T_v = (\mu_k B_k^v) V_k^\dagger, \quad T_v = \sum_{k=0}^{r-1} V_k (\mu_k B_k^v) V_k^\dagger.$$

Partition the tensors B_k into matrix-product-vector phase classes and choose one representative A_α from each class. If $\{A_\alpha\}_\alpha$ is a basis of normal tensors in the spectral sense for the weighted direct sum

$$A_{\text{ret}} = \bigoplus_{k=0}^{r-1} \mu_k B_k,$$

then $\{A_\alpha\}_\alpha$ is an algebraic basis of normal tensors for T .

This is the basis-of-normal-tensors conclusion in [CPGSV16, Proposition 4.13, lines 1863–1921], stated for the grouped decomposition before the final coisometry is formed.

Proof. [✓ Proof](#) The reconstruction and Theorem 21.4.81 imply $r > 0$, so the phase partition has a representative A_{α_0} of positive bond dimension d_{α_0} .

Put $C_k^v = \mu_k B_k^v$. Induction on a word w gives

$$V_k^\dagger T^w = C_k^w V_k^\dagger.$$

For a nonempty word $w = (v, w')$, reconstruction of its first letter, cyclicity of trace, and $V_k^\dagger V_k = \mathbb{1}$ therefore give

$$\begin{aligned} \text{tr}(T^w) &= \sum_k \text{tr}(V_k C_k^v V_k^\dagger T^{w'}) \\ &= \sum_k \text{tr}(C_k^v C_k^{w'}) = \sum_k \mu_k^{|w|} \text{tr}(B_k^w) = V^{(|w|)}(A_{\text{ret}})_w. \end{aligned}$$

Hence the spectral BNT spanning identity for A_{ret} gives the algebraic spanning identity for T at every positive length. At length zero, assign coefficient d/d_{α_0} to A_{α_0} and zero to the other representatives; then

$$d = \frac{d}{d_{\alpha_0}} d_{\alpha_0}.$$

Spectral normality implies algebraic normality for every representative, and the eventual linear independence assertion is unchanged. These are the three algebraic BNT conditions. $\square$

Theorem 21.4.83 (Horizontal canonical form implies vertical canonical form). [Lean](#) [✓ Stmt](#) *A tensor in horizontal canonical form that generates matrix product density operators is also in canonical form in the vertical direction: there exist a basis of normal tensors $\{M_\alpha\}_\alpha$ for the vertically viewed tensor $\widetilde{M}$, positive diagonal matrices μ_α , and a coisometry U from the physical space onto the retained nonzero sectors. Writing $B = \bigoplus_\alpha \mu_\alpha \otimes M_\alpha$, one has*

$$UU^\dagger = \mathbb{1}, \quad U\widetilde{M}U^\dagger = B, \quad \widetilde{M} = U^\dagger BU.$$

This is [CPGSV16, Proposition 4.13].

Proof. [✓ Proof](#) Apply Theorem 21.4.76 once and keep its representatives M_α , multiplicities r_α , exact positive weights $\mu_{\alpha,q}$, and physical sector maps. For these same witnesses, Theorem 21.4.82 proves that $\{M_\alpha\}_\alpha$ is a BNT for $\widetilde{M}$.

Theorem 21.4.63 absorbs each normalized bond gauge into its physical map and gives maps $W_{\alpha,q}$ such that

$$W_{\alpha,q}^\dagger W_{\alpha,q} = \mathbb{1}, \quad W_{\alpha,q}^\dagger W_{\beta,p} = 0 \quad \text{if } (\alpha, q) \neq (\beta, p),$$

$$\widetilde{M}_v W_{\alpha,q} = W_{\alpha,q}(\mu_{\alpha,q} M_\alpha^v), \quad \widetilde{M}_v = \sum_\alpha \sum_q W_{\alpha,q}(\mu_{\alpha,q} M_\alpha^v) W_{\alpha,q}^\dagger.$$

The canonical equivalence between the dependent disjoint union $\coprod_\alpha \{0, \dots, r_\alpha - 1\}$ and its standard finite index set reindexes the nested reconstruction sum without changing any weight or sector. The hypotheses of Theorem 21.4.79 now hold, and that theorem gives the vertical canonical form of M . $\square$

Chapter 22

Pure States: Renormalization of Matrix Product States

This chapter develops renormalization fixed points for matrix product vectors. It compares physical two-site blocking, idempotence of the transfer map, and zero correlation length, and then gives the isometry canonical form of the fixed points. The organization follows the pure-state analysis of [CPGSV16, Section 3].

22.1 Physical blocking and transfer idempotence

Definition 22.1.1 (Physical blocking isometry). Lean ✓ Stmt An MPS tensor A is a *renormalization fixed point* if there is an isometry $V: \mathbb{C}^d \rightarrow \mathbb{C}^{d^2}$, with coefficients $V_{(i_1, i_2), j}$ and $V^\dagger V = 1$, such that, for all physical indices i_1, i_2 ,

$$A^{i_1} A^{i_2} = \sum_{j=0}^{d-1} V_{(i_1, i_2), j} A^j. \quad (22.1)$$

This is the relation $AA = A$ in [CPGSV16, Definition 3.2].

Definition 22.1.2 (Idempotent transfer-map criterion). Lean ✓ Stmt The transfer map of A satisfies the idempotence criterion when $\mathcal{E}_A \circ \mathcal{E}_A = \mathcal{E}_A$.

Theorem 22.1.3 (A physical blocking isometry gives transfer idempotence). Lean ✓ Stmt Suppose there exists an isometry $V: \mathbb{C}^d \rightarrow \mathbb{C}^{d^2}$, written as coefficients $V_{(i_1, i_2), j}$, such that

$$A^{i_1} A^{i_2} = \sum_{j=0}^{d-1} V_{(i_1, i_2), j} A^j \quad (22.2)$$

for all $i_1, i_2 \in \{0, \dots, d-1\}$. Then $\mathcal{E}_A^2 = \mathcal{E}_A$.

Proof. ✓ Proof Substituting the decomposition (22.2) into the double Kraus sum gives, for every

bond matrix X ,

$$\begin{aligned}
\mathcal{E}_A^2(X) &= \sum_{i_1, i_2} (A^{i_1} A^{i_2}) X (A^{i_1} A^{i_2})^\dagger \\
&= \sum_{j, k} \left(\sum_{i_1, i_2} V_{(i_1, i_2), j} \overline{V_{(i_1, i_2), k}} \right) A^j X (A^k)^\dagger \\
&= \sum_{j, k} (V^\dagger V)_{j, k} A^j X (A^k)^\dagger = \sum_{j, k} \delta_{j, k} A^j X (A^k)^\dagger \\
&= \sum_j A^j X (A^j)^\dagger = \mathcal{E}_A(X).
\end{aligned}$$

Hence $\mathcal{E}_A^2 = \mathcal{E}_A$. $\square$

Theorem 22.1.4 (Kraus-isometry criterion for transfer idempotence). [Lean](#) [✓ Stmt](#) *The transfer map of A is idempotent if and only if there exists an isometry $V : \mathbb{C}^d \rightarrow \mathbb{C}^{d^2}$, written as coefficients $V_{(i_1, i_2), j}$ with $V^\dagger V = 1$, such that*

$$A^{i_1} A^{i_2} = \sum_{j=0}^{d-1} V_{(i_1, i_2), j} A^j \quad (22.3)$$

for all $i_1, i_2 \in \{0, \dots, d-1\}$.

Proof. [✓ Proof](#) The backward implication is Theorem 22.1.3. For the forward implication, expanding $\mathcal{E}_A^2 = \mathcal{E}_A$ as a Kraus sum gives, for every X ,

$$\sum_{i_1, i_2} (A^{i_1} A^{i_2}) X (A^{i_1} A^{i_2})^\dagger = \sum_j A^j X (A^j)^\dagger.$$

Theorem 16.4.10 then supplies the isometry V relating the two Kraus families. $\square$

Theorem 22.1.5 (Physical and transfer-map fixed-point criteria). [Lean](#) [✓ Stmt](#) *The transfer map of an MPS tensor is idempotent if and only if the tensor has a physical blocking isometry.*

Proof. [✓ Proof](#) This is Theorem 22.1.4, with the physical blocking relation written as Definition 22.1.1. $\square$

Theorem 22.1.6 (Limits of the renormalization flow). *A tensor appears as a limit of the two-site renormalization flow if and only if it has a physical blocking isometry. This is [CPGSV16, Theorem 3.1].*

$$\begin{aligned}
A^{i_1} A^{i_2} &= \sum_j V_j^{i_1 i_2} A^j \\
\sum_{i_1, i_2} (V^\dagger)_{i_1 i_2}^j A^{i_1} A^{i_2} &= A^j
\end{aligned}$$

Figure 22.1: The fusion identity of the renormalization fixed-point characterization: two contracted copies of A equal a single copy of A with the isometry V merging the summed index j into the physical pair (i_1, i_2) , which is equation (22.3); conversely, applying $V^\dagger$ to the two physical legs of the contracted pair recovers the single tensor. The source draws the isometry as U [CPGSV16, Theorem 3.1, lines 396–412].

22.2 Zero-correlation-length conditions

Definition 22.2.1 (Single-block local orthogonality convention). [Lean](#) [✓ Stmt](#) For the single-block convention used here, local orthogonality means

$$\mathcal{E}_A^2 = \mathcal{E}_A.$$

The source local-orthogonality condition for a BNT family is instead the collection of mixed-sector equations for distinct components,

$$\mathcal{E}_{j,j'} = 0.$$

Thus this definition records the one-block idempotence convention.

Definition 22.2.2 (BNT local orthogonality). [Lean](#) [✓ Stmt](#) Let $(A_j)_j$ be the BNT components of a tensor in canonical form. The family is *locally orthogonal* when, for every pair of distinct components,

$$\mathcal{E}_{j,j'} = \sum_i A_j^i \otimes \overline{A_{j'}^i} = 0.$$

This is the mixed-sector condition of [CPGSV16, Definition 3.5].

Definition 22.2.3 (Virtual-insertion BNT zero correlation length). [Lean](#) [✓ Stmt](#) Let $(A_j)_j$ be a basis of normal tensors for A . The BNT zero-correlation-length condition is that

$$\text{CID}(A)$$

and, for every pair of distinct BNT components,

$$\mathcal{E}_{j,j'} = 0.$$

This virtual-insertion condition is not [CPGSV16, Definition 3.6], which quantifies over physical observables on all disjoint regions.

Definition 22.2.4 (Positive-gap BNT zero correlation length). [Lean](#) [✓ Stmt](#) A family $(A_j)_j$ of MPS tensors has positive-gap BNT zero correlation length for A when $(A_j)_j$ is a basis of normal tensors for A in the sense of Definition 10.1.2, its components are locally orthogonal, and the physical correlations of A are independent of separation whenever both complementary gaps are positive. Adjacent regions are not included.

Lemma 22.2.5 (Positive-gap BNT zero correlation length, unfolded). [Lean](#) [✓ Stmt](#) *Positive-gap BNT zero correlation length is equivalent to the conjunction of the BNT relation, positive-gap physical distance independence, and BNT local orthogonality.*

Proof. [✓ Proof](#) This is the defining conjunction. □

Definition 22.2.6 (BNT zero correlation length). [Lean](#) [✓ Stmt](#) A BNT family has zero correlation length when the associated matrix-product vectors have correlations independent of distance for all disjoint physical regions and the BNT components are locally orthogonal. This is [CPGSV16, Definition 3.6].

Theorem 22.2.7 (Unequal raw weights obstruct the ZCL–RFP equivalence). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Let P be the canonical decomposition with one scalar basis tensor*

$C^0 = (1)$, two one-dimensional copies, and respective weights 1 and $1/2$. Write $A = \mathcal{A}(P)$ for its assembled tensor. Relative to these sector coordinates, A^0 has block form

$$\begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}.$$

The weights $(1, 1/2)$ satisfy the raw-weight BNT canonical-form normalization, and the resulting canonical assembly is exactly A . Moreover,

$$V^{(N)}(A) = 1 + 2^{-N}.$$

The BNT has one component, so local orthogonality is vacuous. Since the physical space has dimension one, all physical correlations are independent of separation. Thus A has zero correlation length in the sense of Definition 22.2.6, but A is not a renormalization fixed point. Consequently [CPGSV16, Theorem 3.8] is false under its stated raw-weight BNT normalization.

Proof. ✓ Proof The $(0, 0)$ and $(1, 1)$ entries of the one-letter blocking equation $(A^0)^2 = vA^0$ would give

$$1 = v \cdot 1 \quad \text{and} \quad \frac{1}{4} = v \cdot \frac{1}{2}.$$

The first equality forces $v = 1$, while substitution in the second gives $1/4 = 1/2$, a contradiction. Hence no such scalar exists. The physical distance-independence assertion follows because every observable on a one-letter physical space is scalar. □

Lemma 22.2.8 (Source BNT zero correlation length implies its positive-gap form). Lean ✓ Stmt
If a BNT family has zero correlation length for all disjoint physical regions, then it has positive-gap BNT zero correlation length.

Proof. ✓ Proof Restrict the physical distance-independence condition to configurations in which both complementary gaps are positive. The BNT relation and local orthogonality are unchanged. □

Theorem 22.2.9 (Direct-sum RFP implies positive-gap BNT zero correlation length). Lean ✓ Stmt
Let $(A_j)_j$ be nonzero-dimensional, irreducible, left-canonical, pairwise gauge-phase-distinct BNT components. If the direct-sum tensor $A = \bigoplus_j A_j$ satisfies $\mathcal{E}_A^2 = \mathcal{E}_A$, then A has positive-gap physical CID and, for all $j \neq j'$,

$$\mathcal{E}_{j,j'} = 0.$$

Thus this explicit direct-sum representative has positive-gap BNT zero correlation length. This does not include the adjacent-region cases in [CPGSV16, Theorem 3.8].

Proof. ✓ Proof Idempotence implies positive-gap physical CID by Theorem 22.5.10. On each off-diagonal bond block, idempotence of $\mathcal{E}_A$ gives

$$\mathcal{E}_{j,j'} \circ \mathcal{E}_{j,j'} = \mathcal{E}_{j,j'}.$$

Distinct irreducible left-canonical blocks satisfy $\rho(\mathcal{E}_{j,j'}) < 1$. Since the spectrum of an idempotent is contained in $\{0, 1\}$,

$$\mathcal{E}_{j,j'}^2 = \mathcal{E}_{j,j'}, \quad \rho(\mathcal{E}_{j,j'}) < 1 \quad \implies \quad \mathcal{E}_{j,j'} = 0.$$

□

22.3 Normal tensors and isometry canonical forms

Theorem 22.3.1 (RFP normal tensor is injective). [Lean](#) [✓ Stmt](#) *If A is a nonzero normal RFP tensor, then A is injective.*

Proof. [✓ Proof](#) The RFP condition $\mathcal{E}_A^2 = \mathcal{E}_A$ gives coefficients $V_{i_1 i_2, j}$ such that

$$A^{i_1} A^{i_2} = \sum_j V_{i_1 i_2, j} A^j.$$

Induction on the word length therefore gives

$$A^{i_1} \dots A^{i_n} \in \text{span}\{A^j : 0 \leq j < d\} \quad (n \geq 1).$$

By normality, the words of some length N span $M_D(\mathbb{C})$. If $N \geq 1$, the preceding inclusion shows that the one-site span is already $M_D(\mathbb{C})$. If $N = 0$, then $\text{span}\{\mathbf{1}\} = M_D(\mathbb{C})$, and any nonzero A^j spans this space. Thus A is injective in either case. $\square$

Theorem 22.3.2 (Left-canonical normal RFP tensor is injective). [Lean](#) [✓ Stmt](#) *Assume $D > 0$. Let A be a normal renormalization fixed point such that*

$$\sum_i (A^i)^\dagger A^i = \mathbf{1}.$$

Then A is injective. This is the left-canonical injectivity step in [CPGSV16, Appendix B].

Proof. [✓ Proof](#) The normalization $\sum_i (A^i)^\dagger A^i = \mathbf{1}$ forces at least one matrix A^i to be nonzero, since otherwise the left-hand side would vanish. Hence A is a nonzero normal renormalization fixed point, and Theorem 22.3.1 applies. $\square$

Theorem 22.3.3 (Unitary diagonal fixed point for a left-canonical normal RFP). [Lean](#) [✓ Stmt](#) *Assume $D > 0$. Let A be a normal renormalization fixed point such that*

$$\sum_i (A^i)^\dagger A^i = \mathbf{1}.$$

Then there exist a unitary U and a diagonal positive definite matrix Λ such that, for $B^i := U^\dagger A^i U$,

$$\sum_i (B^i)^\dagger B^i = \mathbf{1} \quad \text{and} \quad \mathcal{E}_B(\Lambda) = \Lambda.$$

This is the diagonal fixed-point reduction in [CPGSV16, Appendix B].

Proof. [✓ Proof](#) By Theorem 22.3.2, the tensor A is injective. By Theorem 4.2.5, its transfer map $\mathcal{E}_A$ is irreducible, and hence A is irreducible as a tensor. Applying Theorem 9.2.1.9 gives a unitary conjugate B for which the trace-preserving normalization remains valid and a diagonal positive definite matrix Λ with $\mathcal{E}_B(\Lambda) = \Lambda$. $\square$

Theorem 22.3.4 (Rank-one classification for RFP transfer maps). [Lean](#) [✓ Stmt](#) *Let A be an injective left-canonical RFP tensor with positive-definite fixed point ρ of the transfer map $\mathcal{E}_A$. Then $\mathcal{E}_A = P_\rho$, where $P_\rho(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$ is the rank-one fixed-point projection.*

Proof. [✓ Proof](#) The exponential convergence bound gives $\|\mathcal{E}_A^n(X) - P_\rho(X)\| \leq C(1 - \delta)^n \|X\|$. Idempotence of $\mathcal{E}_A$ implies $\mathcal{E}_A^{n+1} = \mathcal{E}_A$ for all $n \geq 0$, so $\|\mathcal{E}_A(X) - P_\rho(X)\| \leq C(1 - \delta)^{n+1} \|X\|$ for all n . Taking $n \rightarrow \infty$ gives $\mathcal{E}_A = P_\rho$. $\square$

Theorem 22.3.5 (Full structural form for RFP tensors). [Lean](#) [✓ Stmt](#) *A normal tensor A in canonical form II that is a renormalization fixed point admits a decomposition $A^i = X\Lambda U^i X^{-1}$, where Λ is diagonal positive, $\sum_i (U^i)^\dagger U^i = I$, and*

$$\sum_i \overline{(U^i)_{\alpha,\beta}} (U^i)_{\alpha',\beta'} = D^{-1} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

Proof. [✓ Proof](#) After the diagonal fixed-point reduction, write the conjugated tensor as B and its positive diagonal fixed point as ρ . The rank-one classification gives

$$\mathcal{E}_B(Y) = P_\rho(Y) = \frac{\text{tr}(Y)}{\text{tr}(\rho)} \rho.$$

Set

$$\Lambda_\alpha = \sqrt{\frac{D\rho_{\alpha,\alpha}}{\text{tr}(\rho)}}, \quad E_{\alpha,\beta} = D^{-1/2} e_{\alpha,\beta}, \quad K_{\alpha,\beta} = \Lambda E_{\alpha,\beta}.$$

Direct multiplication of matrix units yields

$$\sum_{\alpha,\beta} K_{\alpha,\beta} Y K_{\alpha,\beta}^\dagger = P_\rho(Y).$$

Kraus freedom therefore gives an isometry V such that $B^i = \sum_{\alpha,\beta} V_{i,(\alpha,\beta)} K_{\alpha,\beta}$. With $U^i = \sum_{\alpha,\beta} V_{i,(\alpha,\beta)} E_{\alpha,\beta}$, one has $B^i = \Lambda U^i$ and

$$U_{\alpha,\beta}^i = D^{-1/2} V_{i,(\alpha,\beta)}, \quad \sum_i \overline{U_{\alpha,\beta}^i} U_{\alpha',\beta'}^i = D^{-1} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

Conjugating back gives $A^i = X\Lambda U^i X^{-1}$. □

Theorem 22.3.6 (Structural form with diagonal square-sum identity). [Lean](#) [✓ Stmt](#) *The same decomposition as the full structural form (Theorem 22.3.5) additionally records that the diagonal weights satisfy $\sum_\alpha \Lambda_\alpha^2 = D$. This square-sum identity is the trace-normalization seed: the explicit weights are $\Lambda_\alpha = \sqrt{D\rho_{\alpha,\alpha}/\text{tr}\rho}$, and $\sum_\alpha \rho_{\alpha,\alpha} = \text{tr}\rho$ for the diagonal fixed point ρ .*

Proof. [✓ Proof](#) The diagonal fixed-point reduction and the rank-one classification of injective left-canonical renormalization fixed points produce the diagonal weights $\Lambda_\alpha = \sqrt{D\rho_{\alpha,\alpha}/\text{tr}\rho}$, so $\Lambda_\alpha^2 = D\rho_{\alpha,\alpha}/\text{tr}\rho$. Summing over α and using $\sum_\alpha \rho_{\alpha,\alpha} = \text{tr}\rho$ gives $\sum_\alpha \Lambda_\alpha^2 = D$. □

Theorem 22.3.7 (Unit pair-index form for RFP tensors). [Lean](#) [✓ Stmt](#) *A normal tensor A in canonical form II that is a renormalization fixed point admits a decomposition $A^i = X\Lambda U^i X^{-1}$, where Λ is diagonal positive and*

$$\sum_i \overline{(U^i)_{\alpha,\beta}} (U^i)_{\alpha',\beta'} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

This is the unit pair-index convention of [CPGSV16, Section 3]. The statement still does not impose the trace-normalization $\text{tr}(\Lambda) = 1$.

Proof. [✓ Proof](#) Apply Theorem 22.3.11, which gives $A^i = X\sqrt{\Lambda} U^i X^{-1}$ with U already a unit pair-index isometry. Set $\tilde{\Lambda}_\alpha = \sqrt{\Lambda_\alpha}$, so that $A^i = X\tilde{\Lambda} U^i X^{-1}$ with $\tilde{\Lambda}$ diagonal positive; the decomposition and the pair-index condition are immediate. □

Theorem 22.3.8 (Per-block isometry canonical form). [Lean](#) [✓ Stmt](#) When each block A_k of a multi-block tensor is a normal, left-canonical renormalization fixed point, that block admits the isometry decomposition $A_k^i = X_k \Lambda_k U_k^i X_k^{-1}$, with X_k invertible, Λ_k diagonal positive, and $U_k = (U_k^i)$ satisfies $\sum_i (U_k^i)^\dagger U_k^i = I$ and the pair-index orthonormality condition

$$\sum_i \overline{(U_k^i)_{\alpha,\beta}} (U_k^i)_{\alpha',\beta'} = D_k^{-1} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

This is the blockwise form of the isometry canonical form (Theorem 22.3.5); see [CPGSV16, Section 3]. The source additionally imposes the normalization $\text{tr}(\Lambda_k) = 1$; the statement here gives positive Λ_k without it. That normalization is genuine rather than a conjugation gauge, since rescaling $\Lambda_k \mapsto \Lambda_k / \text{tr}(\Lambda_k)$ factors out as an overall scalar on A_k^i (conjugation by X_k preserves the scale); the statement is thus the un-normalized isometry form. Scope restriction (source isometry). Corollary III.cor3 also invokes the joint isometry condition

$$\sum_i (U_k^i)_{\alpha,\beta} \overline{(U_\ell^i)_{\alpha',\beta'}} = \delta_{k,\ell} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

The theorem above records the contracted per-block condition $\sum_i (U_k^i)^\dagger U_k^i = I$ and the diagonal pair-index equation with right-hand side $D_k^{-1} \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}$. It does not impose the trace-normalization $\text{tr}(\Lambda_k) = 1$, and it does not include the cross-block equations for $k \neq \ell$.

Proof. [✓ Proof](#) Apply Theorem 22.3.5 to each block. □

Theorem 22.3.9 (Per-block unit pair-index form). [Lean](#) [✓ Stmt](#) Under the hypotheses of Theorem 22.3.8, each block admits a decomposition $A_k^i = X_k \Lambda_k U_k^i X_k^{-1}$ with Λ_k diagonal positive and

$$\sum_i \overline{(U_k^i)_{\alpha,\beta}} (U_k^i)_{\alpha',\beta'} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

This is a blockwise unit pair-index comparison. It still does not impose the source trace-normalization of Λ_k or the cross-block orthogonality equations for distinct blocks.

Proof. [✓ Proof](#) Apply Theorem 22.3.7 to each block. □

Definition 22.3.10 (Isometry canonical form). [Lean](#) [✓ Stmt](#) A normal tensor A is in *isometry canonical form* when there are an invertible X , a positive diagonal weight Λ with the trace normalization $\sum_\alpha \Lambda_\alpha = 1$, and a residual tensor U satisfying the unit pair-index isometry

$$\sum_i \overline{(U^i)_{\alpha,\beta}} (U^i)_{\alpha',\beta'} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'},$$

such that

$$A^i = X \sqrt{\Lambda} U^i X^{-1}.$$

The square root dresses the unit isometry because the reference tensor of [CPGSV16, Section 3] is $\widehat{A}_{\alpha,\beta}^{(\alpha',\beta')} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'} \sqrt{\Lambda_\alpha}$; the normalization $\text{tr}(\Lambda) = 1$ is imposed on Λ , equivalently on the normalized diagonal fixed point $\rho = \text{diag}(\Lambda)$ of the transfer map. The displayed source form writes $A^i = X \Lambda U^i X^{-1}$ with the bare diagonal; that pairing forces a non-unit residual, so the consistent decomposition keeps the square root $\sqrt{\Lambda}$ while the trace normalization stays on Λ . This records the diagonal $j = j'$ case of the source isometry condition.

Theorem 22.3.11 (Trace-normalized isometry canonical form for RFP tensors). [Lean](#) [✓ Stmt](#) A normal tensor in canonical form Π that is a renormalization fixed point is in isometry canonical form: it admits a decomposition $A^i = X\sqrt{\Lambda}U^iX^{-1}$ with Λ diagonal, positive, trace-normalized ($\sum_{\alpha}\Lambda_{\alpha} = 1$, the source condition $\text{tr}(\Lambda) = 1$), and U a unit pair-index isometry. The diagonal weight is $\Lambda_{\alpha} = \rho_{\alpha,\alpha}/\text{tr}\rho$, the normalized diagonal fixed point of the transfer map; trace-normalization is then the trace identity $\sum_{\alpha}\rho_{\alpha,\alpha} = \text{tr}\rho$. This is the single-block normal-tensor case of the structural characterization of pure-state renormalization fixed points in [CPGSV16, Section 3], recovering the diagonal part of the isometry condition there.

Proof. [✓ Proof](#) The structural form with the square-sum identity (Theorem 22.3.6) supplies a positive diagonal weight $\tilde{\Lambda}$ with $\sum_{\alpha}\tilde{\Lambda}_{\alpha}^2 = D$. Rescaling to the unit pair-index convention by $U^i \mapsto \sqrt{D}U^i$ and setting $\Lambda_{\alpha} = (\tilde{\Lambda}_{\alpha}/\sqrt{D})^2$ gives $\sum_{\alpha}\Lambda_{\alpha} = D^{-1}\sum_{\alpha}\tilde{\Lambda}_{\alpha}^2 = D^{-1} \cdot D = 1$, while $\sqrt{\Lambda_{\alpha}} = \tilde{\Lambda}_{\alpha}/\sqrt{D}$ recovers the decomposition $A^i = X\sqrt{\Lambda}U^iX^{-1}$. $\square$

$$A = X\sqrt{\Lambda}U X^{-1}$$

Figure 22.2: The isometry canonical form of a normal renormalization fixed point: $A^i = X\sqrt{\Lambda}U^iX^{-1}$, with the diagonal weight $\sqrt{\Lambda}$ on the virtual chain and the isometry U emitting the physical index [CPGSV16, Appendix B, lines 1275–1289].

Lemma 22.3.12 (Reference-tensor transfer identity for a unit pair-index isometry). [Lean](#) [✓ Stmt](#) Let U be a family of bond matrices satisfying the unit pair-index isometry condition

$$\sum_i \overline{(U^i)_{\alpha,\beta}} (U^i)_{\alpha',\beta'} = \delta_{\alpha,\alpha'} \delta_{\beta,\beta'}.$$

Then for every bond matrix Z ,

$$\sum_i U^i Z (U^i)^{\dagger} = \text{tr}(Z) I.$$

Proof. [✓ Proof](#) Reading off the (x, y) entry, the pair-index condition gives $\sum_i (U^i)_{x,\alpha} \overline{(U^i)_{y,\beta}} = \delta_{x,y} \delta_{\alpha,\beta}$, so the entry collapses to $\delta_{x,y} \sum_{\alpha} Z_{\alpha,\alpha} = \delta_{x,y} \text{tr}(Z)$, which is the (x, y) entry of $\text{tr}(Z) I$. $\square$

Theorem 22.3.13 (Isometry canonical form implies the renormalization fixed-point property). [Lean](#) [✓ Stmt](#) A tensor in isometry canonical form is a renormalization fixed point. This is the backward direction of the structural characterization of pure-state renormalization fixed points in [CPGSV16, Section 3], which the source states as immediate.

Proof. [✓ Proof](#) Write $A^i = X\sqrt{\Lambda}U^iX^{-1}$. Substituting into the transfer map and pulling the index-independent factors $X\sqrt{\Lambda}$ and $\sqrt{\Lambda}X^{\dagger}$ outside the sum, the unit pair-index isometry identity (Lemma 22.3.12) gives the rank-one form

$$\mathcal{E}_A(Y) = \text{tr}(X^{-1}Y(X^{-1})^{\dagger}) R, \quad R = X \Lambda X^{\dagger}.$$

Idempotence then reduces to $\text{tr}(X^{-1}R(X^{-1})^{\dagger}) = \text{tr}(\Lambda) = 1$, the trace normalization, so $\mathcal{E}_A \circ \mathcal{E}_A = \mathcal{E}_A$. $\square$

Theorem 22.3.14 (Per-block trace-normalized isometry canonical form). Lean ✓ Stmt *Each block of a multi-block tensor whose blocks are normal, left-canonical renormalization fixed points is in isometry canonical form, with a trace-normalized diagonal weight and a unit pair-index isometry. This is the single-block trace-normalized form applied to each block; it omits the cross-block orthogonality between distinct normal-tensor blocks.*

Proof. ✓ Proof Apply Theorem 22.3.11 to each block. □

$$A = X \left(\bigoplus_j \mu_j \otimes \sqrt{\Lambda_j} U_j \right) X^{-1}$$

Figure 22.3: The per-block canonical form $A^i = \bigoplus_{j,q} \mu_{j,q} X_{j,q} \sqrt{\Lambda_j} U_j^i X_{j,q}^{-1}$: the block-index lines j and q thread the gauge, the diagonal weight, the isometry, and the coefficient tensor μ . The picture is per-block, matching the entry above; the cross-block physical orthogonality of [CPGSV16, Section 3.4, lines 543–561] is not displayed.

22.4 Direct sums and residual isometries

Lemma 22.4.1 (Block decomposition of the direct-sum transfer map). Lean ✓ Stmt *Let $(B_k)_k$ be a family of tensors and let $\bigoplus_k B_k$ be their direct sum on the total bond space. For every bond matrix X and every pair of blocks, the (j, j') bond block of the transfer sum*

$$\sum_i \left(\bigoplus_k B_k^i \right) X \left(\bigoplus_k B_k^i \right)^\dagger$$

equals $\mathcal{E}_{j,j'}$ applied to the (j, j') bond block of X .

Proof. ✓ Proof A block-diagonal factor connects only matching blocks, so the (j, j') block of the product is

$$\left(\sum_i \left(\bigoplus_k B_k^i \right) X \left(\bigoplus_k B_k^i \right)^\dagger \right)_{j,j'} = \sum_i B_j^i X_{j,j'} (B_{j'}^i)^\dagger = \mathcal{E}_{j,j'}(X_{j,j'}).$$

□

Lemma 22.4.2 (Pairwise idempotence of the block transfer sum). Lean ✓ Stmt *Let $(B_k)_k$ be a family of tensors. If every mixed transfer operator $\mathcal{E}_{j,j'}$ is idempotent, then the block transfer sum $\mathcal{S}_B$ is idempotent:*

$$\mathcal{S}_B(\mathcal{S}_B(X)) = \mathcal{S}_B(X).$$

Proof. ✓ Proof By Lemma 22.4.1, the (j, j') block satisfies

$$(\mathcal{S}_B^2(X))_{j,j'} = \mathcal{E}_{j,j'}^2(X_{j,j'}) = \mathcal{E}_{j,j'}(X_{j,j'}) = (\mathcal{S}_B(X))_{j,j'}.$$

Equality on every block gives the asserted equality on the total bond space. □

Theorem 22.4.3 (Pairwise mixed-transfer criterion for a direct sum). Lean ✓ Stmt *Suppose every bond dimension is positive. The transfer map of $\bigoplus_k B_k$ is idempotent if and only if, for every pair j, j' , the mixed transfer operator is idempotent:*

$$\mathcal{E}_{\bigoplus_k B_k}^2 = \mathcal{E}_{\bigoplus_k B_k} \iff \mathcal{E}_{j,j'}^2 = \mathcal{E}_{j,j'}.$$

Proof. ✓ Proof By Lemma 22.4.1, the (j, j') block of the direct-sum transfer map depends only on the (j, j') block of its argument and is obtained by applying $\mathcal{E}_{j, j'}$. Thus pairwise idempotence gives

$$(\mathcal{E}_{\oplus_k B_k}^2(X))_{j, j'} = \mathcal{E}_{j, j'}^2(X_{j, j'}) = \mathcal{E}_{j, j'}(X_{j, j'}) = (\mathcal{E}_{\oplus_k B_k}(X))_{j, j'}$$

for every pair of blocks, and hence gives idempotence on the total bond space, as in Lemma 22.4.2. Conversely, every matrix M of the appropriate rectangular size occurs as the (j, j') block of some Y on the total bond space. Applying Lemma 22.4.1 and whole-tensor idempotence gives

$$\mathcal{E}_{j, j'}^2(M) = (\mathcal{E}_{\oplus_k B_k}^2(Y))_{j, j'} = (\mathcal{E}_{\oplus_k B_k}(Y))_{j, j'} = \mathcal{E}_{j, j'}(M).$$

□

Lemma 22.4.4 (Scaling of a rectangular mixed transfer operator). Lean ✓ Stmt For complex scalars c, e and tensors A, B of possibly different bond dimensions,

$$\mathcal{E}_{cA, eB} = c\bar{e} \mathcal{E}_{A, B}.$$

Proof. ✓ Proof Conjugate transposition changes the second scalar to its complex conjugate:

$$\sum_i (cA^i)X(eB^i)^\dagger = c\bar{e} \sum_i A^i X(B^i)^\dagger.$$

□

Theorem 22.4.5 (Phase coherence for literal repeated blocks). Lean ✓ Stmt Let A have positive bond dimension and a nonzero idempotent transfer map. Let μ_q have unit modulus. If the literal bond direct sum $\bigoplus_q \mu_q A$ has an idempotent transfer map, then for every pair q, q' ,

$$\mu_q = \mu_{q'}.$$

Proof. ✓ Proof The pairwise criterion (Theorem 22.4.3) gives that $\mathcal{E}_{\mu_q A, \mu_{q'} A}$ is idempotent. Lemma 22.4.4 and $\mathcal{E}_{A, A} = \mathcal{E}_A$ give

$$\mathcal{E}_{\mu_q A, \mu_{q'} A} = \mu_q \overline{\mu_{q'}} \mathcal{E}_{A, A} = \mu_q \overline{\mu_{q'}} \mathcal{E}_A.$$

Hence $(\mu_q \overline{\mu_{q'}}) \mathcal{E}_A$ is idempotent. Since $\mathcal{E}_A$ is nonzero and $|\mu_q \overline{\mu_{q'}}| = |\mu_q| |\mu_{q'}| = 1$,

$$(\mu_q \overline{\mu_{q'}})^2 \mathcal{E}_A = \mu_q \overline{\mu_{q'}} \mathcal{E}_A \implies \mu_q \overline{\mu_{q'}} (\mu_q \overline{\mu_{q'}} - 1) = 0 \implies \mu_q \overline{\mu_{q'}} = 1.$$

Since $|\mu_{q'}| = 1$ gives $\overline{\mu_{q'}} \mu_{q'} = 1$,

$$\mu_q = \mu_q (\overline{\mu_{q'}} \mu_{q'}) = (\mu_q \overline{\mu_{q'}}) \mu_{q'} = \mu_{q'}.$$

□

Theorem 22.4.6 (Cross-block transfer vanishing for a direct-sum fixed point). Lean ✓ Stmt Let $(A_j)_j$ be a family of distinct irreducible left-canonical blocks, no two of which are gauge-phase equivalent, and suppose the direct sum $\bigoplus_j A_j$ is a renormalization fixed point. Then for every pair of distinct components the mixed transfer operator vanishes,

$$\mathcal{E}_{j, j'} = \sum_i A_j^i \otimes \overline{A_{j'}^i} = 0.$$

This is the cross-block $(\delta_{j, j'})$ content of the isometry condition of [CPGSV16, Theorem III & Corollary III.3] at the level of the normal-tensor blocks; converting it to the residual isometries U_j is a further step left open.

Proof. ✓ Proof By Lemma 22.4.1 the (j, j') bond block of the direct-sum transfer map acts as $\mathcal{E}_{j, j'}$. Whole-tensor idempotence $\mathcal{E}^2 = \mathcal{E}$ therefore restricts to each block as $\mathcal{E}_{j, j'}^2 = \mathcal{E}_{j, j'}$. For distinct irreducible left-canonical blocks, Theorem 7.8.2 gives $\rho_{\text{spec}}(\mathcal{E}_{j, j'}) < 1$ when the bond dimensions agree, because the blocks are not gauge-phase equivalent; Theorem 7.8.4 gives the same inequality when the bond dimensions differ. Consequently, Lemma 7.6.2 gives

$$\mathcal{E}_{j, j'}^2 = \mathcal{E}_{j, j'}, \quad \rho_{\text{spec}}(\mathcal{E}_{j, j'}) < 1 \quad \implies \quad \mathcal{E}_{j, j'} = 0.$$

□

Definition 22.4.7 (Residual isometry family). Lean ✓ Stmt A family of residual tensors $(U_j)_j$ is a *residual isometry family* when it satisfies the source isometry condition

$$\sum_i (U_j^i)_{\alpha, \beta} \overline{(U_{j'}^i)_{\alpha', \beta'}} = \delta_{j, j'} \delta_{\alpha, \alpha'} \delta_{\beta, \beta'} :$$

each block satisfies the within-block pair-index orthonormality, and the cross-block sums between distinct blocks vanish. This is the isometry condition of [CPGSV16, Theorem III], split into its $j = j'$ and $j \neq j'$ cases.

Lemma 22.4.8 (Covariance of the mixed transfer operator). Lean ✓ Stmt For decompositions $A^i = X_A D_A U^i X_A^{-1}$ and $B^i = X_B D_B V^i X_B^{-1}$, the mixed transfer operator of A, B is the mixed transfer operator of the residual tensors U, V conjugated by the outer factors:

$$\mathcal{E}_{A, B}(Y) = X_A D_A \mathcal{E}_{U, V}(X_A^{-1} Y (X_B^{-1})^\dagger) (X_B D_B)^\dagger.$$

Proof. ✓ Proof Substitute the two decompositions into $\mathcal{E}_{A, B}(Y) = \sum_i A^i Y (B^i)^\dagger$ and collect the outer factors, which are independent of the summation index, outside the sum. □

Lemma 22.4.9 (Vanishing of the residual transfer operator). Lean ✓ Stmt With decompositions $A^i = X_A D_A U^i X_A^{-1}$ and $B^i = X_B D_B V^i X_B^{-1}$ whose factors X_A, D_A, X_B, D_B are invertible, if $\mathcal{E}_{A, B} = 0$ then $\mathcal{E}_{U, V} = 0$.

Proof. ✓ Proof The covariance identity Lemma 22.4.8 expresses $\mathcal{E}_{A, B}$ as $\mathcal{E}_{U, V}$ conjugated by the invertible outer factors $X_A D_A$ and $(X_B D_B)^\dagger$, so $\mathcal{E}_{A, B} = 0$ forces $\mathcal{E}_{U, V} = 0$. □

Lemma 22.4.10 (Entrywise residual isometry sum). Lean ✓ Stmt If $\mathcal{E}_{U, V} = 0$, then for all virtual indices

$$\sum_i (U^i)_{\alpha, \beta} \overline{(V^i)_{\alpha', \beta'}} = 0.$$

Proof. ✓ Proof Apply $\mathcal{E}_{U, V} = 0$ to the matrix unit supported at (β, β') and read the (α, α') entry. □

Lemma 22.4.11 (Each block of a direct-sum fixed point is a fixed point). Lean ✓ Stmt If the direct sum $\bigoplus_k B_k$ is a renormalization fixed point, with $\dim_k \geq 1$ for all k , then each block B_j is a renormalization fixed point.

Proof. ✓ Proof The diagonal mixed transfer operator $\mathcal{E}_{j, j}$ is the transfer map of B_j , and whole-tensor idempotence makes it idempotent, which is exactly the renormalization fixed-point condition for B_j . □

Theorem 22.4.12 (Residual-isometry form of the cross-block structure). [Lean](#) [✓ Stmt](#) Let $(A_j)_j$ be a family of normal, irreducible, left-canonical blocks, no two of which are gauge-phase equivalent, with $\dim_j \geq 1$ for all j , and suppose the direct sum $\bigoplus_j A_j$ is a renormalization fixed point. Then there are invertible X_j , positive trace-normalized diagonal weights Λ_j , and residual tensors U_j with

$$A_j^i = X_j \sqrt{\Lambda_j} U_j^i X_j^{-1},$$

such that $(U_j)_j$ is a residual isometry family. This is the residual-isometry content of Corollary III.cor3 [CPGSV16, Corollary III.3] at the level of the normal-tensor blocks.

Proof. [✓ Proof](#) Whole-tensor idempotence makes the diagonal mixed transfer operator of each block its own transfer map, hence each block is a renormalization fixed point; the isometry canonical form (Theorem 22.3.11) supplies the decomposition $A_j^i = X_j \sqrt{\Lambda_j} U_j^i X_j^{-1}$ with the within-block orthonormality, which is the $j = j'$ case. For $j \neq j'$, Theorem 22.4.6 gives $\mathcal{E}_{A_j, A_{j'}} = 0$; by the covariance Lemma 22.4.8 and invertibility of the outer factors $\mathcal{E}_{U_j, U_{j'}} = 0$, and reading the entry of this operator at a matrix unit yields the cross-block sum $\sum_i (U_j^i)_{\alpha, \beta} \overline{(U_{j'}^i)_{\alpha', \beta'}} = 0$. $\square$

Theorem 22.4.13 (A direct sum of isometry-canonical-form blocks is a renormalization fixed point). [Lean](#) [✓ Stmt](#) Let $(B_k)_k$ be a family of tensors, each in isometry canonical form, whose cross-block mixed transfer operators vanish, $\mathcal{E}_{j, j'} = 0$ for $j \neq j'$. Then the direct sum $\bigoplus_k B_k$ is a renormalization fixed point. This is the backward direction of the structural characterization of pure-state renormalization fixed points [CPGSV16, Section 3], in the distinct-blocks case where each normal tensor appears once; the cross-block vanishing is the $j \neq j'$ part of the isometry condition of that characterization, so it belongs to the source isometry form.

Proof. [✓ Proof](#) The direct-sum transfer map decouples block by block: by Lemma 22.4.1 its (j, j') bond block acts as $\mathcal{E}_{j, j'}$ on the (j, j') bond block of the argument. Each diagonal block $\mathcal{E}_{j, j}$ is the transfer map of B_j , which is idempotent because B_j is a renormalization fixed point (Theorem 22.3.13); each off-diagonal block $\mathcal{E}_{j, j'}$ with $j \neq j'$ vanishes by hypothesis. Therefore, for every bond block,

$$[\mathcal{E}_{\bigoplus B}^2(Y)]_{j, j'} = \mathcal{E}_{j, j'}(\mathcal{E}_{j, j'}(Y_{j, j'})) = \mathcal{E}_{j, j'}(Y_{j, j'}) = [\mathcal{E}_{\bigoplus B}(Y)]_{j, j'},$$

so $\mathcal{E}_{\bigoplus B}^2 = \mathcal{E}_{\bigoplus B}$ and the direct sum is a renormalization fixed point. $\square$

Theorem 22.4.14 (Distinct-blocks renormalization fixed-point characterization). [Lean](#) [✓ Stmt](#) Let $(B_k)_k$ be a family of normal, irreducible, left-canonical blocks with $\dim_k \geq 1$ for all k , no two of which are gauge-phase equivalent. Then the direct sum $\bigoplus_k B_k$ is a renormalization fixed point if and only if each block is in isometry canonical form and the mixed transfer operators between distinct blocks vanish:

$$\left(\bigoplus_k B_k \text{ is a RFP} \right) \iff \left(\forall k, B_k \text{ in isometry CF} \right) \wedge \left(\forall j \neq j', \mathcal{E}_{j, j'} = 0 \right).$$

This is the distinct-blocks (multiplicity-one, phase-one) case of the structural characterization of pure-state renormalization fixed points [CPGSV16, Theorem III], combining its forward and backward directions.

Proof. [✓ Proof](#) For the forward direction, whole-tensor idempotence makes each diagonal mixed transfer operator the transfer map of its block (Lemma 22.4.11), so each block is a renormalization

fixed point and hence in isometry canonical form (Theorem 22.3.11), while the off-diagonal operators vanish,

$$\mathcal{E}_{j,j'} = 0 \quad (j \neq j'),$$

by Theorem 22.4.6. Conversely, a direct sum of isometry-canonical-form blocks whose cross-block operators vanish is a renormalization fixed point (Theorem 22.4.13). $\square$

Corollary 22.4.15 (BNT basis direct-sum characterization). Lean Lean ✓ Stmt *Let P be in BNT canonical form, with distinct basis tensors $(B_j)_j$. Then the direct sum containing one copy of each basis tensor is a renormalization fixed point if and only if every B_j is in isometry canonical form and*

$$\mathcal{E}_{j,j'} = 0 \quad (j \neq j').$$

This is the multiplicity-one, phase-one specialization of [CPGSV16, Theorem III and Corollary III.3]. It concerns the direct sum of the basis representatives, not the repeated-copy tensor carrying the weights $\mu_{j,q}$.

Moreover, if the basis direct sum is a renormalization fixed point, there are invertible matrices X_j , positive trace-normalized diagonal weights Λ_j , and residual tensors U_j such that

$$B_j^i = X_j \sqrt{\Lambda_j} U_j^i X_j^{-1},$$

and the family $(U_j)_j$ satisfies the full residual isometry condition, including the cross-block equations for $j \neq j'$.

Proof. ✓ Proof Positive basis dimensions supply the nonzero bond spaces. For each basis tensor, irreducibility and left-canonical normalization are part of the BNT canonical form. If $m_j > 0$ is the peripheral period of B_j , the normalized self-overlap hypothesis and the periodic self-overlap formula used in Theorem 21.4.16 give, along the same subsequence,

$$\lim_{k \rightarrow \infty} O_{B_j B_j}(m_j k) = 1, \quad \lim_{k \rightarrow \infty} O_{B_j B_j}(m_j k) = m_j.$$

Hence $m_j = 1$. The transfer map is primitive by Theorem 4.5.3.6, and normality follows from Theorem 9.6.2.1. Gauge-phase distinctness is also part of the BNT canonical form. The equivalence now follows from Theorem 22.4.14; the same derived hypotheses in Theorem 22.4.12 give the residual isometry family. $\square$

Theorem 22.4.16 (Ambiguity of the literal repeated-phase display). Lean ✓ Stmt *Let B be the one-letter, one-dimensional tensor $B^0 = (1)$, and form two block-diagonal copies with phases 1 and -1 :*

$$A^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The representative B is in isometry canonical form, and both copy coefficients have unit modulus. Thus A satisfies the literal virtual block-diagonal reading of the displayed repeated-copy formula in [CPGSV16, Section 3]. It does not, however, satisfy the accompanying interpretation in which the copy index also belongs to the physical direct sum: there is no scalar v such that $(A^0)^2 = vA^0$. In particular, A is not a renormalization fixed point in the sense of Definition 22.1.2.

Proof. ✓ Proof Taking the witness data of Definition 22.3.10 to be $X = \Lambda = U^0 = (1)$, the scalar representative is in isometry canonical form; moreover, $|1| = |-1| = 1$. Direct multiplication gives $(A^0)^2 = I$. Comparing diagonal entries in $I = vA^0$ would give simultaneously $v = 1$ and $v = -1$, which is impossible. On the off-diagonal matrix unit E_{01} , the transfer map is conjugation by A^0 , so

$$\mathcal{E}_A(E_{01}) = -E_{01}, \quad \mathcal{E}_A^2(E_{01}) = E_{01}.$$

Hence $\mathcal{E}_A^2 \neq \mathcal{E}_A$. $\square$

22.5 Renormalization flow and physical correlations

Theorem 22.5.1 (Canonical-form blocks are injective). [Lean](#) [✓ Stmt](#) For a family of tensors in canonical form, every block is injective. This is part of the canonical-form hypotheses in [CPGSV16, Section 2.3].

Proof. [✓ Proof](#) The canonical-form hypotheses include injectivity of each block. $\square$

Theorem 22.5.2 (RG flow convergence for canonical-form tensors). [Lean](#) [✓ Stmt](#) For a tensor in canonical form with blocks (A_k) and weights (μ_k) , the iterated blocking $\mathcal{E}_{A_k}^{2^n}$ converges pointwise to an idempotent linear map $\mathcal{E}_\infty$ as $n \rightarrow \infty$. That is, there exists $\mathcal{E}_\infty$ with $\mathcal{E}_\infty^2 = \mathcal{E}_\infty$ such that $\mathcal{E}_{A_k}^{2^n}(\rho) \rightarrow \mathcal{E}_\infty(\rho)$ for all density matrices ρ . This is [CPGSV16, Appendix B].

Proof. [✓ Proof](#) Each block is injective and left-canonical, so its transfer map is a primitive channel with a unique positive-definite fixed point ρ_0 . The exponential convergence bound $\|\mathcal{E}^n(X) - P(X)\| \leq C(1 - \delta)^n \|X\|$ (Theorem 14.3.1) then gives pointwise convergence $\mathcal{E}^n(X) \rightarrow P(X)$, and composing with the subsequence $2^n \rightarrow \infty$ yields the result. The witness $\mathcal{E}_\infty = P$ is the rank-one fixed-point projection, which is idempotent. $\square$

Remark 22.5.3. When the transfer map is idempotent ($\mathcal{E}^2 = \mathcal{E}$), connected correlations become separation-independent for all separations $n \geq 1$. Informally, this corresponds to zero correlation length ($\xi = 0$). The full spectral equivalence (idempotence iff vanishing subleading spectrum) is deferred here.

Definition 22.5.4 (Virtual-insertion distance independence). [Lean](#) [✓ Stmt](#) For a positive definite right fixed point $\rho^R > 0$, the virtual insertion expression is independent of distance when for all virtual bond matrices X and Y and all separations $n, m \geq 1$,

$$C(X, Y; n) = C(X, Y; m).$$

Definition 22.5.5 (Physical observable transfer). [Lean](#) [✓ Stmt](#) For an observable O on a block of L physical spins, define the linear map on the virtual matrix space by

$$\mathcal{E}_O(X) = \sum_{i,j} \langle j|O|i \rangle A^i X (A^j)^\dagger.$$

This is the observable transfer map of [CPGSV16, lines 490–496].

Definition 22.5.6 (Two-observable periodic expectation). [Lean](#) [✓ Stmt](#) Let O_1, O_2 act on blocks of L_1, L_2 physical spins, separated around the periodic chain by complementary gaps of lengths n_1, n_2 . Their two-observable expectation is

$$\text{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{n_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{n_1}).$$

The trace is the operator trace on the virtual matrix space, as in [CPGSV16, lines 490–496].

Definition 22.5.7 (Physical correlations independent of distance). [Lean](#) [✓ Stmt](#) Let O_1, O_2 act on two disjoint contiguous regions of positive lengths $L_1, L_2 \geq 1$ in a periodic chain, with complementary gap lengths $n_1, n_2 \geq 0$. Physical correlations are independent of distance when, for every $m_1, m_2 \geq 0$ satisfying $n_1 + n_2 = m_1 + m_2$, one has

$$\text{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{n_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{n_1}) = \text{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{m_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{m_1}).$$

Thus either region may be translated without crossing the other. Zero gaps, corresponding to adjacent regions, are included. This is [CPGSV16, Definition 3.3 and lines 490–496].

Definition 22.5.8 (Positive-gap physical distance independence). [Lean](#) [✓ Stmt](#) For an observable O on a block of L physical spins, write

$$\mathcal{E}_O(X) = \sum_{i,j} \langle j|O|i\rangle A^i X(A^j)^\dagger.$$

Place two observables O_1, O_2 on nonempty finite blocks of a periodic chain, with positive complementary gap lengths n_1, n_2 . The tensor has *positive-gap physical correlations independent of distance* when the expectation is unchanged upon replacing these gaps by positive m_1, m_2 satisfying $n_1 + n_2 = m_1 + m_2$:

$$\mathrm{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{n_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{n_1}) = \mathrm{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{m_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{m_1}).$$

This is the transfer-matrix form of [CPGSV16, Definition 3.3 and the correlation formula preceding Theorem 3.8], restricted to positive complementary gaps. It excludes adjacent regions.

Lemma 22.5.9 (Physical CID implies positive-gap physical CID). [Lean](#) [✓ Stmt](#) *Correlations independent of distance for all disjoint physical regions are independent of distance when both complementary gaps are positive.*

Proof. [✓ Proof](#) Restrict the four gap lengths to positive integers. □

Theorem 22.5.10 (Idempotent transfer implies positive-gap physical CID). [Lean](#) [✓ Stmt](#) *If $\mathcal{E}_A^2 = \mathcal{E}_A$, then physical correlations are independent of distance whenever both complementary gaps are positive.*

Proof. [✓ Proof](#) Idempotence gives $\mathcal{E}_A^n = \mathcal{E}_A$ for every $n \geq 1$. Substituting $\mathcal{E}_A^{n_1} = \mathcal{E}_A^{n_2} = \mathcal{E}_A^{m_1} = \mathcal{E}_A^{m_2} = \mathcal{E}_A$ in the two-observable transfer formula gives

$$\mathrm{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{n_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{n_1}) = \mathrm{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A) = \mathrm{Tr}_{M_D(\mathbb{C})}(\mathcal{E}_{O_2} \circ \mathcal{E}_A^{m_2} \circ \mathcal{E}_{O_1} \circ \mathcal{E}_A^{m_1}).$$

□

Definition 22.5.11 (Zero correlation length). [Lean](#) [✓ Stmt](#) In the single-block convention, an MPS tensor has *zero correlation length* when

$$\mathcal{E}_A^2 = \mathcal{E}_A \quad \text{and} \quad \mathrm{CID}(A)$$

both hold. The source definition is Definition 22.2.6; the present condition uses virtual insertions instead.

Theorem 22.5.12 (Virtual-insertion distance independence implies RFP). [Lean](#) [✓ Stmt](#) *If an MPS tensor admits a positive definite right fixed point and its virtual-insertion correlator is independent of distance, then its transfer map is idempotent.*

Proof. [✓ Proof](#) Let $\rho > 0$ be the fixed point. For every matrix Z , invertibility gives $Z = (Z\rho^{-1})\rho$. Apply distance independence with $X = Z\rho^{-1}$, the second insertion N , and separations 2 and 1. The identical disconnected terms cancel. Thus, for every N ,

$$\mathrm{tr}(N\mathcal{E}_A^2(Z)) = \mathrm{tr}(N\mathcal{E}_A(Z)).$$

Nondegeneracy of the trace pairing yields $\mathcal{E}_A^2(Z) = \mathcal{E}_A(Z)$ for every Z , hence $\mathcal{E}_A^2 = \mathcal{E}_A$. □

Theorem 22.5.13 (Zero correlation length iff idempotent transfer). Lean ✓ Stmt Under the single-block convention above,

$$(\mathcal{E}_A^2 = \mathcal{E}_A \text{ and } \text{CID}(A)) \iff \mathcal{E}_A^2 = \mathcal{E}_A.$$

This is the idempotence/CID part of [CPGSV16, Theorem 3.8]. The full source theorem for canonical-form tensors instead uses the unrestricted physical condition in Definition 22.2.6; under its stated raw-weight normalization, that biconditional is false by Theorem 22.2.7.

Proof. ✓ Proof The forward direction extracts the idempotence hypothesis. Conversely, idempotence gives $\mathcal{E}_A^n = \mathcal{E}_A$ for every $n \geq 1$. Hence, for every positive-definite right fixed point ρ^R , every pair of virtual matrices X, Y , and every $n \geq 1$,

$$\begin{aligned} C(X, Y; n) &= \text{tr}(Y \mathcal{E}_A^n(X \rho^R)) - \text{tr}(X \rho^R) \text{tr}(Y \rho^R) \\ &= \text{tr}(Y \mathcal{E}_A(X \rho^R)) - \text{tr}(X \rho^R) \text{tr}(Y \rho^R) \\ &= C(X, Y; 1). \end{aligned}$$

Thus the connected correlator is independent of every positive separation. □

Theorem 22.5.14 (Single-block RFP iff ZCL inside a canonical form). Lean ✓ Stmt Let $A^i = \bigoplus_k \mu_k A_k^i$ be a tensor in canonical form. For each k , the tensor A_k is a renormalization fixed point if and only if A_k has zero correlation length.

This is the single-block consequence of [CPGSV16, Theorem 3.10]. The ZCL condition in the source theorem is the BNT-family condition: CID together with the local orthogonality equations $\mathbb{E}_{j,j'} = 0$ for all $j \neq j'$, as in Definition 22.2.6.

Proof. ✓ Proof Apply Theorem 22.5.13 to the chosen canonical-form block. □

Chapter 23

Mixed States: Renormalization of Matrix Product Operators

Starting from the MPO, MPDO, and LPDO notation of Chapter 21, this chapter studies renormalization fixed points for density-operator chains. Following the order of the mixed-state analysis in [CPGSV16, Section 4], the discussion passes from the local renormalization maps to zero correlation length and purification fixed points, then to mutual information and saturation of the area law. It concludes with the local and commuting structures of the simple case. The boundary-theory interlude of the source is outside the present MPO development. The general case and its algebraic characterization continue in Chapter 24.

23.1 Renormalization fixed points and pure-state recovery

Definition 23.1.1 (MPO transfer-map idempotence). [Lean](#) [✓ Stmt](#) An MPO tensor M has *transfer-map idempotence* if

$$\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M.$$

Definition 23.1.2 (Trace-preserving completely positive map). [Lean](#) [✓ Stmt](#) A linear map $\mathcal{S}$ between matrix algebras is *trace-preserving completely positive* if it has a Kraus form $\mathcal{S}(X) = \sum_i A_i X A_i^\dagger$ with $\sum_i A_i^\dagger A_i = I$. The Kraus operators may be rectangular, so the input and output dimensions need not agree.

Theorem 23.1.3 (Trace preservation from a Kraus resolution). [Lean](#) [✓ Stmt](#) Every trace-preserving completely positive map in rectangular Kraus form preserves the matrix trace.

Proof. [✓ Proof](#) If $\mathcal{S}(X) = \sum_a A_a X A_a^\dagger$ and $\sum_a A_a^\dagger A_a = I$, cyclicity gives

$$\mathrm{tr} \mathcal{S}(X) = \sum_a \mathrm{tr}(A_a X A_a^\dagger) = \mathrm{tr} \left(\left(\sum_a A_a^\dagger A_a \right) X \right) = \mathrm{tr} X.$$

□

Lemma 23.1.4 (Identity map is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) The identity map $\mathrm{id}(X) = X$ is trace-preserving completely positive, with single Kraus operator $A_0 = I$.

Proof. [✓ Proof](#) Take $r = 1$, $A_0 = I$; then $\mathrm{id}(X) = I X I^\dagger = X$ and $A_0^\dagger A_0 = I$. □

Theorem 23.1.5 (Conjugation by an isometry is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) Let H and K be finite-dimensional complex vector spaces, and let $V : H \rightarrow K$ satisfy $V^\dagger V = I_H$. Then the map

$$\Phi_V : \text{End}_{\mathbb{C}}(H) \longrightarrow \text{End}_{\mathbb{C}}(K), \quad \Phi_V(X) = VXV^\dagger,$$

is trace-preserving and completely positive. This is the general one-isometry form of the local basis change used in [CPGSV16, Appendix C.2, lines 1439 and 1520].

Proof. [✓ Proof](#) Take the sole Kraus operator to be V . Its resolution of the identity is precisely $V^\dagger V = I_H$. $\square$

Definition 23.1.6 (Matrix reindexing along an equivalence). [Lean](#) [✓ Stmt](#) Let $e : I \simeq J$ be an equivalence of index sets. The associated matrix reindexing is the linear map $R_e : \mathbb{C}^{I \times I} \rightarrow \mathbb{C}^{J \times J}$ determined by

$$[R_e(X)]_{j,j'} = X_{e^{-1}(j), e^{-1}(j')}.$$

Theorem 23.1.7 (Equivalence reindexing is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) For every equivalence $e : I \simeq J$ between finite index sets, the reindexing map R_e is trace-preserving and completely positive.

Proof. [✓ Proof](#) Let $P_e : \mathbb{C}^I \rightarrow \mathbb{C}^J$ be the permutation matrix of e . Then

$$R_e(X) = P_e X P_e^\dagger, \quad P_e^\dagger P_e = I_{\mathbb{C}^I}.$$

The claim follows from the preceding isometry lemma. $\square$

Lemma 23.1.8 (Composition of trace-preserving completely positive maps). [Lean](#) [✓ Stmt](#) The composition of two trace-preserving completely positive maps is again trace-preserving completely positive. If $\mathcal{S}$ has Kraus operators A_i and $\mathcal{T}$ has Kraus operators B_j , then $\mathcal{S} \circ \mathcal{T}$ has Kraus operators $A_i B_j$.

Proof. [✓ Proof](#) The Kraus form of the composite is $(\mathcal{S} \circ \mathcal{T})(X) = \sum_{i,j} (A_i B_j) X (A_i B_j)^\dagger$, and the resolution of identity follows from those of $\mathcal{S}$ and $\mathcal{T}$:

$$\sum_{i,j} (A_i B_j)^\dagger (A_i B_j) = \sum_j B_j^\dagger \left(\sum_i A_i^\dagger A_i \right) B_j = \sum_j B_j^\dagger B_j = I.$$

$\square$

Theorem 23.1.9 (Tensor extension preserves trace-preserving complete positivity). [Lean](#) [✓ Stmt](#) Let $\mathcal{S} : \text{End}_{\mathbb{C}}(H) \rightarrow \text{End}_{\mathbb{C}}(K)$ be trace-preserving and completely positive, and let R be another finite-dimensional space. Then

$$\mathcal{S} \otimes \text{id}_R : \text{End}_{\mathbb{C}}(H \otimes R) \longrightarrow \text{End}_{\mathbb{C}}(K \otimes R)$$

is trace-preserving and completely positive. If $(A_a)_a$ is a Kraus family for $\mathcal{S}$, then $(A_a \otimes I_R)_a$ is a Kraus family for its tensor extension.

Proof. [✓ Proof](#) The tensor extension has the Kraus form

$$(\mathcal{S} \otimes \text{id}_R)(X) = \sum_a (A_a \otimes I_R) X (A_a \otimes I_R)^\dagger.$$

Its normalization follows from that of the original family:

$$\sum_a (A_a \otimes I_R)^\dagger (A_a \otimes I_R) = \left(\sum_a A_a^\dagger A_a \right) \otimes I_R = I_H \otimes I_R.$$

□

Definition 23.1.10 (Rectangular Kraus map). [Lean](#) [✓ Stmt](#) Let I be a finite index set, let J be an arbitrary index set, and let $(A_a \in \mathbb{C}^{J \times I})_a$ be a finite family of matrices. Its rectangular Kraus map is

$$\Phi_A(X) = \sum_a A_a X A_a^\dagger.$$

Theorem 23.1.11 (Relabeling a rectangular Kraus family). [Lean](#) [✓ Stmt](#) If $e : I \simeq I'$ and $A'_b = A_{e^{-1}(b)}$, then $\Phi_{A'} = \Phi_A$.

Proof. [✓ Proof](#) Reindexing the finite sum gives

$$\sum_{b \in I'} A_{e^{-1}(b)} X A_{e^{-1}(b)}^\dagger = \sum_{a \in I} A_a X A_a^\dagger.$$

□

Definition 23.1.12 (Orthogonally controlled Kraus map). [Lean](#) [Lean](#) [✓ Stmt](#) Let $H = \bigoplus_k H_k$ and $K = \bigoplus_k K_k$. For each k , let $(A_{k,a} : H_k \rightarrow K_k)_a$ be a finite family of operators, and let $\tilde{A}_{k,a} : H \rightarrow K$ agree with $A_{k,a}$ on H_k and vanish on every other summand. The orthogonally controlled Kraus map is

$$\mathcal{C}(X) = \sum_{k,a} \tilde{A}_{k,a} X \tilde{A}_{k,a}^\dagger.$$

In particular, $\mathcal{C}(X)_{kl} = 0$ for $k \neq l$. This is the sector control in the definitions of $\mathcal{T}_1$ and $\mathcal{S}_1$ in [\[CPGSV16, Appendix C.2, lines 1523–1535 and 1548–1555\]](#).

Theorem 23.1.13 (Diagonal blocks of an orthogonally controlled map). [Lean](#) [✓ Stmt](#) Let $X_{kk} : H_k \rightarrow H_k$ denote the k th diagonal block of X . Then

$$\mathcal{C}(X)_{kk} = \sum_a A_{k,a} X_{kk} A_{k,a}^\dagger = \Phi_{A_k}(X_{kk}).$$

Proof. [✓ Proof](#) If $j \neq k$, then every zero-extended Kraus operator satisfies

$$[\tilde{A}_{j,a} X \tilde{A}_{j,a}^\dagger]_{(k,b),(k,c)} = 0.$$

Hence only the $j = k$ terms survive, and

$$[\mathcal{C}(X)]_{kk} = \sum_a A_{k,a} X_{kk} A_{k,a}^\dagger.$$

□

Theorem 23.1.14 (Off-diagonal blocks of an orthogonally controlled map). [Lean](#) [✓ Stmt](#) If $k \neq l$, then $\mathcal{C}(X)_{kl} = 0$.

Proof. ✓ Proof For every (j, a) and $k \neq l$, the zero-extended operator satisfies

$$[\tilde{A}_{j,a} X \tilde{A}_{j,a}^\dagger]_{(k,b),(l,c)} = 0.$$

Therefore $[\mathcal{C}(X)]_{kl} = \sum_{j,a} 0 = 0$. □

Theorem 23.1.15 (Orthogonal control preserves trace-preserving complete positivity). Lean

✓ Stmt Suppose that for every k the sectorwise Kraus family resolves the identity:

$$\sum_a A_{k,a}^\dagger A_{k,a} = I_{H_k}.$$

Then the orthogonally controlled map $\mathcal{C}$ is trace-preserving and completely positive.

Proof. ✓ Proof Each embedded operator has support in one summand, and therefore

$$\sum_{k,a} \tilde{A}_{k,a}^\dagger \tilde{A}_{k,a} = \bigoplus_k \left(\sum_a A_{k,a}^\dagger A_{k,a} \right) = \bigoplus_k I_{H_k} = I_H.$$

The displayed Kraus form gives complete positivity, while this identity gives trace preservation. □

Definition 23.1.16 (Right partial trace). Lean ✓ Stmt For a matrix X on $A \otimes B$, the right partial trace is the linear map from matrices on $A \otimes B$ to matrices on A given by

$$[\mathrm{tr}_B(X)]_{ij} = \sum_k X_{(i,k),(j,k)}.$$

After the retained and discarded subspins have been regrouped as $A \otimes B$, this is the partial-trace ingredient of the maps $\mathcal{T}_0$ and $\mathcal{S}_0$ in [\[CPGSV16, Appendix C.2, lines 1521–1522 and 1547\]](#).

Theorem 23.1.17 (Right partial trace of a Kronecker product). Lean ✓ Stmt For square matrices X on A and Y on B ,

$$\mathrm{tr}_B(X \otimes Y) = \mathrm{tr}(Y)X.$$

Proof. ✓ Proof For all i, j ,

$$[\mathrm{tr}_B(X \otimes Y)]_{ij} = \sum_t X_{ij} Y_{tt} = \mathrm{tr}(Y) X_{ij}.$$

□

Lemma 23.1.18 (The right partial trace is trace-preserving completely positive). Lean ✓ Stmt

The map tr_B is trace-preserving and completely positive for arbitrary finite-dimensional spaces A and B .

Proof. ✓ Proof For each basis vector e_k of B , define $V_k : A \otimes B \rightarrow A$ by

$$V_k(e_i \otimes e_\ell) = \delta_{k\ell} e_i.$$

Then

$$\sum_k V_k X V_k^\dagger = \mathrm{tr}_B(X), \quad \sum_k V_k^\dagger V_k = I_{A \otimes B}.$$

□

Definition 23.1.19 (State-preparation map). [Lean](#) [✓ Stmt](#) Let ρ be a matrix on a finite-dimensional space B . The state-preparation map from matrices on A to matrices on $A \otimes B$ is

$$\mathcal{P}_\rho(X) = X \otimes \rho.$$

This is the elementary preparation operation used in the maps $\mathcal{T}_1$ and $\mathcal{S}_1$ of [CPGSV16, Appendix C.2, lines 1527–1533 and 1551–1555].

Definition 23.1.20 (Preparation Kraus operators). [Lean](#) [✓ Stmt](#) Let $R = \sqrt{\rho}$. For an orthonormal basis $(e_j)_j$ of B , define rectangular operators $A_j : A \rightarrow A \otimes B$ by

$$A_j(e_a) = e_a \otimes R e_j.$$

Theorem 23.1.21 (Kraus action of state preparation). [Lean](#) [✓ Stmt](#) If $\rho \succeq 0$ and $A_j(e_a) = e_a \otimes \sqrt{\rho} e_j$, then

$$\sum_j A_j X A_j^\dagger = X \otimes \rho.$$

Proof. [✓ Proof](#) Put $R = \sqrt{\rho}$. Since R is Hermitian and $R^2 = \rho$, the $((a, s), (b, t))$ entry of the left-hand side is

$$X_{ab} \sum_j R_{sj} \overline{R_{tj}} = X_{ab} (R^2)_{st} = X_{ab} \rho_{st}.$$

□

Theorem 23.1.22 (Preparation Kraus operators resolve the identity). [Lean](#) [✓ Stmt](#) If $\rho \geq 0$ and $\text{tr}(\rho) = 1$, then

$$\sum_j A_j^\dagger A_j = I_A.$$

Proof. [✓ Proof](#) Since R is Hermitian and $R^2 = \rho$, the diagonal entries of the sum are

$$\sum_{j,t} \overline{R_{tj}} R_{tj} = \text{tr}(R^2) = \text{tr}(\rho) = 1,$$

while its off-diagonal entries vanish. □

Lemma 23.1.23 (State preparation is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) If $\rho \geq 0$ and $\text{tr}(\rho) = 1$, then $\mathcal{P}_\rho : X \mapsto X \otimes \rho$ is trace-preserving completely positive.

Proof. [✓ Proof](#) The operators from Definition 23.1.20 satisfy

$$\sum_j A_j X A_j^\dagger = X \otimes R R = X \otimes \rho,$$

and Theorem 23.1.22 gives their resolution of the identity. □

Definition 23.1.24 (Physical closure maps). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For an MPO tensor M , a length N , and a virtual operator X , define the physical operator $M_N(X)$ by

$$M_N(X)_{i,j} = \text{tr}(M^{i_0 j_0} M^{i_1 j_1} \dots M^{i_{N-1} j_{N-1}} X).$$

This is linear in X . The displayed order chooses the cut of the cyclic virtual contraction immediately before the first site, so that X follows the last site. The one- and two-site operators are constructed in [CPGSV16, lines 638–654 and Definition 4.1]. When $M = \mathcal{K}$, write these operators as $\mathcal{K}_N(X)$; the cases $N = 2, 3$ occur in [CPGSV16, Proposition C.7, lines 1510–1516].

Theorem 23.1.25 (One- and two-site physical closures). [Lean](#) [Lean](#) [✓ Stmt](#) Under the canonical identifications of one-site configurations with physical indices and two-site configurations with pairs of physical indices,

$$M_1(X)_{i,j} = \text{tr}(M^{ij}X), \quad M_2(X)_{(i_0,i_1),(j_0,j_1)} = \text{tr}(M^{i_0j_0}M^{i_1j_1}X).$$

Proof. [✓ Proof](#) Under these identifications, the length-one and length-two word evaluations are M^{ij} and $M^{i_0j_0}M^{i_1j_1}$, respectively. $\square$

Lemma 23.1.26 (Three-site physical closure). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The general three-site physical closure has coefficients

$$M_3(X)_{(i_0,(i_1,i_2),(j_0,(j_1,j_2)))} = \text{tr}(M^{i_0j_0}M^{i_1j_1}M^{i_2j_2}X).$$

For $M = \mathcal{K}$, this is the operator $\mathcal{K}_3(X)$ in [CPGSV16, Proposition C.7, lines 1510–1516].

Proof. [✓ Proof](#) The length-three word evaluation is $M^{i_0j_0}M^{i_1j_1}M^{i_2j_2}$. $\square$

Definition 23.1.27 (Renormalization fixed point via trace-preserving maps). [Lean](#) [✓ Stmt](#) An MPO tensor M is a *renormalization fixed point* if there exist two trace-preserving completely positive maps $\mathcal{S}, \mathcal{T}$ on the physical indices such that $\mathcal{S}[M_2(X)] = M_1(X)$ and $\mathcal{T}[M_1(X)] = M_2(X)$ for every virtual operator X , where $M_1(X)_{ij} = \text{tr}(M^{ij}X)$ and $M_2(X)_{(i_1i_2)(j_1j_2)} = \text{tr}(M^{i_1j_1}M^{i_2j_2}X)$ are the one- and two-site physical operators obtained by closing one or two tensors with X . This condition is distinct from idempotence of the doubled-index transfer map for general mixed states. Its source zero-correlation-length consequence concerns the physical-trace transfer and is proved below. This is [CPGSV16, Definition 4.1, line 657]. The two maps act as follows on every virtual closure X :

$$M_1(X) = \text{---} \bullet \text{---} \xrightarrow[\mathcal{S}]{\mathcal{T}} \text{---} \bullet \text{---} \bullet \text{---} = M_2(X)$$

Theorem 23.1.28 (Renormalization fixed points have idempotent physical-trace transfer). [Lean](#) [✓ Stmt](#) If M is a renormalization fixed point via trace-preserving maps, then

$$\mathcal{T}_M^2 = \mathcal{T}_M.$$

This proves the zero-correlation-length component of implication (i) $\Rightarrow$ (ii) in [CPGSV16, Theorem 4.9, lines 851–893]. The corresponding calculation appears in the proposition “RFP implies ZCL and SAL” in Appendix C, lines 1333–1340; its strong-area-law component is separate.

Proof. [✓ Proof](#) For every virtual matrix X , the one-site and two-site closures satisfy

$$\text{tr } M_1(X) = \text{tr}(\mathcal{T}_M X), \quad \text{tr } M_2(X) = \text{tr}(\mathcal{T}_M^2 X).$$

Write Φ for the one-to-two trace-preserving map $\mathcal{T}$ of Definition 23.1.27; this is distinct from the physical-trace transfer $\mathcal{T}_M$. Then $\Phi[M_1(X)] = M_2(X)$, and hence

$$\text{tr}(\mathcal{T}_M^2 X) = \text{tr } M_2(X) = \text{tr } \Phi[M_1(X)] = \text{tr } M_1(X) = \text{tr}(\mathcal{T}_M X).$$

Nondegeneracy of the trace pairing now gives $\mathcal{T}_M^2 = \mathcal{T}_M$. $\square$

Theorem 23.1.29 (Renormalization fixed points have source zero correlation length). [Lean](#) [✓ Stmt](#) If M is a renormalization fixed point via trace-preserving maps and $\mathcal{T}_M \neq 0$, then M has source zero correlation length.

Proof.  Apply the preceding idempotence theorem and take the positive scalar in Definition 23.2.2 to be 1. $\square$

Theorem 23.1.30 (Transfer-map idempotence iff doubled-index condition).   *MPO transfer-map idempotence is exactly the doubled-index transfer condition of Definition 23.2.5.*

Proof.  Both conditions assert $\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M$; they are definitionally identical. $\square$

Theorem 23.1.31 (Transfer-map idempotence iff doubled-index MPS RFP).   *MPO transfer-map idempotence is equivalent to the pure-state RFP condition for the doubled-index MPS tensor.*

Proof.  Unfold transfer-map idempotence to its definitional equivalent, the doubled-index transfer condition, and apply Theorem 23.2.6. $\square$

Definition 23.1.32 (Diagonal pure-state embedding as an MPO).   An MPS tensor $A = \{A^i\}_{i=0}^{d-1}$ determines an MPO tensor by placing A^i on the diagonal in the bra and ket indices:

$$M^{ij} = \delta_{ij} A^i.$$

Equivalently, $M^{ii} = A^i$ and $M^{ij} = 0$ for $i \neq j$.

Theorem 23.1.33 (Transfer map of the diagonal pure-state embedding).   *The transfer map of the diagonal MPO associated to A agrees with the original MPS transfer map:*

$$\mathcal{E}_{A^{\text{MPO}}} = \mathcal{E}_A.$$

Proof.  In the double sum defining $\mathcal{E}_{A^{\text{MPO}}}$, all off-diagonal terms vanish because $M^{ij} = 0$ for $i \neq j$, while the diagonal terms are exactly $A^i X(A^i)^\dagger$. Thus only the sum over i remains, which is the defining formula for $\mathcal{E}_A$. $\square$

Theorem 23.1.34 (Pure-state recovery of transfer-map idempotence).   *For an MPS tensor A , the diagonal MPO embedding has idempotent transfer map if and only if A is an RFP:*

$$\mathcal{E}_{A^{\text{MPO}}} \circ \mathcal{E}_{A^{\text{MPO}}} = \mathcal{E}_{A^{\text{MPO}}} \iff \mathcal{E}_A \circ \mathcal{E}_A = \mathcal{E}_A.$$

Proof.  Both conditions assert idempotence of the corresponding transfer map. Theorem 23.1.33 identifies those transfer maps, so the two predicates are equivalent. $\square$

Theorem 23.1.35 (Pure-state recovery of zero correlation length).   *For an MPS tensor A , the diagonal MPO embedding has idempotent transfer map if and only if A has zero correlation length.*

Proof.  Combine Theorem 23.1.34 with the characterization of zero correlation length by idempotence of the MPS transfer map (Theorem 22.5.13). $\square$

23.2 Zero correlation length

Definition 23.2.1 (Physical-trace transfer).   The physical-trace transfer of an MPO tensor M is the virtual matrix obtained by contracting the ket and bra physical legs of one tensor:

$$\mathcal{T}_M = \sum_i M^{ii}.$$

This is the transfer object appearing in the zero-correlation-length condition of [CPGSV16, Definition 4.2, lines 736–741].

Definition 23.2.2 (Source zero correlation length for MPO tensors). [Lean](#) [✓ Stmt](#) An MPO tensor has source zero correlation length when, for some real number $\lambda > 0$,

$$\mathcal{T}_M \neq 0 \quad \text{and} \quad \mathcal{T}_M^2 = \lambda \mathcal{T}_M$$

hold. The nonzero condition excludes the degenerate zero transfer, and the positive scalar makes the condition invariant under rescaling of M .

Theorem 23.2.3 (Literal physical-trace idempotence gives source ZCL). [Lean](#) [✓ Stmt](#) If $\mathcal{T}_M \neq 0$ and $\mathcal{T}_M^2 = \mathcal{T}_M$, then M has source zero correlation length.

Proof. [✓ Proof](#) This is the preceding definition with $\lambda = 1$. □

$$\mathcal{T}_M = \mathcal{T}_M^2$$

Here closing the ket and bra legs gives $\mathcal{T}_M$, and juxtaposition is virtual multiplication. Thus the displayed identity is precisely $\mathcal{T}_M = \mathcal{T}_M^2$. This is the normalized identity in [CPGSV16, Definition 4.2, lines 736–741]; Definition 23.2.2 permits the positive factor λ .

Lemma 23.2.4 (Normalized transfer is idempotent under source zero correlation length). [Lean](#) [✓ Stmt](#) If M has source zero correlation length, then there exists $\lambda > 0$ such that $\lambda^{-1}\mathcal{T}_M$ is idempotent.

Proof. [✓ Proof](#) From source zero correlation length, choose $\lambda > 0$ with $\mathcal{T}_M^2 = \lambda \mathcal{T}_M$. Then

$$(\lambda^{-1}\mathcal{T}_M)^2 = \lambda^{-2}\mathcal{T}_M^2 = \lambda^{-1}\mathcal{T}_M.$$

□

Definition 23.2.5 (Doubled-index transfer idempotence for MPO tensors). [Lean](#) [✓ Stmt](#) An MPO tensor M satisfies the doubled-index transfer condition when its completely positive transfer map is idempotent:

$$\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M.$$

This is the condition obtained from the doubled-index MPS view. It is not the physical-trace condition of Definition 23.2.2.

Theorem 23.2.6 (Doubled-index transfer idempotence iff MPS RFP). [Lean](#) [✓ Stmt](#) An MPO tensor satisfies the doubled-index transfer condition if and only if its doubled-index MPS view is a renormalization fixed point.

Proof. [✓ Proof](#) By Lemma 21.2.2, the transfer map of M agrees with the transfer map of its doubled-index MPS view. Thus $\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M$ holds exactly when the transfer map of that doubled-index MPS tensor is idempotent, which is the RFP condition. □

23.3 Purification fixed points

Definition 23.3.1 (Purifying spin–ancilla tensor). [Lean](#) [✓ Stmt](#) Given matrices $A^{(i,k)}$ with a spin index i and an ancillary index k , the associated MPS tensor on the product physical space is

$$\widehat{A}^{(i,k)} = A^{(i,k)}.$$

This is the tensor whose finite-chain vector appears in the purification formula of [CPGSV16, lines 744–751].

Definition 23.3.2 (Ancillary trace of a purifying MPS). [Lean](#) [✓ Stmt](#) For a purifying spin-ancilla tensor A , the reduced spin density matrix is defined by

$$[\rho_p^{(N)}(A)]_{\sigma,\tau} = \sum_{\kappa} \psi_A(\sigma, \kappa) \overline{\psi_A(\tau, \kappa)},$$

where

$$\psi_A(\sigma, \kappa) = \text{tr}(A^{(\sigma_0, \kappa_0)} A^{(\sigma_1, \kappa_1)} \dots A^{(\sigma_{N-1}, \kappa_{N-1})}).$$

This is the coefficient form of the ancillary trace in [\[CPGSV16, eq. \(4.7\), line 751\]](#).

Definition 23.3.3 (Single-site ancillary trace map). [Lean](#) [✓ Stmt](#) For a matrix X on the spin-ancilla space, the ancillary trace map is

$$[\text{tr}_a(X)]_{ij} = \sum_k X_{(i,k),(j,k)}.$$

This is the one-site trace over ancillary degrees of freedom appearing in [\[CPGSV16, lines 751 and 761–764\]](#).

Theorem 23.3.4 (The ancillary trace is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) For every ancillary dimension, the one-site ancillary trace map is trace-preserving and completely positive.

Proof. [✓ Proof](#) This is Lemma [23.1.18](#) with A the spin space and B the ancillary space. $\square$

Definition 23.3.5 (Global purification equation). [Lean](#) [✓ Stmt](#) An MPO tensor M satisfies the global purification equation with a purifying tensor A if, for every positive length N ,

$$\rho^{(N)}(M) = \rho_p^{(N)}(A).$$

Equivalently, for every positive length N ,

$$\rho^{(N)}(M) = \text{tr}_a(|\Psi^{(N)}(A)\rangle\langle\Psi^{(N)}(A)|),$$

as in [\[CPGSV16, line 751\]](#).

Definition 23.3.6 (Trace-preserving spin reduction). [Lean](#) [✓ Stmt](#) The spin reduction obtained by tracing the ancillary index is trace-preserving completely positive: the ancillary trace map tr_a satisfies Definition [23.1.2](#). This is the structure described immediately after Definition 4.3 in [\[CPGSV16, lines 761–764\]](#).

Corollary 23.3.7 (Trace-preserving spin reduction of the ancillary trace). [Lean](#) [✓ Stmt](#) For every ancillary dimension, the spin reduction obtained by tracing the ancillary index is trace-preserving and completely positive.

Proof. [✓ Proof](#) This is exactly Theorem [23.3.4](#), unpacked through Definition [23.3.6](#). $\square$

Definition 23.3.8 (Bare global purification RFP witness). [Lean](#) [✓ Stmt](#) An MPO tensor M has a bare global purification RFP witness if it satisfies the global purification equation for some purifying spin-ancilla tensor A , and $\hat{A}$ is a pure-state renormalization fixed point:

$$\mathcal{E}_{\hat{A}} \circ \mathcal{E}_{\hat{A}} = \mathcal{E}_{\hat{A}}.$$

This records the displayed conditions of Definition 4.3 in [\[CPGSV16, lines 756–764\]](#). It does not include the nondegeneracy implicit when the subsequent theorem speaks of density operators.

Definition 23.3.9 (Bare global purification RFP). [Lean](#) [✓ Stmt](#) An MPO tensor satisfies the bare global purification RFP predicate if it has a positive-length global purification RFP witness. Equivalently, it admits a purifying tensor A such that, for every positive length N ,

$$\rho^{(N)}(M) = \text{tr}_a(|\Psi^{(N)}(A)\rangle\langle\Psi^{(N)}(A)|),$$

and $\hat{A}$ is a pure-state renormalization fixed point. These are the displayed conditions of Definition 4.3 in [CPGSV16, lines 756–764]; the nondegenerate tensor-level predicate is Definition 23.3.13.

Definition 23.3.10 (Trace-preserving spin reduction for a purification RFP). [Lean](#) [✓ Stmt](#) A purification RFP is viewed together with trace-preserving spin reduction by applying Corollary 23.3.7 to the ancillary trace map appearing in its purification equation. This records the structure obtained after tracing the ancillary indices in [CPGSV16, lines 761–764], without adding a separate hypothesis to Definition 4.3.

Lemma 23.3.11 (Forgetting the recorded spin reduction). [Lean](#) [Lean](#) [✓ Stmt](#) *Recording the trace-preserving ancillary reduction does not change the purification RFP condition:*

$$\text{PRFP}_{\text{tp}}(M) \iff \text{PRFP}(M).$$

Proof. [✓ Proof](#) Immediate from the equivalence in Definition 23.3.10. □

Definition 23.3.12 (Local purification RFP condition). [Lean](#) [✓ Stmt](#) An MPO tensor M satisfies the local purification RFP condition if it is an LPDO whose purifying tensor A , viewed as a matrix product state tensor on the combined spin–ancilla index, is a pure-state renormalization fixed point. Thus

$$M^{ij} = \left(\sum_k A^{(i,k)} \otimes \overline{A^{(j,k)}} \right)_{e,e}.$$

This local condition is motivated by the purification tensor formula [CPGSV16, lines 744–747], but it is a one-site tensor-level condition rather than the global finite-chain definition in Definition 23.3.5.

Definition 23.3.13 (Nondegenerate purification RFP). [Lean](#) [✓ Stmt](#) An MPO tensor is a nondegenerate purification renormalization fixed point if it is the local ancillary contraction of a pure-state renormalization fixed point and

$$\mathcal{T}_M = \sum_i M^{ii} \neq 0.$$

The nonzero condition records the nondegeneracy implicit in the density operators of [CPGSV16, lines 744–786] and excludes the zero purifying tensor.

Theorem 23.3.14 (A local purification RFP generates MPDOs). [Lean](#) [✓ Stmt](#) *Every tensor satisfying the local purification RFP condition generates matrix product density operators.*

Proof. [✓ Proof](#) Dropping the renormalization-fixed-point condition on the purifying tensor leaves the local purification structure, so M is an LPDO; the conclusion follows from Theorem 21.3.3. □

Theorem 23.3.15 (Local purification RFP does not imply doubled-index idempotence). [Lean](#) [✓ Stmt](#) *There is an MPO tensor satisfying the local purification RFP condition for which the literal transfer-map idempotence $\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M$ fails.*

Proof. ✓ Proof Take the diagonal purification at $d = d_K = 2$ and $D = D' = 1$ with amplitudes $A = [\frac{1}{\sqrt{2}}, 0, 0, \frac{1}{\sqrt{2}}]$. The purifying tensor is a pure-state renormalization fixed point, since $\sum |A|^2 = 1$ makes its transfer map the identity. The ancilla contraction

$$M^{ij} = \sum_k A^{(i,k)} \overline{A^{(j,k)}}$$

gives the scalar entries $M^{00} = M^{11} = \frac{1}{2}$ (off-diagonal 0), so the induced transfer map is $\frac{1}{2} \cdot \text{id}$ and $\mathcal{E}_M \circ \mathcal{E}_M = \frac{1}{4} \cdot \text{id} \neq \mathcal{E}_M$. The trace contraction has dropped the leading eigenvalue from 1 to $\frac{1}{2}$; the literal zero-correlation-length condition is therefore strictly stronger than the source's normalized one. $\square$

Lemma 23.3.16 (Physical-trace transfer of the maximally mixed witness). Lean ✓ Stmt *For the maximally mixed witness M_w , closing the ket and bra physical legs gives*

$$\mathcal{T}_{M_w} = M_w^{00} + M_w^{11} = \frac{1}{2} + \frac{1}{2} = 1.$$

Proof. ✓ Proof This is the direct computation of the two nonzero diagonal entries of the witness tensor. $\square$

Theorem 23.3.17 (The maximally mixed witness has source zero correlation length). Lean ✓ Stmt *The maximally mixed witness M_w has source zero correlation length.*

Proof. ✓ Proof Lemma 23.3.16 gives $\mathcal{T}_{M_w} = 1$. The identity is nonzero and idempotent, so source zero correlation length holds with $\lambda = 1$. $\square$

Theorem 23.3.18 (Fixed local purification: pure RFP iff normalized physical-trace idempotence). Lean ✓ Stmt *Suppose that M is the ancillary contraction of a fixed spin-ancilla tensor A through a bond identification e :*

$$M^{ij} = \left(\sum_k A^{(i,k)} \otimes \overline{A^{(j,k)}} \right)_{e,e}.$$

Then the purification tensor $\widehat{A}$ is a pure-state renormalization fixed point if and only if

$$\mathcal{T}_M^2 = \mathcal{T}_M.$$

This is the normalized tensor-level equivalence underlying the purification theorem in [CPGSV16, lines 744–786].

Proof. ✓ Proof Let K' be the transfer matrix of $\widehat{A}$, let s interchange the two Kronecker factors, and let K denote the same Kronecker sum in the physical ordering. The local contraction gives

$$\mathcal{T}_M = K_e, \quad K = (K')_s.$$

Since both reindexings are bijective and commute with matrix multiplication,

$$(K')^2 = K' \iff \mathcal{T}_M^2 = \mathcal{T}_M.$$

Injectivity of the transfer-matrix representation gives

$$(K')^2 = K' \iff \mathcal{E}_{\widehat{A}} \circ \mathcal{E}_{\widehat{A}} = \mathcal{E}_{\widehat{A}}.$$

$\square$

Theorem 23.3.19 (Physical-trace transfer is idempotent under the local purification RFP condition). [Lean](#) [✓ Stmt](#) *If an MPO tensor M is the ancilla contraction of a pure-state renormalization fixed point, then its physical-trace transfer $\mathcal{T}_M = \sum_i M^{ii}$ is idempotent,*

$$\mathcal{T}_M \mathcal{T}_M = \mathcal{T}_M.$$

Proof. [✓ Proof](#) Apply the forward implication of Theorem 23.3.18 to the local purifying tensor and bond identification. $\square$

Theorem 23.3.20 (Local purification form and transfer idempotence characterize a local purification RFP). [Lean](#) [✓ Stmt](#) *An MPO tensor satisfies the local purification RFP condition if and only if it has a local purification and its physical-trace transfer is idempotent:*

$$\text{LocalPRFP}(M) \iff \text{LPDO}(M) \text{ and } \mathcal{T}_M^2 = \mathcal{T}_M.$$

Proof. [✓ Proof](#) Choose the local purifying tensor and bond identification. The equivalence is Theorem 23.3.18 for that fixed purification. $\square$

Theorem 23.3.21 (Nondegenerate PRFP equivalence). [Lean](#) [✓ Stmt](#) *An MPO tensor is a nondegenerate purification renormalization fixed point if and only if it is a local purification density operator (Definition 21.3.2) and*

$$\mathcal{T}_M \neq 0, \quad \mathcal{T}_M^2 = \mathcal{T}_M.$$

This is the normalized, nonzero structural characterization used in the forward PRFP-to-ZCL implication of [CPGSV16, lines 744–786].

Proof. [✓ Proof](#) Combine the preceding equivalence with the nonzero clause in Definition 23.3.13. $\square$

Theorem 23.3.22 (Local purification RFP with nonzero physical-trace transfer has source ZCL). [Lean](#) [✓ Stmt](#) *Let M be the ancilla contraction of a pure-state renormalization fixed point, satisfying the local purification renormalization fixed point condition. If the physical-trace transfer $\mathcal{T}_M = \sum_i M^{ii}$ is nonzero, then M has source zero correlation length.*

Proof. [✓ Proof](#) Theorem 23.3.19 gives idempotence of the physical-trace transfer, and the nonzero hypothesis lets Theorem 23.2.3 conclude source zero correlation length with $\lambda = 1$. $\square$

Theorem 23.3.23 (A nondegenerate purification RFP has source ZCL). [Lean](#) [✓ Stmt](#) *Every nondegenerate purification renormalization fixed point has source zero correlation length.*

Proof. [✓ Proof](#) Definition 23.3.13 supplies both the local purification RFP condition and the nonzero physical-trace transfer required by Theorem 23.3.22. $\square$

Theorem 23.3.24 (The tensor purification identity yields the global purification equation). [Lean](#) [✓ Stmt](#) *Let M be the ancilla contraction of a purifying spin-ancilla tensor A through a bond identification e , so that*

$$M^{ij} = \left(\sum_k A^{(i,k)} \otimes \overline{A^{(j,k)}} \right)_{e,e}.$$

Then at every length N the matrix product operator equals the ancillary trace of the pure spin-ancilla matrix product state,

$$\rho^{(N)}(M) = \rho_p^{(N)}(A).$$

This is the coefficient form of the purification equation of [CPGSV16, line 751].

Proof. ✓ Proof Applying the mixed-product property $(A \otimes B)(C \otimes D) = (AC) \otimes (BD)$ repeatedly, the length- N product of the contracted tensor entries distributes over the ancillary index sum:

$$\prod_{l=0}^{N-1} \left(\sum_k A^{(\sigma_l, k)} \otimes \overline{A^{(\tau_l, k)}} \right) = \sum_{\kappa: [N] \rightarrow [d_K]} \left(\prod_l A^{(\sigma_l, \kappa_l)} \right) \otimes \overline{\left(\prod_l A^{(\tau_l, \kappa_l)} \right)}.$$

Tracing this sum and using $\text{tr}(X \otimes \overline{Y}) = \text{tr}(X) \text{tr}(\overline{Y})$ gives the coefficient of the ancillary trace at every σ, τ . The equation holds at every length, so no positive-length restriction is needed. $\square$

Theorem 23.3.25 (The local purification RFP condition implies the purification RFP). Lean ✓ Stmt *Every MPO tensor satisfying the local purification renormalization fixed point condition is a purification renormalization fixed point in the sense of Definition 4.3 of [CPGSV16, lines 756–764].*

Proof. ✓ Proof The local condition supplies a purifying tensor A that is a pure-state renormalization fixed point and whose ancilla contraction is M . By Theorem 23.3.24 this contraction satisfies the global purification equation at every positive length, so the same A is a purification RFP witness. $\square$

Corollary 23.3.26 (A nondegenerate purification RFP satisfies the global PRFP predicate). Lean ✓ Stmt *Every nondegenerate purification renormalization fixed point satisfies the positive-length global purification predicate of Definition 4.3 in [CPGSV16, lines 744–758].*

Proof. ✓ Proof Forget the nonzero condition and apply Theorem 23.3.25. $\square$

Theorem 23.3.27 (The bare global PRFP predicate does not imply source ZCL). Lean ✓ Stmt *There is an MPO tensor satisfying the bare positive-length global purification RFP predicate which does not have source zero correlation length.*

Proof. ✓ Proof Take the zero tensor at spin, ancillary, and bond dimensions one, purified by the zero spin–ancilla tensor. Its transfer map is idempotent and every positive-length density operator is zero, so the global purification equation holds. Its physical-trace transfer is zero, which is excluded by source zero correlation length. $\square$

23.4 Saturation of the area law

Definition 23.4.1 (Block entropy of an MPDO). Lean ✓ Stmt For a fixed system size N , a block length $L \leq N$, and an MPO tensor M such that $\rho^{(N)}(M)$ is positive semidefinite, let

$$\sigma^{(N)}(M) = \frac{\rho^{(N)}(M)}{\text{tr}[\rho^{(N)}(M)]}, \quad \rho_L^{(N)}(M) = \text{tr}_{N \setminus L}[\sigma^{(N)}(M)]$$

denote the normalized state and the reduced state of the first L of N spins. The L -block entropy is

$$S_L^{(N)}(M) = S(\rho_L^{(N)}(M)).$$

Definition 23.4.2 (Mutual information of an MPDO). Lean ✓ Stmt For the same fixed system size N , block length $L \leq N$, and positive semidefiniteness of $\rho^{(N)}(M)$, the *mutual information* between a block of L spins and the rest is $I_L = S_L^{(N)}(M) + S_{N-L}^{(N)}(M) - S_N^{(N)}(M)$.

Definition 23.4.3 (Saturation of the area law). [Lean](#) [✓ Stmt](#) An MPO tensor M that generates an MPDO and whose finite-chain operators $\rho^{(N)}(M)$ all have nonzero trace (so the normalized states are well-defined) *saturates the area law* if $I_L = I_{L+1}$ for all $1 \leq L < \lfloor N/2 \rfloor$ and all N ; that is, the mutual information is constant, $I_1 = I_2 = \dots = I_{\lfloor N/2 \rfloor}$. This is [CPGSV16, Definition 4.6, line 811].

Lemma 23.4.4 (Saturation telescopes to a constant mutual information). [Lean](#) [✓ Stmt](#) If an MPO tensor M saturates the area law, then the mutual informations coincide for all $1 \leq L, L' \leq \lfloor N/2 \rfloor$:

$$I_L = I_{L'}.$$

Proof. [✓ Proof](#) The defining condition supplies the single-step equalities $I_m = I_{m+1}$ for $1 \leq m < \lfloor N/2 \rfloor$. Climbing one step at a time from the smaller of L and L' up to the larger composes these into the stated equality. $\square$

Theorem 23.4.5 (Tripartite marginals as block entropies). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let the N -site operator generated by M be positive semidefinite, and let three consecutive regions have lengths a , b , and c , with $a + b + c \leq N$. After reindexing the reduced state of the first $a + b + c$ sites as a tripartite state on $A \otimes B \otimes C$, its four entropies in strong subadditivity are

$$S(ABC) = S_{a+b+c}, \quad S(AB) = S_{a+b}, \quad S(B) = S_b, \quad S(BC) = S_{b+c}.$$

Proof. [✓ Proof](#) Write $\rho^{(k)}$ for the first- k -site reduced state, with $\cong$ denoting equality up to a permutation of basis indices. Reindexing a matrix does not change its von Neumann entropy, giving the first equality $S(ABC) = S_{a+b+c}$ immediately. Tracing out C leaves the first $a + b$ sites,

$$\mathrm{tr}_C(\rho_{ABC}) \cong \rho^{(a+b)},$$

so $S(AB) = S_{a+b}$. Tracing out A and C leaves the middle b sites, while tracing out A leaves the final $b + c$ sites; translation invariance gives

$$\mathrm{tr}_{A,C}(\rho_{ABC}) \cong \rho^{(b)}, \quad \mathrm{tr}_A(\rho_{ABC}) \cong \rho^{(b+c)}.$$

Hence $S(B) = S_b$ and $S(BC) = S_{b+c}$. $\square$

Proposition 23.4.6 (Monotonicity of the mutual information). [Lean](#) [✓ Stmt](#) Let M generate an MPDO whose finite-chain operator $\rho^{(N)}(M)$ is positive semidefinite with nonzero trace. For every block length L with $2L + 1 \leq N$,

$$I_L \leq I_{L+1}.$$

The implication

$$1 \leq L < \lfloor N/2 \rfloor \implies 2L + 1 \leq N$$

shows that this contains the range of [CPGSV16, Proposition 4.5, line 801].

Proof. [✓ Proof](#) Apply strong subadditivity to the reduced state of the first $N - L$ spins, split into three contiguous segments of lengths 1, L , and $N - 2L - 1$. Tracing the appropriate segments yields the four block entropies S_{N-L} , S_L , S_{L+1} , and S_{N-L-1} , where the cyclic invariance of $\rho^{(N)}(M)$ identifies the entropy of a contiguous block with that of any block of the same length. Strong subadditivity then gives $S_{N-L} + S_L \leq S_{L+1} + S_{N-L-1}$, which rearranges to $I_L \leq I_{L+1}$ after subtracting S_N . $\square$

Lemma 23.4.7 (Nonnegativity of block entropies). [Lean](#) [✓ Stmt](#) For a normalized positive semidefinite finite-chain operator, every block entropy is nonnegative: $0 \leq S_m$.

Proof. [✓ Proof](#) The reduced state of a block is a density matrix, and the von Neumann entropy of a density matrix is nonnegative. $\square$

Lemma 23.4.8 (Nonnegativity of the mutual information). [Lean](#) [✓ Stmt](#) Let M generate an MPDO whose finite-chain operator $\rho^{(N)}(M)$ is positive semidefinite with nonzero trace. For every block length $L \leq N$,

$$0 \leq I_L.$$

Proof. [✓ Proof](#) Apply strong subadditivity with a trivial middle subsystem (equivalently, subadditivity) to the bipartition of the chain into the first L and last $N-L$ spins: $S_N + S_0 \leq S_L + S_{N-L}$. Since $S_0 \geq 0$ (Lemma 23.4.7), rearranging gives $I_L = S_L + S_{N-L} - S_N \geq S_0 \geq 0$. $\square$

Lemma 23.4.9 (Symmetry of the mutual information across the cut). [Lean](#) [✓ Stmt](#) For every block length $L \leq N$, the mutual information of the first L spins equals that of their complement:

$$I_L = I_{N-L}.$$

Proof. [✓ Proof](#) Both sides equal $S_L + S_{N-L} - S_N$ by the symmetric definition $I_L = S_L + S_{N-L} - S_N$, using $N - (N - L) = L$. $\square$

Lemma 23.4.10 (Entropy of the empty block). [Lean](#) [✓ Stmt](#) For a normalized positive semidefinite finite-chain operator, the block entropy of the empty block vanishes: $S_0 = 0$.

Proof. [✓ Proof](#) The reduced state of zero spins is one-dimensional, so its rank is one and the rank bound on the entropy gives $S_0 \leq \log 1 = 0$; with $S_0 \geq 0$ this forces $S_0 = 0$. $\square$

Lemma 23.4.11 (Empty-block mutual information vanishes). [Lean](#) [✓ Stmt](#) For a normalized positive semidefinite finite-chain operator, the mutual information of the empty block vanishes: $I_0 = 0$.

Proof. [✓ Proof](#) By definition $I_0 = S_0 + S_N - S_N = S_0$, which is zero by Lemma 23.4.10. $\square$

Remark 23.4.12 (Boundedness of the mutual information). The monotone sequence I_L is bounded by a constant depending only on the bond dimension, $I_L \leq 4 \log D$, so it converges as $L \rightarrow \infty$. For a general mixed MPDO this bound is not proved in [CPGSV16]; that reference attributes it (line 1319) to the mutual-information area law of Wolf et al., used there as an external result. The pure-state instance is Proposition 23.4.32 below.

Definition 23.4.13 (Block entropy and area-law saturation for pure states). [Lean](#) [Lean](#) [✓ Stmt](#) For an MPS tensor A , a system size N , and a block length $L \leq N$, let

$$\sigma^{(N)}(A) = \frac{|V^{(N)}(A)\rangle\langle V^{(N)}(A)|}{\|V^{(N)}(A)\|^2}, \quad \rho_L^{(N)}(A) = \text{tr}_{N \setminus L}[\sigma^{(N)}(A)]$$

denote the normalized state and the reduced state of the first L spins. The L -block entropy is

$$S_L^{(N)}(A) = S(\rho_L^{(N)}(A)).$$

The tensor saturates the area law if $S_L^{(N)}(A) = S_{L+1}^{(N)}(A)$ for all $1 \leq L < \lfloor N/2 \rfloor$ and all N . This is [CPGSV16, Definition 3.13, line 600].

Definition 23.4.14 (Purification tensor). [Lean](#) [✓ Stmt](#) For an MPS tensor A on bond space $\mathbb{C}^D$, define the associated MPO tensor M on bond space $\mathbb{C}^{D^2}$ by

$$M^{ij} = A^i \otimes \bar{A}^j.$$

This is the single-ancilla purification tensor of [\[CPGSV16\]](#).

Lemma 23.4.15 (Pure state as a purification MPDO). [Lean](#) [✓ Stmt](#) For an MPS tensor A , the pure-state operator $|V^{(N)}(A)\rangle\langle V^{(N)}(A)|$ equals the MPDO generated by the tensor $M^{ij} = A^i \otimes \bar{A}^j$, acting on a bond space of dimension D^2 :

$$\rho^{(N)}(A \otimes \bar{A}) = |V^{(N)}(A)\rangle\langle V^{(N)}(A)|.$$

This is the single-ancilla case of the purification picture of [\[CPGSV16\]](#); it lets the MPDO block-entropy theory apply to pure-state block entropies.

Proof. [✓ Proof](#) The word evaluation of the doubled tensor is the Kronecker product of the word evaluation of A and its conjugate,

$$M^{\sigma,\tau} = A^\sigma \otimes \bar{A}^\tau,$$

so its trace factors as $\text{tr}(M^{\sigma,\tau}) = \text{tr}(A^\sigma) \text{tr}(\bar{A}^\tau)$, i.e.

$$\overline{V^{(N)}(A)}_\tau V^{(N)}(A)_\sigma,$$

the corresponding matrix element of $|V^{(N)}(A)\rangle\langle V^{(N)}(A)|$. □

Lemma 23.4.16 (Block entropy of the purification tensor). [Lean](#) [✓ Stmt](#) For an MPS tensor A , system size N , and block length $L \leq N$, the MPDO block entropy of the tensor $M^{ij} = A^i \otimes \bar{A}^j$ agrees with the pure-state block entropy:

$$S_L^{(N)}(M) = S_L^{(N)}(A).$$

Proof. [✓ Proof](#) Lemma 23.4.15 gives $\rho^{(N)}(M) = |V^{(N)}(A)\rangle\langle V^{(N)}(A)|$, hence

$$\sigma^{(N)}(M) = \frac{\rho^{(N)}(M)}{\text{tr}[\rho^{(N)}(M)]} = \frac{|V^{(N)}(A)\rangle\langle V^{(N)}(A)|}{\|V^{(N)}(A)\|^2} = \sigma^{(N)}(A).$$

Therefore the reduced block states agree:

$$\rho_L^{(N)}(M) = \text{tr}_{N \setminus L}[\sigma^{(N)}(M)] = \text{tr}_{N \setminus L}[\sigma^{(N)}(A)] = \rho_L^{(N)}(A).$$

Thus

$$S_L^{(N)}(M) = S(\rho_L^{(N)}(M)) = S(\rho_L^{(N)}(A)) = S_L^{(N)}(A). \quad \square$$

Lemma 23.4.17 (Schmidt symmetry of the block entropy). [Lean](#) [✓ Stmt](#) For an MPS tensor A and a block length $L \leq N$, the block entropies of a block and its complement coincide:

$$S_L^{(N)}(A) = S_{N-L}^{(N)}(A).$$

Proof. [✓ Proof](#) The reduced state of the first L spins is $\rho_L \propto WW^\dagger$, where W is the matrix of amplitudes $\langle uw | V^{(N)}(A) \rangle$ with u the first L spins and w the remaining $N - L$. By cyclic invariance of the amplitudes, the reduced state of the complement is $\propto \bar{W}^\dagger \bar{W}$. Cyclic invariance of the entropy, $S(WW^\dagger) = S(W^\dagger W)$, together with invariance of the entropy under entrywise conjugation of a Hermitian matrix, $S(\bar{W}^\dagger \bar{W}) = S(W^\dagger W)$, gives $S_L^{(N)}(A) = S_{N-L}^{(N)}(A)$. □

Proposition 23.4.18 (Pure-state area-law monotonicity). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor with $V^{(N)}(A) \neq 0$. For every block length L with $2L + 1 \leq N$ (covering $1 \leq L < \lfloor N/2 \rfloor$),

$$S_L^{(N)}(A) \leq S_{L+1}^{(N)}(A).$$

This is the pure-state area law of [CPGSV16, Section 3, line 599].

Proof. [✓ Proof](#) Apply the block-entropy form of strong subadditivity (the inequality underlying Proposition 23.4.6) to the purification MPDO of A , giving $S_{N-L} + S_L \leq S_{L+1} + S_{N-L-1}$. By the Schmidt symmetry $S_{N-L} = S_L$ and $S_{N-L-1} = S_{L+1}$, this reduces to $S_L \leq S_{L+1}$. $\square$

Lemma 23.4.19 (Operator-Schmidt Gram is a transfer-map power). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let A be an MPS tensor of bond dimension D . For a length- ℓ configuration σ and a length- m configuration τ , write the operator-Schmidt factors as

$$(L_\ell)_{\sigma, (a_1, a_2)} = (A^\sigma)_{a_1 a_2}, \quad (R_m)_{(p_1, p_2), \tau} = (A^\tau)_{p_2 p_1}.$$

The $D^2 \times D^2$ Gram matrix of the left factor equals the ℓ -fold transfer map evaluated on a matrix unit:

$$(L_\ell^\dagger L_\ell)_{(a_1, a_2), (b_1, b_2)} = \sum_\sigma \overline{(A^\sigma)_{a_1 a_2}} (A^\sigma)_{b_1 b_2} = (\mathbb{E}_A^\ell(|b_2\rangle\langle a_2|))_{b_1, a_1},$$

the sum running over length- ℓ configurations σ . The Gram of the right factor R_m of an m -spin complement satisfies the bond-swapped identity

$$(R_m R_m^\dagger)_{(p_1, p_2), (q_1, q_2)} = \sum_\tau (A^\tau)_{p_2 p_1} \overline{(A^\tau)_{q_2 q_1}} = (\mathbb{E}_A^m(|p_1\rangle\langle q_1|))_{p_2, q_2},$$

the sum running over length- m configurations τ .

Proof. [✓ Proof](#) The left Gram entry is $\sum_\sigma \overline{(A^\sigma)_{a_1 a_2}} (A^\sigma)_{b_1 b_2}$ by definition of L_ℓ . Expanding $\mathbb{E}_A^\ell(|b_2\rangle\langle a_2|) = \sum_\sigma A^\sigma |b_2\rangle\langle a_2| (A^\sigma)^\dagger$ and reading off the (b_1, a_1) entry gives the same sum, since $(A^\sigma |b_2\rangle\langle a_2| (A^\sigma)^\dagger)_{b_1, a_1} = (A^\sigma)_{b_1 b_2} \overline{(A^\sigma)_{a_1 a_2}}$. The right-factor formula is the same expansion with the two boundary bonds interchanged: the (p, q) entry of $R_m R_m^\dagger$ is $\sum_\tau (A^\tau)_{p_2 p_1} \overline{(A^\tau)_{q_2 q_1}}$, which is the (p_2, q_2) entry of $\mathbb{E}_A^m(|p_1\rangle\langle q_1|)$. $\square$

Lemma 23.4.20 (RFP collapse of the operator-Schmidt Grams). [Lean](#) [Lean](#) [✓ Stmt](#) Let A be a renormalization fixed point. For every $L \geq 1$ and $m \geq 1$, the operator-Schmidt Grams of the L -block and of the m -site complement are the one-site Grams:

$$L_L^\dagger L_L = L_1^\dagger L_1, \quad R_m R_m^\dagger = R_1 R_1^\dagger.$$

Proof. [✓ Proof](#) By definition, a renormalization fixed point satisfies $\mathbb{E}_A^2 = \mathbb{E}_A$, so $\mathbb{E}_A^k = \mathbb{E}_A$ for every $k \geq 1$. Applying Lemma 23.4.19 to the left and right Gram entries gives the two displayed identities. $\square$

Lemma 23.4.21 (Pure block entropy as a bond-environment charpoly sum). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor, let $L \leq N$, put $m = N - L$, and write $c = (\text{tr } \rho^{(N)})^{-1}$. Then the pure block entropy is the charpoly-root entropy sum of the $D^2 \times D^2$ bond environment:

$$S_L^{(N)}(A) = \sum_{\lambda \in \text{roots } \chi_{c(R_m R_m^\dagger)(L_L^\dagger L_L)}} -\text{Re}(\lambda) \log \text{Re}(\lambda),$$

with roots counted with algebraic multiplicity.

Proof. ✓ Proof The reduced block state has the exact factorization

$$\rho_L^{(N)}(A) = c L_L (R_m R_m^\dagger) L_L^\dagger.$$

Cyclic invariance of the charpoly-root entropy sum gives

$$S(c L_L (R_m R_m^\dagger) L_L^\dagger) = S(c (R_m R_m^\dagger) (L_L^\dagger L_L)),$$

which is the displayed formula. □

Theorem 23.4.22 (A renormalization fixed point saturates the area law). Lean ✓ Stmt *If an MPS tensor A is a renormalization fixed point, then it saturates the area law: for every chain length N and every block length $1 \leq L < \lfloor N/2 \rfloor$, the pure-state block entropy is constant in the block size,*

$$S_L^{(N)}(A) = S_{L+1}^{(N)}(A).$$

The source statement is [CPGSV16, Proposition 3.14, lines 606–608]. It assumes canonical form; the theorem above is the stronger pure-state implication obtained from transfer-map idempotence alone.

Proof. ✓ Proof Write $c = (\text{tr } \rho^{(N)})^{-1}$. Lemma 23.4.21 computes the entropy from the bond environment:

$$S_L^{(N)}(A) = \sum_{\lambda \in \text{roots } \chi_{c(R_{N-L} R_{N-L}^\dagger)(L_L^\dagger L_L)}} -\text{Re}(\lambda) \log \text{Re}(\lambda).$$

Since $1 \leq L < \lfloor N/2 \rfloor$, the four lengths L , $L+1$, $N-L$, and $N-(L+1)$ are positive. The RFP Gram-collapse lemma gives

$$L_L^\dagger L_L = L_1^\dagger L_1, \quad L_{L+1}^\dagger L_{L+1} = L_1^\dagger L_1,$$

and

$$R_{N-L} R_{N-L}^\dagger = R_1 R_1^\dagger, \quad R_{N-(L+1)} R_{N-(L+1)}^\dagger = R_1 R_1^\dagger.$$

Therefore the two bond environments agree:

$$(R_{N-L} R_{N-L}^\dagger)(L_L^\dagger L_L) = (R_1 R_1^\dagger)(L_1^\dagger L_1) = (R_{N-(L+1)} R_{N-(L+1)}^\dagger)(L_{L+1}^\dagger L_{L+1}).$$

The scalar c depends only on N , so the two charpoly-root entropy sums are equal. Hence $S_L^{(N)}(A) = S_{L+1}^{(N)}(A)$. □

Theorem 23.4.23 (A renormalization fixed point has a constant block-entropy chain). Lean ✓ Stmt *If an MPS tensor A is a renormalization fixed point, then all block entropies in the range $1 \leq L \leq \lfloor N/2 \rfloor$ coincide:*

$$S_1^{(N)}(A) = S_2^{(N)}(A) = \dots = S_{\lfloor N/2 \rfloor}^{(N)}(A).$$

This is the explicit constant-entropy chain of the area-law saturation [CPGSV16, Definition 3.13, line 600], under the fixed-point hypothesis.

Proof. ✓ Proof The fixed point saturates the area law (Theorem 23.4.22), so by the telescoping of saturation (Lemma 23.4.30) the block entropies at any two lengths $1 \leq L, L' \leq \lfloor N/2 \rfloor$ agree:

$$S_L^{(N)}(A) = S_{L'}^{(N)}(A).$$

□

Lemma 23.4.24 (Operator-Schmidt rank bound). [Lean](#) [✓ Stmt](#) For an MPS tensor A of bond dimension D and a block length $L \leq N$, the reduced state $\rho_L^{(N)}(A)$ of the first L spins has rank at most D^2 .

Proof. [✓ Proof](#) The reduced state is $\rho_L \propto WW^\dagger$, where the amplitude matrix $W(u, w) = \langle u w | V^{(N)}(A) \rangle = \text{tr}(A^{u_1} \dots A^{u_L} A^{w_1} \dots A^{w_{N-L}})$ factors through the two bond pairs cut by the block boundary. Writing the trace as a sum over the bond index pair $(a, b) \in \{1, \dots, D\}^2$ exhibits $W = L'R'$ with L' having D^2 columns, so $\text{rank } \rho_L \leq \text{rank } W \leq D^2$. $\square$

Lemma 23.4.25 (The reduced pure block state is a density matrix). [Lean](#) [Lean](#) [✓ Stmt](#) For an MPS tensor A with $V^{(N)}(A) \neq 0$, the reduced state $\rho_L^{(N)}(A)$ of the first L spins is a density matrix: it is positive semidefinite and has unit trace.

Proof. [✓ Proof](#) Positive semidefiniteness is preserved by the partial trace of the positive semidefinite normalized pure state, and the partial trace preserves the trace, which the normalization fixes to one. $\square$

Proposition 23.4.26 (Pure-state area-law upper bound). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor of bond dimension $D \geq 1$ with $V^{(N)}(A) \neq 0$. For every block length $L \leq N$,

$$S_L^{(N)}(A) \leq 2 \log D,$$

uniformly in L and N . This is the area law of [CPGSV16, Section 3, line 599]: the block entropy is bounded by a constant independent of the block size.

Proof. [✓ Proof](#) The reduced state $\rho_L^{(N)}(A)$ is a density matrix (Lemma 23.4.25) of rank at most D^2 by Lemma 23.4.24. By the rank bound on the entropy (Theorem 19.1.8), $S_L^{(N)}(A) \leq \log(D^2) = 2 \log D$. $\square$

Lemma 23.4.27 (Nonnegativity of the pure-state block entropy). [Lean](#) [✓ Stmt](#) For an MPS tensor A with $V^{(N)}(A) \neq 0$ and every block length $L \leq N$, the pure-state block entropy is nonnegative: $0 \leq S_L^{(N)}(A)$.

Proof. [✓ Proof](#) The reduced state $\rho_L^{(N)}(A)$ is a density matrix (Lemma 23.4.25), and the von Neumann entropy of a density matrix is nonnegative. $\square$

Lemma 23.4.28 (Full-block pure-state entropy vanishes). [Lean](#) [✓ Stmt](#) For an MPS tensor A with $V^{(N)}(A) \neq 0$, the entropy of the whole chain vanishes:

$$S_N^{(N)}(A) = 0.$$

Proof. [✓ Proof](#) The normalized operator is the rank-one pure state

$$\frac{|V^{(N)}(A)\rangle\langle V^{(N)}(A)|}{\langle V^{(N)}(A), V^{(N)}(A) \rangle},$$

after reindexing the full block. A pure state has von Neumann entropy zero. $\square$

Lemma 23.4.29 (Empty-block pure-state entropy vanishes). [Lean](#) [✓ Stmt](#) For an MPS tensor A with $V^{(N)}(A) \neq 0$, the pure-state block entropy of the empty block vanishes: $S_0^{(N)}(A) = 0$.

Proof. [✓ Proof](#) By the Schmidt symmetry $S_0^{(N)}(A) = S_N^{(N)}(A)$, and the entropy of the full system vanishes for a pure state (Lemma 23.4.28). $\square$

Lemma 23.4.30 (Saturation telescopes to a constant block entropy). [Lean](#) [✓ Stmt](#) If an MPS tensor A saturates the area law, then the block entropies coincide for all $1 \leq L, L' \leq \lfloor N/2 \rfloor$:

$$S_L^{(N)}(A) = S_{L'}^{(N)}(A).$$

Proof. [✓ Proof](#) The defining condition supplies the single-step equalities $S_m^{(N)}(A) = S_{m+1}^{(N)}(A)$ for $1 \leq m < \lfloor N/2 \rfloor$. Climbing one step at a time from the smaller of L and L' up to the larger composes these into the stated equality. $\square$

Lemma 23.4.31 (Pure-state mutual information is twice the block entropy). [Lean](#) [✓ Stmt](#) For an MPS tensor A with $V^{(N)}(A) \neq 0$, the mutual information of the purification MPDO equals twice the pure-state block entropy:

$$I_L = 2 S_L^{(N)}(A).$$

Proof. [✓ Proof](#) By definition $I_L = S_L + S_{N-L} - S_N$ for the generated state. The global entropy of the pure state vanishes (Lemma 23.4.28), and the Schmidt symmetry (Lemma 23.4.17) gives $S_{N-L} = S_L$, so $I_L = 2 S_L$. $\square$

Proposition 23.4.32 (Pure-state mutual-information bound). [Lean](#) [✓ Stmt](#) Let A be an MPS tensor of bond dimension $D \geq 1$ with $V^{(N)}(A) \neq 0$. For the purification MPDO of A , whose generated operator is the pure state $|V^{(N)}(A)\rangle\langle V^{(N)}(A)|$, the mutual information between a block of L spins and the rest satisfies

$$I_L \leq 4 \log D,$$

uniformly in L and N .

Proof. [✓ Proof](#) For a pure state $I_L = 2 S_L^{(N)}(A)$, since the global entropy S_N vanishes and a block and its complement share the same Schmidt spectrum. The block-entropy bound $S_L^{(N)}(A) \leq 2 \log D$ (Proposition 23.4.26) then gives $I_L \leq 4 \log D$. $\square$

23.5 Simple local structure from SAL and ZCL

23.5.1 Markov decomposition and inverse-map sector factorization

For the simple MPDO case in Appendix C.2 of [CPGSV16], the strong-area-law hypothesis is used locally through the equality case of strong subadditivity for a normalized three-site reduced state.

Definition 23.5.1.1 (Local η -structure for a three-site reduced state). [Lean](#) [✓ Stmt](#) For a three-site density operator ρ_{ABC} , the local η -structure is the quantum Markov decomposition on the middle subsystem B produced by equality in strong subadditivity. Concretely, after a unitary change of basis on B , one has a finite direct-sum decomposition

$$B = \bigoplus_k (b_1^{(k)} \otimes b_2^{(k)})$$

such that

$$\rho_{ABC} = \bigoplus_k \rho_{Ab_1}^{(k)} \otimes \rho_{b_2C}^{(k)}.$$

Theorem 23.5.1.2 (Saturation gives equality in strong subadditivity). [Lean](#) [Lean](#) [✓ Stmt](#) Let $N \geq 4$, and divide the first $N - 1$ sites of the normalized periodic state into consecutive regions A, B, C of lengths $1, 1, N - 3$. If the tensor saturates the area law, then

$$S(ABC) + S(B) = S(AB) + S(BC).$$

In particular, the three-site reduced state of the four-site chain satisfies equality in strong subadditivity.

Proof. [✓ Proof](#) By saturation, $I_1 = I_2$. Expanding the two mutual informations and cancelling the entropy of the full chain gives

$$S_{N-1} + S_1 = S_2 + S_{N-2},$$

which is the stated equality for the regions A, B, C . $\square$

Theorem 23.5.1.3 (SAL implies local η -structure). [Lean](#) [✓ Stmt](#) If a normalized three-site reduced state satisfies the equality case of strong subadditivity — the local entropy form of the simple-MPDO strong area law — then it admits the local η -structure of Definition 23.5.1.1.

Proof. [✓ Proof](#) This is exactly the forward implication of the Hayashi equality characterization: equality in strong subadditivity yields a quantum Markov decomposition on the middle subsystem. $\square$

Theorem 23.5.1.4 (Four-site SAL marginal has a Hayashi decomposition). [Lean](#) [✓ Stmt](#) Let K satisfy the strong area law, and let $\sigma_3^{(4)}(K) = \text{tr}_4 \sigma^{(4)}(K)$ be the three-site marginal of its normalized four-site periodic state. Then $\sigma_3^{(4)}(K)$ admits a quantum-Markov decomposition on its middle site.

This assertion is restricted to the four-site chain. It is the $N = 4$ specialization of the three-site decomposition in [CPGSV16, Appendix C.2, lines 1351–1363], which treats all chain lengths in the range of its four-region argument.

Proof. [✓ Proof](#) Let ρ^{flat} denote the three-site marginal with index set $(\{0, \dots, d^1 - 1\})^3$. By Theorem 23.5.1.2,

$$S(\rho_{ABC}^{\text{flat}}) + S(\rho_B^{\text{flat}}) = S(\rho_{AB}^{\text{flat}}) + S(\rho_{BC}^{\text{flat}}).$$

Each singleton block has the canonical identification

$$e_{\text{site}} : \{0, \dots, d-1\} \simeq \{0, \dots, d^1 - 1\}, \quad E = e_{\text{site}} \times e_{\text{site}} \times e_{\text{site}},$$

and reindexing ρ^{flat} along E gives $\sigma_3^{(4)}(K)$. Hence Theorem 19.3.6 preserves the displayed equality. Positivity and unit trace follow from the normalized reduced state, so Theorem 23.5.1.3 gives the required quantum Markov decomposition. $\square$

Definition 23.5.1.5 (Injective inverse map for a simple MPO tensor). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A simple MPO tensor K is called injective if its doubled-index MPS tensor is injective. For such a tensor, the chosen decomposition map of the doubled-index MPS furnishes a concrete inverse tensor K^{-1} together with right- and left-physical realization maps that turn virtual bond insertions back into local physical operators.

Theorem 23.5.1.6 (Inverse-map identities for an injective simple MPO tensor). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For an injective simple MPO tensor, contracting K^{-1} with K recovers the matrix units on the virtual bond space. Likewise, every right virtual insertion admits a physical realization, this realization is multiplicative, and the analogous left-insertion statement also holds.

For an injective simple tensor, contracting K^{-1} with K through the physical index identifies the corresponding virtual legs. In components, this is the matrix-unit identity $\sum_p (K^{-1})_p^{\alpha,\beta} K^p = E_{\alpha,\beta}$:

$$K^{-1}K = \mathbb{1}$$

This is the identity in [CPGSV16, Appendix C.2, lines 1376–1380].

Applying this inverse map to the doubled physical index and closing its virtual legs against a further copy of K recovers K :

$$K K^{-1} K = K$$

The doubled physical index is represented by a single physical line. This is the single-block identity of [CPGSV16, Appendix C.2, lines 1671–1676].

Definition 23.5.1.7 (Normalized fourth-site virtual tail). Lean ✓ Stmt Let K be an MPO tensor and set $Z_4 = \text{tr}[\rho^{(4)}(K)]$. Its normalized fourth-site virtual tail is

$$R_4 := Z_4^{-1} \mathcal{T}_K = \text{tr}[\rho^{(4)}(K)]^{-1} \sum_{i_4=0}^{d-1} K^{i_4 i_4}.$$

Thus the contraction of the fourth ket and bra indices is incorporated into the virtual closing matrix. This is the $N = 4$ specialization of the normalization convention and three-site marginal in [CPGSV16, lines 792–793 and 1343–1348].

Theorem 23.5.1.8 (Three-site marginal closed by the normalized fourth-site tail). Lean ✓ Stmt Let $u = (i_1, i_2, i_3)$ and $v = (j_1, j_2, j_3)$ be physical words of length three. The corresponding entry of the first-three-site marginal of the normalized four-site periodic operator is

$$\sigma_3^{(4)}(K)_{u,v} = \text{tr}(K^{u,v} R_4) = \text{tr}(K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} R_4).$$

$$\text{tr}_4(K_4) = \sum_{\alpha} L_{\alpha} \otimes B_{\alpha} \otimes R_{\alpha}$$

Figure 23.1: Tracing the fourth site of the normalized four-site periodic MPO leaves the physical legs of sites 1, 2, 3 open and replaces the fourth site by the virtual closing operator $R_4 = \text{tr}[\rho^{(4)}(K)]^{-1} \sum_{i_4} K^{i_4 i_4}$; compare the normalization convention and three-site marginal in [CPGSV16, lines 792–793 and 1343–1348].

Proof. ✓ Proof Expanding the partial trace over the fourth site and factoring each length-four word at its last letter gives

$$\sigma_3^{(4)}(K)_{u,v} = Z_4^{-1} \sum_{i_4=0}^{d-1} \text{tr}(K^{u,v} K^{i_4 i_4}).$$

Linearity of the trace now yields

$$Z_4^{-1} \sum_{i_4=0}^{d-1} \text{tr}(K^{u,v} K^{i_4 i_4}) = \text{tr} \left(K^{u,v} \left(Z_4^{-1} \sum_{i_4=0}^{d-1} K^{i_4 i_4} \right) \right) = \text{tr}(K^{u,v} R_4).$$

□

Theorem 23.5.1.9 (Non-vanishing of the normalized fourth-site tail). [Lean](#) [✓ Stmt](#) If $Z_4 = \text{tr}[\rho^{(4)}(K)] \neq 0$, then $R_4 \neq 0$.

Proof. [✓ Proof](#) If $R_4 = 0$, then the tail formula makes every entry of $\sigma_3^{(4)}(K)$ vanish, so

$$\text{tr}[\sigma_3^{(4)}(K)] = 0.$$

On the other hand, $Z_4 \neq 0$ implies

$$\text{tr}[\sigma_3^{(4)}(K)] = 1,$$

a contradiction. □

Theorem 23.5.1.10 (A nonzero entry of the normalized fourth-site tail). [Lean](#) [✓ Stmt](#) If $Z_4 \neq 0$, there exist virtual indices β and α such that

$$(R_4)_{\beta,\alpha} \neq 0.$$

These are the indices selected before the sector factorization in [CPGSV16, Appendix C.2, lines 1431–1437].

Proof. [✓ Proof](#) If every matrix entry of R_4 vanished, then R_4 would be the zero matrix, contrary to the preceding theorem. □

Definition 23.5.1.11 (Three-site inverse-map contraction). [Lean](#) [✓ Stmt](#) Let K be an injective simple MPO tensor, and let $R \in M_D(\mathbb{C})$ be the virtual contraction of the sites following a three-site word. For virtual indices $\alpha_1, \beta_1, \alpha_3, \beta_3$, define

$$C_{\alpha_1, \beta_1, \alpha_3, \beta_3}(p_2; R) = \sum_{p_1, p_3} (K^{-1})_{p_1}^{\alpha_1, \beta_1} (K^{-1})_{p_3}^{\alpha_3, \beta_3} \text{tr}(K^{p_1} K^{p_2} K^{p_3} R).$$

This is the contraction used in the proof of [CPGSV16, Appendix C.2, lines 1415–1438].

Theorem 23.5.1.12 (Evaluation of the three-site inverse-map contraction). [Lean](#) [✓ Stmt](#) For every middle physical index p_2 ,

$$C_{\alpha_1, \beta_1, \alpha_3, \beta_3}(p_2; R) = K_{\beta_1, \alpha_3}^{p_2} R_{\beta_3, \alpha_1}.$$

$$(K^{-1} \otimes \mathbb{1} \otimes K^{-1})K_3 = \sum_{\alpha, \beta} L_\alpha \otimes B_{\alpha, \beta} \otimes R_\beta$$

Figure 23.2: Applying K^{-1} at the first and third sites collapses the three-site contraction to $K_{\beta_1, \alpha_3}^{p_2} R_{\beta_3, \alpha_1}$, where $p_2 = (i_2, j_2)$ is the doubled physical index; compare [CPGSV16, Appendix C.2, lines 1415–1438].

Proof. [✓ Proof](#) Move the two finite sums through the trace. By the inverse-map identity

$$\sum_p (K^{-1})_p^{\alpha, \beta} K^p = E_{\alpha, \beta},$$

the double sum collapses to

$$\text{tr}(E_{\alpha_1, \beta_1} K^{p_2} E_{\alpha_3, \beta_3} R) = K_{\beta_1, \alpha_3}^{p_2} R_{\beta_3, \alpha_1}.$$

□

Definition 23.5.1.13 (Three-site closure against a virtual tail). [Lean](#) [✓ Stmt](#) Let K be an MPO tensor and let $R \in M_D(\mathbb{C})$. A tripartite operator ρ is the three-site closure of K against R if

$$\rho_{i_1 i_2 i_3, j_1 j_2 j_3} = \text{tr}(K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} R).$$

This is the algebraic form of the three-site reduced operator in [CPGSV16, Appendix C.2, lines 1422–1433].

Definition 23.5.1.14 (Physical slice and Hayashi inverse factors). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Fix virtual indices β, α and set

$$\kappa_{\beta, \alpha}(i, j) = K_{\beta, \alpha}^{ij}.$$

For a Hayashi sector k , define the two inverse factors

$$\begin{aligned} A_{\alpha_1, \beta_1}^{(k)}(l, l') &:= \sum_{i_1, j_1} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (\rho_{Ab_1}^{(k)})_{(i_1, l), (j_1, l')}, \\ B_{\alpha_3, \beta_3}^{(k)}(r, r') &:= \sum_{i_3, j_3} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} (\rho_{b_2 C}^{(k)})_{(r, i_3), (r', j_3)}. \end{aligned}$$

These are the two factors in the sectorwise inverse-map calculation of [CPGSV16, Appendix C.2, lines 1407–1428].

Theorem 23.5.1.15 (Outer inverse-map contraction of a closure). [Lean](#) [Lean](#) [✓ Stmt](#) Let K be injective and let ρ be its three-site closure against R . Applying the inverse tensor at the two outer sites collapses the closure to one entry of the middle physical slice and one entry of the tail:

$$\sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} = K_{\beta_1, \alpha_3}^{i_2 j_2} R_{\beta_3, \alpha_1}.$$

Consequently, for any matrix U on the middle site,

$$R_{\beta_3, \alpha_1} (U \kappa_{\beta_1, \alpha_3} U^\dagger)_{b, b'} = \sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} \sum_{i_2, j_2} U_{b, i_2} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} \overline{U_{b', j_2}}.$$

Proof. [✓ Proof](#) Substituting the three-site closure relation

$$\rho_{i_1 i_2 i_3, j_1 j_2 j_3} = \text{tr}(K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} R)$$

converts the left-hand side of the first identity to $C_{\alpha_1, \beta_1, \alpha_3, \beta_3}((i_2, j_2); R)$. Theorem 23.5.1.12 then gives

$$C_{\alpha_1, \beta_1, \alpha_3, \beta_3}((i_2, j_2); R) = K_{\beta_1, \alpha_3}^{i_2 j_2} R_{\beta_3, \alpha_1},$$

which is the first identity. The second identity follows by expanding the matrix product entrywise,

$$(U \kappa_{\beta_1, \alpha_3} U^\dagger)_{b, b'} = \sum_{i_2, j_2} U_{b, i_2} \kappa_{\beta_1, \alpha_3}(i_2, j_2) \overline{U_{b', j_2}},$$

multiplying by R_{β_3, α_1} , substituting the first identity in the form

$$R_{\beta_3, \alpha_1} \kappa_{\beta_1, \alpha_3}(i_2, j_2) = \sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} \rho_{i_1 i_2 i_3, j_1 j_2 j_3},$$

and reordering the finite sums. □

Theorem 23.5.1.16 (Sectorwise inverse-map comparison). [Lean](#) [✓ Stmt](#) Suppose that K is injective, ρ is its three-site closure against R , and a Hayashi decomposition of ρ has been chosen. Write

$$\tilde{\kappa}_{\beta_1, \alpha_3}^{(k)}((l, r), (l', r')) := [U_B \kappa_{\beta_1, \alpha_3} U_B^\dagger]_{(k, l, r), (k, l', r')}.$$

Then every diagonal sector satisfies

$$R_{\beta_3, \alpha_1} \tilde{\kappa}_{\beta_1, \alpha_3}^{(k)}((l, r), (l', r')) = p_k A_{\alpha_1, \beta_1}^{(k)}(l, l') B_{\alpha_3, \beta_3}^{(k)}(r, r').$$

The probability p_k remains explicit because both sector density operators are normalized. No zero-correlation-length or non-vanishing hypothesis is used in this comparison.

The same comparison may be read directly as the following compact identity:

$$R_{\beta_3, \alpha_1} \cdot \begin{array}{c} k; l, r \\ | \\ \beta_1 \text{ --- } \bullet \text{ --- } \alpha_3 \\ | \\ k; l', r' \end{array} \xrightarrow{\tilde{\kappa}^{(k)}} = p_k \boxed{A_{\alpha_1, \beta_1}^{(k)}} \otimes \boxed{B_{\alpha_3, \beta_3}^{(k)}} \begin{array}{c} l, l' \\ | \\ r, r' \end{array}$$

Here the closing entry R_{β_3, α_1} multiplies the physical slice in the basis adapted to sector k ; compare [CPGSV16, Appendix C.2, lines 1415–1438].

Proof. [✓ Proof](#) At fixed indices (k, l, r) and (k, l', r') , the Hayashi block equality is

$$\sum_{i_2, j_2} [U_B]_{(k, l, r), i_2} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} \overline{[U_B]_{(k, l', r'), j_2}} = p_k (\rho_{Ab_1}^{(k)})_{(i_1, l), (j_1, l')} (\rho_{b_2 C}^{(k)})_{(r, i_3), (r', j_3)}.$$

For each i_2, j_2 , the preceding theorem gives

$$\sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} = K_{\beta_1, \alpha_3}^{i_2 j_2} R_{\beta_3, \alpha_1}.$$

Apply $\sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} (-)$ to the Hayashi block equality. Substituting the preceding display and interchanging the finite sums gives

$$\begin{aligned} & \sum_{i_1, j_1, i_3, j_3} (K^{-1})_{(i_1, j_1)}^{\alpha_1, \beta_1} (K^{-1})_{(i_3, j_3)}^{\alpha_3, \beta_3} \\ & \times \left(\sum_{i_2, j_2} [U_B]_{(k, l, r), i_2} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} \overline{[U_B]_{(k, l', r'), j_2}} \right) \\ & = \sum_{i_2, j_2} [U_B]_{(k, l, r), i_2} K_{\beta_1, \alpha_3}^{i_2 j_2} R_{\beta_3, \alpha_1} \overline{[U_B]_{(k, l', r'), j_2}}. \end{aligned}$$

The right-hand side is

$$\begin{aligned} & R_{\beta_3, \alpha_1} \sum_{i_2, j_2} [U_B]_{(k, l, r), i_2} \kappa_{\beta_1, \alpha_3}(i_2, j_2) \overline{[U_B]_{(k, l', r'), j_2}} \\ & = R_{\beta_3, \alpha_1} \tilde{\kappa}_{\beta_1, \alpha_3}^{(k)}((l, r), (l', r')). \end{aligned}$$

The contraction of the right-hand side of the Hayashi block equality is

$$p_k A_{\alpha_1, \beta_1}^{(k)}(l, l') B_{\alpha_3, \beta_3}^{(k)}(r, r').$$

Since the Hayashi block equality holds, equating the two contractions gives

$$R_{\beta_3, \alpha_1} \tilde{\kappa}_{\beta_1, \alpha_3}^{(k)}((l, r), (l', r')) = p_k A_{\alpha_1, \beta_1}^{(k)}(l, l') B_{\alpha_3, \beta_3}^{(k)}(r, r').$$

□

Theorem 23.5.1.17 (The four-site marginal is a three-site closure). [Lean](#) [✓ Stmt](#) *Carried onto the tripartite site index, the three-site marginal of the normalized four-site chain is the three-site closure of K against the normalized fourth-site tail R_4 .*

Proof. [✓ Proof](#) The entry formula for $\sigma_3^{(4)}(K)$, carried onto the tripartite site index (i_1, i_2, i_3) , gives

$$\rho_{i_1 i_2 i_3, j_1 j_2 j_3} = \text{tr}(K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} R_4),$$

which is the three-site closure condition of Definition 23.5.1.13 against R_4 . □

Theorem 23.5.1.18 (Vanishing of the off-diagonal Hayashi sectors). [Lean](#) [✓ Stmt](#) *In the setting of Theorem 23.5.1.16, for all $k \neq k'$, every off-diagonal sector satisfies*

$$R_{\beta_3, \alpha_1} [U_B \kappa_{\beta_1, \alpha_3} U_B^\dagger]_{(k, l, r), (k', l', r')} = 0.$$

Together with the diagonal comparison this gives the direct-sum structure of the factorized tensor: the direct sum over sectors is inherited from the splitting of the middle site.

Proof. [✓ Proof](#) The Hayashi block equality at off-diagonal indices gives

$$\sum_{i_2, j_2} (U_B)_{b, i_2} \rho_{i_1 i_2 i_3, j_1 j_2 j_3} \overline{(U_B)_{b', j_2}} = 0$$

for b in sector k and b' in sector k' with $k \neq k'$. Substituting this vanishing into the expansion of Theorem 23.5.1.15 gives $R_{\beta_3, \alpha_1} (U_B \kappa_{\beta_1, \alpha_3} U_B^\dagger)_{b, b'} = 0$. □

Definition 23.5.1.19 (Sector tensors). [Lean](#) [Lean](#) [✓ Stmt](#) Fix outer virtual indices α_1, β_3 and a Hayashi sector k . The sector tensors are

$$(l_k)_{\beta_1} := R_{\beta_3, \alpha_1}^{-1} p_k A_{\alpha_1, \beta_1}^{(k)}, \quad (r_k)_{\alpha_3} := B_{\alpha_3, \beta_3}^{(k)}.$$

These are the tensors of the factorization displayed in [CPGSV16, Appendix C.2, lines 1435–1448].

Lemma 23.5.1.20 (Zero-weight left sector tensor). [Lean](#) [Lean](#) [✓ Stmt](#) *If $p_k = 0$, then*

$$(l_k)_\beta = 0$$

for every virtual index β . Consequently,

$$\eta_{h, k} = 0$$

for every sector h .

Proof. [✓ Proof](#) The scalar weight p_k is a factor of every matrix entry of $(l_k)_\beta$. Each neighboring operator entering sector k contains $(l_k)_\beta$ as its target factor. □

Theorem 23.5.1.21 (Sector factorization of an injective simple tensor). [Lean](#) [✓ Stmt](#) Suppose that K is injective, ρ is its three-site closure against R , a Hayashi decomposition of ρ has been chosen, and the outer indices satisfy $R_{\beta_3, \alpha_1} \neq 0$. Then for all virtual indices β_1, α_3 ,

$$U_B \kappa_{\beta_1, \alpha_3} U_B^\dagger = \bigoplus_k (l_k)_{\beta_1} \otimes (r_k)_{\alpha_3}.$$

The direct sum refers to the physical indices and is inherited from the splitting of the middle site. This is the factorization displayed at [CPGSV16, Appendix C.2, lines 1435–1448].

In the middle-site basis selected by U_B , the same factorization is displayed as

$$U_B \mathcal{K} U_B^\dagger = \bigoplus_k l_k \otimes r_k$$

Proof. [✓ Proof](#) On a diagonal sector, the sectorwise comparison gives $R_{\beta_3, \alpha_1} \tilde{\kappa}^{(k)} = p_k A^{(k)} B^{(k)}$, and dividing by the nonzero entry R_{β_3, α_1} produces $(l_k)_{\beta_1} \otimes (r_k)_{\alpha_3}$. On an off-diagonal sector, the product with R_{β_3, α_1} vanishes, and since $R_{\beta_3, \alpha_1} \neq 0$ the sector itself vanishes. $\square$

Theorem 23.5.1.22 (Existence of the sector tensors). [Lean](#) [✓ Stmt](#) Let K be injective with nonzero four-site trace, let ρ be a three-site closure of K against the normalized fourth-site tail R_4 , and choose a Hayashi decomposition of ρ . Then there exist sector tensors l_k and r_k such that, for all virtual indices β_1, α_3 ,

$$U_B \kappa_{\beta_1, \alpha_3} U_B^\dagger = \bigoplus_k (l_k)_{\beta_1} \otimes (r_k)_{\alpha_3}.$$

The neighboring operators $\eta_{k,h}$ are assembled from these tensors in Definition 23.5.2.1; the remaining steps of [CPGSV16, Appendix C.2, Lemma C.4] prove the all-length identity and primitivity.

Proof. [✓ Proof](#) A nonzero four-site trace supplies outer indices with $(R_4)_{\beta_3, \alpha_1} \neq 0$, and the sector factorization applies. $\square$

23.5.2 Neighboring operators and commuting bond products

Definition 23.5.2.1 (Neighboring operators). [Lean](#) [✓ Stmt](#) Fix outer virtual indices α_1, β_3 with the chosen Hayashi decomposition, and let l_k, r_k be the sector tensors of Definition 23.5.1.19. For each pair of sectors (k, h) , the neighboring operator contracts the right sector tensor of one site with the left sector tensor of the following site over the virtual bond they share:

$$\eta_{k,h} = \sum_\gamma (r_k)_\gamma \otimes (l_h)_\gamma.$$

It acts on the neighboring bond space $B_k^R \otimes B_h^L$. This is the operator displayed in [CPGSV16, Appendix C.2, lines 1441–1445].

Theorem 23.5.2.2 (Neighboring bond contraction). [Lean](#) [✓ Stmt](#) In the setting of Theorem 23.5.1.21, for all sectors k, h and virtual indices β_1, α_3 , contracting the shared virtual bond of two neighboring factorized slices leaves the neighboring operator between the outer sector tensors:

$$\begin{aligned} & \sum_\gamma [U_B \kappa_{\beta_1, \gamma} U_B^\dagger]_{(k, l_1, r_1), (k, l'_1, r'_1)} \\ & \quad \cdot [U_B \kappa_{\gamma, \alpha_3} U_B^\dagger]_{(h, l_2, r_2), (h, l'_2, r'_2)} \\ & = [(l_k)_{\beta_1}]_{l_1, l'_1} [\eta_{k,h}]_{(r_1, l_2), (r'_1, l'_2)} \\ & \quad \cdot [(r_h)_{\alpha_3}]_{r_2, r'_2}. \end{aligned}$$

This is the single-bond step of the identity assembling $\bigotimes_n \eta_{k_n, k_{n+1}}$ from the factorized chain in [CPGSV16, Appendix C.2, lines 1446–1449].

Proof. ✓ Proof Substituting the sector factorization (Theorem 23.5.1.21) into each slice, the outer sector tensors decouple from the bond sum, and the inner sum over the shared bond γ reads

$$\sum_{\gamma} [(r_k)_{\gamma}]_{r_1, r'_1} [(l_h)_{\gamma}]_{l_2, l'_2} = [\eta_{k, h}]_{(r_1, l_2), (r'_1, l'_2)}$$

by definition of the neighboring operator. □

Definition 23.5.2.3 (Source-minimal physical-sector factorization). Lean Lean Lean Lean Lean Lean ✓ Stmt A physical-sector factorization of an MPO tensor K consists of a finite decomposition

$$\mathbb{C}^d \simeq \bigoplus_k (B_k^L \otimes B_k^R),$$

whose left and right factors are nonzero, an isometry U on the physical space, and matrix families $(l_k)_{\beta}$ and $(r_k)_{\alpha}$ such that every physical slice satisfies

$$U \kappa_{\beta, \alpha} U^{\dagger} = \bigoplus_k (l_k)_{\beta} \otimes (r_k)_{\alpha}.$$

No positivity, trace factorization, or quantum-Markov decomposition is included in this datum. This is the factorization displayed in [CPGSV16, Appendix C.2, lines 1381–1388].

Definition 23.5.2.4 (Algebraic neighboring contraction). Lean Lean ✓ Stmt For sectors k, h , contract the common virtual index of the right and left sector tensors:

$$\eta_{k, h} = \sum_a (r_k)_a \otimes (l_h)_a.$$

This is an operator on $B_k^R \otimes B_h^L$. Positivity is not part of this definition; it is a separate hypothesis in Proposition C.7.

Definition 23.5.2.5 (One-site matrix spaces of sector sets). Lean Lean ✓ Stmt For a sector k and physical matrix indices x, y in its summand, let $A_{k; x, y} \in M_D(\mathbb{C})$ be the corresponding entry of the sector-coordinate tensor. For a set S of sectors, define

$$A_S = \text{span}\{A_{k; x, y} : k \in S\}.$$

These are the matrix spaces associated with the physical projections in [CPGSV16, Appendix C.2, lines 1463–1467].

Theorem 23.5.2.6 (Directed-cut annihilation of one-site matrix spaces). Lean Lean ✓ Stmt Let S and C be sets of sectors. If

$$\eta_{k, h} = 0 \quad (k \in S, h \in C),$$

then

$$XY = 0 \quad (X \in A_S, Y \in A_C).$$

No conclusion in the reverse product order is asserted. This is the directed local-orthogonality implication used in [CPGSV16, Appendix C.2, lines 1463–1470].

Proof. [✓ Proof](#) Expand two generating matrices in sector coordinates. Their product has entries

$$\sum_{\gamma} (l_k)_{\beta} (r_k)_{\gamma} (l_h)_{\gamma} (r_h)_{\alpha}.$$

The middle contraction is a matrix entry of $\eta_{k,h}$ and therefore vanishes. Bilinearity extends the conclusion from the generators to the two linear spans. $\square$

Definition 23.5.2.7 (Vertex rephasing of the sector tensors). [Lean](#) [✓ Stmt](#) Let $z_k \in \mathbb{C}$ satisfy $|z_k| = 1$ for every sector. Replacing

$$(l_k)_a \mapsto z_k (l_k)_a, \quad (r_k)_a \mapsto z_k^{-1} (r_k)_a$$

leaves every physical-slice factorization unchanged.

Theorem 23.5.2.8 (Rephasing identities for the sector data). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Vertex rephasing transforms the neighboring operators by

$$\eta_{k,h} \mapsto z_k^{-1} z_h \eta_{k,h}.$$

It leaves every sector virtual matrix unchanged, preserves the nonzero neighboring support, and leaves every complete cyclic product of neighboring operators unchanged. These are the rephasing identities used in [CPGSV16, Appendix C.2, lines 1434–1450].

Definition 23.5.2.9 (Vanishing right tensors on inactive sectors). [Lean](#) [Lean](#) [✓ Stmt](#) Let S be a set of active sectors in a physical-sector factorization, and suppose that $(l_k)_{\beta} = 0$ for every $k \notin S$ and every β . Retain every physical summand and define

$$\hat{l}_k = l_k, \quad \hat{r}_k = \begin{cases} r_k, & k \in S, \\ 0, & k \notin S. \end{cases}$$

Then $\hat{l}_k \otimes \hat{r}_k = l_k \otimes r_k$ for every k , so the physical-slice factorization is unchanged. The neighboring operators become

$$\hat{\eta}_{k,h} = \begin{cases} \eta_{k,h}, & k \in S, \\ 0, & k \notin S. \end{cases}$$

This is the reparameterization freedom in the factorization and neighboring contraction of [CPGSV16, Appendix C.2, lines 1434–1445].

Definition 23.5.2.10 (Inverse-map physical-sector factorization). [Lean](#) [✓ Stmt](#) Let K be injective, let ρ be its three-site closure against a virtual tail R , and choose a Hayashi decomposition of ρ . Fix outer indices such that $R_{\beta_3, \alpha_1} \neq 0$. The inverse-map tensors l_k, r_k , together with the Hayashi physical decomposition and unitary, determine a physical-sector factorization of K .

The left and right factor spaces are nonzero because the corresponding Hayashi density matrices have trace one; their nonzero dimensions are conclusions rather than hypotheses. No positivity of the neighboring operators is asserted. This gives the factorization in [CPGSV16, Appendix C.2, equations `AppUkU=r1` and `formK`, lines 1383–1387 and 1434–1439].

Theorem 23.5.2.11 (Identification of the neighboring operators). [Lean](#) [✓ Stmt](#) For all sectors k, h , the neighboring contraction of the resulting physical-sector factorization is the inverse-map operator

$$\eta_{k,h} = \sum_{\gamma} (r_k)_{\gamma} \otimes (l_h)_{\gamma}.$$

This is equation `etar1` in [CPGSV16, Appendix C.2, lines 1441–1445].

Definition 23.5.2.12 (Zero-weight inverse-map reparameterization). [Lean](#) [✓ Stmt](#) In the inverse-map factorization, retain every physical sector and its factor spaces, and set

$$\hat{l}_k = l_k, \quad \hat{r}_k = \begin{cases} r_k, & p_k \neq 0, \\ 0, & p_k = 0. \end{cases}$$

The physical-slice factorization is unchanged. This treats the zero-weight freedom left by the Hayashi decomposition used in [\[CPGSV16, Appendix C.2, lines 1415–1445\]](#).

Theorem 23.5.2.13 (Neighboring operators after zero-weight reparameterization). [Lean](#) [Lean](#) [✓ Stmt](#) *The neighboring operators of the reparameterized factorization are*

$$\hat{\eta}_{k,h} = \begin{cases} \eta_{k,h}, & p_k \neq 0, \\ 0, & p_k = 0. \end{cases}$$

Thus a zero-weight sector has no outgoing support edge. Since $p_h = 0$ also gives $l_h = 0$, it has no incoming support edge either.

Proof. [✓ Proof](#) The neighboring contraction uses r_k at its source and l_h at its target. The first assertion follows from the definition of $\hat{r}_k$, and the last assertion follows from Lemma [23.5.1.20](#). $\square$

Corollary 23.5.2.14 (SAL gives a raw physical-sector factorization). [Lean](#) [✓ Stmt](#) *Every injective MPO tensor satisfying SAL admits a physical-sector factorization. This is only the raw algebraic factorization from Lemma C.4 and equations `formK` and `etarl`; coherent positivity of the neighboring operators, recurrence, and primitivity are not part of this conclusion.*

Definition 23.5.2.15 (Finite-chain coordinates of a physical-sector factorization). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $N \geq 1$. For a cyclic sector configuration $k = (k_0, \dots, k_{N-1})$, write

$$I_k = \prod_{n=0}^{N-1} (B_{k_n}^L \times B_{k_n}^R).$$

Applying the one-site sector decomposition at every site gives a bijection

$$\Phi_N : \{0, \dots, d-1\}^N \longrightarrow \coprod_{k_0, \dots, k_{N-1}} I_k.$$

On I_k , define the cyclic neighboring product by

$$\Omega_k(x, y) = \prod_{n=0}^{N-1} \eta_{k_n, k_{n+1}}((x_n^R, x_{n+1}^L), (y_n^R, y_{n+1}^L)),$$

where site labels are read modulo N .

Theorem 23.5.2.16 (Finite-chain decomposition of a physical-sector factorization). [Lean](#) [Lean](#) [✓ Stmt](#) *Suppose that the physical-sector factorization is given. For every $N \geq 1$, the physically transformed MPO is block diagonal in the cyclic sector configurations, and*

$$\text{reindex}_{\Phi_N}(M_N(\tilde{K})) = \bigoplus_{k_0, \dots, k_{N-1}} \Omega_k.$$

This is the all-length algebraic decomposition in [\[CPGSV16, Appendix C.2, lines 1435–1450\]](#), conditional here on the displayed physical-sector factorization. It does not assert positivity of the individual neighboring operators.

Proof. ✓ Proof Fix a sector configuration k and entries $x, y \in I_k$. Expanding the closed horizontal contraction gives

$$\sum_g \prod_n (l_{k_n})_{g_n} (x_n^L, y_n^L) (r_{k_n})_{g_{n+1}} (x_n^R, y_n^R).$$

Cyclically shifting the summation variables and collecting the factors with the same horizontal index turns this expression into $\Omega_k(x, y)$. If the row and column sector configurations differ, one transformed physical slice lies between distinct direct summands and vanishes. Thus all off-diagonal sector blocks vanish. $\square$

Corollary 23.5.2.17 (SAL gives positive cyclic sector products). Lean ✓ Stmt *Every injective MPO tensor satisfying SAL admits a physical-sector factorization such that, for every $N \geq 1$ and every cyclic sector configuration $k = (k_0, \dots, k_{N-1})$,*

$$\bigotimes_{n=0}^{N-1} \eta_{k_n, k_{n+1}} \geq 0, \quad k_N = k_0.$$

This is the projected-chain inequality in [CPGSV16, Appendix C.2, Lemma C.4, lines 1446–1450]. It does not assert positivity of the individual neighboring operators.

Proof. ✓ Proof Choose the raw physical-sector factorization supplied by SAL. The physical basis change sends the positive N -site MPO to a positive congruence. Each cyclic neighboring product is a diagonal sector compression of this congruence and is therefore positive semidefinite. $\square$

Definition 23.5.2.18 (Boundary contraction against a virtual matrix). Lean Lean Lean Lean Lean ✓ Stmt For an arbitrary virtual matrix X , define the outer boundary operator on $B_k^L \otimes B_h^R$ by

$$B_{k,h}(X) = \sum_{a,b} X_{b,a} (l_k)_a \otimes (r_h)_b.$$

The map $X \mapsto B_{k,h}(X)$ is complex-linear. No positivity of X or $B_{k,h}(X)$ is assumed.

Definition 23.5.2.19 (Regrouped two-site sector closure). Lean Lean Lean Lean ✓ Stmt First pass to the physical basis selected by U , so that each slice is written in the sector coordinates of Definition 23.5.2.3. Form the two-site closure of this sector-coordinate tensor, restrict it to the physical sectors (k, h) , and reindex its physical factors by

$$((L_k, R_k), (L_h, R_h)) \mapsto ((L_k, R_h), (R_k, L_h)).$$

Denote the resulting matrix by $K_{2;k,h}(X)$.

Theorem 23.5.2.20 (Two-site closure factorization). Lean ✓ Stmt *For every virtual matrix X and every sector pair (k, h) ,*

$$K_{2;k,h}(X) = B_{k,h}(X) \otimes \eta_{k,h}.$$

Proof. ✓ Proof At the regrouped matrix entry indexed by $((\lambda_k, \rho_h), (\rho_k, \lambda_h))$ and $((\lambda'_k, \rho'_h), (\rho'_k, \lambda'_h))$, the virtual trace expands as the scalar

$$\sum_{a,b,c} X_{b,a} [(l_k)_a]_{\lambda_k, \lambda'_k} [(r_k)_c]_{\rho_k, \rho'_k} [(l_h)_c]_{\lambda_h, \lambda'_h} [(r_h)_b]_{\rho_h, \rho'_h}.$$

Commuting scalar factors and collecting the sums over (a, b) and c gives respectively $B_{k,h}(X)$ and $\eta_{k,h}$. $\square$

Definition 23.5.2.21 (Regrouped three-site sector closure). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) First pass to the physical basis selected by U and form the three-site closure of the resulting sector-coordinate tensor. Restrict this closure to the sectors (k, l, h) and reindex its physical factors by

$$((L_k, R_k), ((L_l, R_l), (L_h, R_h))) \mapsto ((L_k, R_h), ((R_k, L_l), (R_l, L_h))).$$

Denote the resulting matrix by $K_{3;k,l,h}(X)$.

Theorem 23.5.2.22 (Three-site closure factorization). [Lean](#) [✓ Stmt](#) For every virtual matrix X and fixed sectors (k, l, h) ,

$$K_{3;k,l,h}(X) = B_{k,h}(X) \otimes (\eta_{k,l} \otimes \eta_{l,h}).$$

Proof. [✓ Proof](#) At the regrouped matrix entry indexed by $((\lambda_k, \rho_h), ((\rho_k, \lambda_l), (\rho_l, \lambda_h)))$ and its primed counterpart, the left-hand side is the scalar

$$\sum_{a,b,c,e} X_{b,a} [(l_k)_a]_{\lambda_k, \lambda'_k} [(r_k)_c]_{\rho_k, \rho'_k} [(l_l)_c]_{\lambda_l, \lambda'_l} [(r_l)_e]_{\rho_l, \rho'_l} [(l_h)_e]_{\lambda_h, \lambda'_h} [(r_h)_b]_{\rho_h, \rho'_h}.$$

Reordering the four finite sums separates the (a, b) boundary contraction from the neighboring contractions indexed by c and e , which gives the three factors on the right-hand side. $\square$

Definition 23.5.2.23 (Three-site neighboring operator). [Lean](#) [✓ Stmt](#) For every outer-sector pair (k, h) , define

$$\Omega_{k,h} = \bigoplus_l (\eta_{k,l} \otimes \eta_{l,h}).$$

This is the direct sum of the two neighboring contractions appearing in the three-site closure factorization [[CPGSV16](#), Appendix C.2, lines 1510–1516].

$$\mathfrak{R}_2(K_2(X))_{k,h} = a_k b_h B_{k,h}(X) \qquad \mathfrak{R}_3(K_3(X))_{k,l,h} = B_{k,h}(X) \otimes \eta_{k,l} \otimes \eta_{l,h}$$

Figure 23.3: First each slice is conjugated by U and expressed in the corresponding sector coordinates. After restriction to fixed sectors and the displayed regroupings, the arbitrary- X two- and three-site closures separate into the boundary factor and respectively one or two neighboring contractions. The diagrams do not assert positivity, preparation, recovery, or the full conclusion of Proposition C.7. They follow the contractions in [[CPGSV16](#), lines 638–655 and Appendix C.2, lines 1435–1448].

Definition 23.5.2.24 (Neighboring-operator trace factorization). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that every neighboring operator is positive semidefinite and that there are real numbers a_k, b_h such that

$$\text{tr}(\eta_{k,h}) = a_k b_h, \qquad \sum_k a_k b_k = 1.$$

These are precisely the remaining hypotheses used in the construction of Proposition C.7 after the physical-sector factorization has been fixed [[CPGSV16](#), Appendix C.2, lines 1389–1403].

Theorem 23.5.2.25 (Trace of the three-site neighboring operator). [Lean](#) [Lean](#) [✓ Stmt](#) Assume the neighboring-operator trace factorization of Definition 23.5.2.24. Then, for every outer-sector pair (k, h) ,

$$\text{tr}(\Omega_{k,h}) = \sum_l \text{tr}(\eta_{k,l}) \text{tr}(\eta_{l,h}) = a_k b_h.$$

This is the normalization used in the coarse-graining channel of [[CPGSV16](#), Appendix C.2], lines 1547–1555.

Proof. ✓ Proof The trace of a direct sum is the sum of the traces of its blocks, and the trace of a tensor product is the product of the traces. Hence

$$\mathrm{tr}(\Omega_{k,h}) = \sum_l (a_k b_l)(a_l b_h) = a_k b_h \sum_l a_l b_l = a_k b_h.$$

□

Theorem 23.5.2.26 (Partial traces of the sector closures). Lean Lean Lean Lean ✓ Stmt For every virtual matrix X ,

$$\mathrm{tr}_{R_k \otimes L_h}(K_{2;k,h}(X)) = \mathrm{tr}(\eta_{k,h}) B_{k,h}(X) = a_k b_h B_{k,h}(X).$$

Similarly,

$$\mathrm{tr}_{(R_k \otimes L_l) \otimes (R_l \otimes L_h)}(K_{3;k,l,h}(X)) = \mathrm{tr}(\eta_{k,l}) \mathrm{tr}(\eta_{l,h}) B_{k,h}(X),$$

and summing the scalar coefficient over l gives $a_k b_h$.

Proof. ✓ Proof The partial trace of $A \otimes B$ is $\mathrm{tr}(B)A$. The two-site identity follows immediately. For three sites, the coefficient is

$$\sum_l (a_k b_l)(a_l b_h) = a_k \left(\sum_l a_l b_l \right) b_h = a_k b_h.$$

□

Definition 23.5.2.27 (Global two- and three-site sector coordinates). Lean Lean Lean Lean ✓ Stmt The two- and three-site physical spaces are reindexed by their canonical direct sums over sector pairs and triples, with each summand further regrouped into its boundary and neighboring factors. Write $\mathfrak{G}_2$ and $\mathfrak{G}_3$ for these reindexings of matrices.

Theorem 23.5.2.28 (Global direct-sum form of the sector closures). Lean Lean ✓ Stmt After the canonical regrouping of all physical-sector indices, the full closures are

$$\mathfrak{G}_2(K_2(X)) = \bigoplus_{k,h} B_{k,h}(X) \otimes \eta_{k,h}$$

and

$$\mathfrak{G}_3(K_3(X)) = \bigoplus_{k,l,h} B_{k,h}(X) \otimes (\eta_{k,l} \otimes \eta_{l,h}).$$

Every matrix entry between distinct sector tuples vanishes.

Proof. ✓ Proof On a diagonal sector tuple these are the fixed-sector factorizations. For distinct sector labels $k \neq p$, the block structure of the sector-coordinate tensor gives

$$[U \kappa_{\beta,\alpha} U^\dagger]_{k,p} = 0.$$

Consequently, if $(k, h) \neq (p, q)$ or $(k, l, h) \neq (p, m, q)$, respectively, then

$$[\mathfrak{G}_2(K_2(X))]_{(k,h),(p,q)} = 0, \quad [\mathfrak{G}_3(K_3(X))]_{(k,l,h),(p,m,q)} = 0.$$

□

Definition 23.5.2.29 (Two-site sector bond). [Lean](#) [✓ Stmt](#) In the two-site sector coordinates, define

$$B_{\text{sector}} = \bigoplus_{k,h} \left(I_{B_k^L \otimes B_h^R} \otimes \eta_{k,h} \right).$$

Thus the outer factors carry the identity, while the neighboring factors carry the operator $\eta_{k,h}$. This is the local bond in [\[CPGSV16, Appendix C.2, Proposition C.8, lines 1581–1593\]](#).

Theorem 23.5.2.30 (Positivity of the two-site sector bond). [Lean](#) [✓ Stmt](#) Suppose that $\eta_{k,h}$ is positive semidefinite for every pair of sectors. Then B_{sector} is positive semidefinite. The positivity premise is the conclusion at [\[CPGSV16, Appendix C.2, lines 1446–1450\]](#).

Proof. [✓ Proof](#) Each summand $I_{B_k^L \otimes B_h^R} \otimes \eta_{k,h}$ is positive semidefinite. Positivity is preserved by finite block diagonals and by the reindexing from the regrouped direct sum to the two-site sector coordinates. $\square$

Definition 23.5.2.31 (Two-site bond in physical coordinates). [Lean](#) [Lean](#) [✓ Stmt](#) Let V_2 be the two-site tensor product of the project's coordinate map, so that $V_2 X V_2^\dagger$ expresses a physical operator X in sector coordinates. Thus $V_2 = U_{\text{paper}}^\dagger$ in the convention of [\[CPGSV16, Appendix C.2, lines 1581–1583\]](#). Define

$$B_{\text{physical}} = V_2^\dagger B_{\text{sector}} V_2,$$

and identify pairs of physical indices with functions $\{0, 1\} \rightarrow \{0, \dots, d-1\}$. This is the physical two-site bond in [\[CPGSV16, Appendix C.2, Proposition C.8, lines 1581–1593\]](#).

Theorem 23.5.2.32 (Positivity of the physical two-site bond). [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that $\eta_{k,h}$ is positive semidefinite for every pair of sectors. Then B_{physical} is positive semidefinite, both with pair indices and with two-site configuration indices.

Proof. [✓ Proof](#) Apply positivity of B_{sector} to the congruence $V_2^\dagger B_{\text{sector}} V_2$. Reindexing the resulting matrix by two-site configurations preserves positivity. $\square$

Definition 23.5.2.33 (Three-site lifts of a two-site operator). [Lean](#) [Lean](#) [✓ Stmt](#) Let B be an operator on two copies of a finite-dimensional space H . On the right-associated space $H \otimes (H \otimes H)$, define

$$B_{01} = \alpha(B \otimes I_H) \alpha^{-1}, \quad B_{12} = I_H \otimes B,$$

where $\alpha : (H \otimes H) \otimes H \rightarrow H \otimes (H \otimes H)$ is the canonical associator.

Theorem 23.5.2.34 (Three-site translation identities). [Lean](#) [Lean](#) [✓ Stmt](#) Under the canonical identification of three-site configurations with $H \otimes (H \otimes H)$, the translates of the physical bond beginning at sites 0 and 1 are respectively

$$(B_{\text{physical}})_{01} = B_{01}, \quad (B_{\text{physical}})_{12} = B_{12}.$$

Proof. [✓ Proof](#) An entry of the first translate vanishes unless the third indices agree; its remaining entry is that of B_{physical} . Similarly, an entry of the second translate vanishes unless the first indices agree. These are precisely the matrix entries of B_{01} and B_{12} . $\square$

Theorem 23.5.2.35 (Commutativity on a fixed sector triple). [Lean](#) [✓ Stmt](#) Fix sectors (k, l, h) . On the regrouped sector space, set

$$\begin{aligned} C_{01}^{k,l,h} &= I_{B_k^L \otimes B_h^R} \otimes (\eta_{k,l} \otimes I_{B_l^R \otimes B_h^L}), \\ C_{12}^{k,l,h} &= I_{B_k^L \otimes B_h^R} \otimes (I_{B_k^R \otimes B_l^L} \otimes \eta_{l,h}). \end{aligned}$$

Then

$$C_{01}^{k,l,h} C_{12}^{k,l,h} = C_{12}^{k,l,h} C_{01}^{k,l,h}.$$

This is the fixed-sector calculation in [CPGSV16, Appendix C.2, Proposition C.8, lines 1589–1593].

On the sector triple (k, l, h) , the two operators act on distinct neighboring factors, while the outer factors B_k^L and B_h^R remain unchanged. Their commutativity is therefore the local identity

$$\eta_{k,l}^{(12)} \eta_{l,h}^{(23)} = \eta_{l,h}^{(23)} \eta_{k,l}^{(12)}$$

Proof. ✓ Proof The operators $\eta_{k,l}$ and $\eta_{l,h}$ act on the distinct factors $B_k^R \otimes B_l^L$ and $B_l^R \otimes B_h^L$, respectively. The claim follows from the mixed-product identity for tensor products. $\square$

Theorem 23.5.2.36 (Commutativity of adjacent physical bonds). Lean ✓ Stmt The two translates of the physical bond on three sites commute:

$$(B_{\text{physical}})_{01} (B_{\text{physical}})_{12} = (B_{\text{physical}})_{12} (B_{\text{physical}})_{01}.$$

This is the adjacent-bond calculation in [CPGSV16, Appendix C.2, Proposition C.8, lines 1589–1593].

Proof. ✓ Proof Regroup the three-site sector coordinates into the direct sum over triples (k, l, h) . The two bonds become block diagonal, with blocks $C_{01}^{k,l,h}$ and $C_{12}^{k,l,h}$, which commute by Theorem 23.5.2.35. Conjugating by the three-fold physical coordinate unitary preserves the equality. The three-site translation identities then give the displayed physical commutator. $\square$

Theorem 23.5.2.37 (Agreement outside a cyclic window). Lean ✓ Stmt Let $W_i^L = \{i, i+1, \dots, i+L-1\}$ be a cyclic window of length $L \leq N$. Two configurations agree outside W_i^L precisely when their values agree at every site not contained in W_i^L .

Proof. ✓ Proof Replacing the entries in W_i^L does not affect any complementary site. Conversely, if the two configurations agree on the complement, replacing the window of the first by the window of the second reconstructs the second configuration. $\square$

Theorem 23.5.2.38 (Multiplicativity of cyclic local embeddings). Lean ✓ Stmt Let E_i^L embed an operator on L consecutive sites into an N -site periodic chain at the cyclic window beginning at i . Then

$$E_i^L(BC) = E_i^L(B)E_i^L(C).$$

Proof. ✓ Proof Identify a periodic configuration with its restriction to W_i^L and its restriction to the cyclic complement. In these coordinates, $E_i^L(B) = B \otimes I$, so the assertion is the mixed-product identity. $\square$

Theorem 23.5.2.39 (Commutativity on disjoint cyclic windows). Lean ✓ Stmt Let $L \leq N$, and let W_i^L and W_j^L be disjoint cyclic windows in an N -site periodic chain. For arbitrary operators B and C on L sites,

$$E_i^L(B)E_j^L(C) = E_j^L(C)E_i^L(B).$$

Proof. ✓ Proof Apply both sides to a chain amplitude. The first order of application is a double sum over replacements on W_i^L and W_j^L . Since the windows are disjoint, each replacement leaves the other restricted configuration unchanged, and the two replacements commute. Interchanging the two finite sums gives the reverse order. $\square$

Theorem 23.5.2.40 (Adjacent physical bonds on periodic chains). [Lean](#) [✓ Stmt](#) Let $N \geq 3$. For every site $i \in \mathbb{Z}/N\mathbb{Z}$, including $i = N - 1$, the adjacent translates of the physical bond commute:

$$(B_{\text{physical}})_{i,i+1}(B_{\text{physical}})_{i+1,i+2} = (B_{\text{physical}})_{i+1,i+2}(B_{\text{physical}})_{i,i+1}.$$

This is the periodic-chain form of the adjacent-bond calculation in [CPGSV16, Appendix C.2, Proposition C.8, lines 1571–1593]. The case $N = 2$ is treated separately below, because its two translated bonds have the same support with opposite cyclic order.

Proof. [✓ Proof](#) Enclose the two bonds in the cyclic three-site window beginning at i . In these coordinates they are the bonds beginning at sites 0 and 1, respectively. Embedding an operator into a fixed cyclic window preserves products, so the assertion follows from Theorem 23.5.2.36. $\square$

Theorem 23.5.2.41 (Commutativity of the crossed two-site translates). [Lean](#) [✓ Stmt](#) On the periodic chain of length two, the translates beginning at sites zero and one read the physical sites in the opposite cyclic orders $(0, 1)$ and $(1, 0)$. Nevertheless they commute:

$$B_{01}B_{10} = B_{10}B_{01}.$$

This is the length-two instance of the translated-bond commutativity in [CPGSV16, Appendix C.2, Proposition C.8]. The source does not discuss the crossed finite-size ordering separately.

Proof. [✓ Proof](#) Fix sectors (k, h) and regroup the two physical sites as

$$(L_k \otimes R_h) \otimes (R_k \otimes L_h).$$

The translate in the order $(0, 1)$ is

$$I_{L_k \otimes R_h} \otimes \eta_{k,h},$$

while the translate in the order $(1, 0)$ is

$$\eta_{h,k}^{\text{op}} \otimes I_{R_k \otimes L_h},$$

where $\eta_{h,k}^{\text{op}}$ denotes the same matrix after exchanging the factors R_h and L_k . These operators act on complementary factors and commute. Taking the direct sum over (k, h) and conjugating by the two-fold physical coordinate unitary proves the claim. $\square$

Theorem 23.5.2.42 (Disjoint physical bonds on periodic chains). [Lean](#) [✓ Stmt](#) Let $N \geq 2$, and suppose that the cyclic bonds beginning at i and j have disjoint two-site supports. Then their physical bond operators commute:

$$(B_{\text{physical}})_{i,i+1}(B_{\text{physical}})_{j,j+1} = (B_{\text{physical}})_{j,j+1}(B_{\text{physical}})_{i,i+1}.$$

This is the locality part of the pairwise commutation assertion in [CPGSV16, Appendix C.2, Proposition C.8, lines 1571–1593].

Proof. [✓ Proof](#) This is Theorem 23.5.2.39 applied to the same physical bond operator in both windows. No positivity or additional property of the sector factorization is needed for this locality step. $\square$

Theorem 23.5.2.43 (Pairwise commutativity of the physical bonds). [Lean] [✓ Stmt] Suppose that the physical-sector factorization is given. For every periodic chain of length $N \geq 2$ and every pair of sites $i, j \in \mathbb{Z}/N\mathbb{Z}$, the corresponding translates of the physical bond commute:

$$(B_{\text{physical}})_{i,i+1}(B_{\text{physical}})_{j,j+1} = (B_{\text{physical}})_{j,j+1}(B_{\text{physical}})_{i,i+1}.$$

This is the pairwise commutation assertion in [CPGSV16, Appendix C.2, Proposition C.8, lines 1571–1593], conditional here on the physical-sector factorization constructed in the preceding results.

Proof. [✓ Proof] For $N = 2$, equal translates commute trivially, while the two distinct translates commute by Theorem 23.5.2.41. Let $N \geq 3$. If the two cyclic supports are disjoint, apply Theorem 23.5.2.42. Otherwise, Lemma 13.9.11 shows that the two starting sites either coincide or are cyclic neighbors. The coincident case is immediate, and the two neighboring orientations follow from Theorem 23.5.2.40 and its symmetric equality. $\square$

Theorem 23.5.2.44 (Exact product of physical-sector bonds). [Lean] [✓ Stmt] Suppose that K is equipped with a physical-sector factorization F . Let B_i denote the cyclic translate beginning at site i of the physical two-site bond determined by F . For every $N \geq 2$, the N -site periodic operator is

$$\rho_N = B_0 B_1 \cdots B_{N-1}.$$

No positivity, zero-correlation-length, injectivity, or normalization hypothesis is required. This is the product identity in [CPGSV16, Appendix C.2, Proposition C.8, lines 1581–1593], conditional here on the given physical-sector factorization.

Proof. [✓ Proof] For a fixed cyclic sector sequence, contraction of the virtual index between the right tensor at site i and the left tensor at site $i + 1$ produces $\eta_{k_i, k_{i+1}}$. Taking the virtual trace therefore gives the cyclic product of these neighboring operators. The translated sector bonds give the same ordered product. Conjugating by the tensor power of the one-site coordinate unitary gives the stated physical identity. For $N = 2$, the second translate has the opposite cyclic order; the two factors are the ordinary and crossed bonds. $\square$

Theorem 23.5.2.45 (Unit scalar in the physical-sector product). [Lean] [✓ Stmt] Under the physical-sector factorization F of the preceding theorem, for every $N \geq 2$ there is a real number $c > 0$ such that

$$\rho_N = c B_0 B_1 \cdots B_{N-1}.$$

Here the bonds are the canonical bonds determined by F , and one may take $c = 1$. No positivity hypothesis on the neighboring operators $\eta_{k,h}$ is required.

Proof. [✓ Proof] Take $c = 1$ in Theorem 23.5.2.44. $\square$

Theorem 23.5.2.46 (Positive translation-invariant bond datum). [Lean] [✓ Stmt] Suppose that every neighboring operator $\eta_{k,h}$ is positive semidefinite. Then the physical bond is positive semidefinite and all of its cyclic translates commute on every periodic chain of length at least two. Thus it determines a single translation-invariant positive bond datum. This assertion does not include the product formula for the finite-chain density operator.

Proof. [✓ Proof] Positivity is Theorem 23.5.2.32; pairwise commutativity is Theorem 23.5.2.43. $\square$

Theorem 23.5.2.47 (η -local structure from a positive physical-sector factorization). [Lean](#) [✓ Stmt](#) Suppose that a physical-sector factorization of K is given and that every neighboring operator $\eta_{k,h}$ is positive semidefinite. Then the physical two-site bond determines an η -local structure for K . Its translated copies commute pairwise, and for every $N \geq 2$,

$$\rho_N = B_0 B_1 \cdots B_{N-1}.$$

Thus the positive normalization scalar may be chosen to be 1. This is Proposition C.8 conditional on the coherently positive physical-sector factorization that the source derives from SAL.

Proof. [✓ Proof](#) Theorem 23.5.2.46 gives the positive translation-invariant bond and its pairwise commuting translates. Theorem 23.5.2.44 realizes every finite-chain operator with normalization scalar 1. $\square$

Corollary 23.5.2.48 (η -local structure from SAL). [Lean](#) [✓ Stmt](#) Every injective MPO tensor satisfying SAL admits an η -local structure. Thus there is a single positive two-site bond whose cyclic translates commute pairwise and whose ordered product is the finite-chain density operator at every length $N \geq 2$, with normalization scalar one. This is the conclusion of Proposition C.8.

Proof. [✓ Proof](#) Theorem 23.5.3.41 supplies the positive physical-sector factorization. Apply Theorem 23.5.2.47. $\square$

23.5.3 Closed-sector contractions and coherent rephasing

Definition 23.5.3.1 (Closed sector tensors). [Lean](#) [Lean](#) [✓ Stmt](#) Tracing the two physical legs of a sector tensor leaves a vector over the virtual bond:

$$|l_k\rangle_\gamma := \text{tr}[(l_k)_\gamma], \quad (r_k|_\gamma := \text{tr}[(r_k)_\gamma].$$

These are the closed tensors of [CPGSV16, Appendix C.2, lines 1473–1477].

Definition 23.5.3.2 (Closed-sector pairing operator). [Lean](#) [✓ Stmt](#) For sector tensors $(l_k)_\beta$ and $(r_k)_\alpha$, close their physical legs and set

$$S := \sum_k |l_k\rangle (r_k|.$$

This is the pairing operator displayed in [CPGSV16, Appendix C.2, lines 1473–1493].

Theorem 23.5.3.3 (Sector trace pairing). [Lean](#) [✓ Stmt](#) For all sectors k, h , the trace of the neighboring operator is the pairing of the closed sector tensors:

$$T_{k,h} = \text{tr}(\eta_{k,h}) = (r_k|l_h).$$

This is [CPGSV16, Appendix C.2, lines 1452–1455 and 1478–1481].

Proof. [✓ Proof](#) Expanding the definition of the neighboring operator and applying $\text{tr}(A \otimes B) = \text{tr}(A) \text{tr}(B)$,

$$\text{tr}(\eta_{k,h}) = \sum_\gamma \text{tr}((r_k)_\gamma \otimes (l_h)_\gamma) = \sum_\gamma \text{tr}[(r_k)_\gamma] \text{tr}[(l_h)_\gamma] = (r_k|l_h).$$

$\square$

Definition 23.5.3.4 (Concrete closed-sector trace matrix). [Lean](#) [Lean](#) [✓ Stmt](#) For the inverse-map sector tensors, define the complex matrix

$$T_{k,h} := (r_k | l_h).$$

Equivalently,

$$T_{k,h} = \text{tr}(\eta_{k,h}).$$

This is the trace matrix in [\[CPGSV16, Appendix C.2, lines 1478–1481\]](#).

Theorem 23.5.3.5 (Closed-sector form of the physical-trace transfer). [Lean](#) [✓ Stmt](#) More generally, any sector factorization $U_B \kappa_{\beta,\alpha} U_B^\dagger = \bigoplus_k (l_k)_\beta \otimes (r_k)_\alpha$ gives the closed-sector pairing operator

$$\mathcal{T}_K = \sum_k |l_k)(r_k|.$$

This identifies the transfer matrix of [Definition 23.2.1](#) with the operator on the left-hand side of the zero-correlation-length identity in [\[CPGSV16, Appendix C.2, lines 1489–1493\]](#).

Proof. [✓ Proof](#) Trace the sector factorization over the physical index. Unitary invariance gives

$$\text{tr}(U_B \kappa_{\beta,\alpha} U_B^\dagger) = \text{tr}(\kappa_{\beta,\alpha}).$$

The trace of the block diagonal is the sum of the block traces. In each sector,

$$\text{tr}((l_k)_\beta \otimes (r_k)_\alpha) = \text{tr}[(l_k)_\beta] \text{tr}[(r_k)_\alpha] = [|l_k)(r_k|]_{\beta,\alpha}.$$

Summing over k gives the asserted operator identity. □

Corollary 23.5.3.6 (Physical-trace transfer from the inverse-map sectors). [Lean](#) [✓ Stmt](#) Let K be injective. Let R and ρ give a three-site closure of K , choose a Hayashi decomposition of ρ , and fix virtual indices α_1, β_3 such that $R_{\beta_3, \alpha_1} \neq 0$. For the corresponding closed sector tensors,

$$\mathcal{T}_K = \sum_k |l_k)(r_k|.$$

Proof. [✓ Proof](#) The inverse-map sector tensors satisfy the sector factorization of [Theorem 23.5.1.21](#). Apply [Theorem 23.5.3.5](#). □

Theorem 23.5.3.7 (Normalized closed-sector pairing operator). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let K be injective. Let R and ρ give a three-site closure of K , choose a Hayashi decomposition of ρ , and fix virtual indices α_1, β_3 such that $R_{\beta_3, \alpha_1} \neq 0$. Form the corresponding closed sector tensors $|l_k)$ and $(r_k|$. If K has source zero correlation length, set

$$S := \sum_k |l_k)(r_k|.$$

Then there is a real number $\lambda > 0$ such that

$$S^2 = \lambda S, \quad (\lambda^{-1} S)^2 = \lambda^{-1} S.$$

If K is the canonically normalized representative satisfying $\mathcal{T}_K^2 = \mathcal{T}_K$, then the raw pairing operator itself satisfies $S^2 = S$. This is the display preceding [\[CPGSV16, Appendix C.2, Lemma C.5\]](#); the first identity records the rescaling-invariant form of the same zero-correlation-length condition.

Proof. ✓ Proof Substitute $S = \mathcal{T}_K$ from Corollary 23.5.3.6 into the defining relation $\mathcal{T}_K^2 = \lambda \mathcal{T}_K$, and then apply Lemma 23.2.4. Under literal idempotence, the same substitution gives $S^2 = S$ directly. $\square$

Theorem 23.5.3.8 (Normalized closed-sector trace relations). Lean ✓ Stmt *Under the hypotheses of Theorem 23.5.3.7, suppose that K has source zero correlation length. There is a real number $\lambda > 0$ such that, for*

$$\widehat{T} := \lambda^{-1}T,$$

one has

$$\widehat{T}^2 = \widehat{T}^3.$$

For every positive integer N , one also has

$$\text{tr}(\widehat{T}^N) = \text{tr}(\widehat{T}).$$

This is the valid trace-power display in [CPGSV16, Appendix C.2, lines 1490–1497]. No idempotence or rank-one conclusion for $\widehat{T}$ is asserted.

Proof. ✓ Proof Let L be the matrix whose h th column is $|l_h\rangle$, and let Q be the matrix whose k th row is $\langle r_k|$. Then $S = LQ$ and $T = QL$. Place the normalization scalar on L . Since $\lambda^{-1}LQ$ is idempotent, associativity gives

$$(Q\lambda^{-1}L)^2 = Q(\lambda^{-1}LQ)\lambda^{-1}L = Q(\lambda^{-1}LQ)^2\lambda^{-1}L = (Q\lambda^{-1}L)^3.$$

Cyclicity of trace and the idempotence of $\lambda^{-1}LQ$ give

$$\text{tr}(Q\lambda^{-1}L) = \text{tr}(\lambda^{-1}LQ) = \text{tr}((\lambda^{-1}LQ)^2) = \text{tr}((Q\lambda^{-1}L)^2).$$

The preceding square-cube identity then gives $\widehat{T}^N = \widehat{T}^2$ for every $N \geq 2$, proving the stated trace identity. $\square$

Definition 23.5.3.9 (Explicit neighboring η -operator data). Lean Lean ✓ Stmt Fix a Hayashi decomposition witness h_η . An explicit neighboring η -family is a dependent family $(\eta_{k,h})$ indexed by sector pairs, where $\eta_{k,h}$ acts on the neighboring bond space $B_k^R \otimes B_h^L$. Requiring positivity for each pair gives the corresponding positive η -data.

Definition 23.5.3.10 (Neighboring operators from sector tensors). Lean ✓ Stmt Let $(l_k)_a$ and $(r_k)_a$ be sector tensors indexed by the horizontal virtual index a . For neighboring sectors k, h , define an operator on $B_k^R \otimes B_h^L$ by

$$\eta_{k,h}((x_R, x_L), (y_R, y_L)) = \sum_a (r_k)_a(x_R, y_R) (l_h)_a(x_L, y_L).$$

Thus $\eta_{k,h} = r_k l_h$ is the contraction over the shared horizontal index in [CPGSV16, Appendix C.2, equation (etarl)]. Positivity is not part of this definition.

Lemma 23.5.3.11 (Entries of the neighboring operators). Lean ✓ Stmt *For every pair of sectors and every pair of matrix indices, the entry of $\eta_{k,h}$ is the contraction displayed in Definition 23.5.3.10.*

Proof. ✓ Proof This is immediate from the definition of $\eta_{k,h}$. $\square$

Neighboring right and left sector tensors meet in the bond operator $\eta_{k,h} = r_k l_h$:

$$\eta_{k,h} = r_k l_h$$

This is the contraction in [CPGSV16, Appendix C.2, lines 1441–1445].

Theorem 23.5.3.12 (Cyclic contraction of the sector tensors). Lean ✓ Stmt *Let $N \geq 1$ and let $k_0, \dots, k_{N-1}$ be a cyclic sequence of sectors. Fix matrix indices in the left and right factor of every sector, and abbreviate*

$$L_n(a) = (l_{k_n})_a(x_n^L, y_n^L), \quad R_n(a) = (r_{k_n})_a(x_n^R, y_n^R).$$

With all site labels read modulo N ,

$$\sum_{g_0, \dots, g_{N-1}} \prod_{n=0}^{N-1} L_n(g_n) R_n(g_{n+1}) = \prod_{n=0}^{N-1} \eta_{k_n, k_{n+1}}((x_n^R, x_{n+1}^L), (y_n^R, y_{n+1}^L)).$$

This is the entrywise cyclic contraction at [CPGSV16, Appendix C.2, lines 1446–1450].

Proof. ✓ Proof Expand the product of the sums defining the neighboring operators and set $g'_n = g_{n+1}$. Cyclic invariance of the finite product and commutativity of scalar multiplication give

$$\begin{aligned} \sum_{g_0, \dots, g_{N-1}} \prod_{n=0}^{N-1} L_n(g_n) R_n(g_{n+1}) &= \sum_{g'_0, \dots, g'_{N-1}} \prod_{n=0}^{N-1} R_n(g'_n) L_{n+1}(g'_n) \\ &= \prod_{n=0}^{N-1} \sum_a R_n(a) L_{n+1}(a). \end{aligned}$$

The last expression is the product of the corresponding entries of $\eta_{k_n, k_{n+1}}$. □

Theorem 23.5.3.13 (Closed sector word as neighboring contractions). Lean ✓ Stmt *Under the hypotheses of the preceding theorem, define $M_n(a, b) = L_n(a) R_n(b)$. Then*

$$\text{tr}(M_0 M_1 \cdots M_{N-1}) = \prod_{n=0}^{N-1} \eta_{k_n, k_{n+1}}((x_n^R, x_{n+1}^L), (y_n^R, y_{n+1}^L)).$$

$$\text{tr}(M_{k_0} \cdots M_{k_{N-1}}) = \text{tr}(\eta_{k_0, k_1} \otimes \cdots \otimes \eta_{k_{N-1}, k_0})$$

Figure 23.4: Closing the horizontal indices of the factorized sector tensors pairs each right factor with the left factor at the next site. The cycle therefore becomes the product of the neighboring operators $\eta_{k_n, k_{n+1}}$; compare [CPGSV16, Appendix C.2, lines 1446–1450].

Proof. ✓ Proof Expanding the trace of the matrix word gives

$$\text{tr}(M_0 M_1 \cdots M_{N-1}) = \sum_{g_0, \dots, g_{N-1}} \prod_{n=0}^{N-1} M_n(g_n, g_{n+1}).$$

Substituting $M_n(a, b) = L_n(a) R_n(b)$ gives the cyclic sum in Theorem 23.5.3.12; apply that theorem. □

Definition 23.5.3.14 (Sector-adapted chain coordinates). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For a sector configuration $k = (k_0, \dots, k_{N-1})$, let

$$I_k = \prod_{n=0}^{N-1} (\{0, \dots, d_L(k_n) - 1\} \times \{0, \dots, d_R(k_n) - 1\}).$$

Applying the one-site Hayashi decomposition at every site gives a bijection

$$\Phi_N : \{0, \dots, d-1\}^N \longrightarrow \prod_{k_0, \dots, k_{N-1}} I_k.$$

We write $\text{reindex}_{\Phi_N}(M)$ for the matrix obtained by transporting both indices of M along this bijection.

Definition 23.5.3.15 (Physical conjugation of an MPO tensor). [Lean](#) [✓ Stmt](#) Let U be a matrix on the physical space. The physical conjugation of K by U is the tensor $\tilde{K}$ defined by

$$\tilde{\kappa}_{\beta, \alpha}^{ij} = (U \kappa_{\beta, \alpha} U^\dagger)_{ij}.$$

Its horizontal bond indices are unchanged. When U is unitary, this is the corresponding physical basis change.

Definition 23.5.3.16 (Cyclic tensor product of neighboring operators). [Lean](#) [✓ Stmt](#) Let $N \geq 1$ and let k be a sector configuration. Define the matrix Ω_k on I_k by

$$\Omega_k(x, y) = \prod_{n=0}^{N-1} \eta_{k_n, k_{n+1}}((x_n^R, x_{n+1}^L), (y_n^R, y_{n+1}^L)),$$

where the site labels are read modulo N . Thus $\Omega_k = \eta_{k_0, k_1} \otimes \dots \otimes \eta_{k_{N-1}, k_0}$ after the neighboring tensor factors are ordered cyclically.

Theorem 23.5.3.17 (A fixed sector block is the cyclic neighboring product). [Lean](#) [✓ Stmt](#) Suppose that the physical slices in the Hayashi basis satisfy

$$U_B \kappa_{\beta, \alpha} U_B^\dagger = \bigoplus_q (l_q)_\beta \otimes (r_q)_\alpha,$$

and let $\eta_{q, h} = r_q l_h$. For every $N \geq 1$ and every sector configuration k , the corresponding diagonal block of $\text{reindex}_{\Phi_N}(M_N(\tilde{K}))$ is Ω_k .

Proof. [✓ Proof](#) Fix $x, y \in I_k$ and expand the closed horizontal trace of the MPO word. The sector factorization writes each local entry as a product of entries of l_{k_n} and r_{k_n} . Set

$$M_n(\beta, \alpha) = (l_{k_n})_\beta(x_n^L, y_n^L)(r_{k_n})_\alpha(x_n^R, y_n^R).$$

Then

$$\begin{aligned} [\text{reindex}_{\Phi_N}(M_N(\tilde{K}))]_{(k, x), (k, y)} &= \text{tr}(M_0 M_1 \dots M_{N-1}) \\ &= \prod_{n=0}^{N-1} \eta_{k_n, k_{n+1}}((x_n^R, x_{n+1}^L), (y_n^R, y_{n+1}^L)) \\ &= \Omega_k(x, y). \end{aligned}$$

The middle equality is Theorem [23.5.3.13](#). □

Theorem 23.5.3.18 (Sector-adapted MPO decomposition). [Lean](#) [✓ Stmt](#) Under the same sector factorization hypothesis, for every $N \geq 1$,

$$\text{reindex}_{\Phi_N}(M_N(\tilde{K})) = \bigoplus_{k_0, \dots, k_{N-1}} \eta_{k_0, k_1} \otimes \eta_{k_1, k_2} \otimes \cdots \otimes \eta_{k_{N-1}, k_0}.$$

This is the algebraic sector decomposition in [CPGSV16, Appendix C.2, lines 1446–1450]. It does not assert that the individual neighboring operators are positive.

Proof. [✓ Proof](#) If the row and column sector configurations coincide, the corresponding block is Theorem 23.5.3.17. If they differ, choose a site n at which their labels k_n and h_n are unequal. Block diagonality gives, for every pair of horizontal indices,

$$[U_{B\kappa_{\beta, \alpha}} U_B^\dagger]_{(k_n, \cdot), (h_n, \cdot)} = 0.$$

This local factor occurs in every cyclic horizontal-index product, so every summand in the closed trace is zero. $\square$

Theorem 23.5.3.19 (Positive rescaling of Kronecker factors). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let A and B be nonzero complex square matrices whose Kronecker product $A \otimes B$ is positive semidefinite. Then there is a scalar $c \neq 0$ with $cA \geq 0$ and $c^{-1}B \geq 0$. Since $(cA) \otimes (c^{-1}B) = A \otimes B$, the rescaled factors represent the same product.

Proof. [✓ Proof](#) On a product vector $x \otimes y$ the quadratic form of $A \otimes B$ factors, so for all vectors x and y ,

$$(x^\dagger A x)(y^\dagger B y) \geq 0.$$

By polarization a complex matrix is determined by its quadratic form; since A and B are nonzero there are vectors with $a = v_0^\dagger A v_0 \neq 0$ and $c = w_0^\dagger B w_0 \neq 0$. Every value $x^\dagger (cA) x = (x^\dagger A x) c$ is nonnegative, and over the complex numbers a nonnegative quadratic form already forces the matrix to be Hermitian, hence positive semidefinite. For the second factor, $ac > 0$ and $a(y^\dagger B y) \geq 0$ give $c^{-1}(y^\dagger B y) = (a(y^\dagger B y))(ac)^{-1} \geq 0$. $\square$

Lemma 23.5.3.20 (A matrix and its negative cannot both be positive). [Lean](#) [✓ Stmt](#) If X and $-X$ are positive semidefinite complex matrices, then $X = 0$.

Proof. [✓ Proof](#) The quadratic form of X is both nonnegative and nonpositive, hence vanishes identically. Polarization gives $X = 0$. $\square$

Definition 23.5.3.21 (Finite Kronecker product). [Lean](#) [✓ Stmt](#) For square matrices A_n indexed by $n = 0, \dots, N-1$, possibly with different index types, their finite Kronecker product is the matrix with entries

$$\left(\bigotimes_{n=0}^{N-1} A_n \right)_{x,y} = \prod_{n=0}^{N-1} (A_n)_{x_n, y_n}.$$

Lemma 23.5.3.22 (Elementary finite Kronecker identities). [Lean](#) [Lean](#) [✓ Stmt](#) A finite Kronecker product of nonzero matrices is nonzero. Replacing one factor A_i by cA_i multiplies the whole finite product by c .

Proof. [✓ Proof](#) Choose a nonzero entry in each factor and evaluate the product at the resulting row and column. The rescaling identity follows immediately from the entry formula in Definition 23.5.3.21. $\square$

Theorem 23.5.3.23 (Uniqueness of the positive rescaling phase). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Stmnt](#) Let $M \neq 0$. If $c_1, c_2 \neq 0$ and both $c_1 M$ and $c_2 M$ are positive semidefinite, then $c_1 = t c_2$ for some positive real number t . In particular, every nonzero scalar c has a unit-modulus part $u = c/|c|$, positivity of cM implies positivity of uM , and there is at most one such unit scalar when $M \neq 0$.

Proof. [Proof](#) Hermiticity of $c_1 M$ and $c_2 M$ gives $c_2 \overline{c_1} = c_1 \overline{c_2}$, so c_1/c_2 is real. It cannot be negative: otherwise $c_1 M$ and its negative would both be positive semidefinite, forcing $c_1 M = 0$. Thus the quotient is positive. Multiplication by $|c|^{-1} > 0$ replaces c by its unit-modulus part without changing positivity. Two admissible unit scalars differ by a positive real number of modulus one, and are therefore equal. $\square$

Theorem 23.5.3.24 (Positive rescaling of a finite Kronecker product). [Lean](#) [Stmnt](#) Let $N \geq 1$, and let $A_0, \dots, A_{N-1}$ be nonzero complex square matrices, with possibly different dimensions. If their finite Kronecker product is positive semidefinite, then there are nonzero scalars $c_0, \dots, c_{N-1}$ such that

$$\prod_{n=0}^{N-1} c_n = 1, \quad c_n A_n \geq 0 \quad (0 \leq n < N).$$

Proof. [Proof](#) The assertion is proved by induction on N . Separate the first factor from the remaining Kronecker product and apply Theorem 23.5.3.19. Absorb the inverse of the first rescaling coefficient into one factor of the tail, then apply the induction hypothesis. The resulting coefficients are nonzero, their product is one, and every rescaled factor is positive semidefinite. $\square$

Lemma 23.5.3.25 (One-sided product obstruction in a full matrix algebra). [Lean](#) [Stmnt](#) Let $\mathcal{A}_1, \mathcal{A}_2 \subseteq M_D(\mathbb{C})$ be nonzero linear subspaces. If $XY = 0$ for every $X \in \mathcal{A}_1$ and $Y \in \mathcal{A}_2$, then

$$\mathcal{A}_1 + \mathcal{A}_2 \neq M_D(\mathbb{C}).$$

No vanishing condition on products in the reverse order is required. This is the matrix-algebra obstruction used in [CPGSV16, Appendix C.2, lines 1465–1470].

Proof. [Proof](#) Choose nonzero matrices $X \in \mathcal{A}_1$ and $Y \in \mathcal{A}_2$, together with nonzero entries X_{ip} and Y_{qj} . If the two subspaces spanned the full matrix algebra, write the matrix unit $E_{pq} = E_1 + E_2$ with $E_1 \in \mathcal{A}_1$ and $E_2 \in \mathcal{A}_2$. The one-sided product hypothesis gives

$$X E_{pq} Y = X(E_1 Y) + (X E_2) Y = 0.$$

Its (i, j) entry is $X_{ip} Y_{qj} \neq 0$, a contradiction. $\square$

Definition 23.5.3.26 (Nonzero sector support and recurrence). [Lean](#) [Lean](#) [Lean](#) [Stmnt](#) The nonzero support graph of a neighboring-operator family has a directed edge $k \rightarrow h$ precisely when $\eta_{k,h} \neq 0$. Its support is recurrent when every such edge belongs to a directed cycle, equivalently when h can reach k whenever $k \rightarrow h$ is an edge.

Definition 23.5.3.27 (Sector-coordinate virtual matrix families). [Lean](#) [Lean](#) [Lean](#) [Stmnt](#) In sector coordinates, let $A_{k;x,y}$ be the virtual matrix obtained from the (x, y) entry within sector k . The sector-coordinate family contains all transformed physical matrices, while the virtual-matrix family consists of all matrices $A_{k;x,y}$.

Theorem 23.5.3.28 (Sector spanning from injectivity). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Stmnt](#) Let K be injective and let

$$U \kappa_{\beta,\alpha} U^\dagger = \bigoplus_k (l_k)_\beta \otimes (r_k)_\alpha$$

be a physical-sector factorization. Then the virtual matrices $A_{k;x,y}$ obtained from all within-sector physical entries span the full virtual matrix algebra $M_D(\mathbb{C})$.

Proof. [✓ Proof](#) Since U is unitary, every original physical matrix of K is a linear combination of the matrices in the transformed physical coordinates. The off-diagonal sector entries in those coordinates vanish, while each diagonal sector entry is one of the matrices $A_{k;x,y}$. Injectivity says that the original physical matrices span $M_D(\mathbb{C})$, and the conclusion follows. $\square$

Theorem 23.5.3.29 (Triangle closure from sector spanning). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For a physical-sector factorization, let $A_{k;x,y} \in M_D(\mathbb{C})$ be the virtual matrix obtained from the physical matrix entry (x, y) within sector k . Suppose that all such matrices span $M_D(\mathbb{C})$ and that both endpoint sectors of every nonzero neighboring operator contain a nonzero matrix of this form. Then every nonzero $\eta_{k,h}$ admits a two-edge return: there is a sector j for which

$$\eta_{h,j} \neq 0, \quad \eta_{j,k} \neq 0.$$

In particular, the nonzero neighboring support is recurrent.

Proof. [✓ Proof](#) A nonzero entry of $\eta_{k,h}$, together with nonzero sector matrices at k and h , gives matrices A from sector k and B from sector h with $AB \neq 0$. By nondegeneracy of the trace pairing, choose Y with $\text{tr}(ABY) \neq 0$. Expanding Y in the spanning family, some sector matrix C from a sector j satisfies $\text{tr}(ABC) \neq 0$. Cyclicity of the trace gives $BC \neq 0$ and $CA \neq 0$. The sector multiplication identity then forces $\eta_{h,j} \neq 0$ and $\eta_{j,k} \neq 0$. $\square$

Lemma 23.5.3.30 (Equivalent forms of sector reachability). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A vertex h is reachable from k in the nonzero sector support if and only if there is a finite directed walk from k to h . Consequently, the support is recurrent if and only if every directed edge admits a directed return walk.

Proof. [✓ Proof](#) A walk gives an element of the reflexive transitive closure by induction on its edges. Conversely, append the final edge at each induction step in the reflexive transitive closure. $\square$

Lemma 23.5.3.31 (Finite cyclic form of a closed directed walk). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Every nonempty closed directed walk of length N determines a cyclic family of vertices $v_0, \dots, v_{N-1}$. Each $v_n \rightarrow v_{n+1}$ is an edge, where $v_N = v_0$, and for any edge weights in a commutative monoid,

$$\text{wt}(w) = \prod_{n=0}^{N-1} \kappa_{v_n, v_{n+1}}.$$

Proof. [✓ Proof](#) Enumerate the sources of the successive edges. The last edge ends at the initial vertex because the walk is closed. The product identity follows by induction on the number of edges. $\square$

Theorem 23.5.3.32 (Closed-walk criterion for vertex phases). [Lean](#) [✓ Stmt](#) Let E be a directed relation on a set V , and suppose that every edge $a \rightarrow b$ admits a directed return walk from b to a . Let G be a group and assign a weight $\kappa_{a,b} \in G$ to each ordered pair of vertices. If the product of the edge weights along every closed directed walk is one, then there are vertex weights $z_a \in G$ such that

$$\kappa_{a,b} = z_a^{-1} z_b$$

whenever $a \rightarrow b$ is an edge. This is the graph-theoretic coboundary step needed to make the phase choices in [CPGVS16, Appendix C.2, lines 1446–1450] coherent across a recurrent support component.

Proof. ✓ Proof Reachability is an equivalence relation because every edge has a return walk. Choose one vertex in each reachability class and, for every vertex a , choose a walk from the class representative to a . Define z_a as the product of the edge weights along this walk. If two such walks have the same endpoints, append a common return walk; the two resulting closed walks have weight one, so the original walk weights agree. Comparing the chosen walk to b with the chosen walk to a followed by the edge $a \rightarrow b$ gives $z_b = z_a \kappa_{a,b}$. $\square$

Definition 23.5.3.33 (Horizontal bond fiber of a sector cycle). Lean Lean ✓ Stmt For a cyclic sector assignment $(k_n)_{n \in \mathbb{Z}/N\mathbb{Z}}$, the horizontal bond fiber at n is

$$H_n = B_{k_n}^R \times B_{k_{n+1}}^L.$$

Shifting the left component by one site identifies $\prod_n H_n$ with the vertex-indexed sector fiber used in the cyclic neighboring-operator product.

Lemma 23.5.3.34 (A cyclic block is a finite Kronecker product). Lean ✓ Stmt *Reindexing a cyclic neighboring-operator block along its horizontal bond fiber gives the finite Kronecker product*

$$\bigotimes_{n=0}^{N-1} \eta_{k_n, k_{n+1}}.$$

Proof. ✓ Proof Expand both sides entrywise. The shifted left component of the bond fiber at site $n+1$ is precisely the left index paired with the right component at site n . $\square$

Theorem 23.5.3.35 (Positive rescaling around one sector cycle). Lean ✓ Stmt *Let $N \geq 1$, and fix a cyclic sector assignment $k_0, \dots, k_{N-1}$ such that every operator $\eta_{k_n, k_{n+1}}$ is nonzero. If the corresponding cyclic tensor product is positive semidefinite, then there are nonzero scalars c_n satisfying*

$$\prod_{n=0}^{N-1} c_n = 1, \quad c_n \eta_{k_n, k_{n+1}} \geq 0 \quad (0 \leq n < N).$$

This is a choice on one fixed cycle. It does not assert that choices made on two different cycles agree on a common edge.

Proof. ✓ Proof Reindex the cyclic sector block by the shared horizontal-bond indices. Its matrix entries then become the finite Kronecker product of the consecutive neighboring operators. Apply Theorem 23.5.3.24. $\square$

Corollary 23.5.3.36 (Positive rescaling of the concrete sector operators around one cycle). Lean ✓ Stmt *Let K be an injective MPDO, let R give a three-site closure, and fix an η -structure and a nonzero tail entry. Let $N \geq 1$. For a cyclic sector assignment $k_0, \dots, k_{N-1}$, suppose that every concrete operator $\eta_{k_n, k_{n+1}}$ obtained from these data is nonzero. Then there are nonzero scalars c_n such that*

$$\prod_{n=0}^{N-1} c_n = 1, \quad c_n \eta_{k_n, k_{n+1}} \geq 0 \quad (0 \leq n < N).$$

This is the specialization of Theorem 23.5.3.35 to the inverse-map sector operators; positivity of the cyclic product follows from the MPDO property.

Proof. ✓ Proof The fixed-sector cyclic product is positive semidefinite by Theorem 23.5.3.38. Apply Theorem 23.5.3.35. $\square$

Theorem 23.5.3.37 (The conjugated chain is a congruence). [Lean](#) [Lean](#) [✓ Stmt](#) For every matrix U on the physical space and every $N \geq 1$,

$$\rho^{(N)}(\tilde{K}) = U^{\otimes N} \rho^{(N)}(K) (U^{\otimes N})^\dagger.$$

In particular $\rho^{(N)}(\tilde{K}) \geq 0$ whenever $\rho^{(N)}(K) \geq 0$.

Proof. [✓ Proof](#) Expand the closed trace over bond configurations g (with $g_{N+1} = g_1$), and each conjugated local entry over its two physical indices:

$$[\rho^{(N)}(\tilde{K})]_{\sigma,\tau} = \sum_g \prod_{n=1}^N \tilde{\kappa}_{g_n, g_{n+1}}^{\sigma_n \tau_n} = \sum_g \prod_{n=1}^N \sum_{p_n, q_n} U_{\sigma_n p_n} \kappa_{g_n, g_{n+1}}^{p_n q_n} \overline{U_{\tau_n q_n}}.$$

Distributing the site product over the physical sums factors the two U -products out of the bond sum, which leaves the matrix entry of $U^{\otimes N} \rho^{(N)}(K) (U^{\otimes N})^\dagger$. Positivity is preserved by any congruence. $\square$

Theorem 23.5.3.38 (Positivity of the projected chain blocks). [Lean](#) [✓ Stmt](#) Let K be an MPDO whose transformed physical slices satisfy the sector factorization hypothesis. Then for every $N \geq 1$ and every cyclic sector assignment $k_1, \dots, k_N$ (with $k_{N+1} = k_1$),

$$0 \leq [Q_{k_1} \otimes \dots \otimes Q_{k_N}] \tilde{\sigma} [Q_{k_1} \otimes \dots \otimes Q_{k_N}] = \bigotimes_{n=1}^N \eta_{k_n, k_{n+1}}.$$

This is the positivity half of the display at [CPGSV16, Appendix C.2, lines 1446–1450].

Proof. [✓ Proof](#) The cyclic tensor product Ω_k of Definition 23.5.3.16 is a diagonal sector block of the basis-conjugated chain by Theorem 23.5.3.17. The conjugated chain is positive semidefinite by Theorem 23.5.3.37 and the MPDO property, and a principal submatrix of a positive semidefinite matrix is positive semidefinite. $\square$

Theorem 23.5.3.39 (Positivity of the diagonal and paired neighboring operators). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let K be an injective MPDO with the chosen Hayashi decomposition and a nonzero tail entry. Then for all sectors k, h :

1. $\eta_{k,k} \geq 0$;
2. $\eta_{k,h} \otimes \eta_{h,k} \geq 0$;
3. if $\eta_{k,h} \neq 0$ and $\eta_{h,k} \neq 0$, there is a scalar $c \neq 0$ with $c \eta_{k,h} \geq 0$, $c^{-1} \eta_{h,k} \geq 0$, and $(c \eta_{k,h}) \otimes (c^{-1} \eta_{h,k}) = \eta_{k,h} \otimes \eta_{h,k}$.

Proof. [✓ Proof](#) The chain of length one projected to the sector k is $\eta_{k,k}$ up to the exchange of its two tensor factors, and the chain of length two projected to (k, h) is $\eta_{k,h} \otimes \eta_{h,k}$, both positive by Theorem 23.5.3.38. The third item, including the Kronecker-product-preserving equality, is Theorem 23.5.3.19. $\square$

Theorem 23.5.3.40 (Coherent positive rephasing under recurrent support). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $(\eta_{k,h})$ be a neighboring-operator family for which every nonempty cyclic tensor product is positive semidefinite. Suppose, in addition, that every nonzero edge $k \rightarrow h$ admits a directed return path from h to k . Then there are unit complex numbers z_k such that

$$z_k^{-1} z_h \eta_{k,h} \geq 0 \quad \text{for all } k, h.$$

Consequently, if the neighboring operators arise from the inverse-map physical-sector factorization of an injective MPDO and its nonzero support is recurrent, the sector tensors admit a reciprocal rephasing for which all neighboring operators are positive semidefinite. This factorization determines the positive commuting nearest-neighbor product structure of Proposition C.8.

This is a scope-restricted result. Recurrence is not a hypothesis of Lemma C.4 in [CPGSV16, Appendix C.2, lines 1406–1450]. Its derivation for the zero-weight reparameterized inverse-map factorization is treated in Theorem 23.5.3.41 and recorded in `docs/paper-gaps/cpgsv17_mpdo_sal_zcl_eta_local_structure.tex`.

Proof. ✓ Proof For a nonzero edge, choose a directed return path and close it with the edge. Positive rescaling of the corresponding cyclic tensor product supplies a scalar that makes the edge operator positive; retain only its unit-modulus part. On any closed directed walk, positive rescaling of the associated cyclic tensor product gives phases whose product is one. Uniqueness of the unit phase identifies them edge by edge with the chosen phases. Hence the chosen edge phases have product one around every closed walk. The closed-walk criterion gives $\kappa_{k,h} = z_k^{-1} z_h$ on every nonzero edge. Zero edge operators are already positive semidefinite. Finally, reciprocal rephasing of the left and right sector tensors multiplies each neighboring operator by this phase and preserves the physical-slice factorization. The conditional eta-local construction then gives the positive commuting product data. □

Theorem 23.5.3.41 (Coherent positive choice of the neighboring operators). Lean Lean Lean ✓ Stmt Let K be an injective MPO tensor satisfying SAL. Then it admits a physical-sector factorization for which every neighboring operator

$$\eta_{k,h} = \sum_a (r_k)_a \otimes (l_h)_a$$

is positive semidefinite. Zero-weight Hayashi sectors remain in the physical direct sum, but their right tensors may be set to zero without changing the physical slices.

Proof. ✓ Proof First set the right tensor to zero in every zero-weight sector. The left tensor already vanishes there, so this does not alter the factorization and removes every support edge incident to such a sector. Every remaining endpoint has positive weight. If all virtual matrices in a positive-weight sector vanished, then the corresponding middle-site block of the three-site closure would vanish; this contradicts its Hayashi expression as a nonzero weight times two trace-one sector states. Thus every endpoint of a nonzero edge contains a nonzero virtual matrix. Injectivity supplies spanning by all sector virtual matrices, and triangle closure gives recurrence. The projected chain identity makes every cyclic Kronecker product positive semidefinite. Apply Theorem 23.5.3.40 to obtain the coherent vertex rephasing. □

Theorem 23.5.3.42 (Positivity choice with symmetric support). Lean ✓ Stmt In the setting of Theorem 23.5.3.39, assume in addition that the nonzero neighboring operators occur in symmetric pairs: $\eta_{k,h} = 0$ forces $\eta_{h,k} = 0$. Then there is a positive semidefinite family $\eta'_{k,h}$ with $\eta'_{k,k} = \eta_{k,k}$ and, for every pair of sectors, scalars $c_{k,h} \neq 0$ with $\eta'_{k,h} = c_{k,h} \eta_{k,h}$ and $\eta'_{h,k} = c_{k,h}^{-1} \eta_{h,k}$. In particular every two-site block $\eta'_{k,h} \otimes \eta'_{h,k} = \eta_{k,h} \otimes \eta_{h,k}$ of the sector-adapted decomposition is unchanged.

The source asserts at [CPGSV16, Appendix C.2, lines 1446–1450] that the η 's can be chosen positive semidefinite, without the symmetric-support hypothesis. That hypothesis excludes a scalar obstruction: a nonzero neighboring operator whose reverse pair vanishes is constrained only by cycles through at least three sectors, and a coherent choice for the whole family can then fail.

Proof. ✓ Proof The diagonal operators are positive by Theorem 23.5.3.39, and their scalar is taken to be one. For a pair of distinct sectors, either both neighboring operators vanish and any nonzero scalar works, or both are nonzero by symmetric support and Theorem 23.5.3.39 supplies the reciprocal pair of scalars. Choosing the scalar once per unordered pair makes the two assignments reciprocal, and the block-preserving equality of Theorem 23.5.3.19 applied to that scalar gives $\eta'_{k,h} \otimes \eta'_{h,k} = \eta_{k,h} \otimes \eta_{h,k}$. $\square$

23.5.4 Normalized preparations and controlled partial traces

Definition 23.5.4.1 (Trace matrices of explicit η -data). Lean Lean Lean Lean ✓ Stmt Every explicit neighboring η -family determines a sector-by-sector trace matrix. We give both its complex-valued form and the real-part version, which is the direct data for the real Perron–Frobenius matrix T used in the rank-one step.

Definition 23.5.4.2 (Rank-one factorization of neighboring-operator traces). Lean ✓ Stmt Let $(\eta_{k,h})$ be a positive neighboring-operator family. A rank-one trace factorization consists of real families (a_k) and (b_h) such that

$$\mathrm{tr}(\eta_{k,h}) = a_k b_h, \quad \sum_l a_l b_l = 1.$$

These identities occur in [CPGSV16, Appendix C.2, lines 1395–1402].

Definition 23.5.4.3 (Direct-sum neighboring operator). Lean Lean ✓ Stmt For sectors k, h , define the positive-operator candidate

$$\Omega_{k,h} = \bigoplus_l (\eta_{k,l} \otimes \eta_{l,h}).$$

The summand indexed by l acts on $(B_k^R \otimes B_l^L) \otimes (B_l^R \otimes B_h^L)$. This is the operator adjointed by $\mathcal{T}_{k,h}$ in [CPGSV16, Appendix C.2, lines 1527–1535].

Theorem 23.5.4.4 (Positivity of finite dependent block diagonals). Lean ✓ Stmt Let $(A_i)_{i \in I}$ be a finite family of positive semidefinite matrices, where the size of A_i may depend on i . Then the block-diagonal matrix $\bigoplus_{i \in I} A_i$ is positive semidefinite.

Proof. ✓ Proof Write each $A_i = B_i^* B_i$ using its positive square root. The dependent block diagonal then satisfies $\bigoplus_i A_i = (\bigoplus_i B_i)^* (\bigoplus_i B_i)$. $\square$

Theorem 23.5.4.5 (Positivity of the direct-sum neighboring operator). Lean ✓ Stmt Every operator $\Omega_{k,h}$ is positive semidefinite.

Proof. ✓ Proof For every l , positivity is preserved by the Kronecker product:

$$\eta_{k,l} \succeq 0, \quad \eta_{l,h} \succeq 0 \quad \implies \quad \eta_{k,l} \otimes \eta_{l,h} \succeq 0.$$

Apply Theorem 23.5.4.4 to the family $(\eta_{k,l} \otimes \eta_{l,h})_l$. $\square$

Theorem 23.5.4.6 (Trace of the direct-sum neighboring operator). Lean Lean ✓ Stmt The trace is

$$\mathrm{tr}(\Omega_{k,h}) = \sum_l \mathrm{tr}(\eta_{k,l}) \mathrm{tr}(\eta_{l,h}).$$

Under the rank-one trace factorization, this becomes

$$\mathrm{tr}(\Omega_{k,h}) = a_k b_h \sum_l a_l b_l = a_k b_h.$$

This is the trace calculation in [CPGSV16, Appendix C.2, lines 1531–1535].

Proof. ✓ Proof The direct-sum and Kronecker trace identities give

$$\mathrm{tr}(\Omega_{k,h}) = \sum_l \mathrm{tr}(\eta_{k,l} \otimes \eta_{l,h}) = \sum_l \mathrm{tr}(\eta_{k,l}) \mathrm{tr}(\eta_{l,h}).$$

Substitute $\mathrm{tr}(\eta_{k,h}) = a_k b_h$ and $\sum_l a_l b_l = 1$. □

Theorem 23.5.4.7 (Vanishing of zero-weight neighboring operators). Lean Lean Lean ✓ Stmt Every product $a_k b_h$ is nonnegative. If $a_k b_h = 0$, then

$$\eta_{k,h} = 0, \quad \Omega_{k,h} = 0.$$

Proof. ✓ Proof Since $\eta_{k,h} \succeq 0$, its trace is nonnegative, and hence $a_k b_h = \mathrm{tr}(\eta_{k,h}) \geq 0$. If $a_k b_h = 0$, then

$$\eta_{k,h} \succeq 0, \quad \mathrm{tr}(\eta_{k,h}) = a_k b_h = 0 \implies \eta_{k,h} = 0.$$

The same implication gives $\Omega_{k,h} = 0$ because $\Omega_{k,h} \succeq 0$ and $\mathrm{tr}(\Omega_{k,h}) = a_k b_h = 0$. □

Definition 23.5.4.8 (Normalized active neighboring operators). Lean Lean ✓ Stmt With the convention $0^{-1} = 0$, define for every pair (k, h)

$$\hat{\eta}_{k,h} = (a_k b_h)^{-1} \eta_{k,h}, \quad \hat{\Omega}_{k,h} = (a_k b_h)^{-1} \Omega_{k,h}.$$

These operators are normalized when the pair is active, namely when $a_k b_h \neq 0$.

Theorem 23.5.4.9 (Positivity of normalized neighboring operators). Lean Lean ✓ Stmt For every pair (k, h) , the operators $\hat{\eta}_{k,h}$ and $\hat{\Omega}_{k,h}$ are positive semidefinite.

Proof. ✓ Proof By Theorem 23.5.4.7, $a_k b_h \geq 0$, and hence $(a_k b_h)^{-1} \geq 0$. Therefore

$$\eta_{k,h} \succeq 0, \quad (a_k b_h)^{-1} \geq 0 \implies \hat{\eta}_{k,h} = (a_k b_h)^{-1} \eta_{k,h} \succeq 0,$$

and the identical argument applies to $\hat{\Omega}_{k,h}$. □

Theorem 23.5.4.10 (Normalized active preparations). Lean Lean Lean Lean ✓ Stmt If $a_k b_h \neq 0$, then $\hat{\eta}_{k,h}$ and $\hat{\Omega}_{k,h}$ have trace one. Consequently the maps

$$X \mapsto X \otimes \hat{\eta}_{k,h}, \quad X \mapsto X \otimes \hat{\Omega}_{k,h}$$

are trace-preserving and completely positive. These are the individual preparations $\mathcal{S}_{k,h}$ and $\mathcal{T}_{k,h}$ in [CPGSV16, Appendix C.2, lines 1527–1535 and 1551–1555].

Proof. ✓ Proof On an active pair,

$$\mathrm{tr}(\hat{\eta}_{k,h}) = (a_k b_h)^{-1} \mathrm{tr}(\eta_{k,h}) = (a_k b_h)^{-1} a_k b_h = 1.$$

Likewise, $\mathrm{tr}(\hat{\Omega}_{k,h}) = 1$ follows from $\mathrm{tr}(\Omega_{k,h}) = a_k b_h$. Apply Lemma 23.1.23 to each operator. □

Lemma 23.5.4.11 (Trace-one matrices have nonempty index spaces). Lean ✓ Stmt If a matrix indexed by a finite type has trace one, then its index type is nonempty.

Proof. ✓ Proof A matrix on an empty index space has trace zero, contrary to the hypothesis. □

Lemma 23.5.4.12 (Nonemptiness of the Hayashi sector factors). Lean Lean ✓ Stmt Every left and right factor in a Hayashi sector is nonzero-dimensional.

Proof. ✓ Proof A matrix on an empty index space has trace zero. Hence

$$\text{tr}(\rho_k^R) = 1 \implies \dim(B_k^R \otimes C) > 0 \implies \dim B_k^R > 0,$$

and the same argument applied to ρ_k^L gives $d_k^L > 0$. □

Lemma 23.5.4.13 (Existence of a Hayashi sector). Lean ✓ Stmt *The set of Hayashi sectors is nonempty.*

Proof. ✓ Proof The sector weights satisfy

$$\sum_k p_k = 1 \implies m > 0.$$

□

Lemma 23.5.4.14 (Nonemptiness of the neighboring auxiliary spaces). Lean Lean ✓ Stmt *For every pair of sectors (k, h) , the spaces carrying $\eta_{k,h}$ and $\Omega_{k,h}$ are nonzero-dimensional.*

Proof. ✓ Proof The tensor product carrying $\eta_{k,h}$ is nonzero-dimensional by Lemma 23.5.4.12. Choosing an intermediate sector l by Lemma 23.5.4.13 then supplies an element of the direct-sum space carrying $\Omega_{k,h}$. □

Definition 23.5.4.15 (Faithful uniform density matrix). Lean ✓ Stmt On a nonzero finite-dimensional space H , define

$$\tau_H = \frac{1}{\dim H} I_H.$$

Theorem 23.5.4.16 (The uniform density matrix is faithful and normalized). Lean Lean ✓ Stmt *The matrix τ_H is positive definite and has trace one.*

Proof. ✓ Proof Since $\dim H > 0$, the scalar $(\dim H)^{-1}$ is positive. Thus τ_H is positive definite, and $\text{tr}(\tau_H) = (\dim H)^{-1} \dim H = 1$. □

Definition 23.5.4.17 (Completed neighboring-state preparation). Lean Lean Lean ✓ Stmt For every sector pair define

$$\bar{\eta}_{k,h} = \begin{cases} \hat{\eta}_{k,h}, & a_k b_h \neq 0, \\ \tau_{B_k^R \otimes B_h^L}, & a_k b_h = 0, \end{cases} \quad \bar{\mathcal{S}}_{k,h}(X) = X \otimes \bar{\eta}_{k,h}.$$

On an active pair this is precisely the preparation $\mathcal{S}_{k,h}$ of [CPGSV16, Appendix C.2, lines 1551–1555]; the second branch gives a normalized extension on a pair where the displayed source quotient is undefined.

Theorem 23.5.4.18 (Completed neighboring-state preparations are trace-preserving completely positive). Lean Lean Lean Lean Lean ✓ Stmt *For every pair (k, h) , the operator $\bar{\eta}_{k,h}$ is positive semidefinite with trace one. Hence $\bar{\mathcal{S}}_{k,h}$ is trace-preserving and completely positive. If $a_k b_h \neq 0$, then both the density matrix and the preparation equal their normalized active forms.*

Proof. ✓ Proof The two branches give

$$a_k b_h \neq 0 \implies \bar{\eta}_{k,h} = \hat{\eta}_{k,h} \succeq 0, \quad \text{tr}(\bar{\eta}_{k,h}) = 1,$$

and

$$a_k b_h = 0 \implies \bar{\eta}_{k,h} = \tau_{B_k^R \otimes B_h^L} \succ 0, \quad \text{tr}(\bar{\eta}_{k,h}) = 1.$$

The state-preparation result applies in either case. □

Definition 23.5.4.19 (Completed direct-sum preparation). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For every sector pair define

$$\mathcal{H}_{k,h}^\Omega = \bigoplus_l (B_k^R \otimes B_l^L) \otimes (B_l^R \otimes B_h^L), \quad \bar{\Omega}_{k,h} = \begin{cases} \hat{\Omega}_{k,h}, & a_k b_h \neq 0, \\ \tau_{\mathcal{H}_{k,h}^\Omega}, & a_k b_h = 0, \end{cases} \quad \bar{\mathcal{T}}_{k,h}(X) = X \otimes \bar{\Omega}_{k,h},$$

where the second density acts on the full direct-sum index space of $\Omega_{k,h}$. On an active pair this is precisely the preparation $\mathcal{T}_{k,h}$ of [CPGSV16, Appendix C.2, lines 1527–1535]; the inactive branch is a normalized extension.

Theorem 23.5.4.20 (Completed direct-sum preparations are trace-preserving completely positive). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For every pair (k, h) , the operator $\bar{\Omega}_{k,h}$ is positive semidefinite with trace one. Hence $\bar{\mathcal{T}}_{k,h}$ is trace-preserving and completely positive. On every active pair it equals the normalized active preparation.

Proof. [✓ Proof](#) On the active branch,

$$a_k b_h \neq 0 \implies \bar{\Omega}_{k,h} = \hat{\Omega}_{k,h} \succeq 0, \quad \text{tr}(\bar{\Omega}_{k,h}) = 1,$$

while on the inactive branch,

$$a_k b_h = 0 \implies \bar{\Omega}_{k,h} = \tau_{\mathcal{H}_{k,h}^\Omega} \succ 0, \quad \text{tr}(\bar{\Omega}_{k,h}) = 1.$$

The state-preparation result therefore applies to $\bar{\mathcal{T}}_{k,h}$. □

Definition 23.5.4.21 (Globally controlled neighboring-state preparation). [Lean](#) [✓ Stmt](#) Let $H = \bigoplus_{k,h} H_{k,h}$. Orthogonally controlling the completed preparations over the sector pairs gives a map

$$\bar{\mathcal{S}}_1(X) = \sum_{k,h,a} \tilde{A}_{k,h,a} X \tilde{A}_{k,h,a}^\dagger,$$

where $(A_{k,h,a})_a$ is a Kraus family for $\bar{\mathcal{S}}_{k,h}$ and the tilde denotes its extension by zero outside $H_{k,h}$. Thus the map discards inter-sector coherences and applies the completed preparation in each diagonal sector. On active sectors this is the sector-control formula for $\mathcal{S}_1$ in [CPGSV16, Appendix C.2, lines 1548–1555].

Theorem 23.5.4.22 (Diagonal-sector action of the completed neighboring preparation). [Lean](#) [✓ Stmt](#) If $X_{k,h}$ is the (k, h) diagonal block of X , then

$$[\bar{\mathcal{S}}_1(X)]_{k,h} = \bar{\mathcal{S}}_{k,h}(X_{k,h}) = X_{k,h} \otimes \bar{\eta}_{k,h}.$$

Proof. [✓ Proof](#) The diagonal-block formula and invariance under Kraus-index relabeling give

$$[\bar{\mathcal{S}}_1(X)]_{k,h} = \Phi_{\tilde{A}_{k,h}}(X_{k,h}) = \Phi_{A_{k,h}}(X_{k,h}).$$

The preparation Kraus identity then gives

$$\Phi_{A_{k,h}}(X_{k,h}) = X_{k,h} \otimes \bar{\eta}_{k,h}.$$

□

Theorem 23.5.4.23 (The globally controlled neighboring-state preparation is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) The map $\bar{\mathcal{S}}_1$ is trace-preserving and completely positive.

Proof. ✓ Proof If $(A_{k,h,a})_a$ is the sectorwise Kraus family, then

$$\sum_{k,h,a} \tilde{A}_{k,h,a}^\dagger \tilde{A}_{k,h,a} = \bigoplus_{k,h} \left(\sum_a A_{k,h,a}^\dagger A_{k,h,a} \right) = \bigoplus_{k,h} I_{H_{k,h}} = I_H.$$

Thus the controlled Kraus family is trace-preserving. □

Definition 23.5.4.24 (Globally controlled direct-sum preparation). Lean ✓ Stmt Orthogonally controlling the completed direct-sum preparations gives

$$\bar{\mathcal{T}}_1(X) = \sum_{k,h,a} \tilde{B}_{k,h,a} X \tilde{B}_{k,h,a}^\dagger,$$

where $(B_{k,h,a})_a$ is a Kraus family for $\bar{\mathcal{T}}_{k,h}$. This map discards inter-sector coherences and, on active sectors, is the sector-control formula for $\mathcal{T}_1$ in [CPGSV16, Appendix C.2, lines 1523–1535].

Theorem 23.5.4.25 (Diagonal-sector action of the completed direct-sum preparation). Lean

✓ Stmt If $X_{k,h}$ is the (k, h) diagonal block of X , then

$$[\bar{\mathcal{T}}_1(X)]_{k,h} = \bar{\mathcal{T}}_{k,h}(X_{k,h}) = X_{k,h} \otimes \bar{\Omega}_{k,h}.$$

Proof. ✓ Proof The diagonal-block formula and invariance under Kraus-index relabeling give

$$[\bar{\mathcal{T}}_1(X)]_{k,h} = \Phi_{\tilde{B}_{k,h}}(X_{k,h}) = \Phi_{B_{k,h}}(X_{k,h}).$$

The preparation Kraus identity then gives

$$\Phi_{B_{k,h}}(X_{k,h}) = X_{k,h} \otimes \bar{\Omega}_{k,h}.$$

□

Theorem 23.5.4.26 (The globally controlled direct-sum preparation is trace-preserving completely positive). Lean ✓ Stmt The map $\bar{\mathcal{T}}_1$ is trace-preserving and completely positive.

Proof. ✓ Proof For the sectorwise Kraus family $(B_{k,h,a})_a$,

$$\sum_{k,h,a} \tilde{B}_{k,h,a}^\dagger \tilde{B}_{k,h,a} = \bigoplus_{k,h} \left(\sum_a B_{k,h,a}^\dagger B_{k,h,a} \right) = \bigoplus_{k,h} I_{H_{k,h}} = I_H.$$

The orthogonal-control theorem now gives the claim. □

Remark 23.5.4.27 (Scope of the completed preparation). The conclusions above begin with the rank-one trace factorization $\text{tr}(\eta_{k,h}) = a_k b_h$ and $\sum_l a_l b_l = 1$. Deriving this factorization for the inverse-map neighboring operators from the strong area law and zero correlation length is the Perron–Frobenius step of [CPGSV16, Appendix C.2, lines 1404–1410 and 1484–1498]. The completed maps therefore give the global controlled preparation once that factorization is supplied; they do not by themselves prove the full two-map statement [CPGSV16, Appendix C.2, lines 1510–1517]. The regrouping maps, shift maps, and the required closure identities are separate parts of that statement.

Definition 23.5.4.28 (Controlled dependent partial trace). [Lean](#) [✓ Stmt](#) Let

$$H = \bigoplus_{i \in I} (A_i \otimes B_i), \quad K = \bigoplus_{i \in I} A_i.$$

The controlled dependent partial trace discards the off-diagonal blocks between distinct i and applies tr_{B_i} to the i th diagonal block.

Theorem 23.5.4.29 (Diagonal blocks of the controlled dependent partial trace). [Lean](#) [✓ Stmt](#) If X_{ii} denotes the i th diagonal block of X , then

$$[\mathcal{C}_{\text{tr}}(X)]_{ii} = \text{tr}_{B_i}(X_{ii}).$$

Proof. [✓ Proof](#) If $(K_{i,j})_j$ is the Kraus family chosen for tr_{B_i} , then the diagonal-block formula gives

$$[\mathcal{C}_{\text{tr}}(X)]_{ii} = \sum_j K_{i,j} X_{ii} K_{i,j}^\dagger = \text{tr}_{B_i}(X_{ii}).$$

□

Theorem 23.5.4.30 (The controlled dependent partial trace is trace-preserving completely positive). [Lean](#) [✓ Stmt](#) The controlled dependent partial trace is trace-preserving and completely positive.

Proof. [✓ Proof](#) Choose Kraus operators $(V_{i,a})_a$ for tr_{B_i} and extend each $V_{i,a}$ by zero outside the i th summand. Their controlled family satisfies

$$\sum_{i,a} \widetilde{V}_{i,a}^\dagger \widetilde{V}_{i,a} = \bigoplus_i \left(\sum_a V_{i,a}^\dagger V_{i,a} \right) = \bigoplus_i I_{A_i \otimes B_i} = I_H.$$

Its Kraus form gives complete positivity, and the displayed resolution of the identity gives trace preservation. □

Definition 23.5.4.31 (Two-site subspin regrouping). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) In sector-adapted coordinates, regroup two sites by

$$((l_k, r_k), (l_h, r_h)) \mapsto ((l_k, r_h), (r_k, l_h)).$$

Globally this gives the dependent-sum equivalence

$$\left(\bigoplus_k L_k \otimes R_k \right)^{\otimes 2} \simeq \bigoplus_{k,h} (L_k \otimes R_h) \otimes (R_k \otimes L_h).$$

The forward equivalence prepares the neighboring factors $R_k \otimes L_h$ for the partial trace in $\mathcal{T}_0$; its inverse is the shift $\mathcal{S}_2$. These are the reorderings in [CPGSV16, Appendix C.2, lines 1521–1522 and 1555–1559].

Definition 23.5.4.32 (The two-site partial-trace map $\mathcal{T}_0$). [Lean](#) [Lean](#) [✓ Stmt](#) Define $\mathcal{T}_0$ by the two-site regrouping followed, in each pair (k, h) , by the partial trace over $R_k \otimes L_h$. Its output is indexed by the retained outer factors $L_k \otimes R_h$. Thus, writing R_{Φ_2} for the regrouping reindexing,

$$\mathcal{T}_0 = \text{CTr}_{(R_k \otimes L_h)_{k,h}} \circ R_{\Phi_2}.$$

Definition 23.5.4.39 (The three-site shift $\mathcal{T}_2$). [Lean](#) [Lean](#) [✓ Stmt](#) The map $\mathcal{T}_2$ is matrix reindexing along the inverse three-site equivalence. On each summand it sends

$$((l_k, r_h), ((r_k, l_l), (r_l, l_h))) \mapsto ((l_k, r_k), ((l_l, r_l), (l_h, r_h))).$$

Theorem 23.5.4.40 ($\mathcal{T}_2$ is trace-preserving completely positive). [Lean](#) [Lean](#) [✓ Stmt](#) *The shift $\mathcal{T}_2$ is trace-preserving and completely positive.*

Proof. [✓ Proof](#) If Φ_3 is the three-site regrouping, then $\mathcal{T}_2 = R_{\Phi_3^{-1}}$. It is therefore trace-preserving and completely positive by Theorem 23.1.7. $\square$

Definition 23.5.4.41 (Fiberwise partial-trace maps). [Lean](#) [Lean](#) [✓ Stmt](#) For fixed sectors (k, h) , define the fiberwise form of $\mathcal{T}_0$ by regrouping two sites and tracing $R_k \otimes L_h$. For fixed (k, l, h) , define the fiberwise form of $\mathcal{S}_0$ by regrouping three sites and tracing $(R_k \otimes L_l) \otimes (R_l \otimes L_h)$.

Theorem 23.5.4.42 (The fiberwise partial traces are trace-preserving completely positive). [Lean](#) [Lean](#) [✓ Stmt](#) *Every fiberwise form of $\mathcal{T}_0$ and $\mathcal{S}_0$ is trace-preserving and completely positive.*

Proof. [✓ Proof](#) The fiberwise forms factor as

$$\mathcal{T}_0^{(k,h)} = \text{tr}_{R_k \otimes L_h} \circ R_{\phi_{k,h}}, \quad \mathcal{S}_0^{(k,l,h)} = \text{tr}_{(R_k \otimes L_l) \otimes (R_l \otimes L_h)} \circ R_{\phi_{k,l,h}},$$

where $\phi_{k,h}$ is the two-site equivalence of Definition 23.5.4.31 on the (k, h) sector, $\phi_{k,l,h}$ is the three-site equivalence of Definition 23.5.4.36 on the (k, l, h) sector, and R_ϕ denotes equivalence reindexing. Both factors in each composition are trace-preserving and completely positive, so the composition is as well. $\square$

Definition 23.5.4.43 (Fiberwise shifts). [Lean](#) [Lean](#) [✓ Stmt](#) On fixed sector pairs, define $\mathcal{S}_2$ and $\mathcal{T}_2$ by the same inverse regroupings as the corresponding global shifts.

Theorem 23.5.4.44 (The fiberwise shifts are trace-preserving completely positive). [Lean](#) [Lean](#) [✓ Stmt](#) *Every fiberwise shift is trace-preserving and completely positive.*

Proof. [✓ Proof](#) The shifts $\mathcal{S}_2^{(k,h)}$ and $\mathcal{T}_2^{(k,h)}$ are reindexings along the inverse two-site and three-site equivalences of Definitions 23.5.4.31 and 23.5.4.36, respectively. They are therefore trace-preserving and completely positive by Theorem 23.1.7. $\square$

On each fixed (k, h) -summand, the controlled maps $\mathcal{T}_0$ and $\mathcal{S}_0$ discard coherences between distinct outer summands and take the displayed partial trace in each diagonal summand. In particular, $\mathcal{S}_0$ traces the whole space

$$\bigoplus_l [(R_k \otimes L_l) \otimes (R_l \otimes L_h)].$$

The shifts $\mathcal{S}_2$ and $\mathcal{T}_2$ are global reindexings, with all displayed tensor factors present at their inputs [CPGSV16, Appendix C.2, lines 1521–1559].

23.5.5 Refinement and coarse-graining channels

Theorem 23.5.5.1 (Positivity and zero-weight neighboring operators). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The sector set and every middle-subspin space are nonempty. Moreover, $\Omega_{k,h}$ is positive semidefinite, and*

$$a_k b_h = 0 \implies \Omega_{k,h} = 0.$$

$$\begin{array}{c}
\bigoplus_{k,h}; \text{ one } (k,h)\text{-fiber is drawn} \\
\mathcal{T}_0 : \mathfrak{R}_2 \rightarrow L_k \otimes R_h, \quad \mathcal{S}_2 : (L_k \otimes R_h) \otimes (R_k \otimes L_h) \rightarrow \mathfrak{R}_2 \\
\bigoplus_{k,h}; \text{ one } (k,h)\text{-fiber and one } l\text{-summand are drawn} \\
\mathcal{S}_0 : \mathfrak{R}_3 \rightarrow L_k \otimes R_h, \quad \mathcal{T}_2 : (L_k \otimes R_h) \otimes \Omega_{k,h} \rightarrow \mathfrak{R}_3
\end{array}$$

Figure 23.5: Fiberwise partial traces and regrouping maps on two and three physical sites. One (k, h) -summand and one l -summand are drawn.

Definition 23.5.5.2 (The three-site neighboring-state preparation $\mathcal{T}_1$). [Lean](#) [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [✓ Stmt](#) On a sector pair with $a_k b_h \neq 0$, set

$$\widehat{\Omega}_{k,h} = (a_k b_h)^{-1} \Omega_{k,h}.$$

If $a_k b_h = 0$, positivity and vanishing trace imply $\Omega_{k,h} = 0$; choose any density operator on that summand and denote the resulting completed density by $\overline{\Omega}_{k,h}$. The sectorwise preparation is

$$X \mapsto X \otimes \overline{\Omega}_{k,h}.$$

Taking the orthogonal direct sum over the outer sectors (k, h) gives the preparation stage $\mathcal{T}_1$ of [CPGSV16, Appendix C.2, lines 1523–1535]. The zero-weight choice makes the source map total without imposing a nonzero-weight hypothesis.

Theorem 23.5.5.3 (Properties of the three-site neighboring densities). [Lean](#) [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The normalized density is positive semidefinite and has trace one on every active pair. The density chosen on an inactive pair is positive definite and has trace one. Hence $\overline{\Omega}_{k,h}$ is positive semidefinite with trace one for every pair, and agrees with $\widehat{\Omega}_{k,h}$ on active pairs. The corresponding sectorwise and controlled maps prepare these densities on their diagonal summands.*

Theorem 23.5.5.4 ($\mathcal{T}_1$ is trace-preserving completely positive). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#)
Each completed preparation is trace-preserving and completely positive. The same holds for their orthogonally controlled direct sum.

Proof. [✓ Proof](#) On an active pair, $\text{tr}(\widehat{\Omega}_{k,h}) = (a_k b_h)^{-1} a_k b_h = 1$; the chosen density on a zero-weight pair also has trace one. Preparation by a positive operator of trace one is trace-preserving and completely positive. The orthogonal control discards coherences between distinct outer-sector pairs and applies these preparations on the diagonal summands. $\square$

Definition 23.5.5.5 (The refinement channel $\mathcal{T}$). [Lean](#) [✓ Stmt](#) Define the channel from two physical sites to three physical sites by

$$\mathcal{T} = \mathcal{T}_2 \mathcal{T}_1 \mathcal{T}_0.$$

Thus $\mathcal{T}_0$ traces the neighboring subspins, $\mathcal{T}_1$ adjoins the completed three-site neighboring density, and $\mathcal{T}_2$ restores the three complete physical sectors. This is the refinement map in [CPGSV16, Appendix C.2, lines 1522–1545].

Definition 23.5.5.11 (Two- and three-site sector coordinates). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#)

The coordinate equivalences used by the coarse-graining channel expose the direct sums over site-sector pairs and triples. Write $\mathfrak{R}_2$ and $\mathfrak{R}_3$ for the corresponding matrix reindexings. Their inverses send every sector block to the corresponding physical matrix entries. These reindexings preserve the left-right order within every physical site. They differ from the regrouped coordinates $\mathfrak{G}_2$ and $\mathfrak{G}_3$ of Definition 23.5.2.27, which place the two outer subspins before the neighboring subspins. If R_{Φ_2} and R_{Φ_3} denote the two regrouping maps, then

$$\mathfrak{G}_2 = R_{\Phi_2} \mathfrak{R}_2, \quad \mathfrak{G}_3 = R_{\Phi_3} \mathfrak{R}_3.$$

Theorem 23.5.5.12 (Diagonal outer-sector block of the three-site closure). [Lean](#) [✓ Stmt](#) For every virtual matrix X and outer-sector pair (k, h) ,

$$\left[R_{\Phi_3} \mathfrak{R}_3(K_3(X)) \right]_{(k,h),(k,h)} = B_{k,h}(X) \otimes \Omega_{k,h}.$$

Proof. [✓ Proof](#) On the summand labelled by the middle sector l , this is the fixed-sector factorization $B_{k,h}(X) \otimes (\eta_{k,l} \otimes \eta_{l,h})$. Taking the direct sum over l gives the stated block. $\square$

Theorem 23.5.5.13 (Off-diagonal outer-sector entries of the three-site closure). [Lean](#) [✓ Stmt](#) Every matrix entry of $R_{\Phi_3} \mathfrak{R}_3(K_3(X))$ between distinct outer-sector pairs $(k, h) \neq (p, q)$ vanishes.

Proof. [✓ Proof](#) Distinct outer-sector pairs differ either in their first sector or in their third sector. In the corresponding physical slice, the sector-coordinate tensor is block diagonal, so that factor vanishes. $\square$

Theorem 23.5.5.14 (Regrouped two-site closure block). [Lean](#) [✓ Stmt](#) For every virtual matrix X and outer-sector pair (k, h) ,

$$\left[R_{\Phi_2} \mathfrak{R}_2(K_2(X)) \right]_{(k,h),(k,h)} = B_{k,h}(X) \otimes \eta_{k,h}.$$

Proof. [✓ Proof](#) This is the fixed-sector two-site factorization after identifying the sitewise sector coordinates with the regrouped boundary and neighboring coordinates. $\square$

Theorem 23.5.5.15 (Diagonal-block action of $\mathcal{T}_0$). [Lean](#) [✓ Stmt](#) For every two-site matrix Y , outer-sector pair (k, h) , and boundary indices a, b ,

$$[\mathcal{T}_0(Y)]_{(k,h,a),(k,h,b)} = \left[\text{tr}_{R_k \otimes L_h} \left([R_{\Phi_2} Y]_{(k,h),(k,h)} \right) \right]_{a,b}.$$

Proof. [✓ Proof](#) The controlled partial trace acts on a diagonal outer-sector block by the ordinary partial trace over its neighboring factor. $\square$

Theorem 23.5.5.16 (Partial trace of a factorized two-site block). [Lean](#) [✓ Stmt](#) If the (k, h) diagonal block of $R_{\Phi_2} Y$ is $B \otimes \eta_{k,h}$, then

$$[\mathcal{T}_0(Y)]_{(k,h),(k,h)} = a_k b_h B.$$

Proof. [✓ Proof](#) The partial trace of $B \otimes \eta_{k,h}$ is $\text{tr}(\eta_{k,h})B = a_k b_h B$. $\square$

Theorem 23.5.5.17 (Preparation of a scaled boundary block). [Lean](#) [✓ Stmt](#) For every outer-sector pair (k, h) and boundary matrix B , the completed sectorwise preparation satisfies

$$\overline{\mathcal{T}}_{k,h}(a_k b_h B) = B \otimes \Omega_{k,h}.$$

Proof. [✓ Proof](#) If $a_k b_h \neq 0$, the scalar cancels the normalization in $\widehat{\Omega}_{k,h} = (a_k b_h)^{-1} \Omega_{k,h}$. If $a_k b_h = 0$, both sides vanish because $\Omega_{k,h} = 0$. $\square$

Theorem 23.5.5.18 (Diagonal-block action of $\mathcal{T}_1$). [Lean](#) [✓ Stmt](#) *If the (k, h) diagonal block of a boundary matrix Y is $a_k b_h B$, then*

$$[\mathcal{T}_1(Y)]_{(k,h),(k,h)} = B \otimes \Omega_{k,h}.$$

Proof. [✓ Proof](#) On a diagonal outer-sector block, the controlled map is the completed sectorwise preparation, to which the preceding theorem applies. $\square$

Theorem 23.5.5.19 (Off-diagonal action of $\mathcal{T}_1$). [Lean](#) [✓ Stmt](#) *The map $\mathcal{T}_1$ annihilates every matrix entry between distinct outer-sector pairs.*

Proof. [✓ Proof](#) This is the orthogonal control in the definition of $\mathcal{T}_1$. $\square$

Theorem 23.5.5.20 (Reindexing action of $\mathcal{T}_2$). [Lean](#) [✓ Stmt](#) *Let Φ_3 be the three-site regrouping. For every prepared matrix Y and three-site indices p, q ,*

$$[\mathcal{T}_2(Y)]_{p,q} = Y_{\Phi_3(p), \Phi_3(q)}.$$

Proof. [✓ Proof](#) This is matrix reindexing along the inverse regrouping Φ_3^{-1} . $\square$

Theorem 23.5.5.21 (Refinement of the two-site closure). [Lean](#) [✓ Stmt](#) *Assume the neighboring-operator trace factorization of Definition 23.5.2.24. For every virtual matrix X , in the canonical physical-sector coordinates selected by the isometry U , the refinement channel satisfies*

$$\mathcal{T}(\mathfrak{R}_2(K_2(X))) = \mathfrak{R}_3(K_3(X)).$$

This is the refinement half of Proposition C.7 [CPGSV16, Appendix C.2, lines 1510–1516 and 1522–1545].

Proof. [✓ Proof](#) The regrouping contained in $\mathcal{T}_0$ changes $\mathfrak{R}_2$ into $\mathfrak{G}_2$. On the (k, h) diagonal block, the partial trace then gives

$$\mathcal{T}_0(\mathfrak{R}_2(K_2(X)))_{k,h} = a_k b_h B_{k,h}(X).$$

On an active pair, $\mathcal{T}_1$ adjoins $(a_k b_h)^{-1} \Omega_{k,h}$, and hence

$$\mathcal{T}_1 \mathcal{T}_0(\mathfrak{R}_2(K_2(X)))_{k,h} = B_{k,h}(X) \otimes \Omega_{k,h}.$$

If $a_k b_h = 0$, both sides vanish because $\Omega_{k,h} = 0$. After the canonical reassociation of the intermediate sector sum, these are precisely the diagonal blocks of $\mathfrak{G}_3(K_3(X))$; all off-diagonal sector blocks vanish. Finally, the inverse regrouping $\mathcal{T}_2$ changes $\mathfrak{G}_3$ into $\mathfrak{R}_3$, which gives the stated equality. $\square$

Theorem 23.5.5.22 (Coarse-graining of the three-site closure). [Lean](#) [Lean](#) [✓ Stmt](#) *Assume the neighboring-operator trace factorization of Definition 23.5.2.24. For every virtual matrix X , in the canonical physical-sector coordinates selected by the isometry U , the coarse-graining channel satisfies*

$$\mathcal{S}(\mathfrak{R}_3(K_3(X))) = \mathfrak{R}_2(K_2(X)).$$

Proof. ✓ Proof On the (k, h) diagonal block, $\mathcal{S}_0$ traces $\Omega_{k,h}$ and gives $a_k b_h B_{k,h}(X)$. On an active sector pair,

$$\mathcal{S}_1((a_k b_h) B_{k,h}(X)) = B_{k,h}(X) \otimes \eta_{k,h}.$$

If $a_k b_h = 0$, positivity and zero trace give $\eta_{k,h} = 0$, so both sides of this identity vanish. Finally, $\mathcal{S}_2$ regroups the four factors into two physical sites. For distinct sector pairs $(k, h) \neq (p, q)$,

$$(\mathcal{S}_1 Y)_{(k,h),(p,q)} = 0, \quad [\Re_2(K_2(X))]_{(k,h),(p,q)} = 0.$$

□

For the following construction, through Theorem 23.5.5.42, assume the neighboring-operator trace factorization of Definition 23.5.2.24. Write

$$q = \dim \left(\bigoplus_k L_k \otimes R_k \right)$$

for the physical dimension of the sector-coordinate tensor $\widehat{K}$.

Definition 23.5.5.23 (Refinement in ordinary physical coordinates). Lean ✓ Stmt Transporting $\mathcal{T}$ through the canonical two-site and three-site coordinate identifications gives a channel

$$\widehat{\mathcal{T}} : M_{q^2}(\mathbb{C}) \longrightarrow M_{q^3}(\mathbb{C}),$$

where all multiple physical indices are associated to the right.

Theorem 23.5.5.24 (The physical refinement map is a channel). Lean ✓ Stmt The map $\widehat{\mathcal{T}}$ is trace-preserving and completely positive.

Proof. ✓ Proof The coordinate identifications are unitary permutation channels. Composing them with a trace-preserving completely positive map preserves both properties. □

Theorem 23.5.5.25 (Refinement identity in ordinary physical coordinates). Lean ✓ Stmt For every virtual matrix X ,

$$\widehat{\mathcal{T}}(\widehat{K}_2(X)) = \widehat{K}_3(X).$$

Proof. ✓ Proof Transport the sector-coordinate identity through the inverse three-site coordinate identification. □

Definition 23.5.5.26 (Physical coarse-graining). Lean ✓ Stmt Transporting $\mathcal{S}$ through the canonical three-site and two-site coordinate identifications gives a channel

$$\widehat{\mathcal{S}} : M_{q^3}(\mathbb{C}) \longrightarrow M_{q^2}(\mathbb{C}).$$

Theorem 23.5.5.27 (The physical coarse-graining map is a channel). Lean ✓ Stmt The map $\widehat{\mathcal{S}}$ is trace-preserving and completely positive.

Proof. ✓ Proof This follows by composition with the two coordinate-permutation channels. □

Theorem 23.5.5.28 (Coarse-graining identity in ordinary physical coordinates). Lean ✓ Stmt For every virtual matrix X ,

$$\widehat{\mathcal{S}}(\widehat{K}_3(X)) = \widehat{K}_2(X).$$

Proof. ✓ Proof Transport the sector-coordinate identity through the inverse two-site coordinate identification. □

Definition 23.5.5.29 (Localized physical refinement). [Lean](#) [✓ Stmt](#) Reassociating three right-associated sites as (12)3, applying $\widehat{\mathcal{T}} \otimes \text{id}$ to the first two, and reassociating the result back to 1(2(34)) defines the localized refinement channel

$$\widehat{\mathcal{T}}_{(12)3} : M_{q^3}(\mathbb{C}) \longrightarrow M_{q^4}(\mathbb{C})$$

on right-associated physical coordinates.

Theorem 23.5.5.30 (The localized physical refinement is a channel). [Lean](#) [✓ Stmt](#) *The localized refinement map is trace-preserving and completely positive.*

Proof. [✓ Proof](#) Tensoring a channel with the identity preserves complete positivity and trace, as do the two associativity permutations. $\square$

Theorem 23.5.5.31 (Localized refinement of physical closures). [Lean](#) [✓ Stmt](#) *For every virtual matrix X ,*

$$(\widehat{\mathcal{T}} \otimes \text{id})(\widehat{K}_3(X)) = \widehat{K}_4(X),$$

with the displayed map understood in right-associated coordinates.

Proof. [✓ Proof](#) Fixing the last ket and bra indices turns the three-site closure into a two-site closure with the last tensor matrix absorbed into X . Apply the two-to-three-site refinement identity to this virtual matrix. $\square$

Definition 23.5.5.32 (Localized physical coarse-graining). [Lean](#) [✓ Stmt](#) Reassociating four right-associated sites as (123)4, applying $\widehat{\mathcal{S}} \otimes \text{id}$ to the first three, and reassociating the result back to 1(23) defines the localized coarse-graining channel

$$\widehat{\mathcal{S}}_{(123)4} : M_{q^4}(\mathbb{C}) \longrightarrow M_{q^3}(\mathbb{C})$$

on right-associated physical coordinates.

Theorem 23.5.5.33 (The localized physical coarse-graining is a channel). [Lean](#) [✓ Stmt](#) *The localized coarse-graining map is trace-preserving and completely positive.*

Proof. [✓ Proof](#) Tensor the physical coarse-graining channel with the identity and compose with the two associativity permutations. $\square$

Theorem 23.5.5.34 (Localized coarse-graining of physical closures). [Lean](#) [✓ Stmt](#) *For every virtual matrix X ,*

$$(\widehat{\mathcal{S}} \otimes \text{id})(\widehat{K}_4(X)) = \widehat{K}_3(X),$$

in right-associated coordinates.

Proof. [✓ Proof](#) Fixing the last ket and bra indices identifies the four-site closure with a three-site closure having the last tensor matrix absorbed into X . Apply the three-to-two-site coarse-graining identity. $\square$

Definition 23.5.5.35 (Two-step physical refinement). [Lean](#) [✓ Stmt](#) Let $\widehat{K}$ be the sector-coordinate tensor selected by the physical isometry of Proposition C.7. Define

$$\widehat{\mathcal{T}}^2 = \widehat{\mathcal{T}}_{(12)3} \circ \widehat{\mathcal{T}}, \quad \widehat{\mathcal{T}}^2 : M_{q^2}(\mathbb{C}) \longrightarrow M_{q^4}(\mathbb{C}),$$

where the second application acts on the first two sites and leaves the third site unchanged. Thus the superscript denotes two successive overlapping applications, as in the proof of [CPGSV16, Theorem 4.9, (iv) $\Rightarrow$ (v)].

Theorem 23.5.5.36 (The two-step physical refinement is a channel). [Lean](#) [✓ Stmt](#) *The map $\widehat{\mathcal{T}}^2$ is trace-preserving and completely positive.*

Proof. [✓ Proof](#) It is the composition of the physical refinement channel and its localization on the first two sites. $\square$

Theorem 23.5.5.37 (Two-step refinement of the sector-coordinate tensor). [Lean](#) [✓ Stmt](#) *For every virtual matrix X , the sector-coordinate tensor satisfies*

$$\widehat{\mathcal{T}}^2(\widehat{K}_2(X)) = \widehat{K}_4(X).$$

Proof. [✓ Proof](#) The first refinement gives $\widehat{K}_3(X)$; applying the localized refinement to its first two sites gives $\widehat{K}_4(X)$. $\square$

Definition 23.5.5.38 (Two-step physical coarse-graining). [Lean](#) [✓ Stmt](#) Define

$$\widehat{\mathcal{S}}^2 = \widehat{\mathcal{S}} \circ \widehat{\mathcal{S}}_{(123)4}, \quad \widehat{\mathcal{S}}^2 : M_{q^4}(\mathbb{C}) \longrightarrow M_{q^2}(\mathbb{C}),$$

where the first application acts on the first three sites and leaves the fourth site unchanged. The superscript again denotes successive overlapping applications.

Theorem 23.5.5.39 (The two-step physical coarse-graining is a channel). [Lean](#) [✓ Stmt](#) *The map $\widehat{\mathcal{S}}^2$ is trace-preserving and completely positive.*

Proof. [✓ Proof](#) It is the composition of the localized coarse-graining channel and the physical coarse-graining channel. $\square$

Theorem 23.5.5.40 (Two-step coarse-graining of the sector-coordinate tensor). [Lean](#) [✓ Stmt](#) *For every virtual matrix X , the sector-coordinate tensor satisfies*

$$\widehat{\mathcal{S}}^2(\widehat{K}_4(X)) = \widehat{K}_2(X).$$

Proof. [✓ Proof](#) The localized coarse-graining first gives $\widehat{K}_3(X)$; the second coarse-graining gives $\widehat{K}_2(X)$. $\square$

Theorem 23.5.5.41 (Blocked sector-coordinate tensor is a renormalization fixed point). [Lean](#)

[✓ Stmt](#) *Let $\widehat{K}^{[2]}$ be the tensor obtained by blocking two adjacent sites of the sector-coordinate tensor $\widehat{K}$. Then $\widehat{K}^{[2]}$ is a renormalization fixed point in the sense of Definition 23.1.27. More precisely, after the canonical identifications of one blocked site with two original sites and two blocked sites with four original sites, the required channels are $\widehat{\mathcal{S}}^2$ and $\widehat{\mathcal{T}}^2$.*




Figure 23.7: The superscripts in $\widehat{\mathcal{T}}^2$ and $\widehat{\mathcal{S}}^2$ denote successive overlapping applications. The dashed frames identify the two-site and four-site endpoints with one and two sites of the blocked tensor. This is the channel construction in [CPGSV16, Appendix C.2, lines 1821–1825].

Proof. ✓ Proof Decode one blocked index as a pair of physical indices and two blocked indices as four physical indices. The one-site and two-site closures of $\widehat{K}^{[2]}$ thereby become respectively $\widehat{K}_2(X)$ and $\widehat{K}_4(X)$. Transporting the two channels through these permutation identifications preserves complete positivity and trace, and the two closure identities give the required equations in Definition 23.1.27. $\square$

Theorem 23.5.5.42 (Two-site blocking of the original tensor is a renormalization fixed point).
Lean ✓ Stmt Let $K^{[2]}$ be the tensor obtained by blocking two adjacent sites of the original tensor K . Then $K^{[2]}$ is a renormalization fixed point in the sense of Definition 23.1.27. This is [CPGSV16, Theorem 4.9, (iv) $\Rightarrow$ (v)].

Proof. ✓ Proof Transport the refinement and coarse-graining channels for $\widehat{K}^{[2]}$ through the two-site and four-site tensor powers of the physical isometry, as in Figure 23.8. These unitary congruences preserve complete positivity and trace, while the corresponding closure identities carry the two fixed-point equations back to $K^{[2]}$. $\square$

$$U^{\otimes 4} \widehat{\mathcal{T}}^2 (U^\dagger)^{\otimes 2} = \mathcal{T}, \quad U^{\otimes 2} \widehat{\mathcal{S}}^2 (U^\dagger)^{\otimes 4} = \mathcal{S}$$

Figure 23.8: Physical-isometry transport of the blocked channels. The downward arrows conjugate the ket and bra indices by $U^{\otimes n}$ and $(U^\dagger)^{\otimes n}$; the upward arrows give the inverse conjugations. Hence the lower sector-coordinate channels induce the upper ordinary-coordinate channels. This combines Proposition C.7 with [CPGSV16, Appendix C.2, lines 1510–1563 and 1821–1825].

Tracing the ket and bra indices of a sector tensor gives the bond vector $|l_k\rangle$ or the bond functional $\langle r_k|$. Hence a neighboring pair gives

$$T_{k,h} = \langle r_k | l_h \rangle = \text{tr}(\eta_{k,h}),$$

as in [CPGSV16, Appendix C.2, lines 1474–1481].

23.5.6 Primitivity and rank-one trace matrices

Definition 23.5.6.1 (Explicit η -data from the sector tensors). Lean ✓ Stmt If every neighboring operator $\eta_{k,h}$ of Definition 23.5.2.1 is positive semidefinite, the family constitutes explicit neighboring η -data. The source obtains the positivity from the projected chain identity: writing $\tilde{\sigma} = U^{\dagger \otimes N} \sigma^{(N)}(K) U^{\otimes N}$ for the N -site operator in the sector basis and Q_k for the projector onto the k -th sector of the physical splitting, every cyclic sector assignment $k_1, \dots, k_N$ (with $k_{N+1} = k_1$) satisfies

$$0 \leq [Q_{k_1} \otimes \dots \otimes Q_{k_N}] \tilde{\sigma} [Q_{k_1} \otimes \dots \otimes Q_{k_N}] = \bigotimes_{n=1}^N \eta_{k_n, k_{n+1}},$$

and the sector tensors can be chosen so that each factor is positive semidefinite [CPGSV16, Appendix C.2, lines 1446–1450]. The all-length sector identity is Theorem 23.5.3.18. That algebraic identity does not by itself choose coherent positive representatives for all neighboring pairs, so positive semidefiniteness remains a hypothesis on the chosen representatives here. Theorem 23.5.3.39 provides the diagonal and paired positivity, and Theorem 23.5.3.42 a rescaled positive family under symmetric support.

Theorem 23.5.6.2 (Trace matrix of the sector-tensor η -data). [Lean](#) [✓ Stmt](#) Under the hypotheses of Definition 23.5.6.1, the trace matrix of the explicit η -data is the pairing of the closed sector tensors:

$$T_{k,h} = \text{tr}(\eta_{k,h}) = (r_k | l_h).$$

Proof. [✓ Proof](#) Immediate from Theorem 23.5.3.3 and the definition of the trace matrix. $\square$

Theorem 23.5.6.3 (Primitivity from returns of lengths two and three). [Lean](#) [✓ Stmt](#) Let T be a nonnegative irreducible matrix on a finite index set. If

$$(T^2)_{k,k} > 0 \quad \text{and} \quad (T^3)_{k,k} > 0$$

for every index k , then T is primitive.

Proof. [✓ Proof](#) Choose a positive path between every ordered pair of indices, and let L be the sum of the chosen path lengths. Every chosen length is at most L . For a chosen path of length m , write the difference $2L + 2 - m$ as $2a + 3b$ with $a, b \geq 0$. Appending these returns at the terminal index gives a positive path of common length $2L + 2$ between every ordered pair. $\square$

Definition 23.5.6.4 (Nonzero-weight sector trace matrix). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) For sector weights p_k , let $I_* = \{k : p_k \neq 0\}$. The one-site matrices and their spaces on I_* are

$$A_{k;x,y}, \quad A_S = \text{span}\{A_{k;x,y} : k \in S\}, \quad S \subseteq I_*.$$

The real trace matrix on this index set is

$$T_{k,h}^* = \text{Re tr}(\eta_{k,h}), \quad k, h \in I_*.$$

Only the trace matrix is restricted: the physical direct sum still contains every zero-weight summand.

Theorem 23.5.6.5 (Identification and spanning of active one-site matrices). [Lean](#) [Lean](#) [✓ Stmt](#) For every sector entry, the one-site matrix used in the directed-cut argument equals the virtual matrix used in the spanning argument. If all sector virtual matrices $\text{span } M_D(\mathbb{C})$ and the matrices in every zero-weight sector vanish, then

$$\text{span}\{A_{k;x,y} : k \in I_*\} = M_D(\mathbb{C}).$$

Theorem 23.5.6.6 (Complementary active-sector spaces). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that the active one-site matrices $\text{span } M_D(\mathbb{C})$ and each active sector contains a nonzero such matrix. For every $S \subseteq I_*$,

$$A_S + A_{I_* \setminus S} = M_D(\mathbb{C}),$$

and both spaces are nonzero when the corresponding sector sets are nonempty. If $\eta_{k,h} = 0$ for $k \in S$ and $h \in C$, then $A_S A_C = 0$.

Theorem 23.5.6.7 (Strong connectivity of the active neighboring support). [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that the active one-site matrices $\text{span } M_D(\mathbb{C})$ and each active sector contains a nonzero such matrix. Then for every $k, h \in I_*$ there is a directed walk

$$k = k_0 \longrightarrow k_1 \longrightarrow \cdots \longrightarrow k_m = h, \quad \eta_{k_j, k_{j+1}} \neq 0.$$

In particular, every active sector has an outgoing nonzero neighboring operator.

$$P_j \mathcal{K}_i P_{j'} = 0 \quad \text{unless } i = j = j'$$

Figure 23.9: The local-orthogonality contraction of [CPGSV16, Appendix C.2, lines 1457–1470]. After decomposing the sector trace matrix into primitive components, the source assumes that two components occur, obtains two locally orthogonal summands, and chooses orthogonal projectors P_1 and P_2 . The first two diagrams give the cross contractions $P_1 K P_2 = P_2 K P_1 = 0$, while the corresponding two-site contraction in the last diagram also vanishes. For $A_i = \text{span}\{\text{tr}(X P_i K P_i) : X\}$, these vanishings and injectivity imply $A_1 + A_2 = M_D(\mathbb{C})$ and $A_1 A_2 = 0$, a contradiction. Theorem 23.5.6.7 does not construct P_1 or P_2 : an arbitrary reachability cut $I_* = S \sqcup C$ supplies only $A_S A_C = 0$. The common final step is the same one-sided obstruction: two nonzero matrix spaces cannot sum to $M_D(\mathbb{C})$ while their ordered products vanish.

Theorem 23.5.6.8 (Positive entries of the active trace matrix). [Lean](#) [Lean](#) [✓ Stmt](#) *If every neighboring operator is positive semidefinite, then T^* is entrywise nonnegative and*

$$T_{k,h}^* > 0 \quad \Longleftrightarrow \quad \eta_{k,h} \neq 0.$$

Theorem 23.5.6.9 (Two-step return through every active sector). [Lean](#) [Lean](#) [✓ Stmt](#) *Suppose that the active one-site matrices span $M_D(\mathbb{C})$ and each active sector contains a nonzero such matrix. For every $k \in I_*$ there is an $h \in I_*$ such that*

$$\eta_{k,h} \neq 0, \quad \eta_{h,k} \neq 0.$$

If the neighboring operators are positive semidefinite, then

$$(T^{*2})_{k,k} > 0.$$

Theorem 23.5.6.10 (Irreducibility of the active trace matrix). [Lean](#) [✓ Stmt](#) *Suppose that the neighboring operators are positive semidefinite, the active one-site matrices span $M_D(\mathbb{C})$, and each active sector contains a nonzero such matrix. Then the nonnegative matrix T^* is irreducible.*

Theorem 23.5.6.11 (Three-step returns and a common positive power). [Lean](#) [Lean](#) [✓ Stmt](#) *Suppose that the neighboring operators are positive semidefinite, the active one-site matrices span $M_D(\mathbb{C})$, each active sector contains a nonzero such matrix, and every nonzero edge closes through a third active sector. Then $(T^{*3})_{k,k} > 0$ for every $k \in I_*$. Together with the length-two returns and irreducibility, this implies that T^* is primitive.*

Proof. [✓ Proof](#) Triangle closure supplies a positive closed walk of length three through each active sector. Combine these walks with the length-two returns and apply Theorem 23.5.6.3. $\square$

Theorem 23.5.6.12 (Primitivity of the active sector trace matrix). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *Let K be an injective tensor that generates MPDOs and satisfies SAL, and choose the positive neighboring operators of Theorem 23.5.3.41. Restrict the sector index to the positive-weight summands of the Hayashi decomposition. Then positive semidefiniteness makes every neighboring trace real and nonnegative, and the matrix*

$$T_{k,h}^* = \text{Re tr}(\eta_{k,h})$$

is primitive. Zero-weight summands remain in the physical direct sum but are not indices of T^ .*

Proof. ✓ Proof First use the zero-weight reparameterization to make every zero-weight sector isolated. Injectivity makes the nonzero-weight one-site matrices span $M_D(\mathbb{C})$, and every such sector contains a nonzero matrix. Hence Theorem 23.5.6.7 gives strong connectivity of the active neighboring support. Its proof is the one-sided directed-cut argument at [CPGSV16, Appendix C.2, lines 1463–1470].

Finally, nondegeneracy and cyclicity of the trace pairing give a closed support walk of length two through every active sector. Triangle closure gives one of length three. Strong connectivity together with these two coprime return lengths yields a common length at which every entry of the active trace-matrix power is positive. Thus T is primitive. $\square$

Theorem 23.5.6.13 (Zero-correlation-length relations on the primitive active trace matrix).

Lean Lean Lean ✓ Stmt Let K be injective and suppose that it satisfies SAL and source zero correlation length. Then there are a physical-sector factorization, weights $p_k \geq 0$ with $\sum_k p_k = 1$, and positive semidefinite neighboring operators $\eta_{k,h}$ such that the active trace matrix

$$T_{k,h}^* = \operatorname{Re} \operatorname{tr}(\eta_{k,h}), \quad k, h \in I_* := \{j : p_j \neq 0\},$$

is primitive. Moreover, there is a real number $\lambda > 0$ for which, upon setting $\widehat{T}^* := \lambda^{-1}T^*$,

$$(\widehat{T}^*)^2 = (\widehat{T}^*)^3.$$

For every positive integer N ,

$$\operatorname{tr}((\widehat{T}^*)^N) = \operatorname{tr}(\widehat{T}^*).$$

Thus the primitivity and the two normalized identities concern the same active, positively rephased trace matrix. No idempotence, rank-one, or semisimplicity conclusion is asserted.

Proof. ✓ Proof Choose the coherent sector phases from Theorem 23.5.6.12. For the resulting factorization, let L have columns $(\operatorname{tr}((l_h)_\beta))_\beta$ and let Q have rows $(\operatorname{tr}((r_k)_\alpha))_\alpha$, with both indices restricted to I_* . The left tensor vanishes when $p_k = 0$, so the rectangular product LQ is the full physical-trace transfer. Source zero correlation length therefore makes $\lambda^{-1}LQ$ idempotent for some $\lambda > 0$.

Positivity of the neighboring operators makes their traces real, and hence QL is the complexification of T^* . Associativity and cyclicity of trace give

$$(\lambda^{-1}QL)^2 = (\lambda^{-1}QL)^3, \quad \operatorname{tr}((\lambda^{-1}QL)^N) = \operatorname{tr}(\lambda^{-1}QL)$$

for every $N \geq 1$. Injectivity of the real-to-complex inclusion gives the two real matrix identities. $\square$

Theorem 23.5.6.14 (Positivity of bipartite partial traces). Lean Lean ✓ Stmt If ρ is positive semidefinite on a bipartite space $A \otimes B$, then both reduced operators $\operatorname{tr}_A \rho$ and $\operatorname{tr}_B \rho$ are positive semidefinite.

Proof. ✓ Proof Write each partial trace as the finite sum, over the traced index, of a principal submatrix of ρ . Principal submatrices of a positive semidefinite matrix are positive semidefinite, and finite sums of positive semidefinite matrices remain positive semidefinite. $\square$

Definition 23.5.6.15 (Hayashi–Markov extraction of explicit η -operators). Lean ✓ Stmt Every Hayashi decomposition witness h_η canonically produces an explicit neighboring η -family by Kronecker-multiplying the sector-reduced states on the neighboring bond spaces:

$$\eta_{k,h} := \operatorname{tr}_C(\rho_{b_2C}^{(k)}) \otimes \operatorname{tr}_A(\rho_{Ab_1}^{(h)}).$$

Positive semidefiniteness of each $\eta_{k,h}$ follows from the fact that partial traces preserve positivity and that positive semidefiniteness is closed under Kronecker products. This is the sector-reduced extraction; the remaining connection to [CPGSV16, Appendix C.2] is the identification of this family with the inverse-map construction in the original simple-MPDO tensor coordinates.

Theorem 23.5.6.16 (Trace matrix of the Hayashi–Markov η -operators). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The trace matrix induced by the extracted family is entrywise $T_{k,h} = 1$ in both its complex-valued form and its real-valued form. If there are m sectors, then $T = \mathbf{1}\mathbf{1}^\top$ and $\text{tr}(T) = m$.*

Proof. [✓ Proof](#) Each $\rho_{b_2C}^{(k)}$ is a density matrix, hence has unit trace; the partial trace tr_C preserves the trace. The analogous statement holds for $\text{tr}_A \rho_{Ab_1}^{(h)}$. Multiplicativity of the trace under Kronecker products gives $T_{k,h} = 1 \cdot 1 = 1$, and taking real parts gives the real-valued statement. Summing the m diagonal entries gives $\text{tr}(T) = m$. $\square$

Theorem 23.5.6.17 (Positive semidefiniteness of the Hayashi–Markov trace matrix). [Lean](#) [✓ Stmt](#) *The real trace matrix of the sector-reduced family is positive semidefinite.*

Proof. [✓ Proof](#) This trace matrix is the all-ones matrix $\mathbf{1}\mathbf{1}^\top$, hence is positive semidefinite. $\square$

Theorem 23.5.6.18 (Entrywise nonnegativity of the real trace matrix). [Lean](#) [✓ Stmt](#) *Positivity of the neighboring operators gives entrywise nonnegativity of the real trace matrix. Matrix-level positive semidefiniteness of T is a separate structural condition.*

Proof. [✓ Proof](#) The trace of a positive semidefinite neighboring operator is non-negative in the complex order. Taking real parts gives the stated entrywise nonnegativity. $\square$

Definition 23.5.6.19 (Auxiliary matrix predicates for the rank-one step). [Lean](#) [Lean](#) [✓ Stmt](#) The auxiliary predicates state two matrix-theoretic ingredients used in the rank-one step of [CPGSV16, Appendix C.2, Lemma C.5]: constant traces of all positive powers and existence of a rank-one factorization $T = ab^\top$.

The proposed implication from primitivity and constant trace powers to a rank-one factorization is not a valid general theorem. A concrete 3×3 counterexample appears in the archive. The corrected criterion proved here adds positive semidefiniteness and trace normalization, which provide the diagonalizability needed to turn the trace-power condition into a rank-one factorization.

Theorem 23.5.6.20 (Trace square of a Hermitian matrix). [Lean](#) [✓ Stmt](#) *Let T be a finite Hermitian matrix over a real or complex scalar field, with Hermitian eigenvalues λ_i . Then*

$$\text{tr}(T^2) = \sum_i \lambda_i^2.$$

Proof. [✓ Proof](#) The spectral theorem writes T as a unitary conjugate of the diagonal matrix of its Hermitian eigenvalues. Squaring commutes with this conjugation, and cyclicity of trace reduces the identity to the trace of the squared diagonal matrix. $\square$

Theorem 23.5.6.21 (PSD trace-power rank-one criterion). [Lean](#) [✓ Stmt](#) *Let T be a finite real square matrix. If T is positive semidefinite, normalized by $\text{tr}(T) = 1$, and has constant traces on all positive powers, then T has a rank-one factorization.*

Proof. [✓ Proof](#) The spectral theorem diagonalizes the positive semidefinite matrix with non-negative real eigenvalues. The trace gives that their sum is 1, while the second trace moment gives that the sum of their squares is also 1. Hence exactly one eigenvalue is equal to 1 and all others vanish, so the diagonal form is rank one. Unitary conjugation preserves the existence of an outer-product factorization. $\square$

Theorem 23.5.6.22 (Idempotent trace-one criterion). [Lean](#) [✓ Stmt](#) Let T be a finite real square matrix. If

$$T^2 = T, \quad \text{tr}(T) = 1,$$

then there are real vectors a, b such that $T = ab^\top$.

Proof. [✓ Proof](#) For an idempotent endomorphism f one has

$$\text{tr}(f) = \dim(\text{range}(f)).$$

Hence the range of T has dimension one. Choosing a nonzero vector a spanning it expresses every column of T as $b_j a$ for a scalar b_j , and therefore $T = ab^\top$. $\square$

Theorem 23.5.6.23 (Constant trace powers of an idempotent matrix). [Lean](#) [✓ Stmt](#) Let T be a finite real square matrix with $T^2 = T$. Then for every positive integer N ,

$$\text{tr}(T^N) = \text{tr}(T).$$

This recovers the trace identity displayed in the proof of [\[CPGSV16, Appendix C.2, Lemma C.5\]](#) from idempotence.

Proof. [✓ Proof](#) Induction on N : for $N \geq 1$, $T^{N+1} = T^N T = T T = T$, so every positive power of T equals T . $\square$

Theorem 23.5.6.24 (Positive entries in every row and column of a primitive matrix). [Lean](#) [Lean](#) [✓ Stmt](#) Let T be a primitive non-negative matrix. Then every row of T contains a strictly positive entry, and so does every column.

Proof. [✓ Proof](#) If row k of T vanishes, then for every $m \geq 1$,

$$(T^m)_{k,k} = \sum_j T_{k,j} (T^{m-1})_{j,k} = 0,$$

contradicting the strict entrywise positivity of some power of T . Likewise, if column h of T vanishes, then for every $m \geq 1$,

$$(T^m)_{h,h} = \sum_j (T^{m-1})_{h,j} T_{j,h} = 0,$$

again contradicting the strict positivity of some power of T . $\square$

Theorem 23.5.6.25 (Idempotence of the sector trace matrix). [Lean](#) [Lean](#) [✓ Stmt](#) Let V be a module over a commutative ring, let $l_1, \dots, l_n \in V$, let $r_1, \dots, r_n$ be linear functionals on V , and let $T_{k,h} = (r_k | l_h)$ be the pairing matrix; for the tensors of [\[CPGSV16, Appendix C.2\]](#) this is the sector trace matrix. Assume the identity

$$\sum_k |l_k)(r_k| = \sum_{k,h} T_{k,h} |l_k)(r_h|,$$

whose right-hand side is the square of its left-hand side, so that the operator $M = \sum_k |l_k)(r_k|$ satisfies $M^2 = M$. If the vectors $l_1, \dots, l_n$ are linearly independent, then

$$T^2 = T.$$

The same conclusion holds if instead the functionals $r_1, \dots, r_n$ are linearly independent.

Proof. ✓ **Proof** Evaluating $M^2v = Mv$ at an arbitrary $v \in V$ gives

$$\sum_k ((r_k|Mv) - (r_k|v)) l_k = 0,$$

so linear independence of the l_k yields $(r_k|Mv) = (r_k|v)$ for all k and v . Taking $v = l_h$ and expanding $ML_h = \sum_j T_{j,h} l_j$ turns the left-hand side into $\sum_j T_{k,j} T_{j,h} = (T^2)_{k,h}$ and the right-hand side into $T_{k,h}$. For linearly independent functionals, apply the same argument in the dual space, with the r_k as vectors and evaluation at l_k as functionals. The dual pairing matrix therefore satisfies

$$(T^\top)^2 = T^\top.$$

Taking transposes gives

$$T^2 = T.$$

□

Equivalently, the zero-correlation-length identity is the algebraic equality

$$\sum_k |l_k)(r_k| = \sum_{k,h} T_{k,h} |l_k)(r_h|, \quad T_{k,h} = (r_k|l_h),$$

which is the displayed identity of [CPGSV16, Appendix C.2, lines 1490–1493].

Theorem 23.5.6.26 (Unconditional constant trace powers of the pairing matrix). Lean Lean

Lean Lean ✓ **Stmt** Let V be a real vector space, let $l_1, \dots, l_n \in V$, let $r_1, \dots, r_n$ be real linear functionals on V , and let $T_{k,h} = (r_k|l_h)$ be the pairing matrix, as in Theorem 23.5.6.25. Assuming only the identity $M^2 = M$ for the operator $M = \sum_k |l_k)(r_k|$, without assuming linear independence of $l_1, \dots, l_n$ or of $r_1, \dots, r_n$,

$$T^2 = T^3,$$

so $T^N = T^2$ for every $N \geq 2$, and for every positive integer N ,

$$\text{tr}(T^N) = \text{tr}(T).$$

Proof. ✓ **Proof** Multiplying the identity $M^2 = M$ on the left by the functional map and on the right by the coefficient map turns the two sides directly into T^2 and T^3 , with no independence hypothesis; induction then gives $T^N = T^2$ for every $N \geq 2$. For the trace identity, restrict the operator M to the finite-dimensional span of $l_1, \dots, l_n$: there it is an idempotent endomorphism of a finite free module, and cyclicity of the trace of a composition between that span and the coefficient space identifies $\text{tr}(T)$ with the trace of the restriction and $\text{tr}(T^2)$ with the trace of its square, which agree since the restriction is idempotent. □

Theorem 23.5.6.27 (Rank-one factorization of the sector trace matrix). Lean ✓ **Stmt** In the setting of Theorem 23.5.6.25 with real scalars and linearly independent vectors $l_1, \dots, l_n$, if in addition $\text{tr}(T) = 1$, then there are real vectors a, b such that $T = ab^\top$. This is the factorization $T_{k,h} = a_k b_h$ obtained in the proof of [CPGSV16, Appendix C.2, Lemma C.5] for the sector trace matrix, derived here from the displayed zero-correlation-length identity together with the linear independence of the l_k , instead of the Perron–Frobenius trace-power argument of the source.

Proof. ✓ **Proof** Theorem 23.5.6.25 gives $T^2 = T$, and the idempotent trace-one criterion of Theorem 23.5.6.22 gives the factorization $T = ab^\top$. □

Theorem 23.5.6.28 (PSD-corrected rank-one T criterion). [Lean](#) [✓ Stmt](#) Let T be a finite real square matrix. Assume that T is primitive, positive semidefinite, normalized by $\text{tr}(T) = 1$, and has constant traces on all positive powers. Then

$$T = ab^\top$$

for some vectors a and b , with $a \cdot b = 1$.

Proof. [✓ Proof](#) The positive-semidefinite matrix criterion diagonalizes T , uses the first two trace moments to show that exactly one eigenvalue is nonzero, and hence gives the factorization. Taking traces and using $\text{tr}(ab^\top) = a \cdot b$ gives the normalization $a \cdot b = 1$. $\square$

Definition 23.5.6.29 (Sector tensors paired into the sector trace matrix). [Lean](#) [✓ Stmt](#) For a real $n \times n$ matrix T and a real vector space V , this structure consists of vectors $|l_0), \dots, |l_{n-1}) \in V$ and linear functionals $(r_0|, \dots, (r_{n-1}|$ on V satisfying the pairing identity

$$T_{k,h} = (r_k|l_h)$$

of [\[CPGSV16, Appendix C.2\]](#) together with the zero-correlation-length identity

$$\sum_k |l_k)(r_k| = \sum_{k,h} T_{k,h} |l_k)(r_h|$$

displayed in the proof of [\[CPGSV16, Lemma C.5\]](#). The vectors are the closed sector tensors obtained there from the sector splitting of an injective simple tensor.

Theorem 23.5.6.30 (Idempotence of the sector trace matrix from the zero-correlation-length identity). [Lean](#) [Lean](#) [✓ Stmt](#) For sector tensors paired into T as in Definition 23.5.6.29, the operator $M = \sum_k |l_k)(r_k|$ satisfies $M^2 = M$. If in addition the vectors $|l_0), \dots, |l_{n-1})$ are linearly independent, then $T^2 = T$.

Proof. [✓ Proof](#) Substituting $T_{k,h} = (r_k|l_h)$ into the right-hand side of the zero-correlation-length identity gives, for every $v \in V$,

$$\sum_{k,h} T_{k,h} (r_h|v) |l_k) = \sum_{k,h} (r_k|l_h) (r_h|v) |l_k) = \sum_k (r_k|Mv) |l_k) = M^2 v,$$

so $M^2 = M$. Linear independence of the $|l_k)$ then gives $T^2 = T$ by Theorem 23.5.6.25. $\square$

Theorem 23.5.6.31 (Unconditional constant trace powers of the sector trace matrix). [Lean](#) [Lean](#) [✓ Stmt](#) For sector tensors paired into T as in Definition 23.5.6.29, without any independence hypothesis on the closed sector tensors or the functionals,

$$T^2 = T^3,$$

and for every positive integer N ,

$$\text{tr}(T^N) = \text{tr}(T).$$

Proof. [✓ Proof](#) The operator identity $M^2 = M$ established in the proof of Theorem 23.5.6.30 gives $T^2 = T^3$ and the constant trace powers by Theorem 23.5.6.26, with no independence hypothesis. $\square$

Theorem 23.5.6.32 (Nonvanishing and independence of the sector tensors). [Lean](#) [Lean](#) [✓ Stmt](#) Suppose the sector trace matrix T of Definition 23.5.6.29 is primitive. Then every closed sector tensor satisfies $|l_k) \neq 0$. If moreover each $|l_k)$ lies in its own member of an independent family of subspaces of V , then the vectors $|l_0), \dots, |l_{n-1})$ are linearly independent.

Proof. [✓ Proof](#) By Theorem 23.5.6.24, some entry of the k -th column of T is positive, and that entry is the pairing $(r_j|l_k)$, so $|l_k) \neq 0$. Nonzero vectors chosen from an independent family of subspaces, one from each member, are linearly independent. $\square$

Theorem 23.5.6.33 (Rank one from sector-pairing idempotence under linear independence).

[Lean](#) [Lean](#) [✓ Stmt](#) Let T be a sector trace matrix with $\text{tr}(T) = 1$, paired from sector tensors satisfying the zero-correlation-length identity of Definition 23.5.6.29. If the closed sector tensors $|l_0), \dots, |l_{n-1})$ are linearly independent, then

$$T_{k,h} = a_k b_h, \quad \sum_k a_k b_k = 1$$

for some real numbers a_k, b_h . The same conclusion holds if, instead of the independence, T is primitive and the closed sector tensors lie in distinct members of an independent family of subspaces. In contrast with the positive-semidefinite criterion of Theorem 23.5.6.28, the constant trace powers are derived rather than assumed.

Proof. [✓ Proof](#) Theorem 23.5.6.30 gives $T^2 = T$, and Theorem 23.5.6.31 gives the constant trace powers unconditionally. The idempotent trace-one criterion of Theorem 23.5.6.22 yields the factorization $T = ab^\top$, and taking traces gives $\sum_k a_k b_k = \text{tr}(T) = 1$. When the independence is not assumed directly, Theorem 23.5.6.32 derives it from primitivity once each sector tensor lies in its own member of an independent family of subspaces. $\square$

23.6 Commuting form and GSNNCH

Definition 23.6.1 (Commuting-form data for an MPDO). [Lean](#) [✓ Stmt](#) For a fixed chain length $N \geq 2$, this consists of a positive semidefinite two-site operator $B \geq 0$ whose translated copies commute pairwise on the periodic N -site chain:

$$[B_{i,i+1}, B_{j,j+1}] = 0.$$

Definition 23.6.2 (Commuting form). [Lean](#) [✓ Stmt](#) An MPO tensor M has the commuting-form property if for every $N \geq 2$ there is a two-site operator B as in Definition 23.6.1 and a constant $c > 0$ such that

$$\rho^{(N)}(M) = c \prod_{i=0}^{N-1} B_{i,i+1}.$$

Definition 23.6.3 (GSNNCH). [Lean](#) [✓ Stmt](#) An MPO tensor M is a Gibbs state of a nearest-neighbor commuting Hamiltonian if for every $N \geq 2$ its finite-chain operator admits the equivalent positive-operator presentation

$$\rho^{(N)}(M) = c \prod_{i=0}^{N-1} B_{i,i+1}, \quad c > 0, \quad B \geq 0, \quad [B_{i,i+1}, B_{j,j+1}] = 0.$$

The projector-limit convention in [CPGSV16] identifies this with the exponential Gibbs form.

Theorem 23.6.4 (GSNNCH iff commuting form). [Lean](#) [✓ Stmt](#) For an MPO tensor, the GSNNCH condition is equivalent to the existence of commuting-form data on every finite chain.

Proof. [✓ Proof](#) Both directions unfold to the same identity $\rho^{(N)}(M) = c \prod_{i=0}^{N-1} B_{i,i+1}$ with $c > 0$: a commuting-form witness for M at length N together with its constant c is exactly a GSNNCH witness for M at length N with that same constant, and conversely. $\square$

Definition 23.6.5 (GSNNCH with doubled-index transfer idempotence). [Lean](#) [✓ Stmt](#) This is the conjunction of the GSNNCH condition with the doubled-index transfer condition

$$\mathcal{E}_M \circ \mathcal{E}_M = \mathcal{E}_M.$$

It is the doubled-index analogue of the ZCL clause in item (iii) of the simple-MPDO equivalence in [CPGSV16]; the source physical-trace condition is Definition 23.2.2.

Theorem 23.6.6 (GSNNCH with doubled-index idempotence iff commuting form with it).

[Lean](#) [✓ Stmt](#) *The GSNCH branch with doubled-index transfer idempotence is equivalent to the conjunction of the commuting-form property with that same idempotence condition.*

Proof. ✔ Proof Unfold the definition of the GSNNCH branch with doubled-index transfer idempotence and apply Theorem 23.6.4. □

Definition 23.6.7 (Translation-invariant two-site bond). [Lean](#) [✓ Stmt](#) This records a single positive semidefinite two-site bond $B \geq 0$ whose translated copies commute pairwise on the periodic N -site chain at every chain length $N \geq 2$:

$$[B_{i,i+1}, B_{j,j+1}] = 0.$$

Definition 23.6.8 (η -local commuting-form data). [Lean](#) [StmT](#) This defines the eta-local form of [CPGSV16, Appendix C.2]: a two-site bond B as in Definition 23.6.7 together with, for every chain length $N \geq 2$, a constant $c > 0$ such that

$$\rho^{(N)}(M) = c \prod_{i=0}^{N-1} B_{i,i+1}.$$

Definition 23.6.9 (Finite-chain bond form). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A translation-invariant positive two-site bond determines finite-chain commuting-form data at every length $N \geq 2$.

Theorem 23.6.10 (η -local structure carries commuting form). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#)
[Lean](#) [✓ Stmt](#) *The η -local structure itself determines a commuting-form witness for every chain length $N \geq 2$: writing B for its bond, the translated copies commute,*

$$[B_{i,i+1}, B_{j,i+1}] = 0,$$

and there exists $c > 0$ with

$$\rho^{(N)}(M) = c \prod_{i=0}^{N-1} B_{i,i+1}.$$

This is the implication from η -local structure to commuting form in [CPGSV16, Proposition C.8].

Proof. ✓ Proof At length N , apply the commuting two-site bond and realization identity provided by the η -local structure. $\square$

Theorem 23.6.11 (η -local structure gives GSNCH). [Lean](#) [Lean](#) [✓ Stmt](#) *The same η -local structure gives a GSNCH witness on every finite periodic chain.*

Proof. ✓ Proof Evaluate the local structure at each chain length $N \geq 2$.

Theorem 23.6.12 (η -local structure with doubled-index idempotence). [Lean](#) [Lean](#) [✓ Stmt](#)
An η -local structure together with the doubled-index transfer condition gives the corresponding GSNNCH branch.

Proof. ✓ Proof Use the commuting-form witness carried by the η -local structure and combine it with the doubled-index transfer hypothesis. $\square$

Remark 23.6.13 (Positive physical-sector factorization). Suppose that K satisfies $\text{SAL}(K)$ and admits a physical-sector factorization whose neighboring operators $\eta_{k,h}$ are simultaneously positive semidefinite. The associated two-site bond is

$$B = \sum_{k,h} \text{Id}_{H_L^{(k)}} \otimes \eta_{k,h} \otimes \text{Id}_{H_R^{(h)}}, \quad B \geq 0, \quad (23.1)$$

and, for every $N \geq 2$,

$$\sigma^{(N)}(K) = c_N \prod_{n=1}^N B_{n,n+1}, \quad c_N > 0, \quad [B_{i,i+1}, B_{j,j+1}] = 0. \quad (23.2)$$

Under this hypothesis, Theorem 23.5.2.47 gives the η -local structure and proves (23.1)–(23.2) with $c_N = 1$. Theorem 23.6.10 then gives the commuting-form conclusion. Adjacent translates commute on every periodic chain of length at least three by Theorem 23.5.2.40, and the crossed two-site case is Theorem 23.5.2.41. Disjoint translates commute by Theorem 23.5.2.42. These cases give pairwise commutativity at every length $N \geq 2$ by Theorem 23.5.2.43. Given the physical-sector factorization, the all-length product identity is Theorem 23.5.2.44. The doubled-index transfer condition enters only in the subsequent RFP and GSNNCH branch conclusions.

Chapter 24

Mixed States: General Case and Algebraic Structure

This chapter continues the mixed-state renormalization analysis of Chapter 23 with the general case of [CPGSV16, Section 4]. Two-site blocking first reduces the tensor to the local form used by the source. The subsequent algebraic characterization identifies the diagonal χ -matrices governing the trace-power formula and culminates in the fusion-isometry description.

24.1 Local structure and two-site blocking

Definition 24.1.1 (Local simple-MPDO structure). [Lean](#) [✓ Stmt](#) This structure records the local ingredients of [CPGSV16, Appendix C.2]: a normalized three-site reduced state with equality in strong subadditivity, the resulting Markov decomposition, and a primitive matrix T satisfying

$$\mathrm{tr}(T) = 1, \quad \mathrm{tr}(T^m) = \mathrm{tr}(T)$$

for every positive integer m , together with a rank-one factorization

$$T = ab^\top, \quad a \cdot b = 1.$$

Theorem 24.1.2 (Local simple-MPDO structure from corrected rank-one criteria). [Lean](#) [✓ Stmt](#) There are two sufficient forms of the local conclusion associated with [CPGSV16, Lemmas C.2, C.4, and C.5]. In the first, T is positive semidefinite, $\mathrm{tr}(T) = 1$, and $\mathrm{tr}(T^m) = 1$ for every positive integer m ; the spectral theorem then gives

$$T = ab^\top, \quad a \cdot b = 1.$$

In the second, T is the sector trace matrix of linearly independent sector tensors satisfying the zero-correlation-length pairing identity. This identity implies $T^2 = T$, so trace one gives the same rank-one factorization. In this second form the constancy of the positive-power traces is a conclusion rather than an assumption.

The physical supports are locally orthogonal:

$$\begin{array}{c} | \\ \bullet \\ \text{---} \mathcal{K}_y \\ \bullet \\ \text{---} \mathcal{K}_x \\ | \end{array} = 0 \quad (x \neq y)$$

In contraction notation, local orthogonality is $\mathcal{K}_y^* \mathcal{K}_x = 0$ whenever $x \neq y$. This is [CPGSV16, Theorem 4.9(iv), lines 863–868].

Expanding $\sum_j P_j = \mathbb{1}$ on both physical indices and using the local orthogonality relations leaves only the diagonal term:

$$\mathcal{K}_i = \sum_{j,j'} P_j \mathcal{K}_i P_{j'} = P_i \mathcal{K}_i = \mathcal{K}_i P_i$$

Equivalently, each local tensor is supported on its own sector: $\mathcal{K}_i = P_i \mathcal{K}_i = \mathcal{K}_i P_i$. This is the calculation in [CPGSV16, Appendix C.2, lines 1745–1751].

The two-site block is the virtual contraction of two copies of K :

The two ket indices and the two bra indices are grouped into the blocked physical indices, as in [CPGSV16, Theorem 4.9, lines 851–856].

Definition 24.1.3 (Positive-length physical blocking of an MPO tensor). [Lean](#) [Lean](#) [✓ Stmt](#) For a positive integer L , group L adjacent physical sites and denote the resulting local tensor by $M^{[L]}$. It has a length- L ket word $I = (i_1, \dots, i_L)$ and a length- L bra word $J = (j_1, \dots, j_L)$ as its physical indices, and

$$(M^{[L]})^{I,J} = M^{i_1 j_1} \dots M^{i_L j_L}.$$

Thus the ket and bra words are grouped separately. This is the MPO version of the physical blocking used for the basis of normal tensors in [CPGSV16, lines 317–345]. It is a local tensor, not the closed-chain operator $O_L(M)$ of [CPGSV16, lines 962–967].

The algebraic definition also assigns the identity matrix to the empty product at $L = 0$. This total convention has no physical role here; every blocking result below assumes $L > 0$.

Theorem 24.1.4 (Doubled-index form of positive MPO blocking). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let $L > 0$. Pairing the ket and bra letters site by site gives the canonical identification

$$\{0, \dots, d^L - 1\}^2 \simeq \{0, \dots, d^2 - 1\}^L.$$

Under this identification, the doubled-index MPS tensor of $M^{[L]}$ is the physical reindexing of the L -blocked doubled-index MPS tensor of M . In particular, one is injective if and only if the other is.

Proof. [✓ Proof](#) Decode the blocked ket and bra words, pair their letters at each site, and compare the two ordered matrix products. Physical reindexing by a bijection preserves the span of the tensor matrices. $\square$

Theorem 24.1.5 (Positive blocking commutes with the MPO product). [Lean](#) [✓ Stmt](#) For $L > 0$ and two MPO tensors M, N with the same physical dimension, let juxtaposition denote the MPO tensor product obtained by contracting the intermediate physical index. Then

$$(MN)^{[L]} = M^{[L]} N^{[L]}.$$

Proof. [✓ Proof](#) Expand the product tensor and sum over its length- L intermediate physical word. Reindexing this word by one blocked physical index gives the product of the two blocked tensors. $\square$

Definition 24.1.6 (Two-site blocking of an MPO tensor). [Lean](#) [✓ Stmt](#) For an MPO tensor K , its two-site blocking $K^{[2]}$ has physical indices (i_0, i_1) and (j_0, j_1) and matrices

$$(K^{[2]})^{(i_0, i_1), (j_0, j_1)} = K^{i_0 j_0} K^{i_1 j_1}.$$

This is the blocking used in [\[CPGSV16, Theorem 4.9, lines 851–856\]](#).

Definition 24.1.7 (Four-site physical closure). [Lean](#) [Lean](#) [✓ Stmt](#) For a virtual matrix X , the four-site physical closure of K is the operator $K_4(X)$ whose matrix coefficients are

$$K_4(X)_{(i_0, (i_1, (i_2, i_3))), (j_0, (j_1, (j_2, j_3)))} = \text{tr}(K^{i_0 j_0} K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} X).$$

Theorem 24.1.8 (General and right-associated four-site closures). [Lean](#) [✓ Stmt](#) Under the canonical right-associated identification of four-site configurations with quadruples of physical indices,

$$K_4(X)_{(i_0, (i_1, (i_2, i_3))), (j_0, (j_1, (j_2, j_3)))} = \text{tr}(K^{i_0 j_0} K^{i_1 j_1} K^{i_2 j_2} K^{i_3 j_3} X).$$

Proof. [✓ Proof](#) Expand the general four-site closure in the right-associated coordinates. □

Theorem 24.1.9 (One-site closure of the two-site blocking). [Lean](#) [✓ Stmt](#) After identifying one blocked physical index with two original indices,

$$(K^{[2]})_1(X) = K_2(X).$$

Proof. [✓ Proof](#) Expand one blocked tensor matrix and decode its two physical indices. □

Theorem 24.1.10 (Two-site closure of the two-site blocking). [Lean](#) [✓ Stmt](#) After identifying two blocked physical indices with four original indices, one has

$$(K^{[2]})_2(X) = K_4(X).$$

Proof. [✓ Proof](#) Expand the two blocked tensor matrices and reassociate their product. □

24.2 Algebra structure

Definition 24.2.1 (Algebra-structure data for an MPDO). [Lean](#) [✓ Stmt](#) A family of support $*$ -algebras $\{\mathcal{A}_n\}_{n \geq 0}$, one at every blocking size, equipped with blocking maps $m_n : \mathcal{A}_n \otimes \mathcal{A}_n \rightarrow \mathcal{A}_{2n}$ and inclusion maps $\iota_n : \mathcal{A}_n \rightarrow \mathcal{A}_{n+1}$ realized, inside the ambient bond-space matrix algebra, by

$$m_n(x, y) = xy, \quad \iota_n(x) = x.$$

Definition 24.2.2 (Algebra-structure RFP predicate). [Lean](#) [✓ Stmt](#) An MPO tensor M satisfies the *algebra-structure formulation* used here when there exist algebra-structure data such that, for every positive blocking size n ,

$$\mathcal{A}_n = \text{Fix}(E_n^\dagger),$$

where E_n is the blocked transfer map. This is a nontrivial algebraic condition, but it is still weaker than the full coefficient formulation of [\[CPGSV16, Theorem 4.14\(ii\)\]](#); that formulation also includes the coefficient family $c_{\alpha, \beta, \gamma}^{(L)}$.

Theorem 24.2.3 (RFP yields a stationary algebra tower under a faithful fixed point). [Lean](#)

[✓ Stmt](#) Assume the doubled tensor is trace-preserving and the transfer map admits a positive-definite fixed point. If $E_1^2 = E_1$, then there is a stationary family of support algebras

$$\mathcal{A}_n = \text{Fix}(E_1^\dagger),$$

with multiplication $m_n(x, y) = xy$ and inclusion $\iota_n(x) = x$.

Proof. [✓ Proof](#) Under the trace-preserving normalization and the faithful fixed point, Wolf's fixed-point-algebra theorem gives the $*$ -subalgebra $\text{Fix}(E_1^\dagger) \subseteq M_D(\mathbb{C})$. Since E_1 is idempotent, $E_n = E_1^n = E_1$ for every $n \geq 1$, so $\text{Fix}(E_n^\dagger) = \text{Fix}(E_1^\dagger)$ at every positive blocking size, giving the stationary tower $\mathcal{A}_n = \text{Fix}(E_1^\dagger)$. $\square$

The two closure maps have the same local form:

$$M_\alpha(X) = \begin{array}{c} X \quad M_\alpha \\ \text{tr} \end{array} \quad A_\alpha(X) = \begin{array}{c} X \quad A_\alpha \\ \text{tr} \end{array}$$

In each cell, X is inserted on the virtual bond and the virtual indices are closed by tr ; the physical legs remain open. Their spans are the fixed-point algebras compared in [CPGSV16, Appendix C.4, lines 1974–1980].

Theorem 24.2.4 (Fusion yields a stationary algebra tower under a faithful fixed point). [Lean](#)

[✓ Stmt](#) Assume the doubled tensor is trace-preserving and the transfer map admits a positive-definite fixed point. If the transfer-map fusion criterion holds, then the MPO satisfies the algebra-structure formulation of the renormalization fixed-point condition.

Proof. [✓ Proof](#) Combine Theorem 24.4.3 with Theorem 24.2.3. $\square$

Theorem 24.2.5 (Stabilized blocked adjoint fixed points yield algebra data). [Lean](#) [✓ Stmt](#)

Assume the doubled tensor is trace-preserving and the transfer map admits a positive-definite fixed point. If, for every positive blocked size n ,

$$\text{Fix}(E_n^\dagger) = \text{Fix}(E_1^\dagger),$$

then the algebra-structure formulation holds.

Proof. [✓ Proof](#) Take the constant tower $\mathcal{A}_n := \text{Fix}(E_1^\dagger)$ for every n . The stabilization hypothesis $\text{Fix}(E_n^\dagger) = \text{Fix}(E_1^\dagger)$ then gives $\mathcal{A}_n = \text{Fix}(E_n^\dagger)$ at every positive blocking size n , which is compatibility with M . $\square$

This does not give the full converse direction of [CPGSV16, Theorem 4.14]: the compatibility condition above only sees stabilized adjoint fixed-point algebras, not the coefficient/BNT data. In particular, [CPGSV16, Theorem 4.14(ii)] also requires the special diagonal matrices $\chi_{\alpha, \beta, \gamma}$. The coefficient data attached to the algebra tower are: a chosen basis of each support algebra, coordinate maps into a finite scalar family, and the corresponding multiplication / inclusion coefficient arrays.

Definition 24.2.6 (Blocked coefficients for the algebra tower). [Lean](#) [✓ Stmt](#) For each blocked size n , fix a basis $\{b_i\}_{i \in I_n}$ of $\mathcal{A}_n$ over a finite index set I_n . The *blocked coefficients* of $x \in \mathcal{A}_n$ are the coordinates $(x_i)_{i \in I_n} \in \mathbb{C}^{I_n}$ with

$$x = \sum_{i \in I_n} x_i b_i.$$

Theorem 24.2.7 (Reconstruction from blocked coefficients). [Lean](#) [✓ Stmt](#) *Reconstructing a coefficient family gives the corresponding basis expansion $\sum_i a_i b_i$ in $\mathcal{A}_n$.*

Proof. [✓ Proof](#) This is exactly the finite-basis reconstruction formula for the chosen basis of $\mathcal{A}_n$. $\square$

Definition 24.2.8 (Blocked multiplication coefficients). [Lean](#) [✓ Stmt](#) For $i, j \in I_n$, the *blocked multiplication coefficients* are the blocked coefficients $(c_{i,j,k}^{(n)})_{k \in I_{2n}}$ of $m_n(b_i, b_j) \in \mathcal{A}_{2n}$:

$$m_n(b_i, b_j) = \sum_{k \in I_{2n}} c_{i,j,k}^{(n)} b_k.$$

Theorem 24.2.9 (Blocked multiplication reconstructs from its coefficients). [Lean](#) [✓ Stmt](#) *In the ambient bond-space matrix algebra,*

$$b_i b_j = \sum_{k \in I_{2n}} c_{i,j,k}^{(n)} b_k.$$

Proof. [✓ Proof](#) Apply the reconstruction formula of Theorem 24.2.7 to the element $m_n(b_i, b_j) \in \mathcal{A}_{2n}$. $\square$

Definition 24.2.10 (Blocked inclusion coefficients). [Lean](#) [✓ Stmt](#) For $i \in I_n$, the *blocked inclusion coefficients* are the blocked coefficients $(d_{ij})_{j \in I_{n+1}}$ of $\iota_n(b_i) \in \mathcal{A}_{n+1}$:

$$\iota_n(b_i) = \sum_{j \in I_{n+1}} d_{ij} b_j.$$

Theorem 24.2.11 (Blocked inclusion reconstructs from its coefficients). [Lean](#) [✓ Stmt](#) *Reconstructing the inclusion coefficients recovers the image of b_i under ι_n :*

$$\sum_{j \in I_{n+1}} d_{ij} b_j = \iota_n(b_i).$$

Proof. [✓ Proof](#) Coefficient extraction and reconstruction are mutually inverse coordinate maps for the chosen basis of $\mathcal{A}_{n+1}$, so extracting the coefficients of $\iota_n(b_i)$ and reconstructing returns the same element. $\square$

Theorem 24.2.12 (Compatible adjoint fixed points reconstruct from blocked coefficients). [Lean](#) [✓ Stmt](#) *If the algebra tower is compatible with an MPO tensor M and $E_n^\dagger X = X$ for a positive blocking size n , then $X \in \mathcal{A}_n$ and, writing $(X_i)_{i \in I_n}$ for its blocked coefficients,*

$$X = \sum_{i \in I_n} X_i b_i.$$

Proof. [✓ Proof](#) Compatibility identifies $\mathcal{A}_n$ with the adjoint fixed-point algebra of E_n , so the fixed point first becomes an element of $\mathcal{A}_n$ and then Theorem 24.2.7 reconstructs it. $\square$

Theorem 24.2.13 (Adjoint fixed points descend through the algebra tower). [Lean](#) [✓ Stmt](#) *If an MPO tensor satisfies the algebra-structure formulation, then for every positive blocking size n and every X ,*

$$E_n^\dagger X = X \implies E_1^\dagger X = X.$$

Proof. ✓ Proof Compatibility at size n places X in $\mathcal{A}_n$. The inclusion map ι_n lifts X into $\mathcal{A}_{n+1}$, and since ι_n is just an inclusion it does not change the underlying operator, so we continue to denote the lifted element by X . Thus compatibility at size $n+1$ yields $E_{n+1}^\dagger X = X$. Rewriting $E_{n+1} = E_n \circ E_1$ and applying $(A \circ B)^\dagger = B^\dagger \circ A^\dagger$ together with the hypothesis $E_n^\dagger X = X$ extracts $E_1^\dagger X = X$. $\square$

Lemma 24.2.14 (Reverse inclusion of adjoint fixed points). Lean ✓ Stmt For every n ,

$$E_1^\dagger X = X \implies E_n^\dagger X = X.$$

Proof. ✓ Proof Induction on n using $E_{n+1} = E_n \circ E_1$ and $(A \circ B)^\dagger = B^\dagger \circ A^\dagger$; the base case is $E_0 = \text{id}$. This step does not require the algebra-structure data. $\square$

Lemma 24.2.15 (Blocked powers of adjoint eigenvectors). Lean ✓ Stmt If $E = E_1$ and $E^\dagger X = \lambda X$, then $E_n^\dagger X = \lambda^n X$ for every n .

Proof. ✓ Proof The case $n = 0$ is $E_0 = \text{id}$. For the induction step, $E_{n+1} = E_n \circ E_1$, hence

$$E_{n+1}^\dagger X = E_1^\dagger E_n^\dagger X = E_1^\dagger (\lambda^n X) = \lambda^n E_1^\dagger X = \lambda^{n+1} X.$$

$\square$

Theorem 24.2.16 (Finite-order adjoint eigenvalues for algebra towers). Lean ✓ Stmt Assume the algebra-structure formulation holds. If $X \neq 0$, $E_1^\dagger X = \lambda X$, and $\lambda^n = 1$ for some $n > 0$, then $\lambda = 1$.

Proof. ✓ Proof By Lemma 24.2.15, $E_n^\dagger X = \lambda^n X = X$. Theorem 24.2.13 gives $E_1^\dagger X = X$. Thus $\lambda X = X$, so $(\lambda - 1)X = 0$. Since $X \neq 0$, one obtains $\lambda = 1$. $\square$

Theorem 24.2.17 (Adjoint fixed-point equality for algebra towers). Lean ✓ Stmt If an MPO tensor satisfies the algebra-structure formulation, then for every positive blocking size n ,

$$\text{Fix}(E_n^\dagger) = \text{Fix}(E_1^\dagger).$$

Proof. ✓ Proof Combine Theorem 24.2.13 with Lemma 24.2.14. $\square$

Theorem 24.2.18 (Algebra data give the stationary fixed-point tower). Lean ✓ Stmt Under the trace-preserving normalization and a positive-definite fixed point, if M satisfies the algebra-structure formulation, then the stationary tower $\mathcal{A}_n = \text{Fix}(E_1^\dagger)$ is itself compatible with M : for every positive blocking size n ,

$$\text{Fix}(E_1^\dagger) = \text{Fix}(E_n^\dagger).$$

Theorem 24.2.19 (The support-algebra condition does not imply fusion). Lean ✓ Stmt There exists an MPO tensor of physical and bond dimension two such that the doubled tensor is trace-preserving, the transfer map has a positive-definite fixed point, and the support-algebra condition holds, but the fusion condition does not hold.

Proof. ✓ Proof Take the phase-flip channel with Kraus operators $K_0 = \frac{3}{5}I$ and $K_1 = \frac{4}{5}Z$. Its transfer map satisfies

$$E(X)_{ij} = \begin{cases} X_{ij}, & i = j, \\ -\frac{7}{25}X_{ij}, & i \neq j. \end{cases}$$

Hence, for every $n > 0$ and $i \neq j$,

$$E^n(X)_{ij} = \left(-\frac{7}{25}\right)^n X_{ij}.$$

Since $(-7/25)^n \neq 1$ for every $n > 0$ and E is self-adjoint,

$$\text{Fix}((E^n)^\dagger) = \text{Fix}(E^\dagger).$$

Thus the support-algebra condition holds. On the other hand,

$$E^2(X)_{01} = \frac{49}{625} X_{01} \neq -\frac{7}{25} X_{01} = E(X)_{01}.$$

Therefore $E^2 \neq E$, so the fusion condition fails. $\square$

Remark 24.2.20 (Why algebra data do not imply fusion data). Write $E = E_1$ for the one-site transfer map. The present algebra predicate gives, for every $n > 0$,

$$\text{Fix}((E^n)^\dagger) = \text{Fix}(E^\dagger).$$

Hence a vector X with $E^\dagger X = \lambda X$ and $\lambda^n = 1$ lies in $\text{Fix}(E^\dagger)$, so $(\lambda - 1)X = 0$ and $\lambda = 1$ by Theorem 24.2.16. It does not imply $E^2 = E$, as Theorem 24.2.19 shows. More generally, if Π_{diag} is the projection onto diagonal matrices and $0 < \varepsilon < 1$, set

$$E(X) = \varepsilon X + (1 - \varepsilon) \Pi_{\text{diag}}(X),$$

so that

$$E^n(X) = \Pi_{\text{diag}}(X) + \varepsilon^n(X - \Pi_{\text{diag}}(X)), \quad E^2 - E = (\varepsilon^2 - \varepsilon)(\text{id} - \Pi_{\text{diag}}).$$

The fixed-point algebra is diagonal for every $n > 0$, but $E^2 \neq E$.

The missing source step is the coefficient argument in [CPGSV16, Appendix C.4]. One needs the same-length product law

$$O_L(M_\alpha)O_L(M_\beta) = \sum_\gamma \text{tr}(\chi_{\alpha,\beta,\gamma}^L) O_L(M_\gamma) \quad (24.1)$$

and the length-one idempotent trace identity

$$m_\gamma = \sum_{\alpha,\beta} \text{tr}(\chi_{\alpha,\beta,\gamma}) m_\alpha m_\beta. \quad (24.2)$$

The proof of [CPGSV16, Theorem 4.14] uses (24.1) and (24.2), positivity, and the simple-matrix lemma [CPGSV16, lines 1155–1163] to give

$$\{\nu_{\gamma,k}\}_k = \{\mu_{\alpha,k_1} \mu_{\beta,k_2} \chi_{\alpha,\beta,\gamma,k_3}\}_{\alpha,\beta,k_1,k_2,k_3}, \quad \text{tr}(\nu_\gamma) = m_\gamma. \quad (24.3)$$

The reconstruction maps in [CPGSV16, lines 2065–2085] then use

$$\widetilde{R}_2 \left(\bigoplus_\gamma X_\gamma \right) = \bigoplus_\gamma \frac{\nu_\gamma}{m_\gamma} \otimes X_\gamma, \quad \widetilde{R}_1 \left(\bigoplus_\gamma X_\gamma \right) = \bigoplus_\gamma \frac{\mu_\gamma}{m_\gamma} \otimes X_\gamma. \quad (24.4)$$

Thus $\text{tr}(\nu_\gamma) = m_\gamma$ and the defining equality $m_\gamma = \text{tr}(\mu_\gamma)$ normalize the factors ν_γ/m_γ and μ_γ/m_γ , respectively, in $T = \widetilde{R}_2 \widetilde{T} R_1$ and $S = \widetilde{R}_1 \widetilde{S} R_2$. The scalar Newton–Girard power-sum identity needed to obtain the trace-power form is already available as Theorem 10.3.1. A source-faithful completion must state the BNT-label hypotheses of [CPGSV16, Theorem 4.14(ii)] as a separate condition, relate them to the one-site and two-site vertical canonical forms, and then construct the maps in (24.4). These hypotheses cannot be derived from the support-algebra condition, by Theorem 24.2.19.

24.3 Diagonal χ -matrices and the trace-power formula

The following statements isolate the form of [CPGSV16, Theorem 4.14(ii)]: the structure coefficients of the length- L operator algebra are trace-powers of positive diagonal matrices $\chi_{\alpha,\beta,\gamma}$. They state the coefficient identity and the elementary trace-power identity for diagonal matrices.

Definition 24.3.1 (Diagonal χ family). [Lean](#) [✓ Stmt](#) A family of diagonal complex matrices, indexed by ordered triples (α, β, γ) drawn from a common index type, consisting of a size $r_{\alpha,\beta,\gamma} \in \mathbb{N}$ and diagonal entries $\chi_{\alpha,\beta,\gamma,1}, \dots, \chi_{\alpha,\beta,\gamma,r_{\alpha,\beta,\gamma}} \in \mathbb{C}$ for each triple, giving the diagonal matrix

$$\chi_{\alpha,\beta,\gamma} = \text{diag}(\chi_{\alpha,\beta,\gamma,1}, \dots, \chi_{\alpha,\beta,\gamma,r_{\alpha,\beta,\gamma}}).$$

This is the matrix $\chi_{\alpha,\beta,\gamma}$ of [CPGSV16, Theorem 4.14(ii)].

Theorem 24.3.2 (Trace of χ^L equals the sum of L -th powers of entries). [Lean](#) [✓ Stmt](#) For every triple (α, β, γ) , if $\chi_{\alpha,\beta,\gamma}$ has size $r_{\alpha,\beta,\gamma}$, then for every $L \geq 0$,

$$\text{tr}(\chi_{\alpha,\beta,\gamma}^L) = \sum_{k=1}^{r_{\alpha,\beta,\gamma}} \chi_{\alpha,\beta,\gamma,k}^L,$$

where $\chi_{\alpha,\beta,\gamma,k}$ are the diagonal entries.

Proof. [✓ Proof](#) Diagonal matrices raise to powers entrywise and the trace of a diagonal matrix is the sum of its entries. $\square$

Definition 24.3.3 (Trace-power form of a structure-coefficient family). [Lean](#) [✓ Stmt](#) A binary compatibility predicate between an abstract structure-coefficient family $c_{\alpha,\beta,\gamma}^{(L)}$ and a diagonal χ family: the pair (c, χ) is said to be in *trace-power form* when

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L) = \sum_{k=1}^{r_{\alpha,\beta,\gamma}} \chi_{\alpha,\beta,\gamma,k}^L$$

for every $L \geq 0$ and every triple (α, β, γ) , where $r_{\alpha,\beta,\gamma}$ is the size of $\chi_{\alpha,\beta,\gamma}$. This is the formal analog of the target identity of [CPGSV16, Theorem 4.14(ii)]; the existential version—“there exists χ for which (c, χ) has trace-power form”—is obtained by quantifying over χ .

Definition 24.3.4 (BNT-label coefficient family). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The same-length coefficient system $c_{\alpha,\beta,\gamma}^{(L)}$ from [CPGSV16, Theorem 4.14(ii)], indexed by fixed BNT labels α, β, γ . The coefficient is attached to the length- L product $O_L(M_\alpha)O_L(M_\beta)$, not to the blocked-basis multiplication $\mathcal{A}_n \times \mathcal{A}_n \rightarrow \mathcal{A}_{2n}$. A diagonal χ -family canonically determines the coefficient family $c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L) = \sum_r \chi_{\alpha,\beta,\gamma,r}^L$, as in [CPGSV16, Appendix C.4].

Definition 24.3.5 (BNT-label operator family). [Lean](#) [✓ Stmt](#) A BNT-label operator family records, for each chain length L , the operators $O_L(M_\alpha)$ indexed by the fixed BNT labels α . The ambient algebra may depend on L and is not identified with a chosen blocked support algebra. The vertical indices of the L copies are contracted cyclically, while the horizontal legs remain open, as in [CPGSV16, lines 962–967]:

$$O_L(M_\alpha) =$$


Definition 24.3.6 (Same-length BNT product form). [Lean](#) [Lean](#) [✓ Stmt](#) A BNT-label operator family and a BNT-label coefficient system have same-length product form when, for every positive length L ,

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} c_{\alpha,\beta,\gamma}^{(L)} O_L(M_\gamma).$$

This is the product identity appearing in [CPGSV16, Theorem 4.14(ii)], stated independently of the blocked-basis multiplication $\mathcal{A}_n \times \mathcal{A}_n \rightarrow \mathcal{A}_{2n}$.

Definition 24.3.7 (BNT-label trace-scalar family). [Lean](#) [✓ Stmt](#) A BNT-label trace-scalar family records the scalars $m_\alpha = \text{tr}(\mu_\alpha)$ indexed by the fixed BNT labels α , as in the idempotent condition of [CPGSV16, Theorem 4.14(ii)].

Definition 24.3.8 (BNT idempotent coefficient form). [Lean](#) [Lean](#) [✓ Stmt](#) A BNT-label trace-scalar family and a BNT-label coefficient system have idempotent coefficient form when, for every BNT label γ ,

$$m_\gamma = \sum_{\alpha,\beta} c_{\alpha,\beta,\gamma}^{(1)} m_\alpha m_\beta.$$

This is the idempotent condition in [CPGSV16, Theorem 4.14(ii)], stated separately from the trace-power formula.

Definition 24.3.9 (Blocked-basis BNT-label assignment). [Lean](#) [Lean](#) [✓ Stmt](#) The BNT-label assignment consists, for every positive blocked length n , of a source label map σ_n from the chosen basis of $\mathcal{A}_n$ and a target label map τ_n from the chosen basis of $\mathcal{A}_{2n}$ to the fixed BNT labels. It also gives the combined label map $\sigma_n \sqcup \tau_n$ on the disjoint union of these two blocked bases.

Definition 24.3.10 (BNT comparison with blocked coefficients). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) A comparison between the BNT-label coefficients and the chosen blocked-basis coefficients consists of a BNT-label assignment together with the equality

$$c_{i,j,k}^{(n)} = c_{\sigma_n(i),\sigma_n(j),\tau_n(k)}^{(n)}.$$

This is a blocked-basis comparison: the left hand side is expanded in the chosen basis of $\mathcal{A}_{2n}$, so the statement is not itself the same-length BNT product law.

Definition 24.3.11 (Positive-length BNT-label trace-power form). [Lean](#) [✓ Stmt](#) A BNT-label coefficient system and a diagonal χ -family are compatible when, for every positive length L ,

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L).$$

The same matrix $\chi_{\alpha,\beta,\gamma}$ is used for all positive L .

Lemma 24.3.12 (BNT-label coefficient as a positive-length trace power). [Lean](#) [✓ Stmt](#) Under positive-length BNT-label trace-power form, for every $L > 0$,

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L).$$

Proof. [✓ Proof](#) Combine the defining positive-length trace-power identity with the trace formula for powers of a diagonal matrix. □

Lemma 24.3.13 (Canonical BNT coefficients from a χ -family). [Lean](#) [✓ Stmt](#) The coefficient family canonically determined by a diagonal χ -family satisfies the positive-length trace-power condition with respect to the same χ -family.

Proof. ✓ Proof This follows from the definition of the canonical coefficient family. □

Definition 24.3.14 (Positive BNT-label χ trace-power witness). Lean Lean ✓ Stmt A positive BNT-label χ witness consists of a length-independent diagonal $\chi_{\alpha, \beta, \gamma}$ -family, positivity of every diagonal entry, and the positive-length trace-power identity for the BNT-label coefficients. This is the coefficient statement of [CPGSV16, Theorem 4.14(ii)]. If the coefficient family is chosen canonically from the same χ -family, positivity of the diagonal entries alone gives such a witness.

Lemma 24.3.15 (Positive BNT-label witness gives trace powers). Lean ✓ Stmt A positive BNT-label χ witness gives, for every $L > 0$, the formula $c_{\alpha, \beta, \gamma}^{(L)} = \text{tr}(\chi_{\alpha, \beta, \gamma}^L)$.

Proof. ✓ Proof Apply the trace reformulation of the positive-length trace-power identity from the witness. □

Theorem 24.3.16 (Same-length BNT product with chi-trace coefficients). Lean ✓ Stmt If the same-length BNT product identity holds and the BNT-label coefficients come from a positive length-independent χ -witness, then for every $L > 0$,

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} \text{tr}(\chi_{\alpha, \beta, \gamma}^L)O_L(M_\gamma).$$

Proof. ✓ Proof Substitute the positive-length trace-power formula into the same-length BNT product expansion. □

Theorem 24.3.17 (Same-length BNT product for canonical chi coefficients). Lean ✓ Stmt If the same-length BNT product identity holds with the coefficient family canonically determined by a diagonal χ -family, then for every $L > 0$,

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} \text{tr}(\chi_{\alpha, \beta, \gamma}^L)O_L(M_\gamma).$$

Proof. ✓ Proof Substitute the canonical trace-power formula for the coefficients into the same-length BNT product expansion. □

Theorem 24.3.18 (BNT idempotent coefficients as chi traces). Lean ✓ Stmt If the BNT trace scalars satisfy the idempotent coefficient condition and the BNT-label coefficients come from a positive length-independent χ -witness, then

$$m_\gamma = \sum_{\alpha, \beta} \text{tr}(\chi_{\alpha, \beta, \gamma})m_\alpha m_\beta.$$

Proof. ✓ Proof Substitute the length-one trace-power formula into the idempotent coefficient identity. □

Theorem 24.3.19 (BNT idempotent coefficients for canonical chi coefficients). Lean ✓ Stmt If the BNT trace scalars satisfy the idempotent coefficient condition with the coefficient family canonically determined by a diagonal χ -family, then

$$m_\gamma = \sum_{\alpha, \beta} \text{tr}(\chi_{\alpha, \beta, \gamma})m_\alpha m_\beta.$$

Proof. ✓ Proof Substitute the length-one canonical trace formula into the idempotent coefficient identity. □

Lemma 24.3.20 (Blocked coefficients inherit BNT trace powers). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#)
If the chosen blocked-basis coefficients are compared with the fixed BNT-label coefficient system, and that BNT-label system has a positive length-independent χ -witness, then each positive-length blocked coefficient is the trace power of the corresponding BNT-label χ -matrix:

$$c_{i,j,k}^{(n)} = \text{tr}(\chi_{\sigma_n(i), \sigma_n(j), \tau_n(k)}^n).$$

Proof. [✓ Proof](#) Use the comparison equality to replace the blocked-basis coefficient by the corresponding BNT-label coefficient, and then apply the BNT-label trace-power formula. In the special case where the compared BNT-label coefficient family is already the canonical family determined by χ , the same conclusion follows directly from the canonical trace formula. $\square$

Definition 24.3.21 (Blocked χ family for multiplication coefficients). [Lean](#) [✓ Stmt](#) For each blocked multiplication triple (n, i, j, k) , where i and j label basis elements of $\mathcal{A}_n$ and k labels a basis element of $\mathcal{A}_{2n}$, a size $r_{n,i,j,k} \in \mathbb{N}$ and diagonal entries $\chi_{n,i,j,k,1}, \dots, \chi_{n,i,j,k,r_{n,i,j,k}} \in \mathbb{C}$, giving the diagonal matrix

$$\chi_{n,i,j,k} = \text{diag}(\chi_{n,i,j,k,1}, \dots, \chi_{n,i,j,k,r_{n,i,j,k}}).$$

Unlike the uniform BNT-label family in [CPGSV16, Theorem 4.14(ii)], the size and diagonal entries may depend on the blocked length and on the chosen basis indices.

Lemma 24.3.22 (Trace of a blocked χ power). [Lean](#) [✓ Stmt](#) For each blocked multiplication triple (n, i, j, k) and every exponent L ,

$$\text{tr}(\chi_{n,i,j,k}^L) = \sum_r \chi_{n,i,j,k,r}^L.$$

Proof. [✓ Proof](#) Diagonal matrices raise to powers entrywise and the trace of a diagonal matrix is the sum of its entries. $\square$

Definition 24.3.23 (Trace-power form for blocked multiplication coefficients). [Lean](#) [✓ Stmt](#)
The blocked multiplication coefficients are in trace-power form when, for every positive blocked size n and every multiplication triple (i, j, k) , the coefficient $c_{i,j,k}^{(n)}$ equals

$$\sum_r \chi_{n,i,j,k,r}^n.$$

Equivalently, by Lemma 24.3.22, it is the trace of the n -th power of the corresponding diagonal matrix. The exponent n is the source blocking length of the two factors in $\mathcal{A}_n$; the product is expanded in the basis of $\mathcal{A}_{2n}$.

Lemma 24.3.24 (Blocked coefficient as a trace power). [Lean](#) [Lean](#) [✓ Stmt](#) Under trace-power form for the blocked multiplication coefficients, for every positive blocked size n ,

$$c_{i,j,k}^{(n)} = \text{tr}(\chi_{n,i,j,k}^n).$$

Proof. [✓ Proof](#) Combine the defining trace-power identity with the trace formula for powers of a diagonal matrix. $\square$

Definition 24.3.25 (Positive blocked χ trace-power witness). [Lean](#) [✓ Stmt](#) A positive blocked χ trace-power witness consists of a blocked χ family, positivity of all its diagonal entries, and the positive-length trace-power identity for the blocked multiplication coefficients. This is the blocked-basis analog of the positive diagonal matrices in [CPGSV16, Theorem 4.14(ii)], without requiring a single family indexed only by the BNT labels.

together with a description of their excitations [BMW⁺17]. The renormalization fixed-point density operators correspond to the boundary theories of the renormalization fixed-point representatives of those phases, which yields a large family of non-trivial examples. Whether there exist renormalization fixed points whose structure coefficients depend on the length is left open in [CPGSV16, Section 4.5].

Definition 24.3.32 (Linearly independent operator family). Lean Lean ✓ Stmt A BNT-label operator family is *linearly independent at length L* when the operators $O_L(M_\alpha)$ are linearly independent, and *eventually linearly independent* when this holds for every length beyond some threshold. Eventual linear independence is the operator-level form of the third defining property of a basis of normal tensors in [CPGSV16, Section 2.3], invoked for the labelled operators in [CPGSV16, Appendix C.4].

Theorem 24.3.33 (Uniqueness of the structure coefficients). Lean ✓ Stmt *Two coefficient systems satisfying the same-length product law for the same operator family agree at every positive length at which the operators are linearly independent.*

Proof. ✓ Proof Applying the same-length product law to both coefficient systems gives

$$\sum_{\delta} c_{\alpha, \beta, \delta}^{(L)} O_L(M_\delta) = O_L(M_\alpha) O_L(M_\beta) = \sum_{\delta} c'_{\alpha, \beta, \delta}{}^{(L)} O_L(M_\delta).$$

Subtracting yields $\sum_{\delta} (c_{\alpha, \beta, \delta}^{(L)} - c'_{\alpha, \beta, \delta}{}^{(L)}) O_L(M_\delta) = 0$, and linear independence forces every coefficient of the difference to zero. □

Theorem 24.3.34 (Associativity constraint on the structure coefficients). Lean ✓ Stmt *At every positive length L at which the operators are linearly independent, the structure coefficients of the same-length product law satisfy*

$$\sum_{\delta} c_{\alpha, \beta, \delta}^{(L)} c_{\delta, \gamma, \epsilon}^{(L)} = \sum_{\delta} c_{\beta, \gamma, \delta}^{(L)} c_{\alpha, \delta, \epsilon}^{(L)}.$$

These are the restrictions that the associativity of the multiplication imposes on the structure coefficients, noted in [CPGSV16, Section 4.5].

Proof. ✓ Proof Applying the same-length product law twice in each order gives

$$\begin{aligned} (O_L(M_\alpha) O_L(M_\beta)) O_L(M_\gamma) &= \sum_{\epsilon'} \left(\sum_{\delta} c_{\alpha, \beta, \delta}^{(L)} c_{\delta, \gamma, \epsilon'}^{(L)} \right) O_L(M_{\epsilon'}), \\ O_L(M_\alpha) (O_L(M_\beta) O_L(M_\gamma)) &= \sum_{\epsilon'} \left(\sum_{\delta} c_{\beta, \gamma, \delta}^{(L)} c_{\alpha, \delta, \epsilon'}^{(L)} \right) O_L(M_{\epsilon'}). \end{aligned}$$

Associativity of multiplication equates the two left-hand sides, so the right-hand sides agree; linear independence of the operators then forces the corresponding coefficient sums to agree for each ϵ' . □

Theorem 24.3.35 (Associativity constraint in trace-power form). Lean ✓ Stmt *For a coefficient system satisfying the same-length product law for a BNT-label operator family and in trace-power form, at every positive length L at which the labelled operators are linearly independent, the associativity constraint is a restriction on the family of diagonal matrices:*

$$\sum_{\delta} \text{tr}(\chi_{\alpha, \beta, \delta}^L) \text{tr}(\chi_{\delta, \gamma, \epsilon}^L) = \sum_{\delta} \text{tr}(\chi_{\beta, \gamma, \delta}^L) \text{tr}(\chi_{\alpha, \delta, \epsilon}^L).$$

In [CPGSV16, Section 4.5] these restrictions, combined with the isometry statement, lead to a pentagon-like equation for the fusion isometries whose solutions come from unitary fusion categories.

Proof. ✓ Proof Substitute the trace-power form of the coefficients into the associativity constraint. □

Definition 24.3.36 (BNT algebra clause). Lean ✓ Stmt Fix a coefficient family $c_{\alpha,\beta,\gamma}^{(L)}$, BNT-label operators $O_L(M_\alpha)$, and trace scalars $m_\alpha = \text{tr}(\mu_\alpha)$. They satisfy the *BNT algebra clause* when there is a length-independent family of positive diagonal matrices $\chi_{\alpha,\beta,\gamma}$ such that, for every $L > 0$,

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L),$$

the same-length product law holds,

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} c_{\alpha,\beta,\gamma}^{(L)} O_L(M_\gamma),$$

and the trace scalars satisfy

$$m_\gamma = \sum_{\alpha,\beta} c_{\alpha,\beta,\gamma}^{(1)} m_\alpha m_\beta.$$

This is the algebra statement in [CPGSV16, Theorem 4.14(ii), labelled IV.13 in the source, lines 972–985], before comparison with a chosen blocked basis.

Definition 24.3.37 (BNT multiplicity-spectrum comparison). Lean ✓ Stmt Fix the one-site diagonal multiplicity matrices μ_α and the two-site diagonal multiplicity matrices ν_γ in the respective vertical canonical decompositions. A multiplicity-spectrum comparison records

$$\text{tr}(\mu_\alpha) = m_\alpha$$

and, for every γ , equality with multiplicity between the entries of ν_γ and the products

$$\mu_{\alpha,i} \mu_{\beta,j} \chi_{\alpha,\beta,\gamma,k}.$$

This is the conclusion of the vertical BNT comparison and positive power-sum argument in [CPGSV16, Appendix C.4, lines 2048–2058]. The definition does not require this comparison to follow from a BNT algebra clause; constructing it from the MPDO tensor is a separate obligation.

Lemma 24.3.38 (Idempotence for the canonical chi coefficients). Lean ✓ Stmt *The length-one idempotent law of a BNT algebra clause also holds for the coefficient family determined by its diagonal chi matrices:*

$$m_\gamma = \sum_{\alpha,\beta} \text{tr}(\chi_{\alpha,\beta,\gamma}) m_\alpha m_\beta.$$

Proof. ✓ Proof The chi representation at length one gives

$$c_{\alpha,\beta,\gamma}^{(1)} = \text{tr}(\chi_{\alpha,\beta,\gamma}).$$

Substitution into the idempotent law

$$m_\gamma = \sum_{\alpha,\beta} c_{\alpha,\beta,\gamma}^{(1)} m_\alpha m_\beta$$

gives the stated identity. □

BNT-label theorem data consist of a BNT algebra clause together with the comparison with the chosen blocked bases. Explicitly, for every $L > 0$ the product law is

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} c_{\alpha,\beta,\gamma}^{(L)} O_L(M_\gamma),$$

while the idempotent scalar law is

$$m_\gamma = \sum_{\alpha,\beta} c_{\alpha,\beta,\gamma}^{(1)} m_\alpha m_\beta.$$

For every positive blocked length n and blocked-basis indices i, j, k , the source and target label maps σ_n, τ_n give the comparison

$$c_{i,j,k}^{(n)} = c_{\sigma_n(i),\sigma_n(j),\tau_n(k)}^{(n)}.$$

They also determine the pulled-back blocked-basis χ -family. These data are the source-side hypotheses from which the blocked-basis consequences below are derived. In the source-side special case where the coefficient family is canonically determined by the same χ -family, theorem data are built from that χ -family, its positivity, and the product, idempotent, and blocked-comparison statements for the resulting canonical coefficients. The same product and idempotent predicates may also be rephrased using the canonical coefficient family determined by the χ -matrices carried by the data, with the corresponding displayed equations.

Theorem 24.3.42 (Source BNT coefficient and product formulas). [Lean](#) [Lean](#) [Lean](#) [Lean](#)

[Lean](#) [✓ Stmt](#) Let $c_{\alpha,\beta,\gamma}^{(L)}$ be the coefficient family and let $\chi_{\alpha,\beta,\gamma}$ be the positive diagonal matrices carried by the theorem data. For every $L > 0$,

$$c_{\alpha,\beta,\gamma}^{(L)} = c_{\alpha,\beta,\gamma}^{\chi,(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L),$$

where c^χ is the canonical coefficient family determined by χ , and every diagonal entry of every $\chi_{\alpha,\beta,\gamma}$ is strictly positive. Consequently, for every $L > 0$, the same-length product identity reads

$$O_L(M_\alpha)O_L(M_\beta) = \sum_{\gamma} \text{tr}(\chi_{\alpha,\beta,\gamma}^L) O_L(M_\gamma),$$

and, at $L = 1$, the idempotent identity reads

$$m_\gamma = \sum_{\alpha,\beta} \text{tr}(\chi_{\alpha,\beta,\gamma}) m_\alpha m_\beta.$$

Proof. [✓ Proof](#) The positive trace-power witness gives

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L).$$

Substitution into the product law and its length-one idempotent specialization gives the displayed formulas. $\square$

Theorem 24.3.43 (BNT theorem data give a positive blocked χ witness). [Lean](#) [Lean](#) [Lean](#)

[Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) BNT-label theorem data give a positive blocked-basis χ trace-power witness by pulling the length-independent BNT-label χ -family back along the blocked-basis comparison maps: writing $\chi_{n,i,j,k}$ for the resulting blocked-basis diagonal matrix at a positive blocked length n ,

$$\chi_{n,i,j,k} = \chi_{\sigma_n(i),\sigma_n(j),\tau_n(k)}.$$

Thus the two matrices have equal size, diagonal entries, and traces of all powers.

Proof. [✓ Proof](#) Apply the definition at blocked size $n = 1$. If $E_1 = S_1 \circ T_1$ and $T_1 \circ S_1 = \text{id}$, then

$$E_1^2 = S_1 T_1 S_1 T_1 = S_1 (T_1 S_1) T_1 = S_1 T_1 = E_1.$$

Since E_1 is the original transfer map, this is exactly transfer-map idempotence. $\square$

Theorem 24.4.4 (One-site fusion retract criterion). [Lean](#) [✓ Stmt](#) *A one-site transfer-map fusion-isometry datum exists if and only if the MPO transfer map is idempotent.*

Proof. [✓ Proof](#) The forward implication is the retract calculation $E_1^2 = S_1 T_1 S_1 T_1 = S_1 T_1$. Conversely, if E_1 is idempotent, then it factors through its range, giving the required one-site datum. $\square$

Theorem 24.4.5 (Transfer-map fusion criterion for idempotence). [Lean](#) [✓ Stmt](#) *The fusion-isometry formulation is equivalent to transfer-map idempotence.*

Proof. [✓ Proof](#) The forward implication is Theorem 24.4.3. Conversely, if the transfer map E is idempotent, then every positive blocked transfer map equals E and therefore factors through its range. This range-factorization gives the required fusion-isometry data. $\square$

Corollary 24.4.6 (Pure-state recovery). [Lean](#) [✓ Stmt](#) *For the diagonal MPO associated to a pure MPS tensor, the fusion-isometry formulation reduces to the usual pure-state RFP condition.*

Proof. [✓ Proof](#) Combine Theorem 24.4.5 with the diagonal pure-state recovery theorem for the MPO predicate. $\square$

Theorem 24.4.7 (All-blocked fusion reduces to one-site fusion). [Lean](#) [✓ Stmt](#) *The fusion-isometry formulation for all positive blocked sizes is equivalent to the existence of a one-site transfer-map fusion-isometry datum.*

Proof. [✓ Proof](#) Combine the one-site criterion with the equivalence between the all-blocked fusion formulation and transfer-map idempotence. $\square$

Definition 24.4.8 (Operator product of MPO tensors). [Lean](#) [✓ Stmt](#) The *product tensor* of two MPO tensors contracts the bra index of the first factor with the ket index of the second and takes the tensor product of the bond spaces:

$$(M \cdot N)^{ik} = \sum_j M^{ij} \otimes N^{jk}.$$

In the vertical reading of [CPGSV16, Section 4.5] this is the tensor obtained by joining two vertically read tensors along the original horizontal bond, the tensor written $M_\alpha M_\beta$ in [CPGSV16, Theorem 4.14(iii)].

Definition 24.4.9 (Canonical reassociation of three bond spaces). [Lean](#) [Lean](#) [✓ Stmt](#) For three bond spaces, let

$$A_{D_1, D_2, D_3} : \mathbb{C}^{D_1} \otimes (\mathbb{C}^{D_2} \otimes \mathbb{C}^{D_3}) \longrightarrow (\mathbb{C}^{D_1} \otimes \mathbb{C}^{D_2}) \otimes \mathbb{C}^{D_3}$$

be the coordinate pullback induced by canonical reassociation, defined on elementary tensors by $A_{D_1, D_2, D_3}(x_1 \otimes (x_2 \otimes x_3)) = (x_1 \otimes x_2) \otimes x_3$.

Theorem 24.4.10 (Associativity of the product tensor). [Lean](#) [Lean](#) [✓ Stmt](#) *The letters of the two parenthesizations of a triple product are related by the canonical bond reassociation. Equivalently, for every pair of physical indices i, k ,*

$$((M \cdot N) \cdot P)^{ik} A_{D_1, D_2, D_3} = A_{D_1, D_2, D_3}(M \cdot (N \cdot P))^{ik}.$$

Proof. ✓ Proof Expand both product tensors. Reassociation preserves every elementary tensor factor, while exchanging the order of the two finite sums gives the same contraction over the two intermediate physical indices. $\square$

Lemma 24.4.11 (Word evaluation of the product tensor). Lean ✓ Stmt *The word evaluation of the product tensor along a configuration pair (σ, τ) expands as a sum over the contracted middle configurations ρ of tensor products of word evaluations:*

$$(M \cdot N)^{\sigma\tau} = \sum_{\rho} M^{\sigma\rho} \otimes N^{\rho\tau}.$$

Proof. ✓ Proof Induction on the length, using the mixed-product property of the tensor product to merge one letter at a time. $\square$

Theorem 24.4.12 (The operator family is multiplicative). Lean ✓ Stmt *For every chain length L , the operator generated by the product tensor is the product of the generated operators: $O_L(M \cdot N) = O_L(M) O_L(N)$.*

Proof. ✓ Proof Take traces in the word-evaluation expansion of Lemma 24.4.11: the trace of a tensor product is the product of the traces, and the sum over middle configurations is the entrywise form of the operator product. $\square$

Lemma 24.4.13 (Word products of isometrically conjugated letters). Lean ✓ Stmt *For an isometry U (a rectangular matrix with $U^\dagger U = 1$) and a nonempty word of square matrices $F_1, \dots, F_L$,*

$$(U F_1 U^\dagger)(U F_2 U^\dagger) \dots (U F_L U^\dagger) = U F_1 F_2 \dots F_L U^\dagger.$$

Proof. ✓ Proof The inner factors $U^\dagger U$ telescope to the identity. $\square$

Lemma 24.4.14 (Word products of block-diagonal tensor-product letters). Lean ✓ Stmt *For block-diagonal letters whose block of index γ is the tensor product of a fixed matrix X_γ with a varying letter $G_{\gamma,l}$,*

$$\prod_{l=1}^L \bigoplus_{\gamma} X_\gamma \otimes G_{\gamma,l} = \bigoplus_{\gamma} X_\gamma^L \otimes \prod_{l=1}^L G_{\gamma,l}.$$


Proof. ✓ Proof Blocks multiply blockwise and tensor products multiply factorwise, so the fixed factors accumulate into the matrix power X_γ^L . $\square$

Definition 24.4.15 (Fusion-isometry family). Lean ✓ Stmt *A fusion-isometry family over a finite set of labels consists of a family of tensors M_γ with a common physical dimension and per-label bond dimensions, diagonal matrices $\chi_{\alpha,\beta,\gamma}$ with positive entries, and isometries $U_{\alpha,\beta}$ such that, site by site,*

$$U_{\alpha,\beta} (M_\alpha M_\beta)^{ij} U_{\alpha,\beta}^\dagger = \bigoplus_{\gamma} \chi_{\alpha,\beta,\gamma} \otimes M_\gamma^{ij}.$$

This is the isometry statement of [CPGSV16, Theorem 4.14(iii)]; the tensors are the vertically read basis of normal tensors of [CPGSV16, Proposition 4.13], so the physical dimension is the bond dimension of the original tensor. Statement (iii) of the source theorem additionally asserts the idempotent identity for the trace scalars of the vertical decomposition, recorded separately by Definition 24.3.8.

The following compact display depicts the sitewise identity $U_{\alpha,\beta}(M_\alpha M_\beta)^{ij}U_{\alpha,\beta}^\dagger = \bigoplus_\gamma \chi_{\alpha,\beta,\gamma} \otimes M_\gamma^{ij}$ from [CPGSV16, Theorem 4.14(iii), equation (89), lines 986–991]. Its arrangement follows the physical isometry and weighted direct-sum decomposition in [CPGSV16, Proposition 4.13, lines 943–959].



Theorem 24.4.16 (Weighted zipper identities). [Lean](#) [Lean](#) [✓ Stmt](#) For every pair of labels and physical indices, write

$$G_{\alpha,\beta}^{ij} = \bigoplus_\gamma \chi_{\alpha,\beta,\gamma} \otimes M_\gamma^{ij}.$$

The fusion isometry satisfies the two chi-weighted zipper identities

$$U_{\alpha,\beta}(M_\alpha M_\beta)^{ij} = G_{\alpha,\beta}^{ij} U_{\alpha,\beta}, \quad (M_\alpha M_\beta)^{ij} U_{\alpha,\beta}^\dagger = U_{\alpha,\beta}^\dagger G_{\alpha,\beta}^{ij}.$$

The corresponding identities in [BMW⁺17, Equation (21)] have identity matrices on the multiplicity spaces in place of the positive diagonal matrices χ . Their unweighted form requires the length-independent integer specialization and is not a consequence of the present hypotheses. Even after that specialization, comparison of the two triple-fusion orders and final-label separation are still needed to construct an invertible F -move.

Proof. [✓ Proof](#) Multiply the fusion identity $U_{\alpha,\beta}(M_\alpha M_\beta)^{ij}U_{\alpha,\beta}^\dagger = G_{\alpha,\beta}^{ij}$ on the right by $U_{\alpha,\beta}$ and on the left by $U_{\alpha,\beta}^\dagger$, respectively, and use $U_{\alpha,\beta}^\dagger U_{\alpha,\beta} = 1$. $\square$

Theorem 24.4.17 (Length-independent zipper identities and completeness). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Suppose the structure coefficients have positive trace-power form and are independent of the positive chain length. Then $\chi_{\alpha,\beta,\gamma} = 1_{r_{\alpha,\beta,\gamma}}$ for every triple of labels. If

$$H_{\alpha,\beta}^{ij} = \bigoplus_\gamma 1_{r_{\alpha,\beta,\gamma}} \otimes M_\gamma^{ij},$$

the fusion isometries satisfy

$$U_{\alpha,\beta}(M_\alpha M_\beta)^{ij} = H_{\alpha,\beta}^{ij} U_{\alpha,\beta}, \quad (M_\alpha M_\beta)^{ij} U_{\alpha,\beta}^\dagger = U_{\alpha,\beta}^\dagger H_{\alpha,\beta}^{ij},$$

and

$$(M_\alpha M_\beta)^{ij} = U_{\alpha,\beta}^\dagger H_{\alpha,\beta}^{ij} U_{\alpha,\beta}.$$

These are the identity-weight specialization of Equations (19)–(21) in [BMW⁺17, Section 3]. The MPDO fusion identity appears in [CPGSV16, Theorem 4.14(iii), lines 986–991]; its identity-weight specialization follows from the length-independent case at [CPGSV16, line 1010].

Proof. [✓ Proof](#) Length independence makes every positive diagonal entry of $\chi_{\alpha,\beta,\gamma}$ equal to one. Substitute the resulting identity matrices into Theorem 24.4.16. For completeness, insert $U_{\alpha,\beta}^\dagger U_{\alpha,\beta} = 1$ on both sides of the product letter and apply the fusion identity between the two inserted factors. $\square$

Definition 24.4.19 (Operator family of a fusion-isometry family). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) The tensors of a fusion-isometry family generate the concrete BNT-label operator family $O_L(M_\gamma)$, and the positive diagonal matrices give the positive trace-power witness for the coefficient family $c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L)$.




Figure 24.1: Closing an L -site chain of the local fusion identity. On the left, the bra legs of the M_α operator are contracted with the ket legs of the M_β operator. On the right, the multiplicity loop and the M_γ loop are disjoint; the former contributes $\text{tr}(\chi_{\alpha,\beta,\gamma}^L)$. Thus the diagram is the trace-power product law of [CPGSV16, Theorem 4.14(ii)–(iii), equations (87) and (89), lines 962–991].

Theorem 24.4.20 (Fusion isometries give the same-length product law). [Lean](#) [Lean](#) [✓ Stmt](#) The operator family of a fusion-isometry family satisfies the same-length product law with the trace-power coefficients of its diagonal matrices: for every chain length $L \geq 1$,

$$O_L(M_\alpha) O_L(M_\beta) = \sum_{\gamma} \text{tr}(\chi_{\alpha,\beta,\gamma}^L) O_L(M_\gamma).$$

This derives the product law of [CPGSV16, Theorem 4.14(ii)] from the isometry statement of [CPGSV16, Theorem 4.14(iii)].

Proof. [✓ Proof](#) The product of the operators is generated by the product tensor. For a nonempty word W of letters of the product tensor, the conjugated letters multiply to the conjugated word, and the closed trace is unchanged because $U_{\alpha,\beta}$ is an isometry:

$$\text{tr}(U_{\alpha,\beta} W U_{\alpha,\beta}^\dagger) = \text{tr}(W U_{\alpha,\beta}^\dagger U_{\alpha,\beta}) = \text{tr}(W).$$

By the fusion identity the conjugated letters are the block-diagonal tensor-product letters of the weighted direct sum. Their word product splits block by block with the diagonal factors accumulating into $\chi_{\alpha,\beta,\gamma}^L$; taking the trace block by block and factor by factor yields the coefficients $\text{tr}(\chi_{\alpha,\beta,\gamma}^L)$. $\square$

Definition 24.4.21 (BNT algebra clause derived from fusion isometries). [Lean](#) [✓ Stmt](#) Suppose the fusion-isometry identity holds with positive diagonal matrices $\chi_{\alpha,\beta,\gamma}$, and suppose the trace scalars satisfy the length-one idempotent law for the corresponding trace-power coefficients. The resulting BNT algebra clause has coefficients, for $L > 0$,

$$c_{\alpha,\beta,\gamma}^{(L)} = \text{tr}(\chi_{\alpha,\beta,\gamma}^L),$$

and records positivity of every diagonal entry. The product law is

$$O_L(M_\alpha) O_L(M_\beta) = \sum_{\gamma} c_{\alpha,\beta,\gamma}^{(L)} O_L(M_\gamma),$$

and the idempotent condition is the assumed identity

$$m_\gamma = \sum_{\alpha, \beta} c_{\alpha, \beta, \gamma}^{(1)} m_\alpha m_\beta.$$

This construction realizes implication (iii) $\Rightarrow$ (ii) of [CPGSV16, Theorem 4.14]; it does not assert either of the implications involving the renormalization fixed-point condition.




Figure 24.2: A schematic recursive contraction for the structure operator Q . Each four-legged box X is the tensor $X = (\chi_{\alpha, \beta, \gamma, k})_{k, \alpha, \beta, \gamma}$ of diagonal entries of $\chi_{\alpha, \beta, \gamma}$, and the terminal contraction is $\text{tr}(M_\gamma)$. The dots indicate index contractions, while the omitted middle part repeats the displayed pattern. This is only the schematic form of the source display and does not assert that Q is a projector [CPGSV16, lines 999–1008].

The associativity of the algebra product in [CPGSV16, Equation (87)] constrains the diagonal matrices $\chi_{\alpha, \beta, \gamma}$ and, through the isometry statement, induces a compatibility equation between the fusion isometries of the two ways of bracketing a triple product, $(M_\alpha M_\beta)M_\gamma$ and $M_\alpha(M_\beta M_\gamma)$ [CPGSV16, lines 995–999]. The source attributes the construction of that equation, and the classification of its solutions, to [BMW⁺17]. The next definitions and theorems first record the tensor-level content of the two bracketings and then compare the resulting full direct sums. Restriction of that comparison to a fixed final label is a separate step.




Figure 24.3: The two ways of fusing three labels to a fixed final label ε , using the printed- F row/column convention of [BMW⁺17, Section 3.4]. The solid fusion trees are the left- and right-bracketed maps constructed below. The dashed line denotes the invertible fixed-final change of multiplicity basis in the source. The results below first extract its action on the multiplicity space and then prove its two-sided unitarity under the simultaneous-selector and positive-bond-dimension hypotheses.

Definition 24.4.22 (Iterated fusion isometry of a triple product, left bracketing). [Lean](#) [Stmt](#)
For labels α, β, γ , the *left iterated fusion isometry* $U_{\alpha, \beta, \gamma}^L$ is obtained by applying $U_{\alpha, \beta}$ (tensored

with the identity on the bond space of M_γ) to fuse the first two tensors, then applying $U_{\delta,\gamma}$, for each label δ appearing in the resulting direct sum, to fuse the outcome with the third tensor. Its codomain is indexed by pairs of labels (δ, ε) together with the multiplicity indices of $\chi_{\alpha,\beta,\delta}$ and $\chi_{\delta,\gamma,\varepsilon}$.

Theorem 24.4.23 (The left iterated fusion isometry is an isometry). [Lean](#) [✓ Stmt](#) $(U_{\alpha,\beta,\gamma}^L)^\dagger U_{\alpha,\beta,\gamma}^L = 1$.

Proof. [✓ Proof](#) The composite of two isometries is an isometry. $\square$

Theorem 24.4.24 (The left iterated fusion isometry conjugates a triple product onto a doubly weighted direct sum). [Lean](#) [✓ Stmt](#) *Site by site*,

$$U_{\alpha,\beta,\gamma}^L ((M_\alpha M_\beta) M_\gamma)^{ij} (U_{\alpha,\beta,\gamma}^L)^\dagger = \bigoplus_\delta \bigoplus_\varepsilon \chi_{\alpha,\beta,\delta} \otimes (\chi_{\delta,\gamma,\varepsilon} \otimes M_\varepsilon^{ij}).$$

Proof. [✓ Proof](#) Apply the fusion identity for $U_{\alpha,\beta}$ to the first two tensors, summed against the physical index contracting with the third tensor; regrouping the resulting sum block by block over δ exhibits it as $\chi_{\alpha,\beta,\delta}$ tensored with a letter of the product tensor $M_\delta M_\gamma$, to which the fusion identity for $U_{\delta,\gamma}$ applies. $\square$

Definition 24.4.25 (Iterated fusion isometry of a triple product, right bracketing). [Lean](#) [✓ Stmt](#) For labels α, β, γ , the *right iterated fusion isometry* $U_{\alpha,\beta,\gamma}^R$ is obtained by applying $U_{\beta,\gamma}$, tensored with the identity on the bond space of M_α , and then applying $U_{\alpha,\delta}$ for every intermediate label δ . Its codomain is indexed by the intermediate and final labels (δ, ε) and by the multiplicity indices of $\chi_{\beta,\gamma,\delta}$ and $\chi_{\alpha,\delta,\varepsilon}$. This is the right-bracketed composite in the associativity equation of [BMW⁺17, Section “Associativity and the pentagon equation”], before the change of basis by an F -matrix.

Theorem 24.4.26 (The right iterated fusion isometry is an isometry). [Lean](#) [✓ Stmt](#) $(U_{\alpha,\beta,\gamma}^R)^\dagger U_{\alpha,\beta,\gamma}^R = 1$.

Proof. [✓ Proof](#) The two successive fusion maps are isometries on each intermediate sector, so their block-diagonal composite is an isometry. $\square$

Theorem 24.4.27 (The right iterated fusion isometry conjugates a triple product onto a doubly weighted direct sum). [Lean](#) [✓ Stmt](#) *Site by site*,

$$U_{\alpha,\beta,\gamma}^R (M_\alpha (M_\beta M_\gamma))^{ij} (U_{\alpha,\beta,\gamma}^R)^\dagger = \bigoplus_\delta \bigoplus_\varepsilon \chi_{\beta,\gamma,\delta} \otimes (\chi_{\alpha,\delta,\varepsilon} \otimes M_\varepsilon^{ij}).$$

Proof. [✓ Proof](#) Apply the fusion identity for $U_{\beta,\gamma}$ to $M_\beta M_\gamma$. In the summand indexed by δ , exchange the first multiplicity factor with the bond space of M_α and regroup the contraction as

$$\chi_{\beta,\gamma,\delta} \otimes (M_\alpha M_\delta)^{ij}.$$

The fusion identity for $U_{\alpha,\delta}$ then gives the stated direct sum over ε . $\square$

Definition 24.4.28 (Full comparison of the two triple-fusion orders). [Lean](#) [✓ Stmt](#) The canonical comparison from the right-associated full direct sum to the left-associated full direct sum is

$$C_{\alpha,\beta,\gamma} = U_{\alpha,\beta,\gamma}^L A_{D_\alpha, D_\beta, D_\gamma} (U_{\alpha,\beta,\gamma}^R)^\dagger.$$

Both intermediate labels and the final label remain in the direct sums. This definition does not select a fixed final label and makes no invertibility assertion. Its orientation is from the right-associated sum to the left-associated sum; consequently it is adjoint-oriented with respect to the printed- F row/column convention used here for [BMW⁺17, Section 3.4].

Lemma 24.4.29 (Left range projection of the full triple-fusion comparison). [Lean](#) [✓ Stmt](#) The product of the full comparison with its adjoint is the range projection of the left iterated fusion isometry:

$$C_{\alpha,\beta,\gamma} C_{\alpha,\beta,\gamma}^\dagger = U_{\alpha,\beta,\gamma}^L (U_{\alpha,\beta,\gamma}^L)^\dagger.$$

The right side is not asserted to be the identity. Surjectivity of the left iterated fusion map requires completeness of the fusion decomposition, corresponding to the invertible total fusion matrix in [BMW⁺17, lines 181–191].

Proof. [✓ Proof](#) Substitute $C = U^L A (U^R)^\dagger$. The isometry identity $(U^R)^\dagger U^R = 1$ and the two-sided inverse identities for the permutation matrix A leave $U^L (U^L)^\dagger$. $\square$

Lemma 24.4.30 (Right range projection of the full triple-fusion comparison). [Lean](#) [✓ Stmt](#) The product in the opposite order is the range projection of the right iterated fusion isometry:

$$C_{\alpha,\beta,\gamma}^\dagger C_{\alpha,\beta,\gamma} = U_{\alpha,\beta,\gamma}^R (U_{\alpha,\beta,\gamma}^R)^\dagger.$$

The right side is not asserted to be the identity. Surjectivity of the right iterated fusion map likewise requires completeness of the fusion decomposition.

Proof. [✓ Proof](#) Substitute $C = U^L A (U^R)^\dagger$. The isometry identity $(U^L)^\dagger U^L = 1$ and the two-sided inverse identities for A leave $U^R (U^R)^\dagger$. $\square$

Theorem 24.4.31 (The full triple-fusion comparison intertwines the unweighted direct sums). [Lean](#) [✓ Stmt](#) Suppose the positive trace-power coefficients are independent of the positive chain length. For every pair of physical indices i, k , let

$$D_{ik}^L = \bigoplus_{\delta, \varepsilon} 1_{r_{\alpha, \beta}^\delta} \otimes (1_{r_{\beta, \gamma}^\varepsilon} \otimes M_\varepsilon^{ik})$$

and define D_{ik}^R by the right-associated multiplicities $r_{\beta, \gamma}^\delta$ and $r_{\alpha, \delta}^\varepsilon$. Then

$$D_{ik}^L C_{\alpha, \beta, \gamma} = C_{\alpha, \beta, \gamma} D_{ik}^R.$$

This is an identity on the full direct sums. It neither extracts a fixed-final F -transformation nor proves invertibility or the pentagon identity.

Proof. [✓ Proof](#) Length independence makes every positive diagonal factor equal to the identity. Substitute the left and right triple-fusion identities into the definition of $C_{\alpha, \beta, \gamma}$, cancel the two isometries, and apply Theorem 24.4.10. $\square$

Definition 24.4.32 (Common word selectors for the final labels). [Lean](#) [✓ Stmt](#) For some common length S , suppose that for each final label ε there are coefficients $c_\varepsilon(w)$, indexed by the words w of length S in the doubled physical alphabet, such that

$$\sum_{|w|=S} c_\varepsilon(w) M_\varepsilon^w = 1, \quad \sum_{|w|=S} c_{\varepsilon'}(w) M_\varepsilon^w = 0 \quad (\varepsilon' \neq \varepsilon).$$

This is the finite-word form of the simultaneous inverse relation $B_d^+ B_{d'} = \delta_{d, d'} 1$ used in the fixed-channel argument of [BMW⁺17].

Theorem 24.4.33 (Pair fusion completeness from positive-length selectors). [Lean](#) [✓ Stmt](#) Suppose the positive trace-power coefficients are independent of the positive chain length, and suppose that common selectors exist at a positive length. Then every pair fusion isometry is surjective:

$$U_{\alpha,\beta} U_{\alpha,\beta}^\dagger = 1.$$

The common selectors are the finite-word form of the simultaneous inverse used in [BMW⁺17, line 269].

Proof. [✓ Proof](#) For a word w of the positive selector length, the identity $U_{\alpha,\beta}^\dagger U_{\alpha,\beta} = 1$ gives

$$D^w = U_{\alpha,\beta} (M_\alpha M_\beta)^w U_{\alpha,\beta}^\dagger,$$

where $D = \bigoplus_\gamma 1_{r_{\alpha,\beta}^\gamma} \otimes M_\gamma$. Hence the range projection $P = U_{\alpha,\beta} U_{\alpha,\beta}^\dagger$ satisfies $PD^w = D^w$. Summing the selector polynomials over the final label gives the identity on the full direct sum, and therefore $P = 1$. $\square$

Theorem 24.4.34 (Pair fusion completeness from a source BNT). [Lean](#) [Lean](#) [✓ Stmt](#) Suppose that the labelled tensors in the fusion family form a basis of normal tensors and that the positive trace-power coefficients are independent of the positive chain length. Then every pair fusion isometry is surjective:

$$U_{\alpha,\beta} U_{\alpha,\beta}^\dagger = 1.$$

In particular, final-label selectors and coisometry are conclusions, not additional hypotheses. This is a derived consequence of the common BNT word span in [CPGSV16, lines 317–345], the fusion identity at lines 986–993, and length independence at lines 995–1010.

Proof. [✓ Proof](#) The basis-of-normal-tensors argument gives a positive common word length at which the labelled word evaluations span the full product matrix algebra. Choosing the tuple which is the identity in one labelled block and zero in every other block gives common final-label selectors. Apply Theorem 24.4.33. $\square$

Definition 24.4.35 (One fusion layer of the topological projector). [Lean](#) [Lean](#) [✓ Stmt](#) Let P_γ be operators on the terminal bond spaces. Define one fusion layer of the recursive operator Q by

$$Q_{\alpha,\beta} = \bigoplus_\gamma \chi_{\alpha,\beta,\gamma} \otimes P_\gamma.$$

Its transport to the product bond space is

$$\widehat{Q}_{\alpha,\beta} = U_{\alpha,\beta}^\dagger Q_{\alpha,\beta} U_{\alpha,\beta}.$$

Theorem 24.4.36 (Projection property of one fusion layer). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Let P_γ be self-adjoint idempotents on the terminal bond spaces. Suppose that the matrices $\chi_{\alpha,\beta,\gamma}$ have the positive-length trace-power form of the label-coefficient family and that this family is independent of the positive chain length. Then

$$Q_{\alpha,\beta} = \bigoplus_\gamma 1_{r_{\alpha,\beta}^\gamma} \otimes P_\gamma$$

is self-adjoint and idempotent. If the labelled tensors form a basis of normal tensors, then $\widehat{Q}_{\alpha,\beta}$ is an orthogonal projection. This is the single recursive fusion step in [CPGSV16, lines 999–1010].

Proof. ✓ Proof The trace-power hypothesis and length independence give $\chi_{\alpha,\beta,\gamma} = 1$. Hence

$$Q_{\alpha,\beta}^\dagger = \bigoplus_{\gamma} (1 \otimes P_\gamma^\dagger) = Q_{\alpha,\beta}, \quad Q_{\alpha,\beta}^2 = \bigoplus_{\gamma} (1 \otimes P_\gamma^2) = Q_{\alpha,\beta}.$$

By Theorem 24.4.34, $U_{\alpha,\beta} U_{\alpha,\beta}^\dagger = 1$. Therefore

$$\widehat{Q}_{\alpha,\beta}^\dagger = \widehat{Q}_{\alpha,\beta}, \quad \widehat{Q}_{\alpha,\beta}^2 = U_{\alpha,\beta}^\dagger Q_{\alpha,\beta} (U_{\alpha,\beta} U_{\alpha,\beta}^\dagger) Q_{\alpha,\beta} U_{\alpha,\beta} = \widehat{Q}_{\alpha,\beta}.$$

□

Theorem 24.4.37 (Left iterated fusion completeness). Lean ✓ Stmt *Under the hypotheses of Theorem 24.4.33,*

$$U_{\alpha,\beta,\gamma}^L (U_{\alpha,\beta,\gamma}^L)^\dagger = 1.$$

Proof. ✓ Proof Pair fusion completeness gives

$$\sum_{\delta} r_{a,b}^\delta D_\delta = D_a D_b.$$

Applying this equality first to (δ, γ) and then to (α, β) shows that the source and target dimensions of $U_{\alpha,\beta,\gamma}^L$ coincide. A square isometry is surjective. □

Theorem 24.4.38 (Right iterated fusion completeness). Lean ✓ Stmt *Under the same hypotheses,*

$$U_{\alpha,\beta,\gamma}^R (U_{\alpha,\beta,\gamma}^R)^\dagger = 1.$$

Proof. ✓ Proof The right-associated dimension calculation is

$$\sum_{\delta,\varepsilon} r_{\beta,\gamma}^\delta r_{\alpha,\delta}^\varepsilon D_\varepsilon = D_\alpha (D_\beta D_\gamma).$$

The right iterated fusion map is therefore a square isometry. □

Theorem 24.4.39 (Right adjoint inverse for the full triple-fusion comparison). Lean ✓ Stmt *Under the same hypotheses, the comparison of the full direct sums satisfies*

$$C_{\alpha,\beta,\gamma} C_{\alpha,\beta,\gamma}^\dagger = 1.$$

Proof. ✓ Proof The left range identity reduces the product to $U^L (U^L)^\dagger$, which is the identity by Theorem 24.4.37. □

Theorem 24.4.40 (Left adjoint inverse for the full triple-fusion comparison). Lean ✓ Stmt *Under the same hypotheses,*

$$C_{\alpha,\beta,\gamma}^\dagger C_{\alpha,\beta,\gamma} = 1.$$

This proves invertibility on the full direct sums. It does not yet prove invertibility of a fixed-final multiplicity matrix.

Proof. ✓ Proof Apply the right range identity and Theorem 24.4.38. □

Theorem 24.4.41 (Vanishing between distinct final sectors). [Lean](#) [✓ Stmt](#) Suppose the positive trace-power coefficients are independent of the positive chain length and common word selectors exist. If $\varepsilon \neq \varepsilon'$, then the corner of the full comparison from the right final sector ε' to the left final sector ε vanishes:

$$P_\varepsilon^L C_{\alpha,\beta,\gamma} P_{\varepsilon'}^R = 0.$$

The assertion is conditional on the displayed word-selector identities. It neither states that a diagonal corner is invertible nor identifies one with an F -matrix, and it does not assert the pentagon identity.

Proof. [✓ Proof](#) Fix one left multiplicity index and one right multiplicity index. Taking the corresponding bond-space corner of the full letterwise identity gives

$$M_\varepsilon^{ij} X = X M_{\varepsilon'}^{ij}.$$

Induction on the word w gives $M_\varepsilon^w X = X M_\varepsilon^w$. Multiply these identities by $c_\varepsilon(w)$ and sum over all words of length S . The two selector identities give $X = 0$. Since the multiplicity indices were arbitrary, the whole off-diagonal final-sector corner vanishes. $\square$

Definition 24.4.42 (Fixed-final multiplicity spaces). [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) Fix a final label ε . The multiplicity spaces of the two bracketings are

$$\mathcal{H}_\varepsilon^L = \bigoplus_{\delta} \mathbb{C}^{\dim \chi_{\alpha,\beta,\delta}} \otimes \mathbb{C}^{\dim \chi_{\delta,\gamma,\varepsilon}}, \quad \mathcal{H}_\varepsilon^R = \bigoplus_{\delta} \mathbb{C}^{\dim \chi_{\beta,\gamma,\delta}} \otimes \mathbb{C}^{\dim \chi_{\alpha,\delta,\varepsilon}}.$$

These spaces have the bracketing pattern of the (e, μ, ν) and (f, λ, σ) index spaces in the printed- F row/column convention of [BMW⁺17, Section 3.4]. Identifying the dimensions of the positive diagonal matrices with the fusion multiplicities of that equation requires the length-independent integer specialization and is not asserted here. The corresponding fixed-final row spaces are $\mathcal{H}_\varepsilon^L \otimes \mathbb{C}^{D_\varepsilon}$ and $\mathcal{H}_\varepsilon^R \otimes \mathbb{C}^{D_\varepsilon}$.

Theorem 24.4.43 (Equality of fixed-final multiplicity dimensions). [Lean](#) [✓ Stmt](#) Suppose that the positive trace-power coefficients are independent of the positive chain length and that the labelled operators are eventually linearly independent. Then, for each final label ε ,

$$\dim \mathcal{H}_\varepsilon^L = \sum_{\delta} \dim \chi_{\alpha,\beta,\delta} \dim \chi_{\delta,\gamma,\varepsilon} = \sum_{\delta} \dim \chi_{\beta,\gamma,\delta} \dim \chi_{\alpha,\delta,\varepsilon} = \dim \mathcal{H}_\varepsilon^R.$$

Consequently a matrix between the two fixed-final multiplicity spaces is square. This is the numerical associativity statement of [BMW⁺17, lines 237–241]; it does not assert completeness of the fusion maps or invertibility of the fixed-final comparison.

Proof. [✓ Proof](#) Choose a positive length beyond the linear-independence threshold. Associativity of the same-length product law gives equality of the two sums of products of structure coefficients. Length independence and positivity identify every coefficient with the size of its corresponding diagonal χ -matrix. The two sums are precisely the cardinalities of the displayed dependent sums. $\square$

Definition 24.4.44 (Triple-fusion maps with fixed final label). [Lean](#) [Lean](#) [✓ Stmt](#) The maps $U_{\alpha,\beta,\gamma;\varepsilon}^L$ and $U_{\alpha,\beta,\gamma;\varepsilon}^R$ are obtained by restricting the two iterated fusion isometries to the summands with final label ε . Their adjoints, after the canonical reassociation of the triple bond space, have the orientation and bracketing pattern of the weighted analogues of the two sides in the printed- F row/column convention of [BMW⁺17, Section 3.4].

Theorem 24.4.45 (Left triple fusion at a fixed final label). [Lean](#) [✓ Stmt](#) For every final label ε , the left-associated restriction satisfies the positive-diagonal weighted identity

$$U_{\alpha,\beta,\gamma;\varepsilon}^L((M_\alpha M_\beta)M_\gamma)^{ij}(U_{\alpha,\beta,\gamma;\varepsilon}^L)^\dagger = \bigoplus_{\delta} \chi_{\alpha,\beta,\delta} \otimes (\chi_{\delta,\gamma,\varepsilon} \otimes M_\varepsilon^{ij}).$$

Proof. [✓ Proof](#) Restrict Theorem 24.4.24 to the rows and columns whose final label is ε . The outer block-diagonal matrix then retains precisely the displayed ε -blocks. $\square$

Theorem 24.4.46 (Right triple fusion at a fixed final label). [Lean](#) [✓ Stmt](#) For every final label ε , the right-associated restriction satisfies the positive-diagonal weighted identity

$$U_{\alpha,\beta,\gamma;\varepsilon}^R(M_\alpha(M_\beta M_\gamma))^{ij}(U_{\alpha,\beta,\gamma;\varepsilon}^R)^\dagger = \bigoplus_{\delta} \chi_{\beta,\gamma,\delta} \otimes (\chi_{\alpha,\delta,\varepsilon} \otimes M_\varepsilon^{ij}).$$

Proof. [✓ Proof](#) Restrict Theorem 24.4.27 to the rows and columns whose final label is ε . The outer block-diagonal matrix then retains precisely the displayed ε -blocks. $\square$

Theorem 24.4.47 (Injective fixed-final intertwiners). [Lean](#) [✓ Stmt](#) Suppose that M_ε is injective. Let C be a matrix from the right fixed-final-sector space to the left fixed-final-sector space. Under the canonical identifications with $\mathcal{H}_\varepsilon^L \otimes \mathbb{C}^{D_\varepsilon}$ and $\mathcal{H}_\varepsilon^R \otimes \mathbb{C}^{D_\varepsilon}$, assume that, for every pair of physical indices i, j ,

$$(1_{\mathcal{H}_\varepsilon^L} \otimes M_\varepsilon^{ij})C = C(1_{\mathcal{H}_\varepsilon^R} \otimes M_\varepsilon^{ij}).$$

Then there is a rectangular matrix $F : \mathcal{H}_\varepsilon^R \rightarrow \mathcal{H}_\varepsilon^L$ such that

$$C = F \otimes 1_{D_\varepsilon}.$$

This conclusion is conditional on the supplied matrix C . It does not assert the existence or invertibility of C or F , and it does not identify the dimensions of the positive diagonal matrices χ with the fusion multiplicities in the printed- F row/column convention of [BMW⁺17, Section 3.4].

Proof. [✓ Proof](#) Apply Lemma 2.1.6 to the family $\{M_\varepsilon^{ij}\}_{i,j}$. This family spans the full matrix algebra by injectivity, and the assumed relation is precisely the amplified intertwining identity of that lemma. $\square$

Theorem 24.4.48 (Conditional fixed-final Kronecker form and separation). [Lean](#) [✓ Stmt](#) Suppose the positive trace-power coefficients are independent of the positive chain length and common word selectors separate the final labels. Fix labels $\alpha, \beta, \gamma, \varepsilon$, and suppose that M_ε is injective at the present blocking. Write

$$C_{\alpha,\beta,\gamma}^{\varepsilon,\varepsilon'} = P_\varepsilon^L C_{\alpha,\beta,\gamma} P_{\varepsilon'}^R$$

for the corner from the right final sector ε' to the left final sector ε . There is a rectangular matrix

$$F_\varepsilon : \mathcal{H}_\varepsilon^R \rightarrow \mathcal{H}_\varepsilon^L$$

such that, after the canonical index identifications,

$$C_{\alpha,\beta,\gamma}^{\varepsilon,\varepsilon} = F_\varepsilon \otimes 1_{D_\varepsilon}.$$

For every $\varepsilon' \neq \varepsilon$,

$$C_{\alpha,\beta,\gamma}^{\varepsilon,\varepsilon'} = 0.$$

The selector identities and injectivity at the present blocking are additional hypotheses here. The theorem neither asserts that F_ε is square or invertible nor identifies it with the printed F -matrix of [BMW⁺17, Section 3.4]; no pentagon identity is asserted.

Proof. ✓ Proof Taking the ε - ε corner of the full comparison identity gives, for every pair i, j ,

$$(1_{\mathcal{H}_\varepsilon^L} \otimes M_\varepsilon^{ij})C^{\varepsilon, \varepsilon} = C^{\varepsilon, \varepsilon}(1_{\mathcal{H}_\varepsilon^R} \otimes M_\varepsilon^{ij}).$$

Injectivity and the amplified commutant result give $C^{\varepsilon, \varepsilon} = F_\varepsilon \otimes 1_{D_\varepsilon}$. The common word selectors give $C^{\varepsilon, \varepsilon'} = 0$ whenever $\varepsilon' \neq \varepsilon$ by Theorem 24.4.41. $\square$

Theorem 24.4.49 (Unitarity of each fixed-final comparison corner). Lean Lean ✓ Stmt Suppose the positive trace-power coefficients are independent of the positive chain length and common selectors exist at a positive length. For every final label ε , write

$$C_\varepsilon = P_\varepsilon^L C_{\alpha, \beta, \gamma} P_\varepsilon^R.$$

Then

$$C_\varepsilon C_\varepsilon^\dagger = 1, \quad C_\varepsilon^\dagger C_\varepsilon = 1.$$

The conclusion is conditional on the simultaneous selector identities. It is the fixed-output-channel invertibility consequence of the simultaneous inverse used in [BMW⁺17, lines 247–277].

Proof. ✓ Proof Order each full direct sum first by its final label. In the product $CC^\dagger$, the ε -diagonal corner is

$$\sum_{\varepsilon'} C^{\varepsilon, \varepsilon'} (C^{\varepsilon, \varepsilon'})^\dagger.$$

Every term with $\varepsilon' \neq \varepsilon$ vanishes, while $CC^\dagger = 1$. Hence $C_\varepsilon C_\varepsilon^\dagger = 1$. Taking instead the ε -diagonal corner of $C^\dagger C = 1$ gives the second identity. $\square$

Theorem 24.4.50 (Unitarity of the fixed-final multiplicity matrix). Lean ✓ Stmt Assume the hypotheses of Theorem 24.4.49. Suppose in addition that M_ε is injective at the present blocking and that $D_\varepsilon > 0$. There is a matrix

$$F_\varepsilon : \mathcal{H}_\varepsilon^R \longrightarrow \mathcal{H}_\varepsilon^L$$

such that

$$C_\varepsilon = F_\varepsilon \otimes 1_{D_\varepsilon}, \quad F_\varepsilon F_\varepsilon^\dagger = 1, \quad F_\varepsilon^\dagger F_\varepsilon = 1.$$

Thus F_ε is invertible. Its orientation is opposite to the printed- F row/column convention of [BMW⁺17, Section 3.4]; no pentagon identity is asserted.

Proof. ✓ Proof Injectivity gives $C_\varepsilon = F_\varepsilon \otimes 1_{D_\varepsilon}$. Substitute this expression into the two identities for C_ε :

$$(F_\varepsilon F_\varepsilon^\dagger) \otimes 1_{D_\varepsilon} = 1, \quad (F_\varepsilon^\dagger F_\varepsilon) \otimes 1_{D_\varepsilon} = 1.$$

Since $D_\varepsilon > 0$, evaluation at any bond basis vector cancels the identity factor and gives the two asserted identities. $\square$

Theorem 24.4.51 (Simultaneous inverse from a one-letter product-algebra span). Lean Lean

✓ Stmt Let $A_c^q \in M_{D_c}(\mathbb{C})$ be a finite labelled tensor family. If the one-letter tuples

$$q \mapsto (A_c^q)_c$$

span the full product algebra $\prod_c M_{D_c}(\mathbb{C})$, then there are coefficients $C_{(c, x, y), q}$ such that

$$\sum_q C_{(c, x, y), q} (A_d^q)_{x' y'} = \delta_{cd} \delta_{xx'} \delta_{yy'}.$$

For an MPO tensor the letter is the ket–bra pair $q = (i, k)$, so the same coefficients form the common block-letter left inverse used in [BMW⁺17, lines 269–277]. The full product-algebra span is obtained after the common blocking described in [BMW⁺17, lines 427–431].

Proof. ✓ **Proof** Fix (c, x, y) and consider the product-algebra element which is the matrix unit E_{xy} in block c and zero in every other block. By the spanning hypothesis it is a linear combination of the one-letter tuples. Reading its (x', y') entry in block d gives the displayed identity. Reindexing the doubled physical letter by the ket–bra pair gives the MPO form. $\square$




Figure 24.4: The fixed-final comparison in the graphical order of [BMW⁺17, equation (pentagon3) and lines 269–277]. At a positive common length, the simultaneous word selector is the finite-word counterpart of the inverse B_d^+ in the source: it retains the chosen final label and annihilates every other one. Thus the comparison has no off-diagonal final-label corners. On the surviving corner, injectivity gives $C_\varepsilon = F_\varepsilon \otimes 1_{D_\varepsilon}$; two-sided unitarity of C_ε and $D_\varepsilon > 0$ give two-sided unitarity of F_ε . This is the associativity mechanism described in [CPGSV16, lines 995–1010]. No pentagon coherence identity is asserted here.

Definition 24.4.52 (Complete zipper fusion family). Lean ✓ **Stmnt** Let Λ be a finite set. A *complete zipper fusion family* consists of positive bond dimensions D_a , MPO blocks $B_a^{ij} \in M_{D_a}(\mathbb{C})$, fusion multiplicities N_{ab}^c , and matrices

$$X_{ab, \mu}^c : \mathbb{C}^{D_c} \longrightarrow \mathbb{C}^{D_a} \otimes \mathbb{C}^{D_b}, \quad X_{ab, \mu}^{c+} : \mathbb{C}^{D_a} \otimes \mathbb{C}^{D_b} \longrightarrow \mathbb{C}^{D_c}.$$

The matrices $X_{ab, \mu}^{c+}$ are independently chosen left inverses, not adjoints. For all labels and multiplicity indices,

$$X_{ab, \nu}^{d+} X_{ab, \mu}^c = \delta_{cd} \delta_{\mu\nu} 1_{D_c}.$$

The opposite product $P_{ab} = \sum_{c, \mu} X_{ab, \mu}^c X_{ab, \mu}^{c+}$ is the projector onto the product-MPO support; it need not be the identity on the ambient bond space. Every product-MPO letter is reconstructed on this support:

$$B_{ab}^{ik} = \sum_{c, \mu} X_{ab, \mu}^c B_c^{ik} X_{ab, \mu}^{c+}.$$

Writing $B_{ab}^{ik} = \sum_j B_a^{ij} \otimes B_b^{jk}$, the two zipper identities are

$$\begin{aligned} B_{ab}^{ik} X_{ab, \mu}^c &= X_{ab, \mu}^c B_c^{ik}, \\ X_{ab, \mu}^{c+} B_{ab}^{ik} &= B_c^{ik} X_{ab, \mu}^{c+}. \end{aligned}$$

Each block is injective, and the labelled block-letter matrix has a common left inverse: its contraction with B_d^{ij} is $\delta_{cd}E_{xy}$ on the block selected by (c, x, y) . These are the fusion-tensor, reconstruction, zipper, and simultaneous inverse relations of [BMW⁺17, lines 161, 181–200, and 269–277].

Theorem 24.4.53 (Complete zipper family from a source BNT). Lean Lean Lean ✓ Stmt
Suppose that the labelled tensors of a length-independent fusion family form a source basis of normal tensors. There is one common positive blocking at which they determine a complete zipper fusion family. Its bond dimensions are the original labelled bond dimensions, its fusion multiplicities are

$$N_{\alpha\beta}^\gamma = \dim \chi_{\alpha,\beta,\gamma},$$

its synthesis map is the adjoint of $U_{\alpha,\beta}$, and its analysis map is $U_{\alpha,\beta}$, after the canonical identification $\mathbb{C}^{D_\alpha D_\beta} \cong \mathbb{C}^{D_\alpha} \otimes \mathbb{C}^{D_\beta}$. In particular, the two zipper identities, literal reconstruction, injectivity, and the simultaneous block-letter inverse are conclusions; none is an additional hypothesis.

Proof. ✓ Proof Length independence makes every chi matrix the identity. Use the common positive BNT blocking for the labelled tensors and transport the fusion equation through it. Pair completeness gives $U_{\alpha,\beta}U_{\alpha,\beta}^\dagger = 1$, so analysis followed by synthesis is the identity. The other three matrix identities are the two transported zipper equations and the reconstruction equation. The common one-letter product-algebra span supplies both block injectivity and the simultaneous block-letter inverse. These data are exactly those of Definition 24.4.52. $\square$

Theorem 24.4.54 (Printed F-move and inverse). Lean Lean Lean ✓ Stmt *Fix $a, b, c, d \in \Lambda$. There is an invertible matrix*

$$F_d^{abc} : \bigoplus_e \mathbb{C}^{N_{ab}^e} \otimes \mathbb{C}^{N_{ec}^d} \longrightarrow \bigoplus_f \mathbb{C}^{N_{bc}^f} \otimes \mathbb{C}^{N_{af}^d}$$

whose rows are indexed by (f, λ, σ) and whose columns are indexed by (e, μ, ν) . Its entries satisfy

$$(X_{ab,\mu}^e \otimes 1)X_{ec,\nu}^d = \sum_{f,\lambda,\sigma} (F_d^{abc})_{e\mu\nu}^{f\lambda\sigma} (1 \otimes X_{bc,\lambda}^f)X_{af,\sigma}^d.$$

The matrix in the opposite orientation is its two-sided inverse. This is the printed F-move of [BMW⁺17, Section 3.4].

Proof. ✓ Proof Form the full left- and right-associated synthesis matrices L and R and their analysis matrices L^+ and R^+ . Applying the reconstruction equation twice to the two associative decompositions gives

$$B_{abc}^{il} = LD_L^{il}L^+ = RD_R^{il}R^+.$$

Contract this equality with the simultaneous inverse of the final block letters and sum the diagonal final-label matrix units. The two direct sums D_L and D_R both contract to the identity. Consequently,

$$P := LL^+ = RR^+, \quad L^+L = R^+R = 1.$$

Hence R^+L changes left-tree coordinates into right-tree coordinates and

$$R(R^+L) = (RR^+)L = PL = (LL^+)L = L.$$

Its opposite-oriented comparison is L^+R . Using the same support identity,

$$(R^+L)(L^+R) = R^+PR = 1, \quad (L^+R)(R^+L) = L^+PL = 1.$$

$$(R^+L)_d = F_d^{abc} \otimes 1_{D_d}.$$

Restricting $R(R^+L) = L$ to this corner gives the displayed F -move. The corresponding fixed-final corner of L^+R , together with $(R^+L)(L^+R) = 1 = (L^+R)(R^+L)$, gives the two inverse identities. $\square$




Figure 24.5: The source-oriented F -move. The left-associated fusion tensor is expanded in the right-associated fusion basis. The upper indices (f, λ, σ) label the rows of F_d^{abc} , and the lower indices (e, μ, ν) label its columns. This is the printed- F row/column convention used here for [BMW⁺17, Section 3.4].

Definition 24.4.55 (Fourfold complete zipper fusion data). Fix labels a, b, c, d, e . The five fusion trees in equation (33) of [BMW+17] determine five multiplicity spaces and five synthesis maps into the fourfold bond space. These spaces use the integer multiplicities N_{ab}^c of the complete zipper family, rather than the weighted positive-diagonal coordinates introduced earlier.

Each edge of the pentagon carries the printed F -matrix for the affected triple tensor factor and the identity on the untouched multiplicity factor. The three upper edges and the two lower edges define composites in the forward orientation. Replacing each edge by its inverse defines the opposite-oriented composites whose entries occur literally in equation (33).

Theorem 24.4.56 (Complete zipper fusion pentagon). [Lean](#) [Lean](#) [Lean](#) [✓ Stmt](#) *The three-edge and two-edge composites of the forward printed F -matrices are equal. The corresponding composites of the inverse F -matrices are also equal. In the upper/lower convention of [BMW⁺ 17, equation (33)], the latter equality is*

$$\begin{aligned} & \sum_{h,\sigma,\lambda,\omega} (F_g^{abc})_{h\sigma\lambda}^{f\mu\nu} (F_e^{ahd})_{i\omega\kappa}^{g\lambda\rho} (F_i^{bcd})_{j\gamma\delta}^{h\sigma\omega} \\ &= \sum_{\tau} (F_e^{fcd})_{j\gamma\tau}^{g\nu\rho} (F_e^{abj})_{i\delta\kappa}^{f\mu\tau} \end{aligned}$$

where every displayed entry denotes the inverse of the forward matrix in the F -move convention.




Figure 24.6: The five complete zipper fusion trees of [BMW⁺17, Section 3.4, equation (33)]. Each arrow is a printed F -move, with right-tree coordinates indexing rows and left-tree coordinates indexing columns.

Proof. ✓ Proof Apply the elementary three-block F -move along each of the five edges. Both paths then give the same expansion in the right-associated fourfold fusion map. Its analysis map is a left inverse, by three successive applications of fusion-tensor biorthogonality, so the forward composites are equal. The reverse composites are two-sided inverses of this common matrix, hence they coincide. Taking a matrix entry gives the displayed identity. $\square$

Theorem 24.4.57 (Coherence of four bond-coordinate reassociations). Lean Lean ✓ Stmt Fix four labels $\alpha, \beta, \gamma, \delta$ in a BNT fusion-isometry family, and write

$$A = \{1, \dots, D_\alpha\}, \quad B = \{1, \dots, D_\beta\}, \quad C = \{1, \dots, D_\gamma\}, \quad D = \{1, \dots, D_\delta\}$$

for their bond-coordinate sets. Let

$$r_{X,Y,Z} : ((X \times Y) \times Z) \longrightarrow X \times (Y \times Z)$$

denote the canonical reassociation. Define

$$\begin{aligned} r_1 &= r_{A,B,C} \times 1_D, & r_2 &= r_{A,B \times C,D}, & r_3 &= 1_A \times r_{B,C,D}, \\ s_1 &= r_{A \times B,C,D}, & s_2 &= r_{A,B,C \times D}. \end{aligned}$$

The three-edge and two-edge reassociations from $((A \times B) \times C) \times D$ to $A \times (B \times (C \times D))$ coincide:

$$r_3 \circ r_2 \circ r_1 = s_2 \circ s_1.$$

If $P(r)_{x,y} = 1$ when $r(x) = y$ and is zero otherwise, then the corresponding coordinate-permutation matrices satisfy

$$P(r_1)P(r_2)P(r_3) = P(s_1)P(s_2).$$

These are equalities of Cartesian bond-coordinate identifications only; they are distinct from the fusion-multiplicity identity in Theorem 24.4.56.

Proof. ✓ Proof Every element $((a, b), c, d)$ is sent by either composite to $(a, (b, (c, d)))$. The first equality follows. The identity $P(v \circ u) = P(u)P(v)$ then turns it into the stated matrix equality. $\square$




Figure 24.7: The five parenthesizations of four bond-coordinate factors. The upper path is the three-edge composite $r_3 \circ r_2 \circ r_1$, and the lower path is the two-edge composite $s_2 \circ s_1$. This figure depicts only canonical Cartesian reassociation. It does not contain fusion multiplicity spaces or F -matrices.

Bibliography

- [And79] T. Ando. Concavity of certain maps on positive definite matrices and applications to Hadamard products. *Linear Algebra and Its Applications*, 26:203–241, 1979.
- [Bha97] Rajendra Bhatia. *Matrix Analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer, New York, 1997.
- [BMW⁺17] N. Bultinck, M. Mariën, D. J. Williamson, M. B. Şahinoğlu, J. Haegeman, and F. Verstraete. Anyons and matrix product operator algebras. *Ann. Phys.*, 378:183–233, March 2017.
- [Car10] Eric A. Carlen. Trace inequalities and quantum entropies: an introductory course. In *Entropy and the quantum*, volume 529 of *Contemporary Mathematics*, pages 73–140. American Mathematical Society, Providence, RI, 2010.
- [CGW11] Xie Chen, Zheng-Cheng Gu, and Xiao-Gang Wen. Classification of gapped symmetric phases in one-dimensional spin systems. *Physical Review B*, 83(3):035107, 2011.
- [CPGSV16] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete. Matrix product density operators: Renormalization fixed points and boundary theories. Preprint; published version: *Ann. Phys.* **378** (2017) 100–149, 2016.
- [CPGSV17] J. Ignacio Cirac, David Pérez-García, Norbert Schuch, and Frank Verstraete. Matrix product unitaries: Structure, symmetries, and topological invariants. *Journal of Statistical Mechanics: Theory and Experiment*, 2017(8):083105, 2017.
- [CPGSV21] J. Ignacio Cirac, David Pérez-García, Norbert Schuch, and Frank Verstraete. Matrix product states and projected entangled pair states: Concepts, symmetries, theorems. *Reviews of Modern Physics*, 93(4):045003, 2021.
- [EHK78] David E. Evans and Raphael Høegh-Krohn. Spectral properties of positive maps on C^* -algebras. *Journal of the London Mathematical Society*, 17(2):345–355, 1978.
- [Fan49] Ky Fan. On a theorem of Weyl concerning eigenvalues of linear transformations. I. *Proceedings of the National Academy of Sciences of the United States of America*, 35(11):652–655, 1949.
- [FGSW⁺15] Carlos Fernández-González, Norbert Schuch, Michael M. Wolf, J. Ignacio Cirac, and David Pérez-García. Frustration free gapless Hamiltonians for Matrix Product States. *Communications in Mathematical Physics*, 333:299–333, 2015.

- [FNW92] M. Fannes, B. Nachtergaele, and R. F. Werner. Finitely correlated states on quantum spin chains. *Communications in Mathematical Physics*, 144(3):443–490, 1992.
- [Hay06] Masahito Hayashi. *Quantum Information: An Introduction*. Springer, Berlin, Heidelberg, 2006.
- [HJPW04] Patrick Hayden, Richard Jozsa, Dénes Petz, and Andreas Winter. Structure of states which satisfy strong subadditivity of quantum entropy with equality. *Communications in Mathematical Physics*, 246(2):359–374, 2004.
- [HP82] Frank Hansen and Gert K. Pedersen. Jensen’s inequality for operators and lõwner’s theorem. *Mathematische Annalen*, 258(3):229–241, 1982.
- [Jac09] Nathan Jacobson. *Basic Algebra II*. Dover Publications, 2009. Unaltered reprint of the 2nd edition (W. H. Freeman, 1989).
- [KL18] Michael J. Kastoryano and Angelo Lucia. Divide and conquer method for proving gaps of frustration free Hamiltonians. *Journal of Statistical Mechanics: Theory and Experiment*, 2018(3):033105, 2018.
- [Lie73] Elliott H. Lieb. Convex trace functions and the Wigner–Yanase–Dyson conjecture. *Advances in Mathematics*, 11(3):267–288, 1973.
- [LR73] Elliott H. Lieb and Mary Beth Ruskai. Proof of the strong subadditivity of quantum-mechanical entropy. *Journal of Mathematical Physics*, 14(12):1938–1941, 1973.
- [Nac96] Bruno Nachtergaele. The spectral gap for some spin chains with discrete symmetry breaking. *Communications in Mathematical Physics*, 175(3):565–606, 1996.
- [PGVWC07] D. Pérez-García, F. Verstraete, M. M. Wolf, and J. I. Cirac. Matrix product state representations. *Quantum Information & Computation*, 7(5):401–430, 2007.
- [PGWS⁺08] D. Pérez-García, M. M. Wolf, M. Sanz, F. Verstraete, and J. I. Cirac. String order and symmetries in quantum spin lattices. *Physical Review Letters*, 100(16):167202, 2008.
- [Rus02] Mary Beth Ruskai. Inequalities for quantum entropy: A review with conditions for equality. *Journal of Mathematical Physics*, 43(9):4358–4375, 2002.
- [SPGC11] Norbert Schuch, David Pérez-García, and Ignacio Cirac. Classifying quantum phases using matrix product states and projected entangled pair states. *Physical Review B*, 84(16):165139, 2011.
- [SPGWC10] M. Sanz, D. Pérez-García, M. M. Wolf, and J. I. Cirac. A quantum version of Wielandt’s inequality. *IEEE Transactions on Information Theory*, 56(9):4668–4673, 2010.
- [TT88] Kotaro Tanahashi and Jun Tomiyama. Indecomposable positive maps in matrix algebras. *Canadian Mathematical Bulletin*, 31(3):308–317, 1988.
- [Wol12] Michael M. Wolf. Quantum channels & operations: Guided tour. Lecture notes, 2012. Originally distributed via TU Munich foswiki; archival PDF at <https://mediatum.ub.tum.de/doc/1099654/1099654.pdf>.

- [Yam93] Shigeru Yamagami. Cyclic inequalities. *Proceedings of the American Mathematical Society*, 118(2):521–527, 1993.