

Fundamental Theorem of Matrix Product States

A Formalization Blueprint

Authors

March 9, 2026

Contents

1	Introduction	3
2	Matrix Product Vectors	4
2.1	Basic definitions	4
2.2	Gauge equivalence and same MPV	5
2.3	Injectivity and normality	6
2.4	Canonical form	6
2.5	Blocking	7
2.6	MPV overlap	8
3	The Single-Block Fundamental Theorem	9
3.1	Trace pairing	9
3.2	Linear extension	10
3.3	Simplicity and Skolem–Noether	10
3.4	Assembly	11
4	Quantum Channels and Positive Maps	12
4.1	Positive maps and channels	12
4.2	Kadison–Schwarz inequality	13
4.3	Multiplicative domain	14
4.4	Irreducibility	15
4.5	Cesàro fixed-point theorem	15
4.6	Fixed-point projection	16
4.7	Peripheral spectrum and primitivity	16
4.7.1	Periodicity removal by powering	17
4.7.2	Peripheral closure via adjoint fixed point	18
5	Quantum Perron–Frobenius Theory	20
5.1	Positive definiteness	20
5.2	Uniqueness	21
5.3	Existence and assembly	21
5.4	Unital and trace-preserving gauges	21
6	Spectral Gap and Block Separation	23
6.1	Mixed transfer operator	23
6.2	MPV overlap as transfer trace	24
6.3	Eigenvalue bound and spectral gap	25
6.4	Spectral gap — rectangular case	25

6.5	MPV overlap decay	26
6.6	Block separation	26
6.7	Primitive overlap convergence	26
7	Wielandt Bound	28
7.1	Cumulative span	28
7.2	Fitting decomposition	29
7.3	Nonzero trace product	29
7.4	Eigenvector extraction	30
7.5	Eigenvector spreading	30
7.6	Assembly	30
7.7	Lemma 2(b): fixed-length matrix spanning	31
7.8	Blocked tensor and rectangular span	32
7.9	Primitivity and normality	33
7.9.1	Primitivity-to-normality bridge	34
7.9.2	Primitive implies irreducible	34
7.10	Cumulative span to word span	35
8	Canonical Form Reduction	36
8.1	Block-diagonal assembly	36
8.2	Transfer map normalization	36
8.3	Invariant subspace decomposition	37
8.4	Iterated reduction	38
8.4.1	Burnside bridge: irreducibility and spanning	38
8.5	Proportional single-block theorem	39
8.6	Canonical form from primitivity	39
8.7	Canonical form existence pipeline	40
8.8	Perron–Frobenius eigenvector existence	41
8.9	End-to-end pipeline status	42
9	Block Permutation and Separation	44
9.1	Block permutation algebra	44
9.2	Pi-algebra linear extension	45
9.3	Block separation	45
10	BNT and Full Assembly	47
10.1	Basis normal tensors	47
10.2	BNT permutation rigidity	48
10.3	Coefficient convergence	48
10.4	Newton–Girard identities	49
10.5	Primitivity bridge	49
10.6	Full assembly	49

Chapter 1

Introduction

Chapter 2

Matrix Product Vectors

Fix a physical dimension $d \geq 1$ and a bond dimension $D \geq 1$. This chapter studies the tensors that generate matrix product vector (MPV) families. Lean keeps the historical name `MPSTensor` for these tensors, and we follow that name in Lean declaration tags. All tensors in this blueprint are translation-invariant with periodic boundary conditions (PBC).

2.1 Basic definitions

Definition 2.1 (MPV tensor (Lean: MPS tensor)). A (translation-invariant, PBC) *MPV tensor*—called an *MPS tensor* in Lean and much of the literature—with physical dimension d and bond dimension D is a collection of matrices

$$\{A^i\}_{i=0}^{d-1}, \quad A^i \in M_D(\mathbb{C}),$$

indexed by a physical index $i \in \{0, \dots, d-1\}$. Such a tensor defines an MPV family.

Definition 2.2 (Word evaluation). Given a word $w = (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$, the *word evaluation* is the matrix product

$$A^w := A^{i_1} A^{i_2} \dots A^{i_L} \in M_D(\mathbb{C}),$$

with the convention that the empty word evaluates to the identity: $A^\emptyset = \mathbb{1}_D$.

Lemma 2.3 (Word evaluation respects concatenation). *For any words w_1, w_2 , $A^{w_1 \cdot w_2} = A^{w_1} A^{w_2}$.*

Proof. Immediate from the definition of A^w . □

Definition 2.4 (Matrix product vector). The *matrix product vector* (MPV) of a tensor A at system size N is the vector

$$|V^{(N)}(A)\rangle = \sum_{i_1, \dots, i_N=0}^{d-1} \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}) |i_1, \dots, i_N\rangle \in (\mathbb{C}^d)^{\otimes N}.$$

Equivalently, for a configuration $\sigma = (i_1, \dots, i_N) \in \{0, \dots, d-1\}^N$, we write

$$V^{(N)}(A)_\sigma := \text{tr}(A^{i_1} A^{i_2} \dots A^{i_N}).$$

For a general word w , we also write

$$c_w(A) := \text{tr}(A^w),$$

and Lean packages this scalar coefficient function separately as `.tr`. The *MPV family* generated by A is the collection $\mathcal{V}(A) = \{|V^{(N)}(A)\rangle\rangle\}_{N \geq 1}$.

Definition 2.5 (Transfer map). The *transfer map* associated to a tensor A is the linear map $\mathcal{E}_A : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$\mathcal{E}_A(X) = \sum_{i=0}^{d-1} A^i X (A^i)^\dagger.$$

Remark 2.6. The transfer map is automatically positive: if $X \geq 0$ then $\mathcal{E}_A(X) \geq 0$, because each summand $A^i X (A^i)^\dagger$ is positive semidefinite. Lean records this local helper as `.transfer_pos`. The abstract positive-map framework appears in Chapter 4.

2.2 Gauge equivalence and same MPV

Definition 2.7 (Gauge equivalence). Two tensors A and B of the same bond dimension D are *gauge equivalent* if there exists an invertible matrix $X \in \text{GL}_D(\mathbb{C})$ such that

$$B^i = X A^i X^{-1} \quad \text{for all } i \in \{0, \dots, d-1\}.$$

Definition 2.8 (Same MPV). Two tensors A and B (of the same bond dimension) *generate the same MPV family* if $|V^{(N)}(A)\rangle\rangle = |V^{(N)}(B)\rangle\rangle$ for every system size $N \geq 1$.

Definition 2.9 (Same MPV — different bond dimensions). For tensors A and B of possibly different bond dimensions D_1 and D_2 , we write $\mathcal{V}(A) = \mathcal{V}(B)$ if $|V^{(N)}(A)\rangle\rangle = |V^{(N)}(B)\rangle\rangle$ for every $N \geq 1$.

Remark 2.10. Definition 2.8 is the same-bond-dimension specialization of Definition 2.9. We keep both because later Lean declarations use both the homogeneous and heterogeneous forms.

Definition 2.11 (Proportional MPV). Two tensors A and B (of possibly different bond dimensions) generate *proportional MPV families* if for every N there exists $c_N \in \mathbb{C}$ such that $|V^{(N)}(A)\rangle\rangle = c_N |V^{(N)}(B)\rangle\rangle$.

Lemma 2.12 (Word evaluation under scaling). Let ζA denote the tensor with matrices $\{\zeta A^i\}_{i=0}^{d-1}$. Then for any word w ,

$$(\zeta A)^w = \zeta^{|w|} A^w,$$

where $|w|$ is the length of w .

Proof. Induction on w : the base case gives $\zeta^0 = 1$, and each step picks up one factor of ζ . $\square$

Definition 2.13 (Gauge-phase equivalence). Two tensors A and B are *gauge-phase equivalent* if there exist $X \in \text{GL}_D(\mathbb{C})$ and $\zeta \in \mathbb{C} \setminus \{0\}$ such that

$$B^i = \zeta X A^i X^{-1} \quad \text{for all } i.$$

Remark 2.14. After spectral-radius normalization one may restrict ζ to a unit-modulus phase $e^{i\phi}$. We allow an arbitrary nonzero scalar because later canonical-form statements keep the block-scaling factors explicit.

Lemma 2.15 (Word evaluation under conjugation). *If $B^i = X A^i X^{-1}$ for all i , then for every word w ,*

$$B^w = X A^w X^{-1}.$$

Proof. Induction on the word length, using $XX^{-1} = \mathbb{1}$. $\square$

Theorem 2.16 (Gauge invariance of MPVs). *If A and B are gauge equivalent, then they generate the same MPV family.*

Proof. Let $B^i = X A^i X^{-1}$. By Lemma 2.15, $B^w = X A^w X^{-1}$ for every word w . Then $\text{tr}(B^w) = \text{tr}(X A^w X^{-1}) = \text{tr}(A^w)$ by cyclicity of the trace. $\square$

2.3 Injectivity and normality

Definition 2.17 (Injectivity). A tensor A is *injective* if its matrices span the full matrix algebra: $\text{span}_{\mathbb{C}}\{A^i : i = 0, \dots, d-1\} = M_D(\mathbb{C})$.

Definition 2.18 (L -block injectivity). A tensor A is *L -block injective* if $\text{span}_{\mathbb{C}}\{A^{i_1} \dots A^{i_L} : (i_1, \dots, i_L) \in \{0, \dots, d-1\}^L\} = M_D(\mathbb{C})$. An injective tensor is 1-block injective.

Definition 2.19 (Normal tensor). A tensor A is *normal* [CPGSV21] if there exists a finite L_0 such that A is L_0 -block injective.

Remark 2.20. Here “normal” means eventual block injectivity. In [CPGSV21], the NT terminology is spectral: it is phrased in terms of irreducibility and the peripheral spectrum of the transfer map. The bridge to that viewpoint is supplied later by Theorem 8.26, Theorem 7.43, and Theorem 7.50.

2.4 Canonical form

Definition 2.21 (Block-injective canonical form). In this blueprint, a *block-injective canonical form* for a tensor consists of:

1. a number of blocks $r \geq 1$,
2. bond dimensions $D_1, \dots, D_r$,
3. injective tensors $\{A_k^i\}_{i=0}^{d-1}$ with $A_k^i \in M_{D_k}(\mathbb{C})$ for each block $k \in \{1, \dots, r\}$,
4. scaling factors $\mu_1, \dots, \mu_r \in \mathbb{C}$.

The associated tensor is the block-diagonal tensor

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

This is the block-injective version of the canonical-form data used in the Lean development. It is discussed in [PGVWC07] and reviewed in [CPGSV21].

Remark 2.22. This blueprint uses the block-injective (biCF-style) canonical form: each block is injective in the sense of Definition 2.17. This is stronger than the NT-block canonical form of [CPGSV21], where the blocks are only normal in the paper’s spectral sense. The paper-faithful NT/BNT/CFII pipeline is developed later in Chapter 8; the present section records the stronger package that the current Lean development uses downstream.

Definition 2.23 (Block-diagonal tensor from blocks). Given r blocks with bond dimensions $(D_k)_{k=1}^r$, per-block tensors $\{A_k^i\}$, and scaling factors $(\mu_k)_{k=1}^r$, the associated *block-diagonal tensor* is the tensor of total bond dimension $D = \sum_{k=1}^r D_k$ defined by

$$A^i = \bigoplus_{k=1}^r \mu_k A_k^i.$$

Theorem 2.24 (MPV decomposition). *The MPV of a block-diagonal tensor decomposes as*

$$|V^{(N)}(A)\rangle = \sum_{k=1}^r \mu_k^N |V^{(N)}(A_k)\rangle.$$

Proof. For each physical index i , the block-diagonal tensor $A^i = \bigoplus_k \mu_k A_k^i$ is a block-diagonal matrix. By Lemma 2.12, the word evaluation of each block picks up a factor $\mu_k^{|w|}$, and the full word evaluation is block-diagonal with blocks $\mu_k^N A_k^w$. Taking the trace of a block-diagonal matrix gives the sum of the block traces: $\text{tr}(A^w) = \sum_k \mu_k^N \text{tr}(A_k^w)$. $\square$

2.5 Blocking

Definition 2.25 (Blocked tensor). Given a tensor A and a blocking length $L \geq 1$, the *L -blocked tensor* is the tensor with physical dimension d^L and the same bond dimension D , whose matrices are indexed by words $(i_1, \dots, i_L) \in \{0, \dots, d-1\}^L$:

$$(A^{[L]})^{(i_1, \dots, i_L)} = A^{i_1} A^{i_2} \dots A^{i_L}.$$

Remark 2.26 (Blocked word evaluation). If $w = (J_1, \dots, J_m)$ is a word in the blocked alphabet, where each J_k is an L -tuple in $\{0, \dots, d-1\}^L$, then concatenating the block words gives a word $\tilde{w}$ of length mL in the original alphabet and one has $(A^{[L]})^w = A^{\tilde{w}}$. This is the direct compatibility between blocking and word evaluation.

Lemma 2.27 (Blocking preserves MPV coefficients). *For each basis state $\sigma = (\sigma_1, \dots, \sigma_N)$ of the blocked system, where each σ_k is an L -tuple in $\{0, \dots, d-1\}^L$, there exists a basis state $\tilde{\sigma} = (\tilde{\sigma}_1, \dots, \tilde{\sigma}_{NL})$ of the original system such that*

$$V^{(N)}(A^{[L]})_{\sigma} = V^{(NL)}(A)_{\tilde{\sigma}}.$$

Proof. Concatenate the block words and apply the blocked word evaluation lemma (Lemma 2.26). $\square$

Theorem 2.28 (Blocking preserves same MPV). *If A and B generate the same MPV family, then for any blocking length L , the blocked tensors $A^{[L]}$ and $B^{[L]}$ also generate the same MPV family.*

Proof. By Lemma 2.27, each MPV coefficient of the blocked tensor $A^{[L]}$ equals a coefficient of A at the corresponding concatenated configuration. The same holds for $B^{[L]}$ (with the same concatenated configuration). Since A and B agree on all configurations, so do $A^{[L]}$ and $B^{[L]}$. $\square$

2.6 MPV overlap

Definition 2.29 (MPV as Hilbert-space vector). We regard $|V^{(N)}(A)\rangle$ as an element of $\mathbb{C}^{d^N}$ indexed by configurations $\sigma \in \{0, \dots, d-1\}^N$, with σ -component $V^{(N)}(A)_\sigma$ as in Definition 2.4. This allows us to use the standard Euclidean inner product on $\mathbb{C}^{d^N}$.

Definition 2.30 (MPV inner product). The *MPV inner product* between two tensors A and B at system size N is

$$\langle V^{(N)}(A) | V^{(N)}(B) \rangle := \sum_{\sigma \in \{0, \dots, d-1\}^N} \overline{V^{(N)}(A)_\sigma} V^{(N)}(B)_\sigma,$$

i.e. the Euclidean inner product on $\mathbb{C}^{d^N}$ (conjugate-linear in the first argument).

Definition 2.31 (MPV overlap). The *MPV overlap* between two tensors A (of bond dimension D_1) and B (of bond dimension D_2) at system size N is

$$O_{AB}(N) := \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_\sigma \overline{V^{(N)}(B)_\sigma}.$$

This is the bilinear physics-oriented overlap used later. By Lemma 2.32, it is the complex conjugate of the Hilbert-space inner product from Definition 2.30.

Lemma 2.32 (Overlap equals conjugate inner product). $O_{AB}(N) = \overline{\langle V^{(N)}(A) | V^{(N)}(B) \rangle}$.

Proof. The overlap sums $V_\sigma \overline{W_\sigma}$ while the inner product sums $\overline{V_\sigma} W_\sigma$. Complex conjugation swaps these. $\square$

Chapter 3

The Single-Block Fundamental Theorem

This chapter proves that if an injective MPS tensor A generates the same MPV family as another tensor B (of the same bond dimension), then A and B are gauge equivalent. The proof here is purely algebraic. The trace-pairing maps introduced below are auxiliary devices attached to the chosen generating families $\{A^i\}$ and $\{B^i\}$; they are used only to construct the linear map that is later shown to be a gauge transform.

3.1 Trace pairing

Theorem 3.1 (Nondegeneracy of the trace pairing). *For $M \in M_D(\mathbb{C})$, $\text{tr}(MN) = 0$ for all $N \in M_D(\mathbb{C})$ if and only if $M = 0$.*

Proof. Testing against the matrix units E_{ji} (the matrix with 1 in position (j, i) and 0 elsewhere) gives $\text{tr}(E_{ji}M) = M_{ij}$. Hence if $\text{tr}(MN) = 0$ for all N , then every entry of M vanishes. $\square$

Lemma 3.2 (Same MPV implies trace agreement). *If A and B generate the same MPV family, then $\text{tr}(A^w) = \text{tr}(B^w)$ for every word w .*

Proof. Viewing the word w as a configuration of length $|w|$, Definition 2.4 gives the corresponding MPV coefficient as the trace coefficient $c_w(A) = \text{tr}(A^w)$. The same-MPV hypothesis gives equality at every system size. $\square$

Definition 3.3 (Trace pairing map). For an MPS tensor A , attach to the chosen family $\{A^i\}$ the auxiliary linear map $\Phi_A : M_D(\mathbb{C}) \rightarrow \mathbb{C}^d$ defined by

$$\Phi_A(M)_i = \text{tr}(M \cdot A^i).$$

Theorem 3.4 (Injectivity of the trace pairing map). *If A is injective, then $\ker \Phi_A = \{0\}$.*

Proof. If $\Phi_A(M) = 0$, then $\text{tr}(M \cdot A^i) = 0$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, this gives $\text{tr}(MN) = 0$ for all N , so $M = 0$ by Theorem 3.1. $\square$

Lemma 3.5 (Nonvanishing trace on injective tensors). *If $D \geq 1$, A is injective, A and B generate the same MPV family, and $B^i = 0$ for all i , then we have a contradiction.*

Proof. If $B^i = 0$ for all i , then Lemma 3.2 gives $\text{tr}(A^w) = 0$ for every word w , in particular for all length-1 words. Since the $\{A^i\}$ span $M_D(\mathbb{C})$, the trace functional vanishes identically, contradicting $\text{tr}(\mathbb{1}_D) = D \neq 0$. $\square$

3.2 Linear extension

Theorem 3.6 (Existence and uniqueness of the linear extension). *If A is injective and A, B generate the same MPV family, then there exists a unique linear map $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ satisfying $T(A^i) = B^i$ for all i .*

Proof. Write $\ell_A : \mathbb{C}^d \rightarrow M_D(\mathbb{C})$ for the map that sends a coefficient vector c to $\sum_i c_i A^i$ (linear combination with the A^i). Applying Lemma 3.2 to all words of length 2 shows $\Phi_A \circ \ell_A = \Phi_B \circ \ell_B$. Since A is injective, ℓ_A is surjective, giving $\text{range}(\Phi_A) \subseteq \text{range}(\Phi_B)$. By Theorem 3.4, $\ker \Phi_A = \{0\}$, so rank-nullity gives $\dim(\text{range}(\Phi_A)) = \dim(M_D(\mathbb{C}))$. Hence $\dim(\text{range}(\Phi_B)) \geq \dim(M_D(\mathbb{C}))$, and rank-nullity again gives $\ker \Phi_B = \{0\}$. A left inverse g of Φ_B gives $T := g \circ \Phi_A$, and uniqueness follows because the $\{A^i\}$ span $M_D(\mathbb{C})$. $\square$

Theorem 3.7 (Multiplicativity of the linear extension). *The linear extension T from Theorem 3.6 is multiplicative: $T(MN) = T(M)T(N)$ for all $M, N \in M_D(\mathbb{C})$.*

Proof. **Step 1** ($\Phi_B \circ T = \Phi_A$). Applying Lemma 3.2 to words of length 2 gives $\Phi_B(T(A^i)) = \Phi_B(B^i) = \Phi_A(A^i)$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, $\Phi_B \circ T = \Phi_A$.

Step 2 ($T(A^i A^j) = B^i B^j$). Lemma 3.2 applied to the word (i, j, k) gives $\text{tr}(A^i A^j A^k) = \text{tr}(B^i B^j B^k)$ for all i, j, k . So $\Phi_B(T(A^i A^j))_k = \Phi_A(A^i A^j)_k = \text{tr}(A^i A^j A^k) = \text{tr}(B^i B^j B^k) = \Phi_B(B^i B^j)_k$. The same rank-nullity argument used in the proof of Theorem 3.6 gives $\ker \Phi_B = \{0\}$. Hence $T(A^i A^j) = B^i B^j$.

Step 3 (bilinear extension). For fixed j , the maps $M \mapsto T(M \cdot A^j)$ and $M \mapsto T(M) \cdot B^j$ agree on the spanning set $\{A^i\}$, hence on all of $M_D(\mathbb{C})$. Applying this once more with j replaced by a general matrix gives full multiplicativity. $\square$

3.3 Simplicity and Skolem–Noether

Theorem 3.8 (Simplicity of $M_D(\mathbb{C})$). *A nonzero multiplicative $\mathbb{C}$ -linear endomorphism of $M_D(\mathbb{C})$ is bijective.*

Proof. The kernel of a multiplicative linear map is a two-sided ideal. Since $M_D(\mathbb{C})$ is a simple ring (it has no nontrivial two-sided ideals), the kernel is either $\{0\}$ or $M_D(\mathbb{C})$. The latter would force the map to be zero, contradicting the hypothesis. Hence the map is injective, and surjectivity follows from finite-dimensionality. $\square$

Theorem 3.9 (Skolem–Noether). *Every $\mathbb{C}$ -algebra automorphism of $M_D(\mathbb{C})$ is inner: for any $f \in \text{Aut}_{\mathbb{C}\text{-alg}}(M_D(\mathbb{C}))$, there exists $X \in \text{GL}_D(\mathbb{C})$ such that $f(M) = XMX^{-1}$ for all M .*

Proof. Via the canonical identification $M_D(\mathbb{C}) \cong \text{End}_{\mathbb{C}}(\mathbb{C}^D)$, the automorphism f induces an automorphism of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$. Every algebra automorphism of $\text{End}(V)$ is conjugation by an invertible linear map (this is a standard result in the theory of central simple algebras). Translating back to matrices gives the invertible X . $\square$

Lemma 3.10 (Promotion to algebra homomorphism). *If $T : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a multiplicative surjective $\mathbb{C}$ -linear map, then $T(\mathbb{1}) = \mathbb{1}$, and T is a $\mathbb{C}$ -algebra homomorphism.*

Proof. By surjectivity, there exists X with $T(X) = \mathbb{1}$. Then $T(\mathbb{1}) = T(X) \cdot T(\mathbb{1}) = T(X \cdot \mathbb{1}) = T(X) = \mathbb{1}$, using multiplicativity and $T(X) = \mathbb{1}$. With $T(\mathbb{1}) = \mathbb{1}$ established, T preserves the unit and hence is a $\mathbb{C}$ -algebra homomorphism. $\square$

3.4 Assembly

Theorem 3.11 (Single-block Fundamental Theorem). *Let A be an injective MPS tensor and let B be an MPS tensor of the same bond dimension. If A and B generate the same MPV family, then they are gauge equivalent: there exists $X \in \text{GL}_D(\mathbb{C})$ such that $B^i = XA^iX^{-1}$ for all i .*

Proof. By Theorem 3.6, there is a unique linear T with $T(A^i) = B^i$. By Theorem 3.7, T is multiplicative. T is nonzero: if $T = 0$ then $B^i = 0$ for all i , contradicting Lemma 3.5. By Theorem 3.8, T is bijective. By Lemma 3.10, T is a $\mathbb{C}$ -algebra automorphism. Theorem 3.9 gives an invertible X with $T(M) = XMX^{-1}$, and evaluating at $M = A^i$ yields $B^i = XA^iX^{-1}$. $\square$

Chapter 4

Quantum Channels and Positive Maps

This chapter develops the theory of positive maps and quantum channels on matrix algebras, following [Wol12].

4.1 Positive maps and channels

Definition 4.1 (Positive map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *positive* if $E(X) \geq 0$ whenever $X \geq 0$, where $X \geq 0$ means that X is positive semidefinite.

Remark 4.2. We use the Loewner order on matrices: $X \leq Y$ means $Y - X \geq 0$.

Definition 4.3 (Trace-preserving map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *trace-preserving* if $\text{tr}(E(X)) = \text{tr}(X)$ for all X .

Definition 4.4 (Completely positive map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *completely positive* (CP) if it admits a *Kraus representation*: there exist operators $\{K_i\}_{i=0}^{r-1}$ with $K_i \in M_D(\mathbb{C})$ such that

$$E(X) = \sum_{i=0}^{r-1} K_i X K_i^\dagger \quad \text{for all } X \in M_D(\mathbb{C}).$$

We use the Kraus characterisation because it matches the later Kadison–Schwarz arguments. By Choi’s theorem, this is equivalent to $E \otimes \text{id}_n$ being positive for all n .

Theorem 4.5 (CP maps are positive). *Every completely positive map is positive.*

Proof. If $X \geq 0$, then each summand $K_i X K_i^\dagger \geq 0$ (conjugation preserves positive semidefiniteness), so the sum is PSD. $\square$

Definition 4.6 (Quantum channel). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is a *quantum channel* if it is both completely positive and trace-preserving (CPTP), following [Wol12].

Definition 4.7 (Density matrices). The set of *density matrices* in $M_D(\mathbb{C})$ is

$$\mathcal{D}_D = \{\rho \in M_D(\mathbb{C}) : \rho \geq 0, \text{tr}(\rho) = 1\}.$$

Theorem 4.8 (Density matrices are compact). $\mathcal{D}_D$ is compact (in the entry-wise topology on $M_D(\mathbb{C})$).

Proof. The PSD cone is closed (preimage of the closed nonnegative cone under continuous quadratic forms), and if $\rho \geq 0$ with $\text{tr}(\rho) = 1$ then each entry satisfies $|\rho_{ij}|^2 \leq \rho_{ii}\rho_{jj} \leq 1$: the diagonal entries are nonnegative and sum to 1, and the off-diagonal bound is the PSD Cauchy–Schwarz inequality. Hence the matrix entries are uniformly bounded, and Heine–Borel gives compactness. $\square$

Theorem 4.9 (Density matrices are convex). $\mathcal{D}_D$ is convex.

Proof. Convex combination of PSD matrices is PSD, and trace is linear. $\square$

Theorem 4.10 (Density matrices are nonempty). For $D \geq 1$, $\mathcal{D}_D \neq \emptyset$.

Proof. $\frac{1}{D}\mathbb{1}_D$ is a density matrix. $\square$

Theorem 4.11 (Channels preserve density matrices). If E is a quantum channel, then $E(\mathcal{D}_D) \subseteq \mathcal{D}_D$.

Proof. Complete positivity implies positivity (Theorem 4.5), so $E(\rho) \geq 0$; trace preservation gives $\text{tr}(E(\rho)) = \text{tr}(\rho) = 1$. $\square$

4.2 Kadison–Schwarz inequality

A Kraus map is the concrete realisation of a completely positive map (Definition 4.4). The Kadison–Schwarz inequality and the multiplicative domain characterisation are proved for Kraus maps, then applied to transfer maps, which are Kraus maps by construction (Remark 4.25).

Definition 4.12 (Kraus map). Given operators $\{K_i\}_{i=0}^{d-1}$ with $K_i \in M_D(\mathbb{C})$, the *Kraus map* is $E(X) = \sum_{i=0}^{d-1} K_i X K_i^\dagger$. A linear map is completely positive (Definition 4.4) if and only if it can be written in this form.

Definition 4.13 (Adjoint Kraus map). The *adjoint Kraus map* is $E^*(X) = \sum_{i=0}^{d-1} K_i^\dagger X K_i$. When the $\{K_i\}$ are the matrices of an MPS tensor A , the adjoint Kraus map corresponds to the transfer map $\mathcal{E}_A$ with the roles of A^i and $(A^i)^\dagger$ exchanged.

Definition 4.14 (Unital Kraus map). A Kraus map is *unital* if $\sum_{i=0}^{d-1} K_i K_i^\dagger = \mathbb{1}$. Equivalently, $E(\mathbb{1}) = \mathbb{1}$.

Definition 4.15 (Trace-preserving Kraus map). A Kraus map is *trace-preserving* if $\sum_{i=0}^{d-1} K_i^\dagger K_i = \mathbb{1}$. Equivalently, the adjoint Kraus map is unital. This is the standard MPS normalization condition, so later Kadison–Schwarz arguments are often applied to the adjoint map.

Theorem 4.16 (Kadison–Schwarz inequality). Let $E(X) = \sum_i K_i X K_i^\dagger$ be a unital Kraus map (i.e. $\sum_i K_i K_i^\dagger = \mathbb{1}$). Then for all $X \in M_D(\mathbb{C})$,

$$E(X^\dagger X) \geq E(X)^\dagger E(X).$$

This is [Wol12, Eq. (5.2)].

Proof. The 2×2 block matrix $\begin{bmatrix} X^\dagger X & X^\dagger \\ X & \mathbb{1} \end{bmatrix}$ is PSD ($= vv^\dagger$ with $v = [X^\dagger, \mathbb{1}]^T$). Applying the unital CP map blockwise preserves positive semidefiniteness (each block $K_i(\cdot)K_i^\dagger$ preserves PSD, and unitality ensures the $(2, 2)$ -block maps to $\mathbb{1}$). The Schur complement of the $(2, 2)$ -block $\mathbb{1}$ gives the result. $\square$

Theorem 4.17 (Kadison–Schwarz for the adjoint channel). *If $\sum_i K_i^\dagger K_i = \mathbb{1}$ (trace-preserving), then the adjoint Kraus map satisfies $E^*(X^\dagger X) \geq E^*(X)^\dagger E^*(X)$.*

Proof. The adjoint operators $\{K_i^\dagger\}$ satisfy $\sum_i K_i^\dagger (K_i^\dagger)^\dagger = \sum_i K_i^\dagger K_i = \mathbb{1}$, so they form a unital family. Apply Theorem 4.16 to $\{K_i^\dagger\}$. $\square$

Theorem 4.18 (Hilbert–Schmidt contraction). *If a Kraus map is both unital and trace-preserving, then $\text{tr}(E(X)^\dagger E(X)) \leq \text{tr}(X^\dagger X)$ for all $X \in M_D(\mathbb{C})$.*

Proof. By Theorem 4.16, $E(X^\dagger X) - E(X)^\dagger E(X) \geq 0$, so $\text{tr}(E(X^\dagger X)) \geq \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. $\square$

4.3 Multiplicative domain

Theorem 4.19 (KS gap decomposition). *For a unital Kraus map E , the Kadison–Schwarz gap decomposes as*

$$E(X^\dagger X) - E(X)^\dagger E(X) = \sum_{i=0}^{d-1} R_i^\dagger R_i,$$

where $R_i := XK_i^\dagger - K_i^\dagger E(X)$.

Proof. Expand the right-hand side $\sum_i (XK_i^\dagger - K_i^\dagger E(X))^\dagger (XK_i^\dagger - K_i^\dagger E(X))$, distribute, and use unitality $\sum_i K_i K_i^\dagger = \mathbb{1}$ to identify the result with the KS gap. $\square$

Theorem 4.20 (KS equality implies Kraus commutation). *Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$ for some X , then $XK_i^\dagger = K_i^\dagger E(X)$ for all i .*

Proof. By Theorem 4.19, the KS gap is $\sum_i R_i^\dagger R_i$ with $R_i = XK_i^\dagger - K_i^\dagger E(X)$. If the gap vanishes, $\sum_i R_i^\dagger R_i = 0$. Each summand $R_i^\dagger R_i$ is PSD, so each must be zero, giving $R_i = 0$. $\square$

Theorem 4.21 (KS equality for peripheral eigenvectors). *Let E be a Kraus map that is both unital and trace-preserving. If $E(X) = \mu X$ with $|\mu| = 1$, then the KS gap vanishes: $E(X^\dagger X) = E(X)^\dagger E(X)$.*

Proof. The gap $G := E(X^\dagger X) - E(X)^\dagger E(X)$ is PSD by Theorem 4.16. Its trace satisfies $\text{tr}(G) = \text{tr}(E(X^\dagger X)) - \text{tr}(E(X)^\dagger E(X))$. Trace preservation gives $\text{tr}(E(X^\dagger X)) = \text{tr}(X^\dagger X)$. Using $E(X) = \mu X$ and $|\mu| = 1$, $\text{tr}(E(X)^\dagger E(X)) = |\mu|^2 \text{tr}(X^\dagger X) = \text{tr}(X^\dagger X)$. So $\text{tr}(G) = 0$, and a PSD matrix with zero trace must be zero. $\square$

Theorem 4.22 (One-sided multiplicative-domain identity). *Let E be a unital Kraus map. If $E(X^\dagger X) = E(X)^\dagger E(X)$, then $E(X^\dagger Y) = E(X)^\dagger E(Y)$ for all $Y \in M_D(\mathbb{C})$.*

Proof. From Theorem 4.20, $XK_i^\dagger = K_i^\dagger E(X)$ for all i . Taking conjugate transposes: $K_i X^\dagger = E(X)^\dagger K_i$. Then

$$\begin{aligned} E(X^\dagger Y) &= \sum_i K_i X^\dagger Y K_i^\dagger \\ &= \sum_i E(X)^\dagger K_i Y K_i^\dagger \\ &= E(X)^\dagger E(Y). \end{aligned}$$

□

4.4 Irreducibility

Definition 4.23 (Irreducible map). A linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ is *irreducible* if whenever P is an orthogonal projection satisfying $E(PM_D(\mathbb{C})P) \subseteq PM_D(\mathbb{C})P$, then $P = 0$ or $P = \mathbb{1}$. This is [Wol12, Thm. 6.2, condition (1)]. The definition applies to any linear map; complete positivity is not required.

Theorem 4.24 (Injectivity implies irreducibility). *If A is an injective MPS tensor, then its transfer map $\mathcal{E}_A$ is irreducible.*

Proof. If P is an invariant projection for $\mathcal{E}_A$, then $(\mathbb{1} - P)A^i P = 0$ for all i . Since the $\{A^i\}$ span $M_D(\mathbb{C})$, $(\mathbb{1} - P)MP = 0$ for all M , so $P = 0$ or $P = \mathbb{1}$. □

Remark 4.25 (Transfer map is CP). The transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is completely positive: it is already written in Kraus form, with Kraus operators $\{A^i\}$.

Theorem 4.26 (Transfer map is a channel). *If $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, then the transfer map $\mathcal{E}_A(X) = \sum_i A^i X (A^i)^\dagger$ is a quantum channel (CPTP).*

Proof. Complete positivity is the observation of Remark 4.25. Trace preservation: $\text{tr}(\mathcal{E}_A(X)) = \sum_i \text{tr}(A^i X (A^i)^\dagger) = \text{tr}((\sum_i (A^i)^\dagger A^i)X) = \text{tr}(X)$ by cyclicity and the normalization hypothesis. □

4.5 Cesàro fixed-point theorem

Definition 4.27 (Cesàro mean). The *Cesàro mean* of a linear map E at order N , applied to a starting matrix ρ_0 , is

$$\sigma_N = \frac{1}{N} \sum_{n=0}^{N-1} E^n(\rho_0).$$

Theorem 4.28 (Cesàro telescope). $E(\sigma_N) - \sigma_N = \frac{1}{N}(E^N(\rho_0) - \rho_0)$.

Proof. Telescoping: $E(\sigma_N) = \frac{1}{N} \sum_{n=1}^N E^n(\rho_0)$ and subtracting σ_N cancels all but the first and last terms. □

Theorem 4.29 (Hermitian fixed-point decomposition). *Let E be a quantum channel and $X = E(X)$ a Hermitian fixed point with decomposition $X = P_+ - P_-$ into orthogonal positive and negative parts ($P_\pm \geq 0$, $\text{tr}(P_+ P_-) = 0$). Then $E(P_+) = P_+$ and $E(P_-) = P_-$. This is [Wol12, Prop. 6.8].*

Proof. Let Q be the support projection of P_+ . Trace preservation and positivity give $\text{tr}(QE(P_-)) = 0$ and $E(P_-) \geq 0$, so $QE(P_-) = 0$. The fixed-point equation $E(P_+) - E(P_-) = P_+ - P_-$ then identifies $E(P_\pm) = P_\pm$. $\square$

Theorem 4.30 (Cesàro fixed-point theorem). *Every quantum channel $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ (with $D \geq 1$) has a nonzero positive semidefinite fixed point: there exists $\rho \geq 0$, $\rho \neq 0$, with $E(\rho) = \rho$.*

Proof. Start with any $\rho_0 \in \mathcal{D}_D$ (Theorem 4.10). Each Cesàro mean σ_N lies in $\mathcal{D}_D$ (by convexity and channel preservation). By compactness (Theorem 4.8), extract a convergent subsequence $\sigma_{N_k} \rightarrow \rho$. The telescope identity (Theorem 4.28) gives $\|E(\sigma_N) - \sigma_N\| = \frac{1}{N}\|E^N(\rho_0) - \rho_0\| \rightarrow 0$, so $E(\rho) = \rho$ by continuity. In [Wol12, Thm. 6.11], existence is proved via Brouwer's fixed-point theorem; we use the Cesàro argument instead. $\square$

4.6 Fixed-point projection

Definition 4.31 (Fixed-point projection). Given a linear map $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ and a fixed point ρ with $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$, the *fixed-point projection* is the rank-one map

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho.$$

This projects onto the span of ρ along the kernel of the trace functional. At this stage it is a projection onto the chosen fixed point, not yet onto the full fixed-point space; the uniqueness statement is proved later.

Theorem 4.32 (Power decomposition). *If E is trace-preserving and $E(\rho) = \rho$, then for $n \geq 1$,*

$$E^n = P + (E - P)^n,$$

where P is the fixed-point projection.

Proof. P is idempotent ($P^2 = P$), the fixed-point relation $E(\rho) = \rho$ gives $E \circ P = P$, and trace preservation gives $P \circ E = P$. These imply $(E - P) \circ P = 0$ and $P \circ (E - P) = 0$, so in the binomial expansion of $(P + (E - P))^n$ all cross terms vanish, leaving $E^n = P^n + (E - P)^n = P + (E - P)^n$. $\square$

4.7 Peripheral spectrum and primitivity

Definition 4.33 (Peripheral spectrum). The *peripheral spectrum* of a bounded linear operator T is the set of spectral values μ with $|\mu|$ equal to the spectral radius of T .

Definition 4.34 (Peripheral eigenvalues). The *peripheral eigenvalues* of a linear map E on a finite-dimensional space are the eigenvalues λ on the unit circle: $|\lambda| = 1$. For a channel, the spectral radius is 1 [Wol12, Prop. 6.1], so these coincide with the eigenvalues of maximal modulus.

Note: The condition $|\lambda| = 1$ is appropriate for channels, since the spectral radius of a channel is 1. For a general linear map with spectral radius $r \neq 1$ the condition would generalise to $|\lambda| = r$.

Theorem 4.35 (1 is a peripheral eigenvalue). *If E has a nonzero fixed point $\rho \neq 0$ with $E(\rho) = \rho$, then 1 is a peripheral eigenvalue of E .*

Proof. ρ is an eigenvector with eigenvalue 1, and $|1| = 1$. $\square$

Remark 4.36 (Peripheral eigenvalues lie on the unit circle). Every peripheral eigenvalue λ satisfies $|\lambda| = 1$. This is simply the defining condition in Definition 4.34.

Theorem 4.37 (Finiteness of peripheral eigenvalues). *On a finite-dimensional space, the set of peripheral eigenvalues is finite.*

Proof. Peripheral eigenvalues are a subset of all eigenvalues, which is a finite set in finite dimensions. $\square$

Theorem 4.38 (Power-stable peripheral eigenvalues are roots of unity). *Let E be a linear endomorphism on a finite-dimensional space. If μ is a peripheral eigenvalue of E and μ^n is an eigenvalue of E for every $n \geq 1$, then μ is a root of unity.*

Proof. Since μ^n is an eigenvalue for every n , the pigeonhole principle on the finite eigenvalue set gives $\mu^a = \mu^b$ for some $a \neq b$, hence $\mu^{|a-b|} = 1$. $\square$

Theorem 4.39 (Peripheral powers imply root of unity). *If the set of peripheral eigenvalues of a linear endomorphism E on a finite-dimensional space is closed under powers ($\mu \in \text{peripheral}(E)$ and $n \geq 1$ imply $\mu^n \in \text{peripheral}(E)$), then every peripheral eigenvalue is a root of unity.*

Proof. The closure hypothesis provides $\mu^n \in \text{peripheral}(E)$ for all n , which in particular means μ^n is an eigenvalue. Theorem 4.38 then gives $\mu^p = 1$ for some $p > 0$. $\square$

4.7.1 Periodicity removal by powering

Remark 4.40. The results in this subsection are purely spectral: they use only finiteness of a set of complex numbers and the spectral mapping theorem for powers. In particular, they do not rely on complete positivity. For transfer maps, replacing a tensor by its p -blocked tensor replaces $\mathcal{E}_A$ by $\mathcal{E}_A^p$, so this powering step is the operator-theoretic form of blocking.

Lemma 4.41 (Common exponent for a finite set of roots of unity). *Let $s \subseteq \mathbb{C}$ be a finite set. Assume that for every $\mu \in s$ there exists an exponent $p_\mu > 0$ with $\mu^{p_\mu} = 1$. Then there exists $p > 0$ such that $\mu^p = 1$ for all $\mu \in s$.*

Proof. Take p to be the least common multiple of the exponents p_μ . $\square$

Lemma 4.42 (Powering collapses peripheral eigenvalues). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a linear map, and assume that E has a nonzero fixed point $\rho \neq 0$. Let $p > 0$. If every peripheral eigenvalue μ of E satisfies $\mu^p = 1$, then $\text{peripheral}(E^p) = \{1\}$.*

Proof. The fixed point ρ gives $1 \in \text{peripheral}(E^p)$. For the reverse inclusion, let $\nu \in \text{peripheral}(E^p)$. Spectral mapping gives $\nu = \mu^p$ for some $\mu \in \sigma(E)$. From $|\nu| = 1$ and $p > 0$ we obtain $|\mu| = 1$, hence $\mu \in \text{peripheral}(E)$. The hypothesis $\mu^p = 1$ then forces $\nu = 1$. $\square$

For a normalized MPS tensor, the transfer map is naturally trace-preserving but need not be unital. In general one should not expect a single gauge choice to make it both unital and trace-preserving. The next result therefore treats a special doubly-stochastic case; the general MPS situation is handled in Subsection 4.7.2.

Lemma 4.43 (Peripheral eigenvalues closed under powers). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is both unital and trace-preserving, and assume that E is irreducible. Then $\text{peripheral}(E)$ is closed under powers: if $\mu \in \text{peripheral}(E)$ then $\mu^n \in \text{peripheral}(E)$ for every $n \in \mathbb{N}$.*

Proof. Let $E(X) = \mu X$ with $|\mu| = 1$. The Kadison–Schwarz equality for peripheral eigenvectors (Theorem 4.21) shows that $X^\dagger X$ is a positive semidefinite fixed point. By irreducibility, this fixed point is positive definite (Theorem 5.3), hence X is invertible. Multiplicative-domain powers then give $E(X^n) = \mu^n X^n$, so μ^n is again peripheral. $\square$

Theorem 4.44 (Roots of unity for irreducible doubly stochastic maps). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is both unital and trace-preserving, and assume that E is irreducible. Then every peripheral eigenvalue μ of E is a root of unity: there exists $p > 0$ with $\mu^p = 1$.*

Proof. Combine the closure-under-powers result (Lemma 4.43) with Theorem 4.39. $\square$

4.7.2 Peripheral closure via adjoint fixed point

The doubly-stochastic hypothesis in Lemma 4.43 (unital *and* trace-preserving) can be weakened: trace preservation is only used to derive the Kadison–Schwarz equality at peripheral eigenvectors. The following result replaces the TP condition by the existence of a positive definite fixed point for the *adjoint* Kraus map, matching the weighted-trace argument in [CPGSV17, Appendix A].

Lemma 4.45 (Peripheral closure via adjoint fixed point). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital ($\sum_i K_i K_i^\dagger = \mathbb{1}$), and assume:*

1. *the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point $\rho > 0$, and*
2. *E is irreducible.*

Then $\text{peripheral}(E)$ is closed under powers: if $\mu \in \text{peripheral}(E)$ then $\mu^n \in \text{peripheral}(E)$ for every $n \in \mathbb{N}$.

Proof. Let $E(X) = \mu X$ with $|\mu| = 1$. The weighted Kadison–Schwarz equality (using the adjoint fixed point ρ in place of trace preservation) shows $E(X^\dagger X) = E(X)^\dagger E(X)$. Hence $X^\dagger X$ is a PSD fixed point, which is positive definite by irreducibility (Theorem 5.3). So X is invertible, and multiplicative-domain powers give $E(X^n) = \mu^n X^n$, so μ^n is peripheral. $\square$

Theorem 4.46 (Roots of unity via adjoint fixed point). *Let $E(X) = \sum_i K_i X K_i^\dagger$ be a Kraus map that is unital, and assume that the adjoint Kraus map $E^*(X) = \sum_i K_i^\dagger X K_i$ has a positive definite fixed point and that E is irreducible. Then every peripheral eigenvalue of E is a root of unity.*

Proof. Combine the closure-under-powers result (Lemma 4.45) with Theorem 4.39. $\square$

Remark 4.47. Lemma 4.45 and Theorem 4.46 strictly generalise Lemma 4.43 and Theorem 4.44: the doubly-stochastic (unital + TP) condition implies $E^*(\mathbb{1}) = \mathbb{1}$, which is a positive definite fixed point of the adjoint. The adjoint-fixed-point formulation matches the argument in [CPGSV17, Appendix A], which uses a faithful invariant state rather than double stochasticity.

Definition 4.48 (Channel period). The *period* of a channel E is the number of peripheral eigenvalues (counted without multiplicity): $p(E) := |\text{peripheral}(E)|$.

Definition 4.49 (Primitive map). A linear map is *primitive* if its only peripheral eigenvalue is $\lambda = 1$, i.e. $\text{peripheral}(E) = \{1\}$. For channels this corresponds to [Wol12, Thm. 6.7]. This definition applies to any linear endomorphism, not only to channels.

Theorem 4.50 (Primitivity equals period one). *A linear map with a nonzero fixed point is primitive if and only if its period is 1.*

Proof. If $\text{peripheral}(E) = \{1\}$, then $|\text{peripheral}(E)| = 1$. Conversely, if $|\text{peripheral}(E)| = 1$, then since $1 \in \text{peripheral}(E)$ (Theorem 4.35), the only element is 1. $\square$

Theorem 4.51 (Primitive maps have a spectral gap on the complement). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be trace-preserving, and let $\rho \neq 0$ satisfy $E(\rho) = \rho$ and $\text{tr}(\rho) \neq 0$. Let*

$$P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)} \rho$$

be the fixed-point projection associated to ρ . Assume that:

1. *E is primitive;*
2. *every eigenvalue μ of E satisfies $|\mu| \leq 1$;*
3. *whenever $E(X) = X$ and $\text{tr}(X) = 0$, one has $X = 0$.*

Then every eigenvalue ν of $E - P$ satisfies $|\nu| < 1$.

Proof. Let $(E - P)(X) = \nu X$ with $X \neq 0$. Then

$$E(X) = \nu X + P(X).$$

If $\text{tr}(X) = 0$, then $P(X) = 0$, so $E(X) = \nu X$ and hence ν is an eigenvalue of E with $|\nu| \leq 1$. If $|\nu| = 1$, primitivity forces $\nu = 1$, so X is a trace-zero fixed point of E ; by assumption this implies $X = 0$, a contradiction. Thus $|\nu| < 1$.

If instead $\text{tr}(X) \neq 0$, trace preservation and $\text{tr}(P(X)) = \text{tr}(X)$ give

$$\text{tr}(X) = \text{tr}(E(X)) = \nu \text{tr}(X) + \text{tr}(P(X)) = \nu \text{tr}(X) + \text{tr}(X),$$

so $\nu \text{tr}(X) = 0$ and therefore $\nu = 0$. In either case, $|\nu| < 1$. $\square$

Chapter 5

Quantum Perron–Frobenius Theory

This chapter establishes that the transfer map of an injective MPS tensor has a unique positive semidefinite fixed point (up to scaling), and that this fixed point is positive definite. The results follow [EHK78] and [Wol12, Ch. 6].

5.1 Positive definiteness

Remark 5.1. We begin with the injective-tensor proof because it is the shorter, more MPS-specific argument. The irreducible CP-map version follows immediately afterwards.

Theorem 5.2 (PSD fixed point is positive definite). *Let A be an injective MPS tensor and let $\rho \geq 0$, $\rho \neq 0$, be a fixed point of the transfer map $\mathcal{E}_A(\rho) = \rho$. Then ρ is positive definite.*

Proof. Suppose $v^\dagger \rho v = 0$ for some $v \neq 0$. Since $\rho \geq 0$, this implies $\rho v = 0$. The fixed-point equation gives $v^\dagger \rho v = \sum_i ((A^i)^\dagger v)^\dagger \rho ((A^i)^\dagger v)$. Each term is nonneg; if the sum is zero, then $\rho (A^i)^\dagger v = 0$ for all i : the kernel of ρ is invariant under all $(A^i)^\dagger$. Since the $\{A^i\}$ span $M_D(\mathbb{C})$, we have $\rho M v = 0$ for every matrix M . Given any vector w , choose M with $M v = w$. Then $\rho w = 0$ for every w , so $\rho = 0$, contradicting $\rho \neq 0$. In [Wol12, Thm. 6.2], positive definiteness is proved using the growth condition $(\mathbb{1} + \mathcal{E})^{D-1}(X) > 0$ for irreducible maps ([Wol12, Thm. 6.2, condition (2)]). Here injectivity rules out a nontrivial kernel directly. $\square$

Theorem 5.3 (PSD fixed point is PD — irreducible version). *If $\mathcal{E}_A$ is irreducible and $\rho \geq 0$, $\rho \neq 0$ is a fixed point, then ρ is positive definite.*

Proof. Suppose ρ is not positive definite. Then ρ has a nontrivial kernel; let Q be the orthogonal projection onto the support of ρ (the complement of the kernel). A direct computation using the fixed-point equation $\mathcal{E}_A(\rho) = \rho$ shows that Q is an invariant projection for $\mathcal{E}_A$: it satisfies $Q \mathcal{E}_A(QXQ) Q = \mathcal{E}_A(QXQ)$ for all X . By irreducibility of $\mathcal{E}_A$ (Definition 4.23), every invariant projection is 0 or $\mathbb{1}$. If $Q = 0$ then $\rho = 0$, contradicting $\rho \neq 0$. If $Q = \mathbb{1}$ then ρ is positive definite, contradicting our assumption. Hence ρ must be positive definite. $\square$

5.2 Uniqueness

Theorem 5.4 (Uniqueness of PSD fixed point). *Let A be an injective MPS tensor. If $\rho, \sigma \geq 0$ are both nonzero fixed points of $\mathcal{E}_A$, then $\sigma = c\rho$ for some $c \in \mathbb{C}$.*

Proof. Both ρ and σ are positive definite by Theorem 5.2. Write $\rho = SS^\dagger$ via a spectral square root and set $H = (S^\dagger)^{-1}\sigma S^{-1}$, which is Hermitian and positive definite. Let c_0 be the minimal eigenvalue of H . Then $\tau := \sigma - c_0\rho$ is a PSD fixed point which is *not* positive definite (it has a zero eigenvalue by construction of c_0). By Theorem 5.2, τ must be zero, giving $\sigma = c_0\rho$. In [Wol12, Thm. 6.2], uniqueness follows from non-degeneracy of the spectral radius eigenvalue via the $(\mathbb{1} + \mathcal{E})^{D-1}$ growth condition. Here the conclusion follows by subtracting the minimal-eigenvalue multiple of ρ and applying Theorem 5.2. $\square$

Remark 5.5. The proof actually produces a positive real scalar $c_0 > 0$: it is the minimal eigenvalue of the positive definite Hermitian matrix $H = (S^\dagger)^{-1}\sigma S^{-1}$. The Lean theorem records the resulting proportionality in the ambient scalar field $\mathbb{C}$.

5.3 Existence and assembly

Theorem 5.6 (Existence of PSD fixed point). *Assume $D \geq 1$. Let A be an MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ (so that $\mathcal{E}_A$ is trace-preserving). Then there exists $\rho \geq 0$, $\rho \neq 0$, with $\mathcal{E}_A(\rho) = \rho$.*

Proof. Under the normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, the transfer map is a quantum channel (Theorem 4.26). The Cesàro fixed-point theorem (Theorem 4.30) then provides the fixed point; injectivity is not needed for this step. $\square$

Theorem 5.7 (Quantum Perron–Frobenius). *Assume $D \geq 1$. Let A be an injective MPS tensor with $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then the transfer map $\mathcal{E}_A$ has a unique PSD fixed point ρ (up to scaling), and ρ is positive definite.*

Proof. Combine Theorems 5.6, 5.2, and 5.4. $\square$

Remark 5.8 (Dimension-zero edge case). Lean also packages a wrapper theorem removing the explicit hypothesis $D \geq 1$ by adding the formal edge case $D = 0$. We treat this as an implementation detail rather than part of the mathematical narrative: the tensor-network results of interest all concern positive bond dimension.

5.4 Unital and trace-preserving gauges

Theorem 5.9 (Unital gauge). *Let $\mathcal{E}_A(\rho) = \rho$ and let S be an invertible matrix with $SS^\dagger = \rho$. Define the gauged operators $A'^i = S^{-1}A^iS$. Then $\sum_i A'^i(A'^i)^\dagger = \mathbb{1}$, i.e. the gauged Kraus map is unital.*

Proof. Expand each summand: $(S^{-1}A^iS)(S^{-1}A^iS)^\dagger = S^{-1}A^i \underbrace{SS^\dagger}_{\rho} (A^i)^\dagger (S^\dagger)^{-1}$. Summing over i and using $\sum_i A^i \rho (A^i)^\dagger = \rho$ gives $S^{-1} \rho (S^\dagger)^{-1} = S^{-1}SS^\dagger(S^\dagger)^{-1} = \mathbb{1}$. $\square$

Theorem 5.10 (Trace-preserving gauge). *Let σ be a fixed point of the adjoint transfer map $\sum_i (A^i)^\dagger \sigma A^i = \sigma$, and let S be invertible with $S^\dagger S = \sigma$. Define $A'^i = SA^iS^{-1}$. Then $\sum_i (A'^i)^\dagger A'^i = \mathbb{1}$, i.e. the gauged Kraus map is trace-preserving.*

Proof. Expand each summand: $(SA^iS^{-1})^\dagger(SA^iS^{-1}) = (S^\dagger)^{-1}(A^i)^\dagger \underbrace{S^\dagger S}_\sigma A^iS^{-1}$. Summing over i and using $\sum_i (A^i)^\dagger \sigma A^i = \sigma$ gives $(S^\dagger)^{-1} \sigma S^{-1} = (S^\dagger)^{-1} S^\dagger S S^{-1} = \mathbb{1}$. $\square$

Remark 5.11. The gauges of Theorems 5.9 and 5.10 are generally different similarity transforms: one is built from a fixed point of $\mathcal{E}_A$, the other from a fixed point of the adjoint transfer map. This section does not claim that a single gauge makes the transfer map simultaneously unital and trace-preserving.

Remark 5.12 (Eigenvector existence for general positive maps). The trace-preserving fixed-point statements in this chapter are already unconditional: Theorem 5.6 obtains a PSD fixed point from the Cesàro argument of Theorem 4.30. The later Perron–Frobenius and TP-gauge step of Chapter 8, §8.8, is now also formalized: Theorem 8.31 gives a Perron–Frobenius eigenvector for a general positive map, and Theorem 8.35 uses it to put an irreducible tensor into TP gauge. The remaining gap lies later in the end-to-end canonical-form / block-injectivity assembly, not in the Perron–Frobenius existence step itself.

Chapter 6

Spectral Gap and Block Separation

This chapter proves that the MPV overlaps between distinct (non-gauge-phase-equivalent) MPS blocks decay to zero as system size grows, while self-overlaps of primitive blocks converge to 1. The argument is based on a spectral analysis of the *mixed transfer operator*, following [PGVWC07].

Throughout this chapter, all MPS tensors are assumed to satisfy the trace-preserving normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$.

6.1 Mixed transfer operator

Definition 6.1 (Mixed transfer operator). The *mixed transfer operator* (or *cross transfer operator*) for two MPS tensors A and B of the same bond dimension D is the linear map $F_{AB} : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ defined by

$$F_{AB}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Definition 6.2 (Rectangular mixed transfer operator). For MPS tensors A of bond dimension D_1 and B of bond dimension D_2 (possibly $D_1 \neq D_2$), the *rectangular mixed transfer operator* is $F_{AB}^{\text{rect}} : M_{D_1 \times D_2}(\mathbb{C}) \rightarrow M_{D_1 \times D_2}(\mathbb{C})$ defined by

$$F_{AB}^{\text{rect}}(X) = \sum_{i=0}^{d-1} A^i X (B^i)^\dagger.$$

Remark 6.3 (Self-transfer). Setting $B = A$ recovers the ordinary transfer map: $F_{AA} = \mathcal{E}_A$.

Theorem 6.4 (Iterated mixed transfer). For all $X \in M_D(\mathbb{C})$ and $N \geq 0$,

$$F_{AB}^N(X) = \sum_{\sigma \in \{0, \dots, d-1\}^N} A^\sigma X (B^\sigma)^\dagger,$$

where $A^\sigma = A^{\sigma_1} \dots A^{\sigma_N}$ denotes the word evaluation (Definition 2.2).

Proof. Induction on N : the base case is $F^0(X) = X$; the inductive step applies F_{AB} to each summand and re-indexes via $\{0, \dots, d-1\}^{N+1} \cong \{0, \dots, d-1\} \times \{0, \dots, d-1\}^N$. $\square$

Theorem 6.5 (Matrix trace of iterated mixed transfer at the identity). *For $N \geq 0$,*

$$\mathrm{tr}(F_{AB}^N(\mathbb{1})) = \sum_{\sigma} \mathrm{tr}(A^{\sigma}(B^{\sigma})^{\dagger}).$$

This is the matrix trace of F_{AB}^N evaluated at the identity matrix. It should not be confused with the operator trace $\mathrm{Tr}(F_{AB}^N)$, which equals the MPV overlap (Theorem 6.7).

Proof. Apply the matrix trace to each summand in Theorem 6.4 with $X = \mathbb{1}$: $\mathrm{tr}(F_{AB}^N(\mathbb{1})) = \sum_{\sigma} \mathrm{tr}(A^{\sigma} \cdot \mathbb{1} \cdot (B^{\sigma})^{\dagger}) = \sum_{\sigma} \mathrm{tr}(A^{\sigma}(B^{\sigma})^{\dagger})$. $\square$

6.2 MPV overlap as transfer trace

Recall from Definition 2.31 that

$$O_{AB}(N) = \sum_{\sigma \in \{0, \dots, d-1\}^N} V^{(N)}(A)_{\sigma} \overline{V^{(N)}(B)_{\sigma}}.$$

This section rewrites that scalar as the operator trace of the mixed transfer power.

Lemma 6.6 (Trace expansion over matrix units). *For any linear endomorphism T on $M_D(\mathbb{C})$, the operator trace can be expanded as*

$$\mathrm{Tr}(T) = \sum_{p,q=0}^{D-1} (T(E_{pq}))_{pq},$$

where E_{pq} is the matrix with 1 in position (p, q) and 0 elsewhere.

Proof. The operator trace $\mathrm{Tr}(T)$ equals the sum of $\mathrm{tr}(T(E_{pq}) \cdot E_{qp})$ over all (p, q) , and $\mathrm{tr}(M \cdot E_{qp}) = M_{pq}$. $\square$

Theorem 6.7 (Overlap equals transfer trace). *For normalized tensors A, B of bond dimension $D \geq 1$,*

$$O_{AB}(N) = \mathrm{Tr}(F_{AB}^N).$$

Proof. Expand the operator trace using Lemma 6.6 as a sum over matrix units E_{pq} . Apply Theorem 6.4 to evaluate $F_{AB}^N(E_{pq})$, extract the (p, q) -entry, and reassemble. The resulting double sum over (p, q) and σ equals $\sum_{\sigma} V^{(N)}(A)_{\sigma} \overline{V^{(N)}(B)_{\sigma}} = O_{AB}(N)$. $\square$

Theorem 6.8 (Rectangular overlap as transfer trace). *For tensors A of bond dimension $D_1 \geq 1$ and B of bond dimension $D_2 \geq 1$,*

$$O_{AB}(N) = \mathrm{Tr}((F_{AB}^{\mathrm{rect}})^N).$$

Proof. Expand the operator trace over the rectangular matrix-unit basis. Then apply the rectangular iterated-transfer formula and sum over the matrix-unit indices exactly as in Theorem 6.7. $\square$

Lemma 6.9 (Word trace as entrywise inner product). *For a word $\sigma \in \{0, \dots, d-1\}^N$,*

$$\mathrm{tr}(A^{\sigma}(B^{\sigma})^{\dagger}) = \sum_{j,k=0}^{D-1} (A^{\sigma})_{jk} \overline{(B^{\sigma})_{jk}}.$$

Proof. Expand the matrix trace and the (j, j) -entry of $A^{\sigma}(B^{\sigma})^{\dagger}$. $\square$

6.3 Eigenvalue bound and spectral gap

The eigenvalue bound comes from a uniform Frobenius-norm estimate obtained directly from the trace-preserving normalization. For the rigidity theorem, one gauges A and B separately using positive fixed points of their transfer maps to obtain unital Kraus families, and then applies the Kadison–Schwarz equality argument to a block embedding of a mixed-transfer eigenvector.

Theorem 6.10 (Eigenvalue bound). *Let A, B be normalized MPS tensors. Every eigenvalue μ of F_{AB} satisfies $|\mu| \leq 1$.*

Proof. Expand $F_{AB}^n(X)$ as a sum over words and use Cauchy–Schwarz, together with $\sum_i (A^i)^\dagger A^i = \sum_i (B^i)^\dagger B^i = \mathbb{1}$, to obtain a uniform Frobenius-norm bound $\|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for all n , where C_D depends only on D . If $F_{AB}(X) = \mu X$ and $X \neq 0$, then $|\mu|^n \|X\|_{\text{HS}} = \|F_{AB}^n(X)\|_{\text{HS}} \leq C_D \|X\|_{\text{HS}}$ for every n . Hence $|\mu| \leq 1$.

The argument uses only the trace-preserving normalization and does not appeal to Perron–Frobenius theory. $\square$

Theorem 6.11 (Spectral radius bound). *$\rho(F_{AB}) \leq 1$ for normalized tensors.*

Proof. The spectral radius is the supremum of $|\mu|$ over all eigenvalues, and each satisfies $|\mu| \leq 1$ by Theorem 6.10. $\square$

Theorem 6.12 (Modulus-one eigenvalue implies gauge-phase equivalence). *Let A, B be injective normalized MPS tensors. If $\rho(F_{AB}) \geq 1$, then A and B are gauge-phase equivalent.*

Proof. Since $\rho(F_{AB}) \leq 1$ (Theorem 6.11) and $\rho(F_{AB}) \geq 1$ by hypothesis, there exists an eigenvalue μ with $|\mu| = 1$ and eigenvector $X \neq 0$: $F_{AB}(X) = \mu X$.

Choose positive fixed points of the transfer maps of A and B (Theorem 5.7) and gauge the two tensors separately to unital Kraus families (Theorem 5.9). Gauge X accordingly and embed it as an off-diagonal block matrix for the block Kraus family with diagonal blocks A'^i and B'^i . The KS equality for peripheral eigenvectors (Theorem 4.21) and the resulting Kraus commutation relation (Theorem 4.20) yield intertwining identities between A'^i , B'^i , and the gauged eigenvector. Injectivity and the multiplicative-domain argument force this intertwiner to be invertible; Skolem–Noether (Theorem 3.9) then yields $B^i = \bar{\mu} X A^i X^{-1}$ for all i , establishing gauge-phase equivalence (Definition 2.13).

The crucial point is that one gauges the individual transfer maps to unital form; one does not require the mixed transfer map itself to be simultaneously unital and trace-preserving. $\square$

Theorem 6.13 (Strict spectral gap). *Let A, B be injective normalized MPS tensors that are not gauge-phase equivalent. Then $\rho(F_{AB}) < 1$.*

Proof. $\rho(F_{AB}) \leq 1$ by Theorem 6.11. If $\rho(F_{AB}) = 1$, Theorem 6.12 gives gauge-phase equivalence, contradicting the hypothesis. $\square$

6.4 Spectral gap — rectangular case

Theorem 6.14 (Rectangular spectral gap). *Let A be an injective normalized tensor of bond dimension D_1 and B an injective normalized tensor of bond dimension D_2 , with $D_1 \neq D_2$. Then $\rho(F_{AB}^{\text{rect}}) < 1$.*

Proof. The same word-expansion argument as in Theorem 6.10 gives $|\mu| \leq 1$ for every eigenvalue of F_{AB}^{rect} . If $\rho(F_{AB}^{\text{rect}}) = 1$, choose a modulus-one eigenvector X .

Gauge A and B separately to unital Kraus families and apply the same block-embedding Kadison–Schwarz argument as in the square case. It shows that $\ker(X)$ is invariant under all matrices on $\mathbb{C}^{D_2}$, so X is injective and therefore $D_2 \leq D_1$. Applying the same argument to $X^\dagger$ gives $D_1 \leq D_2$. Hence $D_1 = D_2$, contradicting the hypothesis. $\square$

Theorem 6.15 (Rectangular overlap decay). *If $D_1 \neq D_2$ and both tensors are injective and normalized, then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. The spectral gap $\rho(F_{AB}^{\text{rect}}) < 1$ (Theorem 6.14) implies $(F_{AB}^{\text{rect}})^N \rightarrow 0$, so the operator trace converges to 0. By Theorem 6.8, $O_{AB}(N) = \text{Tr}((F_{AB}^{\text{rect}})^N) \rightarrow 0$. $\square$

6.5 MPV overlap decay

Theorem 6.16 (Transfer powers converge to zero). *Let A, B be injective normalized tensors that are not gauge-phase equivalent. Then $F_{AB}^n(X) \rightarrow 0$ for every $X \in M_D(\mathbb{C})$.*

Proof. The spectral gap $\rho(F_{AB}) < 1$ (Theorem 6.13) implies $\|F_{AB}^n\|_{\text{op}} \rightarrow 0$ by the Gelfand formula, so $F_{AB}^n(X) \rightarrow 0$ for every X . $\square$

Theorem 6.17 (Overlap decay). *Let A, B be injective normalized tensors of the same bond dimension that are not gauge-phase equivalent. Then $O_{AB}(N) \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. By Theorem 6.7, $O_{AB}(N) = \text{Tr}(F_{AB}^N)$. Expand the operator trace over matrix units (Lemma 6.6): $\text{Tr}(F_{AB}^N) = \sum_{p,q} (F_{AB}^N(E_{pq}))_{pq}$. By Theorem 6.16, each $F_{AB}^N(E_{pq}) \rightarrow 0$ entry-wise. Since the sum is finite, $O_{AB}(N) \rightarrow 0$. $\square$

6.6 Block separation

These asymptotic facts are the ingredients used later in the canonical-form separation arguments.

Remark 6.18 (Cross-correlation decay). If A and B are injective normalized tensors that are not gauge-phase equivalent, then for every $X \in M_D(\mathbb{C})$ one has $\text{tr}(F_{AB}^N(X)) \rightarrow 0$ as $N \rightarrow \infty$. This is the scalar trace consequence of the operator convergence $F_{AB}^N(X) \rightarrow 0$ from Theorem 6.16.

Remark 6.19 (Self-correlation persists). If ρ is a fixed point of $\mathcal{E}_A$, then $\text{tr}(\mathcal{E}_A^N(\rho)) = \text{tr}(\rho)$ for all $N \geq 0$. In particular this applies to the distinguished QPF fixed point from Theorem 5.7.

6.7 Primitive overlap convergence

Theorem 6.20 (Trace of powers converges to 1). *Let E be trace-preserving with fixed point ρ ($\text{tr}(\rho) \neq 0$) and let P be the fixed-point projection (Definition 4.31). If $\rho_{\text{spec}}(E - P) < 1$, then $\text{Tr}(E^n) \rightarrow 1$ as $n \rightarrow \infty$.*

Proof. By the power decomposition (Theorem 4.32), $E^n = P + (E - P)^n$ for $n \geq 1$. Since $\rho_{\text{spec}}(E - P) < 1$, the Gelfand formula gives $(E - P)^n \rightarrow 0$, so $\text{Tr}(E^n) \rightarrow \text{Tr}(P)$. The operator trace of P equals 1: since $P(X) = \frac{\text{tr}(X)}{\text{tr}(\rho)}\rho$, we have $\text{Tr}(P) = \frac{\text{tr}(\rho)}{\text{tr}(\rho)} = 1$. $\square$

Theorem 6.21 (Primitive overlap convergence). *Let A be a normalized MPS tensor with a nonzero PSD fixed point ρ of the transfer map. If $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$ (where P is the fixed-point projection), then $O_{AA}(N) \rightarrow 1$ as $N \rightarrow \infty$.*

Proof. By Theorem 6.7 (with $A = B$), $O_{AA}(N) = \text{Tr}(\mathcal{E}_A^N)$. By Theorem 6.20, $\text{Tr}(\mathcal{E}_A^N) \rightarrow 1$. $\square$

Chapter 7

Wielandt Bound

This chapter proves a quantum analog of Wielandt's inequality: for a normal MPS tensor with bond dimension D , the cumulative span of matrix products stabilises after at most D^2 steps. The results follow [SPGWC10].

7.1 Cumulative span

Definition 7.1 (Word span). The *word span* at length n is the subspace

$$S_n(A) = \text{span}_{\mathbb{C}}\{A^w : |w| = n\} \subseteq M_D(\mathbb{C}).$$

This is [SPGWC10, Eq. (1)].

Definition 7.2 (Cumulative span). The *cumulative span* at level n is

$$T_n(A) = S_0(A) + S_1(A) + \cdots + S_n(A).$$

Lemma 7.3 (Monotonicity of cumulative span). $T_n(A) \leq T_{n+1}(A)$ for all n .

Proof. T_{n+1} includes all words of length $\leq n+1$, which contains all words of length $\leq n$. $\square$

Lemma 7.4 (Dimension bound). $\dim T_n(A) \leq D^2$ for all n .

Proof. $T_n(A) \subseteq M_D(\mathbb{C})$ and $\dim M_D(\mathbb{C}) = D^2$. $\square$

Theorem 7.5 (Cumulative span stabilisation). If $T_n(A) = T_{n+1}(A)$, then $T_m(A) = T_n(A)$ for all $m \geq n$.

Proof. If $T_n = T_{n+1}$, then $S_{n+1} \subseteq T_n$. Left-multiplying any element of S_{n+1} by A^i gives an element of S_{n+2} , so $S_{n+2} \subseteq T_{n+1} = T_n$. By induction, $S_m \subseteq T_n$ for all $m > n$, hence $T_m = T_n$. $\square$

Theorem 7.6 (Dimension growth). If $T_n(A) \neq T_{n+1}(A)$, then $\dim T_{n+1}(A) > \dim T_n(A)$.

Proof. $T_n \leq T_{n+1}$ (Lemma 7.3) and $T_n \neq T_{n+1}$ imply strict submodule inclusion, hence strict rank increase. $\square$

Lemma 7.7 (Word span characterises block injectivity). $S_L(A) = M_D(\mathbb{C})$ if and only if A is L -block injective.

Proof. Both conditions assert that the span of all products of length L equals $M_D(\mathbb{C})$. $\square$

7.2 Fitting decomposition

Definition 7.8 (Fitting decomposition). A *Fitting decomposition* of a linear endomorphism $f : V \rightarrow V$ on a finite-dimensional vector space over an algebraically closed field consists of:

1. f is nilpotent on the generalized 0-eigenspace,
2. f is invertible on each generalized μ -eigenspace for $\mu \neq 0$,
3. the generalized eigenspaces span V ,
4. the generalized eigenspaces are linearly independent.

Theorem 7.9 (Fitting decomposition exists). *Every linear endomorphism on a finite-dimensional vector space over an algebraically closed field admits a Fitting decomposition.*

Proof. Nilpotency on the zero generalized eigenspace follows from the definition of generalized eigenspaces. Invertibility on nonzero generalized eigenspaces follows because $f - \mu$ is nilpotent there, so $f = \mu(1 - (1 - f/\mu))$ is invertible. Spanning and independence are standard results for generalized eigenspaces over algebraically closed fields. In [SPGWC10] and [Wol12, Lemma 6.3(b)], the Jordan normal form is used directly. The argument is phrased in terms of generalized eigenspaces instead. $\square$

Theorem 7.10 (Nilpotent power bound). *If f is nilpotent on a space of dimension n , then $f^n = 0$.*

Proof. A nilpotent endomorphism on an n -dimensional space has nilpotency index $\leq n$. $\square$

7.3 Nonzero trace product

Theorem 7.11 (Cumulative span reaches $M_D(\mathbb{C})$ — explicit bound). *If A is normal, then there exists $n \leq D^2$ with $T_n(A) = M_D(\mathbb{C})$.*

Proof. By Theorem 7.6, if $T_n \neq T_{n+1}$ then $\dim T_{n+1} > \dim T_n$. Since $\dim T_n \leq D^2$ (Lemma 7.4), after at most D^2 steps either $T_n = M_D(\mathbb{C})$ or T_n has stabilised. Stabilisation before reaching $M_D(\mathbb{C})$ contradicts normality (which requires $S_{L_0} = M_D(\mathbb{C})$ for some L_0 , hence $T_{L_0} = M_D(\mathbb{C})$). $\square$

Theorem 7.12 (Cumulative span reaches $M_D(\mathbb{C})$). *If A is normal, then $T_{D^2}(A) = M_D(\mathbb{C})$.*

Proof. Take n from Theorem 7.11; since $n \leq D^2$ and T is monotone, $T_{D^2} = M_D(\mathbb{C})$. $\square$

Theorem 7.13 (Nonzero trace product). *If A is normal, then there exists a word w of length $\leq D^2$ such that $\text{tr}(A^w) \neq 0$. This is [SPGWC10, Lemma 1].*

Proof. By Theorem 7.12, $\mathbb{1} \in T_{D^2}(A)$, so $\mathbb{1}$ is a linear combination of word products of length $\leq D^2$. At least one has nonzero trace (since $\text{tr}(\mathbb{1}) = D \neq 0$). $\square$

7.4 Eigenvector extraction

Theorem 7.14 (Nonzero trace implies eigenvector). *If $M \in M_D(\mathbb{C})$ has $\text{tr}(M) \neq 0$, then M has a nonzero eigenvalue $\mu \neq 0$ and a corresponding eigenvector $\varphi \neq 0$ with $M\varphi = \mu\varphi$.*

Proof. A matrix with nonzero trace has a nonzero eigenvalue (since trace = sum of eigenvalues counted with multiplicity). The Fitting decomposition (Theorem 7.9) ensures the corresponding generalized eigenspace is nontrivial, yielding an eigenvector. $\square$

Theorem 7.15 (Eigenvalue extraction). *If A is normal, then there exist a word w_0 with $|w_0| \leq D^2$, a nonzero scalar $\mu \neq 0$, and a nonzero vector $\varphi \neq 0$ such that $A^{w_0}\varphi = \mu\varphi$.*

Proof. By Theorem 7.13, there is a word w_0 with $\text{tr}(A^{w_0}) \neq 0$ and $|w_0| \leq D^2$. By Theorem 7.14, A^{w_0} has a nonzero eigenvector. $\square$

7.5 Eigenvector spreading

Definition 7.16 (Cumulative vector span). Given an MPS tensor A and a vector $\varphi \in \mathbb{C}^D$, the *cumulative vector span* is

$$K_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w\varphi : |w| \leq n\} \subseteq \mathbb{C}^D.$$

This is [SPGWC10, proof of Lemma 2(a)].

Theorem 7.17 (Vector span stabilisation). *If $K_n(A, \varphi) = K_{n+1}(A, \varphi)$, then $K_m(A, \varphi) = K_n(A, \varphi)$ for all $m \geq n$.*

Proof. Same argument as Theorem 7.5: left-multiplication by A^i cannot escape the stabilised span. $\square$

Lemma 7.18 (Vector span from matrix span). *If $T_N(A) = M_D(\mathbb{C})$ and $\varphi \neq 0$, then $K_N(A, \varphi) = \mathbb{C}^D$.*

Proof. Every matrix $M \in M_D(\mathbb{C})$ is in $T_N(A)$, so in particular every rank-one matrix sending φ to any target vector v is in T_N . Applying such matrices to φ recovers all of $\mathbb{C}^D$. $\square$

Theorem 7.19 (Eigenvector spreading). *Let A be a normal MPS tensor and let $A^{i_0}\varphi = \mu\varphi$ with $\mu \neq 0$ and $\varphi \neq 0$. Then $K_{D-1}(A, \varphi) = \mathbb{C}^D$. This is [SPGWC10, Lemma 2(a)].*

Proof. The vector span K_n is monotone and $\dim K_n \leq D$. If $K_n = K_{n+1}$ for some $n < D-1$, the span stabilises (Theorem 7.17). But $T_{D^2}(A) = M_D(\mathbb{C})$ by Theorem 7.12, so Lemma 7.18 gives $K_{D^2}(A, \varphi) = \mathbb{C}^D$, contradicting early stabilisation. Hence $\dim K_n$ grows by ≥ 1 at each step, reaching D by step $D-1$. $\square$

7.6 Assembly

Definition 7.20 (Wielandt analysis). A *Wielandt analysis* for a normal MPS tensor A packages: a word w_0 of length $\leq D^2$, a nonzero eigenvalue μ , and a nonzero eigenvector φ of the matrix A^{w_0} , such that $A^{w_0}\varphi = \mu\varphi$.

Theorem 7.21 (Wielandt analysis exists). *Every normal MPS tensor admits a Wielandt analysis.*

Proof. Immediate from Theorem 7.15. $\square$

Theorem 7.22 (Wielandt chain). *Every normal MPS tensor admits a Wielandt analysis consisting of data (w_0, μ, φ) with $K_{D-1}(A, \varphi) = \mathbb{C}^D$.*

Proof. Combine Theorems 7.21 and 7.19. $\square$

Theorem 7.23 (Wielandt bound). *If A is a normal MPS tensor with bond dimension D , then $T_{D^2}(A) = M_D(\mathbb{C})$. In particular, the cumulative span stabilises at index at most D^2 . This is [SPGWC10, Theorem 1] (which gives the sharper bound $D^2 - d + 1$ in certain cases; here we prove the D^2 bound).*

Proof. Immediate from Theorem 7.12. $\square$

Remark 7.24 (Explicit blocking-length estimates). Theorem 7.23 gives the D^2 cumulative-span bound. The sharper explicit blocking-length estimates from [SPGWC10] — at most D^4 for injectivity and at most $3D^5$ for the biCF form — are not currently formalized here.

7.7 Lemma 2(b): fixed-length matrix spanning

The preceding sections establish the cumulative-spanning part of the Lemma 2(a) argument from [SPGWC10]: eigenvector spreading gives a cumulative vector span $K_{D-1}(A, \varphi) = \mathbb{C}^D$. The next stage is to pass to a span at one fixed length, namely to find n_0 such that $S_{n_0}(A) = M_D(\mathbb{C})$ (i.e. word products of *exactly* one length span all matrices). The results below first formalize the assembly step and then derive a blocked fixed-length spanning theorem.

Definition 7.25 (Fixed-length vector span). The *fixed-length vector span* at length n is

$$H_n(A, \varphi) = \text{span}_{\mathbb{C}}\{A^w \varphi : |w| = n\} \subseteq \mathbb{C}^D.$$

This is $S_n(A)|\varphi\rangle$ in the notation of [SPGWC10].

Lemma 7.26 (Word span products). $S_m(A) \cdot S_n(A) \subseteq S_{m+n}(A)$: the product of word spans at lengths m and n is contained in the word span at length $m + n$.

Proof. Every product $A^{w_1} A^{w_2}$ with $|w_1| = m$ and $|w_2| = n$ equals $A^{w_1 w_2}$ with $|w_1 w_2| = m + n$. $\square$

Lemma 7.27 (Eigenvector padding). Suppose $A^{i_0} \varphi = \mu \varphi$ with $\mu \neq 0$. Then for all n ,

$$K_n(A, \varphi) = H_n(A, \varphi).$$

In particular, if $K_n(A, \varphi) = \mathbb{C}^D$ then already $H_n(A, \varphi) = \mathbb{C}^D$.

Proof. Any word w with $|w| \leq n$ can be padded to exact length n by appending $n - |w|$ copies of the index i_0 . The eigenvector relation gives $A^{w \cdot i_0^k} \varphi = \mu^k \cdot A^w \varphi$, so the padded vector is a nonzero scalar multiple of the original. Hence every generator of K_n lies in H_n . The reverse inclusion $H_n \leq K_n$ is immediate. $\square$

Theorem 7.28 (Vector spanning to matrix spanning (assembly step)). Assume:

1. $H_n(A, \varphi) = \mathbb{C}^D$ (length- n word products applied to φ span all of $\mathbb{C}^D$), and
2. for each standard basis vector e_j , the rank-one operator $|\varphi\rangle\langle e_j|$ lies in $S_m(A)$.

Then $S_{n+m}(A) = M_D(\mathbb{C})$.

Proof. Fix a matrix unit E_{ij} . By hypothesis (1), there exists $M_i \in S_n(A)$ with $M_i\varphi = e_i$. By hypothesis (2), $|\varphi\rangle\langle e_j| \in S_m(A)$. Their product satisfies

$$M_i \cdot |\varphi\rangle\langle e_j| = |M_i\varphi\rangle\langle e_j| = |e_i\rangle\langle e_j| = E_{ij},$$

and $E_{ij} \in S_{n+m}(A)$ by Lemma 7.26. Since the matrix units span $M_D(\mathbb{C})$, $S_{n+m}(A) = M_D(\mathbb{C})$. $\square$

Lemma 7.29 (Blocking transfer). *If the blocked tensor $A^{[L]}$ satisfies $S_n(A^{[L]}) = M_D(\mathbb{C})$, then $S_{nL}(A) = M_D(\mathbb{C})$.*

Proof. Each length- n word in the blocked alphabet $\{0, \dots, d^L - 1\}$ decodes to a length- nL word in the original alphabet $\{0, \dots, d - 1\}$, so $S_n(A^{[L]}) \subseteq S_{nL}(A)$. $\square$

Remark 7.30 (Relation with Lemma 2(b)). In [SPGWC10], Lemma 2(b) combines a rank-one extraction statement for fixed-length spans with an assembly argument turning fixed-length vector spanning into fixed-length matrix spanning. Theorem 7.28 formalizes the second step. Theorem 7.35 supplies the rank-one input in blocked form, and Theorem 7.36 then derives a blocked, fixed-length word-span saturation result from that pipeline.

7.8 Blocked tensor and rectangular span

This section extends the Lemma 2(b) infrastructure to the *blocked tensor* $A^{[L]}$ and develops the “rectangular span” used to produce rank-one elements.

Theorem 7.31 (Blocking preserves normality). *If A is normal and $L > 0$, then the blocked tensor $A^{[L]}$ is also normal.*

Proof. If $S_N(A) = M_D(\mathbb{C})$ then $S_{NL}(A) = M_D(\mathbb{C})$ (padding by repeated single-letter words). By Lemma 7.32, $S_{NL}(A) \subseteq S_N(A^{[L]})$, so $S_N(A^{[L]}) = M_D(\mathbb{C})$. $\square$

Lemma 7.32 (Chunking: word span embeds into blocked word span). *$S_{nL}(A) \subseteq S_n(A^{[L]})$ for all n, L .*

Proof. Every length- nL word in the original alphabet can be split into n consecutive chunks of length L , each of which is a single blocked Kraus operator. By induction on n , using the word span product lemma (Lemma 7.26). $\square$

Theorem 7.33 (Word eigenvector gives blocked eigenvector). *If $A^{w_0}\varphi = \mu\varphi$ for a word w_0 of length L , then $(A^{[L]})^{j_0}\varphi = \mu\varphi$, where j_0 is the encoded blocked index corresponding to w_0 .*

Proof. By definition, $(A^{[L]})^{j_0} = A^{w_0}$. $\square$

Theorem 7.34 (Wielandt blocked assembly). *Suppose A is normal and $L > 0$, and we have:*

1. *a word w_0 of length L with eigenvector $A^{w_0}\varphi = \mu\varphi$ ($\mu \neq 0$, $\varphi \neq 0$),*
2. *a word τ_0 of length L with transpose eigenvector $(A^{\tau_0})^T\psi = \nu\psi$ ($\nu \neq 0$, $\psi \neq 0$),*
3. *a rank-one element $\varphi\psi^T \in S_m(A^{[L]})$ for some m .*

Then $S_{((D-1)+m+(D-1)) \cdot L}(A) = M_D(\mathbb{C})$.

Proof. Apply the conditional Lemma 2(b) assembly (Theorem 7.28) to the blocked tensor $A^{[L]}$, which is normal by Theorem 7.31. The eigenvector and transpose eigenvector are transferred to single-index eigenvectors of $A^{[L]}$ by Theorem 7.33. Transfer back to the original tensor via Lemma 7.29. $\square$

Theorem 7.35 (Rank-one extraction for blocked tensor). *Suppose A is normal and $N_0 \geq 1$ with $S_{N_0}(A) = M_D(\mathbb{C})$. Then there exist words σ_0, τ_0 of length N_0 , nonzero vectors φ, ψ , nonzero scalars μ, ν , and $m \in \mathbb{N}$ such that:*

1. $A^{\sigma_0} \varphi = \mu \varphi$,
2. $(A^{\tau_0})^T \psi = \nu \psi$,
3. $\varphi \psi^T \in S_m(A^{[N_0]})$.

Proof. Set $B = A^{[N_0]}$. Since $S_{N_0}(A) = M_D(\mathbb{C})$, a nonzero-trace word exists (Theorem 7.13), yielding a column eigenvector φ via Theorem 7.14. A row eigenvector ψ is obtained analogously from the transpose tensor. Set $P = (B^{j_0})^D$ and $Q = (B^{k_0})^D$, where j_0, k_0 encode σ_0, τ_0 in the blocked alphabet. The Fitting decomposition ensures $\varphi \in \text{range}(P)$ and $\psi \in \text{range}(Q^T)$, so $\varphi \psi^T \in \text{range}(\text{mulLeft}(P) \circ \text{mulRight}(Q)) \subseteq S_{2D}(B)$. $\square$

Theorem 7.36 (Fixed-length word-span saturation via blocking). *If A is a normal MPS tensor with bond dimension D , then there exists N such that $S_N(A) = M_D(\mathbb{C})$. This gives the fixed-length spanning conclusion corresponding to [SPGWC10, Lemma 2(b)].*

Proof. By normality, some N_0 has $S_{N_0}(A) = M_D(\mathbb{C})$. If $N_0 = 0$ we are done. Otherwise, Theorem 7.35 produces eigenvectors and a rank-one element $\varphi \psi^T \in S_m(A^{[N_0]})$. Theorem 7.34 then gives $S_{((D-1)+m+(D-1)) \cdot N_0}(A) = M_D(\mathbb{C})$. $\square$

Remark 7.37 (Blocking route to fixed-length spanning). Theorem 7.36 is obtained by first extracting a rank-one operator in a blocked tensor (Theorem 7.35) and then applying the fixed-length assembly step (Theorem 7.34). Thus the fixed-length span of A is reached through a blocked auxiliary tensor rather than by a direct unblocked argument.

7.9 Primitivity and normality

Theorem 7.38 (Primitive MPS tensors have nonzero word products). *If A is a primitive MPS tensor, then for every $n \geq 0$ there exists a word w of length exactly n with $A^w \neq 0$.*

Proof. The PSD fixed point ρ of $\mathcal{E}_A$ satisfies $\mathcal{E}_A^n(\rho) = \rho \neq 0$. Since $\mathcal{E}_A^n(\rho) = \sum_{|w|=n} A^w \rho (A^w)^\dagger$, at least one summand is nonzero, hence some $A^w \neq 0$. $\square$

Theorem 7.39 (Iterated transfer does not vanish). *If A is a primitive MPS tensor, then $\mathcal{E}_A^n \neq 0$ for all $n \geq 0$.*

Proof. By Theorem 7.38, there exists a word w of length n with $A^w \neq 0$, so $\mathcal{E}_A^n(\rho)$ has a nonzero summand. $\square$

7.9.1 Primitivity-to-normality bridge

The spectral-gap notion of primitivity (unique PSD fixed point ρ with $\rho_{\text{spec}}(\mathcal{E}_A - P_\rho) < 1$) is connected to normality through the following results.

Theorem 7.40 (Fixed-point uniqueness from spectral gap). *If A is a primitive MPS tensor with PSD fixed point ρ , then every fixed point σ of $\mathcal{E}_A$ is proportional to ρ : $\sigma = \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$.*

Proof. Set $\sigma' = \sigma - \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Then $\text{tr}(\sigma') = 0$ and $(E - P_\rho)(\sigma') = \sigma'$. If $\sigma' \neq 0$, then 1 is an eigenvalue of $E - P_\rho$, contradicting the spectral gap $\rho_{\text{spec}}(E - P_\rho) < 1$. $\square$

Theorem 7.41 (Complement powers tend to zero). *If A is a primitive MPS tensor with PSD fixed point ρ , then $(\mathcal{E}_A - P_\rho)^n \rightarrow 0$ in operator norm.*

Proof. The spectral radius of $\mathcal{E}_A - P_\rho$ is strictly less than 1 by the spectral-gap hypothesis. Apply the general result that powers of an operator with spectral radius < 1 tend to zero. $\square$

Remark 7.42 (Primitivity: spectral gap vs. positive definiteness). The definition of primitivity used in §7.9.1 encodes only the spectral-gap condition: there exists a nonzero PSD fixed point ρ and $\rho_{\text{spec}}(\mathcal{E}_A - P_\rho) < 1$. The notion of “primitive” in [SPGWC10, Proposition 3] additionally requires ρ to be positive definite (full rank). Without positive definiteness, the spectral-gap condition alone does *not* imply irreducibility or normality. A counterexample is $D = 2$, $\rho = \text{diag}(1, 0)$.

With the additional hypothesis $\rho > 0$, Theorem 7.43 gives irreducibility. Together with the Burnside bridge (Chapter 8, §8.4.1) and Theorem 7.49, one obtains

$$\text{primitive} + \rho \text{ PosDef} \Rightarrow \text{irreducible} \Rightarrow \text{algSpan}(A) = M_D(\mathbb{C}) \Rightarrow \text{normal}.$$

The last step uses the aperiodicity condition $\mathbb{1} \in S_1(A)$ from Remark 7.44.

7.9.2 Primitive implies irreducible

Theorem 7.43 (Primitive with PosDef fixed point implies irreducible tensor). *If A is a primitive MPS tensor with PSD fixed point ρ and ρ is positive definite, then A is an irreducible tensor. This is part of [SPGWC10, Proposition 3] (see also [CPGSV17, §2.3]).*

Proof. By contrapositive. Assume A has an invariant projection P ($P \neq 0$, $P \neq \mathbb{1}$, $(\mathbb{1} - P)A^i P = 0$ for all i). Set $\sigma = P\rho P$.

1. *Invariance of σ under $\mathcal{E}_A^n$:* the projection relation $(\mathbb{1} - P)A^i P = 0$ implies $(\mathbb{1} - P) \cdot \mathcal{E}_A^n(\sigma) = 0$ for all n (by induction on n).
2. *Convergence:* By Theorem 7.41, $\mathcal{E}_A^n(\sigma) \rightarrow \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho$. Hence $(\mathbb{1} - P) \cdot \frac{\text{tr}(\sigma)}{\text{tr}(\rho)}\rho = 0$.
3. *Nonzero scalar:* Since $P \neq 0$ and ρ is positive definite, $\text{tr}(P\rho P) > 0$, so $\frac{\text{tr}(\sigma)}{\text{tr}(\rho)} \neq 0$. Therefore $(\mathbb{1} - P)\rho = 0$.
4. *Contradiction:* Since ρ is positive definite, it is a unit in $M_D(\mathbb{C})$. From $(\mathbb{1} - P)\rho = 0$ we get $\mathbb{1} - P = 0$, i.e. $P = \mathbb{1}$, contradicting $P \neq \mathbb{1}$.

$\square$

7.10 Cumulative span to word span

Cumulative spanning ($T_N(A) = M_D(\mathbb{C})$, the union of word spans of all lengths $\leq N$) is weaker than normality ($S_L(A) = M_D(\mathbb{C})$ for a single length L). The gap is closed by an aperiodicity hypothesis.

Remark 7.44 (Aperiodicity condition). We say A satisfies the *aperiodicity condition* if $\mathbb{1} \in S_1(A)$, i.e. the identity matrix lies in the span of the Kraus operators $\{A^i\}$. This is an additional hypothesis. The one-sided TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ does *not* imply $\mathbb{1} \in S_1(A)$; it only gives a quadratic identity among the Kraus operators. For genuinely primitive tensors one expects aperiodicity to follow from peripheral-spectrum analysis, but that implication is not used in the present Wielandt bridge, so the condition is kept explicit.

Remark 7.45 (Counterexample: cumulative spanning does not imply normality). In $M_2(\mathbb{C})$, let $A^0 = E_{12}$ and $A^1 = E_{21}$ (the off-diagonal matrix units). Then $\text{algSpan}(A) = M_2(\mathbb{C})$ and $T_2(A) = M_2(\mathbb{C})$, but the word spans alternate: $S_n(A)$ contains only off-diagonal matrices for odd n and only diagonal matrices for even n . Hence $S_n(A) \neq M_2(\mathbb{C})$ for every n , and A is not normal. The aperiodicity condition $\mathbb{1} \in S_1(A)$ fails: $S_1(A) = \text{span}_{\mathbb{C}}\{E_{12}, E_{21}\}$.

Theorem 7.46 (Word span monotonicity from aperiodicity). *If $\mathbb{1} \in S_1(A)$, then $S_n(A) \leq S_{n+1}(A)$ for all n .*

Proof. For any $M \in S_n(A)$, $M = M \cdot \mathbb{1} \in S_n(A) \cdot S_1(A) \subseteq S_{n+1}(A)$ by Lemma 7.26. $\square$

Theorem 7.47 (Cumulative span equals word span under aperiodicity). *If $\mathbb{1} \in S_1(A)$, then $T_n(A) = S_n(A)$ for all n .*

Proof. By Theorem 7.46, the word spans are monotone: $S_0 \leq S_1 \leq \dots$. The supremum $T_n = S_0 + \dots + S_n$ therefore equals the largest term S_n . $\square$

Theorem 7.48 (Cumulative spanning plus aperiodicity implies normality). *If $T_N(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. By Theorem 7.47, $S_N(A) = T_N(A) = M_D(\mathbb{C})$. $\square$

Theorem 7.49 (Algebra span plus aperiodicity implies normality). *If $\text{algSpan}(A) = M_D(\mathbb{C})$ and $\mathbb{1} \in S_1(A)$, then A is normal.*

Proof. Since $\text{algSpan}(A) = M_D(\mathbb{C})$, there exists N with $T_N(A) = M_D(\mathbb{C})$ (by the ascending chain condition on finite-dimensional subspaces). Apply Theorem 7.48. $\square$

Theorem 7.50 (Irreducible tensor plus aperiodicity implies normality). *If A is an irreducible tensor, $D > 0$, and $\mathbb{1} \in S_1(A)$, then A is normal. This assembles the full chain:*

$$\text{irreducible tensor} \xrightarrow{8.20} \text{irreducible action} \xrightarrow{\text{Burnside}} \text{algSpan} = M_D(\mathbb{C}) \xrightarrow{7.49} \text{normal}.$$

This corresponds to [SPGWC10, Theorem 1 / Proposition 3].

Proof. By Theorem 8.20, A acts irreducibly on $\mathbb{C}^D$. By Theorem 8.21, $\text{algSpan}(A) = M_D(\mathbb{C})$. By Theorem 7.49 with the aperiodicity hypothesis, A is normal. $\square$

Chapter 8

Canonical Form Reduction

This chapter reduces a general normal MPS tensor to an injective block-diagonal form via iterated invariant-subspace decomposition. The reduction procedure follows [PGVWC07], [CPGSV21], and [Wol12, Ch. 6].

8.1 Block-diagonal assembly

Definition 8.1 (Block-diagonal tensor from blocks). Given r blocks with bond dimensions $(D_k)_{k=1}^r$, per-block MPS tensors $\{A_k^i\}$, and scaling factors $(\mu_k)_{k=1}^r$, define the block-diagonal tensor $A^i = \bigoplus_{k=1}^r \mu_k A_k^i$ of total bond dimension $D = \sum_k D_k$.

Theorem 8.2 (Per-block single-block FT). *If all block tensors A_k are injective and the per-block same-MPV condition holds (A_k and B_k generate the same MPV family for each k), then each pair (A_k, B_k) is gauge equivalent.*

Proof. Apply the single-block fundamental theorem (Theorem 3.11) to each block independently. $\square$

Theorem 8.3 (Per-block SameMPV implies global SameMPV). *If A_k and B_k generate the same MPV family for each block k , then the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ also generate the same MPV family.*

Proof. By the MPV decomposition (Theorem 2.24), each MPV is $\sum_k \mu_k^N V^{(N)}(A_k)_\sigma$. Per-block equality of MPV coefficients gives global equality. $\square$

Theorem 8.4 (Global gauge from block gauge). *If each pair (A_k, B_k) is gauge equivalent ($B_k^i = X_k A_k^i X_k^{-1}$), then the block-diagonal tensors are gauge equivalent via the block-diagonal matrix $X = \bigoplus_k X_k$.*

Proof. Block-diagonal conjugation: $X(\bigoplus_k \mu_k A_k^i)X^{-1} = \bigoplus_k \mu_k X_k A_k^i X_k^{-1} = \bigoplus_k \mu_k B_k^i$. $\square$

8.2 Transfer map normalization

Theorem 8.5 (Scaling preserves injectivity). *If A is injective and $\zeta \neq 0$, then ζA is injective.*

Proof. Scaling by a nonzero scalar preserves linear independence and hence spanning. $\square$

Theorem 8.6 (Transfer map under scaling). $\mathcal{E}_{\zeta A}(X) = |\zeta|^2 \mathcal{E}_A(X)$.

Proof. Each summand: $(\zeta A^i)X(\zeta A^i)^\dagger = |\zeta|^2 A^i X(A^i)^\dagger$. $\square$

Theorem 8.7 (Phase-scaling preserves normalization). *If $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $|\zeta| = 1$, then $\sum_i (\zeta A^i)^\dagger (\zeta A^i) = \mathbb{1}$.*

Proof. $(\zeta A^i)^\dagger (\zeta A^i) = \bar{\zeta} \zeta (A^i)^\dagger A^i = (A^i)^\dagger A^i$, since $\bar{\zeta} \zeta = |\zeta|^2 = 1$. $\square$

Theorem 8.8 (MPV under block normalization). *For any nonzero scaling factors μ_k , write $\mu_k = |\mu_k| \cdot (\mu_k/|\mu_k|)$. The MPVs of $\bigoplus_k \mu_k A_k$ and $\bigoplus_k |\mu_k| \cdot (\mu_k/|\mu_k|) A_k$ agree.*

Proof. By the MPV decomposition (Theorem 2.24), the k -th term is $\mu_k^N V^{(N)}(A_k)_\sigma$. The scaling identity (Lemma 2.12) absorbs the phase into the word evaluation. $\square$

8.3 Invariant subspace decomposition

Theorem 8.9 (Unitary conjugation preserves SameMPV). *If U is a unitary matrix, then A and $U^\dagger A U$ generate the same MPV family.*

Proof. By cyclicity of the trace: $\text{tr}((U^\dagger A^{i_1} U) \dots (U^\dagger A^{i_N} U)) = \text{tr}(A^{i_1} \dots A^{i_N})$. $\square$

Definition 8.10 (Support projection). For a positive semi-definite matrix $\rho \geq 0$, the *support projection* P is the orthogonal projection onto the range of ρ . It is constructed via the spectral decomposition: if $\rho = U \text{diag}(\lambda_1, \dots, \lambda_D) U^\dagger$, then $P = U \text{diag}(\mathbf{1}_{\lambda_1 > 0}, \dots, \mathbf{1}_{\lambda_D > 0}) U^\dagger$, where $\mathbf{1}_{\lambda_j > 0}$ is 1 if $\lambda_j > 0$ and 0 otherwise.

Definition 8.11 (Invariant projection predicate). An MPS tensor A has an *invariant projection* if there exists an orthogonal projection P with $P \neq 0$, $P \neq \mathbb{1}$, and $(\mathbb{1} - P)A^i P = 0$ for all i . The negation of this predicate defines an *irreducible tensor* (Definition 8.14).

Theorem 8.12 (PSD fixed point gives invariant projection). *Let $\rho \geq 0$ be a fixed point of $\mathcal{E}_A$, and let P be the support projection of ρ . Then for all i , $(\mathbb{1} - P)A^i P = 0$: the range of P is invariant under all A^i .*

Proof. From $\mathcal{E}_A(\rho) = \rho$ and positivity of each summand $A^i \rho (A^i)^\dagger$, $(\mathbb{1} - P)A^i \rho (A^i)^\dagger (\mathbb{1} - P) = 0$ for each i . Since ρ is supported on the range of P , this forces $(\mathbb{1} - P)A^i P = 0$.

This is the finite-dimensional version of [Wol12, Prop. 6.10] (“Stationary subspaces”: the support projection of any stationary state is sub-harmonic, i.e. invariant under the channel). The equivalence between irreducibility and absence of nontrivial invariant projections is [Wol12, Thm. 6.2]; see also [CPGSV21, §IV.A.1]. $\square$

Theorem 8.13 (Two-block decomposition). *If the transfer map has a PSD fixed point ρ that is not positive definite (i.e. ρ has a nontrivial kernel), then A is unitarily equivalent to a two-block upper-triangular tensor. Moreover, the diagonal blocks generate the same MPV as A .*

Proof. The support projection P of ρ is neither 0 nor $\mathbb{1}$ (since $\rho \neq 0$ and ρ is not PD). Theorem 8.12 gives $(\mathbb{1} - P)A^i P = 0$ for all i . A unitary change of basis that block-diagonalizes P puts A^i in upper-triangular block form (Theorem 8.9 ensures the MPV is preserved by the conjugation). The trace of the upper-triangular product equals the sum of the block-diagonal traces, so the MPV decomposes into block contributions. $\square$

8.4 Iterated reduction

Definition 8.14 (Irreducible tensor). An MPS tensor A is *irreducible* (in the sense of [CPGSV21]) if the transfer map $\mathcal{E}_A$ has no nontrivial invariant projection, i.e. $\mathcal{E}_A$ is irreducible (Definition 4.23).

Theorem 8.15 (Irreducible block decomposition). *Every MPS tensor A can be decomposed (up to unitary equivalence) into a block-diagonal tensor $\bigoplus_{k=1}^r A_k$ where each A_k is irreducible. The block-diagonal tensor generates the same MPV as A .*

Proof. By strong induction on the bond dimension D . If A is already irreducible, take $r = 1$. Otherwise, the transfer map has a PSD fixed point that is not PD (by the Cesàro argument), and Theorem 8.13 splits A into two blocks of strictly smaller bond dimension. Apply the induction hypothesis to each block and concatenate.

Both [PGVWC07] (Theorem 4, “TI canonical form”) and [CPGSV21] (§IV.A.1) obtain the block decomposition by the same invariant-projection splitting used here. An alternative route, developed in [Wol12, Thm. 6.14], characterises the fixed-point set of a quantum channel as a $*$ -algebra and then applies the Wedderburn–Artin decomposition ([Wol12, Eq. (1.39)]) to read off the block structure directly. The block decomposition is obtained by iterating the invariant-projection splitting and using strong induction on D to ensure termination. $\square$

Theorem 8.16 (Unitary diagonal positive fixed point in TP gauge). *Let A be an irreducible MPS tensor satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, and assume $D > 0$. Then there exist a unitary U and a diagonal positive definite matrix Λ such that, for $B^i := U^\dagger A^i U$,*

$$\sum_i (B^i)^\dagger B^i = \mathbb{1} \quad \text{and} \quad \mathcal{E}_B(\Lambda) = \Lambda.$$

Proof. The TP hypothesis makes $\mathcal{E}_A$ a channel, so it has a nonzero PSD fixed point. The irreducible tensor condition implies that $\mathcal{E}_A$ is irreducible as a CP map, and Theorem 5.3 upgrades the fixed point to a positive definite one. The spectral theorem diagonalizes this matrix by a unitary conjugation, and unitary conjugation preserves the TP normalization. $\square$

8.4.1 Burnside bridge: irreducibility and spanning

The irreducible-tensor predicate (Definition 8.14) is formulated in terms of invariant *projections*. For Burnside-style arguments it is more natural to work with invariant *subspaces* of $\mathbb{C}^D$ under the Kraus operators.

Definition 8.17 (Algebra span). For an MPS tensor A , let $\text{algSpan}(A)$ be the unital $\mathbb{C}$ -subalgebra of $M_D(\mathbb{C})$ generated by the matrices $\{A^i\}$:

$$\text{algSpan}(A) = \mathbb{C}\langle A^i : i = 0, \dots, d-1 \rangle \subseteq M_D(\mathbb{C}).$$

Definition 8.18 (Invariant submodule). A subspace $W \leq \mathbb{C}^D$ is *A-invariant* if $A^i W \subseteq W$ for every i .

Definition 8.19 (Irreducible action). The Kraus operators of A act *irreducibly* on $\mathbb{C}^D$ if every A -invariant subspace is trivial, i.e. equal to 0 or to $\mathbb{C}^D$.

Theorem 8.20 (Irreducible tensor implies irreducible action). *If A is irreducible as a tensor (Definition 8.14), then the Kraus operators act irreducibly on $\mathbb{C}^D$ (Definition 8.19).*

Proof. Let $W \leq \mathbb{C}^D$ be an A -invariant subspace. In finite dimension the orthogonal projection P onto W exists. Invariance of W implies $(\mathbb{1} - P)A^i P = 0$ for all i . Thus P is a nontrivial invariant projection unless $W = 0$ or $W = \mathbb{C}^D$, contradicting irreducibility of the tensor. $\square$

Theorem 8.21 (Burnside's theorem for Kraus algebras). *Assume $D > 0$. If A acts irreducibly on $\mathbb{C}^D$, then the generated unital subalgebra is the full matrix algebra:*

$$\text{algSpan}(A) = M_D(\mathbb{C}).$$

This is the complex finite-dimensional case of Burnside's theorem (equivalently, Jacobson's density theorem); see [Jac09].

Proof. Transport $\text{algSpan}(A)$ along the algebra equivalence $M_D(\mathbb{C}) \simeq_{\mathbb{C}} \text{End}_{\mathbb{C}}(\mathbb{C}^D)$. The irreducible action hypothesis makes $\mathbb{C}^D$ a simple module for this algebra. Over the algebraically closed field $\mathbb{C}$, Schur's lemma identifies the endomorphisms commuting with the action with scalars, and Jacobson density then forces the action map to be surjective onto all of $\text{End}_{\mathbb{C}}(\mathbb{C}^D)$. $\square$

Theorem 8.22 (Irreducible action implies eventual cumulative spanning). *Assume $D > 0$. If A acts irreducibly on $\mathbb{C}^D$, then there exists N such that $T_N(A) = M_D(\mathbb{C})$.*

Proof. By Theorem 8.21, $\text{algSpan}(A) = M_D(\mathbb{C})$. Every element of the algebra span is a linear combination of word products, hence belongs to $T_n(A)$ for some n . Since $M_D(\mathbb{C})$ is finite-dimensional, the ascending chain $T_0(A) \leq T_1(A) \leq \dots$ stabilizes, so one can choose a uniform N with $T_N(A) = M_D(\mathbb{C})$. $\square$

8.5 Proportional single-block theorem

Theorem 8.23 (Proportional MPV implies gauge-phase equivalence). *Let A and B be injective MPS tensors satisfying $\sum_i (A^i)^\dagger A^i = \mathbb{1}$ and $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. If A and B generate proportional MPV families $(V^{(N)}(A))_\sigma = c_N V^{(N)}(B)_\sigma$ for some $c_N \in \mathbb{C}$ and $O_{AA}(N) \rightarrow 1$, $O_{BB}(N) \rightarrow 1$, then A and B are gauge-phase equivalent.*

Proof. By contradiction. If A and B are not gauge-phase equivalent, then $O_{AB}(N) \rightarrow 0$ (Theorem 6.17). From the proportionality hypothesis, $O_{AB}(N) = c_N O_{BB}(N)$. Since $O_{BB}(N) \rightarrow 1$, it follows that $c_N \rightarrow 0$. But $O_{AA}(N) = |c_N|^2 O_{BB}(N) \rightarrow 0$, contradicting $O_{AA}(N) \rightarrow 1$.

In [PGVWC07], the proportional case is handled as part of the canonical form argument. Thus the proportional case is obtained directly from the spectral gap in Chapter 6. $\square$

8.6 Canonical form from primitivity

The canonical form predicate (Definition 9.8) includes an explicit hypothesis that the self-overlap $O_{A_k A_k}(N) \rightarrow 1$ for each block. The following theorem derives this from per-block primitivity, removing it as an independent assumption.

Theorem 8.24 (Canonical form from primitivity). *Let $(\mu_k, A_k)_{k=1}^r$ be a collection of block tensors and scaling factors satisfying:*

1. *each A_k is injective,*
2. *each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,*
3. *$|\mu_1| > \dots > |\mu_r|$ (strictly decreasing),*

4. all $\mu_k \neq 0$,

5. each A_k is primitive (the transfer map $\mathcal{E}_{A_k}$ has peripheral spectrum $\{1\}$).

Then (μ_k, A_k) satisfies the canonical form predicate (Definition 9.8).

Proof. Conditions (1)–(4) are passed through directly. For condition (5) of the canonical form predicate ($O_{A_k A_k}(N) \rightarrow 1$), primitivity of $\mathcal{E}_{A_k}$ implies $\mathcal{E}_{A_k}^N(\rho) \rightarrow \rho_\infty$ for the unique fixed-point state ρ_∞ , and hence the self-overlap converges to 1. The proof applies primitive overlap convergence to each block. $\square$

Remark 8.25. Theorem 8.24 shows that the overlap convergence hypothesis in the canonical form predicate is redundant once primitivity is known. In the paper [CPGSV21], primitivity is obtained from irreducibility plus periodicity removal (blocking to period one). Theorem 8.26 establishes this periodicity-removal step: under irreducibility and TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$, one can block until the transfer map becomes primitive. This corresponds to [CPGSV17, Appendix A].

Theorem 8.26 (Blocking yields a primitive transfer map). *Let A be an irreducible MPS tensor with $D > 0$. Assume that A satisfies the TP normalization $\sum_i (A^i)^\dagger A^i = \mathbb{1}$. Then there exists a blocking length $p > 0$ such that the transfer map $\mathcal{E}_{A^{[p]}}$ of the blocked tensor $A^{[p]}$ is primitive.*

Proof. Pass to the conjugate-transposed Kraus family $K_i := (A^i)^\dagger$. The TP normalization of A makes K unital, and irreducibility of A gives irreducibility of the associated Kraus map. The TP fixed-point theorem for $\mathcal{E}_A$ provides a positive definite fixed point for the adjoint map of K , so Theorem 4.46 shows that every peripheral eigenvalue is a root of unity. Since $\text{peripheral}(\mathcal{E}_A)$ is finite (Theorem 4.37), Lemma 4.41 gives a common exponent p such that $\mu^p = 1$ for all $\mu \in \text{peripheral}(\mathcal{E}_A)$. Lemma 4.42 then implies $\text{peripheral}(\mathcal{E}_A^p) = \{1\}$. Finally, blocking corresponds to taking a power of the transfer map, so $\mathcal{E}_{A^{[p]}}$ is primitive. $\square$

Remark 8.27. Theorem 8.26 is the “periodicity removal by blocking” step in [CPGSV17, Appendix A] for an irreducible block already placed in TP gauge. The later pipeline wrapper simply repackages this step.

8.7 Canonical form existence pipeline

This section assembles the components of the canonical-form construction following [CPGSV17, §2.3 and Appendix A].

Theorem 8.28 (Irreducible block decomposition (pipeline)). *Every MPS tensor A generates the same MPV family as a block-diagonal tensor $\bigoplus_{k=1}^r A_k$ with each block irreducible.*

Proof. Apply Theorem 8.15. $\square$

Theorem 8.29 (CFII data for irreducible TP blocks). *Let A be an irreducible MPS tensor already in TP gauge ($\sum_i (A^i)^\dagger A^i = \mathbb{1}$) with $D > 0$. Then there exist a unitary U and a diagonal positive definite matrix Λ such that:*

1. $B^i := U^\dagger A^i U$ is TP,
2. $\mathcal{E}_B(\Lambda) = \Lambda$,
3. Λ is diagonal and positive definite,

4. A and B generate the same MPV family.

This is the “Canonical Form II” (CFII) step of [CPGSV17, Appendix A]. The unitary conjugation occurs only after TP normalization has already been achieved; it is a second step inside TP gauge, not a unitary normalization of an arbitrary irreducible block.

Proof. Apply Theorem 8.16 to obtain the unitary and the diagonal PD fixed point. Unitary conjugation preserves trace-preservation and the MPV family (Theorem 8.9). $\square$

Theorem 8.30 (Pipeline handoff: block decomposition with conditional CFII). *For any MPS tensor A , there exist irreducible blocks $(A_k)_{k=1}^r$ that generate the same MPV family as A . Moreover, for each block k , if that block has already been placed in TP gauge and $D_k > 0$, one obtains CFII data (unitary, diagonal PD fixed point, MPV preservation).*

Proof. Combine Theorems 8.28 and 8.29. $\square$

8.8 Perron–Frobenius eigenvector existence

This section establishes the existence of a positive definite eigenvector for the adjoint transfer map of an irreducible MPS tensor, and uses it to construct a TP-gauge normalization. The results correspond to [EHK78], [Wol12, Ch. 6], and [CPGSV17, Appendix A].

Theorem 8.31 (PSD eigenvector for positive maps). *Let $E : M_D(\mathbb{C}) \rightarrow M_D(\mathbb{C})$ be a positive linear map with $D > 0$, and assume that $E(\sigma) \neq 0$ for every nonzero PSD matrix σ . Then there exist a nonzero PSD matrix ρ and a positive real $r > 0$ such that $E(\rho) = r\rho$.*

This is the finite-dimensional Perron–Frobenius theorem for positive maps on matrix algebras ([EHK78], [Wol12, Proposition 6.1]). The Lean proof applies Brouwer’s fixed-point theorem to the normalization map $\rho \mapsto E(\rho)/\text{tr}(E(\rho))$ on the compact convex set of density matrices; the required density-matrix Brouwer theorem is now itself formalized.

Theorem 8.32 (Scaling preserves irreducibility). *If E is an irreducible map and $c \neq 0$, then cE is also irreducible.*

Proof. An invariant projection for cE is an invariant projection for E (scale out the nonzero constant from the commutation relation). $\square$

Theorem 8.33 (Adjoint transfer map is nonzero). *If some Kraus operator $A^i \neq 0$, then the adjoint transfer map $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is nonzero.*

Proof. If $\mathcal{E}_A^\dagger = 0$ then $\mathcal{E}_A^\dagger(\mathbb{1}) = 0$, so $\sum_i (A^i)^\dagger A^i = 0$. Since each summand $(A^i)^\dagger A^i$ is PSD, each $(A^i)^\dagger = 0$ and hence each $A^i = 0$. $\square$

Theorem 8.34 (PosDef adjoint eigenvector for irreducible tensors). *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive definite matrix σ and a positive real $r > 0$ such that $\mathcal{E}_A^\dagger(\sigma) = r\sigma$.*

This is the adjoint-map Perron–Frobenius step of [CPGSV17, Appendix A]: after applying Theorem 8.31 to the adjoint transfer map, one upgrades the resulting PSD eigenvector to a positive definite one.

Proof. By Theorem 8.33, $\mathcal{E}_A^\dagger \neq 0$. Since $\mathcal{E}_A^\dagger(X) = \sum_i (A^i)^\dagger X A^i$ is a Kraus map, it is positive. Irreducibility of A transfers to the adjoint transfer map, and together with $\mathcal{E}_A^\dagger \neq 0$ this implies $\mathcal{E}_A^\dagger(\tau) \neq 0$ for every nonzero PSD matrix τ . Hence Theorem 8.31 applies to $\mathcal{E}_A^\dagger$, giving $\sigma_0 \geq 0$ (PSD, nonzero) and $r > 0$ with $\mathcal{E}_A^\dagger(\sigma_0) = r\sigma_0$. Rescale: set $T^i = r^{-1/2}(A^i)^\dagger$. Then $\mathcal{E}_T(\sigma_0) = \sigma_0$, and $\mathcal{E}_T$ is irreducible (by Theorem 8.32, since irreducibility of $\mathcal{E}_A$ transfers to the adjoint and is preserved under scaling). Since $\mathcal{E}_T$ is irreducible with a PSD fixed point, the PSD-to-PosDef upgrade (Theorem 5.3) gives σ_0 positive definite. $\square$

Theorem 8.35 (TP-gauge existence for irreducible tensors). *Let A be an irreducible MPS tensor with $D > 0$ and some $A^i \neq 0$. Then there exist a positive real r , a positive definite matrix σ , and a gauge-equivalent tensor B such that*

$$B^i = \sigma^{1/2} \cdot (r^{-1/2} A^i) \cdot \sigma^{-1/2}$$

is trace-preserving: $\sum_i (B^i)^\dagger B^i = \mathbb{1}$.

This is the “spectral rescaling + TP gauge” step of [CPGSV17, Appendix A], now formalized using Theorem 8.34 and the explicit TP gauge construction. This similarity is generally non-unitary. The subsequent CFII step applies an additional unitary conjugation to this already TP-normalized representative.

Proof. By Theorem 8.34, obtain σ positive definite and $r > 0$ with $\mathcal{E}_A^\dagger(\sigma) = r\sigma$. Set $c = r^{-1/2}$ and $A'^i = cA^i$. Then $\mathcal{E}_{A'}^\dagger(\sigma) = \sigma$. By the TP gauge construction (Theorem 5.10), $B^i = \sigma^{1/2} A'^i \sigma^{-1/2}$ satisfies $\sum_i (B^i)^\dagger B^i = \mathbb{1}$. $\square$

8.9 End-to-end pipeline status

Remark 8.36 (Canonical form pipeline). The Appendix-A Perron–Frobenius step is no longer the remaining issue: Theorems 8.31 and 8.35 are now formalized, as are the CFII and periodicity-removal components (Theorems 8.29 and 8.26). The missing work is later in the end-to-end assembly from these ingredients to the repository’s final canonical-form predicates.

More precisely, the current pipeline proves the following steps.

1. Irreducible block decomposition (Theorem 8.15): $A \sim \bigoplus_k A_k$ with each A_k irreducible.
2. Perron–Frobenius eigenvector (Theorem 8.31): the adjoint transfer map has a PSD eigenvector.
3. TP gauge (Theorem 8.35): rescale and apply a generally non-unitary similarity to reach TP normalization.
4. CFII data (Theorem 8.29): inside this TP gauge, a further unitary conjugation diagonalizes the positive fixed point.
5. Periodicity removal (Theorem 8.26): block the TP-normalized irreducible representative until the transfer map is primitive.

What is still not assembled automatically is the passage from this primitive blocked TP/CFII output to the stronger hypotheses used by the repository’s canonical-form and FT theorems. In particular, Theorem 8.24 requires injective TP-normalized blocks, while the current pipeline does not yet supply the full primitive $\Rightarrow$ block-injective / normality bridge from Chapter 7. The relevant ingredients there (Theorems 7.43 and 7.49) still need extra hypotheses to turn primitive

blocked tensors into the repository’s stronger canonical-form notion. As a consequence, there is still no theorem taking an arbitrary tensor all the way to the final repository canonical form / biCF output matching [CPGSV17, §2.3 and Appendix A]. Moreover, Chapter 10 still marks an additional strict-dominance / oscillatory coefficient gap in the proportional multi-block assembly.

Chapter 9

Block Permutation and Separation

This chapter establishes the algebraic structure of automorphisms of products of matrix algebras and uses it to prove that the same-MPV condition on block-diagonal tensors descends to per-block gauge equivalence. The approach follows [PGVWC07] and [CPGSV21].

9.1 Block permutation algebra

Theorem 9.1 (Ring isomorphisms permute block ideals). *Let $\{R_k\}_{k=1}^r$ be simple rings. Every ring isomorphism $T : \prod_k R_k \xrightarrow{\sim} \prod_k R_k$ permutes the block ideals: there exists a permutation σ such that T maps the k -th block ideal to the $\sigma(k)$ -th block ideal.*

Proof. Each block ideal $I_k = \{x : x_j = 0 \text{ for } j \neq k\}$ is an atom in the lattice of two-sided ideals (since R_k is simple). A ring isomorphism preserves the ideal lattice and hence permutes the atoms. Injectivity of the induced map on atoms gives the permutation σ . $\square$

Theorem 9.2 (Dimension preservation). *If T is an algebra automorphism of $\prod_{k=1}^r M_{D_k}(\mathbb{C})$, and σ is the induced permutation, then $D_{\sigma(k)} = D_k$ for all k .*

Proof. The restriction of T to the k -th block ideal induces a $\mathbb{C}$ -algebra isomorphism $M_{D_k}(\mathbb{C}) \cong M_{D_{\sigma(k)}}(\mathbb{C})$, which forces $D_k = D_{\sigma(k)}$ (by comparing dimensions as $\mathbb{C}$ -vector spaces). $\square$

Theorem 9.3 (Decomposition of automorphisms of $\prod M_{D_k}(\mathbb{C})$). *Every $\mathbb{C}$ -algebra automorphism of $\prod_{k=1}^r M_{D_k}(\mathbb{C})$ decomposes as a block permutation σ followed by per-block inner automorphisms: there exist $\sigma \in S_r$ with $D_{\sigma(k)} = D_k$ and invertible matrices $X_k \in \text{GL}_{D_k}(\mathbb{C})$ such that $T(M)_k = X_k M_{\sigma(k)} X_k^{-1}$.*

Proof. Theorem 9.1 gives the permutation σ . Theorem 9.2 gives $D_{\sigma(k)} = D_k$. The restriction of T to each block is a $\mathbb{C}$ -algebra automorphism of $M_{D_k}(\mathbb{C})$, which is inner by Skolem–Noether (Theorem 3.9).

This is a standard result in the theory of semisimple algebras; see [CPGSV21, §IV.A] for the MPS context. $\square$

9.2 Pi-algebra linear extension

Definition 9.4 (Per-block linear extension). Given injective tensors A_k, B_k of bond dimension D_k with the same MPV for each block k , the *per-block linear extension* T_k is the unique linear map $M_{D_k}(\mathbb{C}) \rightarrow M_{D_k}(\mathbb{C})$ satisfying $T_k(A_k^i) = B_k^i$ for all i . It exists (and is unique) because $\{A_k^i\}$ spans $M_{D_k}(\mathbb{C})$ by injectivity.

Definition 9.5 (Product algebra automorphism). The per-block linear extensions T_k are promoted to $\mathbb{C}$ -algebra homomorphisms (Lemma 3.10) and assembled into a single $\mathbb{C}$ -algebra automorphism $T : \prod_k M_{D_k}(\mathbb{C}) \xrightarrow{\sim} \prod_k M_{D_k}(\mathbb{C})$.

Theorem 9.6 (Per-block linear extensions assemble). *Let A_k and B_k be injective tensors of bond dimension D_k for $k = 1, \dots, r$, and suppose the per-block same-MPV condition holds. Then the per-block linear extensions $T_k : A_k^i \mapsto B_k^i$ assemble into an algebra automorphism of $\prod_k M_{D_k}(\mathbb{C})$, which decomposes as a permutation σ with $D_{\sigma(k)} = D_k$ and per-block conjugation matrices X_k .*

Proof. Each T_k exists uniquely and is multiplicative by Theorems 3.6 and 3.7. Promotion to algebra homomorphisms (Lemma 3.10) and assembly into the product gives an algebra automorphism. Theorem 9.3 decomposes it into permutation + per-block inner automorphisms. $\square$

Theorem 9.7 (Per-block SameMPV gives global gauge equivalence). *If all block tensors A_k are injective and per-block same-MPV holds, then each pair (A_k, B_k) is gauge equivalent, and the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ are globally gauge equivalent.*

Proof. Per-block gauge equivalence follows from the single-block FT (Theorem 3.11) applied to each block. Global gauge equivalence follows by assembling the per-block gauge matrices into a block-diagonal matrix (Theorem 8.4). $\square$

9.3 Block separation

Definition 9.8 (Canonical form predicate). A collection of scaling factors $(\mu_k)_{k=1}^r$ and block tensors $(A_k)_{k=1}^r$ satisfies the *canonical form predicate* if:

1. each A_k is injective,
2. each A_k satisfies the TP normalization $\sum_i (A_k^i)^\dagger A_k^i = \mathbb{1}$,
3. the moduli are strictly decreasing: $|\mu_1| > |\mu_2| > \dots > |\mu_r|$,
4. all $\mu_k \neq 0$,
5. the self-overlap converges: $O_{A_k A_k}(N) \rightarrow 1$ for each k .

Only this one-sided normalization is part of the predicate; no unital condition is assumed here. Condition (5) is a primitivity hypothesis: for a primitive channel ([Wol12, Thm. 6.7, item 3]), $T^n(\rho) \rightarrow \rho_\infty$ for every initial state, which forces the self-overlap to converge to 1. See also [Wol12, Thm. 6.8] for the completely-positive characterisation and Chapter 6 for the spectral-gap proof.

Remark 9.9. In the literature [CPGSV21], condition (5) is derived from primitivity of the transfer map (peripheral spectrum = $\{1\}$). Theorem 8.24 gives this implication: per-block primitivity implies overlap convergence, so condition (5) is redundant when primitivity holds. The canonical form predicate retains condition (5) explicitly because the overlap limit may also be assumed directly.

Theorem 9.10 (Vandermonde separation). *If $\mu_1, \dots, \mu_r$ are pairwise distinct complex numbers and $\sum_{k=1}^r c_k \mu_k^N = 0$ for $N = 0, 1, \dots, r-1$, then $c_k = 0$ for all k .*

Proof. The coefficient vector c satisfies $Vc = 0$ where V is the Vandermonde matrix of the μ_k . Injectivity of μ makes V invertible. $\square$

Lemma 9.11 (Block separation core). *Under the hypotheses of the canonical form predicate (strict ordering, TP normalization, self-overlap convergence), if $\sum_k \mu_k^N (V^{(N)}(A_k)_\sigma - V^{(N)}(B_k)_\sigma) = 0$ for all N, σ , then A_k and B_k generate the same MPV for each k .*

Proof. By induction on the number of blocks r . In the inductive step, take overlaps with the leading block A_0 ; cross-block contributions from $k \geq 1$ decay exponentially (by the spectral gap), leaving $O_{A_0 A_0}(N) - O_{A_0 B_0}(N) \rightarrow 0$. The proportional FT (Theorem 8.23) gives $A_0 \simeq B_0$. Peeling off the leading block and dividing by μ_0^N , the induction hypothesis applies. $\square$

Theorem 9.12 (Block separation from canonical form). *Let $((\mu_k), (A_k))$ satisfy the canonical form predicate, and let (B_k) be injective tensors satisfying the same TP normalization $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$. If the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family, then each pair (A_k, B_k) generates the same MPV family.*

Proof. By induction on the number of blocks r .

Base case ($r = 1$): the summed identity $\mu_1^N (V^{(N)}(A_1)_\sigma - V^{(N)}(B_1)_\sigma) = 0$ and $\mu_1 \neq 0$ give A_1 and B_1 same MPV.

Inductive step ($r \geq 2$): the leading block 0 has the largest $|\mu_0|$. Taking overlaps with a test tensor A_0 and using the spectral gap (Theorems 6.17 and 6.15), the cross-overlap contributions from non-leading blocks decay to zero. This shows $O_{A_0 A_0}(N) - O_{A_0 B_0}(N) \rightarrow 0$, and since $O_{A_0 A_0}(N) \rightarrow 1$ (by the primitivity hypothesis), the proportional FT (Theorem 8.23) gives A_0 and B_0 gauge-phase equivalent, hence same MPV.

Peeling off the leading block and dividing by μ_0^N , the induction hypothesis applies to the remaining $r - 1$ blocks. $\square$

Theorem 9.13 (Canonical form fundamental theorem). *Under the hypotheses of Theorem 9.12, there exist a permutation σ and per-block gauge matrices X_k such that $B_k^i = X_k A_{\sigma(k)}^i X_k^{-1}$.*

Proof. Theorem 9.12 gives per-block same-MPV. Theorem 9.7 converts this to per-block gauge equivalence with permutation. $\square$

Chapter 10

BNT and Full Assembly

This chapter introduces *Basis Normal Tensors* (BNT), establishes their permutation rigidity, and records the strongest end-stage Fundamental Theorem statements currently assembled in Lean. The proportional theorem is formalized with explicit coefficient-convergence data, and the equal theorem is the same-structure CF-BNT specialization rather than the full paper-level equal-MPV theorem. The results combine the canonical form reduction (Chapter 8), block separation (Chapter 9), spectral gap (Chapter 6), and BNT permutation arguments; the Newton–Girard identities are included as supporting infrastructure for future weight-matching extensions. The main references are [PGVWC07] and [CPGSV21].

10.1 Basis normal tensors

Definition 10.1 (Basis Normal Tensor (BNT)). A *basis normal tensor* (BNT) decomposition for a total tensor A_{tot} consists of g block tensors $(A_1, \dots, A_g)$ with bond dimensions $(D_1, \dots, D_g)$ such that:

1. each A_j is normal (Definition 2.19),
2. at each system size N , the MPV of A_{tot} is a linear combination of the block MPVs:
 $|V^{(N)}(A_{\text{tot}})\rangle = \sum_j c_j |V^{(N)}(A_j)\rangle,$
3. for sufficiently large N , the block MPV vectors $\{|V^{(N)}(A_j)\rangle\}_{j=1}^g$ are linearly independent.

This is [CPGSV21, Definition 4.2].

Definition 10.2 (Canonical form with BNT separation). A *canonical form with BNT separation* extends the canonical form (Definition 9.8) with the additional requirement that distinct blocks are *not* gauge-phase equivalent: for $j \neq k$ with $D_j = D_k$, A_j and A_k are not gauge-phase equivalent.

Theorem 10.3 (Cross-overlap decay for distinct CF-BNT blocks). *If $((\mu_k), (A_k))$ satisfies the CF-BNT predicate and $j \neq k$, then $O_{A_j A_k}(N) \rightarrow 0$.*

Proof. If $D_j \neq D_k$, the rectangular spectral gap theorem (Theorem 6.15) gives $O_{A_j A_k}(N) \rightarrow 0$. If $D_j = D_k$, the blocks are not gauge-phase equivalent (by the CF-BNT separation condition), so the spectral-gap theorem (Theorem 6.17) gives decay to 0. $\square$

Theorem 10.4 (CF-BNT yields BNT). *If $((\mu_k), (A_k))$ satisfies the CF-BNT predicate, then the block tensors form a BNT decomposition for the block-diagonal tensor $\bigoplus_k \mu_k A_k$.*

Proof. **Normality:** each block is injective (hence 1-block injective, hence normal). **Spanning:** the MPV decomposition (Theorem 2.24) gives $V^{(N)}(A_{\text{tot}})_\sigma = \sum_k \mu_k^N V^{(N)}(A_k)_\sigma$. **Linear independence:** the overlap matrix $G_{jk}(N) = O_{A_j A_k}(N)$ converges to the identity (diagonal entries $\rightarrow 1$ by the self-overlap hypothesis; off-diagonal entries $\rightarrow 0$ by Theorem 10.3). A Gram matrix converging to the identity is eventually invertible, giving linear independence. $\square$

10.2 BNT permutation rigidity

Theorem 10.5 (BNT permutation — proportional MPV case). *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be two BNT decompositions such that the block-diagonal tensors generate proportional MPV families with convergent nonzero coefficients. Then $g_A = g_B$, and there is a permutation $\sigma \in S_g$ with $D_{\sigma(k)} = D_k$ and $A_{\sigma(k)}$ gauge-phase equivalent to B_k for each k .*

Proof. From the proportional MPV hypothesis and convergent coefficients, one extracts (for each pair (j, k)) the limit $\lim_{N \rightarrow \infty} O_{A_j B_k}(N)$. By the spectral gap (Chapter 6), cross-block overlaps between non-equivalent blocks decay to 0. The resulting overlap matrix has at most one nonzero entry per row and per column, which defines a permutation. Nonzero overlap forces gauge-phase equivalence.

This is part of [CPGSV21, Theorem 4.1] (the Fundamental Theorem for proportional MPVs), which establishes the permutation and gauge-phase equivalence from the uniqueness of BNT decompositions (characterised in [CPGSV21, Proposition 4.1]). $\square$

Theorem 10.6 (BNT permutation — overlap-orthonormal case). *Let $(A_j)_{j=1}^{g_A}$ and $(B_k)_{k=1}^{g_B}$ be two families of injective tensors satisfying the TP normalization $\sum_i (A_j^i)^\dagger A_j^i = \mathbb{1}$ and $\sum_i (B_k^i)^\dagger B_k^i = \mathbb{1}$, whose self-overlaps converge to 1 and whose cross-overlaps (within each family) converge to 0. If they span the same MPV subspace at every system size, then $g_A = g_B$ and the blocks are related by a permutation and gauge-phase equivalence.*

Proof. The overlap matrix $O_{A_j B_k}(N)$ converges to a matrix with at most one nonzero entry per row/column (by the spectral gap). Thus the overlap orthonormality hypotheses are assumed directly, rather than derived from proportional MPVs together with coefficient convergence. $\square$

10.3 Coefficient convergence

Theorem 10.7 (Coefficient ratio decay). *Under the canonical form predicate, for $k \geq 1$, $(\mu_k/\mu_0)^N \rightarrow 0$ as $N \rightarrow \infty$.*

Proof. Since $|\mu_k| < |\mu_0|$ (strict ordering of moduli), $|\mu_k/\mu_0| < 1$, so the geometric sequence converges to 0. $\square$

Remark 10.8 (Strict-dominance regime). Theorem 10.7 applies when one modulus is strictly dominant: $|\mu_0| > |\mu_k|$ for $k \geq 1$. The oscillatory case $\sum_q \mu_{j,q}^N$ with $|\mu_{j,q}| = 1$ is treated separately in the full paper and is not formalized here.

10.4 Newton–Girard identities

Theorem 10.9 (Newton–Girard trace recursion). *If $A, B \in M_n(\mathbb{C})$ satisfy $\text{tr}(A^k) = \text{tr}(B^k)$ for all $k \geq 1$, then A and B have the same characteristic polynomial.*

Proof. The Newton–Girard identities express the coefficients of the characteristic polynomial recursively in terms of the power-sum traces $\text{tr}(A^k)$. Equal power sums give equal coefficients, hence equal characteristic polynomials. $\square$

Theorem 10.10 (Equal power sums imply equal multisets). *Let $\alpha, \beta : \{1, \dots, n\} \rightarrow \mathbb{C}$. If $\sum_i \alpha_i^k = \sum_i \beta_i^k$ for all $k \geq 1$, then the multisets $\{\alpha_i\}$ and $\{\beta_i\}$ are equal.*

Proof. Construct diagonal matrices $\text{diag}(\alpha)$ and $\text{diag}(\beta)$. The power-sum hypothesis gives $\text{tr}(\text{diag}(\alpha)^k) = \text{tr}(\text{diag}(\beta)^k)$. By Theorem 10.9, the characteristic polynomials agree: $\prod_i (X - \alpha_i) = \prod_i (X - \beta_i)$. Extracting roots gives multiset equality. $\square$

10.5 Primitivity bridge

Definition 10.11 (Primitive MPS tensor). A *primitive MPS tensor* is one equipped with a nonzero PSD fixed point ρ of the transfer map such that the complementary spectral radius $\rho_{\text{spec}}(\mathcal{E}_A - P) < 1$ (where P is the fixed-point projection).

Theorem 10.12 (Primitive overlap convergence). *If A is a primitive MPS tensor, then $O_{AA}(N) \rightarrow 1$.*

Proof. Immediate from the primitive overlap convergence (Theorem 6.21). $\square$

10.6 Full assembly

Theorem 10.13 (Fundamental Theorem — proportional MPV case). *Let A_{tot} and B_{tot} be MPS tensors equipped with CF-BNT families $(\mu_j^A, A_j)_{j=1}^{r_A}$ and $(\mu_k^B, B_k)_{k=1}^{r_B}$. Assume one is given coefficient arrays $a_{N,j}$, $b_{N,k}$ and nonzero limits a_j^∞ , b_k^∞ , c^∞ such that*

$$V^{(N)}(A_{\text{tot}})_\sigma = \sum_{j=1}^{r_A} a_{N,j} V^{(N)}(A_j)_\sigma, \quad V^{(N)}(B_{\text{tot}})_\sigma = \sum_{k=1}^{r_B} b_{N,k} V^{(N)}(B_k)_\sigma,$$

with $a_{N,j} \rightarrow a_j^\infty \neq 0$ and $b_{N,k} \rightarrow b_k^\infty \neq 0$, and suppose also that

$$V^{(N)}(A_{\text{tot}})_\sigma = c_N V^{(N)}(B_{\text{tot}})_\sigma$$

for a sequence $c_N \rightarrow c^\infty \neq 0$. Then $r_A = r_B$, and there is a permutation π such that $D_j^A = D_{\pi(j)}^B$ and A_j is gauge-phase equivalent to $B_{\pi(j)}$ for each j . The coefficient and convergence data are hypotheses of the Lean theorem; they are not yet derived automatically from arbitrary input tensors.

Proof. The CF-BNT hypotheses give BNT decompositions by Theorem 10.4. With the supplied decomposition identities and convergence hypotheses on the coefficients and proportionality constants, Theorem 10.5 yields the equality of block counts, the permutation, and the per-block gauge-phase equivalence. $\square$

Theorem 10.14 (Fundamental Theorem — equal MPV case). *Fix a common block structure: the same number of blocks r , the same bond dimensions $(D_k)_{k=1}^r$, and the same weights $(\mu_k)_{k=1}^r$. Let (μ_k, A_k) and (μ_k, B_k) be two CF-BNT families on this common data. If the block-diagonal tensors $\bigoplus_k \mu_k A_k$ and $\bigoplus_k \mu_k B_k$ generate the same MPV family, then each pair (A_k, B_k) is gauge equivalent, and hence the assembled block-diagonal tensors are gauge equivalent. This is the same-structure equal-MPV specialization formalized in Lean, not the full paper-level equal-MPV theorem with independently given block data.*

Proof. Forget the extra BNT-separation field and retain only the underlying canonical-form data. The canonical-form fundamental theorem (Theorem 9.13) then applies directly to this common (r, D_k, μ_k) structure, yielding per-block gauge equivalence and hence global gauge equivalence. $\square$

Bibliography

- [CPGSV17] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete. Matrix product density operators: Renormalization fixed points and boundary theories. *Ann. Phys.*, 378:100–149, March 2017.
- [CPGSV21] J. Ignacio Cirac, David Pérez-García, Norbert Schuch, and Frank Verstraete. Matrix product states and projected entangled pair states: Concepts, symmetries, theorems. *Reviews of Modern Physics*, 93(4):045003, 2021.
- [EHK78] David E. Evans and Raphael Høegh-Krohn. Spectral properties of positive maps on C^* -algebras. *Journal of the London Mathematical Society*, 17(2):345–355, 1978.
- [Jac09] Nathan Jacobson. *Basic Algebra II*. Dover Publications, 2009. Unaltered reprint of the 2nd edition (W. H. Freeman, 1989).
- [PGVWC07] D. Pérez-García, F. Verstraete, M. M. Wolf, and J. I. Cirac. Matrix product state representations. *Quantum Information & Computation*, 7(5):401–430, 2007.
- [SPGWC10] M. Sanz, D. Pérez-García, M. M. Wolf, and J. I. Cirac. A quantum version of Wielandt’s inequality. *IEEE Transactions on Information Theory*, 56(9):4668–4673, 2010.
- [Wol12] Michael M. Wolf. Quantum channels & operations: Guided tour. Lecture notes, 2012. Originally distributed via TU Munich foswiki; archival PDF at <https://mediatum.ub.tum.de/doc/1099654/1099654.pdf>.