


# Mathematical physics

**Mathematical physics** refers to the development of mathematical methods for application to problems in [physics](#). The *Journal of Mathematical Physics* defines the field as "the application of mathematics to problems in physics and the development of mathematical methods suitable for such applications and for the formulation of physical theories".<sup>[1]</sup>



An example of mathematical physics: solutions of Schrödinger's equation for [quantum harmonic oscillators](#) (left) with their [amplitudes](#) (right).

## Scope

There are several distinct branches of mathematical physics, and these roughly correspond to particular historical periods.

### Classical mechanics

The rigorous, abstract and advanced reformulation of Newtonian mechanics adopting the [Lagrangian mechanics](#) and the [Hamiltonian mechanics](#) even in the presence of constraints. Both formulations are embodied in [analytical mechanics](#) and lead to understanding the deep interplay of the notions of symmetry and conserved quantities during the dynamical evolution, as embodied within the most elementary formulation of [Noether's theorem](#). These approaches and ideas have been extended to other areas of physics as [statistical mechanics](#), [continuum mechanics](#), [classical field theory](#) and [quantum field theory](#). Moreover, they have provided several examples and ideas in [differential geometry](#) (e.g. several notions in [symplectic geometry](#) and [vector bundle](#)).

## Partial differential equations

Following mathematics: the theory of [partial differential equation](#), [variational calculus](#), [Fourier analysis](#), [potential theory](#), and [vector analysis](#) are perhaps most closely associated with mathematical physics. These were developed intensively from the second half of the 18th century (by, for example, [D'Alembert](#), [Euler](#), and [Lagrange](#)) until the 1930s. Physical applications of these developments include [hydrodynamics](#), [celestial mechanics](#), [continuum mechanics](#), [elasticity theory](#), [acoustics](#), [thermodynamics](#), [electricity](#), [magnetism](#), and [aerodynamics](#).

## Quantum theory

The theory of [atomic spectra](#) (and, later, [quantum mechanics](#)) developed almost concurrently with some parts of the mathematical fields of [linear algebra](#), the [spectral theory](#) of [operators](#), [operator algebras](#) and more broadly, [functional analysis](#). Nonrelativistic quantum mechanics includes [Schrödinger](#) operators, and it has connections to [atomic and molecular physics](#). [Quantum information](#) theory is another subspecialty.

## Relativity and quantum relativistic theories


The [special](#) and [general](#) theories of relativity require a rather different type of mathematics. This was [group theory](#), which played an important role in both [quantum field theory](#) and [differential geometry](#). This was, however, gradually supplemented by [topology](#) and [functional analysis](#) in the mathematical description of [cosmological](#) as well as [quantum field theory](#) phenomena. In the mathematical description of this physical areas, some concepts in [homological algebra](#) and [category theory](#) are also important nowadays.

## Statistical mechanics

[Statistical mechanics](#) forms a separate field, which includes the theory of [phase transitions](#). It relies upon the [Hamiltonian mechanics](#) (or its quantum version) and it is closely related with the more mathematical [ergodic theory](#) and some parts of [probability theory](#). There are increasing interactions between [combinatorics and physics](#), in particular statistical physics.

## Usage

---



Relationship between mathematics and physics

The usage of the term "mathematical physics" is sometimes [idiosyncratic](#). Certain parts of mathematics that initially arose from the development of [physics](#) are not, in fact, considered parts of mathematical physics, while other closely related fields are. For example, [ordinary differential equations](#) and [symplectic geometry](#) are generally viewed as purely mathematical disciplines, whereas [dynamical systems](#) and [Hamiltonian mechanics](#) belong to mathematical physics. [John Herapath](#) used the term for the title of his 1847 text on "mathematical principles of natural philosophy"; the scope at that time being

"the causes of heat, gaseous elasticity, gravitation, and other great phenomena of nature".<sup>[2]</sup>

## Mathematical vs. theoretical physics

The term "mathematical physics" is sometimes used to denote research aimed at studying and solving problems in physics or [thought experiments](#) within a mathematically [rigorous](#) framework. In this sense, mathematical physics covers a very broad academic realm distinguished only by the blending of some mathematical aspect and physics theoretical aspect. Although related to [theoretical physics](#),<sup>[3]</sup> mathematical physics in this sense emphasizes the mathematical rigour of the similar type as found in mathematics.

On the other hand, theoretical physics emphasizes the links to observations and [experimental physics](#), which often requires theoretical physicists (and mathematical physicists in the more general sense) to use [heuristic](#), [intuitive](#), and approximate arguments.<sup>[4]</sup> Such arguments are not considered rigorous by mathematicians, but that is changing over time .

Such mathematical physicists primarily expand and elucidate physical [theories](#). Because of the required level of mathematical rigour, these researchers often deal with questions that theoretical physicists have considered to be already solved. However, they can sometimes show that the previous solution was incomplete, incorrect, or simply too naïve. Issues about attempts to infer the second law of [thermodynamics](#) from [statistical mechanics](#) are examples. Other examples concern

the subtleties involved with synchronisation procedures in special and general relativity ([Sagnac effect](#) and [Einstein synchronisation](#)).

The effort to put physical theories on a mathematically rigorous footing not only developed physics but also has influenced developments of some mathematical areas. For example, the development of quantum mechanics and some aspects of [functional analysis](#) parallel each other in many ways. The mathematical study of [quantum mechanics](#), [quantum field theory](#), and [quantum statistical mechanics](#) has motivated results in [operator algebras](#). The attempt to construct a rigorous mathematical formulation of [quantum field theory](#) has also brought about some progress in fields such as [representation theory](#).

## Prominent mathematical physicists

---

### Before Newton

There is a tradition of mathematical analysis of nature that goes back to the ancient Greeks; examples include [Euclid](#) (*Optics*), [Archimedes](#) (*On the Equilibrium of Planes*, *On Floating Bodies*), and [Ptolemy](#) (*Optics*, *Harmonics*).<sup>[5][6]</sup> Later, [Islamic](#) and [Byzantine](#) scholars built on these works, and these ultimately were reintroduced or became available to the West in the [12th century](#) and during the [Renaissance](#).

In the first decade of the 16th century, amateur astronomer [Nicolaus Copernicus](#) proposed [heliocentrism](#), and published a treatise on it in 1543. He retained the [Ptolemaic](#) idea of [epicycles](#), and merely sought to simplify astronomy by constructing simpler sets of epicyclic orbits. Epicycles consist of circles upon circles. According to [Aristotelian physics](#), the circle was the perfect form of motion, and was the intrinsic motion of Aristotle's [fifth element](#)—the quintessence or universal essence known in Greek as [aether](#) for the English *pure air*—that was the pure substance beyond the [sublunary sphere](#), and thus was celestial entities' pure composition. The German [Johannes Kepler](#) [1571–1630], [Tycho Brahe](#)'s assistant, modified Copernican orbits to [ellipses](#), formalized in the equations of Kepler's [laws of planetary motion](#).

An enthusiastic atomist, [Galileo Galilei](#) in his 1623 book *The Assayer* asserted that the "book of nature is written in mathematics".<sup>[7]</sup> His 1632 book, about his telescopic observations, supported heliocentrism.<sup>[8]</sup> Having introduced experimentation, Galileo then refuted geocentric cosmology by refuting Aristotelian physics itself. Galileo's 1638 book *Discourse on Two New Sciences* established the law of equal free fall as well as the principles of inertial motion, founding the central concepts of what would become today's [classical mechanics](#).<sup>[8]</sup> By the Galilean [law of inertia](#) as well as the principle of [Galilean invariance](#), also called Galilean relativity, for any object experiencing inertia,

there is empirical justification for knowing only that it is at *relative* rest or *relative* motion—rest or motion with respect to another object.

[René Descartes](#) famously developed a complete system of heliocentric cosmology anchored on the principle of vortex motion, [Cartesian physics](#), whose widespread acceptance brought the demise of Aristotelian physics. Descartes sought to formalize mathematical reasoning in science, and developed [Cartesian coordinates](#) for geometrically plotting locations in 3D space and marking their progressions along the flow of time.<sup>[9]</sup>

An older contemporary of Newton, [Christiaan Huygens](#), was the first to idealize a physical problem by a set of parameters and the first to fully mathematize a mechanistic explanation of unobservable physical phenomena, and for these reasons Huygens is considered the first [theoretical physicist](#) and one of the founders of modern mathematical physics.<sup>[10][11]</sup>

## Newtonian and post Newtonian

In this era, important concepts in [calculus](#) such as the [fundamental theorem of calculus](#) (proved in 1668 by Scottish mathematician [James Gregory](#)<sup>[12]</sup>) and finding extrema and minima of functions via differentiation using Fermat's theorem (by French mathematician [Pierre de Fermat](#)) were already known before Leibniz and Newton. [Isaac Newton](#) (1642–1727) developed some concepts in [calculus](#) (although [Gottfried Wilhelm Leibniz](#) developed similar concepts outside the context of physics) and [Newton's method](#) to solve problems in physics. He was extremely successful in his application of [calculus](#) to the theory of motion. Newton's theory of motion, shown in his *Mathematical Principles of Natural Philosophy*, published in 1687,<sup>[13]</sup> modeled three Galilean laws of motion along with Newton's [law of universal gravitation](#) on a framework of [absolute space](#)—hypothesized by Newton as a physically real entity of Euclidean geometric structure extending infinitely in all directions—while presuming [absolute time](#), supposedly justifying knowledge of absolute motion, the object's motion with respect to absolute space. The principle of Galilean invariance/relativity was merely implicit in Newton's theory of motion. Having ostensibly reduced the Keplerian celestial laws of motion as well as Galilean terrestrial laws of motion to a unifying force, Newton achieved great mathematical rigor, but with theoretical laxity.<sup>[14]</sup>

In the 18th century, the Swiss [Daniel Bernoulli](#) (1700–1782) made contributions to [fluid dynamics](#), and [vibrating strings](#). The Swiss [Leonhard Euler](#) (1707–1783) did special work in [variational calculus](#), dynamics, fluid dynamics, and other areas. Also notable was the Italian-born Frenchman, [Joseph-Louis Lagrange](#) (1736–1813) for work in [analytical mechanics](#): he formulated [Lagrangian mechanics](#)) and variational methods. A major contribution to the formulation of Analytical Dynamics called [Hamiltonian dynamics](#) was also made by the Irish physicist, astronomer and

mathematician, [William Rowan Hamilton](#) (1805-1865). Hamiltonian dynamics had played an important role in the formulation of modern theories in physics, including field theory and quantum mechanics. The French mathematical physicist [Joseph Fourier](#) (1768 – 1830) introduced the notion of [Fourier series](#) to solve the [heat equation](#), giving rise to a new approach to solving partial differential equations by means of [integral transforms](#).

Into the early 19th century, following mathematicians in France, Germany and England had contributed to mathematical physics. The French [Pierre-Simon Laplace](#) (1749–1827) made paramount contributions to mathematical [astronomy](#), [potential theory](#). [Siméon Denis Poisson](#) (1781–1840) worked in [analytical mechanics](#) and [potential theory](#). In Germany, [Carl Friedrich Gauss](#) (1777–1855) made key contributions to the theoretical foundations of [electricity](#), [magnetism](#), [mechanics](#), and [fluid dynamics](#). In England, [George Green](#) (1793-1841) published *An Essay on the Application of Mathematical Analysis to the Theories of Electricity and Magnetism* in 1828, which in addition to its significant contributions to mathematics made early progress towards laying down the mathematical foundations of electricity and magnetism.

A couple of decades ahead of Newton's publication of a particle theory of light, the Dutch [Christiaan Huygens](#) (1629–1695) developed the wave theory of light, published in 1690. By 1804, [Thomas Young](#)'s double-slit experiment revealed an interference pattern, as though light were a wave, and thus Huygens's wave theory of light, as well as Huygens's inference that light waves were vibrations of the [luminiferous aether](#), was accepted. [Jean-Augustin Fresnel](#) modeled hypothetical behavior of the aether. The English physicist [Michael Faraday](#) introduced the theoretical concept of a field—not action at a distance. Mid-19th century, the Scottish [James Clerk Maxwell](#) (1831–1879) reduced electricity and magnetism to Maxwell's electromagnetic field theory, whittled down by others to the four [Maxwell's equations](#). Initially, optics was found consequent of Maxwell's field. Later, radiation and then today's known [electromagnetic spectrum](#) were found also consequent of this electromagnetic field.

The English physicist [Lord Rayleigh](#) [1842–1919] worked on [sound](#). The Irishmen [William Rowan Hamilton](#) (1805–1865), [George Gabriel Stokes](#) (1819–1903) and [Lord Kelvin](#) (1824–1907) produced several major works: Stokes was a leader in [optics](#) and fluid dynamics; Kelvin made substantial discoveries in [thermodynamics](#); Hamilton did notable work on [analytical mechanics](#), discovering a new and powerful approach nowadays known as [Hamiltonian mechanics](#). Very relevant contributions to this approach are due to his German colleague mathematician [Carl Gustav Jacobi](#) (1804–1851) in particular referring to [canonical transformations](#). The German [Hermann von Helmholtz](#) (1821–1894) made substantial contributions in the fields of [electromagnetism](#), waves, [fluids](#), and sound. In the United States, the pioneering work of [Josiah Willard Gibbs](#) (1839–1903) became the basis for [statistical mechanics](#). Fundamental theoretical results in this area were



achieved by the German [Ludwig Boltzmann](#) (1844-1906). Together, these individuals laid the foundations of electromagnetic theory, fluid dynamics, and statistical mechanics.

## Relativistic

By the 1880s, there was a prominent paradox that an observer within Maxwell's electromagnetic field measured it at approximately constant speed, regardless of the observer's speed relative to other objects within the electromagnetic field. Thus, although the observer's speed was continually lost relative to the electromagnetic field, it was preserved relative to other objects *in* the electromagnetic field. And yet no violation of [Galilean invariance](#) within physical interactions among objects was detected. As Maxwell's electromagnetic field was modeled as oscillations of the [aether](#), physicists inferred that motion within the aether resulted in [aether drift](#), shifting the electromagnetic field, explaining the observer's missing speed relative to it. The [Galilean transformation](#) had been the mathematical process used to translate the positions in one reference frame to predictions of positions in another reference frame, all plotted on [Cartesian coordinates](#), but this process was replaced by [Lorentz transformation](#), modeled by the Dutch [Hendrik Lorentz](#) [1853–1928].

In 1887, experimentalists Michelson and Morley failed to detect aether drift, however. It was hypothesized that motion *into* the aether prompted aether's shortening, too, as modeled in the [Lorentz contraction](#). It was hypothesized that the aether thus kept Maxwell's electromagnetic field aligned with the principle of Galilean invariance across all [inertial frames of reference](#), while Newton's theory of motion was spared.

Austrian theoretical physicist and philosopher [Ernst Mach](#) criticized Newton's postulated absolute space. Mathematician [Jules-Henri Poincaré](#) (1854–1912) questioned even absolute time. In 1905, [Pierre Duhem](#) published a devastating criticism of the foundation of Newton's theory of motion.<sup>[14]</sup> Also in 1905, [Albert Einstein](#) (1879–1955) published his [special theory of relativity](#), newly explaining both the electromagnetic field's invariance and Galilean invariance by discarding all hypotheses concerning aether, including the existence of aether itself. Refuting the framework of Newton's theory—[absolute space and absolute time](#)—special relativity refers to *relative space* and *relative time*, whereby *length* contracts and *time* dilates along the travel pathway of an object.

In 1908, Einstein's former mathematics professor [Hermann Minkowski](#) modeled 3D space together with the 1D axis of time by treating the temporal axis like a fourth spatial dimension—altogether 4D spacetime—and declared the imminent demise of the separation of space and time.<sup>[15]</sup> Einstein initially called this "superfluous learnedness", but later used [Minkowski spacetime](#) with great elegance in his [general theory of relativity](#),<sup>[16]</sup> extending invariance to all reference frames—whether

perceived as inertial or as accelerated—and credited this to Minkowski, by then deceased. General relativity replaces Cartesian coordinates with [Gaussian coordinates](#), and replaces Newton's claimed empty yet Euclidean space traversed instantly by Newton's [vector](#) of hypothetical gravitational force—an instant [action at a distance](#)—with a gravitational *field*. The gravitational field is [Minkowski spacetime](#) itself, the 4D [topology](#) of Einstein aether modeled on a [Lorentzian manifold](#) that "curves" geometrically, according to the [Riemann curvature tensor](#). The concept of Newton's gravity: "two masses attract each other" replaced by the geometrical argument: "mass transform curvatures of [spacetime](#) and free falling particles with mass move along a geodesic curve in the spacetime" ([Riemannian geometry](#) already existed before 1850s, by mathematicians [Carl Friedrich Gauss](#) and [Bernhard Riemann](#) in search for intrinsic geometry and non-Euclidean geometry.), in the vicinity of either mass or energy. (Under special relativity—a special case of general relativity—even massless energy exerts gravitational effect by its [mass equivalence](#) locally "curving" the geometry of the four, unified dimensions of space and time.)

## Quantum

Another revolutionary development of the 20th century was [quantum theory](#), which emerged from the seminal contributions of [Max Planck](#) (1856–1947) (on [black-body radiation](#)) and Einstein's work on the [photoelectric effect](#). In 1912, a mathematician [Henri Poincare](#) published *Sur la théorie des quanta*.<sup>[17][18]</sup> He introduced the first non-naïve definition of quantization in this paper. The development of early quantum physics followed by a heuristic framework devised by [Arnold Sommerfeld](#) (1868–1951) and [Niels Bohr](#) (1885–1962), but this was soon replaced by the [quantum mechanics](#) developed by [Max Born](#) (1882–1970), [Werner Heisenberg](#) (1901–1976), [Paul Dirac](#) (1902–1984), [Erwin Schrödinger](#) (1887–1961), [Satyendra Nath Bose](#) (1894–1974), and [Wolfgang Pauli](#) (1900–1958). This revolutionary theoretical framework is based on a probabilistic interpretation of states, and evolution and measurements in terms of [self-adjoint operators](#) on an infinite-dimensional vector space. That is called [Hilbert space](#) (introduced by mathematicians [David Hilbert](#) (1862–1943), [Erhard Schmidt](#) (1876–1959) and [Frigyes Riesz](#) (1880–1956) in search of generalization of Euclidean space and study of integral equations), and rigorously defined within the axiomatic modern version by [John von Neumann](#) in his celebrated book *Mathematical Foundations of Quantum Mechanics*, where he built up a relevant part of modern functional analysis on Hilbert spaces, the [spectral theory](#) (introduced by [David Hilbert](#) who investigated [quadratic forms](#) with infinitely many variables. Many years later, it had been revealed that his spectral theory is associated with the spectrum of the hydrogen atom. He was surprised by this application.) in particular. Paul Dirac used algebraic constructions to produce a relativistic model for the [electron](#), predicting its [magnetic moment](#) and the existence of its antiparticle, the [positron](#).



## List of prominent contributors to mathematical physics in the 20th century

Prominent contributors to the 20th century's mathematical physics include, (ordered by birth date) [William Thomson \(Lord Kelvin\)](#) [1824–1907], [Oliver Heaviside](#) [1850–1925], [Jules Henri Poincaré](#) [1854–1912], [David Hilbert](#) [1862–1943], [Arnold Sommerfeld](#) [1868–1951], [Constantin Carathéodory](#) [1873–1950], [Albert Einstein](#) [1879–1955], [Max Born](#) [1882–1970], [George David Birkhoff](#) [1884–1944], [Hermann Weyl](#) [1885–1955], [Satyendra Nath Bose](#) [1894–1974], [Norbert Wiener](#) [1894–1964], [John Lighton Synge](#) [1897–1995], [Wolfgang Pauli](#) [1900–1958], [Paul Dirac](#) [1902–1984], [Eugene Wigner](#) [1902–1995], [Andrey Kolmogorov](#) [1903–1987], [Lars Onsager](#) [1903–1976], [John von Neumann](#) [1903–1957], [Sin-Itiro Tomonaga](#) [1906–1979], [Hideki Yukawa](#) [1907–1981], [Nikolay Nikolayevich Bogolyubov](#) [1909–1992], [Subrahmanyan Chandrasekhar](#) [1910–1995], [Mark Kac](#) [1914–1984], [Julian Schwinger](#) [1918–1994], [Richard Phillips Feynman](#) [1918–1988], [Irving Ezra Segal](#) [1918–1998], [Ryogo Kubo](#) [1920–1995], [Arthur Strong Wightman](#) [1922–2013], [Chen-Ning Yang](#) [1922– ], [Rudolf Haag](#) [1922–2016], [Freeman John Dyson](#) [1923–2020], [Martin Gutzwiller](#) [1925–2014], [Abdus Salam](#) [1926–1996], [Jürgen Moser](#) [1928–1999], [Michael Francis Atiyah](#) [1929–2019], [Joel Louis Lebowitz](#) [1930– ], [Roger Penrose](#) [1931– ], [Elliott Hershel Lieb](#) [1932– ], [Sheldon Lee Glashow](#) [1932– ], [Steven Weinberg](#) [1933– ], [Ludvig Dmitrievich Faddeev](#) [1934–2017], [David Ruelle](#) [1935– ], [Yakov Grigorevich Sinai](#) [1935– ], [Vladimir Igorevich Arnold](#) [1937–2010], [Arthur Michael Jaffe](#) [1937– ], [Roman Wladimir Jackiw](#) [1939– ], [Leonard Susskind](#) [1940– ], [Rodney James Baxter](#) [1940– ], [Michael Victor Berry](#) [1941– ], [Giovanni Gallavotti](#) [1941– ], [Stephen William Hawking](#) [1942–2018], [Jerrold Eldon Marsden](#) [1942–2010], [Alexander Markovich Polyakov](#) [1945– ], [Herbert Spohn](#) [1946– ], [John Lawrence Cardy](#) [1947– ], [Giorgio Parisi](#) [1948– ], [Edward Witten](#) [1951– ], [Ashoke Sen](#) [1956–] and [Juan Martín Maldacena](#) [1968– ].

## See also

- [International Association of Mathematical Physics](#)
- [Notable publications in mathematical physics](#)
- [List of mathematical physics journals](#)
- [Gauge theory \(mathematics\)](#)
- [Relationship between mathematics and physics](#)
- [Theoretical, computational and philosophical physics](#)

## Notes

- Definition from the *Journal of Mathematical Physics*. "Archived copy" (<https://web.archive.org/web/200610>

03233339/<http://jmp.aip.org/jmp/staff.jsp>) . Archived from the original (<http://jmp.aip.org/jmp/staff.jsp>) on 2006-10-03. Retrieved 2006-10-03.

2. John Herapath (1847) *Mathematical Physics; or, the Mathematical Principles of Natural Philosophy, the causes of heat, gaseous elasticity, gravitation, and other great phenomena of nature* (<https://catalog.hathi-trust.org/Record/011557061?type%5B%5D=author&lookfor%5B%5D=John%20Herapath&ft=ft>) , Whittaker and company via [HathiTrust](#)
3. Quote: " ... a negative definition of the theorist refers to his inability to make physical experiments, while a positive one... implies his encyclopaedic knowledge of physics combined with possessing enough mathematical armament. Depending on the ratio of these two components, the theorist may be nearer either to the experimentalist or to the mathematician. In the latter case, he is usually considered as a specialist in mathematical physics.", Ya. Frenkel, as related in A.T. Filippov, *The Versatile Soliton*, pg 131. Birkhauser, 2000.
4. Quote: "Physical theory is something like a suit sewed for Nature. Good theory is like a good suit. ... Thus the theorist is like a tailor." Ya. Frenkel, as related in Filippov (2000), pg 131.
5. Pellegrin, P. (2000). Brunschwig, J.; Lloyd, G. E. R. (eds.). *Physics. Greek Thought: A Guide to Classical Knowledge*. pp. 433–451.
6. Berggren, J. L. (2008). "The Archimedes codex" (<https://www.ams.org/notices/200808/tx080800943p.pdf>) (PDF). *Notices of the AMS*. **55** (8): 943–947.
7. Peter Machamer "Galileo Galilei" (<http://plato.stanford.edu/archives/spr2010/entries/galileo>) —sec 1 "Brief biography", in Zalta EN, ed, *The Stanford Encyclopedia of Philosophy*, Spring 2010 edn
8. Antony G Flew, *Dictionary of Philosophy*, rev 2nd edn (New York: St Martin's Press, 1984), p 129 (<https://books.google.com/books?id=MmJHVU9Rv3YC&pg=PA129>)
9. Antony G Flew, *Dictionary of Philosophy*, rev 2nd edn (New York: St Martin's Press, 1984), p 89 (<https://books.google.com/books?id=MmJHVU9Rv3YC&pg=PA89&dq=mathematical+reasoning>)
10. Dijksterhuis, F. J. (2008). Stevin, Huygens and the Dutch republic. *Nieuw archief voor wiskunde*, 5, pp. 100-107. [https://research.utwente.nl/files/6673130/Dijksterhuis\\_naw5-2008-09-2-100.pdf](https://research.utwente.nl/files/6673130/Dijksterhuis_naw5-2008-09-2-100.pdf)
11. Andreessen, C.D. (2005) *Huygens: The Man Behind the Principle*. Cambridge University Press: 6
12. Gregory, James (1668). *Geometriae Pars Universalis* ([https://archive.org/details/gregory\\_universalis](https://archive.org/details/gregory_universalis)) . *Museo Galileo*: Patavii: typis heredum Pauli Frambotti.
13. "The Mathematical Principles of Natural Philosophy" (<http://www.britannica.com/EBchecked/topic/369153/The-Mathematical-Principles-of-Natural-Philosophy>) , *Encyclopædia Britannica*, London
14. Imre Lakatos, auth, Worrall J & Currie G, eds, *The Methodology of Scientific Research Programmes: Volume 1: Philosophical Papers* (Cambridge: Cambridge University Press, 1980), pp 213–214 (<https://books.google.com/books?id=RRniFBI8Gi4C&pg=PA213>) , 220 (<https://books.google.com/books?id=RRniFBI8Gi4C&pg=PA220>)

15. Minkowski, Hermann (1908–1909), "Raum und Zeit" [Space and Time], *Physikalische Zeitschrift*, 10: 75–88
16. Salmon WC & Wolters G, eds, *Logic, Language, and the Structure of Scientific Theories* (Pittsburgh: University of Pittsburgh Press, 1994), p 125 (<https://books.google.com/books?id=Z9K8lQufcMC&pg=PA125&dq=superfluous+learnedness+Einstein+Minkowski+general+relativity>)
17. McCormmach, Russell (Spring 1967). "Henri Poincaré and the Quantum Theory". *Isis*. **58** (1): 37–55. doi:10.1086/350182 (<https://doi.org/10.1086%2F350182>) .
18. Irons, F. E. (August 2001). "Poincaré's 1911–12 proof of quantum discontinuity interpreted as applying to atoms". *American Journal of Physics*. **69** (8): 879–84. Bibcode:2001AmJPh..69..879I (<https://ui.adsabs.harvard.edu/abs/2001AmJPh..69..879I>) . doi:10.1119/1.1356056 (<https://doi.org/10.1119%2F1.1356056>) .

## References

---

- [Zaslow, Eric](#) (2005), *Physmatics*, arXiv:physics/0506153 (<https://arxiv.org/abs/physics/0506153>) , Bibcode:2005physics...6153Z (<https://ui.adsabs.harvard.edu/abs/2005physics...6153Z>)

## Further reading

---

### Generic works

- [Abraham, Ralph](#); [Marsden, Jerrold E.](#) (2008), *Foundations of mechanics: a mathematical exposition of classical mechanics with an introduction to the qualitative theory of dynamical systems* (2nd ed.), Providence: AMS Chelsea Pub., ISBN 978-0-8218-4438-0
- [Courant, Richard](#); [Hilbert, David](#) (1989), *Methods of Mathematical Physics*, New York: Interscience Publishers
- [Kato, Tosio](#) (1995), *Perturbation theory for linear operators* (2nd repr. ed.), Berlin: Springer-Verlag, ISBN 3-540-58661-X (This is a reprint of the second (1980) edition of this title.)
- [Margenau, Henry](#); [Murphy, George Moseley](#) (1976), *The mathematics of physics and chemistry* (2nd repr. ed.), Huntington: R. E. Krieger Pub. Co., ISBN 0-88275-423-8 (This is a reprint of the 1956 second edition.)
- [Morse, Philip McCord](#); [Feshbach, Herman](#) (1999), *Methods of theoretical physics* (repr. ed.), Boston: McGraw Hill, ISBN 0-07-043316-X (This is a reprint of the original (1953) edition of this title.)
- [Reed, Michael C.](#); [Simon, Barry](#) (1972–1977), *Methods of modern mathematical physics*, **4**, New York City: Academic Press, ISBN 0-12-585001-8

- [Titchmarsh, Edward Charles](#) (1939), *The theory of functions* (2nd ed.), London: Oxford University Press (This tome was reprinted in 1985.)
- [Thirring, Walter E.](#); Harrell, Evans M. (tr.) (1978–1983), *A course in mathematical physics / [Lehrbuch der mathematischen Physik]* (4 vol.), New York: Springer-Verlag

## Textbooks for undergraduate studies

- Arfken, George B.; Weber, Hans J. (1995), *Mathematical methods for physicists* (4th ed.), San Diego: Academic Press, [ISBN 0-12-059816-7](#) (pbk.)
- [Boas, Mary L.](#) (2006), *Mathematical Methods in the Physical Sciences* (3rd ed.), Hoboken: John Wiley & Sons, [ISBN 978-0-471-19826-0](#)
- Butkov, Eugene (1968), *Mathematical physics*, Reading: Addison-Wesley
- [Jeffreys, Harold](#); [Swirles Jeffreys, Bertha](#) (1956), *Methods of mathematical physics* (3rd rev. ed.), Cambridge, [England]: Cambridge University Press
- Joos, Georg; Freeman, Ira M. (1987), *Theoretical Physics*, Dover Publications, [ISBN 0-486-65227-0](#)
- Mathews, Jon; Walker, Robert L. (1970), *Mathematical methods of physics* (2nd ed.), New York: W. A. Benjamin, [ISBN 0-8053-7002-1](#)
- Menzel, Donald Howard (1961), *Mathematical Physics*, Dover Publications, [ISBN 0-486-60056-4](#)
- Stakgold, Ivar (c. 2000), *Boundary value problems of mathematical physics* (2 vol.), Philadelphia: Society for Industrial and Applied Mathematics, [ISBN 0-89871-456-7](#) (set : pbk.)

## Textbooks for graduate studies

- Hassani, Sadri (1999), *Mathematical Physics: A Modern Introduction to Its Foundations*, Berlin, Germany: Springer-Verlag, [ISBN 0-387-98579-4](#)
- [Reed, M.](#); [Simon, B.](#) (1972–1977). *Methods of Mathematical Physics*. Vol 1-4. Academic Press.
- [Teschl, G.](#) (2009). *Mathematical Methods in Quantum Mechanics; With Applications to Schrödinger Operators* (<https://www.mat.univie.ac.at/~gerald/ftp/book-schroe/>) . Providence: American Mathematical Society. [ISBN 978-0-8218-4660-5](#).
- Moretti, V. (2018). *Spectral Theory and Quantum Mechanics; Mathematical Foundations of Quantum Theories, Symmetries and Introduction to the Algebraic Formulation 2nd Edition*. Berlin, Milan: Springer. [ISBN 978-3-319-70705-1](#).
- Landsman, K. (2017). *Foundations of Quantum Theory*. Berlin, Milan: Springer. [ISBN 978-3-319-51776-6](#).

- [Whittaker, Edmund Taylor](#); [Watson, George Neville](#) (13 September 1996), *A course of modern analysis: an introduction to the general theory of infinite processes and of analytic functions, with an account of the principal transcendental functions* (1st AMS ed.), Cambridge: Cambridge University Press (published 1927), [ISBN 978-0-521-58807-2](#)

## Specialized texts

- [Arnold, Vladimir I.](#); [Vogtmann, K.](#); [Weinstein, A.](#) (tr.) (1997), *Mathematical methods of classical mechanics / [Matematicheskie metody klassicheskoi mekhaniki]* (<https://archive.org/details/mathematicalmeth0000arno>) (2nd ed.), New York: Springer-Verlag, [ISBN 0-387-96890-3](#)
- [Baez, John C.](#); [Muniain, Javier P.](#) (1994), *Gauge fields, knots, and gravity*, Singapore ; River Edge: World Scientific, [ISBN 981-02-2034-0](#) (pbk.)
- [Hawking, Stephen W.](#); [Ellis, George F. R.](#) (1973), *The large scale structure of space-time*, Cambridge, England: Cambridge University Press, [ISBN 0-521-20016-4](#)
- [Geroch, Robert](#) (1985), *Mathematical physics*, Chicago: University of Chicago Press, [ISBN 0-226-28862-5](#) (pbk.)
- [Glimm, James](#); [Jaffe, Arthur](#) (1987), *Quantum physics: a functional integral point of view* (2nd ed.), New York: Springer-Verlag, [ISBN 0-387-96477-0](#) (pbk.)
- [Haag, Rudolf](#) (1996), *Local quantum physics: fields, particles, algebras* (2nd rev. & enl. ed.), Berlin; New York: Springer-Verlag, [ISBN 3-540-61049-9](#) (softcover)
- [von Neumann, John](#); [Beyer, Robert T.](#) (tr.) (1955), *Mathematical Foundations of Quantum Mechanics*, Princeton: Princeton University Press
- [Weyl, Hermann](#); [Robertson, H. P.](#) (tr.) (1931), *The theory of groups and quantum mechanics / [Gruppentheorie und Quantenmechanik]*, London: Methuen & Co.
- [Ynduráin, Francisco J](#) (2006), *Theoretical and Mathematical Physics. The Theory of Quark and Gluon Interactions*, Berlin: Springer, [ISBN 978-3642069741](#) (pbk.)

## External links

-  Media related to [Mathematical physics](#) at Wikimedia Commons