

1 Observations of Stratospheric Sudden Warmings in **2 Earth Rotation Variations**

Lisa Neef,¹ Sophia Walther,² Katja Matthes,¹ Kunihiro Koder⁴

Corresponding author: Lisa J. Neef, Düsternbrooker Weg 20, D-24105 Kiel, Germany
(lneef@geomar.de)

¹Ocean Circulation and Climate
Dynamics - Marine Meteorology, Helmholtz
Centre for Ocean Research Kiel
(GEOMAR), Kiel, Germany.

²Institute for Meteorology, Free
University of Berlin, Berlin, Germany.

³Solar-Terrestrial Environment
Laboratory, Nagoya University, Nagoya,
Japan

Abstract. Stratospheric sudden warmings (SSWs) are extreme events in the polar stratosphere, that are both caused by, and have effects on, the tropospheric flow. This means that SSWs are associated with changes in the angular momentum of the atmosphere, both before and after their onset. Because these angular momentum changes are transferred to the solid Earth, they can be observed in the rate of the Earth's rotation and the wobble of its rotational pole. By comparing observed Earth rotation variations to reanalysis data, we find that an anomaly in the orientation of the Earth's rotational pole, up to four times as large as the annual polar wobble, typically precedes SSWs by 20-40 days. The polar motion signal is due to pressure anomalies that are typically seen before SSW events, and represents a new type of observable that may aid in the prediction of SSWs. A decline in the length-of-day is also seen, on average, near the time of the SSW wind reversal, and is found to be due to anomalous easterly winds generated in the tropical troposphere around this time, though the structure and timing of this signal seems to vary widely from event to event.

1. Introduction

Stratospheric sudden warmings (SSWs) are extreme events that happen roughly every other year in the polar stratosphere; the usually-cold polar vortex warms up (usually 30°-50°C) over the course of a few days, and the vortex winds reverse from westerly to easterly. Figure 1 shows the (a) temperature and (b) zonal wind anomalies over the polar cap during the warming event of January 2009, which was exceptionally strong and unexpected [Harada *et al.*, 2009; Ayarzagüena *et al.*, 2011]. The reversal of zonal wind at 60°N propagated downward in time, crossing the 10hPa surface on 24 January 2009; this date is defined by Charlton and Polvani [2007] as the *central date* of the warming.

The 2009 SSW was the result of strong tropospheric forcing, in the form of a Rossby wave packet that was excited by a deep ridge over the eastern Pacific region, and a cyclonic anomaly in the North Atlantic region [Ayarzagüena *et al.*, 2011]. It not only affected tropospheric weather but also the rotation of the Earth. Fig. 1(c)-(d) shows observations of three parameters of Earth rotation over the course of the 2009 SSW. The first two parameters, χ_1 and χ_2 , are angles that define the motion of the Earth's rotational pole (after rotating to a terrestrial reference frame, see Section 2.1), and the third is the deviation in the length of a day from its 24-hour period. In all three parameters, we have removed the daily climatology (in order to remove the seasonal cycle), as well as the 151-day average around the central date, in order to remove the interannual variability (due to, e.g., the QBO or ENSO). This leaves the subseasonal fluctuations, which are typically on the order of tens of milliarcseconds for the polar motion angles, and microseconds for the length-of-day anomalies [Salstein and Rosen, 1989; Eubanks *et al.*, 1985; Rosen *et al.*,

1991]. Polar motion angle 2 in particular shows a negative anomaly of 30 mas about 3 weeks before the central date, while the length-of-day anomaly shows a steady decline as the central date is approached and passed. But are these features related to the SSW, and if so, why?

Earth rotation parameters (ERPs) may be an unusual observable for studying SSWs, but can actually serve as a global measure of atmospheric dynamics because they reflect the atmosphere’s angular momentum (AAM). Angular momentum within the Earth system is conserved in the absence of outside torques; therefore, changes in the axial AAM change the Earth’s rotational velocity, and changes in the two equatorial components of AAM the orientation of the Earth’s rotational pole. Of course there are also other sources of angular momentum in the Earth system (the ocean, continental hydrosphere, and solid Earth), but on subseasonal timescales the atmosphere is the dominant source of axial angular momentum [Rosen and Salstein, 1983; Eubanks et al., 1985] and a major source, along with the ocean, of equatorial angular momentum [Dobslaw et al., 2010].

Total AAM is the sum of relative angular momentum of the atmosphere (i.e., winds), and changes in the atmospheric moment of inertia (i.e., the atmospheric mass distribution). For example, the seasonal variation in the extratropical tropospheric jets causes a change in the axial relative AAM, which causes ΔLOD to fluctuate by about 1 ms every year [Hide et al., 1997]. Likewise, the annual appearance of the Siberian high pressure system causes a yearly fluctuation in the two equatorial components of AAM, which results in a polar wobble of several mas [Chao and Au, 1991; Nastula et al., 2009; Dobslaw et al., 2010].

63 In this paper we ask the question of whether SSWs affect AAM and, by extension, the
 64 rotation of the Earth. The effect of stratospheric phenomena on earth rotation variations
 65 has not been studied **much**, primarily because the low mass of the stratosphere typically
 66 makes its contribution to total AAM quite small [*Rosen and Salstein*, 1985; *Zhou et al.*,
 67 2008]. However, SSWs are a stratospheric phenomenon with strong links to the tropo-
 68 sphere; not only do they affect tropospheric weather for 1-2 months after the start of the
 69 warming [*Baldwin and Dunkerton*, 2001; *Thompson et al.*, 2002; *Woollings et al.*, 2010],
 70 they are also, typically, preceded by large-scale **mid- and high-latitude** pressure anomalies
 71 that trigger or enhance upward propagating planetary waves [*Quiroz*, 1986; *Martius et al.*,
 72 2009; *Woollings et al.*, 2010; *Garfinkel et al.*, 2010; *Ayarzagüena et al.*, 2011]. Moreover, it
 73 has been shown that SSWs induce an anomalous global meridional circulation that causes
 74 upwelling in the tropics, cooling of the tropical lower stratosphere, and consequently a
 75 westerly wind anomaly in the north-subtropical stratosphere [*Kodera*, 2006] and increased
 76 convection in the southern tropics [*Kodera et al.*, 2011].

77 Thus it is likely that SSWs might alter the global AAM. In order to identify the footprint
 78 of SSWs in the observed ERP records, we have composited observations of polar motion
 79 and length-of-day variations over the known SSW events over the 48 years since the
 80 beginning of the modern ERP record (1962-2010), and compared these composites to
 81 the corresponding atmospheric excitation of Earth rotation variations that is implied by
 82 reanalysis data.

83 The paper is organized as follows. Section 2 outlines the observational and reanalysis
 84 data used, and the connection between observed Earth rotation variations and geophys-
 85 ically modeled AAM excitation functions. Section 4 examines the impact of SSWs on

polar motion, while Section 5 examines the impact of major SSWS on the rate of Earth's rotation. A discussion and conclusions are given in Section 6.

2. Methods

2.1. Earth Rotation Observations

Earth rotation variations are described by anomalies in the length-of-day and the orientation of the Earth's figure axis. These so-called Earth rotation parameters (ERPs hereafter) are observed by a combination of optical astrometry, lunar and satellite laser ranging, Very Long Baseline Interferometry, and GPS, and are compiled regularly by the International Earth Rotation and Reference Systems Service (IERS). We have used the EOP-CO4 data series, which contains daily measurements over the period 1962 to the present day, and is available online at <http://hpiers.obspm.fr/eop-pc/>. In this data set, solid Earth tides (ranging in period from 5.64 days - 18.6 years) have been removed in postprocessing, while semidiurnal and diurnal ocean tide signals fall away due to the daily resolution of the data.

The angles of polar motion, p_1 and p_2 , represent the location of the Earth's rotational axis in an inertial, celestial reference frame that is fixed in space and defined relative to a group of stars (the so-called celestial ephemeris pole). *Barnes et al.* [1983] and later *Gross* [1992] showed that these vectors can be directly related to unit variations in the equatorial components of the Earth's angular momentum, χ_1 and χ_2 (defined along the Greenwich meridian and the 90°E, respectively) using:

$$p_1 + \frac{\dot{p}_2}{\sigma_0} = \chi_1^{\text{GEO}} \quad (1)$$

$$-p_2 + \frac{\dot{p}_1}{\sigma_0} = \chi_2^{\text{GEO}}, \quad (2)$$

where the overdots represent time derivatives, and 'GEO' denotes that the angular momentum components are observed geodetically, rather than derived from mechanical equations. Note that this equation involves a rotation into an inertial reference frame of the so-called Chandler wobble, a free nutation of the Earth of frequency $\sigma_0 = 2\pi/433\text{d}$, which results from the oblateness of the Earth's figure.

ΔLOD is the difference between the duration of the day that is determined astronomically and the solar day of 86400s. It is simply related to unit changes in the axial component of angular momentum, χ_3 :

$$\frac{\Delta\text{LOD}}{\text{LOD}_0} = \Delta\chi_3, \quad (3)$$

where LOD_0 represents the nominal length-of-day, 86400s.

2.2. Atmospheric Excitation Functions

The angular momentum excitation functions χ_i ($i = 1, 2, 3$) actually represent the net angular momentum of the entire Earth system, including the atmosphere, oceans, continental hydrosphere, and solid Earth. On timescales from a few days to months, fluctuations in the angular momentum of the atmosphere dominate changes in both LOD [Rosen and Salstein, 1983; Rosen *et al.*, 1990] and polar motion, modified by the response of the sea levels to pressure loading from the atmosphere [Eubanks *et al.*, 1988]. The rest of this manuscript will examine only the atmospheric angular momentum excitation functions (AEFs hereafter), with the exception of some oceanic effects covered in Section 4.1.

Each AEF can be separated into contributions from relative angular momentum (hereafter the wind term, χ_i^W), and changes in the atmospheric moment of inertia (hereafter

the mass term, χ_i^M). The wind and mass terms are as follows [Barnes et al., 1983]:

$$\chi_1^M = \frac{-1.10R^4}{(g(C-A))} \int \int p_s \sin \phi \cos^2 \phi \cos \lambda d\lambda d\phi \quad (4)$$

$$\chi_1^W = \frac{-1.61R^3}{\Omega(C-A)g} \int \int \int (u \sin \phi \cos \phi \cos \lambda - v \cos \phi \sin \lambda) d\lambda d\phi dp \quad (5)$$

$$\chi_2^M = \frac{-1.10R^4}{(g(C-A))} \int \int p_s \sin \phi \cos^2 \phi \sin \lambda d\lambda d\phi \quad (6)$$

$$\chi_2^W = \frac{-1.61R^3}{\Omega(C-A)g} \int \int \int (u \sin \phi \cos \phi \sin \lambda + v \cos \phi \cos \lambda) d\lambda d\phi dp \quad (7)$$

$$\chi_3^M = \frac{0.748R^4}{C_m g} \int \int p_s \cos^3 \phi d\lambda d\phi \quad (8)$$

$$\chi_3^W = \frac{0.997R^3}{C_m \Omega g} \int \int \int u \cos^2 \phi d\lambda d\phi dp, \quad (9)$$

where ϕ and λ represent latitude and longitude, respectively, p_s represents the surface pressure, and u and v are the zonal and meridional winds, respectively. $R = 6371.0$ km represents the radius of the Earth, $\Omega = 7.292115 \times 10^{-5}$ rad/s the average rotation rate, and $g = 9.81$ m/s² the acceleration due to gravity. $C = 8.0365 \times 10^{37}$ kgm² and $A = 8.0101 \times 10^{37}$ kgm² are the axial and next-largest principal moments of inertia of the solid Earth, and $C_m = 7.1236 \times 10^{37}$ kgm² is the principal inertia tensor component of the Earth's mantle [Gross, 2009].

Note that the equatorial excitation functions χ_1 and χ_2 are actually defined in radians, while the axial excitation function χ_3 is dimensionless. The trigonometric functions that weight wind and surface pressure in each integral come from the reference frame in which the ERPs are defined, and are illustrated graphically in the supplementary material.

It is also worth noting that χ_3 , which excites Δ LOD, depends only on zonal wind and surface pressure, and is weighted most strongly in the tropics, with uniform zonal weighting. In contrast, the equatorial excitation functions χ_1 and χ_2 also depend on the meridional wind and are weighted most strongly at midlatitudes, with a wave-1 zonal

weighting (see supplementary material). Note also that the wind excitation functions [(5), (7), and (9)] involve integrals over the mass of the atmosphere and are therefore weighted the most at the lowest levels, where the mass is highest.

2.3. ECMWF Reanalysis Data

SSWs are examined using the two major reanalyses of the European Centre for Medium-Range Weather Forecasts (ECMWF), ERA-40 [Uppala *et al.*, 2005] and ERA-Interim [Dee *et al.*, 2011], both at 2.5° horizontal resolution. These data are freely available online at <http://data-portal.ecmwf.int/>. Only ERA-Interim data (1979-2010) were used for the polar motion analysis in Section 4, because this analysis relies heavily on surface pressure data, whereas only sea-level pressure is publicly available in the ERA-40 reanalysis. For the analysis of length-of-day anomalies (Section 5), which focuses on wind excitation, the two datasets were selected for the vertical levels that they have in common, with the top at 1hPa, and joined together at 1.4.1979; this uses as many ERA-Interim data as possible, while keeping the junction away from the major warming event of February 1979.

2.4. Selection of Major Warming Events

SSWs are generally defined by rapidly increasing temperatures in the stratospheric polar vortex, along with an abrupt reversal of the vortex winds. Major midwinter warmings are defined by the WMO as events where the zonal mean zonal wind at 10hPa and 60°N becomes easterly during boreal winter (November-March), and simultaneously the meridional gradient in zonal-mean temperature at 10 hPa and 60-85°N is positive for more than 5 days [Labitzke and Naujokat, 2000].

In this study, major warming events are identified following the method of *Charlton and Polvani* [2007], which identifies SSWs by the wind criterion of the WMO definition. The first day where the wind at 10hPa and 60°N reverses to easterly is defined as the central date of the warming. In order to ensure that events with small westerly-wind fluctuations are not counted twice, no day within 20 days of this central date can also be defined as a central date. Final warmings, i.e. warmings where the vortex does not recover before the onset of the easterly summer circulation, are excluded from our analysis. This procedure is also done following *Charlton and Polvani* [2007], by requiring that winds must return to winter (westerly) wind conditions for at least 10 consecutive days before 30 April for an event to be considered non-final.

The above approach results in 14 major warmings identified in the ERA-40 period (1957-1979), and 22 events in the ERA-Interim period (1980-2010). These events are listed, in order of their central dates, in Table 1. Only the period of overlap between the reanalysis data and the ERP observations (1962 - present day) can be used; thus the SSWs of 1958 and 1960 are excluded. This leaves a total of 34 major SSWs on which to perform our analysis.

The events shown in Table 1 are in general agreement with the long-term meteorological observations performed at the Free University of Berlin (FUB) [*Labitzke and Naujokat*, 2000, and online at <http://www.geo.fu-berlin.de/met/ag/strat/produkte/northpole/index.html>], with the exception of 7 events identified as major warmings in this study but not by the FUB record (see Table caption). These 7 events also qualify as major warmings in the studies of *Charlton and Polvani* [2007] (which used the

NCEP/NCAR reanalysis set) and *Bancalá et al.* [2012] (which used ERA-40 data exclusively), but are generally weaker events without a strong tropospheric effect.

The events shown in Table 1 represent instances where the stratospheric and possibly tropospheric flow was significantly disturbed. Could these events also have influenced Earth rotation, as in the 2009 event (Fig. 1)? In order to answer this question, it is necessary to compute the AAM during these events; this will be discussed in the next section.

3. Observed Earth Rotation Anomalies during SSWS

Fig. 2 is similar to Fig. 1, but here the wind, temperature, and ERP anomalies have all been composited over the 34 major warming events identified in the combined ERA data set, from 1962 to 2010. The composites in each panel are centered on the central date of each event. For the three ERP observations [Fig. 2(c)-(e)], the 96% confidence interval has been estimated using a stationary bootstrap algorithm [Wilks, 1995], and is shown by shading.

Fig. 2 (a) - (b) illustrates the overall patterns common to major warmings, namely that the positive temperature anomalies start in the upper stratosphere several days before the central date, preceding the reversal in zonal wind, and that both the temperature and wind anomalies propagate downward into the lower stratosphere, lasting about 40-60 days after the central date.

The bottom three panels of Figure 2 show the observed ERPs, again rotating the polar motion angles to their respective angular momentum components, and now also compositing over the 34 SSW events. As in Fig. 1, we have removed the 151-day mean around the central date for each rotation parameter. A statistically significant signal can be seen in

χ_2 , and (for a few days around the central date) in ΔLOD , both parameters showing qualitatively the same behavior that was seen in the 2009 event (Fig. 1): χ_2 swings from positive to negative anomalies over the two months preceding the central date and then takes on weak positive anomalies after the central date, while ΔLOD declines rapidly in the two weeks before the central date and then recovers slowly towards zero anomalies over the 50 or so days after the central date.

4. Polar Motion Excitation by Mass Anomalies During SSWS

Figure 1(d) shows that the 2009 SSW was preceded by negative anomalies in χ_2^{GEO} , the atmospheric angular momentum component defined along the Greenwich meridian. This signal can also be seen in the composite over all 34 SSW events, while no clear signal was seen in the other component, χ_1 .

The AAM excitation functions for polar motion [(4)-(7)] are weighted zonally following sine and cosine waves, which means that only zonally-asymmetric wind and mass anomalies result in a net polar motion excitation. Consequently, subseasonal variations in polar motion are not generally excited by wind anomalies, which tend to cancel out in the zonal integral [Barnes *et al.*, 1983; Eubanks *et al.*, 1988], but rather by midlatitude anomalies in the atmospheric mass distribution. Mass anomalies in the mid troposphere are a common precursor of SSWS, because they excite upward-propagating planetary waves that break and thereby weaken the vortex, and SSWS are often preceded by persistent northern European blocking anticyclones [Quiroz, 1986; Martius *et al.*, 2009; Woollings *et al.*, 2010] and positively correlated to warm ENSO events [Garfinkel and Hartmann, 2008]. The impact of these mass variations on polar motion is investigated in the following two subsections.

4.1. Inverted Barometer Response of the Ocean

Figure 3 compares the observed equatorial AAM components, compared to their corresponding mass excitation functions χ_1^M and χ_2^M [(4) and (6)], over 75 days on either side of the central date. Because the excitation functions [(4) and (6)] are integrals of surface pressure, which is not publicly available in ERA-40, the curves in Figure 3 are composites over only the 22 SSWs in ERA-Interim. The blue lines show the pure mass excitation functions computed from (4) and (6). Both χ_1^M and χ_2^M show large fluctuations over the SSW life cycle, but for the observations, only χ_2 shows strong observed polar motion variations.

The difference between the large fluctuation seen in χ_2^{GEO} , and the weak fluctuation seen in χ_1^{GEO} , is explained when the AAM excitation functions are adjusted for the response of the oceans to atmospheric mass loading. This response can be simply modeled by averaging the surface pressure over the oceans globally, the so-called “inverted barometer” approximation [Wunsch and Stammer, 1997]. The adjusted excitation function is shown by the orange curves, which agree much more with the observed polar motion in both cases. The strong variations of χ_1^M over the SSW life cycle are clearly damped out by the response of the ocean, leading to a much weaker observed variation in χ_1^{GEO} . This makes sense, since the weighting function for χ_1^M is maximal at 0° and 180° , i.e. over the oceans. χ_2^M which happens to be weighted more strongly over the continents, clearly excites corresponding variations in χ_2^{GEO} . Therefore, the remainder of this paper will focus only on the angular momentum component χ_2 .

4.2. Polar motion anomalies preceding SSWs

Figure 4 examines the average surface pressure anomaly pattern associated with the SSWs at different points in time around the central date, along with the vertical profiles of geopotential height, averaged between 50N and 80N.

The first row of Figure 4 shows height-longitude slices of the geopotential height, averaged for each time block and over the 50N-80N latitudinal band. Geopotential height anomalies are computed with respect to the zonal mean, and then scaled by the relative mass of each vertical layer in order to emphasize the tropospheric anomalies. The composite geopotential height anomalies extend with a westward tilt into the stratosphere, indicating upward planetary wave propagation, which intensifies in the month before the warming onset [Fig. 4(a)-(b)].

At the surface [Fig. 4(e)], the upward wave propagation is related, on average, to high pressure anomalies over Eurasia and Northern Europe, and low anomalies over the northeastern Pacific. *Garfinkel et al.* [2010] showed that, while the individual pressure anomalies preceding SSWs can vary greatly, SSWs are most efficiently induced by anomalies that project onto the climatological planetary wave-1 that results naturally from orographic and thermal forcing in the Northern Hemisphere. This means that SSWs are often associated with negative tropospheric geopotential height anomalies over the North Pacific, and positive anomalies over Eastern Europe.

The meaning of this surface pressure pattern in terms of the AAM component χ_2 is examined in the bottom row of Figure 4[(g)-(i)], which shows the surface pressure anomalies weighted as in the integrand for the atmospheric moment-of-inertia (including the negative prefactor) in equation (6). The combined result of these two anomalies is that

the mass excitation function χ_2^M [Fig. 4(k)-(l)] becomes extremely negative in the month before the SSW onset.

Note that a recent study by *Mitchell et al.* [2012] pointed out that the surface pressure signals preceding SSWs differ between vortex-displacement and vortex-splitting events, with vortex displacements more strongly associated with a low pressure anomaly over North America, and vortex splits associated with a high pressure anomaly over Western Europe. However, we did not find a statistically significant difference in the polar motion signatures of vortex-splitting and -displacement events, presumably because both types of surface pressure anomalies contribute to negative χ_2 , and because of the relatively low sample size of each type of event.

5. LOD Excitation by Wind Anomalies During SSWs

Returning back to the composite of all three ERPs over the SSW events (Fig. 2), we see that SSWs on average don't just show a polar wobble but also a decline in ΔLOD [Fig. 2(e)] starting roughly a month before the central date. This implies that the atmospheric precursors that give rise to SSW events also change the axial AAM.

The date at which ΔLOD begins to decline varies widely from event to event; for example for the January 2009 event, the LOD decline begins about 50 days before the central date (Fig. 1), while for the February 1979 event, it begins about 25 days before the central date. For the January 1987 event, a noticeable decline in LOD doesn't happen at all (not shown).

The average wind AAM excitation function (χ_3^W) is examined in Figure 5(a), cast in terms of equivalent ΔLOD using (3) and compared to the observed ΔLOD . Variations of ΔLOD on this timescale are almost entirely explained by variations in the wind AAM,

which is why the mass term (8), which is about an order of magnitude smaller [Eubanks
et al., 1985], is omitted.

We can investigate the source of the axial AAM anomaly more closely by decomposing
the angular momentum into contributions from different latitude bands. In Figure 5(b),
the global axial angular momentum (gray) is compared to the angular momentum of the
following latitude bands: the South Polar cap (SP, 90°S-60°S), Southern Midlatitudes
(SH, 60°S-30°S), Tropics (T, 30°S-30°N), Northern Midlatitudes (NH, 30°N-60°N), and
the North Polar cap (NP, 60°N-90°N).

Here it can be seen that the wind reversal associated with the SSW causes a noticeable
decline in angular momentum from the North Polar band (dark blue), starting about 2
weeks before the central date. This angular momentum change contributes to the ob-
served ΔLOD decline but doesn't account for all of it. We also see an angular momentum
signal from the Northern Hemisphere extratropical band (green) that somewhat opposes
the angular momentum from the polar band. The strongest contribution to the observed
 ΔLOD decline actually comes from the tropical band, which shows sharply decreasing an-
gular momentum starting about two weeks before the central date, and a positive anomaly
after the central date. The prominence of the tropical band is not really surprising, since
the tropics are most strongly weighted in the integral (eq. 9), but it is surprising that the
tropical troposphere shows a such strong angular momentum changes during SSWs.

The zonal mean zonal winds behind these AAM changes are shown in the first column
of Figure 6, averaged over four blocks of time around the central date that characterize
the main ΔLOD changes: 60 to 20 days before the central date, when ΔLOD vacillates
around zero; 15 days before to 15 days after the central date, when it reaches its observed

minimum; 20 to 40 days after the central date, when it slowly recovers, and 40 to 60 days
 after the central date, when it has largely returned to zero anomalies. We see that on
 average, the SSWS are associated with tropospheric zonal wind anomalies on the order
 of 1 m/s, which, though weak, is comparable to the response of tropospheric wind to
 temperature anomalies in the tropical lower stratosphere *Haigh et al.* [2005]. Moreover,
 the contribution of these tropical wind anomalies to the axial angular momentum of the
 atmosphere is stronger since lower levels of the atmosphere have exponentially more mass.
 To illustrate this, the righthand column of Fig. 6 shows pressure-latitude slices of daily
 anomalies of $u \cos^2 \phi dp$, i.e. the fractional axial angular momentum at each level. Here
 we see anomalous westerlies forming in the troposphere near the equator during the ± 15
 days around the central date, which was also found by *Kodera* [2006] and attributed to
 the anomalous meridional circulation induced by the warming event at the poles.

The westerly anomalies would imply an increase in the ΔLOD , but are largely cancelled
 out by easterly anomalies at higher latitudes. The real cause of the tropical contribution
 to the declining ΔLOD is that the northern side of the tropical band shows an easterly
 wind anomaly in the SSW precursor period (top row), which is then weakened as the
 central date is approached (second row). We also see tropical easterly wind anomalies
 intensifying in the two months after the central date, though these are partially canceled
 out by positive wind anomalies at midlatitudes.

Thus it seems that SSWS are associated with tropical tropospheric wind anomalies
 throughout their life cycle, which are enough to cause a measurable decline in the observed
 length-of-day. However, since the statistical significance of our composite ΔLOD signal

is quite small, we defer a more thorough investigation of what causes these anomalies to future work.

6. Summary and Conclusions

This study showed that sudden stratospheric warmings are often preceded by strong anomalies in the angular momentum of the atmosphere, which is observable as polar motion, and anomalies in the length-of-day. SSWs are typically preceded by strong anomalies in χ_2 , one of the two equatorial components of the atmospheric angular momentum, which fluctuates by about 30 mas over the life cycle of an SSW, showing a positive anomaly about two months before the 10hPa wind reversal, and a negative anomaly about three weeks before the wind reversal, though only the latter is statistically significant. For individual events (see supplementary material) the total fluctuation of χ_2 can be as high as 60 mas. This is four times the observed annual polar wobble of about 15 mas (e.g. *Dobslaw et al.* [2010]).

The cause of the negative χ_2 anomaly is the surface pressure pattern that is on average associated with planetary waves that eventually induce SSWs [*Garfinkel et al.*, 2010; *Kodera et al.*, 2013]: a positive pressure anomaly over Eurasia and an enhanced Aleutian or Northeast Pacific low. As both of surface pressure patterns contribute negatively to χ_2 , many SSWs are preceded by a negative χ_2 anomaly, even though they may not exhibit the full surface pressure anomaly pattern identified in Fig. 4. A similar signal is not observable in χ_1 , the polar motion angle defined along the Greenwich Meridian, because the response of the oceans to atmospheric mass loading damps out the AAM anomalies in this direction.

Our work suggests that this polar wobble represents a new observable SSW precursor, which may aid in the prediction of SSWs, which is notoriously difficult. To investigate the efficacy of this signal as an observable precursor, Figures 7 and 8 show χ_2^{GEO} for all winters in the 1990s, which were relatively devoid of strong SSW events (Fig. 7), and the 2000s, which exhibited several strong events (Fig. 8). For each winter, the variation of χ_2^{GEO} is compared to the mean and standard deviation of χ_2^{GEO} over the entire epoch 1962-2010, and the central dates of SSW events are indicated by red dots.

It can be seen that strong negative anomalies in χ_2^{GEO} (i.e. anomalies outside of one standard deviation from the mean) are often harbingers of an SSW occurring 30-50 days later. All SSW events shown seem to be preceded by sharp negative values of χ_2^{GEO} in the month or two preceding the wind turnaround. On the other hand, extreme negative values of χ_2^{GEO} are also observed in winters without an SSW, including winters 1989/90, 1991/92, 1994/95, and 1995/96. A possible reason for this is that SSWs are not a simple response to tropospheric forcing, but also depend on the condition of the stratosphere, and whether planetary waves are able to propagate from the troposphere into the stratosphere.

SSWs may also be accompanied by a decline in the rate of the Earth's rotation by a tenth of a millisecond on average (Fig. 2e). For some warmings this effect is much stronger, for example, about a week before the central date the SSW event of February 2001 shows a ΔLOD of -0.6 ms, which is comparable to the 0.3-0.4 ms typically seen for subseasonal ΔLOD fluctuations [Eubanks *et al.*, 1985; Rosen *et al.*, 1991]. The decline in ΔLOD is the combined result of the stratospheric wind reversal from westerly to easterly, and westerly wind anomalies that might precede SSWs tropical troposphere, then decline at the onset of the SSWs, though it is difficult to say whether this is a statistically significant result.

A natural question to ask might be whether the observed atmospheric angular momentum anomalies associated with SSWs differ for warmings where the polar vortex splits, and warmings where the vortex is displaced off the pole, especially since previous studies have suggested that these types of events have different precursors [Martius *et al.*, 2009]). However, we did not find statistically significant differences between vortex splitting and displacement events for either the χ_2 or ΔLOD anomalies, and have omitted a discussion of vortex split and displacement events from this study.

Note also that this study has not discussed the transfer of AAM to the solid Earth, which typically happens by a combination of torques from surface friction and pressure systems around mountains [Egger *et al.*, 2007]. Since the estimated AAM explains most of the observed Earth rotation changes during SSWs, it is not necessary for the purpose of this study to estimate individual torques. However, a calculation of the relative magnitudes of the different torques would be an interesting point of future research.

Acknowledgments. This work has been performed within the Helmholtz-University Young Investigators Group NATHAN, funded by the Helmholtz-Association through the President’s Initiative and Networking Fund, the Helmholtz Centre for Ocean Sciences Kiel, the Helmholtz Centre for Geosciences Potsdam, and the Freie Universität Berlin. We are grateful to Nour-Eddine Omrani for helpful discussions, and to Christian Blume for help with the SSW selection algorithm. For information on how to obtain the data used to produce the results of this paper, please contact Dr. L. Neef.

References

- 397 Ayarzagüena, B., U. Langematz, and E. Serrano (2011), Tropospheric forcing of the strato-
 398 sphere: A comparative study of the two different major stratospheric warmings in 2009
 399 and 2010, *J. Geophys. Res.*, *116*, D18,114, doi:10.1029/2010JD015023.
- 400 Baldwin, M. P., and T. J. Dunkerton (2001), Stratospheric harbingers of anomalous
 401 weather regimes, *Science*, *294*, 581–584.
- 402 Bancalá, S., K. Krüger, and M. Giorgetta (2012), The preconditioning of major sudden
 403 stratospheric warmings, *J. Geophys. Res.*, *117*, D04,101, doi:doi:10.1029/2011JD016769.
- 404 Barnes, R., R. Hide, F.R.S., A. White, and C. Wilson (1983), Atmospheric angular mo-
 405 mentum fluctuations, length-of-day changes and polar motion, *Proc. Roy. Soc. London*,
 406 *387*, 31–73.
- 407 Chao, B., and A. Y. Au (1991), Atmospheric excitation of the Earth’s annual wobble:
 408 1980-1988, *J. Geophys. Res.*, *B4*, 6577–6582.
- 409 Charlton, A. J., and L. M. Polvani (2007), A new look at stratospheric sudden warmings.
 410 Part I: Climatology and modeling benchmarks, *J. Clim.*, *20*, 449–469.
- 411 Dee, D. P., et al. (2011), The ERA-Interim reanalysis: configuration and performance of
 412 the data assimilation system, *Quart. J. Roy. Meteorol. Soc.*, *137*(656), 553–597, doi:
 413 10.1002/qj.828.
- 414 Dobslaw, H., R. Dill, A. Grötsch, A. Brzezinski, and M. Thomas (2010), Seasonal po-
 415 lar motion excitation from numerical models of atmosphere, ocean, and continental
 416 hydrosphere, *J. Geophys. Res.*, *115*, 10,406, doi:10.1029/2009JB007127,2010.
- 417 Egger, J., K. Weickmann, and K.-P. Hoinka (2007), Angular momentum in the global
 418 atmospheric circulation, *Rev. Geophys.*, *45*, RG4007.

- 419 Eubanks, T., J. Steppe, J. Dickey, and P. Callahan (1985), A spectral analysis of the
420 Earth's angular momentum budget, *J. Geophys. Res.*, *90*, 5385–5404.
- 421 Eubanks, T., J. Steppe, J. Dickey, R. Rosen, and D. Salstein (1988), Causes of rapid
422 motions of the Earth's pole, *Nature*, *334*(14 July 1988), 115–119.
- 423 Garfinkel, C. I., and D. L. Hartmann (2008), Different ENSO teleconnections and their
424 effects on the stratospheric polar vortex, *Journal of Geophysical Research*, *113*(D18),
425 D18,114, doi:10.1029/2008JD009920.
- 426 Garfinkel, C. I., D. L. Hartmann, and F. Sassi (2010), Tropospheric Precursors of Anoma-
427 lous Northern Hemisphere Stratospheric Polar Vortices, *Journal of Climate*, *23*(12),
428 3282–3299, doi:10.1175/2010JCLI3010.1.
- 429 Gross, R. S. (1992), Correspondence between theory and observations of polar motion,
430 *Geophys. J. Int.*, *109*, 162–170.
- 431 Gross, R. S. (2009), Earth rotation variations - long period, in *Geodesy, Treatise on*
432 *Geophysics*, edited by T. Herring, pp. 239–294, Elsevier.
- 433 Haigh, J. D., M. Blackburn, and R. Day (2005), The Response of Tropospheric Circulation
434 to Perturbations in Lower-Stratospheric Temperature, *18*, 3672–3685.
- 435 Harada, Y., A. Goto, H. Hasegawa, and N. Fujikawa (2009), A major stratospheric sudden
436 warming event in january 2009, *J. Atmos. Sci.*, *67*, 2052–2069.
- 437 Hide, R., J. Dickey, S. Marcus, R. Rosen, and D. Salstein (1997), Atmospheric angular
438 momentum fluctuations during 1979-1988 simulated by global circulation models, *J.*
439 *Geophys. Res.*, *102*, 16,423–16,438.
- 440 K.Labitzke, and B.Naujokat (2009), On the remarkable arctic winter in 2008/2009, *J.*
441 *Geophys. Res.*, *114*, D00I02.

- 442 Kodera, K. (2006), Influence of stratospheric sudden warming on the equatorial tropo-
443 sphere, *Geophys. Res. Lett.*, *33*, L06,804.
- 444 Kodera, K., N. Eguchi, J. N. Lee, Y. Kuroda, and S. Yukimoto (2011), Sudden change in
445 the tropical stratosphere and troposphere during January 2009, *J. Met. Soc. Japan*, *89*,
446 283–290, doi:10.2151/jmsj.2011-208.
- 447 Kodera, K., H. Mukougawa, and A. Fujii (2013), Influence of the vertical and zonal propa-
448 gation of stratospheric planetary waves on tropospheric blockings, *Journal of Geophys-*
449 *ical Research: Atmospheres*, *118*(15), 8333–8345, doi:10.1002/jgrd.50650.
- 450 Labitzke, K., and B. Naujokat (2000), On the remarkable arctic winter in 2008/2009,
451 *SPARC Newsletter*, *15*, 11–14.
- 452 Martius, O., L. Polvani, and H. Davies (2009), Blocking precursors to stratospheric sudden
453 warming events, *Geophys. Res. Lett.*, *36*, L14,806, doi:10.1029/2009GL038776.
- 454 Mitchell, D. M., L. J. Gray, J. Anstey, M. P. Baldwin, and A. J. Charlton-Perez (2012),
455 The influence of stratospheric vortex displacements and splits on surface climate, *Jour-*
456 *nal of Climate*, *26*, 2668–2682, doi:10.1175/JCLI-D-12-00030.1.
- 457 Nastula, J., D. Salstein, and B. Kolaczek (2009), Patterns of atmospheric excitation
458 functions of polar motion from high-resolution regional sectors, *J. Geophys. Res.*, *114*,
459 B04,407.
- 460 Quiroz, R. S. (1986), The Association of Stratospheric Warmings With Tropospheric
461 Blocking, *J. Geophys. Res.*, *91*(D4), 5277–5285.
- 462 Rosen, R. D., and D. A. Salstein (1983), Variations in atmospheric angular momentum
463 on global and regional scales and the length of day, *J. Geophys. Res.*, *88*, 5451–5470.

- 464 Rosen, R. D., and D. A. Salstein (1985), Contribution of stratospheric winds to annual
465 and semiannual fluctuations in atmospheric angular momentum and the length of day,
466 *J. Geophys. Res.*, *90*, 8033–8041.
- 467 Rosen, R. D., D. A. Salstein, and T. M. Wood (1990), Discrepancies in the Earth-
468 Atmosphere Angular Momentum Budget, *J. Geophys. Res.*, *95*(89), 265–279.
- 469 Rosen, R. D., D. A. Salstein, and T. M. Wood (1991), Zonal contributions to global mo-
470 mentum variations on intraseasonal through interannual timescales, *J. Geophys. Res.*,
471 *96*, 5145–5151.
- 472 Salstein, D. A., and R. D. Rosen (1989), Regional contributions to the atmospheric exci-
473 tations of rapid polar motions, *J. Geophys. Res.*, *94*, 9971–9978.
- 474 Thompson, D. W., M. P. Baldwin, and J. M. Wallace (2002), Stratospheric connection
475 to Northern Hemisphere wintertime weather: Implications for prediction, *J. Clim.*, *15*,
476 14211,428.
- 477 Uppala, S. M., et al. (2005), The ERA-40 re-analysis, *Quart. J. Roy. Meteorol. Soc.*,
478 *131*(612), 2961–3012, doi:10.1256/qj.04.176.
- 479 Wilks, D. S. (1995), *Statistical methods in the atmospheric sciences: An Introduction*, 467
480 pp., Academic Press, Inc., San Diego, California.
- 481 Woollings, T., A. Charlton-Perez, S. Ineson, A. Marshall, and G. Masato (2010), Asso-
482 ciations between stratospheric variability and tropospheric blocking, *J. Geophys. Res.*,
483 *115*, D06,108.
- 484 Wunsch, C., and D. Stammer (1997), Atmospheric loading and the oceanic “inverted
485 barometer” effect, *Rev. Geophys.*, *35*(1), 79–107.

486 Zhou, Y., J. Chen, and D. Salstein (2008), Tropospheric and stratospheric wind contribu-
487 tions to earth's variable rotation from NCEP/NCAR reanalyses (2000-2005), *Geophys.*
488 *J. Int.*, *174*, 453–463, doi:10.1111/j.1365-246X.2008.03843.

Table 1. Major stratospheric sudden warming events identified in ERA-40 (1958-1978) and ERA-Interim (1979-2010).

<i>ERA-40</i>	<i>ERA-Interim</i>
31 Jan 1958 ^b	29 Feb 1980 ^c
15 Jan 1960 ^{b,c}	4 Mar 1981
28 Jan 1963	4 Dec 1981 ^c
16 Dec 1965 ^e	24 Feb 1984 ^c
23 Feb 1966	1 Jan 1985
7 Jan 1968	23 Jan 1987
28 Nov 1968	8 Dec 1987
13 Mar 1969	14 Mar 1988 ^d
2 Jan 1970	21 Feb 1989
18 Jan 1971	15 Dec 1998
20 Mar 1971	26 Feb 1999
31 Jan 1973	20 Mar 2000 ^c
9 Jan 1977	11 Feb 2001
22 Feb 1979 ^d	30 Dec 2001
	18 Jan 2003
	5 Jan 2004
	21 Jan 2006
	24 Feb 2007
	22 Feb 2008
	24 Jan 2009
	9 Feb 2010
	24 Mar 2010

^b Warmings that are excluded because they fall outside of the ERP observation record.

^c Warmings that are not found in the observational record of *K.Labitzke and B.Naujokat* [2009].

^d Warmings that are identified as final in the observational record of *K.Labitzke and B.Naujokat* [2009].

^e Warmings that are identified as Canadian Warmings (i.e., warmings where the anomalous warm temperatures are observed mainly in the lower stratosphere) in the observational record of *K.Labitzke and B.Naujokat* [2009].




Figure 1. First and second panels: altitude-time composites of the polar cap (60°-90°N) temperature and 60 °N zonal wind anomalies, for the January 2009 SSW event. Bottom three panels: the observed anomalies in the three Earth rotation parameters over the same time.




Figure 2. First and second panels: altitude-time composites of the polar cap (60°-90°N) temperature and 60 °N zonal wind anomalies, composited over the SSW events given in Table 1 and centered on the central date. Bottom three panels: the observed anomalies in the three Earth rotation parameters composited over the same events. The shading in ERP composites indicates the 96% bootstrap confidence interval.




Figure 3. Top: Composites of the p_1 anomaly (black) and the corresponding AAM mass excitation function (χ_1^M), with and without the IB approximation (see text), composited over the 22 major warming events in ERA-Interim. Bottom: As for top row but for observed p_2 anomaly and corresponding components of the mass excitation function χ_2^M .




Figure 4. Top row: Height-longitude cross sections of the geopotential height, composited over the 22 major warming events in ERA-Interim, averaged over 3 periods before and after the central date. Second row: the corresponding composite surface pressure anomaly. Third row: the surface pressure anomalies weighted by $\sin \phi \cos^2 \phi \cos \lambda$, as in the χ_1^M integral (eq. 4). Bottom row: the surface pressure anomalies weighted by $\sin \phi \cos^2 \phi \sin \lambda$, as in the χ_2^M integral (eq. 6).




Figure 5. (a) the observed ΔLOD (black) and the corresponding axial wind excitation (χ_3 , gray), composited over the all major warming events in the joint dataset. (b): The composite wind excitation functions integrated over different latitude bands, along with the global value (gray).




Figure 6. Left: Pressure-latitude slices of zonal-mean zonal wind, averaged over four blocks of time around the central date (see text), and composited over all SSW events in the joint dataset. Right: multiplying the wind anomalies by $\cos^2 \phi$ and by the relative mass of each pressure level, such that each gridbox gives the local contribution to the global χ_3 integral.




Figure 7. χ_2^{GEO} for all 10 winters in the 1990s. The mean and standard deviation of χ_2^{GEO} over the period 1962-2010 are shown by the shaded band in each plot, and the central dates of the SSW events during this decade are shown by red dots.




Figure 8. As in Fig. 7, but for the first decade of the 2000s.