

Current perspectives

The role of eosinophils in host defense against helminth parasites

Amy D. Klion, MD, and Thomas B. Nutman, MD Bethesda, Md

The precise function of eosinophils in parasitic infection *in vivo* remains poorly understood despite eosinophils having been shown to be potent effectors in killing parasites *in vitro*. Although it has long been held that the primary function of the eosinophil is protection against helminth parasites, there are little data to prove this unequivocally. Moreover, eosinophils are responsible for a considerable amount of inflammatory pathology accompanying helminth infections. This article will provide an overview of our current knowledge about eosinophils and their role, both protective and pathogenic, in parasitic helminth infections. (J Allergy Clin Immunol 2004;113:30-7)

Key words: Immunity, resistance, cytotoxicity, nematodes

Parasitic infections are caused by unicellular protozoa or multicellular helminths (worms), the extraordinary worldwide prevalence of which exacts a major medical and economic burden. Each parasite has an exceedingly diverse and unique biology. Protozoa are usually a few micrometers in size, whereas helminths are typically centimeters to meters in length. Tissue-dwelling protozoa are often intracellular parasites at some stage of infection, whereas helminths, being larger than most tissue cells, are almost always extracellular pathogens, with the significant exception being *Trichinella spiralis*. Although protozoa usually replicate during infection of a single host, helminths do not reproduce without the passage through either intermediate hosts or through soil or water.

Helminths have many developmental stages present during infection.¹ In the course of a single infection, the host might be repeatedly exposed to larval-, adult-, or egg-stage antigens. For example, free-swimming cercariae of the trematode *Schistosoma mansoni* penetrate the skin of human subjects immersed in infested water and evolve into tissue-stage schistosomula that migrate to the liver and mesenteric veins for further differentiation into sexually dimorphic adult worms. Eggs are laid that migrate through tissues into the lumen of the bowel or

bladder for environmental release. Similarly, filarial infection involves repeated exposure to arthropod-borne infective larvae and parasitization by long-lived adult worms capable of continuously releasing microfilariae that circulate in the bloodstream or migrate through subcutaneous tissues.

Because each stage of parasite development can be antigenically distinct, the host response to helminth infection is often characterized by a series of discrete immune responses that evolve at different times during the course of infection. Protective immunity directed against a single stage might be circumvented by parasite differentiation, assisting in the survival of the pathogen and posing a significant challenge to immune-mediated resistance. Each stage of parasite development might also entail a change in tissue tropism, introducing a compartmental feature to immune or inflammatory responses. For example, a variety of distinct cutaneous, pulmonary, and intestinal inflammatory or hyper-eosinophilic syndromes are associated with different stages of *Ascaris* and *Strongyloides* species as they migrate through the skin and lung before reaching adulthood in the gastrointestinal tract.²

Because complex parasitic life cycles can only be maintained by sequential passage through intermediate and definitive hosts, parasites have adapted methods of optimizing transmission by prolonging infection. Perhaps because of this evolutionary pressure, chronicity, latency, or both are hallmarks of helminth infections. For example, adult schistosomes and filariae can survive in host tissues for as long as 20 to 30 years, continuously producing eggs and larvae. *Strongyloides* species, because of their ability to "autoinfect," maintain their life cycle for decades.³ Chronicity might also reflect pressure toward "true parasitism" in that inducing mortality in the host would disrupt parasite transmission if the host were to die before egg laying or larval release could occur. Adaptations for chronicity can be so successful among parasites that naturally acquired protective immunity is only observed rarely in areas endemic for a given disease. Specifically, schistosomiasis results in an incomplete form of protection in which reinfection is limited but adult worms are tolerated for years (concomitant immunity).⁴

Whereas pathology and protective immunity to most protozoa are believed to reflect T cell-, B cell-, and macrophage-dependent (or perhaps cytokine-mediated) mechanisms, infection with helminth parasites induces immune effector responses that are characterized by IgE

From the Helminth Immunology Section, Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health.

Received for publication October 20, 2003; revised October 20, 2003; accepted for publication October 21, 2003.

Reprint requests: Thomas B. Nutman, MD, Laboratory of Parasitic Diseases, Building 4, Room B1-03, 4 Center Dr, National Institutes of Health, Bethesda, MD 20892-0425.

doi:10.1016/j.jaci.2003.10.050

antibody production, tissue and peripheral blood eosinophilia, and the participation of inflammatory mediator-rich tissue mast cells, qualitatively not unlike that seen in atopic diseases. For the atopic state, these responses have clearly been implicated in the pathogenesis of allergic diseases. In parasitic infection, although these types of responses can certainly induce pathologic reactions, they have also been implicated in mediating protective immunity to the helminth parasites.⁵⁻⁷

GENERAL FEATURES OF THE IMMUNE RESPONSE TO HELMINTHS

Whereas the cytokine response to helminth infection is clearly more complex than initially appreciated,⁸ the eosinophilia and increased serum IgE levels characteristic of helminth infection are associated with the production of IL-4 and IL-5. The critical role of IL-5 in this eosinophilic response has been clearly demonstrated in animal models, including transgenic and knockout mice infected with a wide variety of intestinal and tissue helminths.^{9,10} Whereas IL-5 alone is sufficient to cause helminth infection-mediated eosinophilia and eosinophil activation in most instances, a number of other cytokines (eg, IL-4 and IL-13)^{11,12} and chemokines (eg, RANTES and eotaxin)^{13,14} appear to be necessary for the recruitment of eosinophils to the tissues.

A number of transcription factors involved in the regulation of *IL5* gene expression, including nuclear factor (NF) IL-6 (C/EBP β), NF-AT, and GATA-3, have been recently identified.¹⁵ At least one of these, GATA-3, appears to play a direct role in helminth-induced eosinophilia, as demonstrated by enhanced IL-5 production and eosinophilia in GATA-3-transgenic mice infected with *Heligiosomoides polygyrus*, but not in uninfected GATA-3-transgenic control mice.¹⁶ The role of other transcription factors in the eosinophilia of helminth infection remains to be explored.

Eosinophils in the blood and tissues of helminth-infected patients and experimental animals exhibit morphologic and functional changes associated with eosinophil activation in vitro.¹⁷ These include decreased density,^{18,19} upregulation of surface activation molecules (eg, CD69, CD25, CD44, and HLA-DR),²⁰ enhanced cellular cytotoxicity, and release of granule proteins, cytokines, leukotrienes, and other mediators of inflammation.

The kinetics of eosinophilia after infection with any parasitic infection have been difficult to define because the time of initial infection can be defined only rarely. However, information from both single-source outbreaks^{21,22} and experimental human infection has shed light on the initiation of eosinophilia and its natural regulation. In a study of 5 volunteers infected with the hookworm *Necatur americanus*,²³ it was shown that blood eosinophil counts increased from baseline between 2 and 3 weeks after infection and peaked between weeks 7 and 9 (Fig 1). This increase was accompanied by a parallel increase in total leukocyte counts. In those exposed to lymphatic filariae,²⁴ peak eosinophil levels occur 11 to




FIG 1. Blood eosinophil response in 5 normal volunteers after exposure to *Necatur americanus* third-stage larvae. Each line represents an individual patient's eosinophil levels after experimental infection with hookworm. The arrows indicate the time of anthelmintic treatment. Adapted from data of Maxwell et al.²³

30 weeks after infection, which is consistent with the differences in life cycles between these parasites (Table I). More interesting was a spontaneous decrease in eosinophil numbers in the absence of treatment, suggesting active downregulation of the eosinophilia, a process seen in other experimental volunteer infections.^{25,26} This active (and spontaneous) modulation of eosinophilia generally occurred when migration of the larval forms ceased and the infections became patent (sexually mature adults began egg laying [eg, hookworm²³ or schistosomes]) or larval release (eg, filariae²⁴).

EOSINOPHILS AND PROTECTIVE IMMUNITY

Epidemiologic evidence for the role of eosinophils in the immune response to helminth infections

As early as 1939, eosinophils were postulated to play a role in the immune response to helminth infection.²⁷ Such hypotheses were based primarily on histopathologic evidence of eosinophils surrounding dying parasites in tissue biopsy specimens. Later, in vitro killing of parasites by eosinophils (in the presence of antibodies, complement, or both)²⁸⁻³⁰ and eosinophil granule products^{31,32} was demonstrated. Despite these in vitro findings and epidemiologic evidence correlating high eosinophil counts with resistance to posttreatment reinfection with *Schistosoma haematobium* and *S. mansoni* in human subjects,^{33,34} the in vivo role of eosinophils in immunity to helminth infection has been much more difficult to define. There has been conflicting evidence^{35,36} on the protective efficacy of eosinophils in animals depleted of eosinophils and subsequently challenged with a variety of parasites

TABLE I. Blood eosinophil response in human volunteers infected with lymphatic filariae

Parasite	Peak WBC	Peak eosinophil count (/ μ L)	Time to peak eosinophil count (wk)	Reference
<i>Brugia malayi</i>	20,400	12,840	14	84
<i>B malayi</i>	8800	2640	13	85
<i>B malayi</i>	7600	1234	14	85
<i>Brugia pahangi</i>	8200	1330	12	85
<i>B pahangi</i>	9600	1692	12	85
<i>B malayi</i>	32,000	22,400	26	86
<i>B malayi</i>	10,000	<500	20	87
<i>B malayi</i>	11,880	3020	30	87
<i>B malayi</i>	10,800	1728	11	88

WBC, White blood cell count.

TABLE II. The role of the eosinophil in mediating protection in selected helminth infections

Parasite	Protection		References
	Animal model	Human model	
Cestode			
<i>Mesocostoides corti</i>	—		89-91
<i>Hymenolepis diminuta</i>			92
Trematode			
<i>Fasciola</i> species			35, 92
<i>Schistosoma</i> species	—	++	33, 34, 41, 43, 77, 93, 94
Nematodes			
<i>Angiostrongylus</i> species	++		48, 49, 95
<i>H polygyrus</i>	—		9, 96
<i>Brugia</i> species	++		97
<i>Litomosoides</i> species	++		98
<i>Nippostrongylus</i> species	±		10, 99
<i>Onchocerca</i> species	++	±	50-53, 60, 71, 73, 75, 100, 101
<i>Strongyloides</i> species	++		44-46
<i>Toxocara</i> species	—		102-105
<i>Trichinella</i> species	±		39, 42, 106-108
<i>Trichuris</i> species	—		96, 109-111

—, Shown to play no role; ++, documented role; ±, conflicting information.

(Table II). Overall, the discrepancy seen in the various studies examining the role of eosinophils in protective immunity in different animal models of parasite infections has been attributed to the redundancy of effector immune responses or to the basic differences in the animal models used.³⁷

Animal models

Animal models of helminth infection have allowed for the assessment of the eosinophil's role in vivo. Treatment with anti-IL-5 antibody prevents the tissue and blood eosinophilia that occurs in animal models of gastrointestinal helminth infection.^{10,38} However, such treatment has not been shown to increase susceptibility to infection in murine models of *T spiralis*,³⁹ *Trichuris muris*,⁴⁰ *Nippostrongylus brasiliensis*,¹⁰ *H polygyrus*,⁹ or schistosomiasis.⁴¹ Similarly, IL-5 transgenic mice that have persistent increases of eosinophil levels do not demonstrate enhanced immunity to *T spiralis* or *S mansoni* infection.^{42,43}

Eosinophils do appear to play a role in the killing of some helminths, particularly those with tissue-migratory stages. Larval stages of many of the intestinal helminths, including *Strongyloides*, *Nippostrongylus*, and *Ascaris*

species, migrate through the lungs or skin on their way to the gastrointestinal tract. Although treatment with anti-IL-5 antibody had no effect on primary infection with *Strongyloides venezuelensis* in mice, recovery of worms from the lungs after challenge infection was increased in anti-IL-5-treated mice compared with that in control animals.⁴⁴ The most conclusive in vivo evidence for the role of eosinophils in helminth killing comes from experiments with diffusion chambers containing infective *Strongyloides stercoralis* larvae implanted in the subcutaneous tissue of mice. Not only were eosinophils the only cells that accumulated in the chamber concomitant with larval killing,^{45,46} but such killing could be prevented by means of inhibition of eosinophil migration into the chamber or treatment of the mice with anti-IL-5 antibody.⁴⁵⁻⁴⁷

In *Angiostrongylus cantonensis* infection in mice, immature worms develop in the brain before migrating to the lungs. Prolonged survival of intracranial worms resulting in a larger parasite burden in the lungs has been demonstrated in normal mice treated with anti-IL-5 antibody,⁴⁸ as well as in IL-5 receptor α -deficient mice,⁴⁹ which is again consistent with a role for eosinophils in tissue-based larval killing.

TABLE III. The role of the eosinophil in mediated pathology in selected parasitic helminth infections

Parasite	Host	Organ	Eosinophil role in pathology	Reference
<i>Angiostrongylus</i> species	Mouse	Brain	+	112, 113
	Guinea pig		+	
	Rat		–	
<i>Anisakis</i> species	Human	Intestine	+	69
<i>Baylisascaris</i> species	Human	Brain	+	70, 114
<i>Fasciola</i> species	Mouse	Liver	+	92
<i>Mesostoides</i> species	Mouse	Lung	–	91
<i>Nippostrongylus</i> species	Mouse	Lung	–	10
<i>Onchocerca</i> species	Mouse	Eye	+	60, 71, 73, 100, 115
		Skin	+	
<i>Schistosoma</i> species	Mouse	Liver	–	11, 77, 116, 117
	Human	Bladder	+	
<i>Toxocara</i> species	Mouse	Lung	+	102, 103

Animal models of protective immunity to the exclusively tissue-dwelling nematode *Onchocerca* species further support this hypothesis. In diffusion chamber studies of infective larval survival in immune mice, eosinophils were the only cells that accumulated in the chambers at the time of larval killing.⁵⁰ Furthermore, similar to the results with *Strongyloides* species larvae, eosinophil larval contact was necessary for larval destruction to occur.⁵⁰ In a different murine model system, mice infected with *Onchocerca lienalis* depleted of eosinophils (but not macrophages or neutrophils) with mAb showed delayed clearance of primary infection and abrogation of resistance to secondary infection.⁵¹

Overall, despite the prevailing dogma, the animal models do not strongly support an indispensable role for eosinophils in immunity to most helminth parasites in vivo.

Human studies

In human subjects with onchocerciasis, the picture is somewhat more complex. Some studies of in vitro cytokine production in response to parasite antigens have demonstrated increased levels of IL-5 production by PBMCs from putatively immune individuals living in areas endemic for onchocerciasis in comparison with infected individuals,⁵² whereas others have detected no difference in parasite antigen-induced IL-5 production between these groups.⁵³ The most compelling association between eosinophilia and protective immunity in human subjects comes from the elegant posttreatment reinfection studies in schistosomiasis that demonstrated a direct relationship between the lack of reinfection and the level of peripheral blood eosinophils in patients from areas endemic for both *S. mansoni* and *S. haematobium* infections.^{33,34}

Mechanism of eosinophil-mediated killing of helminth parasites

The mechanism by which eosinophil-mediated protection against helminth infection occurs is incompletely understood, but in most cases it appears to involve antibody-induced release, complement-induced release, or both of toxic granule proteins and reactive oxygen inter-

mediates by activated eosinophils. Interestingly, evidence suggests that there might be significant differences in the susceptibility to and mechanisms of eosinophil-mediated killing between different life-cycle stages of the same parasite.^{54,55} Unlike protozoa, helminths are too big to be engulfed by phagocytes and can only be killed by these cells when the latter have been activated by products of the adaptive immune response.

Eosinophils, which frequently accumulate in tissues soon after worm invasion, might also play a role in innate defense against this type of parasitic pathogen. For example, L3 larvae of *S. stercoralis* are killed within 72 hours when implanted in chambers in susceptible mice,^{45,47} with eosinophils being prominent among the cells infiltrating the chambers. Larval attrition is greatly reduced when worms are transferred instead into IL-5-deficient mice. This rapid and nonspecific eosinophilic response might be a barrier-limiting mechanism active against the invasion of many tissue-dwelling helminths.

EOSINOPHILS AND PATHOGENESIS

Eosinophils have also been implicated in the pathogenesis of helminth infection (Table III). Similarities between the sequelae of hypereosinophilic syndrome, a syndrome characterized by extremely high levels of eosinophilia and eosinophil-mediated end-organ damage,⁵⁶ and the pathologic consequences of infection with helminths, including loiasis and lymphatic filariasis^{57,58} and toxocariasis,⁵⁹ suggest a primary role for eosinophils in the pathogenesis of these infections. The association between the severity of clinical manifestations and the degree of eosinophil activation, as exemplified by increased eosinophil responsiveness to platelet-activating factor in patients with sowda, a hyperreactive form of onchocerciasis associated with severe dermal pathology,⁶⁰ provides additional indirect support for this hypothesis.

Although eosinophils can cause tissue damage through a number of different mechanisms, including direct cellular cytotoxicity, physical damage caused by tissue infiltration (space occupying), and thromboembolic phenomena caused by eosinophil-induced hypercoagulability,⁵⁶ it

is the release of eosinophil granule proteins and other cytotoxic substances that has been best correlated with helminth-induced tissue damage. Eosinophil cationic proteins (including major basic protein, eosinophil cationic protein, and eosinophil-derived neurotoxin) are toxic to various normal tissues and cells both in vitro and in vivo.^{7,61,62} Other potentially toxic inflammatory mediators that are released by activated eosinophils include leukotrienes,⁶³ platelet-activating factor,⁶⁴ reactive oxygen species,⁶⁵ and lysosomal hydrolases.⁶² These mediators not only damage the parasite and surrounding tissues (reactive oxygen intermediates^{66,67} and hydrolases) but play a role in the perpetuation of the local immune response (platelet-activating factor and leukotrienes).

Serum and urine levels of eosinophil granule proteins provide an indirect measure of degranulation in the tissues and are markedly increased in many helminth-infected patients.⁶⁸ Furthermore, deposition of eosinophil granule proteins in the affected tissues of patients with helminth infection has also been described.^{69,70} One of the characteristics of the posttreatment reaction in onchocerciasis is an increase in blood and tissue levels of eosinophils and eosinophil granule proteins that is temporally related to the onset of symptomatology^{71,72} and preceded by an increase in serum levels of IL-5.⁷³ Although localization of eosinophils and their products in the skin is most pronounced at sites of microfilarial degeneration,⁷¹ which is consistent with the hypothesis that eosinophil-mediated parasite killing plays a major role in pathogenesis, the possibility that recruitment of eosinophils occurs secondarily in response to parasite death cannot be excluded.

More direct evidence that eosinophils contribute to the pathogenesis of helminth infection comes from animal models. Eosinophilic infiltration of the cornea and conjunctival tissue, a characteristic component of onchocercal sclerosing keratitis in human subjects, occurs within 72 hours of intrastromal injection of sensitized mice with onchocercal antigens.⁷⁴ Attenuation of both the eosinophil response and ocular inflammation is observed in IL-4 knockout mice on a C57BL/6 background,⁷⁴ whereas eosinophil infiltration and corneal opacification is increased in mice treated with IL-12 at the time of corneal sensitization to onchocercal antigens despite decreased local expression of T_H2 cytokines, including IL-5.⁷⁵ The presence of eosinophil infiltration in a T_H1 environment could be accounted for by a marked increase in corneal chemokines, including RANTES and eotaxin.⁷⁵ Taken together, these data suggest that eosinophils, rather than the cytokine milieu, are the crucial factor in the development of onchocercal keratitis.

Although it seems clear that eosinophils play a role in some models of helminth-induced pathology, they do not appear to be essential to the development of pathologic sequelae in all cases. Liver granulomas in schistosomiasis typically contain a high percentage of activated eosinophils, the major source of T_H2-associated cytokines in the granuloma milieu.⁷⁶ Despite this, anti-IL-5-treated mice infected with schistosomiasis produce normal-sized granulomas, despite the absence of

eosinophils.⁷⁷ Similarly, lung infiltrates in *N. brasiliensis*-infected mice treated with anti-IL-5 differ from those of *N. brasiliensis*-infected control mice in the absence of eosinophils but not in size or number of lesions.¹⁰ Finally, in contrast to the attenuation of ocular disease observed in IL-4^{-/-} C57BL/6J mice, corneal opacification is exacerbated in sensitized IL-4^{-/-} BALB/cJ, STAT6^{-/-} BALB/cJ, and IL-5^{-/-} C57BL/6J^{78,79} mice after intrastromal injection of onchocercal antigen and is accompanied by increased tissue infiltration by neutrophils (but not eosinophils).⁷⁹ Whether these findings reflect a true lack of involvement of eosinophils in the pathologic sequelae of natural infection with these parasites or the induction of alternative mechanisms of pathogenesis in the setting of dysregulation of the cytokine milieu remains to be elucidated.

CONCLUSIONS

Much has been learned about the biology of the eosinophil⁸⁰ and its role in the pathogenesis of allergic,⁸¹ gastrointestinal,⁸² and hypereosinophilic disorders⁸³ since its discovery in 1879. The importance of its role in parasitic infections remains a large question, despite the plethora of studies in animals and human populations. Although we wait for unequivocal evidence for the salutary role of the eosinophil in helminth infections, we are left still with strong epidemiologic support for the eosinophil's benefit. Although unsatisfying as a conclusion, we must endeavor to elucidate its role more definitively if we consider the billions of persons worldwide infected with these multicellular and elusive parasitic organisms.

REFERENCES

- Markell EK, Voge M. Medical parasitology. 4th ed. Philadelphia: W.B. Saunders Company; 1976.
- Neva FA, Ottesen EA. Tropical (filarial) eosinophilia. *N Engl J Med* 1978;298:1129-31.
- Pelletier LJ, Baker CB, Gam AA, Nutman TB, Neva FA. Diagnosis and evaluation of treatment of chronic strongyloidiasis in ex-prisoners of war. *J Infect Dis* 1988;157:573-6.
- Smithers SR, Terry RJ. Concomitant immunity. *Adv Parasitol* 1976;14:399-428.
- Damonville M, Auriault C, Verwaerde C, Delanoye A, Pierce R, Capron A. Protection against experimental *Schistosoma mansoni* schistosomiasis achieved by immunization with schistosomula released products antigens (SRP-A): role of IgE antibodies. *Clin Exp Immunol* 1986;65:244-52.
- Ridel PR, Auriault C, Darcy F, Pierce RJ, Leite P, Santoro F, et al. Protective role of IgE in immunocompromised rat toxoplasmosis. *J Immunol* 1988;141:978-83.
- Gleich GJ, Frigas E, Loegering DA, Wassom DL, Steinmuller D. Cytotoxic properties of the eosinophil major basic protein. *J Immunol* 1979;123:2925-7.
- Maizels RM, Yazdanbakhsh M. Immunoregulation by helminth parasites: cellular and molecular mechanisms. *Nat Rev Immunol* 2003;3:733-44.
- Finkelman FD, Shea-Donohue T, Goldhill J, Sullivan CA, Morris SC, Madden KB, et al. Cytokine regulation of host defense against parasitic gastrointestinal nematodes: lessons from studies with rodent models. *Annu Rev Immunol* 1997;15:505-33.
- Coffman RL, Seymour BW, Hudak S, Jackson J, Rennick D. Antibody to interleukin-5 inhibits helminth-induced eosinophilia in mice. *Science* 1989;245:308-10.

11. Fallon PG, Richardson EJ, McKenzie GJ, McKenzie AN. Schistosome infection of transgenic mice defines distinct and contrasting pathogenic roles for IL-4 and IL-13: IL-13 is a profibrotic agent. *J Immunol* 2000;164:2585-91.
12. Mochizuki M, Bartels J, Mallet AI, Christophers E, Schroder JM. IL-4 induces eotaxin: a possible mechanism of selective eosinophil recruitment in helminth infection and atopy. *J Immunol* 1998;160:60-8.
13. Gurish MF, Humbles A, Tao H, Finkelstein S, Boyce JA, Gerard C, et al. CCR3 is required for tissue eosinophilia and larval cytotoxicity after infection with *Trichinella spiralis*. *J Immunol* 2002;168:5730-6.
14. Cooper PJ, Beck LA, Espinel I, Deyampert NM, Hartnell A, Jose PJ, et al. Eotaxin and RANTES expression by the dermal endothelium is associated with eosinophil infiltration after ivermectin treatment of onchocerciasis. *Clin Immunol* 2000;95:51-61.
15. Lavender P, Cousins D, Lee T. Regulation of Th2 cytokine gene expression. *Chem Immunol* 2000;78:16-29.
16. Watanabe N, Tamauchi H, Ozawa H, Ito M, Ovary Z, Habu S. Th2 immune responses in GATA-3-transgenic mice infected with *Heligmosomoides polygyrus*. *Int Arch Allergy Immunol* 2003;13(suppl 1):11-4.
17. Bochner BS. Systemic activation of basophils and eosinophils: markers and consequences. *J Allergy Clin Immunol* 2000;106(suppl):S292-302.
18. Hua XX, Sugaya H, Yoshimura K. Alteration in density of eosinophils in the cerebrospinal fluid of mice infected with *Angiostrongylus cantonensis*. *Int J Parasitol* 1990;20:681-3.
19. Sugaya H, Abe T, Yoshimura K. Eosinophils in the cerebrospinal fluid of mice infected with *Angiostrongylus cantonensis* are resistant to apoptosis. *Int J Parasitol* 2001;31:1649-58.
20. Mawhorter SD, Stephany DA, Ottesen EA, Nutman TB. Identification of surface molecules associated with physiologic activation of eosinophils: application of whole blood flow cytometry to eosinophils. *J Immunol* 1996;156:4851-8.
21. Hiatt RA, Sotomayor ZR, Sanchez G, Zambrana M, Knight WB. Factors in the pathogenesis of acute schistosomiasis mansoni. *J Infect Dis* 1979;139:659-66.
22. Lawley TJ, Ottesen EA, Hiatt RA, Gazze LA. Circulating immune complexes in acute schistosomiasis. *Clin Exp Immunol* 1979;37:221-7.
23. Maxwell C, Hussain R, Nutman TB, Poindexter RW, Little MD, Schad GA, et al. The clinical and immunologic responses of normal human volunteers to low dose hookworm (*Necator americanus*) infection. *Am J Trop Med Hyg* 1987;37:126-34.
24. Nutman TB. Experimental infection of humans with filariae. *Rev Infect Dis* 1991;13:1018-22.
25. Cline BL, Little MD, Bartholomew RK, Halsey NA. Larvicidal activity of albendazole against *Necator americanus* in human volunteers. *Am J Trop Med Hyg* 1984;33:387-94.
26. Ogilvie BM, Bartlett A, Godfrey RC, Turton JA, Worms MJ, Yeates RA. Antibody responses in self-infections with *Necator americanus*. *Trans R Soc Trop Med Hyg* 1978;72:66-71.
27. Taliaferro WH, Sables MP. The cellular reactions in the skin, lungs, and intestine of normal and immune rats after infection with *Nippostrongylus brasiliensis*. *J Infect Dis* 1939;64:157-92.
28. David JR, Butterworth AE, Vadas MA. Mechanism of interaction mediating killing of *Schistosoma mansoni* by human eosinophils. *Am J Trop Med Hyg* 1980;29:842-8.
29. Haque A, Ouaisi A, Joseph M, Capron M, Capron A. IgE antibody in eosinophil- and macrophage-mediated in vitro killing of *Dipetalonema viteae* microfilariae. *J Immunol* 1981;127:716-25.
30. Kazura JW, Grove DI. Stage-specific antibody-dependent eosinophil-mediated destruction of *Trichinella spiralis*. *Nature* 1978;274:588-9.
31. Hamann KJ, Gleich GJ, Checkel JL, Loegering DA, McCall JW, Barker RL. In vitro killing of microfilariae of *Brugia pahangi* and *Brugia malayi* by eosinophil granule proteins. *J Immunol* 1990;144:3166-73.
32. Butterworth AE, Wassom DL, Gleich GJ, Loegering DA, David JR. Damage to schistosome of *Schistosoma mansoni* induced directly by eosinophil major basic protein. *J Immunol* 1979;122:221-9.
33. Sturrock RF, Kimani R, Cottrell BJ, Butterworth AE, Seitz HM, Siongok TK, et al. Observations on possible immunity to reinfection among Kenyan schoolchildren after treatment for *Schistosoma mansoni*. *Trans R Soc Trop Med Hyg* 1983;77:363-71.
34. Hagan P, Wilkins HA, Blumenthal UJ, Hayes RJ, Greenwood BM. Eosinophilia and resistance to *Schistosoma haematobium* in man. *Parasite Immunol* 1985;7:625-32.
35. Behm CA, Ovington KS. The role of eosinophils in parasitic helminth infections: insights from genetically modified mice. *Parasitol Today* 2000;16:202-9.
36. Meeusen EN, Balic A. Do eosinophils have a role in the killing of helminth parasites? *Parasitol Today* 2000;16:95-101.
37. Dombrowicz D, Capron M. Eosinophils, allergy and parasites. *Curr Opin Immunol* 2001;13:716-20.
38. Rennick DM, Thompson-Snipes L, Coffman RL, Seymour BW, Jackson JD. In vivo administration of antibody to interleukin-5 inhibits increased generation of eosinophils and their progenitors in bone marrow of parasitized mice. *Blood* 1990;76:312-6.
39. Herndon FJ, Kayes SG. Depletion of eosinophils by anti-IL-5 monoclonal antibody treatment of mice infected with *Trichinella spiralis* does not alter parasite burden or immunologic resistance to infection. *J Immunol* 1992;149:3642-7.
40. Urban JFJ, Katona IM, Paul WM, Findelman RD. Interleukin 4 is important in protective immunity to a gastrointestinal nematode infection in mice. *Proc Natl Acad Sci U S A* 1991;88:5513-7.
41. Sher A, Coffman RF, Hieny S, Cheever AW. Ablation of eosinophil and IgE responses with anti-IL-5 or anti-IL-4 antibodies fails to affect immunity against *Schistosoma mansoni* in the mouse. *J Immunol* 1990;145:3911-6.
42. Hokibara S, Takamoto M, Tominaga A, Takatsu K, Sugane K. Marked eosinophilia in interleukin-5 transgenic mice fails to prevent *Trichinella spiralis* infection. *J Parasitol* 1997;83:1186-9.
43. Dent LA, Munro GH, Piper KP, Sanderson CJ, Finlay DA, Dempster RK, et al. Eosinophilic interleukin 5 (IL-5) transgenic mice: eosinophil activity and impaired clearance of *Schistosoma mansoni*. *Parasite Immunol* 1997;19:291-300.
44. Korenaga M, Hitoshi Y, Yamaguchi N, Sato Y, Takatsu K, Tada I. The role of interleukin-5 in protective immunity to *Strongyloides venezuelensis* infection in mice. *Immunol* 1991;72:502-7.
45. Rotman HL, Yutanawiboonchai W, Brigandi RA, Leon O, Gleich GJ, Nolan TJ, et al. *Strongyloides stercoralis*: eosinophil-dependent immune-mediated killing of third stage larvae in BALB/cByJ mice. *Exp Parasitol* 1996;82:267-78.
46. Abraham D, Rotman HL, Haberstroff HF, Yutanawiboonchai W, Brigandi RA, Leon O, et al. *Strongyloides stercoralis*: protective immunity to third-stage larvae in BALB/cByJ mice. *Exp Parasitol* 1995;80:297-307.
47. Herbert DR, Lee JJ, Lee NA, Nolan TJ, Schad GA, Abraham D. Role of IL-5 in innate and adaptive immunity to larval *Strongyloides stercoralis* in mice. *J Immunol* 2000;165:4544-51.
48. Sasaki O, Sugaya H, Ishida K, Yoshimura K. Ablation of eosinophils with anti-IL-5 antibody enhances the survival of intracranial worms of *Angiostrongylus cantonensis* in the mouse. *Parasite Immunol* 1993;15:349-54.
49. Yoshida T, Ikuta K, Sugaya H, Maki K, Takagi M, Kanazawa H, et al. Defective B1-cell development and impaired immunity against *Angiostrongylus cantonensis* in IL-5R alpha-deficient mice. *Immunity* 1996;4:483-94.
50. Lange AM, Yutanawiboonchai W, Scott P, Abraham D. IL-4- and IL-5-dependent protective immunity to *Onchocerca volvulus* infective larvae in BALB/cByJ mice. *J Immunol* 1994;153:205-11.
51. Folkard SG, Hogarth PJ, Taylor MJ, Bianco AE. Eosinophils are the major effector cells of immunity to microfilariae in a mouse model of onchocerciasis. *Parasitology* 1996;112:323-9.
52. Steel C, Nutman TB. Regulation of IL-5 in onchocerciasis. A critical role for IL-2. *J Immunol* 1993;150:5511-8.
53. Elson LH, Calvopina M, Paredes W, Araujo E, Bradley JE, Guderian RH, et al. Immunity to onchocerciasis: putative immune persons produce a Th1-like response to *Onchocerca volvulus*. *J Infect Dis* 1995;171:652-8.
54. Brattig NW, Tischendorf FW, Strote G, Medina-de la Garza CE. Eosinophil-larval-interaction in onchocerciasis: heterogeneity of in vitro adherence of eosinophils to infective third and fourth stage larvae and microfilariae of *Onchocerca volvulus*. *Parasite Immunol* 1991;13:13-22.
55. MacDonald AS, Loke P, Martynoga R, Dransfield I, Allen JE. Cytokine-dependent inflammatory cell recruitment patterns in the peritoneal cavity of mice exposed to the parasitic nematode *Brugia malayi*. *Med Microbiol Immunol (Berl)* 2003;192:33-40.
56. Weller PF, Bubley GJ. The idiopathic hypereosinophilic syndrome. *Blood* 1994;83:2759-79.

57. Nutman TB, Miller KD, Mulligan M, Ottesen EA. Loa loa infection in temporary residents of endemic regions: recognition of a hyperresponsive syndrome with characteristic clinical manifestations. *J Infect Dis* 1986;154:10-8.
58. O'Bryan L, Pinkston P, Kumaraswami V, Vijayan V, Yenokida G, Rosenberg HF, et al. Localized eosinophil degranulation mediates disease in tropical pulmonary eosinophilia. *Infect Immun* 2003;71:1337-42.
59. Kayes SG. Human toxocariasis and the visceral larva migrans syndrome: correlative immunopathology. *Chem Immunol* 1997;66:99-124.
60. de Krömer MT, Medina-De-la-Garza CE, Brattig NW. Differences in eosinophil and neutrophil chemotactic responses in sowda and generalized form of onchocerciasis. *Acta Trop* 1995;60:21-33.
61. Gleich GJ, Schroeter AL, Marcoux JP, Sachs MI, O'Connell EJ, Kohler PF. Episodic angioedema associated with eosinophilia. *N Engl J Med* 1984;310:1621-6.
62. Costa JJ, Weller PF, Galli SJ. The cells of the allergic response: mast cells, basophils, and eosinophils. *JAMA* 1997;278:1815-22.
63. Shaw RJ, Walsh GM, Cromwell O, Moqbel R, Spry CJ, Kay AB. Activated human eosinophils generate SRS-A leukotrienes following IgG-dependent stimulation. *Nature* 1985;316:150-2.
64. Cromwell O, Wardlaw AJ, Champion A, Moqbel R, Osei D, Kay AB. IgG-dependent generation of platelet-activating factor by normal and low density human eosinophils. *J Immunol* 1990;145:3862-8.
65. McCormick ML, Metwali A, Railsback MA, Weinstock JV, Britigan BE. Eosinophils from schistosome-induced hepatic granulomas produce superoxide and hydroxyl radical. *J Immunol* 1996;157:5009-15.
66. Thomas GR, McCrossan M, Selkirk ME. Cytostatic and cytotoxic effects of activated macrophages and nitric acid donors on *Brugia malayi*. *Infect Immun* 1997;65:2732-9.
67. Yazdanbakhsh M, Tai PC, Spry CJ, Gleich GJ, Roos D. Synergism between eosinophil cationic protein and oxygen intermediates in killing of schistosomula of *Schistosoma mansoni*. *J Immunol* 1987;138:3443-7.
68. Tischendorf FW, Brattig NW, Burchard GD, Kubica T, Kreuzpaintner G, Lintzel M. Eosinophils, eosinophil cationic protein and eosinophil-derived neurotoxin in serum and urine of patients with onchocerciasis coinfecting with intestinal nematodes and in urinary schistosomiasis. *Acta Trop* 1999;72:157-73.
69. del Pozo V, Arrieta I, Tunon T, Cortegano I, Gomez B, Cardaba B, et al. Immunopathogenesis of human gastrointestinal infection by *Anisakis simplex*. *J Allergy Clin Immunol* 1999;104:637-43.
70. Hamann KJ, Kephart GM, Kazacos KR, Gleich GJ. Immunofluorescent localization of eosinophil granule major basic protein in fatal human cases of *Baylisascaris procyonis* infection. *Am J Trop Med Hyg* 1989;40:291-7.
71. Ackerman SJ, Kephart GM, Francis H, Awadzi K, Gleich GJ, Ottesen EA. Eosinophil degranulation: an immunologic determinant in the pathogenesis of the Mazzotti reaction in human onchocerciasis. *J Immunol* 1990;144:3961-9.
72. Cooper PJ, Awadzi K, Ottesen EA, Remick D, Nutman TB. Eosinophil sequestration and activation are associated with the onset and severity of systemic adverse reactions following the treatment of onchocerciasis with ivermectin. *J Infect Dis* 1999;179:738-42.
73. Limaye AP, Abrams JS, Silver JE, Awadzi K, Francis HF, Ottesen EA, et al. Interleukin-5 and the posttreatment eosinophilia in patients with onchocerciasis. *J Clin Invest* 1991;88:1418-21.
74. Pearlman E, Lass JH, Bardenstein DS, Kopf M, Hazlett FE, Jr, Diaconu E, et al. Interleukin 4 and T helper type 2 cells are required for the development of onchocercal keratitis (river blindness). *J Exp Med* 1995;182:931-40.
75. Pearlman E, Lass JH, Bardenstein DS, Diaconu E, Hazlett JFE, Albright J, et al. IL-12 exacerbates helminth-mediated corneal pathology by augmenting inflammatory cell recruitment and chemokine expression. *J Immunol* 1997;158:827-33.
76. Rumbley CA, Sugaya H, Zekavat SA, El Refaei M, Perrin PJ, Phillips SM. Activated eosinophils are the major source of Th2-associated cytokines in the schistosome granuloma. *J Immunol* 1999;162:1003-9.
77. Sher A, Coffman RL, Hieny S, Scott P, Cheever AW. Interleukin 5 is required for the blood and tissue eosinophilia but not granuloma formation induced by infection with *Schistosoma mansoni*. *Proc Natl Acad Sci* 1990;87:61-5.
78. Pearlman E, Hall LR, Higgins AW, Bardenstein DS, Diaconu E, Hazlett FE, et al. The role of eosinophils and neutrophils in helminth-induced keratitis. *Invest Ophthalmol Vis Sci* 1998;39:1176-82.
79. Hall LR, Berger RB, Diaconu E, Pearlman E. *Onchocerca volvulus* keratitis (river blindness) is exacerbated in BALB/c IL4 gene knockout mice. *Cell Immunol* 2002;216:1-5.
80. Bochner BS. Verdict in the case of therapies versus the eosinophil: the jury is still out. *J Allergy Clin Immunol* 2004;113:3-9.
81. Hamid Q. Eosinophils in allergic diseases. *J Allergy Clin Immunol* 2004;113:182-4.
82. Rothenberg ME. Eosinophilic gastrointestinal disorders (EGID). *J Allergy Clin Immunol* 2004;113:11-28.
83. Leiferman K, Gleich GJ. Hypereosinophilic syndromes. *J Allergy Clin Immunol* 2004;113:50-8.
84. Buckley JJC. Tropical pulmonary eosinophilia in relation to filarial infections (*Wuchereria* spp.) of animals. Preliminary note. *Trans R Soc Trop Med Hyg* 1958;52:335-6.
85. Edeson JFB, Wilson T, Wharton RH, Laing ABG. Experimental transmission of *Brugia malayi* and *Brugia pahangi* to man. *Trans R Soc Trop Med Hyg* 1960;54:229-34.
86. Buckley JJC, Wharton RH. Anomalous results from an experimental infection of man with *Brugia malayi* (Brug, 1927). *J Helminthol RT Leiper* 1961;(Suppl);17-24.
87. Dondero TJ Jr, Mullin SW, Balasingam S. Early clinical manifestations in filariasis due to *Brugia malayi*: observations on experimental infections in man. *Southeast Asian J Trop Med Public Health* 1972;3:569-75.
88. Liu JY, Wang GQ, Chen YN, Tu ZP, Fang Z. Observations on the clinical manifestations and treatment of an experimental infection with *Brugia malayi* in man. *J Trop Med Hyg* 1989;92:93-6.
89. Strath M, Dent L, Sanderson C. Infection of IL5 transgenic mice with *Mesocostoides corti* induces very high levels of IL5 but depressed production of eosinophils. *Exp Hematol* 1992;20:229-34.
90. Dent LA, Strath M, Mellor AL, Sanderson CJ. Eosinophilia in transgenic mice expressing interleukin 5. *J Exp Med* 1990;172:1425-31.
91. Kopf M, Brombacher F, Hodgkin PD, Ramsay AJ, Milbourne EA, Dai WJ, et al. IL-5-deficient mice have a developmental defect in CD5+ B-1 cells and lack eosinophilia but have normal antibody and cytotoxic T cell responses. *Immunity* 1996;4:15-24.
92. Ovington KS, Behm CA. The enigmatic eosinophil: investigation of the biological role of eosinophils in parasitic helminth infection. *Mem Inst Oswaldo Cruz* 1997;92:93-104.
93. Tischendorf FW, Brattig NW, Buttner DW, Pieper A, Lintzel M. Serum levels of eosinophil cationic protein, eosinophil-derived neurotoxin and myeloperoxidase in infections with filariae and schistosomes. *Acta Trop* 1996;62:171-82.
94. Knopf PM, Nutman TB, Reasoner JA. *Schistosoma mansoni*: resistance to reinfection in the rat. *Exp Parasitol* 1977;41:74-82.
95. Sugaya H, Aoki M, Yoshida T, Takatsu K, Yoshimura K. Eosinophilia and intracranial worm recovery in interleukin-5 transgenic and interleukin-5 receptor alpha chain-knockout mice infected with *Angiostrongylus cantonensis*. *Parasitol Res* 1997;83:583-90.
96. Urban JF Jr, Katona IM, Paul WE, Finkelman FD. Interleukin 4 is important in protective immunity to a gastrointestinal nematode infection in mice. *Proc Natl Acad Sci* 1991;88:5513-7.
97. Rajan TV, Ganley L, Paciorkowski N, Spencer L, Klei TR, Shultz LD. Brugian infections in the peritoneal cavities of laboratory mice: kinetics of infection and cellular responses. *Exp Parasitol* 2002;100:235-47.
98. Martin C, Le Goff L, Ungeheuer MN, Vuong PN, Bain O. Drastic reduction of a filarial infection in eosinophilic interleukin-5 transgenic mice. *Infect Immun* 2000;68:3651-6.
99. Zhou Y, Bao S, Rothwell TLW, Husband AJ. Differential expression of interleukin-5 mRNA+ cells and eosinophils in *Nippostrongylus brasiliensis* infection in resistant and susceptible strains of mice. *Eur J Immunol* 1996;26:2133-9.
100. Pearlman E. Immunopathology of onchocerciasis: a role for eosinophils in onchocercal dermatitis and keratitis. *Chem Immunol* 1997;66:26-40.
101. Hogarth PJ, Bianco AE. IL-5 dominates cytokine responses during expression of protective immunity to *Onchocerca lienalis* microfilariae in mice. *Parasite Immunol* 1999;21:81-8.
102. Takamoto M, Ovington KS, Behm CA, Sugane K, Young IG, Matthaei KI. Eosinophilia, parasite burden and lung damage in *Toxocara canis* infection in C57BL/6 mice genetically deficient in IL-5. *Immunol* 1997;90:511-7.
103. Cookston M, Stober M, Kayes S. Eosinophilic myocarditis in CBA/J mice infected with *Toxocara canis*. *Am J Pathol* 1990;136:1137-45.

104. Sugane K, Kusama Y, Takamoto M, Tominaga A, Takatsu K. Eosinophilia, IL-5 level and recovery of larvae in IL-5 transgenic mice infected with *Toxocara canis*. *J Helminthol* 1996;70:153-8.
105. Dent LA, Daly CM, Mayrhofer G, Zimmerman T, Hallett A, Bignold LP, et al. Interleukin-5 transgenic mice show enhanced resistance to primary infections with *Nippostrongylus brasiliensis* but not primary infections with *Toxocara canis*. *Infect Immun* 1999;67:989-93.
106. Grove DI, Hamburger J, Warren KS. Kinetics of immunological responses, resistance to reinfection, and pathological reactions to infection with *Trichinella spiralis*. *J Infect Dis* 1977;136:562-70.
107. Artis D, Humphreys NE, Potten CS, Wagner N, Muller W, McDermott JR, et al. Beta7 integrin-deficient mice: delayed leukocyte recruitment and attenuated protective immunity in the small intestine during enteric helminth infection. *Eur J Immunol* 2000;30:1656-64.
108. Vallance BA, Blennerhassett PA, Deng Y, Matthaie KI, Young IG, Collins SM. IL-5 contributes to worm expulsion and muscle hypercontractility in a primary *T. spiralis* infection. *Am J Physiol* 1999;277:G400-8.
109. Betts CJ, Else KJ. Mast cells, eosinophils and antibody-mediated cellular cytotoxicity are not critical in resistance to *Trichuris muris*. *Parasite Immunol* 1999;21:45-52.
110. Else KJ, Entwistle GM, Grecis RK. Correlations between worm burden and markers of Th1 and Th2 cell subset induction in an inbred strain of mouse infected with *Trichuris muris*. *Parasite Immunol* 1993;15:595-600.
111. Else KJ, deSchoolmeester ML. Immunity to *Trichuris muris* in the laboratory mouse. *J Helminthol* 2003;77:95-8.
112. Du WY, Liao JW, Fan CK, Su KE. Combined treatment with interleukin-12 and mebendazole lessens the severity of experimental eosinophilic meningitis caused by *Angiostrongylus cantonensis* in ICR mice. *Infect Immun* 2003;71:3947-53.
113. Perez O, Capron M, Lastre M, Venge P, Khalife J, Capron A. *Angiostrongylus cantonensis*: role of eosinophils in the neurotoxic syndrome (Gordon-like phenomenon). *Exp Parasitol* 1989;68:403-13.
114. Moertel CL, Kazacos KR, Butterfield JH, Kita H, Watterson J, Gleich GJ. Eosinophil-associated inflammation and elaboration of eosinophil-derived proteins in 2 children with raccoon roundworm (*Baylisascaris procyonis*) encephalitis. *Pediatrics* 2001;108:e93.
115. Pearlman E, Toe L, Boatn BA, Gilles AA, Higgins AW, Unnasch TR. Eotaxin expression in *Onchocerca volvulus*-induced dermatitis after topical application of diethylcarbamazine. *J Infect Dis* 1999;180:1394-7.
116. Brunet LR, Sabin EA, Cheever AW, Kopf MA, Pearce EJ. Interleukin 5 (IL-5) is not required for expression of a Th2 response or host resistance mechanisms during murine *Schistosoma mansoni* but does play a role in development of IL-4-producing non-T, non-B cells. *Infect Immun* 1999;67:3014-8.
117. Chiamonte MG, Schopf LR, Neben TY, Cheever AW, Donaldson DD, Wynn TA. IL-13 is a key regulatory cytokine for Th2 cell-mediated pulmonary granuloma formation and IgE responses induced by *Schistosoma mansoni* eggs. *J Immunol* 1999;162:920-30.