

Title

Selectivity for Alkynyl or Allenyl Imidamides and Imidates in Copper-Catalyzed Reactions of Terminal 1,3-Diynes and Azides

authors

Sourav Ghorai and Daesung Lee*



publisher

Cite This: *Org. Lett.* 2021, 23, 697–701



other

Read Online

Headline

ACCESS

other

Metrics & More

other

Article Recommendations

other

Supporting Information

image
figure

ABSTRACT: Copper-catalyzed reactions of terminal 1,3-diynes with electron-deficient azides to generate either 3-alkynyl or 2,3-dienyl imidamides and imidates are described. The selectivity depends on the diyne substituents and the nucleophile that reacts with the ketenimine intermediate generated from the corresponding triazole precursor. Reactions of 1,3-diynes containing a propargylic acetate afford [3]cumulenyl imidamides, while reactions using methanol as the trapping agent selectively generate 2,3-dienyl imidates. Five-membered heterocycles were obtained from 1,3-diynes containing a homopropargylic hydroxyl or amine substituent.

Content

Allenes constitute a distinct class of organic compounds with two orthogonal π -bonds. There are numerous natural products and bioactive molecules that contain allene substructures.^{1–3} It has been demonstrated that allene-substituted bioactive compounds, like steroids,^{4,5} prostaglandins,^{6,7} carbacyclins,⁷ nucleosides,⁸ and unnatural amino acids,⁹ display higher potency, increased metabolic stability, and bioavailability. Because of the strained nature of the cumene structure, allenes have been engaged in numerous synthetic transformations as a versatile building block to form a variety of carbo- and heterocyclic frameworks.^{10–19} Axial-to-central chirality transfer is an efficient method for generating chiral compounds containing one or more stereogenic centers from chiral allenes.^{19–22}

Due to the important utility of allenes, it is highly desirable to develop new synthetic methods to generate functionalized allenes from readily available starting materials.^{23–28} One of the traditional approaches to the synthesis of allenes involves 1,2-elimination of vinyl derivatives under strong basic or metal-catalyzed conditions.²⁹ S_N2' type reactions with propargylic alcohol derivatives ($FG-CH_2-C\equiv C$)^{30–33} or isomerization of enyne moieties³⁴ also constitutes an efficient method for generating allenes. Transition metal-catalyzed coupling between terminal alkynes and carbonyl moieties in the presence of a secondary amine is also a well-established method,³⁵ and a Cu(I)-catalyzed cross-coupling of terminal alkynes with diazo compounds is another efficient protocol for generating allenes.³⁶ Trisubstituted allenes can be accessed via metal-catalyzed cationic Heck coupling of alkynes with aryl halide/triflate.^{37–39} Enones and ynones are also explored as a precursor of allenes.^{40,41}


In 2004, Fu reported a Cu(I)-catalyzed coupling of alkynes and diazoacetate under mild conditions to generate 3-alkynyl carboxylate and only small amounts of the corresponding allenates were observed (Scheme 1A).⁴² On the contrary, Lee

Headline


imageDescription

Scheme 1. Two Different Approaches to the Formation of Closely Related Functional Groups

A) Fu's Cu-catalyzed coupling of terminal alkyne and diazo compound



B) This work: Cu-catalyzed coupling of diyne and azide



Content

and others found that the alkynes containing a heteroatom substituent at the propargylic or homopropargylic carbon center preferentially generate the allenate, and in particular in

Other

Received: November 22, 2020

Published: January 14, 2021



publisher
ACS Publications

publisher

© 2021 American Chemical Society

small

<https://dx.doi.org/10.1021/acs.orglett.0c03861>
Org. Lett. 2021, 23, 697–701

the presence of a base such as triethyl amine, allenolate became the exclusive product.⁴³ Fox also reported a similar reaction that selectively generated allenolate as the product.⁴⁴ Although the exact mechanisms of these reactions are not well understood, on the basis of other related reactions, we propose a copper acetylide-mediated mechanism for the formation of the 3-alkynoate and allenolate. Relying on this copper acetylide-mediated mechanism, we envision that the copper-catalyzed coupling of terminal 1,3-diyne **1** and azide would generate 3-alkynyl and allenyl imidamide/imidate **2/3** (Scheme 1B). We surmise that the terminal 1,3-diyne would form the corresponding copper acetylide, which will participate in a [3+2] cycloaddition⁴⁵ with tosylazide to form triazole IN-1. The subsequent loss of molecular nitrogen will lead to two equilibrating organocopper aza-cummulenes IN-2 and IN-3,⁴⁶ which then react with a nucleophile such as an amine or alcohol to generate imidamide⁴⁷ or imidate⁴⁸ containing either an alkyne (**2**) or an allene moiety (**3**). A unique feature of this transformation is the selective transformation of the terminal alkyne moiety for the construction of imidamide and imidate functionality. Although starting from a different set of starting materials, the formation of closely related products from transformations in A and B is noteworthy. Herein, we describe our exploration of copper-catalyzed reactions of terminal 1,3-diyne **1** with tosylazide, which generated functionalized 1,3-disubstituted 2,3-dienyl imidamides and imidates **3** with good selectivity over the corresponding alkyne derivative **2**. Also, it was found that the types of nucleophiles, the base additives, and the substituent patterns of the 1,3-diynes not only affect the ratio of **2** to **3** but also promote the formation of alternative products such as [3]cumulenes and triple-bond-migrated products.


Our exploration commenced with the assessment of the efficiency and selectivity for the formation of 3-alkynyl and allenyl imidamide **2** and **3** (Table 1). Under the conditions that include a copper catalyst (CuI, 10 mol %), an azide (1.2 equiv), and an amine (1.2 equiv), 1,3-diynes containing different substituents were examined. A hexyl-substituted terminal 1,3-diyne **1a** and TsN₃ provided a mixture of 3-alkynyl and allenyl imidamides **2aa** and **3aa** in 75% yield with a

2:1 ratio (entry 1). Reactions of **1a** with other azides such as diphenyl phosphoryl azide⁴⁹ and mesyl azide are also efficient, affording 3-alkynyl imidamide **2ab** and **2ac** as the major products (entries 2 and 3, respectively). On the contrary, employing piperidine as a nucleophile under otherwise identical conditions, **1a** provided **2ad** exclusively (entry 4). 1,3-Diynes with a secondary (**1b**) or tertiary alkyl group (**1c**) provided good yields of **2/3ba** and **2/3ca** but low selectivity with a slight preference for alkynyl products **2ba** and **2ca** (entries 5 and 6, respectively). 1-Cyclohexenyl-substituted 1,3-diyne provided **2/3da** in 48% yield with a preference for compound **3da**. 1,3-Diyne **1e** containing a propargylic hydroxyl group afforded allene derivative **3ea** predominantly (entry 8), whereas 1,3-diyne **1f** containing a trimethylsilyl group provided alkyne derivative **2fa** selectively but in marginal yield (entry 9).

Once a general trend for the efficiency and selectivity between **2** and **3** had been revealed from the entries in Table 1, we next explored the selective formation of 3-alkynyl imidamide **2** (Scheme 2). As piperidine exclusively generated

ImageDescription

Scheme 2. Selective Formation of 3-Alkynyl Imidamides with Assorted Nucleophiles, Azides, and 1,3-Diynes



figure

^aYields in parentheses represent those of the corresponding 2,3-dienyl isomer **3**.

Content

Table 1. Efficiencies and Product Distributions with Assorted Nucleophiles, Azides, and 1,3-Diynes of Different Substituents

entry	R	Nu-H	azide	yield (%) (alkyne:allene) ^a
1	<i>n</i> -Hex	<i>i</i> -Pr ₂ NH	TsN ₃	2/3aa, 75% (2:1)
2	<i>n</i> -Hex	<i>i</i> -Pr ₂ NH	(PhO) ₂ PON ₃	2/3ab, 66% (5:1)
3	<i>n</i> -Hex	<i>i</i> -Pr ₂ NH	MsN ₃	2/3ac, 54% (2:1)
4	<i>n</i> -Hex	piperidine	TsN ₃	2ad, 56% (1:0) ^b
5	<i>i</i> -Pr	<i>i</i> -Pr ₂ NH	TsN ₃	2/3ba, 79% (2:3:1)
6	<i>i</i> -Pr	<i>i</i> -Pr ₂ NH	TsN ₃	2/3ca, 71% (1:3:1)
7	1-cyclohexenyl	<i>i</i> -Pr ₂ NH	TsN ₃	2/3da, 48% (1:2:5)
8	CH ₂ OH	<i>i</i> -Pr ₂ NH	TsN ₃	2/3ea, 73% (1:15)
9	<i>t</i> -BuMe ₃	<i>i</i> -Pr ₂ NH	TsN ₃	2fa, 45% (1:0)

figure

^aIsolated yield. ^bH₂O (1 equiv) and NH₂OH·HCl (1–2 mol %) were used as additives.


Content

3-alkynyl imidamide, reactions of *n*-decyl- and homobenzyl-substituted 1,3-diynes with piperidine were performed, resulting in the corresponding 3-alkynyl imidamides **2ac'** and **2ac''** in 57% and 40% yields, respectively. Reaction with phosphoryl azide and alkyl-substituted diynes predominantly generated 3-alkynyl imidamides (**2ab'**, **2ab''**, and **2bb**) in moderate 45–60% yields. Electron-rich 1,3-diynes with silyl substituents selectively delivered 3-alkynyl imidamides. TES-, TIPS-, and TBS-substituted 1,3-diynes provided the corresponding alkyne derivatives **2fa'–2fa'''** in 85–94% yields. Reactions of TIPS-substituted 1,3-diyne with phosphoryl azide and mesyl azide also generated 3-alkynyl imides **2fb''** and **2fc''** exclusively in 59% and 82% yields, respectively. It is evident from Scheme 2 that reactions with cyclic amine, phosphoryl azide, and electron-rich silyl-substituted 1,3-diynes tend to generate 3-alkynyl imidamide predominantly or exclusively.

Next, we examined reactions of different 1,3-diynes to selectively form 2,3-dienyl imidamide **3** (Scheme 3). On the basis of the initial observation with propargylic alcohol-containing diyne **1e** that selectively generated 2,3-dienyl imidamide, we further tested the reactivity of structurally diversified propargyl alcohols, (thio)ethers, amines, and

ImageDescription

Scheme 3. Formation of Allenyl Imidamides from Diverse 1,3-Terminal Diynes



footnote
“Yields in parentheses represent those of the corresponding 3-alkynyl compound 2.

Content


amides. 1,3-Diynes containing secondary or tertiary alcohols afforded 2,3-dienyl imidamides (3ga , $3\text{ga}'$, $3\text{ga}''$, $3\text{ga}'''$, 3ja , $3\text{ja}'$, $3\text{ja}''$, and 3ka) as a predominant or exclusive product in 65–76% yields. The selectivity between the allenyl and alkynyl isomer depends on the substituent, but no clear trend has been found. 1,3-Diynes with a tertiary alcohol and a cycloalkyl substituent delivered 2,3-dienyl imidamides (3la , $3\text{la}'$, and $3\text{la}''$) in 70–80% yields with a roughly 10:1 selectivity. Carvone-containing 1,3-diyne afforded allene 3ma (7:1 allene:alkyne ratio) in 69% yield with a 1.4:1 diastereomeric ratio. While a vinyl-conjugated 1,3-diyne containing a free hydroxyl group provided allene 3na in 46% yield with a low selectivity (4:1), 1,3-diynes with benzyl-, 3-MeO-Ph-, and THP-protected primary alcohols exclusively generated allene derivatives (3oa , $3\text{oa}'$, and $3\text{oa}''$, respectively) in 54–70% yields. Similarly, a cholesteryl ether-substituted 1,3-diyne generated allene 3pa in 56% yield with an 8:1 allene:alkyne ratio. On the contrary, the corresponding thioether afforded 3qa contaminated with the alkynyl isomer (7.5:1 allene:alkyne). 1,3-Diynes containing aniline and tosylamido substituents at the propargylic position selectively generated allenes 3ra and 3sa in 58% and 88% yields, respectively. A *gem*-dimethyl, however, decreased the yield and selectivity for $3\text{sa}'$ (68%, 8:5:1). An indole-substituted 1,3-diyne provided only allene 3ta in 68% yield, and *N*-allyl tosyl-substituted 1,3-diyne provided single isomer 3ua in 67% yield. On the contrary, phenyl- and 4-MeO-phenyl-substituted 1,3-diynes provided moderate yields and selectivity provided allenes 3va (64%, 4.6:1) and $3\text{va}'$ (64%, 3.6:1).

Content

Although the selectivity of forming allenyl imidamides is good, the formation of alkynyl isomer could not be suppressed in many cases. At this juncture, we surmised that trapping the ketenimine intermediate with alcohols may have different product distributions.^{50,51} Indeed, under identical conditions except for the replacement of *i*-Pr₂NH (1.2 equiv) with MeOH (10 equiv) and Et₃N (2 equiv), the reaction of 1,3-diynes selectively provided 2,3-dienyl imidates without a vestige of the alkynyl isomer (Table 2). The lower stoichiometry of methanol

Table

Table 2. Synthesis of Allenyl Imidates by Trapping with Methanol



Content

led to the lower efficiency of the reaction. Alkyl and alkenyl diynes, which provided a mixture of allenyl and alkynyl imidamides previously, selectively generated allenyl imidates (3ae , 3ce , and 3de) in moderate to good yield (entries 1–3, respectively). Even though 1,3-diyne 1g containing a free secondary alcohol led to decomposition (entry 4), the corresponding acyl-, benzyl-, and *tert*-butyldimethylsilyl-protected 1,3-diynes provided 2,3-dienyl imidates (3he , $3\text{he}'$, and $3\text{he}''$, respectively) in good yield (entries 5–7, respectively). Similarly, 1,3-diyne with benzyl-protected secondary alcohol 1i delivered 2,3-dienyl imidate 3ie in 72% yield.

Subsequently, we observed that under standard conditions, 1,3-diynes 4 containing an an acetoxy or benzoyloxy substituent at the propargylic position provided mono-, di-, and trisubstituted [3]cumulenes (Table 3).^{52–56} For example, 1,3-diynes 4a–4c afforded trisubstituted cumulenes 5a–5c , respectively, in 76–80% yields (entries 1–3, respectively). 1,3-Diynes substituted with a cycloalkyl moiety afforded the corresponding cumulenes 5d–5f in good yields (entries 4–6, respectively). Unexpectedly, while the acetate derivative of tertiary alcohol was afforded, the corresponding primary and secondary acetates provide a mixture of the expected [3]cumulenes and the corresponding acetoxy allene derivatives. However, upon replacement of the acetate with *p*-nitrobenzoate (4g–4i), only cumulenes 5g–5i were obtained (entries 7–9, respectively). We believe this is the consequence of the better leaving group capacity of a benzoate compared to that of an acetate. [3]Cumulenes make up a special class of polyene organic compounds whose synthetic utilities have been little explored.^{57–60} Thus, the current mild protocol to allow the preparation of [3]cumulenes containing various

Table 3. Synthesis of [3]Cumulenyl Imidamides via Eliminating the Acetoxy or Benzoyloxy Group of a Putative Allene Intermediate

Entry	1,3-Diene 4	[3]Cumulene 5	Yield (%)
1	1a, R = Me	5a	76
2	1b, R = Et	5b	79
3	1c, R = Ph	5c	80
4	1d, n = 1	5d	56
5	1e, n = 2	5e	82
6	1f, n = 3	5f	76
7	1g, R = H	5g	68
8	1h, R = Me	5h	74
9	1i, R = Ph	5i	77

substituent patterns from readily available building blocks is of highly synthetically useful.

We envision that with a suitably tethered nucleophile, the conversion of 1,3-dienes to the corresponding conjugated allenyl imidamide and imidates would promote an intramolecular Michael-type addition (Table 4).^{61–65} Under

Table 4. Synthesis of Heterocycles via Intramolecular Trapping of the Putative Allene Intermediate

Entry	1,3-Diene 6	Heterocycle 7	Yield (%)
1	2a, R = H	7a	48
2	2b, R = Me	7b	87
3	2c, n = 1	7c	72
4	2d, n = 2	7d	0
5	2e, X = O	7e	70
6	2f, X = NH	7f	64


Conditions: (A) TsN₃ (1.2 equiv), Et₃N (2 equiv), MeOH (10 equiv), 12 h; (B) TsN₃ (1.2 equiv), *i*-Pr₂NH (1.2 equiv), 4 h.

standard conditions, homopropargyl alcohol-containing 1,3-dienes **6a** and **6b** were smoothly converted to tetrahydrofuran-ylidene imidates **7a** and **7b** in 48% and 87% yields, respectively (entries 1 and 2, respectively). The corresponding homopropargyl sulfonamide **6c** led to 1-tosylpyrrolidinylidene imidamide **7c** in 72% yield (entry 3); however, the formation of a six-membered ring **7d** from **6d** failed (entry 4). 1,3-Dienes **6e** and **6f** substituted with a phenyl group containing an *o*-OH or NH₂ participated in the cascade reaction to generate benzofuranyl imidamide **7e** in 70% yield and indolyl imidamide **7f** in 64% yield (entries 5 and 6, respectively).

Although expected, migration of the triple bond from 3-alkynyl or 2,3-dienyl isomers to the corresponding 2-alkynyl

isomer **8** was not observed under the conditions regardless of the reaction time. However, because of the thermodynamic preference for 2-alkynyl isomers, we surmised that the isomerization of **2** or **3** to **8** would happen if a base stronger than secondary amines is used (Scheme 4).⁶⁶ Indeed, treating a

Scheme 4. Isomerization of 3-Alkynyl and 2,3-Dienyl Imidamides to the Corresponding 2-Alkynyl Isomers



^aDecomposition of starting materials. ^b3-Alkynyl imidamide **2** was recovered.

mixture of **2** and **3** with DBU (0.1 equiv) rapidly induced isomerization to provide 2-alkynyl imidamides **8a–8d**. On the contrary, phenyl- and silyl-substituted 3-alkynyl imidamides **2** and **3** did not isomerize to the corresponding 2-alkynyl isomers **8e** and **8f**, respectively.

In conclusion, we have developed efficient protocols to generate discrete isomers of 2-alkynyl, 3-alkynyl, 2,3-dienyl, and 2,3,4-trienyl imidamides and imidates from copper-catalyzed reactions of 1,3-dienes and tosylazide. The selectivity between 3-alkynyl and 2,3-dienyl imidamides could be controlled by a heteroatom substituent at the propargylic position of the 1,3-dienes and employing different trapping reagents such as amines and alcohols. [3]Cumulene derivatives were also generated by employing 1,3-dienes that contain an acetoxy or bezoyloxy substituent at the propargylic position. While trapping of the putative azacumulene intermediates with amines provided either 3-alkynyl or 2,3-dienyl imidamides depending on the structure of the trapping amines, trapping with methanol selectively generated 2,3-dienyl imidates. It was found that both 3-alkynyl and 2,3-dienyl imidamides could be isomerized to selectively generate the corresponding 2-alkynyl isomers under equilibrating conditions with a stronger base such as DBU. Intramolecular trapping of the putative azacumulene intermediates provided five-membered heterocyclic products if the 1,3-diene substrates contained a homopropargylic hydroxyl or amino substituent. A unique feature of these unprecedented reactions is that under mild reaction conditions, terminal 1,3-dienes could be selectively converted to different unsaturated carboxylic acid derivatives with good selectivity.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c03861>.

Experimental procedures and spectroscopic data for all new compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Institutions and Publishers

Daesung Lee – Department of Chemistry, University of Illinois at Chicago, Chicago, Illinois 60607, United States;

orcid.org/0000-0003-3958-5475; Email: dsunglee@uic.edu

Authors

Institutions and Publishers

Sourav Ghorai – Department of Chemistry, University of Illinois at Chicago, Chicago, Illinois 60607, United States

Institutions

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.orglett.0c03861>

Notes

Content

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

acknowledgements

The authors thank the National Science Foundation (CHE-1764141) for financial support. The Mass Spectrometry Laboratory at the University of Illinois at Chicago is acknowledged.

Headline

REFERENCES

References

- (1) Hoffmann-Röder, A.; Krause, N. *Angew. Chem., Int. Ed.* **2004**, *43*, 1966.
- (2) Krause, N.; Hoffmann-Röder, A. *Modern Allene Chemistry* **2004**, 997.
- (3) Dembitsky, V. M.; Maoka, T. *Prog. Lipid Res.* **2007**, *46*, 328.
- (4) Metcalf, B. W.; Wright, C. L.; Burkhardt, J. P.; Johnston, J. O. *J. Am. Chem. Soc.* **1981**, *103*, 3221.
- (5) Faraj, H.; Aumelas, A.; Claire, M.; Rondot, A.; Auzou, G. *Steroids* **1991**, *56*, 558.
- (6) Baret, P.; Barreiro, E.; Greene, A. E.; Luché, J. L.; Teixeira, M. A.; Crabbe, P. *Tetrahedron* **1979**, *35*, 2931.
- (7) Collins, P. W.; Djuric, S. W. *Chem. Rev.* **1993**, *93*, 1533.
- (8) Phadtare, S.; Zemlicka, J. *J. Am. Chem. Soc.* **1989**, *111*, 5925.
- (9) Casara, P.; Jund, K.; Bey, P. *Tetrahedron Lett.* **1984**, *25*, 1891.
- (10) Ban, H. S.; Onagi, S.; Uno, M.; Nabeyama, W.; Nakamura, H. *ChemMedChem* **2008**, *3*, 1094.
- (11) Maes, M. *Acc. Chem. Res.* **2009**, *42*, 1679.
- (12) Krause, N.; Winter, C. *Chem. Rev.* **2011**, *111*, 1994.
- (13) Ye, S.; Ma, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 3074.
- (14) Kitagaki, S.; Inagaki, F.; Mukai, C. *Chem. Soc. Rev.* **2014**, *43*, 2986.
- (15) Trus, M. A. *Chem. Soc. Rev.* **2014**, *43*, 2979.
- (16) Alcaide, B.; Almendros, P.; Aragoncillo, C. *Chem. Soc. Rev.* **2014**, *43*, 2106.
- (17) Ye, S.; Ma, S. *Acc. Chem. Res.* **2014**, *47*, 989.
- (18) Adams, C. S.; Weatherly, C. D.; Burke, E. G.; Schomaker, J. M. *Chem. Soc. Rev.* **2014**, *43*, 3136.
- (19) Alonso, J. M.; Quirós, M. T.; Muñoz, M. P. *Org. Chem. Front.* **2016**, *3*, 186.
- (20) Campolo, D.; Gastaldi, S.; Roussel, C.; Bertrand, M. P.; Neelab, M. *Chem. Soc. Rev.* **2013**, *42*, 8434.
- (21) Neff, R. K.; Frantz, D. E. *Tetrahedron* **2015**, *71*, 7.
- (22) Xu, Y.; Hong, Y. J.; Tantillo, D. J.; Brown, M. K. *Org. Lett.* **2017**, *19*, 3703.
- (23) Sydes, L. K. *Chem. Rev.* **2003**, *103*, 1133.
- (24) Krause, N.; Hoffmann-Röder, A. *Tetrahedron* **2004**, *60*, 11671.
- (25) Brummond, K. M.; DeForrest, J. E. *Synthesis* **2007**, *2007*, 795.
- (26) Ye, S.; Ma, S. *Chem. Commun.* **2011**, *47*, 5384.
- (27) Neff, R. K.; Frantz, D. E. *ACS Catal.* **2014**, *4*, 519.
- (28) Chu, W. D.; Zhang, Y.; Wang, J. *Catal. Sci. Technol.* **2017**, *7*, 4570.
- (29) Armstrong, R. J. *Curr. Org. Chem.* **2020**, *23*, 3027.

REFERENCES

- (30) Li, Q. H.; Jeng, J. Y.; Gau, H. M. *Eur. J. Org. Chem.* **2014**, *2014*, 7816.
- (31) Kang, J.; Tang, X.; Ma, S. *Org. Chem. Front.* **2015**, *2*, 470.
- (32) Lim, J.; Choi, J.; Kim, H. S.; Kim, I. S.; Nam, K. C.; Kim, J.; Lee, S. *Org. Chem.* **2016**, *81*, 303.
- (33) Boreux, A.; Lonca, G. H.; Riant, O.; Gagosz, F. *Org. Lett.* **2016**, *18*, 5162.
- (34) Ye, S.; Zhang, J. *Adv. Synth. Catal.* **2011**, *353*, 1265.
- (35) Huang, X.; Ma, S. *Acc. Chem. Res.* **2019**, *52*, 1301.
- (36) Mossain, M. L.; Wang, J. *Chem. Rec.* **2018**, *18*, 1548.
- (37) Lv, W.; Chen, Y.; Zhao, Z.; Wen, Si; Cheng, G. *Org. Lett.* **2019**, *21*, 7795.
- (38) Nella, N.; Parker, E.; Hitce, J.; Larini, P.; Jazsar, R.; Baudoin, O. *Chem. Rev.* **2014**, *20*, 13272.
- (39) Neff, R. K.; Frantz, D. E. *J. Am. Chem. Soc.* **2018**, *140*, 17428.
- (40) Tang, M.; Fan, C.-A.; Zhang, F.-M.; Tu, Y.-Q.; Zhang, W.-X.; Wang, A.-X. *Org. Lett.* **2008**, *10*, 5585.
- (41) Maekawa, H.; Takano, A.; Watanabe, M. *Tetrahedron Lett.* **2014**, *55*, 6208.
- (42) Suarez, A.; Fu, G. C. *Angew. Chem., Int. Ed.* **2004**, *43*, 3580.
- (43) Sabbasani, V. R.; Mamidipalli, P.; Lu, H.; Xia, Y.; Lee, D. *Org. Lett.* **2013**, *15*, 1552.
- (44) Hassink, M.; Liu, X.; Fox, J. M. *Org. Lett.* **2011**, *13*, 2388.
- (45) Kim, S. H.; Park, S. H.; Choi, J. H.; Chang, S. *Chem. - Asian J.* **2011**, *6*, 2618.
- (46) Yoo, E. J.; Ahlquist, M.; Kim, S. H.; Bae, I.; Fokin, V. V.; Sharpless, K. B.; Chang, S. *Angew. Chem., Int. Ed.* **2007**, *46*, 1730.
- (47) Bae, I.; Han, H.; Chang, S. *J. Am. Chem. Soc.* **2005**, *127*, 2038.
- (48) Yoo, E. J.; Bae, I.; Cho, S. H.; Han, H.; Chang, S. *Org. Lett.* **2006**, *8*, 1247.
- (49) Kim, S. H.; Jung, D. Y.; Chang, S. *J. Org. Chem.* **2007**, *72*, 9769.
- (50) Cai, S.-L.; Lin, X.-F.; Wang, Y.-G. *Org. Lett.* **2006**, *8*, 4517.
- (51) Husmann, R.; Na, Y. S.; Bolm, C.; Chang, S. *Chem. Commun.* **2010**, *46*, 5494.
- (52) Mizuno, K.; Maeda, H.; Sugita, H.; Nishioka, S.; Hirai, T.; Sugimoto, A. *Org. Lett.* **2001**, *3*, 581–584.
- (53) Leroyer, L.; Maraval, V.; Chauvin, R. *Chem. Rev.* **2012**, *112*, 1210.
- (54) Ueta, S.; Hida, K.; Nishiuchi, M.; Kawamura, Y. *Org. Biomol. Chem.* **2014**, *12*, 2784–2791.
- (55) Wendinger, D.; Tykewinski, R. R. *Acc. Chem. Res.* **2017**, *50*, 1468.
- (56) Ardila-Fierro, K. J.; Bolm, C.; Hernández, J. G. *Angew. Chem., Int. Ed.* **2019**, *58*, 12945–12949.
- (57) Guan, X.-Y.; Shi, M. *J. Org. Chem.* **2009**, *74*, 1977–1981.
- (58) Li, W.; Shi, M. *Org. Biomol. Chem.* **2009**, *7*, 1775–1777.
- (59) Wang, L. F.; Cao, X. P.; Shi, Z. F.; An, P.; Chow, H. F. *Adv. Synth. Catal.* **2014**, *356*, 3383–3390.
- (60) Gawel, P.; Dengiz, C.; Finke, A. D.; Trapp, N.; Boudon, C.; Gisselbrecht, J. P.; Diederich, F. *Angew. Chem., Int. Ed.* **2014**, *53*, 1311.
- (61) Kumaraswamy, G.; Jayaprakash, N.; Balakishan, G. *Org. Biomol. Chem.* **2011**, *9*, 7913–7920.
- (62) Zhou, L.; Shi, Y.; Xiao, Q.; Liu, Y.; Ye, F.; Zhang, Y.; Wang, J. *Org. Lett.* **2011**, *13*, 968–971.
- (63) Shin, Y. H.; Maheswara, M.; Hwang, J. Y.; Kang, E. J. *Eur. J. Org. Chem.* **2014**, *2014*, 2305–2311.
- (64) Zhang, Y. Y.; Hao, J.; Shi, M. *Org. Chem. Front.* **2015**, *2*, 394–397.
- (65) Li, Y.; Hirabayashi, S.; Yoshioka, S.; Aoyama, H.; Murai, K.; Fujino, H.; Arisawa, M. *Org. Lett.* **2019**, *21*, 3501–3504.
- (66) Spence, J. D.; Wyatt, J. K.; Bender, D. M.; Moss, D. K.; Nantz, M. H. *J. Org. Chem.* **1996**, *61*, 4014–4021.