

Too Close to Integrable: Crossover from Normal to Anomalous Heat Diffusion

Stefano Lepri,^{1,2} Roberto Livi,^{3,1,2} and Antonio Politì⁴

¹*Consiglio Nazionale delle Ricerche, Istituto dei Sistemi Complessi,
Via Madonna del Piano 10 I-50019 Sesto Fiorentino, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Firenze,
via G. Sansone 1 I-50019, Sesto Fiorentino, Italy*

³*Dipartimento di Fisica e Astronomia and CSDC, Università di Firenze,
via G. Sansone 1 I-50019, Sesto Fiorentino, Italy*

⁴*Institute for Complex Systems and Mathematical Biology & SUPA
University of Aberdeen, Aberdeen AB24 3UE, United Kingdom*

(Dated: April 15, 2020)

Energy transport in one-dimensional chains of particles with three conservation laws is generically anomalous and belongs to the Kardar-Parisi-Zhang dynamical universality class. Surprisingly, some examples where an apparent normal heat diffusion is found over a large range of length scales were reported. We propose a novel physical explanation of these intriguing observations. We develop a scaling analysis which explains how this may happen in the vicinity of an integrable limit, such as, but not only, the famous Toda model. In this limit, heat transport is mostly supplied by quasi-particles with a very large mean free path ℓ . Upon increasing the system size L , three different regimes can be observed: a ballistic one, an intermediate diffusive range, and, eventually, the crossover to the anomalous (hydrodynamic) regime. Our theoretical considerations are supported by numerical simulations of a gas of diatomic hard-point particles for almost equal masses and of a weakly perturbed Toda chain. Finally, we discuss the case of the perturbed harmonic chain, which exhibits a yet different scenario.

After more than twenty years of theoretical research there is a general consensus that energy transport in one and two-dimensional systems is anomalous, meaning that Fourier's law is invalid [1–3]. Numerics [4–6] as well as hydrodynamic [7, 8] and kinetic [9–11] theories consistently indicate that the nonlinear interactions of fluctuations of conserved quantities yield, in reduced space-dimension, non standard relaxation and transport properties, even in the linear response regime. The main signature of the anomaly is the divergence of the thermal conductivity κ with the system size L , i.e. a superdiffusive heat transport. In one-dimension, although this is a genuine many-body problem, it can be described effectively as an ensemble of Lévy particles, namely random walkers performing free ballistic steps with finite velocity for times that are power-law distributed [12]. This description accounts quantitatively for several non-equilibrium properties, both transient and stationary [6, 13–15]. Remarkably, the phenomenon was shown to belong to the class of the famous Kardar-Parisi-Zhang (KPZ) equation [16, 17], suggesting a universal behavior with implications for the theory of transport in nano-sized objects like individual nanowires [18], nanotubes [19] or polymers [20]. In this general context, nanowires and single-walled nanotubes have been analyzed to look for deviations from the standard Fourier's law [19]. Experimental evidence of such deviations has been reported for single-walled carbon nanotubes [19, 21] (see also Ref. [22]). Non-trivial length dependence of thermal conductance has been also observed in molecular chains [23]. Transport anomalies can be even exploited to achieve optimal efficiency of thermal to electric energy conversion [24, 25].

Although the general framework is pretty well understood, there are still open issues that escaped so far a convincing explanation. For definiteness, we focus on anharmonic chains, represented by a Hamiltonian of the form

$$H = \sum_{n=1}^L \left[\frac{p_n^2}{2m_n} + U(q_{n+1} - q_n) \right], \quad (1)$$

where m_n , q_n and p_n are respectively the mass, displacement and momentum of the n th particle. For a generic potential U , this family of models should show superdiffusive heat transport in the KPZ universality class, as confirmed by several studies [26, 27]. However, there is evidence of significant deviations of the dynamical exponents in some models with hard-core potential [28]. Moreover, chains allowing for bond dissociation (like e.g Lennard-Jones, Morse, and Coulomb potentials) unexpectedly display finite thermal conductivity [29, 30], while other similar potentials closely follow the prediction of anomalous scaling [31]. For the double-well potential, an intermediate-energy regime with almost diffusive transport has been reported [32, 33].

Another, more surprising feature is the (apparent) normal diffusive heat transport observed at low energies in asymmetric potentials [34] like the Fermi-Pasta-Ulam-Tsingou- $\alpha\beta$ (FPUT) chain, where $m_n = m$ and $U_{\alpha\beta}(y) = y^2/2 + \alpha y^3/3 + \beta y^4/4$. Yet, more compelling evidence of a seemingly normal transport has been found in a Toda lattice under the action of an additional conservative noise [35] and in the diatomic Hard Point Gas (HPG) [36]. In the first context, successive studies

demonstrated that the diffusive regime is a finite-size effect, whereby anomalous behavior is recovered for L large enough [37, 38] (see also the discussion based on mode-coupling arguments in Ref. [39]).

In this Letter we present a general explanation of the counter-intuitive normal transport, by tracing it back to the vicinity to an integrable limit. In this context, the kinetics is dominated by quasi-particles (those associated to the integrals of motion of the integrable regime, e.g., solitons) over a broad range of scales. In fact, this range extends beyond the *very large* quasi-particle mean free path, where a purely ballistic behavior is expected. More specifically, we proceed by first devising a finite-size scaling approach and then validating the theoretical arguments with the numerical analysis of a few test models.

Let the parameter ε denote the distance from the integrable limit ($\varepsilon = 0$ corresponding to the perfectly integrable regime). Building upon the intuition contained in Ref. [40], we propose that for small ε , the heat flux $J(L, \varepsilon)$ can be written as the sum of two terms,

$$J(L, \varepsilon) = J_N(L, \varepsilon) + J_A(L, \varepsilon). \quad (2)$$

J_A is the hydrodynamic contribution, arising from the mutual interaction among density, energy, and momentum fluctuations, and responsible for the asymptotic superdiffusion. For $L \rightarrow \infty$, $J_A \approx L^{\eta-1}$ with $\eta = 1/3$ in systems belonging to the KPZ class [16, 17], while $\eta = 1/2$ in some special cases, like models with symmetric interaction potentials (e.g., FPUT- β chain with potential $U_\beta(y) = y^2/2 + y^4/4$) and models subject to conservative noise (e.g., the noisy harmonic [40, 41] or nonlinear [35, 42] chains). J_N is a kinetic contribution, accounting for the energy transported by weakly interacting quasi-particles.

On the basis of standard kinetic arguments, J_N is expected to be a function of a single compound variable, the effective length $\xi = L/\ell$ expressed in units of the mean free path ℓ of the quasi particles,

$$J_N(L, \varepsilon) = j_N(L/\ell). \quad (3)$$

For $\xi \gg 1$, we expect $j_N(\xi) \propto 1/\xi$, meaning that the flux is the result of a standard diffusive process, while $j_N(0)$ is a finite value, meaning that the process is ballistic for system sizes smaller than the mean free path ($\xi \ll 1$). The entire $j_N(\xi)$ dependence is captured by the simple effective formula

$$j_N(\xi) = \frac{j_0}{r + \xi} \quad (4)$$

where r is a constant accounting for the boundary resistance [43] and j_0 is a suitable additional constant.

The vicinity to the integrable limit manifests itself as a divergence of the mean free path, which we account for by assuming $\ell \approx \varepsilon^{-\theta}$, where $\theta > 0$ is a system-dependent

exponent. As long as $J_A(L, \varepsilon)$ does not display any singularity for $\varepsilon \rightarrow 0$ (we return to this point in the final part of the Letter), we can neglect its dependence on ε (for $\varepsilon \ll 1$). Therefore, for large L , Eq. (2) can be rewritten as

$$J(L, \varepsilon) \approx \frac{c_N}{L\varepsilon^\theta} + \frac{c_A}{L^{1-\eta}}, \quad (5)$$

where c_N and c_A are two suitable parameters. Accordingly, the anomalous contribution prevails only above the crossover length $\ell_c \approx \varepsilon^{-\theta/\eta}$. For $L \leq \ell_c$, heat conduction is dominated by the j_N . In particular, within the range $[\ell = \varepsilon^{-\theta}, \ell_c]$ an apparent normal conductivity is expected, which is nothing but a finite size effect.

Now, we start the numerical analysis, focusing on models of the class (1). More specifically, we shall consider the HPG [44–46], and the Toda chain [47, 48] with interaction potential $U_T(y) = (e^{-y} + y - 1)$.

The HPG dynamics consists of successive collisions between neighboring particles according to the kinematic rules

$$\begin{aligned} u'_i &= \frac{m_i - m_{i+1}}{m_i + m_{i+1}}u_i + \frac{2m_{i+1}}{m_i + m_{i+1}}u_{i+1}, \\ u'_{i+1} &= \frac{2m_i}{m_i + m_{i+1}}u_i - \frac{m_i - m_{i+1}}{m_i + m_{i+1}}u_{i+1}, \end{aligned} \quad (6)$$

where $u_n = \dot{q}_n$ and the primed variables denote the values after the collision. Simulations are very efficient since they only require keeping track of the collisions [46].

For equal masses ($m_n = m$), both models are completely integrable: in the HPG, the constants of motion are the initial velocities, while in the Toda model the conserved actions are given functions of positions and momenta [49, 50]. Both models can be seen as gases of quasi-particles: *velocityons* for HPG [51] and *solitons* for Toda [47]. Transport is ballistic: $\kappa(L)$ is proportional to L and the energy-current correlation function does not decay to zero at large times [52, 53]. Note that, according to the classification of Ref. [54], the two models are noninteracting and interacting, respectively.

Here below, we consider two different ways of breaking integrability (i.e. to induce interactions among the quasi-particles): (i) different masses, such as a diatomic arrangement whereby $m_n = m_1 = \frac{M}{2}(1 - \delta)$ ($m_n = m_2 = \frac{M}{2}(1 + \delta)$) for odd (even) n [44–46]; (ii) a conservative noise through random collisions exchanging the momenta of neighboring particles at a given rate γ [35] (see also a related model in Ref. [55]). In the former (latter) case δ (γ) plays the role of the above mentioned closeness parameter ε . In either case, only three conservation laws survive, momentum, energy and stretch, yielding anomalous transport and dynamical scaling [56]. For large enough δ -values, there is overwhelming evidence that both the diatomic HPG and Toda [45] belong to the KPZ universality class. For the randomly perturbed Toda evidence of a diverging conductivity is solid, yielding $\gamma \approx 0.44$ for a large collision rate $\gamma = 1$ [35].




Figure 1. Nonequilibrium simulations: scaled thermal conductivity versus scaled chain lengths for (a) the HPG model $T_L = 6, T_R = 4, M = 1, \delta = 0.05, 0.07, 0.10, 0.14, 0.2$ from bottom to top; (b) the diatomic Toda model (data taken from Ref.[36], Fig.4) $\delta = 0.07, 0.10, 0.14, 0.22, 0.30, 0.50, 1.0$ from bottom to top and (c) the Toda model with conserving noise $\gamma = 0.005, 0.01, 0.02, 0.04, 0.08, 0.16$ from bottom to top.

Here, we are interested in the quasi-integrable regime, $\delta, \gamma \ll 1$. The first issue is the determination of the mean free path ℓ . In the HPG with a diatomic mass arrangement, ℓ corresponds to the average space travelled by a single velociton before the collisions induce a sizeable change of its original velocity. Given the mass arrangement, the collision types (m_1, m_2) and (m_2, m_1) alternate and it is thereby appropriate to look at velocity changes every second iterate ($u \rightarrow u' \rightarrow u''$). From Eq. (6)

$$u'' = (1-\delta)u' + \delta\chi_2 = (1-\delta^2)u + \delta(\chi_2 - \chi_1) + \delta^2\chi_1 \quad (7)$$

where $\chi_{1,2}$ denote the velocities of the quasi-particles encountered by the velociton u , hereby assumed to be uncorrelated Gaussian variables with zero average and variance $\langle \chi_{1,2}^2 \rangle = v^2$ (neglecting the mass difference between




Figure 2. Estimates of dependence of the normal component $J_N(L)$ of the energy flux, as defined by Eq. (3), versus the rescaled system length L . Filled circles are for the HPG model with mass parameter $\delta = 0.05$ and triangles for the Toda with random collisions $\gamma = 0.005$. The solid lines are best fit with the functions $6.18/(3.73 + \xi)$ and $1/(27.0 + 5.62\xi + 7.87\xi^{0.2})$ respectively.

the two particles). To leading order in δ , the map can be turned into the stochastic differential equation

$$\dot{u} = -\delta^2 u + \sqrt{2}\delta v \zeta,$$

where ζ is a unit-variance white noise, while time is measured in 2τ units, where τ is the average collision time. As a result, the distribution of u values diffuses, its variance growing initially as $D_u = 2\delta^2 v^2 t$, so that the time needed for D_u to be approximately equal to v^2 is $t \approx \tau/\delta^2$ (in physical time units) and the corresponding mean free path is $\ell \approx v\tau/\delta^2$. In other words, we expect $\theta = 2$.

Numerical results have been obtained by implementing the standard non-equilibrium procedure [1, 2]. Left and right edges are attached to Maxwellian heat baths at temperatures $T_L = 6$ and $T_R = 4$ ($\Delta T = T_L - T_R$) and the flux J determined from the average energy exchanged in the steady state. For the HPG we employ the thermal-wall method as detailed, e.g., in Ref. [36].

In Fig. 1(a) we plot the rescaled thermal conductivity $\delta^2 \kappa$ of the diatomic HPG, referred to the effective length $\xi = L\delta^2$ [57] (the various curves correspond to different δ values – δ decreases from top to bottom). There is clear evidence of a ballistic regime followed by a diffusive one, as accounted for by Eq. (3). For the smaller δ 's, Fourier-like transport persists up to the maximal available L , while a crossover to the anomalous regime is seen upon increasing δ (see the uppermost curve). Upon decreasing δ , the curves converge from above towards an asymptotic shape, $\kappa_N = j_N(\xi)\xi/\Delta T$. In fact, for $\delta \rightarrow 0$ and fixed ξ , L increases ($L = \xi/\delta^2$), so that the corresponding $J_A(L)$ contribution becomes increasingly negligible.

Then, j_N is estimated from the data for the smallest perturbation amplitude (since J_A is practically negligible over the explored length range). The simple formula (4)

proves remarkably accurate: see the solid upper curve in Fig. 2, to be compared with the circles, which represent the numerical HPG results.

As a second test, we consider the diatomic Toda model [36]. For large energy densities, most of its dynamical properties are basically equal to the HPG [58] and we thus expect again $\ell \approx \delta^{-2}$, i.e. $\theta = 2$. At lower energies, one should estimate the soliton scattering rates due to mass inhomogeneities; however there are no reason to expect a different scaling behavior. Indeed, the typical thermalization time is of order δ^{-2} in a wide energy range [59]. The conductivities taken from Ref. [36], are reported in Fig. 1(b) after a proper rescaling. The data collapse confirms the validity of our arguments.

Finally, we have considered the Toda model with conservative noise. In this case, it is natural to argue that the mean free path scales as the inverse of the collision rate, $\ell \sim \gamma^{-1}$. This intuition is confirmed by the data reported in Fig. 1(c), where we observe the same scenario as for the previous models, after setting $\theta = 1$. For small γ , j_N converges again towards a function which asymptotically decays as $1/\xi$. However, the heuristic formula (4) is not comparably accurate: it is necessary to add a correction term to reproduce the observed data as shown in Fig. 2 (see triangles vs. the corresponding solid curve).

So far we have tested the structure of the first addendum in Eq. (2), by studying a regime where the second contribution is negligible. What about the second addendum? Once j_N has been determined, one can proceed by estimating the anomalous component as $J_A(L) = J(L, \varepsilon) - j_N(L\varepsilon^\theta)$. The data in Fig. 3 indicates that J_A exhibits the expected anomalous scaling already for system sizes where the direct estimates are strongly affected by the diffusive component. For the HPG the fitted slope, about -0.66 , corresponds to $\eta = 0.33$, in excellent agreement with the KPZ prediction $\eta = 1/3$. For Toda we obtain $\eta = 0.52$, consistent with Ref. [35] and even closer to $\eta = 1/2$, the value rigorously proven for harmonic models with momentum-conserving noise [40–42]. We thus conclude that the measurements are in full agreement with the crossover scenario from diffusive to hydrodynamic behavior proposed above.

The same scenario is expected to emerge in the presence of generic momentum-conserving perturbations of the Toda chain, such as a potential of the form $\varepsilon W(q)$. In fact, in this case, it has been already noticed that the energy-flux correlation decays over a time scale inversely proportional to a power of ε [53]. A preliminary evidence of a diffusive regime for small ε was also given in Ref. [60].

More in general, we argue that the crossover from normal to anomalous regimes of the FPUT- $\alpha\beta$ model [37, 38] is fully accounted for by the above described scenario. Indeed, the FPUT- $\alpha\beta$ (at low enough energies) can be regarded as a perturbed Toda chain over very long time scales, on which the Toda actions are only weakly perturbed [61]. We surmise that other model potentials




Figure 3. The anomalous part of the energy flux versus the system length L calculated as $J_A = J(L) - j_N(L)$ with j_N as determined in the previous figure. HPG model with $\delta = 0.14$ (squares) and $\delta = 0.20$ (circles) and for Toda with conservative noise $\gamma = 0.04$ (triangles). The dashed lines are power-law fit on the largest sizes where scaling sets in (values given in the text).

should display the same phenomenology, if their form is “close enough” to U_T .

What can we say about the simplest model of a perturbed harmonic chain? This textbook case deserves a special consideration. Numerical analysis of the FPUT- β model at very low energy, i.e. below the strong stochasticity threshold, does not reveal any signature of an intermediate diffusive regime, but rather a direct crossover from ballistic to anomalous regimes [62]. More compelling evidence of the absence of a diffusive regime comes from the study of the harmonic chain with conservative noise [41] in the limit of vanishing noise, i.e. $\gamma \rightarrow 0$. It has been found analytically [40] and confirmed numerically [63] that $J_A(L, \varepsilon)$, exhibits a singular dependence for $\varepsilon \rightarrow 0$ (here $\gamma \rightarrow 0$) in the form of a divergence of the coefficient c_A in Eq. (5), $c_A \approx \gamma^{-1/2}$, which implies that J_A prevails over j_N across the entire range of system sizes.

Hence, the different behavior displayed by weakly perturbed harmonic oscillators can be traced back to a divergence of the anomalous component, which is itself a consequence of the nonlinearity of the dynamical equations. While this counter-intuitive phenomenon requires further clarifications, it does not however jeopardize the general scaling theory developed in this Letter, which provides a convincing explanation of the seemingly normal diffusion observed in several models of nonlinear chains.

SL and RL acknowledge partial support from project MIUR-PRIN2017 *Coarse-grained description for non-equilibrium systems and transport phenomena (CO-NEST)* n. 201798CZL and the Eurofusion Enabling Research ENR-MFE19.CEA-06 *Emissive divertor*. SL is grateful to S. Ruffo and the program *Collaborazioni di Eccellenza* of SISSA, Trieste, Italy.

-
- [1] S. Lepri, R. Livi, and A. Politi, Phys. Rep. **377**, 1 (2003).
- [2] A. Dhar, Adv. Phys. **57**, 457 (2008).
- [3] S. Lepri, ed., *Thermal transport in low dimensions: from statistical physics to nanoscale heat transfer*, Lect. Notes Phys., Vol. 921 (Springer-Verlag, Berlin Heidelberg, 2016).
- [4] S. Lepri, R. Livi, and A. Politi, Phys. Rev. E **68**, 067102 (2003).
- [5] L. Wang and T. Wang, EPL (Europhysics Letters) **93**, 54002 (2011).
- [6] S. Liu, P. Hänggi, N. Li, J. Ren, and B. Li, Phys. Rev. Lett. **112**, 040601 (2014).
- [7] O. Narayan and S. Ramaswamy, Phys. Rev. Lett. **89**, 200601 (2002).
- [8] L. Delfini, S. Lepri, R. Livi, and A. Politi, Phys. Rev. E **73**, 060201(R) (2006).
- [9] A. Pereverzev, Phys. Rev. E **68**, 056124 (2003).
- [10] B. Nickel, J. Phys. A-Math. Gen. **40**, 1219 (2007).
- [11] J. Lukkarinen and H. Spohn, Communications on Pure and Applied Mathematics **61**, 1753 (2008).
- [12] V. Zaburdaev, S. Denisov, and J. Klafter, Rev. Mod. Phys. **87**, 483 (2015).
- [13] P. Cipriani, S. Denisov, and A. Politi, Phys. Rev. Lett. **94**, 244301 (2005).
- [14] S. Lepri and A. Politi, Phys. Rev. E **83**, 030107 (2011).
- [15] A. Dhar, K. Saito, and B. Derrida, Phys. Rev. E **87**, 010103 (2013).
- [16] H. van Beijeren, Phys. Rev. Lett. **108**, 180601 (2012).
- [17] H. Spohn, J. Stat. Phys. **154**, 1191 (2014).
- [18] M. Upadhyaya and Z. Aksamija, Phys. Rev. B **94**, 174303 (2016).
- [19] C.-W. Chang, in *Thermal Transport in Low Dimensions: From Statistical Physics to Nanoscale Heat Transfer*, edited by S. Lepri (Springer International Publishing, 2016) pp. 305–338.
- [20] A. Crnjar, C. Melis, and L. Colombo, Phys. Rev. Materials **2**, 015603 (2018).
- [21] V. Lee, C.-H. Wu, Z.-X. Lou, W.-L. Lee, and C.-W. Chang, Phys. Rev. Lett. **118**, 135901 (2017).
- [22] Q.-Y. Li, K. Takahashi, and X. Zhang, Phys. Rev. Lett. **119**, 179601 (2017).
- [23] T. Meier, F. Menges, P. Nirmalraj, H. Hölscher, H. Riel, and B. Gotsmann, Phys. Rev. Lett. **113**, 060801 (2014).
- [24] G. Casati, L. Wang, and T. Prosen, J. Stat. Mech.: Theory and Experiment, L03004 (2009).
- [25] G. Benenti, G. Casati, K. Saito, and R. S. Whitney, Physics Reports **694**, 1 (2017).
- [26] C. B. Mendl and H. Spohn, Phys. Rev. Lett. **111**, 230601 (2013).
- [27] S. G. Das, A. Dhar, K. Saito, C. B. Mendl, and H. Spohn, Phys. Rev. E **90**, 012124 (2014).
- [28] P. I. Hurtado and P. L. Garrido, Scientific reports **6**, 38823 (2016).
- [29] A. V. Savin and Y. A. Kosevich, Physical Review E **89**, 032102 (2014).
- [30] O. Gendelman and A. Savin, EPL (Europhysics Letters) **106**, 34004 (2014).
- [31] C. B. Mendl and H. Spohn, Phys. Rev. E **90**, 012147 (2014).
- [32] D. Xiong, J. Stat. Mech.: Theory and Experiment **2016**, 043208 (2016).
- [33] G. R. Archana and D. Barik, Phys. Rev. E **99**, 022103 (2019).
- [34] Y. Zhong, Y. Zhang, J. Wang, and H. Zhao, Phys. Rev. E **85**, 060102 (2012).
- [35] A. Iacobucci, F. Legoll, S. Olla, and G. Stoltz, J. Stat. Phys. **140**, 336 (2010).
- [36] S. Chen, J. Wang, G. Casati, and G. Benenti, Phys. Rev. E **90**, 032134 (2014).
- [37] L. Wang, B. Hu, and B. Li, Phys. Rev. E **88**, 052112 (2013).
- [38] S. Das, A. Dhar, and O. Narayan, J. Stat. Phys.; **154**, 204 (2014).
- [39] G. R. Lee-Dadswell, Phys. Rev. E **91**, 012138 (2015).
- [40] S. Lepri, C. Mejía-Monasterio, and A. Politi, J. Phys. A: Math. Theor. **42**, 025001 (2009).
- [41] G. Basile, C. Bernardin, and S. Olla, Phys. Rev. Lett. **96**, 204303 (2006).
- [42] G. Basile, L. Delfini, S. Lepri, R. Livi, S. Olla, and A. Politi, Eur. Phys. J.-Special Topics **151**, 85 (2007).
- [43] K. Aoki and D. Kusnezov, Phys. Rev. Lett. **86**, 4029 (2001).
- [44] G. Casati, Found. Phys. **16**, 51 (1986).
- [45] T. Hatano, Phys. Rev. E **59**, R1 (1999).
- [46] P. Grassberger, W. Nadler, and L. Yang, Phys. Rev. Lett. **89**, 180601 (2002).
- [47] M. Toda, *Theory of nonlinear lattices*, Vol. 20 (Springer Science & Business Media, 2012).
- [48] M. Toda, Phys. Scr. **20**, 424 (1979).
- [49] H. Flaschka, Physical Review B **9**, 1924 (1974).
- [50] M. Hénon, Phys. Rev. B **9**, 1921 (1974).
- [51] A. Politi, J. Stat. Mech.: Theory and Experiment, P03028 (2011).
- [52] X. Zotos, J. Low. Temp. Phys. **126**, 1185 (2002).
- [53] B. S. Shastry and A. Young, Phys. Rev. B **82**, 104306 (2010).
- [54] H. Spohn, Journal of Mathematical Physics **59**, 091402 (2018), <https://doi.org/10.1063/1.5018624>.
- [55] C. Bernardin and P. Gonçalves, Communications in Mathematical Physics **325**, 291 (2014).
- [56] Perturbations breaking momentum conservation (e.g. an external pinning potential) induce normal heat diffusion in the Toda model, although they may also display strong finite-size effects depending on the type of potential [60, 64].
- [57] Here and in the following, we take into account only the scaling of ℓ with ε , neglecting irrelevant multiplicative factors.
- [58] A. Kundu and A. Dhar, Phys. Rev. E **94**, 062130 (2016).
- [59] W. Fu, Y. Zhang, and H. Zhao, Phys. Rev. E **100**, 052102 (2019).
- [60] P. Di Cintio, S. Iubini, S. Lepri, and R. Livi, Chaos, Solitons & Fractals **117**, 249 (2018).
- [61] G. Benetti, H. Christodoulidi, and A. Ponomarev, Journal of Statistical Physics **152**, 195 (2013).
- [62] S. Lepri, R. Livi, and A. Politi, CHAOS **15**, 015118 (2005).
- [63] L. Delfini, S. Lepri, R. Livi, C. Mejía-Monasterio, and A. Politi, J. Phys. A: Math. Theor. **43**, 145001 (2010).
- [64] A. Dhar, A. Kundu, J. L. Lebowitz, and J. A. Scaramazza, Journal of Statistical Physics **175**, 1298 (2019).