## Kwame Nkrumah University of Science and Technology




### **MATH 166: Statistics and Probability**

Dr. Eric Nimako Aidoo

en.aidoo@yahoo.com 19-01-2018

#### Nature of Statistics



#### Uses of Statistics

Statistical techniques are used extensively by marketing, accounting, quality control, consumers, professional sports people, hospital administrators, educators, politicians, physicians, and many others.



Who Uses Statistics?

### **Descriptive Statistics**: Methods of organizing, summarizing, and presenting data in an informative way.

**EXAMPLE 1: A** Gallup poll found that 49% of the people in a survey knew the name of the first book of the Bible. The statistic 49 describes the number out of every 100 persons who knew the answer.

**EXAMPLE 2: According** to Consumer Reports, **General Electric washing** machine owners reported 9 problems per 100 machines during 2001. The statistic 9 describes the number of problems out of every 100 machines.

**Branches of Statistics** 

Inferential Statistics: A decision, estimate, prediction, or generalization about a population, based on a sample.

A Population is a Collection of all possible individuals, objects, or measurements of interest.



A **Sample** is a portion, or part, of the population of interest

**Branches of Statistics** 

Example 1: TV networks constantly monitor the popularity of their programs by hiring Nielsen and other organizations to sample the preferences of TV viewers.



Example 2: Wine tasters sip a few drops of wine to make a decision with respect to all the wine waiting to be released for sale.

Example 3: The accounting department of a large firm will select a sample of the invoices to check for accuracy for all the invoices of the company.

Branches of Statistics (examples of inferential statistics)

### Qualitative or Attribute Variable the

characteristic being studied is nonnumeric.





Type of car












Types of Variables

# In a **Quantitative Variable** information is reported numerically.

Balance in your checking account



Minutes remaining in class



Number of children in a family



### Quantitative variables can be classified as either **Discrete** or **Continuous**.

Discrete Variables: can only assume certain values and there are usually "gaps" between values.

Example: the number of bedrooms in a house, or the number of hammers sold at the local Home Depot (1,2,3,...,etc).




# A Continuous Variable can assume any value within a specified range.

### The pressure in a tire



The weight of a pork chop





The height of students in a class.

There are four levels of data

Nominal

Ordinal


Interval

Ratio

Levels of Measurement

Nominal level
Data that is
classified into
categories and
cannot be arranged
in any particular
order.







Nominal data

**Ordinal level**: involves data arranged in some order, but the differences between data values cannot be determined or are meaningless.


During a taste test of 4 soft drinks, Coca Cola was ranked number 1, Dr. Pepper number 2, Pepsi number 3, and Root Beer number 4.



Levels of Measurement

### **Interval level**

Similar to the ordinal level, with the additional property that meaningful amounts of differences between data values can be determined. There is no natural zero point.



Temperature on the Fahrenheit scale.

Ratio level: the interval level with an inherent zero starting point. Differences and ratios are meaningful for this level of measurement.

Miles traveled by sales Monthly income representative in a month of surgeons

Levels of Measurement