Technical supplement

Retrospective reconstruction and numerical qualification of historical CMJ recovery research

Owner-review supplement

6 November 2026

This supplement records the detailed formulation, architecture, route-level qualification, synthetic result, uncertainty, and analysis-contract evidence summarized in the main article. Public evidence files are copied from the v1.1.0 release-linked record with their original source hashes in the source manifest. The original neural checkpoints, fitted values, and restricted records are not distributed here. Table values below are rounded for display; the accompanying CSV files retain the full stored precision.

1 Historical formulation identities and ancestry

The eight formulations are distinct identities. A parent is included only when the lineage register records one. "Unresolved" is not a parent edge. Formulation status is descriptive and does not rank scientific validity.

Table 1. Complete formulation matrix from the recorded lineage and methods register. A missing direct parent is shown as unresolved, not inferred.

Identity	Status / parent	Response / observation	Horizon / task	Limitation
`Initial preseason`	SUPERSEDED; parent: Unresolved	event time adaptation recovery; camp history	Force and impulse innovations at H72 / D7	Superseded sample/surface; direct parent is unresolved; task and world law match the canonical state.
`Canonical preseason`	SUPERSEDED; parent: Initial preseason	event time adaptation recovery; camp history	Force and impulse innovations at H72 / D7	Canonical query surface and sample; not a new task or world law.
`Rich-history camp`	RETIRED; parent: Unresolved	fitness fatigue impulse response; rich monitoring	Force and impulse innovations at H72 / D7	Retired formulation; its direct scientific parent is unresolved.
`Preliminary post-exposure`	SUPERSEDED; parent: Unresolved	biexponential episode response; episode summary	Force and net-impulse innovations at H24 / H48 / H72	Direct parent is unresolved; preliminary measurement contract without a frozen force–impulse identity.
`Phase-consistent`	SUPERSEDED; parent: Preliminary post-exposure	biexponential episode response; phase consistent force impulse	Force and net-impulse innovations at H24 / H48 / H72	Observation-law repair; the episode response law is retained.
`Correlated exposure`	SUPERSEDED; parent: Phase-consistent	correlated exposure response; phase consistent force impulse	Force and net-impulse innovations at H24 / H48 / H72	Material correlated exposure-distribution change; later redesigned.
`Threshold response`	PROPOSED; parent: Correlated exposure	threshold response; phase consistent force impulse	Force and net-impulse innovations at H24 / H48 / H72	Proposed alternate, not an accepted successor formulation.
`Fixed-mode discrepancy`	ACTIVE CANDIDATE; parent: Correlated exposure	fixed mode discrepancy response; phase consistent force impulse	Force and net-impulse innovations at H24 / H48 / H72	Active candidate; fixed modes are mathematical anchors, not biological compartments.

2 Original neural architecture and input contracts

The architecture inventory is separate from the eight scientific formulations. Each family has two repeats, r0 and r1. Parameter totals are copied from the sanitized architecture qualification table; no private weights are included.

Table 2. Sanitized architecture and inference contract by family. Repeat-specific private weights and hashes are excluded.

Model / repeats	Inputs	Architecture	Parameters
`M0 (r0/r1)`	39 raw features	39 → 32 → 2; exact GELU	1,346
`M1 (r0/r1)`	39 raw features	39 → 64 → 2; exact GELU	2,690
`M2 (r0/r1)`	39 raw features	39 → 64 → 64 → 2; exact GELU	6,850
`M3 (r0/r1)`	39 raw + 4 × 4 history	Raw backbone + pointwise history encoder; mean/latest pooling → 128 → 2	7,138
`M4 (r0/r1)`	39 raw + 4 × 4 history	Raw backbone + 4 → 32 GRU; mean/latest pooling → 128 → 2	10,626
`M5 (r0/r1)`	39 raw + 4 × 4 history + horizon	M4 + 2 × 8 horizon embedding → 136 → 2	10,658
`M6 (r0/r1)`	39 raw + 4 × 4 history + horizon + 5 basis values	M5 + first two basis values → 16 → 152 → 2	10,738
`M7 (r0/r1)`	39 raw + 4 × 4 history + horizon + 5 basis values	M5 + all five basis values → 16 → 152 → 2	10,786
`M8 (r0/r1)`	39 raw + 4 × 4 history + horizon + 5 basis values	M7 + bias-free 5 → 2 linear basis head	10,796

3 Route-by-route numerical qualification

The stored saved-CUDA rule is $|a_i - b_i| \le 10^{-5} + 10^{-4}|b_i|$ for every compared element. The CPU-source comparison separately requires exact array equality. The reported maximum absolute discrepancy combines two outputs with different physical units and is not a comparable physical effect size. "Not tabulated" is retained when the source record did not report a failing-element count for a passing route.

Table 3. All 18 sanitized model-repeat records. The recorded inference-rerun flag is false for every route. The maximum absolute error pools force and impulse outputs with different units and is descriptive only.

Route	CPU-source	Saved CUDA	ATOL/RTOL	Max abs. error	Failing elements
`M0-r0`	Exact	Pass	1e-05 / 1e-04	1.43051147e-06	0
`M0-r1`	Exact	Pass	1e-05 / 1e-04	1.43051147e-06	0
`M1-r0`	Exact	Pass	1e-05 / 1e-04	1.43051147e-06	not tabulated
`M1-r1`	Exact	Pass	1e-05 / 1e-04	1.43051147e-06	not tabulated
`M2-r0`	Exact	Pass	1e-05 / 1e-04	1.90734863e-06	not tabulated
`M2-r1`	Exact	Pass	1e-05 / 1e-04	1.90734863e-06	not tabulated
`M3-r0`	Exact	Pass	1e-05 / 1e-04	2.38418579e-06	not tabulated
`M3-r1`	Exact	Pass	1e-05 / 1e-04	1.19209290e-06	not tabulated
`M4-r0`	Exact	FAIL	1e-05 / 1e-04	4.75883484e-04	254
`M4-r1`	Exact	FAIL	1e-05 / 1e-04	3.26395035e-04	146
`M5-r0`	Exact	FAIL	1e-05 / 1e-04	3.71456146e-04	443
`M5-r1`	Exact	FAIL	1e-05 / 1e-04	2.70366669e-04	206
`M6-r0`	Exact	FAIL	1e-05 / 1e-04	1.99317932e-04	88
`M6-r1`	Exact	FAIL	1e-05 / 1e-04	2.79307365e-04	140
`M7-r0`	Exact	FAIL	1e-05 / 1e-04	3.17573547e-04	163
`M7-r1`	Exact	FAIL	1e-05 / 1e-04	3.04937363e-04	198
`M8-r0`	Exact	FAIL	1e-05 / 1e-04	2.21252441e-04	98
`M8-r1`	Exact	FAIL	1e-05 / 1e-04	3.04222107e-04	145

All 18 routes have recorded exact CPU-source equality. The eight M0–M3 saved-CUDA comparisons pass and the ten M4–M8 comparisons fail. These are preserved qualification results, not a fresh model inference or GPU replay.

4 Full public synthetic six-cell results

Each row shows a stored cellwise NRMSE, which divides RMSE by that cell's population standard deviation (denominator convention n_c, not $n_c - 1$). Root A, B, and C are three caller-supplied public evaluation roots from the same generator. The mean and sample standard deviation summarize those three roots; they are not a confidence interval or human-population estimate.

Table 4. Stored cellwise NRMSE by public evaluation root. Values are rounded to 6 decimal places for display; `six_cell_results.csv` retains the full stored values.

Model	Outcome-horizon cell	Root A	Root B	Root C	Mean	Sample SD
Zero innovation	H24 / net impulse (m/s)	2.757823	2.759502	2.767942	2.761756	0.005423
Training mean	H24 / net impulse (m/s)	1.053386	1.036381	1.045551	1.045106	0.008512

Continued on next page

Model	Outcome-horizon cell	Root A	Root B	Root C	Mean	Sample SD
Ridge	H24 / net impulse (m/s)	0.651329	0.651082	0.654060	0.652157	0.001653
Structured	H24 / net impulse (m/s)	0.694226	0.698253	0.695487	0.695989	0.002060
Zero innovation	H24 / force innovation (N/kg)	3.043088	3.049200	3.008360	3.033549	0.022028
Training mean	H24 / force innovation (N/kg)	1.054776	1.040758	1.047428	1.047654	0.007012
Ridge	H24 / force innovation (N/kg)	0.630319	0.630139	0.612765	0.624408	0.010083
Structured	H24 / force innovation (N/kg)	0.663259	0.664902	0.644444	0.657535	0.011367
Zero innovation	H48 / net impulse (m/s)	2.292457	2.222562	2.278043	2.264354	0.036904
Training mean	H48 / net impulse (m/s)	1.035905	1.022505	1.036852	1.031754	0.008024
Ridge	H48 / net impulse (m/s)	0.752218	0.765857	0.761406	0.759827	0.006955
Structured	H48 / net impulse (m/s)	0.797196	0.812503	0.804108	0.804603	0.007666
Zero innovation	H48 / force innovation (N/kg)	2.525989	2.550929	2.549185	2.542034	0.013923
Training mean	H48 / force innovation (N/kg)	1.044715	1.028447	1.045208	1.039457	0.009538
Ridge	H48 / force innovation (N/kg)	0.737124	0.737851	0.720804	0.731926	0.009639
Structured	H48 / force innovation (N/kg)	0.767842	0.768971	0.746271	0.761028	0.012792
Zero innovation	H72 / net impulse (m/s)	1.882918	1.844269	1.892503	1.873230	0.025535
Training mean	H72 / net impulse (m/s)	1.028582	1.018947	1.026964	1.024831	0.005159
Ridge	H72 / net impulse (m/s)	0.834328	0.829607	0.836870	0.833602	0.003686
Structured	H72 / net impulse (m/s)	0.877276	0.876384	0.879950	0.877870	0.001856
Zero innovation	H72 / force innovation (N/kg)	2.110591	2.101190	2.115307	2.109029	0.007187
Training mean	H72 / force innovation (N/kg)	1.032853	1.019695	1.029735	1.027428	0.006875
Ridge	H72 / force innovation (N/kg)	0.814940	0.821730	0.810435	0.815702	0.005686
Structured	H72 / force innovation (N/kg)	0.843228	0.850143	0.835237	0.842869	0.007459

5 Recorded camp-bootstrap intervals

The stored percentile intervals use 5,000 paired bootstrap replicates per root. The resampling unit is camp: 16 camps are sampled with replacement and the 250 episodes and three horizons for each selected camp remain together. The paired contrast is model minus zero innovation; negative values favor the named model over that baseline. The receipt did not record a ridge-versus-structured interval.

Table 5. Recorded 95% camp-bootstrap percentile endpoints by root. Delta is model minus zero innovation. No ridge-versus-structured interval is recorded.

Root	Model	NRMSE low	NRMSE high	Delta low	Delta high	Replicates
A	`ridge_full`	0.728033	0.746188	-1.752167	-1.644956	5000
A	`structured_full`	0.763766	0.784347	-1.714653	-1.608125	5000
B	`ridge_full`	0.733825	0.745350	-1.715574	-1.648424	5000
B	`structured_full`	0.773494	0.784623	-1.676622	-1.609237	5000
C	`ridge_full`	0.726382	0.740127	-1.733792	-1.669113	5000
C	`structured_full`	0.761200	0.775063	-1.700474	-1.632528	5000

6 Ablation and nested-camp result detail

The ablation variants are distinct receipt-locked model configurations. The structured-without-history variant is a separate comparator diagnostic. The nested-camp analysis uses 24, 48, and 96 training camps from a single recorded seeded ordering; the alpha selected on full training/development data is held fixed. Root values are descriptive and do not identify causal feature effects.

Table 6. Recorded six-cell NRMSE root values for selected ablation and nested-camp comparisons. Values are rounded to 6 decimal places; source receipt retains full precision.

Variant	Training camps	Evaluation-root NRMSE	Root mean ± sample SD
Full ridge	96	A 0.736710; B 0.739378; C 0.732724	0.736270 ± 0.003349
Ridge without prior history	96	A 0.822457; B 0.823679; C 0.812906	0.819681 ± 0.005899
Ridge without current exposure	96	A 0.868359; B 0.861227; C 0.873620	0.867735 ± 0.006220
Ridge without participant context	96	A 0.736746; B 0.739455; C 0.732837	0.736346 ± 0.003327
Structured without prior history	96	A 0.823035; B 0.824241; C 0.813749	0.820342 ± 0.005741
Ridge, nested 24 camps	24	A 0.736637; B 0.740818; C 0.733212	0.736889 ± 0.003809
Ridge, nested 48 camps	48	A 0.736861; B 0.739726; C 0.733219	0.736602 ± 0.003261
Ridge, nested 96 camps	96	A 0.736710; B 0.739378; C 0.732724	0.736270 ± 0.003349
Structured, nested 24 camps	24	A 0.774179; B 0.778755; C 0.767939	0.773624 ± 0.005429
Structured, nested 48 camps	48	A 0.774324; B 0.778711; C 0.767828	0.773621 ± 0.005475

Continued on next page

Variant	Training camps	Evaluation-root NRMSE	Root mean ± sample SD
Structured, nested 96 camps	96	A 0.773838; B 0.778526; C 0.767583	0.773316 ± 0.005490

7 Public synthetic study protocol

Protocol item	Recorded value
Training split	96 camps; 24,000 episodes; 72,000 horizon rows.
Development split	16 camps; 4,000 episodes; 12,000 horizon rows.
Each evaluation root	16 camps; 4,000 episodes; 12,000 horizon rows; 4,000 rows per outcome-horizon cell.
Generator relation	Three caller-supplied public roots from one fixed-mode synthetic generator; no human data or private historical test root.
Predictor schema	82 public source fields; one-hot horizon expansion to 84 numeric features; grouping keys, target labels, target-time realizations, hidden effects, and oracle states excluded.
Preprocessing	Training-only means and population standard deviations; zero scales replaced by one.
Models	Zero innovation; training-only outcome-by-horizon mean; cellwise ridge; structured comparator using observed predictors and fixed 24 h/84 h mathematical modes.
Alpha selection	Grid $\{0.01, 0.1, 1, 10, 100\}$; minimum development six-cell mean NRMSE; lower alpha breaks ties; all full-data selections choose 0.01.
Registered cells	H24/H48/H72 crossed with the registered force-innovation cell and net-impulse innovation.
Primary aggregate	Equal-weight mean of six cellwise NRMSE values; each cell is normalized separately.
Bootstrap	PCG64; 5,000 paired whole-camp resamples; 16 camps per root; seed 49,505,000; all episode and horizon records stay with the camp.
Sample-size analysis	24, 48, 96 camps from one seeded nested permutation; alpha chosen on the full dataset held fixed; seed 52,610,526.

8 Native measurement and evaluation contract

For the native task, force is reported in $\mathrm{N\,kg^{-1}}$ and gravity-adjusted net impulse in $\mathrm{m\,s^{-1}}$. With a supplied force trace $F_z(t)$, system mass m, gravity g, concentric onset t_{co}, takeoff t_{to}, and duration $T = t_{\mathrm{to}} - t_{\mathrm{co}}$:

$$Y_F = \frac{1}{mT} \int_{t_{\mathrm{co}}}^{t_{\mathrm{to}}} F_z(t)\,\mathrm{d}t, \tag{1}$$

$$Y_J = \frac{1}{m} \int_{t_{\mathrm{co}}}^{t_{\mathrm{to}}} \left(F_z(t) - mg\right)\,\mathrm{d}t. \tag{2}$$

The implementation interpolates force at event boundaries and uses composite trapezoids. It does not detect events or authenticate calibration, mass, or trial validity. An assessment is the arithmetic mean of exactly three distinct valid trials. Baseline is the mean of the two latest distinct valid assessments in the inclusive 24–240 h pre-origin interval. Native target windows are H72 at 66–78 h and D7 at 156–180 h.

The four native analysis cells are force and impulse at H72 and D7. Within each cell, query-weighted RMSE, mean absolute error (MAE), and signed bias are computed separately. Repeats are reported separately; a family metric is the arithmetic mean of repeat metrics, never an ensemble of predictions. Uncertainty resamples participant groups, preserving all participant queries, origins, horizons, candidates, repeats, and baselines. The 95% percentile intervals are marginal.

The native primary sign-flip test uses participant sums of zero-baseline squared error minus mean two-repeat squared error. The one-sided null requires independent participant groups and symmetry of cluster loss differences. At $n \le 16$, all 2^n sign assignments are enumerated; above that threshold, sampled assignments use a plus-one Monte Carlo correction. Holm adjustment applies to four locked primary tests. No athlete-level result from this analysis was generated for this manuscript. The stored six-participant engineering rehearsal uses invented values only; its smallest exact p value is $1/64$, and the four-cell Holm-adjusted value is $4/64 = 0.0625$, so it cannot pass a 0.05 family-wise threshold.

9 Evidence and interpretation boundaries

- The public scientific software identity is release v1.1.0 at commit `401f87aad1e4ef4108011fa0c1d769a24f1ad329`.
- The M8 synthetic-study result and reproduction receipts keep their own package-source, protocol, selection-lock, and result digests. They are not relabeled as a new run of the original neural model campaign.
- Private checkpoint bytes, fitted preprocessing values, and restricted records are excluded. The public-safe route summary supports figure regeneration but not independent replay of those artifacts.
- No new training, fit, checkpoint inference, empirical scoring, athlete-data acquisition, or GPU replay was performed for this manuscript.
- Historical saved-CUDA failures for M4–M8 and rich-history row-count discrepancies remain negative findings. No cause for the CUDA discrepancy is asserted.
- No independent athlete predictive validity, clinical utility, out-of-distribution transportability, or production readiness is established.