

# Tactical Combat Casualty Care and Wilderness Medicine



## Advancing Trauma Care in Austere Environments

Frank K. Butler, MD<sup>a,\*</sup>, Brad Bennett, PhD<sup>b</sup>,  
Colonel Ian Wedmore, MD<sup>c</sup>

### KEYWORDS

- Tactical combat casualty care • TCCC • Wilderness medicine
- Battlefield trauma care • Tourniquets • Hemostatic dressings

### KEY POINTS

- The Joint Trauma System helped the US military achieve unprecedented success in casualty survival during the wars in Afghanistan and Iraq.
- Tactical Combat Casualty Care (TCCC) is the prehospital component of the Joint Trauma System.
- Since most combat fatalities occur before the casualty reaches the care of a surgeon, TCCC plays a key role in ensuring that casualties have a maximal chance of survival.
- The realization that extremity hemorrhage, a leading cause of preventable death on the battlefield, could be effectively and safely addressed with limb tourniquets was the primary driving factor for the Special Operations medical research effort that gave rise to TCCC.
- Organizations advocating for the translation of TCCC concepts to the civilian sector include the Wilderness Medical Society, the National Association of Emergency Medical Technicians, the American College of Surgeons' Hartford Consensus working group, the White House Stop the Bleed campaign, and the Committee on Tactical Emergency Casualty Care.

---

**Disclaimers:** The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

**Release:** This document was reviewed by the Public Affairs Office and the Operational Security Office at the US Army Institute of Surgical Research. It is approved for unlimited public release.

**Disclosures:** The authors have no disclosures.

<sup>a</sup> Committee on Tactical Combat Casualty Care, Joint Trauma System, US Army Institute of Surgical Research, 3698 Chambers Pass, JBSA Fort Sam Houston, TX 78234-6315, USA;

<sup>b</sup> Uniformed Services University of the Health Sciences, 4301 Jones Bridge Rd, Bethesda, MD 20814, USA; <sup>c</sup> Madigan Army Medical Center, 9040 Jackson Ave., Tacoma, WA 98431, USA

\* Corresponding author.

*E-mail address:* [fk064@yahoo.com](mailto:fk064@yahoo.com)

Emerg Med Clin N Am 35 (2017) 391–407

<http://dx.doi.org/10.1016/j.emc.2016.12.005>

0733-8627/17/Published by Elsevier Inc.

[emed.theclinics.com](http://emed.theclinics.com)

## INTRODUCTION

In 1992, the Naval Special Warfare Biomedical Research and Development program undertook a review of battlefield trauma care. The primary driver for this research effort was the realization that, although extremity hemorrhage was a leading cause of preventable death in combat casualties<sup>1,2</sup> and tourniquets could be applied safely for short periods, tourniquet use was universally disparaged in both civilian and military prehospital trauma care.<sup>3-5</sup> A thorough review of battlefield trauma care recommendations at the time resulted in the development of the first set of Tactical Combat Casualty Care (TCCC) Guidelines, a set of evidence-based, best-practice trauma care guidelines designed specifically for use on the battlefield, which were published in *Military Medicine* in 1996.<sup>3</sup> The Wilderness Medical Society (WMS) assisted in reviewing the TCCC recommendations, reevaluating the interpretation of the available evidence that supported these recommendations, and discussing their implications for the battlefield and other austere environments.<sup>6</sup>

After 2001, the lessons learned from the battlefields of Iraq and Afghanistan allowed continuous refinement of the TCCC Guidelines through the efforts of the Committee on Tactical Combat Casualty Care (CoTCCC). Importantly, combat medics, corpsmen, and pararescuemen (PJs) have consistently been a strong presence in the CoTCCC.

Now, in 2016, TCCC has been well-documented to have played a major role in achieving the highest casualty survival rate in the history of modern warfare in military units that train all of their members in TCCC.<sup>4,7-11</sup>

TCCC is presently the standard for battlefield trauma care in the US Military and for many allied nations. CoTCCC members work closely with civilian trauma colleagues in the Hartford Consensus effort and the White House Stop the Bleed campaign to translate trauma care lessons learned on the battlefield to lives saved at home. In cooperation with the National Association of Emergency Medical Technicians (NAEMT), the CoTCCC has helped to develop TCCC-based trauma courses to assist in training civilian emergency medical services systems, fire and rescue, and law enforcement organizations in trauma care.

## TACTICAL COMBAT CASUALTY CARE AND WILDERNESS MEDICINE

The wilderness environment presents some of the challenges experienced by medics on the battlefield: both patient and provider are typically in remote locations where evacuation is neither quick nor easy; there may be ongoing hazards to contend with; equipment is limited; the environment may be cold, hot, or aquatic; and the providers are often not trauma care specialists.<sup>6,12</sup> The extreme diversity of nature that makes the wilderness so alluring complicates the care of patients with trauma in a myriad of ways, from whitewater rescue to high-angle rescue to avalanche rescue. Although many of these challenges are distinct from those encountered on the battlefield, there is a good deal of overlap in the approach to patients in these two settings. A selected subset of the trauma care recommendations in TCCC and their applicability to the wilderness environment are reviewed and discussed later. The current TCCC Guidelines may be found on the Joint Trauma System (JTS) and NAEMT Web sites and in the Prehospital Trauma Life Support (PHTLS) textbook that is published every 3 to 4 years by NAEMT.<sup>13</sup>

## APPROACH TO PATIENTS

TCCC recognizes that battlefield trauma care must combine good medicine with good small unit tactics and divides its recommendations into 3 phases: Care under Fire

(while the unit is actively taking hostile fire), Tactical Field Care (the unit and the casualty are still in a prehospital combat environment but not actively engaging the enemy), and Tactical Evacuation Care (during which time the casualty is being transported to definitive care).<sup>3,13</sup> In the wilderness, the terrain, environment, the location, and the nature of the activity in which the group is engaged provide variation and complexity to prehospital trauma care. An open fracture sustained during a multi-day caving expedition must be approached differently than one that occurs in an urban setting.

In both environments, the well-known ABC (airway, breathing, circulation) sequence of initial steps should be replaced with the acronym MARCH (massive bleeding, airway, respirations, circulation, and head/hypothermia.)

In patients with trauma, control of massive external hemorrhage is the most important initial step. Injury to a major vessel may result in death within as little as 5 to 10 minutes. When the hemorrhage is external, first responders have the ability to effectively intervene to stop the bleeding and should do so.

### **EXTERNAL HEMORRHAGE CONTROL: TOURNIQUETS**

The renewed focus on prehospital tourniquet use is one example of the lifesaving potential of the TCCC Guidelines. Until recently, military medics were taught that tourniquets should be used as a last resort, if at all, to control extremity hemorrhage. During the Vietnam conflict and at the start of the war in Afghanistan, tourniquet use was strongly discouraged. As a result, preventable death from extremity hemorrhage was common. Maughon's<sup>1</sup> study of 2600 combat fatalities in the Vietnam conflict found that 7.4% of the deaths examined resulted from extremity hemorrhage. Kelly and colleagues'<sup>14</sup> 2008 study of 982 combat fatalities sustained during the early years in Afghanistan and Iraq found that the incidence of death from extremity hemorrhage as a percentage of the total was essentially unchanged from the Maughon<sup>1</sup> study at 7.8%. Holcomb and colleagues'<sup>15</sup> 2007 study of Special Operations fatalities noted that 3 of the 12 potentially preventable deaths were from extremity hemorrhage. These findings prompted the US Special Operations Command TCCC Transition Initiative in 2005, which expedited the fielding of tourniquets and hemostatic dressings to deploying Special Operations units.<sup>16</sup> As a result, tourniquet and hemostatic dressing use increased in Afghanistan and Iraq and extremity hemorrhage deaths decreased.<sup>4</sup> Eastridge and colleagues'<sup>9</sup> 2012 study of 4596 US combat fatalities from 2001 to 2011 noted that only 2.6% of total combat fatalities resulted from extremity hemorrhage; a 67% decrease.<sup>4,9,17</sup> Kragh and colleagues'<sup>18,19</sup> landmark studies in 2008 and 2009 documented that extremity tourniquets saved lives and did not result in loss of limbs caused by tourniquet ischemia.

Tourniquets have proved to be lifesaving in the civilian and wilderness sectors as well.<sup>20–22</sup> If a person attacked by an animal or a climber with an open fracture sustains a vascular injury with massive external hemorrhage, immediate application of an extremity tourniquet is required. The first responder then has time to convert the tourniquet to other methods of hemostasis when feasible.<sup>23</sup>

### **EXTERNAL HEMORRHAGE CONTROL: HEMOSTATIC DRESSINGS**

When external hemorrhage occurs at sites that are not amenable to tourniquet use, another modality for achieving control of bleeding is the use of hemostatic dressings. The current TCCC Guidelines recommend Combat Gauze as the hemostatic dressing of choice and Celox Gauze and ChitoGauze as alternative choices. All of these hemostatic dressings should be applied with at least 3 minutes of firm, direct pressure.<sup>13,24–29</sup>

Combat medics on the CoTCCC have consistently expressed a strong preference for gauze-type dressings rather than powdered or granular hemostatic agents, especially for wounds in which the bleeding vessel is at the bottom of a narrow wound tract. Gauze-based hemostatic dressings are more easily packed into the depths of such wounds where they can make direct contact with the bleeding vessel. Further, powdered or granular agents may present an ocular hazard if used in a windy environment or in the presence of rotor wash from helicopters. The Israeli Defense Force has reported excellent success in treating external hemorrhage in their combat wounded with Combat Gauze.<sup>30</sup>

Zietlow and colleagues<sup>20</sup> described the Mayo Clinic experience with Combat Gauze in civilian trauma, reporting a 95% success rate after the failure of standard dressings. Both TCCC-recommended tourniquets and hemostatic dressings should be included in all wilderness first aid kits.

## THE PREHOSPITAL TRAUMA AIRWAY

In trauma, managing the airway entails different considerations than those encountered in medical patients. Most airway fatalities in combat are related to direct maxillofacial trauma.<sup>31</sup> Endotracheal intubation in the traumatized airway is challenging, even for experienced intubationists, and most military medics have little experience in intubating patients with airway trauma. Airway trauma on the battlefield is often best managed by allowing the casualty to maintain the sit-up-and-lean-forward position if the patient is conscious and able to do so. This position allows gravity and the patient's protective reflexes to maintain a patent airway.<sup>13</sup> The use of supraglottic airways in the prehospital setting is increasing, but these devices have not been well studied in trauma.

When the measures discussed above do not provide an adequate airway in a casualty with direct trauma to the maxillofacial region, a surgical airway is the intervention of choice.<sup>13,32,33</sup> Combat medics have been shown to be able to perform this procedure with 100% success in a cadaver model when they are well trained and use a CricKey device, which is the recommended device in TCCC.<sup>32,34</sup>

Should the need to secure the airway arise from unconsciousness secondary to traumatic brain injury (TBI) or hemorrhagic shock, nasopharyngeal airways (NPAs) have proved to be a good option.<sup>13</sup> The NPA is an easily trained intervention and no airway fatalities were identified in the 2010 Mabry and colleagues<sup>31</sup> review of this topic as being caused by NPA failure in nontraumatized airways.

In the wilderness setting, a lower incidence of direct maxillofacial trauma and a higher incidence of unconsciousness caused by TBI would be expected. Thus the use of an NPA is a good option in wilderness settings in which a skilled and equipped intubationist is not typically present. A surgical airway remains the emergent airway of last resort for patients with trauma in the wilderness.<sup>35,36</sup>

## RESPIRATIONS/BREATHING

### *Tension Pneumothorax*

In the Vietnam conflict, tension pneumothorax was reported to be the second leading cause of preventable battlefield death.<sup>2</sup> The incidence of death has decreased with the use of body armor that provides significant (but not complete) protection to the chest and back. In addition, combat medical personnel are now taught to use needle decompression (NDC) aggressively to treat suspected tension pneumothorax. The Eastridge and colleagues<sup>9</sup> study noted that tension pneumothorax comprised only

0.2% of deaths among US combat fatalities, which represents a decrease of more than 90% in preventable deaths from this cause.<sup>13</sup>

Current TCCC Guidelines call for casualties who have progressive respiratory distress following torso trauma to be suspected of having a tension pneumothorax and considered for NDC on the side of the injury with a 14-gauge, 8-cm (3.25-inch) needle/catheter unit. NDC is performed at the second intercostal space at the midclavicular line or the fourth or fifth intercostal space at the anterior axillary line. Needle entry into the chest should not be medial to the nipple line and the needle should not be directed toward the heart.<sup>13</sup>

Use of an 8-cm (3.25-inch) needle rather than the previously used 5-cm (2-inch) needle was introduced after the Harcke and colleagues<sup>37</sup> article in 2008. Those investigators noted that several of the cases in their autopsy series had failed attempts at NDC because the needle/catheter units used for the procedure were too short and did not reach the pleural space. This observation was followed by a virtual autopsy computed tomography (CT) study of chest wall thickness that found a mean chest wall thickness of 5.36 cm in the 100 military fatalities studied. Harcke and colleagues<sup>37</sup> recommended that an 8-cm (3.25-inch) needle/catheter unit be used for NDC in order to achieve a 99% assurance of reaching the pleural space.<sup>37</sup> Other investigators have presented similar findings and concerns.<sup>38,39</sup> Harcke and colleagues'<sup>37</sup> findings led to the TCCC recommendation to use an 8-cm (3.25-inch), 14-gauge needle/catheter unit inserted to the hub.<sup>13</sup> The authors are unaware of any reports of death from tension pneumothorax in US combat casualties caused by failed NDC since this change was made almost a decade ago.<sup>13</sup> Bilateral NDC should be performed before resuscitation efforts are abandoned when a casualty with torso trauma or polytrauma has a prehospital cardiopulmonary arrest.<sup>13</sup>

In the wilderness setting, death from tension pneumothorax is expected to be less common, although penetrating chest trauma may occur from hunting accidents and blunt chest trauma may result from falls, avalanche, or mountain bike accidents. Inclusion of an 8-cm (3.25-inch), 14-gauge needle in wilderness medical kits is a reasonable measure in order to be able to perform NDC in the unlikely event that this uncommon but life-threatening disorder is encountered in the wilderness. Improved success using an 8-cm (3.25-inch) needle (83%) compared with a 5-cm (2-inch) needle (41%) was reported by the Mayo Clinic.<sup>40</sup> No complications were reported with either length needle.

### ***Open Pneumothorax***

An open pneumothorax (also sometimes referred to as a sucking chest wound) may result from a penetrating injury to the chest wall. When the defect in the chest wall is sufficiently large (usually two-thirds or more of the diameter of the trachea), air preferentially flows into the chest cavity via the defect in the chest wall, instead of into the lung via the trachea, as the casualty inhales. Air entering through the defect in the chest wall allows the lung on the affected side to collapse and impairs oxygen exchange.<sup>13</sup>

Although there is little evidence that an open pneumothorax by itself (ie, without injury to underlying lung tissue and major vascular structures), is a potentially lethal injury,<sup>9,41</sup> the impaired pulmonary gas exchange could potentially result in pulmonary compromise in a polytrauma casualty and contribute to secondary hypoxic brain injury in casualties with TBI.<sup>13</sup>

Treatment of an open pneumothorax consists of applying a vented occlusive chest seal over the defect in the chest wall, thus preventing air from entering the pleural space through the defect in the chest wall.<sup>41</sup> The vent in the chest seal prevents the development of a tension pneumothorax in the presence of an underlying lung injury

and air leak. Commercially available vented and nonvented chest seals have been evaluated in animal studies.<sup>42,43</sup> In the Kheirabadi study, chest seals were applied and then 200-mL increments of air were injected into the pleural cavity of thoracotomized swine every 5 minutes until either tension pneumothorax developed or the volume of air injected equaled 100% of the animal's estimated total lung capacity.<sup>42</sup> Tension pneumothorax did not develop in animals treated with vented chest seals (incorporating a 1-way valve that allowed air to leave but not to enter the pleural space). However, tension pneumothorax did occur in the animals with chest seals without valves. Vented chest seals, then, are preferred for the prehospital management of open pneumothorax, to be followed by tube thoracostomy when time, skills, and circumstances allow.<sup>13,41</sup>

For wilderness medicine providers, open pneumothorax and the need for a vented chest seal is an injury pattern that might result from gunshot wounds sustained in hunting accidents. This injury, in the setting of delays to evacuation, is one that should be treated with prophylactic antibiotics, as described later.

## **INTRAVENOUS AND INTRAOSSEOUS ACCESS**

It has been a long-standing prehospital trauma care practice to establish intravenous (IV) access for individuals with significant trauma. TCCC has reconsidered this intervention. Starting IV lines inflicts significant cost in both time and logistics and may delay a combat unit's ability to maneuver when needed. TCCC recommends that IV lines be started only for individuals who require fluid resuscitation as a result of hemorrhagic shock or who need IV medications, especially tranexamic acid (TXA) or analgesics. IV lines are easily dislodged, especially during combat operations. The TCCC curriculum describes a technique developed by medics in the 75th Ranger Regiment that helps to ensure that IV lines are not dislodged during casualty movement.<sup>13,44</sup>

## **INTRAOSSEOUS ACCESS**

Establishing IV access is a fairly easy procedure unless the patient is obese, a small child, or in shock, or the person attempting to start the IV does not perform this procedure routinely. However, the last 2 caveats often apply to the battlefield setting. Another way to achieve vascular access is to use an intraosseous (IO) device.<sup>45,46</sup>

This technique is easily trained<sup>47</sup> and can be performed by prehospital providers with a high rate of success.<sup>48</sup> Because the technique is performed based on bony landmarks rather than by visualizing peripheral veins, it is more easily performed under low light and night vision device conditions.<sup>13</sup> The IO approach was proposed by TCCC as an alternative to IV lines on the battlefield in 2002 and quickly became widely used by combat medical personnel because the procedure is quickly and easily performed. IO access is now a widely used technique in the civilian sector as well.<sup>49</sup> The Pyng FAST-1 and the EZ-IO devices have been the most widely used by the US Military in the recent conflicts.<sup>13</sup> Although the risk of osteomyelitis is very low and the authors are unaware of any reports of this complication from the conflicts in Iraq and Afghanistan, both the potential for infection and the patient discomfort experienced with an IO procedure make using peripheral IV access the option of first choice when circumstances allow.

For practitioners of wilderness medicine, the indications for IV and IO access are basically the same as for the battlefield, except that it may be less common for medical providers in the wilderness setting to be prepared to perform fluid resuscitation from hemorrhagic shock.

## TRANEXAMIC ACID

Noncompressible hemorrhage is the leading cause of preventable death in combat casualties.<sup>9</sup> The large, prospective, randomized Clinical Randomization of an Antifibrinolytic in Significant Hemorrhage (CRASH-2) study examined the effect of TXA administration in trauma patients at risk of bleeding and documented a small but statistically significant survival benefit. Although deep venous thrombosis is a known complication of trauma, there was no increase in the rate of vascular occlusive events in the TXA group.<sup>50</sup> The subsequent subgroup analysis of the CRASH-2 data focused on deaths caused by bleeding rather than all-cause mortality and examined the effect of the timing of TXA administration on outcomes. The greatest benefit of TXA administration was obtained when the medication was given within 1 hour of the time of injury. TXA administered between 1 and 3 hours after the time of injury also reduced the risk of death caused by bleeding. However, TXA administered later than 3 hours after injury was observed to increase the risk of death from exsanguination.<sup>51</sup>

The Military Application of Tranexamic Acid in Trauma Emergency Resuscitation Study (MATTERS) was performed at a role 3 facility in Afghanistan.<sup>52</sup> The MATTERS investigators found a decreased mortality among combat casualties who received TXA, despite their being more seriously injured. In the subgroup of casualties who received massive transfusions (more than 10 units of red cells within the first 24 hours), mortality in the TXA group was markedly lower (14.4%) compared with the control group (28.1%). The CRASH-2 and MATTERS findings both supported the use of TXA in combat casualties who are either in hemorrhagic shock or at significant risk of that condition. As a result of these two studies, TXA was added to the TCCC Guidelines in 2011.<sup>53</sup>

A subsequent report found a benefit from administering TXA in the prehospital setting.<sup>54</sup> Studies examining the effect of giving TXA before elective surgical procedures associated with significant blood loss have consistently found that TXA reduces blood loss in these procedures and does not increase the risk of deep venous thrombosis.<sup>55</sup> TXA is recommended in the TCCC Guidelines for casualties who are anticipated to need significant blood transfusion (eg, casualties with hemorrhagic shock, 1 or more major amputations, penetrating torso trauma, or evidence of severe bleeding).<sup>44</sup> The recommended dose is 1 g of TXA in 100 mL of normal saline or Lactated Ringers solution. TXA should be infused over 10 minutes to avoid the risk of hypotension and should be given as soon as possible after injury but not later than 3 hours after injury. In the event that evacuation to a medical treatment facility is delayed, a second infusion of 1 g of TXA should be administered after fluid resuscitation has been performed.<sup>13</sup>

In the wilderness environment, TXA administration should be considered for trauma patients in shock, as well as those with penetrating torso trauma, severe external hemorrhage, or blunt trauma with suspicion of non-compressible hemorrhage. Falls and TBI are common in the wilderness and the value of TXA in reducing intracranial hemorrhage is currently an area of active investigation.

## FLUID RESUSCITATION FROM HEMORRHAGIC SHOCK

Hemorrhagic shock is the most common cause of potentially preventable death on the battlefield.<sup>9</sup> In the hospital setting, hemorrhagic shock is treated by resuscitation with blood or blood components until adequate tissue perfusion is restored. A large, prospective, randomized study at Ben Taub Hospital in Houston examined the benefit of early, aggressive fluid resuscitation in patients with penetrating torso trauma and shock.<sup>56</sup> This study found that early fluid resuscitation group had decreased survival



as compared to the group of patients in whom fluid resuscitation was delayed until after surgical hemostasis had been achieved.<sup>56</sup> TCCC originally recommended that fluid resuscitation for casualties with shock from noncompressible bleeding sites be delayed until after surgical control of the bleeding and that 1000 mL of Hespan be used to resuscitate casualties in shock from external bleeding sites after hemorrhage control.<sup>3</sup>

However, in 1999, subject matter experts reviewing battlefield trauma care issues from the Battle of Mogadishu recommended that casualties who were experiencing unconsciousness or mental status changes caused by hemorrhagic shock should be fluid resuscitated to an end point of improved mentation, even in the presence of ongoing noncompressible (internal) hemorrhage.<sup>57</sup> Further recommendations from subsequent Department of Defense (DoD) fluid resuscitation workshops on fluid resuscitation led to a single hypotensive fluid resuscitation strategy for combat casualties in shock, whether their hemorrhage was controlled or uncontrolled. The synthetic hetastarch solution Hextend replaced the previously used Hespan because of the former solution's lesser impact on coagulation status.<sup>58,59</sup>

Both of the strategies mentioned earlier made the implicit assumption that blood products would not be available in the tactical field care phase of combat casualty care. In the ensuing years, a number of events have occurred: (1) the value of whole blood as the preferred fluid for fluid resuscitation has been documented<sup>60–63</sup>; (2) transfusing red blood cells in a 1:1 or 1:1:1 ratio with plasma and platelets has been shown to be superior to a transfusion strategy using a predominance of RBCs<sup>64,65</sup>; (3) large volumes of crystalloid have been shown to worsen outcomes in patients with trauma,<sup>66,67</sup>; (4) questions have been raised about the safety of using hetastarch solutions in critically ill patients<sup>68,69</sup>; (5) both military units and civilian trauma systems have shown that blood products can be used in the prehospital setting<sup>61,70–77</sup> and that this practice improves casualty outcomes<sup>74</sup>; and (6) a preponderance of the literature has found that plasma resuscitation is a better choice for resuscitation than either crystalloids or colloids.<sup>78,79</sup> Accordingly, a recent review of fluid resuscitation options in TCCC has recommended a preference list of options for fluid resuscitation, going from the most desirable (whole blood) to the least desirable (crystalloids).

In the wilderness setting, as with ground-based combat medical personnel, blood components are unlikely to be readily available. In this circumstance, the best strategy for treating hemorrhagic shock is hypotensive resuscitation using a reconstituted dried plasma product. Although dried plasma products (French FyLP or German LyoPlas) have been used widely by coalition forces in Afghanistan and Iraq, they are not currently approved by the US Food and Drug Administration for use in the United States (Butler FK: Fluid resuscitation in tactical combat casualty care – yesterday and today. Submitted for publication to *Wilderness and Environmental Medicine*).<sup>78</sup> As long as dried plasma remains unavailable in the United States, if a prolonged evacuation is anticipated, Hextend will provide a more sustained intravascular volume expansion than crystalloids.<sup>80</sup>

## HYPOTHERMIA AND COAGULOPATHY ON THE BATTLEFIELD

As noted previously, hemorrhage is the leading cause of preventable death in combat casualties. Hypothermia may contribute to the coagulopathy of trauma by decreasing platelet function, decreasing the activity of enzymes in the coagulation cascade, and causing alterations of the fibrinolytic system.<sup>81,82</sup> Hypothermia (defined as a core temperature <36°C) was present in 18% of 2848 combat casualties and was an independent predictor of mortality.<sup>83</sup> Hypothermia may occur even in warm environments and



may be exacerbated by long evacuation times, helicopter transport, and wet clothing. Hemorrhagic shock also contributes to hypothermia by reducing the body's ability to produce metabolic heat.<sup>13</sup>

The body temperature in combat casualties should be maintained as close to 37°C as possible.<sup>84</sup> Hypothermia prevention should be started early in the management of combat casualties, after external hemorrhage has been controlled and TXA administered, if indicated. The casualty should be protected against the environment to the greatest extent possible. Clothing should be retained where that is consistent with treating injuries. Conductive heat loss should be prevented by not having the casualty lying directly on the ground or on the floor of an evacuation helicopter. Personal protective equipment should be kept on or with the casualty as feasible. After massive hemorrhage has been controlled, the airway opened, and breathing difficulties addressed, wet clothing should be removed and replaced with dry clothes, if feasible.

TCCC recommends the Hypothermia Prevention and Management Kit (HPMK) to minimize heat loss in combat casualties. The HPMK combines an active heating device (the Ready-Heat blanket) with a passive device (the Heat Reflective Shell) to produce a synergistic heat-conserving effect for the casualty. The HPMK has been shown in United States Army Institute of Surgical Research testing to effectively prevent heat loss.<sup>85</sup> Care should be taken to place the Ready-Heat blanket over the uniform shirt, not directly on the skin, because the device may cause burns if placed on bare skin.

If purposed hypothermia prevention devices are not available, then blankets, ponchos, sleeping bags, or other items at hand should be used to keep the casualty as warm and dry as possible. Controlling hemorrhage and providing adequate fluid resuscitation, with blood components if possible, will restore tissue perfusion and help to maintain the casualty's ability to generate metabolic heat.

The HPMK or similar devices may be very useful in the wilderness, but often are not available. Extra clothing, sleeping bags, insulating blankets, and other items may be more readily available and should be applied in layers to achieve effective hypothermia prevention for patients with trauma in the wilderness.

The WMS Practice Guidelines for the prevention and treatment of hypothermia offer additional treatment options and perspectives.<sup>86</sup>

## EYE TRAUMA

Penetrating eye injuries are frequently encountered on the battlefield, especially when exposed to improvised explosive devices (IEDs) with small rock or metallic items embedded in the device to increase the incidence of fragment injuries from the blast. Providers should have a heightened index of suspicion in the presence of multiple punctate facial wounds (peppering). When injury to the globe is suspected, first-responder management of these injuries includes a rapid field evaluation and documentation of vision, covering the injured eyes with a rigid eye shield, and prompt administration of TCCC-recommended antibiotics. Eye shields are essential in this setting but have been reported to be underused in the management of eye injuries during the recent conflicts.<sup>87</sup> If no rigid eye shield is available, a set of protective eyewear will serve this purpose well. A point of emphasis is that pressure patches and other dressings that may come into direct contact with an injured eye should be avoided.<sup>13</sup> Also, avoid manipulating the eye during evaluation and treatment in any way that might increase intraocular pressure and result in the expulsion of intraocular contents through the corneal or scleral defect.<sup>13,88</sup> Moxifloxacin 400 mg once a day, given immediately after injury, is the antibiotic of choice in the

prehospital environment. This medication has good coverage of the typical causative organisms for intraocular infections and penetrates well into the vitreous cavity. No topical antibiotics (ointments or eye drops) should be used on an unrepaired penetrating injury of the eye.<sup>13,88</sup>

Wilderness medicine practitioners are unlikely to encounter any IEDs, but may have to treat patients with fishhook or impaling thorn injuries to the eye. In both cases, the most important additional point of management is not to try to remove the fishhook or thorn.<sup>88</sup> Wilderness patients with known or suspected penetrating eye injuries should be evacuated as soon as feasible. A more complete discussion of the management of eye injuries and medical disorders in a wilderness setting may be found in the new seventh edition of Auerbach's<sup>88</sup> *Wilderness Medicine* as well as the WMS practice guidelines for eye injuries.<sup>89</sup>

## ANALGESIA

At the onset of the Afghanistan conflict, the state of the art for battlefield analgesia in most of the US Military was the same as it had been at the Battle of Bull Run in the US Civil War.<sup>13,90</sup> Intramuscular (IM) morphine was the primary medication used to relieve the pain of combat wounds despite that fact that it is very slow-acting and prolongs the casualty's pain.<sup>3</sup> This, in turn, increases the likelihood of repeated dosing and eventual overdose.<sup>91</sup> Early adopters of TCCC in 2001 were using morphine administered intravenously for rapidity of analgesia and improved ability to titrate the dosage administered.<sup>3</sup>

After demonstration that oral transmucosal fentanyl (OTFC) could safely provide rapid and powerful analgesia in combat casualties without the need to establish an IV, this medication was added to the TCCC Guidelines.<sup>92,93</sup> Wedmore and colleagues<sup>94</sup> subsequently provided additional documentation of the safety of OTFC on the battlefield. OTFC had been recommended for use in wilderness settings as early as 1999.<sup>95</sup>

Ketamine is an excellent analgesic alternative for casualties either in, or at risk of, hemorrhagic shock because this medication does not cause hemodynamic or respiratory depression. Ketamine was added as an analgesic option in TCCC in 2012.<sup>90</sup>

In 2014, the TCCC triple-option analgesia approach to battlefield analgesia was developed. This concept originated at the direct request from combat medical personnel that the CoTCCC provide them with an inclusive but simplified approach on how best to use the available options for analgesia.<sup>17</sup> Thus, oral medications (acetaminophen and meloxicam, a platelet-sparing nonsteroidal analgesic) are recommended for analgesia when pain is not severe and the casualty can continue to function effectively as a combatant for his or her unit. When there is more severe pain, but the casualty is not in, or at significant risk for, shock, OTFC is the analgesic of choice. If there is moderate to severe pain, and the casualty is in shock, or at risk of shock or pulmonary compromise, ketamine is the agent of choice.<sup>92</sup> Ondansetron is recommended for the nausea and vomiting that may result either from opioid administration or from the combat wounds.<sup>96</sup>

The TCCC triple-option analgesia approach lends itself well to the wilderness environment. A practice guideline that mirrors the TCCC triple-option approach was recently approved by the American College of Emergency Physicians.<sup>97</sup> The WMS practice guidelines for pain management also include a discussion of the analgesic options noted earlier.<sup>98</sup>

## ANTIBIOTICS

Open combat wounds are prone to infection and are a late cause of morbidity and mortality in combat casualties.<sup>99</sup> Antibiotics are indicated for combat trauma and should be

started as soon as feasible. Fourth-generation oral fluoroquinolones are the prehospital antibiotic of choice when the casualty is able to take oral medications.<sup>99</sup> The current recommendation is 400 mg of moxifloxacin when oral medications are feasible and IM or IV ertapenem when the casualty is not able to take oral medications.<sup>13,44,100</sup> Both antibiotics were selected because of their excellent spectrum of activity and their minimal serious adverse effects. In addition, moxifloxacin has excellent bioavailability when taken orally and both medications penetrate well into the vitreous cavity, which is important in treating casualties with penetrating eye injuries.

For injuries that occur in the wilderness environment, as in combat, evacuation is likely to be delayed and oral or parenteral antibiotics are indicated for any trauma that includes infection-prone injuries such as open fractures or animal bites.

## TRAUMATIC CARDIOPULMONARY ARREST

Although cardiopulmonary resuscitation (CPR) has been lifesaving in patients with nontraumatic cardiac arrest, CPR is much less likely to be successful in victims of traumatic cardiac arrest.<sup>101–103</sup> On the battlefield, performing CPR on casualties in cardiac arrest may result in additional lives lost as combat medical personnel are exposed to hostile fire while performing CPR and care is withheld from casualties with potentially survivable wounds. In addition, delays in unit movement in order to perform CPR may result in mission compromise.<sup>3,13</sup> CPR should be considered in the tactical prehospital setting only if the arrest is associated with hypothermia, near drowning, or electrocution, or when the casualty is in the tactical evacuation phase of care and arrival at a medical treatment facility is imminent.<sup>13,104</sup> Unrecognized tension pneumothorax is a potentially reversible cause of traumatic cardiac arrest in combat casualties. Any combat casualty who sustains torso trauma or polytrauma and is found to have no pulse or respirations should have bilateral needle decompression performed to ensure they do not have a tension pneumothorax before discontinuation of care.<sup>13,104</sup>

## DOCUMENTATION OF CARE

The lack of adequate documentation of prehospital care rendered to US casualties in Iraq and Afghanistan is problematic in ensuring optimal care for casualties and has been a clear obstacle to CoTCCC and Joint Trauma System efforts to improve TCCC.<sup>105</sup> The TCCC Card was developed by the 75th Ranger Regiment to document prehospital care and was adopted by TCCC in 2008. This card is simple, cost-effective, and easily fielded. It has been used widely throughout the wars in Iraq and Afghanistan and is now approved for use throughout the DoD. The recently updated TCCC card allows for a casualty Battle Roster Number (to link the casualty to the DoD Trauma Registry), provides better definition of the mechanism of injury, improves the documentation of tourniquet and hemostatic dressing use, and contains a space for types and doses of analgesics used as well as for several other commonly used elements of combat casualty care.

The TCCC card is commercially available on ruggedized paper stock that enables it to be used with reduced incidence of becoming damaged or unreadable in the harsh combat environment. Documentation of care is an important aspect of caring for casualties in the wilderness as well and either a TCCC card or some other durable record of care should be maintained throughout the time that the patient is receiving prehospital care and subsequently should be transported with the patient to the treating hospital.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge ongoing efforts of the CoTCCC, the TCCC Working Group, and the Joint Trauma System to improve the care provided to US casualties on the battlefield and to patients with trauma in the United States.

## REFERENCES

1. Maughon JS. An inquiry into the nature of wounds resulting in killed in action in Vietnam. *Mil Med* 1970;135:8–13.
2. Bellamy RF. The causes of death in conventional land warfare: implications for combat casualty care research. *Mil Med* 1984;149:55–62.
3. Butler FK, Hagmann J, Butler EG. Tactical Combat Casualty Care in Special Operations. *Mil Med* 1996;161(Suppl):3–16.
4. Butler FK. The US military experience with tourniquets and hemostatic dressings in the Afghanistan and Iraq conflicts. *Bull Am Coll Surg* 2015;100(Hartford Consensus Supplement):60–5.
5. Kragh J, Walters T, Westmoreland T, et al. Tragedy into drama: an American history of tourniquet use in the current war. *J Spec Oper Med* 2013;13:5–25.
6. Butler FK, Zafren K. Tactical management of wilderness casualties in Special Operations. *Wilderness Environ Med* 1998;9:62–117.
7. Butler FK. Two decades of saving lives on the battlefield: Tactical Combat Casualty Care turns 20. *Mil Med*, in press.
8. Butler FK, Blackburn LH. Battlefield trauma care then and now: a decade of tactical combat casualty care. *J Trauma* 2012;73:S395–402.
9. Eastridge BJ, Mabry RL, Seguin P, et al. Death on the battlefield (2001–2011): implications for the future of combat casualty care. *J Trauma* 2012;73(6 Suppl 5):S431–7.
10. Kotwal RS, Montgomery HR, Kotwal BM, et al. Eliminating preventable death on the battlefield. *Arch Surg* 2011;146:1350–8.
11. Savage E, Forestier C, Withers N, et al. Tactical Combat Casualty Care in the Canadian Forces: lessons learned from the Afghan War. *Can J Surg* 2011;59: S118–23.
12. Sward D, Bennett BL. Wilderness medicine. *World J Emerg Med* 2014;5:5–15.
13. Butler FK, Giebner SD, McSwain N, et al, editors. Prehospital trauma life support manual. 8th edition (Military). Burlington (MA): Jones and Bartlett Learning; 2014.
14. Kelly JF, Ritenour AE, McLaughlin DF, et al. Injury severity and causes of death from Operation Iraqi Freedom and Operation Enduring Freedom: 2003–2004 versus 2006. *J Trauma* 2008;64(Suppl):S21–6.
15. Holcomb JB, McMullen NR, Pearse L, et al. Causes of death in Special Operations Forces in the Global War on Terror. *Ann Surg* 2007;245:986–91.
16. Butler F, Holcomb J. The tactical combat casualty care transition initiative. *Army Med Department J* 2005;April–June:33–8.
17. Kotwal RS, Butler FK, Edgar EP, et al. Saving lives on the battlefield: a Joint Trauma System review of pre-hospital trauma care in combined joint operating area – Afghanistan. *J Spec Oper Med* 2013;13:77–80.
18. Kragh JF, Walters TJ, Baer DG, et al. Practical use of emergency tourniquets to stop bleeding in major limb trauma. *J Trauma* 2008;64(Suppl):S38–50.
19. Kragh JF, Walters TJ, Baer DG, et al. Survival with emergency tourniquet use to stop bleeding in major limb trauma. *Ann Surg* 2009;249:1–7.

20. Zietlow J, Zietlow S, Morris D, et al. Prehospital use of hemostatic bandages and tourniquets: translation from military experience to implementation in a civilian trauma center. *J Spec Oper Med* 2015;15:48–53.
21. Scerbo MH, Mumm JP, Gates K, et al. Safety and Appropriateness of Tourniquets in 105 Civilians. *Prehosp Emerg Care* 2016;20:712–22.
22. Pons O, Jerome J, McMullen J, et al. The Hartford Consensus on active shooters: implementing the continuum of prehospital trauma response. *J Emerg Med* 2015;49:878–85.
23. Shackelford S, Butler F, Kragh J, et al. Optimizing the use of limb tourniquets in Tactical Combat Casualty Care: TCCC guidelines change 14-02. *J Spec Oper Med* 2015;15:17–31.
24. Bennett BL, Littlejohn LF, Kheirabadi BS, et al. Management of external hemorrhage in Tactical Combat Casualty Care: chitosan-based hemostatic gauze dressings. *J Spec Oper Med* 2014;14:12–29.
25. Bennett BL, Littlejohn L. Review of new topical hemostatic dressings for combat casualty care. *Mil Med* 2014;179:497–514.
26. Littlejohn L, Bennett B, Drew B. Application of current hemorrhage control techniques for backcountry care: part 2: hemostatic dressings and other adjuncts. *Wilderness Environ Med* 2015;26:246–54.
27. Drew B, Bennett B, Littlejohn L. Application of current hemorrhage control techniques for backcountry care: part 1: tourniquets and hemorrhage control adjuncts. *Wilderness Environ Med* 2015;26:236–45.
28. Kheirabadi BS, Edens JW, Terrazas IB, et al. Comparison of new hemostatic granules/powders with currently deployed hemostatic products in a lethal model of extremity arterial hemorrhage in swine. *J Trauma* 2009;66:316–26.
29. Kheirabadi B, Mace J, Terrazas I, et al. Safety evaluation of new hemostatic agents, smectite granules, and kaolin-coated gauze in a vascular injury wound model in swine. *J Trauma* 2010;68:269–78.
30. Shina A, Lipsky A, Nadler R, et al. Prehospital use of hemostatic dressings by the Israel Defense Forces Medical Corps: a case series of 122 patients. *J Trauma* 2015;79:S204–9.
31. Mabry RL, Edens JW, Pearse L, et al. Fatal airway injuries during Operation Enduring Freedom and Operation Iraqi Freedom. *Prehosp Emerg Care* 2010;14:272–7.
32. Mabry R, Frankfurt A, Kharod C, et al. Emergency cricothyroidotomy in Tactical Combat Casualty Care. *J Spec Oper Med* 2015;15:11–9.
33. Hessert MJ, Bennett BL. Optimizing emergent surgical cricothyrotomy for use in austere environments. *Wilderness Environ Med* 2013;24:53–66.
34. Mabry R, Nichols M, Shiner D, et al. A comparison of two open surgical cricothyroidotomy techniques by military medics using a cadaver model. *Ann Emerg Med* 2014;63:1–5.
35. Wharton D, Bennett B. Surgical cricothyrotomy in the wilderness: a case report. *Wilderness Environ Med* 2013;24:12–4.
36. Johnson CA, Goodwine DS, Passier I. Improvised cricothyrotomy on a mountain using hiking gear. *Wilderness Environ Med* 2016;27(4):500–3.
37. Harcke HT, Pearse LA, Levy AD, et al. Chest wall thickness in military personnel: implications for needle thoracentesis in tension pneumothorax. *Mil Med* 2008;172:1260–3.
38. Zengerink I, Brink PR, Laupland KB, et al. Needle thoracostomy in the treatment of tension pneumothorax in trauma patients: what size needle? *J Trauma* 2008;64:111–4.

39. Givens ML, Ayotte K, Manifold C. Needle thoracostomy: implications of computed tomography chest wall thickness. *Acad Emerg Med* 2004;11:211–3.
40. Aho J, Thiels C, El Khatin M, et al. Needle thoracostomy: clinical effectiveness is improved using a longer angiocatheter. *J Trauma* 2016;80:272–7.
41. Butler F, Dubose J, Otten E, et al. Management of open pneumothorax in Tactical Combat Casualty Care: TCCC guidelines change 13-02. *J Spec Oper Med* 2013;13:81–6.
42. Kheirabadi BS, Terrazas IB, Koller A, et al. Vented versus unvented chest seals for treatment of pneumothorax and prevention of tension pneumothorax in a swine model. *J Trauma* 2013;75:150–6.
43. Kotora JG Jr, Henao J, Littlejohn LF, et al. Vented chest seals for prevention of tension pneumothorax in a communicating pneumothorax. *J Emerg Med* 2013;45:686–94.
44. National Association of Emergency Medical Technicians Web site: TCCC guidelines. Available at: [http://www.naemt.org/education/TCCC/guidelines\\_curriculum](http://www.naemt.org/education/TCCC/guidelines_curriculum); Accessed September 27, 2016.
45. Dubick MA, Holcomb JB. A review of intraosseous vascular access: current status and military application. *Mil Med* 2000;165:552–9.
46. Calkins MD, Fitzgerald G, Bentley TB, et al. Intraosseous infusion devices: a comparison for potential use in special operations. *J Trauma* 2000;48:1068–74.
47. Benson G. Intraosseous access to the circulatory system: an under-appreciated option for rapid access. *J Perioper Pract* 2015;25:140–3.
48. Byars DV, Tsuchitani SN, Erwin E, et al. Evaluation of success rate and access time for an adult sternal intraosseous device deployed in the prehospital setting. *Prehospital Disaster Med* 2011;26:127–9.
49. Lewis P, Wright C. Saving the critically injured trauma patient: a retrospective analysis of 1000 uses of intraosseous access. *Emerg Med J* 2015;32:463–7.
50. CRASH-2 Collaborators. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant hemorrhage. (CRASH-2); a randomized, placebo-controlled trial. *Lancet* 2010;376:23–32.
51. CRASH-2 Collaborators. The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomized controlled trial. *Lancet* 2011;377:1096–101.
52. Morrison JJ, Dubose JJ, Rasmussen TE, et al. Military Application of Tranexamic Acid in Trauma Emergency Resuscitation Study (MATTERS). *Arch Surg* 2012; 147:113–9.
53. Dickey NW, Jenkins D. Defense Health Board recommendation for the addition of tranexamic acid to the Tactical Combat Casualty Care guidelines. Defense Health Board Memorandum 2011:1–12. Available at: <http://www.naemt.org/docs/default-source/education-documents/tccc/10-9-15-updates/dhb-memo-110923-txa.pdf?sfvrsn=2>. Accessed January 19, 2017.
54. Wafaisade A, Lefering R, Bouillon B, et al. Prehospital administration of tranexamic acid in trauma patients. *Crit Care* 2016;20:1–9.
55. Huang F, Wu D, Ma G, et al. The use of tranexamic acid to reduce blood loss and transfusion in major orthopedic surgery: a meta-analysis. *J Surg Res* 2014;186:318–27.
56. Bickell WH, Wall MJ, Pepe PE, et al. Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. *N Engl J Med* 1994; 331:1105–9.
57. Butler FK, Hagmann J. Tactical management of urban warfare casualties in special operations. *Mil Med* 2000;165:1–48.

58. Champion HR. Combat fluid resuscitation: introduction and overview of conferences. *J Trauma* 2003;54(Suppl):S7–12.
59. Holcomb JB. Fluid resuscitation in modern combat casualty care: lessons learned from Somalia. *J Trauma* 2003;54(Suppl):S46–51.
60. Spinella P, Pidcock H, Strandenes G, et al. Whole blood transfusion for hemostatic resuscitation of major bleeding. *Transfusion* 2016;56:S190–202.
61. Cap A, Pidcock H, DePasquale M, et al. Blood far forward: time to get moving! *J Trauma* 2015;78:S2–6.
62. Stubbs J, Zielinski M, Jenkins D. The state of the science of whole blood: lessons learned at Mayo Clinic. *Transfusion* 2016;56:S173–81.
63. Spinella PC, Perkins JG, Grathwohl KW, et al. Warm fresh whole blood is independently associated with improved survival for patients with combat-related traumatic injuries. *J Trauma* 2009;66:S69–76.
64. Holcomb J, Spinella P. Optimal use of blood in trauma patients. *Biologicals* 2010;38:72–7.
65. Borgman MA, Spinella PC, Perkins JG, et al. The ratio of blood products transfused in patients receiving massive transfusions at a combat support hospital. *J Trauma* 2007;63:805–13.
66. Ley E, Clond M, Srouf M, et al. Emergency department crystalloid resuscitation of 1.5 L or more is associated with increased mortality in elderly and non-elderly trauma patients. *J Trauma* 2011;70:398–400.
67. Duke MD, Guidry C, Guice J, et al. Restrictive fluid resuscitation in combination with damage control resuscitation: time for adaptation. *J Trauma* 2012;73:674–8.
68. Lissauer ME, Chi A, Kramer ME, et al. Association of 6% hetastarch resuscitation with adverse outcomes in critically ill trauma patients. *Am J Surg* 2011;202:501–8.
69. Zarychanski R, Abou-Setta A, Turgeon A, et al. Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation. *JAMA* 2013;309:678–88.
70. Morrison JJ, Oh J, Dubose JJ, et al. En-Route Care Capability From Point of Injury Impacts Mortality After Severe Wartime Injury. *Ann Surg* 2013;257:330–4.
71. Spinell P, Cap A. Whole blood – back to the future. *Curr Opin Hematol* 2016;23(6):536–42.
72. Strandenes G, Cap A, Cacic D, et al. Blood far forward - a whole blood research and training program for austere environments. *Transfusion* 2013;53:S124–30.
73. Holcomb J, Donathan D, Cotton B, et al. Prehospital transfusion of plasma and red blood cells in trauma patients. *Prehosp Emerg Care* 2015;19:1–9.
74. Powell E, Hinckley W, Gottula A, et al. Shorter times to packed red blood cells transfusion are associated with decreased risk of death in traumatically injured patients. *J Trauma Acute Care Surg* 2016;81(3):458–62.
75. Fisher A, Miles E, Cap A, et al. Tactical damage control resuscitation. *Mil Med* 2015;180:869–75.
76. Wild G, Anderson D, Lund P. Round Afghanistan with a fridge. *J R Army Med Corps* 2013;159:24–9.
77. Kim B, Zielinski M, Jenkins D, et al. The effects of prehospital plasma on patients with injury: a prehospital plasma resuscitation. *J Trauma Acute Care Surg* 2012;73:S49–53.
78. Butler FK, Blackburn LH, Gross KR. The combat medic aid bag – 2025: CoTCCC top ten recommended battlefield trauma care research, development, and evaluation priorities for 2015. *J Spec Oper Med* 2015;15:7–19.



79. Butler FK, Holcomb JB, Kotwal RS, et al. Fluid resuscitation for hemorrhagic shock in Tactical Combat Casualty Care: TCCC guidelines change 14-01. *J Spec Oper Med* 2014;14:13–38.
80. Mortelmans Y, Merckx E, van Nerom C, et al. Effect of an equal volume replacement with 500cc 6% hydroxyethyl starch on the blood and plasma volume of healthy volunteers. *Eur J Anesthesiol* 1995;12:259–64.
81. Wolberg AS, Meng ZH, Monroe DM, et al. A systematic evaluation of the effect of temperature on coagulation enzyme activity and platelet function. *J Trauma* 2004;56:1221–8.
82. Watts DD, Trask A, Soeken K, et al. Hypothermic coagulopathy in trauma: effect of varying levels of hypothermia on enzyme speed, platelet function, and fibrinolytic activity. *J Trauma* 1998;44:846–54.
83. Arthurs Z, Cuadrado D, Beekley A, et al. The impact of hypothermia on trauma care at the 31st combat support hospital. *Am J Surg* 2006;191:610–4.
84. Winkenwerder W. Defense-wide policy on combat trauma casualty hypothermia prevention and treatment. Assistant Secretary of Defense for Health Affairs Memorandum dated February 16, 2006. Available at: [http://www.health.mil/libraries/HA\\_Policies\\_and\\_Guidelines/06-005.pdf](http://www.health.mil/libraries/HA_Policies_and_Guidelines/06-005.pdf). Accessed 14 August 2013.
85. Allen PB, Salyer SW, Dubick MA, et al. Preventing hypothermia: comparison of current devices used by the U.S. Army with an in vitro warmed fluid model. *J Trauma* 2010;69(Suppl):S154–61.
86. Zafren K, Giesbrecht GG, Danzl DF, et al. Wilderness Medical Society practice guidelines for the out-of-hospital evaluation and treatment of accidental hypothermia: 2014 update. *Wilderness Environ Med* 2014;25(Suppl):S66–85.
87. Mazzoli RA, Gross KR, Butler FK. The use of rigid eye shields (fox shields) at the point of injury for ocular trauma in Afghanistan. *J Trauma* 2014;77:S156–62.
88. Butler FK, Chalfin S. The eye in the wilderness. In: Auerbach PS, editor. *Wilderness medicine*. 7th edition. St Louis (MO): Mosby; 2016. p. 1109–27.
89. Paterson R, Drake B, Tabin G, et al. Wilderness Medical Society practice guidelines for treatment of eye injuries and illnesses in the wilderness: 2014 update. *Wilderness Environ Med* 2014;25(Suppl):S19–29.
90. Dickey N, Jenkins D, Butler F. Prehospital use of ketamine in battlefield analgesia. Defense Health Board Memorandum 2012:1–11. Available at: <http://www.naemt.org/docs/default-source/education-documents/tccc/10-9-15-updates/dhb-memo-120308-ketamine.pdf?sfvrsn=2>. Accessed January 19, 2017.
91. Beecher H. Delayed morphine poisoning in battle casualties. *JAMA* 1944;124:1193–4.
92. Butler FK, Kotwal RS, Buckenmaier CC III, et al. A triple-option analgesia plan for Tactical Combat Casualty Care. *J Spec Oper Med* 2014;14:13–25.
93. Kotwal R, O'Connor K, Johnson T, et al. A novel pain management strategy for combat casualty care. *Ann Emerg Med* 2004;44:121–7.
94. Wedmore I, Kotwal R, McManus J, et al. Safety and efficacy of oral transmucosal fentanyl citrate for prehospital pain control on the battlefield. *J Trauma* 2012;73:S490–5.
95. Weiss E. Medical considerations for wilderness and adventure travelers. *Med Clin North Am* 1999;83:885–902.
96. Onifer DJ, Butler FK, Gross KR, et al. Replacement of Phenergan (promethazine) with Zofran (ondansetron) for treatment of opioid and trauma-related nausea and vomiting in Tactical Combat Casualty Care: TCCC guidelines change 14-03. *J Spec Oper Med* 2015;15:17–24.

97. American College of Emergency Physicians Policy Statement. Out of hospital use of analgesia and sedation. *Ann Emerg Med* 2016;67:305–6.
98. Russell KW, Scaife CL, Weber DC, et al. Wilderness Medical Society practice guidelines for the treatment of acute pain in remote environments: 2014 update. *Wilderness Environ Med* 2014;25(Suppl):S96–104.
99. O'Connor K, Butler F. Antibiotics in tactical combat casualty care 2002. *Mil Med* 2003;168:911–4.
100. Hospenthal DR, Murray CK, Anderson RC, et al. Guidelines for prevention of infection after combat-related injuries. *J Trauma* 2008;64(Suppl):S211–20.
101. Battistella FD, Nugent W, Owings JT, et al. Field triage of the pulseless trauma patient. *Arch Surg* 1999;134:742–5.
102. Branney SW, Moore EE, Feldhaus KM, et al. Critical analysis of two decades of experience with post-injury emergency department thoracotomy in a regional trauma center. *J Trauma* 1988;45:87–94.
103. Rosemurgy AS, Norris PA, Olson SM, et al. Prehospital traumatic cardiac arrest: the cost of futility. *J Trauma* 1993;35:468–73.
104. Dickey N, Jenkins D. Needle decompression of tension pneumothorax and cardio-pulmonary resuscitation in Tactical Combat Casualty Care. Defense Health Board Memorandum 2011:1–11. Available at: <http://www.naemt.org/docs/default-source/education-documents/tccc/10-9-15-updates/dhb-memo-120706-needle-decompression.pdf?sfvrsn=2>. Accessed January 19, 2017.
105. Kotwal RS, Butler FK, Montgomery HR, et al. The Tactical Combat Casualty Care Casualty Card. *J Spec Oper Med* 2013;13:82–6.