


Crystal structure of potassium sodium Λ -tris(oxalato)ferrate(III), $K_{2.5}Na_{0.5}Fe(C_2O_4)_3$

R. Wartchow

Universität Hannover, Institut für Anorganische Chemie, Callinstr. 9, D-30167 Hannover, Germany

Received April 4, 1996, transferred to 1st update of database ICSD in 1997, CSD-No. 402454



$C_6FeK_{2.5}Na_{0.5}O_{12}$, cubic, $P4_332$ (No. 212), $a = 13.536(3) \text{ \AA}$, $V = 2480.1 \text{ \AA}^3$, $Z = 8$, $R(F) = 0.028$, $R_w(F^2) = 0.070$, Flack $x = 0.04(5)$.

Table 1. Parameters used for the X-ray data collection

Crystal:	green crystal of irregular shape, size 0.19 x 0.23 x 0.38 mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 $\text{\AA}$)
μ :	20 cm^{-1}
Diffractometer:	Siemens-Stoe
Scan mode:	$\omega/2\theta$
$T_{\text{measurement}}$:	293 K
$2\theta_{\text{max}}$:	49.9°
$N(hkl)_{\text{unique}}$:	739
Criterion for I_o :	$I_o > 2 \sigma(I_o)$
$N(\text{param})_{\text{refined}}$:	68
Program:	SHELXL-93

Source of material: The preparation followed ref. 2, but in the educts 1/6 K was replaced by Na. The product was a mixture of yellow-green crystals of the monoclinic trihydrate and green crystals of both enantiomers of the cubic anhydrous phase. The $Fe(C_2O_4)_3$ group has the Λ configuration. Further programs used: STRUX (Fa.Stoe), PLATON (A.L.Spek)

Table 2. Final atomic coordinates and displacement parameters (in $\text{\AA}^2$)

Atom	Site	x	y	z	U_{11}	U_{22}	U_{33}	U_{12}	U_{13}	U_{23}
Fe(1)	8c	0.98907(3)	x	x	0.0234(2)	U_{11}	U_{11}	-0.0000(2)	U_{12}	U_{12}
K(1)	8c	0.82915(8)	x	x	0.0489(5)	U_{11}	U_{11}	0.0135(5)	U_{12}	U_{12}
K(2)	12d	7/8	0.75693(6)	$y-1/4$	0.0371(7)	0.0310(4)	U_{22}	0.0061(4)	- U_{12}	-0.0006(5)
Na(1)	4b	5/8	5/8	5/8	0.0249(7)	U_{11}	U_{11}	0.0015(8)	U_{12}	U_{12}
C(1)	24e	0.7936(3)	1.0507(3)	1.0200(3)	0.025(2)	0.024(2)	0.027(2)	-0.003(2)	-0.002(2)	0.004(2)
C(2)	24e	0.8588(3)	1.1376(3)	1.0579(2)	0.028(2)	0.023(2)	0.022(2)	-0.001(2)	-0.003(1)	0.003(2)
O(1)	24e	0.8424(2)	0.9743(2)	0.9960(2)	0.024(1)	0.022(1)	0.043(2)	-0.003(1)	0.003(1)	-0.005(1)
O(2)	24e	0.7044(2)	1.0612(2)	1.0143(2)	0.023(2)	0.035(2)	0.056(2)	-0.002(1)	-0.004(1)	0.001(2)
O(3)	24e	0.9524(2)	1.1224(2)	1.0459(2)	0.027(2)	0.024(1)	0.035(2)	-0.002(1)	-0.002(1)	-0.003(1)
O(4)	24e	0.8206(2)	1.2093(2)	1.0955(2)	0.038(2)	0.025(1)	0.041(2)	0.004(1)	0.004(1)	-0.006(1)

Acknowledgments. This work was supported by Prof. Dr. M. Binnewies. Computing facilities of RRZN (Hannover) were used.

References

1. Henneicke, S.; Wartchow, R.: Crystal structure of potassium sodium Λ -tris(oxalato)-ferrate(III), $K_{2.5}Na_{0.5}Fe(C_2O_4)_3$. Z. Kristallogr. **212** (1997) 56.
2. Johnson, R. C: A Convenient Procedure for the Preparation of Potassium Trioxalatoferrate(III). J. Chem. Educ. **47** (1970) 702.
3. Sheldrick, G. M.: SHELXL-93. Program for Crystal Structure Refinement. University of Göttingen, Germany 1993.