

# MAST30025 Linear Statistical Models

## Assignment 3, 2021

- Let  $A$  be an  $n \times p$  matrix with  $n \geq p$ .
  - Show directly that  $r(A^c A) = r(A)$ .
  - Show directly that  $A^c A$  is idempotent.
  - Show directly that  $A(A^T A)^c A^T$  is unique (invariant to the choice of conditional inverse).
- We are interested in examining the lifetime of three different types of bulb (in 1000 hours). A study is conducted and the following data obtained:

Bulb type		
1	2	3
22	16	28
23	18	27
24	19	29
22		29
26		

We fit the model

$$y_{ij} = \mu + \tau_i + \varepsilon_{ij},$$

where  $\mu$  is the overall mean and  $\tau_i$  is the effect of the  $i$ th type of bulb.

**For this question, you may NOT use the `lm` function in R.**

- Find a conditional inverse for  $X^T X$ , using the algorithm given in Theorem 6.2.
  - Find  $s^2$ .
  - Is  $\mu + 2\tau_1 + \tau_2$  estimable?
  - Find a 90% confidence interval for the lifetime of the 2nd type of bulb.
  - Test the hypothesis that there is no difference in lifetime between 1st and 3rd types of bulb, at the 5% significance level.
- Consider a linear model with only categorical predictors, written in matrix form as  $\mathbf{y} = X_1 \boldsymbol{\beta}_1 + \boldsymbol{\varepsilon}_1$ . Suppose we add some continuous predictors, resulting in an expanded model  $\mathbf{y} = X \boldsymbol{\beta} + \boldsymbol{\varepsilon}$ .  
Now consider a quantity  $\mathbf{t}^T \boldsymbol{\beta}$ , where  $\mathbf{t}^T = [\mathbf{t}_1^T | \mathbf{t}_2^T]$  is partitioned according to the categorical and continuous predictors. Show that if  $\mathbf{t}_1^T \boldsymbol{\beta}_1$  is estimable in the first model, then  $\mathbf{t}^T \boldsymbol{\beta}$  is estimable in the second model.

If you write  $X = [X_1 | X_2]$ , you may assume that  $r(X) = r(X_1) + r(X_2)$ .

*Hint: Use Theorems 6.9 and 6.3. For any vectors  $\mathbf{z}_1$  and  $\mathbf{z}_2$ , you can write*

$$\begin{bmatrix} X_1^T X_1 & X_1^T X_2 & X_1^T X_1 \mathbf{z}_1 \\ X_2^T X_1 & X_2^T X_2 & X_2^T X_2 \mathbf{z}_2 \end{bmatrix} = \begin{bmatrix} X_1^T & \mathbf{0} \\ \mathbf{0} & X_2^T \end{bmatrix} \begin{bmatrix} X_1 & X_2 & X_1 \mathbf{z}_1 \\ X_1 & X_2 & X_2 \mathbf{z}_2 \end{bmatrix}.$$

- Data was collected on the world record times (in seconds) for the one-mile run. For males, the records are from the period 1861–1999, and for females, from the period 1967–1996. The data is given in the file `mile.csv`, available on the LMS.
  - Plot the data, using different colours and/or symbols for male and female records. Without drawing diagnostic plots, do you think that this data satisfies the assumptions of the linear model? Why or why not?

- (b) Test the hypothesis that there is no interaction between the two predictor variables. Interpret the result in the context of the study.
  - (c) Write down the final fitted models for the male and female records. Add lines corresponding to these models to your plot from part (a).
  - (d) Calculate a point estimate for the year when the female world record will equal the male world record. Do you expect this estimate to be accurate? Why or why not?
  - (e) Is the year when the female world record will equal the male world record an estimable quantity? Is your answer consistent with part (d)?
  - (f) Calculate a 95% confidence interval for the amount by which the gap between the male and female world records narrow every year.
  - (g) Test the hypothesis that the male world record decreases by 0.4 seconds each year.
5. You wish to perform a study to compare 2 medical treatments (and a placebo) for a disease. Treatment 1 is an experimental new treatment, and costs \$5000 per person. Treatment 2 is a standard treatment, and costs \$2000 per person. Treatment 3 is a placebo, and costs \$1000 per person. You are given \$100,000 to complete the study. You wish to test if the treatments are effective, i.e.,  $H_0 : \tau_1 = \tau_2 = \tau_3$ .
- (a) Determine the optimal allocation of the number of units to assign to each treatment.
  - (b) Perform the random allocation. You must use R for randomisation and include your R commands and output.