

## MAT185 Linear Algebra Assignment 1

### Instructions:

Please read the **MAT185 Assignment Policies & FAQ** document for details on submission policies, collaboration rules and academic integrity, and general instructions.

1. **Submissions are only accepted by Gradescope.** Do not send anything by email. Late submissions are not accepted under any circumstance. Remember you can resubmit anytime before the deadline.
2. **Submit solutions using only this template pdf.** Your submission should be a single pdf with your full written solutions for each question. If your solution is not written using this template pdf (scanned print or digital) then your submission will not be assessed. Organize your work neatly in the space provided. Do not submit rough work.
3. **Show your work and justify your steps** on every question but do not include extraneous information. Put your final answer in the box provided, if necessary. We recommend you write draft solutions on separate pages and afterwards write your polished solutions here on this template.
4. **You must fill out and sign the academic integrity statement below;** otherwise, you will receive zero for this assignment.

### Academic Integrity Statement:

Full Name: \_\_\_\_\_

Student number: \_\_\_\_\_

Full Name: \_\_\_\_\_

Student number: \_\_\_\_\_

I confirm that:

- I have read and followed the policies described in the document **MAT185 Assignment Policies & FAQ**.
- In particular, I have read and understand the rules for collaboration, and permitted resources on assignments as described in subsection II of the the aforementioned document. I have not violated these rules while completing and writing this assignment.
- I understand the consequences of violating the University's academic integrity policies as outlined in the [Code of Behaviour on Academic Matters](#). I have not violated them while completing and writing this assignment.

By signing this document, I agree that the statements above are true.

Signatures: 1) \_\_\_\_\_

2) \_\_\_\_\_

1. A *vector spyce* is a set  $V$  together with two operations called *vector addition* and *scalar multiplication* such that the following nine *axioms* hold:

- (i) For all vectors  $\mathbf{x}, \mathbf{y} \in V$ ,  $\mathbf{x} + \mathbf{y} \in V$
- (ii) For all vectors  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$ ,  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$
- (iii) There exists a vector  $\mathbf{0} \in V$  with the property that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for all vectors  $\mathbf{x} \in V$
- (iv) For each vector  $\mathbf{x} \in V$ , there exists a vector  $-\mathbf{x} \in V$  with the property that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
- (v) For all vectors  $\mathbf{x} \in V$ , and scalars  $c \in \mathbb{R}$ ,  $c\mathbf{x} \in V$
- (vi) For all vectors  $\mathbf{x} \in V$ , and scalars  $c, d \in \mathbb{R}$ ,  $(cd)\mathbf{x} = c(d\mathbf{x})$
- (vii) For all vectors  $\mathbf{x}, \mathbf{y} \in V$ , and scalars  $c \in \mathbb{R}$ ,  $c(\mathbf{x} + \mathbf{y}) = c\mathbf{x} + c\mathbf{y}$
- (viii) For all vectors  $\mathbf{x} \in V$ , and scalars  $c, d \in \mathbb{R}$ ,  $(c + d)\mathbf{x} = c\mathbf{x} + d\mathbf{x}$
- (ix) For all vectors  $\mathbf{x} \in V$ ,  $c\mathbf{x} = \mathbf{0}$  implies that either  $c = 0$  or  $\mathbf{x} = \mathbf{0}$

Prove that a *vector spyce* is a vector space.

Vector spyce axioms  $i$  through  $viii$  equate Medici's vector space axioms  $AI - AIV$  and  $MI - MIII$  where  $vii \Leftrightarrow MIII(b)$  and  $viii \Leftrightarrow MII(a)$ .

Thus, if vector spyce axiom  $ix$  and vector space axiom  $MIV$  can be constructed of the previous 8 (7) axioms; the remaining axioms are identical and a vector spyce will be a vector space by definition.

Vector spyce and Medici vector space axioms, propositions, and theorems will be in the form  $xx$  and  $XX$  respectively

Personal Postulation: let  $v = u \Leftrightarrow u = u$  and by  $i/AI \rightarrow u + a = u + a \Leftrightarrow v + a = u + a$ ; all vector space and vector spyce vectors may be represented by scalar multiples of other existing vectors by  $v/MI$

let $x, y \in \mathcal{V}$ and by axiom $v/MI$ let	$y = 1x$	(1)
Sub	$1x = 1x$	(2)
$\mathbb{R}$	$(1 \cdot 1)x = 1x$	(3)
$vi/MII$	$1(1x) = 1x$	(4)
Sub	$1y = 1x$	(5)
Personal Postulation	$1y + 1(-x) = 1x + 1(-x)$	(6)
$vii/MIII$	$1(y + (-x)) = 1(x + (-x))$	(7)
$iv/AIV$	$1(y + (-x)) = 1(0)$	(8)
$ix$	$1(y + (-x)) = 0$	(9)
$ix$	$1 \neq 0 \implies y + (-x) = 0$	(10)
Personal Postulation	$y + (-x) + x = 0 + x$	(11)
PI: $(-u) + u = 0$	$y + 0 = 0 + x$	(12)
$iii/AIII$	$y = 0 + x$	(13)
PII: $0 + u = u$	$y = x$	(14)
Sub	$1x = x$	(15)

$\therefore$  Axioms  $i$  through  $viii$  equate Medici's axioms  $AI - AIV$  and  $MI - MIII$ ,  $1x = x$  in a vector spyce and vector space, and  $ix$  equates  $PV$ : a proposition constructed of  $MIII(a/b)$ ,  $AII$ ,  $AIII$ , and  $AIV$ ; then vector spyces are constructed of the same axioms as vector spaces and are, as such, vector spaces.

2. Let  $V = \{a_0 + a_1x + a_2x^2 \mid a_0 > 0\}$ . Define vector addition in  $V$  by

$$(a_0 + a_1x + a_2x^2) + (b_0 + b_1x + b_2x^2) = (a_0b_0) + (a_1 + b_1 + 4)x + (a_2 + b_2)x^2$$

and scalar multiplication in  $V$  by

$$c(a_0 + a_1x + a_2x^2) = a_0^c + (ca_1 + 4(c - 1))x + ca_2x^2$$

Verify that axiom MIII. (Medici, pp104) holds in  $V$ .

MIII a:

$$\begin{aligned} (c + d)(a_0 + a_1x + a_2x^2) &= a_0^{c+d} + ((c + d)a_1 + 4(c + d - 1))x + (c + d)a_2x^2 \\ &= a_0^c a_0^d + (ca_1 + da_1 + 4c + 4d - 4)x + (ca_2 + da_2)x^2 \\ &= a_0^c a_0^d + (ca_1 + 4c - 4 + da_1 + 4d - 4 + 4)x + (ca_2 + da_2)x^2 \\ &= a_0^c a_0^d + (ca_1 + 4(c - 1) + da_1 + 4(d - 1) + 4)x + (ca_2 + da_2)x^2 \\ &= (a_0^c + (ca_1 + 4(c - 1))x + ca_2x^2) + (a_0^d + (da_1 + 4(d - 1))x + da_2x^2) \\ &= c(a_0 + a_1x + a_2x^2) + d(a_0 + a_1x + a_2x^2) \end{aligned}$$

MIII b:

$$\begin{aligned} c((a_0 + a_1x + a_2x^2) + (b_0 + b_1x + b_2x^2)) &= c((a_0b_0) + (a_1 + b_1 + 4)x + (a_2 + b_2)x^2) \\ &= (a_0b_0)^c + (c(a_1 + b_1 + 4) + 4(c - 1))x + c(a_2 + b_2)x^2 \\ &= (a_0^c b_0^c) + (ca_1 + cb_1 + 4c + 4c - 4)x + (ca_2 + cb_2)x^2 \\ &= (a_0^c b_0^c) + (ca_1 + 4c - 4 + cb_1 + 4c - 4 + 4)x + (ca_2 + cb_2)x^2 \\ &= (a_0^c b_0^c) + (ca_1 + 4(c - 1) + cb_1 + 4(c - 1) + 4)x + (ca_2 + cb_2)x^2 \\ &= (a_0^c + (ca_1 + 4(c - 1))x + ca_2x^2) + (b_0^c + (cb_1 + 4(c - 1))x + cb_2x^2) \\ &= c(a_0 + a_1x + a_2x^2) + c(b_0 + b_1x + b_2x^2) \end{aligned}$$

3. We asked ChatGPT if  $V = \{f \mid f' \text{ is constant}\}$  is a vector space. It's response is given below.



**You**

Is  $V = \{f \mid f' \text{ is constant}\}$  a vector space?



**ChatGPT**

To determine if the set  $V = \{f \mid f' \text{ is constant}\}$  is a vector space, we need to check the three vector space properties:

1. **Zero vector:** The constant function  $f(x) = c$  has a derivative of  $f'(x) = 0$ , which is a constant. Therefore, the zero function is in  $V$ .
2. **Closure under addition:** Let  $f$  and  $g$  be functions in  $V$  with constant derivatives. Consider the function  $h(x) = f(x) + g(x)$ . The derivative of  $h$  is the sum of the derivatives of  $f$  and  $g$ , which are constants. So,  $h$  also has a constant derivative, and  $V$  is closed under addition.
3. **Closure under scalar multiplication:** Let  $f$  be a function in  $V$  with a constant derivative. If we multiply  $f$  by a scalar  $c$ , the derivative of  $cf$  is  $cf'$ , which is still a constant. Thus,  $V$  is closed under scalar multiplication.

Since  $V$  satisfies all three vector space properties, it is indeed a vector space.



Identify *two* errors in ChatGPT's response and write *two* paragraphs, one for each error, each of which clearly identifies an error, and how you would correct them.

The first error in ChatGPT's response was that it used the three conditions of the subspace test by assuming that  $V$  is a subset of a vector space, but it did not without stating the reasoning behind this assumption. To correct this error, one could first state the reason that since  $f'$  is constant, then  $f$  would be in the form  $f = ax + b, \forall a, b \in \mathbb{R}$ . Therefore  $V \subset \mathbb{F}(\mathbb{R})$ , or  $V$  is a subset of  $\mathbb{F}(\mathbb{R})$ .

The second error in ChatGPT's response was that it wrongly determined the zero vector in  $V$ . The zero vector in  $V$  should be  $f(x) = 0$ , but ChatGPT determined that the zero vector in  $V$  is  $f(x) = c, c \in \mathbb{R}$ . To correct this error, one could first state that the zero vector in  $V$  is  $f(x) = 0$ , and then prove that  $f(x) = 0$  is in  $V$  by stating  $f(x) = 0 \Rightarrow f'(x) = 0$ .

4. Let  $c \in \mathbb{R}$ , and consider the subset  $W_c = \{(x, y, z) \in \mathbb{R}^3 \mid x + y + z = c|z|\}$  of  $\mathbb{R}^3$ . Determine *all* values of  $c$  such that  $W$  is a subspace of  $\mathbb{R}^3$ . Your answer should clearly demonstrate that you've found *all* values of  $c$  such that  $W_c$  is a subspace. This includes demonstrating why  $W_c$  is not a subspace for certain values of  $c$ , if you think no such  $c$  exists.

Since  $W_c$  is a subset of  $\mathbb{R}^3$  and  $\mathbb{R}^3$  is a vector space,  $W_c$  is a subspace of  $\mathbb{R}^3$  if  $W_c$  satisfies the conditions of the subspace test ( $W_c$  is non-empty and contains the zero vector,  $W_c$  is closed under vector addition and scalar multiplication).

The zero vector:

$$\begin{aligned}\forall c \in \mathbb{R}, 0 + 0 + 0 &= c|0| = 0 \\ \forall x \in W_c, x + 0 &= (x_1, x_2, x_3) + (0, 0, 0) = (x_1 + x_2 + x_3) = x \\ \therefore (0, 0, 0) &\in W_c\end{aligned}$$

Therefore  $W_c$  is non-empty and contains the zero vector for all  $c$ .

Inverse: Since the inverse of a vector  $-v = (-1)v$ , the inverse of a vector should also exist in a vector space due to closure of scalar multiplication.

$$\begin{aligned}\forall x = (x_1, x_2, x_3) \in W_c, \exists -x &= (-x_1, -x_2, -x_3) \in W_c \mid x + (-x) = 0 \\ (x_1 + (-x_1), x_2 + (-x_2), x_3 + (-x_3)) &= 0 \\ \therefore x \in W_c, x_1 + x_2 + x_3 &= c|x_3| \\ \therefore -x \in W_c, -x_1 + -x_2 + -x_3 &= c|-x_3| = c|x_3| \\ \therefore x_1 + x_2 + x_3 &= -x_1 + -x_2 + -x_3 \\ x_1 + x_2 + x_3 &= -(x_1 + x_2 + x_3) \\ \therefore x_1 + x_2 + x_3 &= 0 \\ \therefore c|x_3| &= 0 \\ \therefore \forall x_3 \in \mathbb{R} \\ \therefore c &= 0\end{aligned}$$

Therefore only  $c = 0$  satisfies the inverse condition. If  $c \neq 0$ , then  $x_1 + x_2 + x_3 \neq -x_1 + -x_2 + -x_3$ .

Since only  $c = 0$  satisfies the inverse condition, only  $W_0$  will be checked for closure under vector addition and scalar multiplication.

$$\begin{aligned}\forall x = (x_1, x_2, x_3), y = (y_1, y_2, y_3) \in W_0, \forall c \in \mathbb{R} \\ cx + y &= (cx_1 + y_1, cx_2 + y_2, cx_3 + y_3) \\ (cx_1 + y_1) + (cx_2 + y_2) + (cx_3 + y_3) &= cx_1 + cx_2 + cx_3 + y_1 + y_2 + y_3 \\ &= c(x_1 + x_2 + x_3) + (y_1 + y_2 + y_3) \\ \therefore x \in W_0, x_1 + x_2 + x_3 &= 0 \\ \therefore y \in W_0, y_1 + y_2 + y_3 &= 0 \\ \therefore (cx_1 + y_1) + (cx_2 + y_2) + (cx_3 + y_3) &= c(0) + (0) = 0 \\ \therefore cx + y &\in W_0\end{aligned}$$

Therefore  $W_0$  is closed under vector addition and scalar multiplication.

Therefore  $W_c$  is a subspace of  $\mathbb{R}^3$  if and only if  $c = 0$ .