

The Galois Correspondence

Jiang-Hua Lu

The University of Hong Kong

MATH4302, Algebra II

Monday April 28, 2025

In this file: §4.1.5 of lecture notes.

① The main theorem of Galois theory: the Galois Correspondence

Recall

Theorem

For a finite extension $K \subset L$ with $G = \text{Aut}_K(L)$, the following are equivalent:

- ① $K \subset L$ is Galois, i.e., $|G| = [L : K]$;
- ② $K = L^G$;
- ③ The extension $K \subset L$ is normal and separable;
- ④ L is a splitting field over K of some separable polynomial in $K[x]$.

§4.1.5: The Galois correspondence

Let $K \subset L$ be a field extension, and let $G = \text{Aut}_K(L)$.

Definition-Lemma.

- A subfield M of L containing K is called an **intermediate field** of $K \subset L$, and denoted as $K \subset M \subset L$.
- For any intermediate field $K \subset M \subset L$, $\text{Aut}_M(L)$ is a subgroup of G ;
- For any subgroup H of G ,

$$L^H \stackrel{\text{def}}{=} \{a \in L : \sigma(a) = a, \forall \sigma \in H\}$$

is an intermediate field of $K \subset L$, called the **fixed field** of H .

$$\{K \subset \underline{M} \subset L\} \longleftrightarrow \{ \underline{H} \subset G \}$$

§4.1.5: The Galois correspondence

For a field extension $K \subset L$ and $G = \text{Aut}_K(L)$, have

$$\{\text{intermediate fields } K \subset M \subset L\} \begin{matrix} \xrightarrow{\Gamma} \\ \xleftarrow{F} \end{matrix} \{\text{subgroups } H \subset G\},$$

$$\Gamma(M) = \text{Aut}_M(L) \quad \text{and} \quad F(H) = L^H = \{a \in L : \sigma(a) = a, \forall \sigma \in H\}.$$

Lemma. Both F and Γ are **inclusion reversing**:

$$M_1 \subset M_2 \implies \Gamma(M_1) \supset \Gamma(M_2),$$

$$H_1 \subset H_2 \implies F(H_1) \supset F(H_2).$$

Moreover, for all intermediate field M and subgroup H of G , one has

$$M \subset F(\Gamma(M)), \quad H \subset \Gamma(F(H)).$$

If H is a finite subgroup of G , Artin's Theorem gives $H = \Gamma(F(H))$

Proof.

Theorem (Fundamental Theorem of Galois Theory)

Let $K \subset L$ be a finite Galois extension. Then

- 1 for any intermediate $K \subset M \subset L$, the extension $M \subset L$ is Galois;
- 2 the two maps Γ and F are inverses of each other;
- 3 for an intermediate field $K \subset M \subset L$, the extension $K \subset M$ is Galois if and only if $\Gamma(M) = \text{Aut}_M(L)$ is a normal subgroup of $\text{Aut}_K(L)$, and in this case,

$$\text{Aut}_K(M) = \text{Aut}_K(L) / \text{Aut}_M(L).$$

The correspondence between intermediate fields $K \subset M \subset L$ and subgroups of G is called the [The Galois Correspondence](#).

Proof of Fundamental Thm. of Galois Theory. Let $K \subset L$ be finite Galois.

- ① L is splitting field over K of a separable $f \in K[x]$. Then L is also a splitting field over M of the separable $f \in M[x]$. Thus $M \subset L$ is Galois.
- ② Already know that $H = \Gamma(F(H))$ by Artin's Theorem. For any intermediate $K \subset M \subset L$, by (i), $M \subset L$ is Galois, so $M = F(\Gamma(M))$.
- ③ Assume first that $K \subset M$ is Galois.
 - M is the splitting field of some $g(x) \in K[x]$, so $M = K(R_g)$.
 - $\sigma(R_g) = R_g$ for every $\sigma \in \text{Aut}_K(L)$, so $\sigma(M) = M$. Thus have the group homomorphism

$$\phi : \text{Aut}_K(L) \longrightarrow \text{Aut}_K(M), \sigma \longmapsto \sigma|_M$$

with $\ker \phi = \text{Aut}_M(L)$. Thus $\text{Aut}_M(L)$ is a normal subgroup of $\text{Aut}_K(L)$.

Proof continued:

- Have injective group homomorphism

$$[\phi] : \text{Aut}_K(L)/\text{Aut}_M(L) \longrightarrow \text{Aut}_K(M).$$

- As both $K \subset L$ and $K \subset M$ are Galois,

$$\begin{aligned} |\text{Aut}_K(L)/\text{Aut}_M(L)| &= \frac{|\text{Aut}_K(L)|}{|\text{Aut}_M(L)|} = \frac{[L : K]}{[L : M]} = [M : K] \\ &= |\text{Aut}_K(M)|. \end{aligned}$$

Thus $[\phi]$ is a group isomorphism.

Assume now that $\text{Aut}_M(L)$ is a normal subgroup of $\text{Aut}_K(L)$.

- Let $\sigma \in G$. For any $a \in M$ and $\tau \in \text{Aut}_M(L)$, have

$$\sigma^{-1}\tau\sigma \in \text{Aut}_M(L),$$

so $(\sigma^{-1}\tau\sigma)(a) = a$, i.e., $\tau(\sigma(a)) = \sigma(a)$, so $\sigma(a) \in F(\Gamma(M))$. By (ii), $F(\Gamma(M)) = M$, so $\sigma(a) \in M$.

Proof continued:

- Thus again have group homomorphism

$$\phi : \text{Aut}_K(L) \longrightarrow \text{Aut}_K(M), \sigma \longmapsto \sigma|_M,$$

and injective group homomorphism

$$[\phi] : \text{Aut}_K(L)/\text{Aut}_M(L) \longrightarrow \text{Aut}_K(M).$$

- Since $|\text{Aut}_K(M)| \leq [M : K]$, one has $|\text{Aut}_K(M)| = [M : K]$. Thus $K \subset M$ is Galois.
- End of proof.

Corollary

A finite Galois extension $K \subset L$ has finitely many intermediate subfields.

Example. $\mathbb{F}_p \subset \mathbb{F}_{p^n}$:

- $\mathbb{F}_p \subset \mathbb{F}_{p^n}$ is Galois because \mathbb{F}_{p^n} is a splitting field of $x^{p^n} - x \in \mathbb{F}_p[x]$;
- $G = \text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p) \cong \mathbb{Z}/n\mathbb{Z}$, generated by the Frobenius isomorphism $\sigma : a \mapsto a^p$.
- One subgroup of G of order m for each $m|n$, generated by σ^d , where $d = n/m$.
- The fixed field of $\langle \sigma^d \rangle$ is the subfield \mathbb{F}_{p^d} of \mathbb{F}_{p^n} .

Example. $L = \mathbb{Q}(\sqrt[3]{2}, \omega)$, where $\omega = e^{(2\pi i)/3}$.

- Know that $|G = \text{Aut}(\mathbb{Q})(L)| = [L : \mathbb{Q}] = 6$.
- f has exactly three roots, namely

$$r_1 = \sqrt[3]{2}, \quad r_2 = \omega\sqrt[3]{2}, \quad r_3 = \omega^2\sqrt[3]{2},$$

so $G \cong S_3$, permutation group of the three roots.

- Every $g \in G$ must satisfy

$$g(\omega) \in \{\omega, \omega^2\}, \quad g(\sqrt[3]{2}) \in \{r_1, r_2, r_3\}.$$

- Define $\sigma, \tau \in G$ by

$$\sigma(\omega) = \omega, \quad \sigma(\sqrt[3]{2}) = \omega\sqrt[3]{2}, \quad \tau(\omega) = \omega^2, \quad \tau(\sqrt[3]{2}) = \sqrt[3]{2}.$$

Then $\sigma^3 = \tau^2 = \text{Id}$, and

$$G = \{e, \sigma, \sigma^2, \tau, \tau\sigma, \sigma\tau = \tau\sigma^2\}.$$

Among the 6 intermediate fields:

- the extensions

$$\mathbb{Q} \subset \mathbb{Q}, \quad \mathbb{Q} \subset \mathbb{Q}(\omega), \quad \mathbb{Q} \subset L = \mathbb{Q}(\omega, \sqrt[3]{2})$$

are Galois, corresponding to the three normal subgroups

$$\{e\}, \quad \{e, \sigma, \sigma^2\}, \quad G;$$

- the other three extensions

$$\mathbb{Q} \subset \mathbb{Q}(\sqrt[3]{2}), \quad \mathbb{Q} \subset \mathbb{Q}(\omega\sqrt[3]{2}), \quad \mathbb{Q} \subset \mathbb{Q}(\omega^2\sqrt[3]{2})$$

are not Galois.

§4.1.5: The Galois correspondence

