

Homework 1

21-470 Calculus of Variations

Name: Shashank Singh¹

Due: Wednesday, January 29, 2014

Problem 1

[I wasn't able to get very far with the integrals in parts (b) and (c), and wasn't able to determine much about the constants in part (d). Based on numerical computation, I believe (a) offers the slowest solution, followed by (b), (c), and then (d).]

(a) Since $y'(x) = 1$ for all $x \in [0, 1]$,

$$J(y) = \int_0^1 \frac{\sqrt{1 + y'(x)^2}}{\sqrt{y(x)}} dx = \sqrt{2} \int_0^1 x^{-1/2} dx = 2\sqrt{2}x^{1/2} \Big|_{x=0}^{x=1} = 2\sqrt{2}.$$

(b) Since

$$y(x) = \sqrt{x - x^2} \quad \text{and} \quad y'(x) = \frac{1 - x}{\sqrt{2x - x^2}}$$

for all $x \in (0, 1]$,

$$\begin{aligned} J(y) &= \int_0^1 \frac{\sqrt{1 + (1 - x)^2/(2x - x^2)}}{(2x - x^2)^{1/4}} dx = \int_0^1 \frac{\sqrt{2x - x^2 + x^2 - 2x + 1}}{(2x - x^2)^{3/4}} dx \\ &= \int_0^1 (2x - x^2)^{-3/4} dx. \end{aligned}$$

(c) Since $y'(x) = x^{-1/2}/2$ for all $x \in (0, 1]$, using the inequality $\sqrt{a + b} \leq \sqrt{a} + \sqrt{b}$ for $a, b \geq 0$,

$$J(y) = \int_0^1 \frac{\sqrt{1 + 1/(4x)}}{x^{1/4}} dx = \int_0^1 \sqrt{x^{-1/2} + x^{-3/2}/4} dx.$$

(d) Observing that

$$\frac{dx}{d\theta} = \frac{c^2}{2}(1 - \cos \theta) \quad \text{and} \quad \frac{dy}{dx} = \frac{dy}{d\theta} \Big/ \frac{dx}{d\theta} = \frac{\sin \theta}{1 - \cos \theta},$$

so that

$$\left(\frac{dy}{dx}\right)^2 = \left(\frac{\sin \theta}{1 - \cos \theta}\right)^2 = \frac{1 - \cos^2 \theta}{(1 - \cos \theta)^2} = \frac{1 + \cos \theta}{1 - \cos \theta},$$

and changing variables from x to t , $J(y)$ simplifies significantly:

$$\begin{aligned} J(y) &= \int_0^{\theta_1} \frac{\sqrt{1 + \frac{1 + \cos \theta}{1 - \cos \theta}}}{\sqrt{\frac{c^2}{2}(1 - \cos \theta)}} \frac{c^2}{2}(1 - \cos \theta) d\theta = \frac{c}{\sqrt{2}} \int_0^{\theta_1} \sqrt{1 + \frac{1 + \cos \theta}{1 - \cos \theta}} \sqrt{1 - \cos \theta} d\theta \\ &= \frac{c}{\sqrt{2}} \int_0^{\theta_1} \sqrt{1 - \cos \theta + 1 + \cos \theta} d\theta = \frac{c}{\sqrt{2}} \int_0^{\theta_1} \sqrt{2} d\theta = c\theta_1. \end{aligned}$$

¹sssl@andrew.cmu.edu

Problem 2

Since

$$\frac{d^2}{dx^2} \sqrt{1+x^2} = \frac{d}{dx} \frac{x}{\sqrt{1+x^2}} = (1+x^2)^{-3/2} \geq 0,$$

the function $x \mapsto \sqrt{1+x^2}$ is convex. Thus, by Jensen's Inequality and a linear change of variables

$$\begin{aligned} J(y) &= \int_a^b \sqrt{1+y'(x)^2} dx = (b-a) \int_0^1 \sqrt{1+y'((b-a)u+a)^2} du && \text{(Change of Variables)} \\ &\geq (b-a) \sqrt{1 + \left(\int_0^1 y'((b-a)u+a) du \right)^2} && \text{(Jensen's Inequality)} \\ &= \sqrt{(b-a)^2 + \left(\int_a^b y'(x) dx \right)^2} && \text{(Change of Variables)} \\ &= \sqrt{(b-a)^2 + (B-A)^2}, \end{aligned}$$

where the last step uses that $\int_a^b y'(x) dx = B - A$ by the Fundamental Theorem of Calculus.

Problem 3

(a) The bound represents the ‘vertical component’ of the surface area:

$$\begin{aligned} J(y) &= 2\pi \int_0^b y(x) \sqrt{1+y'(x)^2} dx \geq 2\pi \int_0^b y(x) \sqrt{y'(x)^2} dx \\ &\geq \pi \int_0^b 2y(x)y'(x) dx = \pi y(x)^2 \Big|_{x=0}^{x=b} = \pi B^2, \end{aligned}$$

since $y \in \mathcal{Y}$ is non-negative. The bound essentially states that the minimal surface area is at least the surface area of the disc of radius B centered at $(b, 0)$, normal to the x -axis. ■

(b) If, for any $x \in [0, b]$, $y(x) > 0$, then $y(x)\sqrt{1+y'(x)^2} > y(x)\sqrt{y'(x)^2}$. Since y and y' are continuous, it follows that the first inequality in part (a) above can be made strict, so that no $y \in \mathcal{Y}$ achieves $J(y) = \pi B^2$. ■

(c) There is indeed such a sequence. For $n \in \mathbb{N}$, define $y_n \in \mathcal{Y}$ by $y_n(x) = B(x/b)^n, \forall x \in [0, b]$. Then, using the inequality $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ for $a, b \geq 0$,

$$\begin{aligned} J(y_n) &= 2\pi \int_0^b y_n(x) \sqrt{1+y_n'(x)^2} dx = 2\pi \int_0^b B(x/b)^n \sqrt{1 + (Bnx^{n-1}/b^n)^2} dx \\ &\leq 2\pi B \int_0^b (x/b)^n (1 + Bnx^{n-1}/b^n) dx \\ &= 2\pi B \left(\frac{b}{n+1} + \frac{B}{2} \right) \rightarrow \pi B^2 \end{aligned}$$

as $n \rightarrow \infty$. By the result of part (a), it follows that $J(y_n) \rightarrow \pi B^2$ as $n \rightarrow \infty$. ■