

Math 720: Homework.

Do, but don't turn in optional problems. *There is a firm 'no late homework' policy.*

Assignment 1: Assigned Wed 08/28. Due Wed 09/04

Keep in mind there is a firm "no late homework" policy. Starred problems are optional; but I'd recommend looking at them. They often involve results I will use later in class.

- Let μ be a positive measure on (X, Σ) .
 - If $A_i \in \Sigma$ are such that $A_i \subseteq A_{i+1}$, show that $\mu(\bigcup_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$.
 - If $A_i \in \Sigma$ are such that $A_i \supseteq A_{i+1}$, show that $\mu(\bigcap_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$, provided $\mu(A_1) < \infty$. Show by example this is false true if $\mu(A_1) = \infty$.
- Prove any open subset of \mathbb{R}^d is a countable union of cells.
- For each of the following sets, compute the Lebesgue outer measure.
 - Any countable set.
 - The Cantor set.
 - $\{x \in [0, 1] \mid x \notin \mathbb{Q}\}$.
- If $V \subseteq \mathbb{R}^d$ is a subspace with $\dim(V) < d$, then show that $\lambda^*(V) = 0$.
 - If $P \subseteq \mathbb{R}^2$ is a polygon show that $\text{area}(P) = \lambda^*(P)$.
- Does there exist a σ -algebra whose cardinality is countably infinite? Disprove, or find an example.

Optional problems, and details in class I left for you to check.

- * Define $\mu(A)$ to be the number of elements in A . Show that μ is a measure on $(X, \mathcal{P}(X))$. (This is called the counting measure.)
- * Let $x_0 \in X$ be fixed. Define $\delta_{x_0}(A) = 1$ if $x_0 \in A$ and 0 otherwise. Show that δ_{x_0} is a measure on $(X, \mathcal{P}(X))$. (This is called the delta measure at x_0 .)
- * Show that $\lambda^*(a + E) = \lambda^*(E)$ for all $a \in \mathbb{R}^d$, $E \subseteq \mathbb{R}^d$.
- * Show that $\lambda^*(I) = \ell(I)$ for all cells. (I only proved it for closed cells in class.)
- * Show that $\mathcal{B}(\mathbb{R})$ has the same cardinality as \mathbb{R} .
- * (*Challenge*) Suppose $f_n : [0, 1] \rightarrow [0, 1]$ are all Riemann integrable, $0 \leq f_n \leq 1$ and $(f_n) \rightarrow 0$ pointwise. Show that $\lim_{n \rightarrow \infty} \int_0^1 f_n = 0$, using only standard tools from Riemann integration.

Assignment 2: Assigned Wed 09/04. Due Wed 09/11

- Say μ is a *translation invariant* measure on $(\mathbb{R}^d, \mathcal{L})$ (i.e. $\mu(x + A) = \mu(A)$ for all $A \in \mathcal{L}$, $x \in \mathbb{R}^d$) which is finite on bounded sets. Show that $\exists c \geq 0$ such that $\mu(A) = c\lambda(A)$.
 - Let $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a linear transformation, and $A \in \mathcal{L}$. Show that $T(A) \in \mathcal{L}$ and $\lambda(T(A)) = |\det(T)|\lambda(A)$. [HINT: Express T in terms of elementary transformations.]
- Let $\mathcal{E} \subseteq \mathcal{P}(X)$, and $\rho : \mathcal{E} \rightarrow [0, \infty]$ be such that $\emptyset \in \mathcal{E}$, $X \in \mathcal{E}$ and $\rho(\emptyset) = 0$. For any $A \subseteq X$ define

$$\mu^*(A) = \inf \left\{ \sum_1^{\infty} \rho(E_i) \mid E_i \in \mathcal{E}, \text{ and } A \subseteq \bigcup_1^{\infty} E_i \right\}.$$

Show that μ^* is an outer measure.

- Let (X, d) be any metric space, $\delta > 0$, $\alpha \geq 0$ and define

$$\mathcal{E}_{\delta} = \{A \subseteq X \mid \text{diam}(A) < \delta\} \quad \text{and} \quad \rho_{\alpha}(A) = \frac{\pi^{\alpha/2}}{\Gamma(1 + \frac{\alpha}{2})} \left(\frac{\text{diam}(A)}{2} \right)^{\alpha}.$$

Let $H_{\alpha, \delta}^*$ be the outer measure obtained with $\rho = \rho_{\alpha}$ and the collection of sets \mathcal{E}_{δ} . Define $H_{\alpha}^* = \lim_{\delta \rightarrow 0} H_{\alpha, \delta}^*$. Show H_{α}^* is an outer measure and restricts to a measure H_{α} on a σ -algebra that contains all Borel sets. The measure H_{α} is called the *Hausdorff measure of dimension α* .

- If $X = \mathbb{R}^d$, and $\alpha = d$ show that H_d is a non-zero, finite constant multiple of the Lebesgue measure. [In fact $H_d = \lambda$ because of our choice of normalization constant, but the proof is much harder.]
 - Let $S \in \mathcal{B}(X)$. Show that there exists (a unique) $d \in [0, \infty]$ such that $H_{\alpha}(S) = \infty$ for all $\alpha \in (0, d)$, and $H_{\alpha}(S) = 0$ for all $\alpha \in (d, \infty)$. This number is called the *Hausdorff dimension* of the set S .
 - Compute the Hausdorff dimension of the Cantor set.
- Using notation from the previous question, let $\mathcal{S}_{\delta} = \{B(x, r) \mid x \in X, r \in (0, \delta)\}$. Using the collection of sets \mathcal{S}_{δ} and the function $\rho = \rho_{\alpha}$, we obtain an outer measure $S_{\alpha, \delta}^*$. As before one can show that $S_{\alpha}^* = \lim_{\delta \rightarrow 0} S_{\alpha, \delta}^*$ is an outer measure, and gives a Borel measure S_{α} .
 - Show by example $S_{\alpha} \neq H_{\alpha}$ in general.
 - If $X = \mathbb{R}^d$ with the standard metric show that $S_d = \lambda$. [You may assume $\rho_d(B_r) = \lambda(B_r)$.]

Details in class I left for you to check. (Do it, but don't turn it in.)

- * Using notation from the proof of Caratheodory, show that $\mu^*(A \cap (\bigcup_1^{\infty} E_i)) = \sum_1^{\infty} \mu^*(A \cap E_i)$. [We only proved it for $A = X$ in class.]

Assignment 3: Assigned Wed 09/11. Due Wed 09/18

- Let μ, ν be two measures on (X, Σ) . Suppose $\mathcal{C} \subseteq \Sigma$ is a π -system such that $\mu = \nu$ on \mathcal{C} .
 - Suppose $\exists C_i \in \mathcal{C}$ such that $\bigcup_1^\infty C_i = X$ and $\mu(C_i) = \nu(C_i) < \infty$. Show that $\mu = \nu$ on $\sigma(\mathcal{C})$.
 - If we drop the finiteness condition $\mu(C_i) < \infty$ is the previous subpart still true? Prove or find a counter example.
- Let X be a metric space and μ a Borel measure on X . Suppose there exists a sequence of sets $B_n \subseteq X$ such that $\bar{B}_n \subseteq \overset{\circ}{B}_{n+1}$, \bar{B}_n is compact, $X = \bigcup_1^\infty B_n$ and $\mu(B_n) < \infty$. Show that μ is regular.
 - Show directly that for all $A \in \mathcal{L}$, $\lambda(A) = \sup\{\lambda(K)\}$ where $K \subseteq A$ is compact, and $\lambda(A) = \inf\{\lambda(U)\}$ where $U \supseteq A$ is open. [Note: The previous subpart will *only* show this for all $A \in \mathcal{B}(\mathbb{R}^d)$.]
- Find $E \in \mathcal{B}(\mathbb{R})$ so that for all $a < b$, we have $0 < \lambda(E \cap (a, b)) < b - a$.
 - Let $\kappa \in (0, 1/2)$. Does there exist $E \in \mathcal{B}(\mathbb{R})$ such that for all $a < b \in \mathbb{R}$, we have $\kappa(b - a) \leq \lambda(E \cap (a, b)) \leq (1 - \kappa)(b - a)$? Prove it.
- Let $A \in \mathcal{L}(\mathbb{R}^d)$. Prove every subset of A is Lebesgue measurable $\iff \lambda(A) = 0$.
- Prove $\mathcal{B}(\mathbb{R}^{m+n}) = \sigma(\{A \times B \mid A \in \mathcal{B}(\mathbb{R}^m) \& B \in \mathcal{B}(\mathbb{R}^n)\})$.
 - Prove $\mathcal{L}(\mathbb{R}^{m+n}) \supsetneq \sigma(\{A \times B \mid A \in \mathcal{L}(\mathbb{R}^m) \& B \in \mathcal{L}(\mathbb{R}^n)\})$.
 - Show $\mathcal{L}(\mathbb{R}^2) \supsetneq \mathcal{B}(\mathbb{R}^2)$.

Optional problems, and details in class I left for you to check.

- * Let μ be a finite Borel measure on a compact metric space. Let

$$\mathcal{C} = \{A \mid \sup_{\substack{K \subseteq A \\ K \text{ compact}}} \mu(K) = \mu(A) = \inf_{\substack{U \supseteq A \\ U \text{ open}}} \mu(U)\}.$$

We saw in class that \mathcal{C} is closed under countable increasing unions. Show \mathcal{C} is closed under relative compliments.

- * Is any σ -finite Borel measure on \mathbb{R}^d regular?
- * Show that there exists $A \subseteq \mathbb{R}$ such that if $B \subseteq A$ and $B \in \mathcal{L}$ then $\lambda(B) = 0$, and further, if $B \subseteq A^c$ and $B \in \mathcal{L}$ then $\lambda(B) = 0$.

We say $\mathcal{A} \subseteq \mathcal{P}(X)$ is an algebra if $\emptyset \in \mathcal{A}$, and \mathcal{A} is closed under complements and finite unions. We say $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ is a (positive) *pre-measure* on \mathcal{A} if $\mu_0(\emptyset) = 0$, and for any countable disjoint sequence of sets sequence $A_i \in \mathcal{A}$ such that $\bigcup_1^\infty A_i \in \mathcal{A}$, we have $\mu_0(\bigcup_1^\infty A_i) = \sum_1^\infty \mu_0(A_i)$.

Namely, a pre-measure is a finitely additive measure on an algebra \mathcal{A} , which is also countably additive for disjoint unions *that belong to the algebra*.

- * (*Caratheodory extension*) If \mathcal{A} is an algebra, and μ_0 is a pre-measure on \mathcal{A} , show that there exists a measure μ defined on $\sigma(\mathcal{A})$ that extends μ_0 .

Assignment 4: Assigned Wed 09/18. Due Wed 09/25

- Let $C \subseteq \mathbb{R}^d$ be convex. Must C be Lebesgue measurable? Must C be Borel measurable? Prove or find counter examples. [The cases $d = 1$ and $d > 1$ are different.]
- Let (X, Σ, μ) be a measure space. For $A \in \mathcal{P}(X)$ define $\mu^*(A) = \inf\{\mu(E) \mid E \supseteq A \& E \in \Sigma\}$, and $\mu_*(A) = \sup\{\mu(E) \mid E \subseteq A \& E \in \Sigma\}$.
 - Show that μ^* is an outer measure.
 - Let $A_1, A_2, \dots \in \mathcal{P}(X)$ be disjoint. Show that $\mu_*(\bigcup_1^\infty A_i) \geq \sum_1^\infty \mu_*(A_i)$. [The set function μ_* is called an *inner measure*.]
 - Show that for all $A \subseteq X$, $\mu^*(A) + \mu_*(A^c) = \mu(X)$.
 - Let $A \subseteq \mathcal{P}(X)$ with $\mu^*(A) < \infty$. Show that $A \in \Sigma_\mu \iff \mu_*(A) = \mu^*(A)$.
- Let $f : X \rightarrow \mathbb{R}$ be measurable, and $g : \mathbb{R} \rightarrow \mathbb{R}$ be Lebesgue measurable. True or false: $g \circ f : X \rightarrow \mathbb{R}$ is measurable? Prove or find a counter example.
- Let (X, Σ) be a measure space, and $f, g : X \rightarrow [-\infty, \infty]$ be measurable. Suppose whenever $g = 0$, $f \neq 0$, and whenever $f = \pm\infty$, $g \in (-\infty, \infty)$. Show that $\frac{f}{g} : X \rightarrow [-\infty, \infty]$ is measurable. [Note that by the given data you will never get a ‘meaningless’ quotient of the form $\frac{0}{0}$ or $\frac{\pm\infty}{\pm\infty}$. The remainder of the quotients (e.g. $\frac{1}{\infty}$) can be defined in the natural manner.]
- Let $f_n : X \rightarrow \mathbb{R}$ be a sequence of measurable functions such that $(f_n) \rightarrow f$ almost everywhere (a.e.). Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function.
 - If for a.e. $x \in X$, g is continuous at $f(x)$, then show $(g \circ f_n) \rightarrow g \circ f$ a.e.
 - Is the previous part true without the continuity assumption on g ?

Optional problems, and details in class I left for you to check.

- * (*An alternate approach to λ -systems.*) Let $\mathcal{M} \subseteq \mathcal{P}(X)$. We say \mathcal{M} is a *Monotone Class*, if whenever $A_i, B_i \in \mathcal{M}$ with $A_i \subseteq A_{i+1}$ and $B_i \supseteq B_{i+1}$ then $\bigcup_1^\infty A_i \in \mathcal{M}$ and $\bigcap_1^\infty B_i \in \mathcal{M}$. If $\mathcal{A} \subseteq \mathcal{P}(X)$ is an algebra, then show that the *smallest* monotone class containing \mathcal{A} is exactly $\sigma(\mathcal{A})$. [You should also address existence of a smallest monotone class containing \mathcal{A} .]
- * Prove that the completion Σ_μ we defined in class is the smallest μ -complete σ -algebra that contains Σ .
- * Show that $f : X \rightarrow [-\infty, \infty]$ is measurable if and only if any of the following conditions hold

(a) $\{f < a\} \in \Sigma$ for all $a \in \mathbb{R}$.	(c) $\{f \leq a\} \in \Sigma$ for all $a \in \mathbb{R}$.
(b) $\{f > a\} \in \Sigma$ for all $a \in \mathbb{R}$.	(d) $\{f \geq a\} \in \Sigma$ for all $a \in \mathbb{R}$.
- * Let (f_n) is a sequence of real valued measurable functions. Define $f(x) = \lim f_n(x)$ if the limit exists, and $f(x) = \infty$ otherwise. Show that f is measurable.

Assignment 5: Assigned Wed 09/25. Due Wed 10/02

- Let (X, Σ, μ) be a measure space, and $(X, \Sigma_\mu, \bar{\mu})$ its completion. Show that $g : X \rightarrow [-\infty, \infty]$ is Σ_μ -measurable if and only if there exists two Σ -measurable functions $f, h : X \rightarrow [-\infty, \infty]$ such that $f = h$ μ -almost everywhere, and $f \leq g \leq h$ everywhere.
- Let μ be a regular (but not necessarily finite) Borel measure on a metric space X .
 - True or false: For any $f : X \rightarrow \mathbb{R}$ measurable and $\varepsilon > 0$ there exists $g : X \rightarrow \mathbb{R}$ continuous such that $\mu\{f \neq g\} < \varepsilon$? Prove it or find a counter example.
 - Do the previous subpart when $X = \mathbb{R}^d$.
- Let for $n \in \mathbb{N}$ define $A_n = \bigcup_{k \in \mathbb{Z}} [\frac{2k}{2^n}, \frac{2k+1}{2^n})$. If $E \in \mathcal{B}(\mathbb{R})$ does $\lim_{n \rightarrow \infty} \lambda(A_n \cap E)$ exist? Prove it.
- If $f \geq 0$ is measurable show that $\int_X f d\mu = 0 \iff f = 0$ almost everywhere.
- Suppose $I \subseteq \mathbb{R}^d$ is a cell, and $f : I \rightarrow \mathbb{R}$ is Riemann integrable. Show that f is measurable, Lebesgue integrable and that the Lebesgue integral of f equals the Riemann integral.
 - Is the previous subpart true if we only assume that an improper (Riemann) integral of f exists? Prove or find a counter example.

Optional problems, and details in class I left for you to check.

- * Let $f : [0, 1] \rightarrow [0, 1]$ be the Cantor function, and $g(x) = \inf\{f = x\}$. Show that f is (Hölder) continuous, and the range of g is the Cantor set. What is the largest exponent α for which f is Hölder- α continuous?
- * Let μ be the counting measure on \mathbb{N} , and $f : \mathbb{N} \rightarrow \mathbb{R}$ a function.
 - If $\sum_1^\infty |f(n)| < \infty$, then show that $\sum_{n=1}^\infty f(n) = \int_{\mathbb{N}} f d\mu$.
 - If the series $\sum_{n=1}^\infty f(n)$ is conditionally convergent, show that $\int_{\mathbb{N}} f d\mu$ is not defined.
- * Let X be a metric space $C \subseteq X$ be closed and $f : C \rightarrow \mathbb{R}$ be continuous.
 - If $0 \leq f \leq 1$, then show that there exists $F : X \rightarrow \mathbb{R}$ continuous such that $F(c) = f(c)$ for all $c \in C$. [HINT: Let $F(x) = f(x)$ for all $x \in C$, and $F(x) = \inf\{f(c) + \frac{d(x,c)}{d(x,C)} - 1 \mid c \in C\}$ for $x \notin C$.]
 - (Tietze extension theorem in metric spaces) Do the previous subpart without assuming $0 \leq f \leq 1$. [HINT: Put $g = \tan^{-1}(f)$, construct G by the previous subpart and set $F = \tan(G)$.]
- * Finish the proof of Lusin's theorem. (I only proved it for bounded positive functions in class.)
- * Find a Borel measurable function $f : [0, 1] \rightarrow \mathbb{R}$ which is not continuous almost everywhere.
- * Let $0 \leq s \leq t$ be two simple functions. Show $\int_X s \leq \int_X t$.
- * Show directly $\int_X \alpha f = \alpha \int_X f$ for any $\alpha \in \mathbb{R}$ and integrable function f .

Assignment 6: Assigned Wed 10/02. Due Never

*In light of your **MIDTERM** this homework is optional.*

- If f is a bounded measurable function and $\mu(X) < \infty$, then show $\int_X f d\mu = \inf\{\int_X t d\mu \mid t \geq f \text{ is simple}\}$.
 - If f, g are bounded measurable functions and $\mu(X) < \infty$ show directly that $\int_X (f + g) d\mu = \int_X f d\mu + \int_X g d\mu$.
- Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a measurable function. We define the *Laplace Transform* of f to be the function $F(s) = \int_0^\infty \exp(-st)f(t) dt$ wherever defined.
 - If $\int_0^\infty |f(t)| dt < \infty$, show that $F : [0, \infty) \rightarrow \mathbb{R}$ is continuous.
 - If $\int_0^\infty t|f(t)| dt < \infty$, show that $F : [0, \infty) \rightarrow \mathbb{R}$ is differentiable.
 - If f is continuous and bounded, compute $\lim_{s \rightarrow \infty} sF(s)$.
- For $p \in \mathbb{R}$ define $F(y) = \int_0^\infty \frac{\sin(xy)}{1+x^p} dx$.
 - For what $p \in \mathbb{R}$ is F defined? When defined, is F continuous? Prove it.
 - Show that F is differentiable for $p > 2$, and not differentiable when $p = 2$.
- (Push forward measures) Let μ be a measure on (X, Σ) , and $f : X \rightarrow Y$ be any function. Define $\tau \subseteq \mathcal{P}(Y)$ by $\tau = \{A \subseteq Y \mid f^{-1}(A) \in \Sigma\}$. For $A \in \tau$ define $\nu(A) = \mu(f^{-1}(A))$.
 - Show that τ is a σ -algebra, and ν is a measure on (Y, τ) . [The measure ν is called the push-forward of μ under f , and often denoted by $\mu_{f^{-1}}$.]
 - If $g \in L^1(Y, \nu)$, then show that $g \circ f \in L^1(X, \mu)$ and $\int_X g \circ f d\mu = \int_Y g d\nu$.
- (Pull back measures) Say ν is a measure on (Y, τ) and $f : X \rightarrow Y$ is surjective.
 - Show that $\Sigma = \{A \subseteq X \mid f(A) \in \tau\}$ need not be a σ -algebra. If Σ is a σ -algebra, show that $\mu(A) = \nu(f(A))$ need not be a measure on (X, Σ) .
 - Define instead $\Sigma = \{A \subseteq X \mid f^{-1}(f(A)) = A, \& f(A) \in \tau\}$, and $\mu(A) = \nu(f(A))$. Show that Σ is a σ -algebra and μ is a measure.
 - If $g \in L^1(Y, \nu)$, then show that $g \circ f \in L^1(X, \mu)$ and $\int_X g \circ f d\mu = \int_Y g d\nu$.
- (Linear change of variable) Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be integrable.
 - For any $y \in \mathbb{R}^d$ show that $\int_{\mathbb{R}^d} f(x+y) d\lambda(x) = \int_{\mathbb{R}^d} f(x) d\lambda(x)$.
 - If $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ an invertible linear transformation, and $E \in \mathcal{L}(\mathbb{R}^d)$. Show that

$$\int_{T^{-1}(E)} (f \circ T) |\det T| d\lambda = \int_E f d\lambda.$$

Details in class I left for you to check.

- * Check that if s, t are non-negative simple functions then $\int_X (s+t) = \int_X s + \int_X t$.
- * Show that there exists $f : \mathbb{R} \rightarrow [0, \infty)$ Borel measurable such that $\int_a^b f d\lambda = \infty$ for all $a, b \in \mathbb{R}$ with $a < b \in \mathbb{R}$. [HINT: Let $g(x) = \chi_{\{|x|<1\}} |x|^{-1/2}$, and define $h(x) = \sum_{m=-\infty}^\infty \sum_{n=1}^\infty 2^{-m-n} g(x-m/n)$.]

Assignment 7: Assigned Wed 10/09. Due Wed 10/16

- Do questions 3, 5, and 6 from HW6.
- (a) (*Jensen's inequality*) Let $a, b \in [-\infty, \infty]$ with $a < b$ and $\varphi : (a, b) \rightarrow \mathbb{R}$ be a convex function. If $\mu(X) = 1$ and $f : X \rightarrow (a, b)$ is integrable then show

$$\varphi\left(\int_X f d\mu\right) \leq \int_X \varphi \circ f d\mu.$$

- (b) If φ above is strictly convex, when can you have equality?
- (a) Suppose $p, q, r \in [1, \infty]$ with $p < q < r$. Prove that for all $f \in L^p \cap L^r$, $f \in L^q$. Further, find $\theta \in (0, 1)$ such that $\|f\|_q \leq \|f\|_p^\theta \|f\|_r^{1-\theta}$.
(b) If for some $p \in [1, \infty)$, $f \in L^p(X) \cap L^\infty(X)$ show that $\lim_{q \rightarrow \infty} \|f\|_q = \|f\|_\infty$.
[This sort of justifies the notation $\|\cdot\|_\infty$.]
(c) Let $p_0 \in (0, \infty]$, $\mu(X) = 1$ and $f \in L^{p_0}(X)$. Prove $\lim_{p \rightarrow 0^+} \|f\|_p = \exp\left(\int_X \ln|f| d\mu\right)$.

Optional problems, and details in class I left for you to check.

- * Let $g \geq 0$ be measurable, and define $\nu(A) = \int_A g d\mu$. Show that ν is a measure, and $\int_E f d\nu = \int_E fg d\mu$.
- * Prove Hölder's inequality for $p = 1$ and $q = \infty$.
- * If $p_i, q \in [1, \infty]$ with $\sum_1^N \frac{1}{p_i} = \frac{1}{q}$, show that $\|\prod_1^n f_i\|_q \leq \prod \|f_i\|_{p_i}$.
- * Show that L^∞ is a Banach space.
- * For $p \in [0, 1)$ show that you need not have $\|f + g\|_p \leq \|f\|_p + \|g\|_p$.
- * Let $p, q \in (1, \infty)$, $\frac{1}{p} + \frac{1}{q} = 1$, $f \in L^p$ and $g \in L^q$. Show that $\int_X |fg| d\mu = \|f\|_p \|g\|_q$ if and only if there exists constants $\alpha, \beta \geq 0$ such that $\alpha f^p = \beta g^q$.
- (a) If X is σ -finite, then show $\|f\|_\infty = \sup_{g \in L^1 - \{0\}} \frac{1}{\|g\|_1} \int_X fg d\mu$.
(b) Show that the previous subpart is false if X is not σ -finite.

Assignment 8: Assigned Wed 10/16. Due Wed 10/23

- (a) If $\mu(X) < \infty$, $1 \leq p < q \leq \infty$, show $L^q(X) \subseteq L^p(X)$ and the inclusion map from $L^q(X) \rightarrow L^p(X)$ is continuous. Find an example where $L^q(X) \subsetneq L^p(X)$. [HINT: Show $\|f\|_p \leq \mu(X)^{\frac{1}{p} - \frac{1}{q}} \|f\|_q$.]
(b) Let $\ell^p = L^p(\mathbb{N})$ with respect to the counting measure. If $1 \leq p < q$ show that $\ell^p \subsetneq \ell^q$. Is the inclusion map $\ell^p \hookrightarrow \ell^q$ continuous? Prove your answer.
- (a) Suppose $p \in [1, \infty)$, and $f \in L^p(\mathbb{R}^d, \lambda)$. For $y \in \mathbb{R}^d$, let $\tau_y f : \mathbb{R}^d \rightarrow \mathbb{R}$ be defined by $\tau_y f(x) = f(x - y)$. Show that $(\tau_y f) \rightarrow f$ in L^p as $|y| \rightarrow 0$.
(b) What happens for $p = \infty$?
- Suppose $\Sigma = \sigma(\mathcal{C})$, where $\mathcal{C} \subseteq \mathcal{P}(X)$ is countable. If μ is a σ -finite measure and $1 \leq p < \infty$, show that $L^p(X)$ is separable (i.e. has a countable dense subset).
- (a) Suppose $\lim_{\lambda \rightarrow \infty} \sup_n \int_{|f_n| > \lambda} |f_n| d\mu = 0$. Show that there exists an increasing function φ with $\varphi(\lambda)/\lambda \rightarrow \infty$ as $\lambda \rightarrow \infty$, such that $\sup_n \int_X \varphi(|f_n|) < \infty$.
(b) Suppose $\{f_n\}$ is uniformly integrable, and $\sup_n \int |f_n| < \infty$. Show that $\lim_{\lambda \rightarrow \infty} \sup_n \int_{|f_n| > \lambda} |f_n| = 0$.
(c) Show that the previous part fails without the assumption $\sup_n \int |f_n| < \infty$.
- Let $e_n(x) = e^{2\pi i n x}$, $X = [0, 1]$. For what $p \in [1, \infty]$ does $\{e_n\}$ have a convergent subsequence in $L^p(X, \lambda)$? Prove it.

Optional problems, and details in class I left for you to check.

- * In Vitali's convergence theorem prove that the assumption $f \in L^1$ is unnecessary.
- * If $(f_n) \rightarrow f$ in L^1 , show that $\{f_n\}$ is uniformly integrable. [This is part of Vitali's theorem which I didn't have time to prove in class.]
- * Show that if $(f_n) \rightarrow f$ in measure, then (f_n) need not converge to f in L^p .
- * Finish the proof that $C_c(X)$ is dense in L^p . [I only did the case when X is compact in class.]
- * Show that simple functions are dense in L^∞ .
- * Show that $C_c(\mathbb{R})$ is not dense in $L^\infty(\mathbb{R})$.
- * Show that $L^\infty(\mathbb{R})$ is not separable.

Assignment 9: Assigned Wed 10/23. Due Wed 10/30

- Recall we defined the variation of μ by $|\mu| = \mu^+ + \mu^-$, and the total variation by $\|\mu\| = |\mu|(X)$. (You should check that these are well defined.)
 - Let \mathcal{M} be the space of all finite signed measures on (X, Σ) . Show that \mathcal{M} with total variation norm (i.e. with $\|\mu\| = |\mu|(X)$) is a Banach space.
 - Show that $(\mu_n) \rightarrow \mu$ if and only if $(\mu_n(A)) \rightarrow \mu(A)$ uniformly in A , $\forall A \in \Sigma$.
- For a signed measure, we define $\int_X f d\mu = \int_X f d\mu^+ - \int_X f d\mu^-$. Suppose $(f_n) \rightarrow f$, $(g_n) \rightarrow g$, and $|f_n| \leq g_n$ almost everywhere with respect to $|\mu|$. If $\lim \int_X g_n d|\mu| = \int_X g d|\mu| < \infty$, show that $\lim \int_X f_n d\mu = \int_X f d\mu$.
 - Suppose $f, f_n \in L^1$, and $(f_n) \rightarrow f$ almost everywhere. Show that $\lim \int |f_n - f| d|\mu| = 0$ if and only if $\lim \int |f_n| d|\mu| = \int |f| d|\mu|$.
- If μ is a positive σ -finite measure, and ν is a finite signed measure such that $|\nu| \ll \mu$, show that there exists $f \in L^1(X, \mu)$ such that $d\nu = f d\mu$.
 - Compute $\frac{d\nu}{d|\nu|}$ in terms of the Hanh decomposition of ν . [NOTATION: We say $g = \frac{d\nu}{d\mu}$ if $d\nu = g d\mu$.]
- Let ν_1 and ν_2 be two finite signed measures on X . Show that there exists a finite signed measure $\nu_1 \vee \nu_2$ such that $\nu_1 \vee \nu_2(A) \geq \nu_1(A) \vee \nu_2(A)$, and for any other finite signed measure ν such that $\nu(A) \geq \nu_1(A) \vee \nu_2(A)$ we must have $\nu_1 \vee \nu_2 \leq \nu$.
 - If ν_1, ν_2 above are absolutely continuous with respect to a positive σ -finite measure μ , prove $\nu_1 \vee \nu_2 \ll \mu$ and express $\frac{d(\nu_1 \vee \nu_2)}{d\mu}$ in terms of $\frac{d\nu_1}{d\mu}$ and $\frac{d\nu_2}{d\mu}$.
- Let (Ω, \mathcal{F}, P) be a measure space with $P(\Omega) = 1$, and $X \in L^1(\Omega, \mathcal{F}, P)$. [The probabilistic interpretation is that Ω is the sample space, $A \in \mathcal{F}$ is an event, X is a random variable, and $P(X \in B)$ is the chance that $X \in B$, where $B \in \mathcal{B}(\mathbb{R})$.]
 - Suppose $\mathcal{G} \subseteq \mathcal{F}$ is a σ -sub-algebra of \mathcal{F} . Show that there exists a unique \mathcal{G} -measurable function Y such that $\int_A Y dP = \int_A X dP$ for all $A \in \mathcal{G}$. [Y is called the *conditional expectation* of X given \mathcal{G} , and denoted by $E(X|\mathcal{G})$.]
 - (*Tower property*) If $\mathcal{H} \subseteq \mathcal{G}$ is a σ -sub-algebra, show that $E(X|\mathcal{H}) = E(E(X|\mathcal{G})|\mathcal{H})$ almost everywhere.
 - (*Conditional Jensen*) If $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is convex, show that $\varphi(E(X|\mathcal{G})) \leq E(\varphi(X)|\mathcal{G})$ almost everywhere.
 - Suppose $X \in L^2(\Omega, \mathcal{F}, P)$. Show that $E(X|\mathcal{G})$ is the L^2 -orthogonal projection of X onto the subspace $L^2(\Omega, \mathcal{G})$. [Namely show $E(X|\mathcal{G}) \in L^2(\Omega, \mathcal{G})$, and $\int_\Omega (X - E(X|\mathcal{G}))Y dP = 0$ for all $Y \in L^2(\Omega, \mathcal{G})$.]

Optional problems, and details in class I left for you to check.

- * In the proof of the Hanh decomposition, prove the following: Say $\mu(X) > -\infty$, and $\alpha = \inf\{\mu(B)\}$. Let B'_n be a sequence of negative sets such that $\mu(B'_n) \rightarrow \alpha$. Let $N = \cup B'_n$. Show $\mu(N) = \alpha$.
- * Prove the Hanh decomposition is unique up to null sets.
- * Prove uniqueness of the Jordan decomposition.
- * Show that the Radon-Nikodym theorem need not hold if μ, ν are not σ -finite.

Assignment 10: Assigned Wed 10/30. Due Wed 11/06

- Let $p \in [1, \infty)$ and q be conjugate Hölder exponent. If X is σ -finite, show that there exists a bijective linear isometry between $(L^p)^*$ and L^q .
 - The above result is *false* for $p = \infty$ even when $\mu(X) < \infty$. Find where our proof from class (when $\mu(X) < \infty$) fails when $p = \infty$.
 - We can (partially) construct a counter example on ℓ^∞ as follows. The Hanh-Banach theorem shows that there exists $T \in (\ell^\infty)^*$ such that $Ta = \lim a_n$, for all $a = (a_n) \in \ell^\infty$ such that $\lim a_n$ exists and is finite. Show that there does not exist $b \in \ell^1$ such that $Ta = \sum a_n b_n$ for all $a \in \ell^\infty$.
- Suppose $\sum_{m=1}^\infty (\sum_{n=1}^\infty |a_{m,n}|) < \infty$. Show that $\sum_{m=1}^\infty (\sum_{n=1}^\infty a_{m,n}) = \sum_{n=1}^\infty (\sum_{m=1}^\infty a_{m,n})$.
 - Give a counter example to (a) if we only assume $\sum_m \sum_n a_{m,n} < \infty$. Find a counter example where both iterated sums are finite.
- If X and Y are not σ -finite, show that Fubini's theorem need not hold.
 - If $\int_{[-1,1]^2} f d\lambda$ is not assumed to exist (in the extended sense), show that both iterated integrals can exist, be finite, but need not be equal.
- (*Fubini for completions*.) Suppose (X, Σ, μ) and (Y, τ, ν) are two σ -finite, complete measure spaces. Let $\varpi = (\Sigma \otimes \tau)_\pi$ denote the completion of $\Sigma \otimes \tau$ with respect to the product measure $\pi = \mu \times \nu$.
 - Show that $\Sigma \otimes \tau$ need not be π -complete (i.e. $\varpi \not\supseteq \Sigma \otimes \tau$ in general).
 - Suppose $f : X \times Y \rightarrow [-\infty, \infty]$ is \mathcal{F} -measurable. Define as usual the slices $\varphi_{f,x} : Y \rightarrow [0, \infty]$ by $\varphi_{f,x}(y) = f(x, y)$, and similarly $\psi_{f,y}(x) = f(x, y)$. Show that for μ -almost all $x \in X$, $\varphi_{f,x}$ is an τ -measurable, and for ν -almost all y , $\psi_{f,y}$ is an Σ -measurable.
 - Suppose f is integrable on $X \times Y$ in the extended sense. Define $F(x) = \int_Y f(x, y) d\nu(y)$ and $G(y) = \int_X f(x, y) d\mu(x)$. Show F is defined μ -a.e. and Σ -measurable. Similarly show G is defined ν -a.e., and τ -measurable. Further, show and that $\int_X F d\mu = \int_Y G d\nu = \int_{X \times Y} f d(\mu \times \nu)$.
- Let (X, Σ, μ) , (Y, τ, ν) be two σ -finite measure spaces, $p \in [1, \infty]$, and $f : X \times Y \rightarrow \mathbb{R}$ is $\Sigma \otimes \tau$ measurable. Let $F(x) = \int_Y f(x, y) d\nu(y)$, and $\psi_{y,f}$ be the slice of f defined by $\psi_{y,f}(x) = f(x, y)$. Show that $\|F\|_{L^p(X)} \leq \int_Y \|\psi_{y,f}\|_{L^p(X)} d\nu(y)$. [When $Y = \{1, 2\}$ with the counting measure, this is exactly Minkowski's triangle inequality.]

Optional problems, and details in class I left for you to check.

- * Let $\mu(X) < \infty$, $p \in [1, \infty)$ and $T \in (L^p)^*$. Let $\nu(A) = T(\chi_A)$. We've seen in class that $\nu \ll \mu$ and so $d\nu = g d\mu$ for some $g \in L^1(\mu)$.
 - Show that $Tf = \int_X fg d\mu$ for all f simple.
 - If $\frac{1}{p} + \frac{1}{q} = 1$ show $\|g\|_q = \sup\{\int_X sg\}$, where the supremum runs over all simple functions s such that $\|s\|_p \leq 1$. Conclude $g \in L^q$ and $\|g\|_q \leq \|T\|$.
 - Show that $Tf = \int_X fg d\mu$ for all $f \in L^p$, to conclude the proof.
- * Show that the Lebesgue measure on \mathbb{R}^{m+n} is the product of the Lebesgue measures on \mathbb{R}^m and \mathbb{R}^n respectively.