
Milly
INTERNATIONAL CO., LTD

天然植粹 NATURAL / 以人為本 HUMANISM / 創新研發 INNOVATION



維納斯N
賦活時空膠囊

美顏x抗老x煥白x青春


米蕾國際事業有限公司

控制老化的關鍵是什麼？



Elizabeth Helen Blackburn

答案就在每個人身體中的「端粒」。
正是端粒的長短，控制了細胞和人體的壽命以及疾病的發生。




米蕾國際事業有限公司

控制老化的關鍵是什麼？




米蕾國際事業有限公司

端粒-老化的過程



米蕾國際事業有限公司

老化時間表



Milly INTERNATIONAL CO., LTD

米蕾國際事業有限公司


老化的徵兆




老化
真的是
宿命?

米蕾國際事業有限公司

NAD+與衰老




NMN之父David Sinclair

David Sinclair教授曾於2014年，被《時代》雜誌（Time 100）評為世界上最有影響力的100個人之一。於2018年，被《時代》評為50位醫療健康領域最具影響力的人物之一。

哈佛大學醫學院教授David Sinclair團隊發表研究文章稱，NAD+是DNA修復蛋白PARP1的底物，也是長壽基因SIRT1的激活劑，發揮關鍵的生理功能。**但是在年齡增長的過程中，NAD+的含量會逐漸下降進而導致SIRT1(長壽基因)的活性降低，從而導致細胞衰老。**

米蕾國際事業有限公司

NAD+與衰老




米蕾國際事業有限公司

抗老關鍵-NAD+

年齡越高的人

NAD+流失就越多




米蕾國際事業有限公司

諾貝爾得獎主研究認證有效



Richard J. Roberts
1993年諾貝爾醫學獎得主



Eric Kandel
2000年諾貝爾醫學獎得主




Danuel Kahneman
2002年諾貝爾經濟學獎得主



Aaron Ciechanover
2004年諾貝爾化學獎得主



Jack W. Szostak
2009年諾貝爾醫學獎得主



Martin Karplus
2013年諾貝爾化學獎得主



Thomas C. Südhof
2013年諾貝爾醫學獎得主



Paul Modrich
2015年諾貝爾化學獎得主

米蕾國際事業有限公司

長生不老藥大戰

追求長生不老的大亨》貝佐斯阻止細胞衰老 遠離病痛

工商時報 陳穎芃、鍾志恆 2020.10.18



亞馬遜執行長貝佐斯。圖／美聯社

89歲香港首富砸逾七億元研究，還親自試吃

「長生不老藥」大戰　李嘉誠、貝佐斯搶加入

產業風雲｜ 撰文者： 蔡靚萱 ｜出刊日期： 2017-09-28



巴菲特旗下與NAD+前體NMN產品品牌方瑞維拓達成合作



米蕾國際事業有限公司

全民瘋NMN












米蕾國際事業有限公司

NMN是什麼?

中文爲β-菸醯胺單核苷酸，英文則寫成 NMN
功用：NMN可轉變成「NAD+」，而「NAD+」
此物質參與我們的青春、美麗與代謝



隨著年齡、「NAD+」將大趨勢流失


米蕾國際事業有限公司

NAD+可以活化抗衰老基因

- 修復DNA
- 維持粒線體功能
- 促進細胞修復
- 抑制神經凋亡

米蕾國際事業有限公司

NAD+掌管全身生理反應



米蕾國際事業有限公司

補充劑目的是增加體內NAD+

NMN ⟹ NAD+ ⇧

NADH ⟹⟹ NAD+ ⇧⇧⇧

美麗 青春 代謝

息息相關

Milly INTERNATIONAL CO., LTD

米蕾國際事業有限公司

德國專利PANMOL® NADH

成分1

PANMOL®

NADH


專利菸鹼醯胺複合物



素食　零果糖　零乳糖　零麩質　零基改　清真認證　猶太食品認證　專利認證

米蕾國際事業有限公司

合成NAD+的步驟



Milly INTERNATIONAL CO., LTD
米蕾國際事業有限公司

NMN & NADH

專利NADH	VS	NMN
○	產生NAD+	○
較快速	產生NAD+效率	較慢
1	機轉步驟	3
生成NAD+過程，同時生成能量ATP	是否需要消耗能量	必須損耗ATP才能升成NAD+
X	是否需要特定的酶	○
○	產生能量	X
有	臨床實驗	有
無副作用	安全性	待研究

米蕾國際事業有限公司

NMN的不確定性

FDA：抗老保健夯品NMN非膳食補充劑 以藥物審查！

撰文 | 記者 李林璦　　日期 | 2022-11-14



FDA：抗老保健夯品NMN非膳食補充劑 以藥物審查！


FDA 仔細考慮了可獲得的信息，並確定 NMN是一項已經進行了大量臨床調查並且已經公開存在此類調查的臨床新藥物，” FDA官員 Yeager 補充道。

“因此，我們得出結論，NMN不得作為膳食補充劑或在膳食補充劑中銷售。”

"NMN已被授權作為一種新藥進行調查” ，因此根據法規不得作為膳食補充劑或在膳食補充劑中銷售！若此結論確定，NMN將會很快在美國市場上下架！

米蕾國際事業有限公司

微囊化技術




穩定

抗酸

天然

米蕾國際事業有限公司


微囊化技術



Milly INTERNATIONAL CO., LTD

米蕾國際事業有限公司

多國認證



七國專利 **美國、加拿大、歐洲、塞爾維亞、克羅埃西亞、挪威、俄羅斯**

european Patent No: 1562613

US Patent No: 7255813

Russian Patent No: 2338540

canada Patent No: 2503860

Serbian Patent No: 51606

croatian Patent No: P20050573

Norwegian Patent No: 333911

品質認證 KOSHER HALAL NON GMO IFS FOOD

米蕾國際事業有限公司

功效臨床實驗

> Science. 2016 Jun 17;352(6292):1436-43. doi: 10.1126/science.aaf2693. Epub 2016 Apr 28.

NAD+ repletion improves mitochondrial and stem cell function and enhances life span in mice

NAD+可增強粒腺體及幹細胞功能，以減緩衰老，並延長壽命。

PMID: 27127236 DOI: 10.1126/science.aaf2693

Abstract

Adult stem cells (SCs) are essential for tissue maintenance and regeneration yet are susceptible to senescence during aging. We demonstrate the importance of the amount of the oxidized form of

> Cell Metab. 2016 Oct 11;24(4):566-581. doi: 10.1016/j.cmet.2016.09.004.

NAD+ Replenishment Improves Lifespan and Healthspan in Ataxia Telangiectasia Models via Mitophagy and DNA Repair

Evandro Fei Fang [1], Henok Kassahun [2], Deborah L Croteau [1], Morten Scheibye-Knudsen [3], Krisztina Marosi [4], Huiming Lu [1], Raghavendra A Shamanna [1], Sumana Kalyanasundaram [5], Ravi Chand Bollineni [6], Mark A Wilson [4], Wendy B Iser [4], Bradley N Wollman [1], Marya Morevati [3], Jun Li [7], Jesse S Kerr [1], Qiping Lu [1], Tyler B Waltz [1], Jane Tian [1], David A Sinclair [8], Mark P Mattson [9], Hilde Nilsen [2], Vilhelm A Bohr [10]

Affiliations + expand

NAD+可修復神經元DNA，改善神經肌肉，延遲記憶喪失。

Ataxia telangiectasia (A-T) is a rare autosomal recessive disease characterized by progressive neurodegeneration and cerebellar ataxia. A-T is causally linked to defects in ATM, a master regulator

> Cell Metab. 2011 Oct 5;14(4):528-36. doi: 10.1016/j.cmet.2011.08.014.

Nicotinamide mononucleotide, a key NAD(+) intermediate, treats the pathophysiology of diet- and age-induced diabetes in mice

Jun Yoshino [1], Kathryn F Mills, Myeong Jin Yoon, Shin-ichiro Imai

透過恢復NAD+改善葡萄糖耐受性，提高胰島素敏感度

Abstract

Type 2 diabetes (T2D) has become epidemic in our modern lifestyle, likely due to calorie-rich diets overwhelming our adaptive metabolic pathways. One such pathway is mediated by nicotinamide phosphoribosyltransferase (NAMPT), the rate-limiting enzyme in mammalian NAD(+) biosynthesis, and the NAD(+)-dependent protein deacetylase SIRT1. Here, we show that NAMPT-mediated NAD(+) biosynthesis is severely compromised in metabolic organs by high-fat diet (HFD). Strikingly, nicotinamide mononucleotide (NMN), a product of the NAMPT reaction and a key NAD(+) intermediate, ameliorates glucose intolerance by restoring NAD(+) levels in HFD-induced T2D mice. NMN also enhances hepatic insulin sensitivity and restores gene expression related to oxidative stress, inflammatory response, and circadian rhythm, partly through SIRT1 activation. Furthermore, NAD(+) and NAMPT levels show significant decreases in multiple organs during aging, and NMN improves glucose intolerance and lipid profiles in age-induced T2D mice. These findings provide critical insights into a potential nutriceutical intervention against diet- and age-induced T2D.

Copyright © 2011 Elsevier Inc. All rights reserved.

Randomized Controlled Trial > Cell Rep. 2019 Aug 13;28(7):1717-1728.e6.

doi: 10.1016/j.celrep.2019.07.043.

Nicotinamide Riboside Augments the Aged Human Skeletal Muscle NAD+ Metabolome and Induces Transcriptomic and Anti-inflammatory Signatures

Yasir S Elhassan [1], Katarina Kluckova [2], Rachel S Fletcher [2], Mark S Schmidt [3], Antje Garten [2], Craig L Doig [2], David M Carturight [2], Lucy Oakey [2], Claire V Burley [4], Ned Jenkinson [4]

NAD+可改善年長者的肌肉，並減少發炎

Affiliations + expand

PMID: 31412242 PMCID: PMC6702140 DOI: 10.1016/j.celrep.2019.07.043

Free PMC article

Abstract

Nicotinamide adenine dinucleotide (NAD^+) is modulated by conditions of metabolic stress and has been reported to decline with aging in preclinical models, but human data are sparse. Nicotinamide riboside (NR) supplementation ameliorates metabolic dysfunction in rodents. We aimed to establish whether oral NR supplementation in aged participants can increase the skeletal muscle NAD^+ metabolome and if it can alter muscle mitochondrial bioenergetics. We supplemented 12 aged men with 1 g NR per day for 21 days in a placebo-controlled, randomized, double-blind, crossover trial. Targeted metabolomics showed that NR elevated the muscle NAD^+ metabolome, evident by increased nicotinic acid adenine dinucleotide and nicotinamide clearance products. Muscle RNA sequencing revealed NR-mediated downregulation of energy metabolism and mitochondria pathways, without altering mitochondrial bioenergetics. NR also depressed levels of circulating inflammatory cytokines. Our data establish that oral NR is available to aged human muscle and identify anti-inflammatory effects of NR.

米蕾國際事業有限公司

功效臨床實驗

Therapeutic effects of oral NADH on the symptoms of patients with chronic fatigue syndrome

口服NADH對慢性疲倦症候群(CFS)的治療效果:

PMID: 10071523 DOI: 10.1016/S1081-1206(10)62595-1

Free article

[Stabilized NADH (ENADA) improves jet lag-induced cognitive performance deficit]

[Article in German]

G D Birkmayer [1], G G Kay, E Vürre

Affiliations + expand

PMID: 12385067 DOI: 10.1046/j.1563-258x.2002.02072.x

NADH改善時差造成的負面認知及嗜睡:

limited in efficacy and practicality. The efficacy of a stabilized, sublingual form of reduced nicotinamide adenine dinucleotide (NADH, ENADAlert, Menuco Corp.) as a countermeasure for jet lag was examined. Because NADH increases cellular production of ATP and facilitates dopamine synthesis, act the effects of jet lag on cognitive functioning and sleepiness. Thirty-five healthy, jects participated in this double-blind, placebo-controlled study. Training and baseline onducted on the West Coast before subjects flew overnight to the East Coast, where perience a 3-hour time difference. Upon arrival, individuals were randomly assigned to

Oral reduced B-nicotinamide adenine dinucleotide (NADH) affects blood pressure, lipid peroxidation, and lipid profile in hypertensive rats (SHR)

N Bushehri [1], S T Jarrell, S Lieberman, N Mirdamadi-Zonozi, G Birkmayer

Affiliations + expand

PMID: 9893217 DOI: 10.1023/a:1008242900153

NADH對於高血壓的治療效果

are important. Although oral B-nicotinamide adenine dinucleotide (NAD used principally to treat various neurologic disorders, we wished to inve the same potential to lower BP and benefit the cardiovascular system a similar-type agent. As a first approximation, spontaneously hypertensiv determine effects of oral NADH. In a blinded, placebo-controlled study, ten, NADH for ten weeks. Systolic BP was measured by tail plethysmogr terminally, and chemistries were performed by routine methodologies. species (TBARS) (an estimate of lipid peroxidation/free radical formation hepatic tissues. The following was noted: water and food intake were co weight gain of young SHR were similar in the placebo and NADH groups differ between the two groups over the first month, it decreased and sta remainder of study in SHR receiving oral NADH. At the end of 60 days, S 184 mm Hg +/- 2.8 (SEM) compared to 201 mm Hg +/- 2.1 (SEM) in con significant differences were seen in blood levels of glucose, insulin, trigly NADH intake lowered total cholesterol ($p < 0.002$) and LDL ($p < 0.02$). Re significantly lower in SHR receiving NADH ($P < 0.001$). Accordingly, supp coenzyme NADH theoretically could prove to be useful in preventing ag thus, various cardiovascular maladies.

Treatment of Alzheimer's disease with stabilized oral nicotinamide adenine dinucleotide: a randomized, double-blind study

wn etiology, consisting of

lobnik, D Storga-Tomic, G Kay

NADH對阿茲海默症效果

dinucleotide (NADH) on cognitive functioning in patients with Alzheimer's disease (AD). NAD coenzyme that plays a key role in cellular energy production and stimulates dopamine produ previous trials NADH has been shown to improve cognitive functioning in patients with Parki disease, depression and AD. The present trial was a randomized, placebo-controlled, matche double-blind, 6-month clinical study. Patients with probable AD (n = 26) were randomized to either stabilized oral NADH (10 mg/day) or placebo. Twelve pairs of subjects were matched f and baseline total score on the Mattis Dementia Rating Scale (MDRS) and the Mini Mental St Examination. After 6 months of treatment, subjects treated with NADH showed no evidence progressive cognitive deterioration and had significantly higher total scores on the MDRS co with subjects treated with placebo ($p < 0.05$). Analysis of MDRS subscales revealed significan performance by NADH subjects on measures of verbal fluency ($p = 0.019$), visual-constructio ($p = 0.038$) and a trend ($p = 0.08$) to better performance on a measure of abstract verbal rea There were no differences between groups in measures of attention, memory, or in clinician dementia severity (Clinical Dementia Rating). Consistent with earlier studies, the present findi support NADH as a treatment for AD.

Nicotinamide adenine dinucleotide (NADH)--a new therapeutic approach to Parkinson's disease. Comparison of oral and parenteral application

Birkmayer [1], G Vrecko, D Volc, W Birkmayer

iations +

D: 8101414

NADH對帕金森氏症的效果

Abstract

The reduced coenzyme nicotinamide adenine dinucleotide (NADH) has been used as medication in 885 parkinsonian patients in an open label trial. About half of the patients received NADH by intravenous infusion, the other part orally by capsules. In about 80% of the patients a beneficial clinical effect was observed: 19.3% of the patients showed a very good (30-50%) improvement of disability, 58.8% a moderate (10-30%) improvement. 21.8% did not respond to NADH. Statistical analysis of the improvement in correlation with the disability prior to treatment, the duration of the disease and the age of the patients revealed the following results: All these 3 parameters have a significant although weak influence on the improvement. The disability before the treatment has a positive regression coefficient (t value < 0.01). The duration of the disease has a negative regression coefficient (< 0.01) and so has the age a negative regression coefficient (t value < 0.05). In other words younger patients and patients with a shorter duration of disease have a better chance to gain a marked improvement than older patients and patients with longer duration of the disease. The orally applied form of NADH yielded an overall improvement in the disability which was comparable to that

米蕾國際事業有限公司


市售產品原料種類分析



	酵母發酵物	花椰菜	藜麥發芽種子 NADH菸鹼醯胺複合物
含量	0.5%NMN	(含NMN)	10%NADH
有效劑量	250mg	-	50mg
產生NAD+	慢	-	快
進入粒線體	產生NAD+ 再還原成NADH才能進入	-	直接進入粒線體
特色訴求	目前台灣唯一可以 訴求含NMN的原料	-	台灣唯一能有效 吃到足量的原料

米蕾國際事業有限公司

NADH競品分析

	營養師輕食 NMN賦活膠囊	醇耀妍 麗彤生醫NMN超能飲	質妍系 維納斯N賦活時空膠囊
			
抗老因子 種類	NMN	NMN	NADH 菸鹼醯胺複合物
來源	酵母發酵	花椰菜 高麗菜	藜麥發芽種子
功效 複方成分	β-聚葡萄糖	-	緩釋型C 900mg/3顆
含量	1.5mg/2顆 *有效劑量250mg	-	50mg/3顆 *有效劑量50mg
價格	999元/30顆	1680元/20包	2200元/90顆

有效

雙專利

米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

成分2


PUREWAY C™

Antioxidant Super Nutrient

1. 非基改玉米
(維生素C約93%)

2. 植物來源脂質
lipid metabolites

使抗壞血酸可以更快、安全、有效的方式進入細胞，作為抗壞血酸的載體去增加維生素C的腸道吸收及組織分布並增進細胞攝取率



3. 柑橘生物類黃酮
Citrus bioflavonoids

保護維生素C不受體內氧化劑影響，並和維生素C進行協同作用，以支持對抗慢性發炎之疾病、增進免疫系統的功能、幫助傷口復原



one INNOVATION LABS

歷時20年研發結晶

米薗國際事業有限公司

一般型vs緩釋型

一般型

一般型維生素C在
體內約2-4小時
便代謝慢慢流失

高劑量來自於
化學合成

緩釋型



快速吸收
完整送達



超高濃度
有效利用



溫和穩定
持續運作

米薈國際事業有限公司

PUREWAY-C®緩釋型維生素C

國際認證

- 無過敏原
- 素食者可食
- 非基改聲明
- 原料為天然來源
- 不含有害溶劑
- Kosher & Halal 認證
- 美國GRAS
- ISO9001

MADE IN USA

ISO 9001
品質管理系統

FDA GRAS
美國食安食認證

Halal
根據伊斯蘭教義
的食品認證

Kosher
根據猶太教義
的食品認證

米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

產品優勢

◆天然來源

◆美國大廠製造

◆4 項原廠實驗支持

◆獨特成份可提升吸收率與體內保留率

(相較於一般抗壞血酸提升233%)

◆相較一般抗壞血酸對胃更溫和


◆製程專利

◆物性穩定

Milly INTERNATIONAL CO., LTD

米蕾國際事業有限公司

PUREWAY C™



米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

高濃度


實驗方法：

40位健康受試者接受14天的限制維他命C飲食，禁食一個晚上後，隔天服用1000毫克的抗壞血酸(AA)、抗壞血酸鈣(CaA)、PureWay-C® (PWC)、Ester-C® (EC)，抽取受試者的血液樣本作分析

實驗結果：

PureWay-C® 組別在六小時內血液維生素C濃度皆高於其他組別(統計上顯著高於抗壞血酸鈣)

PureWay-C® 吸收較快且較好，優於其他維生素C品牌



Referance:Pancorbo D, Vazquez C, Fletcher MA.
Vitamin C-lipid metabolites: uptake and retention and effect on plasma C-reactive protein and oxidized LDL levels in healthy volunteers. Med Sci Monit. 2008;14(11):CR547–CR551.

米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C


高吸收

- 測量人類淋巴母細胞(免疫細胞)的維生素C含量
- 加入30分後，**PureWay-C® 的免疫細胞維生素C吸收速率提升至233%(相較於抗壞血酸組)**

Table 1 維生素C吸收速率百分比

	Percent Increase		Compared to ascorbic acid
	CaA	Ester-C®	PureWay-C™
Time after addition			
15 min	128	171	214
30 min	122	189	233
45 min	133	192	233

Referance:Weeks BS, Perez PP. Absorption rates and free radical scavenging values of vitamin C-lipid metabolites in human lymphoblastic cells. Med Sci Monit. 2007;13(10):BR205–BR210.




米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

高修復

- 在傷口癒合過程中需要新合成表皮、血管、結締組織，纖維母細胞可合成新的細胞外基質成分(例如：纖連蛋白、膠原蛋白)
- 纖連蛋白是高分子量（~440kDa）的糖蛋白細胞外基質
- 纖維母細胞黏附於細胞外基質成分（例如：纖連蛋白）是合成新真皮和其他結締組織的必要條件
- **PureWay-C® 促進的黏附效果顯著優於Ester C組(約為其三倍)**



米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

高穩定

儲存時間	25°C 60% r.h.	40°C 75% r.h.
0	92.3	92.3
12個月	90.9	90.6
24個月	90.5	90.1


米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

高生長

- 神經生長因子(NGF)：
  屬於神經營養因子，主要參與神經細胞的發育及存活
- 神經突生長代表神經元處於發育狀態 → 神經元訊息傳遞 → 影響人類的思考、感覺、行動
- PC12細胞(神經細胞)可受神經生長因子刺激神經突生長
- **在24小時內PureWay-C® 促進神經突生長效果顯著優於其他維生素C**
- **PureWay-C® 加入3小時後，神經突生長效果優於Ester C組約8倍**






Referance: Weeks BS, Perez PP. A novel vitamin C preparation enhances neurite formation and fibroblast adhesion and reduces xenobiotic-induced T-cell hyperactivation. Med Sci Monit. 2007;13(3):BR51-BR58.

米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

- 神經生長因子(NGF)：
  屬於神經營養因子，主要參與神經細胞的發育及存活

- 神經突生長代表神經元處於發育狀態 → 神經元訊息傳遞 → 影響人類的思考、感覺、行動
- PC12細胞(神經細胞)可受神經生長因子刺激神經突生長
- **在24小時內PureWay-C® 促進神經突生長效果顯著優於其他維生素C**
- **PureWay-C® 加入3小時後，神經突生長效果優於Ester C組約8倍**



Referance: Weeks BS, Perez PP. A novel vitamin C preparation enhances neurite formation and fibroblast adhesion and reduces xenobiotic-induced T-cell hyperactivation. Med Sci Monit. 2007;13(3):BR51-BR58.

米蕾國際事業有限公司

PUREWAY-C®緩釋型維生素C

抗氧化


實驗方法：

40位健康受試者接受14天的限制維他命C飲食，禁食一個晚上後，隔天服用1000毫克的抗壞血酸(AA)、抗壞血酸鈣(CaA)、PureWay-C® (PWC)、Ester-C® (EC)，抽取受試者的血液樣本作分析

實驗結果：

服用PUREWAY-C® 後的氧化LDL下降量顯著大於抗壞血酸下降量

PureWay-C® 對於血液中氧化低密度脂蛋白的降低，效果優於其他維生素C品牌



(註) 一旦血液中含有過多的低密度脂蛋白（LDL）時，過量的LDL會聚集在動脈內壁上，形成脂肪堆積，並且發生氧化；氧化型的低密度脂蛋白（LDL）具有細胞毒性，會啟動血管壁的發炎反應

米蕾國際事業有限公司

PUREWAY-C®檢驗報告



SGS

食品實驗室-高雄
FOOD LAB-KAOHSIUNG
測 試 報 告
Test Report

Page 2 of 4

測試項目	測試方法	測試結果	定量/偵測極限(MDL)	單位
維生素C	衛生福利部食品藥物管理署110年10月01日公開建議檢驗方法TFDA0012.03-修訂開啟試劑食品中水溶性維生素之檢驗方法，以高效液相層析儀二極體陣列偵測器(HPLC/DAD)檢測。（對，實驗室提供原方法進行驗證。）	94.7	0.0500	%(w/w)

備註：
1. 測試報告僅就委託者之委託事項提供測試結果，不對產品合法性做判斷。
2. 本報告共 4 頁，分離使用無效。
3. 若該測試項目是的定量分析以「定量極限」表示；若該測試項目是的定性分析以「偵測極限」表示。
4. 低於定量極限/偵測極限之測定值以"未檢出"或"陰性"表示。
5. 本檢驗報告之所有檢驗內容，均依委託事項執行檢驗，如有不實，願意承擔完全責任。
6. 本次委託測試樣本由SGS抽樣產生，其結果僅對當批樣品(C0021900010)。

- END -

維生素C含量
94.7%



台美檢驗科技股份有限公司
檢驗中心 測試報告

SUPER LAB

委託單位

美隆貿易股份有限公司
台南市南區中華西路一段450號

第 1 頁 / 共 2 頁

檢體名稱：Pure Way-C(Fine Powder)(提供樣品)
製造公司：-
製造日期：2021.04
批　　號：T02-ITF-040721-41
檢體狀態：室溫
送檢方式：顧客送檢
聯 絡 人：品管部
聯絡電話：(06)261-9348
有效期限：2024.04
原 廠 檢：無證
檢體名稱：粉製
檢體數量：25g
報告用途：自主管理

以上檢體資料係由委託單位提供並確認

送檢日期：2021/11/11　檢驗日期：2021/11/12　報告日期：2021/11/19

檢驗項目	檢驗結果	單位	檢驗方法	定量 / 偵測極限
總重金屬（以鉛計）	小於 5 (on (以鉛計)	ppm (ug/g)	衛生福利部 110.11.05 部授食字第1101902451 號公告修正：食品添加物規格檢驗方法食品第一版附錄A 一般試驗法 (A7 第11法)，檢驗原理：比色法，(註：實驗室提供方法適用基質，非屬食品等相關範圍。)	5

以下空白

總重金屬
未檢出

測試報告僅就委託者之委託事項提供檢驗結果，不對產品合法性做判斷

SL 蔡岳廷

米蕾國際事業有限公司

西班牙柑橘萃取物-維生素P

成分3





米蕾國際事業有限公司

西班牙柑橘萃取物-維生素P

抑制維生素C消耗

抑制維生素C消耗

維生素P作用

強化毛細血管



米蕾國際事業有限公司

歐洲萬能草藥-接骨木莓萃取物

成分4

抗氧化

天然營養素

強化免疫



抗病毒

豐富生物活性

抗菌

保護呼吸道

保護心血管

米蕾國際事業有限公司

質妍系-維納斯N 賦活時空膠囊

煥白 Q彈

*PUREWAY-C®緩釋型維生素C

青春回朔

*德國專利PANMOL® NADH

天然逆時配方

歐洲萬能草藥

*接骨木莓萃取物

協同抗氧

*西班牙柑橘維生素P

◆ NADH

全素

開啟話題 引導需求	✓ 人都會變老，但為何有些人看起來特別年輕或特別健康，關鍵就在於端粒，端粒可以保護染色體，讓我們更健康活得更久。 ✓ 延長端粒的關鍵營養素就是NAD+，隨著年齡增長，NAD+斷崖式驟減，且難以從飲食獲得，導致身體機能逐年衰退。 ✓ 頻尿與便祕、五官感受力衰退、睡眠困擾、皮膚變差、肌肉量減少、大腦功能衰退，這些都是開始衰老的徵兆。
F 特色	1. 天然藜麥發芽種子萃取NADH 2. 目前唯一能有效吃到足量的原料 3. 產生抗老因子NAD+速度快、含量足 4. 搭配天然來源專利緩釋型維生素C
A 作用	1. 煥白Q彈：美國專利緩釋型維生素C，促進膠原蛋白合成，減少皺紋。 2. 青春回朔：德國專利NADH，合成抗老因子，啟動體內抗衰老基因，青春抗老。 3. 協同抗氧：西班牙柑橘維生素P，協同維生素C作用，加成抗氧化作用，延長體內維生素C活性。 4. 歐洲藥典：接骨木莓萃取，抗氧化、抗病度、提升免疫，歐洲的萬能草藥。
B 功效	1. 適用目標：青春抗老、養顏美容、強健體力、大腦健康、免疫提升

米薈國際事業有限公司

產品介紹

適用症狀

大腦健康　強化體力　心血管健康　  　青春抗老　免疫提升

目標客群


- 精神壓力大的人
- 注意力不集中/記憶障礙者
- 有永久壓力的人
- 情緒低落的人
- 加速衰老的人
- 時差亂的人

使用方式

保養劑量-每日3粒
加強劑量-每日6粒

主　成　分

- PurewayC複方維生素C
- Pandmol NADH菸鹼醯胺複合物
- 接骨木莓萃取物
- 柑橘萃取物



產 品 規 格

- 硬膠囊
- 90粒/盒
- 維生素C 900mg
- NADH菸鹼醯胺複合物 50mg

攝 取 建 議

- 3粒/天，建議『空腹、餐前』食用。
- 13歲以上可食用。
- 孕期、哺乳期不建議食用。

適 用 族 群

- 精神壓力大
- 注意力不集中
- 記憶障礙者
- 有永久壓力的人
- 情緒低落的人
- 希望延緩衰老者
- 追求青春美麗者
- 時差亂的人

Milly
INTERNATIONAL CO., LTD
米蕾國際事業有限公司

THANKS

感謝您的聆聽