

Global rise in forest fire emissions linked to climate change in the extratropics

Authors: Matthew W. Jones^{*1}, Sander Veraverbeke^{1,2}, Niels Andela³, Stefan H. Doerr⁴, Crystal Kolden⁵, Guilherme Mataveli^{1,6}, M. Lucrecia Pettinari⁷, Corinne Le Quéré¹, Thais M. Rosan⁸, Guido R. van der Werf⁹, Dave van Wees^{2,3}, John T. Abatzoglou⁵

Affiliations:

¹Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia (UEA); Norwich, UK.

²Faculty of Science, Vrije Universiteit Amsterdam; Amsterdam, The Netherlands.

³BeZero Carbon Ltd.; London, UK.

⁴Centre for Wildfire Research, Swansea University; Swansea, UK.

⁵Department of Management of Complex Systems, University of California Merced; Merced, California, USA.

⁶Earth Observation and Geoinformatics Division, National Institute for Space Research (INPE); São José dos Campos, Brazil.

⁷Department of Geology, Geography and the Environment, Universidad de Alcalá; Madrid, Spain

⁸Faculty of Environment, Science and Economy, University of Exeter; Exeter, UK.

⁹Department of Meteorology and Air Quality, Environmental Sciences Group, Wageningen University; Wageningen, The Netherlands

*Corresponding author. Email: matthew.w.jones@uea.ac.uk

Abstract:

Climate change increases fire-favorable weather in forests, but fire trends are also affected by multiple other controlling factors that are difficult to untangle. Here, we use machine learning to systematically group forest ecoregions into twelve global forest pyromes, with each showing distinct sensitivities to climatic, human, and vegetation controls. This delineation revealed that rapidly increasing forest fire emissions in extratropical pyromes, linked to climate change, offset declining emissions in tropical pyromes during 2001-2023. Annual emissions tripled in one extratropical pyrome due to increases in fire-favorable weather, compounded by increased forest cover and productivity. This contributed to a 60% increase in forest fire carbon emissions from forest ecoregions globally. Our results highlight the increasing vulnerability of forests and their carbon stocks to fire disturbance under climate change.

One-Sentence Summary: Extratropical forest fire emissions are increasing as climate change promotes fire-favorable weather and greening.

Fire is a natural ecosystem disturbance that has shaped the global distribution of Earth's forests and controlled carbon (C) storage in vegetation and soils over geological time (1–3).

Nonetheless, anthropogenic climate change has contributed to an increase in fire-favorable weather conditions globally (4–7) and these enhanced risks have translated into increased burned area (BA) and fire C emissions in some forested regions during the past two decades or longer (6–12). Expanding land use or historic fire management policies have variably interacted with the effects of climate change to amplify forest fire activity and emissions (13, 14). The increases in forest fire C emissions observed regionally contrasts with declines in the global savannahs (8, 15).

A series of highly anomalous episodes of extreme forest fire C emissions have recently punctuated longer-term trends (11, 16–19). During the 2019–2020 bushfire season in Australia, the area burned by fires was over double the previous record since 1930, and fire C emissions were also greater than in any other year since 2003 (9, 17). In 2021, a new record was set for pan-boreal fire C emissions amidst a water deficit spanning both Eurasia and North America (16). In the 2023 fire season, fire C emissions from Canadian boreal forests were over nine times the 2001–2022 average (19).

The increased occurrence of fire, and particularly extreme fires, threatens the functioning and resilience of some forests as well as their ecosystem services, including C storage (13, 20, 21). The recovery of C stocks in vegetation and organic soils following forest fires can take decades to centuries, and so increases in annual fire C emissions and extreme emissions events lead to a lasting deficit of terrestrial C storage (8, 22–24). Increased fire C emissions can thus reduce the capacity of global forests to absorb C from the atmosphere, posing a challenge for achieving climate targets. For example, increased fire activity in boreal North America alone is projected to result in net C losses equivalent to 0.3–3% of the remaining C budget necessary to limit global warming to 1.5°C (25).

Beyond their effects on C storage, extreme wildfires also cause major disruption or irreversible loss to society, including deaths, evacuations, reduced air quality, pressures on healthcare systems, and economic losses (26–30). Further, major declines in biodiversity have also been recorded in the wake of several extreme fire events and many of Earth’s most threatened species are afflicted by an altered fire regime (1, 20). Recent extreme wildfire seasons across the globe have demonstrated the power of the most extreme wildfires to affect both the environment and society.

One of the drivers of change in forest fire potential is anthropogenic climate change, which is causing more frequent and extreme periods of drought and fire-favorable weather, often referred to as fire weather (4, 6, 31). Increased hot and dry conditions create periods of low fuel moisture, promoting wildfire potential in ecosystems where ample stocks of fuels (vegetation biomass and organic soils) are available, notably in forests (4, 10, 31). Increased lightning frequency under climate change has also exacerbated the ignition of forest fires in some locations, particularly in ignition-limited forests of the high latitudes (32–34). Increased atmospheric instability has been linked to more erratic and extreme wildfire behavior that enhances fire spread and intensity and challenges the potential for firefighters to suppress fire (35, 36). Several attribution studies have shown that climate change raised the likelihood of extreme fire weather conditions during a range of recent extreme wildfire seasons (5, 37, 38).

Alongside climatic factors, forest fire extent is controlled by various *in situ* human activities and by the ecological traits and productivity of vegetation (6, 39–42). People influence patterns of forest fire in numerous ways, such as by using fire for forest clearing and land use (43), causing unwanted ignitions (accidental or arson) (44), suppressing wildfires via firefighting (45), managing stocks of fuel on the landscape (46, 47), increasing forest edge length through fragmentation (48, 49), or inadvertently amplifying fuel stocks by excluding fire from forests where it is a central element of a functioning ecosystem (14, 50). The composition of forest ecosystems with species that have developed fire-adapted evolutionary traits, such as canopy


1 structure, self-pruning, and leaf waxiness, also influence fire dynamics (e.g. crowning potential)
2 and rates of spread (51, 52). In addition, the productivity of vegetation during the growing season
3 influences fuel availability during subsequent dry seasons (40, 53, 54).

4 While climatic, human, and vegetation factors all affect patterns of fire in forests, the
5 prominence of each control varies regionally (6, 40, 55, 56). The relationships between climate
6 and fire are generally modulated by non-climatic factors, and likewise non-climatic drivers of fire
7 often depend on the episodes of fire-favorable weather. Hence, it has proven remarkably
8 challenging to identify the forest regions where fires are most sensitive to climate change or other
9 facets of environmental change (39, 55, 56). To identify the world regions where responses to
10 future climate change or other environmental stressors are comparatively strong or weak, further
11 study of the temporal and spatial relationships between fire and a comprehensive set of fire controls
12 is required.

13 Here, we used the *k*-means clustering algorithm to group 414 forest ecoregions of the
14 World (57) into twelve forest *pyromes* (**Fig. 1**), within which forest burned area (BA) (58) shares
15 a common set of relationships with climatic, vegetation, and human controls (Fig. 2, Fig. S1-S2;
16 see **Methods**). Having isolated the pyromes with a distinctive strong sensitivity to climatic
17 controls, we analysed trends in annual forest BA (58) and fire C emissions (59) during the period
18 2001-2023 and evaluated their connection with trends in key climate variables.

19 We used a comprehensive set of fire controls to distinguish the pyromes. The climatic
20 controls included fire weather (4, 60, 61), soil moisture (62), atmospheric instability (represented
21 by the continuous Haines index) (35, 36), and lightning frequency (34, 63). The vegetation controls
22 included potential fuel stocks related to land cover (52), vegetation productivity during the
23 growing season (represented by the normalized difference vegetation index) (64, 65), and forest
24 continuity (represented by forest area density) (49). The human controls included population
25 density (15, 66), cropland and pasture cover (15, 67, 68), and road density (69). Terrain roughness
26 (70) is also included for its potential to affect fire behavior (71). These variables have each shown
27 power to explain spatial or temporal variability in BA in at least some world regions (see
28 **Methods**).

29 The concept of the pyrome was first introduced by Archibald et al. (72) as a
30 pyrogeographical counterpart to the biogeographical concept of the biome. Biomes are defined not
31 only by their observable biological characteristics but also by the climatic and other environmental
32 controls that cause particular biological characteristics to arise. In past work, pyromes have been
33 characterized only by observable fire characteristics such as size, duration, intensity, and frequency
34 (72–75). Here, we expand the pyrome concept to include a systematic grouping of ecoregions
35 based on the strength of climatic and other environmental controls on fire. This novel approach
36 enriches the pyrome concept with a dimension that mirrors the complexity inherent in the study of
37 biomes, while also providing critical insights into the varying sensitivity of pyromes to different
38 facets of global change. Delineating the global forest pyromes revealed a rapid increase in
39 emissions from extratropical forest pyromes that exceeded declining emissions from tropical forest
40 pyromes during 2001-2023. This increase demonstrates that climatic controls on forest fire are
41 overwhelming human controls in global-scale emissions trends.



Pyrome	Associated Forest Biomes	Associated N-grams from Ecoregion Descriptions
ExTropF1	Boreal forests/taiga; Temperate broadleaf and mixed forests; Temperate coniferous forests	mixed forests; canadian shield; forests canadian; siberian taiga
ExTropF2	Tropical and subtropical moist broadleaf forests; Tropical and subtropical dry broadleaf forests; Temperate broadleaf and mixed forests; Temperate coniferous forests	atlantic; deciduous forests; evergreen forests; mixed forests; conifer forests; dry forests; sclerophyllous; subtropical
ExTropF3	Temperate broadleaf and mixed forests; Tropical and subtropical moist broadleaf forests; Temperate coniferous forests; Mediterranean forests, woodlands, and scrub or sclerophyll forests	mixed forests; mountains; dry forests; conifer; deciduous forests; moist forests; sclerophyllous
ExTropF4	Tropical and subtropical moist broadleaf forests	moist forests; conifer forests; lowland rain forests; montane rain forests; new; sumatran; swamp forests
SubTropF1	Temperate broadleaf and mixed forests; Temperate coniferous forests; Tropical and subtropical moist broadleaf forests; Mediterranean forests, woodlands, and scrub or sclerophyll forests	mixed forests; montane; broadleaf forests; conifer forests; forest steppe; forests himalayan; kurile; sclerophyllous; swamp forests
SubTropF2	Tropical and subtropical moist broadleaf forests; Tropical and subtropical dry broadleaf forests; Tropical and subtropical coniferous forests; Mediterranean forests, woodlands, and scrub or sclerophyll forests	dry forests; pine oak forests; montane forests; rain forests; sierra madre
SubTropF3	Tropical and subtropical moist broadleaf forests; Tropical and subtropical dry broadleaf forests	dry deciduous forests; rain forests
TropF1	Tropical and subtropical moist broadleaf forests	moist forests; congoian lowland forests; swamp forests
TropF2	Tropical and subtropical moist broadleaf forests; Tropical and subtropical dry broadleaf forests	dry forests; lowland; montane rain forests; coastal forests; moist forests; pine; swamp forests
TropF3	Tropical and subtropical moist broadleaf forests; Tropical and subtropical coniferous forests	moist deciduous forests; subtropical
SupZoF1	Tropical and subtropical moist broadleaf forests; Tropical and subtropical dry broadleaf forests; Tropical and subtropical coniferous forests	dry forests; montane forests; pine
SupZoF2	Mediterranean forests, woodlands, and scrub or sclerophyll forests; Tropical and subtropical moist broadleaf forests; Temperate coniferous forests; Temperate broadleaf and mixed forests	forests; woodlands; coastal

Figure 1: World map of the twelve forest pyromes and a summary of their tendencies to associate with biomes and ecoregion types. 414 forest ecoregions are attributed to one pyrome using k-means clustering, which identifies ecoregions sharing a similar set of correlations between burned area (BA) and 14 predictor variables (Fig. 2 and Fig. S1-S2). Grey areas are not included in the analysis, either because they are not within forest biomes (light grey) or because fire is extremely rare (the mean annual fraction of forest area burned by fire is below 0.01%; dark grey). Fig. S21 shows an alternative mapping of the pyromes for the ecoregions that were clustered most ambiguously. The table shows the most common biome associations for each pyrome and the most common text substrings (n-grams, up to 3 words) that appear in the ecoregion descriptions, based on the ecoregion descriptions in the Terrestrial Ecoregions of the World dataset (ref. (57)).

Results

Twelve global forest pyromes emerged from the clustering analysis (Fig. 1), and this grouping revealed large increases in forest fire C emissions in extratropical forest pyromes. By contrast, forest fire C emissions declined in tropical and subtropical forest pyromes.

Geography and Traits of the Pyromes

To characterize the key controls on fire in each pyrome, we examined the correlations between forest BA and each variable representing the fire controls amongst the constituent ecoregions of the pyromes (Fig. 2, Fig. S1-S2). We consider the mean correlations within each pyrome to indicate sensitivity of the forest BA response to each control, and we consider these relationships to be a trait of a pyrome when at least 75 percent of the constituent ecoregions display a correlation of the same sign (Fig. 2, Fig. S1-S2). Significant differences in correlation were observed in 58% of pairwise comparisons, indicating a robust grouping of the ecoregions into pyromes with distinctive fire controls (see Methods). A more complete description of pyrome characteristics is provided in Supplementary Text 2.

Pyromes in Extratropical Forests

Pyromes ExTropF1 and ExTropF2 encompass the North American and Eurasian boreal forests and some temperate and high-altitude tropical forests (Fig. 1). Forest BA in these pyromes correlates positively with fire weather and atmospheric instability, negatively with seasonal soil moisture, and shows no correlation with population density, agricultural land cover, and road density (Fig. 2, Fig. S1-S2). ExTropF1, more common in North America, has BA strongly correlated with lightning flash density, indicating lightning as a key ignition source (33, 34). In ExTropF2, more common in Eurasia, BA correlates with NDVI from the prior growing season, suggesting that previous climatic conditions impacting vegetation growth and the production of fine fuels bear an influence on subsequent fire extent (76, 77). Forest fire extent in pyromes ExTropF1 and ExTropF2 is governed by different combinations of climatic factors affecting fuel moisture, fuel growth, and natural ignition.

Pyromes ExTropF3 and ExTropF4 include forests of Scandinavia, western Russia, and certain areas of North American, Europe, and China (Fig. 1). While BA in these pyromes correlates with fire weather, the strength of correlations is lower than in ExTropF1 or ExTropF2 and especially weak in ExTropF4 (Fig. 2, Fig. S1-S2). Additionally, in ExTropF4, no correlation with soil moisture further indicates that fires are also relatively insensitive to water deficits (Fig. 2, Fig. S1-S2). Weak correlations with most variables in ExTropF3 and ExTropF4 likely relate to infrequent burning in these stable humid climates (Fig. S8, Table S1), which challenges diagnoses of controls on fire over a two-decade time period.

Pyromes in Tropical Forests

TropF1 and TropF2 are widespread in the tropical deforestation zones of Amazonia, Congo, and equatorial southeast Asia (Fig. 1). Forest BA correlates positively with population density, road density, agricultural land cover, and fire weather, and negatively with forest continuity and soil moisture (Fig. 2, Fig. S1-S2). These traits characterize a region with widespread deforestation and degradation fires that, particularly in TropF2, are facilitated by dry conditions (48, 78, 79). TropF1, primarily in Amazonia and the Congo, shows a strong correlation with pasture cover (Fig. S1), reflecting cattle ranching-driven deforestation (80, 81). TropF2, found in

Sumatra, Kalimantan, Borneo, Guianas, and southeast Russia, has stronger correlations with soil moisture and weaker with population, roads, and pasture cover than TropF1, highlighting particularly prominent role of drought in facilitating peak fire activity (79, 82, 83). Several forest ecoregions in southeast Russia, a global hotspot of extratropical forest loss through fire linked to forestry operations (83), are also grouped with TropF2 (Fig. 1).

TropF3 characteristically maps to older, heavily-fragmented deforestation frontiers in Brazil, Mexico, West Africa, and some southeast Asian islands (Fig. 1) (84, 85). Here, forest BA correlates positively with fire weather and negatively with soil moisture but not with population or agriculture density, most likely caused by saturation of ignition sources in these highly populated regions during fire-favorable weather conditions.

Pyromes in Subtropical Forests

SubTropF1, SubTropF2, and SubTropF3 span subtropical or dry tropical forest ecoregions in regions such as northern Colombia, Madagascar, northeast India, southeast Asia, Sri Lanka, East Africa, and drier parts of the Brazilian atlantic forests (Fig. 1). BA consistently correlates with forest continuity and fuel stocks in these pyromes, in addition to fire weather (Fig. 2, Fig. S1-S2). This indicates a tendency for greater fire extent when meteorological conditions allow in locations where fuel production is greater or flammable natural landscapes are less fragmented (15, 49). In SubTropF1 and SubTropF2, negative correlations with population density, cropland cover, and road density, particularly strong in SubTropF2, suggest reduced fire activity in the areas most fragmented by human activity. Lightning frequency often correlates with BA in SubTropF2, pointing towards a greater frequency of natural ignitions.

In contrast to SubTropF1 and SubTropF2, BA in SubTropF3 shows no consistent correlation with population density, cropland cover, or road density, indicating that natural factors (e.g. topographic and hydrological) are more important controls on fuel loads and continuity than human factors. SubTropF3 also lacks strong correlations between BA and soil moisture, indicating sensitivity to short-term fire-prone weather rather than seasonal moisture deficits (15, 49).

Pyromes in Zones of Fire Suppression

The final two pyromes, **SupZoF1** and **SupZoF2**, span tropical, subtropical, and temperate forest ecoregions and are common in regions with significant fire management efforts, such as the southeast US, western US, southeast and western Australia, and parts of Iberia (Fig. 1). In these pyromes, forest BA negatively correlates with population density, road density, and agriculture, indicating reduced fire extent in proximity to human activities (46, 86). Positive correlations with forest continuity and fire spread suggests that continuous forests facilitate fire spread, especially in topographically complex areas with fewer human activities (46, 87).

Fire suppression, fuel load management, and community programs are in place to reduce fire extent in these areas (46, 88). Despite these efforts, fires can still occur during fire-prone weather (37, 89), and stronger positive correlations with fire weather and negative correlations with soil moisture in **SupZoF2** suggest that climatic factors bear stronger influence on forest BA than in **SupZoF1** (Fig. 2). In **SupZoF1**, forest BA often (but inconsistently) correlates with lightning frequency, highlighting the role of natural ignitions, an effect that is seen most strongly in southeast Australia (90) (Fig. S2). Additionally, **SupZoF1** shows a correlation between forest BA and vegetation productivity from the prior growing season, emphasizing the role of fuel production as a driver of fire extent (9, 42, 91).

1




Figure 2: Variation in the relationship between forest burned area (BA) and nine predictors across the global forest pyromes. The violins plot the kernel density distribution of correlations values (spearman's ρ) for each predictor amongst the constituent ecoregions of each pyrome. White dots mark the median correlation value for the ecoregions of a pyrome, black line ranges mark the interquartile range, and open diamonds mark the mean value. See **Methods** for a description of all correlation analyses and the motivation for including each predictor. Distributions are shown for all predictor variables in Fig. S1. Correlations are mapped for each forest ecoregion in Fig. S2.

Increased Fire Emissions in Extratropical Forest Pyromes

Forest fire C emissions increased in several of the pyromes between 2001 and 2023 (**Fig. 3**), however the most striking trend was a 194% increase in fire C emissions in pyrome **ExTropF2** (+116 Tg C year⁻¹; **Table S1**). This large increase in fire C emissions was driven by a 167% increase in forest BA (+35 thousand km² year⁻¹) and a 58% increase in C combustion rate (C emissions per unit BA; **Fig. 3, Table S1**). Increased forest BA was a widespread feature of the ecoregions in pyrome **ExTropF2**, with over half showing significant increases and fewer than 5% showing significant decreases (**Fig. 4, Fig. S11**). Consequently, the increases in forest fire C emissions were also widespread. For example, forest fire C emissions increased significantly in parts of Russia (east and northeast Siberian taiga), Europe (e.g. Balkan mixed forests, Pindus and Dinaric mountains mixed forests), western North America (e.g. Sierra Nevada forests, North-Central Rockies forests, Muskwa-Slave lake forests, Fraser Plateau and Basin complex, and Northwest Territories taiga), Chile (Valdivian temperate forests), and China (Northeast China Plain deciduous forests and Hengduan Mountains conifer forests; **Fig. 4**).

The increases in forest BA and fire C emissions in pyrome **ExTropF2** align with changes in the variables that control temporal variability in forest BA. During 2001-2023, the annual number of extreme fire weather days increased by 5 days per year on average across the ecoregions of the pyrome (**Fig. 4, Fig. S14, Table S2**). The average soil moisture content during the fire season reduced by around 3% on average, in contrast to other extratropical pyromes where soil moisture either increased or remained level (**Fig. 4, Fig. S15, Table S2**). Mean NDVI during the growing season also increased at a rate comparable to the other extratropical pyromes (**Fig. 4, Fig. S16, Table S2**). These trends were also widespread and consistent. For example, over half of the ecoregions in pyrome **ExTropF2** synchronously experienced an increase in extreme fire weather days, increased NDVI and reduced soil moisture, with over one-quarter of ecoregions showing *significant* changes for all three variables. This evidence suggests that the trends in forest BA and fire C emissions in pyrome **ExTropF2** were driven by changes in the climate of the fire season, which led to reduced fuel moisture, combined with changes in the climate of the growing season, which led to increased vegetation growth and fuel production.

The increase in forest productivity during the growing season in pyrome **ExTropF2** also corresponds with a 30% increase in forest area in the pyrome (+1 million km²; **Fig. S3**) during 2001-2023. This striking rate of forest expansion is consistent with reported rates exceeding 1% per year in some of these regions during 2001-2019 based on MODIS observations (92) and with the accumulating evidence for increased vegetation greenness and biomass stocks in the high-latitudes (77, 93, 94). On the other hand, such a large increase in forest area has not been seen in Landsat-based estimates of change in forest cover, likely due to differences in the resolution (see further discussion in **Methods**) (95–97). Dual increases in forest area and productivity highlight how a warming climate and CO₂ fertilization have enhanced forest growth at higher latitudes and contributed to both a greater forest area available to burn and greater rates of fuel production (98, 99). Nonetheless, growth in forest BA has outpaced growth in forest area, as indicated by a 158% increase in the fraction of forest area that burned annually during 2001-2023 (i.e. the forest BA fraction; **Table S1, Fig. S3**).

Pyrome **ExTropF1** also showed large and significant increases in forest BA (+30%, or 6 thousand km² year⁻¹) and fire C emissions (+65%, or 30 Tg C year⁻¹) during 2001-2023, though these trends were not quite as pronounced as in pyrome **ExTropF2** (**Fig. 3, Table S1**). Trends in forest BA were also more mixed amongst the ecoregions of pyrome **ExTropF1**, with around one-third showing a significant increase in forest BA and around 15% showing a significant decrease (**Fig. 4, Fig. S11, Table S1**). The varied trends in forest BA can be explained by mixed trends in

fire weather and soil moisture across the pyrome. The annual number of extreme fire weather days increased significantly in around 35% of the ecoregions of pyrome **ExTropF1** but also decreased in 15%, while very few ecoregions experienced a significant increase in soil moisture (**Fig. 4, Fig. S14-S15, Table S2**). Only around 10% of ecoregions in pyrome **ExTropF1** showed synchronous significant increases in extreme fire weather, soil moisture and lightning density, which are the three key controls on fire activity that emerged from our clustering analysis in this pyrome (**Fig. 2, Fig. S1-S2**).

Increased Fire Emissions in Pyromes with Fire Suppression

In pyromes **SupZoF1** and **SupZoF2**, forest fire C emissions increased by 43-44% during 2001-2023 (**Fig. 3, Table S1**). In both pyromes **SupZoF1** and **SupZoF2**, the increased fire C emissions were driven primarily by significant 37-79% increases in the C combustion rate, combined with smaller non-significant increases in forest BA of 8-18% (**Fig. 3, Table S1**). Within pyromes **SupZoF1** and **SupZoF2**, significant increases in fire C emissions were spatially concentrated in forest ecoregions of Australia (e.g. Blue Mountains forests, Naracoorte woodlands, Jarrah-Karri forests), southern Europe (e.g. Southwest Iberian Mediterranean sclerophyllous forests, Northwest Iberian montane forests, and Aegean and Western Turkey sclerophyllous forests), the western USA (Klamath-Siskiyou and coastal forests and California interior chaparral and woodlands), Madagascar (subhumid and lowland forests; **Fig. 4**). The large upticks in emissions from forests in the western US and eastern Australia during 2019 and 2020 (17, 18) are clearly visible in the emission time series for pyromes **SupZoF1** and **SupZoF2** and influence the slope of the trends in these pyromes (**Fig. 3**).

Reduced Fire Emissions in Tropical Forests

Forest fire C emissions showed opposing trends in the pyromes occupying the tropical deforestation zones, with the 96% decline ($-26 \text{ Tg C year}^{-1}$) in forest fire C emissions in pyrome **TropF2** outweighing 56% increases ($+24 \text{ Tg C year}^{-1}$) in forest fire C emissions in pyrome **TropF1** (**Fig. 3**). In pyrome **TropF1**, the increase in fire C emissions was caused by an increase in forest fire C combustion rate combined with a 19% increase in forest BA (**Fig. 3, Table S1**). Increased forest BA was widespread throughout the pyrome, with around two-thirds of the constituent ecoregions of pyrome **TropF1** showing an increase during 2001-2023 (**Fig. 4; Table S1**). In pyrome **TropF2**, the decline in forest fire C emissions was predominantly driven by a 56% reduction in forest BA (**Fig. 3; Table S1**). The reductions in fire C emissions were consistent across the ecoregions of pyrome **TropF2**, with around 80% of the constituent ecoregions of the pyrome showing a decrease during 2001-2023 (**Fig. 4, Fig. S11, Table S1**).




Figure 3: Changes in the burned area (BA) in forests and carbon (C) emissions from forest fires during 2001-2023. By row, the panels show (a-c) forest BA, (d-f) fire C emissions per unit burned area of forest, (g-i) C emissions from forest fires, and (j-l) the forest fraction of total (forest plus non-forest) fire C emissions. By column the panels show (a,d,g,j) show annual data (solid lines) and trendlines (dashed lines) for each pyrome, (b,e,h,k) absolute changes during the data period, and (c,f,i,l) relative changes (%) for the same period. Trendlines are fitted using Theil-Sen regression. Fire C emissions are extrapolated for 2001 and 2023 based on the trend in C combustion rate during 2002-2020 and the observed annual BA in 2001 and 2023. Absolute

1 changes are calculated as the difference between the trendline values at the start and end of the
2 period, and relative changes are calculated conservatively as the absolute change divided by the
3 period mean. **Figs. S3-S5** present various aspects of forest and total (forest + non-forest) fire
4 trends, including changes in burned area, C emissions, and combustion rates. **Figs. S6-S11** show
5 mapped trends for individual forest ecoregions and the distribution of these values across the
6 ecoregions of each pyrome.
7




Figure 4: Changes in bioclimatic variables and fire observations at ecoregion level. The panels show relative changes in four bioclimatic variables used to distinguish pyromes(a-d), forest burned area (BA; e), forest burned area fraction (km^2 burned km^{-2} forest; f), forest fire carbon (C) emissions (g) and the forest fire C combustion rate (g C km^{-2} burned) during the period 2001-2023 (2010-2021 for lightning flash density), mapped to forest ecoregions. The climate variables

are (a) days during the fire season with 95th percentile fire weather index (FWI) values relative to all days in the period 1980-2009, (b) average soil moisture content during the fire season, (c) mean normalized difference vegetation index (NDVI) during the prior growing season, and (d) lightning flash density during the fire season. The strength of the relationships between forest BA and fire weather, soil moisture, vegetation growth and lightning varies across pyromes (Fig. 2) and there is lesser variability within the constituent ecoregions of each pyrome (Fig. S1-S2). Changes in forest fire C emissions (g) relate to changes in forest burned area (e) as well and emissions per unit BA (h). Fig. S6-S17 additionally show the mapped mean values of these variables for individual ecoregions and the distribution of trends across the ecoregions of each pyrome.

Discussion

Our mapping of pyromes has revealed variation in the controls on forest fire extent across the world's forest ecoregions. The pyrome boundaries tend to align with the boundaries of climate zones or biomes (Fig. 1), though not precisely due to significant variation in the drivers of fire within those zones. The expected first-order patterns of macro-scale pyrogeography are apparent in the distinctive traits of fire control that emerge from the clustering analysis (Fig. 2, Fig. S1-S2). For example, human controls on fire emerge as a stronger trait of the tropical pyromes than of the extratropical pyromes, consistent with the expectation that the tropical fire regime is dominated by human activities (2, 100, 101).

In the largest extratropical pyromes (ExTropF1 and ExTropF2), forest fires are influenced by climatic factors affecting fuel moisture during the fire season and variably by the production of vegetation fuels in the growing season or by opportunities for lightning ignition. The near tripling of fire C emissions in pyrome ExTropF2 can be explained by pervasive increases in fire-favorable weather during the fire season, increased vegetation productivity in the growing season, and expanding forest cover. Increased forest extent and productivity at higher latitudes have been linked to climate changes that are favorable for vegetation growth and CO₂ fertilization (98), and our results show that these trends have coupled with reduced fuel moisture to drive an increase in forest BA and fire C emissions in pyrome ExTropF2. The weaker increases in fire C emissions in pyrome ExTropF1 can be attributed to less consistent trends in fire-favorable weather across the pyrome.

During 2001-2023, forest fire C emissions grew by 60% across all forest ecoregions globally, principally driven by trends in the extratropical pyromes (Fig. 3, Table S1). Forest BA and fire C emissions were redistributed from tropical and subtropical pyromes to extratropical pyromes (Fig. 5). Amidst these geographical shifts, the C combustion rate of forest fires also increased by 47% across all forest ecoregions globally, reflecting greater fuel consumption per unit of forest BA (Fig. 3, Table S1). Extreme examples of C combustion per unit area have been recorded during recent extreme wildfire episodes and tied to extremes in fire-favorable weather (8, 11, 16, 17, 102), whereas our findings support a more general trend towards increases in fuel consumption in forests. In addition, the contribution of forest fires to total (forest plus non-forest) BA and fire C emissions has also increased globally and in most forest pyromes, with the exception of the subtropical forest pyromes (Fig. 3, Table S1), signaling that the increased susceptibility of forests to fire has generally outpaced that of non-forest environments experiencing similar environmental changes.

Our mapping of the pyromes enabled us to link rising fire C emissions in extratropical forests to climate change. For example, without distinguishing pyrome ExTropF2 and the climatic factors that influence its forest BA, the increase in fire emissions that has occurred there could be

overlooked or masked. The emissions trend in pyrome **ExTropF2** potentially signals that a step-change in the fire regime and a destabilization of forest C stocks is underway in some extratropical ecoregions. Recent studies have identified a rise in forest productivity and fire-favorable weather as compounding drivers of increased fire C emissions in Siberia (16, 103–105), whereas our results indicate that similar dynamics are leading to increased fire C emissions more broadly across the ecoregions of pyrome **ExTropF2**.

Even in the pyromes where non-climatic factors, particularly *in situ* human activities, exert significant control on forest BA, climatic factors remain a key enabler of fire. Increased fire weather under climate change can be expected to increase the windows of opportunity for fires to occur even in regions with significant fire suppression (31, 106, 107). For example, the uptick in the forest BA and fire C emissions in the pyromes **SupZoF1** and **SupZoF2**, which encompass many zones of aggressive fire suppression and management, are consistent with warnings that the effectiveness of wildfire suppression is waning in a warming climate (108, 109). These findings highlight a potential for the relationships between climate and fire to strengthen in future climates.

Forest ecoregions with the most ambiguous cluster assignment (as measured by silhouette width statistics), are scattered globally with little tendency to concentrate in particular world regions (Fig. S21), suggesting that the pyromes arising from our clustering procedure were largely free of regional bias (Supplementary Text 1). One exception is the Iberian Peninsula, where almost all ecoregions showed low silhouette widths, indicating a relatively low parity with other world regions. Among pyromes, clustering ambiguity was highest in SubTropF3, SupZoF1, and SupZoF2, with the most common alternative assignments to pyromes ExTropF1, ExTropF2, ExTropF3, ExTropF4, and SubTropF3, indicating a higher level of confusion between various climate-sensitive and extratropical pyromes (Supplementary Text 1).

Overall, we have contributed a new geographical mapping of forest pyromes based on distinctive fire drivers and discovered significant increases in forest BA and fire C emissions in some of the pyromes where they are most expected. Our work complements previous studies that used machine learning to disentangle the effect of multiple fire controls on global patterns or trends in BA (7, 40, 55, 56). For example, prior studies also indicated that increased vegetation productivity and fire-favorable weather both contributed to increased BA in boreal Eurasia during 2001–2014 (7, 40). Our explicit focus on forest BA and fire C emissions has also provided novel insights. For example, we find strong spatial contrasts in the effect of human activities on forest BA across different tropical and subtropical pyromes, whereas prior work indicated that human activities reduce total BA more uniformly across the tropics (7, 40).

Our work complements prior endeavors to define pyromes based on observable fire characteristics (72–75). While a novel and insightful aspect of our study is its focus on grouping regions with similar fire drivers, by doing so, we concentrated exclusively on BA as a target variable, foregoing information about other observable fire traits that vary geographically and are important aspects of the fire regime. Future work could aim to integrate geographical distinctions in both fire traits and fire drivers to provide a more holistic definition of the pyrome. This approach would further enhance the analogy with the term ‘biome’, which encapsulates both the biological properties and physical presentation of grouped ecosystems, as well as the climatic and other environmental factors that cause those properties to emerge.

Looking forward, our pyrome classification could play a key role in the development of global fire models to better represent observed fire dynamics by creating opportunities to tailor model parameters in regions with distinct fire drivers. For example, parameters that represent the influence of people on fire processes could be optimized by pyrome in DGVMs to better represent the distinct relationships between human activities and fire across pyromes, in a manner akin to

optimizing biological processes across plant functional types. Moreover, the pyromes layer also serves to highlight priority areas for the study of changes in fire weather, drought or vegetation productivity, since some pyromes are distinctly more sensitive to changes in these factors than others.

A caveat of our approach is that it provides a global zonation of fire controls at the macro-scale – a scale that is particularly suited to questions concerning global environmental change, including differential responses to climate change. We do not suggest that all areas within an ecoregion are uniformly sensitive to the same fire controls. For example, differences in land use and management approaches across landowner types can be expected to produce varying relationships between fire and human factors within an ecoregion, as seen between protected areas, Indigenous areas, and private land (110, 111). Hence, our analysis identifies the dominant controls that emerge at the ecoregion scale but omits the local effects associated with specific actors at sub-ecoregion level. The application of similar techniques to smaller (or larger) world regions would provide a finer (or coarser) geography of fire controls to which a different set of environmental questions may apply. In addition, our mapping of pyromes should not be viewed as fixed in time. For example, regional changes in policy, land use or population dynamics or ongoing shifts in climate or vegetation types could all lead to the re-allocation of an ecoregion to a different pyrome in future (75).

Relatedly, although our analysis provides valuable insights into the impacts of climate change on fire dynamics over a two-decade period, it is important to recognize the limitations inherent in using relatively short datasets to interpret fire regimes that operate over much longer intervals. Many forest ecosystems are subject to fire return intervals spanning decades to centuries, which can obscure the detection of longer-term trends. This is particularly the case in pyromes ExTropF3 and ExTropF4, where long fire return intervals (~1,000 years; **Figure S7; Table S1**) likely contributed to low correlations between BA and all explanatory variables and challenged the identification of fire drivers by clustering. Therefore, while our findings indicate significant trends such as the increase in emissions in the extratropics, increased combustion rates, and a shift from savannas and grasslands to forests as major fire emissions sources, these must be interpreted with some caution. Future studies extending beyond the 20-year timeframe are essential to fully understand the long-term fire regimes and validate the persistence of such trends (9, 112).

Both observations and models suggest that extratropical forests are greening and becoming more productive due to a combination of climate change and CO₂ fertilization (98, 99, 113). Dynamic global vegetation models (DGVMs) also generally project that C storage will continue to increase in the future in high latitude forests, although some variability is seen across models and climate scenarios (114, 115). Nonetheless, DGVMs currently show a limited capacity to reproduce historical trends and contemporary spatial patterns of fire (6, 116), raising concerns as to whether future change in fire disturbance is reliably captured in projections of future vegetation distribution and C storage. Additional uncertainty in future C storage stems from the potential for post-fire ecosystem shifts to occur due to increased fire severity not captured by models (117, 118). Our finding of increased forest fire C emissions lends further support to previous warnings that fire could offset projected gains in C storage in extratropical forests (119–121).

As forest fire C emissions grow, so does their relevance to carbon accounting, including the greenhouse gas (GHG) inventories submitted to the United Nations (UN). For example, C emissions from wildfires in Canada during 2023 alone are likely to have overturned a significant portion of the C sink to Canadian forests that accumulated over the prior decade (19, 122, 123). Wildfires in Canada are not free of anthropogenic influence and are becoming more likely due to

anthropogenic climate change (19), yet they are designated as natural disturbances in Canada's national emissions inventory and thus their influence is largely omitted from UN records (122, 123). Prior work has advocated for more comprehensive reporting of fire emissions on both managed and unmanaged land, to facilitate routine assessments of how fires impact national and global inventories of anthropogenic emissions (122). Our work further highlights the importance of this comprehensive reporting by revealing the growing role that forest fire C emissions play in the carbon budget of boreal forests.

Relatedly, enhancing C storage in forests using forestry practices is viewed as a promising strategy for C dioxide removal (CDR) from the atmosphere to offset anthropogenic C emissions (124, 125). One recent study estimated that an additional 60 Pg C could be stored on extratropical land that is highly suitable for forestry (126), or 600-3,000 Tg C year⁻¹ when annualized over a period of 20-100 years representing the time taken for potential C stocks to accumulate (127). The estimates of potential C storage from ref. (126) derive from relationships fitted to the C stocks held in current intact forests, yet these forests were established in historical fire regimes inferring that potential C storage is overestimated in forests where fire regime shifts are underway. For a crude comparison, we estimate that forest fire C emissions grew by 114 Tg C year⁻¹ across all extratropical pyromes between 2001 and 2023. We suggest that a continued increase in forest BA and fire C emissions could reduce the potential for CDR in extratropical forests by a nontrivial margin, particularly in the absence of effective fuel and fire management.

While climatic factors show a varying strength of control on the extent of forest fires across pyromes, their effects are nonetheless pervasive. This result emphasizes the need to address the primary causes of climate change, by reducing emissions from fossil and land use sources, in order to mitigate the increased fire-related risks to C sinks (128, 129). Moreover, our findings inform forest management and Net Zero policies by identifying pyromes where specific human actions can support forest C sinks by reducing C emissions from fires. In tropical pyromes, where fire shows a strong dependence on human ignition patterns, reducing ignitions during extreme fire-favorable weather and preventing forest fragmentation should enhance C retention (30, 130). In pyromes with a history of aggressive wildfire suppression, shifting focus and funds from active fire suppression to managed, ecologically beneficial fires may prevent C sink-to-source conversion (7, 30, 131). In extratropical pyromes where climatic factors have the most direct and unmodulated control on fire extent, monitoring changes in vegetation and productivity can guide the prioritization of areas for forest management (7, 30, 131). In all pyromes, substantial financing is required to support strategic programs of forest management, stakeholder engagement, and public education, all of which represent a meaningful shift of fire management strategy from largely reactive to increasingly proactive (7, 30, 131). Overall, global forest C sinks could be undermined by wildfire without action to address the leading causes of climate change, while forest management strategies for mitigating the problem are likely to be most effective when tailored to pyromes. Cutting anthropogenic emissions is central to securing resilient forests for the future.



Figure 5: Geographical shifts in forest burned area (BA) and fire carbon (C) emissions from the tropics to the extratropics during 2001-2023. The plot shows contributions of groups of pyromes in the tropics, subtropics, extratropics and zones of suppression to (a-b) forest BA in all forest ecoregions globally and (c-d) the fire C emissions in all forest ecoregions globally. By column the panels show (a,c) annual data (solid lines) and trendlines (dashed lines) for each pyrome, (b,d) relative changes during 2001-2023. Trendlines are fitted using Theil-Sen regression. Fire C emissions are extrapolated for 2001 and 2023 based on the trend in C combustion rate during 2002-2020 and the observed annual BA in 2001 and 2023. Absolute changes are calculated as the difference between the trendline values at the start and end of the period, and relative changes are calculated conservatively as the absolute change divided by the period mean.

References and Notes

1. L. T. Kelly, K. M. Giljohann, A. Duane, N. Aquilué, S. Archibald, E. Batllori, et al., Fire and biodiversity in the Anthropocene. *Science* 370, eabb0355 (2020).
2. D. M. J. S. Bowman, J. K. Balch, P. Artaxo, W. J. Bond, J. M. Carlson, M. A. Cochrane, et al., Fire in the Earth System. *Science* 324, 481–484 (2009).
3. G. Lasslop, S. Hantson, S. P. Harrison, D. Bachelet, C. Burton, M. Forkel, et al., Global ecosystems and fire: Multi-model assessment of fire-induced tree-cover and carbon storage reduction. *Glob Change Biol* 26, 5027–5041 (2020).
4. W. M. Jolly, M. A. Cochrane, P. H. Freeborn, Z. A. Holden, T. J. Brown, G. J. Williamson, et al., Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat Commun* 6, 7537 (2015).
5. J. T. Abatzoglou, A. P. Williams, Impact of anthropogenic climate change on wildfire across western US forests. *Proc Natl Acad Sci USA* 113, 11770–11775 (2016).
6. M. W. Jones, J. T. Abatzoglou, S. Veraverbeke, N. Andela, G. Lasslop, M. Forkel, et al., Global and Regional Trends and Drivers of Fire Under Climate Change. *Reviews of Geophysics* 60, e2020RG000726 (2022).
7. United Nations Environment Programme, “Spreading like Wildfire – The Rising Threat of Extraordinary Landscape Fires. A UNEP Rapid Response Assessment.” (Nairobi, Kenya, 2022); <https://www.unep.org/resources/report/spreading-wildfire-rising-threat-extraordinary-landscape-fires>.
8. B. Zheng, P. Ciais, F. Chevallier, E. Chuvieco, Y. Chen, H. Yang, Increasing forest fire emissions despite the decline in global burned area. *Science Advances* 7, eabh2646 (2021).
9. J. G. Canadell, C. P. (Mick) Meyer, G. D. Cook, A. Dowdy, P. R. Briggs, J. Knauer, et al., Multi-decadal increase of forest burned area in Australia is linked to climate change. *Nat Commun* 12, 6921 (2021).
10. A. P. Williams, J. T. Abatzoglou, A. Gershunov, J. Guzman-Morales, D. A. Bishop, J. K. Balch, et al., Observed Impacts of Anthropogenic Climate Change on Wildfire in California. *Earth’s Future* 7, 892–910 (2019).
11. E. Ponomarev, N. Yakimov, T. Ponomareva, O. Yakubailik, S. G. Conard, Current Trend of Carbon Emissions from Wildfires in Siberia. *Atmosphere* 12, 559 (2021).
12. S. A. Parks, J. T. Abatzoglou, Warmer and Drier Fire Seasons Contribute to Increases in Area Burned at High Severity in Western US Forests From 1985 to 2017. *Geophys. Res. Lett.* 47 (2020).
13. L. V. Gatti, L. S. Basso, J. B. Miller, M. Gloor, L. Gatti Domingues, H. L. G. Cassol, et al., Amazonia as a carbon source linked to deforestation and climate change. *Nature* 595, 388–393 (2021).
14. J. R. Marlon, P. J. Bartlein, D. G. Gavin, C. J. Long, R. S. Anderson, C. E. Briles, et al., Long-term perspective on wildfires in the western USA. *Proceedings of the National Academy of Sciences* 109, E535–E543 (2012).
15. N. Andela, D. C. Morton, L. Giglio, Y. Chen, G. R. van der Werf, P. S. Kasibhatla, et al., A human-driven decline in global burned area. *Science* 356, 1356–1362 (2017).
16. B. Zheng, P. Ciais, F. Chevallier, H. Yang, J. G. Canadell, Y. Chen, et al., Record-high CO₂ emissions from boreal fires in 2021. *Science* 379, 912–917 (2023).
17. I. R. van der Velde, G. R. van der Werf, S. Houweling, J. D. Maasakkers, T. Borsdorff, J. Landgraf, et al., Vast CO₂ release from Australian fires in 2019–2020 constrained by satellite. *Nature* 597, 366–369 (2021).
18. P. E. Higuera, J. T. Abatzoglou, Record-setting climate enabled the extraordinary 2020 fire season in the western United States. *Glob. Change Biol.* 27, 1–2 (2021).
19. M. W. Jones, D. I. Kelley, C. A. Burton, F. Di Giuseppe, M. L. F. Barbosa, E. Brambleby, et al., State of Wildfires 2023–2024. *Earth System Science Data* 16, 3601–3685 (2024).
20. M. Ward, A. I. T. Tulloch, J. Q. Radford, B. A. Williams, A. E. Reside, S. L. Macdonald, et al., Impact of 2019–2020 mega-fires on Australian fauna habitat. *Nat Ecol Evol* 4, 1321–1326 (2020).
21. D. B. Lindenmayer, C. Taylor, New spatial analyses of Australian wildfires highlight the need for new fire, resource, and conservation policies. *Proc Natl Acad Sci USA* 117, 12481–12485 (2020).
22. C. Yue, P. Ciais, D. Zhu, T. Wang, S. S. Peng, S. L. Piao, How have past fire disturbances contributed to the current carbon balance of boreal ecosystems? *Biogeosciences* 13, 675–690 (2016).
23. S. P. Harrison, P. J. Bartlein, V. Brovkin, S. Houweling, S. Kloster, I. C. Prentice, The biomass burning contribution to climate–carbon-cycle feedback. *Earth Syst. Dynam.* 9, 663–677 (2018).

24. D. M. J. S. Bowman, G. J. Williamson, O. F. Price, M. N. Ndalila, R. A. Bradstock, Australian forests, megafires and the risk of dwindling carbon stocks. *Plant, Cell & Environment* 44, 347–355 (2021).
25. C. A. Phillips, B. M. Rogers, M. Elder, S. Cooperdock, M. Moubarak, J. T. Randerson, et al., Escalating carbon emissions from North American boreal forest wildfires and the climate mitigation potential of fire management. *Science Advances* 8, eabl7161 (2022).
26. D. M. J. S. Bowman, G. J. Williamson, J. T. Abatzoglou, C. A. Kolden, M. A. Cochrane, A. M. S. Smith, et al., Human exposure and sensitivity to globally extreme wildfire events. *Nat Ecol Evol* 1, 0058 (2017).
27. D. Wang, D. Guan, S. Zhu, M. M. Kinnon, G. Geng, Q. Zhang, et al., Economic footprint of California wildfires in 2018. *Nat Sustain* 4, 252–260 (2021).
28. F. H. Johnston, N. Borchers-Arriagada, G. G. Morgan, B. Jalaludin, A. J. Palmer, G. J. Williamson, et al., Unprecedented health costs of smoke-related PM_{2.5} from the 2019–20 Australian megafires. *Nat Sustain* 4, 42–47 (2021).
29. H. Clarke, R. H. Nolan, V. R. De Dios, R. Bradstock, A. Griebel, S. Khanal, et al., Forest fire threatens global carbon sinks and population centres under rising atmospheric water demand. *Nat Commun* 13, 7161 (2022).
30. World Bank, “World Bank Policy Note: Managing Wildfires in a Changing Climate” (Washington DC, 2020); https://www.profor.info/sites/profor.info/files/PROFOR_ManagingWildfires_2020_final.pdf
31. J. T. Abatzoglou, A. P. Williams, R. Barbero, Global Emergence of Anthropogenic Climate Change in Fire Weather Indices. *Geophys. Res. Lett.* 46, 326–336 (2019).
32. J. T. Abatzoglou, C. A. Kolden, J. K. Balch, B. A. Bradley, Controls on interannual variability in lightning-caused fire activity in the western US. *Environ. Res. Lett.* 11, 045005 (2016).
33. S. Veraverbeke, B. M. Rogers, M. L. Goulden, R. R. Jandt, C. E. Miller, E. B. Wiggins, et al., Lightning as a major driver of recent large fire years in North American boreal forests. *Nature Clim Change* 7, 529–534 (2017).
34. T. A. J. Janssen, M. W. Jones, D. Finney, G. R. van der Werf, D. van Wees, W. Xu, et al., Extratropical forests increasingly at risk due to lightning fires. *Nat. Geosci.* 16, 1136–1144 (2023).
35. G. A. Mills, W. L. McCaw, Centre for Australian Weather and Climate Research, Atmospheric Stability Environments and Fire Weather in Australia: Extending the Haines Index (Centre for Australian Weather and Climate Research, Melbourne, 2010).
36. G. Di Virgilio, J. P. Evans, S. A. P. Blake, M. Armstrong, A. J. Dowdy, J. Sharples, et al., Climate Change Increases the Potential for Extreme Wildfires. *Geophys. Res. Lett.* 46, 8517–8526 (2019).
37. G. J. van Oldenborgh, F. Krikken, S. Lewis, N. J. Leach, F. Lehner, K. R. Saunders, et al., Attribution of the Australian bushfire risk to anthropogenic climate change. *Natural Hazards and Earth System Sciences* 21, 941–960 (2021).
38. M. C. Kirchmeier-Young, F. W. Zwiers, N. P. Gillett, A. J. Cannon, Attributing extreme fire risk in Western Canada to human emissions. *Climatic Change* 144, 365–379 (2017).
39. J. G. Pausas, J. E. Keeley, Wildfires and global change. *Frontiers in Ecology and the Environment* 19, 387–395 (2021).
40. D. I. Kelley, I. Bistinas, R. Whitley, C. Burton, T. R. Marthews, N. Dong, How contemporary bioclimatic and human controls change global fire regimes. *Nat. Clim. Chang.* 9, 690–696 (2019).
41. D. M. J. S. Bowman, C. A. Kolden, J. T. Abatzoglou, F. H. Johnston, G. R. van der Werf, M. Flannigan, Vegetation fires in the Anthropocene. *Nat Rev Earth Environ* 1, 500–515 (2020).
42. N. J. Abram, B. J. Henley, A. Sen Gupta, T. J. R. Lippmann, H. Clarke, A. J. Dowdy, et al., Connections of climate change and variability to large and extreme forest fires in southeast Australia. *Commun Earth Environ* 2, 8 (2021).
43. L. E. O. C. Aragão, L. O. Anderson, M. G. Fonseca, T. M. Rosan, L. B. Vedovato, F. H. Wagner, et al., 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nat Commun* 9, 536 (2018).
44. J. K. Balch, B. A. Bradley, J. T. Abatzoglou, R. C. Nagy, E. J. Fusco, A. L. Mahood, Human-started wildfires expand the fire niche across the United States. *Proc Natl Acad Sci USA* 114, 2946–2951 (2017).

45. P. M. Fernandes, A. P. Pacheco, R. Almeida, J. Claro, The role of fire-suppression force in limiting the spread of extremely large forest fires in Portugal. *Eur J Forest Res* 135, 253–262 (2016).
46. F. Moreira, D. Ascoli, H. Safford, M. A. Adams, J. M. Moreno, J. M. C. Pereira, et al., Wildfire management in Mediterranean-type regions: paradigm change needed. *Environ. Res. Lett.* 15, 011001 (2020).
47. R. A. Bradstock, M. M. Boer, G. J. Cary, O. F. Price, R. J. Williams, D. Barrett, et al., Modelling the potential for prescribed burning to mitigate carbon emissions from wildfires in fire-prone forests of Australia. *Int. J. Wildland Fire* 21, 629–639 (2012).
48. C. H. L. Silva Junior, L. E. O. C. Aragão, L. O. Anderson, M. G. Fonseca, Y. E. Shimabukuro, C. Vancutsem, et al., Persistent collapse of biomass in Amazonian forest edges following deforestation leads to unaccounted carbon losses. *Sci. Adv.* 6, eaaz8360 (2020).
49. T. M. Rosan, S. Sitch, L. M. Mercado, V. Heinrich, P. Friedlingstein, L. E. O. C. Aragão, Fragmentation-Driven Divergent Trends in Burned Area in Amazonia and Cerrado. *Frontiers in Forests and Global Change* 5 (2022).
50. M.-A. Parisien, Q. E. Barber, K. G. Hirsch, C. A. Stockdale, S. Erni, X. Wang, et al., Fire deficit increases wildfire risk for many communities in the Canadian boreal forest. *Nat Commun* 11, 2121 (2020).
51. B. M. Rogers, A. J. Soja, M. L. Goulden, J. T. Randerson, Influence of tree species on continental differences in boreal fires and climate feedbacks. *Nature Geosci* 8, 228–234 (2015).
52. M. L. Pettinari, E. Chuvieco, Generation of a global fuel data set using the Fuel Characteristic Classification System. *Biogeosciences* 13, 2061–2076 (2016).
53. M. Forkel, N. Andela, S. P. Harrison, G. Lasslop, M. van Marle, E. Chuvieco, et al., Emergent relationships with respect to burned area in global satellite observations and fire-enabled vegetation models. *Biogeosciences* 16, 57–76 (2019).
54. J. T. Abatzoglou, A. P. Williams, L. Boschetti, M. Zubkova, C. A. Kolden, Global patterns of interannual climate–fire relationships. *Glob Change Biol* 24, 5164–5175 (2018).
55. M. Forkel, W. Dorigo, G. Lasslop, I. Teubner, E. Chuvieco, K. Thonicke, et al., A data-driven approach to identify controls on global fire activity from satellite and climate observations (SOFIA V1). *Geosci. Model Dev.* 10, 4443–4476 (2017).
56. E. Chuvieco, M. L. Pettinari, N. Koutsias, M. Forkel, S. Hantson, M. Turco, et al., Human and climate drivers of global biomass burning variability. *Science of The Total Environment* 779, 146361 (2021).
57. D. M. Olson, E. Dinerstein, E. D. Wikramanayake, N. D. Burgess, G. V. N. Powell, E. C. Underwood, et al., Terrestrial Ecoregions of the World: A New Map of Life on Earth. *BioScience* 51, 933 (2001).
58. L. Giglio, L. Boschetti, D. P. Roy, M. L. Humber, C. O. Justice, The Collection 6 MODIS burned area mapping algorithm and product. *Remote Sensing of Environment* 217, 72–85 (2018).
59. D. van Wees, G. R. van der Werf, J. T. Randerson, B. M. Rogers, Y. Chen, S. Veraverbeke, et al., Global biomass burning fuel consumption and emissions at 500 m spatial resolution based on the Global Fire Emissions Database (GFED). *Geoscientific Model Development* 15, 8411–8437 (2022).
60. C. E. Van Wagner, Development and Structure of the Canadian Forest Fire Weather Index System, Forestry Technical Report 35 (Canadian Forestry Service, Ottawa, 1987; <http://cfs.nrcan.gc.ca/publications?id=19927>)vol. 35.
61. C. Vitolo, F. Di Giuseppe, C. Barnard, R. Coughlan, J. San-Miguel-Ayanz, G. Libertá, et al., ERA5-based global meteorological wildfire danger maps. *Sci Data* 7, 216 (2020).
62. J. Muñoz-Sabater, E. Dutra, A. Agustí-Panareda, C. Albergel, G. Arduini, G. Balsamo, et al., ERA5-Land: a state-of-the-art global reanalysis dataset for land applications. *Earth System Science Data* 13, 4349–4383 (2021).
63. R. H. Holzworth, J. B. Brundell, M. P. McCarthy, A. R. Jacobson, C. J. Rodger, T. S. Anderson, et al., Lightning in the Arctic. *Geophysical Research Letters* 48, e2020GL091366 (2021).
64. A. Huete, K. Didan, T. Miura, E. P. Rodriguez, X. Gao, L. G. Ferreira, Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment* 83, 195–213 (2002).
65. Didan, Kamel, MODIS/Terra Vegetation Indices Monthly L3 Global 1km SIN Grid V061, NASA EOSDIS Land Processes DAAC (2021); <https://doi.org/10.5067/MODIS/MOD13A3.061>.

66. J. E. Dobson, E. A. Bright, P. R. Coleman, R. C. Durfee, B. A. Worley, LandScan: a global population database for estimating populations at risk. *Photogrammetric engineering and remote sensing* 66, 849–857 (2000).
67. Friedl, Mark, Sulla-Menashe, Damien, MCD12Q1 MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 500m SIN Grid V006, NASA EOSDIS Land Processes DAAC (2019); <https://doi.org/10.5067/MODIS/MCD12Q1.006>.
68. T. P. Robinson, G. R. W. Wint, G. Conchedda, T. P. V. Boeckel, V. Ercoli, E. Palamara, et al., Mapping the Global Distribution of Livestock. *PLOS ONE* 9, e96084 (2014).
69. J. R. Meijer, M. A. J. Huijbregts, K. C. G. J. Schotten, A. M. Schipper, Global patterns of current and future road infrastructure. *Environ. Res. Lett.* 13, 064006 (2018).
70. G. Amatulli, D. McInerney, T. Sethi, P. Strobl, S. Domisch, Geomorpho90m, empirical evaluation and accuracy assessment of global high-resolution geomorphometric layers. *Sci Data* 7, 162 (2020).
71. G. Di Virgilio, J. P. Evans, S. A. P. Blake, M. Armstrong, A. J. Dowdy, J. Sharples, et al., Climate Change Increases the Potential for Extreme Wildfires. *Geophys. Res. Lett.* 46, 8517–8526 (2019).
72. S. Archibald, C. E. R. Lehmann, J. L. Gomez-Dans, R. A. Bradstock, Defining pyromes and global syndromes of fire regimes. *Proceedings of the National Academy of Sciences* 110, 6442–6447 (2013).
73. M. E. Cattau, A. L. Mahood, J. K. Balch, C. A. Wessman, Modern Pyromes: Biogeographical Patterns of Fire Characteristics across the Contiguous United States. *Fire* 5, 95 (2022).
74. M. García, M. L. Pettinari, E. Chuvieco, J. Salas, F. Mouillot, W. Chen, et al., Characterizing Global Fire Regimes from Satellite-Derived Products. *Forests* 13, 699 (2022).
75. C. X. Cunningham, G. J. Williamson, R. H. Nolan, L. Teckentrup, M. M. Boer, D. M. J. S. Bowman, et al., Pyrogeography in flux: Reorganization of Australian fire regimes in a hotter world. *Global Change Biology* 30, e17130 (2024).
76. M. Forkel, K. Thonicke, C. Beer, W. Cramer, S. Bartalev, C. Schmullius, Extreme fire events are related to previous-year surface moisture conditions in permafrost-underlain larch forests of Siberia. *Environ. Res. Lett.* 7, 044021 (2012).
77. V. I. Kharuk, M. L. Dvinskaya, S. T. Im, A. S. Golyukov, K. T. Smith, Wildfires in the Siberian Arctic. *Fire* 5, 106 (2022).
78. P. M. Brando, L. Paolucci, C. C. Ummenhofer, E. M. Ordway, H. Hartmann, M. E. Cattau, et al., Droughts, Wildfires, and Forest Carbon Cycling: A Pantropical Synthesis. *Annual Review of Earth and Planetary Sciences* 47, 555–581 (2019).
79. T. Nikonovas, A. Spessa, S. H. Doerr, G. D. Clay, S. Mezbahuddin, Near-complete loss of fire-resistant primary tropical forest cover in Sumatra and Kalimantan. *Commun Earth Environ* 1, 1–8 (2020).
80. J. Barlow, E. Berenguer, R. Carmenta, F. França, Clarifying Amazonia’s burning crisis. *Global Change Biology* 26, 319–321 (2020).
81. V. R. Pivello, The Use of Fire in the Cerrado and Amazonian Rainforests of Brazil: Past and Present. *fire ecol* 7, 24–39 (2011).
82. R. D. Field, G. R. van der Werf, S. S. P. Shen, Human amplification of drought-induced biomass burning in Indonesia since 1960. *Nature Geosci* 2, 185–188 (2009).
83. P. G. Curtis, C. M. Slay, N. L. Harris, A. Tyukavina, M. C. Hansen, Classifying drivers of global forest loss. *Science* 361, 1108–1111 (2018).
84. M. R. Rosa, P. H. S. Brancalion, R. Crouzeilles, L. R. Tambosi, P. R. Piffer, F. E. B. Lenti, et al., Hidden destruction of older forests threatens Brazil’s Atlantic Forest and challenges restoration programs. *Science Advances* 7, eabc4547 (2021).
85. J. C. Aleman, M. A. Jarzyna, A. C. Staver, Forest extent and deforestation in tropical Africa since 1900. *Nat Ecol Evol* 2, 26–33 (2018).
86. Y. Le Page, D. Oom, J. M. N. Silva, P. Jönsson, J. M. C. Pereira, Seasonality of vegetation fires as modified by human action: observing the deviation from eco-climatic fire regimes. *Global Ecology and Biogeography* 19, 575–588 (2010).
87. M. Castellnou, N. Prat-Guitart, E. Arilla, A. Larrañaga, E. Nebot, X. Castellarnau, et al., Empowering strategic decision-making for wildfire management: avoiding the fear trap and creating a resilient landscape. *fire ecol* 15, 31 (2019).
88. H. Hessel, Wildland Fire Prevention: a Review. *Curr Forestry Rep* 4, 178–190 (2018).

89. J. T. Abatzoglou, D. S. Battisti, A. P. Williams, W. D. Hansen, B. J. Harvey, C. A. Kolden, Projected increases in western US forest fire despite growing fuel constraints. *Commun Earth Environ* 2, 1–8 (2021).
90. H. Clarke, R. Gibson, B. Cirulis, R. A. Bradstock, T. D. Penman, Developing and testing models of the drivers of anthropogenic and lightning-caused wildfire ignitions in south-eastern Australia. *Journal of Environmental Management* 235, 34–41 (2019).
91. R. A. Bradstock, A biogeographic model of fire regimes in Australia: current and future implications: A biogeographic model of fire in Australia. *Global Ecology and Biogeography* 19, 145–158 (2010).
92. R. Rotbarth, E. H. Van Nes, M. Scheffer, J. U. Jepsen, O. P. L. Vindstad, C. Xu, et al., Northern expansion is not compensating for southern declines in North American boreal forests. *Nat Commun* 14, 3373 (2023).
93. L. T. Berner, S. J. Goetz, Satellite observations document trends consistent with a boreal forest biome shift. *Global Change Biology* 28, 3275–3292 (2022).
94. H. Yang, P. Ciais, F. Frappart, X. Li, M. Brandt, R. Fensholt, et al., Global increase in biomass carbon stock dominated by growth of northern young forests over past decade. *Nat. Geosci.* 16, 886–892 (2023).
95. Y. Li, D. Sulla-Menashe, S. Motesharrei, X.-P. Song, E. Kalnay, Q. Ying, et al., Inconsistent estimates of forest cover change in China between 2000 and 2013 from multiple datasets: differences in parameters, spatial resolution, and definitions. *Sci Rep* 7, 8748 (2017).
96. H. Yin, A. Khamzina, D. Pflugmacher, C. Martius, Forest cover mapping in post-Soviet Central Asia using multi-resolution remote sensing imagery. *Sci Rep* 7, 1375 (2017).
97. X. Wei, Y. Liu, L. Qi, J. Chen, G. Wang, L. Zhang, et al., Monitoring forest dynamics in Africa during 2000–2020 using a remotely sensed fractional tree cover dataset. *International Journal of Digital Earth* 16, 2212–2232 (2023).
98. S. Piao, X. Wang, T. Park, C. Chen, X. Lian, Y. He, et al., Characteristics, drivers and feedbacks of global greening. *Nat Rev Earth Environ* 1, 14–27 (2020).
99. Y. He, Y. Liu, L. Lei, C. Terrer, C. Huntingford, J. Peñuelas, et al., CO₂ fertilization contributed more than half of the observed forest biomass increase in northern extra-tropical land. *Global Change Biology* 29, 4313–4326 (2023).
100. S. L. Lewis, D. P. Edwards, D. Galbraith, Increasing human dominance of tropical forests. *Science* 349, 827–832 (2015).
101. S. Archibald, Managing the human component of fire regimes: lessons from Africa. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371, 20150346 (2016).
102. Q. Xu, A. L. Westerling, A. Notohamiprodjo, C. Wiedinmyer, J. J. Picotte, S. A. Parks, et al., Wildfire burn severity and emissions inventory: an example implementation over California. *Environ. Res. Lett.* 17, 085008 (2022).
103. E. A. Kukavskaya, E. G. Shvetsov, L. V. Buryak, P. D. Tretyakov, P. Y. Groisman, Increasing Fuel Loads, Fire Hazard, and Carbon Emissions from Fires in Central Siberia. *Fire* 6, 63 (2023).
104. R. C. Scholten, D. Coumou, F. Luo, S. Veraverbeke, Early snowmelt and polar jet dynamics co-influence recent extreme Siberian fire seasons. *Science* 378, 1005–1009 (2022).
105. E. I. Ponomarev, A. N. Zabrodin, E. G. Shvetsov, T. V. Ponomareva, Wildfire Intensity and Fire Emissions in Siberia. *Fire* 6, 246 (2023).
106. T. D. Hessilt, J. T. Abatzoglou, Y. Chen, J. T. Randerson, R. C. Scholten, G. van der Werf, et al., Future increases in lightning ignition efficiency and wildfire occurrence expected from drier fuels in boreal forest ecosystems of western North America. *Environ. Res. Lett.* 17, 054008 (2022).
107. J. K. Shuman, J. K. Balch, R. T. Barnes, P. E. Higuera, C. I. Roos, D. W. Schwilk, et al., Reimagine fire science for the anthropocene. *PNAS Nexus* 1, pgac115 (2022).
108. M. A. Cochrane, D. M. J. S. Bowman, Manage fire regimes, not fires. *Nat. Geosci.* 14, 455–457 (2021).
109. S. Bloem, A. C. Cullen, L. O. Mearns, J. T. Abatzoglou, The Role of International Resource Sharing Arrangements in Managing Fire in the Face of Climate Change. *Fire* 5, 88 (2022).
110. C. F. Starrs, V. Butsic, C. Stephens, W. Stewart, The impact of land ownership, firefighting, and reserve status on fire probability in California. *Environ. Res. Lett.* 13, 034025 (2018).

111. A. C. M. Pessôa, T. F. Morello R.S., C. H. L. Silva-Junior, J. Doblas, N. S. Carvalho, L. E. O. C. Aragão, et al., Protected areas are effective on curbing fires in the Amazon. *Ecological Economics* 214, 107983 (2023).
112. C. C. Hanes, X. Wang, P. Jain, M.-A. Parisien, J. M. Little, M. D. Flannigan, Fire-regime changes in Canada over the last half century. *Can. J. For. Res.* 49, 256–269 (2019).
113. P. Friedlingstein, M. O’Sullivan, M. W. Jones, R. M. Andrew, D. C. E. Bakker, J. Hauck, et al., Global Carbon Budget 2023. *Earth System Science Data* 15, 5301–5369 (2023).
114. S. Sitch, C. Huntingford, N. Gedney, P. E. Levy, M. Lomas, S. L. Piao, et al., Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs): UNCERTAINTY IN LAND CARBON CYCLE FEEDBACKS. *Global Change Biology* 14, 2015–2039 (2008).
115. C.-E. Yang, J. Mao, F. M. Hoffman, D. M. Ricciuto, J. S. Fu, C. D. Jones, et al., Uncertainty Quantification of Extratropical Forest Biomass in CMIP5 Models over the Northern Hemisphere. *Sci Rep* 8, 10962 (2018).
116. S. Kloster, G. Lasslop, Historical and future fire occurrence (1850 to 2100) simulated in CMIP5 Earth System Models. *Global and Planetary Change* 150, 58–69 (2017).
117. M. C. Mack, X. J. Walker, J. F. Johnstone, H. D. Alexander, A. M. Melvin, M. Jean, et al., Carbon loss from boreal forest wildfires offset by increased dominance of deciduous trees. *Science* 372, 280–283 (2021).
118. J. L. Baltzer, N. J. Day, X. J. Walker, D. Greene, M. C. Mack, H. D. Alexander, et al., Increasing fire and the decline of fire adapted black spruce in the boreal forest. *Proceedings of the National Academy of Sciences* 118, e2024872118 (2021).
119. M. S. Balshi, A. D. McGuire, P. Duffy, M. Flannigan, D. W. Kicklighter, J. Melillo, Vulnerability of carbon storage in North American boreal forests to wildfires during the 21st century. *Global Change Biology* 15, 1491–1510 (2009).
120. S. Schaphoff, C. P. O. Reyer, D. Schepaschenko, D. Gerten, A. Shvidenko, et al., Tamm Review: Observed and projected climate change impacts on Russia’s forests and its carbon balance. *Forest Ecology and Management* 361, 432–444 (2016).
121. M. R. Turetsky, E. S. Kane, J. W. Harden, R. D. Ottmar, K. L. Manies, E. Hoy, et al., Recent acceleration of biomass burning and carbon losses in Alaskan forests and peatlands. *Nature Geosci* 4, 27–31 (2011).
122. J. MacCarthy, A. Tyukavina, M. J. Weisse, N. Harris, E. Glen, Extreme wildfires in Canada and their contribution to global loss in tree cover and carbon emissions in 2023. *Global Change Biology* 30, e17392 (2024).
123. Environment and Climate Change Canada, “National Inventory Report, 1990–2022: Greenhouse Gas Sources and Sinks in Canada, available at: publications.gc.ca/pub?id=9.506002&sl=0, last access: 15 August 2024.” (2024).
124. B. W. Griscom, J. Adams, P. W. Ellis, R. A. Houghton, G. Lomax, D. A. Miteva, et al., Natural climate solutions. *Proceedings of the National Academy of Sciences* 114, 11645–11650 (2017).
125. H. B. Smith, N. E. Vaughan, J. Forster, Long-term national climate strategies bet on forests and soils to reach net-zero. *Commun Earth Environ* 3, 1–12 (2022).
126. W. S. Walker, S. R. Gorelik, S. C. Cook-Patton, A. Baccini, M. K. Farina, K. K. Solvik, et al., The global potential for increased storage of carbon on land. *Proceedings of the National Academy of Sciences* 119, e2111312119 (2022).
127. B. K. Haya, S. Evans, L. Brown, J. Bukoski, V. Butsic, B. Cabiyo, et al., Comprehensive review of carbon quantification by improved forest management offset protocols. *Frontiers in Forests and Global Change* 6 (2023).
128. IPCC, Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, et al.,

- (Eds.]). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, Pp. 3–32, Doi:10.1017/9781009157896.001 (2021).
129. S. Smith, O. Geden, G. Nemet, M. Gidden, W. Lamb, C. Powis, et al., State of Carbon Dioxide Removal - 1st Edition. doi: 10.17605/OSF.IO/W3B4Z (2023).
130. J. Barlow, L. Parry, T. A. Gardner, J. Ferreira, L. E. O. C. Aragão, R. Carmenta, et al., The critical importance of considering fire in REDD+ programs. *Biological Conservation* 154, 1–8 (2012).
131. OECD, Taming Wildfires in the Context of Climate Change (OECD, 2023; https://www.oecd-ilibrary.org/environment/taming-wildfires-in-the-context-of-climate-change_dd00c367-en).
132. M. W. Jones, Supplementary Data: Global rise in forest fire emissions linked to climate change in the extratropics, Zenodo (2024); <https://doi.org/10.5281/zenodo.10036942>.
133. D. van Wees, G. R. van der Werf, J. T. Randerson, B. M. Rogers, Y. Chen, S. Veraverbeke, L. Giglio, D. C. Morton, Model data for “Global biomass burning fuel consumption and emissions at 500-m spatial

- 1 resolution based on the Global Fire Emissions Database (GFED),” version v2, Zenodo (2024);
- 2 <https://doi.org/10.5281/zenodo.12670427>.
- 3 134. G. R. van der Werf, J. T. Randerson, L. Giglio, T. T. van Leeuwen, Y. Chen, B. M. Rogers, et al., Global
- 4 fire emissions estimates during 1997–2016. *Earth Syst. Sci. Data* 9, 697–720 (2017).
- 5 135. C. DiMiceli, J. Townshend, M. Carroll, R. Sohlberg, Evolution of the representation of global vegetation
- 6 by vegetation continuous fields. *Remote Sensing of Environment* 254, 112271 (2021).
- 7 136. DiMiceli, Charlene, Carroll, Mark, Sohlberg, Robert, Kim, Do-Hyung, Kelly, Maggi, Townshend, John,
- 8 MOD44B MODIS/Terra Vegetation Continuous Fields Yearly L3 Global 250m SIN Grid V006, NASA
- 9 EOSDIS Land Processes DAAC (2015); <https://doi.org/10.5067/MODIS/MOD44B.006>.
- 10 137. T. Majasalmi, M. Rautiainen, Representation of tree cover in global land cover products: Finland as a
- 11 case study area. *Environ Monit Assess* 193, 121 (2021).
- 12 138. P. Potapov, M. C. Hansen, A. Pickens, A. Hernandez-Serna, A. Tyukavina, S. Turubanova, et al., The
- 13 Global 2000-2020 Land Cover and Land Use Change Dataset Derived From the Landsat Archive: First
- 14 Results. *Frontiers in Remote Sensing* 3 (2022).
- 15 139. T. J. Hawbaker, M. K. Vanderhoof, Y.-J. Beal, J. D. Takacs, G. L. Schmidt, J. T. Falgout, et al., Mapping
- 16 burned areas using dense time-series of Landsat data. *Remote Sensing of Environment* 198, 504–522
- 17 (2017).
- 18 140. H. Tatli, M. Türkeş, Climatological evaluation of Haines forest fire weather index over the
- 19 Mediterranean Basin. *Meteorological Applications* 21, 545–552 (2014).
- 20 141. B. E. Potter, Quantitative Evaluation of the Haines Index’s Ability to Predict Fire Growth Events.
- 21 *Atmosphere* 9, 177 (2018).
- 22 142. M. N. Ndalila, G. J. Williamson, P. Fox-Hughes, J. Sharples, D. M. J. S. Bowman, Evolution of a
- 23 pyrocumulonimbus event associated with an extreme wildfire in Tasmania, Australia. *Natural Hazards*
- 24 *and Earth System Sciences* 20, 1497–1511 (2020).
- 25 143. H. Hersbach, B. Bell, P. Berrisford, S. Hirahara, A. Horányi, J. Muñoz-Sabater, et al., The ERA5 global
- 26 reanalysis. *Q.J.R. Meteorol. Soc.* 146, 1999–2049 (2020).
- 27 144. Y. Chen, D. M. Romps, J. T. Seeley, S. Veraverbeke, W. J. Riley, Z. A. Mekkonen, et al., Future
- 28 increases in Arctic lightning and fire risk for permafrost carbon. *Nat. Clim. Chang.* 11, 404–410 (2021).
- 29 145. WWLLN, Worldwide Lightning Location Network 1° Detection Efficiency Maps, available at:
- 30 <http://wwlln.net/deMaps/> (2023). <http://wwlln.net/deMaps/>.
- 31 146. M. Zubkova, L. Boschetti, J. T. Abatzoglou, L. Giglio, Changes in Fire Activity in Africa from 2002 to
- 32 2016 and Their Potential Drivers. *Geophysical Research Letters* 46, 7643–7653 (2019).
- 33 147. T. Kitzberger, T. W. Swetnam, T. T. Veblen, Inter-hemispheric synchrony of forest fires and the El
- 34 Niño-Southern Oscillation. *Global Ecology and Biogeography* 10, 315–326 (2001).
- 35 148. Y. Chen, D. C. Morton, N. Andela, G. R. van der Werf, L. Giglio, J. T. Randerson, A pan-tropical
- 36 cascade of fire driven by El Niño/Southern Oscillation. *Nature Clim Change* 7, 906–911 (2017).
- 37 149. R. de Jong, S. de Bruin, A. de Wit, M. E. Schaepman, D. L. Dent, Analysis of monotonic greening and
- 38 browning trends from global NDVI time-series. *Remote Sensing of Environment* 115, 692–702 (2011).
- 39 150. R. D. Ottmar, D. V. Sandberg, C. L. Riccardi, S. J. Prichard, An overview of the Fuel Characteristic
- 40 Classification System — Quantifying, classifying, and creating fuelbeds for resource planning. *Can. J.*
- 41 *For. Res.* 37, 2383–2393 (2007).
- 42 151. T. A. P. West, P. M. Fearnside, Brazil’s conservation reform and the reduction of deforestation in
- 43 Amazonia. *Land Use Policy* 100, 105072 (2021).
- 44 152. S. Pais, N. Aquilué, J. Campos, Â. Sil, B. Marcos, F. Martínez-Freiría, et al., Mountain farmland
- 45 protection and fire-smart management jointly reduce fire hazard and enhance biodiversity and carbon
- 46 sequestration. *Ecosystem Services* 44, 101143 (2020).
- 47 153. D. M. J. S. Bowman, J. Balch, P. Artaxo, W. J. Bond, M. A. Cochrane, C. M. D’Antonio, et al., The
- 48 human dimension of fire regimes on Earth: The human dimension of fire regimes on Earth. *Journal of*
- 49 *Biogeography* 38, 2223–2236 (2011).
- 50 154. M. Gilbert, G. Nicolas, G. Cinardi, T. P. Van Boeckel, S. O. Vanwambeke, G. R. W. Wint, et al., Global
- 51 distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010. *Sci Data*
- 52 5, 180227 (2018).
- 53 155. European Commission Eurostat Portal, Livestock units (LSU); [https://ec.europa.eu/eurostat/statistics-](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Glossary:Livestock_unit_(LSU))
- 54 [explained/index.php?title=Glossary:Livestock_unit_\(LSU\)](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Glossary:Livestock_unit_(LSU)).

156. P. Vogt, K. Riitters, P. Rambaud, R. d'Annunzio, E. Lindquist, A. Pekkarinen, et al., GuidosToolbox Workbench: spatial analysis of raster maps for ecological applications. *Ecography* 2022, e05864 (2022).
157. European Commission. Joint Research Centre., FAO, State of the World's Forests: Forest Fragmentation. (Publications Office, LU, 2019; <https://data.europa.eu/doi/10.2760/145325>).
158. L. Teckentrup, S. P. Harrison, S. Hantson, A. Heil, J. R. Melton, M. Forrest, et al., Response of simulated burned area to historical changes in environmental and anthropogenic factors: a comparison of seven fire models. *Biogeosciences* 16, 3883–3910 (2019).
159. E. Doxsey-Whitfield, K. MacManus, S. B. Adamo, L. Pistolesi, J. Squires, O. Borkovska, et al., Taking Advantage of the Improved Availability of Census Data: A First Look at the Gridded Population of the World, Version 4. *Papers in Applied Geography* 1, 226–234 (2015).
160. D. Nepstad, G. Carvalho, A. C. Barros, A. Alencar, J. P. Capobianco, J. Bishop, et al., Road paving, fire regime feedbacks, and the future of Amazon forests. *Forest Ecology and Management*, 13 (2001).
161. F. X. Catry, F. C. Rego, F. L. Bação, F. Moreira, et al., Modeling and mapping wildfire ignition risk in Portugal. *Int. J. Wildland Fire* 18, 921–931 (2009).
162. G. Narayanaraj, M. C. Wimberly, Influences of forest roads on the spatial pattern of wildfire boundaries. *Int. J. Wildland Fire* 20, 792 (2011).
163. R. A. Bradstock, K. A. Hammill, L. Collins, O. Price, Effects of weather, fuel and terrain on fire severity in topographically diverse landscapes of south-eastern Australia. *Landscape Ecol* 25, 607–619 (2010).
164. G. J. Cary, R. E. Keane, R. H. Gardner, S. Lavorel, M. D. Flannigan, I. D. Davies, et al., Comparison of the Sensitivity of Landscape-fire-succession Models to Variation in Terrain, Fuel Pattern, Climate and Weather. *Landscape Ecol* 21, 121–137 (2006).
165. J. J. Sharples, R. H. D. McRae, S. R. Wilkes, Wind–terrain effects on the propagation of wildfires in rugged terrain: fire channelling. *Int. J. Wildland Fire* 21, 282–296 (2012).
166. R. Durão, C. Alonso, C. Gouveia, The Performance of ECMWF Ensemble Prediction System for European Extreme Fires: Portugal/Monchique in 2018. *Atmosphere* 13, 1239 (2022).
167. A. Huete, C. Justice, W. Van Leeuwen, MODIS Vegetation Index (MOD13) Algorithm Theoretical Basis Document Version 3; https://modis.gsfc.nasa.gov/data/atbd/atbd_mod13.pdf (1999).
168. M. L. Hutchins, R. H. Holzworth, C. J. Rodger, J. B. Brundell, Far-Field Power of Lightning Strokes as Measured by the World Wide Lightning Location Network. doi: 10.1175/JTECH-D-11-00174.1 (2012).
169. J. O. Kaplan, K. H.-K. Lau, The WGLC global gridded lightning climatology and time series. *Earth System Science Data* 13, 3219–3237 (2021).
170. M. Li, P. Wu, Z. Ma, A comprehensive evaluation of soil moisture and soil temperature from third-generation atmospheric and land reanalysis data sets. 40, 5744–5766 (2020).
171. A. Strahler, D. Muchoney, J. Borak, M. Friedl, S. Gopal, E. Lambin, et al., MODIS Land Cover Product (MCD12) Algorithm Theoretical Basis Document (ATBD) Version 5; https://lpdaac.usgs.gov/documents/437/MCD12_ATBD_V5.pdf (1999).
172. E. Chuvieco, F. Mouillot, G. R. van der Werf, J. San Miguel, M. Tanase, N. Koutsias, et al., Historical background and current developments for mapping burned area from satellite Earth observation. *Remote Sensing of Environment* 225, 45–64 (2019).
173. D. van Wees, G. R. van der Werf, J. T. Randerson, B. M. Rogers, Y. Chen, S. Veraverbeke, et al., Global biomass burning fuel consumption and emissions at 500 m spatial resolution based on the Global Fire Emissions Database (GFED). *Geoscientific Model Development* 15, 8411–8437 (2022).
174. G. R. van der Werf, J. T. Randerson, L. Giglio, G. J. Collatz, M. Mu, P. S. Kasibhatla, et al., Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmos. Chem. Phys.* 10, 11707–11735 (2010).
175. T. T. van Leeuwen, G. R. van der Werf, A. A. Hoffmann, R. G. Detmers, G. Rücker, N. H. F. French, et al., Biomass burning fuel consumption rates: a field measurement database. *Biogeosciences* 11, 7305–7329 (2014).
176. J. T. Randerson, Y. Chen, G. R. van der Werf, B. M. Rogers, D. C. Morton, Global burned area and biomass burning emissions from small fires: BURNED AREA FROM SMALL FIRES. *J. Geophys. Res.* 117 (2012).
177. D. van Wees, G. R. van der Werf, Modelling biomass burning emissions and the effect of spatial resolution: a case study for Africa based on the Global Fire Emissions Database (GFED). *Geoscientific Model Development* 12, 4681–4703 (2019).

178. S. Veraverbeke, B. M. Rogers, J. T. Randerson, Daily burned area and carbon emissions from boreal fires in Alaska. *Biogeosciences* 12, 3579–3601 (2015).
179. N. Andela, G. R. van der Werf, J. W. Kaiser, T. T. van Leeuwen, M. J. Wooster, C. E. R. Lehmann, et al., Biomass burning fuel consumption dynamics in the tropics and subtropics assessed from satellite. *Biogeosciences* 13, 3717–3734 (2016).
180. M. Charrad, N. Ghazzali, V. Boiteau, A. Niknafs, NbClust: An R Package for Determining the Relevant Number of Clusters in a Data Set. *Journal of Statistical Software* 61, 1–36 (2014).
181. J. Lizundia-Loiola, M. L. Pettinari, E. Chuvieco, Temporal Anomalies in Burned Area Trends: Satellite Estimations of the Amazonian 2019 Fire Crisis. *Remote Sensing* 12, 151 (2020).
182. M. Tabarelli, L. P. Pinto, J. M. C. Silva, M. Hirota, L. Bede, Challenges and Opportunities for Biodiversity Conservation in the Brazilian Atlantic Forest. *Conservation Biology* 19, 695–700 (2005).
- 183.

Acknowledgments: The authors thank Harry Smith (UEA) for his guidance on the commitment periods of forestry-related CDR. The authors thank the TRopical Ecosystems and Environmental Sciences (TREES) lab at the Brazilian Institute for Space Research (INPE) for their feedback on the inclusion of predictor variables relevant to tropical forest ecoregions.

Funding:

UK Natural Environment Research Council (NERC) grant NE/V01417X/1 (MWJ).

European Commission (EC) Horizon 2020 (H2020) project VERIFY grant 776810 (MWJ).

São Paulo Research Foundation (FAPESP) grants 2019/25701-8, 2020/15230-5 and 2023/03206-0 (GM).

EC H2020 project FirEURisk grant no. 101003890 (SHD, MLP).

NERC project UK-FDRS grant NE/T003553/1 (SHD).

European Space Agency (ESA) Climate Change Initiative (CCI) FireCCI project contract no. 4000126706/19/I-NB (MLP).

Royal Society grant RP\R1\191063 (CLQ).

National Science Foundation grant OAI-2019762 (JTA).

Author contributions:

Conceptualization: MWJ.

Methodology: MWJ, SV, JTA.

Resources/Software: NA, MLP, TR, DvW, GRvdW.

Investigation/Formal analysis: MWJ.

Visualization: MWJ, JTA.

Writing—original draft: MWJ.

Writing—review & editing: All co-authors.

Competing interests: The authors declare no competing interests.

Data and materials availability: Pyromes are provided in three geospatial formats via the Zenodo data repository (ref. (132)): shapefiles; 0.25° grids; and 0.05° grids. Gridded

correlations for all variables are also available via the Zenodo data repository (ref. (132)). R code used for clustering forest ecoregions into pyromes is also archived at the Zenodo repository (ref. (132)). The raw data representing burned area, carbon emissions, and all predictor variables in our analysis are publicly available via refs. (49, 58, 59, 61, 62, 67-70, 133), except for the lightning flash data from the WWLLN (ref. (63)), which are subject to a commercial agreement but can be provided in a gridded and coarsened form upon request.

Supplementary Materials

Materials and Methods

Supplementary Text 1

Supplementary Text 2

Figs. S1 to S22

Tables S1 to S2

References (134–182)